**University of São Paulo
“Luiz de Queiroz” College of Agriculture**

Regulation of the nitrate uptake process in grass species

Luís Henrique Damasceno Serezino

Thesis presented to obtain the degree of Doctor in Science.
Program: International Plant Cell and Molecular Biology

**Piracicaba
2020**

Luís Henrique Damasceno Serezino
Bachelor's in Biotechnology Engineering

Regulation of the nitrate uptake process in grass species

Advisor:

Prof. Dr. **ANTONIO VARGAS DE OLIVERA FIGUEIRA**

Thesis presented to obtain the degree of Doctor in Science.

Program: International Plant Cell and Molecular Biology

Piracicaba
2020

Dados Internacionais de Catalogação na Publicação
DIVISÃO DE BIBLIOTECA – DIBD/ESALQ/USP

Serezino, Luis Henrique Damasceno

Regulation of the nitrate uptake process in grass species / Luis Henrique Damasceno Serezino. - Piracicaba, 2020.

127 p.

Tese (Doutorado) - - USP / Escola Superior de Agricultura "Luiz de Queiroz".

1. Nitrato 2. Transporte HATS 3. Brachypodium 4. Cana-de-açúcar 5. *Saccharum* 6. NUE I. Título

Acknowledgments

First of all, I would like to thank God for giving me health, strength, and wisdom to complete another step of my life.

To Dr. Antonio Figueira for the opportunity to work in his laboratory, for believing and trusting in my work, and for the knowledge given in these over 9 years of partnership and friendship.

To the Lab Managers of the Plant Breeding Laboratory at Center for Nuclear Energy in Agriculture (CENA/USP): Felipe Campana, Wlamir Godoy, and Paulo Cassieri for their attention and help. Without this, experiments would not be possible. Thank you very much!

To my friends who lived with me during the Ph.D. at the Plant Breeding Laboratory: Albania José Patiño Torres, Alessandra Koltun, Anieli Baldo, Bruna Factor, Bruna Queiroz, Daniele Paschoal, Danielle Scotton, Eder Marques da Silva, Fernando Ferreira, Flávia Bento, Gabriel Ragazzo, Jamile Silva, Juliana Leles, Laecio Sampaio, Larissa Costa de Souza, Marília Morandi, Rafael Monteiro do Carmo, Rodolfo Augusto Maniero and Thaísa Pinheiro. It was a great experience to share moments and memories during this period with you. Especially, I would like to thank the undergrad students Gabriel Silvino, Hugo Corocher, and Vitor Ometto, for following me in the experiments, for their understanding and daily friendship. Thank you very much, LAMP!

To the professors and staff of the CENA/USP and the Luiz de Queiroz College of Agriculture (ESALQ/USP) for the support and contribution during this period.

I would like to express my gratitude to the Brazilian fellow scientists who somehow helped in this study: to Dr. Erika Jorge from Federal University of Minas Gerais (UFMG) for all the help with Western blot analysis; to Dr. Keila Duarte for helping me with the protein extraction and ELISA analysis.

I thank all the professors and staff members involved in assembling the Double Degree Ph.D. Program for making it happen. Dr. Helaine Carrer (USP), Megan Francis, and Nalini Kaul for the effortless help with my bureaucratic process during the exchange program.

I gratefully acknowledge all people involved in my internship abroad at Rutgers University: Dr. Eric Lam for accepting me as his Ph.D. student and as a friend, making easy the change of country. My LamLab mates and beloved friends: Buntora Pasaribu, Kenneth Acosta, Sarah Gilbert, and Shawn Sorrels. Thank you so much to accept me with your arms open. I hope to see you soon! And my friend Nimmi: you helped me at all times during the year that I was in New Brunswick. Thank you for the conversations, the laughter, the fellowship, thanks for everything!

I deeply thank the person that I've chosen to be by my side for my life: Natália, for all the unconditional love, for being my safety harbor and my best friend. For the partnership and

patience, for being with me in the tough moments and encouraging me to keep going, for all the love we shared all these years and for the future that we are building together. Love you!

To my beloved parents, Luís Henrique and Márcia, for all their love, kindness, attention, companionship, understanding, and for being who they are: my greatest examples! I love you so much! I hope one day to be able to give back everything you have given me. To my sisters, Bruna and Amanda, for being my eternal partners, for always being by my side. Love you! To all my family for their trust in me and their unconditional support. Thank you!

To my friends from São João da Boa Vista, Piracicaba, UNESP, and from Perdição. Thank you for all your support and friendship. Even though I was not present all the time, I always had you in my heart. I hope to have your friendship forever!

To Brazilian Federal Agency for Support and Evaluation of Graduate Education (CAPES) for granting the Ph.D. scholarship in the Brazil and in the USA (process numbers 88881.135748 / 2016-01).

Epigraphe

“Joy lies in the fight, in the attempt, in the suffering involved, not in the victory itself”

Mahatma Gandhi

TABLE OF CONTENTS

RESUMO	9
ABSTRACT	10
1. INTRODUCTION	11
1.1 N FERTILIZERS AND ENVIRONMENTAL IMPACTS	11
1.1.1 <i>Nitrogen as a threat to water resources</i>	12
1.1.2 <i>N fertilization as a threat to air and atmosphere quality</i>	13
1.1.3 <i>Nitrogen as a threat to soil property</i>	15
1.2 NITROGEN USE EFFICIENCY	16
1.3 NITRATE ACQUISITION	17
1.3.1 <i>Nitrate Transporter 1/Peptide Transporter (NRT1/NPF) members</i>	18
1.3.2 <i>Nitrate Transporter 2 (NRT2) family</i>	19
1.3.3 <i>Nitrate Transporter 3 (NRT3) family</i>	24
1.4 NITRATE UPTAKE REGULATION	26
1.5 CONCLUSION	30
REFERENCES.....	32
2. REGULATION OF NITRATE TRANSPORTERS (NRT2 AND NRT3) GENES EXPRESSION IN THE MODEL GRASS <i>BRACHYPODIUM DISTACHYON</i> UNDER VARIOUS N CONDITIONS	45
2.1 ABSTRACT	45
2.2 INTRODUCTION.....	46
2.3 METHODS	47
2.3.1 <i>Plant material and germination</i>	47
2.3.2 <i>Analysis of nitrogen uptake kinetics in roots by rapid ¹⁵N influx</i>	48
2.3.3 <i>Identification of preferential N source in <i>Brachypodium distachyon</i></i>	48
2.3.4 <i>Understanding the nitrate uptake process in roots upon distinct inorganic N source supply and N resupply</i>	48
2.3.5 <i>RNA extraction and cDNA synthesis</i>	49
2.3.6 <i>Quantitative reversed transcript amplification (RT-qPCR) of the <i>BdNRT2</i> and <i>BdNRT3</i> family members</i>	49
2.4 RESULTS	50
2.4.1 <i>Gene and protein analysis of the NRT2 and NRT3 subfamily in <i>Brachypodium distachyon</i></i>	50
2.4.2 <i><i>Brachypodium distachyon</i> prefers ammonium uptake rather than nitrate as N inorganic source</i>	57
2.4.3 <i>Nitrate uptake process is downregulated by ammonium in N-replete <i>Brachypodium distachyon</i> plants</i>	58
2.4.4 <i>Regulation of nitrate uptake in <i>Brachypodium distachyon</i> roots by N resupply</i> ...	66
2.5 DISCUSSION.....	74
2.6 CONCLUSION	78
REFERENCES.....	79
3. REGULATION OF NITRATE UPTAKE IN SUGARCANE (<i>SACCHARUM SPP.</i>) OCCURS IN POST-TRANSCRIPCIONAL AND POST-TRANSLATIONAL LEVELS AMONG GENOTYPES	85

3.1	ABSTRACT	85
3.2	INTRODUCTION.....	86
3.3	METHODS	88
3.3.1	<i>Plant material</i>	88
3.3.2	<i>Analysis of nitrate uptake kinetics in roots by rapid ¹⁵N influx</i>	89
3.3.3	<i>Understanding the nitrate uptake process in roots upon distinct inorganic N source supply and N resupply</i>	89
3.3.4	<i>RNA extraction and cDNA synthesis</i>	89
3.3.5	<i>Gene expression by quantitative reversed transcript amplification (RT-qPCR)</i> ...	90
3.3.6	<i>Protein extraction</i>	91
3.3.7	<i>NRT2.1 and NRT3.1 Immunodetection</i>	92
3.4	RESULTS	94
3.4.1	<i>Comparing the uptake response of sorghum and ‘SP80-3280’ to nitrate resupply</i> 94	
3.4.2	<i>Comparing sugarcane hybrids in response to nitrate resupply</i>	102
3.4.3	<i>ELISA and Western Blot analysis</i>	109
3.5	DISCUSSION.....	113
3.6	CONCLUSION	120
	REFERENCES.....	121

RESUMO

Regulação do processo de absorção de nitrato em espécies gramíneas

Devido às consequências ambientais e econômicas negativas do uso indiscriminado e excessivo de fertilizantes nitrogenados, há uma necessidade crescente de se desenvolver culturas mais eficientes para absorção de nitrogênio (N). Sendo o nitrato a principal fonte de N inorgânico disponível no solo, há um esforço notável para entender o processo de absorção de nitrato pelas plantas, um fator-chave na melhoria da eficiência no uso de N (NUE - *Nitrogen Use Efficiency*), e os fatores moleculares que a modulam. Assim, o conhecimento adquirido em plantas-modelo sobre os fatores moleculares e os mecanismos reguladores dos genes codificadores de transportadores de nitrato (NRTs) responsáveis pela absorção de nitrato poderia ser usado para melhorar a NUE das culturas. Surge, portanto, como um desafio para permitir que uma agricultura ambientalmente mais amigável que permita sustentar o crescimento da população humana e a consequente demanda por alimentos, uma vez que o uso intensivo de fertilizantes nitrogenados ameaça a sustentabilidade das culturas e a segurança alimentar. A caracterização fisiológica e molecular do processo de aquisição de nitrato em *Brachypodium distachyon*, uma planta modelo de gramíneas, poderia ajudar a melhorar as culturas não modelo, com maiores complexidades biológicas como a cana-de-açúcar. Análises de influxo de ^{15}N -amônio e ^{15}N -nitrato em condições de suficiência e deficiência em N revelaram que o amônio é a fonte inorgânica preferencial de N de *B. distachyon*. Foi observada uma regulação por *feedback* negativo do processo de absorção de nitrato exercido por amônio nas plantas sob tratamento de provisão e ressuplementação de N. Análises de expressão dos genes codificantes para os transportadores NRT2.1 e NRT2.2 em *Brachypodium* sugerem ambas proteínas como os principais transportadores NRT2 responsáveis pela aquisição de nitrato nesta espécie. Ambos os membros da família gênica *BdNRT3* foram coexpressos com os genes *BdNRT2.1* e *BdNRT2.2*. Assim, a análise do influxo de nitrogênio ^{15}N e o acúmulo de transcritos de membros do complexo de dois componentes (NRT2 / NRT3) após os tratamentos de provisão de cloreto de amônio e de ressuplementação de nitrato de amônio sugerem a ocorrência de uma regulação transcricional do HATS (*High Affinity Transport System*) em *B. distachyon*. Por outro lado, estudos utilizando sorgo e cana-de-açúcar revelaram uma regulação distinta do processo de absorção de nitrato na faixa HATS em ambas as espécies sob reabastecimento de nitrato. O influxo de nitrato marcado mostrou alta absorção de nitrato pela cultivar de sorgo 'BTX623' e pelo híbrido de cana-de-açúcar 'IAC87-3396'. Análises de expressão de genes que codificam NRT2 e NRT3 exibiram regulação transcricional distinta entre os genótipos da cana-de-açúcar. A falta de correlação entre o acúmulo de proteínas NRT2.1 e NRT3.1 com o influxo de ^{15}N -nitrato sugere a presença de mecanismo de controle do transporte de nitrato nas raízes de sorgo e cana-de-açúcar em nível pós-traducional. Essa presumível regulação alostérica do complexo HATS seria o principal mecanismo regulador da absorção de nitrato nas raízes de sorgo e cana-de-açúcar. Melhor compreensão dos mecanismos e moléculas envolvidas na regulação pós-traducional do processo de absorção de nitrato e, portanto, na via de sinalização desta fonte de N em gramíneas traria o desenvolvimento de culturas com maior NUE.

Palavras-chave: Nitrato, HATS, *Brachypodium*, Cana-de-açúcar, *Saccharum*, NUE

ABSTRACT

Regulation of the nitrate uptake process in grass species

Due to the negative environmental and economic consequences of excessive and indiscriminate use of N fertilizer, there is an increasing need for developing more efficient crops for N uptake. As nitrate is the main inorganic N source available in most soils, there has been a notable research effort to understand the nitrate uptake process, a key factor in the improvement of Nitrogen Use Efficiency (NUE), and the molecular factors that modulate it. Thus, the knowledge acquired in model plants about the molecular factors and regulatory mechanisms of genes encoding Nitrate Transporters (NRTs) responsible for nitrate uptake could be used to improve NUE in crops. The improvement of NUE is a challenge to allow environmentally friendly agriculture to support the growth of the human population and the consequent food demand, since the intensive use of N fertilizers threat both crop sustainability and food security. The physiological and molecular characterization of the nitrate acquisition process in *Brachypodium distachyon*, a grass model plant, could help to improve non-model crops with higher biological complexity such as sugarcane. ¹⁵N-ammonium and ¹⁵N-nitrate influx analyses in *B. distachyon* upon N-replete and N-deficient conditions revealed that ammonium is the preferential inorganic N source. A negative feedback regulation of nitrate uptake process by ammonium in *Brachypodium* plants under N provision and N resupply was observed. *BdNRT2.1* and *BdNRT2.2* expression analyses suggest that both proteins are the main *B. distachyon* NRT2 transporters responsible for nitrate acquisition. Both identified *BdNRT3* members were co-expressed with *BdNRT2.1* and *BdNRT2.2*. So, the ¹⁵N-nitrate influx analysis and transcript accumulation of members of the two-component complex (NRT2/NRT3) after ammonium chloride provision and ammonium nitrate resupply treatments suggest the transcriptional regulation of HATS in *B. distachyon*. Conversely, studies with sorghum and sugarcane revealed distinct regulation of HATS nitrate uptake process under nitrate resupply. ¹⁵N-nitrate influx assays showed high nitrate uptake by sorghum 'BTX623' cultivar and the sugarcane 'IAC87-3396' genotype. Expression analysis of genes encoding NRT2 and NRT3, HATS complex components, exhibit distinct transcriptional regulation among sugarcane genotypes. The lack of correlation between the accumulation of NRT2.1 and NRT3.1 proteins with the ¹⁵N-nitrate influx suggest the presence of a mechanism controlling nitrate transport in sorghum and sugarcane roots at post-translational level. This putative allosteric regulation of the HATS complex would be the major player on nitrate uptake in roots of sorghum and sugarcane. A better understanding of the mechanisms and molecules involved in the post-translational regulation of the nitrate uptake process and, therefore, in the signaling pathway of this N source in grasses would bring about the development of crops with higher NUE.

Keywords: Nitrate, HATS, *Brachypodium*, Sugarcane, *Saccharum*, NUE

1. INTRODUCTION

1.1 N fertilizers and environmental impacts

Nitrogen (N) is an essential macronutrient for plant growth and development (Fageria and Baligar 2005). N is the main constituent of nucleotides, amino acids, and molecules such as chlorophyll, participating in most of the physiological processes of plant development (Reetz 2016). N is also an important signal, affecting a large number of plant processes, including lateral root growth, biotic and abiotic stress response, seed germination and hormone signaling (Vidal et al. 2010, Alvarez et al. 2012, Wang et al. 2012a, Xu et al. 2012).

Considering the grown agricultural needs in the end of 19th century, biological nitrogen fixation (BNF) and manure recycling were considered insufficient, and fossil nitrogen deposits (Chilean saltpeter and Peruvian guano) were becoming depleted (Sutton et al. 2013). The development of the 'Haber-Bosch' process in the early 20th century enabled a massive boost in N fertilizer production, contributing to the increase of agricultural production to support population growth, enhanced nutrition and diminished hunger in the world (Smil 2002, Erisman et al. 2008, Follett et al. 2010, Erisman et al. 2013). In this process, combination of N₂ with H₂ allowed the large-scale production of NH₃ that has since doubled total global circulation of N compounds (Galloway et al. 2008, Erisman et al. 2008, Fowler et al. 2013). Around 2% of world energy use is dedicated specifically to the industrial manufacture of N fertilizer through the Haber-Bosch process (Sutton et al. 2013), pointing the large energy requirements of this process. Annually, 120 Tg of atmospheric N is applied as nitrogen fertilizer to sustain large agricultural production around the world (Galloway et al. 2008). Added to this amount, 50-70 Tg of N through the FBN is input in agricultural systems (Herridge et al. 2008).

Natural mechanisms of N fixation in the soil alone would not meet the currently global population growth and the global food demand (Crews and Peoples 2004). Crop production would be 48% lower without N fertilizer use (Erisman et al. 2008), confirming its important role in the Green Revolution (Tilman et al. 2002, Mueller et al. 2012, Sinclair and Rufty 2012). On the other hand, the excessive use of N fertilizers generates substantial ecological and environmental impacts, since most of the reactive N added into the agricultural system does not reach its final goal, i.e., protein in the human diet (Robertson and Vitousek, 2009). For every 100 units of N used in agriculture around the world, only 17 units are consumed by humans as crop foods, dairy products, and animal protein (UNEP and WHRC 2007).

N losses from the soil-plant system are estimated to be around 50% and 70% (Peoples et al. 1995, Smil 1999, Hodge et al. 2000, Cassman et al. 2002, Tilman et al. 2002, Balasubramanian et al. 2004, Dobermann 2007, Ciampitti and Vyn 2014). Losses result in the contamination of soils, water bodies and marine ecosystems, mainly due to eutrophication caused by nitrate leaching (Tilman 1999, Jaynes et al. 2001, Galloway et al. 2004, Schröder

et al. 2004, Burt et al. 2008, Vitousek et al. 2009, Foley et al. 2011). Further, N losses result in increased emission of greenhouse gases, such as nitric oxide and ammonia (Tilman et al. 2001, Galloway et al. 2008, Davidson 2009, Burney et al. 2010, Millar et al. 2010, Hoben et al. 2011, Reay et al. 2012); loss of biodiversity (Dirzo and Raven 2003); and the constant increment in reactive N in terrestrial ecosystems (Galloway et al. 2004, Martinelli and Filoso 2008, Austin et al. 2013).

Rockström et al. (2009) proposed the concept of "planetary boundaries" to fix human operational limits in several terrestrial processes. If exceeded, these boundaries could mean threatening climate change to human life. The N global boundary is calculated at 62 Tg N per year from industrial and intentional biological N fixation, fixed to avoid eutrophication of aquatic ecosystems (de Vries et al. 2013, Steffen et al. 2015). The global human population is expected to reach ~9 billion by 2050, requiring the boost in food production demand and the use of N fertilizer (Kearney 2010). Increasing demand for food, the need for more environmental friendly agriculture and future hazards resulting from climate change are all associated with the urgent need for the modern agriculture to improve Nitrogen Use Efficiency (NUE) in plants (Zhang et al. 2015).

In this review, we will discuss important aspects related to the environmental consequences due to low NUE in plants. Added to the updated overview of Nitrate Transporters (NRTs), we will discuss molecular components involved in nitrate uptake and its regulation.

1.1.1 Nitrogen as a threat to water resources

Excessive and indiscriminate N fertilizers use causes serious environmental impacts, driving pollutants to accumulation in water bodies, decreasing quality. Research has shown that approximately 39-95 Tg of reactive nitrogen species (N_r) is released from agriculture soil into freshwater per year (Billen et al. 2013, Bouwman et al. 2013). Part of the leached N reaches coastal waters (Seitzinger et al. 2005, Voss et al. 2011), causing eutrophication and hypoxia, creating oceans dead-zones. The eutrophication of freshwater and marine ecosystems is due mainly to nitrate accumulation, rising as a serious global issue (Rabalais et al. 2002, Selman et al. 2008, Sebilo et al. 2013). Excess of reactive nitrogen leads to increased growth of algae in water bodies. This algae population is consumed by aerobic bacteria, depleting dissolved oxygen and inducing hypoxia, leading to the decline of oxygen-dependent marine life, such as coral reefs, fish, seagrass and shellfish (Rabalais et al. 2002, Canfield et al. 2010, Bouwman et al. 2013). Other phenomena that drive to oxygen depletion in water bodies with low water turnover are sedimentation and decomposition of large mass of phytoplankton resulted from eutrophication of such aquatic bodies (Rabalais et al. 2002). Diaz et al. (2010) identified over 770 eutrophic and hypoxic coastal systems around the globe,

with about 220 areas of attention for developing hypoxia and 540 areas with documented hypoxia (Figure 1).

Figure 1. Eutrophic and hypoxic hotspot areas in the world (Diaz et al. 2010). Eutrophic areas are those with excess nutrients (orange dots), putting them at risk of adverse effects. Hypoxic areas are those where oxygen levels in the water are already depleted and adverse effects expected due to nutrient and or organic pollution (red dots). Blue dots are systems that were hypoxic at one time but now in recovery from management of nutrients and organic matter.

As the contamination of nitrate in aquatic systems is due to a long period of application and N fertilizer retention in agricultural soils (Sebilo et al. 2013), significant steps must be taken to diminish the environmental impact and improve water quality. So, such knowledge highlights the urgency for an immediate reduction and more efficient use of N fertilizers (Grizzetti et al. 2011).

1.1.2 N fertilization as a threat to air and atmosphere quality

Combined with N_r released into water bodies, N-gases are lost to the atmosphere, contributing to air pollution (Figure 2) (Moldanova et al. 2011). An example of gases discharged by the misuse of N fertilizers is N_2O , which contributes to the worsening of the greenhouse effect (Robertson and Vitousek 2009). Agriculture is responsible for 60% of this total (Mosier et al. 1998, Crutzen et al. 2007, Davidson 2009, Syakila and Kroeze 2011, Bouwman et al. 2013). Agriculture-related N_2O emissions are predicted to rise from about 6.4 Tg N_2O per year in 2010 to 7.6 Tg N_2O -N per year in 2030 (US Environmental Protection Agency 2011). The notable anthropic participation in N_2O emissions, mainly due to the agriculture activities, confronts efforts to reduce CO_2 emissions. On a molecule-by-molecule basis, an N_2O molecule can absorb 300-fold more infrared radiation than CO_2 , representing a potent greenhouse gas (Ravishankara et al. 2009). Even contributing only 6% to the greenhouse effect (Forster et al. 2007), decreasing N_2O emissions per unit of N fertilizer

applied is key in alleviating the adverse effects resulting from the indiscriminate use (Ravishankara et al. 2009, Reay et al. 2012, Campbell et al. 2017). Acknowledged as one of the most important N loss pathway in agricultural systems (Freney et al. 1990, Houlton et al. 2006), the denitrification process produces other N-gases besides N_2O . Nitric oxide (NO) and nitrogen dioxide (NO_2), identified collectively as NO_x , contribute to air pollution and acid rain generation (Robertson and Vitousek 2009). Another serious impact caused by NO_x emission is the formation of O_3 in the troposphere (Butterbach-Bahl et al. 2011). The increase in tropospheric ozone harms human health, evidenced by the increase in respiratory diseases (Von Mutius 2000, WHO 2003, 2008). Also, tropospheric ozone has significant effects on ecosystem biodiversity (Wedlich et al. 2012).

Figure 2: Hot spots of modelled atmospheric nitrogen deposition (Galloway et al. 2008), based on Dentener et al. (2006) (kg N per hectare per year). In North America, Europe, South and East Asia and parts of Latin America, the high values of Nr deposition are associated with fossil fuel combustion and intensive agricultural activity. In areas of Central Africa and Latin America with little N input, high levels of deposition are associated with biomass burning. Satellite data suggest that the modelled values are under-estimated in parts of central Asia (Clarisse et al. 2009).

Another relevant form of N loss is the volatilization of ammonia. This N-gas, added to transferring large amounts of reactive N to low turnover ecosystems (Robertson and Vitousek, 2009), causes several environmental impacts, such as acidification and eutrophication of terrestrial and aquatic ecosystems, decreased availability of nutrients, disruption of the N fixation, and biodiversity damage (Bruinsma 2017). NH_3 also causes adverse effects on human health, through the formation of fine particulate matter and photochemical smog (Sutton et al. 2013).

Alongside to the world population growth, and consequently, increased demand for food, feedstock and energy production, the agricultural land area and the N_r emission into the atmosphere will grow continually (Mosier and Kroeze 2000, Van Vuuren et al. 2006, Galloway et al. 2008, Davidson 2009, US Environmental Protection Agency 2011). Thus, improving nitrogen use efficiency in agricultural systems worldwide becomes a key procedure to fit the food, feed and bioenergy demand without provoking major environmental impacts.

1.1.3 Nitrogen as a threat to soil property

Agricultural expansion has had tremendous consequences on habitats, biodiversity, carbon storage, and soil conditions (Tilman et al. 2002, Foley et al. 2005, Millennium Ecosystem Assessment 2005, Steinfeld et al. 2006). N_r accumulation has a spectrum of impacts on soil quality, including soil erosion, acidification, salinization, compaction, contamination, and organic matter decline, affecting N losses to water and air (Velthof et al. 2011).

Recent investigations have identified critical soil acidification derived from N fertilizer application (Guo et al. 2010, Lu et al. 2014). Due to the release of H^+ during hydrolysis after application of N fertilizers causes a decrease in soil pH (Mahler and Harder 1984). This outcome following the application of N fertilizers induces the development of infertile soils non-responsive to crop yields with further application of N fertilizers (Herrero et al. 2010, Schroder et al. 2011), thereby resulting in inefficient use of fertilizers (Liebig et al. 2002). Another consequence of soil acidification is the release of heavy metals in soil solution, which may lead to nutrient disorders and even toxic effects in plants (Sutton et al. 2013).

Soil pH is reported to affect the microbial community in soil, another soil health parameter affected by the indiscriminate use of N fertilizers. Soil biota is negatively affected by N fertilizer over-use by the damage of their habitats and disrupting their functions (Chivenge et al. 2011). Accumulation or depletion of N_r in soils has so far been neglected in future scenarios (Bouwman et al. 2009, 2013), considering that soil organic matter (SOM) is stable and all excessive N_r will volatilize or drain. SOM is a fundamental indicator of soil fitness because of its vital functions that affect soil fertility, productivity, and the environment. N fertilizer application stimulates the decomposition of organic matter in the soil and adversely influence the soil health (Grace et al. 2006, Singh 2018). Changes in the microbial community impact soil processes like organic matter mineralization and nutrient cycling (Velthof et al. 2011), illustrating pathways affected by N fertilizer application.

Negatively effects on biodiversity of aboveground species by deposition of atmospheric N in terrestrial environments is observed to reduce the lichens population and plant diversity (van Herk et al. 2003, Wolseley et al. 2006, Bobbink et al. 2010, Dise et al. 2011). Change of species composition forces competitive interactions, creating composition change and/or

unfavorable conditions for some species. So, added to NUE improvement in crops, soil health parameters should be applied to estimate the sustainability of agricultural development (SDSN 2015).

1.2 Nitrogen Use Efficiency

Increasing the nitrogen use efficiency (NUE) by crop plants represents an urgent strategy in agriculture to reduce the cost of production and negative impacts on the environment derived from the overuse of N fertilizers. Several reviews have attempted to compile aspects and processes involved in the NUE of crops (Glass et al. 1992, Good et al. 2004, Fageria and Baligar 2005, Lea and Azevedo 2006, Fan et al. 2007, Hirel et al. 2007, Garnett et al. 2009, Robertson and Vitousek 2009, Masclaux-Daubresse et al. 2010, Pathak et al. 2011, Xu et al. 2012, Han et al. 2015, Bhattacharya 2019). Overall, two plant physiological components contribute to plant NUE: N utilization efficiency (NUE), describing the assimilation and remobilization efficiency of plant N; and N uptake efficiency (NUpE), which is the N absorption/uptake efficiency focusing on the recovery of N fertilizer from the soil (Moll et al. 1982, Good et al. 2004).

NUE is affected by N uptake, metabolism, allocation, and remobilization (Habash et al. 2001, Tsay et al. 2011, Girondé et al. 2015). Many approaches to improving plant NUE have focused mainly on the genetic manipulation of N metabolic processes (Ameziane et al. 2000, Chichkova et al. 2001, Habash et al. 2001, Yamaya et al. 2002, Seiffert et al. 2004, Good et al. 2007, Shrawat et al. 2008, Peña et al. 2017, Perchlik and Tegeder 2017). As a multigenic trait, NUE involves the coordinated expression of several genes including, but not limited to N assimilation. Transgenics overexpressing genes involved in primary N assimilation did not lead to improvement in NUE (Pathak et al. 2008, Pathak et al. 2011, Krapp et al. 2014, Sinha et al. 2018).

NUpE is also controlled by many aspects including the N soil availability and by plant N status, plant N assimilation, and N metabolite levels (Ruffel et al. 2011, Nacry et al. 2013, Stahl et al. 2016). NUpE is also dependent on nitrate and ammonium transport at the root level (von Wirén et al. 2000, Miller and Cramer 2005, Miller et al. 2007, Xu et al. 2012, Krapp et al. 2014). Low N uptake rates are responsible for the increase in the economic cost of crop production and environmental pollution (Fageria and Baligar 2005). Among NUE-related traits, NUpE appears to be critical for many plants (Dhugga and Waines 1989, Imsande and Touraine 1994, Singh et al. 1998, Le Gouis et al. 2000).

Plants uptake organic and inorganic N sources present in the soil, with ammonium (NH_4^+) and nitrate (NO_3^-) being the major inorganic forms acquired by the roots (Wolt 1994, Miller et al. 2007, Marschner 2012). On average, soil NO_3^- concentration is 10-fold higher than NH_4^+ concentration, making NO_3^- the predominant form of N available to crops (Miller et al.

2007). Because nitrate is negatively charged and water soluble, NO_3^- concentrations in soil fluctuate considerably, from micromolar to millimolar (Glass 2003). Advances of NUE in crops through manipulation of NO_3^- transporters involved in nitrate uptake has recently been successful (Hu et al. 2015, Fan et al. 2016a) suggesting that nitrate uptake processes may be limiting NUE improvement (Griffiths and York 2020).

1.3 Nitrate Acquisition

The N acquisition in plant roots occurs by membrane transporters, classified according to the affinity for their substrate. HATS (High-Affinity Transporter System) and LATS transporters (Low-Affinity Transporter System) are activated in low and high N concentrations, respectively (Glass et al. 2002). Additionally, N transporters are classified for their response to the N presence. Transporters can be constitutive (cHATS), when the respective genes are not regulated by the presence of nutrient, or inducible (iHATS) when the transporters gene is regulated by the N presence (Tsay et al. 1993, Ninnemann et al. 1994, Huang et al. 1999, Li et al. 2007).

To deal with the fluctuation of nitrate concentrations in the soil, plants manage LATS and HATS nitrate uptake systems. HATS is active when NO_3^- availability in the soil is low (concentration lower than 250 μM); whereas LATS predominates at high soil nitrate concentration (>250 μM) (Siddiqi et al. 1990, Kronzucker et al. 1995, Garnett et al. 2003). However, investigations have revealed that the HATS contributes to nitrate uptake in concentrations higher than 2.5 mM, attending the plant N demand (Malagoli et al. 2004, Garnett et al. 2013). Several membrane proteins have been characterized and identified controlling nitrate absorption, translocation, and storage in plants more than 20 years ago. The nitrate transporters are categorized into four families: Nitrate Transporter 1/Peptide Transporter (NRT1/NPF), Nitrate Transporter 2 (NRT2), the Chloride Channel (CLC), and the Slow Anion Associated Channel Homolog (SLC/SLAH) family (Orsel et al. 2002, Tsay 2007, Negi 2008, Barbier-Brygoo et al. 2011, Dechorgnat et al. 2011, Wang et al. 2012, Krapp et al. 2014, Lran et al. 2014, Noguero and Lacombe 2016, O'Brien et al. 2016, Fan et al. 2017, Li et al. 2017).

The tonoplast localized CLC proteins (CLCa and CLCb) present critical role in NO_3^- accumulation in the vacuole by proton- NO_3^- exchange mechanism (De Angeli et al. 2006, Von Der Fecht-Bartenbach et al. 2010). AtCLCa and AtCLCb showed interconnection, but not redundancy, among each nitrate assimilation pathways, exhibiting a crosstalk between both transporters (Monachello et al. 2009). The SLAC/SLAH family in Arabidopsis contains five transporters, SLAC1 and SLAH1–SLAH4 (Krapp et al. 2014). AtSLAC1 and AtSLAH3 are likely nitrate transporters in guard cells involved in stomatal closure (Geiger et al. 2011).

The activity of both LATS and HATS involved in nitrate uptake depends on the cell energy supply (Miller et al. 2007, Siddiqi et al. 1990). Additionally, NRT1/NPF and NRT2 transporter families are known to transport nitrate in a symport proton-coupled mechanism that is driven by the pH gradients across membranes (Espen et al. 2004, Xu et al. 2012, Xia et al. 2015, Fan et al. 2016b). The NPF family (NRT1/PTR family) contains 53 genes in *Arabidopsis* and 93 in rice (Wang et al. 2018a), and up to 139 genes in higher plants genomes, classified into eight subfamilies (Lerán et al. 2014, von Wittgenstein et al. 2014, O'Brien et al. 2016, Noguero and Lacombe 2016). Members of this family share common protein topology, displaying 12 transmembrane domains connected by loops (Tsay et al. 2007). NRT2 transporters have similar transmembrane protein arrangement to the NPF family, but there is no homology between the two families (von Wittgenstein et al. 2014). NRT2 family members, from the major facilitator superfamily (MFS) of transporters, were first identified in *Aspergillus nidulans* and *Chlamydomonas reinhardtii* (Truemann et al. 1996). Seven NRT2 members are present in the *Arabidopsis* genome (Miller et al. 2007). Five NRT2 transporters were identified in the rice genome (Feng et al. 2011). Using *Arabidopsis* orthologous, Plett et al. (2010) identified NRT2 members in maize, rice, sorghum, and *Brachypodium distachyon*. They are described as influx HATS transporters for nitrate uptake in roots (Filleur and Daniel-Vedele 1999), generally unable to transport NO_3^- on their own, requiring forming a heteroligomer with AtNAR2.1 (AtNRT3.1), which belong to the NAR2 (NRT3) family, for a functional HATS in nitrate uptake in root plants (Tong et al. 2005, Kawachi et al. 2006, Okamoto et al. 2006, Orsel et al. 2006, Chopin et al. 2007, Li et al. 2007, Yong et al. 2010, Yan et al. 2011, Kiba et al. 2012, Kotur et al. 2012, Laugier et al. 2012, Lezhneva et al. 2014, Kotur and Glass 2015, Lupini et al. 2016). The first NAR2/NRT3 gene was identified in *Chlamydomonas reinhardtii* (Quesada et al. 1994). Orthologs in plants have been identified in *Arabidopsis*, rice, barley, maize, sorghum and *Brachypodium distachyon* (Tong et al. 2005, Miller et al. 2007, Plett et al. 2010, Feng et al. 2011, Wang et al. 2012b).

Since the focus of this review is the environmental impacts caused by the low NUpE, the major nitrate transporters families involved in the nitrate uptake process (NRT1/NPF, NRT2, and NRT3 families) will be described in detail in the next topics.

1.3.1 Nitrate Transporter 1/Peptide Transporter (NRT1/NPF) members

Although the NPF members are described as acting as the main components of the LATS, some proteins are dual-affinity transporters, involved in both HATS and LATS. AtNPF6.3 (AtNRT1.1) was the first nitrate transporter identified involved in the NO_3^- uptake process (Tsay et al. 1993), performing dual-affinity nitrate transport activity (Figure 3) (Liu et al. 1999). The switch between the HATS and LATS form is determined by the Thr101 phosphorylation of NPF6.3 by the CIPK23 kinase (CBL-INTERACTING PROTEIN KINASE

23; Liu and Tsay 2003, Ho et al. 2009). This post-translational modification occurs under low nitrate concentration in the soil, promoting high-affinity nitrate transport activity. Crystallization of the AtNPF6.3 structure infers that Thr101 phosphorylation switches its affinity for nitrate decoupling the AtNPF6.3 dimer configuration (Parker and Newstead 2014, Sun et al. 2014). The ability to change the nitrate affinity according to NO_3^- availability reveals that AtNPF6.3 behaves as a nitrate sensor rather than a nitrate transporter (Ho and Tsay 2010). Gojon et al. (2011) proposed NPF6.3 as a 'transceptor', i.e., a protein with dual transport/sensing function. In addition to the ability to adapt function in response to external nitrate concentration, studies have indicated that NPF6.3 plays a role in cell signaling. NPF6.3 is responsible to trigger the primary nitrate response in plants, as stimulating the emergence of lateral roots, and modulating *NRT2.1* gene expression upon nitrate starvation (Munõs et al. 2004, Little et al. 2005, Ho et al. 2009, Krouk et al. 2010, Gojon et al. 2011, Xu et al. 2012, Krapp et al. 2014).

Orthologous of AtNPF6.3 in other species have been identified involved in the nitrate uptake process, although monocots and dicots differ in the number of *NRT1.1* genes (Plett et al. 2010). The ortholog of AtNPF6.3 in *Medicago truncatula*, MtNRT1.3, was classified also as dual-affinity transporter involved in nitrate uptake (Morère-LePaven et al. 2011). Genes expression analysis and subcellular localization patterns in rice proposed OsNRT1.1B (OsNPF6.5), the closest orthologs of AtNPF6.3 in rice, as with the same nitrate uptake, transport and signaling functions (Lin et al. 2000, Hu et al. 2015, Xia et al. 2015, Wang et al. 2018a). NRT1.2 (AtNPF4.6) antisense lines exhibited lower nitrate uptake than wild-type plants, indicating its functional role in Arabidopsis (Huang et al. 1999).

NPF2.7, identified as low-affinity nitrate excretion transporter1 (NAXT1), is mainly involved in nitrate efflux in Arabidopsis root (Segonzac et al. 2007). OsNPF2.4 (OsNRT1.6) participates not only in nitrate acquisition in rice roots but also in root-to-shoot nitrate transport and source-to-sink nitrogen remobilization (Xia et al. 2015). In maize, from the 86 NPF genes identified, *ZmNPF6.4* and *ZmNPF6.6* genes are regulated to nitrate exposure, similar to *AtNPF6.3* (Sekhon et al. 2011, Lran et al. 2014, Wen et al. 2017). In *Saccharum spontaneum*, 178 NPF genes were identified spread across all monoploid chromosomes. Several *SsNPF* genes are products of gene fusions of several tandem duplications, providing valuable structural resources for functional characterization of nitrate transporters in sugarcane (Wang et al. 2019a). All other NPFs characterized to date are mainly associated with internal transport processes, such as xylem and phloem nitrate loading, and nitrate translocation to leaves or seed (Dechorgnat et al. 2011, Xu et al. 2012, O'Brien et al. 2016, Wang et al. 2018a).

1.3.2 Nitrate Transporter 2 (NRT2) family

The first NRT2 gene family member in higher plants were identified based on the knowledge obtained in *Aspergillus nidulans* and *Chlamydomonas reinhardtii*. In the algae *C.*

reinhardtii, CrNRT2.1 was identified to participate in nitrate transport by complementation of a deletion mutant (Quesada et al. 1994, Truemann et al. 1996). Seven NRT2 transporters have been characterized in Arabidopsis (Krapp et al. 2014). After the identification of *AtNRT2.1* and *AtNRT2.2* (Filleur and Daniel-Vedele 1999, Zhuo et al. 1999), studies confirmed their role in the nitrate uptake process in Arabidopsis roots (Figure 3) (Cerezo et al. 2001, Filleur et al. 2001, Li et al. 2007). In the deletion mutant *atnrt2.1*, iHATS showed a 72% decrease in nitrate influx, whereas *atnrt2.2* showed a 19% reduction. Double mutants (*atnrt2.1/atnrt2.2*) revealed an 80% reduction of iHATS, demonstrating the major role of *NRT2.1* and *NRT2.2* in the nitrate uptake process in Arabidopsis (Li et al. 2007). Two additional AtNRT2 family members are involved in nitrate influx into roots. NRT2.4 has been shown to participate in nitrate acquisition during long-term N starvation. Compared to *atnrt2.1/atnrt2.2* double mutant, a triple mutant (*atnrt2.1/atnrt2.2/atnrt2.4*) with a loss of function of NRT2.4, showed lower biomass production under low nitrate supply. NRT2.4 is a nitrate transporter that has a role in both roots and shoots under N starvation (Kiba et al. 2012). Similar to AtNRT2.4, AtNRT2.5 also plays a role, contributing to nitrate uptake and nitrate loading in the phloem in N-deficient plant (Lezhneva et al. 2014). *AtNRT2.3*, *AtNRT2.6* and *AtNRT2.7* are insensitive to nitrate supply, and constitutively transcribed. *AtNRT2.3* did not show nitrate uptake activity in oocytes (Orsel et al. 2006). Correlation between *AtNRT2.6* expression and ROS accumulation in response to infection by *Erwinia amylovora* shows this gene is related to plant-microbe interaction (Dechorgnat et al. 2012). *AtNRT2.7* is the only exception among other NRT2 transporters in Arabidopsis since its expression is more accentuated in shoots than roots (Okamoto et al. 2003). Overexpression of *AtNRT2.7* or a simple mutation affect seed nitrate content, showing that this gene plays a specific role in nitrate accumulation in the seed (Chopin et al. 2007). Recently, another study has suggested that *AtNRT2.7* is also involved in seeds proanthocyanidins accumulation and oxidation (David et al. 2014).

Figure 3: Summary of key transporters and regulators mediating nitrate uptake in plant roots. The dual-affinity transporter, NPF6.3 and the high-affinity transporters, NRT2.1 and NRT2.2 are involved in acquiring nitrate from the rhizosphere, while NRT2.5 mediates the loading of nitrate into the transpirational stream in the stele. The root tissue types represented are epidermis (EP), cortex (CO), and stele (ST). Depicted are the transporters (circles), transcription factors (squares), kinases (trapezoids), peptides (triangles) and chromatin regulators (pentagons). Regulation of the transporters which has been established as direct interaction (red arrows) or indirect interaction (or not determined to date) (blue arrows). Transporters are localized to the tissue in which they are most highly expressed, and the area of the transporter circle represents the relative expression level of the genes encoding the respective transporter in either low (left) or high (right) nitrate provision. Top half of the diagram represents mature root tissue, while the bottom represents the root tip region. Source: Plett et al. 2018.

Phylogenetic analysis of NRT2 genes across the plant kingdom are well discussed (Plett et al. 2010, von Wittgenstein et al. 2014). NRT2 grass genes show a significant similarity distance from AtNRT2, implying that proper functional analysis in grasses should be done to examine functions of NRT2 in monocots (Plett et al. 2010). In rice, five high-affinity NRT2 transporters have been identified (Cai et al. 2008, Feng et al. 2011). *OsNRT2.1* and *OsNRT2.2* are highly similar to their corresponding homologs in other monocots, associated with nitrate acquisition. *OsNRT2.3* is transcribed into two spliced isoforms (*OsNRT2.3a* and *OsNRT2.3b*). *OsNRT2.3b* overexpression in rice significantly enhances grain yield and NUE (Fan et al. 2017). *OsNRT2.3a* exhibits a role in long-distance nitrate transport from root to shoot in rice since its downregulation leads to the impairment of xylem loading and decreases plant growth at low nitrate levels (Yan et al. 2011). Thus, both alternative splicing forms play distinct functions in rice development. *OsNRT2.4* was recognized as a dual-affinity nitrate transporter, required for nitrate-regulated root and shoot growth (Wei et al. 2018).

Comprehensive phylogenies reconstruction supplemented with other data (transmembrane domain topology predictions; subcellular localization predictions; and *in silico* expression profiling) recognized seven orthologues of *AtNRT2* genes in *Brachypodium*

distachyon genome (von Wittgenstein et al. 2014). Those *BdNRT2* gene encoding putative high-affinity nitrate transporters were evaluated in this monocot model plant (David et al. 2019, Wang et al. 2019b). The *bdnrt2.1* mutant exhibited a decrease of 30% of nitrate influx, revealing that *BdNRT2.1* acts as the main *BdNRT2* member for the nitrate uptake process in this grass model plant. In the wild type, *BdNRT2.1* and *BdNRT2.2* expressions were strongly induced by nitrate resupply, while the *BdNRT2.5* was suppressed. In *bdnrt2.1* mutant, *BdNRT2.2* and *BdNRT2.6* genes were significantly upregulated. Other *BdNRT2* genes presented constitutive expression in roots, demonstrating that *BdNRT2* gene members have diverse roles in response to various nitrogen conditions (Wang et al. 2019b).

In barley, four cDNAs of *HvNRT2* genes were isolated from roots (Trueman et al. 1996, Vidmar et al. 2000). However, only after 20 years, *HvNRT2* genes were described based on genomic information. Guo et al. (2020) identified ten putative *NRT2* genes in the barley genome. Tandem duplication events could explain the difference in the number of *NRT2* genes among barley, Arabidopsis, rice, and maize (Guo et al. 2020). *HvNRT2.10*, a putative homologous of *AtNRT2.7*, displayed higher expression in the shoots than the roots, and a relatively high expression in seeds, which is consistent with its homologs in Arabidopsis (Chopin et al. 2007). Overexpression of *HvNRT2.1*, a homologous of *OsNRT2.3*, improved yield-related traits in Arabidopsis, indicating that *HvNRT2.1* plays a specific role in nitrate accumulation in barley (Guo et al. 2020).

Previous investigations recognized only 3 *NRT2* orthologous in sorghum (Plett et al. 2010). Following research comparing N-stress tolerant and sensitive sorghum genotypes correlated a higher abundance of *NRT2* transcripts (*SbNRT2.2*, *SbNRT2.3*, *SbNRT2.5*, and *SbNRT2.6*) with N tolerant, adaptation to N deficiency by producing greater root mass for efficient uptake of nutrients, in sorghum (Gelli et al. 2014). Additionally, combining online genome database and identification based on homology, five *SbNRT2* genes in total were identified in the *S. bicolor* genome (Massel et al. 2016).

Until 2008, only two *AtNRT2.1* homologous genes, *ZmNRT2.1* and *ZmNRT2.2*, had been classified in maize (Quaggiotti et al. 2003, Trevisan et al. 2008). Phylogenetic analysis conducted made possible to identify two additional *NRT2* nitrate transporter (Plett et al. 2010). The regulation of *ZmNRT2.1* expression by NO_3^- availability together with preference for expression suggest that this gene is a major component of the iHATS for nitrate uptake along maize root axis (Sorgonà et al. 2011). Garnett et al. (2013) found a redundancy of function between *ZmNRT2.1* and *ZmNRT2.2* transporters, considering that both genes were upregulated at N-limiting conditions. Broad studies should be conducted to functionally evaluate *NRT2* transporters in maize roots. Traits *NRT2*-related, as root system architecture, influencing nitrate uptake and accumulated tissue N content have been shown should be considered to improve NUE in maize (Dechorgnat et al. 2018).

Wang et al. (2019a) were capable to identify 20 NRT2 members across all eight *S. spontaneum* monoploid chromosomes. As observed in many other species, including *Arabidopsis*, *Brachypodium*, rice, maize, and *Medicago truncatula*, *SsNRT2.1* and *SsNRT2.2* are present as tandem repeats in the genome (Orsel et al. 2002, Cai et al. 2008, Garnett et al. 2013, Pellizzaro et al. 2015, Wang et al. 2019a, b). *SsNRT2* members share the same gene structure as their homologs in sorghum and *B. distachyon*, different from *AtNRT2* genes. These results suggest that the *NRT2* gene structures are relatively conserved among monocots, but their functions are likely to differ from the well-characterized *NRT2* genes in *Arabidopsis* (Wang et al. 2019a).

NRT2 transporters were identified in other species. *TaNRT2.1*, the *AtNRT2.1* orthologue in wheat, was firstly characterized by Yin et al. (2007). Further studies dissected more members in wheat and their respective regulation under contrasting N conditions. In plants under high N supply, *TaNRT2.1*, *TaNRT2.2*, and *TaNRT2.3* expressions were downregulated (Wang et al. 2011). Correlation between NO_3^- uptake and *TaNRT2.1* expression suggest that *TaNRT2.1* may be a candidate for the inducible HATS transport in wheat, presumably with the major role in nitrate uptake, up to post-flowering stage (Wang et al. 2011, Taulemesse et al. 2015). Together with *TaNRT2.1*, *TaNRT2.5* was shown to be involved in post-anthesis nitrate acquisition. Mutant defective for *TaNRT2.5* (*tanrt2.5*) showed low grain nitrate concentration and altered seed vigor. So, *TaNRT2.5* has a key role in regulating grain nitrate accumulation and seed vigor (Li et al. 2020).

Heterologous expression of *CmNRT2.1* in *atnrt2.1* mutant was accompanied by an enhanced rate of nitrate uptake, suggesting that *CmNRT2.1* represents a high-affinity nitrate transporter in *Chrysanthemum* root (Gu et al. 2016). Similar complementation of *atnrt2.1* mutant with *MeNRT2.1*, *AtNRT2.1* homolog in cassava revealed that *MeNRT2.1* plays a crucial role in nitrate uptake in transgenic *Arabidopsis* plants under low N availability (Zou et al. 2019). In the *Coffea canephora* genome, the *NRT2* family was represented by three members, two exclusively expressed in roots (dos Santos et al. 2017). The first *NRT2* high-affinity nitrate transporter identified in *Nicotiana plumbaginifolia* was obtained cloning based on homology to transporters from *A. nidulans*, *C. reinhardtii*, and *H. vulgaris*. *Nrt2:1Np* is expressed specifically in roots and it is induced by nitrate (Quesada et al. 1997). Correlation between *NRT2* mRNA accumulation and ^{15}N -nitrate influx rates was the first evidence of its importance in *N. plumbaginifolia* (Krapp et al. 1998). HATS activity was restored in *atnrt2.1* mutant complemented with *Nrt2:1Np* gene, corroborating its role in nitrate uptake process. (Filleur et al. 2001). In the same genus, two *NRT2* orthologs were identified in *N. tabacum*, *NtNRT2.1* and *NtNRT2.2* (Liu et al. 2018). Those genes present a greater similarity with *LeNRT2.3*, a *NRT2* member functionally characterized in *Solanum lycopersicum* (Fu et al. 2015). Sestili et al. (2018) confirmed that, besides *LeNRT2.3* expression in roots, *LeNRT2.1*

presented expression regulated by nitrate availability. Another NRT2.1 of Solanaceae family was identified in Habanero pepper (*Capsicum chinense* Jacq.) *CcNRT2.1*, expressed exclusively in roots, with preferential location in root epidermal and cortical cells, supporting that *CcNRT2.1* belongs to nitrate uptake HATS (Lizama-Gasca et al. 2020).

So, NRT2 transporters play an important role in NO_3^- uptake from the soil. But, unlikely *A. nidulans* NRT2, NRT2 transporters of higher plant require nitrate assimilation-related protein (NAR2/NRT3) to function in nitrate uptake (Okamoto et al. 2006, Orsel et al. 2006). NRT2/NRT3 interactions will be described below.

1.3.3 Nitrate Transporter 3 (NRT3) family

The NAR2 (NRT3) family members are auxiliary proteins for HATS nitrate uptake (Kawachi et al. 2006, Okamoto et al. 2006, Orsel et al. 2006, Yong et al. 2010, Laugier et al. 2012). The requirement for simultaneous *NRT2/NRT3* gene expression to sustain high-affinity transport system activated was first demonstrated in *Chlamydomonas reinhardtii*. The HATS capacity by CrNRT2.1 or CrNRT2.2 was lost in *crnar2.1* (Quesada et al. 1994). In the Arabidopsis genome, two *NAR2* (*NRT3*) genes are present, but only *AtNAR2.1* (*AtNRT3.1*) participates in the two-component high-affinity nitrate uptake system. iHATS is reduced by as much as 95% in *atnar2.1* (*atnrt3.1*) mutant (Okamoto et al. 2006, Orsel et al. 2006, Li et al. 2007). Several reports have shown that *NRT3.1* expression is regulated as *NRT2.1* expression, although with a much- less-pronounced amplitude in the transcript-level changes (Krouk et al. 2006, Okamoto et al. 2006, Orsel et al. 2006). The operational assembly for the HATS complex was proposed to be a tetramer consisting of two AtNRT2.1 subunits bound to two subunits of AtNRT3.1, displaying a complex of ~150 kDa (Yong et al. 2010). This hypothesis differs from a previous work by Wirth et al. (2007), who suggested that monomeric AtNRT2.1 could be the active form for nitrate transport. Except for AtNRT2.7, all Arabidopsis NRT2s proteins interact strongly with AtNRT3.1, meaning those can arrange the HATS functional complex. Recently, a 150 kDa complex of AtNRT2.5 and AtNAR2.1 was shown to be involved in cHATS (Kotur and Glass 2015).

Regarding nitrate transporters in rice, two *OsNRT3* genes have been identified so far. The OsNAR2.1 (OsNRT3.1), induced by high and low N concentration, associates with three OsNRT2 transporters (OsNRT2.1, OsNRT2.2, and OsNRT2.3a) for nitrate uptake in rice roots. Both OsNRT2.1 and OsNRT2.2 high-affinity transporters need to interact with OsNAR2.1 (OsNRT3.1) for a functional nitrate uptake in rice roots (Yan et al. 2011). The long-distance nitrate transporter OsNRT2.3a requires OsNRT3.1 to operate, although showing a 10-fold lower affinity for nitrate than OsNRT2.1 and OsNRT2.2. So, depending on the partner transporter, OsNRT3.1 can provide a switch to allow a prompt response over the dynamic variation of external nitrate concentrations. OsNRT2.3b and OsNRT2.4 can function

independently of OsNRT3 (Araki and Hasegawa 2006, Cai et al. 2008, Feng et al. 2011, Yan et al. 2011, Tang et al. 2012). The Arginine 100 and Aspartic acid 109 of OsNRT3.1 are the key amino acids in the interaction with OsNRT2.3a for plasma membrane localization and nitrate transporter activity (Liu et al. 2014).

By Reciprocal Best Hit (RBH) approach, used to identify putative NRT3 Arabidopsis orthologues, were identified a pair of co-orthologous genes in maize, sorghum and Brachypodium (Plett et al. 2010). Both *BdNRT3* genes were expressed mainly in the roots, with a *BdNRT3.1* constitutive, whereas *BdNRT3.2* was repressed by high nitrate concentration. Since *BdNRT3* gene expression pattern follows the HATS activity, David et al. (2019) proposed *BdNRT3* as potentially involved in HATS complex formation. To date, no functional study has been conducted to show the NRT2 and NRT3 interaction in Brachypodium.

NRT3 genes have been identified in other species besides model plants. In the barley genome, three *HvNRT3* (*HvNAR2.1-2.3*) have been identified (Ishikawa et al. 2009, Guo et al. 2020). *Xenopus* oocytes co-injected with HvNRT2.1 and HvNAR2.3 showed an increase of ¹⁵N-nitrate uptake, suggesting a highly specific interaction between them (Tong et al. 2005). In the maize genome, two copies of the *ZmNRT3* family are present (Plett et al. 2010). The *ZmNRT2/NRT3s* transcriptional and protein profiles suggested that the iHATS regulation is mainly based on the transcriptional/translational modulation of *ZmNRT3.1A* (Pii et al. 2016). Further, *ZmNRT2.1* and *ZmNRT3.1A* appear to be associated in a ~150 kDa oligomer, as AtNRT2/NRT3 complex (Yong et al. 2010). Previous investigations recognized a pair of co-orthologous genes in sorghum (Plett et al. 2010). Based on *Sorghum bicolor* transcriptome (Makita et al. 2015), three *SbNRT3* genes in total were identified (Massel et al. 2016), but no functional studies with *SbNRT3* members has been conducted so far. Comprehensive analysis at genome-wide scale in *Saccharum spontaneum* identified six NRT3 genes, which exhibit high level of diversity, as opposed to NRT2 members which are more evolutionarily conserved (Wang et al. 2019a). In *Chrysanthemum*, protein-protein interaction approaches showed that CmNRT2.1 interacts at the plasma membrane with CmNAR2 (Gu et al. 2016). In *Medicago truncatula*, *MtNRT2.1* is suggested to be the major contributor to the HATS due to the same spatial expression profiles with *MtNAR2.1* and *MtNAR2.2*, members of NRT3 family (Pellizzaro et al. 2015). The co-expression of *CcNRT2.1* and *CcNAR2* strongly supports that both belong to a HATS in *Capsicum chinense*, being the first components of a HATS characterized in *Capsicum* (Lizama-Gasca et al. 2020).

In the *atnar2.1* mutants, NRT2.1 protein is absent in plasma membrane and nitrate HATS is deficient (Okamoto et al. 2006, Orsel et al. 2006), although mRNA encoding NRT2.1 is present (Wirth et al. 2007, Yong et al. 2010). It leads to the hypothesis that, unless NAR2.1 (NRT3.1) is available to generate the NRT2.1/NRT3.1 complex, NRT2.1 protein is degraded.

In barley, changes in HvNRT2.1 were not correlated with those of *HvNRT2.1* transcript accumulation during induction by nitrate (Ishikawa et al. 2009). Research using Arabidopsis transgenic lines 35S–*NRT2.1* has revealed post-transcriptional regulation of nitrate uptake by ammonium or dark treatment (Laugier et al. 2012). In the wild type, both the *AtNRT2.1* mRNA level and the HATS are downregulated by ammonium resupply, while the mRNA level in the Arabidopsis 35S–*NRT2.1* plant remained unchanged, but the HATS and NRT2.1 protein level were reduced. These results indicate, in addition to transcriptional control, the presence of a post-transcriptional regulation mechanism affecting HATS activity in Arabidopsis (Laugier et al. 2012).

With all the work indicating the importance of NRT2/NRT3 interaction for nitrate uptake across species, it is tempting to speculate that this interaction in the plasma membrane and the putative involvement of the membrane trafficking system may be important for regulating this interaction, thus providing a quick response method of adapting plant nitrate uptake capacity (Plett et al. 2018). So, this information highlights the influence of post-translational control mechanisms on the nitrate uptake system in plants, although, nitrate is also controlled at transcriptional and translational levels. In the next topic, we will review the nitrate uptake regulation processes.

1.4 Nitrate Uptake Regulation

Inorganic N use by plants is regulated by highly complex and sophisticated signaling pathways, activated by sensing of both external N availability and internal N status of the whole plant (Gojon et al. 2011, Krapp et al. 2014, O'Brien et al. 2016, Bellegarde et al. 2017, Fan et al. 2017, Li et al. 2017, Undurraga et al. 2017, Kant 2018, Wang et al. 2018b, Zhao et al. 2018, Fredes et al. 2019). Nitrate, as signaling molecule, triggers a wide range of molecular, physiological and developmental responses, known as primary nitrate response (PNR). One of the earliest events of the PNR involves NPF6.3, a nitrate transceptor (Ho et al. 2009, Gojon et al. 2011). Loss-of-function mutants of NPF6.3 strongly attenuates the upregulation of nitrate responsive genes, such as *NRT2.1* (Ho et al. 2009, Wang et al. 2009). *NPF6.3* mutant with defective nitrate uptake function shows a typical induction of nitrate genes, clearly demonstrating that NPF6.3-dependent induction of transcription occurs independently of its transporter activity (Bouguyon et al. 2015). In contrast to wild-type plants, several *atnrt1.1* mutants showed a lack of down-regulation of *AtNRT2.1* expression under high N conditions. This result evidenced that NPF6.3 is also associated with the signaling pathway triggering feedback repression of HATS genes, in response to high N supply (Muños et al. 2004, Krouk et al. 2006).

Important advances on the role of nitrate in modulating cellular, growth and developmental processes have been made in recent years which allow the postulation of a

canonical nitrate signaling pathway (Fredes et al. 2019). It includes the increase of cytoplasmic Ca^{2+} levels, a downstream of NPF6.3/NRT1.1 regulation event, controlling the activity of different Ca^{2+} -SENSOR PROTEIN KINASES (CPK) (Riveras et al. 2015, Liu et al. 2017). CPKs have been implicated to modulate the activity of the transceptor NPF6.3/ NRT1.1 and to regulate the function regulatory factors of the nitrate-signaling pathway. ANR1 (ARABIDOPSIS NITRATE REGULATED 1) was the first putative MADS-box transcription factor characterized with a role in response to N (Zhang and Forde 1998). ANR1 promotes lateral root development in nitrate-rich patches, in an NPF6.3-dependent manner (Remans et al. 2006). NLP7 (NIN-LIKE PROTEIN 7) is one of the master regulators of nitrate signaling (Figure 3) (Liu et al. 2017). NLP7 directly regulates many steps of the primary N assimilation and nitrate signaling pathways, such as *NRT2.1* expression (Castaings et al. 2009). Recently, the signaling mechanism by which NLP7 rapidly induces the PNR has been elucidated. Under nitrate deficiency, NLP7 is maintained in the cytoplasm by nuclear exclusion. With NLP7 nuclear retention mechanism regulated by nitrate perception, NLP7 is able to reach its genomic target loci (Marchive et al. 2013). In the presence of ammonium, genetic and molecular analyses indicate that NLP7 works upstream of NPF6.3 in nitrate regulation, and it may bind to specific regions of the promoter. In the absence of ammonium, NLP7 functions in nitrate signaling independently of NRT1.1, evidencing the regulation of NLP7 role in nitrate signaling by the N inorganic source exposure (Zhao et al. 2018). Other NLPs, in particular NLP6, can bind the nitrate responsive *cis*-element (NRE) present in the promoter region of some nitrate responsive genes (Konishi and Yanagisawa 2013). ZmNLP6 and ZmNLP8 overexpression in *atnlp7-4* recovered the nitrate assimilation and the induction of nitrate-responsive genes to WT levels, indicating that ZmNLP6 and ZmNLP8 exert major role in nitrate signaling and metabolism (Cao et al. 2017). Additionally, a functional characterization of ZmNLP5 defined its importance for the molecular network associated with N metabolism in maize (Ge et al. 2019). Overexpression of OsNLP1 and OsNLP4 in Arabidopsis *nlp7* mutant recovered its phenotype and increased the biomass, demonstrating that both OsNLPs are master regulators of NUE in rice (Alfatih et al. 2020, Wu et al. 2020). A protein with unknown molecular function, named NITRATE-REGULATED GENE2 (NRG2), has been identified as another signaling component of the PNR (Xu et al. 2016). NRG2 physically interacts with NLP7 in the nucleus, suggesting its importance as a player in the transcriptional response to nitrate (Xu et al. 2016). The identification of an antagonistic regulator by NLP primary transcription factors for nitrate signaling provided a resolution to the complexity of nitrate-induced transcriptional regulations in Arabidopsis. NIGT1 acts repressing the NLP-transcriptional cascade-mediated, such as *AtNRT2.1* expression downregulation (Maeda et al. 2018).

Protein-protein interactions between TFs (e.g., NLP7-TCP20 and NRG2-NLP7) have been found to be important for gene expression of nitrate responsive genes. The transcription

factor TCP20 (TEOSINTE BRANCHED1/CYCLO-IDEA/ PROLIFERATING CELL FACTOR1-20) plays a key role in nitrate root-foraging by controlling lateral root development under heterogeneous environments (Guan et al. 2014). In plants under N deficiency, TCP20 physically interacts with NLP6 and NLP7 controlling the expression of key nitrate-responsive genes, such as *AtNRT2.1* (Guan et al. 2017). In Arabidopsis, ATAF2 and NAP were identified as new TFs that positively regulate the expression of NIR and NRT2.1 in response to nitrate, respectively. This study revealed that NAP interacts with TGA1, which is known to be also involved in nitrate induction of NRT2.1 (Alvarez et al. 2014). NAP and TGA1 footprints in the NRT2.1 promoter region suggest that NAP-TGA1 interaction has a functional importance for NRT2.1 expression (Alvarez et al. 2019). Like TGA1, another bZIP transcription factor gene *TGA4* is able to bind in both NRT2.1 and NRT2.2 loci and activate their expression. In fact, about 97% of the genes were altered in *tga1/tga4* double mutant (Alvarez et al. 2014).

HRS1 (HYPERSENSITIVITY TO LOW PI-ELICITED PRIMARY ROOT SHORTENING 1), is rapidly induced by nitrate, in an NPF6.3- and NLP7-dependent manner, regulating a genomic response involved in the repression of primary root growth, specifically under nitrate provision (Medici et al. 2015). The LBD37/LBD38/LBD39 transcription factors (LATERAL ORGAN BOUNDARY DOMAIN) were shown to regulate many N-responsive genes (Rubin et al. 2009). They are induced by nitrate, repressing several *NRTs* genes (*NPF6.3*, *NRT2.1*, *NRT2.2*, and *NRT2.5*), as well as genes involved in nitrate assimilation, indicating that they may be involved in the feedback down-regulation by high N supply. LBD39 induction by high N supply was found to depend on phosphorylated form of NPF6.3, suggesting that it acts downstream of NPF6.3 to trigger repression of *NRT2.1* expression (Bouguyon et al. 2015). As LBDs, SLP9 also modulates the expression of many genes responsible for nitrate transport and assimilation (Krouk et al. 2010). NAC protein family members (NAC1, NAC2 and NAC4) were recognized as key regulatory element controlling a nitrate-responsive network in an auxin-related pathway, by regulating lateral root growth in the context of the nitrate response (Xie et al. 2000, He et al. 2005, Vidal et al. 2013). TaNAC2 is involved in grain nitrate accumulation and seed vigor in wheat through TaNRT2.5 gene expression regulation (Li et al. 2020).

MicroRNAs (miRNAs) are molecules known to regulate cell-signaling mechanisms in plants, controlling key targets at post-transcriptional and/or translational levels. Many of them act as key regulators in N metabolism, acting on the expression of specific genes for N uptake (Zuluaga et al. 2019). Members of the miR169 family are reported in many species to regulate the expression of N transporters under N-starvation conditions. In Arabidopsis, miR169 targets NFY (NUCLEAR FACTOR Y), a ubiquitous transcription factor that binds to the promoter regions and regulates the nitrate transporters *AtNRT2.1/AtNPF6.3* (Zhao et al. 2011). A study under N deficiency and resupply in maize highlighted that over 100 conserved miRNAs were

differentially expressed. In particular, miR169, miR1214, miR2199, miR398, miR408, and miR827 families might participate in the regulation of N metabolism in maize (Yang et al. 2019). Analysis of miRNA expression in root tissues of several wheat genotypes showed that miR159a, miR159b, miR399, and miR408 are differentially expressed in response to N availability (Sinha et al. 2015).

A research showed that long noncoding RNAs (lncRNAs) are crucial regulators of nitrate signaling network. Liu et al. (2019) identified 695 lncRNAs containing 480 previously known and 215 novel lncRNAs in Arabidopsis seedling roots. Six of these lncRNAs were significantly altered after nitrate treatment. Among these, the lncRNA showing the highest nitrate induction, T5120, was induced by nitrate directly. Overexpression of T5120 in Arabidopsis promoted the response to nitrate, enhanced nitrate assimilation and improved biomass and root development (Liu et al. 2019). Additionally, T5120 was regulated by NLP7 through direct binding to the T5120 promoter region. *nlp7* mutant had its phenotype restored by T5120 overexpression, indicating that T5120 functions in the nitrate regulatory pathway mediated by NLP7. Moreover, expression of T5120 was regulated by NRT1.1, suggesting that T5120 is modulated by NRT1.1 to regulate nitrate regulation. So, these results indicated that lncRNA T5120 plays an important role in nitrate regulation (Liu et al. 2019).

Mobile signals involved in the regulation of root N acquisition by the N status of the plant have been identified. HY5 (ELONGATED HYPOCOTYL 5), a light-responsive bZIP TF found in Arabidopsis, works as a shoot-to-root mobile signal that activates AtNRT2.1 expression and root nitrate uptake in response to changes in the light (Chen et al. 2016). CEPs (C-TERMINALLY ENCODED PEPTIDES) have been shown to act through CEP receptors (CEPRs) to control N signaling and lateral root development (Talestki et al. 2018). These peptides are root-to-shoot mobile signals that trigger expression of two glutaredoxin genes (CEPD1 and CEPD2) in response to N starvation. Both CEPD1 and CEPD2 travel back to enhance *NRT2.1* expression and lateral root growth, specifically in nitrate-rich zones, to ensure N acquisition (Ohkubo et al. 2017, Xuan et al. 2017).

Several studies have suggested that chromatin modifications could play an important role in the response to several abiotic constraints, such as N contrasting conditions in the soil. Using high N-insensitive mutants (*hni9-1*) it was possible to propose a systemic feedback repression of *NRT2.1* by high N supply, correlated with an enrichment of trimethylation of the Lys27 of the histone H3 (H3K27me3) at the *NRT2.1* locus (Widiez et al. 2011). All NLPs share a PB1 domain (Phox and Bem1 domain at the carboxy terminal), necessary for full induction of nitrate-dependent expression of target genes in planta via homo- and hetero-oligomerization of chromosomal DNA in the presence of nitrate (Konishi and Yanagisawa 2019). PRC2 (Polycomb Repressive Complex 2), a major regulator of gene expression in eukaryotes, catalyzes the repressive chromatin mark H3K27me3, which leads to very low

expression of target genes, such as *NRT2.1* (Zhang et al. 2007, Roudier et al. 2011). A mutation in a PRC2 subunit, leading to a loss of H3K27me3 at *NRT2.1*, results in an increase of *NRT2.1* expression. This indicates that PRC2 controls the *NRT2.1* expression in a context of upregulation (Bellegarde et al. 2018). Another H3K27me3 methylation recruiter was identified in rice. Increased N supply enhanced transcription and abundance of the rice APETALA2-domain transcription factor encoded by an *NGR5* (NITROGEN- MEDIATED TILLER GROWTH RESPONSE 5) allele. The modulation of H3K27me3 methylation via *NGR5*-dependent recruitment of PRC2 enables enhanced grain yield in elite rice varieties despite reduced N fertilizer inputs (Wu et al. 2020).

Knock-out mutation of *NPF6.3* does not lead to an impaired PNR (Wang et al. 2009), suggesting the presence of other nitrate sensors in plants. Since the sensing function of *NPF6.3* depends on the phosphorylation at T101 (Ho et al. 2009, Bouguyon et al. 2015), post-translational modifications of *NRT2.1* could be involved in the sensing function of *NRT2.1*. In fact, phosphorylation sites have been found in *NRT2.1* in two phospho-proteomic approaches in response to N (Engelsberger and Schulze 2012, Menz et al. 2016). Among them, *NRT2.1* seems to be phosphorylated at S28 when plants are N starved and rapidly dephosphorylated upon nitrate resupply (Engelsberger and Schulze 2012). Phosphorylation of *NRT2.1* at S21, driven by *NURK1* (NITRATE UPTAKE REGULATORY KINASE1), resulted in the inactivation of *NRT2.1*, decreasing the interaction with *NRT3.1*. Oppositely, phosphorylation of *NRT2.1* at the S28 residue enhanced the *NRT2.1/NRT3.1* interaction (Li et al. 2020). Experiments with transgenic plants with truncated forms of *NRT2.1* revealed an essential sequence for activity, located between the residues 494-513. More specifically, a S501 phosphorylation was shown as an essential mechanism for the regulation of *NRT2.1* activity (Jacquot et al. 2019). N-terminal region of *NRT2.1* seems to be modified in response to N availability. It suggests that posttranslational modifications through phosphorylation are involved to control *NRT2.1* activity in response to environmental cues.

1.5 Conclusion

Due to the environmental and economic consequences of indiscriminate and excessive use of N fertilizer, there is an increasing need for more N efficient crops suited for sustainable agriculture. Being nitrate the principal inorganic N source available in the soil, there has been a notable research effort to understand plant nitrate uptake, a key factor in NUE, and the molecular players that modulate it. This review, besides to explicate the environmental impacts surrounding the use of N fertilizer, aimed to display the wide view of the NRTs identified so far, and their regulation on different levels in response to N availability. Also, it was presented the recent advances in the nitrate signaling network.

HATS transporters, responsible for the larger proportion of nitrate uptake, are regulated in different levels by N-plant status and nitrate availability (Laugier et al. 2012, Nacry et al. 2013, Jacquot et al. 2019). Chromatin modifications (Bellegarde et al. 2018, Wu et al. 2020) and phospho-proteome analysis (Engelsberger and Schulze 2012, Menz et al. 2016, Jacquot et al. 2019, Li et al. 2020) recently seem to drag much attention among researchers. There is the huge potential to manipulate TFs targeting to improve NUpE in crops since those regulators can control a cascade of genes involved in nitrate uptake, rather than modulating a single or handful of NRTs genes simultaneously (Cao et al. 2017, Ge et al. 2019, Alfatih et al. 2020, Wu et al. 2020). Thus, the knowledge acquired in model plants about molecular players and regulatory factors interacting with NRTs involved in nitrate uptake could be used to improve NUE in crops. It emerges as an attractive step to allow eco-friendly agriculture to support the growth of the human population and food demand.

References

- ALFATIH A, WU J, ZHANG ZS, et al. (2020) Rice NIN-LIKE PROTEIN 1 rapidly responds to nitrogen deficiency and improves yield and nitrogen use efficiency. *BioRxiv*. DOI: 10.1101/2020.02.22.961193
- ALVAREZ JM, VIDAL EA, and GUTIÉRREZ RA (2012) Integration of local and systemic signaling pathways for plant N responses. *Curr Opin Plant Biol*, 15:185–191. DOI: 10.1016/j.pbi.2012.03.009
- ALVAREZ JM, RIVERAS E, VIDAL EA, et al. (2014) Systems approach identifies TGA1 and TGA4 transcription factors as important regulatory components of the nitrate response of *Arabidopsis thaliana* roots. *Plant J*, 80: 1–13. DOI: 10.1111/tpj.12618
- ALVAREZ JM, MOYANO TC, ZHANG T, et al. (2019) Local changes in chromatin accessibility and transcriptional networks underlying the nitrate response in *Arabidopsis* roots. *Mol Plant*, 12: 1545–1560. DOI: 10.1016/j.molp.2019.09.002
- AMEZIANE R, BERNHARD K, and LIGHTFOOT D (2000) Expression of the bacterial *gdhA* gene encoding a NADPH glutamate dehydrogenase in tobacco affects plant growth and development. *Plant Soil*, 22: 147–157. DOI: 10.1023/A:1004794000267
- ARAKI R and HASEGAWA H (2006) Expression of rice (*Oryza sativa* L.) genes involved in high-affinity nitrate transport during the period of nitrate induction. *Breed Sci*, 56: 295–302. DOI: 10.1270/jsbbs.56.295
- AUSTIN AT, BUSTAMANTE MMC, NARDOTO GB, et al. (2013) Latin America's Nitrogen Challenge. *Science*, 340(6129):149. DOI: 10.1126/science.1231679
- BALASUBRAMANIAN V, ALVES B, AULAKH MS, et al. (2004) Crop environmental and management factors affecting nitrogen use efficiency. In, "Agriculture and nitrogen cycle. Assessing the impact of nitrogen fertilizer use on food production and the environment". *Scope*, 65: 19–33.
- BARBIER-BRYGOO H, DE ANGELI A, FILLEUR S, et al. (2011) Anion channels/transporters in plants: from molecular bases to regulatory networks. *Annu Rev Plant Biol*, 62: 25–51. DOI: 10.1146/annurev-arplant-042110-103741
- BELLEGARDE F, GOJON A, and MARTIN A (2017) Signals and players in the transcriptional regulation of root responses by local and systemic N signaling in *Arabidopsis thaliana*. *J Exp Bot*, 68(10): 2553–2565. DOI: 10.1093/jxb/erx062
- BELLEGARDE F, HERBERT L, SÉRÉ D et al. (2018) Polycomb Repressive Complex 2 attenuates the very high expression of the *Arabidopsis* gene *NRT2.1*. *Sci Rep*, 8: 7905. DOI: 10.1038/s41598-018-26349-w
- BHATTACHARYA A (2019) Nitrogen-use efficiency under changing climatic conditions. *Changing climate and resource use efficiency in plants*. Chapter 4, pp: 181–240. DOI: 10.1016/C2017-0-04681-5
- BILLEN G, GARNIER J, and LASSALET L (2013) The nitrogen cascade from agricultural soils to the sea: Modelling N transfers at regional watershed and global scales. *Philos Trans Royal Soc*, 368(1621). DOI: 10.1098/rstb.2013.0123
- BOBBINK R, HICKS K, GALLOWAY J, et al. (2010) Global assessment of nitrogen deposition effects on terrestrial plant diversity: a synthesis. *Ecol Appl*, 20: 30–59. DOI: 10.1890/08-1140.1
- BOUGUYON E, BRUN F, MEYNARD D, et al. (2015) Multiple mechanisms of nitrate sensing by *Arabidopsis* nitrate transceptor *NRT1.1*. *Nature Plants*, 1: 15015. DOI: 10.1038/nplants.2015.15
- BOUWMAN AF, BEUSEN A, and BILLEN G (2009) Human alteration of the global nitrogen and phosphorus soil balances for the period 1970–2050. *Global Biogeochem Cy*, 23: 1–15. DOI: 10.1029/2009GB003576
- BOUWMAN L, GOLDEWIJK KK, VAN DER HOEK KW, et al. (2013) Exploring global changes in nitrogen and phosphorus cycles in agriculture induced by livestock production over the 1900–2050 period. *Proc Natl Acad Sci USA*, 110(52): 21195. DOI: 10.1073/pnas.1012878108
- BRUINSMA J (2017) *World Agriculture: Towards 2015/2030: An FAO Study*; Routledge: Abingdon, UK, 2017.
- BURNEY JA, DAVIS SJ, and LOBELL DB (2010) Greenhouse gas mitigation by agricultural intensification. *Proc Natl Acad Sci USA*, 107: 12052–12057. DOI: 10.1073/pnas.0914216107
- BURT TP, HOWDEN NJK, WORRALL F, and WHELAN MJ (2008) Importance of long-term monitoring for detecting environmental change: Lessons from a lowland river in south east England. *Biogeosciences*, 5:1529–1535. DOI: 10.5194/bg-5-1529-2008
- BUTTERBACH-BAHL K, NEMITZ E, ZAEHLE S, et al. (2011) Nitrogen as threat to European greenhouse balance. In: *The European Nitrogen Assessment: Sources, Effects and Policy Perspectives*. Cambridge University Press, pp. 434–462. ISBN: 9780511976988

- CAI CH, WANG JY, ZHU YG, et al. (2008) Gene structure and expression of the high-affinity nitrate transport system in rice roots. *J Integr Plant Biol*, 50: 443-451. DOI: 10.1111/j.1744-7909.2008.00642.x.
- CAO HR, QI SD, SUN MW, et al. (2017) Overexpression of the maize ZmNLP6 and ZmNLP8 can complement the arabidopsis nitrate regulatory mutant nlp7 by restoring nitrate signaling and assimilation. *Front Plant Sci*, 8: 1703. DOI: 10.3389/fpls.2017.01703
- CAMPBELL BM, BEARE DJ, BENNETT EM, et al. (2017) Agriculture production as a major driver of the Earth system exceeding planetary boundaries. *Ecol Soc*, 22(4): 8. DOI: 10.5751/ES-09595-220408
- CANFIELD DE, GLAZER AN, and FALKOWSKI PG (2010) The evolution and future of earth's nitrogen cycle. *Science*, 330(6001): 192-196. DOI: 10.1126/science.1186120
- CASSMAN KG, DOBERMANN D, and WALTERS DT (2002) Agroecosystems, nitrogen-use efficiency, and nitrogen management. *Ambio*, 31: 132-140. DOI: 10.1579/0044-7447-31.2.132
- CASTAINGS L, CAMARGO A, POCHOLLE D, et al. (2009) The nodule inception-like protein 7 modulates nitrate sensing and metabolism in Arabidopsis. *Plant J*, 57: 426-435. DOI: 10.1111/j.1365-313X.2008.03695.x
- CEREZO M, TILLARD P, FILLEUR S, et al. (2001) Major alterations of the regulation of root NO₃⁻ uptake are associated with the mutation of Nrt2.1 and Nrt2.2 genes in Arabidopsis. *Plant Physiol*, 127: 262-271. DOI: 10.1104/pp.127.1.262
- CIAMPITTI IA and VYN TJ (2014) Understanding global and historical nutrient use efficiencies for closing maize yield gaps. *Agron J*, 106: 2107-2117. DOI: 10.2134/agronj14.0025
- CHEN X, YAO Q, GAOET X, et al. (2016) Shoot-to-root mobile transcription factor HY5 coordinates plant carbon and nitrogen acquisition. *Curr Biol*, 26: 640-646. DOI: 10.1016/j.cub.2015.12.066
- CHICHKOVA S, ARELLANO J, VANCE CP, and HERNÁNDEZ G (2001) Transgenic tobacco plants that overexpress alfalfa NADH-glutamate synthase have higher carbon and nitrogen content. *J Exp Bot*, 52: 2079-2087. DOI: 10.1093/jexbot/52.364.2079
- CHIVENG P, VANLAUWE B, GENTILE R, and SIX J (2011) Comparison of organic versus mineral resource effects on short-term aggregate carbon and nitrogen dynamics in a sandy soil versus a fine textured soil. *Agric Ecosyst Environ*, 140: 361-371. DOI: 10.1016/j.agee.2010.12.004
- CHOPIN F, ORSEL M, DORBE MF, et al. (2007). The Arabidopsis ATNRT2.7 nitrate transporter controls nitrate content in seeds. *Plant Cell*, 19:1590-1602. DOI: 10.1105/tpc.107.050542
- CLARISSE L, CLERBAUX C, DENTENER F, et al. (2009) Global ammonia distribution derived from infrared satellite observations. *Nat Geosci*, 2: 479-483. DOI: 10.1038/ngeo551
- CREWS TE and PEOPLES MB (2004) Legume versus fertilizer sources of nitrogen: Ecological tradeoffs and human needs. *Agric Ecosyst Environ*, 102: 279-297. DOI: 10.1016/j.agee.2003.09.018
- CRUTZEN PJ, MOSIER AR, SMITH KA, and WINIWARTER W (2007) N₂O release from agro-biofuel production negates global warming reduction by replacing fossil fuels. *Atmos Chem Phys*, 8: 389-395. hal-00296423
- DAVID LC, DECHORGNAT J, BERQUIN P, et al. (2014) Proanthocyanidin oxidation of Arabidopsis seeds is altered in mutant of the high-affinity nitrate transporter NRT2.7. *J Exp Bot*, 65: 885-893. DOI: 10.1093/jxb/ert481
- DAVID LC, GIRIN T, FLEURISSON E, et al. (2019) Developmental and physiological responses of *Brachypodium distachyon* to fluctuating nitrogen availability. *Sci Rep*, 9: 3824. DOI: 10.1038/s41598-019-40569-8 <https://doi.org/10.1038/s41598-019-40569-8>
- DAVIDSON EA (2009) The contribution of manure and fertilizer nitrogen to atmospheric nitrous oxide since 1860. *Nat Geosci*, 2: 659-662. DOI: 10.1038/ngeo608
- DE ANGELI A, MONACHELLO D, EPHRITIKHINE G, et al. (2006) The nitrate/proton antiporter AtCLCa mediates nitrate accumulation in plant vacuoles. *Nature*, 442: 939-942. DOI: 10.1038/nature05013
- DECHORGNAT J, NGUYEN CT, ARMENGAUD P, et al. (2011) From the soil to the seeds: The long journey of nitrate in plants. *J Exp Bot*, 62:1349-1359. DOI: 10.1093/jxb/erq409
- DECHORGNAT J, PATRIT O, KRAPP A, et al. (2012) Characterization of the *Nrt2.6* Gene in Arabidopsis thaliana: A Link with Plant Response to Biotic and Abiotic Stress. *PLoS ONE* 7(8): e42491. DOI: 10.1371/journal.pone.0042491
- DECHORGNAT J, FRANCIS KL, DHUGGA KS, et al. (2018) Root Ideotype Influences Nitrogen Transport and Assimilation in Maize. *Front. Plant Sci*, 9:531. DOI: 10.3389/fpls.2018.00531
- DENTENER F, DREVET J, LAMARQUE JF, et al. (2006) Nitrogen and sulfur deposition on regional and global scales: A multi-model evaluation. *Global Biogeochem Cy*, 20: GB4003. DOI: 10.1029/2005GB002672

- DE VRIES W, KROS J, KROEZE CP, and SEITZINGER SP (2013) Assessing planetary and regional nitrogen boundaries related to food security and adverse environmental impacts. *Curr Opin Env Sust*, 5(3-4): 392-402. DOI: 10.1016/j.cosust.2013.07.004
- DHUGGA KS and WAINES JG (1989) Analysis of nitrogen accumulation and use in bread and durum-wheat. *Crop Science*, 29: 1232-1239. DOI: 10.2135/cropsci1989.0011183X0029000500029x
- DIAZ R, SELMAN M, and CHIQUE C (2010) Global Eutrophic and Hypoxic Coastal Systems. World Resources Institute. Eutrophication and Hypoxia: Nutrient Pollution in Coastal Waters. <https://www.wri.org/resources/data-sets/eutrophication-hypoxia-map-data-set>
- DISE NB, ASHMORE M, BELYAZID S, et al. (2011) Nitrogen as a threat to European terrestrial biodiversity. Chapter 20, in "The European Nitrogen Assessment". Cambridge University Press, pp: 463-494.
- DIRZO R and RAVEN PH (2003) Global state of biodiversity and loss. *Annu Rev Env Resour*, 28: 137-167. DOI: 10.1146/annurev.energy.28.050302.105532
- DOBERMANN A (2007) Nutrient use efficiency - measurement and management. In "IFA International Workshop on Fertilizer Best Management Practices", p: 1-28.
- DOS SANTOS TB, LIMA JE, FELICIO MS, et al. (2017) Genome-wide identification, classification and transcriptional analysis of nitrate and ammonium transporters in *Coffea*. *Genet Mol Biol*, 40(1): 346-359. DOI: 10.1590/1678-4685-GMB-2016-0041
- ENGELSBERGER WR and SCHULZE WX (2012) Nitrate and ammonium lead to distinct global dynamic phosphorylation patterns when resupplied to nitrogen-starved *Arabidopsis* seedlings. *Plant J*, 69: 978-995. DOI: 10.1111/j.1365-313X.2011.04848.x
- ERISMAN JW (2008) How a century of ammonia synthesis changed the world. *Nat Geosci*, 1: 636-639. DOI: 10.1038/ngeo325
- ERISMAN JW, GALLOWAY JN, SEITZINGER S, et al. (2013) Consequences of human modification of global nitrogen cycle. *Philos Trans R Soc Lond B Biol Sci*, 368(1621). DOI: 10.1098/rstb.2013.0116
- ESPEN L, NOCITO FF, and COCUCCI M (2004) Effect of NO₃⁻ transport and reduction on intracellular pH: an in vivo NMR study in maize roots. *J Exp Bot*, 55: 2053-2061. DOI: 10.1093/jxb/erh231
- FAGERIA NK and BALIGAR VC (2005) Enhancing nitrogen use efficiency in crop plants. *Adv Agron*, 88: 97-185. DOI: 10.1016/S0065-2113(05)88004-6
- FAN X, JIA L, and LI Y (2007) Comparing nitrate storage and remobilization in two rice cultivars that differ in their nitrogen use efficiency. *J Exp Bot*, 58(7): 1729-1740. DOI:10.1093/jxb/erm033
- FAN XR, FENG HM, TAN YW, et al. (2016a) A putative 6-transmembrane nitrate transporter OsNRT1.1b plays a key role in rice under low nitrogen. *J Integr Plant Biol*, 58(6): 590-599. DOI: 10.1111/jipb.12382
- FAN XR, TANG Z, TAN Y, et al. (2016b) Overexpression of a pH-sensitive nitrate transporter in rice increases crop yields. *Proc Natl Acad Sci USA*, 113(26): 7118-7123. DOI: 10.1073/pnas.1525184113
- FAN X, NAZ M, FAN X, et al. (2017) Plant nitrate transporters: from gene function to application. *J Exp Bot*, 68(10): 2463-2475. DOI:10.1093/jxb/erx011
- FENG HM, YAN M, FAN XR, et al. (2011) Spatial expression and regulation of rice high-affinity nitrate transporters by nitrogen and carbon status. *J Exp Bot*, 62(7): 2319-2332. DOI: 10.1093/jxb/erq403
- FILLEUR S and DANIEL-VEDELE F (1999) Expression analysis of a high-affinity nitrate transporter isolated from *Arabidopsis thaliana* by differential display. *Planta*, 207(3): 461-469. DOI: 10.1007/s004250050505
- FILLEUR S, DORBE MF, CEREZO M, et al. (2001) An *Arabidopsis* T-DNA mutant affected in Nrt2 genes is impaired in nitrate uptake. *FEBS Lett*, 489: 220-224. DOI: 10.1016/S0014-5793(01)02096-8
- FOLEY JA, DEFRIES R, ASNER GP, et al. (2005) Global consequences of land use. *Science*, 309(5734): 570-574. DOI: 10.1126/science.1111772
- FOLEY JA, RAMANKUTTY N, BRAUMAN KA, et al. (2011) Solutions for a cultivated planet. *Nature*, 478: 337-342. DOI: 10.1038/nature10452
- FOLLETT JR, FOLLETT RF, and HERZ W C (2010) Environmental and Human Impacts of Reactive Nitrogen. *Adv Nitr Managem Water Qual, SWCS*.
- FORSTER P, RAMASWAMY V, ARTAXO P, et al. (2007) Chapter 2: Changes in Atmospheric Constituents and in Radiative Forcing. In: *Climate Change 2007: The Physical Science Basis*. Cambridge University Press, pp:129-234. ISBN 978-0521-70596-7
- FOWLER D, COYLE M, SKIBA U, et al. (2013) The global nitrogen cycle in the twenty-first century. *Phil Trans R Soc B*, 368: 20130164. DOI: 10.1098/rstb.2013.0164

- FREDES I, MORENO S, DÍAZ FP, and GUTIÉRREZ RA (2019) Nitrate signaling and the control of Arabidopsis growth and development. *Curr Opin Plant Biol*, 47: 112-118. DOI: 10.1016/j.pbi.2018.10.004
- FRENEY JR, TREVITT ACF, DE DATTA SK, et al. (1990) The interdependence of ammonia volatilization and denitrification as nitrogen loss processes in flooded rice fields in the Philippines. *Biol Fertil Soils*, 9:31-36. DOI: 10.1007/BF00335858
- FU Y, YI H, BAO J, and GONG J (2015) LeNRT2.3 functions in nitrate acquisition and long-distance transport in tomato. *FEBS Letters*, 589: 1072-1079. DOI: 10.1016/j.febslet.2015.03.016
- GALLOWAY JN, DENTENER FJ, CAPONE DG, et al. (2004) Nitrogen Cycles: Past, Present, and Future. *Biogeochem*, 70: 153-226. DOI: 10.1007/s10533-004-0370-0
- GALLOWAY JN, TOWNSEND AR, ERISMAN JW, et al. (2008) Transformation of the nitrogen cycle: recent trends, questions, and potential solutions. *Science*, 320(5878): 889-892. DOI: 10.1126/science.1136674
- GARNETT TP, SHABALA SN, SMETHURST PJ, and NEWMAN IA (2003) Kinetics of ammonium and nitrate uptake by eucalyptus roots and associated proton fluxes measured using ion selective microelectrodes. *Funct Plant Biol*, 30: 1165-1176. DOI: 10.1071/fp03087
- GARNETT T, CONN V, and KAISER VN (2009) Root based approaches in improving nitrogen use efficiency in plants. *Plant Cell Environ*, 32:1272-1283. DOI: 10.1111/j.1365-3040.2009.02011.x
- GARNETT T, CONN V, PLETT D, et al. (2013) The response of the maize nitrate transport system to nitrogen demand and supply across the lifecycle. *New Phytol*, 198: 82-94. DOI: 10.1111/nph.12166
- GE M, WANG Y, LIU Y, et al. (2019) The NIN-like protein 5 (ZmNLP5) transcription factor is involved in modulating the nitrogen response in maize. *Plant J*, 5: 1-16. DOI: 10.1111/tpj.14628
- GEIGER D, MAIERHOFER T, AL-RASHEID KA, et al. (2011) Stomatal closure by fast abscisic acid signaling is mediated by the guard cell anion channel SLAH3 and the receptor RCAR1. *Sci Signal*, 4(173): ra32. DOI: 10.1126/scisignal.2001346
- GELLI M, DUO Y, KONDA AR, et al. (2014) Identification of differentially expressed genes between sorghum genotypes with contrasting nitrogen stress tolerance by genome-wide transcriptional profiling. *BMC Genomics*, 15: 179. DOI: 10.1186/1471-2164-15-179
- GIRONDE A, ETIENNE P, TROUVERIE J, et al. (2015) The contrasting N management of two oilseed rape genotypes reveals the mechanisms of proteolysis associated with leaf N remobilization and the respective contributions of leaves and stems to N storage and remobilization during seed filling. *BMC Plant Biol* 15: 59. DOI: 10.1186/s12870-015-0437-1
- GLASS ADM, SHAFF JE, and KOCHIAN LV (1992). Studies of the uptake of nitrate in barley. IV. Electrophysiology. *Plant Physiol*, 99: 456-463. DOI: 10.1104/pp.99.2.456
- GLASS ADM, BRITTO DT, KAISER BN, et al. (2002) The regulation of nitrate and ammonium transport systems in plants. *J Exp Bot*, 53: 855-864. DOI: 10.1093/jexbot/53.370.855
- GLASS ADM (2003) Nitrogen use efficiency of crop plants: physiological constraints upon nitrogen absorption. *Crit Rev Plant Sci*, 22(5): 453-470. DOI: 10.1080/07352680390243512
- GRACE PR, COLUNGA-GARCIA M, GAUGE SH, et al. (2006) The potential impact of agricultural management and climate change on soil organic carbon resources in terrestrial ecosystems of the North Central region of the United States. *Ecosystems*, 9:816-827. DOI: 10.1007/s10021-004-0096-9
- GRIFFITHS M and YORK LM (2020) Targeting root ion uptake kinetics to increase plant productivity and nutrient use efficiency. *Plant Physiol*, 182(4): 1854-1868. DOI: 10.1104/pp.19.01496
- GRIZZETTI B, BOURAOUI F, BILLEN G, et al. (2011) Nitrogen processes in aquatic ecosystems. Nitrogen as a threat to European water quality. In: *The European Nitrogen Assessment*, ed. M. A. Sutton, C. M. Howard, J. W. Erisman et al. Cambridge University Press, pp: 379-404.
- GOJON A, KROUK G, PERRINE-WALKER F, and LAUGIER E (2011) Nitrate transceptor(s) in plants. *J Exp Bot*, 62(7): 2299-2308. DOI: 10.1093/jxb/erq419
- GOOD AG, SHRAWAT AK, and MUENCH DG (2004) Can less yield more? Is reducing nutrient input into the environment compatible with maintaining crop production? *Trends Plant Sci*, 9: 1360-1385. DOI: 10.1016/j.tplants.2004.10.008
- GOOD A, JOHNSON S, DE PAUW M, et al. (2007) Engineering nitrogen use efficiency with alanine aminotransferase. *Can J Bot*, 85: 252-262. DOI: 10.1139/B07-019
- GU CS, SONG AP, ZHANG XX, et al. (2016) Cloning of chrysanthemum high affinity nitrate transporter family (CmNRT2) and characterization of CmNRT2.1. *Sci Rep*, 6: 23462. DOI: 10.1038/srep23462

- GUAN PZ, WANG RC, NACRY P, et al. (2014) Nitrate foraging by Arabidopsis roots is mediated by the transcription factor TCP20 through the systemic signaling pathway. *Proc Natl Acad Sci USA*, 111: 15267-15272. DOI: 10.1073/pnas.1411375111
- GUAN PZ, RIPOLL J-J, WANG R, et al. (2017) Interacting TCP and NLP transcription factors control plant responses to nitrate availability. *Proc Natl Acad Sci USA*, 114: 2419-2424. DOI: 10.1073/pnas.1615676114
- GUO B, LI Y, WANG S, et al. (2020) Characterization of the Nitrate Transporter gene family and functional identification of HvNRT2.1 in barley (*Hordeum vulgare* L.). *PLoS ONE*, 15(4): e0232056. DOI: 10.1371/journal.pone.0232056
- GUO JH, LIU XJ, ZHANG Y, et al. (2010) Significant acidification in major Chinese crop lands. *Science*, 327(5968): 1008-1010. DOI: 10.1126/science.1182570
- HABASH DZ, MASSIAH AJ, RONG HL, et al. (2001) The role of cytosolic glutamine synthetase in wheat. *Ann Appl Biol*, 138: 83-89. DOI: 10.1111/j.1744-7348.2001.tb00087.x
- HAN M, OKAMOTO M, and BEATTY PH (2015) The Genetics of Nitrogen Use Efficiency in Crop Plants. *Annu Rev Genet*, 49: 9.1-9.21. DOI: 10.1146/annurev-genet-112414-055037
- HE XJ, MU RL, CAO WH, et al. (2005) AtNAC2, a transcription factor downstream of ethylene and auxin signaling pathways, is involved in salt stress response and lateral root development. *Plant J*, 44(6): 903-916. DOI: 10.1111/j.1365-313X.2005.02575.x
- HERRERO M, THORNTON PK, NOTENBAERT AM, et al. (2010) Smart investments in sustainable food production: revisiting mixed crop-livestock systems. *Science*, 327(5967):822-825. DOI: 10.1126/science.1183725
- HERRIDGE DF, PEOPLES MB, and BODDEY RM (2008) Global inputs of biological nitrogen fixation in agricultural systems. *Plant Soil*, 311: 1-18. DOI: 10.1007/s11104-008-9668-3
- HIREL B, LE GOUIS J, NEY B, and GALLAIS A (2007) The challenge of improving nitrogen use efficiency in crop plants: towards a more central role for genetic variability and quantitative genetics within integrated approaches. *J Exp Bot*, 58: 2369-2387. DOI: 10.1093/jxb/erm097
- HO CH, LIN SH, HU HC, and TSAY YF (2009) CHL1 functions as a nitrate sensor in plants. *Cell*, 138(6): 1184-1194. DOI: 10.1016/j.cell.2009.07.004
- HO CH and TSAY YF (2010) Nitrate, ammonium, and potassium sensing and signaling. *Curr Opin Plant Biol*, 13(5): 604-610. DOI: 10.1016/j.pbi.2010.08.005
- HOBEN JP, GEHL RJ, MILLAR N, et al. (2011) Nonlinear nitrous oxide (N₂O) response to nitrogen fertilizer in on-farm corn crops of the US Midwest. *Glob Change Biol*, 17: 1140-1152. DOI: 10.1111/j.1365-2486.2010.02349.x
- HODGE A, ROBINSON D, and FITTER A (2000) Are microorganisms more effective than plants at competing for nitrogen? *Trends Plant Sci*, 5: 304-308. DOI: 10.1016/s1360-1385(00)01656-3
- HOULTON BZ, SIGMAN DM, and HEDIN LO (2006) Isotopic evidence for large gaseous nitrogen losses from tropical rainforests. *Proc Natl Acad Sci USA*, 103: 8745-8750. DOI: 10.1073/pnas.0510185103
- HU B, WANG W, OU S, et al. (2015) Variation in *NRT1.1B* contributes to nitrate-use divergence between rice subspecies. *Nature Genet*, 47: 834-838. DOI: 10.1038/ng.3337
- HUANG NC, LIU KH, LO HJ, and TSAY YF (1999) Cloning and functional characterization of an Arabidopsis nitrate transporter gene that encodes a constitutive component of low-affinity uptake. *Plant Cell*, 11: 1381-1392. DOI: 10.1105/tpc.11.8.1381
- IMSANDE J and TOURAINE B (1994) N demand and the regulation of nitrate uptake. *Plant Physiol*, 105: 3-7. DOI: 10.1104/pp.105.1.3
- ISHIKAWA S, ITO Y, SATO Y, et al. (2009) Two-component high-affinity nitrate transport system in barley: membrane localization, protein expression in roots and a direct protein-protein interaction. *Plant Biotechnol*, 26: 197-205. DOI: 10.5511/plantbiotechnology.26.197
- JACQUOT A, CHAPUT V, MAURIES A, et al. (2019) NRT2.1 phosphorylation prevents root high affinity nitrate uptake activity in *Arabidopsis thaliana*. *BioRxiv*. DOI: 10.1101/583542
- JAYNES DB, COLVIN TS, KARLEN DL, et al. (2001) Nitrate loss in subsurface drainage as affected by nitrogen fertilizer rate. *J Environ Qual*, 30: 1305-1314. DOI: 10.2134/jeq2001.3041305x
- KAWACHI T, SUNAGA Y, EBATO M, et al. (2006) Repression of nitrate uptake by replacement of Asp105 by asparagine in AtNRT3.1 in *Arabidopsis thaliana* L. *Plant Cell Physiol*, 47(10): 1437-1441. DOI: 10.1093/pcp/pcl010
- KANT S (2018) Understanding nitrate uptake, signaling and remobilization for improving plant nitrogen use efficiency. *Semin Cell Dev Biol*, 74: 89-96. DOI: 10.1016/j.semcdb.2017.08.034
- KEARNEY J (2010) Food consumption trends and drivers. *Philos Trans R Soc Lond B Biol Sci*, 365(1554): 2793-2807. DOI: 10.1098/rstb.2010.0149

- KIBA T, FERIA-BOURRELLIER AB, LAFOUGE F, et al. (2012) The Arabidopsis nitrate transporter NRT2.4 plays a double role in roots and shoots of nitrogen-starved plants. *Plant Cell*, 24: 245-258. DOI: 10.1105/tpc.111.092221
- KONISHI M and YANAGISAWA S (2013) Arabidopsis NIN-like transcription factors have a central role in nitrate signaling. *Nat Commun*, 4: 1617. DOI: 10.1038/ncomms2621
- KONISHI M and YANAGISAWA S (2019) The role of protein-protein interactions mediated by the PB1 domain of NLP transcription factors in nitrate-inducible gene expression. *BMC Plant Biol*, 19: 90. DOI: 10.1186/s12870-019-1692-3
- KOTUR Z, MACKENZIE N, RAMESH S, et al. (2012) Nitrate transport capacity of the Arabidopsis thaliana NRT2 family members and their interactions with AtNAR2.1. *New Phytol*, 194: 724–731. DOI: 10.1111/j.1469-8137.2012.04094.x
- KOTUR Z and GLASS ADM (2015) A 150 kDa plasma membrane complex of AtNRT2.5 and AtNAR2.1 is the major contributor to constitutive high-affinity nitrate influx in Arabidopsis thaliana. *Plant Cell Environ*, 38: 1490-1502. DOI: 10.1111/pce.12496
- KRAPP A, FRAISIER V, SCHEIBLE WR, et al. (1998) Expression studies of Nrt2:1NP, a putative high-affinity nitrate transporter: evidence for its role in nitrate uptake. *Plant J*, 14: 723-731. DOI: 10.1046/j.1365-313x.1998.00181.x
- KRAPP A, DAVID LC, CHARDIN C, et al. (2014) Nitrate transport and signaling in Arabidopsis. *J Exp Bot*, 65(3): 789–798. DOI: 10.1093/jxb/eru001.
- KROUK G, TILLARD P, and GOJON A (2006) Regulation of the high-affinity NO₃⁻ uptake system by NRT1.1-mediated NO₃⁻ demand signaling in Arabidopsis. *Plant Physiol*, 142(3): 1075-1086. DOI: 10.1104/pp.106.087510
- KROUK G, LACOMBE B, BIELACH A, et al. (2010) Nitrate-regulated auxin transport by NRT1.1 defines a mechanism for nutrient sensing in plants. *Dev Cell*, 18(6): 927-937. DOI: 10.1016/j.devcel.2010.05.008
- KRONZUCKER HJ, SIDDIQI MY, and GLASS ADM (1995) Kinetics of NO₃⁻ influx in spruce. *Plant Physiol*, 109: 319-326. DOI: 10.1104/pp.109.1.319
- LAUGIER E, BOUGUYON E, MAURIES A, et al. (2012) Regulation of high-affinity nitrate uptake in roots of Arabidopsis depends predominantly on posttranscriptional control of the NRT2.1/NAR2.1 transport system. *Plant Physiol*, 158: 1067-1078. DOI: 10.1104/pp.111.188532
- LEA PJ and AZEVEDO RA (2006) Nitrogen use efficiency. 1. Uptake of nitrogen from the soil. *Ann Appl Biol*, 149: 243-247. DOI: 10.1111/j.1744-7348.2006.00101.x
- LE GOUIS J, BEGHIN D, HEUMEZ E, and PLUCHARD P (2000). Genetic differences for nitrogen uptake and nitrogen utilization efficiencies in winter wheat. *Eur J Agron*, 12: 163-173. DOI: 10.1016/S1161-0301(00)00045-9
- LÉRAN S, VARALA K, BOYER JC, et al. (2014) A unified nomenclature of NITRATE TRANSPORTER 1/PEPTIDE TRANSPORTER family members in plants. *Trends Plant Sci*, 19: 5-9. DOI: 10.1016/j.tplants.2013.08.008
- LEZHNEVA L, KIBA T, FERIA-BOURRELLIER AB, et al. (2014) The Arabidopsis nitrate transporter NRT2.5 plays a role in nitrate acquisition and remobilization in nitrogen-starved plants. *Plant J*, 80: 230-241. DOI: 10.1111/tpj.12626
- LI H, HU B, and CHU C (2017) Nitrogen use efficiency in crops: lessons from Arabidopsis and rice. *J Exp Bot*, 68(10): 2477-2488. DOI: 10.1093/jxb/erx101
- LI W, WANG Y, OKAMOTO M, et al. (2007) Dissection of the AtNRT2.1: AtNRT2.2 inducible high-affinity nitrate transporter gene cluster. *Plant Physiol*, 143: 425-433. DOI: 10.1104/pp.106.091223
- LI W, HE X, CHEN Y, et al. (2020) A wheat transcription factor positively sets seed vigour by regulating the grain nitrate signal. *New Phytol*, 225: 1667-1680. DOI: 10.1111/nph.16234
- LI Z, WU XN, JAQUOT A, et al. (2020). A phospho-switch in the N-terminus of NRT2.1 affects nitrate uptake by controlling the interaction of NRT2.1 with NAR2.1. *BioRxiv*. DOI: 10.1101/2020.01.08.898254
- LIEBIG MA, VARVEL GE, DORAN JW, and WIENHOLD BJ (2002) Crop sequence and nitrogen fertilization effects on soil properties in the western Corn Belt. *Soil Sci Soc Am J*, 66: 596-601. DOI: 10.2136/sssaj2002.5960
- LIN CM, KOH S, STACEY G, et al. (2000) Cloning and functional characterization of a constitutively expressed nitrate transporter gene, OsNRT1, from rice. *Plant Physiol*, 122: 379-388. DOI: 10.1104/pp.122.2.379
- LITTLE DY, RAO H, OLIVA S, et al. (2005) The putative high-affinity nitrate transporter NRT2.1 represses lateral root initiation in response to nutritional cues. *Proc Natl Acad Sci USA*, 102: 13693-13698. DOI: 10.1073/pnas.0504219102

- LIU F, XU Y, CHANG K, et al. (2019) The long noncoding RNA T5120 regulates nitrate response and assimilation in Arabidopsis. *New Phytol*, 224: 117-131. DOI: 10.1111/nph.16038
- LIU KH, HUANG CY, and TSAY YF (1999) CHL1 is a dual-affinity nitrate transporter of Arabidopsis involved in multiple phases of nitrate uptake. *Plant Cell*, 11: 865-874. DOI: 10.1105/tpc.11.5.865
- LIU KH and TSAY YF (2003) Switching between the two action modes of the dual-affinity nitrate transporter CHL1 by phosphorylation. *EMBO J*, 22(5): 1005-1013. DOI: 10.1093/emboj/cdg118
- LIU KH, NIU Y, KONISHI M, et al. (2017) Discovery of nitrate–CPK–NLP signaling in central nutrient–growth networks. *Nature*, 545(7654): 311-316. DOI: 10.1038/nature22077
- LIU LH, FAN TF, SHI DX, et al. (2018) Coding-sequence identification and transcriptional profiling of nine AMTs and four NRTs from tobacco revealed their differential regulation by developmental stages, nitrogen nutrition, and photoperiod. *Front Plant Sci*, 9:210. DOI: 10.3389/fpls.2018.00210
- LIU X, HUANG D, TAO J, et al. (2014) Identification and functional assay of the interaction motifs in the partner protein OsNAR2.1 of the two-component system for high-affinity nitrate transport. *New Phytol*, 204: 74-80. DOI: 10.1111/nph.12986
- LIZAMA-GASCA MG, ESTRADA-TAPIA G, ESCALANTE-MAGAÑA CA, et al. (2020) Cloning and molecular characterization of CcNRT2.1/CcNAR2, a putative inducible high affinity nitrate transport system in *Capsicum chinense* Jacq. roots. *Trop Plant Biol*, 13(1): 73-90. DOI: 10.1007/s12042-019-09248-w
- LUPINI A, MERCATI F, ARANITI F, et al. (2016) NAR2.1/NRT2.1 functional interaction with NO_3^- and H^+ fluxes in high-affinity nitrate transport in maize root regions. *Plant Physiol Biochem*, 102: 107-114. DOI: 10.1016/j.plaphy.2016.02.022
- LU X, MAO Q, GILLIAM FS, et al. (2014) Nitrogen deposition contributes to soil acidification in tropical ecosystems. *Glob Chang Biol*, 20: 3790-3801. DOI: 10.1111/gcb.12665
- MAEDA Y, KONISHI M, KIBA T, et al. (2018) A NIGT1-centred transcriptional cascade regulates nitrate signaling and incorporates phosphorus starvation signals in Arabidopsis. *Nat Commun*, 9: 1-14. DOI: 10.1038/s41467-018-03832-6
- MAHLER RL and HARDER RW (1984) The influence of tillage methods, cropping sequence, and N rates on the acidification of a northern Idaho soil. *Soil Sci*, 137: 52-60.
- MAKITA Y, SHIMADA S, KAWASHIMA M, et al. (2015). MOROKOSHI: transcriptome database in *Sorghum bicolor*. *Plant Cell Physiol*, 56: e6. DOI: 10.1093/pcp/pcu187
- MALAGOLI P, LAINE P, LE DEUNFF E, et al. (2004) Modeling nitrogen uptake in oilseed rape cv Capitol during a growth cycle using influx kinetics of root nitrate transport systems and field experimental data. *Plant Physiol*, 134: 388-400. DOI: 10.1104/pp.103.029538
- MARCHIVE C, ROUDIER F, CASTAINGS L, et al. (2013) Nuclear retention of the transcription factor NLP7 orchestrates the early response to nitrate in plants. *Nat Commun*, 4: 1713. DOI: 10.1038/ncomms2650
- MARTINELLI LA and FILOSO S (2008) expansion of sugarcane ethanol production in Brazil: environmental and social challenges. *Ecol Appl*, 18(4): 885–898. DOI: 10.1890/07-1813.1
- MARSCHNER H (2012) Marschner's Mineral Nutrition of Higher Plants (Vol. 89). Academic Press, 672 p.
- MASCLAUX-DAUBRESSE C, DANIEL-VEDELE F, DECHORGNAT J, et al. (2010) Nitrogen uptake, assimilation and remobilization in plants: challenges for sustainable and productive agriculture. *Ann Bot*, 105: 1141-1157. DOI: 10.1093/aob/mcq028
- MASSEL K, CAMPBELL BC, MACE ES, et al. (2016) Whole genome sequencing reveals potential new targets for improving nitrogen uptake and utilization in *Sorghum bicolor*. *Front Plant Sci*, 7: 1544. DOI: 10.3389/fpls.2016.01544
- MEDICI A, MARSHALL-COLON A, RONZIER E, et al. (2015). AtNIGT1/HRS1 integrates nitrate and phosphate signals at the Arabidopsis root tip. *Nat. Commun*, 6: 6274. DOI: 10.1038/ncomms7274
- MENZ J, LI Z, SCHULZE WX, and LUDEWIG U (2016). Early nitrogen-deprivation responses in Arabidopsis roots reveal distinct differences on transcriptome and (phospho-) proteome levels between nitrate and ammonium nutrition. *Plant J*, 88: 717-734. DOI: 10.1111/tpj.13272
- MILLAR N, ROBERTSON GP, GRACE PR, et al. (2010) Nitrogen fertilizer management for nitrous oxide (N_2O) mitigation in intensive corn (Maize) production: an emissions reduction protocol for US Midwest agriculture. *Mitig Adapt Strat Gl*, 15: 185–204. DOI 10.1007/s11027-010-9212-7
- MILLENNIUM ECOSYSTEM ASSESSMENT (2005) Ecosystems and Human Well-being: Opportunities and Challenges for Business and Industry. World Resources Institute. ISBN: 1-56973-592-1
- MILLER AJ and CRAMER MD (2005) Root nitrogen acquisition and assimilation. *Plant Soil*, 274: 1-36. DOI: 10.1007/s11104-004-0965-1

- MILLER AJ, FAN X, ORSEL M, et al. (2007) Nitrate transport and signaling. *J Exp Bot*, 58: 2297-2306. DOI: 10.1093/jxb/erm066
- MOLDANOVA J, GRENNFELT P, JONSSON A, et al. (2011) Nitrogen as a threat to European air quality, in: *The European Nitrogen Assessment: Sources, Effects and Policy Perspectives*. Cambridge University Press, pp 405-433. DOI: 10.1017/CBO9780511976988.021
- MOLL RH, KAMPRATH EJ, and JACKSON WA (1982) Analysis and interpretation of factors which contribute to efficiency of nitrogen utilization. *Agron J*, 74: 562-564. DOI: 10.2134/agronj1982.00021962007400030037x
- MONACHELLO D, ALLOT M, OLIVA S et al. (2009) Two anion transporters AtCICa and AtCICe fulfil interconnecting but not redundant roles in nitrate assimilation pathways. *New Phytol*, 183: 88-94. DOI: 10.1111/j.1469-8137.2009.02837.x
- MORÈRE-LE PAVEN MC, VIAU L, HAMON A, et al. (2011) Characterization of a dual-affinity nitrate transporter MtNRT1.3 in the model legume *Medicago truncatula*. *J Exp Bot*, 62: 5595-5605. DOI: 10.1093/jxb/err243
- MOSIER A, DUXBURY J, FRENEY J, et al. (1998) Assessing and mitigating N₂O emissions from agricultural soils. *Clim Change*, 40: 7-38. DOI: 10.1023/A:1005386614431
- MOSIER A and KROEZE C (2000) Potential impact on the global atmospheric N₂O budget of the increased nitrogen input required to meet future global food demands. *Chemosphere* 2, 465-473. DOI: 10.1016/S1465-9972(00)00039-8
- MUELLER ND, GERBER JS, JOHNSTON M, et al. (2012) Closing yield gaps through nutrient and water management. *Nature*, 490: 254-257. DOI: 10.1038/nature11420
- MUNÕS S, CAZETTES C, FIZAMES C, et al. (2004) Transcript profiling in the chl1-5 mutant of *Arabidopsis* reveals a role of the nitrate transporter NRT1.1 in the regulation of another nitrate transporter, NRT2.1. *Plant Cell*, 16(9): 2433-2447. DOI: 10.1105/tpc.104.024380
- NACRY P, BOUGUYON E, and GOJON A (2013) Nitrogen acquisition by roots: physiological and developmental mechanisms ensuring plant adaptation to a fluctuating resource. *Plant Soil*, 370: 1-29. DOI: 10.1007/s11104-013-1645-9
- NEGI J, MATSUDA O, NAGASAWA T, et al. (2008) CO₂ regulator SLAC1 and its homologues are essential for anion homeostasis in plant cells. *Nature*, 452: 483-486. DOI: 10.1038/nature06720
- NINNEMANN O, JAUNIAUX JC, and FROMMER WB (1994) Identification of a high affinity NH₄⁺ transporter from plants. *EMBO J*, 13: 3464-3471. PMID: PMC395249
- NOGUERO M and LACOMBE B (2016) Transporters involved in root nitrate uptake and sensing by *Arabidopsis*. *Front Plant Sci*, 7: 1391. DOI: 10.3389/fpls.2016.01391
- O'BRIEN JA, VEGA A, BOUGUYON E, et al. (2016) Nitrate transport, sensing, and responses in plants. *Mol Plant*, 9: 837-856. DOI: 10.1016/j.molp.2016.05.004
- OHKUBO Y, TANAKA M, TABATA R, et al. (2017) Shoot-to-root mobile polypeptides involved in systemic regulation of nitrogen acquisition. *Nat Plants*, 3:1 7029. DOI: 10.1038/nplants.2017.29
- OKAMOTO M, VIDMAR JJ, and GLASS ADM (2003). Regulation of NRT1 and NRT2 gene families of *Arabidopsis thaliana*: responses to nitrate provision. *Plant Cell Physiol*, 44: 304-317. DOI: 10.1093/pcp/pcg036
- OKAMOTO M, KUMAR A, LI WB, et al. (2006) High-affinity nitrate transport in roots of *Arabidopsis* depends on expression of the NAR2-like gene AtNRT3.1. *Plant Physiol*, 140(3): 1036-1046. DOI: 10.1104/pp.105.074385
- ORSEL M, KRAPP A, and DANIEL-VEDELE F (2002) Analysis of the NRT2 nitrate transporter family in *Arabidopsis*. Structure and gene expression. *Plant Physiol*, 129: 886-896. DOI: 10.1104/pp.005280
- ORSEL M, CHOPIN F, LELEU O, et al. (2006) Characterization of a two-component high-affinity nitrate uptake system in *Arabidopsis*: Physiology and protein-protein interaction. *Plant Physiol*, 142(3): 1304-1317. DOI: 10.1104/pp.106.085209
- PARKER JL and NEWSTEAD S (2014) Molecular basis of nitrate uptake by the plant nitrate transporter NRT1.1. *Nature*, 507: 68-72. DOI: 10.1038/nature13116
- PATHAK RR, AHMAD A, LOCHAB S, and RAGHURAM N (2008) Molecular physiology of plant N-use efficiency and biotechnological options for its enhancement. *Curr Sci*, 94(11): 1394-1403.
- PATHAK RR, LOCHAB S, and RAGHURAM N (2011) Improving Plant Nitrogen-Use Efficiency in *Comprehensive Biotechnology*, 2nd Edition Elsevier. Oxford. 4, 209-218.
- PELLIZZARO A, CLOCHARD T, PLANCHET E, et al. (2015) Identification and molecular characterization of *Medicago truncatula* NRT2 and NAR2 families. *Physiol Plant*, 154: 256-269. DOI: 10.1111/ppl.12314
- PEÑA PA, QUACH T, SATO S, et al. (2017) Expression of the maize Dof1 transcription factor in wheat and sorghum. *Front Plant Sci*, 8: 434. DOI: 10.3389/fpls.2017.00434

- PEOPLES MB, MOSIER AR, and FRENEY JR (1995) Minimizing gaseous losses of nitrogen. In: "Nitrogen fertilization in the environment", p: 505-602. <http://hdl.handle.net/102.100.100/230910?index=1>
- PERCHLIK M and TEGEDER M (2017) Improving plant nitrogen use efficiency through alteration of amino acid transport processes. *Plant Physiol*, 175: 235-247. DOI: 10.1104/pp.17.00608
- PII Y, ALESSANDRINI M, DALL'OSTO L, et al. (2016) Time-resolved investigation of molecular components involved in the induction of NO₃⁻ High Affinity Transport System in Maize Roots. *Front Plant Sci*, 7:1657. DOI: 10.3389/fpls.2016.01657
- PLETT D, TOUBIA J, GARNETT T, et al. (2010) Dichotomy in the NRT gene families of dicots and grass species. *PLoS One*, 5: e15289. DOI: 10.1371/journal.pone.0015289
- PLETT DC, HOLTHAM LR, OKAMOTO M, and GARNETT TP (2018) Nitrate uptake and its regulation in relation to improving nitrogen use efficiency in cereals. *Semin Cell Dev Biol*, 74: 97-104. DOI:10.1016/j.semcdb.2017.08.027
- QUAGGIOTTI S, RUPERTI B, BORSA P, et al. (2003). Expression of a putative high-affinity NO₃⁻ transporter and of an H⁺-ATPase in relation to whole plant nitrate transport physiology in two maize genotypes differently responsive to low nitrogen availability. *J Exp Bot*, 54: 1023-1031. DOI: 10.1093/jxb/erg106
- QUESADA A, GALVÁN A, and FERNÁNDEZ E (1994) Identification of nitrate transporter genes in *Chlamydomonas reinhardtii*. *Plant J*, 5: 407-419. DOI: 10.1111/j.1365-3113x.1994.00407.x
- QUESADA A, KRAPP A, TRUEMAN LJ, et al. (1997) PCR-identification of a *Nicotiana plumbaginifolia* cDNA homologous to the high-affinity nitrate transporters of the crnA family. *Plant Mol Biol*, 34: 265-274. DOI: 10.1023/a:1005872816881
- RABALAIS NN, TURNER RE, and SCAVIA D (2002) Beyond science into policy: Gulf of Mexico hypoxia and the Mississippi river. *Bioscience*, 52: 129-142. DOI: 10.1641/0006-3568(2002)052[0129:BSIPGO]2.0.CO;2
- RAVISHANKARA AR, DANIEL JS, and PORTMANN RW (2009) Nitrous Oxide (N₂O): The Dominant Ozone-Depleting Substance Emitted in the 21st Century. *Science*, 326: 123-125. DOI: 10.1126/science.1176985
- REAY DS, DAVIDSON EA, SMITH KA, et al. (2012) Global agriculture and nitrous oxide emissions. *Nat Clim Change*, 2: 410-416. DOI: 10.1038/NCLIMATE1458
- REETZ H (2016) Fertilizers and their efficient use. IFA - International Fertilizer Industry Association. ISBN 979-10-92366-04-4
- REMANS T, NACRY P, PERVENT M, et al. (2006) The Arabidopsis NRT1.1 transporter participates in the signaling pathway triggering root colonization of nitrate rich patches. *Proc Natl Acad Sci USA*, 103: 19206-19211. DOI: 10.1073/pnas.0605275103
- RIVERAS E, ALVAREZ JM, VIDAL EA, et al. (2015) The calcium ion is a second messenger in the nitrate signaling pathway of Arabidopsis. *Plant Physiol*, 169: 1397-1404. DOI: 10.1104/pp.15.00961
- ROBERTSON GP and VITOUSEK PM (2009) Nitrogen in agriculture: balancing the cost of an essential resource. *Annu Rev Environ Resour*, 34: 97-125. DOI: 10.1146/annurev.environ.032108.105046
- ROCKSTRÖM J, STEFFEN W, NOONE K, et al. (2009) A safe operating space for humanity. *Nature*, 461(7263):472-475. DOI: 10.1038/461472a
- ROUDIER F, AHMED I, BÉRARD C, et al. (2011) Integrative epigenomic mapping defines four main chromatin states in Arabidopsis. *EMBO J*, 30(10): 1928-1938. DOI: 10.1038/emboj.2011.103
- RUBIN G, TOHGE T, MATSUDA F, et al. (2009). Members of the LBD family of transcription factors repress anthocyanin synthesis and affect additional nitrogen responses in Arabidopsis. *Plant Cell*, 21: 3567-3584. DOI: 10.1105/tpc.109.067041
- RUFFEL S, KROUK G, RISTOVA D, et al. (2011) Nitrogen economics of root foraging, transitive closure of the nitrate–cytokinin relay and distinct systemic signaling for N supply vs. demand. *Proc Natl Acad Sci USA*, 108: 18524-18529. DOI: 10.1073/pnas.1108684108
- SCHRÖDER JJ, SCHOLEFIELD D, CABRAL F, and HOFMAN G (2004) The effects of nutrient losses from agriculture on ground and surface water quality: the position of science in developing indicators for regulation. *Environ Sci Policy*, 7: 15-23. DOI: 10.1016/j.envsci.2003.10.006
- SCHRÖDER JL, ZHANG H, GIRMA H, et al. (2011) Soil acidification from long-term use of nitrogen fertilizers on winter wheat. *Soil Sci Soc Am J*, 75: 957-964. DOI: 10.2136/sssaj2010.0187
- SDSN - Leadership Council of the Sustainable Development Solutions Network (2015) Indicators and a monitoring framework for sustainable development goals. <http://unsdsn.org/resources>
- SEBILO M, MAYER B, NICOLARDOT B, et al. (2013) Long-term fate of nitrate fertilizer in agricultural soils. *Proc Natl Acad Sci USA*, 110: 18185-18189. DOI: 10.1073/pnas.1305372110

- SEGONZAC C, BOYER JC, IPOTESI E, et al. (2007) Nitrate efflux at the root plasma membrane: identification of an Arabidopsis excretion transporter. *Plant Cell*, 19: 3760-3777. DOI: 10.1105/tpc.106.048173
- SEKHON RS, LIN H, CHILDS KL, et al. (2011) Genome-wide atlas of transcription during maize development. *Plant J*, 66: 553-563. DOI: 10.1111/j.1365-313X.2011.04527.x
- SEIFFERT B, ZHOU Z, WALLBRAUN M, et al. (2004) Expression of a bacterial asparagine synthetase gene in oilseed rape (*Brassica napus*) and its effect on traits related to nitrogen efficiency. *Physiol Plant*, 121: 656-665. DOI: 10.1111/j.1399-3054.2004.00361.x
- SEITZINGER SP, HARRISON JA, DUMONT E, et al. (2005) Sources and delivery of carbon, nitrogen, and phosphorus to the coastal zone: An overview of Global Nutrient Export from Watersheds (NEWS) models and their application. *Global Biogeochem Cy*, 19: GB4S01. DOI: 10.1029/2005GB002606
- SELMAN M, SUGG Z, GREENHALGH S, and DIAZ R (2008) Eutrophication and hypoxia in coastal areas: a global assessment of the state of knowledge. WRI Report. <https://www.wri.org/publication/eutrophication-and-hypoxia-coastal-areas>
- SESTILI F, ROUPHAEL Y, CARDARELLI M, et al. (2018) Protein hydrolysate stimulates growth in tomato coupled with N-dependent gene expression involved in N assimilation. *Front Plant Sci*, 9: 1233. DOI: 10.3389/fpls.2018.01233
- SHRAWAT AK, CARROLL RT, DEPAUW M, et al. (2008) Genetic engineering of improved nitrogen use efficiency in rice by the tissue-specific expression of alanine aminotransferase. *Plant Biotechnol J*, 6: 722-732. DOI: 10.1111/j.1467-7652.2008.00351.x
- SIDDIQI MY, GLASS ADM, RUTH TJ and RUFTY TW (1990) Studies of the uptake of nitrate in barley. I: Kinetics of $^{13}\text{NO}_3^-$ influx. *Plant Physiol*, 93:1426-1432. DOI: 10.1104/pp.93.4.1426
- SINCLAIR TR and RUFTY TW (2012) Nitrogen and water resources commonly limit crop yield increases, not necessarily plant genetics. *Glob Food Secur*, 1: 94-98. DOI: 10.1016/j.gfs.2012.07.001
- SINGH U, LADHA JK, CASTILLO EG, et al. (1998) Genotypic variation in nitrogen use efficiency. I. Medium-and long-duration rice. *Field Crops Res*, 58: 35-53. DOI: 10.1016/S0378-4290(98)00084-7
- SINGH B (2018) Are nitrogen fertilizers deleterious to soil health? *Agronomy*, 8. DOI:10.3390/agronomy8040048
- SINHA SK, RANI M, BANSAL N, et al. (2015) Nitrate starvation induced changes in root system architecture, carbon: nitrogen metabolism, and miRNA expression in nitrogen-responsive wheat genotypes. *Appl Biochem Biotechnol*, 177: 1299-1312. DOI: 10.1007/s12010-015-1815-8
- SINHA VB, JANGAM AP, RAGHURAM N (2018) Biological determinants of crop N use efficiency and biotechnological avenues for improvement. In: Masso C, Bleeker A, Raghuram N, Bekunda M, Sutton M (eds) *Proceedings of the N2013*. Springer.
- SMIL V (1999) Nitrogen in crop production: An account of global flows. *Global Biogeochem Cy*, 13: 647-662. DOI: 10.1029/1999GB900015
- SMIL V (2002) Nitrogen and food production: proteins for human diets. *Ambio*, 31: 126-131. DOI: 10.1579/0044-7447-31.2.126
- SORGONÀ A, LUPINI A, MERCATI F, et al. (2011). Nitrate uptake along the maize primary root: an integrated physiological and molecular approach. *Plant Cell Environ*, 34: 1127-1140. DOI: 10.1111/j.1365-3040.2011.02311.x
- STAHL A, FRIEDT W, WITTKOP B, and SNOWDON RJ (2016) Complementary diversity for nitrogen uptake and utilization efficiency reveals broad potential for increased sustainability of oilseed rape production. *Plant Soil*, 400: 245-262. DOI: 10.1007/s11104-015-2726-8
- STEFFEN W, RICHARDSON K, ROCKSTRÖM J, et al. (2015) Planet boundaries: Guiding human development on a changing planet. *Science*, 347: 1259855. DOI: 10.1126/science.1259855
- STEINFELD H, GERBER P, WASSENAAR T, et al. (2006). *Livestock's Long Shadow – Environmental Issues and Options*. Food and Agriculture Organization. ISBN 92-5-105571-8.
- SUN J, BANKSTON JR, PAYANDEH J, et al. (2014) Crystal structure of the plant dual-affinity nitrate transporter NRT1.1. *Nature*, 507: 73-77. DOI: 10.1038/nature13074
- SUTTON MA, BLEEKER A, HOWARD CM, et al. (2013) Our Nutrient World: The challenge to produce more food and energy with less pollution. *Global Overview of Nutrient Management*. Centre for Ecology and Hydrology, Edinburgh on behalf of the Global Partnership on Nutrient Management and the International Nitrogen Initiative. ISBN: 978-1-906698-40-9
- SYAKILA A and KROEZE C (2011) The global nitrous oxide budget revisited. *Greenhouse Gas Measurement & Management*, 1: 17-26. DOI: 10.3763/ghgmm.2010.0007

- TALESKI M, IMIN N, and DJORDJEVIC MA (2018) CEP peptide hormones: key players in orchestrating nitrogen-demand signaling, root nodulation, and lateral root development. *J Exp Bot*, 69(8): 1829-1836. DOI: 10.1093/jxb/ery037
- TANG Z, FAN X, LI Q, et al. (2012) Knockdown of a rice stelar nitrate transporter alters long-distance translocation but not root influx. *Plant Physiol*, 160: 2052-2063. DOI: 10.1104/pp.112.204461
- TAULEMESSE F, LE GOUIS J, GOUACHE D, et al. (2015) Post flowering nitrate uptake in wheat is controlled by N status at flowering, with a putative major role of root nitrate transporter NRT2.1. *PLoS One*, 10(3): e0120291. DOI: 10.1371/journal.pone.0120291
- TILMAN D (1999) Global environmental impacts of agricultural expansion: The need for sustainable and efficient practices. *Proc Natl Acad Sci USA*, 96: 5995-6000. DOI: 10.1073/pnas.96.11.5995
- TILMAN D, FARGIONE J, WOLFF B, et al. (2001) Forecasting Agriculturally Driven Global Environmental Change. *Science*, 292: 281-284. DOI: 10.1126/science.1057544
- TILMAN D, CASSMAN KG, MATSON PA, et al. (2002) Agricultural sustainability and intensive production practices. *Nature*, 418: 671-677. DOI: 10.1038/nature01014
- TONG Y, ZHOU J-J, LI Z, and MILLER AJ (2005) A two-component high-affinity nitrate uptake system in barley. *Plant J*, 41: 442-450. DOI: 10.1111/j.1365-3113X.2004.02310.x
- TREVISAN S, BORSA P, BOTTON A, et al. (2008) Expression of two maize putative nitrate transporters in response to nitrate and sugar availability. *Plant Biol*, 10: 462-475. DOI: 10.1111/j.1438-8677.2008.00041.x
- TRUEMAN LJ, RICHARDSON A, and FORDE BG (1996) Molecular cloning of higher plant homologues of the high-affinity nitrate transporters of *Chlamydomonas reinhardtii* and *Aspergillus nidulans*. *Gene*, 175: 223-231. DOI: 10.1016/0378-1119(96)00154-0
- TSAY YF, SCHROEDER JI, FELDMANN KA, and CRAWFORD NM (1993) The herbicide sensitivity gene CHL1 of Arabidopsis encodes a nitrate-inducible nitrate transporter. *Cell*, 72: 705-713. DOI: 10.1016/0092-8674(93)90399-b
- TSAY YF, CHIU CC, TSAI CB, HO CH, and HSU PK (2007) Nitrate transporters and peptide transporters. *FEBS Lett*, 581: 2290-2300. DOI: 10.1016/j.febslet.2007.04.047
- TSAY YF, FAN SC, CHEN HY, and CHEN KE (2011) Method for changing nitrogen utilization efficiency in plants. US Patent Application No. 12/832, 234
- UNDURRAGA SF, IBARRA-HENRÍQUEZ C, FREDES I, et al. (2017) Nitrate signaling and early responses in Arabidopsis roots. *J Exp Bot*, 68 (10) 2541-2551. DOI: 10.1093/jxb/erx041
- UNEP and WHRC (2007) Reactive Nitrogen in the Environment: Too Much or Too Little of a Good Thing. United Nations Environment Programme. ISBN: 978 92 807 2783 8
- US Environmental Protection Agency (2011) Global Anthropogenic Non-CO₂ Greenhouse Gas Emissions 1990–2030. https://www.epa.gov/sites/production/files/2016-08/documents/summary_global_nonco2_projections_dec2012.pdf
- VAN HERK CM, MATHIJSEN-SPIEKMAN EAM, and DE Zwart D (2003). Long distance nitrogen air pollution effects on lichens in Europe. *Lichenologist*, 35: 347-359. DOI: 10.1016/S0024-2829(03)00036-7
- VAN VUUREN DP, WEYANT J, and DE LA CHESNAYE F (2006) Multi-gas scenarios to stabilize radiative forcing. *Energy Eco*, 28: 102-120. DOI: 10.1016/j.eneco.2005.10.003
- VELTHOF G, BAROT S, BLOEM J, et al. (2011) Nitrogen as a threat to European soil quality. In "The European Nitrogen Assessment". Cambridge University Press, pp: 495-510. ISBN: 9781107006126
- VIDAL EA, ARAUS V, LU C, et al. (2010) Nitrate-responsive miR393/AFB3 regulatory module controls root system architecture in *Arabidopsis thaliana*. *Proc Natl Acad Sci USA*, 107:4477–4482. DOI: 10.1073/pnas.0909571107
- VIDAL EA, MOYANO TC, RIVERAS E, et al. (2013) Systems approaches map regulatory networks downstream of the auxin receptor AFB3 in the nitrate response of *Arabidopsis thaliana* roots. *Proc Natl Acad Sci USA*, 110: 12840-12845. DOI: 10.1073/pnas.1310937110
- VIDMAR JJ, ZHUO D, SIDDIQI MY, et al. (2000) Regulation of high-affinity nitrate transporter genes and high-affinity nitrate influx by nitrogen pools in roots of barley. *Plant Physiol*, 123: 307-318. DOI: 10.1104/pp.123.1.307
- VITOUSEK PM, NAYLOR R, CREWS T, et al. (2009) Nutrient imbalances in agricultural development. *Science*, 324: 1519–1520. DOI: 10.1126/science.1170261
- VON DER FECHT-BARTENBACH J, BOGNER M, DYNOWSKI M and LUDEWIG U (2010) CLC-b-mediated NO₃⁻/H⁺ exchange across the tonoplast of Arabidopsis vacuoles. *Plant Cell Physiol*, 51(6): 960-968. DOI: 10.1093/pcp/pcq062
- VON MUTIUS E. (2000) The environmental predictors of allergic disease. *J Allergy Clin Immunol*, 105: 9-19. DOI: 10.1016/S0091-6749(00)90171-4

- VON WIRÉN N, GAZZARRINI S, GOJON A, and FROMMER WB (2000) The molecular physiology of ammonium uptake and retrieval. *Curr Opin Plant Biol*, 3: 254-261. DOI: 10.1016/S1369-5266(00)80074-6
- VON WITTGENSTEIN NJ, LE CH, HAWKINS BJ and EHLTING J (2014) Evolutionary classification of ammonium, nitrate, and peptide transporters in land plants. *BMC Evol Biol*, 14:11. DOI: 10.1186/1471-2148-14-11
- VOSS M, BAKER A, BANGE HW, et al. (2011) Nitrogen processes in coastal and marine systems. In: "The European Nitrogen Assessment", pp. 147-176. DOI: 10.1017/CBO9780511976988.011
- WANG J, LI Y, ZHU F, et al. (2019a) Genome-Wide Analysis of Nitrate Transporter (NRT/NPF) Family in Sugarcane *Saccharum spontaneum* L. *Trop Plant Biol*, 12:133-149. DOI: 10.1007/s12042-019-09220-8
- WANG J, HÜNER N, and TIAN L (2019b) Identification and molecular characterization of the *Brachypodium distachyon* NRT2 family, with a major role of BdNRT2.1. *Physiol Plant*, 165: 498-510. DOI: 10.1111/ppl.12716
- WANG P, WANG Z, CAI R, et al. (2011) Physiological and molecular response of wheat roots to nitrate supply in seedling stage. *Agric Sci China*, 10: 695-704. DOI: 10.1016/S1671-2927(11)60052-7
- WANG R, XING X, WANG Y, et al. (2009) A genetic screen for nitrate regulatory mutants captures the nitrate transporter gene NRT1.1. *Plant Physiol*, 151: 472-478. DOI: 10.1104/pp.109.140434
- WANG W, HU B, YUAN D, et al. (2018a) Expression of the nitrate transporter gene OsNRT1.1A/OsNPF6.3 confers high yield and early maturation in rice. *Plant Cell*, 30: 638-651. DOI: 10.1105/tpc.17.00809
- WANG X, BIAN Y, CHENG K, et al. (2012a) A comprehensive differential proteomic study of nitrate deprivation in *Arabidopsis* reveals complex regulatory networks of plant nitrogen responses. *J Proteome Res*, 11: 2301-2315. DOI: 10.1021/pr2010764
- WANG YY, HSU PK, and TSAY YF (2012b) Uptake, allocation and signaling of nitrate. *Trends Plant Sci*, 17: 458-467. DOI: 10.1016/j.tplants.2012.04.006
- WANG YY, CHENG YH, CHEN KE, and TSAY YF (2018b) Nitrate Transport, Signaling, and Use Efficiency. *Annu Rev Plant Biol*, 69: 85-122. DOI: 10.1146/annurev-arplant-042817-040056
- WEDLICH K, RINTOUL N, PEACOCK S, et al. (2012) Effects of ozone on species composition in an upland grassland. *Oecologia*, 168(4): 1137-1146. DOI: 10.1007/s00442-011-2154-2
- WEI J, ZHENG Y, FENG H, et al. (2018) OsNRT2.4 encodes a dual-affinity nitrate transporter and functions in nitrate-regulated root growth and nitrate distribution in rice. *J Exp Bot*, 69(5): 1095-1107. DOI: 10.1093/jxb/erx486
- WEN ZY, TYERMAN SD, DECHORGNAT J, et al. (2017) Maize NPF6 proteins are homologs of *Arabidopsis* CHL1 that are selective for both nitrate and chloride. *Plant Cell*, 29(10): 2581-2596. DOI: 10.1105/tpc.16.00724
- WHO (2003) Health aspects of air pollution with particulate matter, ozone and nitrogen dioxide. Report on a WHO Working Group; Bonn, Germany; 13-15 January 2003. World Health Organization, Copenhagen. <https://apps.who.int/iris/handle/10665/107478>
- WHO (2008) Health risks of ozone from long-range transboundary air pollution. World Health Organization, Regional Office for Europe, Copenhagen, Denmark. http://www.euro.who.int/__data/assets/pdf_file/0005/78647/E91843.pdf
- WIDIEZ T, EL KAFABI ES, GIRIN T, et al. (2011) High nitrogen insensitive 9 (HNI9)-mediated systemic repression of root NO₃⁻ uptake is associated with changes in histone methylation. *Proc Natl Acad Sci USA*, 108: 13329-13334. DOI: 10.1073/pnas.1017863108
- WIRTH J, CHOPIN F, SANTONI V, et al. (2007) Regulation of root nitrate uptake at the NRT2.1 protein level in *Arabidopsis thaliana*. *J Biol Chem*, 282: 23541-23552. DOI: 10.1074/jbc.M700901200
- WOLSELEY PA, JAMES PW, THEOBALD MR, and SUTTON MA (2006) Detecting changes in epiphytic lichen communities at sites affected by atmospheric ammonia from agricultural sources. *Lichenologist*, 38: 161-176. DOI: 10.1017/S0024282905005487
- WOLT JD (1994) Soil solution chemistry: applications to environmental science and agriculture. John Wiley & Sons Inc, 345p. ISBN: 0471585548.
- WU J, ZHANG ZS, XIA JQ, et al. (2020) Rice NIN-LIKE PROTEIN 4 is a master regulator of nitrogen use efficiency. *BioRxiv*. DOI: 10.1101/2020.01.16.908558
- WU K, WANG S, SONG W, et al. (2020) Enhanced sustainable green revolution yield via nitrogen-responsive chromatin modulation in rice. *Science*, 367(6478): eaaz2046. DOI: 10.1126/science.aaz2046
- XIA X, FAN X, WEI J, et al. (2015) Rice nitrate transporter OsNPF2.4 functions in low-affinity acquisition and long-distance transport. *J Exp Bot*, 66: 317-331. DOI: 10.1093/jxb/eru425

- XIE Q, FRUGIS G, COLGAN D, and CHUA NH (2000) Arabidopsis NAC1 transduces auxin signal downstream of TIR1 to promote lateral root development. *Genes Dev*, 14(23): 3024-3036. DOI: 10.1101/gad.852200
- XU G, FAN X, and MILLER AJ (2012) Plant nitrogen assimilation and use efficiency. *Annu Rev Plant Biol*, 63: 153-82. DOI: 10.1146/annurev-arplant-042811-105532
- XU N, WANG R, ZHAO L, et al. (2016) The Arabidopsis NRG2 protein mediates nitrate signaling and interacts with and regulates key nitrate regulators. *Plant Cell*, 28: 485-504. DOI: 10.1105/tpc.15.00567
- XUAN W, BEECKMAN T, and XU G (2017) Plant nitrogen nutrition: sensing and signaling. *Curr Opin Plant Biol*, 39: 57-65. DOI: 10.1016/j.pbi.2017.05.010
- YAMAYA T, OBARA M, NAKAJIMA H, et al. (2002) Genetic manipulation and quantitative-trait loci mapping for nitrogen recycling in rice. *J Exp Bot*, 53: 917-925. DOI: 10.1093/jexbot/53.370.917
- YAN M, FAN X, FENG H, et al. (2011) Rice OsNAR2.1 interacts with OsNRT2.1, OsNRT2.2 and OsNRT2.3a nitrate transporters to provide uptake over high and low concentration ranges. *Plant Cell Environ*, 34: 1360-1372. DOI: 10.1111/j.1365-3040.2011.02335.x
- YANG Z, WANG Z, YANG C, et al. (2019) Physiological responses and small RNAs changes in maize under nitrogen deficiency and resupply. *Genes Genom*, 41(10): 1183-1194. DOI: 10.1007/s13258-019-00848-0
- YIN LP, LI P, WEN B, et al (2007) Characterization and expression of a high-affinity nitrate system transporter gene (TaNRT2.1) from wheat roots, and its evolutionary relationship to other NRT2 genes. *Plant Sci*, 172: 621-631. DOI: 10.1016/j.plantsci.2006.11.014
- YONG Z, KOTUR Z, and GLASS AMD (2010) Characterization of an intact two-component high- affinity nitrate transporter from Arabidopsis roots. *Plant J*, 63: 739-748. DOI: 10.1111/j.1365-313X.2010.04278.x
- ZHANG H and FORDE BG (1998) An Arabidopsis MADS box gene that controls nutrient-induced changes in root architecture. *Science*, 279: 407-409. DOI: 10.1126/science.279.5349.407
- ZHANG X, GERMANN S, BLUS BJ, et al. (2007) The Arabidopsis LHP1 protein colocalizes with histone H3 Lys27 trimethylation. *Nat Struct Mol Biol*, 14(9): 869-871. DOI: 10.1038/nsmb1283
- ZHANG X, DAVIDSON EA, MAUZERALL DL, et al. (2015) Managing nitrogen for sustainable development. *Nature*, 528(7580): 51-59. DOI: 10.1038/nature15743
- ZHAO L, LIU F, CRAWFORD NM, and WANG Y (2018) Molecular regulation of nitrate responses in plants. *Int J Mol Sci*, 19: 2039. DOI: 10.3390/ijms19072039
- ZHAO M, DING H, ZHU JK, et al. (2011) Involvement of miR169 in the nitrogen-starvation responses in Arabidopsis. *New Phytol*, 190: 906-915. DOI: 10.1111/j.1469-8137.2011.03647.x
- ZHUO D, OKAMOTO M, VIDMAR JJ, and GLASS ADM (1999) Regulation of a putative high-affinity nitrate transporter (Nrt2;1At) in roots of *Arabidopsis thaliana*. *Plant J*, 17: 563-568. DOI: 10.1046/j.1365-313x.1999.00396.x
- ZOU L, QI D, SUN J, et al. (2019) Expression of the cassava nitrate transporter NRT2.1 enables Arabidopsis low nitrate tolerance. *J Genet*, 98: 74. DOI: 10.1007/s12041-019-1127-9
- ZULUAGA DL and SONNANTE G (2019) The use of nitrogen and its regulation in cereals: structural genes, transcription factors, and the role of miRNAs. *Plants*, 8: 294. DOI: 10.3390/plants8080294

2. Regulation of Nitrate Transporters (NRT2 and NRT3) genes expression in the model grass *Brachypodium distachyon* under various N conditions

2.1 Abstract

Nitrogen (N), as a major limiting nutrient in the field for crop production, has been extensively used in modern agriculture to increase crop production. The indiscriminate use of N fertilizer and the low average Nitrogen Use Efficiency (NUE) in crops results in many negative environmental outcomes. The intensive use of N fertilizers raises questions regarding to crop sustainability and food security. An increase in the global food and energy demand implies that the requirement for N fertilizers is likely to grow substantially. Thus, N management and NUE improvement are critical to maintain a growing population, while decreasing environmental and health consequences. *Brachypodium distachyon* is a genetic model system for gene function studies of grasses. Despite the exponential growth in research using *B. distachyon* as a model plant, little has been explored in relation to investigating N uptake, and particularly NUE. The physiological and molecular characterization of the nitrate acquisition process in this species could help to improve non-model crops, such as sugarcane. ¹⁵N-ammonium and ¹⁵N-nitrate influx analyses upon N-replete and N-deficient conditions revealed that ammonium is the preferential inorganic N source of *B. distachyon*. A negative feedback regulation of nitrate uptake process by ammonium in plants under N provision and N resupply treatments was observed. Gene expression analysis by RT-qPCR indicated *BdNRT2.1* and *BdNRT2.2* as the main *B. distachyon* genes responsible for nitrate acquisition. The expression of *BdNRT3.1* and *BdNRT3.2* were inducible, revealing the co-expression of members from both NRT subfamilies in response to either N treatments. There was no correlation between ¹⁵N-nitrate influx and transcript accumulation for members of the two-component complex (NRT2/ NRT3) after ammonium chloride provision and ammonium nitrate resupply treatments, corroborating the occurrence of a transcriptional regulation of HATS (High Affinity Transport System) in *B. distachyon*. Therefore, even though ammonium is a preferential N inorganic source in *B. distachyon*, nitrate HATS is repressed by this N inorganic source.

Keyword: Nitrate, NUE, Preferential N Source, HATS, Ammonium

2.2 Introduction

Nitrogen (N) is a major limiting nutrient in the field for crop production. As a result, N fertilizers are extensively used in modern agriculture to improve crop yields. The large scale of N fertilization raises questions about crop sustainability and food security since both are related to fertilizer and nutrient use efficiency. The indiscriminate use of N fertilizer on fields and the low Nitrogen Use Efficiency (NUE) in crops may result in many environmental hazards, such as water and air pollution, loss of biodiversity, coastal marine waters and watersheds contamination, N_2O and reactive N gases accumulation in the troposphere, leading to greenhouse effect and climate change (Robertson and Vitousek 2009, Canfield et al. 2010, Howarth et al. 2011, Boudry et al. 2012, Swaney et al. 2012, Sutton et al. 2013, Zhang et al. 2015). By 2050, there will be a 70% increase in the global food demand, as the world population will increase to over 9.7 billion (Yang et al. 2012, York et al. 2016, Chen and Liao 2017), implying that the demand for N fertilizers and agricultural land are likely to grow substantially (Foley et al. 2011, Mueller et al. 2012). So, N management and NUE improvement will be required to support a growing population, while reducing adverse environmental and health consequences.

Several definitions have been proposed for NUE (Moll et al. 1982, Good et al. 2004, Pathak et al. 2011, Xu et al. 2012, Yu et al. 2016). Overall, NUE is composed of two plant physiological components: N utilization efficiency (NUE), describing the assimilation and remobilization efficiency of plant N; and N uptake efficiency (NUpE), which is the N absorption/uptake efficiency focusing on the recovery of N fertilizer from the soil (Moll et al. 1982, Good et al. 2004). In simple terms, plants can either increase the amount of N uptaken from the soil (Nitrogen uptake efficiency), or plants can incorporate the absorbed N more efficiently (Nitrogen utilization efficiency). Thus, to improve NUE in crops, it is important to understand the key mechanisms involved in N uptake and N utilization processes. Plants absorb soluble organic and inorganic N sources present in the soil, with ammonium and nitrate the main inorganic forms taken up by the roots (Miller et al. 2007, Marschner 2012). Nitrate is the most abundant inorganic N form in aerobic soils, and it represents the main N source for several crops (Miller and Cramer 2005, Miller et al. 2007, Marschner 2012, Xu et al. 2012). The nitrate uptake by plants roots relies on transmembrane transporter, that can be classified as High or Low-Affinity Transporter Systems (HATS and LATS, respectively). HATS are responsible for the acquisition of nitrate in concentrations below 1 mM, while LATS act when nitrate concentrations are above 1 mM (Glass et al. 2002). LATS involve most NRT1/NPF family members with the exception of NRT1.1/NPF6.3. NRT1.1/NPF6.3 can operate as LATS or HATS, depending on the phosphorylation–dephosphorylation of the threonine residue T101, responsible for the change from a low to a high-affinity uptake form (Liu and Tsay 2003). In contrast to the NRT1/NPF transporters, the NRT2 family is composed of high-affinity nitrate

transporters, regulated by nitrate availability (Tsay et al. 2007). Most NRT2 high-affinity transporters are active in the plasma membrane where NAR2/NRT3 family members are co-localized, forming a multi-component system (Tong et al. 2005, Kawachi et al. 2006, Okamoto et al. 2006, Orsel et al. 2006, Yong et al. 2010, Kotur et al. 2012, Laugier et al. 2012).

HATS have been extensively characterized at the physiological and molecular levels mostly in the dicot *Arabidopsis* (Dechorgnat et al. 2011, Xu et al. 2012, Havé et al. 2017). Nevertheless, the most important crop plants for human and livestock nutrition are members of the grass family (*Poaceae*), including rice, wheat, maize, sorghum, and sugarcane. Differences in gene number, family structure and phylogeny of NRT2 and NRT3 families have been confirmed between *Arabidopsis* and monocots species (Plett et al. 2010, Girin et al. 2014). *Setaria* is proposed as a genetic model system for functional genomics of grasses species with inherent biological difficulties, particularly C4 species. However, there are fewer genetic resources available for *Setaria* than *Brachypodium*. To our knowledge, there are no characterized insertional mutant lines available, but an EMS-mutagenized M1 population has been developed and used to map-clone phytoene-desaturase gene (Sun et al. 2019).

On the other hand, *Brachypodium distachyon*, with a simple diploid sequenced genome, short life cycle, well-established genetic transformation protocols, and genetic resources such as T-DNA mutant lines, become an important experimental model for grasses (Draper et al. 2001, Bevan et al. 2010, Brkljacic et al. 2011, Mur et al. 2011, Mochida and Shinozaki 2013). A set of indexed T-DNA insertion lines was recently released, providing a powerful genetic resource for functional genomics approaches (Bragg et al. 2012, Hsia et al. 2017). In this chapter of my thesis work, I will explore the use of *Brachypodium distachyon* as a genetic model species to investigate the regulation mechanisms of *NRT2* and *NRT3* gene family members in response to nitrate availability in order to deepen the understanding how NUE is affected in grass crops, particularly those less amenable to genetic studies, such as sugarcane.

2.3 Methods

2.3.1 Plant material and germination

The *Brachypodium distachyon* seeds (Bd21) were gently provided by Dr. Eric Lam and Dr. Rong Di at the Department of Plant Biology, School of Environmental and Biological Sciences, Rutgers - The State University of New Jersey. Prior to germination, seeds were placed in the dark at 4°C for 7 days. Seeds were sown in each pot (3 x 3 cm) with optimal spacing and cover with a thin layer of soil. The pots were covered by plastic lid to maintain high humidity. After cold stratification, pots were moved to a growth chamber with 14 h light/ 10 h dark at 21°C. When the cotyledons emerged completely (one week later), pot covers

were removed. The plants were watered and fertilized once a week with Hoagland solution (Hoagland and Arnon, 1950). After the full cycle, when plants were completely dry, the panicles were harvested, and seeds were separated and stored. The seeds were stored in airtight containers at room temperature for further experiments.

2.3.2 Analysis of nitrogen uptake kinetics in roots by rapid ^{15}N influx

After seed germination, the plants were grown in complete N nutrient solution (2 mM NH_4NO_3) for 3 weeks. After, plants were subjected to various N conditions (sufficiency, deficiency or resupply treatments). For the *in vivo* N uptake kinetics analysis, we used the rapid influx assay with ^{15}N (Short Term Uptake; Loqué et al. 2006). To estimate the rapid influx, first, the roots are washed in 1 mM CaSO_4 to remove apoplastic N. Then, roots are placed in 0.2 L of nutrient solution, containing 0.2 mM of ^{15}N in 1 mM MES (2- (N-morpholino) ethanesulfonic acid monohydrate), pH 5.8 for 10 min, followed by 1 min in 1 mM CaSO_4 solution to wash the roots (Loqué et al., 2006). To quantify the ^{15}N -nitrate influx, roots samples are collected, dried and ground for subsequent analysis by mass spectrometer ionization by electron impact IRMS (GSL 20/20 ANCA or ANCA-SL of Sercon, England) in the Stable Isotopes Laboratory, CENA, by Prof. Dr. José Albertino Bendassolli.

2.3.3 Identification of preferential N source in *Brachypodium distachyon*

To define the preferential inorganic N source (nitrate or ammonium) of *B. distachyon*, plants grown for one week in soil were transferred to nutrient solution for three weeks. Then, the plants were submitted to N sufficient (+N; 2 mM de NH_4NO_3) or N absence for three days. After this period, we conducted the rapid ^{15}N -influx analysis using ^{15}N labeled nitrate or ammonium [0.2 mM K^{15}NO_3 , 60% ^{15}N atoms or 0.1 mM $(^{15}\text{NH}_4)_2\text{SO}_4$, 60% ^{15}N atoms]. Five biological replicates were sampled for each treatment.

2.3.4 Understanding the nitrate uptake process in roots upon distinct inorganic N source supply and N resupply

The experiments were conducted to characterize the regulation of nitrate transport in roots. Plants were grown for three weeks in a N sufficient nutrient solution (N+; 2 mM NH_4NO_3), when exposed to nitrate (4 mM KNO_3) or ammonium (4 mM NH_4Cl). Subsequently, we performed the rapid ^{15}N -nitrate influx analysis (0.2 mM K^{15}NO_3) at 0, 2, 4 and 24 h after the N provision. For the N resupply experiment, plants grown under N sufficiency were subjected to restriction N (-N) for one week. After this period, the plants were exposed to nitrate (10 mM KNO_3) or ammonium nitrate (5 mM NH_4NO_3), followed by analyzing the rapid ^{15}N -nitrate influx (0.2 mM K^{15}NO_3) at 0, 2, 4, 24 and 72 h after N resupply. For both experiments, five biological replicates were sampled at each time point. Samples of *B. distachyon* roots were collected, frozen and used for total RNA extraction to quantify the transcripts via RT-qPCR (Quantitative

Reverse Transcription Polymerase Chain Reaction) of NRT2 and NRT3 gene family members of *B. distachyon*, the key plant gene families responsible for nitrate uptake process (*NRT - NITRATE TRANSPORTERS*).

2.3.5 RNA extraction and cDNA synthesis

Samples were collected and immediately frozen in liquid N₂ for subsequent RNA extraction. Samples (100 mg) were extracted using the method described by Leal et al (2007). Briefly, the root samples were ground in liquid N and transferred to a 2 mL microtube, and 1 mL of extraction buffer (2% CTAB; 2 M NaCl; 2% PVP; 100 mM Tris; 25 mM EDTA; 2% β -mercaptoethanol) warmed to 55°C was added. After homogenization, samples were incubated at 55°C for 60 min. Afterwards, the samples were placed immediately on ice for 1 min, followed by addition of 1 mL of CIA (chloroform isoamyl-alcohol; 24:1 v/v) and spun at 9000 *g* for 10 min at 4°C. Then, the supernatant was transferred to a new 1.5 mL microtube, adding an equal volume of CIA, followed by centrifugation at 9000 *g* for 10 min at 4°C. Once again, the supernatant was transferred to a new 1.5 mL microtube for subsequent precipitation of the total RNA using a 10 M solution of LiCl and incubation at -80 °C for 12 h. Subsequently, the samples were spun at 9000 *g* for 30 min at 4°C, followed by discarding the supernatant. The pellet was washed twice with 500 μ L of cold 75% ethanol and centrifuged at 9000 *g* for 10 min at 4°C. Finally, the pellet was resuspended in 30 μ L of 0.01% inactive DEPC water. Total RNA was quantified in NanoDrop One (Thermo Scientific; Wilmington, DE, USA) and the integrity was confirmed by gel electrophoresis.

About 2 μ g of RNA from each sample was treated with 1 U of DNase I in appropriate buffer, 2 U of RNaseOUT (Invitrogen) and DEPC ultrapure water in reaction, incubated at 37°C for 30 min. The reaction was stopped by addition of EDTA (1 mM) and then incubated at 65°C for 10 min. Samples of 1 μ g of total RNA and oligo dT (0.5 μ g μ L⁻¹) are denatured at 70°C for 10 min, and then incubated at 4°C for 2 min, followed by addition of 4 μ L 5x buffer-MuLV RT, 1 mM dNTP, 40 U RNaseOUT, and 20 U of the enzyme SuperScript III (Thermo Fisher Scientific). The reaction was then incubated at 55°C for 1 h, then at 70°C for 10 min and at 4°C for 2 min. Finally, the reaction was cooled to 4°C and stored at -80°C.

2.3.6 Quantitative reversed transcript amplification (RT-qPCR) of the BdNRT2 and BdNRT3 family members

The cDNA was then diluted [1:10 (v/v)], and used in quantitative PCR reactions, using 1 μ L of cDNA, 5 μ L of *Taq* Platinum (Kapa Biosystems, Wilmington, MA, USA), and 0.5 μ L of both forward and reverse primers (0.25 mM) with the reactions performed in a RotorGene Q 6000 thermocycler (Qiagen; Hilden, Germany), programmed with an initial incubation at 95°C for 2 min, followed by 40 cycles of 95°C for 20 s and 60°C-62°C for 20 s, and 72°C for 25 s.

After cycling, the melting curve for each amplicon was determined between 72°C and 95°C. Transcript-specific primers are listed below (Table 1). The RT-qPCR included for each run three independent biological sample, each sample with two replicates, and an additional negative control (no template DNA). Data acquisition was performed in a Rotor-Gene Q Real-Time Analysis 6.0 (Qiagen). *BdSamDC* was used as the reference gene (Table 1). C_q (quantification cycle) was used to determine differences in expression according to the $2^{-\Delta(\Delta C_q)}$ method (Livak and Schmittgen 2001).

Table 1: Primer sequences related to genes encoding transporters proteins (*NRTs* - *NITRATE TRANSPORTERS*), responsible for the nitrate uptake in *Brachypodium distachyon* roots, and the reference gene (*BdSamDC*), used for RT-qPCR.

Gene name	Locus ID	Primers
<i>BdNRT2.1</i>	<i>BRADI_3g01270v3</i>	F: 5'- TTTCTCTGTCCCAGCTACG -3' R: 5'- CGCAGCAAATGTGTAGTACCTC -3'
<i>BdNRT2.2</i>	<i>BRADI_3g01250v3</i>	F: 5'- AAGCCGGCGCTCATGGAGAC -3' R: 5'- GAGGTTGTCGCGGATGATAG -3'
<i>BdNRT2.3</i>	<i>BRADI_3g01277v3</i>	F: 5'- AATACTCCACCGCCAAAGG-3' R: 5'- TTCTCGGCAAACCTTCTGTCC-3'
<i>BdNRT2.4</i>	<i>BRADI_3g01290v3</i>	F: 5'- GTCGCCCCGAGCACATATAA-3' R: 5'- GTCATGTAGTGTGGGAAAACCA-3'
<i>BdNRT2.5</i>	<i>BRADI_2g47640v3</i>	F: 5'- CCGACGTGTGAAGAAAATCC-3' R: 5'- TCGCGAAAAGAAAAGTGGAG-3'
<i>BdNRT2.6</i>	<i>BRADI_2g26210v3</i>	F: 5'- GCACGTCTTTGGAACATCTG-3' R: 5'- ATGCCGATTGCCGAGGGA-3'
<i>BdNRT2.7</i>	<i>BRADI_2g40740v3</i>	F: 5'-TCTCCTTCACGGGCCTCA -3' R: 5'-GTGCAAGCACCGATCAATTT-3'
<i>BdNRT3.1</i>	<i>BRADI_3g47710v3</i>	F: 5'- TACGGCCAGACCAAGCTC -3' R: 5'- CCTTGGCAAGTTGCATTATC -3'
<i>BdNRT3.2</i>	<i>BRADI_3g47720v3</i>	F: 5'- TCGCCTTCTTCTTCTTCATC -3' R: 5'- GCCTTGCAATAGATTGCTGT -3'
<i>BdSamDC</i>	<i>BRADI_5g14640v3</i>	F: 5'- TGCTAATCTGCTCCAATGGC-3' R: 5'- GACGCAGCTGACCACCTAGA-3'

2.4 Results

2.4.1 Gene and protein analysis of the NRT2 and NRT3 subfamily in *Brachypodium distachyon*

The importance of the NRT2 and NRT3 gene family members concerning nitrate uptake have been described for many species (Trueman et al. 1996, Vidmar et al. 2000, Ruffel et al. 2008, Plett et al. 2010, Feng et al. 2011). The high-affinity nitrate uptake process in *B. distachyon* might be due to additive/redundant protein activities or it might depend on a major member of this subfamily, or even each member could play organ-specific or N status regulation. To understand the biological function of both families, the gene structures genomic, phylogeny, transmembrane domains and gene expression profile analysis in response to N provision and resupply are required.

A schematic representation of BdNRT2 gene structures was performed using Gene Structure Display Server 2.0 (GSDS 2.0) web-based tool. According to the number of exons and relative positions of introns within the coding sequence, the *BdNRT2* genes can be classified into two different groups. Five members of the first group (*BdNRT2.1*, *BdNRT2.2*, *BdNRT2.3*, *BdNRT2.4*, and *BdNRT2.5*) are characterized by the absence of introns (Figure 1). The second group (*BdNRT2.6* and *BdNRT2.7*) is characterized by the presence of one intron (Figure 1). The *BdNRT2* genomic distribution on *B. distachyon* chromosomes was also analyzed. *BdNRT2.1*, *BdNRT2.2*, *BdNRT2.3* and *BdNRT2.4* are clustered on chromosome 3, presented in a tail-to-tail configuration for *BdNRT2.1* and *BdNRT2.2*, or in a head-to-tail configuration for *BdNRT2.3* and *BdNRT2.4* (Figure 2). The other members of the NRT2 gene family (*BdNRT2.5*, *BdNRT2.6*, and *BdNRT2.7*) are located on chromosome 2. However, unlike the other members, these genes are not grouped at the same genomic region (Figure 2).

Figure 1: Gene structure of the *BdNRT2* and *BdNRT3* family. Sequences were aligned using the Gene Structure Display Server 2.0 (GSDS 2.0) web-based tool. Exons and introns were shown by filled blue boxes and single lines, respectively. UTRs were displayed by thick green boxes at both ends.

Figure 2: Genomic localization and orientation of *BdNRT2* and *BdNRT3* genes on chromosomes of *Brachypodium distachyon*. The genes shown in red are transcribed in the positive orientation; the genes shown in blue are transcribed in the negative orientation.

Not only the importance of AtNRT2.1 and AtNRT2.2 concerning the nitrate uptake process has been reported, but also the *AtNRT2.1* and *AtNRT2.2* genomic localization (Cerezo et al. 2001, Plett et al. 2010). Both are neighboring genes at opposite orientation, as in the *B. distachyon* genome. Therefore, the similarity of the structural genomic organization between both species could provide a hint about *BdNRT2.1* and *BdNRT2.2* function.

Table 2: Protein BdNRT2 sequence identity matrix

Seq->	BdNRT2.1	BdNRT2.2	BdNRT2.3	BdNRT2.4	BdNRT2.5	BdNRT2.6	BdNRT2.7
BdNRT2.1	ID	0.931	0.825	0.807	0.549	0.660	0.364
BdNRT2.2	0.931	ID	0.798	0.788	0.553	0.644	0.361
BdNRT2.3	0.825	0.798	ID	0.897	0.535	0.597	0.364
BdNRT2.4	0.807	0.788	0.897	ID	0.520	0.584	0.353
BdNRT2.5	0.549	0.553	0.535	0.520	ID	0.413	0.409
BdNRT2.6	0.660	0.644	0.597	0.584	0.413	ID	0.315
BdNRT2.7	0.364	0.361	0.364	0.353	0.409	0.315	ID

An alignment of the seven BdNRT2 protein sequences, beginning and ending at the putative ATG and stop codons was performed using Blosom62 pairwise alignment (Table 2, Figure 3). Except for BdNRT2.7, the other members of the BdNRT2 family show great protein sequence similarity (Table 2). The highest levels of nucleotide sequence identity are found between the BdNRT2.1 and BdNRT2.2 (93.1%) and between the BdNRT2.3 and BdNRT2.4 (89.7%) proteins. Phylogenetic analysis comparing *B. distachyon* NRT2 members with orthologues of arabidopsis, rice, maize, and sorghum was also performed. Except for BdNRT2.5 and BdNRT2.7, BdNRT2 group at the same clade with other grass NRT2, being separated from the Arabidopsis NRT2 (Figure 4).

Figure 3: Alignment of NRT2 protein sequences from *Brachypodium distachyon* using ClustalW software and displayed with BioEdit program.

Figure 4. Phylogenetic analysis of the NRT2 family. Evolutionary analysis was inferred by using the Maximum Likelihood method and Poisson correction model comparing NRT2 sequences of Arabidopsis, rice, sorghum, maize and *Brachypodium distachyon*. The percentage of trees in which the associated taxa clustered together is shown next to the branches. Initial tree(s) for the heuristic search were obtained automatically by applying Neighbor-Join and BioNJ algorithms to a matrix of pairwise distances estimated using a JTT model, and then selecting the topology with superior log likelihood value. The tree is drawn to scale, with branch lengths measured in the number of substitutions per site. Evolutionary analyses were conducted in MEGA X.

The topology of BdNRT2 proteins were simulated using the webtool Protter (Omasits et al. 2014) to correlate their respective structures and arrangements in the plasma membrane with their role in the nitrate uptake process. According to the number of transmembrane

domains, BdNRT2 family proteins could be classified into three groups. Group I is comprised of BdNRT2.1 and BdNRT2.2, considering the presence of 12 transmembrane domains and N-terminal and C-terminal cytosolic regions (Figure 5). Group II contains proteins with 11 transmembrane domains. In this group, the BdNRT2.3 and BdNRT2.4 proteins have N-terminal regions in the cytosolic portion and the C-terminal regions at the extracellular portion. The BdNRT2.5 protein, belonging to this group, with the opposite configuration, *i.e.*, N-terminal regions predicted to be at the extracellular portion and C-terminal regions at the cytosolic portion (Figure 5). Group III, composed of the proteins BdNRT2.6 and BdNRT2.7, holds members proteins with 10 transmembrane domains. However, there are differences between Group II proteins about the amino and carboxylic ends display. BdNRT2.6 displays both ends at the extracellular portion, while BdNRT2.7 has both terminal ends in the cytosolic portion (Figure 5). In summary, BdNRT2.1 and BdNRT2.2 share a common transmembrane conformation, suggesting that both proteins are plasma membrane high-affinity NO_3^- transporters in *B. distachyon*, similar to orthologues in other species (Forde et al. 2000, Plett et al. 2010).

Figure 5: Topology illustration of BdNRT2 showing divergency in the number of predicted transmembrane domains among BdNRT2 members. Peptides sequences were visualized by Protter.

The same approach was used for both *BdNRT3.1* and *BdNRT3.2* characterization. Both members have one intron (Figure 1). Both are located on chromosome 3, in a head-to-tail configuration (Figure 2; Plett et al. 2010), distinct from the orthologous *AtNRT3* in Arabidopsis. BdNRT3 sequence alignment showed a substantial similarity between both members *BdNRT3.1* and *BdNRT3.2* (Table 3, Figure 6), with 97% of amino acid sequence identity. Phylogenetic analysis with NRT3 members with orthologues from Arabidopsis, rice, maize, and sorghum revealed that BdNRT3 proteins are at the same clade with other grass NRT3, revealing evolution divergence between dicots and monocots NRT3 (Figure 7).

Table 3: Protein BdNRT3 sequence identity matrix

Seq->	BdNRT3.1	BdNRT3.2
BdNRT3.1	ID	0.970
BdNRT3.2	0.970	ID

Figure 6: Alignment of NRT3 protein sequences from *Brachypodium distachyon* using ClustalW software and displayed with BioEdit program.

Figure 7. Phylogenetic analysis of the NRT3 family. Evolutionary analysis was inferred by using the Maximum Likelihood method and Poisson correction model comparing NRT3 sequences of Arabidopsis, rice, sorghum, maize and *Brachypodium distachyon*. The percentage of trees in which the associated taxa clustered together is shown next to the branches. Initial tree(s) for the heuristic search were obtained automatically by applying Neighbor-Join and BioNJ algorithms to a matrix of pairwise distances estimated using a JTT model, and then selecting the topology with superior log likelihood value. The tree is drawn to scale, with branch lengths measured in the number of substitutions per site. Evolutionary analyses were conducted in MEGA X.

Figure 8: Topology illustration of BdNRT3 showing divergency in the number of predicted transmembrane domains among BdNRT3 members. Peptides sequences were visualized by Protter.

The structure of BdNRT3 proteins were simulated to investigate the structure across the plasma membrane to compare with BdNRT2, as described for other plant species, such as, *Arabidopsis thaliana*, barley (*Hordeum vulgare*) and rice (*Oryza sativa*) (Okamoto et al. 2006, Orsel et al. 2006, Ishikawa et al. 2009, Yong et al. 2010, Feng et al. 2011). BdNRT3.1 and

BdNRT3.2 proteins have identical presumed arrangements across the plasma membrane, exhibiting a single transmembrane domain, an extracellular N-terminus, and a cytosolic C-terminus (Figure 8). All together, these *in silico* predictions of the BdNRT3.1 and BdNRT3.2 members suggest that both comply with the formation of a NRT2/NRT3 complex formation, similar to the one described for *Arabidopsis*, and consequently in the nitrate uptake process, in *Brachypodium distachyon*.

2.4.2 *Brachypodium distachyon* prefers ammonium uptake rather than nitrate as N inorganic source

Few studies have been conducted to define the preference of the inorganic N source for several species (Robinson et al. 2011, Boudsocq et al. 2012, Boschiero et al. 2018). To identify the N source preference in *B. distachyon* roots, physiological characterization of the uptake process was conducted under N contrasting conditions. *Brachypodium distachyon* (Bd21) plants grown for three weeks under N sufficiency conditions were subjected to sufficiency treatment (+N; 2 mM NH_4NO_3) or N absence (-N) for three days. Rapid influx assay (Short-Term Uptake; Loqué et al. 2006) using ^{15}N -nitrate or ^{15}N -ammonium were performed at the HATS activation concentration range (0.2 mM), given the importance of HATS regulation under cultivation conditions in crops (Malagoli et al. 2004), and the LATS activity dependence of HATS (Wang et al. 1993a, 1993b).

Figure 9. Influx of $^{15}\text{N}\text{-NO}_3^-$ or $^{15}\text{N}\text{-NH}_4^+$ in *Brachypodium distachyon* 'Bd21' roots submitted to contrasting N availability. Plants cultivated in solution under N sufficient conditions (+N; 2 mM NH_4NO_3) were submitted to absence of N (-N) or with N sufficiency (2 mM NH_4NO_3) for 3 days. Uptake kinetic analyses were conducted using ^{15}N -nitrate or ^{15}N -ammonium at 0.2 mM total N total. Values are means of 5 biological replicates $\pm$ SD, and significant differences at $P < 0.05$ are indicated by different letters. Asterisk represents significant differences at $P < 0.05$ in each N condition.

Short-term uptake analysis indicated that ^{15}N -ammonium influx was 1.7-fold higher in plants under N absence ($105.27 \mu\text{mol h}^{-1} \text{g}^{-1} \text{DW}$) compared to N sufficient ($61.92 \mu\text{mol h}^{-1} \text{g}^{-1} \text{DW}$), showing HATS induction under N limitation (Figure 9). The same HATS induction was observed for nitrate uptake. The ^{15}N -nitrate influx in plants under N absence ($62.45 \mu\text{mol h}^{-1} \text{g}^{-1} \text{DW}$) was approximately 2.9-fold higher than under N sufficient ($21.70 \mu\text{mol h}^{-1} \text{g}^{-1} \text{DW}$), indicating the regulation of nitrate absorption according to N status (Figure 9). These results demonstrate that ammonium is the preferential inorganic N source for *B. distachyon* roots. Similar preference of ammonium has been observed for other crops, such as citrus, coffee and sugarcane (Camanès 2009, Robinson et al. 2011, Serezino 2015, Santos et al. 2017). HATS in *B. distachyon* for both inorganic sources of N are induced under low external levels of N and/or the low N status of the plant. The HATS induction mechanism is conserved in many organisms, even monocots and dicots species (von Wirén et al. 2000, Subbarao et al. 2009, Robinson et al. 2011, Serezino 2015, Santos et al. 2017).

2.4.3 Nitrate uptake process is downregulated by ammonium in N-replete *Brachypodium distachyon* plants

By the availability of inorganic N in the soil, plant roots respond promptly by regulating nitrate or ammonium transporters, by generally inducing N influx under N limitation or inhibiting N influx under excess of this nutrient (Glass et al. 2002, Britto and Kronzucker 2013). However, the plant N status is not the only mechanism for regulating the acquisition of this nutrient. The inhibitory effect of ammonium upon nitrate uptake has been described in several species (Krapp et al. 1998, Cerezo et al. 2001, Krouk et al. 2006). Conversely, short term provision of nitrate can cause an induction of nitrate HATS activity in roots (Gansel et al. 2001, Okamoto et al. 2003, Krouk et al. 2006). To determine the effect of ammonium or nitrate supply in root nitrate uptake, *Brachypodium* plants were grown in the presence of NH_4NO_3 and then transferred to nutrient solution containing either nitrate or ammonium as the sole inorganic N source.

Bd21 roots of both treatments maintained the same level of ^{15}N -nitrate influx at the initial period (up to 4 h) (Figure 10). In the presence of ammonium, the ^{15}N -nitrate influx ($6.29 \mu\text{mol h}^{-1} \text{g}^{-1} \text{DW}$) decreased significantly after 24 h compared to 0 h ($21.38 \mu\text{mol h}^{-1} \text{g}^{-1} \text{DW}$). Bd21 plants exposed to exogenous nitrate increased the ^{15}N -nitrate influx significantly at 24 h ($69.24 \mu\text{mol h}^{-1} \text{g}^{-1} \text{DW}$), being 3.23-fold higher compared to 0 h (Figure 10). Contrary to what was observed in the presence of nitrate, the addition of ammonium did not upregulate the ^{15}N -nitrate uptake process. Therefore, there is a negative feedback regulation of nitrate uptake process by ammonium in roots of N-replete *Brachypodium distachyon* plants.

Figure 10. $^{15}\text{NO}_3$ influx in N replete *Brachypodium distachyon* roots after nitrate or ammonium supply. Plants roots were grown under adequate N supply (2 mM NH_4NO_3) were transferred to 4 mM NO_3^- (4 mM KNO_3) or 4 mM NH_4^+ (4 mM NH_4Cl) for 0, 2, 4 or 24 h. Uptake rates were measured for 10 min. into *B. distachyon* roots at the external concentration of 200 μM ^{15}N -labeled NO_3^- (K^{15}NO_3). Values are means of 5 biological replicates $\pm$ SD, and significant differences at $P < 0.05$ are indicated by different letters.

To better understand the negative regulation of the nitrate uptake process and the time-dependent response of major nitrate transporter genes by ammonium supply, expression analyses of the *BdNRT2* and *BdNRT3* subfamily genes, responsible for the absorption of nitrate in the HATS system, were carried out in *Bd21* plants under both supply treatments. There is a rapid and strong upregulation of *BdNRT2.1* expression by supplying nitrate (Figure 11). After 2 h of nitrate exposure, *BdNRT2.1* expression is 9-fold higher than at 0 h. Such upregulation remains increasing, reaching a relative expression level 500-fold higher compared to the initial time (0 h) after 24 h (Figure 11). On the other hand, ammonium supply led to a weak and transient induction of *BdNRT2.1* (Figure 11). A short increase in the transcript accumulation was observed after 2 h after the exposure to ammonium. In the subsequent periods, there was a significant decrease of *BdNRT2.1* transcript accumulation. After 24 h, the expression level of *BdNRT2.1* was lower than observed at the initial period (0 h) (Figure 11). Comparing the *BdNRT2.1* expression profile upon nitrate or ammonium supply, it becomes evident that the accumulation of *BdNRT2.1* transcripts by ammonium was minimal when compared to the upregulation after nitrate treatment (Figure 11). Studies using *Arabidopsis* showed that *NRT2.1* and *NAR2.1* (*NRT3.1*) are repressed in the presence of ammonium, reducing the role of HATS for nitrate acquisition (Krouk et al. 2006), corroborating the results obtained for *B. distachyon* roots.

Figure 11. Regulation of *BdNRT2.1* and *BdNRT2.2* in N replete *Brachypodium distachyon* roots after nitrate or ammonium supply. Plants roots grown under adequate N supply (2 mM NH_4NO_3) were transferred to 4 mM NO_3^- (4 mM KNO_3) or 4 mM NH_4^+ (4 mM NH_4Cl) for 0, 2, 4 or 24 h for estimating the relative level of transcripts (RT-qPCR). Expression was normalized for mean expression values for plants under N replete (0 h). *BdSamDC* was used gene reference. Bars represent $\pm$ SE; $n=5$ plants per time-point. Significant differences at $P < 0.05$ are indicated by different letters.

Due to the importance of *NRT2.2* in the nitrate uptake process (Cerezo et al. 2001), the regulation of its expression in *B. distachyon* in response to N inorganic sources supply was also evaluated. The same pattern of gene expression regulation was observed for *BdNRT2.2* (Figure 11). There is a significant *BdNRT2.2* transcript accumulation, reaching the highest level after 24 h of exposure to nitrate, 70-fold higher than at 0 h (Figure 11). As observed for

BdNRT2.1, the ammonium supply led to a weak and transient induction of *BdNRT2.2* gene expression (Figure 11), with a small accumulation of transcripts after 4 h of treatment. Later, the level of transcript accumulation was the same observed at the initial period (0 h) (Figure 11). The expression data show that, the induction of *BdNRT2.2* by nitrate was more significant and persistent than observed under ammonium supply (Figure 11). So, compared to nitrate supply treatment, the weak upregulation of *BdNRT2.1* and *BdNRT2.2* expression in Bd21 roots exposed to ammonium show their downregulation by this N inorganic source.

NRT2.3 gene has been described as involved in the nitrate uptake process in other species, such as *Arabidopsis*, rice and barley (Vidmar et al. 2000, Orsel et al. 2002, Tang et al. 2012). In sorghum, tolerance to N deficiency can be associated with higher expression of *NRT2.3* in roots (Gelli et al. 2014). To identify the role of the *NRT2.3* in *B. distachyon* roots, its expression was evaluated. No regulation of this gene was observed in response to ammonium supply (Figure 12). Conversely, nitrate supply led to a transcript accumulation 15-fold higher compared to plants with N sufficiency (0 h) after 4 h. At 24 h, a significant difference was noted in *BdNRT2.3* transcript accumulation comparing both N supply treatments. In the presence of nitrate, *BdNRT2.3* expression was approximately 8-fold higher when compared to plants under ammonium supply (Figure 12). Therefore, it can be concluded that the *BdNRT2.3* gene is downregulated in the presence of ammonium.

Functional analysis in *Xenopus laevis* oocytes and *in planta* revealed that *AtNRT2.4* is a nitrate transporter functioning in the high-affinity range for nitrate uptake (Kiba et al. 2012). *Arabidopsis* triple mutants (*atnrt2.1*, *atnrt2.2* and *atnrt2.4*) demonstrated lower biomass production compared to double mutants (*atnrt2.1* and *atnrt2.2*), evidencing another role of *AtNRT2.4* in shoots (Kiba et al. 2012). Even though it is predominantly expressed in *arabidopsis* leaves (Orsel et al. 2002), its expression in *B. distachyon* roots under N supply treatments was evaluated. A short, but not significant, induction of *BdNRT2.4* expression was observed after 4 h of nitrate supply, when compared to 0 h (Figure 12). At 2 and 24 h after imposition of nitrate treatment, *BdNRT2.4* transcript accumulation was not significantly different from at 0 h. For the ammonium supply, it was not possible to observe a significant difference in *BdNRT2.4* transcript accumulation (Figure 12). Thus, comparing the relative expression of the *BdNRT2.4* in response to supplying the two different inorganic N sources, there is no apparent significant difference in expression in response to either nitrate and/or ammonium exposure (Figure 12).

Figure 12. Regulation of *BdNRT2.3* and *BdNRT2.4* in N replete *Brachypodium distachyon* roots after nitrate or ammonium supply. Plants roots grown under adequate N supply (2 mM NH₄NO₃) were transferred to 4 mM NO₃⁻ (4 mM KNO₃) or 4 mM NH₄⁺ (4 mM NH₄Cl) for 0, 2, 4 or 24 h for estimating the relative level of transcripts (RT-qPCR). Expression was normalized for mean expression values for plants under N replete (0 h). *BdSamDC* was used gene reference. Bars represent $\pm$ SE; $n=5$ plants per time-point. Significant differences at $P < 0.05$ are indicated by different letters.

Analysis of tissue-specific expression of *BdNRT2.5* in *B. distachyon* showed that it is predominantly expressed in roots and spikelets (Wang et al. 2019). Here, comparing both N supply treatments, there is a similar downregulation of *BdNRT2.5* in response to either N provision (Figure 13). Therefore, *BdNRT2.5* appears to not be involved in the nitrate uptake in response to N provision in *B. distachyon*. A similar result was observed in *Arabidopsis*, in

which *AtNRT2.5* expression was strongly repressed by nitrate provision in roots (Okamoto et al. 2003).

Figure 13. Regulation of *BdNRT2.5* and *BdNRT2.6* in N replete *Brachypodium distachyon* roots after nitrate or ammonium supply. Plants roots grown under adequate N supply (2 mM NH₄NO₃) were transferred to 4 mM NO₃⁻ (4 mM KNO₃) or 4 mM NH₄⁺ (4 mM NH₄Cl) for 0, 2, 4 or 24 h for estimating the relative level of transcripts (RT-qPCR). Expression was normalized for mean expression values for plants under N replete (0 h). *BdSamDC* was used gene reference. Bars represent $\pm$ SE; $n=5$ plants per time-point. Significant differences at $P < 0.05$ are indicated by different letters.

Studies revealed that *AtNRT2.6* is weakly expressed in most plant organs (Dechorgnat et al. 2012). Conversely, *bdnrt2.1* mutants showed upregulation of *BdNRT2.6* in roots, suggesting a possible role in reestablishing nitrate absorption in *B. distachyon*. No significant

difference was found in the *BdNRT2.6* transcript accumulation in either N supply treatment (Figure 13). Comparing both treatments, it was only possible to recognize a significant difference after 4 h of treatments; *BdNRT2.6* expression in plants exposed to nitrate was approximately 3.2-fold higher than in those supplied with ammonium (Figure 13).

Despite being predominantly expressed in *Arabidopsis* seeds (Chopin et al. 2007), strong expression of the *NRT2.7* gene has been observed in *B. distachyon* roots (Wang et al. 2019), suggesting a possible function in nitrate uptake. As noted in Figure 14, there was a significant difference in the *BdNRT2.7* expression in *B. distachyon* roots submitted to different N resupply treatments. The nitrate treatment did not lead to any regulation since the expression levels in response to nitrate were similar to those observed in N replete plants. On the other hand, supplementation of ammonium led to a decrease in transcript accumulation, without detection of *BdNRT2.7* expression after 4 h of treatment (Figure 14). Therefore, it can be concluded that ammonium supplementation led to transcription inhibition of *BdNRT2.7* in Bd21 roots.

Figure 14. Regulation of *BdNRT2.7* in N replete *Brachypodium distachyon* roots after nitrate or ammonium supply. Plants roots grown under adequate N supply (2 mM NH₄NO₃) were transferred to 4 mM NO₃⁻ (4 mM KNO₃) or 4 mM NH₄⁺ (4 mM NH₄Cl) for 0, 2, 4 or 24 h for estimating the relative level of transcripts (RT-qPCR). Expression was normalized for mean expression values for plants under N replete (0 h). *BdSamDC* was used gene reference. Bars represent $\pm$ SE; $n=5$ plants per time-point. Significant differences at $P < 0.05$ are indicated by different letters.

HATS has been described in several species as mediated by NRT2 transporters in association with NRT3 (Okamoto et al. 2006, Orsel et al. 2006, Yong et al. 2010, Feng et al. 2011, Yan et al. 2011, Laugier et al. 2012). Two members of the NRT3 family were identified in *B. distachyon* (Plett et al. 2010), which are mainly expressed in roots (David et al. 2019). *BdNRT3.1* expression analysis shows an increase accumulation of transcripts in response to nitrate supply (Figure 15). *BdNRT3.1* transcript accumulation is observed after 2 h of nitrate

provision (2.5-fold), and significantly higher after 4 h of treatment (5-7-fold) in comparison to 0 h (Figure 15). In response to ammonium supply, there is an induction of *BdNRT3.1* expression after 2 h of treatment. The expression level remains stable for the following point analyzed (4 h).

Figure 15. Regulation of *BdNRT3.1* and *BdNRT3.2* in N replete *Brachypodium distachyon* roots after nitrate or ammonium supply. Plants roots grown under adequate N supply (2 mM NH₄NO₃) were transferred to 4 mM NO₃⁻ (4 mM KNO₃) or 4 mM NH₄⁺ (4 mM NH₄Cl) for 0, 2, 4 or 24 h for estimating the relative level of transcripts (RT-qPCR). Expression was normalized for mean expression values for plants under N replete (0 h). *BdSamDC* was used gene reference. Bars represent $\pm$ SE; $n=5$ plants per time-point. Significant differences at $P < 0.05$ are indicated by different letters.

So, comparing both supply treatments, there is no significant difference in *BdNRT3.1* expression over the periods 0, 2 and 4 h; although significant upregulation is observed

comparing its expression in 0 and 4 h (Figure 15). After 24 h, there was a significant distinction of *BdNRT3.1* expression in both N supply treatment. While there is an accumulation of *BdNRT3.1* transcripts in nitrate presence in this period, the expression level under ammonium supply is the same as that observed initially (0 h) (Figure 15). The other member of NRT3 subfamily, *BdNRT3.2* showed the same gene expression regulation as *BdNRT3.1* for both N provision treatments (Figure 15). A significant difference in expression between the N sources was observed only at 24 h, when nitrate supply induces transcript accumulation at 7-fold rate greater when compared to 0 h. In the ammonium supply, the *BdNRT3.2* expression level does not differ from 0 h (Figure 15). Hence, both members of the NRT3 family display a down-regulation in response to ammonium supplementation in Bd21 roots, but not to nitrate.

2.4.4 Regulation of nitrate uptake in *Brachypodium distachyon* roots by N resupply

Physiological studies of nitrate transport in roots of many species demonstrate the importance of HATS uptake this inorganic form of N after nitrate resupply. After the addition of nitrate, there is a rapid induction of the nitrate influx by HATS, denominated inducible HATS (iHATS), which remains induced for a few hours, followed by a rapid drop of nitrate influx in roots (Tompkins et al. 1978, Siddiqi et al. 1990, Tischner et al. 1993, Zhuo et al. 1999, Glass 2003). This HATS induction by exogenous addition of nitrate occurs both in plants pre-cultivated under N sufficiency and in plants grown under deficiency and/or low N (Glass et al., 2002). Differences between species have been identified related to iHATS. For example, in the coniferous *Picea glauca*, the induction of influx of ^{13}N -nitrate occurs after three days of nitrate supply (Kronzucker et al. 1995), whereas, in barley and wheat roots, the largest acquisition of nitrate occurs after 3 h (Tompkins et al. 1978, Siddiqi et al. 1990, Tischner et al. 1993).

To determine the effect of ammonium nitrate or nitrate resupply in root nitrate uptake, *B. distachyon* plants grown in the presence of NH_4NO_3 were submitted to N deficiency for 7 d, when the plants were transferred to a nutrient solution containing either potassium nitrate or ammonium nitrate as the inorganic N source. For the nitrate resupply, temporal analysis of ^{15}N -nitrate influx showed rapid and significant induction of nitrate uptake, since the influx after 2 h ($56.58 \mu\text{mol h}^{-1} \text{g}^{-1} \text{DW}$) was approximately 2.5-fold higher in Bd21 plants submitted to N absence, equivalent to 0 h ($22.36 \mu\text{mol h}^{-1} \text{g}^{-1} \text{DW}$) (Figure 16). ^{15}N -nitrate uptake was significantly higher compared to N restricted plants after 4 h of nitrate resupply treatment ($43.29 \mu\text{mol h}^{-1} \text{g}^{-1} \text{DW}$), reaching approximately 2-fold labeled nitrate absorbed. After 4 h, the influx of ^{15}N -nitrate declines, reaching similar influx levels observed initially (Figure 16). The influx of ^{15}N -nitrate was induced after 2 and 4 h of ammonium nitrate supply, but it was not significantly different from N deficiency. At 24 and 72 h after treatment, ^{15}N -nitrate influx

reached same levels as observed in plants under N restriction. So, ammonium nitrate resupply did not lead to ^{15}N -nitrate influx upregulation in *B. distachyon* roots.

Figure 16. ^{15}N -nitrate influx by *Brachypodium distachyon* submitted to N resupply. Plants grown in N sufficient solution (2 mM NH_4NO_3) were subjected to N restriction for 7 d. Then, plants were cultivated in the presence of nitrate (10 mM KNO_3) or ammonium nitrate (5 mM NH_4NO_3), for 0, 2, 4, 24 and 72 h. Uptake rates were measured for 10 min. into sugarcane roots at the external concentration of 200 μM ^{15}N -labeled NO_3^- (K^{15}NO_3). Values are means of 5 biological replicates $\pm$ SD, and significant differences at $P < 0.05$ are indicated by different letters.

Considering that the HATS regulation directly correlates with the nitrate high-affinity transporters transcription of several species (Lejay et al. 1999, Filleur et al. 2001, Orsel et al. 2006, Li et al. 2007), the analysis of the expression of ortholog genes in *B. distachyon* was performed in root sampled in the previous experiment. *BdNRT2.1* relative expression analysis in response to nitrate resupply exhibits strong upregulation. After 2 h, *BdNRT2.1* transcript accumulation was approximately 110-fold greater compared to at 0 h (Figure 17). Although some decrease in transcript accumulation occurred, this level of transcripts remains significantly higher up to 4 h (approx. 24-fold), while returning to the initial expression level after 24 h. At 72 h after treatment, there is an induction of *BdNRT2.1* expression when compared to the in plants in the absence of N. For the ammonium nitrate treatment resupply, a rapid induction (2 h, 24-fold higher than 0 h) of the *BdNRT2.1* expression is observed (Figure 17), but at a lower intensity compared to the nitrate resupply treatment. In subsequent periods, the transcript accumulation reaches similar levels observed initially (Figure 17). Therefore, the expressive induction of *BdNRT2.1* correlates with the response to nitrate resupply in *B. distachyon* roots.

AtNRT2.1 and *AtNRT2.2* are known to play a key role in regulation of the high-affinity nitrate uptake in Arabidopsis, and in the adaptive responses of the plant to N availability (Cerezo et al. 2001). No significant difference for *BdNRT2.2* transcript accumulation between

both treatments of N resupply was observed (Figure 17). There is a rapid (2 h) mRNA accumulation, since the accumulation was significantly higher when compared to Bd21 plants submitted to N restriction at 0 h. In the following periods, the expression of this gene decreases, reaching after 4 h and 72 h of treatment, levels similar to those observed in plants under N deficiency, (Figure 17). Therefore, *BdNRT2.2* appears to be upregulated briefly by N resupply in *B. distachyon* roots.

Figure 17: Regulation of *BdNRT2.1* and *BdNRT2.2* in *Brachypodium distachyon* (Bd21 genotype) roots submitted to N resupply. Plants grown in N sufficient solution (2 mM NH₄NO₃) were subjected to N restriction for 7 d. Then, plants were cultivated in the presence of nitrate (10 mM KNO₃) or ammonium nitrate (5 mM NH₄NO₃), and root samples were collected 0, 2, 4, 24 and 72 h after treatment. for estimating the relative level of transcripts (RT-

qPCR). Expression was normalized for mean expression values for plants under N deficiency for 7 d (-N, 0 h). *BdSamDC* was used gene reference. Bars represent $\pm$ SE; $n=5$ plants per time-point. Significant differences at $P < 0.05$ are indicated by different letters.

Nitrate resupply led to the late induction of *BdNRT2.3* expression (Figure 18). The expression level remains constant, up to 24 h of resupply of nitrate (Figure 18). After 72 h, the *BdNRT2.3* transcript accumulation was significantly higher (5-fold) than in plants under N deficiency (Figure 18). In Bd21 plants submitted to ammonium nitrate resupply, there is a repression of *BdNRT2.3* expression in the analyzed periods. It was not possible to detect the *BdNRT2.3* mRNA accumulation after 2, 4 and 24 h of treatment. After 72 h, the relative expression was 10-fold lower than observed in Bd21 roots under N restriction (Figure 18). Thus, a strong repression of the *BdNRT2.3* expression occurs in response to ammonium nitrate resupply in Bd21 roots.

The other member of the NRT2 subfamily of *B. distachyon*, *BdNRT2.4*, was also evaluated for the regulation in response to N resupply. *BdNRT2.4* showed a similar pattern of regulation as *BdNRT2.3* in responses to the tested treatments. *BdNRT2.4* was upregulated later after exogenous nitrate supplying, with a 5.5-fold higher expression after 72 h of treatment when compared to Bd21 plants under N deficiency (0 h) (Figure 18). A strong and significant repression of *BdNRT2.4* was observed in plants resupplied with ammonium nitrate (Figure 18). Therefore, *BdNRT2.4* might be down regulated in the presence of ammonium in roots of *B. distachyon*.

Figure 18. Regulation of *BdNRT2.3* and *BdNRT2.4* in *Brachypodium distachyon* (Bd21 genotype) roots submitted to N resupply. Plants grown in N sufficient solution (2 mM NH₄NO₃) were subjected to N restriction for 7 d. Then, plants were cultivated in the presence of nitrate (10 mM KNO₃) or ammonium nitrate (5 mM NH₄NO₃), and root samples were collected 0, 2, 4, 24 and 72 h after treatment. For estimating the relative level of transcripts (RT-qPCR). Expression was normalized for mean expression values for plants under N deficiency for 7 d (-N, 0 h). *BdSamDC* was used gene reference. Bars represent $\pm$ SE; n=5 plants per time-point. Significant differences at $P < 0.05$ are indicated by different letters.

For *BdNRT2.5*, a significant distinction in gene expression between both N resupply treatments is observed (Figure 19). After 2 h of exposure to the treatments, ammonium nitrate resupply showed no significant change in *BdNRT2.5* expression, whereas the presence of nitrate led to a 10-fold increase in *BdNRT2.5* expression, compared to the initial moment (0 h) (Figure 19). At 4 h, in both treatments, the *BdNRT2.5* expression level was similar as those seen in plants under N deficiency. Compared to the 0 h sample, *BdNRT2.5* showed a down-regulation after 24 h of ammonium nitrate treatment, while the expression level remained constant in the nitrate resupply. At 72 h, the *BdNRT2.5* expression in both treatments was similar to that observed initially (Figure 19). Therefore, there was a significant divergence in the *BdNRT2.5* expression regulation in response to N resupply in *B. distachyon* roots.

Gene expression analysis revealed that nitrate resupply positively regulated *BdNRT2.6* expression (Figure 19). After 2 h of treatment, there was a significant upregulation of *BdNRT2.6* expression. At 2 h, the expression was 5-fold higher than observed in plants under N deficiency (Figure 19). Such induction was maintained at 4 and 72 h, although it does not diverge significantly from the initial period analyzed (0 h). On the other hand, *BdNRT2.6* expression was not regulated in *B. distachyon* roots under ammonium nitrate resupply treatment. There is a negative regulation of *BdNRT2.6* expression in Bd21 plants resupplied in the presence of ammonium.

Figure 19. Regulation of *BdNRT2.5* and *BdNRT2.6* in *Brachypodium distachyon* (Bd21 genotype) roots submitted to N resupply. Plants grown in N sufficient solution (2 mM NH₄NO₃) were subjected to N restriction for 7 d. Then, plants were cultivated in the presence of nitrate (10 mM KNO₃) or ammonium nitrate (5 mM NH₄NO₃), and root samples were collected 0, 2, 4, 24 and 72 h after treatment. for estimating the relative level of transcripts (RT-qPCR). Expression was normalized for mean expression values for plants under N deficiency for 7 d (-N, 0 h). *BdSamDC* was used gene reference. Bars represent $\pm$ SE; $n=5$ plants per time-point. Significant differences at $P < 0.05$ are indicated by different letters.

The last member of the NRT2 subfamily evaluated for the regulation of gene expression in response to N resupply was *BdNRT2.7*. After exclusive nitrate resupply, *BdNRT2.7* mRNA accumulation was rapidly induced (2 h), reaching 5-fold higher than initially observed (0 h) (Figure 20). After this early induction, *BdNRT2.7* expression was suppressed after 4 h of

treatment, without detection in the following periods. In plants submitted to ammonium nitrate resupply, strong and continuous repression of the *BdNRT2.7* gene was perceived, without detection of expression after 24 h of this treatment. At 72 h, *BdNRT2.7* transcript accumulation returned to the same level observed in plants under N deficiency (Figure 20). As reported previously for other *BdNRT2* members, such results show a negative regulation of *BdNRT2.7* expression in *Bd21* plants resupplied in the presence of ammonium.

Figure 20. Regulation of *BdNRT2.7* in *Brachypodium distachyon* (*Bd21* genotype) roots submitted to N resupply. Plants grown in N sufficient solution (2 mM NH₄NO₃) were subjected to N restriction for 7 d. Then, plants were cultivated in the presence of nitrate (10 mM KNO₃) or ammonium nitrate (5 mM NH₄NO₃), and root samples were collected 0, 2, 4, 24 and 72 h after treatment. for estimating the relative level of transcripts (RT-qPCR). Expression was normalized for mean expression values for plants under N deficiency for 7 d (-N, 0 h). *BdSamDC* was used gene reference. Bars represent $\pm$ SE; $n=5$ plants per time-point. Significant differences at $P < 0.05$ are indicated by different letters.

Expression analyses of the *NRT3* subfamily members, *BdNRT3.1* and *BdNRT3.2*, were also performed in response to N resupply in *B. distachyon*. *BdNRT3.1* was promptly up regulated after 2 h of exposure to both treatments, but at different intensities (Figure 21). While the induction in response to ammonium nitrate resupply was around 2-fold higher when compared to the one at 0 h, for plants under nitrate resupply the *BdNRT3.1* transcript accumulation was approximately 50-fold higher (Figure 21). Comparing with *BdNRT3.1* expression observed in N deficient plants (0 h), *BdNRT3.1* expression in the presence of nitrate remains elevated (15-fold) after 4 h of treatment (Figure 21). After 4 h of ammonium nitrate treatment, the *BdNRT3.1* expression level returned to the level at 0 h. The *BdNRT3.1* transcript accumulation remained significantly high (2.5-fold) in plants submitted to nitrate resupply, when compared to time 0 h, after 24 h. For the same period, *BdNRT3.1* was down regulated (10-fold) in ammonium nitrate resupply. For the last period analyzed (72 h),

induction of *BdNRT3.1* expression could be recognized in nitrate resupply, whereas in Bd21 plants submitted to ammonium nitrate treatment, transcript accumulation was similar to observed in the 0 h period (Figure 21).

Figure 21. Regulation of *BdNRT3.1* and *BdNRT3.2* in *Brachypodium distachyon* (Bd21 genotype) roots submitted to N resupply. Plants grown in N sufficient solution (2 mM NH₄NO₃) were subjected to N restriction for 7 d. Then, plants were cultivated in the presence of nitrate (10 mM KNO₃) or ammonium nitrate (5 mM NH₄NO₃), and root samples were collected 0, 2, 4, 24 and 72 h after treatment for estimating the relative level of transcripts (RT-qPCR). Expression was normalized for mean expression values for plants under N deficiency for 7 d (-N, 0 h). *BdSamDC* was used gene reference. Bars represent $\pm$ SE; $n=5$ plants per time-point. Significant differences at $P < 0.05$ are indicated by different letters.

Expression analyses of *BdNRT3.2* display the same pattern of regulation as *BdNRT3.1* in response to both treatments of N resupply (Figure 21). Compared to expression levels

present in Bd21 plants under N deficiency (0 h), *BdNRT3.2* was strongly and rapidly upregulated (2 h) in response to nitrate resupply, maintaining significant higher relative expression levels in the following periods. The treatment ammonium nitrate resupply led to a weak induction of *BdNRT3.2* expression after 2 h. After that, there was a downregulation of *BdNRT3.2* expression, with transcript accumulation after 4 h similar to at 0 h, or significantly lower levels afterwards (24 and 72 h) (Figure 21). Therefore, it can be concluded that both members of the NRT3 subfamily, *BdNRT3.1* and *BdNRT3.2*, are regulated equally in response to N resupply in *Brachypodium distachyon*.

2.5 Discussion

Brachypodium distachyon has been established as a genetic model plant for studies of grass species due to the available biological resources available and its evolutionary proximity to the top global agricultural commodities (Draper et al. 2001, Bevan et al. 2010, Brkljacic et al. 2011, Bragg et al. 2012, Brutnell et al. 2015, Kellogg 2015b, Lyons and Scholthof 2015, Lyons and Scholthof 2016, Vogel 2016, Hsia et al. 2017, Sibout et al. 2017, Scholthof et al. 2018). With the immediate need for eco-friendly food production, studies with *B. distachyon* aiming at the improvement of nitrogen use efficiency (NUE) nicely match this requirement. The nitrogen uptake efficiency from soils is one of the essential processes for NUE, since N is the agent of relevant environmental impacts caused by agriculture (Dirzo and Raven 2003, Martinelli and Filoso 2008, Vitousek et al. 2009, Burney et al. 2010, Foley et al. 2011).

Despite the significance of N as a limiting nutrient for plant growth, few investigations have been conducted to clearly define the preferential inorganic N source for *B. distachyon* (Boudsocq et al. 2012). This information is basic to adopt *Brachypodium distachyon* as a model species in functional genomic studies of genes associated with N uptake from related grasses. Short-term uptake analysis using ^{15}N -ammonium and ^{15}N -nitrate under contrasting N conditions in *Brachypodium distachyon* roots revealed its preference for ammonium instead of nitrate as inorganic N source, similarly to other species (Figure 9). Another similarity found was that HATS induction in *B. distachyon* roots under N deficiency and/or low N status is a conserved mechanism similar to other monocots and dicots species (von Wirén et al. 2000, Camanès 2009, Subbarao et al. 2009, Robinson et al. 2011, Serezino 2015, Santos et al. 2017).

NRT2 and NRT3 transporters and their regulation have been extensively characterized in several species (Crawford and Glass 1998, Glass et al. 2002, Hildebrandt et al. 2002, Okamoto et al. 2003, Orsel et al. 2006, Li et al. 2007, Cai et al. 2008, Plett et al. 2010, Feng et al. 2011, Dechorgnat et al. 2011, Yan et al. 2011; Laugier et al. 2012, Xu et al. 2012, Bai et al. 2013, Garnett et al. 2013, Lupini et al. 2016, Havé et al. 2017, Bajgain et al. 2018). However, *NRT2* and *NRT3* genes from grasses seem to be relatively distant phylogenetically

from those in *A. thaliana* (Figure 4 and 7), making it difficult to speculate about possible orthologues of the NRT2 and NRT3 transporters in grasses to those that have been functionally characterized in Arabidopsis. Among the BdNRT2 family members, BdNRT2.1 and BdNRT2.2 displayed important similarities in terms of their gene structure and amino acid sequence (Figure 1, Figure 3, and Table 2). Furthermore, as described in Arabidopsis, BdNRT2.1 and BdNRT2.2 appear as clustered genes in opposite orientation in the genome, which could suggest a common gene expression pattern and/or evolutionary origin (Figure 2; Cerezo et al. 2001). Additionally, both proteins exhibit common transmembrane structure (Figure 5), inferring that both are high-affinity NO_3^- transporters in *B. distachyon* roots, similar to their orthologous in other species (Forde et al. 2000, Plett et al. 2010). Both members of the BdNRT3 family also share high similarities of gene structures, amino acid sequences and protein topology with orthologues in other species (Figures 1, 6, and 8). However, different from Arabidopsis, *BdNRT3.1* and *BdNRT3.2* have a head-to-tail configuration regarding their chromosomal position (Figure 2; Plett et al. 2010). Therefore, this could indicate distinct functions and regulation of *BdNRT3.1* and *BdNRT3.2* genes compared to their orthologous in Arabidopsis, suggesting both genes are involved in nitrate uptake process.

Analyzing the expression levels for members of the *BdNRT2* gene family in response to inorganic sources of N provision, it was possible to observe significant repression of *BdNRT2.1*, *BdNRT2.2*, *BdNRT2.3* and *BdNRT2.7* in plants under ammonium supply treatment. Upregulation of *NRT2.1* transcript abundance by nitrate and its downregulation by the presence of ammonium have been described in many species (Quesada et al. 1997, Krapp et al. 1998, Zhuo et al. 1999, Cerezo et al. 2001, Nazoa et al. 2003, Okamoto et al. 2003, Laugier et al. 2012). Our data reveal that not only *BdNRT2.1* but also the *BdNRT2.2* are strongly repressed in the presence of ammonium, since there was no increase in transcript accumulation similarly noticed in plants exposed to nitrate supply (Figure 11). *NRT2.3* has been described as involved in the nitrate uptake in other species, such as arabidopsis, rice and barley (Vidmar et al. 2000, Orsel et al. 2002, Tang et al. 2012). In tomato, *LeNRT2.3* expression was remarkably repressed upon ammonium exposure (Hildebrandt et al. 2002). Roots of *B. distachyon* plants exposed to nitrate supply showed upregulation of *BdNRT2.3* (Figure 12). This result is the opposite to what was described by Wang et al. (2019), who defined *BdNRT2.3* as nitrate constitutive gene. Still, according to Wang et al (2019), another nitrate constitutive gene is *BdNRT2.7*. The expression of *BdNRT2.7* in *B. distachyon* roots may suggest a role in nitrate uptake, distinct from *AtNRT2.7* function (Chopin et al. 2007). *BdNRT2.7* was strongly downregulated in Bd21 plants submitted to ammonium supply, showing that this gene is differently regulated according to the inorganic N source (Figure 14).

For both members of the NRT3 family, *BdNRT3.1* and *BdNRT3.2*, there is a significant distinction in their gene expression profile only after 24 h provision of the different sources of

N. While both genes were significantly induced in the presence of nitrate, ammonium supply treatment led to their downregulation, returning *BdNRT3.1* and *BdNRT3.2* transcripts abundance similar to those noted in plants under N deficiency (Figure 15). These results indicate the negative co-regulation exerted by ammonium on both *NRT2* and *NRT3* gene family members of *B. distachyon* in response to the provision of this source. As reported for other species, an inhibitory effect of ammonium on ^{15}N -nitrate uptake in *B. distachyon* roots had been also observed (Figure 10; Krapp et al. 1998, Cerezo et al. 2001, Krouk et al. 2006). Correlating the ^{15}N influx with the expression analyses of *BdNRT2* and *BdNRT3* gene family members, it is possible to suggest that nitrate uptake inhibition occurs by the repression of the expression of both family members in N-sufficient *B. distachyon* under ammonium supply. The occurrence of this nitrate high-affinity transport downregulation by ammonium has been described in many species as an essential mechanism to mitigate the toxic effects caused by ammonium in plant development, even in species that favor ammonium than nitrate uptake (Givan 1979, Hageman 1984, Salsac et al. 1987, Bloom et al. 1992, Raven et al. 1992, Britto et al. 2001, Kronzucker et al. 2001, Britto and Kronzucker 2002, Rahayu et al. 2005). The repression of the ^{15}N -nitrate influx is directly correlated with the *BdNRT2.1*, *BdNRT2.2*, *BdNRT3.1*, and *BdNRT3.2* expression downregulation in response to ammonium supply treatment (Figures 10, 11, and 15). Unlikely the mechanism proposed in arabidopsis, in which ammonium inhibits nitrate HATS transport via *AtNRT2.1* and *AtNAR2.1* (*AtNRT3.1*), in *B. distachyon* it is possible to recognize the redundancy of these genes in each subfamily (Krouk et al. 2006). In other words, not only the *BdNRT2.1* and *BdNRT3.1* genes would be essential for nitrate uptake in roots, but also the *BdNRT2.2* and *BdNRT3.2* genes may also play a significant role in this process.

BdNRT2 gene expression analysis in N-deficient *B. distachyon* plants under N resupply treatments revealed strong and fast (2 h) upregulation of *BdNRT2.1* and *BdNRT2.2* expressions in response to both means of N resupply (Figure 17). However, unlikely what was observed for *BdNRT2.2*, such gene expression induction was significantly distinct comparing the *BdNRT2.1* transcript accumulation in both N resupply treatments. In the presence of ammonium, the expression after 2 h of treatment was significantly lower compared to the exposure to nitrate only (Figure 17). Levels of *AtNRT2.1* transcript also decreases in response to the coexistence of nitrate and ammonium, as compared to what is observed with nitrate alone (Muños et al. 2004, Laugier et al. 2012), showing a conserved downregulation mechanism of *NRT2.1* expression by ammonium among both species. *NRT2.5* was described as the main nitrate transporter in roots of Arabidopsis plants under N deficiency (Lezhneva et al. 2014). A possible role of *BdNRT2.5* in nitrate uptake in *Brachypodium distachyon* roots was proposed by Wang et al. (2019). Contrary to what was shown, under our hands *BdNRT2.5* expression was promptly upregulated in plants submitted to nitrate resupply. In addition, we

observed *BdNRT2.5* downregulation in response to ammonium resupply treatment (Figure 19). These results suggest a possible role of *BdNRT2.5* in nitrate uptake in plants under N resupply. *BdNRT2.6* expression evaluation demonstrated a significant upregulation in *Brachypodium distachyon* roots after 2 h of nitrate resupply treatment. Upon ammonium exposure, *BdNRT2.6* expression was constitutive (Figure 19). Due to the expression pattern of *BdNRT2.6* in response to N resupply, it can be concluded that *BdNRT2.6* expression regulation differs from its ortholog in *Arabidopsis* (Okamoto et al. 2003, Dechorgnat et al. 2012), proposing a role of this transporter in nitrate uptake in roots of *B. distachyon*.

The first grass *NRT2.7*-like gene described exhibited differential regulation of expression comparing both N resupply treatments. It was possible to observe not only the rapid induction of *BdNRT2.7* expression in response to nitrate resupply, but also its repression under ammonium resupply treatment in *Brachypodium* plants (Figure 20). Interestingly, after 72 h of ammonium nitrate resupply, the *BdNRT2.7* expression restored to that observed in N-deficient plants. The regulation of *BdNRT2.7* expression, added to its specific tissue expression pattern not only in leaves, but also in roots (Wang et al. 2019), indicates a possible function of *BdNRT2.7* in nitrate uptake in *B. distachyon* plants under conditions of nitrate resupply. Also, as no correlation with the influx of ^{15}N -nitrate in plants submitted to ammonium nitrate treatment was observed, these results suggest that the *BdNRT2.7* may be acting in the nitrate remobilization/translocation to the aerial part of *Brachypodium distachyon* plants submitted to ammonium resupply. Considering that its ortholog in *Arabidopsis* participates exclusively in the storage of nitrate in seeds (Chopin et al. 2007, David et al. 2014), *BdNRT2.7* may participate in a different and/or additional process in this species.

BdNRT3 expression evaluation in *B. distachyon* roots in response to N resupply treatments revealed a rapid (2 h) and significant upregulation, while nitrate was the sole source of N. In the presence of ammonium in resupply treatment, *BdNRT3.1* and *BdNRT3.2* also exhibited a quick induction of expression, however at a lower rate than observed in plants exposed only to nitrate (Figure 13). Thus, there is a downregulation mechanism of *BdNRT3.1* and *BdNRT3.2* transcript accumulation by ammonium in plants submitted to N resupply. Reduction of mRNA accumulation of *NAR2.1* (*NRT3.1*) in plants under NH_4NO_3 resupply has been described in *Arabidopsis* and rice, revealing a conserved regulation mechanism among monocots and dicots species (Yan et al. 2011, Laugier et al. 2012). Similar to what was observed in N-sufficient *B. distachyon* plants, negative co-regulation of *BdNRT2* and *BdNRT3* gene expression exerted by ammonium was observed in N-deficient plants subjected to resupply in the presence of this N inorganic source. ^{15}N -nitrate influx analyzes revealed that in the presence of ammonium there was no significant change of the nitrate uptake process from that seen in plants submitted to nitrate-only resupply (Figure 16).

Therefore, as already described for other species, the transcript accumulation of NRT2 and NRT3 family gene members is directly correlated with the high-affinity nitrate transport system regulation in *B. distachyon* (Lejay et al. 1999, Filleur et al. 2001, Orsel et al. 2006, Li et al. 2007). Under field growing conditions, the use of N-fertilizer in the form of urea and ammonium nitrate, is a common practice adopted to increase sugarcane yield (Wood et al. 2010). Hence, the identification of ammonium-dependent repression of nitrate uptake in this model grass plant could explain the low efficiency of nitrate acquisition in roots of sugarcane compared to other species (Robinson et al. 2011), and the process of post-transcriptional regulation of HATS transport in sugarcane would affect nitrate acquisition in this grass roots.

2.6 Conclusion

Overall, this work highlights the preference for ammonium as a source of inorganic N by *B. distachyon*, similar to what was observed in other grasses. Besides, this study contributes to the identification and characterization of candidate genes involved in the HATS nitrate uptake process in this species. *BdNRT2.1* and *BdNRT2.2* exhibit a similar expression regulation pattern, according to the N treatment exposure. This suggests the redundancy of both genes in the nitrate uptake process. The same was observed in the *BdNRT3.1* and *BdNRT3.2* suggesting, in addition to the redundancy of *BdNRT3*, but also the co-regulation of the NRT2 and NRT3 families in *B. distachyon*.

In N supply and N resupply treatments, possible repression of *BdNRT2.1*, *BdNRT2.2*, *BdNRT3.1*, and *BdNRT3.2* expression was observed in Bd21 plants exposed to ammonium. Correlation between the corresponding patterns of the *BdNRT2/BdNRT3* gene expressions and the patterns of the $^{15}\text{NO}_3^-$ influx in response to N availability demonstrates that these gene members are likely responsible for HATS nitrate uptake process in *B. distachyon* roots. The knowledge obtained from this work identified potential gene targets for functional analysis of these conserved transporters in the model grass *B. distachyon* Bd21 and could help to elucidate the specific roles of each transporter in monocot plants as well as subsequently, to be used to improve NUE in grass species.

References

- BAI H, EURING D, VOLMER K, et al. (2013) The nitrate transporter (NRT) gene family in poplar. *PLoS One*, 8: e72126. DOI: 10.1371/journal.pone.0072126
- BAJGAIN P, RUSSELL B, and MOHAMMADI M (2018) Phylogenetic analyses and in-seedling expression of ammonium and nitrate transporters in wheat. *Sci Rep*, 8: 7082. DOI: 10.1038/s41598-018-25430-8
- BARRERO JM, JACOBSEN JV, TALBOT MJ, et al. (2012). Grain dormancy and light quality effects on germination in the model grass *Brachypodium distachyon*. *New Phytol*, 193: 376–386. DOI: 10.1111/j.1469-8137.2011.03938.x
- BEVAN MW, GARVIN DF, and VOGEL JP (2010) *Brachypodium distachyon* genomics for sustainable food and fuel production. *Curr Opin Biotech*, 21: 211–217. DOI: 10.1016/j.copbio.2010.03.006
- BLOOM AJ, SUKRAPANNA SS, and WARNER RL (1992) Root respiration associated with ammonium and nitrate absorption and assimilation by barley. *Plant Physiol*, 99: 1294–1301. DOI: 10.1104/pp.99.4.1294
- BODIRSKY BL, POPP A, WEINDL I, et al. (2012) N₂O emissions from the global agricultural nitrogen cycle - current state and future scenarios. *Biogeosciences*, 9(10): 4169–4197. DOI: 10.5194/bg-9-4169-2012
- BOUDSOCQ S, NIBOYET A, LATA JC, et al. (2012) Plant preference for ammonium versus nitrate: a neglected determinant of ecosystem functioning? *Am Nat*, 180: 60–69. DOI: 10.1086/665997
- BOSCHIERO BN, MARIANO E, and TRIVELIN PCO (2018) Preferential ammonium uptake by sugarcane does not increase the 15N recovery of fertilizer sources. *Plant Soil*, 429: 253–269. DOI: 10.1007/s11104-018-3672-z
- BRAGG JN, WU J, GORDON SP, GUTTMAN ME, et al. (2012) Generation and characterization of the Western Regional Research Center *Brachypodium* T-DNA insertional mutant collection. *PLoS One*, 7: e41916. DOI: 10.1371/journal.pone.0041916
- BRITTO DT, SIDDIQI MY, GLASS ADM, and KRONZUCKER HJ (2001) Futile transmembrane NH₄⁺ cycling: a cellular hypothesis to explain ammonium toxicity in plants. *Proc Natl Acad Sci USA*, 98: 4255–4258. DOI: 10.1073/pnas.061034698
- BRITTO DT and KRONZUCKER HJ (2002) NH₄⁺ toxicity in higher plants: a critical review. *J Plant Physiol*, 159: 567–584. DOI: 10.1078/0176-1617-0774
- BRITTO DT and KRONZUCKER HJ (2013) Ecological significance and complexity of N-source preference in plants. *Ann Bot*, 112: 957–963. DOI:10.1093/aob/mct157
- BRKLJACIC J, GROTEWOLD E, SCHOLL R, et al. (2011) *Brachypodium* as a model for the grasses: today and the future. *Plant Physiol*, 157: 3–13. DOI: 10.1104/pp.111.179531
- BRUTNELL TP, BENNETZEN JL, and VOGEL JP (2015). *Brachypodium distachyon* and *Setaria viridis*: Model genetic systems for the grasses. *Annu Rev Plant Biol*, 66: 465–485. DOI: 10.1146/annurev-arplant-042811-105528
- BURNEY JA, DAVIS SJ, and LOBELL DB (2010) Greenhouse gas mitigation by agricultural intensification. *Proc Natl Acad Sci USA*, 107: 12052–12057. DOI: 10.1073/pnas.0914216107
- CAI CH, WANG JY, ZHU YG, et al. (2008) Gene structure and expression of the high-affinity nitrate transport system in rice roots. *J Integr Plant Biol*, 50: 443–451. DOI: 10.1111/j.1744-7909.2008.00642.x.
- CAMAÑES G, CEREZO M, PRIMO-MILLO E, et al. (2009) Ammonium transport and CitAMT1 expression are regulated by N in Citrus plants. *Planta*, 229: 331–342. DOI 10.1007/s00425-008-0833-y
- CANFIELD DE, GLAZER AN, and FALKOWSKI PG (2010) The Evolution and Future of Earth's Nitrogen Cycle. *Science*, 330: 192–196. DOI: 10.1126/science.1186120
- CEREZO M, TILLARD P, FILLEUR S, et al. (2001) Major alterations of the regulation of root NO₃⁻ uptake are associated with the mutation of NRT2.1 and NRT2.2 genes in *Arabidopsis*. *Plant Physiol*, 127: 262–271. DOI: 10.1104/pp.127.1.262
- CHEN L and LIAO H (2017) Engineering crop nutrient efficiency for sustainable agriculture. *J Integr Plant Biol*, 59(10): 710–735. DOI: 10.1111/jipb.12559
- CHOPIN F, ORSEL M, DORBE MF, et al. (2007). The *Arabidopsis* ATNRT2.7 nitrate transporter controls nitrate content in seeds. *Plant Cell*, 19:1590–1602. DOI: 10.1105/tpc.107.050542
- CRAWFORD NM and GLASS ADM (1998) Molecular and physiological aspects of nitrate uptake in plants. *Trends Plant Sci*, 3: 389–395. DOI: 10.1016/S1360-1385(98)01311-9
- DAVID LC, DECHORGNAT J, BERQUIN P, et al. (2014) Proanthocyanidin oxidation of *Arabidopsis* seeds is altered in mutant of the high-affinity nitrate transporter NRT2.7. *J Exp Bot*, 65: 885–893. DOI: 10.1093/jxb/ert481

- DAVID LC, GIRIN T, FLEURISSEON E, et al. (2019) Developmental and physiological responses of *Brachypodium distachyon* to fluctuating nitrogen availability. *Sci Rep*, 9: 3824. DOI: 10.1038/s41598-019-40569-8
- DECHORGNAT J, NGUYEN CT, ARMENGAUD P, et al. (2011) From the soil to the seeds: The long journey of nitrate in plants. *J Exp Bot*, 62:1349-1359. DOI: 10.1093/jxb/erq409
- DECHORGNAT J, PATRIT O, KRAPP A, et al. (2012) Characterization of the NRT2.6 Gene in *Arabidopsis thaliana*: A Link with Plant Response to Biotic and Abiotic Stress. *PLoS One* 7(8): e42491. DOI:10.1371/journal.pone.0042491
- DIRZO R and RAVEN PH (2003) Global State of Biodiversity and Loss. *Annu Rev Env Resour*, 28:137-167. DOI: 10.1146/annurev.energy.28.050302.105532
- DRAPER J, MUR LAJ, JENKINS G, et al. (2001) *Brachypodium distachyon*: A new model system for functional genomics in grasses. *Plant Physiol*, 127: 1539–1555. DOI: 10.1104/pp.010196
- FENG H, YAN M, FAN X, LI B, et al. (2011) Spatial expression and regulation of rice high-affinity nitrate transporters by nitrogen and carbon status. *J Exp Bot*, 62: 2319–2332. DOI: 10.1093/jxb/erq403
- FILLEUR S, DORBE MF, CERESO M, et al. (2001) An *Arabidopsis* T-DNA mutant affected in NRT2 genes is impaired in nitrate uptake. *FEBS Lett*, 489: 220-224. DOI: 10.1016/S0014-5793(01)02096-8
- FOLEY J, RAMANKUTTY N, BRAUMAN K, et al. (2011) Solutions for a cultivated planet. *Nature*, 478: 337-342. DOI:10.1038/nature10452
- FORDE BG (2000) Nitrate transporters in plants: structure, function and regulation. *Biochim Biophys Acta*, 1465: 219–235. DOI: 10.1016/S0005-2736(00)00140-1
- GANSEL X, MUNOS S, TILLARD P, and GOJON A (2001) Differential regulation of the NO_3^- and NH_4^+ transporter genes AtNRT2.1 and AtAMT1.1 in *Arabidopsis*: relation with long-distance and local controls by N status of the plant. *Plant J*, 26: 143-155. DOI: 10.1046/j.1365-313x.2001.01016.x
- GARNETT T, CONN V, PLETT D, et al. (2013) The response of the maize nitrate transport system to nitrogen demand and supply across the lifecycle. *New Phytol*, 198:82-94. DOI: 10.1111/nph.12166
- GELLI M, DUO Y, KONDA AR, et al. (2014) Identification of differentially expressed genes between sorghum genotypes with contrasting nitrogen stress tolerance by genome-wide transcriptional profiling. *BMC Genomics*, 15: 179. DOI: 10.1186/1471-2164-15-179
- GIRIN T, DAVID LC, CHARDIN C, et al. (2014) *Brachypodium*: a promising hub between model species and cereals. *J Exp Bot*, 65: 5683–5696. DOI: 10.1093/jxb/eru376
- GIVAN CV (1979) Metabolic Detoxification of Ammonia in Tissues of Higher-Plants. *Phytochemistry*, 18: 375-382. DOI: 10.1016/S0031-9422(00)81870-1
- GLASS ADM, BRITTO DT, KAISER BN, et al. (2002) The regulation of nitrate and ammonium transport systems in plants. *J Exp Bot*, 53: 855–864. DOI: 10.1093/jexbot/53.370.855
- GLASS ADM (2003) Nitrogen Use Efficiency of Crop Plants: Physiological Constraints upon Nitrogen Absorption. *Crit Rev Plant Sci*, 22: 453-470. DOI: 10.1080/07352680390243512
- GOOD AG, SHRAWAT AK, and MUENCH DG (2004) Can less yield more? Is reducing nutrient input into the environment compatible with maintaining crop production? *Trends Plant Sci*, 9: 597–605. DOI: 10.1016/j.tplants.2004.10.008
- HAGEMAN RH (1984) Ammonium versus nitrate nutrition of higher plants. *Nitrogen in Crop Production*, p 67-85. DOI: 10.2134/1990.nitrogenincropproduction.c4
- HAVÉ M, MARMAGNE A, CHARDON F, and MASCLAUX-DAUBRESSE C (2017) Nitrogen remobilization during leaf senescence: lessons from *Arabidopsis* to crops. *J Exp Bot*, 68 (10): 2513–2529. DOI: 10.1093/jxb/erw365
- HILDEBRANDT U, SCHMELZER E, and BOTHE H (2002). Expression of nitrate transporter genes in tomato colonized by an arbuscular mycorrhizal fungus. *Physiol Plant*, 115: 125-136. DOI: 10.1034/j.1399-3054.2002.115.0115.x
- HOAGLAND DR and ARNON DI (1950) The water culture method for growing plants without soil. University of California, Agriculture Experimental Station (Circular 347).
- HOWARTH R, SWANEY D, BILLEN G, GARNIER J, et al. (2011) Nitrogen fluxes from the landscape are controlled by net anthropogenic nitrogen inputs and by climate. *Front Ecol Environ*, 10(1): 37-43. DOI: 10.1890/100178
- HSIA MM, O'MALLEY R, CARTWRIGHT A, et al. (2017) Sequencing and functional validation of the JGI *Brachypodium distachyon* T-DNA collection. *Plant J*, 91: 361–370. DOI: 10.1111/tpj.13582
- ISHIKAWA S, ITO Y, SATO Y, et al. (2009) Two-component high-affinity nitrate transport system in barley: membrane localization, protein expression in roots and a direct protein–protein interaction. *Plant Biotechnol*, 26: 197–205. DOI: 10.5511/plantbiotechnology.26.197

- KAWACHI T, SUNAGA Y, EBATO M, et al. (2006) Repression of nitrate uptake by replacement of Asp105 by asparagine in AtNRT3.1 in *Arabidopsis thaliana* L. *Plant Cell Physiol*, 47: 1437–1441. DOI: 10.1093/pcp/pcl010
- KELLOGG EA (2015b). *Brachypodium distachyon* as a genetic model system. *Annu Rev Genet*, 49: 1–20. DOI: 10.1146/annurev-genet-112414-055135
- KIBA T, FERIA-BOURRELLIER AB, LAFOUGE F, et al. (2012) The Arabidopsis nitrate transporter NRT2.4 plays a double role in roots and shoots of nitrogen-starved plants. *Plant Cell*, 24: 245–258. DOI: 10.1105/tpc.111.092221
- KOTUR Z, MACKENZIE N, RAMESH S, et al. (2012) Nitrate transport capacity of the Arabidopsis thaliana NRT2 family members and their interactions with AtNAR2.1. *New Phytol*. 194: 724–731. DOI: 10.1111/j.1469-8137.2012.04094.x
- KRAPP A, FRAISIER V, SCHEIBLE WR, et al. (1998) Expression studies of Nrt2;1Np, a putative high-affinity nitrate transporter: evidence for its role in nitrate uptake. *Plant J*, 14: 723–731. DOI: 10.1046/j.1365-313x.1998.00181.x
- KRONZUCKER HJ, SIDDIQI MY, and GLASS ADM (1995) Kinetics of NO_3^- influx in spruce. *Plant Physiol*, 109: 319–326. DOI: 10.1104/pp.109.1.319
- KRONZUCKER HJ, BRITTO DT, DAVENPORT RJ, and TESTER M (2001) Ammonium toxicity and the real cost of transport. *Trends Plant Sci*, 6: 335–337. DOI: 10.1016/S1360-1385(01)02022-2
- KROUK G, TILLARD P, and GOJON A (2006) Regulation of the high-affinity NO_3^- uptake system by NRT1.1-mediated NO_3^- demand signaling in Arabidopsis. *Plant Physiol*, 142: 1075–1086. DOI: 10.1104/pp.106.087510
- LAUGIER E, BOUGUYON E, MAURIES A, et al. (2012) Regulation of high-affinity nitrate uptake in roots of Arabidopsis depends predominantly on posttranscriptional control of the NRT2.1/NAR2.1 transport system. *Plant Physiol*. 158, 1067–1078. DOI: 10.1104/pp.111.188532
- LEAL GA, ALBUQUERQUE PSB, and FIGUEIRA A (2007) Genes differentially expressed in *Theobroma cacao* associated with resistance to witches' broom disease caused by *Crinipellis pernicioso*. *Mol Plant Pathol*, 8: 279–292. DOI: 10.1111/j.1364-3703.2007.00393.x
- LEJAY L, TILLARD P, LEPETIT M et al. (1999) Molecular and functional regulation of two NO_3^- uptake systems by N- and C-status of Arabidopsis plants. *Plant J*, 18: 509–519. DOI: 10.1046/j.1365-313X.1999.00480.x
- LEZHNEVA L, KIBA T, FERIA-BOURRELLIER AB, et al. (2014) The Arabidopsis nitrate transporter NRT2.5 plays a role in nitrate acquisition and remobilization in nitrogen-starved plants. *Plant J*, 80: 230–241. DOI: 10.1111/tpj.12626
- LI W, WANG Y, OKAMOTO M, et al. (2007) Dissection of the AtNRT2.1: AtNRT2.2 inducible high-affinity nitrate transporter gene cluster. *Plant Physiol*, 143: 425–433. DOI: 10.1104/pp.106.091223
- LIU KH and TSAY YF (2003) Switching between the two action modes of the dual-affinity nitrate transporter CHL1 by phosphorylation. *EMBO J*, 22 (5): 1005–1013. DOI: 10.1093/emboj/cdg118
- LIVAK KJ and SCHMITTGEN TD (2001) Analysis of relative gene expression data using real-time quantitative PCR and $2^{-\Delta(\Delta C_q)}$ method. *Methods*, 25:402–8. DOI: 10.1006/meth.2001.1262
- LOQUÉ D, YUAN L, KOJIMA S, et al. (2006) Additive contribution of AMT1;1 and AMT1;3 to high-affinity ammonium uptake across the plasma membrane of nitrogen-deficient Arabidopsis roots. *Plant J*, 48: 522–534. DOI: 10.1111/j.1365-313X.2006.02887.x
- LUPINI A, MERCATI F, ARANITI F, et al. (2016). NAR2.1/NRT2.1 functional interaction with NO_3^- and H^+ fluxes in high-affinity nitrate transport in maize root regions. *Plant Physiol Biochem*, 102: 107–114. DOI: 10.1016/j.plaphy.2016.02.022
- LYONS CW and SCHOLTHOF KBG (2015). Watching grass grow: The emergence of *Brachypodium distachyon* as a model for the *Poaceae*. In *New Perspectives on the History of Life Sciences and Agriculture*, 40: 479–501. DOI: 10.1007/978-3-319
- LYONS CW and SCHOLTHOF KBG (2016) *Brachypodium* as an Arabidopsis for the grasses: Are we there yet? In *Genetics and Genomics of Brachypodium*, pp. 327–341. DOI: 10.1007/7397_2015_8
- MALAGOLI P, LAINE P, LE DEUNFF E, et al. (2004) Modeling nitrogen uptake in oilseed rape cv Capitol during a growth cycle using influx kinetics of root nitrate transport systems and field experimental data. *Plant Physiol*, 134: 388–400. DOI: 10.1104/pp.103.029538
- MARSCHNER H (2012) Marschner's Mineral Nutrition of Higher Plants. Academic Press, Vol. 89, 672 p. eBook ISBN: 9780080571874
- MARTINELLI LA and FILOSO S (2008) Expansion of sugarcane ethanol production in Brazil: Environmental and social challenges. *Ecol Appl*, 18: 885–898. DOI: 10.1890/07-1813.1

- MILLER AJ and CRAMER MD (2005) Root nitrogen acquisition and assimilation. *Plant Soil*, 274: 1–36. DOI: 10.1007/s11104-004-0965-1
- MILLER AJ, FAN X, ORSEL M, et al. (2007) Nitrate transport and signaling. *J Exp Bot*, 58: 2297–2306. DOI: 10.1093/jxb/erm066
- MOCHIDA K and SHINOZAKI K (2013) Unlocking Triticeae genomics to sustainably feed the future. *Plant Cell Physiol*, 54: 1931–1950. DOI: 10.1093/pcp/pct163
- MOLL RH, KAMPRATH EJ, and JACKSON WA (1982) Analysis and interpretation of factors which contribute to efficiency of nitrogen utilization. *Agron J*, 74: 562–564. DOI: 10.2134/agronj1982.00021962007400030037x
- MUELLER ND, GERBER JS, JOHNSTON M, et al. (2012) Closing yield gaps through nutrient and water management. *Nature*, 490: 254–257. DOI: 10.1038/nature11420
- MUÑOS S, CAZETTES C, FIZAMES C, et al. (2004) Transcript profiling in the chl1-5 mutant of *Arabidopsis* reveals a role of the nitrate transporter NRT1.1 in the regulation of another nitrate transporter, NRT2.1. *Plant Cell*, 16: 2433–2447. DOI: 10.1105/tpc.104.024380
- MUR LAJ, ALLAINGUILLAUME J, CATALÁN P, et al. (2011) Exploiting the Brachypodium Toolbox in cereal and grass research. *New Phytol*, 191: 334–347. DOI: 10.1111/j.1469-8137.2011.03748.x
- NAZOA P, VIDMAR JJ, TRANBARGER TJ, et al. (2003) Regulation of the nitrate transporter gene *AtNRT2.1* in *Arabidopsis thaliana*: responses to nitrate, amino acids and developmental stage. *Plant Mol Biol*, 52: 689–703. DOI: 10.1023/a:1024899808018
- OKAMOTO M, VIDMAR JJ, and GLASS ADM (2003) Regulation of NRT1 and NRT2 gene families of *Arabidopsis thaliana*: responses to nitrate provision. *Plant Cell Physiol*, 44: 304–317. DOI: 10.1093/pcp/pcg036
- OKAMOTO M, KUMAR A, LI W, WANG Y, et al. (2006) High-affinity nitrate transport in roots of *Arabidopsis* depends on expression of the NAR2-like gene *AtNRT3.1*. *Plant Physiol* 140: 1036–1046. DOI: 10.1104/pp.105.074385
- OMASITS U, AHRENS CH, MÜLLER S, and WOLLSCHIED B (2014) Protter: interactive protein feature visualization and integration with experimental proteomic data. *Bioinformatics*, 30: 884–888. DOI: 10.1093/bioinformatics/btt607
- ORSEL M, KRAPP A, and DANIEL-VEDELE F (2002) Analysis of the NRT2 nitrate transporter family in *Arabidopsis*. Structure and gene expression. *Plant Physiol*, 129: 886–896. DOI: 10.1104/pp.005280
- ORSEL M, CHOPIN F, LELEU O, et al. (2006) Characterization of a two-component high-affinity nitrate uptake system in *Arabidopsis*. Physiology and protein-protein interaction. *Plant Physiol* 142: 1304–1317. DOI: 10.1104/pp.106.085209
- PATHAK RR, LOCHAB S, and RAGHURAM N (2011) Improving nitrogen use efficiency. *Compr Biotechnol*, 4: 209–218. DOI: 10.1016/B978-0-08-088504-9.00472-4
- PLETT D, TOUBIA J, GARNETT T, et al. (2010) Dichotomy in the NRT gene families of dicots and grass species. *PLoS One* 5: 15289. DOI: 10.1371/journal.pone.0015289
- QUESADA A, KRAPP A, TRUEMAN LJ, et al. (1997) PCR-identification of a *Nicotiana plumbaginifolia* cDNA homologous to the high-affinity nitrate transporters of the crnA family. *Plant Mol Biol*, 34: 265–274. DOI: 10.1023/a:1005872816881
- RAHAYU YS, WALCH-LIU P, NEUMANN G, et al. (2005) Root-derived cytokinins as long-distance signals for NO₃⁻ induced stimulation of leaf growth. *J Exp Bot*, 56: 1143–1152. DOI: 10.1093/jxb/eri107
- RAVEN JA, WOLLENWEBER B, and HANDLEY L (1992) A comparison of ammonium and nitrate as nitrogen sources for photolithotrophs. *New Phytol*, 121: 19–32. DOI: 10.1111/j.1469-8137.1992.tb01088.x
- ROBERTSON GP and VITOUSEK PM (2009) Nitrogen in Agriculture: Balancing the Cost of an Essential Resource. *Annu Rev Environ Resour*, 34:97–125. DOI: 10.1146/annurev.enviro.032108.105046
- ROBINSON N, BRACKIN R, VINALL K, et al. (2011) Nitrate Paradigm Does Not Hold Up for Sugarcane. *PLoS One*, 6: e19045. DOI: 10.1371/journal.pone.0019045
- RUFFEL S, FREIXES S, BALZERGUE S, et al. (2008) Systemic signaling of the plant nitrogen status triggers specific transcriptome responses depending on the nitrogen source in *Medicago truncatula*. *Plant Physiol*, 146: 2020–2035. DOI: 10.1104/pp.107.115667
- SALSAC L, CHAILLOU S, MOROT-GAUDRY JF, et al. (1987) Nitrate and ammonium nutrition in plants. *Plant Physiol Biochem*, 25: 805–812.
- SANTOS TB, LIMA JE, FELICIO MS, et al. (2017) Genome-wide identification, classification and transcriptional analysis of nitrate and ammonium transporters in *Coffea*. *Genet Mol Biol*, 40: 346–359. DOI: 10.1590/1678-4685-gmb-2016-0041.

- SCHOLTHOF KBG, IRIGOYEN S, CATALAN P, and MANDADI KK (2018) Brachypodium: A Monocot Grass Model Genus for Plant Biology. *Plant Cell*, 30: 1673-1694. DOI: 10.1105/tpc.18.00083
- SEREZINO LHD (2015) Physiological and transcriptional characterization of nitrate acquisition and remobilization processes in sugarcane (*Saccharum spp.*) (Master's dissertation, CENA/USP, São Paulo, Brazil)
- SIBOUT R, PROOST S, HANSEN BO, et al. (2017) Expression atlas and comparative coexpression network analyses reveal important genes involved in the formation of lignified cell wall in *Brachypodium distachyon*. *New Phytol*, 215: 1009-1025. DOI: 10.1111/nph.14635
- SIDDIQI MY, GLASS ADM, RUTH TJ, and RUFTY TW (1990) Studies of the uptake of nitrate in barley. I. Kinetics of $^{13}\text{NO}_3^-$ influx. *Plant Physiol*, 93: 1426-1432. DOI: 10.1104/pp.93.4.1426
- SUBBARAO GV, NAKAHARA K, HURTADO MP, et al. (2009) Evidence for biological nitrification inhibition in Brachiaria pastures. *Proc Natl Acad Sci USA*, 106: 17302-17307. DOI: 10.1073/pnas.0903694106
- SUN J, LUU NS, CHEN Z, et al. (2019) Generation and Characterization of a Foxtail Millet (*Setaria italica*) Mutant Library. *Front Plant Sci*, 10: 369. DOI: 10.3389/fpls.2019.00369
- SUTTON MA, BLEEKER A, HOWARD CM, et al. (2013) Our nutrient world: the challenge to produce more food and energy with less pollution. *Global Overview of Nutrient Management*, Centre for Ecology and Hydrology, on behalf of the Global Partnership on Nutrient Management and the International Nitrogen Initiative
- SWANEY DP, HONG B, TI C, et al. (2012) Net anthropogenic nitrogen inputs to watersheds and riverine N export to coastal waters: a brief overview. *Curr Opin Env Sust*, 4:203-211. DOI: 10.1016/j.cosust.2012.03.004
- TANG Z, FANX, LI Q, FENG H, et al. (2012) Knockdown of a rice stelar nitrate transporter alters long-distance translocation but not root influx. *Plant Physiol*, 160: 2052-2063. DOI: 10.1104/pp.112.204461
- TISCHNER R, WALDECK B, GOYAL SS, and RAINS WD (1993). Effect of nitrate pulses on the nitrate-uptake rate, synthesis of mRNA coding for nitrate reductase, and nitrate- reductase activity in the roots of barley seedlings. *Planta*, 189: 533-537. DOI: 10.1007/BF00198216
- TOMPKINS GA, JACKSON WA, and VOLK RJ (1978). Accelerated nitrate uptake in wheat seedlings: effects of ammonium and nitrite treatments and of six methyl purine and puromycin. *Physiol Plant*, 43: 166-171. DOI: 10.1111/j.1399-3054.1978.tb01586.x
- TONG Y, ZHOU JJ, LI Z, and MILLER AJ (2005) A two-component high-affinity nitrate uptake system in barley. *Plant J*, 41: 442-450. DOI: 10.1111/j.1365-313X.2004.02310.x
- TRUEMAN LJ, RICHARDSON A, and FORDE BG (1996) Molecular cloning of higher plant homologues of the high-affinity nitrate transporters of *Chlamydomonas reinhardtii* and *Aspergillus nidulans*. *Gene*, 175: 223-231. DOI: 10.1016/0378-1119(96)00154-0
- TSAY YF, CHIU CC, TSAI CB, HO CH, and HSU PK (2007) Nitrate transporters and peptide transporters. *FEBS Lett*, 581: 2290-2300. DOI: 10.1016/j.febslet.2007.04.047
- VIDMAR JJ, ZHUO D, SIDDIQI MY, et al. (2000) Regulation of high-affinity nitrate transporter genes and high-affinity nitrate influx by nitrogen pools in roots of barley. *Plant Physiol*, 123: 307-318. DOI: 10.1104/pp.123.1.307
- VITOUSEK PM, NAYLOR R, CREWS T, et al. (2009) Nutrient Imbalances in Agricultural Development. *Science*, 324:1519-1520. DOI: 10.1126/science.1170261
- VOGEL JP (2016). The rise of Brachypodium as a model system. In *Genetics and Genomics of Brachypodium*, p. 1-7. DOI: 10.1007/978-3-319-26944-3
- VON WIRÉN N, GAZZARRINI S, GOJON A, and FROMMER WB (2000). The molecular physiology of ammonium uptake and retrieval. *Curr Opin Plant Biol*, 3:254-261. DOI: 10.1016/S1369-5266(00)80074-6
- WANG J, NORMAN H, and LINING T (2019) Identification and molecular characterization of the *Brachypodium distachyon* NRT2 family, with a major role of BdNRT2.1. *Physiol Plant*, 165: 498-510. DOI: 10.1111/pp.12716
- WANG MY, SIDDIQI MY, RUTH TJ, and GLASS ADM (1993a): Ammonium uptake by rice roots. (I. Fluxes and subcellular distribution of $^{13}\text{NH}_4^+$). *Plant Physiol*, 103: 1249-1258. DOI: 10.1104/pp.103.4.1249
- WANG MY, SIDDIQI MY, RUTH TJ, and GLASS ADM (1993b): Ammonium uptake by rice roots. (II. Kinetics of $^{13}\text{NH}_4^+$ across the plasmalemma). *Plant Physiol*, 103: 1259-1267. DOI: 10.1104/pp.103.4.1259
- WOOD AW, SCHROEDER BL, and DWYER R (2010) Opportunities for improving the efficiency of use of nitrogen fertilizer in the Australian sugar industry. *Proceedings of the Australian Society of Sugar Cane Technologists*, 32: 221-233.

- XU G, FAN X, and MILLER AJ (2012) Plant nitrogen assimilation and use efficiency. *Annu Rev Plant Biol*, 63:153-182. DOI: 10.1146/annurev-arplant-042811-105532
- YANG JC, ZHANG H, and ZHANG JH (2012) Root morphology and physiology in relation to the yield formation of rice. *J Integr Agri*, 11(6): 920–926. DOI: 10.1016/S2095-3119(12)60082-3
- YAN M, FAN X, FENG H, et al. (2011) Rice OsNAR2.1 interacts with OsNRT2.1, OsNRT2.2 and OsNRT2.3a nitrate transporters to provide uptake over high and low concentration ranges. *Plant Cell Environ*, 34: 1360-1372. DOI: 10.1111/j.1365-3040.2011.02335.x
- YONG Z, KOTUR Z, and GLASS AMD (2010) Characterization of an intact two-component high- affinity nitrate transporter from *Arabidopsis* roots. *Plant J* 63: 739-748. DOI: 10.1111/j.1365-313X.2010.04278.x
- YORK LM, SILBERBUSH M, and LYNCH JP (2016) Spatiotemporal variation of nitrate uptake kinetics within the maize (*Zea mays* L.) root system is associated with greater nitrate uptake and interactions with architectural phenes. *J Exp Bot*, 67(12): 3763–3775. DOI: 10.1093/jxb/erw133
- ZHANG X, DAVIDSON EA, MAUZERALL DL, et al. (2015) Managing nitrogen for sustainable development. *Nature*, 528: 51-59. DOI: 10.1038/nature15743
- ZHUO D, OKAMOTO M, VIDMAR JJ, and GLASS ADM (1999) Regulation of a putative high-affinity nitrate transporter (Nrt2;1At) in roots of *Arabidopsis thaliana*. *Plant J*, 17: 563-568. DOI: 10.1046/j.1365-313X.1999.00396.x

3. Regulation of nitrate uptake in sugarcane (*Saccharum spp.*) occurs in post-transcriptional and post-translational levels among genotypes

3.1 Abstract

Sugarcane (*Saccharum spp.*) is a crop of great economic importance in Brazil, where it is grown mainly for the production of sugar and bioethanol, a replacement fossil fuels. The increasing demand has motivated the constant expansion of cultivated area to marginal soils, leading to an increased in the use of nitrogen fertilizers (N) to maintain productivity. However, to maintain the competitiveness, the energy balance in sugarcane production should be positive, and the high financial and energy costs for the production of nitrogen fertilizers, in addition to the consequent environmental impacts should be considered. Thus, there is a requirement to find new solutions to increase yield and the N use efficiency (NUE). Sugarcane response to nitrogen fertilizers is low compared to other crops, and losses of N applied to the soil can reach 50%. Compared with sorghum and maize, sugarcane has the lowest uptake efficiency and the lowest nitrate storage capacity tissues, partially explaining the large loss of N in the soil and the low NUE. However, the physiological and genetic causes of low efficiencies are still obscure. In this work, we identified a likely mechanism of post-transcriptional regulation of genes encoding nitrate transporters in roots subjected to resupply of nitrate, suggesting a mechanism responsible for the low nitrate use efficiency in sugarcane. Studies using sorghum and sugarcane revealed distinct regulation of HATS nitrate uptake process in plants of both species under nitrate resupply. ¹⁵N-nitrate influx showed high nitrate uptake by sorghum 'BTX623' and sugarcane hybrid 'IAC87-3396'. Expression analysis of genes coding NRT2 and NRT3, coding HATS complex components, exhibit different transcriptional regulation among sugarcane genotypes. The lack of correlation between the accumulation of NRT2.1 and NRT3.1 proteins with the ¹⁵N-nitrate influx suggest the presence of a mechanism controlling nitrate transport in sorghum and sugarcane roots in post-translational level. This putative allosteric regulation of HATS complex would be the major actor on nitrate uptake in roots of sorghum and sugarcane.

Keywords: NUE, Sorghum, Sugarcane, HATS Regulation, Putative Post-translational Regulation

3.2 Introduction

The application of inorganic Nitrogen (N) fertilizer has maintained high yields in agriculture, allowing food production to be ahead of the growing human population (Vitousek et al. 2009). Large energy inputs are required for the synthesis of N fertilizers, which, together with agricultural biological fixation, incorporates over 120 million tons of N year⁻¹ from the atmosphere into the plant-soil system in a reactive form (Tilman et al. 2001, Galloway et al. 2008, Rockström et al. 2009). The excessive application of N fertilizer, which increased more than 8-fold in the last 50 years, have led to environment eutrophication by nitrate leaching and run-off (Tilman 1999, Jaynes et al. 2001, Galloway et al. 2004, Schröder et al. 2004, Burt et al. 2008, Vitousek et al. 2009, Foley et al. 2011), together with the emission of the potent greenhouse gases nitrous oxide and ammonia (Tilman et al. 2001, Galloway et al. 2008, Davidson 2009, Burney et al. 2010, Millar et al. 2010, Hoben et al. 2011, Reay et al. 2012). Therefore, reduction of N input is urgently required to decrease environmental impacts derived from agricultural systems (Mueller et al. 2012).

Sugarcane (*Saccharum* spp.) is a perennial C₄ grass with a remarkable ability to produce sucrose-storing culms, suitable for sugar and bioethanol production, turning this species in one of the most efficient crop for bioenergy feedstock (Somerville et al. 2010, Waclawovsky et al. 2010). High sugarcane crop yields are maintained by the supply of considerable amounts of N fertilizers (Thorburn et al. 2010; Franco et al. 2011). However, the high N requirement to maintain yields increases the carbon footprint and affects the sustainability and competitiveness of bioenergy production derived from sugarcane (Miller 2010, van der Wiejde et al. 2013, Khoo 2015). The effective N fertilizer recovery by sugarcane has been estimated to be between 20 to 40% (Meyer et al. 2007, Kingston et al. 2008, Franco et al. 2011, Vieira-Megda et al. 2015). This low N recovery has been associated with a low nitrate uptake ability of sugarcane roots, which differs from the major C₄ grain crops, such as sorghum and maize (Robinson et al. 2011). Sugarcane favors ammonium uptake under N-sufficient condition and exhibit a reduced capacity to acquire nitrate by roots (Robinson et al. 2011). Nitrate is the predominant inorganic N form available in most agricultural soils (Miller et al. 2007). Consequently, sugarcane commercial cultivars have a limited ability to benefit from the nitrate abundance in soils after fertilization at the beginning of the crop cycle, which contributes to nitrate losses to the environment (Robinson et al. 2011).

Plant nitrate uptake and transport are constantly regulated according to the soil inorganic N availability and plant N status to fulfill the growth and development demand (Glass 2003). In higher plants, the nitrate uptake process is mediated by proteins from two gene families, namely *NITRATE TRANSPORTER 1/PEPTIDE TRANSPORTER FAMILY* (*NPF/NRT1*) and *NITRATE TRANSPORTERS* (*NRT2*) (Tsay et al. 2007, Krapp et al. 2014).

Those transporters ensure the ability of root cells to uptake through high-affinity transport systems (HATS) and low-affinity transport systems (LATS), depending on nitrate concentrations in the soil solution. Homologues of *NPF* have been identified in the genome of important crop species (Ruffel et al. 2008, Plett et al. 2010, Garnet et al. 2013, Lerán et al. 2014, von Wittgenstein et al. 2014, Wang et al. 2019a). Family members have been functionally characterized in *Arabidopsis* for root and shoot nitrate transport (Chiu et al. 2004, Segonzac et al. 2007, Lin et al. 2008, Fan et al. 2009, Li et al. 2010, Wang and Tsay 2011, Hsu and Tsay 2013).

In roots, nitrate uptake at levels requiring HATS is accomplished by the activity of members from a small gene family *NRT2* (Cerezo et al. 2001, Filleur et al. 2001). The investigation of spatial-temporal expression indicates that mostly of the *NRT2* genes present in the *A. thaliana* genome are expressed in roots to support growth of N-limited plants by ensuring an efficient uptake of nitrate (Lezhneva et al. 2014). Growth analyses of multiple *Arabidopsis* mutants revealed that *AtNRT2.1* and *AtNRT2.2* are responsible for up to 67% of HATS nitrate uptake (Filleur et al. 2001, Kiba et al. 2012, Lezhneva et al. 2014). In addition, the rapid co-induction of *AtNRT2.5* in the root epidermis also contributes for nitrate uptake during prolonged N starvation (Lezhneva et al. 2014, Kotur and Glass 2014). In *Brachypodium distachyon*, the high affinity transport system was reduced by 30% in the *bdnrt2.1* mutant, highlighting the major role of *BdNRT2.1* in this grass species (Wang et al. 2019b).

The transport function of most *NRT2* proteins in higher plants depends on the direct interaction with the *NAR2* protein (*NRT3*) (Orsel et al. 2006, Ishikawa et al. 2009, Feng et al. 2011, Kotur et al. 2012). Disruption of *NAR2.1* (*NRT3.1*) in the *atnar2.1-1* mutant caused an almost complete loss of inducible high-affinity nitrate influx, indicating that *AtNRT2.1* activity in the plasma membrane occurs in association with *AtNAR2.1* (*AtNRT3.1*) (Yong et al. 2010). The identification of *NRT2* and *NRT3* subfamily members in other plant species, such as *Hordeum vulgare*, *Oryza sativa*, *Zea mays*, *Sorghum bicolor*, *B. distachyon*, *Medicago truncatula*, *Lotus japonicus* and *S. spontaneum* suggests a crucial function of these gene families in coordinating high affinity nitrate acquisition in plant roots (Vidmar et al. 2000, Ruffel et al. 2008, Plett et al. 2010, Feng et al. 2011, Kotur et al. 2012, Pellizzaro et al. 2015, Wang et al. 2019a, Wang et al. 2019b).

The relative contributions of HATS and LATS to the overall nitrate uptake in plant roots is dynamically affected according to heterogeneous soil N availability and whole plant demand (Malagoli et al. 2004). Local and systemic N-signaling pathways coordinate uptake responses. A local response is observed in plant roots exposed to nitrate or by resupply after a prolonged N-deprivation period, in which nitrate itself acts as local signal for induction, within minutes, of the expression of several genes involved in nitrate transport and assimilation (Wang et al. 2004). In general, this response is transient, since the nitrate provision for several days leads

to a local inhibitory effect on nitrate uptake (Muños et al. 2004, Krouk et al. 2006). Conversely, systemic signals can communicate the N status of the whole plant to repress root nitrate uptake (Crawford and Glass 1998, Gansel et al. 2001, Alvarez et al. 2012). The main components targeted by local and systemic signals are the expression of *NRT2.1*, which is induced by local nitrate supply (Li et al. 2006) or repressed by systemic feedback signals exerted by high N status (Cerezo et al. 2001, Gansel et al. 2001), and *NRT2.1* protein stabilization during nitrate-limited conditions (Zou et al. 2019).

Over the past few years, key regulatory components of N-signaling pathways controlling root nitrate uptake were identified in model plants (Gutiérrez et al. 2008, Lejay et al. 2008, Krouk et al. 2010, Kiba et al. 2011, Zhao et al. 2011, Marchive et al. 2013, Konishi and Yanagisawa 2013, Tabata et al. 2014). Despite that, fundamental research of transport mechanisms in crop plants remains scarce (Gutiérrez 2012, Garnett et al. 2013). Therefore, a more comprehensive understanding of nitrate uptake ability and its regulation in the context of a changing environment will be important for improving sugarcane crop yields towards the development of plants with enhanced N uptake and use efficiency (Gutiérrez 2012, Xu et al. 2012).

In an attempt to understand the physiological and molecular features of the low nitrate uptake of commercial sugarcane varieties (Robinson et al. 2011), Serezino (2015) revealed a presumed post-transcriptional regulation of HATS nitrate uptake in *in vitro* sugarcane plantlets from some genotypes. This regulation in relation to the physiological N status of the plants is likely to contribute to the lower nitrate use efficiency in sugarcane. Thus, to confirm the presumed post-transcriptional regulation of HATS, nitrate resupply experiments with the same cultivars were conducted upon greenhouse conditions, adding one parental genotype of *Saccharum officinarum* and a sorghum genotype. ¹⁵N influx, gene expression and western blot analysis were carried out to answer this question.

3.3 Methods

3.3.1 Plant material

The *S. officinarum* ('NG57213') and the sugarcane commercial hybrids 'SP80-3280', 'IAC87-3396' and 'IACSP96-2042' were used as indicated in the experiments. Micropropagated plantlets were obtained for the four genotypes starting from *in vitro* culture of shoot apices (Lee 1987), to standardize root and plant size among replicates. Sorghum 'BTX-623' seeds obtained from the "EMBRAPA Milho e Sorgo" were germinated in sterilized filtered paper in gerbox. The gerboxes were incubated at 43 °C under high relative humidity for 120 h. The plantlets or seedlings were hardened in 5 L containers, containing half strength

nutrient solution (Hoagland and Arnon 1950) for three weeks in a greenhouse at $28\text{ }^{\circ}\text{C} \pm 4\text{ }^{\circ}\text{C}$. Individual plants were then transferred to 25 L containers (2 plants per container) containing a vigorously aerated full strength nutrient solution at pH 5.8, connected to 300 L tanks containing nutrient solution, air-pumped for 15 min per hour. For N-free solutions, $(\text{NH}_4)_2\text{SO}_4$ and KNO_3 were replaced by equimolar concentrations of K_2SO_4 . The pH was adjusted to 5.8 daily, and the nutrient solutions were renewed every week. Plants were grown under these experimental conditions in a greenhouse for 3 months.

3.3.2 Analysis of nitrate uptake kinetics in roots by rapid ^{15}N influx

For the *in vivo* N uptake kinetic analyses, we used the 'Short-Term Uptake' assay using ^{15}N (Loqué et al. 2006). The sugarcane plantlet and sorghum seedling roots, grown in nutrient solution were used to estimate the rapid influx. The roots were first rinsed in 1 mM CaSO_4 for 1 min, and subsequently exposed to nutrient solution (1.3 L) supplemented with 1 mM of 2-(N-morpholino) ethanesulfonic acid monohydrated (MES) at pH 5.8 containing $^{15}\text{N}\text{-NO}_3^-$ (60% atoms % ^{15}N) as sole N source for 10 min, followed by another 1 min wash in 1 mM CaSO_4 . To quantify the ^{15}N -nitrate influx, roots were collected, freeze dried for one week and weighted for total ^{15}N content using a continuous-flow isotope ratio mass spectrometry (ANCA SL, Sercon, Chesshire, UK) in the Stable Isotopes Laboratory, CENA, by Prof. Dr. José Albertino Bendassolli.

3.3.3 Understanding the nitrate uptake process in roots upon distinct inorganic N source supply and N resupply

The experiments were conducted to characterize the regulation of the nitrate root transport according to the presence of nitrate as the source of N. Plants grown for three months in a N sufficient nutrient solution (N+; 2 mM NH_4NO_3), were subjected to restriction N (-N) for three days. After this period, the plants were exposed to nitrate (4 mM KNO_3) and analyzed for the ^{15}N -nitrate influx (0.2 mM K^{15}NO_3) at 0, 4, 24 and 72 h after nitrate resupply. Four to five biological replicates were sampled at each time point. Samples of sugarcane and sorghum roots were collected, frozen and used for total RNA extraction to quantify expression of the genes responsible for nitrate acquisition process (NRT2 and NRT3 subfamilies) in both sugarcane and sorghum species by relative quantification of reversed transcripts (RT-qPCR). Besides, the frozen root samples were used in NRT2.1 and NRT3.1 immunodetection (ELISA and Western blot analysis) assays to investigate the levels of NRT2.1 and NRT3.1 proteins in roots of both species under nitrate resupply treatment.

3.3.4 RNA extraction and cDNA synthesis

Root samples were collected and immediately frozen in liquid nitrogen for subsequent RNA extraction. Samples (100 mg) were extracted using the method described by Leal et al

(2007). Briefly, samples were ground in liquid N and transferred to a 2 mL microtube, and 1 mL of extraction buffer (2% CTAB; 2% PVP; 2 M NaCl; 100 mM Tris; 25 mM EDTA pH 8.0; 2% β -mercaptoethanol) heated to 55 °C was added. After homogenization, samples were incubated at 55 °C for about 60 min. Afterwards, the samples were placed immediately on ice for 1 min, followed by addition of 1 mL of CIA (chloroform isoamyl-alcohol; 24:1 v/v) and spun at 9000 *g* for 10 min at 4 °C. Then, the supernatant was transferred to a new 1.5 mL microtube, adding an equal volume of CIA, followed by centrifugation at 9000 *g* for 10 min at 4 °C. Once again, the supernatant was transferred to a new 1.5 mL microtube for precipitation of the total RNA using a 10 M solution of LiCl and incubation at -80 °C for 12 h. Subsequently, the samples were spun at 9000 *g* for 30 min at 4 °C, followed by discarding the supernatant. The pellet was washed twice with 500 μ L of cold 75% ethanol and centrifuged at 9000 *g* for 10 min at 4 °C. Finally, the pellet was resuspended in 30 μ L of 0.01% inactive DEPC water. The total RNA was quantified in a NanoDrop One (Thermo Scientific; Wilmington, DE, USA) and the integrity was confirmed by gel electrophoresis. About 2 μ g of RNA from each sample was treated with 1 U of DNase I in appropriate buffer, 2 U of RNaseOUT (Invitrogen) and DEPC ultrapure water in a reaction incubated at 37 °C for 30 min. The reaction was stopped by adding of EDTA (1 mM), and then incubated at 65 °C for 10 min. Samples of 1 μ g of total RNA and oligo dT (0.5 μ g μ L⁻¹) were denatured at 70 °C for 10 min, and then incubated at 4 °C for 2 min, followed by the addition of 4 μ L 5x buffer-MuLV RT, 1 mM dNTP, 40 U RNaseOUT, and 20 U of the enzyme SuperScript III (Invitrogen; Carlsbad, CA, USA). The reaction was incubated at 55 °C for 1 h, then at 70 °C for 10 min and at 4 °C for 2 min. Finally, the reaction was cooled to 4 °C and stored at -80°C.

3.3.5 Gene expression by quantitative reversed transcript amplification (RT-qPCR)

After reverse transcription, the cDNA was diluted [3:7 (v/v) dilution], and used in quantitative PCR reaction containing 9 μ L of Kapa Sybr Fast (Kapa Biosystems, Wilmington, MA, USA) and 1 μ L of cDNA, with the reactions performed in a RotorGene Q 6000 thermocycler (Qiagen; Hilden, Germany), programmed with an initial incubation at 50°C for 10 min and 95°C for 2 min, and followed by 40 cycles of 95°C for 20 s and 60° or 62°C for 25 s, and 72°C for 25 s. After cycling, the melting curve for each amplicon was determined between 72°C and 95°C. Transcript-specific primers were used (Table 1). Amplifications experiments included for each RT-qPCR run was performed with three independent tissue samples, each sample having two technical replicates, and additional negative control (no template DNA). Data acquisition was performed in a Rotor-Gene Q Real-Time Analysis 6.0 (Qiagen). Amplification efficiency of each primer-pair was determined by standard curve from three serial dilutions of cDNA (10^{-1} , 10^{-2} , 10^{-3}). *ScUBQ2* and *SbUBQ* were used as the reference

gene (Table 1). C_q (quantification cycle) was used to determine differences in expression according to the $2^{-\Delta(\Delta C_q)}$ method (Livak and Schmittgen 2001).

Table 1: Primer sequences related to genes encoding transporters proteins (*NRTs* - *NITRATE TRANSPORTERS*), responsible for the nitrate uptake in sugarcane and sorghum roots, and the reference genes used for RT-qPCR.

Species	Gene name	Primers
<i>Sugarcane</i> spp.	<i>ScNRT2.1</i>	F: 5'- GAGCATCCGTCTCTTCTCCTTC -3' R: 5'- GCCTTGGTGAGGTTGAGGTT -3'
	<i>ScNRT2.5</i>	F: 5'- ATCGTGCCGTTTCGTCTCC -3' R: 5'- ATCCCCGTCTCCGTCTTGT -3'
	<i>ScNRT3.1</i>	F: 5'- GCTGGCGGTGCTTCTCTT -3' R: 5'- GGCTCGGTGGTGTTTCAGG -3'
	<i>ScUBQ2</i>	F: 5'- CTTCTTCTGTCCCTCCGATG -3' R: 5'- TCCAACCAAAGTCTGCTC -3'
<i>Sorghum bicolor</i>	<i>SbNRT2.1</i>	F: 5'- GTCATCCCCTTCGTCTCCCG -3' R: 5'- ATGCCCATGTACTCCAGCCC -3'
	<i>SbNRT2.2</i>	F: 5'- ATCCATGTTCTTCCCGGCCA -3' R: 5'- AACTTGAGGCTGGCGGCAT -3'
	<i>SbNRT2.5</i>	F: 5'- ATCCTGTTTCAGCCTGTTTCGT -3' R: 5'- TGATCTACTTCCCCCAGTGG -3'
	<i>SbNRT3.1</i>	F: 5'- GGAGCCTGAAGAGCACGGAG -3' R: 5'- CGCCTTGTCCTTGCTGAGGT -3'
	<i>SbNRT3.2</i>	F: 5'- ACGACCTGAGCAAGGACAAG -3' R: 5'- GCGACGCGGAAGTAGTAGG -3'
	<i>SbUBQ</i>	F: 5'- CAAGGAGTGCCCCAACAC -3' R: 5'- GGTAGGCGGGTAAAGCAAA -3'

3.3.6 Protein extraction

For the total protein extraction, it was used the protocol described by Duarte et al. (2002). Five g of frozen and ground roots samples from sorghum and sugarcane genotypes from the experiment described above were reground in PBS buffer (137mM NaCl, 2.7mM KCl, 10mM Na₂PO₄, 1.8mM KH₂PO₄), and placed into cooled 50 mL centrifuge tubes. Twenty mL of PBS buffer (137mM NaCl, 2.7mM KCl, 10mM Na₂PO₄, 1.8mM KH₂PO₄) were added and mixed with a homogenizer (UNI-Vortex, Uniscience, Osasco, São Paulo, Brazil) The extracts were spun for 15 min at 25,000 *g* at 4°C in to remove the insoluble material. The supernatant was collected in fresh tube and placed on ice, and the pellet was discarded. All the procedures were conducted on ice to inhibit sample degradation.

Total protein extracts were quantified using Bradford. Samples of 20 μ L of the extracts were placed into 96-well plate. A solution of Bovine Serum Albumin at 1 mg mL⁻¹ was used to prepare the standard curve. Then 200 μ L of the 5X Bradford stock solution (Bio-Rad) was added to each well. After 5 min, the samples were read at 595 nm of absorbance in an absorbance plate reader Accuris SmartReader 96 (model MR9600, Accuris, Edison USA).

3.3.7 NRT2.1 and NRT3.1 Immunodetection

For the NRT2.1 and NRT3.1 proteins assays, polyclonal antibodies for both proteins were produced and purified by Abbomax (California, USA). The polyclonal antibody was developed for NRT2.1 in sorghum and sugarcane root samples against the synthetic peptide RSERGKRNVIQADAGATPEHV, located in the C-terminal region (Figure 1, in orange). The anti-NRT3.1 antisera was developed against the synthetic peptide LAFFFVIENRKKNK, also located in the C-terminal region (Figure 2, in yellow).

Figure 1: Membrane topology of NRT2.1 transporters in sugarcane and sorghum using Protter, with the epitope corresponding to NRT2.1 antibody (orange). Peptides sequences were visualized by Protter.

Figure 2: Membrane topology of NRT3.1 transporters in sugarcane and sorghum using Protter, with the epitope corresponding to NRT3.1 antibody (yellow). Peptides sequences were visualized by Protter.

3.3.7.1 ELISA tests

ELISA assays were performed according to Crowther (2009), with some adaptations. Samples of 100 μ L of total protein (100 μ g/mL) from each time-point and genotype were distributed into micro wells, in triplicates, and incubated overnight at 4 $^{\circ}$ C. Blocking were done using 1% BSA Phosphate-Buffered Saline (137mM NaCl, 2.7mM KCl, 10mM Na₂PO₄, 1.8mM KH₂PO₄) for one hour at 37 $^{\circ}$ C. Plates were dried and the anti-sera dilution (1:500) were applied, and incubated for another hour at 37 $^{\circ}$ C. Two antibodies for each protein were used: anti-NRT2.1 for sorghum and sugarcane; anti-NRT3.1 against five genotypes. Plates were washed three times with PBS-T-G buffer (PBS, Tween 20 0.05% and 0.25 % Gelatin) and then anti-rabbit conjugated AP (Alkaline Phosphatase) serum was added. After one hour at 37 $^{\circ}$ C, plates were washed three times (PBS-T-G) and revealed using pNPP (p-Nitrophenylphosphate) at 450 nm of absorbance in an absorbance plate reader Accuris SmartReader 96 (model MR9600, Accuris, Edison USA, Farias et al. 2012).

3.3.7.2 Western blot analysis

Samples of total proteins (60 µg) extracted from sorghum and sugarcane genotypes under various nitrate treatments were electrophoretically separated in a 15-20% SDS-polyacrylamide denaturing gel and transferred to Hybond-ECL nitrocellulose membrane (GE Healthcare) a 100 mA for 75 min. Membrane blocking was performed with PBST (PBS and 0.05% Tween 20) buffer, containing 5% BSA for 90 min. The incubation with the primary anti-NRT2.1 or anti-NRT3.1 antibodies (1: 500, Abmax) were performed overnight at 4 °C. After washing with PBST, the membranes were incubated for 60 min with biotinylated secondary antibodies at RT. The development was carried out with 3,30-diaminobenzidine (DAB; Sigma-Aldrich), chloronaftol, and 30% hydrogen peroxide for 1 min at RT. The images were obtained using an Epson Perfection 4990 scanner.

3.4 Results

3.4.1 Comparing the uptake response of sorghum and 'SP80-3280' to nitrate resupply

Sugarcane exhibit a reduced capacity to acquire nitrate and to store nitrate in culms, differing from the major C₄ grain crops, such as sorghum (Robinson et al. 2011). Physiology studies focusing in nitrate uptake in model plant species demonstrated that after the addition of nitrate, there is a rapid induction of nitrate high-affinity transport, called iHATS (inducible HATS), which remains induced for some hours, followed by a rapid decrease in the nitrate influx (Tompkins et al. 1978, Siddiqi et al. 1990, Tischner et al. 1993, Zhuo et al. 1999, Glass 2003). This HATS induction by the addition of nitrate (re-supply) occurs when plants are cultivated in sufficient N and in deficiency and/or low N (Glass et al. 2002). To investigate the effect of nitrate resupply on high affinity transport activity in the grasses, sorghum (cv. 'BTX623'), *S. officinarum* (cv. 'NG57213'), and sugarcane hybrids cultivars ('SP80-3280') plants grown upon sufficient N (+ N; 2 mM NH₄NO₃) were transferred to N-free nutrient solution for 72 h, and then transfer to nitrate as the sole inorganic N source. ¹⁵N-nitrate Short-Term Uptake analysis (Loqué et al. 2006) were conducted at 0, 4, 24 and 72 h after the nitrate resupply treatment, at HATS range (0.2 mM ¹⁵N-nitrate). In this topic, it will be present the results comparing sorghum and SP80-3280 genotypes plants under nitrate resupply treatment, to facilitate the comparison and the understanding of nitrate uptake process both species. This sugarcane hybrid was chosen for the comparison with sorghum due to its wide application on molecular investigations in this species, such as BAC library assembly (Figueira et al. 2012) and genome sequencing and assembly (Riaño-Pachón and Mattiello 2017, Souza et al. 2019). For the next topic, it will be shown and analyzed the results obtained in all

sugarcane genotypes used in this experiment (*S. officinarum* and hybrids), to understand the regulation of nitrate uptake among different genotypes this biomass crop.

Figure 3: $^{15}\text{NO}_3$ influx in sorghum and sugarcane roots submitted to nitrate resupply. Sorghum ('BTX-623') and sugarcane ('SP80-3280') plants roots were grown under adequate N supply (2 mM NH_4NO_3), followed by N (-N) deficiency for 72 h, when they were supplemented with nitrate (4 mM KNO_3) for 4, 24 and 72 hours. Uptake rates were measured for 10 min in sorghum and sugarcane roots at the external concentration of 200 μM ^{15}N -labeled NO_3^- (K^{15}NO_3). Values represent means of 4 biological replicates $\pm$ SD, and significant differences at $P < 0.05$ are indicated by different letters.

The temporal ^{15}N influx analysis showed that HATS transport is not regulated in sugarcane roots under nitrate resupply treatment, even with a light increase of ^{15}N -nitrate influx after 72 h (Figure 3). In sorghum, there was a rapid (4 h) and late (72 h) response to nitrate resupply. After 4 h of treatment, the nitrate influx observed in sorghum roots was $6.56 \mu\text{mol h}^{-1} \text{g}^{-1} \text{DW}$, approximately 3.4-fold higher than observed in plants under N absence (0 h) (Figure 3). There is a non-significant decrease in the following period (24 h), with HATS being strongly induced after 72 h nitrate resupply, with ^{15}N -nitrate influx of $17.30 \mu\text{mol h}^{-1} \text{g}^{-1} \text{DW}$, 8.95-fold higher than N-deficient plants (Figure 3). Thus, ^{15}N -influx analysis showed a low nitrate uptake capacity by sugarcane roots compared to sorghum, corroborating the results presented by Robinson et al. (2011) and Serezino (2015), indicating the low NUE of nitrate transport in sugarcane.

Among the seven *NRT2* genes, *NRT2.1* appears to be the most critical for high-affinity nitrate uptake in *Arabidopsis* roots. Knock-out mutants for *NRT2.1* (*atnrt2.1*) show strongly reduced HATS activity under various experimental conditions (Cerezo et al. 2001, Filleur et

al. 2001, Li et al. 2007). *NRT2.1* orthologs have been characterized in other species. In *Brachypodium distachyon*, the high affinity transport system was reduced by 30% in *bdnrt2.1*, highlighting the major role of BdNRT2.1 in this species (Wang et al. 2019b). Considering that HATS induction correlates directly with the transcription of high-affinity nitrate transporters in several species (Lejay et al. 1999, Filleur et al. 2001, Orsel et al. 2006, Li et al. 2007), the expression analysis of *NRT2* genes involved in nitrate uptake in sorghum and sugarcane were carried out, using root samples from plants grown in nitrate resupply conditions.

Sugarcane 'SP80-3280' plants showed an increase in the *NRT2.1* transcript accumulation in response to nitrate resupply treatment (Figure 4). However, this increase was not significantly distinct from that observed in plants under N deficiency (0 h) (Figure 4). In roots of 'BTX-623' sorghum plants, the *NRT2.1* expression regulation was different from that observed in sugarcane roots, exhibiting fast (4 h) and late (72 h) induction in response to nitrate exposure. However, this upregulation was not significantly different from that observed in N-deficient sorghum plants.

Figure 4: Regulation of the *NRT2.1* gene in sorghum and sugarcane roots submitted to nitrate resupply. The transcript levels were determined in roots of sorghum 'BTX-623' and sugarcane 'SP80-3280' genotypes cultivated for three months under normal conditions of N supplementation (+ N; 2 mM NH_4NO_3), followed by N (-N) deficiency for 72 hours. After this period, they were supplemented with nitrate (4 mM KNO_3) for 4, 24 and 72 hours. Expression was normalized for mean expression values for plants under sufficient restriction for 72 h (-N; time 0 h). *ScUBQ2* and *SbUBQ* were used as a reference for sugarcane and sorghum, respectively. Bars indicate $\pm$ SE, $n = 4$ plants per time-point. Significant differences at $P < 0.05$ are indicated by different letters.

In addition to *AtNRT2.1*, the gene encoding the *AtNRT2.2* transporter is upregulated in *Arabidopsis* resupplied with nitrate, displaying an additive role in the nitrate uptake process in *Arabidopsis* roots (Okamoto et al. 2003, Li et al. 2007). Recognizing that plants co-regulate the expression of the main genes encoding for HATS transporters as an adaptation strategy for the increase in the nitrate acquisition in roots, according to N demand, *NRT2.2* expression analysis was also conducted (Siddiqi et al. 1990, Wang et al. 1993a, Wang et al. 1993b, Forde and Clarkson 1999, Fraiser et al. 2000, Cerezo et al. 2001, Filleur et al. 2001, Krouk et al. 2006).

Figure 5: Regulation of the *SbNRT2.2* gene in sorghum roots submitted to nitrate resupply. The transcript levels were determined in roots of sorghum 'BTX-623' genotype cultivated for three months under normal conditions of N supplementation (+ N; 2 mM NH_4NO_3), followed by N (-N) deficiency for 72 hours. After this period, they were supplemented with nitrate (4 mM KNO_3) for 4, 24 and 72 hours. Expression was normalized for mean expression values for plants under sufficient restriction for 72 h (-N; time 0 h). *SbUBQ* was used as a reference. Bars indicate $\pm$ SE, $n = 4$ plants per time-point. Significant differences at $P < 0.05$ are indicated by different letters.

Increased expression of the *SbNRT2.2* gene was detected in roots of sorghum plants under low N concentrations, evidencing its essential role in the nitrate acquisition process in the high-affinity range (Gelli et al. 2014). To verify whether this gene is also important in response to nitrate resupply, *SbNRT2.2* expression was analyzed in plants exposed to this N inorganic source. The sorghum 'BTX623' showed a rapid (4 h) and late (72 h) upregulation of *SbNRT2.2* expression (Figure 5). Only after 4 h of nitrate resupply treatment, *SbNRT2.2* upregulation was significantly distinct with transcript accumulation 4-fold higher than in N-

deficient plants (Figure 5). *SbNRT2.2* expression pattern was similar to that observed for the *SbNRT2.1* (Figure 4), showing that both genes may act additively in the nitrate uptake process in sorghum roots under nitrate exposure. These results demonstrate that the regulatory mechanism of *NRT2.1* and *NRT2.2* gene expression might be conserved between dicots and monocots, considering the additive role of these transporters in *Arabidopsis* (Okamoto et al. 2003, Li et al. 2007). Despite the importance of the *NRT2.2* gene, it was not possible to identify orthologs in the BAC genomic library of sugarcane available in Dr. Marie Anne Van Sluys Laboratory, at the Institute of Biosciences at the University of São Paulo (IB / USP) (Dr. Joni Lima, personal communication).

The gene encoding *NRT2.5* is induced in *Arabidopsis* roots under N starvation, becoming the most abundant transcript amongst the seven *AtNRT2* family members after long-term starvation and affecting high-affinity nitrate uptake (Lezhneva et al. 2014). Thus, the expression of *NRT2.5* orthologues was evaluated in sugarcane and sorghum roots in response to nitrate resupply.

Figure 6: Regulation of the *NRT2.5* gene in sorghum and sugarcane roots submitted to nitrate resupply. The transcript levels were determined in roots of sorghum 'BTX-623' and sugarcane 'SP80-3280' genotypes cultivated for three months under normal conditions of N supplementation (+ N; 2 mM NH_4NO_3), followed by N (-N) deficiency for 72 hours. After this period, they were supplemented with nitrate (4 mM KNO_3) for 4, 24 and 72 hours. Expression was normalized for mean expression values for plants under sufficient restriction for 72 h (-N; time 0 h). *ScUBQ2* and *SbUBQ* were used as a reference for sugarcane and sorghum, respectively. Bars indicate $\pm$ SE, $n = 4$ plants per time-point. Significant differences at $P < 0.05$ are indicated by different letters.

In sorghum, the relative expression levels of *NRT2.5* remained stable until 24 h after nitrate resupply treatment. After 72 h it is possible to notice a decrease of transcript accumulation, evidencing a late *SbNRT2.5* downregulation in response to nitrate exposure (Figure 6). In contrast, a quick and strong upregulation of *ScNRT2.5* expression in ‘SP80-3280’ roots under nitrate resupply treatment was observed. After 4 h, the *NRT2.5* transcript accumulation was significantly 4.5-fold higher than in N-deficient sugarcane plants (0 h). The upregulation maintains high, even after 72 h of nitrate treatment (Figure 6). Therefore, *NRT2.5* transporter could show different roles among grasses, since its differential regulation in sorghum and sugarcane roots in response to nitrate resupply.

Most *NRT2* high-affinity transporters are active in the plasma membrane when *NRT3.1* transporters are coexpressed (Tong et al. 2005, Okamoto et al. 2006), forming the two-component high-affinity transport complex. Arabidopsis mutants for the accessory protein *NAR2/NRT3* showed a drastic decrease in the activity of HATS (80% decrease in nitrate influx) and in the expression of *NRT2* transporters, mainly *AtNRT2.1*, with a decrease in expression by 90% (Orsel et al. 2006). A similar physiological function has been identified for *NRT3* in rice, since the two-component high-affinity transport system also contributes to the absorption of nitrate in the low and high-affinity range (Yan et al. 2011). Therefore, the *NAR2/NRT2.1* protein complex formed in the plasma membrane of root cells is essential for the absorption of nitrate in the HATS range in several species (Okamoto et al. 2003, Tong et al. 2005, Okamoto et al. 2006, Yan et al. 2011). In Arabidopsis, except for *AtNRT2.7*, all *NRT2*s restored growth and β -galactosidase activity in the yeast split-ubiquitin system when these proteins were co-expressed with *AtNAR2.1*. Thus, except for *AtNRT2.7*, all other *NRT2* transporters interact strongly with *AtNAR2.1* (*AtNRT3.1*) (Kotur et al. 2012). Thus, the expression of the gene encoding the *NAR2.1* (*NRT3.1*) protein was evaluated in both grasses species to correlate with the expression of *NRT2.1* and *NRT2.5* genes. Both ‘SP80-3280’ and sorghum displayed upregulation of *NRT3.1* gene in plants roots after 4h of nitrate resupply treatment. In the following evaluated period (24 h), a differential regulation between both species was observed. In ‘BTX623’ roots, the *NRT3.1* mRNA accumulation returned to the same levels as observed in N-deficient plants. Sugarcane roots maintained high *NRT3.1* expression levels after 24 and 72 h of nitrate exposure (Figure 7).

Figure 7: Regulation of the *NRT3.1* gene in sorghum and sugarcane roots submitted to nitrate resupply. The transcript levels were determined in roots of sorghum 'BTX-623' and sugarcane 'SP80-3280' genotypes cultivated for three months under normal conditions of N supplementation (+ N; 2 mM NH_4NO_3), followed by N (-N) deficiency for 72 hours. After this period, they were supplemented with nitrate (4 mM KNO_3) for 4, 24 and 72 hours. Expression was normalized for mean expression values for plants under sufficient restriction for 72 h (-N; time 0 h). *ScUBQ2* and *SbUBQ* were used as a reference for sugarcane and sorghum, respectively. Bars indicate $\pm$ SE, $n = 4$ plants per time-point. Significant differences at $P < 0.05$ are indicated by different letters.

NAR2/NRT3 genes have been characterized in many species. However, not all *NRT3* family members demonstrates a function in two-component nitrate HATS (Tong et al. 2005, Feng et al. 2011 Yan et al. 2011). For example, only *HvNAR2.3* formed a functional unit with *HvNRT2.1* in barley (Tong et al. 2005). In rice *OsNAR2.1* (*OsNRT3.1*), but not *OsNAR2.2* (*OsNRT3.2*), functions as part of the two-component nitrate HATS in rice (Feng et al. 2011). *NAR2/NRT3* members were also identified in other grass species, as maize, sorghum and Brachypodium (Plett et al. 2010). To identify possible *SbNRT3.2* regulation in response to nitrate resupply treatment, gene expression was evaluated in sorghum roots. There was a small but not significant increase of *SbNRT3.2* mRNA accumulation after 4 h of nitrate resupply treatment comparing to N-deficient plants. In the following periods (24 and 72 h), a significant downregulation of *SbNRT3.2* was noticed comparing with the first period analyzed after nitrate exposure (4 h), but not significant comparing with 'BTX-623' plants under absence of N (0 h) (Figure 8). *ScNRT3.2* ortholog in sugarcane was not evaluated because it was mainly expressed in shoots (Dr. Joni Lima, personal communication).

Figure 8: Regulation of the *SbNRT3.2* gene in sorghum roots submitted to nitrate resupply. The transcript levels were determined in roots of sorghum 'BTX-623' genotype cultivated for three months under normal conditions of N supplementation (+ N; 2 mM NH_4NO_3), followed by N (-N) deficiency for 72 hours. After this period, they were supplemented with nitrate (4 mM KNO_3) for 4, 24 and 72 hours. Expression was normalized for mean expression values for plants under sufficient restriction for 72 h (-N; time 0 h). *SbUBQ* gene was used as a reference. Bars indicate $\pm$ SE, $n = 4$ plants per time-point. Significant differences at $P < 0.05$ are indicated by different letters.

Therefore, although not significant, the induction of *SbNRT2.1* and *SbNRT2.2* genes expression correlate with the regulation of the nitrate uptake process in roots of the sorghum 'BTX-623' under nitrate resupply treatment (Figures 3, 4, and 5). In contrast, there is a lack of correlation between the expression of *ScNRT2* and *ScNRT3* members with the ^{15}N -nitrate influx in sugarcane roots during nitrate resupply. *ScNRT2.1* and *ScNRT3.1* genes were co-regulated in response to nitrate resupply, since both showed increasing upregulation in their expression (Figures 4 and 7). However, no regulation of the nitrate uptake process was observed, considering the ^{15}N influx analyses that showed a constant level of nitrate acquisition in sugarcane roots under nitrate exposure (Figure 3). Equivalent behavior occurs for the *ScNRT2.5* gene (Figure 6), since its regulation does not correlate with the ^{15}N -nitrate influx in plants exposed to this treatment. This lack of correlation may indicate the existence of a post-transcriptional regulation of HATS transporter genes in sugarcane roots, or that the genotype used in this work, 'SP80-3280', is defective in the induction HATS system under nitrate resupply.

NRT2.1 regulation has been mostly studied at the transcriptional level. However, previous studies revealed the level of NRT2.1 protein is very stable compared to transcripts level and HATS activity, suggesting activation/inactivation of the NRT2.1/NRT3.1 transport system at the plasma membrane, whereas much faster changes in NRT2.1 mRNA or nitrate HATS activity have been demonstrated (Girin et al. 2007, Wirth et al. 2007, Laugier et al. 2012). The analysis of NRT2.1 and NRT3.1 protein stability in the plasma membrane in wild-type *Arabidopsis* roots indicated that short-term high N provision has no effect in the protein accumulation, which differs from long-term N supply (Wirth et al. 2007, Laugier et al. 2012). Before elucidating the existence of a post-transcriptional mechanism regulation in sugarcane roots, and if that response occurs in sugarcane in general or only in the cultivar 'SP80-3280', ^{15}N -nitrate influx and gene expression analysis were conducted using *S. officinarum* ('NG57213') and other sugarcane hybrids cultivars ('IACSP96-2042' and 'IAC87-3396') under nitrate resupply treatments.

3.4.2 Comparing sugarcane hybrids in response to nitrate resupply

While there are several definitions for NUE (Good et al. 2004), the most commonly applied reflects the amount of biomass (or grain yield) produced per unit of supplied N (Moll et al. 1982). Based on this definition, it is now well established that NUE in sugarcane is genotype-dependent and may be apparent as differences in internal N-use efficiency (Miller and Cramer 2005, Robinson et al. 2007). To understand the regulation of the nitrate uptake process after nitrate resupply among the sugarcane genotypes used in the experiment described above, ^{15}N influx and NRT2/NRT3 gene expression analysis for *S. officinarum* ('NG57213') and sugarcane hybrids cultivars ('SP80-3280', 'IACSP96-2042', and 'IAC87-3396') will be presented below.

The influx of ^{15}N -nitrate demonstrated that in N-deficient plants (0 h), 'NG57213' showed less ($0.974 \mu\text{mol h}^{-1} \text{g}^{-1} \text{DW}$), but not significant, nitrate uptake compared to the other sugarcane genotypes (Figure 9). After 4 h of nitrate resupply treatment, *S. officinarum*, 'SP80-3280', and 'IACSP96-2042' presented equal ^{15}N -nitrate influx level and comparable to the ones from the previous analyzed period (0 h), showing that there was no rapid response to nitrate resupply treatment in these genotypes. On the other hand, the sugarcane cultivar 'IAC87-3396' increased the nitrate uptake after 4 h ($9.142 \mu\text{mol h}^{-1} \text{g}^{-1} \text{DW}$) (Figure 9). This result shows that 'IAC87-3396' responds more quickly to nitrate resupply treatment than the other genotypes. Equivalent results were observed after 24 h of nitrate exposure, with the genotypes 'NG57213', 'SP80-3280', and 'IACSP96-2042' showing similar low rates of ^{15}N -nitrate influx and 'IAC87-3396' significantly higher ^{15}N -nitrate influx compared with other genotypes (Figure 9). 'IACSP96-2042' showed a nitrate uptake induction after 72 h of nitrate resupply treatment, reaching $4.712 \mu\text{mol h}^{-1} \text{g}^{-1} \text{DW}$ of ^{15}N -nitrate influx. Although the evident

increase of ^{15}N -nitrate at 72 h, the increase was not significantly different to the one by *S. officinarum* and 'SP80-3280'. A strong nitrate uptake induction was observed in 'IAC87-3396' since its ^{15}N -nitrate influx was $22.422 \mu\text{mol h}^{-1} \text{g}^{-1} \text{DW}$ after 72 h of nitrate treatment (Figure 9). Therefore, the temporal analysis of the ^{15}N -nitrate influx demonstrates the genetic variation in the nitrate uptake process between commercial hybrids and the ancestral sugarcane genotype. *S. officinarum* ('NG57213') and 'SP80-3280' showed no regulation of nitrate uptake in response to nitrate resupply considering that the ^{15}N -nitrate influx levels observed after 4, 24 and 72 h were not significantly different from the one observed in N-deficient plants (0 h). The same was observed for 'IACSP96-2042', even with an increase of ^{15}N -nitrate influx after 72 h. The genotype 'IAC87-3396' showed regulation of nitrate uptake process as the ^{15}N -nitrate influx rates after nitrate resupply treatment were significantly higher compared with plants under N deficiency (Figure 9).

Figure 9: $^{15}\text{NO}_3$ influx in in sugarcane roots submitted to nitrate resupply. Plants of 'NG57213', 'SP80-3280', 'IACSP96-2042' and 'IAC87-3396' sugarcane genotypes cultivated for three months under normal conditions of N supplementation (+ N; 2 mM NH_4NO_3), followed by N (-N) deficiency for 72 hours. After this period, they were supplemented with nitrate (4 mM KNO_3) for 4, 24 and 72 hours. Uptake rates were measured for 10 min. into sorghum and sugarcane genotypes roots at the external concentration of $200 \mu\text{M}$ ^{15}N -labeled NO_3^- (K^{15}NO_3). Values are means of 4 biological replicates $\pm$ SD, and significant differences at $P < 0.05$ are indicated by different letters.

In order to establish a correlation with the ^{15}N -nitrate influx, the analysis of expression of the main genes encoding the NRT2s and NRT3.1 involved in nitrate uptake process were conducted in *S. officinarum* and sugarcane hybrids cultivars under nitrate resupply treatment. After 4h of nitrate resupply, the commercial sugarcane hybrids 'SP80-3280' and 'IAC87-3396' displayed similar *ScNRT2.1* expression levels, both similar to plants of these genotypes under

N deficiency (0 h) (Figure 10). Despite the *ScNRT2.1* expression upregulation in *S. officinarum* genotype observed at 4 h of nitrate resupply, the transcript accumulation was not significantly distinct from what was observed in 0 h (Figure 10). For the 'IACSP96-2042', nitrate resupply treatment led to the repression of *ScNRT2.1* expression (Figure 10). After 24 h of nitrate resupply treatment, the *ScNRT2.1* mRNA accumulation in 'SP80-3280' roots was induced, although it does not differ from that observed in 'NG57213', which exhibited similar *ScNRT2.1* expression level to that observed in 0 h. 'IACSP96-2042' and 'IAC87-3396' roots displayed significantly lower levels of *ScNRT2.1* expression compared to the other genotypes after 24 h of nitrate resupply (Figure 10). In the last analyzed period (72 h), 'SP80-3280' and 'NG57213' showed the highest *ScNRT2.1* transcript accumulation, followed by 'IACSP96-2042' and 'IAC87-3396'. The *ScNRT2.1* relative expression observed in 'SP80-3280' after 72 h of treatment was significantly higher than those observed in 'IACSP96-2042' and 'IAC87-3396' in this period. 'NG57213' genotype showed an accumulation of *ScNRT2.1* transcripts significantly higher than that observed in genotype 'IAC87-3396' (Figure 10).

Figure 10: Regulation of the *ScNRT2.1* gene in sugarcane roots submitted to nitrate resupply. The transcript levels were determined in roots of 'NG57213', 'SP80-3280', 'IACSP96-2042' and 'IAC87-3396' sugarcane genotypes cultivated for three months under normal conditions of N supplementation (+ N; 2 mM NH_4NO_3), followed by N (-N) deficiency for 72 hours. After this period, they were supplemented with nitrate (4 mM KNO_3) for 4, 24 and 72 hours. Expression was normalized for mean expression values for plants under sufficient restriction for 72 h (-N; time 0 h). *ScUBQ2* gene was used as a reference. Bars indicate $\pm$ SE, $n = 4$ plants per time-point. Significant differences at $P < 0.05$ are indicated by different letters.

In summary, there was a progressive increase in the accumulation of *ScNRT2.1* transcripts in roots of plants of the 'SP80-3280' after exogenous nitrate application, reaching the maximum level 72 h after treatment, with 8.3-fold greater than that observed in plants

under N deficiency (Figure 10). Although it was possible to observe *ScNRT2.1* expression induction in the *S. officinarum* genotype 'NG57213' at 4 h and 72 h after nitrate resupply, this upregulation was not significantly distinct from that observed in N-deficient plants (Figure 10). For the 'IACSP96-2042' genotype, there was a reduction in the accumulation of transcripts in response to nitrate exposure, reaching the lowest value after 24 h of treatment. In the last analyzed period (72 h), there was an increase in the *ScNRT2.1* expression, reaching transcript accumulation similar to that observed in plants under N absence (Figure 10). *ScNRT2.1* expression in 'IAC87-3396' remained stable to nitrate resupply treatment until 24 h of treatment, when a reduction of approximately 50% in *ScNRT2.1* expression was observed; this low level remaining constant until 72 h after exogenous exposure of nitrate (Figure 10). Therefore, gene expression analyzes of the encoding gene for NRT2.1 transporter showed the sugarcane genotypes have different responses under nitrate resupply treatment.

Figure 11: Regulation of the *ScNRT2.5* gene in sugarcane roots submitted to nitrate resupply. The transcript levels were determined in roots of 'NG57213', 'SP80-3280', 'IACSP96-2042' and 'IAC87-3396' sugarcane genotypes cultivated for three months under normal conditions of N supplementation (+ N; 2 mM NH_4NO_3), followed by N (-N) deficiency for 72 hours. After this period, they were supplemented with nitrate (4 mM KNO_3) for 4, 24 and 72 hours. Expression was normalized for mean expression values for plants under sufficient restriction for 72 h (-N; time 0 h). *ScUBQ2* gene was used as a reference. Bars indicate $\pm$ SE, $n = 4$ plants per time-point. Significant differences at $P < 0.05$ are indicated by different letters.

Recent studies have identified the NRT2.5 protein in *Arabidopsis* as an important nitrate transporter in roots exposed to long periods of N deficiency (Lezhneva et al. 2014). In addition, NRT2.5 transporter, as NRT2.1, is able to establish the two-component high-affinity transport complex with the transmembrane protein NAR2.1, forming cHATS (constitutive HATS) transport in *Arabidopsis* roots (Kotur et al. 2012, Kotur and Glass 2015). Strong upregulation

of *ScNRT2.5* gene expression was observed in 'NG57213' roots after 4 h of nitrate resupply treatment, being approximately 20-fold higher than observed in N-deficient plants (0 h) (Figure 11). In the following periods (24 and 72 h), a decrease of *ScNRT2.5* transcript accumulation was noticed compared to that observed previously, even though the expression level was 4.5-fold higher than that observed in plants under the absence of N (Figure 11). For 'SP80-3280', nitrate exposure led to *ScNRT2.5* gene upregulation after 4 h. In this period, transcript accumulation was approximately 4.4-fold higher than it was observed at 0 h. This high relative expression level of *ScNRT2.5* maintained at similar levels on the next periods analyzed (24 and 72 h) (Figure 11). 'IACSP96-2042' plants under nitrate resupply exhibited quick induction of *ScNRT2.5* expression, approximately 4-fold higher compared to what was observed initially (0 h). In the following periods (24 and 72 h), the *ScNRT2.5* transcript accumulation declined, returning to what was observed in plants under N deficiency. 'IAC87-3396' showed rapid (4 h) and late (72 h) upregulation of the *NRT2.5* expression, with 5 and 3-fold higher in the respective periods when compared to that observed initially (0 h) (Figure 11). Therefore, differently from what was described in *Arabidopsis* (Lezhneva et al. 2014), *ScNRT2.5* expression analysis showed induction in response to nitrate resupply treatment, suggesting a fundamental role in the inducible HATS (iHATS) nitrate uptake in roots of parental and hybrids sugarcane genotypes.

ScNRT3.1 expression analysis showed significant and rapid (4 h) upregulation in plant roots of all sugarcane genotypes under nitrate resupply treatment, being substantially induced in the *S. officinarum* genotype, exhibiting approximately 23-fold more transcript accumulation than in plants under N deficiency (Figure 12). 'SP80-3280' showed the lowest *ScNRT3.1* induction in this period, but still significantly 3-fold higher than observed in N-deficient plants. The 'IACSP96-2042' and 'IAC87-3396' genotypes showed *ScNRT3.1* expression levels 10-fold and 6-fold higher than observed at 0 h (Figure 12). In the following period (24 h), *ScNRT3.1* transcript accumulation in 'IACSP96-2042' and 'IAC87-3396' roots reached similar levels as observed initially (0 h). After 24 h of treatment, 'SP80-3280' and 'NG57214' roots showed downregulation of *ScNRT3.1* expression to levels compared with observed in the previous period (4 h), but still were approximately 3 and 5-fold, respectively higher than displayed in plants under N deficiency (Figure 12). In the last period analyzed (72 h), all sugarcane genotypes showed similar and higher *ScNRT3.1* mRNA accumulation compared to the ones observed at 0 h. In summary, *S. officinarum* presented a rapid (4 h) and strong *ScNRT3.1* upregulation in response to nitrate resupply. The transcript accumulation maintained high during this treatment (Figure 12). 'IACSP96-2042' and 'IAC87-3396' showed rapid (4 h) and late (72 h) induction of *ScNRT3.1* expression in plants exposed to exogenous nitrate. Therefore, there is a distinct *ScNRT3.1* gene regulation among parental and hybrids sugarcane genotypes.

Figure 12: Regulation of the *ScNRT3.1* gene in sugarcane roots submitted to nitrate resupply. The transcript levels were determined in roots of 'NG57213', 'SP80-3280', 'IACSP96-2042' and 'IAC87-3396' sugarcane genotypes cultivated for three months under normal conditions of N supplementation (+ N; 2 mM NH_4NO_3), followed by N (-N) deficiency for 72 hours. After this period, they were supplemented with nitrate (4 mM KNO_3) for 4, 24 and 72 hours. Expression was normalized for mean expression values for plants under sufficient restriction for 72 h (-N; time 0 h). *ScUBQ2* gene was used as a reference. Bars indicate $\pm$ SE, $n = 4$ plants per time-point. Significant differences at $P < 0.05$ are indicated by different letters.

In summary, the differences in nitrate acquisition between the sugarcane genotypes occurred when plants were resupplied with nitrate, demonstrating the genetic variation for nitrate acquisition between sugarcane parental genotype and hybrid cultivars. The *S. officinarum* roots did not display a significant increment in the ^{15}N influx in response to nitrate resupply treatment (Figure 9). There was induction in the expression of *ScNRT2.5*, and *ScNRT3.1* after treatment for this genotype (Figures 11 and 12). Even with the *ScNRT3.1* expression upregulation in roots of 'NG57213' in response to nitrate resupply, there was no increase in the ^{15}N influx in the roots of this sugarcane parent. Such data can be supported by the non-regulation of *ScNRT2.1* considering it was not significantly induced under nitrate exposure treatment. As *ScNRT2.5* upregulation did not lead to the increase of ^{15}N -nitrate influx, this transporter would be acting in nitrate remobilization process in 'NG57213' plants under nitrate resupply treatment. So, the data suggest that the abundance of NRT2.1/NRT3.1 complex might remain stable in the plasma membrane of root cells in *S. officinarum* plants exposed to exogenous nitrate treatment. In 'IACSP96-2042' plants under exogenous nitrate exposure, *ScNRT2.1* expression was initially (4 and 24 h) downregulated. In the following period, the expression returned to a level similar to observed in N-deficient plants (Figure 10). *ScNRT2.5* and *ScNRT3.1* expression were upregulated in response to the nitrate resupply

treatment. The *ScNRT2.5* gene was induced in 'IACSP96-2042' plants exposed 4 h to this treatment (Figure 11). The *ScNRT3.1* expression was readily (4 h) and lately (72 h) induced compared to plants under N deficiency (0 h) (Figure 12). There was an increase of ^{15}N -nitrate influx only after 72 h. However, this increase was not significantly higher compared to 'IACSP96-2042' plants under the absence of N (Figure 9). Thus, as 'NG57213', nitrate acquisition process was not affected in plants of 'IACSP96-2042' by nitrate resupply treatment. The upregulation of the genes associated with the nitrate acquisition was observed in 'IAC87-3396' plants under nitrate resupply treatment. The ^{15}N -nitrate uptake was induced in response to treatment, reaching the highest influx value after 72 h of nitrate exposure (Figure 9). *ScNRT2.1* transcript accumulation was suppressed over the course of treatment (Figure 10), distinct from what was observed for *ScNRT2.5* and *ScNRT3.1* expression, which were induced at 4 and 72 h after treatment (Figures 11 and 12). Therefore, the results presented for this genotype suggest that the accessory protein NRT3.1 would be complexed with the NRT2.5 transporter, forming a heterooligomer. So, different from what was observed in the other sugarcane genotypes ('NG57213', 'IACSP96-2042', and 'SP80-3280'), nitrate transport may be regulated in roots of 'IAC87-3396' plants after nitrate exposure. Among the genotypes developed by the Agronomic Institute of Campinas (IAC), the sugarcane hybrid 'IAC87-3396' is the most planted in Brazil. Its planting area is spread over approximately 70,000 hectares in the Cerrado region, mainly due to its production stability in periods of drought (UDOP - National Bioenergy Union, 2020). Among the cultivars evaluated by Tasso Jr (2007), 'IAC87-3396' stood out in terms of yield and straw production. Besides, screening carried out by Dr. Joni using 22 hybrid genotypes from some breeding programs in Brazil showed that cultivar 'IAC87-3396' presented great values of root and shoot biomass production, and N uptake (under N sufficiency and N deficiency), so is considered as high NUE sugarcane genotype. So, the ^{15}N influx and gene expression analyses suggest that 'IAC87-3396' sugarcane hybrid presents high NUE due to capacity to regulate the nitrate uptake process after nitrate resupply.

In an overview of transcriptional and physiological regulation of HATS nitrate uptake process in sorghum and sugarcane, nitrate uptake was strongly induced in sorghum 'BTX623' and the sugarcane 'IAC87-3396' cultivar under nitrate resupply. For *S. officinarum* 'NG57213' and sugarcane hybrid 'IACSP96-2042', no regulation of nitrate uptake process occurred presumably due to unaffected *ScNRT2.1* expression in roots submitted to nitrate exposure. Only in 'SP80-3280' plants there was a lack of correlation between ^{15}N -nitrate influx and expression analysis of gene encoding HATS transporters. Since *ScNRT2.1* was strongly upregulated, this suggest the existence of a post-transcriptional regulation in 'SP80-3280' roots.

3.4.3 ELISA and Western Blot analysis

Compared to *NRT2.1* mRNA level and HATS activity, the level of NRT2.1 protein has been shown to be very stable in response to repressive environmental conditions, such as high N or darkness (Girin et al. 2007, Wirth et al. 2007, Laugier et al. 2012). To investigate the level of control of nitrate uptake in sorghum and sugarcane genotypes, if transcriptional, post-transcriptional or translational, the level of NRT2.1 protein was investigated first by ELISA to determine specificity of the antibodies, and by Western blotting on root total protein extracts from the experiment described above. Rabbit antisera raised against the synthetic peptides based on the presumed cytoplasmic C-terminal end of the NRT2.1 protein was developed (Figure 1). The PTA (Plate Trapped Antigen) ELISA was employed (Farias et al. 2012). In this assay, the two specific antibodies anti-NRT2.1 and anti-NRT3.1 were used against either sorghum ('BTX623') and sugarcane ('SP80-3280') genotypes root samples.

Figure 13: Cross reaction percentage, on time zero, from genotypes 'BTX623' (100%) in comparison to genotype 'SP80-3280' both antibodies (NRT2.1 and NRT3.1).

After titration, cross reaction was calculated based on sorghum 'BTX623' in comparison to 'SP80-3280' against both NRT2.1 and NRT3.1 antibodies. There were a high percentage of cross reaction among all antibodies and proteins extracted. So, ELISA test was not suitable to distinguish the proteins from genotypes. The test showed specificity of the antisera, but there wasn't discrimination between species. In terms of amino acid sequence, both species shared 90% and 92% of identity for the target region of anti-NRT2.1 and anti-NRT3.1, respectively (Figure 14).

Figure 14: NRT2.1 (A) and NRT3.1 (B) C-terminus region alignment performed using ClustalW software and displayed with BioEdit program to display the sequences similarity in the antibodies target regions.

ELISA is an easy and fast assay for screening proteins, genotypes and so on. The quality of this assay is due to the quality of each reagent, specially the antibody used (Crowther 2009). In case of polyclonal antibodies, which were used in this study, cross reaction can occur. Still, proteins expressed used to raise up the antibodies came from similar genetic events, which is expected to have similar antibodies, causing cross reaction for different genotypes (Duarte et al. 2006).

Western blot analysis of total proteins extracted from roots of 'SP80-3280' plants showed an increment in the NRT2.1 protein accumulation only after 72 h of nitrate resupply treatment, reaching protein levels 2-fold higher than in plants under N deficiency (0 h) (Figure 15). Quantification of protein bands showed a decrease in NRT2.1 protein accumulation in *S. officinarum* genotype roots after 24 h of treatment, reaching 60% of the total protein observed initially (0 h). After 72 h of nitrate exposure, there was an increase in the NRT2.1 abundance in 'GN57213' roots (Figure 15). There was a decrease in NRT2.1 protein accumulation in roots of 'IACSP96-2042' under nitrate exposure. Compared to that initially observed (0 h) for this genotype, the abundance of NRT2.1 protein was 75% lower after 24 h of treatment. In the last period (72 h), there was an increase in NRT2.1 accumulation, even less than that observed in plants under N absence (Figure 15). In 'IAC87-3396' roots, there was a strong accumulation of NRT2.1 protein in response to nitrate resupply treatment. At all times analyzed, the abundance of NRT2.1 was higher than that observed initially (0 h) (Figure 15). For the sorghum, a greater accumulation of NRT2.1 was observed in 'BTX623' roots after 24 and 72 h of exogenous nitrate application (Figure 15). Therefore, NRT2.1 accumulation presented by sorghum and sugarcane genotypes under nitrate resupply treatment display differential abundance of this transporter in all genotypes roots evaluated.

Figure 15: Western blot analysis and gel quantification of NRT2.1 protein accumulation in total protein extract of 'NG57213', 'SP80-3280', 'IACSP96-2042', 'IAC87-3396', and 'BTX623' genotypes cultivated for three months under normal conditions of N supplementation (+ N; 2 mM NH_4NO_3), followed by N (-N) deficiency for 72 hours. After this period, they were supplemented with nitrate (4 mM KNO_3) for 4, 24 and 72 hours. Total proteins from roots of these genotypes were separated by SDS-PAGE, transferred to a polyvinylidene fluoride membrane, and hybridized with a NRT2.1-specific polyclonal antibody. Each lane was loading equal quantity of proteins (60 μg).

Figure 16: Western blot analysis and gel quantification of NRT3.1 protein accumulation in total protein extract of 'NG57213', 'SP80-3280', 'IACSP96-2042', 'IAC87-3396', and 'BTX623' genotypes cultivated for three months under normal conditions of N supplementation (+ N; 2 mM NH_4NO_3), followed by N (-N) deficiency for 72 hours. After this period, they were supplemented with nitrate (4 mM KNO_3) for 4, 24 and 72 hours. Total proteins from roots of these genotypes were separated by SDS-PAGE, transferred to a polyvinylidene fluoride membrane, and hybridized with a NRT3.1-specific polyclonal antibody. Each lane was loading equal quantity of proteins (60 μg).

To determine if the level of NRT3.1 protein could explain the HATS nitrate uptake activity in sugarcane and sorghum root under resupply treatment, NRT3.1 protein levels was estimated after 4, 24, and 72 h of treatment. Plants of sugarcane 'SP80-3280' submitted to nitrate exposure exhibited a decrease of NRT3.1 protein accumulation. Compared to the initially observed (0 h), the abundance of NRT3.1 protein was 44% lower after 24 h of

treatment. In the last period (72 h), there was an increase in NRT3.1 accumulation, even less than its abundance observed in plants under N absence (Figure 16). A decrease of NRT3.1 abundance was observed in *S. officinarum* genotype roots after 4 h of nitrate resupply treatment. In the following periods, it was possible to notice an increase of NRT3.1 accumulation, reaching a level 13% higher (72 h) than observed in plants under N deficiency (0 h) (Figure 16). The NRT3.1 abundance in roots of 'IACSP96-2042' under nitrate exposure drastically decreased. The lowest level of NRT3.1 protein accumulation observed was 24 h after nitrate exposure, which was 80% lower compared to N-deficient plants (Figure 16). In 'IAC87-3396' roots, there was an increase in NRT3.1 protein accumulation after 24 and 72 h of nitrate resupply treatment (Figure 16). The pattern of NRT3.1 protein abundance in sorghum was the same as observed in *S. officinarum*. After a small decrease of NRT3.1 protein accumulation, 'BTX623' roots displayed an increase in its abundance, reaching approximately 50% higher level than observed in 0 h (Figure 16). Therefore, as observed for NRT2.1 accumulation presented by sorghum and sugarcane genotypes under nitrate resupply treatment, the genotypes displayed distinct NRT3.1 abundance in roots.

To conclude, Western blot analysis in roots exhibited distinct NRT2.1 and NRT3.1 protein accumulation among sorghum and sugarcane genotypes under nitrate resupply treatment. Plants of sugarcane cultivar 'SP80-3280' showed a late (72 h) increase of NRT2.1 accumulation, and a decrease of NRT3.1 protein abundance in response to nitrate exposure. *S. officinarum* roots displayed a light and late increase in the accumulation for both NRT2.1 and NRT3.1 protein in 'NG57213' plants submitted to nitrate exposure. Exogenous nitrate application in roots of 'IACSP96-2042' plants led to a decrease of both protein accumulation in root cells. Among all genotypes, the sugarcane hybrid 'IAC87-3396' showed distinct NRT2.1 and NRT3.1 abundance, with high abundance of both protein in response to nitrate resupply treatment. Compared with 'BTX623' roots under N deficiency, plants submitted to nitrate exposure presented an increase of both NRT2.1 and NRT3.1 protein accumulation.

3.5 Discussion

The modern sugarcane cultivars are interspecific hybrids derived from crosses between *Saccharum officinarum* ($2n = 8x = 80$) and *Saccharum spontaneum* ($2n = 5x-16x = 40-128$), followed by various backcrossing to *S. officinarum* (Grivet et al. 2004). In consequence, commercial sugarcane hybrids have a complex chromosome set, ranging between $2n = 100-130$, with 80% of the chromosomes originated from *S. officinarum*, 10–15% from *S. spontaneum* origin, and the remaining are recombined chromosomes (Jannoo et al. 1999, D'Hont et al. 2005). The extreme polyploid genome of interspecific hybrids possesses irregular genetic characteristics that are passed from both parental species, making it more complicated than that of its precursors (D'Hont et al. 1996). Because of the random sorting of

chromosomes and chromatids in each crossing, the number of chromosomes varies between genotypes (aneuploids) and contributes to the high level of heterozygosity (Ming et al. 1998). Therefore, the mixed genetic composition of modern sugarcane hybrids also results in complex control of important agronomic traits, such as nitrate uptake process.

Temporal analysis revealed distinct regulation of HATS nitrate uptake process in roots of sorghum and sugarcane under nitrate resupply (Table 2). Sorghum 'BTX623' displayed high influx of labeled nitrate in response to the resupply, confirming the low nitrate uptake capacity of sugarcane roots and corroborating the results presented by Robinson et al. (2011), indicating the low NUE in sugarcane. However, increased expression of genes from the *SbNRT2* (Figures 4 and 5) and *SbNRT3* (Figures 6 and 7) subfamily after 4 h of nitrate resupply did not lead to more accumulation of these proteins in sorghum (Figures 15 and 16). I.e., there is a correlation of *SbNRT2* and *SbNRT3* expression with the nitrate labeled influx in this period, but there is no correspondence between transcript accumulation with the accumulation of NRT2.1 and NRT3.1 proteins, which form the HATS complex responsible for the nitrate uptake in roots of sorghum. The late upregulation of the expression of the *NRT2.1* and *NRT3.1* genes matches with the increase in ^{15}N -nitrate influx and the abundance of these proteins, showing that the NRT2.1/NRT3.1 complex would be active in sorghum roots after 72 h of resupply of nitrate.

In 'SP80-3280' plants, there was a complete lack of correlation between ^{15}N -nitrate influx and expression analysis of genes encoding HATS transporters in roots under nitrate resupply (Table 2). Even with the upregulation of *ScNRT2.1* and *ScNRT3.1* (Figures 10 and 12), no regulation was observed in nitrate uptake process since the ^{15}N -nitrate influx remained at the same rates as observed in N-deficient plants (Figures 3 and 9). It is important to highlight that the upregulation of *ScNRT2.1* led to an increase in protein accumulation 72 h after the treatment (Figures 10 and 14), but the higher expression of *ScNRT3.1* did not lead to a high abundance of NRT3.1 protein in 'SP80-3280' roots (Figures 12 and 16). Therefore, these results suggest the existence of post-translational regulation of NRT2.1 protein and post-transcriptional control of *ScNRT3.1* in 'SP80-3280' plants under nitrate resupply treatment. These controls affect the efficiency in nitrate uptake by this cultivar.

Table 2: Review of ^{15}N influx, gene expression of the NRT2 and NRT3 members, and NRT2.1 and NRT3.1 proteins abundance analysis in roots of sorghum and sugarcane plants submitted to nitrate resupply treatment. Significant increase and decrease obtained in the analysis follow the legend below.

Analysis	Sorghum			Sugarcane											
	'BTX623'			'SP80-3280'			'NG57213'			'IACSP96-2042'			'IAC87-3396'		
	Treatments (hours)			Treatments (hours)			Treatments (hours)			Treatments (hours)			Treatments (hours)		
	4	24	72	4	24	72	4	24	72	4	24	72	4	24	72
^{15}N -nitrate influx	+++++	=	+++++	=	=	=	=	=	=	=	=	=	++++	++++	+++++
NRT2.1 expression	=	=	=	=	=	+++++	=	=	=	-	--	=	=	-	-
NRT2.2 expression	+++	=	=												
NRT2.5 expression	=	=	-	++++	+++++	++++	+++++	++++	+++++	+++	=	=	+++++	=	++++
NRT3.1 expression	++	=	=	++	++	++++	+++++	+++++	+++++	+++++	=	+++++	+++++	=	++
NRT3.2 expression	=	=	=												
NRT2.1 protein abundance	-	+	+	-	-	+	-	-	+	-	--	-	++	++	+
NRT3.1 protein abundance	-	+	+	-	-	-	-	-	+	-	-----	--	-	+	+

Significant Increased or decreased from 0 h (N-deficient plants):

+ or - : 1-2 fold

++ or -- : 2-3 fold

+++ or --- : 3-4 fold

++++ or ---- : 4-5fold

+++++ or ----- : > 5 fold

= : no significant difference from 0 h

For *S. officinarum* genotype 'NG57213', there was an induction of *ScNRT2.1* and *ScNRT3.1* expression in all periods after exogenous nitrate application (Figures 10 and 12). However, only after 72 h, it was noticed a higher abundance of both NRT2.1 and NRT3.1 proteins compared to that observed in plants under N deficiency (0 h) (Figures 15 and 16). Even with the upregulation of both genes and the late (72 h) increase in NRT2.1 and NRT3.1 proteins accumulation, it was not possible to see a significant difference in the ^{15}N -nitrate influx between plants of *S. officinarum* submitted to the absence of N deficiency and nitrate resupply treatments (Figure 9 and Table 2). Such results suggest the presence of post-transcriptional and translational regulation in the nitrate uptake process in the roots of *S. officinarum* under nitrate exposure. This control did not resemble any of the hybrid commercial cultivar suggesting that either there is genetic diversity for these controls in *S. officinarum* or some of these regulations derive from the other sugarcane genitor, *S. spontaneum*.

No regulation of nitrate uptake process was observed in roots of 'IACSP96-2042' plants under nitrate resupply (Table 2). No induction of *ScNRT2.1* expression was determined in response to nitrate application (Figure 10). However, expression of the gene coding for NRT3.1 auxiliary protein was induced after 4 and 72 h (Figure 11), showing that there was no co-expression of both genes in response to nitrate resupply treatment. A decrease of both NRT2.1 and NRT3.1 abundance was observed in 'IACSP96-2042' roots under nitrate exposure (Figures 15 and 16). Therefore, the results revealed that there was a transcriptional control of the *ScNRT2.1* gene due to inhibition of its expression, and post-transcriptional regulation of the *ScNRT3.1* since its transcript accumulation did not lead to an increase in NRT3.1 abundance in roots of this genotype.

Among the sugarcane genotypes, 'IAC87-3396' highlighted the strong induction of nitrate uptake under nitrate resupply treatment (Table 2). The rates of ^{15}N -nitrate were the same, or even higher than showed by sorghum 'BTX623' (Figures 3 and 9). In 'IAC87-3396' roots under nitrate exposure, there was co-expression between *ScNRT2.5* and *ScNRT3.1* (Figures 11 and 12), suggesting that the accessory protein NRT3.1 would be complexed with the NRT2.5 transporter, forming an activated HATS complex. Although the *ScNRT2.1* mRNA accumulation was suppressed over the course of treatment (Figure 10), Western blot analysis revealed an increase of NRT2.1 protein abundance after 'IAC87-3396' N-deficient were submitted to exogenous nitrate addition (Figure 15). Further, it was observed a high abundance of NRT3.1 protein in roots submitted to this treatment (Figure 16). Except for AtNRT2.7, all other NRT2 transporters interact strongly with AtNRT3.1 (Kotur et al. 2012). Therefore, besides the presence of NRT2.1/NRT3.1 complex in 'IAC87-3396' roots under nitrate resupply treatment, the previous results suggest the

presence of other HATS complex composed by NRT2.5 instead of NRT2.1 in this sugarcane cultivar. Different from what was described by Lezhneva et al. (2014), the upregulation of *ScNRT2.5* expression in 'IAC87-3396' plants suggest a fundamental role of NRT2.5 protein in the inducible HATS (iHATS) nitrate uptake in roots under nitrate resupply treatment.

To face the changing N demand during plant growth as well as environmental variation, nitrate HATS activity is rapidly and accurately modulated. It has been assumed that nitrate uptake by HATS is predominantly regulated by the transcriptional control of *NRT2.1* expression. Studies have shown that *NRT2.1* expression is induced upon nitrate supply (Lejay et al. 1999, Girin et al. 2007), repressed by N metabolites or high nitrate provision (Lejay et al. 1999, Gansel et al. 2001, Muños et al. 2004, Krouk et al. 2006, Girin et al. 2007), and upregulated by light and sugars (Lejay et al. 1999, Lejay et al. 2003, Lejay et al. 2008). Nevertheless, previous studies evidencing a lack of correlation between *NRT2* gene expression, NRT2 protein abundance and HATS activity, suggest that in addition to transcriptional regulation, protein-protein interactions and posttranslational regulation of NRT2.1 might play an essential role in modulating the activity of this nitrate transporter.

Using anti-NRT2.1 antibodies, researchers have described that changes in NRT2.1 levels do not match fluctuations in *NRT2.1* transcript levels (Wirth et al. 2007). When NRT2.1 is constitutively overexpressed under the control of a 35S promoter, the nitrate HATS activity is still inhibited under high N supply (Laugier et al. 2012). Thus, the abundance of NRT2.1 protein in the plasma membrane shows much slower changes than in *NRT2.1* transcript accumulation or nitrate HATS activity, suggesting activation/inactivation of the NRT2.1/NRT3.1 transport system at the plasma membrane. Another aspect, which could involve the occurrence of post-translational modifications in NRT2.1 transporter concerns its role in the control of root development in a way that is independent from NRT2.1 transport activity (Little et al. 2005, Remans et al. 2006). This rise the idea that, like NRT1.1 (NPF6.3), NRT2.1 protein may also act as a nitrate sensor, or a signal transducer, although the mechanisms involved are still not known (Little et al. 2005). Since the sensing function of NPF6.3 depends on its T101 phosphorylation (Ho et al. 2009, Bouguyon et al. 2015), it supports the hypothesis that post-translational modifications of NRT2.1 could also be involved in the sensing function of NRT2.1.

In fact, some recent studies have revealed that phosphorylation would affect the activity and stability of NRT2.1 in plasma membrane of root cells. Phosphoproteomic approach using *Arabidopsis* plants under nitrate stimulation (Engelsberger and Schulze 2012, Wang et al. 2013) or nitrate deprivation (Menz et al. 2016), indicated several phosphorylation sites for NRT2.1. Zou et al (2019) observed that in response to low nitrate availability, the NRT2.1 stability increased

rapidly and prior to the transcriptional stimulation of NRT2.1. Also, at the same time scale, the phosphorylation level of NRT2.1 serine 28 (Ser28) increased substantially. Using Arabidopsis transgenic plants expressing a phosphor-mimicking mutant (NRT2.1^{S28E}), and a loss-of-phosphorylation mimicking mutant (NRT2.1^{S28A}) they confirmed that phosphorylation at Ser28 is crucial for NRT2.1 protein stabilization and accumulation under nitrate limitation (Zou et al. 2019). Additionally, Li et al (2020) identified the Nitrate Uptake Regulatory Kinase 1 (NURK1), found to interact with NRT2.1 at dephosphorylated S28. Upon that interaction, NURK1 can phosphorylate NRT2.1 at S21. This phosphorylation would result in low interaction of NRT2.1 transporter with the auxiliary protein NRT3.1. By contrast, phosphorylation of NRT2.1 at S28 by a yet unknown kinase enhanced the interaction with NRT3.1 by the inhibition of NURK1 interaction (Li et al. 2020). Thus, the presence of kinases interacting with NRT2.1 and controlling the formation of two component complex with NRT3.1 protein is one more proof for the presence of a post-translational regulation of HATS activity in the nitrate uptake process in Arabidopsis roots. Ser28 is not conserved among monocotyledonous and dicotyledonous species (Figure 17), being present only in *Arabidopsis thaliana*, *Chrysanthemum morifolium*, and *Manihot esculenta*. Therefore, this multiple alignment suggests that a similar mechanism identified by Zou et al. (2019) and Li et al. (2020) would not be regulating the activity of the NRT2.1 transporter in sorghum and sugarcane.

Figure 17: NRT2.1 N-terminus region alignment performed using ClustalW software and displayed with BioEdit program.

Using a high-accuracy mass spectrometric phosphopeptide detection, Jacquot et al. (2020) found another residue in NRT2.1 protein sequence, this time at the C-terminal, target of phosphorylation. Experiments with Arabidopsis transgenic plants with truncated forms of NRT2.1 showed that its S501 residue phosphorylation has a strong impact on NRT2.1 transporter activity independently of the environmental conditions. The alignment of NRT2.1 C-terminus reveals that the serine corresponding to S501 in Arabidopsis is remarkably conserved across dicotyledons and monocotyledons species, except for wheat and Brachypodium (Figure 18). These similarity between essential amino acids for either NRT2.1 activity and/or interaction with partner proteins

among different plant species, could suggest the presence of a post-translational regulation mechanism at NRT2.1 transporter in sorghum and sugarcane roots.

Figure 18: NRT2.1 C-terminus region alignment performed using ClustalW software and displayed with BioEdit program.

But NRT2.1 phosphorylation does not prevent its interaction with NRT3.1 (Kotur et al. 2017). Yong et al. (2010) found a HATS complex of only ~ 150kDa that would be a result of the interaction of two subunits each of NRT2.1 and NRT3.1. Jacquot et al. (2020) found a large protein complex (~ 480kDa), suggesting that HATS system could contain several subunits of NRT2.1 and NAR2.1 associated with other proteins. In fact, inactivation of a transporter through phosphorylation of the C-terminal region of the protein has already been demonstrated for the ammonium transporter AMT1.1 (Loqué et al. 2007). Indeed, AtAMT1.1 in plants works as a trimer, which activity is controlled by the spatial positioning of the C-terminus. When ammonium is added in the growth medium, it triggers rapid phosphorylation of a conserved threonine residue (T460) in the C-terminus of AMT1.1 in a time- and concentration-dependent manner (Loqué et al. 2007, Lanquar et al. 2009). This phosphorylation of T460 in response to an increase in external ammonium correlates with a reduction of ammonium uptake activity in roots. These results lead to a model in which T460 phosphorylation induces a conformational change and that a single phosphorylation event in the C-terminus of one monomer is sufficient for cooperative closure of the trimer (Lanquar et al. 2009, Lanquar and Frommer 2010).

Therefore, the lack of correlation between the accumulation of NRT2.1 and NRT3.1 proteins with the ¹⁵N-nitrate influx suggest the existence of a mechanism controlling nitrate transport in sorghum and sugarcane roots at the post-translational level. I.e., it does not matter if there was an increase in the accumulation of NRT2.1 and NRT3.1 proteins in roots after nitrate resupply treatment. What matters is whether both proteins present in the root cells are forming the active HATS complex. This conclusion is because it can be recognized that in some genotypes the accumulation of NRT2.1 and NRT3.1 after plants were submitted to nitrate resupply did not lead

to an increase of nitrate-labeled uptake. Another situation observed was that even though the abundance of both proteins remained similar to that observed in plants under N deficiency, there was an increase in the influx of ^{15}N -nitrate. NRT2.1 transporter in both species, due to the presence of S501, could have its activity regulated by an unknown kinase, affecting not more a dual component complex, but a multi-component HATS complex formation. This putative allosteric regulation of HATS complex would be the major actor on nitrate uptake in roots of sorghum and sugarcane.

3.6 Conclusion

Overall, this work revealed distinct regulation of HATS nitrate uptake process in roots of sorghum and sugarcane plants under nitrate resupply. ^{15}N -nitrate influx showed high nitrate uptake by sorghum 'BTX623' and sugarcane hybrid 'IAC87-3396'. Expression analysis of genes coding for NRT2 and NRT3, protein that form HATS nitrate uptake complex, exhibit different transcriptional regulation among sugarcane genotypes under nitrate exposure. The lack of correlation between the accumulation of NRT2.1 and NRT3.1 proteins with the ^{15}N -nitrate influx suggest that HATS complex would not be fully active, proposing the presence of a mechanism controlling nitrate transport in sorghum and sugarcane roots at the post-translational level. The knowledge obtained from this work may help to elucidate which is the critical step control in both species. Further approaches could be used to confirm the phosphorylation site (S501) in NRT2.1 transporter and the real HATS complex in sorghum and sugarcane genotypes.

References

- ALVAREZ JM, VIDAL EA, and GUTIÉRREZ RA (2012) Integration of local and systemic signaling pathways for plant N responses. *Curr Opin Plant Biol*, 15: 185-191. DOI: 10.1016/j.pbi.2012.03.009
- BOUGUYON E, BRUN F, MEYNARD D, et al. (2015). Multiple mechanisms of nitrate sensing by *Arabidopsis* nitrate transceptor NRT1.1. *Nature Plants*, 1: 15015. DOI: 10.1038/nplants.2015.15
- BURNEY JA, DAVIS SJ, and LOBELL DB (2010) Greenhouse gas mitigation by agricultural intensification. *Proc Natl Acad Sci USA*, 107: 12052-12057. DOI: 10.1073/pnas.0914216107
- BURT TP, HOWDEN NJK, WORRALL F, and WHELAN MJ (2008) Importance of long-term monitoring for detecting environmental change: Lessons from a lowland river in south east England. *Biogeosciences*, 5: 1529-1535. DOI: 10.5194/bg-5-1529-2008
- CEREZO M, TILLARD P, FILLEUR S, et al. (2001) Major alterations of the regulation of root NO₃⁻ uptake are associated with the mutation of Nrt2.1 and Nrt2.2 genes in *Arabidopsis*. *Plant Physiol*, 127(1): 262-271. DOI: 10.1104/pp.127.1.262
- CHIU CC, LIN CS, HSIA AP, et al. (2004) Mutation of a nitrate transporter, AtNRT1:4, results in a reduced petiole nitrate content and altered leaf development. *Plant Cell Physiol*, 45: 1139-1148. DOI: 10.1093/pcp/pch143
- CRAWFORD NM and GLASS ADM (1998) Molecular and physiological aspects of nitrate uptake in plants. *Trends Plant Sci*, 3: 389-395. DOI: 10.1016/S1360-1385(98)01311-9
- CROWTHER JR (2009) The ELISA Guidebook 2nd Ed. Series Springer Protocols. Methods Mol Biol, 516, 566 p. DOI: 10.1007/978-1-60327-254-4
- DAVIDSON EA (2009) The contribution of manure and fertilizer nitrogen to atmospheric nitrous oxide since 1860. *Nat Geosci*, 2: 659-662. DOI: 10.1038/ngeo608
- D'HONT A, GRIVET L, FELDMANN P, et al. (1996) Characterization of the double genome structure of modern sugarcane cultivars (*Saccharum spp.*) by molecular cytogenetics. *Molec Gen Genet*, 250: 405-413. DOI: 10.1007/BF02174028
- D'HONT A (2005). Unraveling the genome structure of polyploids using FISH and GISH; examples of sugarcane and banana. *Cytogenet Genome Res*, 109: 27-33. DOI: 10.1159/000082378
- DUARTE KMR, GOMES LH, ANDRINO FG, et al. (2002) Identificação do vírus do mosaico do tomateiro (tomv) *Tobamovirus*, por meio de anticorpos monoclonais. *Sci Agri*, 59(1): 107-112. DOI: 10.1590/S0103-90162002000100016
- DUARTE KMR, GOMES LH, PASCHOAL JAR, and MEIRELLES CF (2006) A fast immunoassay for the screening of alfa-trembolone in bovine urine. *Braz J Vet Res An Sci*, 43: 74-79. DOI: 10.11606/issn.1678-4456.bjvras.2006.26538
- ENGELSBERGER WR and SCHULZE WX (2012) Nitrate and ammonium lead to distinct global dynamic phosphorylation patterns when resupplied to nitrogen-starved *Arabidopsis* seedlings. *Plant J*, 69: 978-995. DOI: 10.1111/j.1365-313X.2011.04848.x
- FAN SC, LIN CS, HSU PK, et al. (2009) The *Arabidopsis* Nitrate Transporter NRT1.7, expressed in phloem, is responsible for source-to-sink remobilization of nitrate. *Plant Cell*, 21(9): 2750-2761. DOI: 10.1105/tpc.109.067603
- FARIAS JF, DUARTE KMR, GOMES LH, et al. (2012) Polyclonal antibodies against isoflavones: fast screening method for Forage legume species. *IOSR J Pharm*, 2: 537-539. ISSN: 2250-3013
- FENG H, YAN M, FAN X, et al. (2011) Spatial expression and regulation of rice high-affinity nitrate transporters by nitrogen and carbon status. *J Exp Bot*, 62(7): 2319-32. DOI: 10.1093/jxb/erq403
- FIGUEIRA TR, OKURA V, RODRIGUES DA SILVA F, et al. (2012) A BAC library of the SP80-3280 sugarcane variety (*Saccharum sp.*) and its inferred microsynteny with the *Sorghum* genome. *BMC Res*, 5:185. DOI: 10.1186/1756-0500-5-185
- FILLEUR S, DORBE MF, CEREZO M, et al. (2001) An *Arabidopsis* T-DNA mutant affected in Nrt2 genes is impaired in nitrate uptake. *FEBS Lett*, 489(2-3): 220-224. DOI: 10.1016/S0014-5793(01)02096-8
- FOLEY JA, RAMANKUTTY N, BRAUMAN KA, et al. (2011) Solutions for a cultivated planet. *Nature*, 478: 337-342. DOI: 10.1038/nature10452
- FORDE BG and CLARKSON DT (1999) Nitrate and ammonium nutrition of plants: physiological and molecular perspectives. *Adv Bot Res*, 30: 1-90. DOI: 10.1016/S0065-2296(08)60226-8
- FRAISIER V, GOJON A, TILLARD P, and DANIEL-VEDELE F (2000) Constitutive expression of a putative high-affinity nitrate transporter in *Nicotiana plumbaginifolia*: evidence for post-transcriptional

- regulation by a reduced nitrogen source. *Plant J*, 23(4): 489-496. DOI: 10.1046/j.1365-313x.2000.00813.x
- FRANCO HCJ, OTTO R, FARONI CE, et al. (2011) Nitrogen in sugarcane derived from fertilizer under Brazilian field conditions. *Field Crops Res*, 121: 29-41. DOI: 10.1016/j.fcr.2010.11.011
- GALLOWAY JN, DENTENER FJ, CAPONE DG, et al. (2004) Nitrogen Cycles: Past, Present, and Future. *Biogeochem*, 70: 153-226. DOI: 10.1007/s10533-004-0370-0
- GALLOWAY JN, TOWNSEND AR, ERISMAN JW, et al. (2008) Transformation of the nitrogen cycle: recent trends, questions, and potential solutions. *Science*, 320(5878): 889-892. DOI: 10.1126/science.1136674
- GANSEL X, MUÑOS S, TILLARD P, and GOJON A (2001) Differential regulation of the NO_3^- and NH_4^+ transporter genes *AtNrt2.1* and *AtAmt1.1* in *Arabidopsis*: relation with long-distance and local controls by N status of the plant. *Plant J*, 26: 143-155. DOI: 10.1046/j.1365-313x.2001.01016.
- GARNETT T, CONN V, PLETT D, et al. (2013) The response of the maize nitrate transport system to nitrogen demand and supply across the lifecycle. *New Phytol*, 198(1): 82-94. DOI: 10.1111/nph.12166
- GELLI M, DUO Y, KONDA AR, et al. (2014) Identification of differentially expressed genes between sorghum genotypes with contrasting nitrogen stress tolerance by genome-wide transcriptional profiling. *BMC Genomics*, 15: 179. DOI: 10.1186/1471-2164-15-179
- GIRIN T, LEJAY L, WIRTH J, et al. (2007) Identification of a 150 bp cis-acting element of the *AtNRT2.1* promoter involved in the regulation of gene expression by the N and C status of the plant. *Plant Cell Environ*, 30: 1366-1380. DOI: 10.1111/j.1365-3040.2007.01712.x
- GOOD AG, SHRAWAT AK, and MUENCH DG (2004) Can less yield more? Is reducing nutrient input into the environment compatible with maintaining crop production? *Trends Plant Sci*, 9: 597-605. DOI: 10.1016/j.tplants.2004.10.008
- GLASS ADM, BRITTO DT, KAISER BN, et al. (2002) The regulation of nitrate and ammonium transport systems in plants. *J Exp Bot*, 53: 855-864. DOI: 10.1093/jexbot/53.370.855
- GLASS ADM (2003) Nitrogen use efficiency of crop plants: physiological constraints upon nitrogen absorption. *Crit Rev Plant Sci*, 22(5): 453-470. DOI: 10.1080/07352680390243512
- GRIVET L, DANIELS C, GLASZMANN JC, and D'HONT A (2004) A review of recent molecular genetics evidence for sugarcane evolution and domestication. *Ethnobot Res Appl*, 2: 9-17. Accessed in www.ethnobotanyjournal.org/vol2/i1547-3465-02-009.pdf
- GUTIÉRREZ RA, STOKES TL, THUM K, et al. (2008) Systems approach identifies an organic nitrogen-responsive gene network that is regulated by the master clock control gene *CCA1*. *Proc Natl Acad Sci USA*, 105(12): 4939-4944. DOI: 10.1073/pnas.0800211105
- GUTIÉRREZ RA (2012) Systems biology for enhanced plant nitrogen nutrition. *Science*, 336(6089): 1673-1675. DOI: 10.1126/science.1217620
- HO CH, LIN SH, HU HC, and TSAY YF (2009) *CHL1* functions as a nitrate sensor in plants. *Cell*, 138: 1184-1194. DOI: 10.1016/j.cell.2009.07.004
- HOAGLAND DR and ARNON DI (1950) The water culture method for growing plants without soil. University of California, Agriculture Experimental Station. Circular 347.
- HOBEN JP, GEHL RJ, MILLAR N, et al. (2011) Nonlinear nitrous oxide (N_2O) response to nitrogen fertilizer in on-farm corn crops of the US Midwest. *Glob Change Biol*, 17: 1140-1152. DOI: 10.1111/j.1365-2486.2010.02349.x
- HSU PK and TSAY YF (2013) Two phloem nitrate transporters, *NRT1.11* and *NRT1.12*, are important for redistributing xylem-borne nitrate to enhance plant growth. *Plant Physiol*, 163(2): 844-856. DOI: 10.1104/pp.113.226563
- ISHIKAWA S, ITO Y, SATO Y, et al. (2009) Two-component high-affinity nitrate transport system in barley: membrane localization, protein expression in roots and a direct protein-protein interaction. *Plant Biotechnol*, 26: 197-205. DOI: 10.5511/plantbiotechnology.26.197
- JACQUOT A, CHAPUT V, MAURIÈS A, et al. (2020) *NRT2.1* C-terminal phosphorylation prevents root high affinity nitrate uptake activity in *Arabidopsis thaliana*. *New Phytol*. DOI: 10.1111/nph.16710
- JANNOO N, GRIVET L, SEGUIN M, et al. (1999) Molecular investigation of the genetic base of sugarcane cultivars. *Theor Appl Genet*, 99: 171-84. DOI: 10.1007/s001220051222
- JAYNES DB, COLVIN TS, KARLEN DL, et al. (2001) Nitrate loss in subsurface drainage as affected by nitrogen fertilizer rate. *J Environ Qual*, 30: 1305-1314. DOI: 10.2134/jeq2001.3041305x

- KHOO HH (2015) Review of bio-conversion pathways of lignocellulose-to-ethanol: Sustainability assessment based on land footprint projections. *Renew Sust Energ Rev*, 46: 100-119. DOI: 10.1016/j.rser.2015.02.027
- KIBA T, KUDO T, KOJIMA M, and SAKAKIBARA H (2011) Hormonal control of nitrogen acquisition: roles of auxin, abscissic acid, and cytokinin. *J Exp Bot*, 62(4): 1399-1409. DOI: 10.1093/jxb/erq410
- KIBA T, FERIA-BOURRELLIER AB, LAFOUGE F, et al. (2012) The Arabidopsis nitrate transporter NRT2.4 plays a double role in roots and shoots of nitrogen-starved plants. *Plant Cell*, 24(1): 245-258. DOI: 10.1105/tpc.111.092221
- KINGSTON G, ANINK MC, and ALLEN D (2008) Acquisition of nitrogen by ratoon crops of sugarcane as influenced by waterlogging and split applications. *Proc Conf Aust Soc Sugar Cane Technol*, 30: 202-211.
- KONISHI M and YANAGISAWA S (2013) Arabidopsis NIN-like transcription factors have a central role in nitrate signalling. *Nat Commun*, 4: 1617. DOI: 10.1038/ncomms2621
- KOTUR Z, MACKENZIE N, RAMESH S, et al. (2012) Nitrate transport capacity of the *Arabidopsis thaliana* NRT2 family members and their interactions with AtNAR2.1. *New Phytol*, 194(3): 724-731. DOI: 10.1111/j.1469-8137.2012.04094.x
- KOTUR Z and GLASS ADM (2015) A 150 kDa Plasma membrane complex of AtNRT2.5 and AtNAR2.1 is the major contributor to constitutive high-affinity nitrate influx in *Arabidopsis thaliana*. *Plant Cell Environ*, 38(8): 1490-1502. DOI: 10.1111/pce.12496
- KOTUR Z, UNKLES SE, and GLASS ADM (2017). Comparisons of the *Arabidopsis thaliana* high-affinity nitrate transporter complex AtNRT2.1/AtNAR2.1 and the *Aspergillus nidulans* AnNRTA: structure function considerations. *Isr J Plant Sci*, 64: 21-31. DOI: 10.1080/07929978.2016.1246147
- KRAPP A, DAVID LC, CHARDIN C, et al. (2014) Nitrate transport and signalling in Arabidopsis. *J Exp Bot*, 65: 789-798. DOI: 10.1093/jxb/eru001
- KROUK G, TILLARD P, and GOJON A (2006) Regulation of the high-affinity NO₃⁻ uptake system by NRT1.1-mediated NO₃⁻ demand signaling in Arabidopsis. *Plant Physiol*, 142(3): 1075-1086. DOI: 10.1104/pp.106.087510
- KROUK G, LACOMBE B, BIELACH A, et al. (2010) Nitrate-regulated auxin transport by NRT1.1 defines a mechanism for nutrient sensing in plants. *Dev Cell*, 18(6): 927-937. DOI: 10.1016/j.devcel.2010.05.008
- LANQUAR V, LOQUE D, HORMANN F, et al. (2009) Feedback inhibition of ammonium uptake by a phospho-dependent allosteric mechanism in Arabidopsis. *Plant Cell*, 21: 3610-3622. DOI: 10.1105/tpc.109.068593
- LANQUAR V and FROMMER WB (2010) Adjusting ammonium uptake via phosphorylation. *Plant Signal Behav*, 5: 736-738. DOI: 10.4161/psb.5.6.11696
- LAUGIER E, BOUGUYON E, MAURIÈS A, et al. (2012) Regulation of high-affinity nitrate uptake in roots of Arabidopsis depends predominantly on posttranscriptional control of the NRT2.1/NAR2.1 transport system. *Plant Physiol*, 158(2): 1067-1078. DOI: 10.1104/pp.111.188532
- LEAL GA, ALBUQUERQUE PSB, and FIGUEIRA A (2007) Genes differentially expressed in *Theobroma cacao* associated with resistance to witches' broom disease caused by *Crinipellis pernicioso*. *Mol Plant Pathol*, 8(3): 279-292. DOI: 10.1111/j.1364-3703.2007.00393.x
- LEE TSG (1987) Micropropagation of sugarcane (*Saccharum spp.*). *Plant Cell Tiss Org*, 10: 47-55. DOI: 10.1007/BF00037496
- LEJAY L, TILLARD P, LEPETIT M et al. (1999) Molecular and functional regulation of two NO₃⁻ uptake systems by N- and C-status of Arabidopsis plants. *Plant J*, 18: 509-519. DOI: 10.1046/j.1365-313X.1999.00480.x
- LEJAY L, GANSEL X, CEREZO M, et al. (2003) Regulation of root ion transporters by photosynthesis: Functional importance and relation with hexokinase. *Plant Cell*, 15: 2218-2232. DOI: 10.1105/tpc.013516
- LEJAY L, WIRTH J, PERVENT M, et al. (2008) Oxidative pentose phosphate pathway-dependent sugar sensing as a mechanism for regulation of root ion transporters by photosynthesis. *Plant Physiol*, 146(4): 2036-2053. DOI: 10.1104/pp.107.114710
- LERÁN S, VARALA K, BOYER JC, et al. (2014) A unified nomenclature of NITRATE TRANSPORTER 1/PEPTIDE TRANSPORTER family members in plants. *Trends Plant Sci*, 19(1): 5-9. DOI: 10.1016/j.tplants.2013.08.008

- LEZHNEVA L, KIBA T, FERIA-BOURRELLIER AB, et al. (2014) The Arabidopsis nitrate transporter NRT2.5 plays a role in nitrate acquisition and remobilization in nitrogen-starved plants. *Plant J*, 80(2): 230-241. DOI: 10.1111/tbj.12626
- LI B, XIN W, SUN S, et al. (2006) Physiological and molecular responses of nitrogen-starved rice plants to re-supply of different nitrogen sources. *Plant Soil*, 287(1-2): 145-159. DOI 10.1007/s11104-006-9051-1
- LI JY, FU YL, PIKE SM, et al. (2010) The Arabidopsis nitrate transporter NRT1.8 functions in nitrate removal from the xylem sap and mediates cadmium tolerance. *Plant Cell*, 22(5): 1633-1646. DOI: 10.1105/tpc.110.075242
- LI W, WANG Y, OKAMOTO M, et al. (2007) Dissection of the AtNRT2.1: AtNRT2.2 inducible high-affinity nitrate transporter gene cluster. *Plant Physiol*, 143: 425-433. DOI: 10.1104/pp.106.091223
- LI Z, WU XN, JACQUOT A, et al. (2020) A phospho-switch in the N-terminus of NRT2.1 affects nitrate uptake by controlling the interaction of NRT2.1 with NAR2.1. *BioRxiv*. DOI: 10.1101/2020.01.08.898254
- LIN SH, KUO HF, CANIVENC G, et al. (2008) Mutation of the Arabidopsis NRT1.5 nitrate transporter causes defective root-to-shoot nitrate transport. *Plant Cell*, 20(9): 2514-2528. DOI: 10.1105/tpc.108.060244
- LITTLE DY, RAO H, OLIVA S, et al. (2005) The putative high-affinity nitrate transporter NRT2.1 represses lateral root initiation in response to nutritional cues. *Proc Natl Acad Sci USA*, 102: 13693-13698. DOI: 10.1073/pnas.0504219102
- LIVAK KJ and SCHMITTGEN TD (2001) Analysis of relative gene expression data using Real-Time Quantitative PCR and the $2^{-\Delta\Delta CT}$ method. *Methods*, 25(4): 402-408. DOI: 10.1006/meth.2001.1262
- LOQUÉ D, YUAN L, KOJIMA S, et al. (2006) Additive contribution of AMT1;1 and AMT1;3 to high-affinity ammonium uptake across the plasma membrane of nitrogen-deficient Arabidopsis roots. *Plant J*, 48(4): 522-534. DOI: 10.1111/j.1365-313X.2006.02887.x
- LOQUÉ D, LALONDE S, LOOGER LL, et al. (2007) A cytosolic trans-activation domain essential for ammonium uptake. *Nature*. 446: 195-198. DOI: 10.1038/nature05579
- MALAGOLI P, LAINÉ P, DEUNFF EL, et al. (2004) Modeling nitrogen uptake in oilseed rape cv Capitot during a growth cycle using influx kinetics of root nitrate transport systems and field experimental data. *Plant Physiol*, 134(1): 388-400. DOI: 10.1104/pp.103.029538
- MARCHIE C, ROUDIER F, CASTAINGS L, et al. (2013) Nuclear retention of the transcription factor NLP7 orchestrates the early response to nitrate in plants. *Nat Commun*, 4: 1713. DOI: 10.1038/ncomms2650.
- MENZ J, LI Z, SCHULZE WX, and LUDEWIG U (2016). Early nitrogen-deprivation responses in Arabidopsis roots reveal distinct differences on transcriptome and (phospho-) proteome levels between nitrate and ammonium nutrition. *Plant J*, 88: 717-734. DOI: 10.1111/tbj.13272
- MEYER J, SCHMANN A, WOOD R, et al. (2007) Recent advances to nitrogen use efficiency of sugarcane in the South African sugar industry. *Proc Int Soc Sugar Cane Technol*, 26: 238-246.
- MILLAR N, ROBERTSON GP, GRACE PR, et al. (2010) Nitrogen fertilizer management for nitrous oxide (N₂O) mitigation in intensive corn (Maize) production: an emissions reduction protocol for US Midwest agriculture. *Mitig Adapt Strat Gl*, 15: 185-204. DOI 10.1007/s11027-010-9212-7
- MILLER AJ and CRAMER MD (2005) Root nitrogen acquisition and assimilation. *Plant Soil*, 274: 1-36. DOI: 10.1007/s11104-004-0965-1
- MILLER AJ, FAN X, ORSEL M, et al. (2007) Nitrate transport and signalling. *J Ex Bot*, 58(9): 2297-2306. DOI: 10.1093/jxb/erm066
- MILLER SA (2010) Minimizing land use and nitrogen intensity of bioenergy. *Environ Sci Technol*, 44(10): 3932-3939. DOI: 10.1021/es902405a.
- MING R, LIU SC, LIN YR, et al. (1998) Detailed alignment of saccharum and sorghum chromosomes: comparative organization of closely related diploid and polyploid genomes. *Genetics*, 150(4): 1663-1682.
- MOLL RH, KAMPRATH EJ, and JACKSON WA (1982) Analysis and interpretation of factors which contribute to efficiency of nitrogen utilization. *Agron J*, 74: 562-564. DOI: 10.2134/agronj1982.00021962007400030037x
- MUELLER ND, GERBER JS, JOHNSTON M, et al. (2012) Closing yield gaps through nutrient and water management. *Nature*, 490: 254-257. DOI: 10.1038/nature11420

- MUNÕS S, CAZETTES C, FIZAMES C, et al. (2004) Transcript profiling in the chl1-5 mutant of *Arabidopsis* reveals a role of the Nitrate Transporter NRT1.1 in the regulation of another nitrate transporter, NRT2.1. *Plant Cell*, 16(3): 2433-2447. DOI: 10.1105/tpc.104.024380
- OKAMOTO M, VIDMAR JJ, and GLASS ADM (2003) Regulation of NRT1 and NRT2 gene families of *Arabidopsis thaliana*: responses to nitrate provision. *Plant Cell Physiol*, 44: 304-317. DOI: 10.1093/pcp/pcg036
- OKAMOTO M, KUMAR A, LI W, et al. (2006) High-affinity nitrate transport in roots of *Arabidopsis* depends on expression of the NAR2-like gene AtNRT3.1. *Plant Physiol*, 140: 1036-1046. DOI: 10.1104/pp.105.074385
- ORSEL M, CHOPIN F, LELEU O, et al. (2006) Characterization of a two-component high-affinity nitrate uptake system in *Arabidopsis*. Physiology and protein-protein interaction. *Plant Physiol*, 142(3): 1304-1317. DOI: 10.1104/pp.106.085209
- PELLIZZARO A, CLOCHARD T, PLANCHET E, et al. (2015) Identification and molecular characterization of *Medicago truncatula* NRT2 and NAR2 families. *Physiol Plant*, 154(2): 256-269. DOI: 10.1111/ppl.12314
- PLETT D, TOUBIA J, GARNETT T, et al. (2010) Dichotomy in the NRT Gene families of dicots and grass species. *PLoS One*, 5(12): e15289. DOI: 10.1371/journal.pone.0015289
- REAY DS, DAVIDSON EA, SMITH KA, et al. (2012) Global agriculture and nitrous oxide emissions. *Nat Clim Change*, 2: 410-416. DOI: 10.1038/NCLIMATE1458
- REMANS T, NACRY P, PERVENT M, et al. (2006) A central role for the nitrate transporter NRT2.1 in the integrated morphological and physiological responses of the root system to nitrogen limitation in *Arabidopsis*. *Plant Physiol*, 140: 909-921. DOI: 10.1104/pp.105.075721
- RIAÑO-PACHÓN D and MATTIELLO L (2017) Draft genome sequencing of the sugarcane hybrid SP80-3280. *F1000Res*, 6:861. DOI: 10.12688/f1000research.11859.2
- ROBINSON N, FLETCHER A, WHAN A, et al. (2007) Sugarcane genotypes differ in internal nitrogen use efficiency. *Funct Plant Biol*, 34(12): 1122-1129. DOI: 10.1071/FP07183
- ROBINSON N, BRACKIN R, VINALL K, et al. (2011) Nitrate Paradigm Does Not Hold Up for Sugarcane. *PLoS One*, 6(4): e19045. DOI: 10.1371/journal.pone.0019045
- ROCKSTRÖM J, STEFFEN W, NOONE K, et al. (2009) A safe operating space for humanity. *Nature*, 461(7263):472-475. DOI: 10.1038/461472a
- RUFFEL S, FREIXES S, BALZERGUE S, et al. (2008) Systemic signaling of the plant nitrogen status triggers specific transcriptome responses depending on the nitrogen source in *Medicago truncatula*. *Plant Physiol*, 146(4): 2020-2035. DOI: 10.1104/pp.107.115667
- SCHRÖDER JJ, SCHOLEFIELD D, CABRAL F, and HOFMAN G (2004) The effects of nutrient losses from agriculture on ground and surface water quality: the position of science in developing indicators for regulation. *Environ Sci Policy*, 7: 15-23. DOI: 10.1016/j.envsci.2003.10.006
- SEGONZAC C, BOYER JC, IPOTESI E, et al. (2007) Nitrate efflux at the root plasma membrane: identification of an *Arabidopsis* excretion transporter. *Plant Cell*, 19(11): 3760-3777. DOI: 10.1105/tpc.106.048173
- SEREZINO LHD (2015) Physiological and transcriptional characterization of nitrate acquisition and remobilization processes in sugarcane (*Saccharum spp.*) (Master's dissertation, CENA/USP, São Paulo, Brazil)
- SIDDIQI MY, GLASS ADM, RUTH TJ, and RUFTY TW (1990) Studies of the uptake of nitrate in barley. I. Kinetics of $^{13}\text{NO}_3^-$ influx. *Plant Physiol*, 93: 1426-1432. DOI: 10.1104/pp.93.4.1426
- SOMERVILLE C, YOUNGS H, TAYLOR C, et al. (2010) Feedstocks for lignocellulosic biofuels. *Science*, 329(5993):790-792. DOI: 10.1126/science.1189268
- SOUZA GM, VAN SLUYS MA, LEMBKE CG, et al. (2019) Assembly of the 373k gene space of the polyploid sugarcane genome reveals reservoirs of functional diversity in the world's leading biomass crop. *GigaScience*, 8(12): 1-18. DOI: 10.1093/gigascience/giz129
- TABATA R, SUMIDA K, YOSHII T, et al. (2014) Perception of root-derived peptides by shoot LRR-RKS mediates systemic N-demand signaling. *Science*, 346(6207): 343-346. DOI: 10.1126/science.1257800
- TASSO JÚNIOR LC (2007) Agrotechnological characterization of sugarcane (*Saccharum spp.*) cultivars in the north-central region of the state of São Paulo (Unpublished doctoral thesis, UNESP, São Paulo, Brazil)

- THORBURN PJ, BIGGS JS, COLLINS K, and PROBERT ME (2010) Using the APSIM model to estimate nitrous oxide emissions from diverse Australian sugarcane production systems. *Agric Ecosyst Environ*, 136: 343-350. DOI: 10.1016/j.agee.2009.12.014
- TILMAN D (1999) Global environmental impacts of agricultural expansion: The need for sustainable and efficient practices. *Proc Natl Acad Sci USA*, 96: 5995-6000. DOI: 10.1073/pnas.96.11.599
- TILMAN D, FARGIONE J, WOLFF B, et al. (2001) Forecasting agriculturally driven global environmental change. *Science*, 292(5515): 281-284. DOI: 10.1126/science.1057544
- TISCHNER R, WALDECK B, GOYAL SS, and RAINS WD (1993) Effect of nitrate pulses on the nitrate-uptake rate, synthesis of mRNA coding for nitrate reductase, and nitrate- reductase activity in the roots of barley seedlings. *Planta*, 189: 533-537. DOI: 10.1007/BF00198216
- TOMPKINS GA, JACKSON WA, and VOLK RJ (1978) Accelerated nitrate uptake in wheat seedlings: effects of ammonium and nitrite treatments and of six methyl purine and puromycin. *Physiol Plant*, 43: 166-171. DOI: 10.1111/j.1399-3054.1978.tb01586.x
- TONG Y, ZHOU JJ, LI Z, and MILLER AJ (2005) A two-component high-affinity nitrate uptake system in barley. *Plant J*, 41: 442-450. DOI: 10.1111/j.1365-313X.2004.02310.x
- TSAY YF, CHIU CC, TSAI CB, et al. (2007) Nitrate transporters and peptide transporters. *FEBS Lett*, 581(12): 2290-2300. DOI: 10.1016/j.febslet.2007.04.047
- UDOP - National Bioenergy Union. https://www.udop.com.br/index.php?item=variedades_iac. Accessed in June 16th, 2020.
- VAN DER WEIJDE T, KAMEI CLA, TORRES AF, et al. (2013) The potential of C₄ grasses for cellulosic biofuel production. *Front Plant Sci*, 4. DOI: 10.3389/fpls.2013.00107
- VIDMAR JJ, ZHUO D, SIDDIQI MY, et al. (2000) Regulation of HvNRT2 expression and high-affinity nitrate influx in roots of *Hordeum vulgare* by ammonium and amino acids. *Plant Physiol*, 123(1): 307-318. DOI: 10.1104/pp.123.1.307
- VIEIRA-MEGDA MX, MARIANO E, LEITE JM et al (2015) Contribution of fertilizer nitrogen to the total nitrogen extracted by sugarcane under Brazilian field conditions. *Nutr Cycl Agroecosystems*, 101: 241-257. DOI: 10.1007/s10705-015-9676-7
- VITOUSEK PM, NAYLOR R, CREWS T, et al. (2009) Agriculture nutrient imbalances in agricultural development. *Science*, 324: 1519-1520. DOI: 10.1126/science.1170261
- VON WITTGENSTEIN NJJB, LE CH, HAWKINS BJ, and EHLTING J (2014) Evolutionary classification of ammonium, nitrate, and peptide transporters in land plants. *BMC Evol Biol*, 14(3): 11. DOI: 10.1186/1471-2148-14-11
- WACLAWOVSKY AJ, SATO PM, LEMBKE CG, et al. (2010) Sugarcane for bioenergy production: an assessment of yield and regulation of sucrose content. *Plant Biotechnol J*, 8(3): 263-276. DOI: 10.1111/j.1467-7652.2009.00491
- WANG J, LI Y, ZHU F, et al. (2019a) Genome-Wide Analysis of Nitrate Transporter (NRT/NPF) Family in Sugarcane *Saccharum spontaneum* L. *Trop Plant Biol*, 12(3): 133-149. DOI: 10.1007/s1204
- WANG J, HÜNER N, and TIAN L (2019b) Identification and molecular characterization of the *Brachypodium distachyon* NRT2 family, with a major role of BdNRT2.1. *Physiol Plant*, 165(3): 498-510. DOI: 10.1111/pp.12716
- WANG MY, SIDDIQI MY, RUTH TJ, and GLASS ADM (1993a): Ammonium uptake by rice roots. (I. Fluxes and subcellular distribution of ¹³NH₄⁺). *Plant Physiol*, 103: 1249-1258. DOI: 10.1104/pp.103.4.1249
- WANG MY, SIDDIQI MY, RUTH TJ, and GLASS ADM (1993b): Ammonium uptake by rice roots. (II. Kinetics of ¹³NH₄⁺ across the plasmalemma). *Plant Physiol*, 103: 1259-1267. DOI: 10.1104/pp.103.4.1259
- WANG R, TISCHNER R, GUTIÉRREZ RA, et al. (2004) Genomic analysis of the nitrate response using a nitrate reductase-null mutant of Arabidopsis. *Plant Physiol*, 136(1): 2512-2522. DOI: 10.1104/pp.104.044610
- WANG X, BIAN Y, CHENG K, et al. (2013) A large-scale protein phosphorylation analysis reveals novel phosphorylation motifs and phosphoregulatory networks in Arabidopsis. *J Proteomics*, 14: 486-498. DOI: 10.1016/j.jprot.2012.10.018
- WANG YY and TSAY YF (2011) Arabidopsis nitrate transporter NRT1.9 is important in phloem nitrate transport. *Plant Cell*, 23(5): 1945-1957. DOI: 10.1105/tpc.111.083618
- WIRTH J, CHOPIN F, SANTONI V, et al. (2007). Regulation of root nitrate uptake at the NRT2.1 protein level in *Arabidopsis thaliana*. *J Biol Chem*, 282: 23541-23552. DOI: 10.1074/jbc.M700901200
- XU G, FAN X, and MILLER AJ (2012) Plant nitrogen assimilation and use efficiency. *Annu Rev Plant Biol*, 63: 153-182. DOI: 10.1146/annurev-arplant-042811-105532

- YAN M, FAN X, FENG H, et al. (2011) Rice OsNAR2.1 interacts with OsNRT2.1, OsNRT2.2 and OsNRT2.3a nitrate transporters to provide uptake over high and low concentration ranges. *Plant Cell Environ*, 34: 1360-1372. DOI: 10.1111/j.1365-3040.2011.02335.x
- YONG Z, KOTUR Z, and GLASS ADM (2010) Characterization of an intact two-component high- affinity nitrate transporter from *Arabidopsis* roots. *Plant J*, 63(5): 739-748. DOI: 10.1111/j.1365-313X.2010.04278.x
- ZHAO M, DING H, ZHU JK, et al. (2011) Involvement of miR169 in the nitrogen-starvation responses in *Arabidopsis*. *New Phytol*, 190(4): 906-915. DOI: 10.1111/j.1469-8137.2011.03647.x
- ZHUO D, OKAMOTO M, VIDMAR JJ, and GLASS ADM (1999) Regulation of a putative high-affinity nitrate transporter (Nrt2;1At) in roots of *Arabidopsis thaliana*. *Plant J*, 17: 563-568. DOI: 10.1046/j.1365-313X.1999.00396.x
- ZOU X, LIU MY, WU WH, and WANG Y (2019) Phosphorylation at Ser28 stabilizes the *Arabidopsis* nitrate transporter NRT2.1 in response to nitrate limitation. *J Integr Plant Bio*, 62(6): 865-876. DOI: 10.1111/jipb.12858