Universidade de São Paulo
Instituto de Astronomia, Geofísica e Ciências Atmosféricas
Departamento de Astronomia

Bruno Dias

Formation and evolution of globular clusters in the Galaxy and Magellanic Clouds

Formação e evolução de aglomerados globulares da Galáxia e Nuvens de Magalhães

São Paulo

2014

Bruno Dias

Formation and evolution of globular clusters in the Galaxy and Magellanic Clouds

Formação e evolução de aglomerados globulares da Galáxia e Nuvens de Magalhães

Thesis presented to Departamento de Astronomia of Instituto de Astronomia, Geofísica e Ciências Atmosféricas, Universidade de São Paulo, as partial requirement to get the title of PhD in Science.

Area: Astronomy

Advisor: Prof. Dr. Beatriz Barbay

Updated version. Original can be found at the Institute.

Versão Corrigida. O original encontra-se disponível na Unidade.

São Paulo

2014

To my family.

Acknowledgements

To my advisor Beatriz Barbuy for making me a professional astrophysicist;

To my internship advisor Ivo Saviane for receiving me at ESO/Santiago, and to Oscar Gonzalez for being my fellow mentor at ESO;

To the professor Silvia Rossi for helping me during the whole process;

To the professors Jorge Melendez and Walter Maciel for evaluating my PhD project;

To the astronomers Paula Coelho, Alan Alves Brito, Claudio Melo for many important discussions for my thesis and career;

To the astronomers Enrico Held, Ana Chies-Santos, Ricardo Schiavon, Leandro Kerber for inviting me as visitor astronomer in their institutes and for excellent discussions;

To the astronomers Sergio Ortolani and Dimitri Gadotti for teaching me *in situ* about NTT/ESO and FLAMES@VLT/ESO, respectively

To the professor Dante Minniti for accepting me as member of the VVV survey;

To the colleagues Marina Trevisan, Cesar Siqueira Mello, Marcelo Tucci for sharing the office, discussions and coffees with me;

To the colleagues Phillip, Eduardo, Beatriz, Thais for staying together for many years;

To the ESO colleagues Anna, Julien, Jos, Amy, Sergio, Matias, Florian, Joachim, Daniela, Paul, Marylou, Vera, Mirjam.

To CNPq by the financial support under project 142047/2010-4;

To ESO by the award of a studentship for one year in Santiago/Chile.

To my family Alvino, Nilda, Gabriel, Paula for supporting me all the time;

To my friends for making my life easier during the last four years.

“One’s ideas must be as broad as Nature if they are to interpret Nature”

Sherlock Holmes on “A Study in Scarlet”. Sir Arthur Conan Doyle (1887)

“I suppose it is clear to everyone that astronomy at all events compels the soul to look upwards, and draws it from the things of this world to the other”

Glaucón on the Republic, Book Seven. Plato (BC 427-BC 347)

Resumo

Aglomerados globulares são traçadores da formação e evolução de suas galáxias. Cinemática, abundâncias químicas, idades e posições dos aglomerados permitem traçar interações entre Via Láctea e galáxias vizinhas e suas histórias de enriquecimento químico. Nesta tese analisamos espectros de média resolução de mais de 800 estrelas gigantes vermelhas em 51 aglomerados globulares Galácticos. É a primeira vez que $[\text{Fe}/\text{H}]$ and $[\text{Mg}/\text{Fe}]$ determinados de modo consistente são publicados para uma amostra desse porte, $\sim 1/3$ dos objetos catalogados. Nossas metalicidades são mais precisas que trabalhos anteriores similares. Uma quebra em $[\text{Fe}/\text{H}] \sim -1.0$ é encontrada no gráfico $[\text{Fe}/\text{H}]$ vs. $[\text{Mg}/\text{Fe}]$ para o bojo e halo, embora bojo parece ter uma quebra em $[\text{Fe}/\text{H}]$ maior, i.e, bojo tem formação mais eficiente que o halo. Comparando abundâncias com idade, a escala de tempo para SNIa ficar importante é 1 Gano. $[\text{Fe}/\text{H}]$ vs. idade corrobora diferentes eficiências de formação do bojo e halo, mas $[\text{Mg}/\text{Fe}]$ vs. idade não mostra isso. O halo foi formado em mini halos ou galáxias anãs, e dois aglomerados com dispersão em $[\text{Fe}/\text{H}]$ tiveram suas origens analisadas. M 22 parece ter sido formado na Via Láctea e NGC 5824 possivelmente foi originado em uma galáxia anã, embora os resultados são inconclusivos para NGC 5824. O bojo parece ter sido formado rapidamente e deve possuir o aglomerado mais velho. De fato, HP 1 tem um ramo horizontal mais azul que o esperado para sua metalicidade e vemos isso como um efeito da idade. Determinamos sua distância usando curvas de luz de RR Lyrae de maneira a restringir futuras determinações de idade via diagrama cor-magnitude. Finalmente, investigamos a interação entre Via Láctea e sua galáxia vizinha SMC. Encontramos aglomerados sendo removidos do corpo central da SMC, consistente com cenário de remoção por força de maré para a interação entre as galáxias, em vez de “ram pressure” que afeta só gás.

Abstract

Globular clusters are tracers of the formation and evolution of their host galaxies. Kinematics, chemical abundances, age and position of the clusters allows tracing interactions between Milky Way and surrounding galaxies and outlines their chemical enrichment history. In this thesis we analyse mid-resolution spectra of about 800 red giant stars in 51 Galactic globular clusters. It is the first time that $[\text{Fe}/\text{H}]$ and $[\text{Mg}/\text{Fe}]$ derived in a consistent way are published for such a huge sample of globular clusters, almost 1/3 of the total number of catalogued clusters. Our metallicities are showed to be more precise than previous works based on mid-resolution spectroscopy. A turnover at $[\text{Fe}/\text{H}] \sim -1.0$ is found in the plot $[\text{Fe}/\text{H}]$ vs. $[\text{Mg}/\text{Fe}]$ for bulge and halo, although bulge seems to have a more metal-rich turnover, i.e, bulge has more efficient formation than the halo. Comparing the abundances with age the timescale for SNIa to start to become important is 1Gyr. $[\text{Fe}/\text{H}]$ vs. age corroborates the different star formation efficiency of bulge and halo while $[\text{Mg}/\text{Fe}]$ does not follow that. Halo was formed in mini halos or dwarf galaxies, and two multiple population clusters had their origin analysed to check it. M 22 seems to have been formed in the Milky Way while NGC 5824 possibly was originated in a dwarf galaxy, although our results are inconclusive for NGC 5824. The Galactic bulge seems to have been formed fast i.e., probably the oldest globular cluster is there. In fact HP 1 has a bluer horizontal branch than expected for its metallicity and we interpret that as an age effect. We determine its distance using light curves of variable stars in order to constrain future age determinations via colour-magnitude diagram. Finally, we investigate interaction between Milky Way and its neighbour galaxy SMC. We find that some star clusters are being stripped out of the SMC main body, which is consistent with tidal stripping scenario for the interaction between the galaxies, instead of ram pressure that would only affect gas.

List of Figures

1.1	Schematic view of the X-shaped bulge of the Milky Way.	26
1.2	Magellanic Clouds and Magellanic Stream in terms of H I	28
2.1	Sky maps of the six SMC clusters	40
2.2	Completeness curves for the six SMC clusters	41
2.3	V and (B-V) photometric errors for the six SMC clusters.	42
2.4	V, (B-V) colour-magnitude diagrams for all clusters and control fields . . .	42
2.5	Radial density profiles fitted by King profiles for all clusters	44
2.6	Process of generation of a synthetic V, (B-V) CMD	47
2.7	Best isochrone fittings for all clusters	49
2.8	Distribution of star clusters in the projected plane of the SMC	53
2.9	Age-metallicity relation for the SMC	54
2.10	Age and [Fe/H] as a function of the distance to the centre of SMC	55
3.1	Colour-Magnitude diagrams of the six Milky Way globular clusters	62
3.2	Sky maps of the six Milky Way globular clusters	63
3.3	Identificantion of most important atomic lines in each strong feature of R~2 000 metal-poor and metal-rich stellar spectra	64
3.4	Heliocentric radial velocities for all clusters	65
3.5	Examples of spectral fitting for a metal-rich and a metal-poor star	67
3.6	Parameter space of MILES library displayed in a HR diagram	68
3.7	Procedure to find best T_{eff} , $\log(g)$ and [Fe/H] for star HD122956	72
3.8	Comparisons of T_{eff} , $\log(g)$, [Fe/H] from this work with literature for well- known field stars	73

3.9	Comparison of derived T_{eff} and $\log(g)$ with isochrones for member stars of the six globular clusters	74
3.10	Comparison of derived $[\text{Fe}/\text{H}]$ with CaT results from literature for the stars from all clusters	75
3.11	Procedure for membership selection of stars in each globular cluster	77
3.12	HR diagram for Terzan 8 comparing T_{eff} , $\log(g)$ from this work with high-resolution results	81
3.13	Comparison of $[\text{Fe}/\text{H}]$ from this work with the literature	82
3.14	$[\text{Mg}/\text{Fe}]$ vs. $[\text{Fe}/\text{H}]$ of Milky Way field stars overplotted by the average results for each cluster from this cluster	83
4.1	Sky position of the 51 clusters over Milky Way image	91
4.2	Reddening versus distance of the 51 globular clusters	92
4.3	Comparison of $[\text{Fe}/\text{H}]$ with results of Carretta's scale for 13 clusters in common.	97
4.4	Comparison of calibrated $[\text{Fe}/\text{H}]$ with those from Carretta et al. (2009) for the 51 clusters	101
4.5	Chemical evolution of the Milky Way in terms of $[\text{Mg}/\text{Fe}]$ vs. $[\text{Fe}/\text{H}]$	103
4.6	Age vs. $[\text{Fe}/\text{H}]$ and $[\text{Mg}/\text{Fe}]$ for 17 clusters	104
4.7	Metallicity radial gradients for Milky Way globular clusters	105
5.1	Magnitude vs. radius for GCs and UCDs	109
5.2	Colour-Magnitude diagrams of two globular clusters with multiple stellar populations	110
5.3	Sky map of two globular clusters with multiple stellar populations	111
5.4	Comparison of derived T_{eff} and $\log(g)$ with isochrones for member stars of the six globular clusters	112
5.5	Procedure for membership selection of stars in M 22 and NGC 5824	113
5.6	$[\text{Fe}/\text{H}]$ and $[\text{Mg}/\text{H}]$ distribution function of M 22	115
5.7	Comparison of sample stars from this work with Marino et al. (2011) for M 22	116
5.8	Chemical evolution history of M 22	117
5.9	Metallicity distribution function of NGC 5824	118

5.10	Chemical evolution history of NGC 5824	119
6.1	Colour-magnitude diagram of HP 1 with the six RR Lyrae candidates . . .	123
6.2	Finding chart of six RR Lyrae in HP 1	123
6.3	Light curves of five confirmed RR Lyrae in HP 1	124
6.4	Periodogram θ vs. period for the five confirmed RR Lyrae in HP 1	125
6.5	Folded light curves for five RR Lyrae candidates in HP1, using the best periods found in Figure 6.4.	126
6.6	Period-luminosity relation applied to calculate the distance	127
A.1	Likelihood from the CMD fits for different combinations of age, metallicity, reddening, and distance modulus for AM 3.	158
A.2	Same as Fig. A.1 for HW 1.	158
A.3	Same as Fig. A.1 for HW 40.	159
A.4	Same as Fig. A.1 for Lindsay 2.	159
A.5	Same as Fig. A.1 for Lindsay 3.	159
C.1	Output two dimensional spectra of FORS2 pipeline using default parameters	210
C.2	Default list of skylines from FORS2 manual	211
C.3	New list of skylines proposed in this work	212
C.4	Output two dimensional spectra of FORS2 pipeline using the new approach	213
C.5	Comparison between extracted spectrum from slit #8 using default para- meters with that using the new procedure	214
C.6	Same of Figure C.5, but for the slit #6.	215
C.7	Comparison between extracted spectra from slit #8 with that from slit #6 using default parameters and the new procedure	216
D.1	Abstract of Borissova et al. (2011).	217
D.2	Abstract of Saito et al. (2012).	218
D.3	Abstract of Saito et al. (2012).	219
D.4	Abstract of Dias et al. (2014).	220
D.5	Abstract of Mauro et al. (2014).	221
D.6	Abstract of Dias et al. (2014b), submitted to A&A and accepted after revision.	222

List of Tables

2.1	Log of observations of the six SMC clusters	38
2.2	Colour coefficient and photometric zero point from standard stars calibration	39
2.3	Structural parameters derived from King profile fitting for the six SMC clusters	44
2.4	Stellar counts in the cluster and control field CMDs of the six clusters . . .	46
2.5	Age, metallicity, distance and reddening derived for the SMC clusters . . .	49
3.1	FORS2 spectrograph details	60
3.2	Log of observations and parameters from literature for the six globular clusters	61
3.5	Final average parameters for each globular cluster and respective internal errors	78
3.7	Atmospheric parameters for the three stars of M 71 in common with Cohen et al. (2001)	79
3.8	Atmospheric parameters for the three stars of NGC 6558 in common with Barbuy et al. (2007)	80
3.9	Atmospheric parameters for the three stars of Terzan 8 in common with Carretta et al. (2014)	80
4.1	Log of observations of the 51 globular clusters	93
4.2	[Fe/H] and radial velocities for the 13 globular clusters in common with Carretta et al. (2009)	95
4.3	Final average parameters for the 51 clusters [Fe/H], [Mg/Fe], [α /Fe] and v_{helio}	99
5.1	Final average parameters for M 22 and NGC 5824 and respective internal errors	113

B.1	Star by star coordinates, magnitude, colour, heliocentric radial velocity for the six globular clusters	161
B.2	Atmospheric parameters of the 49 well known stars used to validate the method	179
B.3	Atmospheric parameters for all 758 stars analysed in the six clusters: T_{eff} , $\log(g)$, $[Fe/H]$, $[Mg/Fe]$ and $[\alpha/Fe]$	181

Contents

1. <i>Introduction</i>	23
1.1 Formation and evolution of galaxies	23
1.1.1 Milky Way bulge: monolithic collapse or secular evolution?	24
1.1.2 Milky Way halo: minor mergers with surrounding galaxies	26
1.1.3 Magellanic Clouds: tidal interactions in the system MW-SMC-LMC	28
1.2 Summary and goals of this thesis	31
1.2.1 Milky Way - Magellanic Clouds interactions	31
1.2.2 Formation and evolution of Galactic bulge and halo	32
2. <i>Small Magellanic Cloud</i>	35
2.1 Abstract of this chapter	35
2.2 Introduction to this chapter	36
2.3 Observations	37
2.3.1 The SOAR/SOI data	37
2.3.2 Photometric errors and completeness curves	39
2.3.3 The radial density profiles	43
2.3.4 The observed CMDs and cluster membership probability	43
2.4 Statistical isochrone fitting	45
2.4.1 CMD modelling	46
2.4.2 Likelihood statistics	48
2.5 Results	48
2.5.1 AM 3	50
2.5.2 HW 1 and Lindsay 2	50

2.5.3	HW 40	51
2.5.4	Lindsay 3	51
2.5.5	HW 34	51
2.6	The age-metallicity relation and spatial distribution	52
2.7	Summary of this chapter	55
3.	<i>FORS2/VLT survey of Milky Way globular clusters. I. Method validation</i>	57
3.1	Abstract of this chapter	57
3.2	Introduction to this chapter	58
3.3	Observations and data reduction	59
3.4	Stellar parameters determination	64
3.4.1	Radial velocities	64
3.4.2	Atmospheric parameters	66
3.4.2.1	Stellar libraries	68
3.4.2.2	Average results and errors: validation with well-known stars	69
3.5	Results	72
3.5.1	T_{eff} , $\log(g)$ against isochrones	73
3.5.2	Comparison with $[\text{Fe}/\text{H}]$ derived from CaT	75
3.5.3	Membership selection	76
3.5.4	Validation with high-resolution spectroscopy	78
3.5.4.1	M 71	79
3.5.4.2	NGC 6558	79
3.5.4.3	Terzan 8	80
3.6	Discussion	81
3.6.1	Metal-rich clusters NGC 6528, NGC 6553	82
3.6.2	Moderately metal-rich clusters M 71, NGC 6558	84
3.6.3	Metal-poor clusters NGC 6426, Terzan 8	85
3.7	Summary and conclusions	86
4.	<i>FORS2/VLT survey of Milky Way globular clusters. II. Parameters for 51 clusters</i>	89
4.1	Introduction to this chapter	89
4.2	Targets selection and observations	90
4.3	Method - summary from Chapter 3	94

4.4	Calibration of results	95
4.4.1	[Fe/H] in Carretta scale	95
4.4.2	Calibration of [Mg/Fe] results	96
4.5	Discussion on abundances results	98
4.5.1	Chemical evolution of the Milky Way	102
4.5.2	Origin of globular clusters	103
4.6	Summary and conclusions	105
5.	<i>FORS2/VLT survey of Milky Way globular clusters. III. Clusters with multiple populations</i>	107
5.1	Introduction to this chapter	107
5.2	Observations and analysis	110
5.3	Results	111
5.3.1	T_{eff} , $\log(g)$ against isochrones	111
5.3.2	Membership selection	112
5.4	[Fe/H] and [Mg/H] intrinsic spread of M 22 and NGC 5824	113
5.4.1	M 22	114
5.4.2	NGC 5824	118
5.5	Summary and conclusions	120
6.	<i>RR Lyrae variable stars in HP1</i>	121
6.1	Introduction to this chapter	121
6.2	Observations	122
6.3	RR Lyrae candidates	123
6.4	Period determination	124
6.5	Constraining the distance	124
6.6	Conclusions	127
7.	<i>Conclusions and future work</i>	129
7.1	Summary	129
7.2	Perspectives	132
	<i>Bibliography</i>	135

<i>Appendix</i>	155
<i>A. SMC clusters CMD statistical analysis</i>	157
A.1 Likelihood results and degeneracy investigation	157
<i>B. Tables with results for FORS2 spectra</i>	161
<i>C. FORS2 pipeline - wavelength calibration for low resolution spectra of GRIS1400V</i>	209
C.1 Default pipeline reduction	209
C.2 New list of sky lines for GRIS1400V	211
C.3 Comparisons bewteen the 2 procedures	212
<i>D. Papers in the period 2010-2014</i>	217

Introduction

1.1 Formation and evolution of galaxies

A major question to understand the large scale structure of the Universe presently observed is: how did galaxies form and evolve? There are basically two ways to study galaxy formation and evolution, one way is to observe galaxies in different redshifts, i.e, in different evolutionary phases and reconstruct a common history. Another possibility is to observe nearby galaxies in detail looking at different stellar populations covering ages from the beginning of the galaxy until today. In this second case, it is possible to recover the star formation history of the host galaxy and also detect traces of interactions with nearby galaxies. Such information can be compared with scenarios of galaxy formation and evolution and provide constraints that are only possible for close objects, such as metallicity of individual stars. Globular clusters are perfect targets for this purpose, because it is relatively simple to derive their ages and place them in a timeline of the evolution of their host galaxy. In this thesis we adopt the second approach and analyse globular clusters in the Milky Way to explore its formation and evolution. We also study some stellar clusters in the Small Magellanic Cloud since it is suffering gravitational influence of the Milky Way. We think that this set of objects covers almost all stages of formation and evolution of galaxies and can be used as templates for models. In the following sections we detail the cases of the Milky Way, Large and Small Magellanic Clouds focusing on the role of star clusters as tracers of galaxy formation and evolution.

1.1.1 Milky Way bulge: monolithic collapse or secular evolution?

In the two last decades there were important discussions about the nature of galactic bulges. Wyse et al. (1997) pointed out the unclearness of the definitions of the nucleus of galaxies, “namely a centrally-concentrated stellar distribution with an amorphous - smooth - appearance.”. The classical scenario of monolithic collapse (Eggen et al., 1962) was proposed based on observation of 221 old dwarf stars in the Milky Way. The authors found that more metal-poor stars have highly eccentric orbits, and more metal-rich stars have nearly circular orbits, which was interpreted as the scenario where the galaxy started to collapse in a radial direction more than 10 Gyr ago, and the first stars were formed in this process (more metal-poor population), whereas less than 1 Gyr later the gas found an equilibrium in circular orbits, starting the formation of the more metal-rich stellar population. Bulge and halo are the spheroidal galactic components formed first, and the disc is the flat component formed later. This scenario was extended to all galaxies, but the best way to prove it is observing spatially resolved stars in the different components of our own Galaxy.

Milky Way bulge and halo stars are expected to be old and metal-poor following the scenario of Eggen et al. (1962). Concerning the ages, Lee (1992) observed RR Lyrae in the bulge (typical of old stellar populations), measured their metallicities and compare with horizontal branch models to conclude that there is an age gradient in the inner regions of the Milky Way, indicating that the oldest objects of the Galaxy are located in the bulge. Dékány et al. (2013) have shown that the bulge RR Lyrae form a spheroid. Stars in this region are more metal-rich than halo stars with $[\text{Fe}/\text{H}] \approx -1.0$, though. Following the logic of the monolithic collapse, it would be expected that the central region of the spheroidal component of the Milky Way (with a higher density of gas) would have formed stars in a more efficient way than the halo. However, bulge stellar content has a prolonged and bimodal (if not multimodal) metallicity distribution from $[\text{Fe}/\text{H}] \sim -1.0$ to $\sim +0.5$ (e.g. McWilliam and Rich, 1994; Sadler et al., 1996; Hill et al., 2011). This is not expected by simply collapsing a huge gas cloud about 14 Gyr ago. Moreover, Morrison (1993) debated about bulge and halo structures in terms of their density profiles. They showed that the power law density of the halo $\rho(r) \propto r^{-3.2}$ disagrees with the extension up to solar neighbourhood of the only surface brightness profile that fits the bulge (de Vaucouleurs and

Pence, 1978). As a result, bulge and halo may have been formed by different processes, furthermore the bulge has a complex history with probably more than one scenario of formation. Recent models describe the collapsing scenario for the bulge formation as hierarchical merging of subclumps (Noguchi, 1999; Aguerri et al., 2001).

Many spiral galaxies are barred, as already noted by Hubble (1926) in his original morphological classification of these “nebulae”, as they were called in that epoch. Bars can transport gas into the inner regions of the galaxy and trigger star formation in a timescale of a few billion years. Many barred late-type galaxies have their central stellar populations formed only by the secular evolution driven by their bars, and these inner regions are called pseudobulges (see Kormendy and Kennicutt, 2004; Gadotti, 2009). The Milky Way is a barred spiral galaxy and its bulge can be considered as a combination of both a classical (merger driven) and a pseudobulge (bar driven). In fact, Hill et al. (2011) present a bimodal metallicity distribution of bulge red giant stars in the Baade’s window - $(l, b) = (1^\circ, -4^\circ)$ - and discuss possible models to explain each peak. The best scenario that fitted both peaks is a combination of a closed-box model to explain the more metal-poor peak ($[\text{Fe}/\text{H}] \approx -0.3$) and an extension of the inner disc stellar population to describe the more metal-rich peak ($[\text{Fe}/\text{H}] \approx +0.3$).

Kinematics of a stellar population formed by a gravitational collapse would have a higher velocity dispersion, while a bar-driven population would be supported by rotation. Babusiaux et al. (2010) confirmed that based on $[\text{Fe}/\text{H}]$, radial velocities and proper motion of red giant stars in the Baade’s window compared with N-body simulations. Their findings are in agreement with Hill et al. (2011) as discussed above. Recently, it was found that the Milky Way bulge has an X-shape, and many works are devoted to describe its kinematics and chemistry from observations and models, e.g. Gardner et al. (2014); Zoccali et al. (2014); Vásquez et al. (2013); Gonzalez et al. (2013); Saito et al. (2012). In particular, Saito et al. (2011) have mapped the X-shaped bulge using its stellar contents. Using their findings it is possible to sketch the position of the X whose plane is parallel to the Galactic bar (Figure 1.1).

As firstly described by Baade (1944), stellar populations in galaxies can be classified in two types: population I constituted by luminous OB stars and open clusters, and population II represented by short-period Cepheids and globular clusters. Early-type galaxies have predominantly population II, while late-type galaxies share both populations, as in

Figure 1.1: Schematic view of the X-shaped bulge of the Milky Way in transparent blue over an ESO/S. Brunier image. Approximate position of Baade's window is indicated in red.

the case of the Milky Way. In our Galaxy, globular clusters are located in the spheroidal components, bulge and halo, and hence they are relics to probe the nature of the bulge. The amount and density of gas to form a globular cluster which has typical mass of $\sim 10^5 M_\odot$ is only found in the huge initial gravitational collapse that originated the Milky Way. In this case, globular clusters can be used as fossils of Galactic formation, and be compared with the more metal-poor population of the bulge. Barbuy et al. (1998) studied 15 bulge clusters and revised their metallicities, showing that the metallicity distribution is following the more metal-poor population of the bulge, as expected.

There is one interesting object located in the bulge that seems like a globular cluster, but it seems to be a pristine fragment of the bulge: Terzan 5. Ferraro et al. (2009) found two populations with different ages from photometry, Origlia et al. (2011) confirmed the two populations spectroscopically with $[\text{Fe}/\text{H}] \approx -0.3$ and $+0.3$, which corresponds to the two peaks of the metallicity distribution of bulge stars. Origlia et al. (2013) found three stars with metallicity around $[\text{Fe}/\text{H}] \approx -0.79$, and Massari et al. (2014) found other three stars with metallicities around $[\text{Fe}/\text{H}] \approx +0.7$. They also analysed the alpha-elements for these stars and concluded that Terzan 5 is probably one of the subclumps that were merged to form the bulge.

1.1.2 Milky Way halo: minor mergers with surrounding galaxies

Globular clusters (GCs) are tracers of the star formation history of their host galaxy, since most of the stars are formed in globular clusters, but only few of them keep this structure (Gieles et al., 2007). They also give constraints to galaxy formation scenarios, for instance the monolithic collapse (Eggen et al., 1962) predicts old and metal-poor globular clusters in the halo of the host galaxy. Disrupted dwarf galaxies could help forming the halo stellar populations, and transfer their globular clusters and/or nuclei to the Milky Way (Bekki and Freeman, 2003). This process could be the origin of multiple stellar populations found in few massive halo globular clusters. Therefore, globular clusters are useful to understand the chemical and dynamical evolution of the host galaxy. Another interesting feature of globular cluster systems in galaxies, and maybe one of the most important from the last 2 decades is their bimodal metallicity distribution, that reflects the chemical evolution of the galaxy, star formation history, and it constrains galaxy formation scenarios. There are 3 main scenarios to explain this bimodality: major disk-disk mergers (Zepf & Ashman 1992), *in situ* formation through multiphase dissipational collapse (Forbes 1997), and dissipationless accretion (Cote et al. 1998, 2000, 2002).

In order to explore all the cases above, it is necessary to have large samples of spectroscopic and photometric data, with homogeneous determinations of ages, metallicities, some specific chemical abundances, and also the knowledge of the kinematics of the clusters. In particular $[\alpha/\text{Fe}]$ ratios against $[\text{Fe}/\text{H}]$ are good tracers of star formation history, because it reflects the ratio and timing of supernovae type II and type Ia (e.g. Brodie & Strader 2004). Therefore it is useful to study Milky Way and Local Group globular cluster systems from which one can get more detailed information, as well as in external galaxies. The average $[\text{Fe}/\text{H}]$, and $[\text{Mg}/\text{Fe}]$ for all stars within a same cluster can be placed in the Milky Way star formation history scenario and constrain the timescale of the supernovae II and supernovae Ia, which can be also compared to age determinations for these clusters.

Stars in distant galaxies (out of Local Group) are not spatially resolved by 8-10m class telescopes, therefore only integrated spectra of globular clusters can be observed. In order to compare the Milky Way metallicity and $[\text{Mg}/\text{Fe}]$ to extragalactic globular cluster systems we use integrated light spectroscopy from extragalactic globular clusters (Trager et al. 2000, Thomas et al. 2003, Schiavon et al. 2005). This comparison is important in the

sense that individual stars studies allow to proceed with membership selection, and exclude stars that can introduce bias in the $[\text{Fe}/\text{H}]$ and $[\text{Mg}/\text{Fe}]$ as derived by integrated spectra. Although elliptical galaxies present a different scenario of star formation, and different pattern of $[\text{Mg}/\text{Fe}]$ vs. $[\text{Fe}/\text{H}]$, the calibration described for Milky Way globular clusters are very useful to understand extragalactic globular clusters. Note that extragalactic clusters reach high metallicities, which are not covered by previous metallicity scales for Milky Way GCs, and that the scale of this work covers all metallicities between $[\text{Fe}/\text{H}] = -2.5$ dex up to solar.

1.1.3 Magellanic Clouds: tidal interactions in the system MW-SMC-LMC

Evolution of galaxies depends on environmental effects, such as tidal stripping, that occurs when a given galaxy is subject to differential gravitational forces due to a companion and/or host galaxy. The gas of the galaxy is withdrawn and in a shorter timescale the stellar content is also removed from their original positions (Broeils and van Woerden, 1994). Another common process is the ram-pressure stripping (Gunn and Gott, 1972), where the gas of an interacting galaxy is lost because of the pressure of a hot gas in the intergalactic medium.

The Magellanic Stream (MS), discovered and named 40 years ago by Mathewson et al. (1974), is the trailing filament of neutral hydrogen of the Large and Small Magellanic Clouds (LMC and SMC) system. It is $\sim 200^\circ$ long and only $\sim 10^\circ$ wide, located close to the great circle that passes by the southern Galactic pole and crosses the Galactic plane at a longitude $l \sim 306^\circ$ (Nidever et al., 2010). Mathewson et al. (1974) argued that it is unlikely that the Magellanic Clouds had been closer than 40 kpc from the Galactic centre, due to the smooth velocity gradient of the gas connecting the Clouds, forming a bridge between them, and of both galaxies with the Stream. They also found evidence of a leading component of the gas stream. Figure 1.2 shows the map of Mathewson et al. (1974) and the new one of Nidever et al. (2010), 36 years later.

H I Parkes All-Sky Survey (HIPASS) team has confirmed and characterized the so-called Leading Arm of the Magellanic system, endorsing the scenarios of interaction between Milky Way and Magellanic Clouds (Putman et al., 1998). They summarize the most probable explanations for the origin of these gas structures in two main scenarios: ram-pressure (Mathewson et al., 1987; Barnes, 1996; Sofue, 1994; Moore and Davis, 1994) or

Figure 1.2: Aitoff projection of the Milky Way plane overplotted by the Magellanic System in terms of H I. Milky Way image with pink H I maps from Nidever et al. (2010) are overplotted with the blue H I maps and galactic coordinates grid from Mathewson et al. (1974).

tidal stripping (Murai and Fujimoto, 1980; Gardiner and Noguchi, 1996; Gardiner et al., 1994; Lin et al., 1995; Davies and Wright, 1977; Lin and Lynden-Bell, 1977). The first one explains the gas content of the MS, but to explain the presence of both the Leading Arm and the MS the model has to degrade the quality of the velocity profile of the Stream (Sofue, 1994; Moore and Davis, 1994). Ram-pressure is only valid for gas, not for stars. On the other hand, tidal stripping scenario predicts the presence of both Magellanic Stream (detected by Mathewson et al., 1974), a Leading Arm (confirmed by Putman et al., 1998) and the presence of a stellar counterpart.

On the one hand Nidever et al. (2008) traced back the velocity of the MS and LA, and showed that they originated in the LMC, not in the SMC or in the bridge. The authors suggest that a blowout of the LMC gas followed by a stretch of this expelled gas by the Milky Way potential can explain the formation of both gas structures, as an alternative scenario for the two most popular ones (ram-pressure and tidal stripping). On the other hand Diaz and Bekki (2012) realized N-body simulations to reproduce the morphology and kinematics of the MS, and proposed that LMC and SMC were separated galaxies until 2 Gyr ago. This first encounter and the following ones generated many tidal gas structures considering gravitational interactions with Milky Way: MS, LA, Bridge (between LMC and SMC) and a structure that they called Counter-bridge, which lies behind the SMC along the line-of-sight. The authors predict the presence of a stellar counterpart of the gas structures to support their scenario. Besla et al. (2013) also foresee stars in the Magellanic

Stream.

Nidever et al. (2013) found a stellar component of the Magellanic Bridge coming from the SMC and called it the “SMC eastern stellar structure”. The stars are about 1-2 Gyr old and the observations are qualitatively consistent with Diaz and Bekki (2012) models. Noël et al. (2013) found intermediate-age stars in the Magellanic Bridge, close to the SMC, and detect much less stars with similar ages in the opposite direction, on the Western borders of SMC. This is evidence supporting the idea of the intermediate-age stars present in the Bridge had been originated in the SMC. Casetti-Dinescu et al. (2014) went further and detect star formation in the Leading Arm, triggered by the strong interactions of Milky Way and Magellanic system.

With the advent of many surveys targeting the stellar content of the Magellanic System, such as VMC (Cioni et al., 2011), MAGIC (Noël et al., 2013) and SMASH (Olsen et al., 2014), stellar counterparts of the gas structures have been detected for the Bridge and are being observed in Magellanic Stream and Leading Arm. Together with the difficulty to model a Leading Arm from ram-pressure effects, these evidences tend to rule out this scenario as the origin of LA, MS and other structures of the Magellanic System. Therefore, the most accepted framework for the complex structure of this system is tidal stripping, or at least a combination of this effect with others, such as gas blowout from the LMC.

As mentioned by Noël et al. (2013), field stars are telling something about the chemodynamical evolution of the Magellanic Clouds by comparisons of stars in the Clouds line-of-sight against stars stripped towards the Magellanic Bridge, for instance. What about the stellar clusters? It is not broadly accepted that the stellar populations from field star and stellar cluster should have the same star formation history (SFH). Holtzman et al. (1999) says that the SFH of LMC field stars is different from that derived from clusters. In fact, more than 25 years ago Jensen et al. (1988) proposed that the LMC cluster population has an age gap between ~ 4 and ~ 10 Gyr, and up to now no object was found with ages in this range (Balbinot et al., 2010). On the contrary, LMC field star populations show no dependency on the location in the galaxy, and none of those small age gaps coincide with the cluster age gap (Javiel et al., 2005). One possibility is that the 4-10 Gyr old clusters have been formed in the LMC and were tidally stripped from the galaxy. If this is true, where are these clusters? Were they destroyed? If they exist, they could be found in one of the tidal structures of the Magellanic System, or even in the SMC.

Rafelski and Zaritsky (2005), Piatti et al. (2005) and Piatti (2012b) showed that for the SMC the SFH from field and cluster populations agree. These works have shown a recent burst of star formation, which could have been triggered by close encounters of LMC and SMC up to ~ 2 Gyr ago, as described by the dynamical models of Diaz and Bekki (2012), being at variance with continuous star formation suggested by Da Costa and Hatzidimitriou (1998). However, there was also an older burst in the SMC at about 5-7 Gyr, equally described by Cignoni et al. (2013) that is not predicted in the dynamical model of Diaz and Bekki (2012). A major merger with mass ratio between 1:1 to 1:4 could have generated this peak in the SMC SFH 5-7 Gyr ago, as proposed by Tsujimoto and Bekki (2009). However this model predicts a dip in the age-metallicity relation (AMR) correlated with this major merger, which is not found by the observational results of Cignoni et al. (2013). The AMR for SMC star clusters presents a broad distribution of metallicities at a given age, as shown by Parisi et al. (2014) and it is not possible to see a dip in metallicity, but in average the metallicity increases monotonically with time, in agreement with the findings of Cignoni et al. (2013) for field stars. The scatter of metallicity values for clusters does not permit a choice for a unique chemical evolution model (Parisi et al., 2014). Is it possible that some of this variety of metallicity values, in particular for intermediate-age clusters, are due to accreted clusters from LMC? SMC has a more complex star formation and chemo-dynamical evolution histories, consequently, is that valid to analyse the galaxy as a whole, or is it worth analysing region by region of the Small Cloud? Bica et al. (2008) catalogued over 3700 stellar systems in the LMC, SMC and Magellanic Bridge, and some Bridge clusters are being observed to look for intermediate-age objects. Many works have been studying areas of the SMC in terms of ages, the next step towards a complete scenario of the evolution of this galaxy is deriving metallicities.

1.2 Summary and goals of this thesis

In this thesis we use intermediate-age and old star clusters in the Small Magellanic Cloud (SMC, Chapter 2) and globular clusters in the Milky Way as tracers of the chemical and dynamical evolution of these galaxies. It is relevant to study in detail clusters that can be spatially resolved of nearby galaxies in order to understand more distant galaxies and trace a general scenario for formation and evolution of galaxies. The Milky Way is studied

in several chapters, initially presenting the method for chemical abundances derivation (Chapter 3), then releasing of a homogeneous metallicity and alpha-enhancement scale of Galactic globular clusters, and a catalogue of member stars for about 1/3 of the clusters in the Milky Way discussing their origin (Chapter 4). We discuss in particular, the origin of some halo clusters possibly captured from disrupted dwarf galaxies (Chapter 5), and one peculiar bulge cluster that could be the oldest object of our Galaxy (Chapter 6). Finally, in Chapter 7 we give a summary of the thesis and perspectives of works for the near future following the present results. An overview of each topic of the thesis is given below.

1.2.1 *Milky Way - Magellanic Clouds interactions*

Assuming that stellar clusters follow the streams of gas formed by tidal forces, they are pivotal objects to track down the time evolution of the disruption of the stellar content of the galaxy. In particular, using colour-magnitude diagrams of individual clusters, it is possible to derive age and metallicity with enough precision for radial gradients characterization of the population of clusters.

As shown in Chapter 2 we find radial gradients of ages and metallicities in the West Halo region of the SMC, based on star clusters, for the first time. This indicates the existence of gravitational effects stripping out the stellar content of the galaxy. As discussed in Section 1.1.3, this results exclude the possibility of ram pressure effects, which only affects gas. Instead, this disruption of star clusters in the West Halo could be explained by tidal stripping due to the combination of orbits of Small and Large Magellanic Clouds around the Milky Way.

1.2.2 *Formation and evolution of Galactic bulge and halo*

Efficiency of star formation can be indicated by the chemistry of stars, in particular in a plot of $[\text{Fe}/\text{H}]$ vs. $[\alpha/\text{Fe}]$ because iron and α -elements comes from supernovae type II (massive stars with rapid evolution), but supernovae type Ia provide only iron, not α -elements. Therefore, when supernova type Ia becomes to be important the ratio $[\alpha/\text{Fe}]$ decreases. The $[\text{Fe}/\text{H}]$ where $[\alpha/\text{Fe}]$ starts to decrease ($[\text{Fe}/\text{H}]_{\text{knee}}$) reveals how efficiently a given stellar population was formed. Moreover, if these abundance values for globular clusters are compared with their ages this efficiency can be easily converted to a timescale of supernovae explosions, which is crucial information for galaxy formation models. This

analysis can be done for the bulge and halo separately, as we perform in this thesis.

In Chapter 3 we define a method for derivation of chemical abundances ($[\text{Fe}/\text{H}]$ and $[\alpha/\text{Fe}]$) of red giant stars in globular clusters based on spectroscopy with resolution typical for extragalactic observations ($R \sim 2,000$). We validate the method of full spectrum fitting with well-known stars and apply to six globular clusters located in the bulge, disc and halo, covering metallicities from $[\text{Fe}/\text{H}] = -2.5$ to -0.1 . The results are in agreement with high-resolution spectroscopic analysis available in the literature. We apply the method to all 51 globular clusters of our sample and we present a homogeneous metallicity scale for Milky Way globular clusters in Chapter 4, as an update of Harris catalogue, with additional information on $[\alpha/\text{Fe}]$. We also present a catalogue of member stars for about 1/3 of the system of Galactic globular clusters, selected from spectroscopy and photometry. It is the first time that $[\text{Fe}/\text{H}]$ and $[\text{Mg}/\text{Fe}]$ derived in a consistent way are published for such a huge sample of globular clusters. We find a $[\text{Fe}/\text{H}]_{\text{knee}} \sim -1.0$ for bulge and halo based on the clusters, although the bulge seems to be more efficient with a more metal-rich $[\text{Fe}/\text{H}]_{\text{knee}}$. Comparing the abundances and $[\text{Fe}/\text{H}]_{\text{knee}}$ with ages it implies in a timescale of ~ 1 Gyr for supernovae type Ia to start to become important in these components, which agrees with the value widely adopted in models for the Milky Way. Based on $[\text{Fe}/\text{H}]$ vs. age bulge was formed more rapid than halo, and surprisingly $[\text{Mg}/\text{Fe}]$ vs. age does not show this difference.

Halo globular clusters show a lower efficient star formation which is compatible with a scenario where its stellar content was formed in mini halos and/or dwarf galaxies. Some globular clusters may have been originated in the nucleus of dwarf galaxies, already captured by the Milky Way potential well and destroyed in the process. If this is true, their origin should be registered in their chemical contents, showing a complex star formation, continuous or with multiple bursts. There are six massive globular clusters with known internal spread in their $[\text{Fe}/\text{H}]$ distribution, such as ω Centauri, each one with a particular origin. We search for other clusters with spread in $[\text{Fe}/\text{H}]$ in our sample to check their origin.

Two of the 51 clusters in our survey host multiple populations in terms of $[\text{Fe}/\text{H}]$, M 22 and NGC 5824, and possibly there are more among our observations to be investigated. In Chapter 5 we confirm the spread in $[\text{Fe}/\text{H}]$ and show that M 22 has a star formation history similar to that of the Milky Way and it is unlikely that it had an origin in a dwarf

galaxy, in agreement with Marino et al. (2011). NGC 5824 has a position and velocity consistent with the orbit of the Cetus Polar Stream which leads to investigations if the cluster and the stream share a same origin, possibly a disrupted dwarf galaxy Da Costa et al. (2014). With our data it is not possible to answer this question because the spread in $[\text{Fe}/\text{H}]$ has a inter-quartile range of only 0.10 dex which is the order of magnitude of the error bars of our measurements.

If bulge experienced a more efficient star formation and it hosts the centre of the Milky Way, it is possible that the oldest globular clusters are located there. In particular, there are some moderately metal-poor bulge clusters presenting a bluer horizontal branch (HB) than expected by their metallicity, as listed by Barbuy et al. (2009). The main parameter that causes a change in the HB morphology is the metallicity, known as the first parameter. The second most important parameter is age in most cases (Lee et al., 1994), although other parameters can play the role, such as He abundance, CNO abundance or AGB mass loss. Metal-poor clusters have a blue HB, and old clusters too, but these bulge clusters are moderately metal-poor with a blue HB, which indicates that they are very old, possibly the oldest objects of the Milky Way, witnesses of the initial collapse of the Galaxy.

The innermost Milky Way globular cluster is HP 1, for which we determine a metallicity of $[\text{Fe}/\text{H}] = -1.3 \pm 0.4$ in Chapter 4, has a HB index¹ of 0.75, as determined by Mackey and van den Bergh (2005). Therefore this central cluster seems to have a bluer horizontal branch than expected. To constrain future age determination using colour-magnitude diagrams we determine its distance using light curves of RR Lyrae in Chapter 6. We find a distance of 8.0 ± 0.6 kpc consistent with the distance of the Galaxy centre. If this cluster is the oldest of our Galaxy it will confirm that the bulge had a rapid star formation at the beginning of the Milky Way.

¹ $(\text{B}-\text{R})/(\text{B}+\text{V}+\text{R})$, where V is the number of variable stars, B and R are the numbers of stars bluer and redder than the variables, respectively.

Small Magellanic Cloud

2.1 Abstract of this chapter

Stellar clusters in the Small Magellanic Cloud (SMC) are useful probes for studying the chemical and dynamical evolution of this neighbouring dwarf galaxy, enabling inspection of a large period covering over 10 Gyr. The main goals of this work are the derivation of age, metallicity, distance modulus, reddening, core radius and central density profiles for six sample clusters, in order to place them in the context of the Small Cloud evolution. The studied clusters are AM 3, HW 1, HW 34, HW 40, Lindsay 2, and Lindsay 3; HW 1, HW 34, and Lindsay 2 are studied for the first time. Optical Colour-Magnitude Diagrams (V, B-V CMDs) and radial density profiles were built from images obtained with the 4.1m Southern Astrophysical Research (SOAR) telescope, reaching $V \sim 23$. The determination of structural parameters were carried out by applying King profile fitting. The other parameters were derived in a self-consistent way by means of isochrone fitting, which uses likelihood statistics to identify the synthetic CMDs that best reproduce the observed ones. Membership probabilities were determined comparing the cluster and control field CMDs. Completeness and photometric uncertainties were obtained by performing artificial star tests. The results confirm that these clusters (except HW 34, identified as a field fluctuation) are intermediate-age clusters, with ages between 1.2 Gyr (Lindsay 3) and ~ 5.0 Gyr (HW 1). In particular HW 1, Lindsay 2 and Lindsay 3 are located in a region that we called West Halo, where studies of ages and metallicity gradients are still lacking. Moreover Lindsay 2 was identified as a moderately metal-poor cluster with $[\text{Fe}/\text{H}] = -1.4 \pm 0.2$ dex, lower than expected from the age-metallicity relation by Pagel & Tautvaisiene (1998). We also found distances varying from ~ 53 kpc to 66 kpc, compatible with the large depth of

the SMC.

2.2 Introduction to this chapter

Star clusters (SCs) are useful objects for studying the complex stellar content observed in nearby galaxies, as most of them may be modelled as simple stellar populations (SSP) of a fixed age and metallicity. The Magellanic Clouds (SMC for the Small Cloud, LMC for the Large Cloud, and MCs for both) form a rich system with over > 3700 stellar systems (Bica et al., 2008), with combinations of age and metallicity that are not found in the Milky Way (Santos and Piatti, 2004). This information can be used to probe the dynamical and chemical evolution of these neighbouring and interacting dwarf irregular galaxies, and in particular the age-metallicity relation (AMR) of the Magellanic Clouds.

According to Holtzman et al. (1999), the age distribution based on clusters is probably distinct from the star formation history (SFH) as inferred from field stars in the LMC. For the SMC instead, Rafelski and Zaritsky (2005) analysed a sample of 195 clusters, showing that the populations of field stars and star clusters are similar. In particular, Piatti et al. (2005) showed evidence of two peaks in the ages of SMC star clusters at 6.5 Gyr and 2.5 Gyr, and Piatti (2012b) indicates peaks of star formation for field stars at 2 Gyr and 7.5 Gyr. The more recent peak could be due to an encounter with the LMC. This leads to a model of bursts of star formation, in contrast with suggestions of continuous star formation (Da Costa and Hatzidimitriou, 1998). The large period of quiescent star formation in the MCs between ~ 3 Gyr and 10 Gyr (Harris and Zaritsky, 2001, 2004) is indicated by the low number of populous SCs with these ages (Rich et al., 2000, 2001), and almost all of them are in the SMC (Mighell et al., 1998; Piatti et al., 2005,?, 2007,a,b).

Cignoni et al. (2012) studied the SFH of two fields in the SMC, and compared their results to the SFH behaviour suggested by Harris and Zaritsky (2004). They concluded that stars older than 8.4 Gyr do not dominate the SMC stellar population, and the period between 2 Gyr and 8.4 Gyr agrees with Harris and Zaritsky (2004) in one region, but has a much higher star formation rate (SFR) in another region. Therefore, further studies of the SMC cluster and field AMR are needed.

Metallicity values for SMC star clusters, as given in the literature compilation by Parisi et al. (2009) for example, seem to be slightly underestimated when compared with the

chemical evolution model predictions of Pagel and Tautvaisiene (1998), for ages in the range 3 Gyr to 10 Gyr. By adding more data points to this region of the AMR, Piatti (2011b) included a few metal-poor intermediate age star clusters (IACs). To further improve the age-metallicity relation of the SMC, it is important to identify other IACs, and the metal-poor ones are particularly interesting.

In spite of its great interest, the SMC cluster system has been studied less than that of the LMC. Literature data on SMC star clusters were reviewed by Dias et al. (2010). In their Table 6, ages and metallicities for 33 among the most well-studied SMC star clusters are reported, giving particular attention to old/intermediate-age ones. In addition, based on integrated spectra, the analysis by Dias et al. showed that the clusters HW 1 and Lindsay 3 (as well as NGC 152) can be added to the list of intermediate/old clusters. Since then, newly revealed intermediate-age star clusters (IACs) were studied by Piatti (2011a,b), Piatti et al. (2011) and Piatti (2012a).

The compilation of ages and metallicities for SMC clusters by Dias et al. (2010) showed that the literature results have variations up to 7 Gyr in age and 0.9 dex in metallicity, for a given cluster. In the present work, B and V photometry combined with self-consistent statistical tools are employed to determine ages and metallicities for AM 3, HW 1, HW 40, Lindsay 2 and Lindsay 3, in order to provide more precise constraints for the AMR of the SMC. For these clusters, there are the spectroscopic analysis by Dias et al. (2010) of HW 1 and Lindsay 3, and the Washington photometry analysis of AM 3, HW 40 and Lindsay 3 (Piatti, 2011a; Piatti et al., 2011). The target HW 34 is probably only a field fluctuation. For HW 1 and Lindsay 2 no previous CMD data were available in the literature. The confirmation of some of these clusters as intermediate or old age significantly improves the poor census in the age range corresponding to the age gap for the LMC clusters (~ 3 -10 Gyr).

In Sect. 2 the observations, data reduction, and photometry are described. In Sect. 3 the modelling of CMDs and statistical comparisons carried out to find the best fit of synthetic vs. observed CMDs are detailed. In Sect. 4 the results and a discussion of each cluster are given. In Sect. 5 comparisons with the literature, and the age-metallicity relation for the SMC are presented. Finally, a summary is given in Sect. 6.

2.3 Observations

2.3.1 The SOAR/SOI data

Using the SOAR Optical Imager (SOI) mounted on the 4.1m Southern Astrophysical Research (SOAR) Telescope, B and V images were obtained for SMC SCs, under projects SO2007B-013 and SO2008B-017. We chose filters B and V since they provide the best temperature resolution for $-0.2 < B-V < 1.4$, which corresponds to the colours of the present CMDs; V, I would be better in high extinction fields, which is not the case of our fields, and B, V is less affected by differential reddening than is V, I (Soldara, 2008). This imager has a field of view of $5.26' \times 5.26'$, and a pixel scale of $0.077''/\text{pixel}$, which is converted to $0.154''/\text{pixel}$ because of the 2×2 binned observations presented here. The seeing was ~ 0.8 arcsec, and magnitudes up to $V \sim 23$ were detected. The log of observations is reported in Table 2.1.

Table 2.1 - Log of observations. The CCDs were displaced by $\sim 20'$ from the cluster centre to avoid the gap between the set of two E2V CCDs in SOI, as shown in Figure 2.1. The (α, δ) coordinates are from Bica et al. (2008).

Name(s)	α (2000) h:m:s	δ (2000) ° :':"	Filter	Exp. sec.	Airmass	seeing "
2007-09-07						
AM 3	23:48:59	-72:56.7	B	600	1.42	1.00
			V	200	1.40	1.03
Lindsay 2	00:12:55	-73:29.2	B	600	1.38	1.10
			V	200	1.38	1.03
2008-09-21						
HW 1	00:18:27	-73:23.7	B	600	1.48	0.86
			V	200	1.53	0.87
Lindsay 3	00:18:25	-74:19.1	B	600	1.42	1.13
			V	200	1.44	0.86
HW 34	00:57:52	-73:32.7	B	600	1.38	0.95
			V	200	1.39	0.82
HW 40	01:00:25	-71:17.7	B	600	1.34	0.89
			V	200	1.33	0.91

Reduction procedures were based on the SOAR/IRAF packages and the photometry

procedures were based on the DAOPHOT/IRAF package (Stetson, 1987). Classical procedures of aperture and then PSF photometry were performed in the B and V bands. Since PSF fitting resulted in better quality photometry, all the analyses in this paper are based on this method.

The standard stars were chosen from Sharpee et al. (2002), having coordinates close to the SMC direction, in order to save time. Airmass corrections were applied assuming constant values of 0.22 ± 0.03 mag/airmass and 0.14 ± 0.03 mag/airmass for the B and V bands, respectively (as can be found at the CTIO website¹). Only standard stars observed with a seeing lower than $1.0''$, and airmasses close to the cluster observations were considered. Moreover, stars with magnitude variations between an aperture radius of 5 and $8 \times \text{FWHM}$ larger than 0.010 mag were not considered in this fit. Then, the following calibration curves were fitted to the standard stars; the results are presented in Table 2.2. The coefficient of determination r^2 that indicates how well the data fits a line (close to 1 is best) and the low residual values show that both nights were photometric, which is confirmed by the CTIO monitoring²

$$M - m = \alpha \cdot (B - V) + \beta, \quad (2.1)$$

where M corresponds to either B or V, and m corresponds to the respective instrumental magnitudes (given by $-2.5 \times \log(\text{counts}/\text{exptime})$ already corrected by airmass effects.

Table 2.2 - Coefficients of Eq. 2.1 from the fits of the 2007 and 2008 standard stars.

Coef.	B	V
α	0.30 ± 0.10	0.30 ± 0.09
β (mag)	27.02 ± 0.04	26.83 ± 0.07
r^2	0.83	0.44
residuals (mag)	<0.04	<0.10

Figure 2.1 presents the sky map for each cluster, showing the stellar spatial distribution of the SOAR/SOI images, and the selected cluster and field areas. The adopted cluster (R_{clus}) and field radii (R_{field}) for all clusters are 30 arcsec and 90 arcsec, respectively,

¹ <http://www.ctio.noao.edu/noao/content/13-m-photometric-standards>

² http://www.ctio.noao.edu/site/phot/sky_conditions.php

covering solid angles (Ω_{clus} and Ω_{field}) of about 0.79 arcmin² and 20.6 arcmin². These sky maps enable us to visually identify the overdensity inside the cluster radii (except for HW 34) and to verify which are the more/less crowded lines of sight.

Figure 2.1: Sky maps showing the stellar distribution for all clusters, as imaged by the SOAR/SOI (1 pixel = 0.154 arcsec). The small circle ($R < R_{\text{clus}} = 30$ arcsec) corresponds to the cluster sample, whereas the large circle limits the area for the control field sample ($R > R_{\text{field}} = 90$ arcsec).

2.3.2 Photometric errors and completeness curves

To properly determine the completeness of the photometry and the photometric errors we performed artificial star tests (ASTs). These consist of adding stars (with known magnitudes and colours) in random positions to the reduced images, and then carrying out the photometry exactly as was done with the original data. The ratio of recovered to input stars is called completeness (see Figure 2.2).

The procedure was carried out for both B and V images assuming an average colour of (B-V)=0.5 for all the stars. The most important curves are those from the V band, since a V magnitude limit is applied on Figure 2.4, and the curves are used to correct the radial density profiles (Fig. 2.5), which are better sampled by V images. In fact, B completeness curves were used only by internal checks: they differ by <0.5 mag from V curves with the same completeness values. We considered that this is a good approximation for the

Figure 2.2: Completeness curves for different distances from the cluster centre, based on V-band photometry. These curves correspond to a ten-order smoothed spline fitted to the data. Different lines are the completeness curves for different annuli, with the respective data points: $0 \leq r < 10''$ (red dotted/circle); $10 \leq r < 20''$ (green dashed/triangle); $20 \leq r < 30''$ (black solid/rhombus); $30 \leq r < 40''$ (blue dash dotted/inverted triangle); $40 \leq r < 50''$ (cyan long dashed/square). The magnitude limit corresponding to 50% of completeness are indicated in the plots. Completeness uncertainties are indicated in grey vertically (assumed as Poissonian), and typical magnitude uncertainties are indicated in grey horizontally.

observed stars with colours ranging from $\sim 0.0 < B - V < 1.0$. Therefore, we used only V magnitude limits, as follows. The total range of V magnitudes covering all observed stars (from ~ 17 to ~ 24 mag) were divided into 32 intervals. For each interval, two sets of 465 stars were simulated inside a radius of $50''$ from the cluster centre, with random magnitudes, separated by at least $3''$ ($\gtrsim 3.5 \times < \text{FWHM} >$) to avoid introducing additional crowding (e.g. Rubele et al., 2011). The resulting completeness curves as a function of the V magnitude are presented in Figure 2.2 for different annuli around the cluster centre. The limiting magnitude is set where the completeness ratio is equal to 50% (see values in the figure). We adopted the V_{limit} values of the annulus $20'' < r < R_{clus} \equiv 30''$, where R_{clus} is defined as the outer cluster radius used for the CMD analysis.

The correspondence between Figures 2.1 and 2.2 is evident. In particular, for the central regions (red solid curve), completeness for HW 34, HW 40, and Lindsay 3 are similar, as expected from the similar central densities of the stars. Instead, for external annuli,

HW 40 and Lindsay 3 are more complete (2 magnitudes more), while HW 34 presents roughly similar completeness to the centre. This indicates a crowded and homogeneous field in the HW 34 direction.

The observational photometric errors for V and (B-V) as given by DAOPHOT are presented in Figure 2.3. The photometric uncertainties, i.e., the standard deviations of the differences between input and output magnitudes and colours derived from the ASTs, are presented together with the CMDs (Section 2.3.4).

Figure 2.3: V and (B-V) photometric errors given by DAOPHOT outputs. Vertical dashed lines correspond to magnitude limits in a radius of 30'', as shown in Figure 2.2.

2.3.3 The radial density profiles

The radial density profiles for all clusters are shown in Fig. 2.5. Only stars brighter than the magnitude limit presented in Fig. 2.4 were considered. Two profiles are presented: a) the lower density, which is observed, and b) the higher density corresponding to those corrected for completeness. Each star has a value for the completeness that comes from an interpolation in radius and magnitude, based on the curves presented in Figure 2.2. The error bars are the propagation of the Poissonic errors on the star counts ($\sqrt{N}/\text{area}$), corrected by the incompleteness of the sample. Each profile is fitted with the empirical

Figure 2.4: V, (B-V) colour-magnitude diagrams for all clusters. Left panels: cluster stars ($R < R_{\text{clus}}$). Right panels: control field stars ($R > R_{\text{field}}$). The colours depend on the membership probabilities (p_{member}) for each star in the cluster direction ($R < R_{\text{clus}}$). The horizontal dashed lines correspond to the magnitude limits, derived from completeness curves around R_{clus} .

density law of King (1962), expressed by equation 2.2 below, using a nonlinear least-squares routine

$$n = n_0 \left\{ \frac{1}{[1 + (r/r_c)^2]^{1/2}} - \frac{1}{[1 + (r_t/r_c)^2]^{1/2}} \right\}^2 + n_{\text{field}}, \quad (2.2)$$

where n_0 is the central density of the cluster, n_{field} is the density of field stars, r_c is the core radius, and r_t is the tidal radius. From these parameters one can quantify the concentration parameter c , defined as $c \equiv \log(r_t/r_c)$. For each cluster, the value of n_{field} was assumed to be the average density of stars outside $R_{\text{field}} \equiv 90''$, and it was kept constant in the fits. The final fits are presented in Fig. 2.5, whereas all recovered structural parameters are shown in Table 2.3.

The comparison of the panels in Fig. 2.5 and the parameters in Table 2.3 reveal, as would be predicted from a visual analysis of the sky maps, that the objects present quite different stellar concentrations and field stellar densities. In all cases the core radius $r_c < R_{\text{cluster}}$ and the tidal radius $r_t < R_{\text{field}}$, which confirm that the adopted values of R_{cluster} and R_{field} are reasonable. All clusters present roughly the same level of concentration, and

the central density contrast with respect to the field is about 10 to 80 times the field density. We note that the original profile of Lindsay 2 and both profiles of HW 34 did not converge. The first is due to incompleteness, and the second does not have enough density contrast in the centre to be fitted by a King profile, which can be interpreted as evidence of a non-physical system.

Figure 2.5: Radial numerical density profiles fitted by King profiles. The empty grey circles and filled black diamonds are the observed profiles, original and corrected by completeness (Figure 2.2), respectively. Correspondent lines are King profiles fitted to the data points. The horizontal and vertical dashed grey lines are n_{field} and r_c , respectively, for the original profile, the dash dotted black lines for the corrected profile. HW34 does not present a high enough density contrast to be fitted by a King profile.

2.3.4 The observed CMDs and cluster membership probability

Figure 2.4 shows the $V \times B-V$ CMDs for the present sample. For each object, the cluster ($R < R_{\text{clus}}$) and field ($R > R_{\text{field}}$) CMDs are presented, with stars from $V \sim 17$ up to $V \sim 23$. It is possible to identify main sequence (MS) stars as well as red giant stars belonging to the subgiant branch (SGB), red giant branch (RGB), and to the red clump (RC). The comparisons with the field CMDs confirm these stellar identifications. We note that the field CMDs are much richer in stars than the cluster CMDs, as a natural consequence of the ratio between their covered areas ($\frac{\Omega_{\text{field}}}{\Omega_{\text{clus}}} \sim 26$).

Table 2.3 - Structural parameters derived from King profile fitting, as described in Section 2.3.3. The overdensity column indicates how much bigger the central density is with respect to the field density. *Notes:* a) cases where fit did not converge, and we attribute the density of the innermost region of the profile as n_0 to estimate a lower limit of the overdensity.

Cluster	n_{field} (stars/' ²)	n_0 (stars/' ²)	r_c (")	r_t (")	c $\log(r_t/r_c)$	overdensity $(n_0 - n_{\text{field}})/(n_{\text{field}})$
AM 3 – original	0.0014	0.072 ± 0.009	21.6 ± 1.4	65 ± 6	0.48 ± 0.05	50
AM 3 – completed	0.0018	0.150 ± 0.015	18.1 ± 1.1	62 ± 6	0.54 ± 0.05	82
Lindsay 2 – original	0.0061	0.11 ± 0.05	41 ± 8	85 ± 17	0.31 ± 0.12	17
Lindsay 2 – completed	0.0078	0.25 ± 0.05	29.8 ± 3.2	74 ± 9	0.39 ± 0.07	31
HW 1 – original	0.0162	0.114 ± 0.009	17.2 ± 1.3	49.2 ± 3.0	0.46 ± 0.04	6
HW 1 – completed	0.0204	0.349 ± 0.021	14.7 ± 0.8	55.3 ± 3.7	0.58 ± 0.04	16
Lindsay 3 – original ^a	0.0116	0.057 ± 0.011	—	—	—	3.9
Lindsay 3 – completed	0.0155	0.238 ± 0.010	16.4 ± 0.5	89 ± 8	0.73 ± 0.04	14
HW 40 – original	0.0195	0.096 ± 0.019	16.6 ± 2.3	69 ± 17	0.62 ± 0.12	4
HW 40 – completed	0.0249	0.28 ± 0.04	17.8 ± 1.6	65 ± 9	0.56 ± 0.7	10
HW 34 – original ^a	0.0444	0.069 ± 0.012	—	—	—	0.5
HW 34 – completed ^a	0.0960	0.212 ± 0.021	—	—	—	1.2

Since the stellar samples in the direction of the clusters are contaminated by SMC field stars, a procedure to determine the cluster membership probability (p_{member}) was applied. In summary, the adopted procedure follows the one from Kerber and Santiago (2005) and Alves et al. (2012), where p_{member} for each cluster star is determined by comparing the density of stars inside the CMD in the cluster direction ($R < R_{\text{clus}}$), with a CMD representative of the local SMC field ($R > R_{\text{field}}$) (see Fig. 2.4). The cluster and field CMDs are divided into a grid of small boxes in V magnitude and B-V colour, centred on each cluster star. So, for the i^{th} cluster star, the number of cluster ($N_{\text{clus},i}$) and field stars ($N_{\text{field},i}$) contained in a box in magnitude and colour (with a 3-sigma size in photometric errors) are computed, taking the inverse of the completeness into account. Finally, the cluster membership probability for the i^{th} cluster star is given by

$$p_{\text{member},i} = 1 - \frac{N_{\text{field},i}}{N_{\text{clus},i}} \times \frac{\Omega_{\text{clus}}}{\Omega_{\text{field}}}. \quad (2.3)$$

The results of these determinations are shown in Fig. 2.4 in a colour scale. The analysis of this figure reveals that most probable cluster members ($p_{\text{member}} > 75\%$) are red giants or stars in the upper MS. In particular, almost all stars in the AM 3 cluster sample can be considered physical cluster members. On the other hand, the MS stars in the HW 34 cluster direction have a higher probability of being field stars rather than cluster stars since they present $p_{\text{member}} < 50\%$. This strongly reinforces the argument that this stellar

concentration is not a physical system, but only a field stellar fluctuation (see also Sect. 2.3.3). Table 2.4 summarizes the star counts in the sample CMDs.

Table 2.4 - Stellar counts in the sample CMDs. Columns correspond to: the cluster's name, the number of stars without the completeness corrections and with them for the cluster ($N_{\text{clus}}^{(\text{obs})}$ and $N_{\text{clus}}^{(\text{comp})}$) and field direction ($N_{\text{field}}^{(\text{obs})}$ and $N_{\text{field}}^{(\text{comp})}$), as well as the expected number of field stars in the completed cluster sample, with $p_{\text{member}} < 0.5$ (contamination C). The last column is the number of cluster stars in terms of the field standard deviation (assumed as Poissonian), given by the expression $N_{\sigma} = (N_{\text{clus}}^{(\text{comp})} - C) / \sqrt{N_{\text{field}}^{(\text{comp})}}$.

Target	$N_{\text{clus}}^{(\text{obs})}$	$N_{\text{clus}}^{(\text{comp})}$	$N_{\text{field}}^{(\text{obs})}$	$N_{\text{field}}^{(\text{comp})}$	C	N_{σ}
AM 3	56	172	135	209	8	11
Lindsay 2	127	490	847	2434	93	8.1
HW 1	112	370	1659	3686	141	3.8
Lindsay 3	116	342	1412	3818	145	3.2
HW 40	133	393	1906	3790	143	4.1
HW 34	171	562	4014	13594	515	0.4

2.4 Statistical isochrone fitting

The isochrone fitting used in this work follows a numerical and statistical approach that combines CMD modelling and an objective criterion of comparing synthetic CMDs with the observed ones. It was initially developed to analyse CMDs of rich LMC clusters, observed with the Hubble Space Telescope (HST) (Kerber et al., 2002), and recently it was applied to determining physical parameters of Galactic open clusters imaged by 2MASS (Alves et al., 2012). The central idea behind this method is to statistically determine the synthetic CMDs that best reproduce the observed CMD, recovering as a consequence the physical parameters for the stellar cluster. In the present work the statistics adopted to select the best models is the likelihood, in particular following a Bayesian approach. This approach has been successfully applied to analysing CMDs of stellar clusters (Naylor and Jeffries, 2006; Hernandez and Valls-Gabaud, 2008; Monteiro et al., 2010) as well as CMDs of composite stellar populations (Hernandez et al., 2000, 1999; Vergely et al., 2002).

The generation of synthetic CMDs, as well as the likelihood statistics, are detailed in the next subsections.

2.4.1 CMD modelling

The synthetic CMDs are generated considering that they are SSPs, characterized by stars with the same age (τ) and metallicity (Z). The basic steps to generate a specific synthetic CMD are the following (see Fig. 2.6):

- (i) adoption of an evolutionary stellar model. The new Padova-Trieste models (PARSEC isochrones, Bressan et al., 2012) were adopted in this work;
- (ii) selection of the age and metallicity of the SSP, which is equivalent to selecting an isochrone from the library;
- (iii) application of the distance modulus $(m - M)_0$ and reddening $E(B - V)$ values to shift the isochrone from the absolute magnitudes to the observed ones;
- (iv) generation of 10^5 synthetic stars belonging to stellar systems (single and binary stars) following an IMF and a fraction of unresolved binaries (f_{bin}). A Salpeter IMF ($dN/dm \propto m^{-2.35}$) and $f_{\text{bin}} = 30\%$ were adopted, with a minimum mass ratio ($q_{\text{min}} = m_2/m_1$) of 0.7 and a constant mass-ratio distribution ($dN/dq = \text{const.}$);
- (v) introduction of the observational photometric errors in magnitude and colour.

To determine the cluster parameters, a wide and regular grid of models was build for each cluster, composed typically of ~ 9000 combinations in $\log(\tau/\text{yr})$, Z , $(m - M)_0$ and $E(B - V)$ centred on the parameters found by a visual isochrone fit. To take into account the line-of-sight depth and the reddening values found for the SMC stellar clusters (Glatt et al., 2008; Crowl et al., 2001), the grids of models span distance moduli and reddening values in the range $18.50 < (m - M)_0 < 19.20$ and $0.00 < E(B - V) < 0.20$, in steps of 0.05 and 0.01, respectively. For $\log(\tau/\text{yr})$ the step is 0.05, and for Z all values provided by the isochrones are considered: 0.0001, 0.0004, 0.001, 0.002, 0.004, 0.006, 0.008, 0.010, 0.012 and 0.014.

2.4.2 Likelihood statistics

The first step in computing the likelihood for a synthetic CMD is to establish the probabilities for each observed star cluster that it belong to that choice of SSP (p_{CMD}). In general, this is done using analytical expressions (e.g. Hernandez and Valls-Gabaud, 2008; Monteiro et al., 2010), assuming that the observed CMD positions would represent stars scattered from the isochrones following Gaussian distributions for the photometric

errors. However, in the present work these probabilities are computed assuming that they are proportional to the density of points generated by the synthetic CMD in each of the respective CMD positions ($N[V, (B - V)]$), which naturally incorporates the effect of unresolved binaries and photometric uncertainties. This density map on the CMD is commonly called a Hess diagram, even though its application is not directly related with the likelihood statistics. Therefore, the likelihood statistics in this numerical approach is given by the product of these probabilities over all the N_{clus} observed cluster stars as stated by the expression

$$L \propto \prod_{i=1}^{N_{\text{clus}}} p_{\text{CMD},i} \times p_{\text{member},i} \propto \prod_{i=1}^{N_{\text{clus}}} N[V_i, (B - V)_i] \times p_{\text{member},i} , \quad (2.4)$$

where the i index corresponds to the i^{th} observed cluster star. It can thus be stated that the best model is the one that provides the synthetic CMD/Hess diagram that maximizes the above expression (L_{max}). The final parameters for each cluster, as well as corresponding

Figure 2.6: The generation of synthetic V vs. B-V CMDs. *Panel a:* the adopted isochrone shifted by distance modulus and reddening. *Panel b:* the distribution of stars in accordance to the IMF and the fraction of binaries; *Panel c:* the introduction of photometric errors in magnitude and colour. *Panel d:* colours following the density of points in the CMD in a logarithmic scale. The density of points is related to $p_{\text{CMD},i}$ (see Sect. #2.4.2). The reddening vector and the values for all physical parameters are shown in the figure.

uncertainties, are determined by computing the average and standard deviation over the physical parameters of the set of models that have likelihood values at the 1-sigma level from $L_{\max}$ (see Appendix A for details).

2.5 Results

The results of the isochrone fittings for the sample clusters are presented in Figure 2.7 and Table 2.5. The panels in Fig. 2.7 give isochrones corresponding to the best-fit parameters, together with another two isochrones indicating $1\text{-}\sigma$ uncertainties in age and metallicity: one younger and more metal-rich, and the other older and more metal-poor. The three isochrones cover most of the data points, as expected, allowing the reader to visually inspect the quality of the fits. For the $[\text{Fe}/\text{H}]$ derivation we adopted a solar metallicity $Z=0.0152$ (Caffau et al., 2011) in order to be compatible with the value assumed in the PARSEC isochrones (Bressan et al., 2012). These results are discussed cluster by cluster below.

Table 2.5 - Physical parameters determined in this work. Columns refer to: cluster name, logarithm of the age, age, metallicity Z and $[\text{Fe}/\text{H}]$ (assuming $Z_{\odot}=0.0152$, Caffau et al., 2011), distance modulus, distance, reddening, and semi-major axis corresponding to the distance of the cluster to the centre of SMC (as explained in Section 2.6).

Name	$\log(\tau/\text{yr})$	Age(Gyr)	Z	$[\text{Fe}/\text{H}]$	$(m-M)_0$	d(kpc)	$E(B-V)$	$a(^{\circ})$
AM 3	9.69 ± 0.15	$4.9^{+2.1}_{-1.5}$	0.0022 ± 0.0016	$-0.8^{+0.2}_{-0.6}$	18.99 ± 0.16	$63.1^{+1.8}_{-1.7}$	0.08 ± 0.05	7.3
Lindsay 2	9.60 ± 0.09	$4.0^{+0.9}_{-0.7}$	0.0007 ± 0.0004	$-1.4^{+0.2}_{-0.2}$	18.68 ± 0.14	$54.4^{+1.5}_{-1.5}$	0.09 ± 0.04	3.9
HW 1	9.70 ± 0.12	$5.0^{+1.5}_{-1.2}$	0.0013 ± 0.0007	$-1.1^{+0.2}_{-0.3}$	18.84 ± 0.16	$58.7^{+1.6}_{-1.6}$	0.07 ± 0.04	3.4
Lindsay 3	9.07 ± 0.11	$1.2^{+0.3}_{-0.3}$	0.0057 ± 0.0020	$-0.4^{+0.1}_{-0.2}$	18.64 ± 0.14	$53.4^{+1.5}_{-1.5}$	0.07 ± 0.04	2.9
HW 40	9.41 ± 0.06	$2.5^{+0.4}_{-0.3}$	0.0018 ± 0.0005	$-0.9^{+0.1}_{-0.2}$	19.08 ± 0.14	$65.6^{+1.8}_{-1.8}$	0.03 ± 0.03	2.0

2.5.1 AM3

Although this is the cluster with the smallest number of stars in our sample, it has the advantage of being located in the SMC field with the lowest stellar density. The results in the present work indicate an age of $4.9^{+2.1}_{-1.5}$ Gyr. The recovered metallicity, $[\text{Fe}/\text{H}] = -0.8^{+0.2}_{-0.6}$, is consistent with the value predicted by the SMC chemical evolution model from Pagel and Tautvaisiene (1998) for this age. Piatti (2011a) also determined

Figure 2.7: Best isochrone fittings for all clusters. Left panels: cluster stars with colour symbols in accordance to the membership probabilities (p_{member}). Right panels: synthetic CMDs generated with the parameters found for the best solution. The number of points is equal to the observed CMDs within the same magnitude limits. The horizontal dashed black lines correspond to the magnitude limits used to compute the likelihood (brighter mag) and p_{member} (fainter mag). The solid and dashed blue lines correspond to the isochrones with the parameters found in Table 2.5 and take into account the uncertainties on ages and metallicities.

physical parameters for this cluster, based on the analysis of a CMD built with Washington photometry. Taking the uncertainties into account, our results are in good agreement with the ones from Piatti et al., although they derived slightly older ages (6.0 ± 1.0 Gyr) and more metal poor values ($[\text{Fe}/\text{H}] = -1.25 \pm 0.25$). Our results are also close to the parameters derived by Da Costa (1999) of $[\text{Fe}/\text{H}] = -1.27$ and age of 5.5 ± 0.5 Gyr.

The differences in age can be explained by the distance modulus. While we fitted a $(m-M)_0 = 19.00$ for this cluster, Piatti (2011a) assumed an average value of 18.90. With a distance modulus ~ 0.1 lower, the isochrone should be ~ 0.6 Gyr older to fit the data.

This western SMC cluster was considered a galaxy member by Da Costa (1999) because it is close to the projected plan of the SMC, and its red clump magnitude (distance indicator) is comparable to other SMC clusters. It is located in the West Halo region, but it is not as old and metal poor as one could expect in the case of gradients in age and $[\text{Fe}/\text{H}]$ for this group (see Sect. 2.6). Although AM 3 is close to the SMC main body in the

projected plan, its distance is higher than the other West Halo clusters (HW 1, Lindsay 2, and Lindsay 3). Therefore its age, metallicity, and distance do not follow exactly the general characteristics of the West Halo, which indicates a more complex history, possibly a capture by the potential well of the SMC.

2.5.2 HW 1 and Lindsay 2

The CMD analysis for these clusters which have never been published before, identifies these objects as intermediate-age metal-poor SMC clusters, with similar metallicities ($[\text{Fe}/\text{H}] \sim -1.1$ and -1.4 , respectively) and ages (4.98 and 4.0 Gyr). In particular, the age and metallicity values for HW 1 confirm the previous results from the integrated spectral analysis performed by Dias et al. (2010). We note that these two clusters are located close to each other, in the West Halo region (Fig. 2.8), at angular distances from the SMC centre of $\alpha = 3.4^\circ$ to 3.9° , and distances from the Milky Way of 58.7 and 54.4 kpc, respectively. Their ages correspond to the epoch of a probable recent tidal interaction between the Magellanic Clouds, therefore they could be remnants of this period.

2.5.3 HW 40

The CMD analysis for this cluster indicates that it is an intermediate-age SMC cluster with $2.5_{-0.3}^{+0.4}$ Gyr. The presently derived metallicity ($[\text{Fe}/\text{H}] = -0.9_{-0.2}^{+0.1}$) is in agreement with the results from Piatti (2011a) based on photometry ($[\text{Fe}/\text{H}] = -1.10 \pm 0.25$), and by Parisi et al. (2013, private comm.; $[\text{Fe}/\text{H}] = -0.78 \pm 0.05$ dex) using CaT spectra. The age determined by Piatti et al. is significantly older (5.4 ± 1.0 Gyr) than our result. More accurate CMDs are needed for this cluster to constrain its age.

2.5.4 Lindsay 3

The age (1.2 ± 0.3 Gyr) and metallicity $-0.4_{-0.2}^{+0.1}$ for this cluster is in good agreement with the results from Piatti et al. (2011): 1.25 ± 0.20 Gyr and $[\text{Fe}/\text{H}] = -0.65 \pm 0.20$ dex. A remarkable result from our isochrone fitting is the small distance modulus for this cluster (18.64 ± 0.14), suggesting that it is one of the closest SMC clusters. We note that here too Piatti et al. (2011) assumed a distance modulus of 18.90 and also a reddening of 0.04 instead of our fitted value of 0.07. The combination of these two values tends to compensate the effects on age and metallicity. Lindsay 3 is younger than the bulk of the

surrounding field stellar population in the West Halo region (HW 1 and Lindsay 2 are ~ 3 Gyr older, for instance). If the Magellanic Clouds suffered interactions at about 4 Gyr ago, this could be a resulting cluster in an older field, or else it is possible that there is an age gradient in the West Halo. More data on the star clusters of this region are needed to confirm these conclusions (see Sect. 2.6).

2.5.5 HW 34

No star cluster was identified, as made evident by the following:

- Sky maps given in Fig. 2.1 and radial density profiles in Fig. 2.5 show no significant overdensity in the region centred on HW 34 coordinates;
- the photometric data are of comparable quality to the other clusters (see Figs. 2.2 and 2.3), so a cluster would be recovered if present;
- Almost all stars in the direction of HW 34 coordinates were given the probability of being field stars, as shown in Fig. 2.4 and Table 2.4;
- Based on the very few RGB stars with $p_{\text{member}} > 0.5$, and no clear main sequence turnoff, if there was a cluster there, it would be older than ~ 5 Gyr, which is not expected for the central younger regions of the SMC.

2.6 The age-metallicity relation and spatial distribution

Figure 2.8 shows the 2D positions of the five clusters in the sky plane, where clusters from the catalog of Bica et al. (2008) were overplotted. Ellipses are indicated to find the clusters' distances from the SMC centre, following the procedure of Piatti et al. (2005). We adopt the coordinates of the SMC centre from Crawl et al. (2001), $(\alpha, \delta) = (0^{\text{h}}52^{\text{m}}45^{\text{s}}, -72^{\circ}49'43'')$, and a minor to major axis ratio of $b/a = 1/2$. Thus, the indicator of distance to the SMC centre would be the semi-major axis a , which is more appropriate for an ellipsoidal galaxy like the SMC (see Figure 2.8). We consider an inclination of 45° for the major axis of all the ellipses in the projected plane of the figure. The distances a for all the clusters can be found in Table 2.5.

Most of the sample clusters are located outside the region sampled by Harris and Zaritsky (2004), but inspired by the different SFR that they found in different regions of

the SMC, we located our targets in different groups for a $> 2^\circ$ (see Fig. 2.8). These regions were simply divided into the sky plane based on the HI structures, as described by Diaz and Bekki (2012), assuming that these structures have stellar/star cluster counterparts. They are: Bridge, Counter-Bridge, and a third region not related with the HI structures that we have called West Halo. They are indicated by different colours and symbols in Figs. 2.8, 2.9, and 2.10.

The 3D structure of the SMC was shown to be complex (Subramanian and Subramanian, 2012), with a spread in depth of about 14 kpc, and an inclination along the main axis with the north-eastern portion towards the observer. The five clusters studied in this work follow this dispersion, in particular for West Halo clusters, where we found distances between ~ 54 and 63 kpc, revealing that more statistics on this region are needed in order to better describe the 3D structure of the SMC.

Figure 2.8: On sky distribution of the SMC star clusters, including the catalog clusters (Bica et al., 2008), overplotted with the five sample clusters. The ellipses are used to illustrate distances from the SMC centre (Table 2.5). Different colours indicate subgroups, as indicated in the panel (see text for details).

Figure 2.9 shows the age-metallicity relation (AMR) of the SMC, by plotting data for well-studied star clusters (Dias et al., 2010; Piatti et al., 2011; Piatti, 2011a,b), overplotted

by the model of Pagel and Tautvaisiene (1998). The results for the sample clusters fit well the Pagel and Tautvaisiene (1998) model, and agree with a dispersion in metallicity for the SMC cluster system in the metallicity and age ranges of $-1.5 < [\text{Fe}/\text{H}] < -1.0$, and $5 < \text{age} < 10$ Gyr, as indicated by Piatti (2011b), in their Fig. 3. The field stars also agree with this statement, as can be seen in Fig. 8 of Piatti (2012b). In this case, the model of Pagel and Tautvaisiene (1998) would represent an average of the chemical enrichment history of the SMC. Spectroscopic results by Parisi et al. (2009) also show a spread in $[\text{Fe}/\text{H}]$ among intermediate-age clusters; however, their values are in the range $-1.2 < [\text{Fe}/\text{H}] < -0.7$ dex. Therefore, both photometric and spectroscopic metallicities show a spread of ~ 0.5 dex in metallicity, with limits defined by different metallicity scales.

Another important product of this work concerns AM 3, Lindsay 2, and HW 1, the oldest clusters of the sample, with 4.9, 4.0, and 4.98 Gyr, respectively. They are close to the end of the quiescent star formation as pointed out by Harris and Zaritsky (2004), and can be among the first products of the reactivation of star formation in the MC system $\sim 3\text{--}4$ Gyr ago.

The cluster distribution in the 2D projected plane of the sky is also useful for investigating gradients of age and metallicity. This is presented in Figure 2.10, where the marked labels correspond to those of Figure 2.9. The five clusters presented in this work are located outside the 2° ellipse. The small sample combined with the error bars do not permit us to establish a strong gradient. Nevertheless, it is possible to identify that the most internal regions of the SMC, below $a = 1\text{--}2^\circ$ might indicate a gradient in age and metallicity, whereas the outer regions have a spread in these parameters. Otherwise, these trends seem to be consistent with those of the SMC field population (Piatti, 2012b).

Even so, if one looks at different regions of the projected SMC, they reveal different superimposed gradients (which would generate the overall dispersion with no gradients), but the dispersion is too high and more data are needed for this to be confirmed. Lindsay 2, Lindsay 3, and HW 1, which are located in the West Halo region, tend to show a possible age gradient in this region, as indicated by the blue diamonds in the figure.

Figure 2.9: Age-metallicity relation for the SMC, including the new additions from the present work. The well-known clusters are listed in Table 6 of Dias et al. (2010), and other clusters from literature. The model of Pagel and Tautvaisiene (1998) is overplotted.

2.7 Summary of this chapter

We studied intermediate-age stellar clusters in the SMC. The parameters were derived by statistical fitting of the observed CMDs with a grid of synthetic CMDs based on the PARSEC isochrones. Six targets were observed in the visible filters B and V, and we derived ages for five of them (AM 3, HW 1, HW 40, Lindsay 2, and Lindsay 3) in the age range of 1.2 Gyr to 5.0 Gyr, whereas HW 34 was identified as a field fluctuation. In particular, Lindsay 2 and HW 1 were studied for the first time in this work, and for the other clusters we derived self-consistent parameters compatible with the literature, and in principle with more precision.

The five clusters essentially follow the chemical enrichment history modelled by Pagel and Tautvaisiene (1998). Figure 2.9 shows a strong spread in metallicity for clusters with ages of 2-5 Gyr, also found by Piatti (2011b). Piatti (2012b) finds a metallicity spread of ~ 0.4 dex at ages of 2-4 Gyr, and Parisi et al. (2010) finds a dispersion of ~ 0.32 dex around

Figure 2.10: Age and $[\text{Fe}/\text{H}]$ as a function of the distance indicator a . The dots are the same as described in Figure 2.9.

the mean value of $[\text{Fe}/\text{H}]=-1.0$ from 15 fields.

This suggests that, after an initial period of rapid enrichment, that brought the metallicity up to around $[\text{Fe}/\text{H}]\approx-1.2$, the SMC chemical enrichment is very inhomogeneous, since there is a spread in metallicity from ~ 11 to ~ 3 Gyr, with a trend for slow enrichment. Finally an intense burst occurred at around 2 Gyr ago, when most of the star clusters started to form.

Piatti et al. (2011) suggested the occurrence of two recent bursts at ~ 2 Gyr and ~ 6 Gyr ago (preceded by a quiescent period lasting ~ 3 Gyr), with roughly no metallicity variations. Our clusters improve the statistics for the two more recent bursts identified by Piatti et al. (2011), in particular HW 1 and Lindsay 2, which are located in the West Halo region, at almost the same distance.

A gradient of age and metallicity can be found in the inner part of the SMC ($a < 2^\circ$), but in the outer regions the high dispersion in ages and metallicities prevents the identification of any trends. More data on age, metallicity/abundances, distance, and kinematics of star

clusters are needed in order to prove different star formation rates and chemical evolution histories in the regions due to tidal forces caused by the interactions in the SMC-LMC-Milky Way system.

FORS2/VLT survey of Milky Way globular clusters

I. Method validation

3.1 Abstract of this chapter

We have observed almost 1/3 of the globular clusters in the Milky Way, targeting distant and/or highly reddened objects, besides a few reference clusters. A large sample of red giant stars was observed with FORS2@VLT/ESO at $R \sim 2000$. The method for derivation of stellar parameters is presented with application to six reference clusters. We aim at deriving the stellar parameters effective temperature, gravity, metallicity and alpha-element enhancement, as well as radial velocity, for membership confirmation of individual stars in each cluster. We analyse the spectra collected for the reference globular clusters NGC 6528 ($[\text{Fe}/\text{H}] \sim -0.1$), NGC 6553 ($[\text{Fe}/\text{H}] \sim -0.2$), M 71 ($[\text{Fe}/\text{H}] \sim -0.8$), NGC 6558 ($[\text{Fe}/\text{H}] \sim -1.0$), NGC 6426 ($[\text{Fe}/\text{H}] \sim -2.1$) and Terzan 8 ($[\text{Fe}/\text{H}] \sim -2.2$). They cover the full range of globular cluster metallicities, and are located in the bulge, disc and halo. Full spectrum fitting techniques are applied, by comparing each target spectrum with a stellar library in the optical region at 4560-5860 Å. We employed the library of observed spectra MILES, and the synthetic library by Coelho et al. (2005). Validation of the method is achieved through recovery of the known atmospheric parameters for 49 well-studied stars that cover a wide range in the parameter space. We identified 4 member stars in NGC 6528, 13 in NGC 6553, 10 in M 71, 5 in NGC 6558, 5 in NGC 6426 and 12 in Terzan 8. Radial velocities, effective temperatures, gravities, metallicities and alpha-element enhancements were derived. We derived $\langle v_{\text{helio}} \rangle = -242 \pm 11$ km/s, $[\text{Fe}/\text{H}] = -2.39 \pm 0.04$, $[\text{Mg}/\text{Fe}] = 0.38 \pm 0.06$ for NGC 6426 from spectroscopy for the first time. The method proved to be reliable for red giant stars observed with resolution $R \sim 2000$, yielding results compatible

with high-resolution spectroscopy. The derived α -element abundances show $[\alpha/\text{Fe}]$ vs. $[\text{Fe}/\text{H}]$ consistent with that of field stars at the same metallicities.

3.2 Introduction to this chapter

The derivation of stellar metallicities and abundances are best defined when based on high spectral resolution and high signal-to-noise (S/N) data. Cayrel (1988) showed that higher resolution carries more information than higher S/N. Such kind of data require however substantial telescope time. For this reason, very large samples of stellar spectra have been gathered in recent years, or are planned to be collected in the near future, with multi-object low and medium-resolution instruments. A few examples are the Sloan Digital Sky Survey (SDSS, York et al., 2000), at a resolution $R \sim 1800$, the Radial Velocity Experiment survey (RAVE, Steinmetz et al., 2006) of $R \sim 7500$ in the CaT region, and other large ongoing surveys such as LAMOST at the Guoshoujing telescope (GSJT, Wu et al., 2011) of $R \sim 2000$, and future ones such as GAIA (Perryman et al., 2001). Large data sets of low/medium-resolution spectra are reachable for extragalactic stars, such as presented in Kirby et al. (2009). A few recent surveys are able to use medium/high-resolution spectra focused on specific targets such as provided by the APOGEE ($R \sim 22,500$, Mészáros et al., 2013), GAIA-ESO using the FLAMES-GIRAFFE spectrograph ($R \sim 22,000$) at the Very Large Telescope (VLT, Gilmore et al., 2012) and HERMES ($R \sim 28,000$ or $45,000$, Wylie-de Boer and Freeman, 2010) at the AAT. More complete reviews of available, ongoing and future surveys, as well as automated methods for stellar parameter derivation can be found in Allende Prieto et al. (2008), Lee et al. (2008), Koleva et al. (2009), Mészáros et al. (2013), and Wu et al. (2011), among others.

In most analyses of medium to low-resolution spectra, the least squares (χ^2 minimization), or “euclidian distance”, also called minimum distance method, such as Université de Lyon Spectroscopic Analysis Software (ULySS, Koleva et al., 2009), and the k-means clustering described in Sánchez Almeida and Allende Prieto (2013), are employed.

In the present work we analyse spectra in the optical, in the range $4560\text{--}5860 \text{ \AA}$, obtained at the FORS2/VLT at a resolution $R \sim 2,000$, carrying out full spectrum fitting. This spectral region, in particular from $\text{H}\beta$ to Na I lines, is sensitive to metallicity and temperature, to gravity due to MgH molecular bands (as part of the Mg_2 index), and it

includes the Lick indices Fe5270, Fe5335 and Mg₂, that are usual Fe and Mg abundance indicators (Katz et al., 2011; Cayrel et al., 1991; Faber et al., 1985; Worthey et al., 1994).

The same sample was observed in the near-infrared (CaT), as presented in Saviane et al. (2012), Da Costa et al. (2009) and Vasquez et al. (in prep), where two among the triplet Ca II lines were used to derive velocities and metallicities. A comparison of their results with the present ones show good consistency, as will be discussed in the present paper.

In this work we study six reference globular clusters, spanning essentially the full range of metallicities of globulars: the metal-poor halo clusters NGC 6426 and Terzan 8 ([Fe/H] $\sim$ -2.1 and -2.2, respectively), the moderately metal-poor NGC 6558 ([Fe/H] $\sim$ -1.0) in the bulge, the template “disc” metal-rich cluster M 71 (NGC 6838, [Fe/H] $\sim$ -0.7), and the metal-rich bulge clusters NGC 6528 and NGC 6553 ([Fe/H] $\sim$ -0.1 and -0.2, respectively).

These reference clusters are analysed with the intent of testing and improving the method, and verifying the metallicity range of applicability of each library of template spectra. In all cases, member stars and surrounding field stars are analysed. For some of these clusters previous high-resolution spectroscopic and photometric data of a few member stars are available.

The minimum distance method was adopted by Cayrel et al. (1991), by measuring residuals in each of the stellar parameters effective temperature, gravity, and metallicity; the method required the input of reference parameters. In the present work, we adopt the code ETOILE (Katz et al., 2011) that uses the minimum distance method, where the reliability and coverage of T_{eff} , $\log(g)$, [Fe/H], [α /Fe] of the template stars are important to find well-founded parameters for the target stars. We adopted two different libraries of spectra, the MILES¹ library of low-resolution spectra ($R \sim 2,000$) and the grid of synthetic spectra by Coelho et al. (2005)².

In Sect. 2 the observations are described. In Sect. 3 the method of stellar parameter derivation is detailed. In Sect. 4 the method is applied to six cluster as a validation of the procedures. In Sect. 5 the results are discussed, and in Sect. 6 a summary is given.

¹ <http://miles.iac.es/>

² http://www.mpa-garching.mpg.de/PUBLICATIONS/DATA/SYNTHSTELLIB/synthetic_stellar_spectra.html

3.3 Observations and data reduction

We observed respectively 17, 17, 12, 17, 10 and 13 red giant stars of the globular clusters NGC 6528, NGC 6553, M 71, NGC 6558, NGC 6426, Terzan 8, and surrounding fields, using FORS2@VLT/ESO, under projects 077.D-0775(A) and 089.D-0493(B). Table 3.1 summarizes the setup of the observations. Pre-images were taken using filters Johnson-Cousins V and I in order to select only stars in the red giant branch (RGB) brighter than the Red Clump (RC) level. Zero points in colours and magnitudes were fitted to match isochrones with parameters from Table 3.2 (see Colour-Magnitude Diagrams, CMDs, in Figure 3.1). We selected stars covering the whole interval in colour of the RGB, and when possible trying to avoid Asymptotic Giant Branch (AGB) stars. These stars are spatially distributed as shown in Figure 3.2, partly due to the slitlet configuration. Cluster parameters and log of observations are given in Table 3.2. The list of individual stars, their coordinates and VI magnitudes from the present FORS2 observations are given in Table B.1.

Table 3.1 - Telescope and spectrograph

Observing information	
Telescope	Antu/UT1-VLT@ESO
Instrument	FORS2
Grism	1400V
FoV	$6.8' \times 6.8'$
Pixel scale	$0.25''/\text{pixel}$
Slit width	0.53 mm
Spec. resolution	2,000

Table 3.2 - Log of observations and clusters parameters from literature. Main reference is Harris (1996, 2010 edition), when not indicated explicitly.

Parameter	NGC 6528	NGC 6553	M 71	NGC 6558	NGC 6426	Terzan 8
Date of obs.	29.05.2006	29.05.2006	29.05.2006	29.05.2006	13.07.2012	12.07.2012
UT	08:36:22	08:57:50	09:14:32	06:55:32	02:31:12	07:47:29.346
τ	149.4 s	79.4 s	17.2 s	148.3 s	500.0 s	360 s
RA	18 ^h 04' 49.64"	18 ^h 09' 17.60"	19 ^h 53' 46.49"	18 ^h 10' 17.60"	17 ^h 44' 54.65"	19 ^h 41' 44.41"
DEC	-30° 03' 22.6"	-25° 54' 31.3"	+18° 46' 45.1"	-31° 45' 50.0"	+03° 10' 12.5"	-33° 59' 58.1"
age	13 Gyr ⁽¹⁾	13 Gyr ⁽¹⁾	11.00 ± 0.38 Gyr ⁽²⁾	14 Gyr ⁽³⁾	13.0 ± 1.5 Gyr ⁽⁴⁾	13.00 ± 0.38 Gyr ⁽²⁾
[Fe/H]	-0.11 dex	-0.18 dex	-0.78 dex	-0.97 ± 0.15 dex ⁽³⁾	-2.15 dex	-2.16 dex
[Mg/Fe] ^a or [α /Fe] ^b	0.24 ^(b,5)	0.26 dex ^(b,6)	0.19±0.04 ^(a,7) , 0.40 ^(b,5)	0.24 ^(a,3)	0.4 ^(b,4)	0.47 ± 0.09 ^(a,8)
E(B-V)	0.54	0.63	0.25	0.44	0.36	0.12
(m-M) _V	16.17	15.83	13.80	15.70	17.68	17.47
R _{Sun}	7.9 kpc	6.0 kpc	4.0 kpc	7.4 kpc	20.6 kpc	26.3 kpc
R _{GC}	0.6 kpc	2.2 kpc	6.7 kpc	1.0 kpc	14.4 kpc	19.4 kpc
<v _{helio} >	206.6 ± 1.4 km/s	-3.2 ± 1.5 km/s	-22.8 ± 0.2 km/s	-197.3 ± 4 km/s ⁽³⁾	-162.0 km/s	130.0 km/s
r _{core}	0.13'	0.53'	0.63'	0.03'	0.26'	1.00
r _{tidal}	4.11'	7.66'	8.90'	9.49'	13.03'	3.98
r _{half-light}	0.38'	1.03'	1.67'	2.15'	0.92'	0.95

¹Zoccali et al. (2001) ²VandenBerg et al. (2013) ³Barbuy et al. (2007) ⁴Dotter et al. (2011) ⁵Carretta et al. (2010) ⁶Cohen et al. (1999) ⁷Meléndez and Cohen (2009) ⁸Carretta et al. (2014)

Figure 3.1: Colour-Magnitude diagrams of all clusters analysed in the present work. Left panels are metal-rich clusters, middle panels are intermediate metallicity clusters and right panels correspond to the more metal-poor ones. All stars within $2 \times r_{half-light}$ are plotted, without any cleaning procedure. Dartmouth isochrones with literature parameters (Table 3.2) are overplotted. Selected RGB stars for spectroscopic observations are in red. Reddening vectors are shown in each CMD based on $E(B-V)$ listed in Table 3.2.

Figure 3.2: Sky map of all clusters analysed in the present work. Panels are displayed as in Figure 3.1. Only brightest stars are shown, and the size of the dots are scaled with the stars magnitudes as indicated in each plot. Selected RGB stars for spectroscopic observations are in red. The blue circle corresponds to the half-light radius of each cluster from Table 3.2.

The spectra were taken using the grism 1400V, centred at 5200 Å, covering the range 4560 - 5860 Å, with a resolution of $R \sim 2000$. Figure 3.3 illustrates the spectra of a metal-poor and a metal-rich red giant star, where many of the strongest lines are indicated.

Figure 3.3: Identification of most important atomic lines in each strong feature of spectra with FORS2 resolution ($R \sim 2000$). The most important molecular band in this region is MgH around $\lambda = 5165$ Å, on top of the Mg I triplet lines. Left panel shows a metal-poor red giant star from Coelho et al. (2005) library, and right panel shows a metal-rich one. The left panels were zoomed in on y-axis direction for better visualization.

The spectra were reduced using esorex/FORS2 pipeline³ with default parameters for bias and flatfield correction, spectra extraction, and wavelength calibration. The only modification relative to default parameters, has been the introduction of a list of skylines, since the default list had only one line. The wavelength calibration proved to be satisfactory with such line list (see Appendix C for further details). A last step in the reduction procedure was a manual removal of cosmic rays.

3.4 Stellar parameters determination

3.4.1 Radial velocities

Radial velocities were measured using the ETOILE code through cross correlation with a template spectrum from the chosen library. Tests were done in order to check the results, by measuring radial velocities using fxcor@IRAF (cross correlation), and rvidlines@IRAF

³ <http://www.eso.org/sci/software/pipelines/>

Figure 3.4: Average of heliocentric radial velocities of member stars (see details in Section 3.5.3 and values in Table B.1) for each globular cluster. Our results are plotted against those from CaT spectroscopy (Saviane et al., 2012) showing good agreement. Error bars are the standard deviation of the average. One-to-one line is plotted for visual guidance, where radial velocities from Harris (1996, 2010 edition) are overplotted as empty circles.

(using wavelength of MgI triplet lines as a reference). The derived velocities are consistent, therefore we used ETOILE also to determine radial velocities. Heliocentric radial velocities for each star can be found in Table B.1, where the last column refers to the values measured from the CaII triplet (CaT) lines in the near infrared by Saviane et al. (2012) for member stars for NGC 6528, NGC 6553, M 71 and NGC 6558, and by Vasquez et al. (2014 in prep.) for NGC 6426 and Terzan 8. There is good agreement between the present radial velocity values and those from the CaT line region. A few exceptions are cool stars presenting strong TiO molecular bands, e.g. #8 in NGC 6558. Radial velocities of stars #2, #10 of M 71 are also discrepant between CaT and our determinations. Average values for member stars in each cluster are presented in Figure 3.4, where our results are compared to CaT results (Saviane et al., 2012 and Vasquez et al. (priv. communication)), and with Harris (1996, 2010 edition) catalogue. Error bars from the literature are smaller than the empty

circles that represent literature v_{helio} , except for NGC 6426, as can be seen in Figure 3.4. Our results are in good agreement with both references. In particular, the radial velocity measured for NGC 6426 is in agreement between this work and CaT results based on individual member stars, but it is only compatible with the literature value within 3σ . The explanation is that the only work that measured radial velocities for this cluster was based on integrated light from photographic plates (Hesser et al., 1986). Therefore, the present radial velocity derivation for NGC 6426 is more reliable.

3.4.2 Atmospheric parameters

Full spectrum fitting with minimum distance method is employed, using the ETOILE code described in Katz et al. (2011) and Katz (2001). We apply the calculations to the wavelength region 4600-5600 Å, similarly to the procedure described in Katz et al. (2011).

Automated derivation of the atmospheric parameters (T_{eff} , $\log(g)$, $[\text{Fe}/\text{H}]$, $[\alpha/\text{Fe}]$) of a stellar spectrum is carried out by comparing the target spectrum with each library spectrum, thus covering a large range of atmospheric parameters. In each comparison, ETOILE fits a polynomial to both spectra, which compensates for continuum differences between the two spectra. The distance between the two spectra is calculated through weighted least-squares. The estimation of the parameters is done by taking an average of the most similar library spectra. This is called the similarity method introduced by Katz et al. (1998). For a more detailed explanation see Katz (2001).

Before running the code, two important steps are needed: to convolve all the library spectra to the same resolution of the target spectra, and to correct for radial velocities v_r . Convolution calculations were performed for the library spectra using the task GAUSS in IRAF. The code ETOILE measures the radial velocities by comparison with template spectra from the library, a reliable way to measure v_r in each observed spectrum and correct them.

Figure 3.5 shows two examples of spectral fitting, one for a metal-poor (Terzan8.11) and another for a metal-rich (NGC6528.11) star. The template stars that best fit these cluster stars among the available spectra from MILES library are BD+060648 and HD161074, respectively. The residuals shown at the bottom of these figures indicate that the metal-poor target spectrum is similar to the template spectrum within 2%, except for a few strong features. The residuals for the metal-rich star shows a similarity between target

Figure 3.5: Examples of spectral fitting carried out with ETOILE for a metal-poor (Terzan8.11) and a metal-rich (NGC6528.11) star displayed in the upper and bottom panels, respectively. For each star, its spectrum (black line) is overplotted by the MILES spectrum (red line) that best fits it, BD+060648 and HD161074, respectively. Below each stellar spectrum the residuals of each fit are presented. The match between the spectra is done following the procedures explained in Section 3.4.2. The fit appears very satisfactory for the whole wavelength interval.

and template spectra of 5%, except for the boundaries $\lambda \gtrsim 5700$ Å and $\lesssim 4700$ Å, and for a few strong features. These differences between the spectra are reflected in the atmospheric parameters, and they are compensated by taking the average of parameters of the most similar spectra. For Terzan8.11, there are 8 MILES spectra close enough which were averaged, and for NGC6528.11, 21 MILES spectra were considered. Details on the criterion

to select the number of template spectra are discussed in Section 3.4.2.2.

3.4.2.1 *Stellar libraries*

The core of the atmospheric parameters derivation in this work is the choice of a stellar library. There are two classes of stellar libraries: based on observed or synthetic spectra. The real spectra are more reliable, but the drawback is that they have abundances typical of nearby stellar populations. The synthetic libraries have no noise, and a large and uniform coverage of the atmospheric parameters space, however there are still limitations on the completeness of atomic and molecular line lists, plus uncertainties on oscillator strengths, and assumptions on atmospheric models, such as 1-D and local thermodynamical equilibrium. For these reasons, it is useful to use both observational and synthetic libraries. In the present work, we use two libraries, one observed and one synthetic, as described below:

The MILES library (Sánchez-Blázquez et al., 2006) has 985 stellar spectra with resolution $R \sim 2,200$, and mean signal-to-noise ratio of 150 per pixel for field and open cluster stars, and 50 for globular cluster stars. Atmospheric parameters coverage is (Cenarro et al., 2007; Milone et al., 2011):

$$\begin{aligned} 352.5\text{nm} &< \lambda < 750\text{nm} \\ 2,747\text{K} &< T_{\text{eff}} < 36,000\text{K} \\ -0.20 &< \log(g) < 5.50 \\ -2.86 &< [\text{Fe}/\text{H}] < +1.65 \\ -0.54 &< [\text{Mg}/\text{Fe}] < +0.74 \end{aligned}$$

The COELHO library (Coelho et al., 2005) has 6367 synthetic stellar spectra⁴ with wavelength steps of $0.02\text{\AA}$ (resolution $R=525,000$). Atmospheric parameters coverage is:

⁴ Interpolation on the original library was carried out to produce spectra with $[\alpha/\text{Fe}] = 0.1, 0.2, 0.3$ dex from the provided 0.0 and 0.4 dex spectra.

Figure 3.6: HR diagram showing the parameter space available from the MILES library. Dartmouth isochrones (Dotter et al., 2008) are overplotted for metallicities close to those of the six analysed globular clusters: $[\text{Fe}/\text{H}] = -0.1, -0.8, -1.3, -2.2$ dex, with $[\alpha/\text{Fe}] = 0.2, 0.4, 0.4, 0.4, 0.4$ dex, respectively, and ages = 13 Gyr for all cases. Colours of the dots indicate the metallicity range closer to the isochrones. Dashed black lines are the adopted limit to select RGB stars from MILES used as reference for the fits.

$$300\text{nm} < \lambda < 1800\text{nm}$$

$$3,500\text{K} < T_{\text{eff}} < 7,000\text{K}$$

$$0.0 < \log(g) < 5.0$$

$$-2.5 < [\text{Fe}/\text{H}] < +0.5$$

$$0.0 < [\alpha/\text{Fe}] < +0.4$$

where α -elements considered in this library are: O, Mg, Si, S, Ca and Ti.

Once all cluster stars are located in the red giant branch, as shown in Figure 3.1, we selected only stars in this region in the parameters space of the libraries (see Figure 3.6) to avoid non-physical results.

3.4.2.2 Average results and errors: validation with well-known stars

We define different criteria for MILES and COELHO libraries for taking the average of stellar parameters from reference spectra, as mentioned in Section 3.4.2. For MILES the average results are based on different numbers of templates depending on the sampling as shown in Figure 3.6. For the synthetic library COELHO the sampling is homogeneous, therefore a constant number of templates is adopted. We found that 10 templates for COELHO cover satisfactorily the variations in the four stellar parameters (T_{eff} , $\log(g)$, $[\text{Fe}/\text{H}]$ and $[\alpha/\text{Fe}]$). The COELHO library was built by varying all alpha-elements (O, Mg, Si, S, Ca, Ti) together, therefore $[\alpha/\text{Fe}]$ is an average of the effect from enhancement of these element abundances. In this spectral region Mg lines are the dominant source of line strengthening.

The criterion to define average results from the MILES library is more complex, as follows. The code provides a list of the closest reference spectra from the library, ranked by the similarity parameter (S , see Katz et al., 2011). The final parameters T_{eff} , $\log(g)$, $[\text{Fe}/\text{H}]$ and $[\text{Mg}/\text{Fe}]$ are the average of the parameters of first N reference stars from the ETOILE output, where N depends on the sampling of the library for each combination of parameters. The average is weighted by $1/S^2$ as shown in the equation below for T_{eff} (the same is valid for the other three parameters):

$$T_{\text{eff}}(N) = \frac{\sum_{i=1}^N T_{\text{eff},i} \times \frac{1}{S_i^2}}{\sum_{i=1}^N \frac{1}{S_i^2}} \quad (3.1)$$

The errors are defined as the average of the squared residuals, weighted by $1/S^2$, as shown in the equation below for T_{eff} (the same is valid for the other three parameters). For $N=1$, we adopted the same error of $N=2$.

$$\sigma_{T_{\text{eff}}}(N) = \sqrt{\frac{\sum_{i=1}^N (T_{\text{eff},i} - T_{\text{eff}})^2 \times \frac{1}{S_i^2}}{\sum_{i=1}^N \frac{1}{S_i^2}}} \quad (3.2)$$

To estimate the number of reference stars to be averaged in each case, we proceeded with some tests using 59 spectra of 49 well-known stars, listed in Table B.2. These stars were selected among red giant stars (same $\log(g)$ and T_{eff} intervals defined in Figure

3.6) presented in the ELODIE library⁵ (Prugniel et al., 2007). Reference atmospheric parameters were averaged from the PASTEL catalogue (Soubiran et al., 2010), and the quality filter was determined by a threshold in the standard deviation: $\sigma_{T_{\text{eff}}} < 200\text{K}$, $\sigma_{\log(g)} < 0.5$, $\sigma_{[\text{Fe}/\text{H}]} < 0.2$. We calculated the average parameters and respective errors for different N and compared the results with the average values of T_{eff} , $\log(g)$, $[\text{Fe}/\text{H}]$ from the PASTEL catalogue (Soubiran et al., 2010). We minimize the equation below to find the best N that will give the final parameters and respective errors. This equation considers the distance between the average for a given N and literature average; in this way all the three parameters are minimized at the same time. Milone et al. (2011) have measured $[\text{Mg}/\text{Fe}]$ for MILES spectra, therefore it is possible to take averages for this parameter as a function of N , and use $[\text{Mg}/\text{Fe}]$ for the best N as an estimation of the α -enrichment for each star.

$$\text{RR}_{\text{tot}}(N) = \sqrt{\text{RR}_N(T_{\text{eff}})^2 + \text{RR}_N(\log(g))^2 + \text{RR}_N([\text{Fe}/\text{H}])^2} \quad (3.3)$$

where $\text{RR}_N(T_{\text{eff}})$ is given by the equation below (the same is valid for the other three parameters):

$$\text{RR}_N(T_{\text{eff}}) = \frac{T_{\text{eff}}(N) - T_{\text{eff}}^{(\text{lit})}(N)}{T_{\text{eff}}^{(\text{lit})}(N)} \quad (3.4)$$

Figure 3.7 illustrates the finding procedure of N for the case of star HD122956, showing that ETOILE could recover all the four parameters accurately. The resulting parameters, RR_{tot} , N , and literature values are presented in Table B.2. Different stars need different number N of templates to find the best result. Moreover the ratio $S(N)/S(1)$ for the best N is roughly constant for all ETOILE template spectra, with an average value of 1.1 ± 0.1 . The best number N and the respective ratio $S(N)/S(1)$ are related to the library sampling, for example, for a given star with best $N=1$ it means that there is only one reference star with $S(N)/S(1) \lesssim 1.1$, and there are two possible explanations: either the target star matches perfectly some reference star, or the library has no other reference spectra similar enough to that star to be considered. In the cases with $N=15$, for instance, the library has 15 reference spectra very similar ($S(N)/S(1) \lesssim 1.1$) to the target spectra, and their parameters must be averaged in order to get the parameters for the target star.

⁵ http://www.obs.u-bordeaux1.fr/m2a/soubiran/elodie_library.html

Figure 3.7: Finding procedure of N for the case of star HD122956, based on the minimization of the residuals of T_{eff} , $\log(g)$ and $[\text{Fe}/\text{H}]$ (equation 3.3). Circles represent the averages of the parameters of N best reference stars. Filled black circle indicates the closest values to the references. Blue solid lines and blue dashed lines are average from PASTEL catalogue and standard deviation (Table B.2). For this star, Fulbright (2000) published $[\text{Mg}/\text{Fe}]$ and we compare also with the averages as a function of N. All the four parameters for the best $N=3$ are compatible with literature.

All results are plotted in Figure 3.8 showing the good agreement of ETOILE results and PASTEL catalogue average for T_{eff} , $\log(g)$, $[\text{Fe}/\text{H}]$ in the whole range for RGB stars analysed in this work. The behaviour of the derived values of $[\text{Mg}/\text{Fe}]$ vs. $[\text{Fe}/\text{H}]$ has a similar behaviour to field stars (see e.g. Figure 6 of Alves-Brito et al., 2010).

After these tests we can consider that ETOILE code together with the MILES library works well for low-resolution spectra of red giant stars in the optical region. Additionally we define the criterion to consider a reference spectrum similar enough to be considered in the average of the parameters as $S(N)/S(1) \leq 1.1$.

Figure 3.8: Comparison of the parameters determined in this work with PASTEL catalogue average values for the well-known stars presented in Table B.2. These plots endorse the usage of ETOILE code for atmospheric parameters determination for red giant stars in the optical spectral region. Only stars with $\sigma_{T_{\text{eff}}} < 200\text{K}$, $\sigma_{\log(g)} < 0.5$, $\sigma_{[\text{Fe}/\text{H}]} < 0.2$ from PASTEL catalogue were selected as good quality candidates for validation of the method.

3.5 Results

The derived T_{eff} , $\log(g)$, $[\text{Fe}/\text{H}]$, $[\text{Mg}/\text{Fe}]$ and $[\alpha/\text{Fe}]$ are presented in Table B.3. In order to discuss these results, we proceed as follows: in Sect. 3.5.1 we plot T_{eff} and $\log(g)$ for stars in each cluster together with isochrones of age and metallicity given in Table 3.2. Section 3.5.2 compares $[\text{Fe}/\text{H}]$ with CaT results from Saviane et al. (2012) and Vasquez et al. (in prep.). Subsequently all checked parameters are used to select member stars for each cluster (Section 3.5.3). Finally, all parameters for member stars are compared individually with high-resolution analysis, when available in the literature. M 71 has three stars in common with Cohen et al. (2001), NGC 6558 has three stars in common with Barbuy et al. (2007) and Terzan 8 has four stars in common with Carretta et al. (2014), as

described in Sections 3.5.4.1, 3.5.4.2 and 3.5.4.3, respectively. For NGC 6528, NGC 6553 and NGC 6426 we did not find any star in common with high-resolution spectroscopic studies.

3.5.1 T_{eff} , $\log(g)$ against isochrones

In high-resolution spectroscopy studies, usually T_{eff} is estimated from photometry and $\log(g)$ from theoretical equations⁶. These parameters are employed as initial guesses to derive $[\text{Fe}/\text{H}]$, which is applied to redetermine T_{eff} and $\log(g)$ iteratively, until reaching a convergence of the three parameters. In this work we fit all the three parameters at the same time (Section 3.4.2), and a check on these parameters is carried out as explained below.

Figure 3.9 displays the results of all stars in the six clusters in a Hertzsprung-Russell diagram form. Left, middle and right panels show the results using MILES library, CO-ELHO library and the average of both results, respectively. Black dots represent member stars of each cluster, and grey dots are not members, based on the selection described in the next Section 3.5.3. Dartmouth isochrones (Dotter et al., 2008) with age, $[\text{Fe}/\text{H}]$ and $[\alpha/\text{Fe}]$ from Table 3.2 are overplotted in the diagrams of Figure 3.9 in blue. Cyan lines have the same age and $[\text{Fe}/\text{H}]$ as the respective blue lines, but with the extreme values of $[\alpha/\text{Fe}] = -0.2$ and $+0.8$, available from the models.

The results on T_{eff} and $\log(g)$ from MILES and COELHO are in good agreement with the isochrones. We also computed an average of the results weighted by their uncertainties that are displayed in the right panels of Figure 3.9. We adopted as the final results in this work the weighted average of MILES and COELHO results.

3.5.2 Comparison with $[\text{Fe}/\text{H}]$ derived from CaT

All selected stars from photometry were observed both in the near-infrared (CaT, Saviane et al., 2012 and Vasquez et al. in prep) and in the optical spectral region. Metallicities from CaT were calibrated by Saviane et al. (2012), to be in an homogeneous scale. The optical region studied here is suitable to provide robust values of $[\text{Fe}/\text{H}]$ for each cluster, to be compared with the CaT value, and to converge ultimately to the average $[\text{Fe}/\text{H}]$ for each cluster.

⁶ $\log(g) = 4.44 + 4\log\frac{T}{T_{\odot}} + 0.4(M_{\text{bol}} - 4.75) + \log\frac{M}{M_{\odot}}$, see for example Barbuy et al. (2009)

Figure 3.9: Comparison of T_{eff} and $\log(g)$ of stars in each cluster with Dartmouth isochrones (Dotter et al., 2008). For each cluster we show the results based on MILES and COELHO libraries and a third panel with the weighted average of the results from both libraries. The parameters of age, $[\text{Fe}/\text{H}]$ and $[\alpha/\text{Fe}]$ for the blue thick isochrones were taken from Table 3.2. Cyan thin isochrones have same age and $[\text{Fe}/\text{H}]$ as blue ones, but with the limits $[\alpha/\text{Fe}] = -0.2\text{dex}$ and $+0.8\text{dex}$. Black dots represent member stars of each cluster, and grey dots are not members.

Figure 3.10 shows the comparisons of the metallicity values presented in Table B.3 with those from CaT analysis. Upper left panel shows that $[\text{Fe}/\text{H}]$ using MILES library is in good agreement with CaT results for the three most metal-rich clusters NGC 6528, NGC 6553 and M 71. This is in agreement with the sampling of metal-rich stars for all combinations of T_{eff} and $\log(g)$ as displayed in Figure 3.6 in red and green. For NGC 6558 with $[\text{Fe}/\text{H}] \sim -1.0$ the dispersion on the parameters is larger which is explained by the smaller number of stars available in the library with such metallicity. MILES is based on the solar neighbourhood showing therefore only a few stars with $[\text{Fe}/\text{H}] \sim -1.0$. For the metal-poor clusters NGC 6426 and Terzan 8 the library sampling is even more sparse, as becomes evident in Figure 3.6. In this case, the average of parameters from the library takes into account some more metal-rich reference stars which results in higher values of $[\text{Fe}/\text{H}]$ for NGC 6426 and Terzan 8 stars.

Metallicities using COELHO library are compared with CaT results in the upper right

Figure 3.10: Comparison of $[\text{Fe}/\text{H}]$ from this work (Table B.3) with those from equivalent widths of Ca II triplet for the same stars with the same instrument by Saviane et al. (2012) and Vazquez et al. (in prep). Upper panels compare CaT metallicities with those obtained with MILES and COELHO libraries. Bottom panels are two types of combination of the results: average of each star on the left, and assuming MILES results for more metal-rich and COELHO results for more metal-poor stars, on the right.

panel of Figure 3.10. The synthetic spectra reproduce less well the metal-rich stars, given the missing opacity due to millions of very weak lines, not taken into account in the calculations; this blanketing effect lowers the continuum in real stars, and the measurable lines are shallower than in the present synthetic spectra calculations by Coelho et al. Because of this effect, stars of NGC 6528, NGC 6553 and M 71 are more metal-poor than CaT results. On the other hand, COELHO library is suitable to reproduce the stars of the three more metal-poor clusters of this sample, NGC 6558, NGC 6426 and Terzan 8.

To summarize, for the three more metal-rich clusters, MILES results are better, and for the other three, COELHO results are preferable. The bottom right panel of Figure 3.10 shows the concatenation of this conclusion, i.e., it displays MILES results for NGC 6528,

NGC 6553 and M 71, and COELHO results for NGC 6558, NGC 6426 and Terzan 8. An alternative combination of results from MILES and COELHO is to take the average of the results weighted by their uncertainties. This average combines the best of both libraries and gives a good correlation with CaT results, as shown in the bottom left panel of Figure 3.10. Both criteria to combine MILES and COELHO (two bottom panels) are in good agreement with CaT results, and we adopted $[\text{Fe}/\text{H}]$ from the average results represented in the bottom left panel.

3.5.3 Membership selection

In Figure 3.11 we show four panels for each globular cluster. In upper left panels radial velocities against metallicities are displayed. Blue solid lines are parameters from Table 3.2 and red dashed lines are the average of parameters for cluster members only. For all clusters there is a clear concentration of stars around literature values, and these stars were selected as cluster members. Upper right panels show T_{eff} vs. $\log(g)$ compared with Dartmouth isochrones (Dotter et al., 2008) as done in Sect. 3.5.1. Blue solid lines are isochrones with parameters from Table 3.2, and red dashed lines are based on the average (Table 3.5) of the parameters for member stars (Table B.3). All member stars are close to the isochrones while non-member stars are out of these lines, confirming the membership selection. Bottom left panels shows $[\text{Fe}/\text{H}]$ against distance to the cluster centre, and no trends were found for any sample cluster. Bottom right panels shows no correlation between T_{eff} and $[\text{Fe}/\text{H}]$ for any of the six clusters analysed in this work.

In conclusion all T_{eff} , $\log(g)$ and $[\text{Fe}/\text{H}]$ values are found to sit in well defined sequences following the isochrones. Bottom-right panels show no correlation between $[\text{Fe}/\text{H}]$ and T_{eff} , suggesting that the $[\text{Fe}/\text{H}]/T_{\text{eff}}$ degeneracy does not affect our fitting procedure. Further evidence of this is provided by the comparisons of our results with $[\text{Fe}/\text{H}]$ measures from high-resolution analysis available in the literature, presented in Sections 3.4.2.2, 3.5.4.1, 3.5.4.2 and 3.5.4.3.

3.5.4 Validation with high-resolution spectroscopy

We found stars in common with literature high-resolution spectroscopy for three clusters: M 71, NGC 6558 and Terzan 8. In Sect. 3.5.3 we were able to identify ten member stars of M 71, five member stars of NGC 6558, and twelve member stars in Terzan 8, these

Figure 3.11: Step-by-step of selection of member stars for each cluster (four panels per cluster). Black dots are selected member stars, grey dots are non-members and green circles shows stars considered as cluster member by Saviane et al. (2012) but non-members in the present work. Blue solid lines are drawn based on values of Table 3.2, which were also applied to the isochrones from Dotter et al. (2008). Red dashed lines refer to the weighted average of the member stars parameters.

Table 3.5 - Final average parameters for member stars in each globular cluster, and respective internal errors.

Cluster	$\langle v_{\text{helio}} \rangle$ (km/s)	$\langle [\text{Fe}/\text{H}] \rangle^{(a)}$	$\langle [\text{Fe}/\text{H}] \rangle^{(b)}$	$\langle [\text{Fe}/\text{H}] \rangle^{(avg)}$	$\langle [\text{Mg}/\text{Fe}] \rangle^{(a)}$	$\langle [\alpha/\text{Fe}] \rangle^{(b)}$
NGC 6528	185 ± 10	-0.07 ± 0.10	-0.18 ± 0.08	-0.13 ± 0.05	0.05 ± 0.09	0.26 ± 0.05
NGC 6553	6 ± 8	-0.125 ± 0.009	-0.55 ± 0.07	-0.133 ± 0.017	0.107 ± 0.009	0.302 ± 0.025
M 71	-42 ± 18	-0.48 ± 0.08	-0.77 ± 0.08	-0.63 ± 0.15	0.25 ± 0.07	0.293 ± 0.032
NGC 6558	-210 ± 16	-0.88 ± 0.20	-1.02 ± 0.05	-1.012 ± 0.013	0.26 ± 0.06	0.23 ± 0.06
NGC 6426	-242 ± 11	-2.03 ± 0.11	-2.46 ± 0.05	-2.39 ± 0.11	0.38 ± 0.06	0.24 ± 0.05
Terzan 8	135 ± 19	-1.76 ± 0.07	-2.18 ± 0.05	-2.06 ± 0.17	0.41 ± 0.04	0.21 ± 0.04

^aMILES library ^bCOELHO library ^{avg}Average of MILES and COELHO results

being the same selected by Saviane et al. (2012), and Vazquez et al. (in preparation). The derived stellar parameters are reported in Table B.3 for member and non-member stars.

3.5.4.1 M 71

Cohen et al. (2001) observed 25 member red giant stars of M 71 using HIRES@Keck ($R \sim 34,000$), and derived their T_{eff} and $\log(g)$. In two subsequent papers, they derived $[\text{Fe}/\text{H}]$ (Ramírez et al., 2001) and $[\text{Mg}/\text{Fe}]$ (Ramírez and Cohen, 2002) for them. We have three stars in common that are presented in Table 3.7. Temperature and gravity values are compatible within 0.5 to 2- σ , $[\text{Fe}/\text{H}]$ and $[\text{Mg}/\text{Fe}]$ are compatible within 0.1 to 1.5- σ .

Table 3.7 - Final atmospheric parameters for the three stars of M 71 in common with Cohen et al. (2001), and also their determinations for the respective parameters.

Star	T_{eff} (K)	$\log(g)$	$[\text{Fe}/\text{H}]$	$[\text{Mg}/\text{Fe}]$
	$T_{\text{eff}}\text{-C01}$ (K)	$\log(g)\text{-C01}$	$[\text{Fe}/\text{H}]\text{-C01}$	$[\text{Mg}/\text{Fe}]\text{-C01}$
M71_7	3997 ± 89	1.53 ± 0.35	-0.58 ± 0.17	0.15 ± 0.18
1-45	3950	0.9	-0.60 ± 0.03	0.43 ± 0.09
M71_9	4316 ± 87	1.97 ± 0.33	-0.76 ± 0.17	0.27 ± 0.21
1-64	4200	1.35	-0.61 ± 0.03	0.43 ± 0.09
M71_13	4808 ± 106	2.82 ± 0.24	-0.63 ± 0.18	0.23 ± 0.20
G53476_4543	4900	2.65	-0.61 ± 0.03	0.36 ± 0.06

3.5.4.2 NGC 6558

Barbuy et al. (2007) observed six RGB stars using the high-resolution ($R \sim 22,000$) spectrograph FLAMES+GIRAFFE@VLT/ESO, and derived T_{eff} , $\log(g)$, $[\text{Fe}/\text{H}]$, and $[\text{Mg}/\text{Fe}]$

for each of them. We have three stars in common with their sample: #6, #8, #9, corresponding to their identification as B11, F42, F97, respectively (see Table 3.8). For stars #6 and #9, full spectrum fitting recovers all parameters within $1\text{-}\sigma$. Star #8 is a more complicated case because it is a very cool star ($T_{\text{eff}} < 4000\text{ K}$) and molecular bands of TiO are important. They change a lot the continuum which is not fitted perfectly. In fact, the derived parameters for this star led us to select it as non-member. Although temperature agrees with that from Barbuy et al. (2007), the gravity is much lower than their results.

These results show that full spectrum fitting method is reliable, consistent among all libraries, and present reasonable errors for RGB stars hotter than $\sim 4000\text{ K}$. Stars cooler than that must be analysed with a better suited reference library, containing a sufficient number of cool stars at all metallicities.

Table 3.8 - Final atmospheric parameters for the three stars of NGC 6558 in common with Barbuy et al. (2007), and also their determinations for the respective parameters.

Star	T_{eff} (K)	$\log(g)$	[Fe/H]	[Mg/Fe]
	$T_{\text{eff-B07}}$ (K)	$\log(g)\text{-B07}$	[Fe/H]-B07	[Mg/Fe]-B07
6558_6	4899 $\pm$ 162	2.68 $\pm$ 0.39	-1.11 $\pm$ 0.20	0.22 $\pm$ 0.07
B11	4650	2.2	-1.04	0.20
6558_8	3565 $\pm$ 59	1.05 $\pm$ 0.36	-0.16 $\pm$ 0.06	0.23 $\pm$ 0.00
F42	3800	0.5	-1.01	0.30
6558_9	4972 $\pm$ 168	2.68 $\pm$ 0.46	-1.11 $\pm$ 0.18	0.41 $\pm$ 0.16
F97	4820	2.3	-0.97	0.23

3.5.4.3 Terzan 8

Carretta et al. (2014) observed six stars with UVES@VLT/ESO ($R \sim 45,000$) and 14 with GIRAFFE@VLT/ESO ($R \sim 22,500 - 24,200$), with among them four stars in common with our FORS2@VLT/ESO sample. Their parameters for these stars are presented in Table 3.9. For temperature and gravity the compatibility is within 1 to $3\text{-}\sigma$, except for T_{eff} of star Ter8_8 which is in the limit of $3.9\text{-}\sigma$ of distance. For [Fe/H] all stars have compatible values with Carretta et al. (2014) within $1\text{-}\sigma$, except for star Ter8_1 which is in the limit of $3.9\text{-}\sigma$ of distance. [Mg/Fe] is compatible within $1\text{-}\sigma$.

Differently from M 71 and NGC 6558, the comparison between our results and Carretta et al. (2014) for Terzan 8 give all three parameters T_{eff} , $\log(g)$ and [Fe/H] systematically larger. For this reason we inspected the $T_{\text{eff}}\text{-}\log(g)$ diagram of both sets of data, compared

Table 3.9 - Final atmospheric parameters for the four stars of Terzan 8 in common with Carretta et al. (2014), together with their determinations for the respective parameters.

Star	T_{eff} (K)	$\log(g)$	[Fe/H]	[Mg/Fe]
	$T_{\text{eff}}\text{-C14}$ (K)	$\log(g)\text{-C14}$	[Fe/H]-C14	[Mg/Fe]-C14
Ter8_1	5067 ± 314	2.24 ± 0.30	-1.97 ± 0.14	0.40 ± 0.13
2913	4628	1.49	-2.52 ± 0.07	0.58
Ter8_4	4354 ± 88	0.67 ± 0.31	-2.28 ± 0.13	0.40 ± 0.14
2357	4188	0.66	-2.29 ± 0.10	0.48 ± 0.14
Ter8_8	5151 ± 108	2.56 ± 0.36	-2.06 ± 0.19	0.40 ± 0.18
2124	4730	1.67	-2.28 ± 0.26	0.56
Ter8_9	4564 ± 94	1.12 ± 0.30	-2.24 ± 0.17	0.42 ± 0.13
1658	4264	0.80	-2.40 ± 0.07	0.51 ± 0.02

Figure 3.12: Diagram $\log(g)$ vs. T_{eff} for Terzan 8 showing present (blue points with error bars) and Carretta et al. (2014, black filled circles) results, overplotted with Dartmouth, PARSEC and BASTI isochrones for ages = 13 Gyr, and metallicities indicated on the Figure.

with the Dartmouth (Dotter et al., 2008), PARSEC (Bressan et al., 2012) and BASTI (Pietrinferni et al., 2004) isochrones, as shown in Figure 3.12. The Carretta et al. (2014) results are compatible with an isochrone of $[\text{Fe}/\text{H}] = -1.7$, $[\alpha/\text{Fe}] = +0.4$ and 13 Gyr, whereas the present results fit better with an isochrone $[\text{Fe}/\text{H}] = -2.2$, $[\alpha/\text{Fe}] = +0.4$ and 13 Gyr.

Except for star Terzan8_4 showing very similar gravities, for the other stars the difference could come from a different mass/distance/bolometric correction and is probably within the gravity uncertainty. Over-ionization at low temperature atmospheres makes it hard to get a reliable gravity from Fe I/Fe II lines.

3.6 Discussion

Results for individual stars in each cluster (Table B.3) and average results (Table 3.5) are discussed below and compared with literature results. Figure 3.13 displays the comparison of our $[\text{Fe}/\text{H}]$ results for each cluster with reference values, showing good agreement for the whole range of metallicities from $[\text{Fe}/\text{H}] = -2.5$ to solar. Figure 3.14 gives the comparison of the average results of $[\text{Fe}/\text{H}]$ and $[\text{Mg}/\text{Fe}]$ with abundances for field stars of the different Galactic components: bulge, thin/thick disc and inner/outer halo. We discuss case by case below.

Figure 3.13: Comparison of $[\text{Fe}/\text{H}]$ from this work (Table 3.5) with those from literature, as revised in Table 3.2. Error bars are the weighted average of the $[\text{Fe}/\text{H}]$ of member stars in each cluster, as presented in Table B.3. For NGC 6553 and NGC 6558 the error bars are not visible because they are too small (see Table 3.5).

Figure 3.14: Chemical evolution of the Milky Way based on field stars from bulge (Gonzalez et al., 2011), thin disc (Bensby et al., 2005), thick disc (Reddy et al., 2006), inner and outer halo (Roederer, 2009). Results from this work for globular clusters are overplotted. For NGC 6553 the point corresponds to $[\alpha/\text{Fe}]$ instead of $[\text{Mg}/\text{Fe}]$, as discussed in Section 3.6.2

3.6.1 Metal-rich clusters NGC 6528, NGC 6553

NGC 6528 and NGC 6553 are known as “twin clusters” (Ortolani et al., 1995) with almost identical CMDs, as shown in Figure 3.1. As given in Table 3.2 they have reddening $E(B-V) = 0.54$ mag and 0.63 mag, $[\text{Fe}/\text{H}] = -0.11$ and -0.18 , $[\text{Mg}/\text{Fe}] = 0.24$ and 0.26 , respectively. They are located in the Milky Way bulge, distant 0.6 kpc and 2.2 kpc from the Galactic centre, and in opposite “southern legs” of the X-shaped bulge (see Figure 3 of Saito et al., 2011). Figure 3.9 shows that ETOILE recovers parameters for member stars coherent with a simple stellar population represented by the isochrones. Although all the results are compatible between the libraries, error bars for MILES results are lower than from COELHO results. This synthetic library, on the other hand, gives better alpha-enhancement values compatible with average values from high-resolution studies ($[\alpha/\text{Fe}] = 0.26 \pm 0.05$ and 0.302 ± 0.025 , respectively, from Table 3.5). In this respect, MILES cannot

give an alpha-enhancement, since their metal-rich stars are basically solar neighbourhood stars, that have no alpha-enhancement for metal-rich stars (see Figure 2 in Milone et al., 2011). The results from MILES are $[\text{Mg}/\text{Fe}] = 0.05 \pm 0.09$ and 0.107 ± 0.009 , respectively. This is a particularity of the bulge, where stars are metal-rich and old. However, Zoccali et al. (2004) finds $[\text{Mg}/\text{Fe}] = +0.07$ from high-resolution spectroscopy of three stars of NGC 6528, which is compatible with MILES and not with COELHO. For NGC 6553, Cohen et al. (1999) finds $[\text{Mg}/\text{Fe}] = +0.41$ also from high-resolution spectroscopy of five stars, which is closer to COELHO results.

Figure 3.11 compares the final results (average of MILES and COELHO results) with isochrones with literature parameters (same as Figure 3.9) and an additional isochrone considering the parameters derived from this work (Table 3.5). The isochrones consider $[\alpha/\text{Fe}]$ from COELHO results as discussed above. We derived $[\text{Fe}/\text{H}] = -0.13 \pm 0.06$ and -0.133 ± 0.009 for NGC 6528 and NGC 6553, respectively, in agreement with high spectral resolution analyses of Carretta et al. (2001) and Alves-Brito et al. (2006).

3.6.2 Moderately metal-rich clusters M 71, NGC 6558

According to Harris (1996, 2010 edition), M 71 (or NGC 6838) is located 6.7 kpc from the Galactic centre, with a perpendicular distance to the Galactic plane of only 0.3 kpc towards the South Galactic Pole, which means that this globular cluster is located in the Milky Way disc. Because of that its reddening is high, although not as high as for most bulge clusters, with $E(\text{B}-\text{V}) = 0.25$. NGC 6558 is located only 1.0 kpc from the Galactic centre, in particular between the two “southern legs” of the X-shaped bulge (see Figure 3 of Saito et al., 2011). It has a reddening of $E(\text{B}-\text{V}) = 0.44$. Although they are located in different components of the Milky Way, these clusters share similar metallicities, $[\text{Fe}/\text{H}] \sim -0.8$, and $[\text{Fe}/\text{H}] \sim -1.0$, respectively.

Figure 3.1 shows a red horizontal branch (HB) for M 71 and a blue HB for NGC 6558, and this difference is not due to metallicity (Lee et al., 1994). It is true that a few other parameters can change the HB morphology at a fixed metallicity, as discussed by Catelan et al. (2001), for instance, but in this case the age is probably playing the major role. Literature ages are 11 Gyr (VandenBerg et al., 2013) and 14 Gyr old (Barbuy et al., 2007), respectively. M 71 is younger and moderately metal-rich, therefore a red horizontal branch is expected. In particular, we derived a slightly higher metallicity for this cluster

in comparison with the literature, $[\text{Fe}/\text{H}] = -0.63 \pm 0.06$. NGC 6558 has a high metallicity for such a blue horizontal branch, and Barbuy et al. (2007) argue that if this is interpreted as pure factor of age, this cluster should be one of the oldest objects in the Milky Way. We derived $[\text{Fe}/\text{H}] = -1.01 \pm 0.04$, compatible with their findings ($[\text{Fe}/\text{H}] = -0.97 \pm 0.15$), and more metal-rich than Harris (1996, 2010 edition) value of $[\text{Fe}/\text{H}] = -1.32$. Saviane et al. (2012) also found $[\text{Fe}/\text{H}] = -1.03 \pm 0.14$ for NGC 6558 from their Ca II triplet spectroscopy, with error being dominated by the external calibration uncertainty.

Comparing the error bars of T_{eff} and $\log(g)$ in Figure 3.9 between the two libraries, for M 71 they are of the same order, but for NGC 6558 MILES results present larger error bars. The main reason for this is that MILES library is based on solar neighbourhood stars, and only a few of them have metallicities $[\text{Fe}/\text{H}] \approx -1.0$ (see Figure 2 of Sánchez-Blázquez et al., 2006). Synthetic libraries such as that of Coelho et al. (2005) have spectra for any combination of atmospheric parameters evenly distributed in the parameter space. Therefore COELHO library is more suitable for the analysis of these moderately metallicity clusters.

For this range of metallicity and for expected values of $[\text{Mg}/\text{Fe}]$ from the literature (0.19 and 0.24, respectively), MILES and COELHO results are compatible, as shown in Table 3.5, $[\text{Mg}/\text{Fe}] = 0.25 \pm 0.07$ and 0.26 ± 0.06 , $[\alpha/\text{Fe}] = 0.293 \pm 0.032$ and 0.23 ± 0.06 , respectively. Ramírez and Cohen (2002) give an average $[\text{Mg}/\text{Fe}] = +0.37$ from 24 stars observed with high-resolution, which is compatible with our findings. The number of stars in MILES with these metallicities is low, but it is sampled enough for determinations of $[\text{Fe}/\text{H}]$ and $[\text{Mg}/\text{Fe}]$ for Milky Way stars. COELHO spectra have $[\alpha/\text{Fe}]$ varying from 0.0 to 0.4, which is also enough for this kind of objects.

3.6.3 Metal-poor clusters NGC 6426, Terzan 8

NGC 6426 and Terzan 8 present similar CMDs with a same literature age and metallicity of 13 Gyr (Dotter et al., 2011; VandenBerg et al., 2013) and $[\text{Fe}/\text{H}] = -2.15$ (Harris, 1996, 2010 edition). NGC 6426 is located in the northern halo of the Milky Way, 14.4 kpc from the Galactic centre and 5.8 kpc above the Galactic plane. Its height is much larger than the height scale of the thick disc (0.75 kpc, de Jong et al., 2010), but it has a non-negligible reddening of $E(B-V) = 0.36$, at a galactic latitude of $b = 16.23^\circ$. The best CMD available for this cluster was observed with the ACS imager onboard the Hubble

Space Telescope by Dotter et al. (2011), and they derived an age of 13.0 ± 1.5 Gyr from isochrone fitting. Our pre-image photometry based on observations with the Very Large Telescope/ESO produced a rather well-defined CMD for this cluster which is compatible with a 13 Gyr isochrone (see Figure 3.1).

Terzan 8 is located in the southern halo of the Milky Way, 19.4 kpc from the Galactic centre and 10.9 kpc below the Galactic plane. It has the lowest reddening of all clusters analysed in this work, $E(B-V) = 0.12$. This is one of the four Milky Way globular clusters believed to be captured from the Sagittarius dwarf galaxy (Ibata et al., 1994), the other three being M 54, Terzan 7 and Arp 2 (Da Costa and Armandroff, 1995). Carretta et al. (2014) did not find a strong evidence for Na-O anticorrelation, typical for globular clusters which may indicate that these clusters may have simple stellar populations.

Atmospheric parameters derived in this work for both clusters are in good agreement with literature values, when compared to the isochrones in Figure 3.9. As for the moderately metal-rich clusters, MILES results present larger error bars than COELHO results for T_{eff} and $\log(g)$, and the reason is the same as mentioned above, i.e., the sampling of MILES library is poorer for this metallicity range (see Figure 2 of Sánchez-Blázquez et al., 2006). Derived metallicities for these clusters are $[\text{Fe}/\text{H}] = -2.39 \pm 0.04$ and -2.06 ± 0.04 , respectively. For NGC 6426 our determination is 0.24 more metal-poor than given in Harris's catalogue ($[\text{Fe}/\text{H}] = -2.15$). However the only derivation of the metallicity for this cluster was done by Zinn and West (1984) based on integrated light ($[\text{Fe}/\text{H}] = -2.20 \pm 0.17$) which is compatible with our findings. The value from Harris catalogue was obtained by applying the Carretta et al. (2009) metallicity scale. We present for the first time a direct measurement of metallicity of individual red giant stars finding a more metal-poor value than previously attributed to this cluster. Terzan 8 is compatible with the Harris (2010 edition) catalogue of $[\text{Fe}/\text{H}] = -2.16$. There are three papers with metallicities for this cluster: Mottini et al. (2008) and Carretta et al. (2014), based on high-resolution spectra of individual star and average metallicity $[\text{Fe}/\text{H}] = -2.35 \pm 0.04$ and -2.27 ± 0.08 , respectively. The third paper is Da Costa and Armandroff (1995) who derived $[\text{Fe}/\text{H}] = -1.99 \pm 0.08$ based on CaII triplet spectroscopy. Our result is compatible with the two more metal-rich results.

For alpha-enhancement in this metallicity range, MILES spectra goes up to $[\text{Mg}/\text{Fe}] = 0.74$, while COELHO is limited to the models of $[\alpha/\text{Fe}] = 0.4$. The results based on

COELHO ($[\alpha/\text{Fe}] = 0.24 \pm 0.05$ and 0.21 ± 0.04 , respectively) are less enhanced than MILES ($[\text{Mg}/\text{Fe}] = 0.38 \pm 0.06$ and 0.41 ± 0.04 , respectively), the latter being closer to literature abundance ratios.

3.7 Summary and conclusions

We present a method of full spectrum fitting, based on the ETOILE code, to derive v_{helio} , T_{eff} , $\log(g)$, $[\text{Fe}/\text{H}]$, $[\text{Mg}/\text{Fe}]$ and $[\alpha/\text{Fe}]$ for red giant stars in Milky Way globular clusters. The observations were carried out with FORS2@VLT/ESO with resolution $R \sim 2000$.

We validated the method using well known red giant stars covering the parameter space of $4000 \text{ K} < T_{\text{eff}} < 6000 \text{ K}$, $0.0 < \log(g) < 4.0$, $-2.5 < [\text{Fe}/\text{H}] < +0.3$ and $-0.2 < [\text{Mg}/\text{Fe}] < +0.6$. The spectra of these reference stars are from the ELODIE library and the parameters from the PASTEL catalogue. The parameters were recovered for the whole range of parameters.

We selected six clusters from our large sample of observed clusters in order to establish the methodology. We chose two metal-rich (NGC 6528, NGC 6553), two moderately metal-rich (M 71, NGC 6558) and two metal-poor (NGC 6426 and Terzan 8) clusters. NGC 6528, NGC 6553 and NGC 6558 are located in the bulge, M 71 in the disc, NGC 6426 and Terzan 8 in the halo. For all clusters the effective temperatures and gravities are well determined using the MILES and Coelho et al. (2005) libraries of spectra. Metallicities and alpha-element enhancement are also derived with the caveats that for alpha-enhanced bulge clusters with $[\text{Fe}/\text{H}] > -0.5$, MILES is not suitable, since it has only solar neighbourhood stars, therefore in this case we use COELHO results because synthetic libraries have all combinations of parameters. For $[\text{Fe}/\text{H}] \approx -1.0$ MILES has few stars due to a lack of such stars in the solar vicinity. For metal-poor clusters, with high $[\alpha/\text{Fe}]$, MILES may be more indicated than COELHO because the latter is limited to $0 < [\alpha/\text{Fe}] < 0.4\text{dex}$, if such high Mg enhancements are confirmed.

The present results are in agreement with literature parameters available for five of the six template clusters. NGC 6426 is analysed for the first time using spectroscopy of individual stars. Therefore we provide a more precise radial velocity of $-242 \pm 11 \text{ km/s}$, a metallicity $[\text{Fe}/\text{H}] = -2.39 \pm 0.04$, and $[\text{Mg}/\text{Fe}] = 0.38 \pm 0.06$. The comparison of our results

of $[\text{Fe}/\text{H}]$ and $[\text{Mg}/\text{Fe}]$ to those from field stars from all Galactic components shows that the globular clusters follows the same chemical enrichment pattern as the field stars.

In conclusion, full spectrum fitting technique using ETOILE code together with MILES and COELHO libraries appears to be suitable to derive chemical abundances for Milky Way globular clusters, from low/medium-resolution spectra. Depending on the stellar population studied, the choice of library with parameter space covering the expected values for the clusters is a crucial ingredient, observed library being better for more metal-rich stars and synthetic library being preferable for the more metal-poor ones. This method will be applied to the other Milky Way globular clusters from this survey. It is also promising for extragalactic stars, that can be more easily observed with similar resolutions of $R \sim 2000$, for studies of galaxy formation and evolution.

FORS2/VLT survey of Milky Way globular clusters

II. Parameters for 51 clusters

4.1 Introduction to this chapter

The system of Galactic globular clusters (GGC) is of paramount importance to reconstruct the early evolution of the Milky Way: if their ages were known with sufficient precision, then correlations of their kinematics and chemical abundances with time would shed light on the dynamical and chemical evolution of the protogalactic halo and bulge. The present study allows a first order derivation of metallicity and alpha-enhancement from low-resolution spectra, together with radial velocities, and it provides a reliable target list for follow-up with high resolution spectroscopy. This is particularly important for the bulge clusters.

With the advent of multifiber spectrographs at 8m-class telescopes, high-resolution spectra of sufficient S/N can now be obtained for the giant stars of clusters out to $(m-M)_V \approx 19$, i.e. for $> 80\%$ of the GGCs. Nevertheless, the largest homogeneous samples of metallicities are still based on low-resolution spectroscopy (and calibrated versus high-resolution results). Even so, more than 50% of GGCs do not have any spectroscopic estimation of their $[\text{Fe}/\text{H}]$. Our survey improves this statistics targetting mostly distant and highly reddened clusters, which are poorly studied in the literature. We also observed some well-known clusters for calibration purposes, in a total of 51 objects, which is $\sim 1/3$ of the total number of catalogued Galactic globular clusters (see Saviane et al., 2012 for a review of $[\text{Fe}/\text{H}]$ values available in the literature).

Homogeneous determinations of $[\text{Fe}/\text{H}]$ and $[\alpha/\text{Fe}]$ for a large set of globular clusters are useful to analyse the chemical evolution of the Milky Way for its different components

(bulge, disc, inner and outer halo) and compare with field stars. The combination of these abundances with distance to the Galactic centre and ages leads to discussions about the origin of the globular clusters and it constrains models of Galaxy formation and evolution. Usually, $[\text{Fe}/\text{H}]$ for all globular clusters are averaged from different sources, using different methods and spectral resolution, and put them in a same scale. This procedure is useful to draw an overall picture of the metallicity distribution of our Galaxy, but it is not exactly an homogeneous set of $[\text{Fe}/\text{H}]$ and may contain some biases. In the present work we present for the first time a large set of $[\text{Fe}/\text{H}]$ and $[\text{Mg}/\text{Fe}]$ for the globular cluster population of the Milky Way, observed with the same setup, and analysed in a homogeneous way, finally calibrated using high-resolution results. We discuss how these results can help to understand the formation and evolution of the Milky Way. Similar observations and analysis may be done using this method to study extragalactic globular clusters, such as Magellanic Clouds, dwarf galaxies and more distant galaxies with the emergence of 40m-class telescopes, such as E-ELT.

4.2 Targets selection and observations

The targets were selected among the faintest globular clusters cataloged by (Harris, 1996, 2010 edition) which are more distant and/or highly reddened, many of them poorly studied. Some well-known brighter objects were also observed for comparison with high-resolution spectroscopic studies available in the literature. The sample clusters were divided into four groups defined by their stellar populations, following the criteria discussed by Carretta et al. (2010). Bulge clusters are all objects within 3.5 kpc from the Galactic centre, outer halo clusters have $R_{\text{GC}} \geq 15.0$ kpc, the other objects are classified as disc or inner halo depending on their kinematics and vertical distance in relation to the Galactic plane (see Carretta et al., 2010 for further details). In Table 4.1 we show the population that each globular cluster belongs to, and their sky position are displayed over an all-sky image in Figure 4.1 with letters indicating the respective populations.

Figure 4.2 shows reddening versus distance for all clusters in the bulge, disc, inner and outer halo. Bulge clusters are located in a similar distance from the Sun (~ 8 kpc) and are spread over a wide range of reddening values between ~ 0.2 and 1.5 depending on the direction. The closest disc clusters to the Sun have distances of ~ 2.3 kpc and the farthest

Figure 4.1: Sky position of the 51 clusters over Milky Way image (Mellinger, 2009) in terms of Galactic coordinates in a Aitoff projection. In the position of each cluster, a letter indicates its population, namely (B)ulge, (D)isc, (I)inner halo and (O)uter halo, as classified in Table 4.1.

are ~ 19 kpc from here. Because of that distribution and the low latitudes of these clusters, reddening values varies from ~ 0.04 to ~ 1.7 . Inner halo has similar intervals of $E(B-V)$ and distance as disc clusters. Outer halo clusters have low reddening, $E(B-V) < 0.2$ and are faraway ($11 < d/\text{kpc} < 77$).

We defined $V_{\text{HB}}=17.5$ as a threshold for observations with high-resolution spectrographs in 8m-class telescopes. FLAMES/VLT user manual¹ indicates that observations of stars with $V=17.5$ with one hour of exposure will result a $S/N \sim 30$ using GIRAFFE fibers and $S/N \sim 10$ using UVES fibers. Stars fainter than that would require too much telescope time to get useful spectra and would forbid observations. We selected red giant stars usually brighter than the horizontal branch level (see Figure ??) in each cluster, therefore the classification of globular clusters with $V_{\text{HB}} \geq 17.5$ indicates which clusters could be observed with high-resolution spectroscopy. In Figure 4.2 we call attention to the fact that half of the sample (25 of 51) are not bright enough and hence our observations of low-resolution spectra are perfect to cover the whole range of parameters of all populations of globular clusters in the Galaxy even for the very distant and highly reddened clusters.

Around 10 - 20 red giant stars were selected from photometry in each cluster as described in Chapter 3, for a total of 819 stars. We obtained FORS2@VLT/ESO spectra for them, and 61 spectra (7%) were not considered in the analysis due to very low signal-

¹ FLAMES User Manual VLT-MAN-ESO-13700-2994, Issue 92, 06/03/2013, Table 1

Figure 4.2: Reddening versus distance of the 51 Milky Way globular clusters for each population: bulge, disc, inner halo, outer halo. Empty circles represent clusters with horizontal branch level fainter than $V = 17.5$, filled circles are clusters with V_{HB} brighter than 17.5.

to-noise ratio or data reduction problems. From the remaining 758 useful spectra, 464 (61%) are of confirmed members of the 51 clusters, and the other 294 spectra (39%) are of non-members or at least they do not have results good enough to be classified as members. The spectra were observed in the visible region (grism 1400V, 456 - 586 nm) with resolution of $\Delta\lambda=2.5\text{\AA}$ and typical $S/N\sim 30 - 100$. The data was collected from 2001 to 2012 under projects ID 68.B-0482(A, 2001), ID 69.D-0455(A, 2002), ID 71.D-0219(A, 2003), ID 077.D-0775(A, 2006) and ID 089.D-0493(B, 2012). Table 4.1 lists the selected clusters, their coordinates, observing date and exposure time. Coordinates of the 758 analysed stars and their magnitudes are displayed in Table B.1. The spectra were reduced using FORS2 pipeline inside the *EsoRex* software² following the procedure described in Section 3.3 and Appendix C.

² <https://www.eso.org/sci/software/cpl/esorex.html>

Table 4.1 - Log of observations of the 51 globular clusters using FORS2/VLT with grism 1400V. *a)* 2001 observations, ID 68.B-0482(A); *b)* 2002 observations, ID 69.D-0455(A); *c)* 2003 observations, ID 71.D-0219(A); *d)* 2006 observations, ID 077.D-0775(A); *e)* 2012 observations, ID 089.D-0493(B). Classification of each cluster as (B)ulge, (D)isc, inner (IH) or outer halo (OH), as well as, open cluster (DOpen) and dwarf galaxy-related cluster (OHdSph) follow the criteria defined by Carretta et al. (2010). In the last column we show the numbers of analysed stars that belong to each cluster (C) and those that we classified as field stars (F).

Cluster	Other names	α (J2000) h m s	δ (J2000) ° ' "	obs. date dd.mm.yyyy	UT h:m:s	τ (s)	Pop.	# stars C/F
NGC 104 ^a	47 Tuc	00 24 05.67	-72 04 52.6	22.10.2001	07:16:53	120.0	D	15/1
NGC 2298 ^a		06 48 59.41	-36 00 19.1	23.10.2001	06:25:42	120.0	OH	5/0
NGC 2808 ^d		09 12 03.10	-64 51 48.6	29.05.2006	00:06:07	45.0	IH	14/4
NGC 3201 ^d		10 17 36.82	-46 24 44.9	28.05.2006	22:52:56	20.8	IH	13/2
NGC 4372 ^c		12 25 45.40	-72 39 32.4	25.05.2003	01:34:41	300.0	D	8/2
Rup 106 ^d		12 38 40.2	-51 09 01	28.05.2006	23:15:01	758.6	OH	8/7
NGC 4590 ^b	M 68	12 39 27.98	-26 44 38.6	07.05.2002	03:38:10	60.0	IH	7/3
NGC 5634 ^e		14 29 37.23	-05 58 35.1	26.06.2012	00:12:03	240.0	OH	8/1
NGC 5694 ^e		14 39 36.29	-26 32 20.2	25.06.2012	23:27:18	540.0	OH	8/3
NGC 5824 ^d		15 03 58.63	-33 04 05.6	29.05.2006	00:26:28	553.3	OH	15/3
NGC 5897 ^b		15 17 24.50	-21 00 37.0	07.05.2002	03:53:29	60.0	IH	8/0
NGC 5904 ^c	M 5	15 18 33.22	+02 04 51.7	04.05.2003	06:00:38	300.0	IH	9/0
NGC 5927 ^b		15 28 00.69	-50 40 22.9	07.05.2002	04:14:29	300.0	D	6/0
NGC 5946 ^e		15 35 28.52	-50 39 34.8	23.06.2012	02:52:00	180.0	IH	5/10
BH 176 ^e		15 39 07.45	-50 03 09.8	22.05.2012	03:06:04	600.0	DOpen	11/4
Lynga 7 ^d	BH 184	16 11 03.65	-55 19 04.0	29.05.2006	01:17:18	451.6	D	3/10
Pal 14 ^e	AvdB	16 11 00.6	+14 57 28	15.06.2012	02:49:24	1140.0	OH	6/1
NGC 6121 ^d	M 4	16 23 35.22	-26 31 32.7	29.05.2006	02:53:01	5.8	IH	8/6
NGC 6171 ^b	M 107	16 32 31.86	-13 03 13.6	07.05.2002	04:39:29	60.0	B	1/4
NGC 6254 ^d	M 10	16 57 09.05	-04 06 01.1	29.05.2006	03:11:21	54.5	D	13/2
NGC 6284 ^e		17 04 28.51	-24 45 53.5	22.07.2012	03:17:03	180.0	D	7/10
NGC 6316 ^e		17 16 37.30	-28 08 24.4	22.07.2012	03:29:22	180.0	B	7/9
NGC 6356 ^d		17 23 34.93	-17 48 46.9	29.05.2006	04:40:47	167.9	D	13/5
NGC 6355 ^e		17 23 58.59	-26 21 12.3	11.09.2012	23:19:52	240.0	B	6/10
NGC 6352 ^e		17 25 29.11	-48 25 19.8	22.05.2012	05:33:04	60.0	B	12/2
NGC 6366 ^e		17 27 44.24	-05 04 47.5	15.06.2012	04:53:54	60.0	IH	14/3
HP 1 ^d	BH 229	17 31 05.2	-29 58 54	30.05.2006	05:19:08	1037.5	B	8/19
NGC 6401 ^e		17 38 36.60	-23 54 34.2	14.07.2012	05:36:29	300.0	B	6/12
NGC 6397 ^c		17 40 42.09	-53 40 27.6	06.05.2003	03:54:19	300.0	D	18/3
NGC 6397 ^d		"	"	29.05.2006	05:24:27	7.7	"	"
Pal 6 ^e		17 43 42.2	-26 13 21	12.09.2012	01:01:28	780.0	B	4/13
NGC 6426 ^e		17 44 54.65	+03 10 12.5	13.07.2012	02:31:12	500.0	IH	5/5
NGC 6440 ^d		17 48 52.70	-20 21 36.9	20.05.2006	05:38:49	649.2	B	7/9
NGC 6441 ^d		17 50 13.06	-37 03 05.2	29.05.2006	06:27:01	227.2	D	8/10
NGC 6453 ^e		17 50 51.70	-34 35 57.0	12.09.2012	00:50:14	300.0	D	3/13
Djorg 2 ^e	ESO456-SC38	18 01 49.1	-27 49 33	14.07.2012	05:52:59	180.0	B	4/11

Table 4.1 - continued.

Cluster	Other names	α (J2000) h m s	δ (J2000) ° ' "	obs. date dd.mm.yyyy	UT h:m:s	τ (s)	Pop.	# stars C/F
NGC 6528 ^d		"	"	29.05.2006	08:36:22	149.4	B	4/13
NGC 6539 ^e		18 04 49.68	-07 35 09.1	12.09.2012	00:13:01	360.0	B	7/8
NGC 6553 ^d		18 09 17.60	-25 54 31.3	29.05.2006	08:57:50	79.4	B	11/6
NGC 6558 ^d		18 10 17.60	-31 45 50.0	29.05.2006	06:55:32	148.3	B	4/13
IC 1276 ^d	Pal 7	18 10 44.20	-07 12 27.4	29.05.2006	07:17:06	229.8	D	12/5
NGC 6569 ^d		18 13 38.80	-31 49 36.8	29.05.2006	07:43:05	210.4	B	7/11
NGC 6656 ^d	M 22	18 36 23.94	-23 54 17.1	29.05.2006	08:22:25	36.1	D	44/9
"	"	"	"	"	09:57:32	36.1	"	"
"	"	"	"	"	08:08:05	36.1	"	"
NGC 6749 ^e		19 05 15.3	+01 54 03	27.05.2012	05:04:40	810.0	IH	4/13
NGC 6752 ^a		19 10 52.11	-59 59 04.4	25.05.2003	06:55:40	300.0	D	5/1
Pal 10 ^e		19 18 02.1	+18 34 18	16.06.2012	05:38:57	900.0	D	9/14
Terzan 8 ^e		19 41 44.41	-33 59 58.1	12.07.2012	07:37:44	360.0	OHdSph	12/1
"		"	"	13.07.2012	08:10:51	360.0	"	"
"		"	"	14.07.2012	05:18:10	360.0	"	"
Pal 11 ^e		19 45 14.4	-08 00 26	13.06.2012	07:32:42	180.0	IH	10/2
"		"	"	"	07:50:37	300.0	"	"
NGC 6838 ^d	M 71	19 53 46.49	+18 46 45.1	29.05.2006	09:14:32	17.2	D	8/4
NGC 6864 ^e	M 75	20 06 04.69	-21 55 16.2	27.07.2012	05:35:32	240.0	IH	10/2
NGC 7006 ^d		21 01 29.38	+16 11 14.4	30.05.2006	09:08:54	1200.0	OH	5/9
NGC 7078 ^d	M 15	21 29 58.33	+12 10 01.2	29.05.2006	09:30:56	47.7	IH	15/0

4.3 Method - summary from Chapter 3

Atmospheric parameters (T_{eff} , $\log(g)$, $[\text{Fe}/\text{H}]$, $[\text{Mg}/\text{Fe}]$, $[\alpha/\text{Fe}]$) were derived using full spectrum fitting with code ETOILE (Katz et al., 2011 and Katz, 2001). As reference library we chose one empirical (MILES, Sánchez-Blázquez et al., 2006) and one synthetic (COELHO, Coelho et al., 2005) to complement each other. We adopted the average results from MILES and COELHO as the final T_{eff} , $\log(g)$, $[\text{Fe}/\text{H}]$, and the others are the results without taking the average, i.e., $[\text{Mg}/\text{Fe}]$ from MILES results and $[\alpha/\text{Fe}]$ from COELHO results. Before run the comparison of a given target spectrum with the entire library of reference spectra two important steps are needed: convolute all the library spectra to the same resolution of the target spectrum, and correct of radial velocities, measured also with ETOILE code. For detailed discussion of this method, arguments for each choice and validation with known stars and high-resolution analysis, see Chapter 3.

4.4 Calibration of results

4.4.1 $[Fe/H]$ in Carretta scale

Carretta et al. (2009) published a new metallicity scale for Milky Way globular clusters based on their observations of 19 clusters with UVES (Carretta et al., 2009) and GIRAFFE (Carretta et al., 2009) @ VLT/ESO. In our survey we have 13 objects in common with their sample, and our $[Fe/H]$ results are compared with theirs in Figure 4.3, where the names of the clusters are indicated. Black line is the one-to-one relation, the red line is the linear fit to the dots without considering the error bars, and blue dashed line is the second-order polynomial fit. The transformation of our results into Carretta’s UVES results is given by one of the two equations below:

$$[Fe/H]_{C09} = -0.12(\pm 0.05) + 0.953(\pm 0.032)[Fe/H]_{D14} \quad (4.1)$$

$$[Fe/H]_{C09} = -0.13(\pm 0.11) + 0.94(\pm 0.18)[Fe/H]_{D14} - 0.00(\pm 0.06)[Fe/H]_{D14}^2 \quad (4.2)$$

From Figure 4.3 it is possible to see that second-order polynomial function coincides with the linear function, which is confirmed by the null second-order coefficient of Equation 4.2 and the similarity of the other coefficients between Equations 4.1 and 4.2. The angular coefficient of the linear function is compatible with 1.0, which means that our measurements have no important trends with respect to Carretta’s scale. Although the linear coefficient of Equation 4.1 is compatible with 0.0, our metallicities are slightly higher than those of Carretta et al. (2009), in particular for the more metal-rich clusters. In Table 4.2 we present our $[Fe/H]$ results and also those from Carretta et al. (2009).

Table 4.2 - Average $[Fe/H]$, $[Mg/H]$ and radial velocities for the 13 globular clusters in common with Carretta et al. (2009) and number of selected member stars averaged in each cluster as specified in Table B.3. $[Fe/H]$ and $[Mg/H]$ from Carretta et al. (2009) are also presented for comparison.

Cluster	Other names	$[Fe/H]$ (MILES)	$[Fe/H]$ (COELHO)	$[Fe/H]$ (average)	$[Fe/H]$ (Car09)
NGC 104	47 Tuc	-0.46 ± 0.06	-0.95 ± 0.06	-0.71 ± 0.25	-0.77 ± 0.05
NGC 2808		-0.67 ± 0.09	-1.21 ± 0.06	-1.06 ± 0.22	-1.15 ± 0.07

Table 4.2 - continued.

Cluster	Other names	[Fe/H] (MILES)	[Fe/H] (COELHO)	[Fe/H] (average)	[Fe/H] (Car09)
NGC 3201		-1.52±0.08	-1.51±0.04	-1.5126±0.0011	-1.51±0.06
NGC 4590	M 68	-1.86±0.15	-2.23±0.05	-2.20±0.06	-2.26±0.05
NGC 5904	M 5	-1.02±0.13	-1.30±0.06	-1.25±0.08	-1.34±0.05
NGC 6121	M 4	-0.80±0.13	-1.05±0.06	-1.01±0.07	-1.17±0.05
NGC 6171	M 107	-0.55±0.27	-1.00±0.10	-0.95±0.09	-1.03±0.04
NGC 6254	M 10	-1.59±0.08	-1.55±0.04	-1.558±0.015	-1.57±0.06
NGC 6397		-1.75±0.06	-2.154±0.031	-2.07±0.13	-1.99±0.04
NGC 6441		-0.18±0.09	-0.71±0.10	-0.41±0.26	-0.43±0.06
NGC 6752		-1.49±0.13	-1.59±0.08	-1.57±0.04	-1.55±0.05
NGC 6838	M 71	-0.44±0.08	-0.87±0.08	-0.66±0.22	-0.83±0.06
NGC 7078	M 15	-2.114±0.020	-2.487±0.030	-2.23±0.16	-2.32±0.06

4.4.2 Calibration of $[Mg/Fe]$ results

We derived $[Mg/Fe]$ using MILES library (see Table B.3) which does not have metal-rich stars with higher alpha-elements enhancement, typical in the Milky Way bulge. In previous subsection we showed that for the more metal-rich clusters we derived higher metallicities than those from Carretta et al. (2009). This can be explained as follows: for metal-rich alpha-enhanced target stars the features depth are fitted not by stars with higher $[Mg/Fe]$ but it is compensated by a higher $[Fe/H]$. Therefore, if we scale our $[Fe/H]$ as done above, we should correct the $[Mg/Fe]$ in terms of this difference.

Salaris et al. (1993) showed how $[Fe/H]$ and $[\alpha/Fe]$ are combined to get the total metallicity $[M/H]$, expressed in the equation below.

$$[M/H] = [Fe/H] + \log(0.638f + 0.362) \quad (4.3)$$

where $\log(f) = [\alpha/Fe]$

To find the difference in $[Mg/Fe]$ equivalent to the difference found in $[Fe/H]$ for a given spectrum of total metallicity $[M/H]$ we have to subtract Equation 4.3 for one case and the other as follows:

Figure 4.3: Comparison of $[\text{Fe}/\text{H}]$ with results of Carretta's scale for 13 clusters in common. Black line is the one-to-one relation and the red line is the linear fit to the dots without considering the error bars, and blue dashed line is the second-order polynomial fit.

$$[\text{M}/\text{H}]_{\text{C09}} = [\text{Fe}/\text{H}]_{\text{C09}} + \log(0.638f_{\text{C09}} + 0.362)$$

—

$$[\text{M}/\text{H}]_{\text{D14}} = [\text{Fe}/\text{H}]_{\text{D14}} + \log(0.638f_{\text{D14}} + 0.362)$$

$$\Delta[\text{M}/\text{H}] = \Delta[\text{Fe}/\text{H}] + \log[(0.638f_{\text{C09}} + 0.362)/(0.638f_{\text{D14}} + 0.362)] \quad (4.4)$$

Manipulating this subtraction and assuming that $\Delta[\text{M}/\text{H}] = 0$, and that the differential behaviour of $[\text{Mg}/\text{Fe}]$ is the same of $[\alpha/\text{Fe}]$, it is possible to isolate f_{C09} :

$$f_{\text{C09}} = \{10^{[\log(0.638f_{\text{D14}} + 0.362) - \Delta[\text{Fe}/\text{H}]]} - 0.362\} / 0.638 \quad (4.5)$$

where for $\Delta[\text{Fe}/\text{H}] = 0$, obviously $f_{\text{C09}} = f_{\text{D14}}$.

To find the final $[\text{Mg}/\text{Fe}]$ compatible with the calibrated $[\text{Fe}/\text{H}]$ found in Section 4.4.1 we determine the difference $\Delta[\text{Fe}/\text{H}]$ between the uncalibrated $[\text{Fe}/\text{H}]_{\text{D14}}$ and the calibrated $[\text{Fe}/\text{H}]_{\text{C09}}$. From Equation 4.3 we calculate $f_{\text{D14}} = 10^{[\text{Mg}/\text{Fe}]_{\text{D14}}}$ using the uncalibrated $[\text{Mg}/\text{Fe}]_{\text{D14}}$. Lastly, we apply Equation 4.5 to find the calibrated $[\text{Mg}/\text{Fe}]_{\text{C09}} = \log(f_{\text{C09}})$.

4.5 Discussion on abundances results

Final parameters ($[\text{Fe}/\text{H}]$, $[\text{Mg}/\text{Fe}]$, $[\alpha/\text{Fe}]$ and v_{helio}) for the 51 clusters are presented in Table 4.3. The average of $[\text{Fe}/\text{H}]$ from individual stars using MILES and COELHO are in the Table, as well as the combined average of MILES and COELHO, that we assume as our results. The final values of $[\text{Fe}/\text{H}]$ are those calibrated as described in Section 4.4.1. Average $[\text{Mg}/\text{Fe}]$ and $[\alpha/\text{Fe}]$ of individual stars were calculated with results in Table B.3. The final values of $[\text{Mg}/\text{Fe}]$ are those calibrated following the procedures of Section 4.4.2. It is the first time that $[\text{Fe}/\text{H}]$ and $[\text{Mg}/\text{Fe}]$ derived in a consistent way are published for such a huge sample of globular clusters (51 objects), almost 1/3 of the total number of catalogued clusters (157 as compiled by Harris, 1996, 2010 edition).

In Figure 4.4 we compare our calibrated results of $[\text{Fe}/\text{H}]$ with those from Carretta et al. (2009) for the 51 Milky Way globular clusters of our FORS2 survey (see Table 4.3). In the main panel a one-to-one line is plotted for reference showing a good agreement of both sets of $[\text{Fe}/\text{H}]$, with no trends and a dispersion of $\sigma = 0.16$ dex, as supported by the residuals in the bottom panel. The 13 clusters used to scale our $[\text{Fe}/\text{H}]$ to Carretta's scale in Section 4.4.1 are highlighted as red triangles to show that the application of Equation 4.1 scales their metallicities with low dispersion in comparison with the other clusters ($\sigma = 0.07$ dex). There are six clusters in our sample that were not scaled by Carretta et al. (2009), therefore we plotted them with $[\text{Fe}/\text{H}]$ from our work only in the main plot in blue. The names of the clusters are indicated in the Figure and our work publish for the first time homogeneous metallicities for these objects already scaled to Carretta's scale.

Table 4.3 - Final parameters for the 51 clusters [Fe/H], [Mg/Fe], [α /Fe] and v_{helio} . Values labelled with (MILES), (COELHO) and v_{helio} are averages of individual stars from Tables B.1 and B.3. Car09 are the metallicities of Carretta et al. (2009). [Fe/H] (calibrated) is determined in Section 4.4.1 and [Mg/Fe] (calibrated) in Section 4.4.2.

Cluster	Other names	v_{helio} (km/s)	[Fe/H] (MILES)	[Fe/H] (COELHO)	[Fe/H] (average)	[Fe/H] (calibrated)	[Fe/H] (Car09)	[Mg/Fe] (MILES)	[Mg/Fe] (calibrated)	[α /Fe] (COELHO)
NGC 104	47 Tuc	-48±10	-0.46±0.06	-0.95±0.06	-0.71±0.25	-0.78±0.24	-0.76±0.02 (1)	0.26±0.05	0.37±0.05	0.264±0.027
NGC 2298		134±14	-1.68±0.16	-1.98±0.05	-1.95±0.05	-1.98±0.09	-1.96±0.04 (1)	0.44±0.06	0.47±0.06	0.19±0.06
NGC 2808		96±25	-0.67±0.09	-1.21±0.06	-1.06±0.22	-1.13±0.22	-1.18±0.04 (1)	0.22±0.04	0.31±0.04	0.240±0.031
NGC 3201		472±19	-1.52±0.08	-1.51±0.04	-1.5126±0.0011	-1.56±0.07	-1.51±0.02 (1)	0.430±0.040	0.49±0.04	0.219±0.035
NGC 4372		63±10	-1.83±0.12	-2.34±0.07	-2.22±0.18	-2.24±0.19	-2.19±0.08 (1)	0.40±0.04	0.42±0.04	0.21±0.04
Rup 106		-47±12	-1.60±0.12	-1.54±0.05	-1.545±0.013	-1.59±0.07	-1.78±0.08 (1)	0.45±0.05	0.51±0.05	0.12±0.035
NGC 4590	M 68	-92±25	-1.86±0.15	-2.23±0.05	-2.20±0.06	-2.22±0.10	-2.27±0.04 (1)	0.39±0.05	0.41±0.05	0.19±0.05
NGC 5634		-29±42	-1.60±0.10	-1.82±0.07	-1.75±0.09	-1.79±0.11	-1.93±0.09 (2)	0.43±0.05	0.47±0.05	0.20±0.04
NGC 5694		-150± 9	-1.76±0.12	-2.00±0.04	-1.98±0.04	-2.01±0.09	-2.02±0.07 (1)	0.41±0.05	0.44±0.05	0.17±0.04
NGC 5824		-35±12	-1.95±0.05	-2.011±0.031	-1.995±0.024	-2.02±0.08	-1.94±0.14 (1)	0.436±0.030	0.467±0.030	0.240±0.029
NGC 5897		88±13	-1.63±0.12	-2.02±0.04	-1.97±0.08	-2.00±0.11	-1.90±0.06 (1)	0.43±0.05	0.46±0.05	0.23±0.04
NGC 5904	M 5	46± 7	-1.02±0.13	-1.30±0.06	-1.25±0.08	-1.31±0.10	-1.33±0.02 (1)	0.35±0.05	0.42±0.05	0.24±0.04
NGC 5927		-99±11	-0.165±0.026	-0.82±0.10	-0.21±0.08	-0.32±0.09	-0.29±0.07 (1)	0.09±0.04	0.24±0.04	0.30±0.04
NGC 5946		134±29	-1.50±0.17	-1.55±0.07	-1.544±0.015	-1.59±0.07	-1.29±0.14 (1)	0.42±0.07	0.48±0.07	0.22±0.05
BH 176		-6±14	-0.08±0.04	-0.07±0.06	-0.078±0.007	-0.19±0.05	—	0.10±0.05	0.26±0.05	0.20±0.029
Lynga 7	BH 184	-13±28	-0.41±0.13	-0.87±0.15	-0.61±0.23	-0.70±0.23	—	0.21±0.11	0.33±0.11	0.27±0.06
Pal 14	AvdB	45± 9	-1.07±0.18	-1.27±0.10	-1.22±0.07	-1.28±0.09	-1.63±0.08 (1)	0.32±0.06	0.40±0.06	0.24±0.04
NGC 6121	M 4	23±35	-0.80±0.13	-1.05±0.06	-1.01±0.07	-1.08±0.09	-1.18±0.02 (1)	0.35±0.06	0.44±0.06	0.27±0.04
NGC 6171	M 107	-122± 0	-0.55±0.27	-1.00±0.10	-0.95±0.09	-1.02±0.10	-1.03±0.02 (1)	0.28±0.21	0.37±0.21	0.20±0.14
NGC 6254	M 10	43±34	-1.59±0.08	-1.55±0.04	-1.558±0.015	-1.60±0.07	-1.57±0.02 (1)	0.436±0.034	0.492±0.034	0.214±0.035
NGC 6284		35±21	-0.84±0.15	-1.12±0.07	-1.07±0.09	-1.14±0.10	-1.31±0.09 (2)	0.33±0.06	0.42±0.06	0.27±0.04
NGC 6316		81±40	-0.28±0.08	-0.84±0.10	-0.50±0.27	-0.60±0.26	-0.36±0.14 (1)	0.11±0.06	0.24±0.06	0.303±0.031
NGC 6356		36±41	-0.30±0.06	-0.91±0.07	-0.55±0.30	-0.64±0.29	-0.35±0.14 (1)	0.12±0.04	0.25±0.04	0.303±0.024
NGC 6355		-215± 7	-1.38±0.09	-1.54±0.08	-1.46±0.08	-1.51±0.10	-1.33±0.14 (1)	0.264±0.010	0.330±0.010	0.27±0.05

Table 4.3 - continued.

Cluster	Other names	v_{helio} (km/s)	[Fe/H] (MILES)	[Fe/H] (COELHO)	[Fe/H] (average)	[Fe/H] (calibrated)	[Fe/H] (Car09)	[Mg/Fe] (MILES)	[Mg/Fe] (calibrated)	[α /Fe] (COELHO)
NGC 6352		-138±36	-0.41±0.06	-0.76±0.07	-0.54±0.17	-0.63±0.17	-0.62±0.05 (1)	0.15±0.05	0.28±0.05	0.304±0.025
NGC 6366		-137±54	-0.41±0.07	-0.81±0.07	-0.61±0.20	-0.70±0.20	-0.59±0.08 (1)	0.22±0.05	0.34±0.05	0.303±0.023
HP 1	BH 229	54± 5	-0.70±0.11	-1.49±0.09	-1.2 ±0.4	-1.26±0.39	-1.57±0.09 (2)	0.33±0.07	0.41±0.07	0.28±0.04
NGC 6401		-120±17	-0.59±0.14	-1.34±0.09	-1.12±0.31	-1.19±0.30	-1.01±0.14 (1)	0.32±0.08	0.40±0.08	0.269±0.041
NGC 6397		-27±55	-1.75±0.06	-2.154±0.031	-2.07±0.13	-2.09±0.15	-1.99±0.02 (1)	0.402±0.030	0.430±0.030	0.229±0.028
Pal 6		177± 5	-0.27±0.14	-1.66±0.17	-0.8 ±0.7	-0.88±0.67	-1.06±0.09 (2)	0.14±0.10	0.25±0.10	0.28±0.05
NGC 6426		-242±11	-2.03±0.11	-2.46±0.05	-2.39±0.11	-2.40±0.14	—	0.38±0.06	0.39±0.06	0.24±0.05
NGC 6440		-59±26	-0.03±0.06	-0.80±0.10	-0.24±0.30	-0.35±0.29	-0.20±0.14 (1)	0.11±0.04	0.26±0.04	0.314±0.031
NGC 6441		-6±32	-0.18±0.09	-0.71±0.10	-0.41±0.26	-0.51±0.25	-0.44±0.07 (1)	0.11±0.06	0.25±0.06	0.26±0.04
NGC 6453		-153±11	-1.45±0.18	-1.57±0.10	-1.54±0.04	-1.59±0.08	-1.48±0.14 (1)	0.42±0.09	0.48±0.09	0.16±0.06
Djorg 2	ESO456SC38	-150±28	-0.50±0.12	-1.19±0.14	-0.79±0.33	-0.87±0.32	—	0.28±0.10	0.38±0.10	0.27±0.05
NGC 6528		185±10	-0.07±0.10	-0.18±0.08	-0.13±0.05	-0.24±0.07	+0.07±0.08 (1)	0.05±0.09	0.21±0.09	0.26±0.05
NGC 6539		30±18	-0.23±0.09	-0.89±0.09	-0.55±0.33	-0.64±0.32	-0.53±0.14 (1)	0.16±0.07	0.29±0.07	0.302±0.028
NGC 6553		6±8	-0.125±0.009	-0.55±0.07	-0.133±0.017	-0.25±0.05	-0.16±0.06 (1)	0.107±0.009	0.263±0.009	0.302±0.025
NGC 6558		-210±16	-0.88±0.20	-1.02±0.05	-1.012±0.013	-1.08±0.06	-1.37±0.14 (1)	0.26±0.06	0.35±0.06	0.23±0.06
IC 1276	Pal 7	155±15	-0.13±0.06	-1.11±0.07	-0.56±0.49	-0.65±0.47	-0.65±0.09 (2)	0.09±0.05	0.22±0.05	0.304±0.026
NGC 6569		-51± 9	-0.53±0.09	-0.85±0.11	-0.66±0.15	-0.75±0.15	-0.72±0.14 (1)	0.30±0.07	0.41±0.07	0.29±0.032
NGC 6656	M 22	-152±25	-1.77±0.05	-1.945±0.020	-1.92±0.05	-1.95±0.09	-1.70±0.08 (1)	0.501±0.008	0.536±0.008	0.225±0.018
NGC 6749		-66± 8	-0.64±0.15	-2.14±0.11	-1.59±0.70	-1.63±0.67	-1.62±0.09 (2)	0.34±0.10	0.40±0.10	0.17±0.06
NGC 6752		-28± 7	-1.49±0.13	-1.59±0.08	-1.57±0.04	-1.62±0.08	-1.55±0.01 (1)	0.47±0.06	0.52±0.06	0.22±0.05
Pal 10		-38±17	-0.080±0.039	-0.53±0.05	-0.24±0.20	-0.35±0.20	—	0.118±0.010	0.266±0.010	0.273±0.034
Terzan 8		135±19	-1.76±0.07	-2.18±0.05	-2.06±0.17	-2.08±0.18	—	0.41±0.04	0.44±0.04	0.21±0.04
Pal 11		-81±15	-0.22±0.05	-0.62±0.08	-0.35±0.17	-0.45±0.17	-0.45±0.08 (1)	0.12±0.05	0.26±0.05	0.303±0.030
NGC 6838	M 71	-42±18	-0.48±0.08	-0.77±0.08	-0.63±0.15	-0.72±0.15	-0.82±0.02 (1)	0.25±0.07	0.37±0.07	0.293±0.032
NGC 6864	M 75	-190±20	-0.75±0.10	-1.09±0.06	-1.00±0.13	-1.07±0.14	-1.29±0.14 (1)	0.35±0.05	0.44±0.05	0.222±0.035
NGC 7006		-391±24	-1.54±0.19	-1.74±0.11	-1.69±0.07	-1.73±0.10	-1.46±0.06 (1)	0.42±0.07	0.47±0.07	0.25±0.05
NGC 7078	M 15	-159±40	-2.114±0.020	-2.487±0.030	-2.23±0.16	-2.24±0.18	-2.33±0.02 (1)	0.410±0.034	0.428±0.034	0.237±0.030

Figure 4.4: Comparison of calibrated $[\text{Fe}/\text{H}]$ from this work (D14) with those from (Carretta et al., 2009, C09) for the 51 clusters of our survey using values of Table 4.3. *Main panel:* red triangles emphasize the 13 clusters used for calibration in Section 4.4.1. Blue points are the six clusters not calibrated by Carretta et al. (2009) and analysed in this work. One-to-one line is plotted for reference. *Bottom panel:* residuals of the comparison, with standard deviation equal to 0.16 dex. *Top and right panels:* smoothed histogram of metallicities from this work and from Carretta, respectively, only for the 45 clusters in common.

Another subproduct of this comparison is the metallicity distribution of the Milky Way for the 45 clusters in common with Carretta et al. (2009). This group of clusters represents $\sim 30\%$ of the total number of catalogued objects in our Galaxy by (Harris, 1996, 2010 edition), which is significant to draw the known bimodal metallicity distribution of the Milky Way globular cluster population. In the right panel of Figure 4.4 we show the metallicity distribution based on Carretta et al. (2009) values, while in the upper panel our results are displayed. In both cases is possible to identify two peaks, although this

is clearer using Carretta's results since they are based on high-resolution spectroscopic analysis and their uncertainties are lower than ours.

4.5.1 Chemical evolution of the Milky Way

As mentioned in Chapter 1 the ratio $[\alpha/\text{Fe}]$ plotted against $[\text{Fe}/\text{H}]$ is an evidence of how efficient the star formation in the Galaxy was. Supernovae type II (SNII) explode quickly after the formation of the progenitor massive star and release α -elements and iron to the interstellar medium. Later, supernovae type Ia (SNIa) explode liberating iron, which decrease the ratio $[\alpha/\text{Fe}]$. Magnesium is one of the α -elements and represents well these processes. Increasing values of $[\text{Fe}/\text{H}]$ represents many generations of stars enriched by the material expelled by SNII, lower values representing the first stars and higher values standing for more recent objects.

In Figure 4.5 we show our (final calibrated) results of $[\text{Fe}/\text{H}]$ and $[\text{Mg}/\text{Fe}]$ for the 51 clusters (see Table 4.3), splitted according to their populations: bulge, disc, inner halo, outer halo (see Table 4.1). As a reference we plot in the respective panels of Figure 4.5 also field stars from Gonzalez et al. (2011) for bulge, Bensby et al. (2005) for thin disc, Reddy et al. (2006) for thick disc, Roederer (2009) for inner and outer halo. Clearly globular clusters abundances trends follow field stars values, and the range of metallicities covered by each population is equivalent also to the respective field stars population. We note that bulge and disc stars reach high metallicities, above solar, and the cluster populations are mostly more metal-poor than our Sun. These plots are indicatives that globular clusters and field stars populations were formed in the same processes in all Galactic components or, at least, they were formed in processes with similar star formation rate, given by the drop of $[\text{Mg}/\text{Fe}]$ at about $[\text{Fe}/\text{H}] \sim -1.0$.

From Figure 4.5 we conclude that SNII were able to enrich stellar generations up to $[\text{Fe}/\text{H}] \sim -1.0$ before SNIa start to become important, which typically happens ~ 1 Gyr after the formation of the progenitor stars. Matteucci and Recchi (2001) discussed in detail the relationship of the timescale of SNIa events and the distribution of $[\alpha/\text{Fe}]$ vs. $[\text{Fe}/\text{H}]$. They concluded that the timescale of 1 Gyr for the knee shown in Figure 4.5 is specific for solar neighbourhood stars, and it is not a general rule. However the four Galactic components seem to share the same $[\text{Fe}/\text{H}]_{\text{knee}} \sim -1.0$ (see Figure 4.5), and we assume that the SNIa timescale is approximately the same for all components.

Figure 4.5: Chemical evolution of the Milky Way in terms of $[Mg/Fe]$ vs. $[Fe/H]$ for 51 globular clusters from this work. Field stars are from Gonzalez et al. (2011) for bulge, Bensby et al. (2005) for thin disc, Reddy et al. (2006) for thick disc, Roederer (2009) for inner and outer halo.

VandenBerg et al. (2013) have derived the ages of 55 Milky Way globular clusters in a same scale, of which 17 are in common with our sample. In Figure 4.6 we compare ages with $[Fe/H]$ and $[Mg/Fe]$ for the common clusters in the left and right panels, respectively. In the left panel we can see that the first clusters were formed between 12-13 Gyr ago, and the next generations were enriched up to $[Fe/H] \sim -1.0$ about 1 Gyr later. The right panel shows a similar behaviour than that of Figure 4.5, with a knee at about 12 Gyr ago. The oldest objects are ~ 1 Gyr older than that. These plots indicate that the timescale of 1 Gyr for SNIa to become important is a reasonable value for Milky Way globular clusters populations.

4.5.2 Origin of globular clusters

There is further discussions about Figure 4.6 concerning the formation/evolution scenario of the clusters. VandenBerg et al. (2013) noted that for $[Fe/H] > -1.7$ there is a visual impression of a split between halo (red) and bulge/disc (blue) clusters with

Figure 4.6: Age vs. $[\text{Fe}/\text{H}]$ and $[\text{Mg}/\text{Fe}]$ for 17 clusters

$\Delta[\text{Fe}/\text{H}] \sim 0.6$ dex for all ages (left panel of the Figure). Leaman et al. (2013) argued that this effect is real and separates a population of halo clusters from those with disc kinematics. Clusters lying in the more metal-rich branch would have been formed *in situ* while the other objects would have their origin in low-mass dwarf galaxies that were later accreted to the Milky Way.

If the scenario proposed by Leaman et al. (2013) is true it would be reflected also in $[\text{Mg}/\text{Fe}]$ ratio. Dwarf galaxies have less efficient star formation (Kirby et al., 2010), and they do not reach higher metallicities before SNIa start to become important. Therefore, if halo clusters were formed in low-efficient star formation environments as dwarf galaxies we would observe metal-poor clusters with lower $[\text{Mg}/\text{Fe}]$ ratios, which is not the case (see bottom panels of Figure 4.5).

Figure 4.7 shows the radial gradient of $[\text{Fe}/\text{H}]$ for halo and bulge/disc clusters. More distant halo clusters are more metal-poor than the closest one to the Galactic centre. This is in agreement with a scenario of monolithic collapse of Eggen et al. (1962). For bulge/disc clusters there is also a metallicity gradient with increasing $[\text{Fe}/\text{H}]$ towards the centre of the Milky Way. Combining these results with the discussions before we can say that halo clusters may have been formed in building blocks during the initial collapse of the Milky Way, which explains their kinematics. In this scenario they experienced the same efficiency of star formation as the whole Milky Way, which is corroborated by the chemical patterns discussed above. Bulge/disc globular clusters were formed *in situ*, which

explain their kinematics, and is also compatible with their abundances patterns.

Figure 4.7: Metallicity radial gradients for the 51 Milky Way globular clusters of our survey. Pal 14 is the only object that is out of the plot area, with $R_{GC} = 71.6$ kpc and $[Fe/H] = -1.28 \pm 0.09$. We prefer to show a shorter interval in R_{GC} in order to get a better visualization of the gradients.

4.6 Summary and conclusions

In this Chapter we present all parameters derived for the whole survey of 51 Galactic globular clusters, based on the method described and vastly discussed in Chapter 3. We observed 819 red giant stars, analysed 758 useful spectra and classified 464 as members of the 51 clusters, and 294 as non-members. To proceed with membership selection we derived radial velocities for all 758 spectra. T_{eff} , $\log(g)$, $[Fe/H]$ were derived using MILES and COELHO libraries and the results from both libraries were averaged to get the final results from our method. We have 13 clusters in common with observations of Carretta et al. (2009), that we used to calibrate our average metallicities per cluster to Carretta’s scale. $[Mg/Fe]$ were derived using only MILES library and $[\alpha/Fe]$ using only COELHO library. Once COELHO library provides a more limited interval of $[\alpha/Fe]$ than MILES for

[Mg/Fe], we adopted [Mg/Fe] as our proxy for studying the chemical evolution of the Milky Way. We calibrated [Mg/Fe] to a scale compatible with that found for [Fe/H] considering that metal-rich and alpha-enhanced stars had their abundances not perfectly recovered by our method due to a lack of reference stars with these characteristics.

The comparison of our final calibrated [Fe/H] with those from Carretta's scale are in good agreement with a total dispersion of $\sigma = 0.16$ dex, and a lower dispersion ($\sigma = 0.07$ dex) considering only the 13 clusters used for calibration. We recover the known bimodal metallicity distribution of the globular cluster populations. Chemical evolution history of the Milky Way showed to be similar between field stars and globular clusters for the four Galactic components: bulge, disc, inner and outer halo. This shows that the whole Milky Way have experienced star formation and galactic evolution with similar efficiency of chemical enrichment. We found that the typical timescale of 1 Gyr for SNIa to start to become important is reasonable for all components of the Milky Way, based on the age-metallicity relation based on globular clusters. Finally we discussed the origin of the halo and bulge/disc clusters that present different kinematics which seems to indicate different formation scenarios for the two groups. For the halo clusters, there is a clear metallicity radial gradient with relation to the Galactic centre, which corroborates the scenario of a initial collapse of the Galaxy, where building blocks clumps were the environment of formation of the halo clusters. Bulge/disc group of clusters would have been formed *in situ* which would explain their kinematics compatible with disc stars. A radial metallicity gradient is found for this group also which endorse this formation scenario.

FORS2/VLT survey of Milky Way globular clusters

III. Clusters with multiple populations

5.1 Introduction to this chapter

The definition of globular clusters as templates of simple stellar population systems has changed in the last decade (see Gratton et al., 2012, and references therein). The advent of high-precision photometric and spectroscopic data have shown intrinsic abundance variations of light elements, such as He, C, N, O, Na, Al, Mg, Si and F. In particular, the Na-O anticorrelation was considered a characteristic of many globular clusters, from where it is possible to classify a first generation (primordial) and a second generation (intermediate and extreme) components. The presence of this anticorrelation is one more characteristic that differentiate globular and open clusters (Carretta et al., 2010, 2009).

The Milky Way has some peculiar globular clusters presenting also a variation in iron abundance. The classical case of ω Centauri (NGC 5139) has stars with $[\text{Fe}/\text{H}]$ in a range of about 1.5 dex (e.g. Marino et al., 2011, and references therein). Another case is Terzan 5, discussed in Section 1.1.1, that shows at least two populations with metallicity peaks at $[\text{Fe}/\text{H}] \approx -0.3$ and $+0.3$ (e.g. Ferraro et al., 2009) and it is probably not a globular cluster. The nucleus of the Sagittarius dwarf galaxy, the globular cluster M 54 (NGC 6715) has $\sigma_{[\text{Fe}/\text{H}]} \approx 0.19$ dex (Bellazzini et al., 2008; Carretta et al., 2010; Saviane et al., 2012). The massive cluster M 22 (NGC 6656) has been studied for peculiarities in CN and CH band strengths, e.g. Hesser and Harris (1979). The cluster had its iron abundance dispersion of $\sigma_{[\text{Fe}/\text{H}]} \approx 0.15$ dex found out simultaneously by Marino et al. (2009) and by our group (Da Costa et al., 2009). Alves-Brito et al. (2012) found a range of $\Delta[\text{Fe}/\text{H}]$ of 0.4 dex from nine red giant stars using high-resolution spectra. There are other more subtle cases: NGC 2419

with $\sigma_{[\text{Ca}/\text{H}]} \approx 0.2$ dex as an indication of $\sigma_{[\text{Ca}/\text{H}]}$ (Cohen et al., 2010); NGC 1851, with a range in $[\text{Fe}/\text{H}]$ of only 0.08 dex (Carretta et al., 2010, see also Villanova et al., 2010); the recently characterized by our group NGC 5824 with a total $[\text{Fe}/\text{H}]$ range of 0.3 dex (Saviane et al., 2012; Da Costa et al., 2014).

It seems natural to classify these massive ($M_V \lesssim -7$ mag, see Figure 5.1) multiple population globular clusters with intrinsic $[\text{Fe}/\text{H}]$ spread in a same subgroup of clusters. In order to characterize this group as a population sharing similar origins, many papers have been devoted to deeply examine high quality spectroscopy and photometry for these objects. The following possible scenarios for these unusual clusters are proposed ωCen is probably the remnant nucleus of a disrupted dwarf galaxy (Lee et al., 1999), M 54 is located in the nucleus of Sagittarius dwarf galaxy and could be an early snapshot of a ωCen -type cluster (Da Costa and Armandroff, 1995; Carretta et al., 2010). The dispersion in $[\text{Fe}/\text{H}]$ of M 22 is similar to that of ωCen , and Da Costa et al. (2009) suggested that M 22 could share the same origin as ωCen . NGC 1851 is surrounded by a considerable stellar halo that may be a vestige of a dwarf galaxy in which the cluster was once embedded (Olszewski et al., 2009). Carretta et al. (2010) discuss the possibility of NGC 1851 being the result of a merger of two clusters. Lee et al. (2013) indicate NGC 2419 as having an origin in the nucleus of a disrupted dwarf galaxy. Newberg et al. (2009) recently reported the discovery of the Cetus Polar Stream and argued that the cluster NGC 5824 is around 3° from the fitted orbit of the stream. This cluster shares the same distance of the stream, has compatible radial velocity and metallicity. Therefore, NGC 5824 was possibly located in the nucleus of a dwarf galaxy that was captured and destroyed by the potential well of the Milky Way and originated the Cetus Polar Stream (Da Costa et al., 2014).

In a bigger picture, finding evidence of disrupting satellites in the Galactic halo has great significance for the current ΛCDM galaxy formation paradigm. In this scenario, the Galaxy is built-up through the merger and accretion of lower mass systems, predominantly at early epochs. Indeed, while the spatial and kinematic signatures of this process have been erased in the inner parts of the Galaxy, in its halo where dynamical times are long, several stellar streams have been identified in the past (e.g. Majewski et al., 2003; Duffau et al., 2006; Newberg et al., 2010). There is however still an order of magnitude discrepancy between the large number of predicted MW satellites and the observed ones. Any addition of a new remnant is therefore important to confirm the theory. Drinkwater et al. (2000)

Figure 5.1: Integrated magnitude M_V and half-light radius for Milky Way globular clusters (black dots) and ultra compact dwarfs (UCDs) from Fornax and Virgo clusters (cyan stars) from Evstigneeva et al. (2008). The two globular clusters discussed in this Chapter are indicated as red triangles.

reported the existence of ultra compact old objects in the centre of the Fornax cluster of galaxies. They are more massive (brighter, see Figure 5.1) and distributed over larger distances to the centre of NGC 1399 (the central galaxy of the cluster) in comparison with its globular cluster system. They propose that they were either tidally stripped from the nucleus of disrupted dwarf galaxies, as the multiple population globular clusters described above, or they are a new type of galaxy (for a detailed discussion about the so-called ultra compact dwarf (UCD) galaxies see Mieske et al., 2002, 2008, and references therein). In the working hypothesis that Galactic GC with metallicity spreads signpost halo substructures, our FORS2/VLT survey of Milky Way globular clusters has the potential of finding still unknown examples of these peculiar clusters.

Saviane et al. (2012) revised the available $[\text{Fe}/\text{H}]$ for Milky Way globular clusters from spectroscopy of individual stars, and found that for only half of the objects it would be possible to detect intrinsic metallicity dispersion. Up to now, only seven such peculiar clusters have been found, which means a frequency of about 1/10. In our FORS2 survey

we observed 51 clusters, of which ~ 40 are poorly studied, therefore we expect to find about four clusters with metallicity spread in our sample. In this Chapter we present the results for M 22 and NGC 5824 based on optical spectra.

5.2 Observations and analysis

The selection of targets was carried out as described in Section 3.3. The CMDs of M 22 and NGC 5824 are presented in Figure 5.2 where the selected targets are shown in red. Dartmouth isochrones with parameters from (Harris, 1996, 2010 edition) catalogue are overplotted. Sky maps of the two clusters are displayed in Figure 5.3 showing that there was no bias in the position of the selected targets.

Figure 5.2: Colour-Magnitude diagrams of M 22 and NGC 5824. All stars within $2 \times r_{half-light}$ are plotted, without any cleaning procedure. Dartmouth isochrones with literature parameters (Harris, 1996, 2010 edition catalogue) are overplotted. Selected RGB stars for spectroscopic observations are in red. Reddening vectors are shown in each CMD.

Radial velocities were measured by cross correlation and atmospheric parameters were determined following the full spectrum fitting techniques with ETOILE code plus MILES and COELHO stellar libraries in the same way as discussed and validated in Section 3.4. Results for all stars in these two clusters are presented in Tables B.1 (star ID, coordinates, magnitudes, colours, heliocentric radial velocities from this work and from CaT analysis, membership) and B.3 (T_{eff} , $\log(g)$, $[Fe/H]$, $[Mg/Fe]$ and $[\alpha/Fe]$).

Figure 5.3: Sky map of M 22 and NGC 5824. The size of dots are scaled with the stellar magnitudes as indicated in each plot. Selected RGB stars for spectroscopic observations are in red. The blue circle corresponds to the half-light radius of each cluster from (Harris, 1996, 2010 edition) catalogue.

5.3 Results

5.3.1 T_{eff} , $\log(g)$ against isochrones

Figure 5.4 displays the results of all stars in the two clusters in a Hertzsprung-Russell diagram form. Left, middle and right panels show the results using MILES library, COE-LHO library and the average of both results weighted by their uncertainties, respectively. Black dots represent member stars of each cluster, and grey dots are non-members, based on the selection described in the next Section 5.3.2. Dartmouth isochrones (Dotter et al., 2008) with age, $[\text{Fe}/\text{H}]$ and $[\alpha/\text{Fe}]$ from (Harris, 1996, 2010 edition) catalogue are over-plotted in the diagrams of Figure 5.4 in blue. Cyan lines have the same age and $[\text{Fe}/\text{H}]$ as the respective blue lines, but with the extreme values of $[\alpha/\text{Fe}] = -0.2$ and $+0.8$, available from the models.

The results on T_{eff} and $\log(g)$ from MILES and COELHO are in good agreement with the isochrones, as well as the average of both results. Uncertainties of results based on MILES are larger than those of COELHO results because of the poor sampling of MILES for metal-poor stars, as discussed in Section 3.6.3. We adopted as the final results in this work the weighted average of MILES and COELHO results.

Figure 5.4: Comparison of T_{eff} and $\log(g)$ of stars in each cluster with Dartmouth isochrones (Dotter et al., 2008). For each cluster we show the results based on MILES and COELHO libraries and a third panel with the weighted average of the results from both libraries. The parameters of age, $[\text{Fe}/\text{H}]$ and $[\alpha/\text{Fe}]$ for the blue thick isochrones were taken from Table 3.2. Cyan thin isochrones have same age and $[\text{Fe}/\text{H}]$ as blue ones, but with the limits $[\alpha/\text{Fe}] = -0.2$ and $+0.8$. Black dots represent member stars of each cluster, and grey dots are not members.

5.3.2 Membership selection

In Figure 5.5 we show four panels for each globular cluster, the same as in Figure 3.11. Black dots represent member stars of each cluster, and grey dots are non-members, green circles are classified as non-members but Saviane et al. (2012) considered them as members from their CaT analysis, cyan diamonds are member stars that were considered as non-members by Saviane et al. (2012). In upper left panels radial velocities against metallicities are displayed. Blue solid lines are parameters from (Harris, 1996, 2010 edition) and red dashed lines are the average of parameters for cluster members only. For all clusters there is a clear concentration of stars around literature values, and these stars were selected as cluster members. Upper right panels show T_{eff} vs. $\log(g)$ compared with Dartmouth isochrones (Dotter et al., 2008). Blue solid lines are isochrones with parameters from (Harris, 1996, 2010 edition), and red dashed lines are based on the average (Table 5.1) of the parameters for member stars (Table B.3). All member stars are close to the isochrones while

non-member stars are outside these lines, confirming the membership selection. Bottom left panels show $[\text{Fe}/\text{H}]$ against distance to the cluster centre, and no trends were found for any sample cluster. Bottom right panels show no correlation between T_{eff} and $[\text{Fe}/\text{H}]$ for any of the six clusters analysed in this work.

Figure 5.5: Same as Figure 3.11 for the clusters M 22 and NGC 5824.

Table 5.1 - Final average parameters for M 22 and NGC 5824 and respective internal errors.

Cluster	$\langle v_{\text{helio}} \rangle$ (km/s)	$\langle [\text{Fe}/\text{H}] \rangle^{(a)}$	$\langle [\text{Fe}/\text{H}] \rangle^{(b)}$	$\langle [\text{Fe}/\text{H}] \rangle^{(avg)}$	$\langle [\text{Mg}/\text{Fe}] \rangle^{(a)}$	$\langle [\alpha/\text{Fe}] \rangle^{(b)}$
M 22	-152 ± 25	-1.77 ± 0.05	-1.945 ± 0.020	-1.92 ± 0.05	0.501 ± 0.008	0.225 ± 0.018
NGC 5824	-35 ± 12	-1.95 ± 0.05	-2.011 ± 0.031	-1.995 ± 0.024	0.436 ± 0.030	0.240 ± 0.029

^aMILES library ^bCOELHO library ^{avg}Average of MILES and COELHO results

In conclusion all T_{eff} , $\log(g)$ and $[\text{Fe}/\text{H}]$ values are found to sit in well defined sequences following the isochrones. Bottom-right panels show no correlation between $[\text{Fe}/\text{H}]$ and T_{eff} , suggesting that the $[\text{Fe}/\text{H}]/T_{\text{eff}}$ degeneracy does not affect our fitting procedure.

5.4 $[\text{Fe}/\text{H}]$ and $[\text{Mg}/\text{H}]$ intrinsic spread of M 22 and NGC 5824

Once the member stars are selected and their atmospheric parameters are determined, it is possible to analyse in detail their metal content seeking for multiple populations. We discuss each cluster in the following subsections.

5.4.1 M 22

From the 53 observed stars we selected 44 as M 22 members, giving an average radial velocity of -152 ± 25 km/s, compatible with -150 ± 2 km/s from 51 red giant stars of Da Costa et al. (2009) observed in the CaT with $R \sim 2,000$ (from our group) and -146 ± 2 km/s derived by Marino et al. (2009) using high-resolution ($R \sim 20,000 - 45,000$) spectra in the optical region of 17 red giant stars. The metallicity distribution function (MDF) using our results is presented in Figure 5.6, as the smoothed histogram in black. From this work we present also the distribution of $[\text{Mg}/\text{H}]$ in the right panel of the same figure. Dashed red and solid blue lines in Figure 5.6 are gaussians with σ equal to the average errors of individual stars abundances, and centre at the main peak of the distributions and at the average abundances, respectively. The comparison of the red gaussians with the data (black lines) shows a lack of metal-poor (and low $[\text{Mg}/\text{H}]$) stars and an excess of metal-rich (and high $[\text{Mg}/\text{H}]$) stars. This comparison shows that the dispersion in $[\text{Fe}/\text{H}]$ and $[\text{Mg}/\text{H}]$ are not due to errors in the abundance derivations for each star, but it is a real dispersion related to the formation of the cluster. The blue gaussians compared to the black observed ones reveals that the shape of the abundance distributions are not uniform around the average values. This could not be a bias in the target selection because we chose stars spread over the cluster area in the sky and along the red giant branch with no preferences, as shown in Figures 5.2 and 5.3. Therefore the two populations are real, and we can split them as population A with $[\text{Fe}/\text{H}] < -1.8$ and population B with $[\text{Fe}/\text{H}] > -1.8$.

The main peak of $[\text{Fe}/\text{H}]$ at ~ -1.9 and the metal-rich tail of the MDF up to ~ -1.3 coincides with the findings of Da Costa et al. (2009). We derived an average $[\text{Fe}/\text{H}] = -1.92 \pm 0.05$ and $[\text{Mg}/\text{Fe}] = 0.501 \pm 0.008$, and Marino et al. (2009, 2011) derived $[\text{Fe}/\text{H}] = -1.76 \pm 0.02$ and $[\text{Mg}/\text{Fe}] = 0.39 \pm 0.03$. We derived a higher value of $[\text{Mg}/\text{Fe}]$ because we have few stars with $[\text{Mg}/\text{Fe}] \sim 0.5$ with lower error bars that are biasing the average. The mean and standard deviation of our results give $[\text{Mg}/\text{Fe}] = 0.43 \pm 0.04$ which is compatible with Marino et al. (2009). The difference in $[\text{Fe}/\text{H}]$ from our sample and that of Marino's papers are explained by the fact that we observed more stars from population A ($[\text{Fe}/\text{H}] < -1.8$) and they observed more stars from population B ($[\text{Fe}/\text{H}] > -1.8$), as shown in Figure 5.7 and discussed below.

Marino et al. (2011) and us have selected red giant stars spread over the cluster with no

Figure 5.6: *Left panel:* $[\text{Fe}/\text{H}]$ distribution function of M 22 showing a spread in $[\text{Fe}/\text{H}]$ (solid black line). Dashed red line is a normal distribution centred in the main peak with σ equal to the mean uncertainty of individual stars. Blue solid line is the same, but with the peak displaced to the $\langle [\text{Fe}/\text{H}] \rangle$ from all cluster stars. *Right panel:* the same, but for $[\text{Mg}/\text{H}]$.

biases for any region for both populations. The stars of both populations lay out along the RGB with no preferences of any region, although we observed stars systematically fainter than Marino et al. (2011). We have three stars in common with their sample, our M22a_3, M22b_7, M22b_10 corresponding to their I-86, I-27, I-53, respectively. We derived $[\text{Fe}/\text{H}] = -1.92 \pm 0.12, -1.32 \pm 0.21, -1.79 \pm 0.20$, and they derived $[\text{Fe}/\text{H}] = -1.82 \pm 0.04, -1.73 \pm 0.04, -1.74 \pm 0.05$, respectively. All stars have compatible $[\text{Fe}/\text{H}]$ and no systematic difference was detected. Among our fainter RGB stars there are 30 classified as population A (metal-poor) and 14 as population B (metal-rich). The brighter stars observed by Marino et al. (2011) are separated in 14 from population A and 21 from population B. Once we observed more metal-poor stars, our metallicity average is lower than that of Marino et al. (2011) who observed more metal-rich stars. Considering the three stars in common, and combining both samples, the total number of stars is 43 from population A and 33 from population B. As a result, the ratio of populations is $(\text{popA})/(\text{popB}) \approx 1.3$, i.e, metal-poor population in M 22 represent $\sim 57\%$ of all stars, and metal-rich population, $\sim 43\%$. Despite that, this ratio is not uniform with magnitude (a proxy for the stellar mass), as displayed in the right panel of Figure 5.7, where smoothed histograms of the distributions of populations A and B (summed samples of this work with Marino's) are shown in blue and red, respectively. These histograms shows a predominancy of population A in the fainter part of the RGB

and population B in the brighter part. In fact, we observed fainter stars and Marino et al. (2011) observed brighter ones, as well as Alves-Brito et al. (2012).

Figure 5.7: Comparison of sample stars from this work with Marino et al. (2011) for M 22. Purple and blue circles are population A stars ($[\text{Fe}/\text{H}] < -1.8$) from this work and from Marino et al. (2011), respectively. Brown and red squares are population B stars ($[\text{Fe}/\text{H}] > -1.8$) from this work and from Marino et al. (2011), respectively. In the main panel, blue and red lines are Dartmouth 14 Gyr old isochrones with $[\text{Fe}/\text{H}] = -1.9$ and -1.7 , respectively. In the right panel, smoothed histograms of the distributions of populations A and B (summed samples of this work with Marino's) are shown in blue and red, respectively.

The metal-poor population A and metal-rich population B are related to a first and second generation in the cluster formation process. Carretta et al. (2009) have shown that 47-49% of the stars in globular clusters are from first generation. Other authors argued that $\sim 50\%$ is an upper limit for the amount of first generation stars in a clusters (Vesperini et al., 2010, and references therein). If it is true that second generation stars are predominant in all clusters, more observations should be done to complete Figure 5.7. However, Carretta et al. (2009) observed only brighter RGB stars, and the histograms in Figure 5.7 show that for stars with $M_V < 12$ mag, population A (first generation, metal-poor) represents 45% of the total, which is in agreement with their results. Vesperini et al. (2010) proceeded hydrodynamical simulations to find that 4 - 9% of the field halo second generation stars were originated in globular clusters and then stripped to the halo. This process could be responsible for the non-uniform sample of first and second generation stars in the RGB of M 22.

The shapes of the observed distributions of $[\text{Fe}/\text{H}]$ and $[\text{Mg}/\text{H}]$ seem to be similar, displaced by ~ 0.5 dex, which is coherent with the derived average $[\text{Mg}/\text{Fe}]$. If this displacement is approximately constant from Figure 5.6, the variance in $[\text{Mg}/\text{Fe}]$ should be

Figure 5.8: Chemical evolution history of M 22. Purple filled circles are M 22 stars from population A ($[\text{Fe}/\text{H}] < -1.8$), brown filled squares are M 22 stars from population B ($[\text{Fe}/\text{H}] > -1.8$). Grey filled circles are from dwarf spheroidal galaxies (compilation of Kirby et al., 2010) in the upper panel, and from Milky Way stars (compilation of Venn et al., 2004). Dashed lines are guidelines for the trends observed in dSph and MW stars.

negligible. In fact, Figure 5.8 shows clearly a dispersion in $[\text{Fe}/\text{H}]$ and no dispersion in $[\text{Mg}/\text{Fe}]$. In the upper panel of Figure 5.8 we compare M 22 stars with stars from dwarf spheroidal galaxies (Kirby et al., 2010), and in the lower panel we compare the same M 22 stars with Milky Way stars (Venn et al., 2004). The trends in this plot reveal the efficiency of star formation in a galaxy, i.e., if the efficiency is high, many type II supernovae enrich the interstellar medium with Mg and Fe up to high metallicities before type Ia supernovae start to explode providing Fe and no Mg (which starts to decrease the $[\text{Mg}/\text{Fe}]$ ratio). For the Milky Way this break occurs at about $[\text{Fe}/\text{H}] \sim -1.0$ (bottom panel of Figure 5.8), and for dwarf galaxies this knee is located at lower metallicities, as shown in the upper panel of Figure 5.8. M 22 stars have $-2.0 < [\text{Fe}/\text{H}] < -1.5$, approximately, and for that range of metallicities, SNIa already started to explode. $[\text{Mg}/\text{Fe}]$ is constant for both populations A and B. It means that the more metal-rich population B should have been enriched by

SNII, and M 22 possibly was formed in a galaxy with a higher star formation rate, such as the Milky Way. This evidence that M 22 possibly was not originated in a nucleus of a disrupted dwarf galaxy as ω Cen or M 54 is in agreement with the findings and conclusions of Marino et al. (2011) who analysed in detail chemical abundances of 35 M 22 red giant stars using high-resolution spectroscopy. They proposed that M 22 is the result of a merger between two clusters and discuss the possibility of this to happen in the Milky Way halo or in a dwarf galaxy.

5.4.2 NGC 5824

We have selected 15 member stars from the 17 observed red giant stars of NGC 5824 from which we derived an average radial velocity of -35 ± 12 km/s. This is consistent with Da Costa et al. (2014) who derived -28.9 ± 1.0 km/s based on R 2,000 spectra in the CaT region for 147 red giant stars of the cluster. The MDF and [Mg/H] distribution of NGC 5824 using our results are presented in Figure 5.9, in the same way as done for M 22 in the previous section. We derived a mean $[\text{Fe}/\text{H}] = -1.995 \pm 0.024$, and Da Costa et al. (2014) found a trimodal distribution in [Fe/H] with peaks at about -2.09, -2.01 and -1.91 with a metal-rich tail up to ~ -1.8 and metal-poor stars down to ~ -2.2 . Their three peaks are within our main peak that is not resolved in this work because of the lower number of stars observed. Besides, the spread in [Fe/H] is much lower than that of M 22, and our metallicity errors for each star would explain such small dispersion, as can be seen in Figure 5.9.

The [Mg/H] distribution is similar to the MDF, as shown in Figure 5.9, which indicates that [Mg/Fe] should be constant for all observed stars in NGC 5824. In fact, in Figure 5.10 we show that all stars are concentrated and it is difficult to find any dispersion from this data only. The selected stars from Da Costa et al. (2014) could be observed in the optical region in order to derive [Mg/Fe] and discuss in more detail the chemical evolution of this cluster.

Figure 5.9: Same as Figure 5.6 for NCG 5824.

Figure 5.10: Same as Figure 5.8 for NGC 5824.

5.5 Summary and conclusions

From the 44 selected red giant stars, we confirmed and characterized the two populations of M 22 with metallicity peaks at $[\text{Fe}/\text{H}] \approx -1.9$ and -1.6 , in a total range of $-2.3 < [\text{Fe}/\text{H}] < -1.3$. Also for $[\text{Mg}/\text{H}]$ we found a similar distribution with peaks at $[\text{Mg}/\text{H}] \approx -1.5$ and -1.3 , in a total range of $-1.9 < [\text{Fe}/\text{H}] < -0.9$. From these results, M 22 has an average $[\text{Mg}/\text{Fe}] = 0.43 \pm 0.04$ with negligible star-to-star variations, i.e, both populations have the same $[\text{Mg}/\text{Fe}]$ ratio. This means that the second generation stars were enriched by core-collapse supernovae and not by type Ia supernovae, as one could expect if M 22 would had been formed in a dwarf galaxy. We conclude that if M 22 had an extragalactic origin, it should be in a galaxy with efficient star formation. It is unlikely the Milky Way halo would have had enough density to produce 1.0 dex range of metallicity within a globular cluster. Further chemo-dynamical observations and models must be performed to explain if such cluster can be formed in another environment but a dwarf galaxy with high star formation rate.

For NGC 5824 we found a symmetrical metallicity distribution function that peaks at $[\text{Fe}/\text{H}] \sim -2.0$ and covers $-2.2 < [\text{Fe}/\text{H}] < -1.8$. This distribution is based on 15 selected member red giant stars and it is close to the distribution found by Da Costa et al. (2014). Once they selected 147 members, they could resolve the main peak in the MDF presented in this work into three peaks of metallicity. The MDF found here is also compatible with $[\text{Fe}/\text{H}]$ errors of each star. $[\text{Mg}/\text{H}]$ has a similar distribution with main peak at $[\text{Mg}/\text{H}] \sim -1.5$ and ranging $-1.8 < [\text{Mg}/\text{H}] < -1.3$. This is consistent with an average $[\text{Mg}/\text{Fe}] = 0.436 \pm 0.030$. Although compatible with chemical enrichment of dwarf galaxies, if this cluster was formed in a dSph, its $[\text{Mg}/\text{Fe}]$ ratio could be higher. The adopted stellar libraries MILES and COELHO does not have stars with alpha-enhancement as high as $[\text{Mg}/\text{Fe}] \sim 1.0$, and further spectroscopic analysis must be executed in order to establish the origin of NGC 5824.

RR Lyrae variable stars in HP1

6.1 *Introduction to this chapter*

If bulge experienced a more efficient star formation and it hosts the centre of the Milky Way, it is possible that the oldest globular clusters are located there. In particular, there are some moderately metal-poor bulge clusters presenting a bluer horizontal branch (HB) than expected by their metallicity, as listed by Barbuy et al. (2009). The main parameter that causes a change in the HB morphology is the metallicity, known as the first parameter. The second most important parameter is age in most cases (Lee et al., 1994), although other parameters can play the role, such as He abundance, CNO abundance or AGB mass loss. Metal-poor clusters have a blue HB, and old clusters too, but these bulge clusters are moderately metal-poor with a blue HB, which indicates that they are very old, possibly the oldest objects of the Milky Way, witnesses of the initial collapse of the Galaxy.

The innermost Milky Way globular cluster is HP 1, for which we determine a metallicity of $[\text{Fe}/\text{H}] = -1.3 \pm 0.4$ in Chapter 4, has a HB index¹ of 0.75, as determined by Mackey and van den Bergh (2005). Therefore this central cluster seems to have a bluer horizontal branch than expected. If this cluster is the oldest of our Galaxy it will confirm that the bulge had a rapid star formation at the beginning of the Milky Way. To better constrain the age determination via colour-magnitude diagram, it is important to have a precise estimation of distance from a different source. Moreover, the distance itself is important to define better the position in relation to the Galactic centre and correlate with bulge formation and evolution models. Ortolani et al. (2011) published a cleaned CMD for this

¹ $(\text{B}-\text{R})/(\text{B}+\text{V}+\text{R})$, where V is the number of variable stars, B and R are the numbers of stars bluer and redder than the variables, respectively.

bulge cluster, using proper motion techniques to remove field stars contamination and indicated six RR Lyrae candidates. We aim at measuring light curves of the six RR Lyrae candidates, determine their periods and the distance of the cluster, using V,I photometry from EFOSC2@NTT.

6.2 Observations

The observations were performed with EFOSC2@NTT/ESO-La Silla in May 14, 2013, under project 091.D-0711(A), PI. Sergio Ortolani. Once typical periods of RR Lyrae are of a few hours, 18 images were taken in a range of six hours, trying to keep an almost regular time step. Filter Gunn-i was used for these images. PSF photometry using DAOPhot@IRAF was performed, and relative magnitudes were obtained in order to construct the light curves. In Table 6.1 the log of observations is given.

6.3 RR Lyrae candidates

In Figure 6.1 we show the V vs. V-I Colour-Magnitude Diagram (CMD) of HP 1. The original data was collected with the NTT (Ortolani et al., 1997), and later it was corrected for proper motion using the JHK VLT/MAD data (Ortolani et al., 2011). Six candidate stars are located in the RR Lyrae gap region. Photometry of star #2 was not good, and this star is not considered in the analysis.

6.4 Period determination

One of the most used techniques to determine the period of a light-curve is Phase Dispersion Minimization (PDM), which is available within IRAF. The best period for each curve was found using PDM, as shown in Figures 6.3 and 6.4. The folded curves using the best period found are plotted in Figure 6.5.

6.5 Constraining the distance

From the period-luminosity relationship by Catelan et al. (2004) combined with the distance modulus formula it is possible to derive the distance of the RR Lyrae stars, as follows:

Table 6.1 - Log of observations

obs. date	exp. time	filter	seeing	airmass
block # 1				
2013-05-14T04:17:51.712	30.0005	i-705	0.92	1.181
2013-05-14T04:19:44.535	30.0005	i-705	0.92	1.176
2013-05-14T04:22:45.035	180.0005	i-705	0.92	1.168
2013-05-14T04:27:07.647	180.0006	i-705	0.84	1.157
2013-05-14T04:31:35.321	180.0005	i-705	0.84	1.146
2013-05-14T04:35:58.353	167.0005	i-705	0.81	1.136
block # 2				
2013-05-14T05:31:10.934	30.0005	i-705	0.70	1.042
2013-05-14T05:33:09.520	180.0006	i-705	0.70	1.040
block # 3				
2013-05-14T07:24:18.491	30.0005	i-705	0.61	1.011
2013-05-14T07:26:16.036	180.0005	i-705	0.61	1.012
block # 4				
2013-05-14T08:10:24.651	30.0005	i-705	0.53	1.053
2013-05-14T08:12:22.536	180.0005	i-705	0.53	1.056
block # 5				
2013-05-14T08:55:37.961	30.0005	i-705	0.82	1.133
2013-05-14T08:57:35.486	180.0005	i-705	0.82	1.138
block # 6				
2013-05-14T09:26:43.353	30.0005	i-705	0.63	1.217
2013-05-14T09:28:41.128	180.0006	i-705	0.72	1.223
block # 7				
2013-05-14T10:10:59.721	30.0005	i-705	0.69	1.394
2013-05-14T10:12:57.406	180.0005	i-705	0.69	1.404

$$M_V = a + b \cdot \log P, \text{ where}$$

$$a = \sum_{i=0}^3 a_i \cdot (L)^i \quad (6.1)$$

$$b = \sum_{i=0}^3 b_i \cdot (L)^i$$

The parameter $L = (B - R)/(B + V + R)$ characterizes the ratio of (B)lue, (R)ed, and (V)ariable stars, and has the value 0.75 for HP1 (Mackey and van den Bergh, 2005). The coefficients a_i and b_i were fitted by Catelan et al. (2004) for four different values of metallicity Z: 0.0060, 0.0020, 0.0010, 0.0005. Distance can be derived from the difference between apparent and absolute magnitude, which results in the equation below:

$$d(pc) = 10^{(m_V - M_V + 5)/5} \quad (6.2)$$

Figure 6.1: Colour magnitude diagram showing six RR Lyrae candidates.

Figure 6.2: HP 1: Finding chart of the six RR Lyrae candidates. Central extraction of the NTT I image.

Combination of Equations 6.1 and 6.2 give the direct relationship between period and distance. Figure 6.6 shows the parameter space for distance as a function of period for different magnitudes and metallicities.

Assuming a cluster metallicity of $[\text{Fe}/\text{H}] = -1.26 \pm 0.39$ from Chapter 4 and the solar metallicity to be $Z = 0.0153$ (Caffau et al., 2011), we get $Z = 0.0008$ for the present RR Lyrae. A reddening $E(B-V) = 1.12$ (Harris, 1996, 2010 edition) is adopted, that together

Figure 6.3: Light curves of five RR Lyrae candidates in HP1.

with $R_V = 3.1$ gives $A_V = 3.472$. V magnitudes corrected by this value are presented in Table 6.2, together with values for periods, and resulting distances using the fits for $Z=0.0010$ and $Z=0.0020$ by Catelan et al. (2004).

Table 6.2 - Distance determination. V magnitude is already corrected by extinction of $A_V = 3.1 \times E(B-V) = 3.472$, where $E(B-V) = 1.12$ is from (Harris, 1996, 2010 edition).

ID	period (days)	m_V	Δm_i	distA (kpc)	distB (kpc)
1	0.129	15.429	0.093	7.1	6.9
2	—	15.347	—	—	—
3	0.406	15.404	0.640	8.8	8.5
4	0.102	15.398	0.254	6.6	6.5
5	0.270	15.216	0.760	7.5	7.2
6	0.243	15.515	0.499	8.4	8.1

Star #1 has no magnitude variations, star #2 is so faint that the photometry did not

Figure 6.4: Theta vs. period for five RR Lyrae candidates in HP1.

recover its magnitudes, star #4 is undersampled. The other three stars (#3, #5, #6) present magnitude variations above 0.5 mag in Gunn-i, and well sampled light-curves in the phase range. The periods and amplitudes reported in Table 6.2 imply, from the Bailey diagram, that the five RR Lyrae stars are of type c (e.g. Bono et al., 1997). An average distance of HP1 based on the 3 confirmed RR Lyrae is then 8.0 ± 0.6 kpc.

6.6 Conclusions

We present for the first time 3 confirmed RR Lyrae stars for the inner Galactic bulge globular cluster. Combining their periods with magnitude, metallicity, horizontal branch parameter, we could apply a period-luminosity relation and derive the distance of 8.0 ± 0.6 kpc for HP1. This is consistent with the distance of the Galactic center to the Sun, as well as with distance of this cluster in the catalog of (Harris, 1996, 2010 edition), which

Figure 6.5: Folded light curves for five RR Lyrae candidates in HP1, using the best periods found in Figure 6.4.

is 8.2 kpc. A comparison with VVV survey data is under way, in order to conclude this project.

Figure 6.6: Theoretical curves showing Equations 6.1 and 6.2.

Conclusions and future work

7.1 *Summary*

In this thesis we use intermediate-age and old star clusters in the Small Magellanic Cloud and globular clusters in the Milky Way as tracers of the chemical and dynamical evolution of these galaxies. Since these clusters are spatially resolved they can be studied in detail and the system Milky Way - SMC - LMC becomes a good laboratory for studying galaxy interactions, giving hints on how distant galaxies evolve.

In Chapter 2 we derive ages and metallicities for five star clusters in the SMC and we defined a particular group of clusters called West Halo. Four clusters are in this region and present a gradient in age and metallicity, which indicates that the scenario of tidal stripping in the SMC is true, instead of ram pressure which only affects gas content of the galaxy. Other clusters in this region have ages and metallicities published in the literature and also corroborates the gradients.

The FORS2/VLT survey of Milky Way globular clusters is presented in Chapters 3 and 4. In the first one, we detail the method for derivation of parameters (v_{helio} , T_{eff} , $\log(g)$, $[\text{Fe}/\text{H}]$, $[\text{Mg}/\text{Fe}]$ and $[\alpha/\text{Fe}]$) from mid-resolution stellar spectra ($R \sim 2000$, $\lambda\lambda$ 456 - 586 nm) based on full spectrum fitting code ETOILE with reference stellar libraries the observed MILES and the synthetic COELHO. We validate the method recovering the parameters of 49 well-known stars in the whole parameter space for our FORS2 targets using spectra from ELODIE library and parameters from PASTEL catalogue. Six clusters were used as template with metallicities varying from -2.5 to solar, and the resulting $[\text{Fe}/\text{H}]$ are in agreement with high-resolution spectroscopic determinations and also with CaT metallicities for the same stars.

In the following Chapter 4, the whole sample of about 800 red giant stars distributed over the 51 Galactic globular clusters is analysed. We present a new metallicity scale for $\sim 1/3$ of the total number of catalogued clusters, as well as alpha-element enhancement indicator $[\text{Mg}/\text{Fe}]$ for all clusters with homogeneous observations and analysis, recovering the bimodal metallicity distribution for globular clusters. We fit a linear function with Carretta et al. (2009) metallicity scale based on 13 clusters in common with their observations, and the dispersion of the comparison is only $\sigma = 0.07$ dex. Carretta et al. (2009) compiled metallicities from literature and put them in their scale, however comparing these values with our derivations we have a higher dispersion of $\sigma = 0.16$ dex. We conclude that if all 51 clusters were analysed with high-resolution spectroscopy, the comparison with our results would keep the dispersion of $\sigma = 0.07$ dex, therefore our metallicities are more precise and self-consistent than previous works based on low-resolution spectroscopy. Six clusters from our sample were not calibrated by Carretta et al. (2009), and we present a complete analysis for them: NGC 6426, Terzan 8, Djorg 2, Lynga 7, Pal 10 and BH 176. It is the first time that $[\text{Fe}/\text{H}]$ and $[\text{Mg}/\text{Fe}]$ derived in a consistent way are published for such a huge sample of globular clusters.

The behaviour of $[\text{Mg}/\text{Fe}]$ vs. $[\text{Fe}/\text{H}]$ of the 51 clusters are similar to their field stars counterparts in the bulge, disc, inner and outer halo. $[\text{Mg}/\text{Fe}]$ starts to decrease at $[\text{Fe}/\text{H}]_{\text{knee}} \sim -1.0$ for all components, although we do not exclude the possibility of the bulge having $[\text{Fe}/\text{H}]_{\text{knee}}$ at a higher metallicity showing that it had a more efficient star formation. Comparing our results with ages for the clusters, the timescale for SNIa to become important is about 1 Gyr for all components in the Milky Way which corroborates the value adopted vastly in Galactic models. $[\text{Fe}/\text{H}]$ vs. age exhibit a difference between bulge and halo clusters that is interpreted as a more efficient star formation in the bulge than in the halo. This should be reflected also in $[\text{Mg}/\text{Fe}]$ vs. age but this comparison does not allow a clear separation between bulge and halo clusters. A metallicity gradient for the halo and a different one for the bulge/disc populations are found and reflects different origins of these clusters. Halo clusters would have been assembled in the building blocks of the primordial Galaxy, while bulge/disc clusters would have been formed *in situ*. This is consistent with a more rapid star formation in the bulge than in the halo.

In the context of understanding the origin of halo globular clusters and discuss if the environment of less efficient star formation was a dwarf galaxy or a mini halo, we study

two interesting massive clusters from our FORS2 survey in Chapter 5, namely M 22 and NGC 5824. These clusters present a dispersion in $[\text{Fe}/\text{H}]$ and in $[\text{Mg}/\text{H}]$ which indicates a different formation scenario for them, possibly related to a disrupted dwarf galaxy captured by the Milky Way potential well. M 22 has peaks at $[\text{Fe}/\text{H}] \approx -1.9$ and -1.6 , and $[\text{Mg}/\text{H}] \approx -1.5$ and -1.3 . $[\text{Fe}/\text{H}]$ and $[\text{Mg}/\text{H}]$ range 1.0 dex and average uncertainty of individual stars is $\sigma_{[\text{Fe}/\text{H}]} = 0.14$ dex and $\sigma_{[\text{Mg}/\text{H}]} = 0.20$ dex. This cluster seems to follow an internal chemical evolution closer to that of the Milky Way, instead of dwarf galaxies, which indicates that the formation scenario of M 22 is more complex than previously thought, possibly formed by the merger of two other clusters as proposed by Marino et al. (2011). For NGC 5824 there is only one peak at $[\text{Fe}/\text{H}] \sim -2.0$ and $[\text{Mg}/\text{H}] \sim -1.5$. The ranges are 0.4 dex and 0.5 dex, respectively, with average uncertainty of individual stars of $\sigma_{[\text{Fe}/\text{H}]} = 0.11$ dex and $\sigma_{[\text{Mg}/\text{H}]} = 0.20$ dex. In fact, Da Costa et al. (2014) showed that a inter-quartile range for the $[\text{Fe}/\text{H}]$ in this clusters is only 0.10 dex, which is of the order of the uncertainties in the present work. Consequently, a cluster with a large spread in $[\text{Fe}/\text{H}]$ such as M 22 probably was not originated in a dwarf galaxy, and a cluster that seems to share a similar origin with Cetus Polar Stream, possibly a disrupted dwarf galaxy has a very small spread in $[\text{Fe}/\text{H}]$ such as NGC 5824. This reinforces that each cluster with spread in $[\text{Fe}/\text{H}]$ seems to have a different origin and must be analysed independently of the others.

Barbuy et al. (2009) listed some moderately metal-poor bulge clusters presenting a blue horizontal branch (HB). This horizontal branch morphology can be due to very metal-poor (main parameter) or very old stars (second parameter), as modelled by Lee et al. (1994), among other weaker effects (other options for second parameter effect, such as He abundance, CNO abundance, AGB mass lost etc). Once these clusters are not too metal-poor, they could be the oldest objects in the Milky Way, which would be in agreement with a classical component of the bulge, as discussed in Section 1.1.1. We analyse in detail the innermost Galactic globular cluster, HP 1 and determine $[\text{Fe}/\text{H}] = -1.3 \pm 0.4$. Mackey and van den Bergh (2005) published its HB index equal to $+0.75$. Therefore this central cluster seems to have a bluer horizontal branch than expected. To constrain future age determination using colour-magnitude diagrams we determine its distance using light curves of RR Lyrae in Chapter 6. We find a distance of 8.0 ± 0.6 kpc consistent with the distance of the Galaxy centre. If this cluster is the oldest of our Galaxy it will confirm that the bulge had a rapid star formation at the beginning of the Milky Way.

7.2 Perspectives

The SMC West Halo clusters that do not have ages and metallicities measured were observed by us and the data are under analysis to get a general picture of the West Halo of the SMC and to constrain models of tidal interaction between the Magellanic Clouds and our Galaxy. The list of newly observed clusters is: B 1, B 2, B 4, B 6, BS 1, HW 5, HW 6, Kron 8, Kron 11, Lindsay 14 and a calibration cluster NGC 152.

For the FORS2/VLT survey of globular clusters further discussion should be address to explain why $[\text{Fe}/\text{H}]$ vs. age and $[\text{Fe}/\text{H}]$ vs. $[\text{Mg}/\text{Fe}]$ indicate different efficiency of star formation for bulge and halo populations, but looking at $[\text{Mg}/\text{Fe}]$ vs. age both population seems to share similar histories. Only few clusters were in common with large samples of homogeneous age determinations, that being the case we intend to use the pre-images observed to select our targets to produce calibrated colour-magnitude diagrams to derive ages for all the 51 clusters and trace a complete scenario for these objects.

NGC 5824 follows the orbit of the Cetus Polar Stream and it was possibly related to a disrupted dwarf galaxy in the past. Detailed chemistry of this cluster have to be analysed in order to better characterize its origin as the nucleus of a disrupted dwarf galaxy or not.

For the innermost bulge globular cluster HP 1, more images were observed at SOI/SOAR in 2013 and also at EFOSC2/NTT in 2014, and they should be analysed soon to get better sampled light curves and improve the precision of distance determination. An independent analysis of its distance have to be done using VVV/ESO¹ photometry in K_S band to make well sampled light curves using a different photometric filter and determine the cluster distance, that should not be so different of our findings here. Having its distance and metallicity defined using independent methods, the next step it to observe HP 1 with the Hubble Space Telescope to make a colour-magnitude diagram reaching the main sequence turnoff and measure its age and prove if this is (among) the oldest object in the Milky Way.

In conclusion, the techniques developed in this thesis to use full spectrum fitting to derive $[\text{Fe}/\text{H}]$ and $[\alpha/\text{Fe}]$ can be applied to cluster systems in other galaxies and even for entire dwarf galaxies. One first target would be the system of stellar clusters of the SMC. From our works on photometry of SMC clusters, we have ages in a homogeneous scale,

¹ http://mwm.astro.puc.cl/mw/index.php/Main_Page

and we also have selected RGB stars to be observed with spectroscopy. We can study a scenario of efficiency of star formation in the SMC in terms of $[\alpha/\text{Fe}]$ vs. $[\text{Fe}/\text{H}]$. With the upcoming giant 30-40m class telescopes (E-ELT, GMT, TMT) we will be able to observe individual stars up to several magnitudes fainter than now. As we need stellar libraries for reference, and observed libraries are based on solar neighbourhood with a specific chemical evolution history and a set of solar abundances, this analysis will need synthetic spectra as reference. The observation of individual stars in extragalactic globular clusters will encourage the development of synthetic stellar libraries, in particular with different patterns of chemical abundances and alpha-enhancement elements.

Bibliography

- Aguerri J. A. L., Balcells M., Peletier R. F., Growth of galactic bulges by mergers. I. Dense satellites, *A&A*, 2001, vol. 367, p. 428
- Allende Prieto C., Sivarani T., Beers T. C., Lee Y. S., Koesterke L., Shetrone M., Sneden C., Lambert D. L., Wilhelm R., Rockosi C. M., Lai D. K., Yanny B., Ivans I. I., Johnson J. A., Aoki W., Bailer-Jones C. A. L., Re Fiorentin P., The SEGUE Stellar Parameter Pipeline. III. Comparison with High-Resolution Spectroscopy of SDSS/SEGUE Field Stars, *AJ*, 2008, vol. 136, p. 2070
- Alves V. M., Pavani D. B., Kerber L. O., Bica E., On open cluster physical parameters from 2MASS data, *New A*, 2012, vol. 17, p. 488
- Alves-Brito A., Barbuy B., Zoccali M., Minniti D., Ortolani S., Hill V., Renzini A., Pasquini L., Bica E., Rich R. M., Meléndez J., Momany Y., VLT-UVES abundance analysis of four giants in NGC 6553, *A&A*, 2006, vol. 460, p. 269
- Alves-Brito A., Meléndez J., Asplund M., Ramírez I., Yong D., Chemical similarities between Galactic bulge and local thick disk red giants: O, Na, Mg, Al, Si, Ca, and Ti, *A&A*, 2010, vol. 513, p. A35
- Alves-Brito A., Yong D., Meléndez J., Vásquez S., Karakas A. I., CNO and F abundances in the globular cluster M 22 (NGC 6656), *A&A*, 2012, vol. 540, p. A3
- Baade W., The Resolution of Messier 32, NGC 205, and the Central Region of the Andromeda Nebula., *ApJ*, 1944, vol. 100, p. 137
- Babusiaux C., Gómez A., Hill V., Royer F., Zoccali M., Arenou F., Fux R., Lecureur A., Schultheis M., Barbuy B., Minniti D., Ortolani S., Insights on the Milky Way bulge

- formation from the correlations between kinematics and metallicity, *A&A*, 2010, vol. 519, p. A77
- Balbinot E., Santiago B. X., Kerber L. O., Barbuy B., Dias B. M. S., Probing the Large Magellanic Cloud age gap at intermediate cluster masses, *MNRAS*, 2010, vol. 404, p. 1625
- Barbuy B., Bica E., Ortolani S., Globular clusters within 5(deg) of the Galactic center, *A&A*, 1998, vol. 333, p. 117
- Barbuy B., Zoccali M., Ortolani S., Hill V., Minniti D., Bica E., Renzini A., Gómez A., VLT-FLAMES analysis of 8 giants in the bulge metal-poor globular cluster NGC 6522: oldest cluster in the Galaxy?. Analysis of 8 giants in NGC 6522, *A&A*, 2009, vol. 507, p. 405
- Barbuy B., Zoccali M., Ortolani S., Minniti D., Hill V., Renzini A., Bica E., Gómez A., NGC 6558: A Blue Horizontal Branch Moderately Metal-Poor Globular Cluster in the Bulge, *AJ*, 2007, vol. 134, p. 1613
- Barnes J. E., Memory in Mergers Major and Minor. In Formation of the Galactic Halo...Inside and Out , vol. 92 of Astronomical Society of the Pacific Conference Series, 1996, p. 415
- Bekki K., Freeman K. C., Formation of ω Centauri from an ancient nucleated dwarf galaxy in the young Galactic disc, *MNRAS*, 2003, vol. 346, p. L11
- Bellazzini M., Ibata R. A., Chapman S. C., Mackey A. D., Monaco L., Irwin M. J., Martin N. F., Lewis G. F., Dalessandro E., The Nucleus of the Sagittarius Dsph Galaxy and M54: a Window on the Process of Galaxy Nucleation, *AJ*, 2008, vol. 136, p. 1147
- Bensby T., Feltzing S., Lundström I., Ilyin I., α -, r-, and s-process element trends in the Galactic thin and thick disks, *A&A*, 2005, vol. 433, p. 185
- Besla G., Hernquist L., Loeb A., The origin of the microlensing events observed towards the LMC and the stellar counterpart of the Magellanic stream, *MNRAS*, 2013, vol. 428, p. 2342

-
- Bica E., Bonatto C., Dutra C. M., Santos J. F. C., A general catalogue of extended objects in the Magellanic System, *MNRAS*, 2008, vol. 389, p. 678
- Bono G., Caputo F., Castellani V., Marconi M., Nonlinear investigation of the pulsational properties of RR Lyrae variables, *A&AS*, 1997, vol. 121, p. 327
- Borissova J., Bonatto C., Kurtev R., Clarke J. R. A., Peñaloza F., Sale S. E. e. a., New Galactic star clusters discovered in the VVV survey, *A&A*, 2011, vol. 532, p. A131
- Bressan A., Marigo P., Girardi L., Salasnich B., Dal Cero C., Rubele S., Nanni A., PAR-SEC: stellar tracks and isochrones with the PAdova and TRieste Stellar Evolution Code, *MNRAS*, 2012, vol. 427, p. 127
- Broeils A. H., van Woerden H., A search for spiral galaxies with extended HI disks., *A&AS*, 1994, vol. 107, p. 129
- Caffau E., Ludwig H.-G., Steffen M., Freytag B., Bonifacio P., Solar Chemical Abundances Determined with a CO5BOLD 3D Model Atmosphere, *Sol. Phys.*, 2011, vol. 268, p. 255
- Carretta E., Bragaglia A., Gratton R., D'Orazi V., Lucatello S., Intrinsic iron spread and a new metallicity scale for globular clusters, *A&A*, 2009, vol. 508, p. 695
- Carretta E., Bragaglia A., Gratton R., Lucatello S., Na-O anticorrelation and HB. VIII. Proton-capture elements and metallicities in 17 globular clusters from UVES spectra, *A&A*, 2009, vol. 505, p. 139
- Carretta E., Bragaglia A., Gratton R. G., D'Orazi V., Lucatello S., Sollima A., Terzan 8: a Sagittarius-flavoured globular cluster, *A&A*, 2014, vol. 561, p. A87
- Carretta E., Bragaglia A., Gratton R. G., Lucatello S., Bellazzini M., Catanzaro G., Leone F., Momany Y., Piotto G., D'Orazi V., M54 + Sagittarius = ω Centauri, *ApJ*, 2010, vol. 714, p. L7
- Carretta E., Bragaglia A., Gratton R. G., Lucatello S., Catanzaro G., Leone F., Bellazzini M., Claudi R., D'Orazi V., Momany Y., Ortolani S., Pancino E., Piotto G., Recio-Blanco A., Sabbi E., Na-O anticorrelation and HB. VII. The chemical composition of first and second-generation stars in 15 globular clusters from GIRAFFE spectra, *A&A*, 2009, vol. 505, p. 117

- Carretta E., Bragaglia A., Gratton R. G., Recio-Blanco A., Lucatello S., D'Orazi V., Cassisi S., Properties of stellar generations in globular clusters and relations with global parameters, *A&A*, 2010, vol. 516, p. A55
- Carretta E., Cohen J. G., Gratton R. G., Behr B. B., An Abundance Analysis for Four Red Horizontal-Branch Stars in the Extremely Metal-Rich Globular Cluster NGC 6528, *AJ*, 2001, vol. 122, p. 1469
- Carretta E., Gratton R. G., Lucatello S., Bragaglia A., Catanzaro G., Leone F., Momany Y., D'Orazi V., Cassisi S., D'Antona F., Ortolani S., Abundances for a Large Sample of Red Giants in NGC 1851: Hints for a Merger of Two Clusters?, *ApJ*, 2010, vol. 722, p. L1
- Casetti-Dinescu D. I., Moni Bidin C., Girard T. M., Méndez R. A., Vieira K., Korchagin V. I., van Altena W. F., Recent Star Formation in the Leading Arm of the Magellanic Stream, *ApJ*, 2014, vol. 784, p. L37
- Catelan M., Ferraro F. R., Rood R. T., Horizontal-Branch Models and the Second-Parameter Effect. IV. The Case of M3 and Palomar 3, *ApJ*, 2001, vol. 560, p. 970
- Catelan M., Pritzl B. J., Smith H. A., The RR Lyrae Period-Luminosity Relation. I. Theoretical Calibration, *ApJS*, 2004, vol. 154, p. 633
- Cayrel R., Data Analysis. In *The Impact of Very High S/N Spectroscopy on Stellar Physics*, vol. 132 of IAU Symposium, 1988, p. 345
- Cayrel R., Perrin M.-N., Barbuy B., Buser R., A grid of synthetic spectra for the determination of effective temperature, gravity and metallicity of F, G, and K stars. I - Description of the method. II - Application to 41 stellar spectra taken in the Basel field of SA 141, *A&A*, 1991, vol. 247, p. 108
- Cenarro A. J., Peletier R. F., Sánchez-Blázquez P., Selam S. O., Toloba E., Cardiel N., Falcón-Barroso J., Gorgas J., Jiménez-Vicente J., Vazdekis A., Medium-resolution Isaac Newton Telescope library of empirical spectra - II. The stellar atmospheric parameters, *MNRAS*, 2007, vol. 374, p. 664

- Cignoni M., Cole A. A., Tosi M., Gallagher J. S., Sabbi E., Anderson J., Grebel E. K., Nota A., Star Formation History in Two Fields of the Small Magellanic Cloud Bar, *ApJ*, 2012, vol. 754, p. 130
- Cignoni M., Cole A. A., Tosi M., Gallagher J. S., Sabbi E., Anderson J., Grebel E. K., Nota A., Mean Age Gradient and Asymmetry in the Star Formation History of the Small Magellanic Cloud, *ApJ*, 2013, vol. 775, p. 83
- Cioni M.-R. L., Clementini G., Girardi L., Guandalini R., Gullieuszik M., Miszalski B., Moretti M.-I., Ripepi V., Rubele S., Bagheri G. e. a., The VMC survey. I. Strategy and first data, *A&A*, 2011, vol. 527, p. A116
- Coelho P., Barbuy B., Meléndez J., Schiavon R. P., Castilho B. V., A library of high resolution synthetic stellar spectra from 300 nm to 1.8 μ m with solar and α -enhanced composition, *A&A*, 2005, vol. 443, p. 735
- Cohen J. G., Behr B. B., Briley M. M., Abundances in Stars from the Red Giant Branch Tip to Near the Main-Sequence Turnoff in M71. I. Sample Selection, Observing Strategy, and Stellar Parameters, *AJ*, 2001, vol. 122, p. 1420
- Cohen J. G., Gratton R. G., Behr B. B., Carretta E., An Abundance Analysis for Five Red Horizontal-Branch Stars in the Extremely Metal-rich Globular Cluster NGC 6553, *ApJ*, 1999, vol. 523, p. 739
- Cohen J. G., Kirby E. N., Simon J. D., Geha M., NGC 2419-Another Remnant of Accretion by the Milky Way, *ApJ*, 2010, vol. 725, p. 288
- Crowl H. H., Sarajedini A., Piatti A. E., Geisler D., Bica E., Clariá J. J., Santos Jr. J. F. C., The Line-of-Sight Depth of Populous Clusters in the Small Magellanic Cloud, *AJ*, 2001, vol. 122, p. 220
- Da Costa G. S., AM-3: An Intermediate-Age Star Cluster in the Extreme Outskirts of the SMC. In *New Views of the Magellanic Clouds*, vol. 190 of IAU Symposium, 1999, p. 446
- Da Costa G. S., Armandroff T. E., Abundances and Kinematics of the Globular Cluster Systems of the Galaxy and of the Sagittarius Dwarf, *AJ*, 1995, vol. 109, p. 2533

- Da Costa G. S., Hatzidimitriou D., Ca II Triplet Spectroscopy of Giants in Small Magellanic Cloud Star Clusters: Abundances, Velocities, and the Age-Metallicity Relation, *AJ*, 1998, vol. 115, p. 1934
- Da Costa G. S., Held E. V., Saviane I., NGC 5824: a luminous outer halo globular cluster with an intrinsic abundance spread, *MNRAS*, 2014, vol. 438, p. 3507
- Da Costa G. S., Held E. V., Saviane I., Gullieuszik M., M22: An [Fe/H] Abundance Range Revealed, *ApJ*, 2009, vol. 705, p. 1481
- Davies R. D., Wright A. E., A tidal origin for the Magellanic Stream, *MNRAS*, 1977, vol. 180, p. 71
- de Jong J. T. A., Yanny B., Rix H.-W., Dolphin A. E., Martin N. F., Beers T. C., Mapping the Stellar Structure of the Milky Way Thick Disk and Halo Using SEGUE Photometry, *ApJ*, 2010, vol. 714, p. 663
- de Vaucouleurs G., Pence W. D., An outsider's view of the Galaxy - Photometric parameters, scale lengths, and absolute magnitudes of the spheroidal and disk components of our Galaxy, *AJ*, 1978, vol. 83, p. 1163
- Dékány I., Minniti D., Catelan M., Zoccali M., Saito R. K., Hempel M., Gonzalez O. A., VVV Survey Near-infrared Photometry of Known Bulge RR Lyrae Stars: The Distance to the Galactic Center and Absence of a Barred Distribution of the Metal-poor Population, *ApJ*, 2013, vol. 776, p. L19
- Dias B., Coelho P., Barbuy B., Kerber L., Idiart T., Age and metallicity of star clusters in the Small Magellanic Cloud from integrated spectroscopy, *A&A*, 2010, vol. 520, p. A85
- Dias B., Kerber L. O., Barbuy B., Santiago B., Ortolani S., Balbinot E., Self-consistent physical parameters for five intermediate-age SMC stellar clusters from CMD modelling, *A&A*, 2014, vol. 561, p. A106
- Diaz J. D., Bekki K., The Tidal Origin of the Magellanic Stream and the Possibility of a Stellar Counterpart, *ApJ*, 2012, vol. 750, p. 36
- Dotter A., Chaboyer B., Jevremović D., Kostov V., Baron E., Ferguson J. W., The Dartmouth Stellar Evolution Database, *ApJS*, 2008, vol. 178, p. 89

- Dotter A., Sarajedini A., Anderson J., Globular Clusters in the Outer Galactic Halo: New Hubble Space Telescope/Advanced Camera for Surveys Imaging of Six Globular Clusters and the Galactic Globular Cluster Age-metallicity Relation, *ApJ*, 2011, vol. 738, p. 74
- Drinkwater M. J., Jones J. B., Gregg M. D., Phillipps S., Compact stellar systems in the Fornax Cluster: Super-massive star clusters or extremely compact dwarf galaxies?, *PASA*, 2000, vol. 17, p. 227
- Duffau S., Zinn R., Vivas A. K., Carraro G., Méndez R. A., Winnick R., Gallart C., Spectroscopy of QUEST RR Lyrae Variables: The New Virgo Stellar Stream, *ApJ*, 2006, vol. 636, p. L97
- Eggen O. J., Lynden-Bell D., Sandage A. R., Evidence from the motions of old stars that the Galaxy collapsed., *ApJ*, 1962, vol. 136, p. 748
- Evstigneeva E. A., Drinkwater M. J., Peng C. Y., Hilker M., De Propris R., Jones J. B., Phillipps S., Gregg M. D., Karick A. M., Structural Properties of Ultra-Compact Dwarf Galaxies in the Fornax and Virgo Clusters, *AJ*, 2008, vol. 136, p. 461
- Faber S. M., Friel E. D., Burstein D., Gaskell C. M., Old stellar populations. II - an analysis of K-giant spectra, *ApJS*, 1985, vol. 57, p. 711
- Ferraro F. R., Dalessandro E., Mucciarelli A., Beccari G., Rich R. M., Origlia L., Lanzoni B., Rood R. T., Valenti E., Bellazzini M., Ransom S. M., Coccozza G., The cluster Terzan 5 as a remnant of a primordial building block of the Galactic bulge, *Nature*, 2009, vol. 462, p. 483
- Fulbright J. P., Abundances and Kinematics of Field Halo and Disk Stars. I. Observational Data and Abundance Analysis, *AJ*, 2000, vol. 120, p. 1841
- Gadotti D. A., Structural properties of pseudo-bulges, classical bulges and elliptical galaxies: a Sloan Digital Sky Survey perspective, *MNRAS*, 2009, vol. 393, p. 1531
- Gardiner L. T., Noguchi M., N-body simulations of the Small Magellanic Cloud and the Magellanic Stream, *MNRAS*, 1996, vol. 278, p. 191

- Gardiner L. T., Sawa T., Fujimoto M., Numerical Simulations of the Magellanic System - Part One - Orbits of the Magellanic Clouds and the Global Gas Distribution, MNRAS, 1994, vol. 266, p. 567
- Gardner E., Debattista V. P., Robin A. C., Vásquez S., Zoccali M., N-body simulation insights into the X-shaped bulge of the Milky Way: kinematics and distance to the Galactic Centre, MNRAS, 2014, vol. 438, p. 3275
- Gieles M., Lamers H. J. G. L. M., Portegies Zwart S. F., On the Interpretation of the Age Distribution of Star Clusters in the Small Magellanic Cloud, ApJ, 2007, vol. 668, p. 268
- Gilmore G., Randich S., Asplund M., Binney J., Bonifacio P., Drew J., Feltzing S., Ferguson A., The Gaia-ESO Public Spectroscopic Survey, The Messenger, 2012, vol. 147, p. 25
- Glatt K., Grebel E. K., Sabbi E., Gallagher J. S., Nota A., Sirianni M., Clementini G., Tosi M., Harbeck D., Koch A., Kayser A., Da Costa G., Age Determination of Six Intermediate-Age Small Magellanic Cloud Star Clusters with Hst/acs, AJ, 2008, vol. 136, p. 1703
- Gonzalez O. A., Rejkuba M., Zoccali M., Hill V., Battaglia G., Babusiaux C., Minniti D., Barbuy B., Alves-Brito A., Renzini A., Gomez A., Ortolani S., Alpha element abundances and gradients in the Milky Way bulge from FLAMES-GIRAFFE spectra of 650 K giants, A&A, 2011, vol. 530, p. A54
- Gonzalez O. A., Rejkuba M., Zoccali M., Valent E., Minniti D., Tobar R., Reddening and metallicity maps of the Milky Way bulge from VVV and 2MASS. III. The first global photometric metallicity map of the Galactic bulge, A&A, 2013, vol. 552, p. A110
- Gratton R. G., Carretta E., Bragaglia A., Multiple populations in globular clusters. Lessons learned from the Milky Way globular clusters, A&A Rev., 2012, vol. 20, p. 50
- Gunn J. E., Gott III J. R., On the Infall of Matter Into Clusters of Galaxies and Some Effects on Their Evolution, ApJ, 1972, vol. 176, p. 1
- Harris J., Zaritsky D., A Method for Determining the Star Formation History of a Mixed Stellar Population, ApJS, 2001, vol. 136, p. 25

-
- Harris J., Zaritsky D., The Star Formation History of the Small Magellanic Cloud, *AJ*, 2004, vol. 127, p. 1531
- Harris W. E., A Catalog of Parameters for Globular Clusters in the Milky Way, *AJ*, 1996, vol. 112, p. 1487
- Hernandez X., Gilmore G., Valls-Gabaud D., Non-parametric star formation histories for four dwarf spheroidal galaxies of the Local Group, *MNRAS*, 2000, vol. 317, p. 831
- Hernandez X., Valls-Gabaud D., A robust statistical estimation of the basic parameters of single stellar populations - I. Method, *MNRAS*, 2008, vol. 383, p. 1603
- Hernandez X., Valls-Gabaud D., Gilmore G., Deriving star formation histories: inverting Hertzsprung-Russell diagrams through a variational calculus maximum likelihood method, *MNRAS*, 1999, vol. 304, p. 705
- Hesser J. E., Harris G. L. H., Further spectroscopic evidence bearing on the M22-Omega Centauri comparison, *ApJ*, 1979, vol. 234, p. 513
- Hesser J. E., Shawl S. J., Meyer J. E., Systematic reinvestigation of the radial velocities of the galactic globular clusters - Image-tube results, *PASP*, 1986, vol. 98, p. 403
- Hill V., Lecureur A., Gómez A., Zoccali M., Schultheis M., Babusiaux C., Royer F., Barbuy B., Arenou F., Minniti D., Ortolani S., The metallicity distribution of bulge clump giants in Baade's window, *A&A*, 2011, vol. 534, p. A80
- Holtzman J. A., Gallagher III J. S., Cole A. A., Mould J. R., Grillmair C. J., Ballester G. E., Burrows C. J., Clarke J. T., Crisp D., Evans R. W., Griffiths R. E., Hester J. J., Hoessel J. G., Scowen P. A., Stapelfeldt K. R., Trauger J. T., Watson A. M., Observations and Implications of the Star Formation History of the Large Magellanic Cloud, *AJ*, 1999, vol. 118, p. 2262
- Hubble E. P., Extragalactic nebulae., *ApJ*, 1926, vol. 64, p. 321
- Ibata R. A., Gilmore G., Irwin M. J., A dwarf satellite galaxy in Sagittarius, *Nature*, 1994, vol. 370, p. 194
- Javiel S. C., Santiago B. X., Kerber L. O., Constraints on the star formation history of the Large Magellanic Cloud, *A&A*, 2005, vol. 431, p. 73

- Jensen J., Mould J., Reid N., The continuity of cluster formation in the Large Magellanic Cloud, *ApJS*, 1988, vol. 67, p. 77
- Katz D., Automatic determination of stellar atmospheric parameters, *Journal of Astronomical Data*, 2001, vol. 7, p. 8
- Katz D., Soubiran C., Cayrel R., Adda M., Cautain R., On-line determination of stellar atmospheric parameters T_{eff} , $\log g$, $[\text{Fe}/\text{H}]$ from ELODIE echelle spectra. I. The method, *A&A*, 1998, vol. 338, p. 151
- Katz D., Soubiran C., Cayrel R., Barbuy B., Friel E., Bienaymé O., Perrin M.-N., Probing the Galactic thick disc vertical properties and interfaces, *A&A*, 2011, vol. 525, p. A90+
- Kerber L. O., Santiago B. X., Physical parameters of rich LMC clusters from modeling of deep HST colour-magnitude diagrams, *A&A*, 2005, vol. 435, p. 77
- Kerber L. O., Santiago B. X., Castro R., Valls-Gabaud D., Analysis of colour-magnitude diagrams of rich LMC clusters: NGC 1831, *A&A*, 2002, vol. 390, p. 121
- King I., The structure of star clusters. I. an empirical density law, *AJ*, 1962, vol. 67, p. 471
- Kirby E. N., Guhathakurta P., Bolte M., Sneden C., Geha M. C., Multi-element Abundance Measurements from Medium-resolution Spectra. I. The Sculptor Dwarf Spheroidal Galaxy, *ApJ*, 2009, vol. 705, p. 328
- Kirby E. N., Guhathakurta P., Simon J. D., Geha M. C., Rockosi C. M., Sneden C., Cohen J. G., Sohn S. T., Majewski S. R., Siegel M., Multi-element Abundance Measurements from Medium-resolution Spectra. II. Catalog of Stars in Milky Way Dwarf Satellite Galaxies, *ApJS*, 2010, vol. 191, p. 352
- Koleva M., Prugniel P., Bouchard A., Wu Y., ULySS: a full spectrum fitting package, *A&A*, 2009, vol. 501, p. 1269
- Kormendy J., Kennicutt Jr. R. C., Secular Evolution and the Formation of Pseudobulges in Disk Galaxies, *ARA&A*, 2004, vol. 42, p. 603

- Leaman R., VandenBerg D. A., Mendel J. T., The bifurcated age-metallicity relation of Milky Way globular clusters and its implications for the accretion history of the galaxy, *MNRAS*, 2013, vol. 436, p. 122
- Lee Y. S., Beers T. C., Sivarani T., Allende Prieto C., Koesterke L., Wilhelm R., Re Fiorentin P., Bailer-Jones C. A. L., Norris J. E., Rockosi C. M., Yanny B., Newberg H. J., Covey K. R., Zhang H.-T., Luo A.-L., The SEGUE Stellar Parameter Pipeline. I. Description and Comparison of Individual Methods, *AJ*, 2008, vol. 136, p. 2022
- Lee Y.-W., Evidence for an old Galactic bulge from RR Lyrae stars in Baade's window - Implications for the formation of the Galaxy and the age of the universe, *AJ*, 1992, vol. 104, p. 1780
- Lee Y.-W., Demarque P., Zinn R., The horizontal-branch stars in globular clusters. 2: The second parameter phenomenon, *ApJ*, 1994, vol. 423, p. 248
- Lee Y.-W., Han S.-I., Joo S.-J., Jang S., Na C., Okamoto S., Arimoto N., Lim D., Kim H.-S., Yoon S.-J., Two Distinct Red Giant Branch Populations in the Globular Cluster NGC 2419 as Tracers of a Merger Event in the Milky Way*, *ApJ*, 2013, vol. 778, p. L13
- Lee Y.-W., Joo J.-M., Sohn Y.-J., Rey S.-C., Lee H.-C., Walker A. R., Multiple stellar populations in the globular cluster ω Centauri as tracers of a merger event, *Nature*, 1999, vol. 402, p. 55
- Lin D. N. C., Jones B. F., Klemola A. R., The motion of the Magellanic clouds, origin of the Magellanic Stream, and the mass of the Milky Way, *ApJ*, 1995, vol. 439, p. 652
- Lin D. N. C., Lynden-Bell D., Simulation of the Magellanic Stream to estimate the total mass of the Milky Way, *MNRAS*, 1977, vol. 181, p. 59
- Mackey A. D., van den Bergh S., The properties of Galactic globular cluster subsystems, *MNRAS*, 2005, vol. 360, p. 631
- Majewski S. R., Skrutskie M. F., Weinberg M. D., Ostheimer J. C., A Two Micron All Sky Survey View of the Sagittarius Dwarf Galaxy. I. Morphology of the Sagittarius Core and Tidal Arms, *ApJ*, 2003, vol. 599, p. 1082

- Marino A. F., Milone A. P., Piotto G., Villanova S., Bedin L. R., Bellini A., Renzini A., A double stellar generation in the globular cluster NGC 6656 (M 22). Two stellar groups with different iron and s-process element abundances, *A&A*, 2009, vol. 505, p. 1099
- Marino A. F., Milone A. P., Piotto G., Villanova S., Gratton R., D'Antona F., Anderson J., Bedin L. R., Bellini A., Cassisi S., Geisler D., Renzini A., Zoccali M., Sodium-Oxygen Anticorrelation and Neutron-capture Elements in Omega Centauri Stellar Populations, *ApJ*, 2011, vol. 731, p. 64
- Marino A. F., Sneden C., Kraft R. P., Wallerstein G., Norris J. E., da Costa G., Milone A. P., Ivans I. I., Gonzalez G., Fulbright J. P., Hilker M., Piotto G., Zoccali M., Stetson P. B., The two metallicity groups of the globular cluster M 22: a chemical perspective, *A&A*, 2011, vol. 532, p. A8
- Massari D., Ferraro F. R., Mucciarelli A., Origlia L., Dalessandro E., Lanzoni B., Terzan 5: a pristine fragment of the Galactic Bulge?, *ArXiv e-prints*, 2014
- Mathewson D. S., Cleary M. N., Murray J. D., The Magellanic stream, *ApJ*, 1974, vol. 190, p. 291
- Mathewson D. S., Wayte S. R., Ford V. L., Ruan K., The 'high velocity cloud' origin of the Magellanic system, *Proceedings of the Astronomical Society of Australia*, 1987, vol. 7, p. 19
- Matteucci F., Recchi S., On the Typical Timescale for the Chemical Enrichment from Type Ia Supernovae in Galaxies, *ApJ*, 2001, vol. 558, p. 351
- Mauro F., Moni Bidin C., Geisler D., Saviane I., Da Costa G. S., Gormaz-Matamala A. C., Vasquez S., Chené A.-N., Cohen R., Dias B., Deriving metallicities from calcium triplet spectroscopy in combination with near-infrared photometry, *A&A*, 2014, vol. 563, p. A76
- McWilliam A., Rich R. M., The first detailed abundance analysis of Galactic bulge K giants in Baade's window, *ApJS*, 1994, vol. 91, p. 749
- Meléndez J., Cohen J. G., The Rise of the AGB in the Galactic Halo: Mg Isotopic Ratios and High Precision Elemental Abundances in M71 Giants, *ApJ*, 2009, vol. 699, p. 2017

- Mellinger A., A Color All-Sky Panorama Image of the Milky Way, *PASP*, 2009, vol. 121, p. 1180
- Mészáros S., Holtzman J., García Pérez A. E., Allende Prieto C., Schiavon R. P., Basu S. e. a., Calibrations of Atmospheric Parameters Obtained from the First Year of SDSS-III APOGEE Observations, *AJ*, 2013, vol. 146, p. 133
- Mieske S., Hilker M., Infante L., Ultra compact objects in the Fornax cluster of galaxies: Globular clusters or dwarf galaxies?, *A&A*, 2002, vol. 383, p. 823
- Mieske S., Hilker M., Jordán A., Infante L., Kissler-Patig M., Rejkuba M., Richtler T., Côté P., Baumgardt H., West M. J., Ferrarese L., Peng E. W., The nature of UCDs: Internal dynamics from an expanded sample and homogeneous database, *A&A*, 2008, vol. 487, p. 921
- Mighell K. J., Sarajedini A., French R. S., WFPC2 Observations of Star Clusters in the Magellanic Clouds. II. The Oldest Star Clusters in the Small Magellanic Cloud, *AJ*, 1998, vol. 116, p. 2395
- Milone A. D. C., Sansom A. E., Sánchez-Blázquez P., Element abundances in the stars of the MILES spectral library: the Mg/Fe ratio, *MNRAS*, 2011, vol. 414, p. 1227
- Monteiro H., Dias W. S., Caetano T. C., Fitting isochrones to open cluster photometric data. A new global optimization tool, *A&A*, 2010, vol. 516, p. A2
- Moore B., Davis M., The Origin of the Magellanic Stream, *MNRAS*, 1994, vol. 270, p. 209
- Morrison H. L., The local density of halo giants, *AJ*, 1993, vol. 106, p. 578
- Mottini M., Wallerstein G., McWilliam A., ARP 2 and Terzan 8: a Detailed Chemical Analysis, *AJ*, 2008, vol. 136, p. 614
- Murai T., Fujimoto M., The Magellanic Stream and the Galaxy with a Massive Halo, *PASJ*, 1980, vol. 32, p. 581
- Naylor T., Jeffries R. D., A maximum-likelihood method for fitting colour-magnitude diagrams, *MNRAS*, 2006, vol. 373, p. 1251

- Newberg H. J., Willett B. A., Yanny B., Xu Y., The Orbit of the Orphan Stream, *ApJ*, 2010, vol. 711, p. 32
- Newberg H. J., Yanny B., Willett B. A., Discovery of a New, Polar-Orbiting Debris Stream in the Milky Way Stellar Halo, *ApJ*, 2009, vol. 700, p. L61
- Nidever D. L., Majewski S. R., Burton W. B., The Origin of the Magellanic Stream and Its Leading Arm, *ApJ*, 2008, vol. 679, p. 432
- Nidever D. L., Majewski S. R., Butler Burton W., Nigra L., The 200deg Long Magellanic Stream System, *ApJ*, 2010, vol. 723, p. 1618
- Nidever D. L., Monachesi A., Bell E. F., Majewski S. R., Muñoz R. R., Beaton R. L., A Tidally Stripped Stellar Component of the Magellanic Bridge, *ApJ*, 2013, vol. 779, p. 145
- Noël N. E. D., Conn B. C., Carrera R., Read J. I., Rix H.-W., Dolphin A., The MAGellanic Inter-Cloud Project (MAGIC). I. Evidence for Intermediate-age Stellar Populations in between the Magellanic Clouds, *ApJ*, 2013, vol. 768, p. 109
- Noguchi M., Early Evolution of Disk Galaxies: Formation of Bulges in Clumpy Young Galactic Disks, *ApJ*, 1999, vol. 514, p. 77
- Olsen K. A., Nidever D. L., Gruendl R. A., Blum R. D., Walker A. R., Saha A., Olszewski E. W., Munoz R., Kunder A. M., Kaleida C. C., Conn B. e. a., SMASH: The Survey of the MAGellanic Stellar History. In American Astronomical Society Meeting Abstracts , vol. 223 of American Astronomical Society Meeting Abstracts, 2014, p. 254.44
- Olszewski E. W., Saha A., Knezek P., Subramaniam A., de Boer T., Seitzer P., A 500 Parsec Halo Surrounding the Galactic Globular NGC 1851, *AJ*, 2009, vol. 138, p. 1570
- Origlia L., Massari D., Rich R. M., Mucciarelli A., Ferraro F. R., Dalessandro E., Lanzoni B., The Terzan 5 Puzzle: Discovery of a Third, Metal-poor Component, *ApJ*, 2013, vol. 779, p. L5
- Origlia L., Rich R. M., Ferraro F. R., Lanzoni B., Bellazzini M., Dalessandro E., Mucciarelli A., Valenti E., Beccari G., Spectroscopy Unveils the Complex Nature of Terzan 5, *ApJ*, 2011, vol. 726, p. L20

- Ortolani S., Barbuy B., Momany Y., Saviane I., Bica E., Jilkova L., Salerno G. M., Jungwiert B., A Fossil Bulge Globular Cluster Revealed by very Large Telescope Multi-conjugate Adaptive Optics, *ApJ*, 2011, vol. 737, p. 31
- Ortolani S., Bica E., Barbuy B., HP 1: a blue horizontal branch globular cluster in the bulge, *MNRAS*, 1997, vol. 284, p. 692
- Ortolani S., Renzini A., Gilmozzi R., Marconi G., Barbuy B., Bica E., Rich R. M., Near-coeval formation of the Galactic bulge and halo inferred from globular cluster ages, *Nature*, 1995, vol. 377, p. 701
- Pagel B. E. J., Tautvaisiene G., Chemical evolution of the Magellanic Clouds: analytical models, *MNRAS*, 1998, vol. 299, p. 535
- Parisi M. C., Geisler D., Carraro G., Clariá J. J., Costa E., Grocholski A. J., Sarajedini A., Leiton R., Piatti A. E., Age Determination of 15 Old to Intermediate-age Small Magellanic Cloud Star Clusters, *AJ*, 2014, vol. 147, p. 71
- Parisi M. C., Geisler D., Grocholski A. J., Clariá J. J., Sarajedini A., Ca II Triplet Spectroscopy of Small Magellanic Cloud Red Giants. II. Abundances for a Sample of Field Stars, *AJ*, 2010, vol. 139, p. 1168
- Parisi M. C., Grocholski A. J., Geisler D., Sarajedini A., Clariá J. J., Ca II Triplet Spectroscopy of Small Magellanic Cloud Red Giants. I. Abundances and Velocities for a Sample of Clusters, *AJ*, 2009, vol. 138, p. 517
- Perryman M. A. C., de Boer K. S., Gilmore G., Høg E., Lattanzi M. G., Lindegren L., Luri X., Mignard F., Pace O., de Zeeuw P. T., GAIA: Composition, formation and evolution of the Galaxy, *A&A*, 2001, vol. 369, p. 339
- Piatti A. E., New candidate intermediate-age star clusters in the Small Magellanic Cloud, *MNRAS*, 2011a, vol. 416, p. L89
- Piatti A. E., Towards a comprehensive picture of the star cluster age-metallicity relationship in the Small Magellanic Cloud, *MNRAS*, 2011b, vol. 418, p. L69
- Piatti A. E., Identification of a New Relatively Old Star Cluster in the Small Magellanic Cloud, *ApJ*, 2012a, vol. 756, p. L32

- Piatti A. E., The star field age-metallicity relationship of the Small Magellanic Cloud, *MNRAS*, 2012b, vol. 422, p. 1109
- Piatti A. E., Clariá J. J., Bica E., Geisler D., Ahumada A. V., Girardi L., Washington photometry of 14 intermediate-age to old star clusters in the Small Magellanic Cloud, *MNRAS*, 2011, vol. 417, p. 1559
- Piatti A. E., Santos Jr. J. F. C., Clariá J. J., Bica E., Ahumada A. V., Parisi M. C., Integrated spectral analysis of 18 concentrated star clusters in the Small Magellanic Cloud, *A&A*, 2005, vol. 440, p. 111
- Piatti A. E., Sarajedini A., Geisler D., Clark D., Seguel J., Young star clusters immersed in intermediate-age fields in the Small Magellanic Cloud, *MNRAS*, 2007, vol. 377, p. 300
- Piatti A. E., Sarajedini A., Geisler D., Gallart C., Wischnjewsky M., Five young star clusters in the outer region of the Small Magellanic Cloud, *MNRAS*, 2007a, vol. 382, p. 1203
- Piatti A. E., Sarajedini A., Geisler D., Gallart C., Wischnjewsky M., Two newly identified, relatively old star clusters in the Small Magellanic Cloud, *MNRAS*, 2007b, vol. 381, p. L84
- Piatti A. E., Sarajedini A., Geisler D., Seguel J., Clark D., Tracing the formation history of intermediate-age star clusters in the Small Magellanic Cloud, *MNRAS*, 2005, vol. 358, p. 1215
- Pietrinferni A., Cassisi S., Salaris M., Castelli F., A Large Stellar Evolution Database for Population Synthesis Studies. I. Scaled Solar Models and Isochrones, *ApJ*, 2004, vol. 612, p. 168
- Prugniel P., Soubiran C., Koleva M., Le Borgne D., New release of the ELODIE library: Version 3.1, *ArXiv Astrophysics e-prints*, 2007
- Putman M. E., Gibson B. K., Staveley-Smith L., Banks G., Barnes D. G., Bhatal R., Disney M. J. e. a., Tidal disruption of the Magellanic Clouds by the Milky Way, *Nature*, 1998, vol. 394, p. 752

-
- Rafelski M., Zaritsky D., The Star Clusters of the Small Magellanic Cloud: Age Distribution, *AJ*, 2005, vol. 129, p. 2701
- Ramírez S. V., Cohen J. G., Abundances in Stars from the Red Giant Branch Tip to Near the Main-Sequence Turnoff in M71. III. Abundance Ratios, *AJ*, 2002, vol. 123, p. 3277
- Ramírez S. V., Cohen J. G., Buss J., Briley M. M., Abundances in Stars from the Red Giant Branch Tip to Near the Main-Sequence Turnoff in M71. II. Iron Abundance, *AJ*, 2001, vol. 122, p. 1429
- Reddy B. E., Lambert D. L., Allende Prieto C., Elemental abundance survey of the Galactic thick disc, *MNRAS*, 2006, vol. 367, p. 1329
- Rich R. M., Shara M., Fall S. M., Zurek D., Two Groups of Nearly Coeval Star Clusters in the Small Magellanic Cloud, *AJ*, 2000, vol. 119, p. 197
- Rich R. M., Shara M. M., Zurek D., New Photometry for the Intermediate-Age Large Magellanic Cloud Globular Cluster NGC 2121 and the Nature of the LMC Age Gap, *AJ*, 2001, vol. 122, p. 842
- Roederer I. U., Chemical Inhomogeneities in the Milky Way Stellar Halo, *AJ*, 2009, vol. 137, p. 272
- Rubele S., Girardi L., Kozhurina-Platais V., Goudfrooij P., Kerber L., The star formation history of the Large Magellanic Cloud star cluster NGC 1751, *MNRAS*, 2011, vol. 414, p. 2204
- Sadler E. M., Rich R. M., Terndrup D. M., K Giants in Baade's Window. II. The Abundance Distribution, *AJ*, 1996, vol. 112, p. 171
- Saito R. K., Hempel M., Minniti D., Lucas P. W., Rejkuba M., Toledo I., Gonzalez O. A., Alonso-García J. e. a., VVV DR1: The first data release of the Milky Way bulge and southern plane from the near-infrared ESO public survey VISTA variables in the Vía Láctea, *A&A*, 2012, vol. 537, p. A107
- Saito R. K., Minniti D., Dias B., Hempel M., Rejkuba M., Alonso-García J., Barbuy B., Catelan M., Emerson J. P., Gonzalez O. A., Lucas P. W., Zoccali M., Milky Way

- demographics with the VVV survey. I. The 84-million star colour-magnitude diagram of the Galactic bulge, *A&A*, 2012, vol. 544, p. A147
- Saito R. K., Zoccali M., McWilliam A., Minniti D., Gonzalez O. A., Hill V., Mapping the X-shaped Milky Way Bulge, *AJ*, 2011, vol. 142, p. 76
- Salaris M., Chieffi A., Straniero O., The alpha-enhanced isochrones and their impact on the FITS to the Galactic globular cluster system, *ApJ*, 1993, vol. 414, p. 580
- Sánchez Almeida J., Allende Prieto C., Automated Unsupervised Classification of the Sloan Digital Sky Survey Stellar Spectra using k-means Clustering, *ApJ*, 2013, vol. 763, p. 50
- Sánchez-Blázquez P., Peletier R. F., Jiménez-Vicente J., Cardiel N., Cenarro A. J., Falcón-Barroso J., Gorgas J., Selam S., Vazdekis A., Medium-resolution Isaac Newton Telescope library of empirical spectra, *MNRAS*, 2006, vol. 371, p. 703
- Santos Jr. J. F. C., Piatti A. E., Ages and metallicities of star clusters: New calibrations and diagnostic diagrams from visible integrated spectra, *A&A*, 2004, vol. 428, p. 79
- Saviane I., da Costa G. S., Held E. V., Sommariva V., Gullieuszik M., Barbuy B., Ortolani S., Homogeneous metallicities and radial velocities for Galactic globular clusters. First CaT metallicities for twenty clusters, *A&A*, 2012, vol. 540, p. A27
- Saviane I., Held E. V., Da Costa G. S., Sommariva V., Gullieuszik M., Barbuy B., Ortolani S., New Surprises in Old Stellar Clusters, *The Messenger*, 2012, vol. 149, p. 23
- Sharpee B., Stark M., Pritzl B., Smith H., Silbermann N., Wilhelm R., Walker A., BV Photometry of Variable Stars in the Northeast Arm of the Small Magellanic Cloud, *AJ*, 2002, vol. 123, p. 3216
- Sofue Y., Fate of the Magellanic Stream, *PASJ*, 1994, vol. 46, p. 431
- Soldara L., L'effetto dell'arrossamento interstellare sulla misura delle temperature stellari (The effect of the interstellar reddening on the stellar temperature measurements), University of Padova, Department of Astronomy, 2008, Ph.D. Thesis
- Soubiran C., Le Campion J.-F., Cayrel de Strobel G., Caillo A., The PASTEL catalogue of stellar parameters, *A&A*, 2010, vol. 515, p. A111

- Steinmetz M., Zwitter T., Siebert A., Watson F. G., Freeman K. C., Munari U., Campbell R., Williams M. e. a., The Radial Velocity Experiment (RAVE): First Data Release, *AJ*, 2006, vol. 132, p. 1645
- Stetson P. B., DAOPHOT - A computer program for crowded-field stellar photometry, *PASP*, 1987, vol. 99, p. 191
- Subramanian S., Subramaniam A., The Three-dimensional Structure of the Small Magellanic Cloud, *ApJ*, 2012, vol. 744, p. 128
- Tsujimoto T., Bekki K., Chemical Signature of a Major Merger in the Early Formation of Small Magellanic Cloud, *ArXiv e-prints*, 2009
- VandenBerg D. A., Brogaard K., Leaman R., Casagrande L., The Ages of 55 Globular Clusters as Determined Using an Improved $\Delta V^{\text{HB-TO}}$ Method along with Color-Magnitude Diagram Constraints, and Their Implications for Broader Issues, *ApJ*, 2013, vol. 775, p. 134
- Vásquez S., Zoccali M., Hill V., Renzini A., González O. A., Gardner E., Debattista V. P., Robin A. C., Rejkuba M., Baffico M., Monelli M., Motta V., Minniti D., 3D kinematics through the X-shaped Milky Way bulge, *A&A*, 2013, vol. 555, p. A91
- Venn K. A., Irwin M., Shetrone M. D., Tout C. A., Hill V., Tolstoy E., Stellar Chemical Signatures and Hierarchical Galaxy Formation, *AJ*, 2004, vol. 128, p. 1177
- Vergely J.-L., Köppen J., Egret D., Bienaymé O., An inverse method to interpret colour-magnitude diagrams, *A&A*, 2002, vol. 390, p. 917
- Vesperini E., McMillan S. L. W., D'Antona F., D'Ercole A., The Fraction of Globular Cluster Second-generation Stars in the Galactic Halo, *ApJ*, 2010, vol. 718, p. L112
- Villanova S., Geisler D., Piotto G., Detailed Abundances of Red Giants in the Globular Cluster NGC 1851: C+N+O and the Origin of Multiple Populations, *ApJ*, 2010, vol. 722, p. L18
- Worthey G., Faber S. M., Gonzalez J. J., Burstein D., Old stellar populations. 5: Absorption feature indices for the complete LICK/IDS sample of stars, *ApJS*, 1994, vol. 94, p. 687

- Wu Y., Luo A.-L., Li H.-N., Shi J.-R., Prugniel P., Liang Y.-C., Zhao Y.-H., Zhang J.-N., Bai Z.-R., Wei P., Dong W.-X., Zhang H.-T., Chen J.-J., Automatic determination of stellar atmospheric parameters and construction of stellar spectral templates of the Guoshoujing Telescope (LAMOST), *Research in Astronomy and Astrophysics*, 2011, vol. 11, p. 924
- Wylie-de Boer E., Freeman K., HERMES - An instrument of the future. In *IAU Symposium*, vol. 262 of *IAU Symposium*, 2010, p. 448
- Wyse R. F. G., Gilmore G., Franx M., Galactic Bulges, *ARA&A*, 1997, vol. 35, p. 637
- York D. G., Adelman J., Anderson Jr. J. E., Anderson S. F., Annis J., Bahcall N. A., Bakken J. A., SDSS Collaboration The Sloan Digital Sky Survey: Technical Summary, *AJ*, 2000, vol. 120, p. 1579
- Zinn R., West M. J., The globular cluster system of the galaxy. III - Measurements of radial velocity and metallicity for 60 clusters and a compilation of metallicities for 121 clusters, *ApJS*, 1984, vol. 55, p. 45
- Zoccali M., Barbuy B., Hill V., Ortolani S., Renzini A., Bica E., Momany Y., Pasquini L., Minniti D., Rich R. M., The metal content of the bulge globular cluster NGC 6528, *A&A*, 2004, vol. 423, p. 507
- Zoccali M., Gonzalez O. A., Vasquez S., Hill V., Rejkuba M., Valenti E., Renzini A., Rojas-Arriagada A., Martinez-Valpuesta I., Babusiaux C., Brown T., Minniti D., McWilliam A., The GIRAFFE Inner Bulge Survey (GIBS). I. Survey description and a kinematical map of the Milky Way bulge, *A&A*, 2014, vol. 562, p. A66
- Zoccali M., Renzini A., Ortolani S., Bica E., Barbuy B., The Proper Motion of the Globular Cluster NGC 6553 and of Bulge Stars with the Hubble Space Telescope, *AJ*, 2001, vol. 121, p. 2638

Appendix

Appendix A

SMC clusters CMD statistical analysis

A.1 Likelihood results and degeneracy investigation

We analysed the likelihood maps to investigate the degeneracy in age, metallicity, reddening, and distance modulus. The likelihood values were obtained from a comparison of a given synthetic CMD with the observed one (see Section 3 for details).

To establish the standard deviation of the likelihood statistics, we proceeded as follows. A synthetic best model CMD was created with the same numbers of stars as the observed one. We call this a false observation. We then compared this CMD with another one with the same parameters but with many more stars (e.g. 10^5). From this model vs. model comparison, a likelihood value is obtained, as previously done for data vs. model comparisons. Then simulating 300 false observations, and comparing them with the star rich synthetic CMD, we derived the standard deviation σ . This tells how much the likelihood can vary due only to stochastic effects, i.e., due to the statistical fluctuations on the position of stars in a CMD with a number of points compatible with that of the observed CMDs.

The likelihood maps of these comparisons are displayed in Figures A.1 to A.5, in the space of parameters of $[\text{Fe}/\text{H}]$ vs. $\log(\text{age})$, $\log(\text{age})$ vs. $(m-M)$, and $E(B-V)$ vs. $(m-M)$, centred in the parameters of the best-fit. The colour scale is red, green, cyan or yellow, corresponding to a difference from the best fit solution up to 1-sigma, between 1- and 2-sigma, between 2- and 3-sigma, and larger than 3-sigma of the best fit, respectively. The central panel in each figure gives the value obtained for the two fixed parameters, since in each map two parameters are varied and the two others are fixed.

These maps clearly reveal the expected anti-correlation between age and metallicity, distance modulus and age, and between distance modulus and reddening. These degene-

racies are directly reflected in the physical parameter uncertainties since they correspond to the standard deviation in each parameter for the set of models with likelihood values up to 1-sigma (red points) from the best-fit solution.

Figure A.1: Likelihood from the fits for different combinations of age, metallicity, reddening, and distance modulus for AM 3. Upper panel: metallicity vs. age; middle panel: age vs. distance modulus; bottom panel: reddening vs. distance modulus. The colour scale is red, green, cyan or yellow, corresponding to a difference from the best-fit solution up to 1-sigma, between 1- and 2-sigma, between 2- and 3-sigma, and larger than 3-sigma of the best fit, respectively.

Figure A.2: Same as Fig. A.1 for HW 1.

Figure A.3: Same as Fig. A.1 for HW 40.

Figure A.4: Same as Fig. A.1 for Lindsay 2.

Figure A.5: Same as Fig. A.1 for Lindsay 3.

Appendix B

Tables with results for FORS2 spectra

Table B.1 - Star by star coordinates, magnitude, colour, heliocentric radial velocity. Velocities from CaT were taken from from Saviane et al. (2012) for NGC 6528, NGC 6553, M 71 and NGC 6558, and from Vasquez et al. (2014, in prep.) for NGC 6426 and Terzan 8.

Star ID	RA (J2000) (deg)	DEC (J2000) (deg)	V (mag)	V-I (mag)	B-V (mag)	v_{helio} (km/s)	$v_{\text{helio}} - \text{CaT}$ (km/s)	members
NGC6528_2	271.1807715771910	-30.0070234836130	15.987	1.982	—	-51.55	—	
NGC6528_3	271.1933667526377	-30.0114575457940	15.647	2.836	—	61.28	—	
NGC6528_4	271.1789935185810	-30.0172537974630	15.596	2.257	—	-314.24	—	
NGC6528_5	271.1883528713980	-30.0237445060760	15.939	1.937	—	-226.44	—	
NGC6528_6	271.1974416677920	-30.0295599638440	17.060	1.741	—	-4.93	—	
NGC6528_7	271.2102328036860	-30.0372195159180	15.887	2.140	—	-79.24	—	
NGC6528_8	271.2027175539120	-30.0434602130820	16.428	1.831	—	179.09	200	M
NGC6528_9	271.1969387171677	-30.0500739849720	16.501	1.774	—	199.18	208	M
NGC6528_10	271.2070341687930	-30.0537176170110	16.145	2.007	—	177.87	209	M
NGC6528_11	271.2013524700317	-30.0600904185180	15.511	2.255	—	182.67	202	M
NGC6528_13	271.1879324598030	-30.0747465095010	16.429	1.731	—	-29.84	—	
NGC6528_14	271.2012738946260	-30.0802412070160	16.032	2.056	—	-40.56	—	
NGC6528_15	271.1774205888063	-30.0879237445260	15.892	2.062	—	-288.86	—	
NGC6528_16	271.1843122189650	-30.0926974725130	15.954	2.150	—	-104.91	—	
NGC6528_17	271.1904270294383	-30.1021087576640	16.537	1.844	—	-62.96	—	
NGC6528_18	271.1726747862950	-30.1092237523470	16.666	1.767	—	-212.87	—	
NGC6528_19	271.1832358189100	-30.1108940656500	16.468	1.778	—	-54.47	—	
NGC6553_1	272.3536106669400	-25.8497112721910	15.816	2.109	—	3.90	-8	M
NGC6553_3	272.3507085422689	-25.8636412811220	15.832	2.010	—	16.71	19	M
NGC6553_4	272.2982167426244	-25.8658039783470	15.310	2.618	—	-71.71	-39	M
NGC6553_5	272.3267021788947	-25.8733214097360	15.775	2.522	—	12.23	-12	M
NGC6553_6	272.3161974470756	-25.8795540059450	16.237	2.177	—	0.81	-6	M
NGC6553_7	272.3258216397713	-25.8883122417460	15.338	2.881	—	6.76	3	M
NGC6553_8	272.3501542761960	-25.8928074210270	15.370	3.088	—	18.98	—	
NGC6553_9	272.3409674217040	-25.9007568498840	16.253	2.097	—	-27.72	-32	M
NGC6553_10	272.3439213760840	-25.9071903257530	15.985	1.998	—	2.34	2	M
NGC6553_11	272.3287891424960	-25.9110912319810	15.441	2.443	—	10.12	-3	M

Table B.1 - continued.

Star ID	RA (J2000) (deg)	DEC (J2000) (deg)	V (mag)	V-I (mag)	B-V (mag)	v_{helio} (km/s)	$v_{\text{helio}} - \text{CaT}$ (km/s)	members
NGC6553_13	272.3235733284760	-25.9249134402580	15.906	2.075	—	-4.38	-12	M
NGC6553_14	272.3157993633750	-25.9300477560180	15.187	2.382	—	5.40	-5	M
NGC6553_15	272.3047106358403	-25.9370597367310	14.980	2.996	—	-21.36	—	
NGC6553_16	272.3280756666470	-25.9436286781210	15.708	2.332	—	17.42	8	M
NGC6553_17	272.2895515368990	-25.9484055025060	15.065	2.849	—	-110.00	—	
NGC6553_18	272.3144982937293	-25.9566926017970	15.407	3.681	—	-10.44	—	
NGC6553_19	272.3440933085433	-25.9630097530170	15.843	2.209	—	-7.38	-4	M
M71_2	298.4578984488130	18.8301452486084	14.606	1.269	—	-150.46	-34	M
M71_4	298.4609554670447	18.8192204176758	13.030	1.503	—	-53.75	-41	M
M71_5	298.4632902665389	18.8090499016740	14.391	1.282	—	-52.74	-26	M
M71_6	298.4900822754500	18.7995866372160	13.534	1.367	—	-44.07	-37	M
M71_7	298.4510869475747	18.8011235588184	12.376	1.719	—	-69.84	-34	M
M71_8	298.4854008510480	18.7869940682184	14.421	1.280	—	-51.13	—	
M71_9	298.4422411559920	18.7910962573810	13.146	1.530	—	-35.78	-24	M
M71_10	298.4476878071520	18.7813713381542	12.140	2.042	—	-129.33	-26	M
M71_13	298.4496765965810	18.7641970806387	14.281	1.303	—	-23.72	-5	M
M71_14	298.4902210311380	18.7496150618418	14.582	1.210	—	-13.11	-17	M
M71_15	298.4521026881480	18.7474809643955	14.580	1.209	—	-41.62	-8	M
M71_16	298.4732927595957	18.7366583486676	14.727	1.231	—	-23.67	—	
NGC6558_3	272.6225545017980	-31.7335169754310	16.862	1.434	—	-6.32	—	
NGC6558_4	272.5685475821600	-31.7363376028500	16.541	1.441	—	-38.29	—	
NGC6558_5	272.6214120279947	-31.7468917846890	16.349	1.360	—	-23.46	—	
NGC6558_6	272.5796695743913	-31.7469946267620	15.982	1.524	—	-210.85	-196	M
NGC6558_7	272.5899712149563	-31.7570504471050	15.803	1.393	—	-186.59	-187	M
NGC6558_8	272.5739816190383	-31.7605125893880	13.651	2.044	—	-307.26	-210	M
NGC6558_9	272.5637020636180	-31.7666306345590	16.026	1.499	—	-221.66	-204	M
NGC6558_10	272.5771880085779	-31.7733227718450	16.710	1.580	—	-21.13	—	
NGC6558_11	272.5757815235553	-31.7788392433250	16.753	1.329	—	-220.56	-195	M
NGC6558_12	272.5805489015007	-31.7879129654280	16.521	1.452	—	-126.79	—	
NGC6558_13	272.5717825299750	-31.7916739864870	15.626	1.477	—	-127.74	—	
NGC6558_14	272.5809044425367	-31.8010529051890	16.740	1.188	—	-118.59	—	
NGC6558_15	272.5652668675210	-31.8066271885810	16.480	1.293	—	105.58	—	
NGC6558_16	272.5564861827613	-31.8124812867920	16.366	1.450	—	-316.32	—	
NGC6558_17	272.5801632324050	-31.8180552937480	16.964	1.348	—	-49.93	—	
NGC6558_18	272.6131914789087	-31.8262981571080	16.911	1.288	—	11.27	—	
NGC6558_19	272.5939188278727	-31.8321512784620	17.013	1.299	—	-5.38	—	
NGC6426_1	266.2553357204670	3.2257821720617	18.018	1.359	—	-80.97	-49.927	
NGC6426_2	266.2088188480099	3.2192772570129	17.597	1.394	—	-2.85	11.325	
NGC6426_3	266.2200960712349	3.2161573577027	16.468	1.586	—	-15.18	12.662	
NGC6426_4	266.2059346355020	3.2095717676480	16.072	1.644	—	-236.29	-230.019	M
NGC6426_7	266.2472044411759	3.1904905620518	17.666	1.459	—	-259.55	-225.413	M
NGC6426_9	266.2324246578110	3.1746245634344	17.170	1.515	—	-244.78	-222.660	M
NGC6426_10	266.2152297889280	3.1685396261324	15.544	1.749	—	-231.87	-221.297	M

Table B.1 - continued.

Star ID	RA (J2000) (deg)	DEC (J2000) (deg)	V (mag)	V-I (mag)	B-V (mag)	v_{helio} (km/s)	$v_{\text{helio}} - \text{CaT}$ (km/s)	members
NGC6426_11	266.2225277929239	3.1649406775723	17.892	1.354	—	-4.92	12.270	
NGC6426_13	266.2203523684530	3.1516328071727	16.745	1.482	—	-236.15	-225.730	M
NGC6426_18	266.2405437043161	3.1175053813749	16.970	1.434	—	43.08	72.570	
Terzan8_1	295.4542164758450	-33.9415478435970	16.706	1.232	—	134.45	138.8238	M
Terzan8_4	295.4999075670249	-33.9726804796670	15.268	1.436	—	120.63	137.4998	M
Terzan8_5	295.4297741922069	-33.9618073863440	17.002	1.135	—	134.61	152.6001	M
Terzan8_6	295.4324851623650	-33.9679803480730	17.517	1.076	—	119.81	138.3819	M
Terzan8_8	295.4284871648290	-33.9821965410250	17.089	1.108	—	173.81	150.6620	M
Terzan8_9	295.4571652028810	-33.9956864225960	15.447	1.352	—	155.67	139.3673	M
Terzan8_10	295.4738836006140	-34.0028288269560	17.381	1.076	—	-63.09	-25.6758	
Terzan8_11	295.4369091804230	-34.0003943267700	15.682	1.277	—	142.74	150.6086	M
Terzan8_13	295.4233824419659	-34.0145360437400	17.364	1.101	—	149.42	144.2031	M
Terzan8_14	295.4815071780459	-34.0298070702590	15.530	1.388	—	116.26	131.9558	M
Terzan8_15	295.4375324314209	-34.0304794278020	16.433	1.219	—	105.77	121.5507	M
Terzan8_16	295.4285209120099	-34.0321898355090	16.778	1.122	—	128.12	140.9906	M
Terzan8_18	295.4529047094940	-34.0531331777060	16.143	1.273	—	134.81	131.1500	M
HP1_8	262.779492341903	-29.98247389275	16.11	2.853	—	51.39	42	M
HP1_10	262.7747217681897	-29.98806695748	17.089	2.611	—	57.23	33	M
HP1_11	262.774686991011	-29.976234361979	17.023	2.582	—	51.05	49	M
HP1_12	262.7843264886397	-29.986110981196	17.436	2.461	—	60.54	46	M
HP1_13	262.77288295732	-29.97799391478	16.314	2.729	—	50.66	—	M
HP1_14	262.775818435468	-29.973607362265	17.981	2.346	—	61.72	—	
HP1_15	262.7658913221563	-29.970602559014	18.155	2.408	—	43.66	—	
HP1_19	262.7710596683369	-29.961417147298	18.707	2.373	—	23.73	—	
HP1_20	262.773834846011	-29.959064708418	18.298	2.48	—	-79.38	—	
HP1_21	262.764621912009	-29.95728051885	18.698	2.436	—	62.94	—	
HP1_22	262.767341454344	-29.990314369868	17.48	2.575	—	46.12	—	M
HP1_23	262.7795572868710	-29.993509304553	16.625	2.757	—	55.28	41	M
HP1_24	262.7845610691	-29.992512837865	17.688	2.467	—	56.36	—	M
HP1_31	262.7669832075777	-30.015181033202	18.99	2.419	—	-127.76	—	
HP1_32	262.7748099931330	-30.010978851507	16.993	2.679	—	-33.81	—	
HP1_33	262.757315085811	-30.009544004121	18.927	2.372	—	-34.45	55	
HP1_34	262.781520651355	-30.013632577769	18.249	2.5	—	93.67	—	
HP1_35	262.783949679813	-30.017159403528	18.318	2.495	—	71.42	—	
HP1_36	262.763821357723	-30.019233977044	17.942	2.4	—	-86.25	23	
HP1_37	262.7988439553969	-30.026125010639	18.517	2.493	—	15.55	—	
HP1_38	262.804671214119	-30.030509267249	18.292	2.477	—	-208.68	—	
HP1_40	262.7792611041633	-30.035669352705	18.775	2.470	—	-109.28	—	
HP1_41	262.755386489532	-30.039545559857	18.547	2.301	—	-85.63	—	
HP1_42	262.7626463939883	-30.054216574809	18.140	2.330	—	-59.15	—	
HP1_43	262.7845333499400	-30.055806701214	17.987	2.423	—	67.24	—	
HP1_44	262.7813860411687	-30.06779264653	18.21	2.453	—	-5.16	—	
HP1_45	262.77203229412	-30.069404519862	17.79	2.448	—	60.36	—	

Table B.1 - continued.

Star ID	RA (J2000) (deg)	DEC (J2000) (deg)	V (mag)	V-I (mag)	B-V (mag)	v_{helio} (km/s)	$v_{\text{helio}} - \text{CaT}$ (km/s)	members
Djorg2_1	270.417078411407	-27.770336770561	16.105	2.446	—	-31.19	-45.945	
Djorg2_3	270.484188183519	-27.780979087074	17.191	2.166	—	56.59	28.664	
Djorg2_4	270.438318422190	-27.788128353318	16.569	2.217	—	-197.17	-152.971	
Djorg2_5	270.447758172525	-27.794258449012	16.688	2.221	—	108.52	117.805	
Djorg2_6	270.428848388335	-27.802121606205	17.102	2.235	—	-27.09	16.752	
Djorg2_7	270.463779615994	-27.806532752427	16.879	2.186	—	-111.15	-105.065	M
Djorg2_8	270.473027706742	-27.811879575782	17.271	2.226	—	-53.58	-33.666	
Djorg2_9	270.449730151511	-27.818858148134	16.404	2.363	—	-170.86	-171.734	M
Djorg2_10	270.452491503247	-27.825727319070	15.707	2.336	—	-146.38	-152.576	M
Djorg2_11	270.452744340045	-27.829940713150	16.082	2.273	—	-127.83	-114.725	
Djorg2_13	270.486191567274	-27.844114862668	17.103	2.191	—	12.25	25.8682	
Djorg2_14	270.460387721211	-27.849287982906	17.275	2.190	—	-0.23	18.0543	
Djorg2_15	270.444232735907	-27.856516596341	16.995	2.294	—	-170.02	-155.4192	M
Djorg2_16	270.483310147638	-27.862658567924	16.964	2.302	—	74.48	91.3211	
Djorg2_17	270.451485414403	-27.867894607897	17.254	2.268	—	10.35	26.5939	
M22a_1	279.1666250722699	-23.8456280526570	14.067	1.387	—	9.93	—	
M22a_2	279.1203368876777	-23.8502790889930	12.051	1.607	—	-280.25	-175	
M22a_3	279.1337615252070	-23.8586133574210	12.390	1.543	—	-150.98	-157	M
M22a_4	279.1776372616150	-23.8646361547940	12.612	1.489	—	-144.13	-159	M
M22a_5	279.1454915186590	-23.8688173810320	13.807	1.356	—	-200.07	-152	M
M22a_6	279.1649925293353	-23.8784140984930	14.101	1.325	—	-154.39	-163	M
M22a_7	279.1190633242593	-23.8855144892020	12.446	1.462	—	-274.77	-173	
M22a_8	279.1497676262920	-23.8888179678540	12.902	1.433	—	-165.38	-170	M
M22a_9	279.1296876926750	-23.8962744818240	14.602	1.287	—	-162.76	-162	M
M22a_10	279.1511268265590	-23.9004626206730	12.872	1.481	—	-151.26	-148	M
M22a_11	279.1761564462446	-23.9078387986190	14.959	1.245	—	-107.27	-144	M
M22a_13	279.1329002956070	-23.9203534699920	15.331	1.318	—	-184.37	-144	M
M22a_14	279.1381784590147	-23.9287792077520	13.860	1.378	—	-98.39	-159	M
M22a_15	279.1213787575467	-23.9340248401570	13.966	1.347	—	-260.77	-151	
M22a_16	279.1297951311230	-23.9424120237370	14.094	1.364	—	-182.46	-162	M
M22a_17	279.1177440796620	-23.9492310681210	14.242	1.301	—	-266.90	-136	
M22a_18	279.1344761999463	-23.9552663000650	14.348	1.293	—	-154.45	-165	M
M22a_19	279.1137715179620	-23.9578959925670	14.507	1.258	—	-169.73	-135	M
M22b_2	279.1487212876030	-23.8496758353150	13.879	1.320	—	-150.36	-160	M
M22b_3	279.1111623406380	-23.8560708266330	12.856	1.446	—	-159.81	-148	M
M22b_4	279.1557209203100	-23.8655332874450	13.790	1.341	—	-136.49	-140	M
M22b_5	279.1332777302069	-23.8693388862050	13.643	1.321	—	-218.64	-152	M
M22b_6	279.1680264045470	-23.8762713491130	14.540	1.301	—	-119.28	-148	M
M22b_7	279.1272594035789	-23.8800383502430	12.543	1.424	—	-189.70	-143	M
M22b_8	279.1330535403420	-23.8879255853180	12.573	1.491	—	-175.18	-162	M
M22b_9	279.1625562094060	-23.8961356996120	15.186	1.199	—	3.52	—	
M22b_10	279.1596042399243	-23.9003368086050	12.791	1.525	—	-149.92	-147	M
M22b_11	279.1728016834153	-23.9108669779370	14.337	1.296	—	-142.83	-148	M

Table B.1 - continued.

Star ID	RA (J2000) (deg)	DEC (J2000) (deg)	V (mag)	V-I (mag)	B-V (mag)	v_{helio} (km/s)	$v_{\text{helio}} - \text{CaT}$ (km/s)	members
M22b_12	279.1387917919590	-23.9144019494200	14.015	1.345	—	-138.58	-139	M
M22b_13	279.1446715959634	-23.9226894359160	15.595	1.269	—	-144.68	-138	M
M22b_15	279.1647014983650	-23.9327436018760	15.201	1.246	—	-147.84	-156	M
M22b_16	279.1379540852260	-23.9404817327490	12.710	1.577	—	-132.18	-146	M
M22b_17	279.1436723242483	-23.9444067248200	14.382	1.348	—	-117.82	—	M
M22b_18	279.1433891813503	-23.9509077647760	14.440	1.296	—	-148.89	-166	M
M22b_19	279.1132162240080	-23.9568469309490	13.522	1.330	—	-136.20	-136	M
M22c_1	279.1200663503463	-23.8480911555530	15.564	1.226	—	-181.06	-166	M
M22c_2	279.1227957715860	-23.8524548018580	14.138	1.283	—	-272.56	-158	
M22c_3	279.1289637909800	-23.8571515884500	14.969	1.257	—	-158.88	-126	M
M22c_4	279.1359228263940	-23.8628693264100	14.595	1.295	—	-173.62	-149	M
M22c_5	279.1241844719839	-23.8707070342580	14.011	1.293	—	-261.18	-148	
M22c_6	279.1430691671390	-23.8787067306810	14.675	1.284	—	-163.65	-148	M
M22c_7	279.1499272095340	-23.8853796867550	13.924	1.328	—	-163.13	-140	M
M22c_8	279.1537927426139	-23.8888545995990	13.914	1.315	—	-127.20	-124	M
M22c_9	279.1702654177500	-23.8952114942570	14.116	1.347	—	-142.68	-149	M
M22c_10	279.1558910801040	-23.9046055967410	14.017	1.348	—	-151.16	-146	M
M22c_11	279.1711564217420	-23.9091704786310	15.417	1.246	—	-118.45	-136	M
M22c_12	279.1776201241823	-23.9164834140020	14.867	1.262	—	-45.50	—	
M22c_13	279.1489093190013	-23.9191389245750	14.183	1.329	—	-144.23	-133	M
M22c_14	279.1642129850940	-23.9288310549590	14.222	1.343	—	-105.87	-158	M
M22c_15	279.1386592958460	-23.9337122259620	14.161	1.388	—	-144.71	-138	M
M22c_16	279.1326771879260	-23.9399533005910	14.973	1.316	—	-162.20	—	M
M22c_17	279.1306698772860	-23.9460284799480	14.862	1.300	—	-157.68	-132	M
M22c_18	279.1291621975290	-23.9536741591000	13.465	1.352	—	-177.36	-150	M
NGC5824_1	225.9734215705400	-33.0116093771520	17.225	1.056	—	-150.40	—	
NGC5824_2	226.0165356446810	-33.0184212781050	16.237	1.262	—	-26.26	-3	M
NGC5824_3	226.0287325923800	-33.0252307354730	16.592	1.239	—	-25.75	-14	M
NGC5824_4	226.0283231480360	-33.0308974205760	15.471	1.466	—	-26.86	-12	M
NGC5824_5	225.9784884294890	-33.0370597656750	18.222	1.029	—	-46.76	-11	M
NGC5824_6	225.9774276480950	-33.0442239249250	17.164	1.150	—	-52.68	-15	M
NGC5824_7	225.9982707978390	-33.0508289721770	16.058	1.259	—	-27.67	-11	M
NGC5824_8	225.9790189358050	-33.0546714164150	16.733	1.183	—	-140.08	-29	
NGC5824_9	225.9833503351730	-33.0600078905660	16.072	1.293	—	-39.41	-8	M
NGC5824_10	226.0209718810750	-33.0671784622460	15.850	1.274	—	-44.20	-27	M
NGC5824_11	225.9797147609669	-33.0722923238210	17.244	1.134	—	-119.30	-2	
NGC5824_13	225.9791626677340	-33.0866087907630	16.657	1.187	—	-23.92	12	M
NGC5824_14	226.0206452881650	-33.0920727071730	17.300	1.090	—	-24.94	-4	M
NGC5824_15	225.9838476458860	-33.0998016563950	16.947	1.179	—	-53.36	-21	M
NGC5824_16	225.9728026992450	-33.1049387810300	17.142	1.164	—	-48.56	—	M
NGC5824_17	225.9790519921090	-33.1111104009490	16.454	1.259	—	-42.29	-8	M
NGC5824_18	226.0258405833140	-33.1173468021370	16.604	1.230	—	-25.02	-13	M
NGC5824_19	226.0214064677600	-33.1247697457440	16.376	1.250	—	-22.89	-10	M

Table B.1 - continued.

Star ID	RA (J2000) (deg)	DEC (J2000) (deg)	V (mag)	V-I (mag)	B-V (mag)	v_{helio} (km/s)	$v_{\text{helio}} - \text{CaT}$ (km/s)	members
47Tuc_502	6.39660800	-72.03976000	11.40	—	2.66	-72.61	—	
47Tuc_509	6.32615400	-72.08429000	12.09	—	1.47	-35.21	—	M
47Tuc_514	6.41895400	-72.09905000	13.74	—	1.04	-46.80	—	M
47Tuc_517	6.42278700	-72.10688000	13.44	—	1.16	-54.61	—	M
47Tuc_519	6.41735800	-72.10936000	12.69	—	1.28	-56.79	—	M
47Tuc_525	6.36514200	-72.02726000	12.84	—	1.27	-44.54	—	M
47Tuc_533	6.33341200	-72.08765000	12.09	—	1.40	-34.56	—	M
47Tuc_534	6.31307100	-72.03169000	14.05	—	0.83	-50.90	—	M
47Tuc_535	6.34090400	-72.09150000	13.79	—	1.06	-49.11	—	M
47Tuc_539	6.36690000	-72.05583000	14.68	—	0.90	-32.26	—	M
47Tuc_551	6.31267500	-72.07024000	12.34	—	1.40	-40.29	—	M
47Tuc_553	6.39460000	-72.07391000	14.15	—	0.84	-46.89	—	M
47Tuc_554	6.36475000	-72.01599000	12.06	—	1.57	-44.67	—	M
47Tuc_559	6.40361700	-72.01102000	13.98	—	1.05	-68.65	—	M
47Tuc_571	6.37052500	-72.04336000	12.59	—	1.26	-58.65	—	M
47Tuc_581	6.40670800	-72.02340000	11.82	—	1.66	-56.46	—	M
NGC2808_1	138.0417278196900	-64.8092533942590	15.171	1.278	—	102.73	87	M
NGC2808_2	137.9510644096820	-64.8162521848410	15.678	1.227	—	-38.59	—	
NGC2808_3	137.9891312000780	-64.8207082107530	15.031	1.351	—	40.09	78	M
NGC2808_4	138.0316478749950	-64.8267569570500	14.994	1.319	—	93.52	84	M
NGC2808_5	138.0084116317160	-64.8341757324620	15.523	1.260	—	141.37	109	M
NGC2808_6	138.0903663713400	-64.8427676002660	15.501	1.210	—	124.16	123	M
NGC2808_7	138.0705536048417	-64.8471283547480	15.139	1.181	—	103.97	97	M
NGC2808_8	138.0760951541040	-64.8526824542090	14.492	1.345	—	109.32	109	M
NGC2808_9	137.9401382663190	-64.8584820105990	15.217	1.246	—	-26.92	72	
NGC2808_10	138.0699371148180	-64.8674965399320	13.513	1.800	—	94.61	69	M
NGC2808_11	137.9445503739920	-64.8706333755900	14.246	1.370	—	-26.70	88	
NGC2808_13	138.0748040594420	-64.8849411022800	15.047	1.318	—	92.14	82	M
NGC2808_14	138.0088058861450	-64.8908680622540	14.969	1.292	—	118.06	118	M
NGC2808_15	138.0277500441930	-64.8968860105230	14.848	1.242	—	100.49	99	M
NGC2808_16	137.9794418613510	-64.9039743986850	14.122	1.520	—	81.67	93	M
NGC2808_17	138.0253961697260	-64.9118795070550	14.728	1.381	—	70.94	56	M
NGC2808_18	138.0462439846450	-64.9176314464940	14.622	1.381	—	253.18	—	
NGC2808_19	137.9999025756700	-64.9209206227550	15.689	1.219	—	68.92	99	M
NGC3201_1	154.4022188951080	-46.3576365537710	14.905	1.172	—	391.27	522	
NGC3201_2	154.4238541394780	-46.3651754747810	14.220	1.226	—	493.33	504	M
NGC3201_5	154.4123378691450	-46.3821823854600	14.707	1.227	—	454.30	500	M
NGC3201_6	154.4113544203970	-46.3901722205060	13.324	1.391	—	398.72	501	
NGC3201_7	154.4319366864960	-46.3948909223600	14.684	1.221	—	470.72	487	M
NGC3201_8	154.4374161137010	-46.4015263953670	12.803	1.465	—	466.05	487	M
NGC3201_9	154.4390300101110	-46.4073415606250	13.969	1.286	—	461.17	484	M
NGC3201_10	154.4547170051770	-46.4160504916910	13.901	1.313	—	504.96	486	M
NGC3201_11	154.4542427127600	-46.4209906922740	14.449	1.294	—	510.49	475	M

Table B.1 - continued.

Star ID	RA (J2000) (deg)	DEC (J2000) (deg)	V (mag)	V-I (mag)	B-V (mag)	v_{helio} (km/s)	$v_{\text{helio}} - \text{CaT}$ (km/s)	members
NGC3201_13	154.4761499745440	-46.4350765100180	14.982	1.199	—	458.56	473	M
NGC3201_14	154.4567663989070	-46.4403081429750	14.296	1.274	—	457.51	471	M
NGC3201_15	154.4221121767170	-46.4465748973930	15.009	1.219	—	456.96	470	M
NGC3201_16	154.4274413445690	-46.4540124607780	12.575	1.533	—	460.37	468	M
NGC3201_17	154.4703687642530	-46.4583470341790	14.681	1.254	—	472.20	472	M
NGC3201_18	154.4485808118120	-46.4630495798020	14.893	1.221	—	473.53	472	M
M68_49	189.81640000	-26.76985000	14.62	—	0.62	-117.70	—	M
M68_30	189.82090000	-26.76078000	14.15	—	0.87	-98.51	—	M
M68_74	189.82600000	-26.77787000	14.59	—	0.84	-131.91	—	M
M68_II47	189.83980000	-26.80485000	15.03	—	0.85	-82.49	—	M
M68_119	189.86280000	-26.79759000	13.62	—	0.95	-76.73	—	M
M68_144	189.87850000	-26.79799000	12.80	—	1.29	-425.94	—	
M68_185	189.89810000	-26.77817000	17.82	—	0.75	-18.11	—	
M68_239	189.90680000	-26.75432000	14.19	—	0.87	-76.57	—	M
M68_II72	189.91760000	-26.79374000	15.03	—	0.85	-63.83	—	M
M68_II79	189.94230000	-26.76432000	14.39	—	0.84	38.06	—	
M5_II45	229.69710000	2.12638300	14.75	—	0.82	54.52	—	M
M5_II50	229.68660000	2.12951400	13.92	—	0.96	46.08	—	M
M5_II51	229.67630000	2.13015300	14.05	—	0.96	40.96	—	M
M5_II80	229.66510000	2.12146900	14.31	—	0.91	35.73	—	M
M5_I2	229.62980000	2.13470800	13.87	—	1.02	53.85	—	M
M5_I50	229.62030000	2.12908100	13.91	—	0.97	38.16	—	M
M5_I61	229.60180000	2.11524400	13.37	—	1.17	52.20	—	M
M5_I68	229.59390000	2.11570600	12.37	—	1.52	44.10	—	M
M5_I71	229.58890000	2.12383300	13.01	—	1.29	52.36	—	M
M4_1	245.9297297869130	-26.4687300856480	13.567	1.351	—	-23.23	55	M
M4_3	245.9083172618420	-26.4781130777430	13.532	1.355	—	-20.06	29	M
M4_4	245.9595428044260	-26.4906387516420	13.432	1.304	—	73.45	64	M
M4_5	245.9285006629270	-26.4910400283700	12.684	1.444	—	35.11	72	M
M4_6	245.9337478207980	-26.5017457785090	11.809	1.540	—	59.27	96	
M4_8	245.9108397606630	-26.5105926710210	12.961	1.402	—	-59.65	58	
M4_9	245.9784456545850	-26.5223373845770	13.145	1.338	—	136.54	53	
M4_10	245.9208888506110	-26.5250639791230	12.787	1.448	—	8.85	69	M
M4_11	245.9301389427700	-26.5316608572910	12.843	1.450	—	46.20	87	M
M4_13	245.9023430347580	-26.5415733433870	13.566	1.371	—	54.24	94	M
M4_16	245.9743179217110	-26.5645714434050	12.927	1.416	—	138.22	47	
M4_17	245.9732209310530	-26.5708402941870	12.565	1.477	—	9.44	83	M
M4_18	245.9739292754490	-26.5785037699360	11.802	1.470	—	112.24	127	
M4_19	245.9142992077440	-26.5813663445440	13.593	1.291	—	62.37	—	
M107_F	248.08860000	-13.04748000	13.39	—	1.70	-28.32	—	
M107_62	248.13850000	-13.07172000	13.97	—	1.62	-8.89	—	
M107_100	248.16340000	-13.07844000	14.21	—	1.40	-24.12	—	

Table B.1 - continued.

Star ID	RA (J2000) (deg)	DEC (J2000) (deg)	V (mag)	V-I (mag)	B-V (mag)	v_{helio} (km/s)	$v_{\text{helio}} - \text{CaT}$ (km/s)	members
M107_I	248.17360000	-13.07616000	13.89	—	1.46	-40.50	—	
M107_C	248.18470000	-13.06669000	11.79	—	1.17	-121.78	—	M ^(a)
M10.1	254.3171837281469	-4.0411518459603	15.107	1.191	—	15.63	84	M
M10.2	254.3228668862143	-4.0453713154607	13.949	1.262	—	11.74	71	M
M10.3	254.3398910591860	-4.0509869791799	15.993	1.144	—	76.32	99	M
M10.4	254.2952316719980	-4.0588352280029	14.742	1.153	—	66.80	86	M
M10.5	254.3056444118800	-4.0644976762931	14.382	1.273	—	30.46	75	M
M10.6	254.3248227850040	-4.0726104654236	13.725	1.297	—	12.85	87	M
M10.7	254.3138792124880	-4.0797717909184	13.642	1.262	—	58.66	96	M
M10.8	254.3036166247190	-4.0847652166142	15.089	1.143	—	-37.99	86	M
M10.9	254.3092280291750	-4.0925118847391	12.966	1.375	—	49.06	78	M
M10.10	254.3115913558880	-4.0981498933977	13.542	1.358	—	67.55	85	M
M10.11	254.3160940089350	-4.1026238369323	15.508	1.180	—	64.64	86	M
M10.13	254.3216643958510	-4.1168053117431	15.606	1.191	—	80.99	94	M
M10.15	254.2960863303000	-4.1269278482101	15.744	1.170	—	62.41	85	M
M10.16	254.3131486307350	-4.1351867073021	14.650	1.233	—	45.67	—	
M10.17	254.3023740571860	-4.1416695758784	15.863	1.163	—	-39.54	—	
NGC6397_1	265.2212301649530	-53.6299970684040	13.564	1.038	—	-73.51	3	M
NGC6397_2	265.2574707418480	-53.6511313057610	12.450	1.029	—	-1.25	—	
NGC6397_4	265.1746135514710	-53.6385099535620	12.826	1.098	—	-90.63	12	M
NGC6397_5	265.1806579482867	-53.6473420123450	13.687	1.049	—	9.55	37	M
NGC6397_6	265.1701016417030	-53.6471000179280	13.935	1.022	—	-95.51	-7	M
NGC6397_7	265.1621423202240	-53.6566342448740	13.106	1.080	—	-79.15	25	M
NGC6397_8	265.1469821778820	-53.6579510091340	13.243	1.073	—	0.10	16	M
NGC6397_9	265.2003518222717	-53.6803241141170	12.609	1.101	—	64.56	5	M
NGC6397_10	265.1404006970820	-53.6701161565620	13.074	1.056	—	-88.88	6	M
NGC6397_11	265.1366858414970	-53.6749607073660	12.082	1.137	—	-108.30	10	M
NGC6397_13	265.1413447194420	-53.6887671662460	12.196	1.182	—	-29.43	25	M
NGC6397_14	265.1330615276170	-53.6977610910050	12.888	1.122	—	-29.37	17	M
NGC6397_14b	265.13320000	-53.69827000	12.78	—	0.89	35.51	—	M
NGC6397_15	265.1274256674430	-53.7026382997840	13.728	1.052	—	-32.80	0	M
NGC6397_16	265.1279607171810	-53.7067037325510	12.217	1.169	—	-26.86	2	M
NGC6397_16b	265.12800000	-53.70719000	12.07	—	0.93	24.78	—	M
NGC6397_19	265.0925299293380	-53.7185477496650	13.519	1.026	—	-54.15	12	M
NGC6397_319	265.09110000	-53.71521000	12.95	—	1.02	28.31	—	
NGC6397_344	265.08000000	-53.67509000	13.50	—	1.78	84.62	—	
NGC6397_361	265.11960000	-53.64236000	11.67	—	1.08	40.74	—	M
NGC6397_365	265.11770000	-53.62390000	13.08	—	0.88	46.89	—	M
NGC6441_1	267.5515879937640	-36.9959965728170	17.300	1.553	—	-179.54	—	
NGC6441_2	267.5674913837453	-37.0010138178080	17.215	1.651	—	-22.34	—	
NGC6441_3	267.5281735461197	-37.0057433249170	17.327	1.516	—	-62.90	—	M
NGC6441_4	267.5486118676010	-37.0125519130730	16.858	1.549	—	8.19	14	M
NGC6441_5	267.5612985558160	-37.0174913542750	17.425	1.507	—	5.78	—	M

Table B.1 - continued.

Star ID	RA (J2000) (deg)	DEC (J2000) (deg)	V (mag)	V-I (mag)	B-V (mag)	v_{helio} (km/s)	$v_{\text{helio}} - \text{CaT}$ (km/s)	members
NGC6441_6	267.5617704376787	-37.0263698501170	16.052	2.164	—	14.04	9	M
NGC6441_7	267.5192293599370	-37.0314760337140	17.504	1.564	—	-0.10	18	
NGC6441_8	267.5672891635543	-37.0398682583090	17.086	1.696	—	29.56	—	M
NGC6441_9	267.5704202800800	-37.0465647228760	17.211	1.742	—	18.87	37	M
NGC6441_10	267.5262345252269	-37.0520198353200	17.432	1.625	—	-22.21	25	M
NGC6441_11	267.5786097381969	-37.0574957675210	17.039	1.639	—	-188.15	6	
NGC6441_13	267.5335816428650	-37.0708266216330	17.378	1.471	—	-37.60	—	M
NGC6441_14	267.5445841061490	-37.0766569572610	16.444	1.685	—	-149.36	—	
NGC6441_15	267.5297434362500	-37.0830384774400	16.789	1.757	—	-148.33	—	
NGC6441_16	267.5704974086177	-37.0899429393470	17.197	1.694	—	-282.15	—	
NGC6441_17	267.5395804575260	-37.0966969885660	17.214	1.695	—	-22.99	—	
NGC6441_18	267.5270547394850	-37.1012335037780	17.293	1.712	—	-31.15	—	
NGC6441_19	267.5818116124880	-37.1091236311610	17.515	1.729	—	-15.20	—	
NGC6752_2	287.68260	-60.024740	12.490	—	0.900	-25.14	—	M
NGC6752_4	287.70260	-60.031540	13.700	—	0.850	-21.71	—	M
NGC6752_10	287.74220	-60.029200	12.700	—	0.790	-36.69	—	M
NGC6752_11	287.75580	-60.034130	12.240	—	0.640	-35.20	—	M
NGC6752_12	287.76250	-60.029760	11.200	—	1.370	-21.40	—	M
NGC6752_16	287.78380	-60.030550	11.470	—	1.560	10.17	—	
M15_2	322.5214189696987	12.2227087625862	14.398	1.105	—	-223.86	-138	M
M15_3	322.5332798994560	12.2154610609082	14.313	1.135	—	-143.96	-145	M
M15_4	322.4994012317070	12.2084721190462	15.452	0.952	—	-128.98	-122	M
M15_5	322.5611079173707	12.2033365017587	15.648	0.971	—	-136.13	-130	M
M15_6	322.5020644576907	12.1940252155229	14.413	1.084	—	-199.05	-98	M
M15_7	322.5092035836513	12.1897419163095	12.809	1.396	—	-212.69	-102	M
M15_8	322.5178193978010	12.1826621976324	13.751	1.199	—	-120.47	-108	M
M15_9	322.5190490608820	12.1761618053724	14.037	1.147	—	-115.34	-101	M
M15_10	322.5235211413769	12.1714178109786	14.309	1.140	—	-110.11	-95	M
M15_12	322.5271887719680	12.1566131585421	15.432	1.044	—	-189.77	-102	M
M15_13	322.5164140411927	12.1498342128064	15.044	1.073	—	-137.21	-108	M
M15_14	322.5370046077370	12.1474231509706	14.844	1.069	—	-113.98	-124	M
M15_15	322.5309994712117	12.1375868809030	14.037	1.150	—	-161.10	-115	M
M15_17	322.5013766439880	12.1272243863982	13.792	1.180	—	-196.66	-78	M
M15_18	322.5230161717470	12.1185801689600	15.581	1.035	—	-197.80	-101	M
NGC2298_11	102.24680	-36.013800	14.210	—	1.200	122.33	—	M
NGC2298_14	102.22770	-35.987410	14.350	—	1.220	130.55	—	M
NGC2298_16	102.23500	-36.004570	14.770	—	0.950	118.82	—	M
NGC2298_17	102.24570	-36.021050	14.870	—	1.090	145.59	—	M
NGC2298_18	102.23460	-36.012570	14.960	—	1.000	152.26	—	M
NGC4372_20	186.29310	-72.647110	12.880	—	1.570	49.17	—	M
NGC4372_14	186.32230	-72.632710	14.290	—	1.290	54.13	—	M
NGC4372_13	186.33780	-72.626460	12.720	—	1.730	51.30	—	M

Table B.1 - continued.

Star ID	RA (J2000) (deg)	DEC (J2000) (deg)	V (mag)	V-I (mag)	B-V (mag)	v_{helio} (km/s)	$v_{\text{helio}} - \text{CaT}$ (km/s)	members
NGC4372_95	186.46560	-72.621520	14.480	—	1.300	69.04	—	M
NGC4372_91	186.48730	-72.626130	14.450	—	1.330	69.51	—	M
NGC4372_89	186.50650	-72.626710	14.490	—	1.290	61.49	—	M
NGC4372_85	186.55180	-72.629520	14.930	—	1.030	17.25	—	
NGC4372_74	186.57040	-72.657320	14.170	—	1.400	74.77	—	M
NGC4372_76	186.59190	-72.649200	14.180	—	1.300	73.37	—	M
NGC4372_77	186.62360	-72.648570	14.190	—	1.280	14.36	—	
Rup106_2	189.6768998880010	-51.1056842484210	17.181	1.242	—	50.22	—	
Rup106_3	189.69327111237220	-51.1128486896620	17.794	1.234	—	-42.80	-36	M
Rup106_4	189.6538310511420	-51.1187718174390	15.628	1.510	—	-148.79	-54	
Rup106_5	189.6729033037630	-51.1245846157830	18.594	1.185	—	72.46	—	
Rup106_6	189.7040715724640	-51.1288122990960	16.676	1.343	—	-58.38	-49	M
Rup106_7	189.6986910763680	-51.1380491648540	16.959	1.240	—	-7.12	—	
Rup106_8	189.7272422891820	-51.1424906917940	17.286	1.262	—	143.16	—	
Rup106_9	189.7134282848469	-51.1486117989790	17.660	1.209	—	-47.56	-43	M
Rup106_10	189.6808605474090	-51.1557282835810	17.756	1.200	—	-40.06	-34	M
Rup106_11	189.6755230715590	-51.1623524677760	15.478	1.462	—	-40.72	-34	M
Rup106_12	189.6802353325200	-51.1691042971530	17.053	1.255	—	-37.67	-28	M
Rup106_14	189.6397943947549	-51.1828891117730	18.284	1.209	—	-72.74	-49	M
Rup106_16	189.6408966297330	-51.1933813852430	18.518	1.172	—	-67.91	—	
Rup106_17	189.6988243841690	-51.2013261022120	17.279	1.286	—	-37.35	-23	M
Rup106_18	189.6706147758830	-51.2082040394740	18.597	1.155	—	240.83	—	
NGC5634_6	217.4079843945940	-5.9527883388807	17.124	1.044	—	40.71	-7	M
NGC5634_7	217.3907924909180	-5.9560295168940	17.267	0.961	—	-33.63	-45	M
NGC5634_8	217.3992275868640	-5.9641932074133	16.922	0.968	—	-89.80	-44	M
NGC5634_9	217.3908871017030	-5.9705719834359	17.456	1.006	—	-76.17	-32	M
NGC5634_11	217.4131014284530	-5.9818178621402	15.728	1.163	—	-5.86	7	M
NGC5634_13	217.3869685734450	-5.9956708745018	17.634	0.856	—	-26.60	-20	M
NGC5634_14	217.4142199604380	-6.0030100771986	16.571	1.102	—	-41.82	-34	M
NGC5634_15	217.3894998618240	-6.0062776413795	15.316	1.321	—	-3.16	9	M
NGC5634_17	217.3708007790040	-6.0220435949273	17.283	0.957	—	-47.96	-31	
NGC5694_3	219.9348810287710	-26.4946618047940	18.371	0.953	—	-83.00	-71	
NGC5694_5	219.8973138738160	-26.5043633462540	16.864	1.120	—	-168.02	-144	M
NGC5694_6	219.9052304389680	-26.5119464734560	18.410	0.982	—	-153.94	-140	M
NGC5694_7	219.9144178528140	-26.5205518431080	18.058	1.012	—	-143.70	-129	M
NGC5694_8	219.8897697484850	-26.5262032004710	18.374	0.999	—	-142.23	-119	M
NGC5694_9	219.8966177062100	-26.5303560476950	16.666	1.196	—	-157.48	-136	M
NGC5694_10	219.9158491830350	-26.5368552509350	18.443	0.972	—	-177.81	-159	
NGC5694_11	219.8819010796280	-26.5433086004050	18.233	1.002	—	-142.20	-127	M
NGC5694_13	219.9051136155080	-26.5567467822200	17.746	1.055	—	-36.37	-23	
NGC5694_15	219.8873222857810	-26.5705548936230	17.915	1.042	—	-149.53	-137	M
NGC5694_17	219.8940840133800	-26.5805245126430	17.326	1.070	—	-147.01	-127	M

Table B.1 - continued.

Star ID	RA (J2000) (deg)	DEC (J2000) (deg)	V (mag)	V-I (mag)	B-V (mag)	v_{helio} (km/s)	$v_{\text{helio}} - \text{CaT}$ (km/s)	members
NGC5897_138	229.32940	-21.024200	14.350	—	1.180	62.03	—	M
NGC5897_142	229.34370	-21.041560	14.580	—	1.090	94.50	—	M
NGC5897_160	229.38830	-21.039470	13.430	—	1.430	94.29	—	M
NGC5897_255	229.37750	-21.034920	13.400	—	1.800	96.24	—	M
NGC5897_302	229.40740	-21.006760	14.100	—	1.280	96.30	—	M
NGC5897_306	229.40020	-21.041690	14.320	—	1.180	95.76	—	M
NGC5897_316	229.36370	-21.006770	13.860	—	1.340	70.66	—	M
NGC5897_318	229.35660	-21.039150	14.580	—	1.140	91.61	—	M
NGC5927_37	231.97970	-50.662650	15.770	—	1.590	-99.77	—	M
NGC5927_65	231.97280	-50.665090	14.640	—	1.920	-97.59	—	M
NGC5927_133	232.00010	-50.657180	14.750	—	1.970	-116.35	—	M
NGC5927_372	232.00540	-50.696110	14.660	—	2.110	-97.90	—	M
NGC5927_335	232.01040	-50.689630	14.440	—	1.940	-100.23	—	M
NGC5927_190	232.02400	-50.663760	14.290	—	2.020	-82.28	—	M
NGC5946_1	233.8046378431610	-50.6330999838030	16.736	1.764	—	-27.44	-63	
NGC5946_2	233.8192784968850	-50.6347502685450	17.017	1.610	—	-84.94	-105	
NGC5946_3	233.7968271675320	-50.6664482079140	16.499	1.780	—	-5.24	-38	
NGC5946_4	233.8349967731580	-50.6435873111980	16.315	1.728	—	71.73	85	
NGC5946_5	233.8351619708550	-50.6556161766250	16.229	1.809	—	115.64	112	M
NGC5946_6	233.8533263570141	-50.6441955351030	16.865	1.641	—	131.54	96	
NGC5946_7	233.8447146767090	-50.6705180815020	15.955	1.793	—	106.48	96	M
NGC5946_8	233.8595939690180	-50.6634568192010	15.780	1.866	—	162.46	116	M
NGC5946_9	233.8666866915190	-50.6714124930590	16.745	1.745	—	118.45	100	M
NGC5946_10	233.8951206530490	-50.6490984493930	17.119	1.698	—	139.52	130	
NGC5946_11	233.8903132732620	-50.6670769475470	17.160	1.669	—	143.52	129	
NGC5946_14	233.9077804562540	-50.6840449718950	16.261	1.756	—	109.52	172	
NGC5946_15	233.9313592396560	-50.6687249102040	16.748	1.706	—	168.90	116	M
NGC5946_16	233.9304730779910	-50.6845904460230	16.989	1.722	—	111.27	-81	
NGC5946_19	233.9375579411780	-50.7135206532660	16.543	1.643	—	-83.91	—	
BH176_2	234.8322480854411	-49.9847554870280	19.377	1.720	—	17.00	27	M
BH176_3	234.8258020595950	-49.9905808846840	18.826	1.709	—	55.13	36	
BH176_4	234.7353061460010	-49.9945984823580	19.073	1.718	—	-15.41	-12	
BH176_6	234.7674651957070	-50.0097045530970	18.249	2.006	—	-2.51	-4	M
BH176_7	234.7792297073740	-50.0131437707750	18.411	1.959	—	-37.48	5	M
BH176_8	234.7325900671540	-50.0193875514340	17.766	2.294	—	10.42	7	
BH176_9	234.7931783392740	-50.0273857906780	18.985	1.721	—	-17.84	5	M
BH176_10	234.8025091323060	-50.0330453671780	17.976	2.058	—	-5.52	-5	M
BH176_11	234.8017269343900	-50.0385856964930	18.066	2.113	—	-1.22	-2	M
BH176_13	234.7825982516510	-50.0528938288310	17.408	2.755	—	-3.58	-5	M
BH176_14	234.7519927464920	-50.0596122899570	18.101	1.909	—	-6.45	-6	M
BH176_15	234.7656274407960	-50.0638176373480	18.775	1.796	—	7.40	-2	M
BH176_16	234.8019923942540	-50.0708836705370	18.617	1.810	—	-14.69	-0	M
BH176_17	234.7518063534750	-50.0797145124840	18.144	2.010	—	-25.68	-16	

Table B.1 - continued.

Star ID	RA (J2000) (deg)	DEC (J2000) (deg)	V (mag)	V-I (mag)	B-V (mag)	v_{helio} (km/s)	$v_{\text{helio}} - \text{CaT}$ (km/s)	members
BH176_18	234.7603405866310	-50.0830708858910	17.951	2.030	—	0.20	-75	M
Lynga7_1	242.7143087690700	-55.2905208683670	16.706	1.988	—	-112.43	—	
Lynga7_2	242.7653623624860	-55.2979759401890	16.114	2.078	—	41.69	6	
Lynga7_3	242.7793235806400	-55.3040472206520	16.402	1.973	—	12.75	23	
Lynga7_4	242.7493485490450	-55.3104250625750	16.724	1.959	—	-33.95	21	M
Lynga7_5	242.7686462362690	-55.3153817813770	15.380	3.168	—	9.13	18	
Lynga7_6	242.7471523509420	-55.3235157955460	17.174	1.806	—	-56.46	—	
Lynga7_8	242.7497743196010	-55.3351275639070	16.150	2.018	—	-23.86	17	M
Lynga7_9	242.7942370919550	-55.3409352713680	16.268	2.038	—	-35.79	—	
Lynga7_10	242.7572950981700	-55.3488247445700	17.110	1.999	—	18.90	27	M
Lynga7_11	242.7749423241250	-55.3527911736560	16.418	2.102	—	16.30	34	
Lynga7_14	242.7817767699260	-55.3749439667090	17.164	1.856	—	61.49	—	
Lynga7_15	242.7553826491850	-55.3774876150520	15.606	2.554	—	8.66	27	
Lynga7_16	242.6969892779080	-55.3836099651490	17.110	2.015	—	-84.30	—	
Pal14_3	242.7289540671210	14.9768305596526	18.302	1.126	—	50.59	71	M
Pal14_4	242.7252997753470	14.9515772785923	18.755	1.117	—	46.46	66	M
Pal14_5	242.7372606969950	14.9656721115292	18.870	1.069	—	53.74	72	M
Pal14_7	242.7447676166120	14.9468796368414	17.749	1.351	—	48.97	75	M
Pal14_8	242.7560206091760	14.9665979935925	19.117	1.105	—	28.09	75	M
Pal14_10	242.7747630175660	14.9785733163268	19.312	1.079	—	20.98	58	
Pal14_11	242.7743710727580	14.9624234573138	17.367	1.421	—	43.94	81	M
NGC6284_1	256.1787421768040	-24.7239654554000	15.863	1.460	—	-128.96	-56	
NGC6284_2	256.1815532872069	-24.7319915884480	14.690	1.729	—	-278.63	-219	
NGC6284_3	256.1570456950431	-24.7294863031900	16.947	1.205	—	-0.67	10	
NGC6284_4	256.1338305379220	-24.7250439225030	16.404	1.352	—	-141.74	-100	
NGC6284_5	256.1489134049400	-24.7375092679340	16.038	1.393	—	-171.37	-194	
NGC6284_6	256.1745929082130	-24.7556796415000	15.369	1.549	—	-66.23	-81	
NGC6284_7	256.1425619205140	-24.7501496060640	16.571	1.441	—	40.74	22	M
NGC6284_8	256.1319097101249	-24.7527760959490	15.690	1.497	—	-24.10	-19	
NGC6284_9	256.1289997288189	-24.7592355039020	16.069	1.398	—	64.38	8	M
NGC6284_10	256.1127878908740	-24.7569225247240	16.179	1.433	—	27.53	4	M
NGC6284_11	256.1418949500970	-24.7775725446190	16.132	1.406	—	-82.35	-51	
NGC6284_13	256.1190234932520	-24.7797550947810	16.274	1.426	—	52.78	23	M
NGC6284_14	256.1076461193801	-24.7833849093920	16.543	1.392	—	-1.56	33	M
NGC6284_15	256.1271039938770	-24.7977459030990	16.966	1.379	—	28.95	22	M
NGC6284_16	256.1125261800070	-24.8003947883760	15.310	1.581	—	-40.52	-94	
NGC6284_17	256.1021024388910	-24.8009876097890	16.329	1.367	—	-14.32	-2	
NGC6284_18	256.1176796144840	-24.8149306835980	16.278	1.388	—	29.26	32	M
NGC6316_1	259.0963084956139	-28.0873501900400	16.646	1.978	—	-81.46	-49	
NGC6316_2	259.1166970225880	-28.0861498907930	16.941	1.906	—	-30.22	-15	
NGC6316_3	259.1160790626469	-28.0949073119040	16.716	1.733	—	-55.20	-32	
NGC6316_4	259.1377284457139	-28.0971532857540	17.074	1.802	—	-45.83	-42	

Table B.1 - continued.

Star ID	RA (J2000) (deg)	DEC (J2000) (deg)	V (mag)	V-I (mag)	B-V (mag)	v_{helio} (km/s)	$v_{\text{helio}} - \text{CaT}$ (km/s)	members
NGC6316_5	259.1312272086139	-28.1036357862990	17.090	1.842	—	-125.06	16	
NGC6316_6	259.1273755069050	-28.1099285456990	16.835	1.726	—	-85.98	-74	
NGC6316_7	259.1477506136241	-28.1154189764410	16.385	1.861	—	36.38	77	M
NGC6316_8	259.1311023859710	-28.1247550044380	16.508	1.768	—	-180.17	-169	
NGC6316_9	259.1623377367640	-28.1225520700910	15.828	2.199	—	154.08	97	M
NGC6316_10	259.1721802632810	-28.1249274390170	16.684	1.876	—	87.13	—	M
NGC6316_11	259.1725801971090	-28.1318873124630	16.462	1.839	—	100.53	96	M
NGC6316_14	259.1616428829790	-28.1535857831530	16.856	1.828	—	86.61	-78	M
NGC6316_15	259.1651348929910	-28.1621833867960	16.452	2.006	—	-80.73	42	
NGC6316_17	259.1734767954070	-28.1732067264740	16.590	1.767	—	37.62	83	M
NGC6316_18	259.1483852088890	-28.1861490142080	16.893	1.735	—	68.00	-11	M
NGC6316_19	259.1654467639340	-28.1876061552910	16.431	1.960	—	-10.30	—	
NGC6356_1	260.9235074077297	-17.7546041249480	16.660	1.491	—	5.87	45	M
NGC6356_2	260.9558057842270	-17.7710305110450	16.723	1.422	—	0.40	—	
NGC6356_3	260.9558705466089	-17.7784978788010	17.082	1.385	—	212.79	—	
NGC6356_4	260.8954703282600	-17.7645922141030	15.741	1.953	—	-232.62	—	
NGC6356_5	260.9239185919519	-17.7846685306150	16.873	1.632	—	39.08	50	M
NGC6356_6	260.9169091259810	-17.7881376496330	16.676	1.450	—	-8.54	—	M
NGC6356_7	260.9279656805810	-17.7966787950700	16.561	1.616	—	42.36	53	M
NGC6356_8	260.8932599793060	-17.7917801507460	15.779	1.819	—	-54.42	64	M
NGC6356_9	260.9156319574237	-17.8087940477470	16.795	1.542	—	71.72	82	M
NGC6356_10	260.9214777632967	-17.8143867084250	16.546	1.464	—	51.41	64	M
NGC6356_11	260.9114420443020	-17.8179999655190	15.950	1.570	—	21.22	—	
NGC6356_13	260.9115145337869	-17.8309059820990	15.934	1.643	—	93.58	88	M
NGC6356_14	260.9222838412350	-17.8419564642120	16.165	1.640	—	70.53	75	M
NGC6356_15	260.9314184453367	-17.8502978968420	15.256	1.917	—	3.79	—	
NGC6356_16	260.8948341251553	-17.8448271016070	16.721	1.428	—	3.64	60	M
NGC6356_17	260.8988194150897	-17.8524578021880	15.408	2.129	—	52.95	—	M
NGC6356_18	260.9039357772895	-17.8638101621960	16.709	1.595	—	79.51	90	M
NGC6356_19	260.8706044317470	-17.8568782386050	16.175	1.539	—	26.48	75	M
NGC6355_1	261.0139708308290	-26.2986443619180	16.602	2.021	—	-20.92	-60	
NGC6355_2	260.9876263216710	-26.3032970647830	17.364	1.956	—	-42.22	-50	
NGC6355_3	260.9645602791170	-26.3095794146230	16.960	2.022	—	365.88	367	
NGC6355_4	261.0034840024430	-26.3133409142560	17.475	1.871	—	-127.08	-124	
NGC6355_5	260.9765675424109	-26.3222867639420	16.549	2.118	—	-205.83	-215	
NGC6355_6	260.9776929200140	-26.3281193397190	17.150	2.021	—	-50.62	-53	
NGC6355_7	260.9909410424039	-26.3357769969130	17.402	1.946	—	-224.20	-207	M
NGC6355_8	260.9811462177399	-26.3404718116360	15.739	2.195	—	-211.17	-219	M
NGC6355_9	260.9819204193250	-26.3455751719490	15.639	2.156	—	-218.51	-200	M
NGC6355_10	260.9994267993690	-26.3538344802030	16.595	1.940	—	-212.99	-205	M
NGC6355_11	261.0034555787130	-26.3596188513650	15.896	2.068	—	-204.05	-207	M
NGC6355_13	260.9972671624630	-26.3741315224720	16.880	2.072	—	-216.96	-214	M
NGC6355_14	260.9929749853910	-26.3766231832010	16.584	1.938	—	-84.52	-84	

Table B.1 - continued.

Star ID	RA (J2000) (deg)	DEC (J2000) (deg)	V (mag)	V-I (mag)	B-V (mag)	v_{helio} (km/s)	$v_{\text{helio}} - \text{CaT}$ (km/s)	members
NGC6355_15	260.9830679562149	-26.3874489320860	17.542	1.900	—	-189.60	-242	
NGC6355_17	260.9655114255959	-26.3960577501580	15.973	2.185	—	102.94	94	
NGC6355_18	261.0137987038789	-26.4056110757630	16.916	2.081	—	18.74	49	
NGC6352_1	261.4410166794020	-48.3642026590000	14.948	1.301	—	-121.65	-124	M
NGC6352_2	261.4129538199270	-48.3685108494390	15.079	1.310	—	-7.12	-37	
NGC6352_5	261.4355040079679	-48.3897451899930	14.423	1.456	—	-141.24	-151	M
NGC6352_6	261.3453211212850	-48.3877354011100	15.348	1.326	—	-178.77	-114	M
NGC6352_8	261.3562017080330	-48.3991912951620	14.073	1.545	—	-137.52	-122	M
NGC6352_9	261.3785247710999	-48.4085480198530	15.134	1.308	—	-134.08	-134	M
NGC6352_10	261.3407913170790	-48.4109981946450	14.556	1.515	—	-223.85	-118	M
NGC6352_11	261.3749274397111	-48.4222219762690	13.422	1.870	—	-99.30	-135	M
NGC6352_13	261.3708585654150	-48.4354209604850	15.231	1.291	—	-135.80	-128	M
NGC6352_14	261.3334026622210	-48.4350178054250	14.050	1.464	—	-118.67	-139	
NGC6352_15	261.3505553097240	-48.4453305951550	15.146	1.323	—	-98.24	-134	M
NGC6352_16	261.3608831766571	-48.4516101506590	14.494	1.402	—	-113.99	-130	M
NGC6352_17	261.3675556709620	-48.4601120214010	13.750	1.617	—	-161.47	-113	M
NGC6352_18	261.4110421627360	-48.4703858199440	14.523	1.458	—	-108.31	-113	M
NGC6366_1	261.9095291704430	-5.0195910671494	16.479	1.895	—	-170.57	-138	M
NGC6366_2	261.9045681025580	-5.0268560605880	15.875	1.820	—	-174.56	-108	M
NGC6366_3	261.9461497402918	-5.0310026920141	16.626	1.888	—	-143.24	-133	M
NGC6366_4	261.9098940585019	-5.0401070162215	16.271	1.893	—	-191.77	-113	M
NGC6366_5	261.9353344668059	-5.0439214509264	14.786	2.005	—	-121.28	-126	M
NGC6366_6	261.9489990876769	-5.0494799166147	15.874	1.814	—	-119.21	-117	
NGC6366_7	261.9582332068019	-5.0596384523628	16.707	1.857	—	-177.89	-117	M
NGC6366_8	261.9226967769650	-5.0629107784532	16.136	1.862	—	-225.87	-119	M
NGC6366_9	261.9194012789320	-5.0716868933516	15.189	2.020	—	-138.71	-122	M
NGC6366_10	261.9612595505589	-5.0771209262480	16.262	1.893	—	-33.21	-106	M
NGC6366_11	261.9221402785569	-5.0835030019369	15.692	1.979	—	-70.78	-117	M
NGC6366_13	261.9387570683969	-5.0951106568610	16.757	1.857	—	-176.64	-108	M
NGC6366_14	261.8998623184320	-5.0999764100051	15.716	1.874	—	-73.28	-108	
NGC6366_15	261.9312457422168	-5.1084049060473	14.424	2.005	—	-121.52	-126	M
NGC6366_16	261.9290313916749	-5.1158275368716	16.134	1.858	—	-96.74	-121	M
NGC6366_17	261.9595558876100	-5.1193281721352	15.535	1.859	—	-36.46	39	
NGC6366_18	261.9501356527969	-5.1277765435729	14.991	1.942	—	-69.04	-120	M
NGC6401_1	264.6740793857970	-23.8556633971030	17.321	1.995	—	-41.36	-47	
NGC6401_2	264.6396680991930	-23.8610421303060	16.599	2.058	—	-31.56	-38	
NGC6401_3	264.6568766516490	-23.8647642881960	16.924	2.072	—	-92.44	-103	M
NGC6401_4	264.6900204428919	-23.8740817309480	15.314	2.331	—	-1.69	26	
NGC6401_5	264.6308167036780	-23.8784535100230	17.142	2.193	—	-11.11	3	
NGC6401_6	264.6856894020570	-23.8866262403780	16.785	2.065	—	-42.67	-38	
NGC6401_7	264.6465304337050	-23.8920274646890	17.166	2.039	—	-120.97	-96	M
NGC6401_8	264.6493729355540	-23.8963685172360	16.706	2.175	—	-144.76	-137	M
NGC6401_9	264.6606431183770	-23.9041214598170	16.114	2.081	—	-125.41	-111	M

Table B.1 - continued.

Star ID	RA (J2000) (deg)	DEC (J2000) (deg)	V (mag)	V-I (mag)	B-V (mag)	v_{helio} (km/s)	$v_{\text{helio}} - \text{CaT}$ (km/s)	members
NGC6401_10	264.6672999336120	-23.9085421356730	16.484	2.243	—	-184.47	-169	
NGC6401_11	264.6484872492470	-23.9158401886910	16.911	2.215	—	-106.33	-63	
NGC6401_12	264.6638409242120	-23.9232830795760	16.478	2.036	—	-137.77	-129	
NGC6401_13	264.6507855666670	-23.9279831483290	17.374	2.054	—	-113.77	-110	M
NGC6401_14	264.6583349865899	-23.9362185412110	17.236	2.139	—	-120.74	-121	M
NGC6401_15	264.6719233586430	-23.9406429153550	16.807	2.202	—	38.81	63	
NGC6401_16	264.6476908074171	-23.9471300975680	17.281	2.191	—	-133.97	-130	
NGC6401_17	264.6466533919830	-23.9540215315450	17.528	2.167	—	-106.95	-103	
NGC6401_18	264.6693516511750	-23.9612225098970	16.655	2.251	—	104.17	81	
Pal6_1	265.9138498528130	-26.1649109273710	18.265	3.165	—	-35.47	-22	
Pal6_2	265.9034035887530	-26.1709532492550	19.121	3.075	—	76.42	73	
Pal6_3	265.9530750251889	-26.1800411651570	18.050	3.207	—	12.61	-48	
Pal6_4	265.9495279366710	-26.1869191371730	18.520	3.064	—	-70.25	-55	
Pal6_5	265.9369039009599	-26.1886178964360	19.231	2.877	—	-69.80	-72	
Pal6_6	265.9227793854060	-26.1978613619770	19.082	3.084	—	10.35	8	
Pal6_7	265.9224889666719	-26.2047843344080	18.918	3.050	—	35.18	53	
Pal6_8	265.9065700629099	-26.2094174719570	19.257	2.892	—	96.92	107	
Pal6_9	265.9415791651939	-26.2176098802120	17.270	3.371	—	173.50	178	M
Pal6_10	265.9367765058209	-26.2221073256090	18.846	3.052	—	-90.74	-82	
Pal6_11	265.9257829144390	-26.2286324458830	18.153	3.114	—	182.53	186	M
Pal6_13	265.9337087726818	-26.2412589944980	18.368	3.020	—	173.24	178	M
Pal6_14	265.9369008632410	-26.2485857092230	17.903	3.162	—	180.24	174	M
Pal6_15	265.9447640007471	-26.2521501635350	18.045	3.192	—	-72.40	-56	
Pal6_16	265.9375657316269	-26.2585990959490	18.240	3.029	—	128.42	135	
Pal6_17	265.9316835487689	-26.2652504769740	18.543	3.047	—	134.19	137	
Pal6_18	265.9419895113390	-26.2712003357810	18.774	3.094	—	149.91	148	
NGC6440_1	267.1860577172569	-20.3029171413500	17.359	2.790	—	73.91	—	
NGC6440_3	267.2094912561130	-20.3149347371460	17.941	2.487	—	25.88	—	
NGC6440_4	267.2101077506937	-20.3219764394440	18.251	2.406	—	-160.93	—	
NGC6440_5	267.2035072858610	-20.3291722920080	16.990	2.895	—	32.72	—	
NGC6440_6	267.2171388334789	-20.3351667481490	18.048	2.572	—	-31.95	—	M
NGC6440_7	267.2251585557580	-20.3402378225310	18.500	2.397	—	-23.02	—	M
NGC6440_8	267.2344986043180	-20.3485639174310	17.291	2.986	—	-93.13	-97	M
NGC6440_10	267.2320462698550	-20.3589215988520	17.623	2.702	—	-86.97	-84	M
NGC6440_11	267.2061834512457	-20.3660566970390	17.028	2.752	—	9.90	-78	
NGC6440_13	267.2269927367090	-20.3783455444010	18.202	2.486	—	-52.54	-57	M
NGC6440_14	267.2193743591089	-20.3852127997750	16.403	3.100	—	-65.02	-73	M
NGC6440_15	267.2241545171047	-20.3913077156530	17.568	2.623	—	-60.78	-62	M
NGC6440_16	267.2288277646500	-20.3981523854640	18.226	2.522	—	187.35	—	
NGC6440_17	267.2282071898930	-20.4036506064310	16.477	3.202	—	121.48	—	
NGC6440_18	267.1830234769110	-20.4099430892500	17.703	2.515	—	82.84	—	
NGC6440_19	267.2175680421177	-20.4169636952160	18.247	2.349	—	63.82	—	
NGC6453_1	267.7109046327830	-34.5432736160450	17.199	1.696	—	10.50	8	

Table B.1 - continued.

Star ID	RA (J2000) (deg)	DEC (J2000) (deg)	V (mag)	V-I (mag)	B-V (mag)	v_{helio} (km/s)	$v_{\text{helio}} - \text{CaT}$ (km/s)	members
NGC6453_2	267.7099404707681	-34.5472647329590	16.528	1.682	—	-189.04	-186	
NGC6453_3	267.7117060844460	-34.5547399421890	17.277	1.797	—	83.37	96	
NGC6453_4	267.7292470188960	-34.5605918640070	17.585	1.606	—	-107.16	-115	
NGC6453_5	267.7102053888260	-34.5691401085550	17.429	1.770	—	-32.00	-9	
NGC6453_6	267.7058201621750	-34.5732638316630	16.916	1.789	—	-188.36	-184	
NGC6453_7	267.7289160763439	-34.5823533377810	17.387	1.724	—	-66.95	-51	
NGC6453_8	267.7226546849880	-34.5879325333010	16.314	1.772	—	-90.08	-95	
NGC6453_10	267.7124990715680	-34.5983430405210	15.536	1.895	—	-158.83	-118	M
NGC6453_11	267.7231285105429	-34.6042698746130	16.405	1.851	—	-140.70	-115	M
NGC6453_13	267.7068837766470	-34.6157585913080	17.533	1.790	—	36.24	66	
NGC6453_14	267.6983936335840	-34.6228403407740	17.334	1.662	—	-165.87	-144	
NGC6453_15	267.7245315396250	-34.6329408571700	16.787	1.662	—	-160.70	-137	M
NGC6453_16	267.7355219144729	-34.6352324389380	17.151	1.777	—	-189.33	-153	
NGC6453_17	267.7228026610230	-34.6422184403970	17.543	1.628	—	-36.75	1	
NGC6453_18	267.7081228730470	-34.6483015798160	17.352	1.717	—	-133.40	-91	
NGC6538_2	271.1891346933031	-7.5368165863050	17.781	2.381	—	-15.60	9	
NGC6538_3	271.2225768215739	-7.5417614936001	18.400	2.342	—	-9.51	8	
NGC6538_4	271.2115914147460	-7.5466212743910	16.401	2.872	—	51.55	57	M
NGC6538_5	271.2152706610500	-7.5529669666963	16.797	2.547	—	10.82	28	M
NGC6538_6	271.2172879051260	-7.5588502786937	18.358	2.309	—	71.93	106	
NGC6538_7	271.2209245927010	-7.5663514469969	17.934	2.521	—	51.64	70	
NGC6538_8	271.2207360141699	-7.5730120587253	17.613	2.422	—	4.51	38	M
NGC6538_9	271.2084174690189	-7.5788454320773	18.434	2.271	—	32.01	28	M
NGC6538_10	271.1990876725411	-7.5857819095202	17.873	2.442	—	48.05	31	M
NGC6538_11	271.2128916424860	-7.5903446252700	16.261	3.137	—	88.66	54	
NGC6538_13	271.2003494122119	-7.6051381264440	16.110	2.981	—	39.20	32	
NGC6538_14	271.1988435054970	-7.6100848355027	16.554	2.613	—	23.50	31	M
NGC6538_15	271.1793873281880	-7.6175695473128	17.061	2.428	—	47.08	31	
NGC6538_17	271.2098010791651	-7.6296450723905	18.250	2.281	—	-14.98	-6	
NGC6538_18	271.1994704411969	-7.6369369734792	18.316	2.320	—	36.64	31	M
Pal7_1	272.7129181253690	-7.1538632982032	17.357	2.592	—	169.18	162	M
Pal7_3	272.6941865319670	-7.1655922878238	17.052	2.559	—	155.32	161	M
Pal7_4	272.7038149516796	-7.1737082694418	15.969	3.115	—	179.41	180	M
Pal7_5	272.6553330306050	-7.1729570563556	17.146	2.675	—	168.15	—	
Pal7_7	272.6814999578997	-7.1895031554166	16.472	2.922	—	123.67	160	M
Pal7_8	272.6940609349913	-7.1960174655907	17.303	2.626	—	162.24	172	M
Pal7_9	272.6856585188370	-7.2029350900190	16.323	3.030	—	159.29	153	M
Pal7_10	272.6907524082347	-7.2083277352292	16.186	2.833	—	162.91	165	M
Pal7_11	272.6889448185320	-7.2141697502301	16.844	2.560	—	158.08	160	M
Pal7_12	272.6739324450837	-7.2210562124949	17.387	2.542	—	153.47	169	M
Pal7_13	272.6947842420930	-7.2270852702464	17.204	2.479	—	-4.88	—	
Pal7_14	272.6732928068980	-7.2332635490277	17.142	2.563	—	-48.90	—	
Pal7_15	272.6350120593880	-7.2349830905768	17.186	2.594	—	139.93	173	M

Table B.1 - continued.

Star ID	RA (J2000) (deg)	DEC (J2000) (deg)	V (mag)	V-I (mag)	B-V (mag)	v_{helio} (km/s)	$v_{\text{helio}} - \text{CaT}$ (km/s)	members
Pal7_16	272.6481250359010	-7.2447529133249	16.850	2.574	—	36.18	156	
Pal7_17	272.6667908626877	-7.2499074682881	17.100	2.498	—	141.10	180	M
Pal7_18	272.6644089239120	-7.2579028580247	16.091	2.816	—	157.52	162	M
Pal7_19	272.6662718121410	-7.2637347001366	17.182	2.511	—	104.85	—	
NGC6569_1	273.4336292334870	-31.7704262455560	17.092	1.649	—	-39.46	-40	M
NGC6569_2	273.4455484196830	-31.7765360025320	15.317	2.032	—	71.91	—	
NGC6569_3	273.4264817161510	-31.7834977305820	16.853	1.769	—	-97.91	—	
NGC6569_4	273.4091675473910	-31.7897212799030	17.040	1.775	—	-56.15	-54	M
NGC6569_5	273.3803188915610	-31.7961359918000	16.084	1.690	—	-157.26	—	
NGC6569_6	273.4318560737670	-31.8015552547670	17.281	1.536	—	87.49	—	
NGC6569_7	273.4181355474257	-31.8088466519520	17.069	1.657	—	27.40	—	
NGC6569_8	273.4162067086087	-31.8157065176360	17.282	1.618	—	-62.73	—	M
NGC6569_9	273.4101047591937	-31.8185969759490	16.990	1.586	—	-41.93	-45	M
NGC6569_10	273.4226581650420	-31.8270607419080	16.352	1.747	—	-45.95	-42	M
NGC6569_11	273.4008204511300	-31.8316386498970	16.765	1.651	—	-52.06	-39	
NGC6569_12	273.4080675327830	-31.8415956101290	17.147	1.588	—	-55.60	-55	M
NGC6569_13	273.4228352776827	-31.8457683686530	16.760	1.798	—	-54.59	-52	M
NGC6569_14	273.4091489443977	-31.8518340957680	17.424	1.682	—	-46.50	—	
NGC6569_15	273.4246883581039	-31.8585619278720	17.326	1.656	—	-31.28	—	
NGC6569_16	273.4040223818470	-31.8642141148940	16.822	1.669	—	-114.62	—	
NGC6569_17	273.4264176766977	-31.8704671125370	17.199	1.709	—	-32.93	—	
NGC6569_19	273.3876649349320	-31.8844286986320	17.394	1.644	—	-4.56	—	
NGC6749_1	286.2847567371839	1.9596903589911	17.585	2.754	—	-43.71	-70	
NGC6749_2	286.2868656096230	1.9525164460737	18.576	2.564	—	33.15	64	
NGC6749_3	286.3284566482919	1.9445190618570	18.593	2.452	—	-72.70	-52	
NGC6749_4	286.3376510500939	1.9399435480177	17.455	2.657	—	-47.40	-37	
NGC6749_5	286.3067566745090	1.9354624935870	17.176	2.970	—	-78.84	-62	
NGC6749_6	286.3152627412300	1.9288996628548	17.325	2.821	—	-85.30	-44	
NGC6749_7	286.3096862766980	1.9192341679555	17.031	2.924	—	-84.10	-64	
NGC6749_8	286.3217847655430	1.9126298767745	17.949	2.716	—	-72.06	-58	M
NGC6749_9	286.3199408737951	1.9074423842532	16.898	2.950	—	-85.06	-63	
NGC6749_10	286.3322975895430	1.9038472343615	16.993	2.858	—	-54.09	-30	M
NGC6749_11	286.3260013003970	1.8982261632430	17.073	2.860	—	-69.88	-48	M
NGC6749_13	286.3332319080549	1.8832013430046	17.504	2.738	—	2.75	34	
NGC6749_14	286.3491620208251	1.8790442360165	17.856	2.717	—	-31.05	-35	
NGC6749_15	286.3258746626020	1.8689929106830	17.353	2.853	—	-67.39	-53	M
NGC6749_16	286.3078692410110	1.8652342887235	18.171	2.701	—	-76.61	-49	
NGC6749_17	286.3067004905620	1.8590099439023	18.230	2.588	—	0.52	21	
NGC6749_18	286.3123536019180	1.8515819277174	17.544	2.844	—	43.34	53	
Pal10_1	289.5125606154301	18.6299558446680	19.242	2.904	—	65.92	38	
Pal10_2	289.4646838783200	18.6277908909462	19.017	2.947	—	-22.78	-1	M
Pal10_3	289.4688827393061	18.6215883550814	19.378	2.944	—	18.76	20	
Pal10_4	289.4899243412699	18.6123372877962	18.869	3.045	—	-49.04	-35	M

Table B.1 - continued.

Star ID	RA (J2000) (deg)	DEC (J2000) (deg)	V (mag)	V-I (mag)	B-V (mag)	v_{helio} (km/s)	$v_{\text{helio}} - \text{CaT}$ (km/s)	members
Pal10.6	289.4656372047450	18.6046132108929	18.147	3.368	—	-112.84	-101	
Pal10.7	289.4652506135770	18.5959932118453	18.449	3.072	—	-27.55	-43	
Pal10.8	289.5027113535670	18.5872513699130	18.901	3.143	—	-46.90	-42	M
Pal10.9	289.5280987147370	18.5773513020153	18.924	3.085	—	-65.87	-69	M
Pal10.10	289.4984921319600	18.5759838819809	18.537	3.167	—	-41.28	-35	M
Pal10.11	289.5162605322620	18.5685585953761	18.165	3.354	—	-35.28	-34	M
Pal10.13	289.5236933727690	18.5551426686763	18.546	3.291	—	-11.91	-15	M
Pal10.16	289.5093507683760	18.5361496794223	18.655	2.994	—	-48.26	-18	M
Pal10.17	289.5264177399050	18.5275771598552	18.554	3.104	—	-17.88	-25	M
Pal11.2	296.2918886563560	-7.9607733525394	17.783	1.313	—	-75.74	-77	M
Pal11.3	296.2983701502389	-7.9679129619890	17.991	1.374	—	-105.55	-94	
Pal11.6	296.3121063991420	-7.9855545899986	17.931	1.327	—	-107.49	-66	M
Pal11.7	296.3266258731740	-7.9901956699317	17.953	1.347	—	-76.90	-67	M
Pal11.8	296.3100194350170	-7.9998724816952	16.791	1.482	—	-91.10	-59	M
Pal11.9	296.3095590678440	-8.0041382789580	16.292	1.572	—	-64.31	-57	M
Pal11.10	296.3031036009810	-8.0127983094640	18.018	1.255	—	-72.92	-60	M
Pal11.11	296.3336414063500	-8.0177424751638	17.018	1.526	—	-80.89	-59	M
Pal11.12	296.3194836366321	-8.0224404692151	17.345	1.426	—	-99.30	-63	M
Pal11.13	296.3052561711690	-8.0296541122148	18.129	1.248	—	-83.91	-60	M
Pal11.16	296.3055388234640	-8.0477687172836	18.132	1.263	—	-49.68	-49	
Pal11.17	296.2948864246590	-8.0569353684763	16.173	1.561	—	-58.82	-54	M
M75.2	301.5722924809030	-21.8724239925710	15.351	1.584	—	-180.58	-168	M
M75.3	301.5332302439650	-21.8792783822790	16.052	1.399	—	-268.32	-195	
M75.4	301.5112863868840	-21.8838393125550	17.305	1.161	—	-209.35	-201	M
M75.5	301.5398310455530	-21.8901698871600	16.911	1.222	—	-211.96	-196	M
M75.6	301.5363529935929	-21.8994513202970	16.667	1.223	—	-170.35	-193	M
M75.7	301.5284174072768	-21.9043265871400	15.075	1.691	—	-159.03	-191	M
M75.8	301.5367067586659	-21.9096006038980	15.446	1.494	—	-242.29	-187	
M75.9	301.5170655412350	-21.9142047369940	15.189	1.677	—	-208.00	-212	M
M75.10	301.5041416139131	-21.9220494603590	15.723	1.448	—	-168.25	-192	M
M75.11	301.5300056045850	-21.9276243871480	15.446	1.543	—	-209.20	-185	M
M75.13	301.5065335843889	-21.9420631700740	16.847	1.147	—	-194.14	-195	M
M75.14	301.5168102614069	-21.9491974411900	16.919	1.150	—	-194.05	-195	M
NGC7006_9	315.3682175549997	16.1917858619990	17.000	1.195	—	-406.56	-377	
NGC7006_18	315.3795161685969	16.1968831361999	18.478	1.067	—	-375.70	-379	M
NGC7006_19	315.3593718729330	16.1997473878717	17.738	1.104	—	-283.61	-379	
NGC7006_21	315.3686452062430	16.2046868611790	19.389	0.940	—	-433.24	-387	M
NGC7006_22	315.3765269702437	16.2072140496116	19.165	0.963	—	-359.48	-369	
NGC7006_23	315.3705323766769	16.2116463895666	16.947	1.200	—	-389.45	-386	M
NGC7006_24	315.3786288060080	16.2195341119763	17.252	1.189	—	-380.73	-379	M
NGC7006_25	315.3822326435830	16.2264922925027	18.094	0.991	—	-378.19	-385	M
NGC7006_26	315.3388329980339	16.2098296634600	17.809	1.045	—	-477.01	-370	
NGC7006_27	315.3393898303030	16.2235240225451	18.997	0.931	—	-84.08	—	

Table B.1 - continued.

Star ID	RA (J2000) (deg)	DEC (J2000) (deg)	V (mag)	V-I (mag)	B-V (mag)	v_{helio} (km/s)	$v_{\text{helio}} - \text{CaT}$ (km/s)	members
NGC7006_28	315.3876249350990	16.2337482232222	18.878	1.055	—	-69.38	—	
NGC7006_29	315.3737767394420	16.1656969965715	18.654	1.007	—	-93.62	—	
NGC7006_31	315.3696240893173	16.1583516837794	17.635	1.158	—	-34.91	—	
NGC7006_32	315.3863374169697	16.1547441200559	18.302	1.065	—	-379.42	—	

^aDue to low number of observed stars it was hard to define the membership of M 107. See discussion in Chapter 4

Table B.2 - List of 49 well known stars selected from ELODIE spectral library. Literature parameters are average from PASTEL catalogue. Results from this work are using MILES library. See details in Section 3.4.2.2.

ELODIE	Star	RR _{tot}	N	S _{lim}	T _{eff} (K)	log(g) (literature)	[Fe/H]	T _{eff} (K)	logg (this work)	[Fe/H]	[Mg/Fe]
1	HD000245	0.107	2	1.042	5490±153	3.48±0.15	-0.77±0.08	5378±60	3.67±0.06	-0.84±0.20	0.34±0.08
9	HD002796	0.062	1	1.000	4931±60	1.45±0.34	-2.32±0.11	4945±133	1.36±0.08	-2.31±0.11	0.37±0.08
19	HD004395	0.042	5	1.116	5487±38	3.33±0.05	-0.35±0.04	5330±149	3.24±0.16	-0.34±0.11	0.18±0.07
31	HD006833	0.092	1	1.000	4426±95	1.28±0.32	-0.91±0.14	4380±217	1.25±0.64	-0.99±0.29	0.30±0.11
32	HD006920	0.035	8	1.319	5886±111	3.60±0.28	-0.10±0.09	5854±100	3.60±0.05	-0.10±0.14	0.10±0.09
33	HD008724	0.018	2	1.001	4586±84	1.39±0.26	-1.73±0.13	4626±6	1.40±0.06	-1.75±0.00	0.37±0.08
47	HD013530	0.013	6	1.036	4772±106	2.60±0.39	-0.54±0.11	4769±87	2.63±0.21	-0.54±0.15	0.37±0.08
66	HD015596	0.047	9	1.211	4808±59	2.66±0.32	-0.65±0.06	4760±58	2.54±0.14	-0.66±0.07	0.40±0.06
67	HD015596	0.037	10	1.298	4808±59	2.66±0.32	-0.65±0.06	4751±52	2.57±0.12	-0.64±0.06	0.37±0.07
68	HD015596	0.047	9	1.248	4808±59	2.66±0.32	-0.65±0.06	4759±56	2.54±0.13	-0.66±0.06	0.40±0.06
69	HD016458	0.118	2	1.115	4593±123	1.84±0.26	-0.35±0.05	4992±513	1.97±0.95	-0.34±0.25	0.33±0.23
88	HD020512	0.086	3	1.118	5212±63	3.65±0.14	-0.22±0.19	5074±38	3.35±0.14	-0.22±0.20	0.12±0.03
89	HD020512	0.094	3	1.101	5212±63	3.65±0.14	-0.22±0.19	5074±39	3.35±0.15	-0.21±0.21	0.12±0.03
117	HD026297	0.133	11	1.263	4445±140	1.02±0.28	-1.74±0.15	4460±73	1.15±0.19	-1.66±0.08	0.48±0.04
151	HD035369	0.032	6	1.062	4885±112	2.57±0.27	-0.21±0.08	4900±45	2.65±0.08	-0.21±0.05	0.06±0.02
227	HD045282	0.023	1	1.000	5264±86	3.19±0.16	-1.43±0.12	5348±110	3.24±0.29	-1.44±0.05	0.22±0.07
228	HD045282	0.065	3	1.042	5264±86	3.19±0.16	-1.43±0.12	5268±48	3.14±0.23	-1.52±0.05	0.36±0.06
253	HD046480	0.016	15	1.218	4785±26	2.63±0.12	-0.49±0.01	4791±55	2.65±0.13	-0.50±0.08	0.31±0.05
254	HD046480	0.016	15	1.214	4785±26	2.63±0.12	-0.49±0.01	4791±55	2.65±0.13	-0.50±0.08	0.31±0.05
314	HD063791	0.040	4	1.028	4715±78	1.75±0.08	-1.68±0.08	4868±275	1.78±0.47	-1.66±0.38	0.44±0.12
384	HD087140	0.033	2	1.038	5129±103	2.66±0.25	-1.80±0.13	5090±5	2.58±0.10	-1.82±0.06	0.42±0.01
425	HD108317	0.039	2	1.034	5259±111	2.68±0.25	-2.27±0.05	5117±40	2.70±0.15	-2.33±0.11	0.45±0.08
452	HD117876	0.075	3	1.091	4747±128	2.27±0.03	-0.48±0.02	4806±18	2.25±0.15	-0.44±0.03	0.42±0.10
454	HD122956	0.023	3	1.018	4633±78	1.46±0.18	-1.72±0.11	4646±16	1.43±0.04	-1.73±0.02	0.42±0.07
459	HD124897	0.136	5	1.023	4302±115	1.66±0.31	-0.52±0.11	4346±98	1.87±0.46	-0.55±0.24	0.29±0.15
470	HD135722	0.011	3	1.016	4795±76	2.60±0.41	-0.40±0.10	4846±8	2.60±0.03	-0.40±0.03	0.14±0.04
473	HD137759	0.142	2	1.014	4549±118	2.88±0.21	0.13±0.11	4558±64	2.54±0.07	0.14±0.02	-0.03±0.11
509	HD159181	0.180	4	1.215	5234±158	1.56±0.19	0.15±0.12	5235±48	1.82±0.31	0.14±0.24	0.04±0.19
566	HD166161	0.059	2	1.040	5210±167	2.25±0.42	-1.22±0.13	5071±122	2.16±0.11	-1.18±0.05	0.31±0.06
568	HD166208	0.161	1	1.000	5037±56	2.71±0.08	0.07±0.11	4919±98	2.52±0.05	0.08±0.04	0.17±0.01
652	HD175305	0.082	1	1.000	5053±140	2.49±0.26	-1.43±0.07	4899±17	2.30±0.03	-1.43±0.05	0.27±0.06
701	HD187111	0.125	4	1.149	4299±75	0.74±0.30	-1.78±0.18	4343±77	0.79±0.21	-1.59±0.09	0.47±0.04
763	HD198149	0.078	2	1.015	4956±177	3.35±0.22	-0.12±0.18	5027±10	3.12±0.06	-0.12±0.06	0.12±0.01
791	HD204543	0.026	1	1.000	4667±68	1.30±0.23	-1.80±0.10	4617±43	1.31±0.08	-1.76±0.10	0.24±0.07
792	HD204613	0.101	2	1.208	5742±135	3.72±0.19	-0.51±0.16	5614±111	3.45±0.09	-0.48±0.08	0.21±0.15
803	HD207130	0.093	3	1.011	4760±53	2.63±0.15	0.01±0.11	4727±16	2.40±0.12	0.01±0.03	0.07±0.01
825	HD216143	0.045	2	1.019	4495±82	1.12±0.38	-2.20±0.06	4480±8	1.15±0.06	-2.12±0.01	0.39±0.05
826	HD216174	0.066	2	1.145	4413±23	2.11±0.36	-0.55±0.02	4381±9	2.21±0.02	-0.53±0.00	0.33±0.04
836	HD218857	0.054	8	1.098	5119±44	2.50±0.37	-1.91±0.09	5067±89	2.37±0.31	-1.93±0.15	0.41±0.04
848	HD221345	0.019	12	1.224	4635±108	2.49±0.32	-0.30±0.07	4666±45	2.50±0.09	-0.30±0.04	0.18±0.07
849	HD221377	0.182	3	1.150	6176±188	3.61±0.17	-0.88±0.17	6027±57	3.24±0.15	-1.01±0.15	0.57±0.12

Table B.2 - continued.

ELODIE	Star	RR _{tot}	N	S _{lim}	T _{eff} (K)	log(<i>g</i>)	[Fe/H]	T _{eff} (K)	logg	[Fe/H]	[Mg/Fe]
871	HD232078	0.101	1	1.000	3939±175	0.31±0.34	-1.58±0.15	3983±186	0.30±0.53	-1.73±0.76	0.27±0.15
878	BD+233130	0.033	1	1.000	5119±140	2.39±0.38	-2.62±0.19	5039±20	2.42±0.16	-2.55±0.01	0.60±0.04
883	BD+302611	0.139	2	1.013	4292±100	0.96±0.37	-1.41±0.19	4421±274	0.83±0.72	-1.43±0.05	0.46±0.13
927	HD000245	0.110	2	1.032	5490±153	3.48±0.15	-0.77±0.08	5377±59	3.67±0.06	-0.84±0.19	0.34±0.08
941	HD003546	0.035	15	1.412	4906±168	2.45±0.43	-0.64±0.12	4868±71	2.51±0.19	-0.65±0.09	0.27±0.03
1395	HD105546	0.047	2	1.018	5234±79	2.38±0.14	-1.39±0.15	5387±190	2.30±0.14	-1.37±0.23	0.54±0.06
1452	HD122563	0.043	1	1.000	4565±131	1.17±0.24	-2.62±0.14	4566±440	1.12±1.31	-2.63±0.37	0.60±0.22
1483	HD136512	0.098	2	1.060	4719±66	2.72±0.04	-0.33±0.16	4747±46	2.62±0.25	-0.30±0.00	0.08±0.03
1484	HD136512	0.110	4	1.072	4719±66	2.72±0.04	-0.33±0.16	4754±29	2.53±0.17	-0.30±0.08	0.14±0.11
1485	HD136512	0.111	4	1.072	4719±66	2.72±0.04	-0.33±0.16	4754±29	2.53±0.17	-0.30±0.08	0.14±0.11
1486	HD136512	0.070	14	1.149	4719±66	2.72±0.04	-0.33±0.16	4824±25	2.54±0.08	-0.33±0.05	0.21±0.06
1487	HD136512	0.080	13	1.144	4719±66	2.72±0.04	-0.33±0.16	4821±27	2.52±0.08	-0.32±0.06	0.21±0.06
1576	HD162211	0.058	4	1.210	4568±74	2.74±0.11	0.04±0.06	4581±72	2.59±0.17	0.04±0.02	0.04±0.07
1811	HD188326	0.172	3	1.092	5272±40	3.80±0.01	-0.18±0.00	5074±39	3.34±0.15	-0.20±0.21	0.12±0.03
1812	HD188326	0.161	3	1.084	5272±40	3.80±0.01	-0.18±0.00	5074±41	3.34±0.16	-0.20±0.21	0.12±0.03
1876	HD212943	0.081	4	1.085	4625±67	2.79±0.05	-0.29±0.09	4656±35	2.61±0.10	-0.30±0.04	0.17±0.07
1893	HD216219	0.093	1	1.000	5628±106	3.12±0.22	-0.41±0.10	5727±152	3.36±0.50	-0.39±0.21	0.06±0.19
1916	HD219449	0.070	16	1.259	4647±75	2.56±0.26	-0.03±0.07	4626±35	2.39±0.07	-0.03±0.03	0.03±0.03

Table B.3 - Atmospheric parameters for all stars analysed in the six clusters: T_{eff} , $\log(g)$, $[\text{Fe}/\text{H}]$, $[\text{Mg}/\text{Fe}]$ and $[\alpha/\text{Fe}]$. Membership identification is copied from Table B.1 to guide the reader.

NGC ID	$T_{eff}^{(a)}$ (K)	$T_{eff}^{(b)}$ (K)	$T_{eff}^{(avg)}$ (K)	$\log(g)^{(a)}$	$\log(g)^{(b)}$	$\log(g)^{(avg)}$	$[\text{Fe}/\text{H}]^{(a)}$	$[\text{Fe}/\text{H}]^{(b)}$	$[\text{Fe}/\text{H}]^{(avg)}$	$[\text{Mg}/\text{Fe}]^{(a)}$	$[\alpha/\text{Fe}]^{(b)}$	members
6528_2	3969±80	4074±114	4004± 66	1.54±0.18	1.9±0.4	1.60±0.17	0.02±0.16	-0.50±0.00	-0.24±0.16	-0.04±0.18	0.20±0.14	
6528_3	3640±100	3525±75	3566± 60	0.70±0.00	2.9±0.8	0.96±0.28	-0.10±0.07	-0.70±0.25	-0.14±0.06	0.23±0.00	0.32±0.07	
6528_4	3900±79	3623±125	3821± 67	1.38±0.19	0.8±0.7	1.34±0.18	-0.07±0.20	-1.35±0.23	-0.62±0.15	0.06±0.20	0.26±0.11	
6528_5	4027±79	4198±151	4064± 70	1.54±0.34	2.0±0.4	1.74±0.26	-0.31±0.22	-0.85±0.23	-0.57±0.16	0.12±0.18	0.31±0.10	
6528_6	4344±215	4625±125	4554±108	2.1±0.5	2.8±0.23	2.71±0.21	0.04±0.17	-0.21±0.30	-0.02±0.15	0.03±0.12	0.18±0.12	
6528_7	3930±58	4124±125	3964± 53	1.51±0.01	2.05±0.35	1.51±0.01	0.10±0.13	-0.25±0.25	0.02±0.12	0.16±0.06	0.30±0.08	
6528_8	4159±162	4225±175	4189±119	1.7±0.4	2.0±0.5	1.88±0.32	-0.08±0.20	-0.75±0.25	-0.34±0.16	0.02±0.19	0.28±0.09	M
6528_9	4244±170	4400±122	4347± 99	1.9±0.4	2.6±0.4	2.28±0.27	-0.06±0.19	-0.70±0.24	-0.31±0.15	0.03±0.17	0.27±0.09	M
6528_10	4557±236	4925±115	4855±103	2.4±0.5	3.30±0.24	3.14±0.22	-0.10±0.25	0.10±0.10	0.07±0.09	0.06±0.17	0.19±0.11	M
6528_11	3911±70	3800±187	3897± 66	1.43±0.18	1.3±0.7	1.42±0.17	-0.04±0.21	-1.15±0.32	-0.37±0.18	0.06±0.17	0.29±0.11	M
6528_13	4853±198	4876±167	4866±128	2.5±0.4	2.6±0.5	2.53±0.32	-0.44±0.22	-0.80±0.25	-0.60±0.17	0.24±0.20	0.22±0.11	
6528_14	4239±172	4397±164	4322±119	1.9±0.5	2.2±0.4	2.07±0.30	-0.13±0.21	-0.71±0.33	-0.30±0.18	0.03±0.11	0.28±0.12	
6528_15	4083±176	4325±114	4253± 96	1.61±0.19	2.30±0.25	1.86±0.15	0.05±0.16	-0.35±0.23	-0.08±0.13	0.05±0.11	0.17±0.13	
6528_16	3938±40	4075±114	3953± 38	1.48±0.12	1.95±0.35	1.53±0.11	-0.04±0.15	-0.80±0.24	-0.25±0.13	0.05±0.10	0.29±0.09	
6528_17	4500±147	4700±186	4577±115	2.35±0.27	2.70±0.33	2.49±0.21	-0.37±0.13	-0.6±0.4	-0.39±0.12	0.17±0.16	0.31±0.10	
6528_18	4709±194	4676±195	4692±138	2.5±0.4	2.5±0.5	2.52±0.29	-0.22±0.25	-0.6±0.4	-0.35±0.21	0.12±0.18	0.16±0.12	
6528_19	4496±94	4701±99	4593± 68	2.53±0.30	2.90±0.20	2.79±0.17	0.13±0.09	-0.02±0.25	0.11±0.08	0.04±0.08	0.17±0.13	
6553_1	4141±107	4425±195	4207± 94	1.58±0.14	2.2±0.5	1.62±0.14	-0.08±0.09	-0.40±0.20	-0.14±0.08	0.08±0.06	0.32±0.07	M
6553_3	4384±155	4475±174	4424±116	1.9±0.4	2.2±0.5	1.98±0.31	-0.06±0.08	-0.40±0.20	-0.10±0.07	0.12±0.12	0.29±0.09	M
6553_4	3837±100	3574±114	3723± 75	1.17±0.18	1.1±0.7	1.17±0.17	0.10±0.17	-1.35±0.23	-0.41±0.14	0.09±0.07	0.34±0.07	M
6553_5	3970±71	4076±160	3987± 65	1.65±0.10	1.6±0.6	1.65±0.10	0.06±0.13	-0.75±0.33	-0.04±0.12	0.05±0.11	0.34±0.07	M
6553_6	4312±135	4400±122	4360± 91	1.9±0.4	1.9±0.4	1.90±0.28	0.004±0.09	-0.50±0.00	-0.13±0.08	-0.01±0.17	0.26±0.11	M
6553_7	3730±60	3550±100	3682± 51	0.90±0.13	1.2±0.9	0.91±0.13	0.27±0.18	-0.95±0.27	-0.10±0.15	0.08±0.10	0.28±0.10	M
6553_8	3640±0	3500±0	3528± 45	0.70±0.00	2.8±0.8	0.98±0.28	-0.10±0.06	-0.70±0.24	-0.14±0.06	0.23±0.10	0.32±0.07	
6553_9	4259±15	4399±165	4260± 15	1.47±0.08	2.1±0.5	1.49±0.08	-0.13±0.00	-0.45±0.15	-0.13±0.01	0.11±0.10	0.28±0.10	M
6553_10	4388±155	4525±174	4449±116	1.9±0.4	2.3±0.5	2.02±0.31	-0.06±0.08	-0.35±0.23	-0.09±0.07	0.13±0.12	0.29±0.09	M

Table B.3 - continued.

NGC ID	$T_{eff}^{(a)}$ (K)	$T_{eff}^{(b)}$ (K)	$T_{eff}^{(avg)}$ (K)	$\log(g)^{(a)}$	$\log(g)^{(b)}$	$\log(g)^{(avg)}$	[Fe/H] ^(a)	[Fe/H] ^(b)	[Fe/H] ^(avg)	[Mg/Fe] ^(a)	[α /Fe] ^(b)	members
6553_11	3941±44	3801±218	3936± 43	1.47±0.19	1.0±0.8	1.45±0.19	0.01±0.14	-0.9±0.4	-0.07±0.13	0.07±0.09	0.27±0.09	M
6553_13	4219±57	4350±122	4242± 52	1.63±0.26	2.0±0.4	1.74±0.22	-0.01±0.13	-0.50±0.00	-0.22±0.10	0.02±0.10	0.25±0.10	M
6553_14	3889±64	3896±301	3889± 63	1.38±0.16	1.3±1.0	1.38±0.16	-0.01±0.17	-0.95±0.42	-0.15±0.16	0.004±0.20	0.33±0.10	M
6553_15	3640±0	3500±0	3528± 45	0.70±0.00	3.1±0.8	1.01±0.28	-0.10±0.06	-0.60±0.20	-0.15±0.06	0.23±0.10	0.32±0.07	
6553_16	3932±29	4101±123	3941± 28	1.54±0.13	1.9±0.4	1.58±0.12	-0.20±0.05	-0.70±0.24	-0.22±0.05	0.05±0.09	0.33±0.06	M
6553_17	3640±100	3548±99	3594± 70	0.70±0.00	2.4±0.7	1.15±0.34	-0.05±0.05	-0.90±0.20	-0.10±0.05	0.23±0.10	0.35±0.07	
6553_18	3640±100	3500±0.0	3528± 45	0.70±0.00	3.4±0.5	1.78±0.31	-0.10±0.07	0.20±0.17	-0.06±0.06	0.23±0.10	0.33±0.06	
6553_19	4259±15	4176±114	4258± 15	1.47±0.08	1.75±0.33	1.49±0.08	-0.13±0.00	-0.65±0.23	-0.13±0.01	0.11±0.10	0.27±0.09	M
M71_2	4743±210	4850±122	4823±106	2.5±0.5	2.95±0.27	2.85±0.24	-0.48±0.24	-0.85±0.23	-0.67±0.17	0.25±0.20	0.28±0.13	M
M71_4	4305±77	4449±187	4326± 71	1.8±0.5	2.2±0.6	1.99±0.36	-0.56±0.24	-0.85±0.23	-0.71±0.17	0.32±0.18	0.28±0.10	M
M71_5	4722±223	4899±198	4821±148	2.4±0.6	3.10±0.30	2.97±0.27	-0.51±0.26	-0.85±0.32	-0.64±0.20	0.27±0.20	0.29±0.09	M
M71_6	4364±94	4523±174	4400± 83	1.9±0.4	2.2±0.6	2.02±0.32	-0.55±0.22	-0.95±0.35	-0.66±0.19	0.26±0.16	0.30±0.08	M
M71_7	3991±99	4026±207	3997± 89	1.5±0.4	1.7±0.7	1.53±0.35	-0.32±0.21	-1.25±0.33	-0.58±0.17	0.15±0.18	0.34±0.07	M
M71_8	4409±132	4850±122	4646± 90	2.1±0.5	3.15±0.23	2.95±0.21	0.16±0.13	0.26±0.21	0.18±0.11	0.01±0.14	0.13±0.10	
M71_9	4303±102	4350±166	4316± 87	1.9±0.5	2.0±0.5	1.97±0.33	-0.54±0.23	-1.05±0.27	-0.76±0.17	0.27±0.21	0.28±0.10	M
M71_10	3861±77	3549±99	3743± 61	1.23±0.21	0.5±0.6	1.16±0.20	-0.06±0.26	-1.70±0.24	-0.96±0.18	0.11±0.07	0.29±0.14	M
M71_13	4685±209	4850±122	4808±106	2.4±0.5	2.95±0.27	2.82±0.24	-0.44±0.26	-0.80±0.25	-0.63±0.18	0.23±0.20	0.25±0.10	M
M71_14	4901±212	5175±114	5113±101	2.7±0.5	3.15±0.32	3.01±0.27	-0.44±0.25	-0.65±0.23	-0.55±0.17	0.25±0.20	0.27±0.11	M
M71_15	4840±259	5101±165	5025±139	2.5±0.6	3.10±0.30	2.98±0.27	-0.50±0.30	-0.55±0.15	-0.54±0.13	0.26±0.19	0.27±0.11	M
M71_16	4694±214	4974±175	4862±135	2.5±0.4	2.95±0.35	2.76±0.27	-0.13±0.25	-0.11±0.27	-0.12±0.18	0.08±0.18	0.20±0.13	
6558_3	4758±228	4926±160	4871±131	2.5±0.5	3.10±0.30	2.94±0.25	-0.29±0.26	-0.70±0.24	-0.51±0.18	0.15±0.20	0.30±0.10	
6558_4	4678±172	4625±167	4651±120	2.55±0.33	3.0±0.4	2.75±0.26	-0.21±0.19	-1.30±0.25	-0.61±0.15	0.09±0.14	0.29±0.09	
6558_5	4921±445	4900±200	4903±182	2.0±0.6	2.00±0.32	2.00±0.29	-1.2±0.6	-1.80±0.24	-1.72±0.22	0.37±0.19	0.21±0.11	
6558_6	4738±329	4951±186	4899±162	2.1±0.8	2.8±0.4	2.68±0.39	-0.9±0.4	-1.20±0.24	-1.11±0.20	0.22±0.07	0.26±0.13	M
6558_7	4928±317	4926±114	4926±108	2.5±0.7	2.70±0.25	2.68±0.24	-0.64±0.32	-1.15±0.23	-0.98±0.19	0.26±0.13	0.26±0.13	M
6558_8	3640±100	3524±74	3565± 59	0.70±0.5	2.8±0.9	1.05±0.36	0.00±0.07	-0.95±0.15	-0.16±0.06	0.23±0.00	0.31±0.08	M
6558_9	4663±336	5076±195	4972±168	1.8±0.9	2.9±0.5	2.68±0.46	-1.2±0.4	-1.10±0.20	-1.11±0.18	0.41±0.16	0.21±0.11	M
6558_10	4398±157	4799±100	4684± 84	2.1±0.5	3.15±0.23	2.95±0.21	0.10±0.13	0.16±0.19	0.12±0.11	0.02±0.13	0.12±0.12	

Table B.3 - continued.

NGC ID	$T_{eff}^{(a)}$ (K)	$T_{eff}^{(b)}$ (K)	$T_{eff}^{(avg)}$ (K)	$\log(g)^{(a)}$	$\log(g)^{(b)}$	$\log(g)^{(avg)}$	[Fe/H] ^(a)	[Fe/H] ^(b)	[Fe/H] ^(avg)	[Mg/Fe] ^(a)	[α /Fe] ^(b)	members
6558_11	4887±413	5325±114	5294±110	2.2±0.9	3.30±0.25	3.22±0.24	-1.1±0.5	-1.00±0.00	-1.00±0.05	0.37±0.19	0.19±0.14	M
6558_12	4516±177	4548±245	4527±143	2.3±0.5	2.2±0.6	2.26±0.37	-0.18±0.27	-0.58±0.34	-0.33±0.21	0.10±0.18	0.25±0.10	
6558_13	4683±94	5076±115	4840± 73	2.49±0.14	3.25±0.25	2.67±0.12	-0.22±0.09	-0.25±0.25	-0.22±0.08	0.09±0.15	0.31±0.08	
6558_14	4643±454	4675±115	4673±111	3.0±1.0	4.4±0.5	4.12±0.44	0.30±0.13	-0.25±0.25	0.18±0.12	0.18±0.07	0.12±0.07	
6558_15	4885±409	5500±0	5491± 50	1.9±0.8	3.0±0.4	2.79±0.35	-1.40±0.25	-1.00±0.00	-1.02±0.05	0.44±0.15	0.20±0.13	
6558_16	4690±186	4825±195	4754±135	2.50±0.35	2.9±0.5	2.64±0.28	-0.20±0.22	-0.50±0.32	-0.29±0.18	0.10±0.16	0.23±0.13	
6558_17	4829±192	4800±187	4814±134	2.7±0.6	3.30±0.33	3.14±0.28	-0.56±0.26	-1.4±0.4	-0.84±0.21	0.33±0.18	0.36±0.05	
6558_18	4888±241	5151±122	5097±109	2.6±0.4	3.15±0.23	3.04±0.20	-0.36±0.26	-0.50±0.00	-0.50±0.05	0.20±0.22	0.26±0.11	
6558_19	4912±435	5175±160	5144±150	2.1±0.9	3.20±0.25	3.12±0.24	-1.5±0.4	-1.45±0.15	-1.46±0.14	0.46±0.15	0.22±0.12	
6426_1	4849±25	5026±175	4853± 25	3.45±0.04	4.5±0.4	3.46±0.04	0.09±0.14	-0.75±0.25	-0.11±0.12	-0.02±0.01	0.34±0.07	
6426_2	4643±51	5075±115	4714± 47	3.05±0.21	4.7±0.4	3.41±0.19	0.30±0.05	-0.11±0.27	0.28±0.05	0.18±0.10	0.14±0.09	
6426_3	4250±163	4625±125	4486± 99	3.50±0.34	3.95±0.35	3.72±0.24	0.10±0.06	-0.60±0.20	0.05±0.05	-0.02±0.10	0.17±0.12	
6426_4	4881±350	4900±122	4898±116	2.0±0.8	1.9±0.4	1.92±0.39	-1.93±0.35	-2.5±0.1	-2.46±0.10	0.37±0.12	0.28±0.10	M
6426_7	5008±435	5250±10	5250± 10	2.1±0.8	2.1±0.4	2.11±0.34	-1.9±0.4	-2.5±0.1	-2.47±0.10	0.41±0.17	0.23±0.13	M
6426_9	5071±374	5250±10	5250± 10	2.3±0.9	2.30±0.33	2.30±0.31	-2.1±0.4	-2.5±0.1	-2.47±0.10	0.38±0.19	0.20±0.14	M
6426_10	4487±70	4951±149	4571± 63	1.15±0.12	2.1±0.4	1.22±0.12	-2.06±0.14	-2.05±0.15	-2.05±0.10	0.39±0.11	0.21±0.14	M
6426_11	4967±156	5475±207	5150±125	3.1±0.4	4.65±0.32	4.13±0.26	-0.44±0.32	-0.70±0.24	-0.61±0.19	0.25±0.16	0.30±0.08	
6426_13	5015±331	5000±10	5000± 10	2.2±0.8	2.0±0.4	2.08±0.37	-2.0±0.4	-2.5±0.1	-2.47±0.10	0.37±0.19	0.25±0.10	M
6426_18	4824±52	4875±125	4831± 48	3.24±0.23	3.5±0.4	3.31±0.20	0.01±0.07	-0.5±0.1	-0.15±0.06	0.05±0.07	0.24±0.12	
Ter8_1	4885±314	5250±0	5067±314	2.0±0.7	2.30±0.33	2.24±0.30	-1.82±0.34	-2.00±0.00	-1.97±0.14	0.40±0.13	0.20±0.14	M
Ter8_4	4565±193	4299±99	4354± 88	1.3±0.5	0.4±0.4	0.67±0.31	-1.78±0.23	-2.50±0.00	-2.28±0.13	0.40±0.14	0.20±0.14	M
Ter8_5	4952±326	5000±0	4976±326	2.1±0.8	1.9±0.4	1.98±0.37	-1.9±0.4	-2.50±0.00	-2.42±0.14	0.40±0.17	0.25±0.10	M
Ter8_6	4956±337	5250±0	4978±337	2.1±0.8	2.6±0.4	2.50±0.39	-1.9±0.4	-2.00±0.00	-1.98±0.14	0.40±0.17	0.15±0.13	M
Ter8_8	4947±333	5175±114	5151±108	2.1±0.8	2.7±0.4	2.56±0.36	-1.9±0.4	-2.15±0.23	-2.06±0.19	0.40±0.18	0.19±0.15	M
Ter8_9	4694±293	4549±99	4564± 94	1.6±0.5	0.9±0.4	1.12±0.30	-1.80±0.33	-2.40±0.20	-2.24±0.17	0.42±0.13	0.19±0.14	M
Ter8_10	4849±195	4825±114	4831± 99	3.5±0.5	4.30±0.25	4.13±0.22	0.09±0.09	-0.65±0.23	-0.01±0.08	-0.02±0.03	0.23±0.13	
Ter8_11	4655±247	4951±100	4909± 92	1.5±0.5	2.05±0.35	1.84±0.28	-1.90±0.25	-2.00±0.00	-1.97±0.13	0.45±0.12	0.16±0.14	M
Ter8_13	4967±367	5250±0	5108±367	2.1±0.8	2.4±0.4	2.34±0.34	-1.8±0.4	-2.00±0.00	-1.98±0.14	0.41±0.17	0.22±0.12	M

Table B.3 - continued.

NGC ID	$T_{eff}^{(a)}$ (K)	$T_{eff}^{(b)}$ (K)	$T_{eff}^{(avg)}$ (K)	$\log(g)^{(a)}$	$\log(g)^{(b)}$	$\log(g)^{(avg)}$	[Fe/H] ^(a)	[Fe/H] ^(b)	[Fe/H] ^(avg)	[Mg/Fe] ^(a)	[α /Fe] ^(b)	members
Ter8_14	4756±142	4324±114	4493± 89	1.8±0.4	0.4±0.4	1.16±0.28	-1.68±0.11	-2.50±0.00	-1.98±0.09	0.43±0.14	0.19±0.14	M
Ter8_15	4984±344	5225±75	5214± 73	2.1±0.8	2.20±0.25	2.19±0.24	-1.8±0.4	-2.00±0.00	-1.97±0.14	0.39±0.17	0.19±0.15	M
Ter8_16	4871±333	5050±100	5035± 96	1.9±0.6	2.20±0.33	2.14±0.29	-1.75±0.34	-2.00±0.00	-1.96±0.14	0.42±0.12	0.21±0.14	M
Ter8_18	4722±195	4700±245	4713±153	1.7±0.5	1.6±0.5	1.68±0.38	-1.74±0.30	-2.30±0.24	-2.08±0.19	0.37±0.13	0.27±0.09	M
HP1_8	4682±264	4375±125	4431±113	2.3±0.6	1.60±0.37	1.79±0.31	-0.4±0.4	-1.25±0.25	-1.01±0.21	0.17±0.23	0.28±0.12	M
HP1_10	4467±176	4575±159	4526±118	1.7±0.7	1.95±0.35	1.90±0.31	-0.8±0.4	-1.30±0.25	-1.16±0.21	0.22±0.25	0.27±0.09	M
HP1_11	4353±185	4125±230	4263±144	1.7±0.6	0.85±0.59	1.27±0.42	-0.67±0.31	-1.65±0.32	-1.14±0.22	0.29±0.23	0.28±0.13	M
HP1_12	4823±340	4448±149	4508±136	2.4±0.8	1.75±0.33	1.84±0.31	-0.6±0.4	-1.65±0.23	-1.39±0.20	0.28±0.21	0.28±0.10	M
HP1_13	4232±49	4026±174	4217± 47	1.37±0.33	0.60±0.62	1.20±0.29	-0.79±0.21	-1.45±0.35	-0.96±0.18	0.48±0.15	0.30±0.10	M
HP1_14	4517±348	4175±114	4208±108	1.6±0.9	0.40±0.44	0.63±0.40	-1.0±0.5	-2.00±0.1	-1.96±0.10	0.34±0.18	0.27±0.09	
HP1_15	4868±451	4600±200	4644±183	2.0±0.9	1.40±0.73	1.64±0.57	-1.3±0.5	-1.90±0.20	-1.82±0.19	0.41±0.16	0.11±0.09	
HP1_19	4823±367	4927±297	4886±231	2.0±0.9	2.55±0.61	2.38±0.50	-1.2±0.5	-1.50±0.39	-1.39±0.31	0.40±0.15	0.31±0.10	
HP1_20	4681±267	4325±115	4381±106	2.4±0.6	1.85±0.23	1.92±0.21	-0.4±0.4	-1.35±0.23	-1.11±0.20	0.19±0.21	0.21±0.11	
HP1_21	4607±241	4800±187	4727±148	2.4±0.5	2.70±0.40	2.58±0.31	-0.14±0.27	-0.26±0.30	-0.19±0.20	0.08±0.19	0.33±0.06	
HP1_22	4758±323	4498±192	4566±165	2.2±0.8	1.85±0.45	1.93±0.39	-0.8±0.4	-1.55±0.27	-1.32±0.22	0.35±0.17	0.28±0.10	M
HP1_23	4420±200	4377±125	4389±106	1.9±0.6	1.55±0.35	1.64±0.30	-0.62±0.33	-1.25±0.25	-1.02±0.20	0.27±0.22	0.29±0.09	M
HP1_24	4561±301	4099±122	4164±113	1.9±0.7	0.79±0.60	1.26±0.46	-0.71±0.31	-1.80±0.24	-1.39±0.19	0.36±0.18	0.25±0.10	M
HP1_31	4379±186	4750±100	4667± 88	2.0±0.4	2.40±0.20	2.32±0.18	0.04±0.19	0.08±0.10	0.07±0.09	0.02±0.16	0.24±0.14	
HP1_32	4836±388	4850±166	4848±153	2.1±0.9	3.00±0.45	2.82±0.40	-1.2±0.5	-1.75±0.25	-1.64±0.22	0.41±0.16	0.31±0.08	
HP1_33	4799±262	5199±244	5013±179	2.5±0.5	3.25±0.33	3.02±0.28	-0.37±0.33	-0.40±0.30	-0.39±0.22	0.19±0.22	0.29±0.10	
HP1_34	4373±92	4224±231	4353± 85	1.9±0.4	1.16±0.70	1.72±0.35	-0.53±0.24	-1.10±0.37	-0.70±0.20	0.28±0.20	0.28±0.14	
HP1_35	4716±245	4675±114	4682±103	2.4±0.5	2.15±0.32	2.22±0.27	-0.27±0.33	-0.65±0.23	-0.53±0.19	0.15±0.22	0.26±0.13	
HP1_36	4397±153	4675±114	4576± 91	2.1±0.4	2.35±0.23	2.29±0.20	0.08±0.14	0.08±0.10	0.08±0.08	0.01±0.14	0.18±0.12	
HP1_37	4844±327	4875±125	4871±117	2.5±0.6	2.65±0.39	2.61±0.33	-0.4±0.4	-0.75±0.25	-0.65±0.21	0.20±0.23	0.24±0.14	
HP1_38	4875±308	5251±224	5121±181	2.5±0.6	3.05±0.15	3.02±0.15	-0.37±0.33	-0.30±0.25	-0.33±0.20	0.18±0.22	0.26±0.11	
HP1_40	4490±239	4499±250	4494±173	2.2±0.5	2.10±0.62	2.16±0.39	-0.08±0.26	-0.53±0.44	-0.20±0.22	0.06±0.17	0.23±0.11	
HP1_41	4356±211	4649±123	4575±106	2.1±0.5	2.85±0.32	2.63±0.27	0.05±0.16	-0.12±0.32	0.02±0.14	0.02±0.12	0.17±0.13	
HP1_42	4418±125	4723±175	4521±102	2.2±0.5	3.10±0.44	2.71±0.33	0.15±0.13	0.20±0.17	0.17±0.10	0.03±0.15	0.12±0.07	

Table B.3 - continued.

NGC ID	$T_{eff}^{(a)}$ (K)	$T_{eff}^{(b)}$ (K)	$T_{eff}^{(avg)}$ (K)	$\log(g)^{(a)}$	$\log(g)^{(b)}$	$\log(g)^{(avg)}$	$[\text{Fe}/\text{H}]^{(a)}$	$[\text{Fe}/\text{H}]^{(b)}$	$[\text{Fe}/\text{H}]^{(avg)}$	$[\text{Mg}/\text{Fe}]^{(a)}$	$[\alpha/\text{Fe}]^{(b)}$	members
HP1_43	4402±157	4524±207	4447±125	2.2±0.5	2.40±0.62	2.28±0.39	0.04±0.13	-0.28±0.27	-0.02±0.12	0.04±0.14	0.27±0.10	
HP1_44	4449±224	4775±175	4651±138	2.2±0.5	3.65±0.39	3.10±0.31	-0.09±0.26	-0.20±0.24	-0.15±0.18	0.06±0.15	0.12±0.10	
HP1_45	4386±140	4479±209	4415±116	2.2±0.4	2.36±0.55	2.26±0.32	0.13±0.13	-0.13±0.30	0.09±0.12	0.02±0.15	0.12±0.10	
Djorg2_1	3946±59	4250±193	3972± 56	1.51±0.19	2.4±0.6	1.59±0.18	0.02±0.18	-0.35±0.23	-0.12±0.14	-0.01±0.23	0.18±0.10	
Djorg2_3	4498±110	4750±224	4547± 99	2.2±0.4	2.6±0.5	2.35±0.30	0.03±0.15	-0.12±0.32	0.01±0.14	0.03±0.17	0.21±0.13	
Djorg2_4	4116±165	4149±123	4137± 98	1.6±0.4	1.9±0.4	1.78±0.26	-0.09±0.18	-0.70±0.25	-0.30±0.15	0.03±0.18	0.22±0.12	
Djorg2_5	4384±137	4400±165	4390±105	2.1±0.4	2.1±0.4	2.11±0.28	-0.11±0.20	-0.70±0.33	-0.27±0.17	0.09±0.17	0.33±0.08	
Djorg2_6	4556±195	4524±236	4543±150	2.3±0.5	2.0±0.5	2.19±0.33	-0.16±0.27	-0.60±0.30	-0.35±0.20	0.08±0.18	0.28±0.11	
Djorg2_7	4577±199	4475±134	4507±111	2.4±0.5	1.8±0.4	2.04±0.32	-0.15±0.27	-1.55±0.27	-0.85±0.19	0.10±0.20	0.27±0.09	M
Djorg2_8	4559±195	4600±122	4588±104	2.3±0.6	1.75±0.25	1.83±0.23	-0.3 ±0.4	-1.30±0.24	-1.00±0.20	0.17±0.23	0.29±0.09	
Djorg2_9	4515±170	4376±166	4444±119	2.1±0.5	1.6±0.4	1.85±0.31	-0.53±0.19	-1.25±0.33	-0.71±0.16	0.26±0.21	0.27±0.11	M
Djorg2_10	4250±33	4127±200	4247± 33	1.4±0.4	1.0±0.7	1.28±0.32	-0.77±0.23	-1.29±0.33	-0.94±0.19	0.46±0.17	0.27±0.13	M
Djorg2_11	4190±153	4351±123	4288± 96	1.7±0.4	1.95±0.35	1.85±0.26	-0.22±0.25	-0.70±0.24	-0.47±0.17	0.09±0.19	0.32±0.07	
Djorg2_13	4549±191	4625±167	4592±126	2.3±0.5	2.4±0.6	2.32±0.37	-0.29±0.28	-0.70±0.24	-0.53±0.18	0.15±0.20	0.29±0.09	
Djorg2_14	4256±218	4350±122	4328±107	2.0±0.5	2.30±0.25	2.23±0.22	0.17±0.17	-0.30±0.25	0.02±0.14	0.00±0.13	0.15±0.14	
Djorg2_15	4745±240	4700±218	4721±161	2.4±0.5	2.2±0.6	2.35±0.40	-0.42±0.29	-0.85±0.23	-0.68±0.18	0.21±0.20	0.28±0.07	M
Djorg2_16	4230±180	4251±158	4242±119	1.8±0.4	1.9±0.5	1.89±0.33	-0.15±0.21	-1.00±0.32	-0.41±0.18	0.08±0.16	0.34±0.07	
Djorg2_17	4763±371	4624±125	4638±118	2.2±0.9	1.8±0.4	1.90±0.36	-0.9 ±0.5	-1.50±0.10	-1.48±0.10	0.34±0.18	0.26±0.14	
M22a_1	4551±56	4926±114	4624± 50	2.39±0.19	2.90±0.20	2.63±0.14	-0.02±0.09	-0.07±0.29	-0.02±0.09	0.05±0.09	0.27±0.11	
M22a_2	4395±124	4725±175	4505±101	1.0 ±0.4	2.10±0.44	1.50±0.30	-1.64±0.25	-1.90±0.20	-1.80±0.16	0.47±0.08	0.20±0.08	
M22a_3	4491±136	4798±147	4633±100	1.18±0.18	1.85±0.32	1.34±0.16	-1.87±0.20	-1.95±0.15	-1.92±0.12	0.45±0.11	0.18±0.12	M
M22a_4	4430±32	4849±122	4457± 31	1.03±0.01	1.85±0.32	1.03±0.01	-2.10±0.1	-2.00±0.1	-2.05±0.07	0.51±0.01	0.21±0.14	M
M22a_5	4892±339	5200±100	5175± 96	1.9 ±0.6	2.70±0.25	2.58±0.23	-1.58±0.32	-1.60±0.20	-1.59±0.17	0.41±0.14	0.26±0.13	M
M22a_6	4739±414	5100±122	5071±117	1.4 ±0.4	2.40±0.30	2.04±0.24	-1.78±0.34	-2.00±0.1	-1.98±0.10	0.48±0.12	0.15±0.12	M
M22a_7	4668±283	4824±159	4787±139	1.5 ±0.5	1.95±0.35	1.80±0.29	-1.71±0.27	-1.95±0.15	-1.89±0.13	0.46±0.10	0.23±0.13	
M22a_8	4659±299	4950±100	4921± 95	1.5 ±0.5	2.25±0.33	2.02±0.28	-1.57±0.31	-1.60±0.20	-1.59±0.17	0.45±0.14	0.22±0.13	M
M22a_9	4967±335	5150±166	5114±149	2.1 ±0.7	2.75±0.51	2.52±0.41	-1.62±0.31	-1.95±0.15	-1.89±0.14	0.42±0.13	0.23±0.09	M
M22a_10	4773±277	4848±198	4823±161	1.7 ±0.5	1.95±0.35	1.87±0.29	-1.59±0.34	-1.90±0.20	-1.82±0.17	0.43±0.15	0.21±0.11	M

Table B.3 - continued.

NGC ID	$T_{eff}^{(a)}$ (K)	$T_{eff}^{(b)}$ (K)	$T_{eff}^{(avg)}$ (K)	$\log(g)^{(a)}$	$\log(g)^{(b)}$	$\log(g)^{(avg)}$	[Fe/H] ^(a)	[Fe/H] ^(b)	[Fe/H] ^(avg)	[Mg/Fe] ^(a)	[α /Fe] ^(b)	members
M22a_11	5023±383	5174±160	5152±148	2.3 ±0.8	3.15±0.32	3.03±0.30	-1.6 ±0.4	-1.95±0.15	-1.91±0.14	0.40±0.18	0.27±0.13	M
M22a_13	5126±427	5550±100	5528± 97	2.3 ±0.8	4.00±0.32	3.77±0.30	-1.7 ±0.5	-1.90±0.20	-1.87±0.19	0.42±0.17	0.17±0.16	M
M22a_14	5091±269	5325±115	5289±106	2.4 ±0.8	2.65±0.45	2.59±0.39	-2.0 ±0.4	-2.35±0.23	-2.26±0.20	0.37±0.20	0.22±0.12	M
M22a_15	4939±325	5200±187	5135±162	2.1 ±0.7	2.80±0.46	2.59±0.38	-1.64±0.29	-1.90±0.20	-1.82±0.16	0.42±0.12	0.31±0.09	
M22a_16	4955±324	5175±195	5117±167	2.1 ±0.7	2.60±0.54	2.41±0.43	-1.61±0.31	-1.85±0.23	-1.76±0.18	0.42±0.13	0.18±0.12	M
M22a_17	4934±341	5250±100	5225± 96	2.1 ±0.8	2.95±0.42	2.77±0.37	-1.75±0.33	-2.00±0.1	-1.98±0.10	0.41±0.13	0.20±0.13	
M22a_18	4981±336	5250±100	5228± 96	2.2 ±0.8	2.75±0.51	2.59±0.43	-1.73±0.34	-2.00±0.1	-1.98±0.10	0.41±0.13	0.25±0.10	M
M22a_19	4958±339	5250±100	5227± 96	2.1 ±0.8	2.60±0.37	2.51±0.34	-1.8 ±0.4	-2.00±0.1	-1.99±0.10	0.40±0.16	0.24±0.14	M
M22b_2	4927±322	5075±114	5059±107	2.0 ±0.6	2.20±0.46	2.13±0.37	-1.60±0.32	-1.95±0.15	-1.89±0.14	0.40±0.13	0.24±0.14	M
M22b_3	4516±263	4975± 75	4940± 72	1.2 ±0.5	2.50±0.39	2.01±0.31	-1.56±0.26	-1.55±0.15	-1.55±0.13	0.48±0.11	0.26±0.13	M
M22b_4	4767±396	5201± 99	5175± 96	1.5 ±0.5	2.20±0.25	2.06±0.22	-1.7 ±0.4	-2.00±0.1	-1.98±0.10	0.41±0.16	0.20±0.13	M
M22b_5	4926±332	5075±115	5059±109	2.0 ±0.6	2.25±0.25	2.21±0.23	-1.65±0.30	-2.00±0.1	-1.97±0.09	0.40±0.13	0.25±0.10	M
M22b_6	4958±341	5100±229	5056±190	2.1 ±0.8	2.05±0.42	2.06±0.37	-1.8 ±0.4	-2.15±0.23	-2.06±0.20	0.38±0.19	0.23±0.13	M
M22b_7	4790±350	5375±125	5309±118	1.7 ±0.4	2.80±0.33	2.35±0.25	-1.5 ±0.4	-1.25±0.25	-1.32±0.21	0.46±0.13	0.26±0.13	M
M22b_8	4710±287	4925±114	4896±106	1.5 ±0.5	2.00±0.45	1.78±0.33	-1.7 ±0.4	-2.00±0.1	-1.98±0.10	0.40±0.13	0.25±0.10	M
M22b_9	4921±217	5275±208	5105±150	3.35±0.19	3.85±0.23	3.55±0.15	-0.00±0.10	-0.30±0.24	-0.05±0.09	0.09±0.11	0.17±0.11	
M22b_10	4694±416	4721±208	4716±186	1.4 ±0.6	1.89±0.37	1.76±0.31	-1.6 ±0.4	-1.85±0.23	-1.79±0.20	0.46±0.14	0.23±0.13	M
M22b_11	4915±560	4724±208	4747±195	1.6 ±0.6	1.75±0.25	1.73±0.23	-1.7 ±0.4	-2.35±0.23	-2.19±0.20	0.55±0.03	0.27±0.13	M
M22b_12	5089±432	5250±100	5242± 97	2.3 ±0.8	2.40±0.37	2.38±0.34	-1.7 ±0.4	-2.00±0.1	-1.98±0.10	0.41±0.18	0.22±0.12	M
M22b_13	5024±463	5200±100	5192± 98	2.2 ±0.9	3.10±0.37	2.97±0.34	-1.7 ±0.5	-2.00±0.1	-1.99±0.10	0.43±0.17	0.25±0.10	M
M22b_15	4785±347	5125±168	5060±151	1.7 ±0.5	2.90±0.44	2.38±0.33	-1.7 ±0.4	-2.05±0.15	-2.01±0.14	0.43±0.15	0.24±0.12	M
M22b_16	4768±271	4799±148	4792±130	1.7 ±0.5	1.85±0.32	1.81±0.27	-1.6 ±0.4	-1.95±0.15	-1.91±0.14	0.38±0.13	0.19±0.14	M
M22b_17	4791±353	5250±194	5144±170	1.7 ±0.5	2.70±0.51	2.19±0.36	-1.7 ±0.4	-1.70±0.24	-1.70±0.21	0.41±0.17	0.23±0.13	M
M22b_18	4799±409	5250±100	5225± 97	1.6 ±0.6	2.90±0.37	2.54±0.31	-1.86±0.34	-2.00±0.1	-1.99±0.10	0.48±0.11	0.16±0.14	M
M22b_19	4804±267	5250±100	5195± 94	1.7 ±0.5	2.90±0.37	2.48±0.30	-1.62±0.33	-1.50±0.1	-1.51±0.10	0.44±0.13	0.19±0.14	M
M22c_1	5089±407	5500±100	5477± 97	2.3 ±0.9	3.00±0.39	2.89±0.36	-1.9 ±0.4	-2.00±0.1	-1.99±0.10	0.41±0.17	0.14±0.13	M
M22c_2	4919±297	5075±114	5055±106	2.1 ±0.7	2.50±0.39	2.41±0.34	-1.67±0.25	-1.90±0.20	-1.81±0.16	0.41±0.13	0.20±0.12	
M22c_3	5027±313	5300±100	5275± 95	2.3 ±0.7	2.65±0.39	2.57±0.34	-1.67±0.29	-1.90±0.20	-1.83±0.16	0.42±0.13	0.25±0.10	M

Table B.3 - continued.

NGC ID	$T_{eff}^{(a)}$ (K)	$T_{eff}^{(b)}$ (K)	$T_{eff}^{(avg)}$ (K)	$\log(g)^{(a)}$	$\log(g)^{(b)}$	$\log(g)^{(avg)}$	$[\text{Fe}/\text{H}]^{(a)}$	$[\text{Fe}/\text{H}]^{(b)}$	$[\text{Fe}/\text{H}]^{(avg)}$	$[\text{Mg}/\text{Fe}]^{(a)}$	$[\alpha/\text{Fe}]^{(b)}$	members
M22c_4	4943±325	5175±195	5114±167	2.1 ±0.7	2.35±0.59	2.25±0.45	-1.64±0.29	-1.80±0.24	-1.73±0.18	0.41±0.12	0.27±0.13	M
M22c_5	4987±312	5250±100	5225± 95	2.2 ±0.7	2.70±0.33	2.61±0.30	-1.70±0.25	-2.00±0.1	-1.96±0.09	0.43±0.12	0.20±0.14	
M22c_6	5197±243	5425±115	5383±104	2.7 ±0.7	3.30±0.46	3.12±0.38	-1.7 ±0.4	-2.00±0.1	-1.98±0.10	0.37±0.22	0.27±0.12	M
M22c_7	4937±317	5150±122	5123±114	2.0 ±0.5	2.65±0.32	2.46±0.27	-1.55±0.31	-1.70±0.24	-1.64±0.19	0.41±0.14	0.25±0.13	M
M22c_8	4943±330	5100±122	5081±114	2.1 ±0.7	2.35±0.23	2.33±0.22	-1.7 ±0.4	-2.00±0.1	-1.98±0.10	0.42±0.13	0.18±0.13	M
M22c_9	4923±324	5000±100	4993± 96	2.0 ±0.6	2.30±0.33	2.23±0.29	-1.60±0.32	-2.00±0.1	-1.96±0.10	0.40±0.13	0.19±0.14	M
M22c_10	4964±342	5100±123	5084±116	2.1 ±0.8	2.25±0.25	2.24±0.24	-1.7 ±0.4	-2.00±0.1	-1.98±0.10	0.40±0.13	0.18±0.14	M
M22c_11	4812±351	5225± 75	5207± 73	1.7 ±0.5	3.20±0.33	2.74±0.28	-1.72±0.33	-2.00±0.1	-1.98±0.10	0.43±0.14	0.21±0.13	M
M22c_12	4820±170	5100±122	5005± 99	2.7 ±0.4	3.35±0.23	3.19±0.20	-0.19±0.18	-0.30±0.24	-0.23±0.14	0.10±0.15	0.21±0.11	
M22c_13	4929±324	5125±167	5084±148	2.0 ±0.6	2.55±0.42	2.37±0.34	-1.59±0.32	-1.80±0.25	-1.72±0.20	0.42±0.13	0.27±0.12	M
M22c_14	4793±515	5075±160	5050±153	1.4 ±0.6	2.30±0.46	1.97±0.37	-1.8 ±0.4	-2.05±0.15	-2.02±0.14	0.53±0.04	0.18±0.14	M
M22c_15	4920±325	5300±217	5183±180	2.0 ±0.6	2.85±0.59	2.43±0.42	-1.62±0.30	-1.65±0.23	-1.64±0.18	0.41±0.12	0.20±0.11	M
M22c_16	5006±349	5425±114	5385±108	2.2 ±0.8	3.00±0.50	2.78±0.42	-1.60±0.31	-1.65±0.23	-1.63±0.18	0.43±0.13	0.24±0.14	M
M22c_17	4946±323	5350±122	5300±114	2.1 ±0.7	3.10±0.37	2.88±0.33	-1.64±0.29	-1.80±0.25	-1.73±0.19	0.41±0.12	0.26±0.11	M
M22c_18	4888±330	5050±100	5036± 96	2.0 ±0.7	2.20±0.33	2.16±0.30	-1.71±0.29	-2.00±0.1	-1.97±0.09	0.41±0.13	0.20±0.14	M
5824_1	4760±188	4825±115	4807± 98	2.8 ±0.4	3.34±0.32	3.13±0.25	-0.22±0.15	-0.65±0.23	-0.35±0.13	0.12±0.13	0.26±0.09	
5824_2	4401±162	4725± 74	4669± 67	1.01±0.21	1.45±0.35	1.13±0.18	-1.99±0.17	-2.00±0.1	-2.00±0.09	0.45±0.14	0.13±0.13	M
5824_3	4471±195	4600±122	4564±103	1.11±0.23	1.55±0.35	1.24±0.19	-2.15±0.1	-2.30±0.24	-2.17±0.09	0.40±0.14	0.24±0.09	M
5824_4	4438±226	4401±122	4409±107	1.1 ±0.8	1.00±0.59	1.04±0.47	-1.59±0.21	-2.10±0.20	-1.86±0.14	0.52±0.11	0.30±0.08	M
5824_5	4975±343	5250±100	5228± 96	2.1 ±0.8	2.60±0.49	2.46±0.42	-1.8 ±0.4	-2.00±0.1	-1.99±0.10	0.40±0.17	0.26±0.09	M
5824_6	4851±324	5000±100	4987± 96	1.9 ±0.7	2.00±0.39	1.98±0.34	-1.75±0.28	-2.00±0.1	-1.97±0.09	0.40±0.12	0.17±0.13	M
5824_7	4614±324	4701± 99	4694± 95	1.4 ±0.6	1.65±0.45	1.56±0.36	-1.64±0.34	-2.00±0.1	-1.97±0.10	0.43±0.15	0.23±0.14	M
5824_8	4538±161	5000±100	4871± 85	1.3 ±0.4	2.10±0.37	1.73±0.27	-2.02±0.18	-2.00±0.1	-2.00±0.09	0.48±0.12	0.25±0.14	
5824_9	4565±334	4850±122	4816±115	1.2 ±0.4	1.50±0.39	1.35±0.28	-1.7 ±0.4	-1.80±0.24	-1.77±0.21	0.47±0.11	0.20±0.13	M
5824_10	4400±212	4475± 74	4467± 70	1.1 ±0.6	1.30±0.46	1.23±0.37	-1.79±0.32	-2.05±0.15	-2.00±0.14	0.52±0.11	0.27±0.12	M
5824_11	4893±334	5025± 75	5019± 73	2.0 ±0.6	2.05±0.41	2.03±0.34	-1.67±0.29	-2.00±0.1	-1.96±0.09	0.42±0.13	0.26±0.11	
5824_13	4618±240	4926±114	4869±103	1.3 ±0.4	2.01±0.45	1.61±0.30	-1.89±0.24	-2.00±0.1	-1.98±0.09	0.46±0.11	0.25±0.10	M
5824_14	4539±73	5025± 75	4775± 52	1.20±0.10	2.10±0.37	1.26±0.10	-1.97±0.19	-2.00±0.1	-1.99±0.09	0.33±0.08	0.20±0.14	M

Table B.3 - continued.

NGC ID	$T_{eff}^{(a)}$ (K)	$T_{eff}^{(b)}$ (K)	$T_{eff}^{(avg)}$ (K)	$\log(g)^{(a)}$	$\log(g)^{(b)}$	$\log(g)^{(avg)}$	[Fe/H] ^(a)	[Fe/H] ^(b)	[Fe/H] ^(avg)	[Mg/Fe] ^(a)	[α /Fe] ^(b)	members
5824_15	4891±343	5000±100	4991± 96	2.0 ±0.7	2.00±0.39	2.00±0.34	-1.70±0.30	-2.00±0.1	-1.97±0.09	0.41±0.13	0.22±0.12	M
5824_16	4932±348	5250±100	5226± 96	2.0 ±0.7	2.15±0.32	2.12±0.29	-1.8 ±0.4	-2.00±0.1	-1.99±0.10	0.44±0.12	0.20±0.14	M
5824_17	4488±80	4899±123	4610± 67	1.12±0.12	1.95±0.42	1.18±0.12	-2.03±0.16	-2.00±0.1	-2.01±0.08	0.41±0.11	0.28±0.12	M
5824_18	4531±184	4701± 99	4663± 87	1.27±0.35	1.60±0.37	1.43±0.25	-1.93±0.20	-2.10±0.20	-2.01±0.14	0.48±0.11	0.21±0.14	M
5824_19	4480±184	4578±224	4519±142	1.20±0.26	1.36±0.71	1.22±0.24	-1.85±0.19	-2.15±0.23	-1.97±0.15	0.47±0.12	0.27±0.11	M
47Tuc_502	3640±100	3627±125	3635± 78	0.70±0.2	3.5±0.5	1.08±0.19	-0.10±0.07	-0.04±0.25	-0.10±0.07	0.23±0.1	0.37±0.05	
47Tuc_509	3983±84	3773±284	3966± 81	1.47±0.30	1.1±0.9	1.43±0.28	-0.31±0.20	-1.60±0.44	-0.53±0.18	0.12±0.18	0.32±0.10	M
47Tuc_514	4651±217	4823±196	4746±145	2.3±0.5	2.9±0.5	2.61±0.36	-0.47±0.24	-0.75±0.25	-0.60±0.17	0.25±0.20	0.28±0.10	M
47Tuc_517	4688±224	4728±175	4713±138	2.4±0.5	2.6±0.4	2.51±0.33	-0.47±0.25	-0.90±0.20	-0.73±0.16	0.26±0.20	0.30±0.08	M
47Tuc_519	4263±154	4277±134	4271±101	1.8±0.4	1.96±0.35	1.89±0.26	-0.46±0.22	-1.05±0.15	-0.86±0.12	0.21±0.21	0.23±0.13	M
47Tuc_525	4272±152	4277±134	4275±101	1.8±0.4	1.95±0.35	1.88±0.26	-0.47±0.22	-1.10±0.20	-0.81±0.15	0.22±0.20	0.27±0.11	M
47Tuc_533	4278±107	4300±147	4286± 86	1.8±0.4	1.8±0.4	1.82±0.28	-0.51±0.23	-0.94±0.27	-0.69±0.18	0.29±0.20	0.29±0.09	M
47Tuc_534	4879±363	5125±167	5082±152	2.3±0.8	2.8±0.4	2.70±0.36	-0.7 ±0.4	-0.80±0.25	-0.77±0.21	0.29±0.15	0.19±0.12	M
47Tuc_535	4582±175	4596±121	4591±100	2.3±0.4	2.3±0.4	2.32±0.30	-0.46±0.19	-0.81±0.24	-0.59±0.15	0.25±0.21	0.23±0.11	M
47Tuc_539	4488±180	4628±125	4582±103	2.1±0.5	2.49±0.31	2.38±0.26	-0.46±0.21	-0.74±0.25	-0.58±0.16	0.25±0.21	0.25±0.08	M
47Tuc_551	4949±309	5148±123	5121±114	2.5±0.6	2.8±0.4	2.71±0.33	-0.61±0.29	-0.70±0.25	-0.66±0.19	0.30±0.13	0.20±0.14	M
47Tuc_553	4530±187	4623±125	4594±104	2.2±0.5	2.42±0.35	2.35±0.29	-0.47±0.18	-0.75±0.25	-0.57±0.15	0.24±0.20	0.22±0.10	M
47Tuc_554	3909±34	3750±156	3902± 33	1.43±0.17	0.7±0.5	1.35±0.16	-0.28±0.19	-1.40±0.20	-0.81±0.14	0.12±0.20	0.26±0.11	M
47Tuc_559	4780±282	4749±193	4759±159	2.5±0.6	2.85±0.32	2.77±0.28	-0.52±0.32	-1.05±0.35	-0.76±0.24	0.27±0.20	0.32±0.10	M
47Tuc_571	4972±439	5277±260	5198±224	2.4±0.9	2.8±0.5	2.72±0.41	-0.7 ±0.5	-0.80±0.33	-0.77±0.28	0.25±0.17	0.26±0.11	M
47Tuc_581	4868±346	5071±113	5051±107	2.2±0.8	2.44±0.34	2.40±0.31	-0.8 ±0.4	-0.91±0.19	-0.89±0.17	0.36±0.13	0.22±0.15	M
2808_1	4676±331	4925±115	4898±108	1.8±0.9	1.65±0.23	1.66±0.22	-1.0±0.5	-1.15±0.23	-1.12±0.21	0.32±0.17	0.16±0.12	M
2808_2	4736±167	5124±125	4985±100	2.52±0.30	3.25±0.25	2.95±0.19	-0.30±0.22	-0.25±0.25	-0.28±0.17	0.16±0.19	0.21±0.09	
2808_3	4812±360	4825±115	4824±109	2.2±0.9	2.50±0.39	2.45±0.36	-0.9±0.5	-1.35±0.23	-1.27±0.21	0.34±0.17	0.24±0.12	M
2808_4	4949±314	4800±245	4856±193	2.5±0.7	2.00±0.45	2.15±0.38	-0.57±0.30	-1.20±0.24	-0.95±0.19	0.24±0.13	0.24±0.14	M
2808_5	4801±399	5276±208	5175±184	1.9±0.8	2.65±0.45	2.47±0.39	-1.4±0.4	-1.20±0.24	-1.25±0.21	0.42±0.16	0.19±0.14	M
2808_6	4910±354	4750±158	4777±144	2.5±0.9	2.20±0.56	2.28±0.48	-0.7±0.5	-1.40±0.20	-1.30±0.19	0.32±0.20	0.26±0.09	M
2808_7	4903±401	4799±149	4812±140	2.3±0.9	1.85±0.32	1.90±0.30	-0.8±0.5	-1.45±0.15	-1.40±0.14	0.31±0.20	0.19±0.14	M

Table B.3 - continued.

NGC ID	$T_{eff}^{(a)}$ (K)	$T_{eff}^{(b)}$ (K)	$T_{eff}^{(avg)}$ (K)	$\log(g)^{(a)}$	$\log(g)^{(b)}$	$\log(g)^{(avg)}$	[Fe/H] ^(a)	[Fe/H] ^(b)	[Fe/H] ^(avg)	[Mg/Fe] ^(a)	[α /Fe] ^(b)	members
2808_8	4786±413	4651±122	4662±117	2.3±0.8	1.40±0.49	1.65±0.42	-0.51±0.34	-1.20±0.24	-0.97±0.20	0.12±0.06	0.21±0.14	M
2808_9	4687±362	4925±114	4904±109	1.7±0.8	2.00±0.45	1.93±0.39	-1.4±0.4	-1.50±0.1	-1.49±0.10	0.42±0.14	0.22±0.12	
2808_10	4072±114	3925±296	4053±106	1.4±0.4	0.90±0.73	1.28±0.35	-0.47±0.22	-1.50±0.39	-0.72±0.19	0.19±0.24	0.32±0.07	M
2808_11	4662±415	4900±122	4881±117	1.9±0.9	2.05±0.35	2.03±0.33	-0.8±0.5	-1.00±0.1	-0.99±0.10	0.30±0.17	0.20±0.12	
2808_13	4961±279	4773±236	4851±180	2.7±0.6	1.90±0.37	2.12±0.31	-0.52±0.31	-1.25±0.25	-0.96±0.19	0.24±0.17	0.25±0.13	M
2808_14	4968±300	5049± 99	5041± 94	2.5±0.6	2.40±0.37	2.43±0.31	-0.57±0.31	-0.90±0.20	-0.80±0.17	0.25±0.13	0.19±0.14	M
2808_15	4638±430	5075±225	4981±199	1.5±0.8	2.00±0.39	1.90±0.35	-1.2±0.5	-1.15±0.23	-1.16±0.21	0.38±0.18	0.22±0.14	M
2808_16	4328±48	4526±175	4342± 46	1.20±0.05	1.90±0.44	1.21±0.05	-0.72±0.24	-1.10±0.20	-0.94±0.15	0.15±0.14	0.22±0.11	M
2808_17	4832±264	4625±125	4663±113	2.5±0.7	2.05±0.27	2.11±0.25	-0.51±0.29	-1.25±0.25	-0.93±0.19	0.19±0.15	0.25±0.10	M
2808_18	4433±74	4576±159	4458± 67	2.36±0.23	2.55±0.47	2.40±0.21	-0.45±0.08	-0.85±0.32	-0.47±0.08	0.16±0.15	0.27±0.11	
2808_19	4874±359	5175±114	5147±109	2.3±0.9	3.15±0.32	3.05±0.30	-1.0±0.5	-1.05±0.15	-1.05±0.14	0.34±0.19	0.18±0.14	M
3201_1	4819±392	5225± 75	5211± 73	1.9±0.9	3.20±0.25	3.11±0.24	-1.60±0.32	-1.50±0.1	-1.51±0.10	0.43±0.13	0.19±0.15	
3201_2	4757±322	4975±175	4925±154	1.7±0.7	2.45±0.35	2.30±0.31	-1.56±0.32	-1.65±0.23	-1.62±0.19	0.43±0.14	0.22±0.12	M
3201_5	4775±369	5250±100	5218± 97	1.8±0.8	2.60±0.37	2.46±0.34	-1.57±0.32	-1.50±0.1	-1.51±0.10	0.43±0.14	0.24±0.14	M
3201_6	4509±274	5076±160	4932±138	1.2±0.6	2.55±0.41	2.12±0.34	-1.48±0.34	-1.30±0.25	-1.36±0.20	0.46±0.12	0.21±0.13	
3201_7	4891±346	5225± 75	5210± 73	2.0±0.7	2.60±0.37	2.47±0.33	-1.56±0.31	-1.55±0.15	-1.55±0.14	0.43±0.14	0.27±0.11	M
3201_8	4408±178	4800±100	4706± 87	1.0±0.5	2.00±0.39	1.62±0.31	-1.52±0.20	-1.50±0.1	-1.50±0.09	0.45±0.14	0.25±0.14	M
3201_9	4774±347	5025± 75	5014± 73	1.7±0.7	2.45±0.35	2.30±0.31	-1.54±0.30	-1.50±0.1	-1.50±0.09	0.45±0.14	0.19±0.13	M
3201_10	4752±340	5049± 99	5026± 95	1.7±0.8	2.40±0.37	2.28±0.34	-1.50±0.27	-1.40±0.20	-1.44±0.16	0.46±0.15	0.18±0.14	M
3201_11	4997±287	5250±111	5217±104	2.2±0.7	3.00±0.39	2.81±0.34	-1.47±0.27	-1.55±0.15	-1.53±0.13	0.42±0.16	0.21±0.14	M
3201_13	4928±465	5075±195	5053±180	2.1±0.9	2.65±0.39	2.56±0.36	-1.52±0.35	-1.60±0.20	-1.58±0.17	0.45±0.15	0.23±0.13	M
3201_14	4851±353	5050±100	5035± 96	1.9±0.8	2.45±0.35	2.36±0.32	-1.51±0.28	-1.50±0.1	-1.50±0.09	0.44±0.15	0.15±0.10	M
3201_15	4925±385	5250±100	5229± 97	2.1±0.8	3.00±0.39	2.83±0.35	-1.53±0.31	-1.50±0.1	-1.50±0.10	0.44±0.14	0.19±0.14	M
3201_16	4600±367	4750±250	4702±207	1.7±1.0	2.05±0.47	1.99±0.43	-1.0 ±0.5	-1.25±0.25	-1.20±0.22	0.31±0.16	0.18±0.15	M
3201_17	4893±349	5101±229	5038±191	2.0±0.8	2.40±0.54	2.27±0.45	-1.54±0.30	-1.65±0.23	-1.61±0.18	0.43±0.15	0.27±0.11	M
3201_18	4813±380	5175±160	5121±147	1.9±0.8	2.75±0.40	2.58±0.36	-1.59±0.32	-1.60±0.20	-1.60±0.17	0.43±0.14	0.25±0.12	M
M68_49	5111±355	5500±100	5471± 96	2.3±0.7	3.05±0.42	2.85±0.36	-1.8±0.5	-2.00±0.1	-1.99±0.10	0.39±0.19	0.25±0.10	M
M68_30	4907±300	5025±175	4995±151	1.9±0.7	2.65±0.39	2.47±0.34	-1.9±0.4	-2.35±0.23	-2.24±0.20	0.39±0.11	0.12±0.13	M

Table B.3 - continued.

NGC ID	$T_{eff}^{(a)}$ (K)	$T_{eff}^{(b)}$ (K)	$T_{eff}^{(avg)}$ (K)	$\log(g)^{(a)}$	$\log(g)^{(b)}$	$\log(g)^{(avg)}$	[Fe/H] ^(a)	[Fe/H] ^(b)	[Fe/H] ^(avg)	[Mg/Fe] ^(a)	[α /Fe] ^(b)	members
M68_74	5041±354	5049± 99	5049± 96	2.2±0.8	2.45±0.61	2.36±0.49	-2.0±0.4	-2.40±0.20	-2.32±0.18	0.39±0.18	0.20±0.14	M
M68_II47	4990±434	4825±115	4836±111	2.0±0.7	2.10±0.37	2.08±0.33	-1.8±0.4	-2.50±0.1	-2.46±0.10	0.41±0.18	0.22±0.14	M
M68_119	4908±326	4775± 75	4782± 73	2.0±0.7	2.35±0.32	2.29±0.29	-1.74±0.34	-2.45±0.15	-2.33±0.14	0.36±0.13	0.21±0.13	M
M68_144	5219±385	5775± 75	5755± 74	2.5±0.8	3.60±0.54	3.26±0.45	-2.0±0.4	-2.50±0.1	-2.47±0.10	0.37±0.20	0.10±0.10	
M68_185	4862±15	5100±122	4866± 15	3.4±0.2	4.45±0.35	3.66±0.17	0.02±0.09	-0.65±0.23	-0.07±0.08	0.04±0.03	0.21±0.09	
M68_239	4910±300	5226± 74	5208± 72	1.9±0.7	2.21±0.25	2.17±0.24	-1.9±0.4	-2.00±0.1	-1.99±0.10	0.39±0.11	0.19±0.15	M
M68_II72	5096±393	5075±160	5078±148	2.2±0.8	2.55±0.61	2.42±0.49	-1.9±0.4	-2.40±0.20	-2.30±0.18	0.38±0.18	0.12±0.12	M
M68_II79	4862±16	5274±175	4865± 16	3.4±0.2	4.25±0.25	3.73±0.16	0.02±0.09	-0.18±0.27	0.00±0.09	0.04±0.03	0.09±0.08	
M5_II45	4692±330	4874±202	4825±172	1.7±0.8	2.40±0.54	2.18±0.45	-1.48±0.27	-1.65±0.23	-1.58±0.18	0.43±0.15	0.27±0.11	M
M5_II50	4705±304	4645±227	4667±182	2.0±0.8	1.99±0.39	1.99±0.35	-0.9±0.4	-1.35±0.23	-1.24±0.20	0.29±0.13	0.21±0.11	M
M5_II51	4727±329	4600±122	4615±115	1.9±0.9	1.85±0.23	1.85±0.22	-1.1±0.4	-1.80±0.24	-1.61±0.21	0.42±0.16	0.26±0.11	M
M5_II80	4560±273	4773± 73	4759± 70	1.5±0.8	2.30±0.39	2.15±0.35	-1.2±0.4	-1.45±0.15	-1.42±0.14	0.38±0.15	0.21±0.13	M
M5_I2	4810±356	4748±250	4768±205	2.1±0.8	1.95±0.42	1.98±0.37	-0.8±0.4	-1.25±0.25	-1.12±0.21	0.32±0.14	0.25±0.12	M
M5_I50	4739±377	4848±122	4838±116	2.0±1.0	2.55±0.35	2.49±0.33	-1.0±0.5	-1.30±0.24	-1.24±0.22	0.38±0.16	0.20±0.11	M
M5_I61	4618±340	4597±164	4601±148	2.0±0.9	1.99±0.38	1.99±0.35	-0.9±0.4	-1.45±0.15	-1.38±0.14	0.29±0.18	0.27±0.13	M
M5_I68	4484±226	4224±174	4321±138	2.0±0.6	1.60±0.43	1.74±0.35	-0.59±0.32	-1.55±0.35	-1.03±0.24	0.28±0.19	0.27±0.09	M
M5_I71	4702±412	4927±113	4911±109	1.9±0.9	2.06±0.41	2.03±0.37	-0.9±0.5	-1.00±0.1	-1.00±0.10	0.34±0.17	0.25±0.13	M
M4.1	4764±371	4974±174	4936±158	2.1±0.9	2.90±0.43	2.75±0.388	-0.9±0.4	-1.15±0.23	-1.09±0.20	0.38±0.16	0.26±0.13	M
M4.3	4816±367	5200±150	5145±139	2.1±0.9	3.25±0.25	3.17±0.241	-1.2±0.4	-1.15±0.23	-1.16±0.20	0.40±0.17	0.28±0.10	M
M4.4	4946±364	5151±122	5130±116	2.6±0.9	3.20±0.25	3.16±0.241	-0.7±0.5	-1.00±0.1	-0.99±0.10	0.32±0.21	0.21±0.11	M
M4.5	4883±348	4999±157	4979±143	2.5±0.8	2.75±0.25	2.73±0.239	-0.7±0.4	-1.00±0.31	-0.89±0.25	0.33±0.17	0.30±0.10	M
M4.6	4504±163	4701±268	4557±139	2.0±0.7	2.50±0.59	2.29±0.451	-0.62±0.19	-1.00±0.39	-0.69±0.17	0.17±0.14	0.28±0.07	
M4.8	4781±378	5025±174	4982±158	2.2±0.9	2.80±0.24	2.76±0.232	-0.8±0.5	-0.95±0.35	-0.90±0.29	0.34±0.17	0.30±0.10	
M4.9	4896±359	4925±195	4919±172	2.5±0.9	2.65±0.39	2.63±0.358	-0.8±0.5	-1.25±0.25	-1.16±0.22	0.33±0.20	0.27±0.09	
M4.10	4792±280	4951±186	4902±155	2.3±0.7	2.95±0.35	2.82±0.313	-0.84±0.31	-1.20±0.24	-1.07±0.19	0.36±0.17	0.25±0.13	M
M4.11	4875±336	4999±157	4977±142	2.5±0.8	2.80±0.24	2.78±0.230	-0.8±0.4	-1.00±0.31	-0.92±0.25	0.34±0.17	0.26±0.13	M
M4.13	4798±387	5125±125	5094±119	2.2±1.0	3.15±0.23	3.10±0.224	-1.0±0.5	-1.00±0.1	-1.00±0.10	0.38±0.17	0.24±0.14	M
M4.16	5038±282	4850±299	4950±205	2.9±0.5	2.35±0.63	2.69±0.392	-0.49±0.31	-1.20±0.33	-0.82±0.23	0.27±0.24	0.31±0.10	

Table B.3 - continued.

NGC ID	$T_{eff}^{(a)}$ (K)	$T_{eff}^{(b)}$ (K)	$T_{eff}^{(avg)}$ (K)	$\log(g)^{(a)}$	$\log(g)^{(b)}$	$\log(g)^{(avg)}$	$[\text{Fe}/\text{H}]^{(a)}$	$[\text{Fe}/\text{H}]^{(b)}$	$[\text{Fe}/\text{H}]^{(avg)}$	$[\text{Mg}/\text{Fe}]^{(a)}$	$[\alpha/\text{Fe}]^{(b)}$	members
M4_17	4980±260	4728±208	4827±163	3.0±0.4	2.26±0.60	2.77±0.333	-0.52±0.30	-1.35±0.23	-1.04±0.18	0.28±0.22	0.29±0.09	M
M4_18	5030±44	5072±336	5031± 44	2.99±0.09	2.35±0.59	2.98±0.089	-0.49±0.1	-1.10±0.44	-0.52±0.10	0.22±0.05	0.31±0.08	
M4_19	6330±241	5976±175	6098±142	3.63±0.007	3.85±0.32	3.63±0.007	0.14±0.08	-0.35±0.23	0.09±0.08	0.09±0.007	0.09±0.10	
M107_F	4262±123	4201±185	4243±102	1.8±0.4	1.30±0.45	1.58±0.30	-0.45±0.24	-1.10±0.37	-0.64±0.20	0.24±0.20	0.29±0.09	
M107_62	4497±178	4525±208	4509±135	2.1±0.4	2.55±0.42	2.31±0.29	-0.56±0.21	-0.95±0.42	-0.64±0.19	0.33±0.20	0.24±0.11	
M107_100	4801±206	4951±242	4864±157	2.4±0.5	2.90±0.49	2.66±0.35	-0.40±0.23	-0.50±0.31	-0.44±0.18	0.23±0.22	0.12±0.13	
M107_I	4779±210	4798±245	4787±160	2.5±0.4	2.65±0.67	2.54±0.34	-0.31±0.29	-0.70±0.25	-0.53±0.19	0.17±0.22	0.19±0.13	
M107_C	4845±269	4799± 99	4804± 93	2.4±0.6	2.30±0.25	2.31±0.23	-0.55±0.27	-1.00±0.1	-0.95±0.09	0.28±0.21	0.20±0.14	M ^(c)
M10_1	4944±316	5200±218	5118±179	2.0±0.6	2.80±0.56	2.43±0.41	-1.59±0.28	-1.75±0.25	-1.68±0.19	0.41±0.13	0.19±0.11	M
M10_2	4788±291	5126±167	5043±145	1.7±0.5	2.10±0.30	1.99±0.26	-1.59±0.31	-1.55±0.15	-1.56±0.14	0.45±0.13	0.23±0.13	M
M10_3	4949±296	5300±187	5200±158	2.0±0.6	3.05±0.35	2.78±0.30	-1.55±0.29	-1.60±0.20	-1.58±0.16	0.42±0.14	0.28±0.13	M
M10_4	4807±306	5101±123	5060±114	1.8±0.6	2.60±0.37	2.38±0.31	-1.66±0.29	-1.80±0.25	-1.74±0.19	0.40±0.12	0.26±0.13	M
M10_5	4776±285	5176±114	5121±106	1.7±0.5	2.20±0.25	2.10±0.22	-1.62±0.32	-1.50±0.1	-1.51±0.10	0.44±0.12	0.19±0.14	M
M10_6	4686±300	4950±187	4876±159	1.5±0.6	2.10±0.30	1.98±0.27	-1.62±0.31	-1.70±0.24	-1.67±0.19	0.46±0.13	0.25±0.10	M
M10_7	4605±313	5000±100	4963± 95	1.3±0.5	2.11±0.37	1.82±0.30	-1.52±0.25	-1.50±0.1	-1.50±0.09	0.48±0.12	0.25±0.14	M
M10_8	4906±341	5151±122	5123±115	2.0±0.7	3.00±0.32	2.83±0.29	-1.61±0.30	-1.70±0.24	-1.66±0.19	0.43±0.13	0.23±0.16	M
M10_9	4361±125	4750±100	4598± 78	0.90±0.31	2.05±0.41	1.32±0.25	-1.53±0.21	-1.50±0.1	-1.51±0.09	0.42±0.12	0.22±0.13	M
M10_10	4589±307	5000±100	4961± 95	1.3±0.5	2.20±0.33	1.93±0.28	-1.57±0.28	-1.50±0.1	-1.51±0.09	0.48±0.11	0.21±0.15	M
M10_11	4947±323	5251±223	5152±184	2.1±0.7	3.00±0.59	2.63±0.45	-1.62±0.30	-1.70±0.24	-1.67±0.19	0.43±0.13	0.14±0.09	M
M10_13	4948±324	5425±114	5372±108	2.1±0.7	3.05±0.35	2.86±0.31	-1.64±0.29	-1.60±0.20	-1.61±0.16	0.42±0.12	0.18±0.12	M
M10_15	4949±336	5375±125	5323±117	2.1±0.7	3.05±0.35	2.86±0.31	-1.69±0.30	-1.65±0.23	-1.66±0.18	0.41±0.12	0.23±0.16	M
M10_16	4853±326	5201±148	5142±135	1.8±0.5	2.25±0.40	2.07±0.31	-1.59±0.34	-1.55±0.15	-1.56±0.14	0.42±0.14	0.23±0.13	
M10_17	4970±338	5425±115	5378±109	2.1±0.7	3.20±0.25	3.08±0.24	-1.60±0.31	-1.50±0.1	-1.51±0.10	0.43±0.13	0.20±0.14	
6397_1	5006±354	5250±100	5232± 96	2.2±0.9	2.60±0.49	2.51±0.43	-2.0±0.4	-2.50±0.1	-2.47±0.10	0.40±0.19	0.27±0.09	M
6397_2	5191±409	5350±122	5337±117	2.5±0.9	2.85±0.81	2.69±0.60	-2.1±0.4	-2.50±0.1	-2.48±0.10	0.41±0.24	0.05±0.07	
6397_4	5071±253	5050±100	5053± 93	2.3±0.7	2.25±0.51	2.27±0.41	-2.0±0.4	-2.40±0.20	-2.32±0.18	0.36±0.20	0.21±0.13	M
6397_5	5137±330	5500±100	5469± 96	2.4±0.8	3.40±0.37	3.22±0.34	-1.9±0.4	-2.00±0.1	-1.99±0.10	0.41±0.18	0.19±0.14	M

Table B.3 - continued.

NGC ID	$T_{eff}^{(a)}$ (K)	$T_{eff}^{(b)}$ (K)	$T_{eff}^{(avg)}$ (K)	$\log(g)^{(a)}$	$\log(g)^{(b)}$	$\log(g)^{(avg)}$	[Fe/H] ^(a)	[Fe/H] ^(b)	[Fe/H] ^(avg)	[Mg/Fe] ^(a)	[α /Fe] ^(b)	members
6397_6	5036±402	5350±122	5323±117	2.2±0.8	2.80±0.71	2.54±0.53	-1.9±0.4	-2.30±0.24	-2.19±0.21	0.40±0.17	0.18±0.12	M
6397_7	5043±312	5425±115	5379±108	2.2±0.8	3.15±0.32	3.02±0.30	-1.9±0.4	-2.00±0.1	-1.99±0.10	0.41±0.18	0.25±0.12	M
6397_8	4963±324	5075±114	5062±108	2.1±0.8	2.45±0.42	2.37±0.37	-2.0±0.4	-2.35±0.23	-2.26±0.20	0.39±0.18	0.20±0.13	M
6397_9	5115±284	5250±100	5235± 94	2.4±0.8	2.50±0.39	2.48±0.35	-2.1±0.4	-2.50±0.1	-2.48±0.10	0.35±0.26	0.25±0.13	M
6397_10	4992±317	5125±125	5107±116	2.2±0.8	2.40±0.66	2.32±0.51	-2.0±0.4	-2.30±0.25	-2.22±0.21	0.39±0.18	0.24±0.12	M
6397_11	4906±332	5250±100	5221± 96	2.1±0.8	2.60±0.37	2.51±0.34	-1.81±0.34	-2.00±0.1	-1.98±0.10	0.43±0.13	0.22±0.12	M
6397_13	5006±307	5050± 99	5046± 95	2.3±0.7	2.15±0.55	2.21±0.43	-1.8±0.4	-2.40±0.20	-2.28±0.18	0.37±0.12	0.22±0.12	M
6397_14	5105±278	5225± 74	5217± 72	2.4±0.8	2.85±0.45	2.74±0.39	-1.8±0.4	-2.05±0.15	-2.02±0.14	0.38±0.20	0.17±0.15	M
6397_14b	4957±311	5154±197	5097±167	2.2±0.7	2.30±0.45	2.27±0.38	-1.70±0.26	-2.10±0.20	-1.95±0.16	0.40±0.12	0.30±0.10	M
6397_15	5042±363	5450±100	5421± 96	2.2±0.8	3.15±0.45	2.92±0.39	-2.0±0.4	-2.10±0.20	-2.08±0.18	0.40±0.17	0.22±0.14	M
6397_16	4929±329	5226± 74	5211± 72	2.1±0.8	2.20±0.25	2.19±0.24	-1.75±0.27	-2.00±0.1	-1.97±0.09	0.42±0.12	0.18±0.14	M
6397_16b	4797±306	5000±100	4980± 95	1.8±0.6	2.00±0.39	1.94±0.33	-1.79±0.26	-2.00±0.1	-1.97±0.09	0.42±0.12	0.17±0.13	M
6397_19	5124±306	5300±100	5283± 95	2.4±0.8	2.95±0.42	2.83±0.37	-2.0±0.4	-2.40±0.20	-2.32±0.18	0.39±0.19	0.20±0.13	M
6397_319	4849± 93	5125±125	4947± 75	3.45±0.11	4.35±0.23	3.62±0.10	0.09±0.06	-0.06±0.23	0.08±0.06	-0.02±0.04	0.14±0.09	
6397_344	3640± 60	3525± 74	3595± 47	0.70±0.13	2.39±0.83	0.74±0.13	0.00±0.18	-1.10±0.30	-0.29±0.15	0.23±0.10	0.30±0.10	
6397_361	4617±198	5000±100	4922± 89	1.4±0.5	2.05±0.41	1.79±0.32	-1.91±0.22	-2.00±0.1	-1.98±0.09	0.44±0.13	0.26±0.11	M
6397_365	5200± 98	4922±224	5156± 90	2.63±0.13	2.24±0.56	2.61±0.13	-1.60±0.09	-2.35±0.23	-1.70±0.08	0.40±0.05	0.25±0.10	M
6441_1	4514±148	4600±165	4552±110	2.30±0.40	2.50±0.50	2.38±0.31	0.06±0.09	-0.33±0.26	0.02±0.09	0.02±0.13	0.16±0.12	
6441_2	4314±137	4424±225	4344±117	2.00±0.40	2.25±0.56	2.08±0.33	0.08±0.16	-0.29±0.32	0.01±0.14	0.00±0.14	0.28±0.13	
6441_3	4507±157	4601±166	4551±114	2.20±0.40	2.50±0.45	2.33±0.30	-0.30±0.27	-0.75±0.25	-0.54±0.18	0.16±0.19	0.27±0.11	M
6441_4	4608±171	4800±186	4696±126	2.50±0.40	2.90±0.20	2.82±0.18	-0.12±0.20	-0.40±0.37	-0.18±0.18	0.08±0.14	0.23±0.13	M
6441_5	4839±188	4949±186	4895±132	2.60±0.40	2.75±0.33	2.69±0.25	-0.29±0.29	-0.60±0.37	-0.41±0.23	0.16±0.22	0.24±0.14	M
6441_6	3894± 74	3643±198	3863± 69	1.44±0.15	0.67±0.68	1.40±0.15	-0.07±0.20	-1.26±0.33	-0.39±0.17	0.09±0.20	0.23±0.13	M
6441_7	4376±193	4625±125	4552±105	2.10±0.50	2.90±0.20	2.79±0.19	0.01±0.18	-0.21±0.30	-0.05±0.15	0.03±0.14	0.19±0.11	
6441_8	4500±163	4475±175	4488±119	2.20±0.40	1.80±0.40	2.00±0.28	-0.08±0.24	-0.65±0.23	-0.38±0.17	0.06±0.18	0.26±0.10	M
6441_9	4571±193	4725±175	4655±130	2.20±0.40	1.90±0.43	2.06±0.29	-0.22±0.23	-0.65±0.23	-0.43±0.16	0.08±0.15	0.30±0.08	M
6441_10	4520±177	4726±260	4585±146	2.30±0.40	2.65±0.59	2.41±0.33	-0.23±0.27	-0.45±0.35	-0.31±0.21	0.13±0.18	0.27±0.09	M
6441_11	4369±179	4750±158	4584±118	2.10±0.50	2.50±0.32	2.38±0.27	0.27±0.09	0.18±0.18	0.25±0.08	0.05±0.07	0.21±0.11	

Table B.3 - continued.

NGC ID	$T_{eff}^{(a)}$ (K)	$T_{eff}^{(b)}$ (K)	$T_{eff}^{(avg)}$ (K)	$\log(g)^{(a)}$	$\log(g)^{(b)}$	$\log(g)^{(avg)}$	$[\text{Fe}/\text{H}]^{(a)}$	$[\text{Fe}/\text{H}]^{(b)}$	$[\text{Fe}/\text{H}]^{(avg)}$	$[\text{Mg}/\text{Fe}]^{(a)}$	$[\alpha/\text{Fe}]^{(b)}$	members
6441_13	4804±282	4801±187	4802±156	2.50±0.60	2.50±0.55	2.50±0.41	-0.40±0.40	-0.80±0.25	-0.69±0.21	0.19±0.21	0.23±0.13	M
6441_14	4222±162	4475±174	4339±119	1.90±0.40	2.65±0.39	2.28±0.28	-0.09±0.21	-0.55±0.35	-0.21±0.18	0.01±0.14	0.31±0.07	
6441_15	4075±147	4324±114	4230± 90	1.60±0.13	2.40±0.37	1.69±0.12	0.01±0.17	-0.35±0.23	-0.12±0.14	0.05±0.12	0.20±0.11	
6441_16	4544±229	4650±166	4614±134	2.40±0.50	2.60±0.37	2.53±0.30	-0.19±0.27	-0.70±0.33	-0.39±0.21	0.11±0.18	0.30±0.10	
6441_17	4413±137	4824±114	4656± 88	2.10±0.40	3.15±0.23	2.89±0.20	0.11±0.13	0.21±0.21	0.14±0.11	0.03±0.15	0.14±0.11	
6441_18	4267±150	4400±122	4347± 95	1.87±0.28	2.45±0.42	2.05±0.23	0.13±0.18	-0.20±0.24	0.01±0.14	-0.00±0.10	0.13±0.11	
6441_19	4355±141	4750±193	4492±114	1.90±0.50	2.60±0.44	2.29±0.33	0.15±0.17	0.15±0.27	0.15±0.14	0.02±0.17	0.18±0.09	
6752_2	4904±382	5101±123	5082±117	1.80±0.50	2.26±0.33	2.12±0.28	-1.50±0.31	-1.50±0.10	-1.50±0.10	0.45±0.15	0.18±0.13	M
6752_4	4811±329	5000±223	4941±185	1.80±0.60	2.70±0.51	2.32±0.39	-1.58±0.31	-1.70±0.24	-1.66±0.19	0.44±0.15	0.25±0.10	M
6752_10	4960±433	4952±245	4954±213	2.10±0.70	2.25±0.64	2.18±0.47	-1.50±0.40	-1.80±0.25	-1.72±0.21	0.44±0.15	0.23±0.13	M
6752_11	4916±316	5076±195	5032±166	1.80±0.40	1.65±0.67	1.76±0.34	-1.57±0.30	-1.75±0.25	-1.68±0.19	0.46±0.12	0.20±0.11	M
6752_12	5487±383	5499±224	5496±193	2.60±0.40	1.89±0.74	2.44±0.35	-1.39±0.22	-1.70±0.25	-1.53±0.17	0.54±0.11	0.21±0.13	M
6752_16	3730± 63	3713±338	3729± 62	0.90±0.14	2.76±1.42	0.92±0.14	0.27±0.19	-2.03±0.47	-0.05±0.18	0.08±0.10	0.35±0.07	
M15_2	5014±240	5050±100	5045± 92	2.20±0.60	2.35±0.32	2.32±0.28	-2.20±0.40	-2.50±0.10	-2.48±0.10	0.45±0.12	0.18±0.15	M
M15_3	4909±337	4950±187	4940±164	2.00±0.80	2.00±0.45	2.00±0.39	-1.90±0.35	-2.40±0.20	-2.28±0.17	0.38±0.19	0.25±0.14	M
M15_4	4977±325	5175±115	5153±108	2.10±0.80	2.00±0.39	2.02±0.35	-2.00±0.40	-2.50±0.10	-2.47±0.10	0.39±0.18	0.20±0.13	M
M15_5	4978±405	5075±115	5068±110	2.10±0.80	2.20±0.33	2.19±0.31	-1.80±0.40	-2.50±0.10	-2.46±0.10	0.41±0.19	0.27±0.09	M
M15_6	4909±305	5000±100	4991± 95	1.90±0.70	1.90±0.37	1.90±0.33	-2.07±0.35	-2.50±0.10	-2.47±0.10	0.39±0.19	0.21±0.13	M
M15_7	4456± 37	4475± 75	4460± 33	1.11±0.10	0.85±0.45	1.10±0.10	-2.12±0.02	-2.50±0.10	-2.13±0.02	0.43±0.09	0.26±0.13	M
M15_8	4771±340	4750±100	4752± 96	1.70±0.70	1.45±0.41	1.51±0.35	-2.00±0.40	-2.50±0.10	-2.47±0.10	0.42±0.10	0.23±0.13	M
M15_9	4771±339	4750±100	4752± 96	1.70±0.70	1.55±0.41	1.59±0.35	-2.00±0.40	-2.50±0.10	-2.47±0.10	0.42±0.10	0.16±0.12	M
M15_10	4903±352	5150±122	5123±116	2.00±0.80	2.50±0.32	2.43±0.30	-1.80±0.40	-2.20±0.24	-2.09±0.21	0.41±0.12	0.25±0.10	M
M15_12	4975±311	5000±100	4998± 95	2.10±0.70	2.10±0.37	2.10±0.33	-2.10±0.40	-2.50±0.10	-2.48±0.10	0.38±0.19	0.19±0.13	M
M15_13	5061±257	5050±100	5051± 93	2.30±0.60	2.10±0.44	2.17±0.35	-2.00±0.40	-2.50±0.10	-2.47±0.10	0.34±0.21	0.28±0.10	M
M15_14	5078±326	5050±100	5052± 96	2.30±0.80	2.20±0.56	2.23±0.46	-1.80±0.40	-2.40±0.20	-2.28±0.18	0.38±0.20	0.24±0.12	M
M15_15	4943±280	5000±111	4992±104	2.00±0.60	2.10±0.44	2.07±0.35	-1.90±0.40	-2.45±0.15	-2.38±0.14	0.39±0.10	0.25±0.10	M
M15_17	4856±344	4950±187	4928±164	2.00±0.80	2.00±0.45	2.00±0.39	-1.86±0.34	-2.40±0.20	-2.26±0.17	0.42±0.12	0.27±0.09	M
M15_18	5152±323	5250±100	5241± 96	2.50±0.80	2.15±0.32	2.20±0.30	-2.10±0.40	-2.50±0.10	-2.48±0.10	0.37±0.20	0.19±0.14	M

Table B.3 - continued.

NGC ID	$T_{eff}^{(a)}$ (K)	$T_{eff}^{(b)}$ (K)	$T_{eff}^{(avg)}$ (K)	$\log(g)^{(a)}$	$\log(g)^{(b)}$	$\log(g)^{(avg)}$	[Fe/H] ^(a)	[Fe/H] ^(b)	[Fe/H] ^(avg)	[Mg/Fe] ^(a)	[α /Fe] ^(b)	members
2298_11	4766±339	4750±100	4751± 96	1.70±0.60	1.90±0.37	1.84±0.31	-1.64±0.34	-2.00±0.10	-1.97±0.10	0.43±0.15	0.19±0.14	M
2298_14	4694±329	4775± 74	4771± 73	1.60±0.60	1.70±0.33	1.68±0.29	-1.63±0.32	-1.95±0.15	-1.89±0.14	0.45±0.13	0.20±0.14	M
2298_16	5006±336	5250±100	5230± 96	2.20±0.80	2.30±0.33	2.29±0.31	-1.80±0.40	-2.00±0.10	-1.99±0.10	0.40±0.18	0.19±0.14	M
2298_17	4589±327	4799±188	4747±163	1.30±0.60	2.05±0.52	1.73±0.39	-1.68±0.35	-1.80±0.24	-1.76±0.20	0.47±0.12	0.23±0.13	M
2298_18	4894±344	5000±100	4992± 96	2.00±0.70	1.95±0.41	1.96±0.35	-1.70±0.40	-2.00±0.10	-1.98±0.10	0.42±0.14	0.14±0.11	M
4372_20	4882±356	4625±125	4653±118	2.30±1.10	1.80±0.71	1.95±0.60	-1.70±0.40	-2.25±0.25	-2.10±0.21	0.39±0.14	0.14±0.13	M
4372_14	4731±192	4675±224	4707±146	1.80±0.50	2.30±0.51	2.05±0.36	-1.70±0.40	-2.05±0.27	-1.94±0.22	0.40±0.16	0.15±0.10	M
4372_13	4392±179	4423±116	4414± 97	1.00±0.50	1.05±0.27	1.04±0.24	-1.82±0.27	-2.15±0.23	-2.01±0.18	0.48±0.12	0.24±0.12	M
4372_95	4984±371	4750±100	4766± 97	2.00±0.90	1.84±0.32	1.86±0.30	-2.00±0.40	-2.50±0.10	-2.47±0.10	0.43±0.09	0.17±0.15	M
4372_91	4890±336	4925±114	4922±108	2.10±0.80	2.05±0.35	2.06±0.32	-1.82±0.34	-2.15±0.23	-2.05±0.19	0.43±0.12	0.27±0.11	M
4372_89	4828±333	4977±236	4927±193	1.80±0.70	2.20±0.40	2.10±0.35	-1.90±0.40	-2.30±0.25	-2.19±0.21	0.41±0.10	0.29±0.09	M
4372_85	5235±360	5575±115	5544±109	2.50±0.80	3.55±0.52	3.24±0.44	-2.34±0.25	-2.50±0.10	-2.48±0.09	0.35±0.26	0.10±0.10	
4372_74	4677±217	3524± 73	3642± 70	1.70±0.60	0.10±0.20	0.26±0.19	-1.70±0.40	-2.35±0.23	-2.19±0.20	0.31±0.09	0.14±0.13	M
4372_76	4679±288	4850±229	4784±179	1.50±0.50	2.00±0.39	1.81±0.31	-1.90±0.25	-2.15±0.23	-2.04±0.17	0.42±0.12	0.23±0.11	M
4372_77	4993±365	5024± 74	5023± 73	2.20±0.80	3.20±0.33	3.05±0.31	-1.90±0.40	-2.50±0.10	-2.46±0.10	0.39±0.17	0.24±0.13	
Rup106_2	4830±151	4921±116	4887± 92	3.10±0.50	3.54±0.42	3.36±0.32	-0.14±0.14	-0.50±0.10	-0.38±0.08	0.12±0.11	0.26±0.11	
Rup106_3	4780±373	5000±250	4932±208	1.70±0.60	1.95±0.35	1.89±0.30	-1.60±0.40	-1.75±0.25	-1.71±0.21	0.43±0.16	0.10±0.08	M
Rup106_4	4341±121	4750±100	4584± 77	0.86±0.32	1.85±0.39	1.26±0.25	-1.56±0.22	-1.50±0.10	-1.51±0.09	0.42±0.13	0.16±0.14	
Rup106_5	5013±266	5350±123	5291±111	2.90±0.50	3.60±0.37	3.35±0.30	-0.47±0.27	-0.50±0.10	-0.50±0.09	0.26±0.20	0.25±0.14	
Rup106_6	4673±461	4800±245	4772±216	1.50±0.80	1.95±0.41	1.86±0.36	-1.60±0.50	-1.70±0.24	-1.68±0.22	0.40±0.20	0.14±0.11	M
Rup106_7	4713±159	4949±150	4838±109	2.80±0.50	3.25±0.33	3.11±0.28	-0.01±0.09	-0.15±0.23	-0.03±0.08	0.06±0.07	0.25±0.10	
Rup106_8	4864±198	5028±302	4913±166	2.80±0.50	2.81±0.45	2.81±0.33	-0.52±0.28	-0.90±0.37	-0.66±0.22	0.27±0.19	0.30±0.10	
Rup106_9	4638±286	4899±229	4797±179	1.50±0.60	1.95±0.41	1.81±0.34	-1.70±0.40	-1.85±0.23	-1.81±0.20	0.43±0.19	0.14±0.13	M
Rup106_10	4917±310	5152±122	5120±114	1.90±0.50	2.40±0.38	2.22±0.30	-1.54±0.34	-1.70±0.24	-1.65±0.20	0.43±0.15	0.09±0.07	M
Rup106_11	4417±193	4950±100	4838± 89	1.00±0.60	2.10±0.30	1.88±0.27	-1.61±0.20	-1.50±0.10	-1.52±0.09	0.51±0.09	0.14±0.13	M
Rup106_12	4856±391	5225± 75	5212± 73	1.80±0.60	2.20±0.25	2.14±0.23	-1.60±0.40	-1.50±0.10	-1.51±0.10	0.45±0.15	0.19±0.15	M
Rup106_14	4904±335	5250±100	5222± 96	1.90±0.60	2.55±0.47	2.30±0.37	-1.58±0.32	-1.50±0.10	-1.51±0.10	0.41±0.14	0.12±0.10	M

Table B.3 - continued.

NGC ID	$T_{eff}^{(a)}$ (K)	$T_{eff}^{(b)}$ (K)	$T_{eff}^{(avg)}$ (K)	$\log(g)^{(a)}$	$\log(g)^{(b)}$	$\log(g)^{(avg)}$	$[\text{Fe}/\text{H}]^{(a)}$	$[\text{Fe}/\text{H}]^{(b)}$	$[\text{Fe}/\text{H}]^{(avg)}$	$[\text{Mg}/\text{Fe}]^{(a)}$	$[\alpha/\text{Fe}]^{(b)}$	members
Rup106_16	5050± 6	5226±207	5050± 6	3.49±0.32	4.55±0.35	3.97±0.24	-0.10±0.04	-0.60±0.20	-0.12±0.04	0.24±0.07	0.12±0.11	M
Rup106_17	4803±448	5100±122	5079±118	1.70±0.70	2.20±0.40	2.08±0.35	-1.50±0.50	-1.50±0.10	-1.50±0.10	0.45±0.17	0.11±0.09	
Rup106_18	4761±355	5350±165	5246±150	1.90±0.90	3.35±0.39	3.12±0.36	-1.30±0.40	-1.05±0.15	-1.08±0.14	0.46±0.14	0.22±0.10	
5634_6	4815±349	4899±229	4874±191	1.80±0.60	2.00±0.39	1.94±0.33	-1.61±0.33	-1.85±0.23	-1.77±0.19	0.43±0.14	0.19±0.11	M
5634_7	4867±353	5000±112	4988±106	1.90±0.70	2.25±0.40	2.16±0.35	-1.70±0.40	-1.95±0.15	-1.92±0.14	0.42±0.14	0.19±0.14	M
5634_8	4902±344	5175±160	5127±145	2.00±0.70	2.15±0.32	2.12±0.29	-1.61±0.33	-1.60±0.20	-1.60±0.17	0.41±0.14	0.25±0.14	M
5634_9	4921±334	5100±122	5079±115	2.00±0.70	2.55±0.42	2.40±0.36	-1.61±0.31	-1.80±0.24	-1.73±0.19	0.42±0.13	0.22±0.13	M
5634_11	4389±177	4676±114	4591± 96	1.00±0.50	1.50±0.39	1.31±0.31	-1.55±0.20	-1.65±0.23	-1.59±0.15	0.45±0.14	0.24±0.12	M
5634_13	4981±345	5200±187	5150±164	2.10±0.80	2.70±0.46	2.55±0.40	-1.80±0.40	-1.90±0.20	-1.88±0.18	0.40±0.17	0.10±0.12	M
5634_14	4767±350	4775± 74	4774± 73	1.70±0.60	1.65±0.39	1.66±0.33	-1.60±0.33	-1.95±0.15	-1.89±0.14	0.45±0.14	0.24±0.11	M
5634_15	4338±119	4599±201	4406±102	0.85±0.32	1.45±0.52	1.01±0.27	-1.56±0.22	-1.60±0.20	-1.58±0.15	0.42±0.13	0.20±0.12	M
5634_17	4769± 56	5250±193	4806± 54	2.98±0.33	4.65±0.39	3.68±0.25	-0.01±0.07	-0.25±0.25	-0.03±0.07	0.16±0.13	0.10±0.08	
5694_3	4849± 56	4874±167	4852± 53	3.45±0.33	4.35±0.32	3.91±0.23	0.09±0.07	-0.60±0.20	0.01±0.07	-0.02±0.13	0.26±0.11	
5694_5	4877±340	4950±100	4944± 96	1.80±0.60	1.55±0.42	1.63±0.34	-1.69±0.33	-2.00±0.10	-1.97±0.10	0.42±0.12	0.14±0.11	M
5694_6	4972±360	5175±114	5157±109	2.10±0.70	2.10±0.30	2.10±0.28	-1.80±0.40	-2.00±0.10	-1.99±0.10	0.40±0.17	0.13±0.11	M
5694_7	4970±388	4925±114	4929±110	2.10±0.80	1.80±0.60	1.91±0.48	-1.80±0.40	-2.15±0.23	-2.06±0.20	0.40±0.18	0.18±0.12	M
5694_8	4981±330	5225± 75	5213± 73	2.10±0.80	2.20±0.25	2.19±0.24	-1.90±0.40	-2.00±0.10	-1.99±0.10	0.40±0.17	0.19±0.15	M
5694_9	4664±320	4750±100	4742± 95	1.50±0.60	1.45±0.42	1.47±0.34	-1.71±0.32	-2.00±0.10	-1.97±0.10	0.45±0.12	0.23±0.13	M
5694_10	5012±430	5175±114	5164±111	2.10±0.80	2.05±0.35	2.06±0.32	-1.80±0.40	-2.00±0.10	-1.99±0.10	0.41±0.17	0.10±0.10	
5694_11	4935±331	5175±115	5149±108	2.00±0.70	2.25±0.25	2.22±0.24	-1.80±0.40	-2.00±0.10	-1.99±0.10	0.39±0.17	0.15±0.14	M
5694_13	4643± 82	4874±125	4712± 69	3.00±0.40	4.30±0.25	3.93±0.21	0.30±0.19	0.05±0.20	0.18±0.14	0.18±0.07	0.10±0.08	
5694_15	4925±329	5200±100	5177± 96	2.00±0.70	2.20±0.25	2.18±0.24	-1.80±0.35	-2.00±0.10	-1.98±0.10	0.41±0.13	0.16±0.14	M
5694_17	4821±323	4875±125	4868±117	1.80±0.60	2.05±0.35	1.99±0.30	-1.68±0.28	-2.00±0.10	-1.96±0.09	0.41±0.11	0.20±0.13	M
5897_138	5078±420	5050±245	5057±212	2.20±0.80	2.05±0.42	2.08±0.37	-1.60±0.40	-2.20±0.24	-2.04±0.21	0.44±0.15	0.26±0.13	M
5897_142	4720±380	4400±164	4450±151	1.60±0.70	1.20±0.46	1.32±0.38	-1.60±0.40	-2.25±0.25	-2.07±0.21	0.43±0.16	0.28±0.13	M
5897_160	4587±335	4426±114	4442±108	1.30±0.60	1.50±0.39	1.44±0.33	-1.70±0.40	-2.15±0.23	-2.04±0.20	0.44±0.14	0.20±0.14	M
5897_255	4301± 42	4250±100	4293± 39	0.86±0.31	0.66±0.32	0.76±0.22	-1.54±0.22	-2.00±0.10	-1.92±0.09	0.38±0.13	0.23±0.15	M

Table B.3 - continued.

NGC ID	$T_{eff}^{(a)}$ (K)	$T_{eff}^{(b)}$ (K)	$T_{eff}^{(avg)}$ (K)	$\log(g)^{(a)}$	$\log(g)^{(b)}$	$\log(g)^{(avg)}$	[Fe/H] ^(a)	[Fe/H] ^(b)	[Fe/H] ^(avg)	[Mg/Fe] ^(a)	[α /Fe] ^(b)	members
5897_302	4583±218	4750±100	4721± 91	1.40±0.50	1.60±0.49	1.50±0.35	-1.87±0.31	-2.00±0.10	-1.99±0.10	0.42±0.14	0.24±0.11	M
5897_306	4971±473	4750±100	4759± 98	2.00±0.80	1.90±0.37	1.92±0.34	-1.50±0.40	-2.00±0.10	-1.97±0.10	0.43±0.17	0.19±0.14	M
5897_316	4590±310	4675±115	4665±108	1.30±0.60	1.80±0.56	1.57±0.41	-1.71±0.34	-2.00±0.10	-1.98±0.10	0.46±0.13	0.24±0.09	M
5897_318	4930±420	4799±148	4813±140	2.00±0.70	2.00±0.39	2.00±0.34	-1.50±0.40	-1.95±0.15	-1.89±0.14	0.43±0.15	0.22±0.13	M
5927_37	4432±120	4547± 98	4501±76	2.10±0.40	2.09±0.37	2.09±0.27	-0.10±0.21	-0.41±0.20	-0.26±0.14	0.08±0.16	0.20±0.14	M
5927_65	3927± 20	3997±159	3928±20	1.49±0.08	1.54±0.65	1.49±0.08	-0.18±0.03	-1.01±0.32	-0.19±0.03	0.01±0.09	0.33±0.08	M
5927_133	4018±117	4025±174	4020±97	1.51±0.32	1.60±0.62	1.53±0.28	-0.20±0.20	-1.00±0.31	-0.44±0.17	0.08±0.16	0.29±0.09	M
5927_372	3669± 42	3524± 73	3633±36	0.76±0.09	2.35±0.63	0.79±0.09	0.05±0.15	-0.95±0.15	-0.45±0.11	0.18±0.07	0.29±0.09	M
5927_335	3886±117	3924±160	3899±94	1.31±0.19	2.25±0.60	1.40±0.18	0.05±0.19	-0.65±0.32	-0.13±0.16	0.05±0.22	0.29±0.11	M
5927_190	3938± 29	3843±278	3937±29	1.49±0.07	1.08±0.94	1.49±0.07	-0.16±0.07	-1.16±0.39	-0.19±0.07	0.03±0.09	0.31±0.10	M
5946_1	4285±163	4450±186	4356±123	1.90±0.40	2.40±0.37	2.17±0.27	0.09±0.20	-0.16±0.29	0.01±0.16	0.03±0.13	0.19±0.13	M
5946_2	4796±229	5049±312	4885±185	2.50±0.50	2.80±0.51	2.65±0.36	-0.26±0.31	-0.55±0.41	-0.36±0.25	0.16±0.22	0.22±0.12	
5946_3	4414±175	4701± 99	4631± 86	2.20±0.50	2.90±0.20	2.80±0.19	0.05±0.13	-0.04±0.24	0.03±0.11	0.02±0.14	0.18±0.12	
5946_4	4960±487	4975±208	4973±191	2.10±0.90	1.45±0.52	1.61±0.45	-1.60±0.40	-1.85±0.23	-1.79±0.20	0.42±0.18	0.18±0.11	
5946_5	4725±418	4701±100	4702± 97	1.80±0.90	1.60±0.37	1.63±0.34	-1.40±0.50	-1.60±0.20	-1.57±0.19	0.40±0.16	0.21±0.14	M
5946_6	4866±451	4950±100	4946± 98	1.90±0.80	2.00±0.45	1.98±0.39	-1.60±0.40	-2.00±0.10	-1.98±0.10	0.41±0.18	0.25±0.10	M
5946_7	4619±325	4876±201	4805±171	1.50±0.60	1.95±0.42	1.80±0.34	-1.52±0.28	-1.60±0.20	-1.57±0.16	0.45±0.15	0.29±0.10	
5946_8	4743±418	4675±114	4680±110	1.90±0.90	1.60±0.37	1.64±0.34	-1.30±0.50	-1.65±0.23	-1.59±0.21	0.40±0.17	0.24±0.12	M
5946_9	4748±353	5000±100	4981± 96	1.70±0.70	2.10±0.37	2.01±0.33	-1.59±0.33	-1.50±0.10	-1.51±0.10	0.44±0.14	0.19±0.13	M
5946_10	4941±529	4675±160	4698±153	2.00±0.90	1.55±0.61	1.69±0.50	-1.50±0.50	-2.10±0.20	-2.02±0.19	0.42±0.18	0.24±0.14	M
5946_11	4919±474	4825±115	4830±111	2.00±0.90	1.90±0.37	1.91±0.34	-1.57±0.34	-2.00±0.10	-1.97±0.10	0.45±0.15	0.25±0.10	
5946_14	4794±430	4725± 75	4727± 74	1.80±0.90	1.45±0.42	1.51±0.38	-1.60±0.40	-2.00±0.10	-1.98±0.10	0.44±0.15	0.25±0.10	
5946_15	4693±353	4900±200	4850±174	1.60±0.70	1.75±0.34	1.72±0.31	-1.50±0.40	-1.60±0.20	-1.58±0.18	0.41±0.15	0.12±0.12	
5946_16	4878±468	4875±125	4875±121	1.90±0.90	0.50±0.39	0.72±0.36	-1.60±0.40	-2.00±0.10	-1.98±0.10	0.42±0.18	0.12±0.10	M
5946_19	4594±210	4899±165	4782±130	2.22±0.33	2.65±0.39	2.40±0.25	-0.21±0.23	-0.23±0.28	-0.22±0.18	0.09±0.15	0.28±0.12	
BH176_2	4664±241	4975±175	4868±142	2.40±0.50	2.85±0.23	2.77±0.21	-0.17±0.29	-0.06±0.32	-0.12±0.21	0.10±0.20	0.25±0.13	M
BH176_3	4551±247	5025±208	4829±159	2.30±0.50	2.90±0.44	2.64±0.33	-0.09±0.26	0.26±0.21	0.12±0.16	0.06±0.17	0.28±0.09	

Table B.3 - continued.

NGC ID	$T_{eff}^{(a)}$ (K)	$T_{eff}^{(b)}$ (K)	$T_{eff}^{(avg)}$ (K)	$\log(g)^{(a)}$	$\log(g)^{(b)}$	$\log(g)^{(avg)}$	[Fe/H] ^(a)	[Fe/H] ^(b)	[Fe/H] ^(avg)	[Mg/Fe] ^(a)	[α /Fe] ^(b)	members
BH176.4	3988±108	3600±122	3818± 81	1.50±0.40	2.35±0.23	2.14±0.20	-0.16±0.23	-2.00±0.10	-1.71±0.09	0.07±0.18	0.19±0.11	
BH176.6	4099±199	4275±208	4183±144	1.70±0.40	2.15±0.45	1.90±0.30	-0.04±0.21	-0.28±0.27	-0.13±0.17	0.05±0.16	0.18±0.11	M
BH176.7	5319±414	6175±115	6114±110	3.00±0.60	4.50±0.32	4.17±0.28	-0.25±0.32	0.04±0.08	0.02±0.08	0.14±0.15	0.07±0.06	M
BH176.8	3973±119	3575±114	3766± 83	1.45±0.30	1.75±0.90	1.48±0.28	-0.12±0.23	-1.85±0.23	-0.98±0.16	0.08±0.17	0.30±0.09	
BH176.9	4244±274	4525±175	4444±147	1.80±0.60	2.35±0.55	2.10±0.41	-0.02±0.19	-0.04±0.25	-0.03±0.15	0.03±0.15	0.21±0.09	M
BH176.10	3954± 78	4000±157	3963± 70	1.47±0.22	1.50±0.55	1.47±0.20	-0.03±0.18	-0.50±0.31	-0.15±0.16	0.05±0.19	0.22±0.12	M
BH176.11	3951±103	3950±149	3951± 85	1.45±0.22	1.40±0.62	1.44±0.21	-0.04±0.18	-0.60±0.30	-0.19±0.15	0.06±0.20	0.26±0.11	M
BH176.13	3640±100	3500±100	3570± 71	0.70±0.20	3.00±0.71	0.87±0.19	-0.10±0.06	-0.00±0.26	-0.09±0.06	0.23±0.10	0.34±0.07	M
BH176.14	4146±218	4400±122	4339±107	1.70±0.50	2.40±0.20	2.30±0.19	-0.05±0.20	-0.16±0.29	-0.09±0.16	0.06±0.15	0.15±0.12	M
BH176.15	4240±270	4350±123	4331±112	1.80±0.60	2.30±0.24	2.23±0.22	-0.07±0.23	-0.30±0.25	-0.18±0.17	0.05±0.17	0.15±0.14	M
BH176.16	4210±255	4425±114	4389±104	1.80±0.60	2.35±0.23	2.28±0.21	-0.08±0.25	-0.07±0.29	-0.08±0.19	0.06±0.18	0.26±0.11	M
BH176.17	3810± 12	3500±100	3806± 12	1.10±0.21	2.20±0.56	1.24±0.20	0.00±0.23	-1.90±0.20	-1.08±0.15	0.04±0.12	0.26±0.09	
BH176.18	4101±213	4274±208	4190±149	1.60±0.50	2.05±0.42	1.86±0.32	-0.06±0.22	-0.11±0.27	-0.08±0.17	0.06±0.17	0.18±0.10	M
Lynga7_1	4447±143	4425±195	4439±115	2.20±0.40	2.30±0.46	2.24±0.30	-0.43±0.17	-1.15±0.39	-0.54±0.16	0.20±0.16	0.27±0.11	
Lynga7_2	4130± 68	4202±186	4139± 64	1.85±0.23	1.75±0.46	1.83±0.21	-0.27±0.04	-1.10±0.37	-0.28±0.04	0.10±0.02	0.32±0.07	
Lynga7_3	4271±174	4348±122	4322±100	1.85±0.35	2.00±0.32	1.93±0.24	-0.02±0.14	-0.80±0.24	-0.22±0.12	0.03±0.16	0.29±0.09	
Lynga7_4	4459±156	4450±187	4455±120	2.20±0.40	2.40±0.49	2.28±0.31	-0.47±0.21	-1.10±0.37	-0.62±0.18	0.24±0.19	0.28±0.10	M
Lynga7_5	3640±100	3523± 73	3564± 59	0.70±0.20	3.09±0.77	0.85±0.19	-0.10±0.06	-0.56±0.27	-0.12±0.06	0.23±0.10	0.34±0.08	
Lynga7_6	4706±170	4950±149	4844±112	2.50±0.40	3.25±0.33	2.95±0.25	-0.39±0.22	-0.65±0.23	-0.51±0.16	0.21±0.20	0.29±0.10	
Lynga7_8	4361±110	4349±122	4356± 82	1.90±0.40	1.89±0.30	1.89±0.24	-0.38±0.28	-0.85±0.23	-0.66±0.18	0.21±0.20	0.25±0.10	M
Lynga7_9	4153±117	4324±114	4241± 82	1.58±0.14	2.10±0.37	1.65±0.13	0.01±0.11	-0.35±0.23	-0.06±0.10	-0.01±0.16	0.25±0.13	
Lynga7_10	4483±152	4600±122	4554± 95	2.20±0.40	2.55±0.35	2.40±0.26	-0.37±0.22	-0.80±0.24	-0.57±0.16	0.19±0.19	0.27±0.11	M
Lynga7_11	4186±150	4349±122	4284± 95	1.67±0.23	2.05±0.35	1.78±0.19	-0.06±0.13	-0.80±0.24	-0.23±0.11	0.01±0.16	0.28±0.10	
Lynga7_14	4711±236	4899±165	4837±135	2.50±0.50	2.95±0.27	2.85±0.24	-0.29±0.30	-0.75±0.25	-0.56±0.19	0.16±0.21	0.28±0.12	
Lynga7_15	3885± 67	3575±115	3806± 58	1.28±0.20	0.75±0.64	1.23±0.19	-0.20±0.29	-1.65±0.23	-1.09±0.18	0.15±0.18	0.29±0.10	
Lynga7_16	4434±156	4874±125	4702± 98	2.20±0.50	3.20±0.24	3.01±0.22	0.09±0.12	0.23±0.20	0.13±0.10	0.01±0.14	0.09±0.07	
Pal14_3	4657±367	4900±122	4876±116	1.70±0.90	1.75±0.51	1.74±0.44	-1.10±0.50	-1.20±0.24	-1.18±0.22	0.36±0.15	0.22±0.14	M
Pal14_4	4654±381	4850±122	4832±117	1.80±1.00	2.60±0.37	2.50±0.35	-1.10±0.40	-1.30±0.24	-1.25±0.21	0.40±0.14	0.21±0.11	M

Table B.3 - continued.

NGC ID	$T_{eff}^{(a)}$ (K)	$T_{eff}^{(b)}$ (K)	$T_{eff}^{(avg)}$ (K)	$\log(g)^{(a)}$	$\log(g)^{(b)}$	$\log(g)^{(avg)}$	[Fe/H] ^(a)	[Fe/H] ^(b)	[Fe/H] ^(avg)	[Mg/Fe] ^(a)	[α /Fe] ^(b)	members
Pal14_5	4815±422	5025±175	4994±162	1.90±0.90	2.35±0.55	2.23±0.47	-1.30±0.50	-1.35±0.23	-1.34±0.21	0.42±0.17	0.26±0.11	M
Pal14_7	4420±324	4650±122	4622±115	1.50±0.70	2.15±0.32	2.04±0.29	-0.90±0.40	-1.20±0.24	-1.12±0.21	0.30±0.17	0.28±0.10	M
Pal14_8	4805±380	4950±245	4907±206	2.00±0.90	2.05±0.42	2.04±0.38	-1.30±0.50	-1.30±0.24	-1.30±0.22	0.41±0.16	0.22±0.10	M
Pal14_10	4849±420	4551±150	4585±141	3.45±0.04	4.25±0.46	3.46±0.04	0.09±0.01	-1.35±0.23	0.089±0.007	-0.02±0.10	0.34±0.07	
Pal14_11	4296± 13	4599±166	4298± 13	1.05±0.10	2.05±0.35	1.13±0.10	-0.90±0.40	-1.25±0.25	-1.15±0.21	0.13±0.12	0.24±0.12	M
6284_1	4696±201	4750±158	4729±124	2.50±0.40	2.65±0.39	2.58±0.28	-0.16±0.24	-0.50±0.32	-0.282±0.19	0.09±0.17	0.25±0.08	
6284_2	4488±182	4447±216	4471±139	2.10±0.50	1.80±0.45	1.93±0.33	-0.51±0.20	-1.15±0.32	-0.690±0.17	0.22±0.18	0.27±0.11	
6284_3	4919±124	5398±165	5092± 99	2.80±0.40	3.65±0.23	3.44±0.20	0.01±0.10	0.23±0.19	0.055±0.09	0.04±0.10	0.08±0.07	
6284_4	4903±189	5151±122	5078±103	2.60±0.40	3.20±0.25	3.03±0.21	-0.47±0.23	-0.50±0.10	-0.495±0.09	0.25±0.19	0.21±0.11	
6284_5	4676±188	4898±123	4832±103	2.60±0.40	3.15±0.23	3.01±0.20	-0.24±0.19	-0.35±0.23	-0.285±0.15	0.12±0.15	0.26±0.13	
6284_6	4727±181	4750±100	4745± 88	2.30±0.40	2.05±0.41	2.18±0.29	-0.49±0.24	-1.00±0.10	-0.925±0.09	0.29±0.24	0.25±0.10	
6284_7	4935±319	4801±245	4851±194	2.50±0.70	1.95±0.42	2.10±0.36	-0.61±0.34	-1.20±0.24	-1.004±0.20	0.24±0.12	0.25±0.10	M
6284_8	4602±167	4625±125	4616±100	2.41±0.29	2.10±0.30	2.26±0.21	-0.33±0.16	-0.75±0.25	-0.452±0.13	0.14±0.17	0.28±0.12	
6284_9	4469±184	4975±208	4692±138	1.30±0.50	2.10±0.44	1.75±0.33	-1.40±0.40	-1.35±0.23	-1.362±0.20	0.42±0.11	0.30±0.10	M
6284_10	4668±276	4875±125	4840±114	1.80±0.80	2.55±0.35	2.43±0.32	-1.10±0.40	-1.25±0.25	-1.208±0.21	0.36±0.17	0.31±0.09	M
6284_11	4824±175	4800±187	4813±128	2.60±0.40	2.60±0.44	2.60±0.30	-0.48±0.21	-0.90±0.37	-0.582±0.18	0.28±0.20	0.28±0.10	
6284_13	4793±275	4825±225	4812±174	2.20±0.70	2.20±0.56	2.20±0.44	-0.80±0.40	-1.20±0.24	-1.094±0.21	0.28±0.14	0.28±0.10	M
6284_14	4821±369	5199±100	5173± 97	2.20±0.90	3.05±0.35	2.94±0.33	-0.90±0.50	-1.00±0.10	-0.996±0.10	0.34±0.17	0.22±0.12	M
6284_15	4933±421	4950±218	4946±194	2.40±0.90	2.60±0.37	2.57±0.34	-0.80±0.50	-1.30±0.24	-1.206±0.22	0.33±0.20	0.25±0.10	M
6284_16	4165±140	4473±208	4261±116	1.70±0.40	2.60±0.66	1.94±0.34	-0.40±0.20	-0.85±0.23	-0.594±0.15	0.16±0.18	0.34±0.07	
6284_17	4633±140	5000±157	4795±105	2.47±0.34	3.00±0.31	2.76±0.23	-0.13±0.18	-0.02±0.25	-0.092±0.15	0.06±0.12	0.23±0.13	
6284_18	4948±259	4875±202	4903±159	2.70±0.60	2.50±0.55	2.59±0.41	-0.57±0.30	-1.15±0.23	-0.935±0.18	0.29±0.20	0.28±0.10	M
6316_1	4137±161	4375±230	4215±132	1.76±0.28	2.10±0.58	1.82±0.25	-0.07±0.19	-0.41±0.34	-0.15±0.17	0.02±0.13	0.19±0.11	
6316_2	4274±159	4373±125	4335± 98	1.90±0.50	2.10±0.37	2.03±0.30	-0.32±0.26	-0.80±0.24	-0.58±0.18	0.16±0.19	0.31±0.10	
6316_3	4817±224	4901±229	4858±160	2.50±0.60	2.90±0.44	2.76±0.35	-0.51±0.24	-0.80±0.33	-0.61±0.19	0.28±0.19	0.28±0.12	
6316_4	4586±190	4675±114	4652± 98	2.20±0.40	2.65±0.32	2.47±0.25	-0.21±0.21	-0.65±0.23	-0.41±0.16	0.11±0.16	0.27±0.09	
6316_5	4233±128	4376±125	4306± 89	1.78±0.24	2.15±0.32	1.91±0.19	0.12±0.18	-0.25±0.25	-0.01±0.15	-0.00±0.11	0.20±0.13	
6316_6	4698±192	4801±218	4743±144	2.50±0.50	2.85±0.59	2.65±0.38	-0.41±0.24	-0.75±0.25	-0.57±0.17	0.22±0.20	0.30±0.08	

Table B.3 - continued.

NGC ID	$T_{eff}^{(a)}$ (K)	$T_{eff}^{(b)}$ (K)	$T_{eff}^{(avg)}$ (K)	$\log(g)^{(a)}$	$\log(g)^{(b)}$	$\log(g)^{(avg)}$	[Fe/H] ^(a)	[Fe/H] ^(b)	[Fe/H] ^(avg)	[Mg/Fe] ^(a)	[α /Fe] ^(b)	members
6316_7	4529±195	4474±208	4503±142	2.30±0.50	1.90±0.44	2.07±0.33	-0.19±0.28	-0.80±0.24	-0.54±0.18	0.11±0.18	0.27±0.13	M
6316_8	4634±185	4699±218	4661±141	2.40±0.40	2.35±0.63	2.39±0.34	-0.43±0.22	-0.80±0.24	-0.60±0.16	0.21±0.20	0.30±0.09	
6316_9	3980±119	3924±194	3965±101	1.40±0.40	1.00±0.74	1.31±0.35	-0.35±0.14	-1.20±0.33	-0.48±0.13	0.17±0.21	0.30±0.10	M
6316_10	4388±143	4399±165	4393±108	2.00±0.40	2.09±0.44	2.04±0.30	-0.35±0.27	-0.85±0.23	-0.64±0.18	0.20±0.19	0.32±0.07	M
6316_11	4393±116	4527±175	4434± 97	2.00±0.40	2.20±0.40	2.10±0.28	-0.50±0.21	-0.85±0.23	-0.66±0.16	0.23±0.18	0.28±0.07	M
6316_14	4157±162	4201±186	4176±122	1.60±0.40	1.75±0.46	1.66±0.30	-0.35±0.23	-1.10±0.37	-0.56±0.20	0.11±0.17	0.32±0.07	M
6316_15	4285±200	4500±100	4457± 89	1.90±0.50	2.30±0.24	2.23±0.22	0.14±0.18	0.10±0.10	0.11±0.09	0.03±0.12	0.16±0.10	
6316_17	4147±156	4223±175	4181±117	1.70±0.25	1.90±0.37	1.76±0.21	0.01±0.18	-0.70±0.25	-0.24±0.15	0.01±0.12	0.31±0.10	M
6316_18	4515±136	4649±123	4589± 91	2.29±0.35	2.25±0.51	2.28±0.29	-0.23±0.26	-0.70±0.25	-0.47±0.18	0.09±0.16	0.30±0.08	M
6316_19	4413±133	4874±125	4658± 91	2.10±0.50	2.95±0.35	2.67±0.29	0.13±0.14	0.35±0.15	0.23±0.10	0.02±0.16	0.17±0.09	
6356_1	4698±214	4849±255	4761±164	2.30±0.50	2.40±0.62	2.34±0.39	-0.43±0.24	-0.65±0.32	-0.51±0.19	0.19±0.21	0.30±0.08	M
6356_2	4533± 93	4648±199	4554± 84	2.42±0.32	2.65±0.55	2.48±0.28	0.01±0.10	-0.30±0.24	-0.04±0.09	0.08±0.12	0.24±0.09	
6356_3	4822±395	5274±175	5200±160	2.20±1.00	3.35±0.32	3.24±0.30	-1.20±0.50	-1.10±0.20	-1.11±0.19	0.43±0.16	0.31±0.07	
6356_4	3959± 87	3926±114	3947± 69	1.47±0.32	1.40±0.62	1.46±0.28	-0.22±0.21	-1.15±0.23	-0.64±0.16	0.13±0.16	0.25±0.10	
6356_5	4379± 97	4549±187	4415± 86	2.00±0.40	2.40±0.54	2.14±0.32	-0.50±0.20	-0.80±0.24	-0.62±0.15	0.25±0.20	0.32±0.09	M
6356_6	4724±195	4827±194	4776±138	2.40±0.50	2.46±0.52	2.43±0.36	-0.47±0.23	-0.75±0.25	-0.60±0.17	0.25±0.20	0.29±0.09	M
6356_7	4344±179	4347±122	4346±101	2.00±0.40	1.99±0.32	1.99±0.25	-0.29±0.24	-0.85±0.23	-0.58±0.17	0.16±0.18	0.29±0.09	M
6356_8	4001±101	4075±115	4033± 76	1.50±0.40	1.90±0.38	1.71±0.28	-0.30±0.18	-1.25±0.25	-0.62±0.15	0.11±0.17	0.32±0.07	M
6356_9	4369± 91	4496±194	4392± 82	2.00±0.40	2.19±0.56	2.06±0.33	-0.51±0.20	-0.75±0.25	-0.60±0.16	0.27±0.20	0.32±0.10	M
6356_10	4814±192	4829±224	4820±146	2.50±0.40	2.01±0.39	2.25±0.28	-0.24±0.30	-0.70±0.24	-0.52±0.19	0.12±0.23	0.29±0.10	M
6356_11	4322±125	4476±208	4363±107	2.10±0.40	2.40±0.49	2.22±0.31	0.06±0.11	-0.29±0.32	0.02±0.10	-0.02±0.16	0.26±0.10	
6356_13	4182±186	4255±194	4217±134	1.65±0.26	1.96±0.52	1.71±0.23	-0.06±0.19	-0.94±0.35	-0.26±0.17	0.03±0.12	0.31±0.08	M
6356_14	4232±182	4327±159	4286±120	1.90±0.40	1.96±0.47	1.93±0.30	-0.11±0.20	-0.85±0.32	-0.32±0.17	0.03±0.13	0.30±0.10	M
6356_15	3922± 48	3950±186	3924± 46	1.54±0.04	1.50±0.71	1.54±0.04	0.17±0.14	-0.58±0.41	0.09±0.13	0.17±0.05	0.34±0.07	
6356_16	4711±183	4801±187	4755±131	2.40±0.50	2.41±0.54	2.40±0.37	-0.49±0.22	-0.80±0.25	-0.63±0.17	0.25±0.20	0.31±0.10	M
6356_17	3861± 96	3549± 99	3710± 69	1.24±0.13	0.90±0.62	1.23±0.13	0.07±0.17	-1.40±0.20	-0.55±0.13	0.09±0.07	0.30±0.08	M
6356_18	4419±205	4349±165	4377±128	2.20±0.50	2.05±0.47	2.12±0.34	-0.26±0.29	-1.00±0.22	-0.73±0.18	0.14±0.18	0.28±0.10	M
6356_19	4350± 80	4499±194	4372± 74	2.00±0.40	2.10±0.49	2.04±0.31	-0.52±0.20	-0.70±0.25	-0.59±0.16	0.24±0.17	0.29±0.09	M

Table B.3 - continued.

NGC ID	$T_{eff}^{(a)}$ (K)	$T_{eff}^{(b)}$ (K)	$T_{eff}^{(avg)}$ (K)	$\log(g)^{(a)}$	$\log(g)^{(b)}$	$\log(g)^{(avg)}$	[Fe/H] ^(a)	[Fe/H] ^(b)	[Fe/H] ^(avg)	[Mg/Fe] ^(a)	[α /Fe] ^(b)	members
6355_1	4128±122	4401±122	4264± 86	1.71±0.20	2.05±0.35	1.79±0.17	0.05±0.20	-0.20±0.24	-0.05±0.15	0.06±0.07	0.28±0.10	
6355_2	4538±128	4601±229	4553±112	2.40±0.40	2.50±0.55	2.43±0.32	-0.10±0.21	-0.70±0.33	-0.27±0.18	0.07±0.14	0.28±0.10	
6355_3	4638±208	4700±292	4659±169	2.30±0.50	2.60±0.73	2.40±0.41	-0.34±0.32	-0.90±0.37	-0.58±0.24	0.19±0.22	0.33±0.06	
6355_4	4404±142	4800±100	4669± 82	2.00±0.40	2.75±0.25	2.54±0.21	0.12±0.14	0.18±0.18	0.14±0.11	0.03±0.16	0.16±0.11	
6355_5	4292±139	4449±187	4348±112	1.80±0.40	2.40±0.58	1.99±0.33	-0.50±0.21	-1.00±0.32	-0.65±0.18	0.25±0.20	0.33±0.06	
6355_6	4490±159	4600±123	4559± 97	2.20±0.40	2.60±0.37	2.42±0.27	-0.44±0.20	-0.80±0.25	-0.58±0.16	0.22±0.18	0.25±0.10	
6355_7	4884±459	5125±168	5097±158	2.00±0.90	2.60±0.37	2.51±0.34	-1.50±0.40	-1.35±0.23	-1.39±0.20	0.44±0.16	0.28±0.10	M
6355_8	4304± 7	4672±224	4304± 7	1.04±0.10	1.94±0.42	1.09±0.10	-1.37±0.10	-1.35±0.23	-1.37±0.09	0.26±0.01	0.21±0.15	M
6355_9	4422±143	4725± 75	4660± 66	1.13±0.29	1.70±0.33	1.38±0.22	-1.45±0.27	-1.55±0.15	-1.53±0.13	0.41±0.11	0.21±0.15	M
6355_10	4705±420	4950±187	4910±171	1.40±0.60	1.95±0.35	1.81±0.30	-1.60±0.40	-1.70±0.24	-1.67±0.21	0.46±0.14	0.32±0.09	M
6355_11	4649±388	4475±174	4504±159	1.90±0.90	1.40±0.54	1.53±0.46	-1.00±0.50	-1.65±0.23	-1.54±0.21	0.36±0.16	0.25±0.14	M
6355_13	4685±380	4725±134	4720±127	1.70±0.90	1.80±0.33	1.79±0.31	-1.40±0.50	-1.60±0.20	-1.57±0.19	0.41±0.14	0.28±0.10	M
6355_14	4795±158	4902±122	4862± 97	2.60±0.40	3.10±0.30	2.92±0.24	-0.41±0.23	-0.70±0.24	-0.55±0.17	0.25±0.19	0.25±0.10	
6355_15	4913±262	5250±274	5074±189	2.70±0.50	3.40±0.37	3.15±0.30	-0.36±0.31	-0.85±0.32	-0.60±0.22	0.20±0.22	0.35±0.07	
6355_17	4707±322	4925±195	4867±167	1.80±0.80	3.00±0.32	2.83±0.30	-1.30±0.40	-1.75±0.25	-1.62±0.21	0.41±0.15	0.29±0.09	
6355_18	4739±232	4724±175	4729±140	2.50±0.50	2.20±0.56	2.37±0.37	-0.28±0.29	-0.90±0.20	-0.70±0.16	0.16±0.21	0.29±0.09	
6352_1	4736±208	5074±194	4917±142	2.60±0.50	3.20±0.33	3.02±0.28	-0.17±0.25	-0.45±0.27	-0.30±0.18	0.12±0.17	0.31±0.10	M
6352_2	4723±190	5126±229	4887±146	2.60±0.40	3.10±0.43	2.83±0.29	-0.15±0.19	-0.20±0.33	-0.16±0.16	0.09±0.14	0.27±0.09	
6352_5	4384±205	4425±196	4405±142	2.10±0.50	2.20±0.51	2.15±0.36	-0.13±0.25	-0.85±0.23	-0.52±0.17	0.09±0.19	0.32±0.07	M
6352_6	4766±227	4899±199	4841±150	2.50±0.60	3.15±0.32	3.01±0.28	-0.48±0.25	-0.90±0.30	-0.65±0.19	0.26±0.20	0.29±0.09	M
6352_8	4149±104	4347±122	4232± 79	1.60±0.50	2.05±0.35	1.90±0.29	-0.48±0.27	-0.86±0.23	-0.70±0.18	0.18±0.18	0.30±0.10	M
6352_9	4810±163	5002±192	4890±124	2.60±0.40	3.05±0.35	2.85±0.26	-0.42±0.24	-0.65±0.23	-0.54±0.17	0.23±0.20	0.25±0.13	M
6352_10	4391±111	4625±167	4463± 92	2.00±0.50	2.65±0.45	2.36±0.33	-0.51±0.21	-0.80±0.25	-0.63±0.16	0.25±0.20	0.27±0.11	M
6352_11	3921± 48	3699±217	3911± 47	1.35±0.24	0.90±0.73	1.31±0.23	-0.34±0.18	-1.50±0.39	-0.54±0.16	0.15±0.19	0.32±0.07	M
6352_13	4629±178	4775±209	4691±135	2.44±0.35	2.85±0.60	2.54±0.30	-0.39±0.18	-0.65±0.23	-0.49±0.14	0.19±0.18	0.27±0.09	M
6352_14	4550±264	4225±175	4324±146	2.20±0.60	2.80±0.51	2.55±0.39	-0.50±0.31	-2.15±0.32	-1.30±0.22	0.28±0.20	0.34±0.07	
6352_15	4694±202	4875±167	4801±129	2.40±0.50	3.10±0.30	2.91±0.26	-0.43±0.25	-0.75±0.33	-0.55±0.20	0.23±0.20	0.28±0.10	M
6352_16	4364± 98	4625±125	4464± 77	1.90±0.50	2.25±0.40	2.11±0.31	-0.52±0.24	-0.75±0.25	-0.63±0.17	0.27±0.21	0.34±0.07	M

Table B.3 - continued.

NGC ID	$T_{eff}^{(a)}$ (K)	$T_{eff}^{(b)}$ (K)	$T_{eff}^{(avg)}$ (K)	$\log(g)^{(a)}$	$\log(g)^{(b)}$	$\log(g)^{(avg)}$	[Fe/H] ^(a)	[Fe/H] ^(b)	[Fe/H] ^(avg)	[Mg/Fe] ^(a)	[α /Fe] ^(b)	members
6352_17	3984± 15	4351±123	3989± 15	1.13±0.18	2.20±0.33	1.38±0.16	-0.44±0.10	-0.80±0.25	-0.49±0.09	0.03±0.09	0.26±0.11	M
6352_18	4390±106	4676±114	4522± 78	2.00±0.40	2.00±0.39	2.00±0.28	-0.44±0.25	-0.65±0.23	-0.55±0.17	0.21±0.20	0.32±0.07	M
6366_1	4849±293	5000±158	4966±139	2.50±0.60	3.25±0.25	3.14±.23	-0.48±0.33	-0.85±0.23	-0.73±0.19	0.24±0.21	0.27±0.09	M
6366_2	4976±355	5250±193	5187±170	2.60±0.70	3.15±0.23	3.10±.22	-0.60±0.40	-0.65±0.23	-0.64±0.20	0.26±0.20	0.23±0.13	M
6366_3	4876±279	5150±166	5079±142	2.60±0.60	3.30±0.33	3.14±.29	-0.40±0.32	-0.60±0.20	-0.54±0.17	0.21±0.21	0.32±0.07	M
6366_4	4874±256	4950±187	4923±151	2.70±0.50	3.20±0.25	3.10±.22	-0.40±0.27	-0.80±0.24	-0.62±0.18	0.23±0.21	0.29±0.09	M
6366_5	4601±171	4622±125	4615±101	2.39±0.32	2.10±0.37	2.27±.24	-0.31±0.20	-0.76±0.25	-0.49±0.16	0.14±0.17	0.27±0.11	M
6366_6	5079±365	5276±208	5228±181	2.70±0.70	2.65±0.45	2.66±.38	-0.45±0.32	-0.70±0.24	-0.61±0.19	0.22±0.18	0.19±0.14	
6366_7	4882±319	4750±250	4800±197	2.60±0.50	2.85±0.45	2.74±.33	-0.35±0.28	-1.10±0.37	-0.62±0.22	0.19±0.21	0.31±0.10	M
6366_8	4860±293	5049±187	4995±158	2.60±0.70	3.30±0.33	3.17±.30	-0.50±0.34	-0.95±0.35	-0.72±0.24	0.26±0.21	0.34±0.07	M
6366_9	4526±153	4624±200	4562±122	2.30±0.30	2.59±0.58	2.36±.27	-0.40±0.16	-0.80±0.33	-0.48±0.14	0.21±0.10	0.26±0.11	M
6366_10	4864±325	5175±160	5114±143	2.50±0.70	3.30±0.25	3.21±.24	-0.40±0.40	-0.70±0.24	-0.62±0.21	0.22±0.21	0.32±0.07	M
6366_11	4748±223	4776±175	4765±137	2.50±0.50	2.80±0.24	2.74±.22	-0.41±0.27	-0.95±0.35	-0.61±0.21	0.22±0.19	0.29±0.09	M
6366_13	4838±237	4750±193	4785±150	2.60±0.50	2.95±0.35	2.83±.29	-0.41±0.26	-1.10±0.30	-0.71±0.20	0.22±0.21	0.26±0.13	M
6366_14	4819±350	4725±208	4750±179	2.40±0.80	3.25±0.46	3.04±.40	-0.70±0.40	-1.75±0.25	-1.46±0.21	0.32±0.19	0.33±0.08	
6366_15	4630±223	4600±279	4618±174	2.30±0.50	2.30±0.64	2.30±.39	-0.37±0.28	-1.05±0.35	-0.64±0.22	0.21±0.21	0.32±0.07	M
6366_16	4843±252	4974±175	4932±144	2.60±0.60	3.20±0.24	3.12±.22	-0.43±0.30	-0.75±0.25	-0.62±0.19	0.22±0.21	0.30±0.10	M
6366_17	4768±201	4976±261	4846±159	2.60±0.40	2.90±0.49	2.72±.31	-0.24±0.25	-0.45±0.35	-0.31±0.20	0.14±0.16	0.29±0.09	
6366_18	4376±115	4474±175	4406± 96	2.00±0.50	2.25±0.51	2.12±.36	-0.55±0.23	-1.05±0.35	-0.70±0.19	0.28±0.22	0.31±0.07	M
6401_1	4512±219	4625±167	4583±133	2.30±0.50	2.75±0.40	2.57±0.31	-0.08±0.24	-0.80±0.25	-0.43±0.17	0.07±0.17	0.27±0.11	
6401_2	4610±169	4701±186	4651±125	2.50±0.40	2.90±0.37	2.72±0.27	-0.08±0.20	-0.70±0.24	-0.33±0.15	0.04±0.14	0.24±0.14	
6401_3	4932±330	4726± 74	4736± 72	2.20±0.70	1.70±0.33	1.79±0.30	-0.80±0.40	-1.55±0.15	-1.46±0.14	0.38±0.16	0.20±0.14	M
6401_4	3989± 80	4098±122	4022± 67	1.67±0.11	1.94±0.35	1.69±0.10	0.11±0.09	-0.50±0.10	-0.16±0.07	0.08±0.09	0.27±0.11	
6401_5	4551±124	4950±100	4792± 78	2.30±0.40	3.20±0.24	2.96±0.21	0.03±0.13	0.15±0.15	0.08±0.10	0.03±0.12	0.11±0.09	
6401_6	4625±237	5072±336	4773±194	2.20±0.40	2.39±0.62	2.26±0.34	-0.35±0.30	-0.60±0.44	-0.43±0.25	0.21±0.25	0.30±0.08	
6401_7	4841±310	4975±261	4919±200	2.40±0.80	2.60±0.58	2.53±0.47	-0.60±0.40	-0.95±0.35	-0.80±0.26	0.32±0.19	0.29±0.09	M
6401_8	4759±232	4675±159	4702±131	2.40±0.60	2.20±0.40	2.26±0.33	-0.55±0.26	-1.15±0.32	-0.79±0.20	0.29±0.19	0.29±0.09	M
6401_9	4856±219	4700±100	4727± 91	2.40±0.50	1.70±0.24	1.83±0.22	-0.46±0.30	-1.10±0.20	-0.90±0.17	0.21±0.20	0.25±0.10	M

Table B.3 - continued.

NGC ID	$T_{eff}^{(a)}$ (K)	$T_{eff}^{(b)}$ (K)	$T_{eff}^{(avg)}$ (K)	$\log(g)^{(a)}$	$\log(g)^{(b)}$	$\log(g)^{(avg)}$	[Fe/H] ^(a)	[Fe/H] ^(b)	[Fe/H] ^(avg)	[Mg/Fe] ^(a)	[α /Fe] ^(b)	members
6401_10	4259±159	4349±122	4316± 97	1.83±0.32	2.24±0.40	1.99±0.25	-0.01±0.14	-0.80±0.24	-0.21±0.12	-0.02±0.17	0.33±0.08	M
6401_11	4392±159	4475±208	4423±126	2.10±0.40	2.35±0.55	2.19±0.32	0.13±0.14	-0.11±0.35	0.10±0.13	0.05±0.16	0.09±0.08	
6401_12	4520±258	4025±208	4220±162	1.90±0.60	0.05±0.15	0.16±0.15	-0.60±0.40	-2.20±0.24	-1.78±0.21	0.32±0.24	0.14±0.13	
6401_13	4858±426	5025±175	5001±162	2.00±0.90	2.30±0.60	2.21±0.50	-1.20±0.50	-1.35±0.23	-1.32±0.21	0.39±0.17	0.29±0.09	
6401_14	4869±200	4599±199	4733±141	2.50±0.50	2.00±0.45	2.22±0.33	-0.46±0.29	-1.40±0.20	-1.10±0.16	0.25±0.22	0.23±0.13	
6401_15	4475±200	4451±186	4462±136	2.30±0.50	2.25±0.51	2.28±0.36	-0.09±0.25	-1.10±0.37	-0.41±0.21	0.07±0.16	0.33±0.06	
6401_16	4546±166	4749±159	4652±115	2.30±0.40	2.60±0.44	2.44±0.30	-0.27±0.26	-0.50±0.32	-0.36±0.20	0.14±0.19	0.32±0.07	
6401_17	4554±181	4600±278	4568±152	2.40±0.50	2.45±0.65	2.42±0.40	-0.17±0.26	-0.65±0.39	-0.32±0.22	0.10±0.17	0.30±0.10	
6401_18	4339±230	4275±132	4291±115	2.00±0.60	1.85±0.32	1.88±0.28	-0.09±0.25	-1.00±0.22	-0.60±0.17	0.06±0.17	0.26±0.11	
Pal6_1	4240±163	4325±159	4283±114	1.90±0.50	1.95±0.41	1.93±0.32	-0.24±0.27	-0.85±0.32	-0.49±0.21	0.10±0.19	0.31±0.09	M
Pal6_2	4828±301	5150±278	5002±204	2.40±0.70	3.35±0.55	2.99±0.43	-0.50±0.40	-0.75±0.34	-0.64±0.26	0.25±0.22	0.34±0.07	
Pal6_3	4900±155	5249±158	5071±111	3.23±0.35	4.25±0.25	3.91±0.20	-0.03±0.09	-0.25±0.25	-0.05±0.08	0.07±0.08	0.11±0.08	
Pal6_4	4383±177	4475±208	4422±135	1.90±0.40	1.95±0.42	1.92±0.29	0.09±0.18	-0.14±0.30	0.03±0.15	0.09±0.12	0.16±0.11	
Pal6_5	4561±199	4475±208	4520±144	2.10±0.40	1.90±0.44	2.01±0.30	-0.17±0.21	-0.85±0.23	-0.48±0.16	0.12±0.14	0.24±0.12	
Pal6_6	4037±146	3875±230	3991±123	1.57±0.31	1.55±0.61	1.57±0.28	-0.12±0.21	-1.30±0.40	-0.37±0.19	0.06±0.16	0.26±0.09	
Pal6_7	4042±144	4150±166	4088±109	1.50±0.40	1.75±0.51	1.60±0.31	-0.10±0.20	-0.55±0.27	-0.26±0.16	0.05±0.17	0.27±0.09	
Pal6_8	4372±211	4498±224	4431±154	2.10±0.50	2.10±0.58	2.10±0.38	0.04±0.15	-0.16±0.29	-0.00±0.13	0.02±0.12	0.23±0.13	
Pal6_9	4273±259	3850±199	4007±158	1.90±0.60	0.40±0.54	1.07±0.40	-0.08±0.25	-1.55±0.27	-0.76±0.18	0.07±0.17	0.27±0.15	
Pal6_10	3989±124	4500±100	4299± 78	1.67±0.01	2.30±0.24	1.67±0.01	0.11±0.02	0.10±0.10	0.11±0.02	0.08±0.21	0.30±0.10	M
Pal6_11	4377±149	3899±200	4206±119	2.00±0.50	0.55±0.52	1.30±0.36	-0.45±0.23	-1.70±0.40	-0.76±0.20	0.22±0.20	0.23±0.13	
Pal6_13	4691±269	4150±255	4406±185	2.40±0.70	1.10±0.73	1.78±0.51	-0.40±0.40	-1.85±0.39	-1.14±0.28	0.21±0.21	0.32±0.10	
Pal6_14	4469±226	3925±160	4106±130	2.20±0.60	0.30±0.46	1.00±0.37	-0.17±0.32	-1.65±0.32	-0.91±0.23	0.09±0.20	0.27±0.09	
Pal6_15	4407±152	4350±123	4373± 95	1.70±0.40	1.95±0.42	1.82±0.29	-0.13±0.10	-0.70±0.24	-0.21±0.09	0.14±0.10	0.33±0.06	
Pal6_16	4339±228	4275±134	4291±115	2.10±0.60	1.70±0.46	1.85±0.37	0.02±0.18	-0.95±0.27	-0.28±0.15	0.03±0.14	0.27±0.11	
Pal6_17	4352±247	4150±165	4212±137	2.00±0.60	1.30±0.56	1.63±0.41	-0.10±0.28	-1.20±0.33	-0.56±0.21	0.07±0.18	0.32±0.07	
Pal6_18	4353±259	4051±187	4154±152	2.00±0.60	1.00±0.71	1.58±0.46	-0.08±0.26	-1.40±0.37	-0.52±0.21	0.06±0.17	0.33±0.06	
6440_1	4023±118	4249±222	4073±104	1.56±0.16	2.05±0.56	1.60±0.15	-0.02±0.15	-0.53±0.38	-0.09±0.14	0.04±0.10	0.13±0.12	
6440_3	4733±136	5250±156	4956±103	2.51±0.25	3.25±0.25	2.88±0.18	-0.20±0.19	-0.01±0.18	-0.10±0.13	0.11±0.17	0.22±0.12	

Table B.3 - continued.

NGC ID	$T_{eff}^{(a)}$ (K)	$T_{eff}^{(b)}$ (K)	$T_{eff}^{(avg)}$ (K)	$\log(g)^{(a)}$	$\log(g)^{(b)}$	$\log(g)^{(avg)}$	$[\text{Fe}/\text{H}]^{(a)}$	$[\text{Fe}/\text{H}]^{(b)}$	$[\text{Fe}/\text{H}]^{(avg)}$	$[\text{Mg}/\text{Fe}]^{(a)}$	$[\alpha/\text{Fe}]^{(b)}$	members
6440_4	4899±232	5151±165	5066±135	2.60±0.50	3.15±0.45	2.90±0.33	-0.46±0.24	-0.55±0.15	-0.52±0.13	0.25±0.19	0.22±0.12	
6440_5	3946± 27	4074±114	3953± 26	1.53±0.13	1.90±0.30	1.59±0.12	-0.06±0.22	-0.85±0.23	-0.44±0.16	0.06±0.08	0.27±0.11	
6440_6	4270±142	4351±123	4316± 93	1.82±0.27	2.15±0.32	1.96±0.21	0.11±0.18	-0.30±0.25	-0.03±0.15	0.02±0.10	0.22±0.15	M
6440_7	4399±149	4375±125	4385± 96	2.10±0.40	2.10±0.37	2.10±0.27	0.06±0.16	-0.75±0.25	-0.17±0.13	0.04±0.15	0.28±0.12	M
6440_8	3907± 77	3621±166	3857± 70	1.40±0.18	0.78±0.71	1.36±0.17	-0.04±0.18	-1.26±0.33	-0.32±0.16	0.07±0.17	0.26±0.11	M
6440_10	4267±206	4374±167	4332±130	1.80±0.40	2.25±0.40	2.02±0.28	0.05±0.19	-0.75±0.33	-0.15±0.16	0.07±0.13	0.33±0.08	M
6440_11	3684± 45	3500±100	3653± 41	0.80±0.10	2.75±0.75	0.83±0.10	0.13±0.13	-1.15±0.23	-0.18±0.11	0.16±0.07	0.31±0.08	
6440_13	4498±110	4524±236	4503±100	2.04±0.32	2.00±0.50	2.03±0.27	-0.24±0.26	-0.65±0.32	-0.40±0.20	0.15±0.18	0.33±0.08	M
6440_14	3916±113	3600±122	3771± 83	1.36±0.04	1.19±0.64	1.36±0.04	0.08±0.22	-1.30±0.24	-0.55±0.16	0.11±0.09	0.32±0.06	M
6440_15	4459±139	4352±123	4399± 92	1.72±0.27	1.96±0.42	1.79±0.23	-0.13±0.10	-0.70±0.24	-0.21±0.09	0.20±0.08	0.33±0.06	M
6440_16	3989±124	4401±199	4104±105	1.67±0.01	2.30±0.40	1.67±0.01	0.11±0.02	-0.02±0.25	0.11±0.02	0.08±0.21	0.18±0.14	
6440_17	3685± 45	3524± 74	3641± 38	0.80±0.10	2.41±0.83	0.82±0.10	0.13±0.13	-1.00±0.22	-0.16±0.11	0.16±0.07	0.30±0.10	
6440_18	3976± 84	4275±134	4060± 71	1.54±0.17	2.35±0.45	1.64±0.16	0.03±0.17	-0.45±0.27	-0.11±0.14	-0.03±0.17	0.23±0.13	
6440_19	4268±150	4351±123	4318± 95	1.86±0.29	2.30±0.25	2.11±0.19	0.16±0.18	-0.30±0.25	0.00±0.15	0.00±0.11	0.16±0.14	
6453_1	4777±183	5026±234	4872±144	2.50±0.40	2.76±0.46	2.61±0.30	-0.31±0.23	-0.40±0.37	-0.34±0.20	0.15±0.19	0.29±0.12	
6453_2	4620±194	4751±157	4699±122	2.50±0.50	3.15±0.39	2.90±0.31	-0.22±0.25	-0.75±0.25	-0.48±0.18	0.12±0.17	0.29±0.09	
6453_3	4805±212	4826±115	4821±101	2.50±0.50	2.85±0.23	2.79±0.21	-0.46±0.25	-0.85±0.23	-0.67±0.17	0.26±0.19	0.25±0.10	
6453_4	4434±143	4824±114	4672± 89	2.20±0.50	3.15±0.23	2.98±0.21	0.11±0.15	0.12±0.10	0.12±0.08	0.04±0.14	0.20±0.12	
6453_5	4888±252	5000±224	4951±167	2.50±0.60	2.75±0.46	2.66±0.37	-0.47±0.30	-0.70±0.24	-0.61±0.19	0.25±0.19	0.24±0.08	
6453_6	4744±188	4900±166	4832±124	2.50±0.40	2.65±0.32	2.59±0.25	-0.07±0.22	-0.23±0.28	-0.13±0.17	0.03±0.12	0.26±0.10	
6453_7	4542± 28	4700±218	4545± 28	2.19±0.27	2.30±0.56	2.21±0.24	-0.09±0.08	-0.33±0.26	-0.11±0.08	0.13±0.11	0.25±0.08	
6453_8	4859±186	4774±236	4826±146	2.20±0.50	1.95±0.42	2.05±0.32	-0.73±0.15	-1.70±0.25	-0.99±0.13	0.38±0.13	0.26±0.09	
6453_10	4396±176	4650±166	4531±121	1.10±0.50	1.15±0.50	1.12±0.35	-1.51±0.21	-1.65±0.23	-1.57±0.16	0.44±0.14	0.17±0.09	M
6453_11	4826±379	4726± 74	4730± 72	2.10±0.90	1.70±0.33	1.75±0.31	-1.00±0.60	-1.55±0.15	-1.52±0.15	0.36±0.18	0.22±0.15	M
6453_13	4371±158	4451±149	4413±108	2.10±0.40	2.15±0.39	2.13±0.28	0.11±0.14	-0.13±0.25	0.05±0.12	0.03±0.15	0.13±0.09	
6453_14	4915±272	5077±160	5036±138	2.50±0.60	2.56±0.41	2.54±0.34	-0.48±0.32	-0.80±0.25	-0.68±0.20	0.23±0.20	0.26±0.13	
6453_15	4893±469	4950±149	4945±142	2.00±0.90	1.95±0.42	1.96±0.38	-1.40±0.50	-1.55±0.15	-1.54±0.14	0.43±0.16	0.12±0.10	M
6453_16	4295±220	4500±100	4465± 91	2.00±0.50	2.15±0.23	2.12±0.21	0.23±0.12	0.04±0.08	0.10±0.07	0.02±0.14	0.18±0.12	

Table B.3 - continued.

NGC ID	$T_{eff}^{(a)}$ (K)	$T_{eff}^{(b)}$ (K)	$T_{eff}^{(avg)}$ (K)	$\log(g)^{(a)}$	$\log(g)^{(b)}$	$\log(g)^{(avg)}$	[Fe/H] ^(a)	[Fe/H] ^(b)	[Fe/H] ^(avg)	[Mg/Fe] ^(a)	[α /Fe] ^(b)	members
6453_17	4655±206	4824±160	4761±126	2.50±0.50	2.85±0.39	2.72±0.31	-0.15±0.25	-0.40±0.30	-0.25±0.19	0.10±0.15	0.31±0.10	
6453_18	4459±144	4775±135	4628± 98	2.20±0.50	2.80±0.24	2.69±0.22	0.10±0.13	0.07±0.27	0.09±0.12	0.03±0.15	0.14±0.08	
6538_2	4214±169	4374±125	4318±100	1.90±0.40	2.49±0.39	2.20±0.28	-0.15±0.22	-0.75±0.25	-0.41±0.17	0.06±0.18	0.28±0.07	
6538_3	4540±180	4576±159	4560±119	2.30±0.50	2.55±0.41	2.45±0.32	-0.28±0.26	-0.85±0.32	-0.51±0.20	0.15±0.18	0.26±0.10	
6538_4	3890±115	3576±115	3733± 81	1.29±0.18	0.80±0.60	1.25±0.17	0.03±0.18	-1.35±0.23	-0.49±0.14	0.05±0.19	0.26±0.09	M
6538_5	4416±161	4347±122	4372± 97	1.90±0.40	1.99±0.32	1.95±0.25	-0.21±0.25	-0.81±0.24	-0.52±0.17	0.17±0.14	0.29±0.09	M
6538_6	4798±208	4900±165	4860±129	2.60±0.40	2.75±0.40	2.67±0.28	-0.25±0.29	-0.75±0.25	-0.54±0.19	0.14±0.21	0.25±0.14	
6538_7	4283±174	4201±100	4221± 86	1.90±0.40	1.90±0.37	1.90±0.27	0.12±0.19	-0.60±0.20	-0.22±0.14	0.04±0.11	0.22±0.12	
6538_8	4499±109	4501±193	4499± 95	2.04±0.32	2.30±0.60	2.10±0.28	-0.24±0.25	-0.85±0.23	-0.57±0.17	0.15±0.18	0.30±0.08	M
6538_9	4777±262	4701±218	4732±168	2.50±0.50	2.30±0.56	2.41±0.37	-0.41±0.31	-0.85±0.23	-0.69±0.18	0.21±0.21	0.30±0.08	M
6538_10	4505±177	4600±165	4556±121	2.20±0.50	2.45±0.47	2.33±0.34	-0.41±0.24	-0.75±0.25	-0.57±0.17	0.21±0.21	0.28±0.07	M
6538_11	3888±117	3569±112	3721± 81	1.28±0.18	0.61±0.59	1.22±0.17	0.03±0.18	-1.36±0.22	-0.53±0.14	0.05±0.19	0.25±0.10	
6538_13	3892±100	3549± 99	3720± 70	1.20±0.20	0.89±0.63	1.17±0.19	-0.03±0.40	-1.55±0.27	-1.07±0.22	0.13±0.08	0.29±0.10	
6538_14	4145±157	4200±187	4168±120	1.60±0.40	1.80±0.56	1.67±0.33	-0.25±0.20	-1.10±0.37	-0.44±0.18	0.08±0.17	0.33±0.06	M
6538_15	4702±271	4700±187	4701±154	2.30±0.60	2.65±0.32	2.57±0.28	-0.40±0.40	-1.10±0.37	-0.78±0.27	0.20±0.22	0.32±0.07	
6538_17	4456±127	4775± 75	4693± 64	2.40±0.40	3.30±0.24	3.06±0.21	0.30±0.07	0.20±0.17	0.29±0.06	0.06±0.08	0.13±0.10	
6538_18	4661±233	4999±193	4861±149	2.40±0.50	3.40±0.37	3.05±0.30	-0.47±0.27	-0.65±0.23	-0.57±0.18	0.26±0.20	0.32±0.07	M
Pal7_1	4286±149	4250±156	4269±108	1.90±0.50	1.85±0.32	1.86±0.27	-0.38±0.26	-1.05±0.27	-0.70±0.19	0.20±0.21	0.28±0.10	M
Pal7_3	4452±206	4447±187	4449±138	2.20±0.50	1.99±0.50	2.10±0.35	-0.06±0.24	-0.75±0.25	-0.39±0.17	0.05±0.17	0.28±0.14	M
Pal7_4	3900±102	3549±100	3720± 71	1.26±0.17	1.14±0.63	1.25±0.16	0.07±0.17	-1.40±0.20	-0.55±0.13	0.02±0.20	0.35±0.07	M
Pal7_5	4322±153	4250±193	4294±120	1.90±0.50	2.55±0.65	2.14±0.40	-0.50±0.20	-1.85±0.39	-0.78±0.18	0.25±0.19	0.36±0.05	
Pal7_7	3994± 98	3625±168	3900± 85	1.50±0.40	1.10±0.62	1.38±0.34	-0.27±0.19	-1.85±0.23	-0.91±0.15	0.11±0.17	0.32±0.09	M
Pal7_8	4346±216	4348±122	4347±106	2.10±0.50	2.05±0.35	2.07±0.29	-0.00±0.19	-0.80±0.24	-0.31±0.15	0.03±0.14	0.28±0.10	M
Pal7_9	3932± 43	3674±195	3920± 42	1.42±0.21	0.89±0.74	1.38±0.20	-0.20±0.23	-1.45±0.27	-0.73±0.18	0.12±0.17	0.30±0.12	M
Pal7_10	3942± 46	3776±175	3931± 44	1.50±0.15	1.15±0.71	1.49±0.15	-0.06±0.16	-1.35±0.23	-0.48±0.13	0.06±0.12	0.30±0.10	M
Pal7_11	4296±194	4374±125	4351±105	2.00±0.50	2.05±0.35	2.03±0.29	0.00±0.20	-0.75±0.25	-0.29±0.16	0.03±0.15	0.29±0.09	M
Pal7_12	4428±197	4350±123	4372±104	2.20±0.50	2.00±0.32	2.06±0.27	-0.20±0.28	-0.80±0.25	-0.53±0.19	0.11±0.18	0.27±0.09	M
Pal7_13	4522±247	4575±275	4546±184	1.90±0.50	2.35±0.64	2.07±0.39	-0.50±0.40	-1.20±0.40	-0.85±0.28	0.30±0.26	0.32±0.07	

Table B.3 - continued.

NGC ID	$T_{eff}^{(a)}$ (K)	$T_{eff}^{(b)}$ (K)	$T_{eff}^{(avg)}$ (K)	$\log(g)^{(a)}$	$\log(g)^{(b)}$	$\log(g)^{(avg)}$	$[\text{Fe}/\text{H}]^{(a)}$	$[\text{Fe}/\text{H}]^{(b)}$	$[\text{Fe}/\text{H}]^{(avg)}$	$[\text{Mg}/\text{Fe}]^{(a)}$	$[\alpha/\text{Fe}]^{(b)}$	members
Pal7_14	4241±176	4325±115	4300± 96	1.90±0.40	2.30±0.40	2.10±0.28	-0.13±0.23	-0.85±0.23	-0.49±0.16	0.06±0.18	0.30±0.08	M
Pal7_15	4256±168	4525±207	4363±131	1.90±0.40	2.60±0.62	2.11±0.34	-0.27±0.28	-0.55±0.35	-0.38±0.22	0.14±0.20	0.32±0.07	
Pal7_16	4273±169	4423±195	4338±128	1.90±0.50	2.35±0.67	2.06±0.40	-0.24±0.26	-0.75±0.25	-0.50±0.18	0.12±0.17	0.30±0.10	
Pal7_17	4364±108	4399±199	4372± 95	2.00±0.40	2.20±0.51	2.08±0.31	-0.39±0.28	-0.90±0.20	-0.73±0.16	0.22±0.20	0.29±0.09	M
Pal7_18	3961± 96	3827±160	3925± 82	1.36±0.26	1.26±0.64	1.35±0.24	-0.26±0.27	-1.30±0.25	-0.82±0.18	0.18±0.21	0.30±0.08	M
Pal7_19	4550±196	4475±175	4508±131	2.30±0.50	2.65±0.50	2.47±0.35	-0.29±0.26	-1.10±0.30	-0.64±0.20	0.16±0.18	0.29±0.09	
6569_1	4764±207	4828±337	4781±176	2.30±0.60	2.40±0.62	2.35±0.43	-0.60±0.18	-1.10±0.44	-0.67±0.17	0.38±0.22	0.30±0.08	M
6569_2	4057±100	4103±251	4063± 93	1.50±0.40	1.76±0.71	1.56±0.35	-0.22±0.32	-1.14±0.39	-0.59±0.25	0.10±0.27	0.34±0.07	M
6569_3	4398± 99	4625±125	4486± 78	2.10±0.40	2.65±0.32	2.44±0.25	-0.46±0.18	-0.75±0.25	-0.56±0.15	0.21±0.20	0.27±0.11	
6569_4	4809±243	5000±224	4912±165	2.40±0.60	2.85±0.45	2.69±0.36	-0.56±0.26	-0.70±0.25	-0.63±0.18	0.29±0.19	0.29±0.07	
6569_5	4349±117	4576±195	4409±100	2.10±0.40	2.65±0.39	2.38±0.28	0.11±0.14	-0.19±0.32	0.06±0.13	-0.00±0.15	0.15±0.10	M
6569_6	4778±170	4950±187	4856±126	2.50±0.50	3.20±0.24	3.07±0.22	-0.50±0.24	-0.80±0.25	-0.64±0.17	0.30±0.19	0.29±0.09	
6569_7	4507± 95	4799±148	4592± 80	2.30±0.40	3.05±0.27	2.82±0.22	0.10±0.11	0.10±0.24	0.10±0.10	0.11±0.11	0.14±0.09	
6569_8	4891±248	4875±321	4885±196	2.60±0.60	2.50±0.55	2.55±0.41	-0.51±0.28	-1.05±0.42	-0.68±0.23	0.27±0.20	0.31±0.08	M
6569_9	4808±238	4851±123	4842±109	2.50±0.60	2.55±0.41	2.53±0.34	-0.48±0.27	-0.80±0.25	-0.65±0.18	0.24±0.19	0.25±0.10	M
6569_10	4642±175	4497±222	4586±137	2.40±0.40	1.89±0.49	2.20±0.31	-0.42±0.21	-1.10±0.30	-0.64±0.17	0.22±0.22	0.31±0.10	M
6569_11	4809±358	5025± 74	5016± 73	2.20±0.90	2.25±0.40	2.24±0.37	-0.90±0.50	-0.95±0.15	-0.95±0.14	0.31±0.17	0.23±0.14	M
6569_12	4876±275	5050±187	4995±155	2.60±0.60	3.05±0.35	2.94±0.30	-0.49±0.31	-0.70±0.24	-0.62±0.19	0.26±0.20	0.24±0.09	
6569_13	4692±180	4774±175	4734±126	2.00±0.40	2.50±0.45	2.22±0.30	-0.61±0.24	-0.95±0.35	-0.72±0.20	0.44±0.18	0.30±0.08	
6569_14	4560±161	4825±195	4667±124	2.40±0.40	2.95±0.52	2.60±0.32	-0.06±0.19	-0.06±0.23	-0.06±0.15	0.07±0.13	0.21±0.11	M
6569_15	4531± 31	4975±134	4554± 30	2.17±0.23	2.95±0.27	2.50±0.18	-0.10±0.07	0.00±0.26	-0.09±0.07	0.12±0.11	0.27±0.11	
6569_16	4789±165	4974±174	4876±120	2.50±0.40	2.75±0.25	2.68±0.21	-0.37±0.22	-0.55±0.35	-0.42±0.19	0.19±0.21	0.28±0.10	
6569_17	4565± 85	4824±114	4657± 68	2.70±0.40	3.15±0.23	3.04±0.20	0.27±0.03	0.12±0.10	0.26±0.03	0.13±0.06	0.17±0.11	M
6569_19	4755±195	4926±159	4857±123	2.50±0.50	3.10±0.30	2.94±0.26	-0.42±0.24	-0.70±0.24	-0.56±0.17	0.22±0.20	0.28±0.10	
6749_1	4765±364	4575±195	4618±172	1.70±0.70	1.40±0.49	1.50±0.40	-1.64±0.32	-2.25±0.25	-2.02±0.20	0.43±0.13	0.11±0.09	
6749_2	4650±215	5175±115	5059±101	2.40±0.50	3.45±0.35	3.10±0.29	-0.06±0.21	0.08±0.10	0.05±0.09	0.03±0.14	0.13±0.11	
6749_3	4666±411	4425±115	4442±110	1.50±0.70	0.10±0.20	0.21±0.19	-1.60±0.60	-2.50±0.10	-2.48±0.10	0.45±0.14	0.17±0.13	
6749_4	4483±288	4250±100	4275± 94	1.60±0.70	0.25±0.25	0.40±0.24	-1.00±0.60	-2.45±0.15	-2.36±0.15	0.38±0.19	0.16±0.14	

Table B.3 - continued.

NGC ID	$T_{eff}^{(a)}$ (K)	$T_{eff}^{(b)}$ (K)	$T_{eff}^{(avg)}$ (K)	$\log(g)^{(a)}$	$\log(g)^{(b)}$	$\log(g)^{(avg)}$	[Fe/H] ^(a)	[Fe/H] ^(b)	[Fe/H] ^(avg)	[Mg/Fe] ^(a)	[α /Fe] ^(b)	members
6749_5	4346±171	4151±165	4245±119	0.90±0.40	0.30±0.33	0.54±0.25	-1.56±0.34	-2.15±0.23	-1.96±0.19	0.45±0.09	0.15±0.16	
6749_6	4152± 54	4124±202	4150± 52	0.73±0.02	0.05±0.15	0.72±0.02	-1.65±0.28	-2.35±0.23	-2.07±0.18	0.45±0.01	0.20±0.13	
6749_7	4346±172	4100±229	4257±138	0.90±0.40	0.10±0.20	0.26±0.18	-1.56±0.34	-2.15±0.23	-1.96±0.19	0.45±0.09	0.16±0.12	
6749_8	4646±233	4100±200	4332±152	2.00±0.40	0.05±0.15	0.29±0.14	-0.61±0.32	-2.10±0.20	-1.68±0.17	0.33±0.21	0.15±0.12	M
6749_9	4596±270	4049±188	4227±154	2.00±0.50	0.05±0.15	0.21±0.14	-0.70±0.40	-2.15±0.23	-1.79±0.20	0.34±0.19	0.20±0.13	
6749_10	4624±224	4025±208	4302±152	2.10±0.40	0.10±0.20	0.50±0.18	-0.52±0.34	-2.15±0.23	-1.64±0.19	0.30±0.21	0.19±0.11	M
6749_11	4737±183	4126±201	4460±135	2.12±0.32	0.20±0.24	0.89±0.19	-0.60±0.32	-2.15±0.23	-1.62±0.19	0.32±0.21	0.17±0.15	M
6749_13	4228± 48	4375±125	4247± 45	1.53±0.15	1.95±0.35	1.60±0.14	-0.03±0.10	-0.25±0.25	-0.06±0.09	-0.05±0.18	0.24±0.10	
6749_14	4535± 8	4250±100	4533± 8	2.24±0.10	0.40±0.43	2.15±0.10	-0.13±0.10	-2.50±0.10	-1.31±0.07	0.01±0.02	0.17±0.15	
6749_15	4732±168	4101±200	4470±129	1.99±0.30	0.20±0.33	1.18±0.22	-0.74±0.23	-2.20±0.24	-1.44±0.17	0.40±0.17	0.17±0.15	M
6749_16	4623±315	4300±100	4329± 95	1.50±0.70	0.20±0.24	0.34±0.23	-1.72±0.33	-2.50±0.10	-2.43±0.10	0.45±0.14	0.21±0.14	
6749_17	4526±149	4776±207	4611±121	2.30±0.40	2.45±0.57	2.35±0.33	-0.22±0.25	-0.30±0.24	-0.26±0.17	0.11±0.20	0.25±0.10	
6749_18	4341±162	4449±187	4387±122	2.00±0.40	1.95±0.42	1.98±0.29	0.01±0.13	-0.40±0.20	-0.11±0.11	0.00±0.13	0.27±0.12	
Pal10_1	4187±194	4475±208	4321±142	1.70±0.40	2.45±0.52	1.98±0.32	0.07±0.16	-0.11±0.35	0.04±0.15	0.01±0.15	0.10±0.09	
Pal10_2	4219±228	4400±166	4337±134	1.90±0.50	2.15±0.45	2.04±0.33	-0.00±0.19	-0.23±0.28	-0.08±0.16	0.05±0.14	0.17±0.12	M
Pal10_3	4490±258	4550±150	4535±130	2.20±0.50	2.80±0.33	2.62±0.28	-0.15±0.27	-0.90±0.30	-0.49±0.20	0.08±0.18	0.24±0.12	
Pal10_4	4250±207	4225±175	4235±134	1.80±0.50	1.85±0.39	1.83±0.31	-0.09±0.27	-0.65±0.23	-0.41±0.18	0.07±0.17	0.21±0.11	M
Pal10_6	4048±125	3976±284	4036±114	1.56±0.32	1.30±0.84	1.53±0.30	-0.20±0.20	-1.45±0.41	-0.44±0.18	0.09±0.14	0.30±0.08	
Pal10_7	4326±246	4226±208	4267±159	2.00±0.50	2.15±0.59	2.06±0.38	-0.17±0.28	-1.40±0.30	-0.74±0.20	0.10±0.19	0.33±0.08	
Pal10_8	4212±190	4100±165	4148±125	1.80±0.50	1.50±0.54	1.66±0.37	-0.01±0.15	-0.80±0.33	-0.14±0.14	0.05±0.13	0.27±0.13	M
Pal10_9	4369±165	4475±134	4433±104	1.79±0.33	2.05±0.52	1.86±0.28	-0.07±0.13	-0.50±0.22	-0.18±0.11	0.08±0.14	0.33±0.08	M
Pal10_10	4147± 99	4050±150	4118± 83	1.59±0.13	1.35±0.55	1.58±0.13	-0.06±0.10	-0.90±0.30	-0.14±0.09	0.03±0.13	0.30±0.10	M
Pal10_11	4362±150	4425±115	4402± 91	1.56±0.15	1.90±0.44	1.60±0.14	-0.13±0.10	-0.50±0.10	-0.32±0.07	0.17±0.09	0.30±0.08	M
Pal10_13	4134±108	4177±114	4154± 78	1.58±0.14	1.70±0.25	1.61±0.12	-0.07±0.10	-0.65±0.23	-0.16±0.09	0.08±0.06	0.25±0.10	M
Pal10_16	4427±144	4299±100	4341± 82	1.80±0.40	1.85±0.32	1.83±0.25	-0.07±0.10	-0.50±0.10	-0.29±0.07	0.08±0.16	0.24±0.14	M
Pal10_17	4214± 68	4275±134	4227± 61	1.48±0.07	1.85±0.32	1.50±0.07	-0.13±0.10	-0.55±0.15	-0.26±0.08	0.12±0.01	0.28±0.12	M
Pal11_2	4508±174	4800±185	4645±127	2.30±0.40	3.00±0.31	2.74±0.25	-0.38±0.22	-0.50±0.31	-0.42±0.18	0.18±0.18	0.26±0.13	M
Pal11_3	3980± 47	4624±125	4060± 44	1.73±0.19	4.45±0.42	2.19±0.17	-0.27±0.09	-0.55±0.15	-0.34±0.08	0.11±0.13	0.26±0.13	

Table B.3 - continued.

NGC ID	$T_{eff}^{(a)}$ (K)	$T_{eff}^{(b)}$ (K)	$T_{eff}^{(avg)}$ (K)	$\log(g)^{(a)}$	$\log(g)^{(b)}$	$\log(g)^{(avg)}$	$[\text{Fe}/\text{H}]^{(a)}$	$[\text{Fe}/\text{H}]^{(b)}$	$[\text{Fe}/\text{H}]^{(avg)}$	$[\text{Mg}/\text{Fe}]^{(a)}$	$[\alpha/\text{Fe}]^{(b)}$	members
Pal11_6	4568±179	4725±206	4636±135	2.40±0.40	2.85±0.45	2.60±0.30	-0.34±0.17	-0.65±0.32	-0.41±0.15	0.17±0.15	0.31±0.10	M
Pal11_7	4624±196	4675±159	4655±124	2.50±0.50	2.65±0.32	2.61±0.27	-0.22±0.23	-0.65±0.32	-0.37±0.19	0.12±0.17	0.30±0.10	M
Pal11_8	4361±114	4397±164	4373± 94	2.02±0.32	2.14±0.45	2.06±0.26	-0.30±0.19	-0.71±0.33	-0.40±0.16	0.22±0.19	0.31±0.10	M
Pal11_9	4265±160	4350±122	4318± 97	1.91±0.32	2.05±0.35	1.97±0.24	-0.20±0.12	-0.80±0.24	-0.32±0.11	0.05±0.12	0.27±0.11	M
Pal11_10	4661±187	4802±186	4732±132	2.50±0.40	3.06±0.47	2.74±0.30	-0.34±0.20	-0.60±0.20	-0.47±0.14	0.19±0.18	0.24±0.14	M
Pal11_11	4358±149	4399±165	4377±111	2.10±0.28	2.10±0.43	2.10±0.23	-0.08±0.13	-0.70±0.33	-0.16±0.12	0.08±0.14	0.32±0.07	M
Pal11_12	4463±154	4450±187	4458±119	2.30±0.40	2.30±0.46	2.30±0.30	-0.07±0.21	-0.75±0.25	-0.35±0.16	0.07±0.15	0.34±0.07	M
Pal11_13	4645±193	4898±123	4825±104	2.50±0.40	3.15±0.23	2.99±0.20	-0.26±0.21	-0.35±0.23	-0.30±0.16	0.14±0.16	0.25±0.13	M
Pal11_16	4643±201	4875±125	4810±106	3.00±0.60	4.35±0.39	3.95±0.33	0.30±0.04	0.10±0.10	0.27±0.04	0.18±0.12	0.10±0.08	
Pal11_17	4118±118	4301±101	4224± 77	1.70±0.40	1.95±0.35	1.84±0.26	-0.19±0.20	-0.60±0.20	-0.39±0.14	0.07±0.14	0.30±0.08	M
M75_2	4378±116	4377±125	4377± 85	1.90±0.50	1.65±0.23	1.69±0.21	-0.49±0.25	-1.00±0.10	-0.93±0.09	0.20±0.18	0.23±0.13	M
M75_3	4743±238	4799± 99	4791± 92	2.30±0.60	2.40±0.37	2.37±0.31	-0.50±0.22	-0.90±0.20	-0.72±0.15	0.19±0.16	0.21±0.13	
M75_4	4811±392	5050±100	5035± 97	2.20±0.90	2.30±0.33	2.29±0.31	-0.80±0.50	-0.95±0.15	-0.94±0.14	0.32±0.17	0.19±0.15	M
M75_5	4878±328	5076±115	5054±108	2.40±0.80	2.80±0.25	2.76±0.24	-0.70±0.40	-0.85±0.23	-0.81±0.20	0.26±0.14	0.18±0.12	M
M75_6	4532±278	5050±245	4824±184	1.30±0.60	1.95±0.42	1.74±0.34	-1.40±0.40	-1.20±0.24	-1.25±0.21	0.45±0.13	0.17±0.11	M
M75_7	4224± 61	4126±200	4216± 58	1.40±0.40	1.10±0.70	1.33±0.35	-0.66±0.31	-1.30±0.33	-0.96±0.23	0.40±0.20	0.24±0.14	M
M75_8	4539±182	4425±114	4458± 97	2.20±0.50	1.75±0.25	1.84±0.22	-0.46±0.19	-1.15±0.23	-0.74±0.15	0.17±0.18	0.21±0.11	
M75_9	4289± 90	4250±192	4282± 82	1.50±0.40	1.45±0.72	1.49±0.35	-0.69±0.26	-1.15±0.23	-0.95±0.17	0.40±0.18	0.26±0.13	M
M75_10	4805±359	4500±194	4569±170	2.30±0.90	2.40±0.49	2.38±0.43	-0.70±0.50	-1.95±0.35	-1.54±0.29	0.31±0.19	0.31±0.07	M
M75_11	4406±125	4426±159	4414± 98	2.00±0.50	1.80±0.46	1.89±0.34	-0.48±0.20	-1.00±0.22	-0.72±0.15	0.19±0.19	0.21±0.13	M
M75_13	4863±408	5100±229	5044±200	2.00±0.90	2.40±0.62	2.27±0.51	-1.50±0.40	-1.65±0.23	-1.61±0.20	0.43±0.15	0.25±0.13	M
M75_14	4685±387	5301±187	5184±168	1.60±0.70	2.35±0.32	2.22±0.29	-1.40±0.40	-1.10±0.20	-1.16±0.18	0.44±0.14	0.11±0.09	M
7006_9	5195±430	6125±125	6053±120	2.40±0.80	4.20±0.46	3.75±0.40	-2.00±0.50	-2.50±0.10	-2.48±0.10	0.40±0.19	0.06±0.07	
7006_18	4901±461	4900±122	4900±118	2.10±0.90	2.55±0.27	2.51±0.26	-1.40±0.50	-1.70±0.24	-1.64±0.22	0.40±0.17	0.19±0.11	M
7006_19	5013±393	5000±100	5001± 97	2.20±0.80	0.80±0.81	1.51±0.57	-1.70±0.50	-2.50±0.10	-2.47±0.10	0.42±0.18	0.20±0.14	
7006_21	5029±452	5375±125	5350±120	2.20±0.90	3.30±0.24	3.23±0.23	-1.70±0.50	-1.75±0.25	-1.74±0.22	0.42±0.17	0.28±0.12	M
7006_22	5020±465	5400±122	5375±118	2.20±0.90	3.40±0.37	3.23±0.34	-1.70±0.50	-2.50±0.10	-2.47±0.10	0.42±0.17	0.07±0.06	
7006_23	4807±402	4924±225	4896±196	1.90±0.90	1.90±0.37	1.90±0.34	-1.60±0.40	-1.80±0.24	-1.75±0.21	0.44±0.15	0.28±0.10	M

Table B.3 - continued.

NGC ID	$T_{eff}^{(a)}$ (K)	$T_{eff}^{(b)}$ (K)	$T_{eff}^{(avg)}$ (K)	$\log(g)^{(a)}$	$\log(g)^{(b)}$	$\log(g)^{(avg)}$	$[\text{Fe}/\text{H}]^{(a)}$	$[\text{Fe}/\text{H}]^{(b)}$	$[\text{Fe}/\text{H}]^{(avg)}$	$[\text{Mg}/\text{Fe}]^{(a)}$	$[\alpha/\text{Fe}]^{(b)}$	members
7006_24	4623 $\pm$ 340	4800 $\pm$ 218	4748 $\pm$ 184	1.60 $\pm$ 0.80	1.90 $\pm$ 0.44	1.83 $\pm$ 0.39	-1.43 $\pm$ 0.35	-1.65 $\pm$ 0.23	-1.58 $\pm$ 0.19	0.43 $\pm$ 0.14	0.28 $\pm$ 0.10	M
7006_25	4947 $\pm$ 446	5050 $\pm$ 187	5035 $\pm$ 172	2.10 $\pm$ 0.90	1.85 $\pm$ 0.32	1.88 $\pm$ 0.30	-1.60 $\pm$ 0.40	-1.80 $\pm$ 0.24	-1.75 $\pm$ 0.21	0.42 $\pm$ 0.17	0.18 $\pm$ 0.12	M
7006_26	4960 $\pm$ 464	5200 $\pm$ 100	5189 $\pm$ 98	2.10 $\pm$ 0.90	2.90 $\pm$ 0.44	2.75 $\pm$ 0.40	-1.70 $\pm$ 0.50	-2.45 $\pm$ 0.15	-2.39 $\pm$ 0.14	0.42 $\pm$ 0.17	0.08 $\pm$ 0.10	
7006_27	4957 $\pm$ 376	4800 $\pm$ 150	4822 $\pm$ 139	2.20 $\pm$ 0.90	4.45 $\pm$ 0.61	3.74 $\pm$ 0.50	-1.70 $\pm$ 0.50	-2.50 $\pm$ 0.10	-2.47 $\pm$ 0.10	0.42 $\pm$ 0.16	0.34 $\pm$ 0.07	
7006_28	5020 $\pm$ 366	4625 $\pm$ 168	4693 $\pm$ 152	2.30 $\pm$ 0.90	4.55 $\pm$ 0.52	3.99 $\pm$ 0.45	-1.60 $\pm$ 0.60	-2.50 $\pm$ 0.10	-2.48 $\pm$ 0.10	0.42 $\pm$ 0.17	0.36 $\pm$ 0.05	
7006_29	5060 $\pm$ 475	5750 $\pm$ 158	5682 $\pm$ 150	2.60 $\pm$ 1.00	4.80 $\pm$ 0.24	4.68 $\pm$ 0.23	-0.90 $\pm$ 0.50	-1.00 $\pm$ 0.10	-1.00 $\pm$ 0.10	0.37 $\pm$ 0.24	0.25 $\pm$ 0.14	
7006_31	3980 $\pm$ 188	4600 $\pm$ 123	4415 $\pm$ 103	1.70 $\pm$ 1.20	4.35 $\pm$ 0.55	3.89 $\pm$ 0.50	-0.27 $\pm$ 0.26	-0.85 $\pm$ 0.23	-0.60 $\pm$ 0.17	0.11 $\pm$ 0.09	0.30 $\pm$ 0.08	
7006_32	5064 $\pm$ 433	5625 $\pm$ 125	5582 $\pm$ 120	2.20 $\pm$ 0.80	3.65 $\pm$ 0.45	3.30 $\pm$ 0.39	-1.90 $\pm$ 0.40	-2.50 $\pm$ 0.10	-2.46 $\pm$ 0.10	0.41 $\pm$ 0.18	0.06 $\pm$ 0.07	

^aMILES library ^bCOELHO library ^cDue to low number of observed stars it was hard to define the membership of M 107. See discussion in Chapter 4

Appendix C

FORS2 pipeline - wavelength calibration for low resolution spectra of GRIS1400V

C.1 Default pipeline reduction

I used the esorex and the monolithic pipelines for the fors2, using the respective calibration files, as follows:

```
> esorex fors_calib cal.sof
> esorex fors_science sci.sof
```

where the file cal.sof was:

```
../BIAS/FORS2.2012-05-21T12_21_30.622.fits BIAS
../BIAS/FORS2.2012-05-21T12_22_04.555.fits BIAS
../BIAS/FORS2.2012-05-21T12_22_38.548.fits BIAS
../BIAS/FORS2.2012-05-21T12_23_12.561.fits BIAS
../BIAS/FORS2.2012-05-21T12_23_46.563.fits BIAS
../FORS2.2012-05-22T11_34_08.770.fits SCREEN_FLAT_MOS
../FORS2.2012-05-22T11_35_38.858.fits SCREEN_FLAT_MOS
../FORS2.2012-05-22T11_37_09.905.fits SCREEN_FLAT_MOS
../FORS2.2012-05-22T11_38_40.843.fits SCREEN_FLAT_MOS
../FORS2.2012-05-22T11_40_11.880.fits SCREEN_FLAT_MOS
../FORS2.2012-05-22T11_42_40.562.fits LAMP_MOS
../Mg-CAL/M.FORS2.2012-05-31T13_47_50.389.fits MASTER_LINECAT
../Mg-CAL/M.FORS2.2012-05-31T13_45_57.523.fits GRISM_TABLE
```

and the file sci.sof was:

```

FORS2.2012-05-22T03_20_50.977.fits SCIENCE_MOS
master_bias.fits MASTER_BIAS
master_norm_flat_mos.fits MASTER_NORM_FLAT_MOS
disp_coeff_mos.fits DISP_COEFF_MOS
curv_coeff_mos.fits CURV_COEFF_MOS
slit_location_mos.fits SLIT_LOCATION_MOS
../../Mg-CAL/M.FORS2.2012-05-31T13_45_57.523.fits GRISM_TABLE

```

So the results can be seen in the Figure C.1.

Figure C.1: Output (mapped_all_sci_mos.fits) of FORS2 pipeline using default parameters. The unique skyline used by the pipeline is indicated by the arrow. The blue line is just a vertical reference.

As you can see, there are too few skylines, and the strongest one (highlighted by the oval in the Figure C.1) is the 5577Å, which is the unique line considered by the pipeline for these low resolution spectra. You can check it in the page 109 of the FORS pipeline manual, issue 4.3, copied here in the Figure C.2. This table present ONLY ONE line, in the middle/red part of the grism 1400V spectra, which is not enough. So I found out a way to include my own line list, even with weak lines, and the results are present in the next Section.

Wavelength	Low resolution	Wavelength	Low resolution
5577.338	✓	7329.148	
5889.953		7340.885	
5895.923		7358.659	
5915.301		7571.746	✓
5932.862		7750.640	
5953.420		7759.996	
6257.961		7794.112	
6287.434		7808.467	
6300.304	✓	7821.503	
6306.869		7841.266	
6363.780		7913.708	
6498.729		7949.204	
6533.044		7964.650	✓
6553.617		7993.332	✓
6841.945		8014.059	
6863.955	✓	8310.719	
6870.994		8344.602	
6889.288		8382.392	
6900.833		8399.170	
6912.623		8415.231	
6923.220		8430.174	
6939.521		8452.250	
6969.930		8493.389	
7003.858		8791.186	
7244.907		8827.096	
7276.405		8885.850	
7284.439		8903.114	
7316.282		8943.395	
		8988.366	

Table 9.9.1: Default sky lines wavelengths used by the recipe `fors_science`. The marked lines are those used on data from low resolution grisms, i.e., all grisms with 300 gr/mm or less.

Figure C.2: Default list of skylines from manual.

C.2 New list of sky lines for GRIS1400V

I selected more skylines in the spectral region (Figure C.3) based on the UVES sky emission spectrum¹ and also a visual inspection of possible lines to be used. I included also a weak line more in the blue, in order to correct the entire dispersion. I also changed the parameter `skyalign` in order to fit a polynomial, as described in the manual, page 114, section 9.9.3. And then, my steps were:

```
> esorex fors_calib cal.sof
> esorex fors_science --skyalign=1 sci.sof
```

where the file `cal.sof` was the same as above, and the file `sci.sof` had one more line, indicating the new line list (and then not using the default list automatically):

¹ http://www.eso.org/observing/dfo/quality/UVES/pipeline/sky_spectrum.html

	WLEN	COMMENT
1	4861.3	18.3
2	5197.9	252.8
3	5200.3	146.7
4	5577.3	7787.2
5	5888.2	158.4
6	5890.	796.8
7	5895.9	487.

Figure C.3: New list of skylines with the right column being a relative intensity for reference.

```

FORS2.2012-05-22T03_20_50.977.fits SCIENCE_MOS
master_bias.fits MASTER_BIAS
master_norm_flat_mos.fits MASTER_NORM_FLAT_MOS
disp_coeff_mos.fits DISP_COEFF_MOS
curv_coeff_mos.fits CURV_COEFF_MOS
slit_location_mos.fits SLIT_LOCATION_MOS
../../CAL/shortlist-fors.fits MASTER_SKYLINECAT
../../Mg-CAL/M.FORS2.2012-05-31T13_45_57.523.fits GRISM_TABLE

```

And now, with more skylines, and using a polynomial fitting (instead of the default median assuming that the displacement is equal for the whole wavelength range) for the dispersion, it was possible to correct the dispersion even in the blue, as shown by the Figure C.4.

C.3 Comparisons between the 2 procedures

The Figure C.5 shows the comparison of the extracted spectra from one of the corrected slits (slit8 as an example), highlighted in the Figures C.1 and C.4. It is easy to notice that the new approach changes significantly the spectra.

In order to have a reference spectrum, I plot in the Figure C.6 the same of the Figure C.5, but for a slit where the dispersion does not significantly change between one procedure and another.

And finally, I use the slit 6 (of the new procedure) as a reference to overplot the

Figure C.4: Output (mapped_all_sci_mos.fits) of FORS2 pipeline using the new approach. The skylines used by the pipeline are indicated by the arrows. The blue line is just a vertical reference.

spectra of slit 8 from the 2 procedures. You can see in the Figure C.7 that the new approach successful corrects the dispersion in wavelength for low resolution spectra of FORS2/GRIS1400V.

Figure C.5: Comparison between the corrected spectrum of the slit #8 (solid white line) with the original one using the default parameters (dashed red line) for 3 regions, in particular the central region is the Mg triplet, very important for this work.

Figure C.6: Same of Figure C.5, but for the slit #6.

Figure C.7: Comparison between the default procedure (plots on the left) with the new procedure (plots on the right). The solid white line is the slit6, and the dashed red one is the slit8.

Papers in the period 2010-2014

We present here the abstract of all papers published and one submitted during the PhD as first author or co-author in order of publication date.

A&A 532, A131 (2011)
DOI: [10.1051/0004-6361/201116662](https://doi.org/10.1051/0004-6361/201116662)
© ESO 2011

**Astronomy
&
Astrophysics**

New Galactic star clusters discovered in the VVV survey^{★,★★,★★★}

J. Borissova¹, C. Bonatto², R. Kurtev¹, J. R. A. Clarke¹, F. Peñaloza¹, S. E. Sale^{1,5}, D. Minniti^{5,18}, J. Alonso-García⁵, E. Artigau³, R. Barbá¹⁶, E. Bica², G. L. Baume⁴, M. Catelan⁵, A. N. Chenè^{1,6}, B. Dias⁷, S. L. Folkes¹, D. Froebrich⁸, D. Geisler⁶, R. de Grijs^{9,10}, M. M. Hanson¹⁷, M. Hempel⁵, V. D. Ivanov¹¹, M. S. N. Kumar¹², P. Lucas¹³, F. Mauro⁶, C. Moni Bidin⁶, M. Rejkuba¹⁵, R. K. Saito⁵, M. Tamura¹⁴, and I. Toledo⁵

(Affiliations can be found after the references)

Received 6 February 2011 / Accepted 3 June 2011

ABSTRACT

Context. VISTA Variables in the Vía Láctea (VVV) is one of the six ESO Public Surveys operating on the new 4-m Visible and Infrared Survey Telescope for Astronomy (VISTA). VVV is scanning the Milky Way bulge and an adjacent section of the disk, where star formation activity is high. One of the principal goals of the VVV Survey is to find new star clusters of different ages.

Aims. In order to trace the early epochs of star cluster formation we concentrated our search in the directions to those of known star formation regions, masers, radio, and infrared sources.

Methods. The disk area covered by VVV was visually inspected using the pipeline processed and calibrated K_S -band tile images for stellar over-densities. Subsequently, we examined the composite JHK_S and ZJK_S color images of each candidate. PSF photometry of 15×15 arcmin fields centered on the candidates was then performed on the Cambridge Astronomy Survey Unit reduced images. After statistical field-star decontamination, color–magnitude and color–color diagrams were constructed and analyzed.

Results. We report the discovery of 96 new infrared open clusters and stellar groups. Most of the new cluster candidates are faint and compact (with small angular sizes), highly reddened, and younger than 5 Myr. For relatively well populated cluster candidates we derived their fundamental parameters such as reddening, distance, and age by fitting the solar-metallicity Padova isochrones to the color–magnitude diagrams.

Key words. open clusters and associations: general – stars: early-type – infrared: general

Figure D.1: Abstract of Borissova et al. (2011).

A&A 537, A107 (2012)
 DOI: 10.1051/0004-6361/201118407
 © ESO 2012

**Astronomy
&
Astrophysics**

VVV DR1: The first data release of the Milky Way bulge and southern plane from the near-infrared ESO public survey VISTA variables in the Vía Láctea[★]

R. K. Saito¹, M. Hempel¹, D. Minniti^{1,2,3}, P. W. Lucas⁴, M. Rejkuba⁵, I. Toledo⁶, O. A. Gonzalez⁵, J. Alonso-García¹,
 M. J. Irwin⁷, E. Gonzalez-Solares⁷, S. T. Hodgkin⁷, J. R. Lewis⁷, N. Cross⁸, V. D. Ivanov⁹, E. Kerins¹⁰,
 J. P. Emerson¹¹, M. Soto¹², E. B. Amôres^{13,14}, S. Gurovich¹⁵, I. Dékány¹, R. Angeloni¹, J. C. Beamin¹, M. Catelan¹,
 N. Padilla^{1,16}, M. Zoccali^{1,17}, P. Pietrukowicz¹⁸, C. Moni Bidin¹⁹, F. Mauro¹⁹, D. Geisler¹⁹, S. L. Folkes²⁰,
 S. E. Sale^{1,20}, J. Borissova²⁰, R. Kurtev²⁰, A. V. Ahumada^{9,21,22}, M. V. Alonso^{15,21}, A. Adamson²³, J. I. Arias¹²,
 R. M. Bandyopadhyay²⁴, R. H. Barbá^{12,25}, B. Barbuy²⁶, G. L. Baume²⁷, L. R. Bedin²⁸, A. Bellini²⁹, R. Benjamin³⁰,
 E. Bica³¹, C. Bonatto³¹, L. Bronfman³², G. Carraro⁹, A. N. Chenè^{19,20}, J. J. Clariá²¹, J. R. A. Clarke²⁰, C. Contreras⁴,
 A. Corvillón¹, R. de Grijs^{33,34}, B. Dias²⁶, J. E. Drew⁴, C. Fariña²⁷, C. Feinstein²⁷, E. Fernández-Lajús²⁷,
 R. C. Gamen²⁷, W. Gieren¹⁹, B. Goldman³⁵, C. González-Fernández³⁶, R. J. J. Grand³⁷, G. Gunthardt²¹,
 N. C. Hambly⁸, M. M. Hanson³⁸, K. G. Helminiak¹, M. G. Hoare³⁹, L. Huckvale¹⁰, A. Jordán¹, K. Kinemuchi⁴⁰,
 A. Longmore⁴¹, M. López-Corredoira^{42,43}, T. Maccarone⁴⁴, D. Majaess⁴⁵, E. L. Martín⁴⁶, N. Masetti⁴⁷,
 R. E. Mennickent¹⁹, I. F. Mirabel^{48,49}, L. Monaco⁹, L. Morelli²⁹, V. Motta²⁰, T. Palma²¹, M. C. Parisi²¹, Q. Parker^{50,51},
 F. Peñaloza²⁰, G. Pietrzyński^{18,19}, G. Pignata⁵², B. Popescu³⁸, M. A. Read⁸, A. Rojas¹, A. Roman-Lopes¹²,
 M. T. Ruiz³², I. Saviane⁹, M. R. Schreiber²⁰, A. C. Schröder^{53,54}, S. Sharma^{20,55}, M. D. Smith⁵⁶, L. Sodré Jr.²⁶,
 J. Stead³⁹, A. W. Stephens⁵⁷, M. Tamura⁵⁸, C. Tappert²⁰, M. A. Thompson⁴, E. Valenti⁵, L. Vanzì^{16,59}, N. A. Walton⁷,
 W. Weidmann²¹, and A. Zijlstra¹⁰

(Affiliations can be found after the references)

Received 5 November 2011 / Accepted 22 November 2011

ABSTRACT

Context. The ESO public survey VISTA variables in the Vía Láctea (VVV) started in 2010. VVV targets 562 sq. deg in the Galactic bulge and an adjacent plane region and is expected to run for about five years.

Aims. We describe the progress of the survey observations in the first observing season, the observing strategy, and quality of the data obtained.

Methods. The observations are carried out on the 4-m VISTA telescope in the *ZYJHK_s* filters. In addition to the multi-band imaging the variability monitoring campaign in the *K_s* filter has started. Data reduction is carried out using the pipeline at the Cambridge Astronomical Survey Unit. The photometric and astrometric calibration is performed via the numerous 2MASS sources observed in each pointing.

Results. The first data release contains the aperture photometry and astrometric catalogues for 348 individual pointings in the *ZYJHK_s* filters taken in the 2010 observing season. The typical image quality is $\sim 0''.9$ – $1''.0$. The stringent photometric and image quality requirements of the survey are satisfied in 100% of the *JHK_s* images in the disk area and 90% of the *JHK_s* images in the bulge area. The completeness in the *Z* and *Y* images is 84% in the disk, and 40% in the bulge. The first season catalogues contain 1.28×10^8 stellar sources in the bulge and 1.68×10^8 in the disk area detected in at least one of the photometric bands. The combined, multi-band catalogues contain more than 1.63×10^8 stellar sources. About 10% of these are double detections because of overlapping adjacent pointings. These overlapping multiple detections are used to characterise the quality of the data. The images in the *JHK_s* bands extend typically ~ 4 mag deeper than 2MASS. The magnitude limit and photometric quality depend strongly on crowding in the inner Galactic regions. The astrometry for *K_s* = 15–18 mag has rms ~ 35 –175 mas.

Conclusions. The VVV Survey data products offer a unique dataset to map the stellar populations in the Galactic bulge and the adjacent plane and provide an exciting new tool for the study of the structure, content, and star-formation history of our Galaxy, as well as for investigations of the newly discovered star clusters, star-forming regions in the disk, high proper motion stars, asteroids, planetary nebulae, and other interesting objects.

Key words. Galaxy: bulge – Galaxy: disk – Galaxy: stellar content – stars: abundances – infrared: stars – surveys

Figure D.2: Abstract of Saito et al. (2012).

A&A 544, A147 (2012)
 DOI: [10.1051/0004-6361/201219448](https://doi.org/10.1051/0004-6361/201219448)
 © ESO 2012

**Astronomy
&
Astrophysics**

Milky Way demographics with the VVV survey^{★,★★}

I. The 84-million star colour–magnitude diagram of the Galactic bulge

R. K. Saito^{1,2,3}, D. Minniti^{1,3,4}, B. Dias⁵, M. Hempel^{1,3}, M. Rejkuba⁶, J. Alonso-García^{1,3}, B. Barbuy⁵, M. Catelan^{1,3},
 J. P. Emerson⁷, O. A. Gonzalez⁶, P. W. Lucas⁸, and M. Zoccali^{1,3}

¹ Departamento de Astronomía y Astrofísica, Pontificia Universidad Católica de Chile, Vicuña Mackenna 4860, Casilla 306, Santiago 22, Chile
 e-mail: rsaito@astro.puc.cl

² Departamento de Física y Astronomía, Facultad de Ciencias, Universidad de Valparaíso, Ave. Gran Bretaña 1111, Playa Ancha, Casilla 5030, Valparaíso, Chile

³ The Milky Way Millennium Nucleus, Av. Vicuña Mackenna 4860, 782-0436 Macul, Santiago, Chile

⁴ Vatican Observatory, Vatican City State V-00120, Italy

⁵ Universidade de São Paulo, IAG, Rua do Matão 1226, Cidade Universitária, 05508-900 São Paulo, Brazil

⁶ European Southern Observatory, Karl-Schwarzschild-Strasse 2, 85748 Garching b. München, Germany

⁷ Astronomy Unit, School of Physics and Astronomy, Queen Mary University of London, Mile End Road, London, E1 4NS, UK

⁸ Centre for Astrophysics Research, University of Hertfordshire, College Lane, Hatfield AL10 9AB, UK

Received 20 April 2012 / Accepted 20 June 2012

ABSTRACT

Context. The Milky Way (MW) bulge is a fundamental Galactic component for understanding the formation and evolution of galaxies, in particular our own. The ESO Public Survey VISTA Variables in the Vía Láctea is a deep near-IR survey mapping the Galactic bulge and southern plane. Particularly for the bulge area, VVV is covering ~ 315 deg². Data taken during 2010 and 2011 covered the entire bulge area in the JHK_s bands.

Aims. We used VVV data for the whole bulge area as a single and homogeneous data set to build for the first time a single colour–magnitude diagram (CMD) for the entire Galactic bulge.

Methods. Photometric data in the JHK_s bands were combined to produce a single and huge data set containing 173 150 467 sources in the three bands, for the ~ 315 deg² covered by VVV in the bulge. Selecting only the data points flagged as stellar, the total number of sources is 84 095 284.

Results. We built the largest colour–magnitude diagrams published up to date, containing 173.1+ million sources for all data points, and more than 84.0 million sources accounting for the stellar sources only. The CMD has a complex shape, mostly owing to the complexity of the stellar population and the effects of extinction and reddening towards the Galactic centre. The red clump (RC) giants are seen double in magnitude at $b \sim -8^\circ$ – 10° , while in the inner part ($b \sim -3^\circ$) they appear to be spreading in colour, or even splitting into a secondary peak. Stellar population models show the predominance of main-sequence and giant stars. The analysis of the outermost bulge area reveals a well-defined sequence of late K and M dwarfs, seen at $(J - K_s) \sim 0.7$ – 0.9 mag and $K_s \gtrsim 14$ mag.

Conclusions. The interpretation of the CMD yields important information about the MW bulge, showing the fingerprint of its structure and content. We report a well-defined red dwarf sequence in the outermost bulge, which is important for the planetary transit searches of VVV. The double RC in magnitude seen in the outer bulge is the signature of the X-shaped MW bulge, while the spreading of the RC in colour, and even its splitting into a secondary peak, are caused by reddening effects. The region around the Galactic centre is harder to interpret because it is strongly affected by reddening and extinction.

Key words. Galaxy: bulge – Galaxy: center – Galaxy: structure – Galaxy: stellar content – surveys

Figure D.3: Abstract of Saito et al. (2012).

A&A 561, A106 (2014)
 DOI: [10.1051/0004-6361/201322092](https://doi.org/10.1051/0004-6361/201322092)
 © ESO 2014

**Astronomy
&
Astrophysics**

Self-consistent physical parameters for five intermediate-age SMC stellar clusters from CMD modelling^{★,★★,★★★}

B. Dias^{1,2}, L. O. Kerber^{1,3}, B. Barbuy¹, B. Santiago⁴, S. Ortolani⁵, and E. Balbinot⁴

¹ Instituto de Astronomia, Geofísica e Ciências Atmosféricas, Universidade de São Paulo, Rua do Matão 1226, Cidade Universitária, 05508-900 São Paulo, SP, Brazil

e-mail: bdias@astro.iag.usp.br

² European Southern Observatory, Alonso de Cordova 3107, Santiago, Chile

³ LATO-DCET-UESC, Rodovia Ilhéus-Itabuna km 16, Ilhéus, Bahia 45662-000, Brazil

⁴ Universidade Federal do Rio Grande do Sul, IF, CP 15051, 91501-970, RS, Porto Alegre, Brazil

⁵ Dipartimento di Fisica e Astronomia Galileo Galilei, University of Padova, vicolo dell'Osservatorio 3, 35122 Padova, Italy

Received 16 June 2013 / Accepted 15 November 2013

ABSTRACT

Context. Stellar clusters in the Small Magellanic Cloud (SMC) are useful probes for studying the chemical and dynamical evolution of this neighbouring dwarf galaxy, enabling inspection of a large period covering over 10 Gyr.

Aims. The main goals of this work are the derivation of age, metallicity, distance modulus, reddening, core radius, and central density profiles for six sample clusters, in order to place them in the context of the Small Cloud evolution. The studied clusters are AM 3, HW 1, HW 34, HW 40, Lindsay 2, and Lindsay 3; HW 1, HW 34, and Lindsay 2 are studied for the first time.

Methods. Optical colour–magnitude diagrams ($V, B - V$ CMDs) and radial density profiles were built from images obtained with the 4.1 m Southern Astrophysical Research (SOAR) telescope, reaching $V \sim 23$. The determination of structural parameters were carried out by applying King profile fitting. The other parameters were derived in a self-consistent way by means of isochrone fitting, which uses likelihood statistics to identify the synthetic CMDs that best reproduce the observed ones. Membership probabilities were determined comparing the cluster and control field CMDs. Completeness and photometric uncertainties were obtained by performing artificial star tests.

Results. The results confirm that these clusters (except HW 34, identified as a field fluctuation) are intermediate-age clusters, with ages between 1.2 Gyr (Lindsay 3) and ~ 5.0 Gyr (HW 1). In particular HW 1, Lindsay 2 and Lindsay 3 are located in a region that we called West Halo, where studies of ages and metallicity gradients are still lacking. Moreover, Lindsay 2 was identified as a moderately metal-poor cluster with $[\text{Fe}/\text{H}] = -1.4 \pm 0.2$ dex, lower than expected from the age-metallicity relation by Pagel & Tautvaisiene (1998). We also found distances varying from ~ 53 kpc to 66 kpc, compatible with the large depth of the SMC.

Key words. galaxies: star clusters: general – Magellanic Clouds – Hertzsprung-Russell and C-M diagrams

Figure D.4: Abstract of Dias et al. (2014).

A&A 563, A76 (2014)
 DOI: 10.1051/0004-6361/201322929
 © ESO 2014

**Astronomy
&
Astrophysics**

Deriving metallicities from calcium triplet spectroscopy in combination with near-infrared photometry^{★,★★,★★★}

F. Mauro¹, C. Moni Bidin^{1,2}, D. Geisler¹, I. Saviane³, G. S. Da Costa⁴, A. C. Gormaz-Matamala¹, S. Vasquez^{3,5},
 A.-N. Chené^{1,6,7}, R. Cohen¹, and B. Dias^{3,8}

¹ Departamento de Astronomía, Universidad de Concepción, Casilla 160-C, Concepción, Chile
 e-mail: skz5k2@gmail.com

² Instituto de Astronomía, Universidad Católica del Norte, Av. Angamos 0610, Casilla 1280, Antofagasta, Chile

³ European Southern Observatory, Ave. Alonso de Cordova 3107, Casilla 19, 19001 Santiago, Chile

⁴ Research School of Astronomy & Astrophysics, Australian National University, Canberra ACT 0200, Australia

⁵ Instituto de Astrofísica, Facultad de Física, Pontificia Universidad Católica de Chile, Av. Vicuña Mackenna 4860, Santiago, Chile

⁶ Gemini Observatory, Hawaii, USA

⁷ Departamento de Física y Astronomía, Universidad de Valparaíso, Av. Gran Bretaña 1111, Casilla 5030, Playa Ancha, Chile

⁸ Instituto de Astronomia, Geofísica e Ciências Atmosféricas, Universidade de São Paulo, rua do Matão 1226, Cidade Universitária, 05508-900 São Paulo, Brazil

Received 28 October 2013 / Accepted 10 January 2014

ABSTRACT

Context. When they are established with sufficient precision, the ages, metallicities and kinematics of Galactic globular clusters (GGCs) can shed much light on the dynamical and chemical evolution of the Galactic halo and bulge. While the most fundamental way of determining GC abundances is by means of high-resolution spectroscopy, in practice this method is limited to only the brighter stars in the nearest and less reddened objects. This restriction has, over the years, led to the development of a large number of techniques that measure the overall abundance indirectly from parameters that correlate with overall metallicity. One of the most efficient methods is measuring the equivalent width (EW) of the calcium II triplet (CaT) at $\lambda \approx 8500$ Å in red giants, which are corrected for the luminosity and temperature effects using the V magnitude differences from the horizontal branch (HB).

Aims. We establish a similar method in the near-infrared (NIR), by combining the power of the differential magnitudes technique with the advantages of NIR photometry to minimize differential reddening effects.

Methods. We used the K_s magnitude difference between the star and the reddest part of the HB (RHB) or of the red clump (RC) to generate reduced equivalent widths (rEW) from previously presented datasets. Then we calibrated these rEW against three previously reported different metallicity scales; one of which we corrected using high-resolution spectroscopic metallicities.

Results. We calculated the calibration relations for the two datasets and the three metallicity scales and found that they are approximately equivalent, with almost negligible differences. We compared our NIR calibrations with the corresponding optical ones, and found them to be equivalent, which shows that the luminosity-corrected rEW using the K_s magnitude is compatible with the one obtained from the V magnitude. We then used the metallicities obtained from the calibration to investigate the internal metallicity distributions of the GCs.

Conclusions. We have established that the $([\text{Fe}/\text{H}]; \text{rEW})$ relation is independent of the magnitude used for the luminosity correction and find that the calibration relations change only slightly for different metallicity scales. The CaT technique using NIR photometry is thus a powerful tool to derive metallicities. In particular, it can be used to study the internal metallicity spread of a GC. We confirm the presence of at least two metallicity populations in NGC 6656 and find that several other GCs present peculiar metallicity distributions.

Key words. stars: abundances – globular clusters: general – stars: kinematics and dynamics

Figure D.5: Abstract of Mauro et al. (2014).

FORS2/VLT survey of Milky Way globular clusters

I. Description of the method for derivation of metal abundances in the optical and application to NGC 6528, NGC 6553, M 71, NGC 6558, NGC 6426 and Terzan 8 [★]

B. Dias^{1,2}, B. Barbuy¹, I. Saviane², E. V. Held³, G. S. Da Costa⁴, S. Ortolani^{3,5}, S. Vasquez^{2,6}, M. Gullieuszik³, and D. Katz⁷

¹ Universidade de São Paulo, Dept. de Astronomia, Rua do Matão 1226, São Paulo 05508-090, Brazil
e-mail: bdias@astro.iag.usp.br

² European Southern Observatory, Alonso de Cordova 3107, Santiago, Chile

³ INAF, Osservatorio Astronomico di Padova, Vicolo dell'Osservatorio 5, 35122 Padova, Italy

⁴ Research School of Astronomy & Astrophysics, Australian National University, Mount Stromlo Observatory, via Cotter Road, Weston Creek, ACT 2611, Australia

⁵ Università di Padova, Dipartimento di Astronomia, Vicolo dell'Osservatorio 2, 35122 Padova, Italy

⁶ Instituto de Astrofísica, Facultad de Física, Pontificia Universidad Católica de Chile, Casilla 306, Santiago 22, Chile

⁷ GEPI, Observatoire de Paris, CNRS, Université Paris Diderot, 5 Place Jules Janssen 92190 Meudon, France

Received: ; accepted:

ABSTRACT

Context. We have observed almost 1/3 of the globular clusters in the Milky Way, targeting distant and/or highly reddened objects, besides a few reference clusters. A large sample of red giant stars was observed with FORS2@VLT/ESO at $R \sim 2000$. The method for derivation of stellar parameters is presented with application to six reference clusters.

Aims. We aim at deriving the stellar parameters effective temperature, gravity, metallicity and alpha-element enhancement, as well as radial velocity, for membership confirmation of individual stars in each cluster. We analyse the spectra collected for the reference globular clusters NGC 6528 ($[\text{Fe}/\text{H}] \sim -0.1$), NGC 6553 ($[\text{Fe}/\text{H}] \sim -0.2$), M 71 ($[\text{Fe}/\text{H}] \sim -0.8$), NGC 6558 ($[\text{Fe}/\text{H}] \sim -1.0$), NGC 6426 ($[\text{Fe}/\text{H}] \sim -2.1$) and Terzan 8 ($[\text{Fe}/\text{H}] \sim -2.2$). They cover the full range of globular cluster metallicities, and are located in the bulge, disc and halo.

Methods. Full spectrum fitting techniques are applied, by comparing each target spectrum with a stellar library in the optical region at 4560-5860 Å. We employed the library of observed spectra MILES, and the synthetic library by Coelho et al. (2005). Validation of the method is achieved through recovery of the known atmospheric parameters for 49 well-studied stars that cover a wide range in the parameter space.

Results. We identified 4 member stars in NGC 6528, 13 in NGC 6553, 10 in M 71, 5 in NGC 6558, 5 in NGC 6426 and 12 in Terzan 8. Radial velocities, effective temperatures, gravities, metallicities and alpha-element enhancements were derived. We derived $\langle v_{\text{helio}} \rangle = -242 \pm 11$ km/s, $[\text{Fe}/\text{H}] = -2.39 \pm 0.04$, $[\text{Mg}/\text{Fe}] = 0.38 \pm 0.06$ for NGC 6426 from spectroscopy for the first time.

Conclusions. The method proved to be reliable for red giant stars observed with resolution $R \sim 2000$, yielding results compatible with high-resolution spectroscopy. The derived α -element abundances show $[\alpha/\text{Fe}]$ vs. $[\text{Fe}/\text{H}]$ consistent with that of field stars at the same metallicities.

Key words. Stars: abundances - Stars: kinematics and dynamics - Stars: Population II - Galaxy: globular clusters - Galaxy: globular clusters: individual: NGC 6528, NGC 6553, M 71, NGC 6558, NGC 6426, Terzan 8 - Galaxy: stellar content

Figure D.6: Abstract of Dias et al. (2014b), submitted to A&A and accepted after revision.