UNIVERSIDADE DE SÃO PAULO
FACULDADE DE MEDICINA DE RIBEIRÃO PRETO

Patrícia Alves de Castro

Identificação dos mecanismos de letalidade da própolis em
Saccharomyces cerevisiae e *Candida albicans*

Ribeirão Preto

2012

Patrícia Alves de Castro

Identificação dos mecanismos de letalidade da própolis em
Saccharomyces cerevisiae e *Candida albicans*

Dissertação de Mestrado apresentada à
Faculdade de Medicina de Ribeirão Preto da
Universidade de São Paulo para obtenção do título
de Mestre em Ciências.

Área de Concentração: Bioquímica

Orientador: Prof. Dr. Gustavo Henrique Goldman

Ribeirão Preto

2012

FICHA CATALOGRÁFICA

AUTORIZO A REPRODUÇÃO E DIVULGAÇÃO TOTAL OU PARCIAL DESTE TRABALHO, POR QUALQUER MEIO CONVENCIONAL OU ELETRÔNICO, PARA FINS DE ESTUDO E PESQUISA, DESDE QUE CITADA A FONTE.

de Castro, Patrícia Alves

Identificação dos mecanismos de letalidade da própolis em *Saccharomyces cerevisiae* e *Candida albicans*. Ribeirão Preto, 2012.

140p.: il.; 30 cm

Dissertação de Mestrado, apresentada à Faculdade de Medicina de Ribeirão Preto/USP – Área de Concentração: Bioquímica.

Orientador: Goldman, Gustavo Henrique.

1. Própolis. 2. Morte celular. 3. *Saccharomyces cerevisiae*. 4. *Candida albicans*. 5. Atividade antifúngica. 6. Microarray. 7. Biologia de sistemas.

Patrícia Alves de Castro

Identificação dos mecanismos de letalidade da própolis em *Saccharomyces cerevisiae* e *Candida albicans*.

Dissertação de Mestrado apresentada à
Faculdade de Medicina de Ribeirão Preto da
Universidade de São Paulo para obtenção do título
de Mestre em Ciências.

Área de Concentração: Bioquímica

Orientador: Prof. Dr. Gustavo Henrique Goldman

Aprovado em: ____/____/____

Banca Examinadora

Prof(a) Dr(a). _____

Instituição: _____ Assinatura: _____

Prof(a) Dr(a). _____

Instituição: _____ Assinatura: _____

Prof(a) Dr(a). _____

Instituição: _____ Assinatura: _____

Dedicatória

*Dedico este trabalho aos meus queridos pais Ângela e Edson
pelo exemplo de vida, confiança, dedicação, apoio e amor
incondicionais e também pela educação, valores ensinados e por
sempre me incentivarem a estudar.*

*Ao Gesiel, meu namorado, pelo amor, confiança, compreensão e
incentivo.*

À minha irmã, Daniela, pela sua alegria, amizade, amor...

Vocês são muito importantes para mim!

Agradecimentos Especiais

Ao meu orientador, Professor Dr. **Gustavo Henrique Goldman**, idealizador deste trabalho, obrigada pela grande oportunidade, por todo apoio profissional, intelectual e pelos ensinamentos compartilhados desde a minha Iniciação científica.

À Fundação de Amparo à Pesquisa do Estado de São Paulo (**FAPESP**), pelo auxílio financeiro ao laboratório e pelas bolsas de iniciação científica e de mestrado concedidas a mim.

Ao Conselho Nacional de Desenvolvimento Científico e Tecnológico (**CNPq**) e à Financiadora de Estudo e Projetos (**FINEP**) pelo apoio financeiro ao laboratório.

À empresa **Apis Flora**, por acreditar neste projeto e por todo apoio fornecido, em especial à Dra. **Andresa A. Berretta e Silva** pela sua colaboração profissional e sua constante gentileza.

Agradecimentos

Agradeço imensamente a todos que profissionalmente, intelectualmente e também emocionalmente contribuíram para meu crescimento profissional e pessoal. Tenho todos em meu coração!

Aos meus pais, **Edson** e **Ângela**, que sempre me ofereceram o máximo de si mesmos, torceram por mim e acreditaram nos meus sonhos. Sem dúvidas, essa conquista foi devido ao grande amor, incentivo, carinho, compreensão, ensinamentos, preocupação que sempre recebi dos meus dois anjos... amarei vocês eternamente!!! E você também **Dani**... minha companheirinha, a melhor irmã do mundo e que eu adoro muito!!! Não há palavras para agradecer a vocês tudo o que fazem por mim.

Ao meu namorado **Gesiel**, pelo seu carinho, atenção, respeito, dedicação... por acreditar, apoiar e incentivar sempre meus objetivos e sonhos. Pela sua compreensão e paciência (e quanta, não é mesmo Ge!!) e também por todo amor ao longo desses anos. Amo muito você!!!

Aos meus avôs **Sebastião** e **Edison** (que me olha lá do céu) e às minhas avós **Elena**, **Elza** e à minha bisã **Rosa** pelo grande amor, dedicação e apoio. Aos meus tios, tias, primos e primas que torceram e me incentivaram durante mais esta etapa da minha vida. Tio **Norberto**, muito obrigada pelas eternas ajudas em química!!!

À minha segunda família D. **Creusa**, Sr. **Antônio** e aos meus cunhados e cunhadas pelo apoio e carinho.

Desde que cheguei ao laboratório conheci pessoas maravilhosas, sempre dispostas a ajudar e, por isso, gostaria de agradecer muito...

... ao Dr. **Iran Malavazi** e à Dra. **Marcela Savoldi** pela amizade, paciência e dedicação em ensinar desde a minha entrada no laboratório em 2006, e também pela grande ajuda com os microarrays...

... aos amigos do laboratório, deste e de outros tempos, Prof^a. Dra. **Márcia F. Balieiro**, Prof^a. Dra. **Márcia Kress**, Prof^a. Dra. **Taisa M. Dinamarco**, Dr. **Frederico Soriani**, Ms. **Paula F. G. Bizzi**, **Ana C. Colabardini**, Ms. **Bárbara C. P. Figueiredo**, Dr. **Wagner R. de Souza**, Ms. **Lívia M. Alfaya**, Prof. Dr. **Ricardo S. C. de Almeida** (pelo ensaio de inibição da transição dimórfica em *C. albicans*!), Dra. **Thaila F. dos Reis**, Dra. **Nádia Krohn**, Dr. **Neil A. Brown**, **José L. Capellaro**, Dr. **Luciano A. Bernardes**, Dra. **Mônica Campiteli**, Dr. **André Mota Jr.**, Ms **Joel F. Lima**, **Sara R. Carvalho**, **Angélica Batista**, **Paulo S. de Carvalho**, **Luciano A. Takami**. Obrigada por proporcionarem um ótimo convívio, pela amizade e imensas ajudas!!!

Ao Prof. Dr. **Mário Henrique de Barros**, Instituto de Ciências Biomédicas - USP, por fornecer as linhagens "petite" de *S. cerevisiae* para este trabalho, pela colaboração no ensaio de consumo de oxigênio e pela ajuda nos estudos de fracionamento mitocondrial.

Ao Prof. Dr. **Diego Bonatto**, Universidade Federal do Rio Grande do Sul, pela grande contribuição e gentileza em me ensinar as análises por Biologia de Sistemas.

À Prof^a. Dr. **Maria Helena de Souza Goldman**, Faculdade de Filosofia Ciências e Letras - USP, Ribeirão Preto pelo apoio e por todo suporte técnico.

A todos os professores que contribuíram com o envio de cepas, plasmídeos, troca de informações e sugestões.

À Faculdade de Medicina de Ribeirão Preto, USP e aos professores do Departamento de Bioquímica, pelas aulas e seminários ministrados, aos funcionários da seção de Pós-Graduação, por me ajudarem a resolver os problemas burocráticos e, em especial, à secretária do Departamento, a **Ivone**, que sempre foi muito atenciosa, gentil e paciente comigo.

Às minhas amigas **Luciana**, **Clévia** e **Nathália**, obrigada pelos momentos de alegria que passamos juntas e pelo carinho desde a graduação e também à super **Barbarela**, metade podre da minha laranja!!!

À **Deus**, por estar comigo em todos os momentos da minha vida...

... Tudo acontece na hora certa.

Tudo acontece, exatamente quando deve acontecer....

(Albert Einstein)

Obrigada a todos que contribuíram de forma intensa e carinhosa, direta ou indiretamente, para a realização deste projeto!!!!

Não me lembro em que momento percebi que viver deveria ser reinvenção de nós mesmos – para não morrermos soterrados na poeira da banalidade embora pareça que ainda estamos vivos. (...)

Para viver de verdade, pensando e repensando a existência, para que ela valha a pena, é preciso ser amado; e amar; e amar-se.

Ter esperança; qualquer esperança.

Questionar o que nos é imposto, sem rebeldias insensatas, mas sem demasiada sensatez.

Saborear o bom, mas aqui e ali enfrentar o ruim. Suportar sem se submeter, aceitar sem se humilhar, entregar-se sem renunciar a si mesmo e à possível dignidade.

Sonhar, porque se desistirmos disso apaga-se a última claridade e nada mais valerá a pena (...)

E que o mínimo que a gente faça seja, a cada momento, o melhor que afinal se conseguiu fazer.

Lya Luft

RESUMO

A própolis é uma mistura resinosa complexa de várias substâncias coletada de plantas pelas abelhas. Ela tem atraído atenção devido à variedade de suas propriedades biológicas e terapêuticas. Diversos estudos têm mostrado a conexão existente entre a morte celular tipo-apoptose em fungos e importantes processos biológicos como desenvolvimento, envelhecimento, resposta a estresse e patogênese. Neste contexto, este projeto avaliou a atividade antifúngica da própolis, com o objetivo de ampliar os conhecimentos acerca das vias metabólicas de morte celular em fungos, como *S. cerevisiae* e *C. albicans*, e também em relação à utilização da própolis como uma terapia antifúngica mais efetiva. Inicialmente, utilizou-se *S. cerevisiae* com o objetivo de compreender como a própolis afeta fungos ao nível celular. Foi observado que ela é capaz de induzir uma resposta de morte celular apoptótica. No entanto, a exposição aumentada à própolis promove um correspondente aumento na resposta tipo necrose. Verificou-se ainda que o citocromo c, mas não a endonuclease G Nuc1p, está envolvido na morte celular mediada por própolis em *S. cerevisiae*. Também foi observado que o gene da metacaspase *YCA1* é importante para a morte celular induzida por esta substância natural. Para elucidar as funções dos genes que poderiam ser necessários para a sensibilidade à própolis em eucariotos, realizou-se um *screening* da coleção completa com cerca de 4.800 cepas haplóides com genes únicos deletados de *S. cerevisiae*. Foram identificadas 138 cepas que apresentaram diferentes graus de sensibilidade à própolis quando comparadas com a cepa do tipo selvagem correspondente. Na análise deste screening por biologia de sistemas e também através do perfil transcricional de *S. cerevisiae* exposto à própolis, foram observados genes envolvidos na cadeia de transporte de elétrons mitocondrial, da acidificação vacuolar, da regulação negativa da transcrição do promotor da RNA polimerase II, da regulação da macroautofagia associada com a proteína alvo para vacúolo e da resposta celular à privação de nutrientes. Os estudos de validação indicaram que a sensibilidade da própolis é dependente da função mitocondrial e que a acidificação vacuolar e autofagia são importantes para a morte causada por própolis em leveduras. Para o fungo patogênico *C. albicans*, foi observado que a própolis induz uma morte celular do tipo necrose; observou-se ainda que o gene *IPF4847* (homólogo para o gene da metacaspase *YCA1* de *S. cerevisiae*) é importante para a morte celular mediada por própolis. Além disso, com o objetivo de tentar esclarecer algumas funções de genes que poderiam estar envolvidos na sensibilidade à própolis de *C. albicans*, 800 mutantes deletados de *C. albicans* foram escaneados e destes, 51 apresentaram maior sensibilidade a própolis quando comparados com as cepas do tipo selvagem correspondente. Vários genes observados em nosso “screening” estão envolvidos na transição dimórfica em *C. albicans*. Desta forma, foi realizado um ensaio a fim de se verificar o papel da própolis na inibição da transição dimórfica e foi observado que este composto inibe não só a transição dimórfica como também o crescimento de todos os morfotipos (levedura, hifa, e pseudohifa) de *C. albicans*. Desta maneira, o uso da própolis para o tratamento clínico da candidíase pode ter grande aplicação, principalmente por afetar um mecanismo importante para a patogenicidade deste fungo.

ABSTRACT

Propolis is a complex mixture of several resinous substances which are collected from plants by bees. Propolis has attracted the attention of researchers because of its variety of biological and therapeutic properties. Studies have shown the connection between Propolis and apoptosis-like cell death in fungi and other important biological processes such as development, aging, stress response and pathogenesis. In this context, this project evaluated the antifungal activity of Propolis, with the aim of expanding the knowledge about the metabolic pathways of cell death in fungi such as *S. cerevisiae* and *C. albicans*, and also in relation to the use of propolis as an effective antifungal therapy. For this, initially we utilised *S. cerevisiae* as a model organism to study the genetics, cell biology and genomics that determine how propolis affects fungi at the cellular level. Propolis is able to induce an apoptosis cell death response. However, increased exposure to propolis provides a corresponding increase in the necrosis response. We showed that cytochrome c, but not endonuclease G (Nuc1p), is involved in propolis-mediated cell death in *S. cerevisiae*. We also observed that the metacaspase *YCA1* gene is important for propolis-mediated cell death. To elucidate the gene functions that may be required for propolis sensitivity in eukaryotes, the full collection of approximately 4,800 haploid *S. cerevisiae* deletion strains was screened for propolis sensitivity. We were able to identify 138 deletion strains that have different degrees of propolis sensitivity compared to the corresponding wild-type strains. Systems biology revealed enrichment for genes involved in the mitochondrial electron transport chain, vacuolar acidification, negative regulation of transcription from RNA polymerase II promoter, regulation of macroautophagy associated with protein targeting to vacuoles, and cellular responses to starvation. Validation studies indicated that propolis sensitivity is dependent on mitochondrial function and that vacuolar acidification and autophagy are important for *S. cerevisiae* cell death caused by propolis. For the pathogenic fungus *C. albicans* it was observed that propolis induces cell death like necrosis. It was also observed that the *IPF4847* gene (homologous to the *YCA1* metacaspase gene of *S. cerevisiae*) was also important for cell death mediated by propolis in *C. albicans*. Furthermore, aiming to clarify some of the functions of genes that could be involved in *C. albicans* sensitivity to propolis, 800 *C. albicans* deletion mutants were screened, and 51 showed greater sensitivity to propolis compared with the corresponding wild-type strains. Several genes found in our screening were involved in the dimorphic transition of *C. albicans*. Thus, an assay was performed in order to verify the role of propolis in the inhibition of the dimorphic switch. It was observed that propolis can inhibit the dimorphic transition and the growth of all its morphotypes (budding hyphal and pseudohyphal) in *C. albicans*. So, the use of propolis for clinical treatment of Candidiasis may have a wide application, principally by affecting an important mechanism for the pathogenicity of fungi.

LISTA DE FIGURAS

Figura 1: Perfil cromatográfico de (a) Extrato Padronizado de Própolis (EPP) Lote 1440110; (b) EPP Lote 1400410 e (c) EPP Lote 010/08.....	61
Figura 2: A própolis induz apoptose e necrose tardia em <i>S. cerevisiae</i>	65
Figura 3: Ensaio do TUNEL (<i>Terminal Deoxynucleotidyl Transferase–mediated dUTP Nick End Labeling</i>)..	66
Figura 4: A própolis induz a produção de ROS.	67
Figura 5: O mutante da metacaspase <i>YCA1</i> é mais tolerante a própolis que a linhagem selvagem correspondente.....	68
Figura 6: Exposição de células de levedura a própolis induz a liberação do citocromo c para o citoplasma.....	69
Figura 7: Nuc1p não está envolvido na sensibilidade à própolis.	70
Figura 8: Sistemas biológicos para a sensibilidade à própolis em <i>S. cerevisiae</i>	72
Figura 9: (A) Sub-redes de proteínas (cluster 6) associadas à divisão celular, estabelecimento da polaridade celular, cadeia de transporte de elétrons e acidificação vacuolar. (B) Sub-redes de proteínas <i>Unclustered</i>	74
Figura 10: Mutantes “ <i>Petite</i> ” são mais tolerantes a própolis.	78
Figura 11: Própolis induz acidificação vacuolar em <i>S. cerevisiae</i>	80
Figura 12: Própolis induz autofagia em <i>S. cerevisiae</i>	82
Figura 13: Análise da Ontologia Gênica dos genes com abundância de RNAm aumentada no ensaio de hibridação de microarray de <i>S. cerevisiae</i> exposta a própolis.....	89
Figura 14: Análise da Ontologia Gênica dos genes com abundância de RNAm diminuída no ensaio de hibridação de microarray de <i>S. cerevisiae</i> exposta a própolis.....	92
Figura 15: PCR em tempo real para os genes selecionados da análise de hibridação de microarray.	95
Figura 16: Análise, por Diagrama de Venn, do grau de sobreposição dos nós das redes PPPI para os genes induzidos e reprimidos na análise do microarray.....	98

Figura 17: Ensaio de diluição para cepas de <i>C. albicans</i> em meio YPD complementado com diferentes concentrações de própolis.....	103
Figura 18: Viabilidade de células de <i>C. albicans</i> em diferentes fases do ciclo de crescimento e expostas a própolis (0,125; 0,250; 0,500 e 1%), por 20 minutos.	104
Figura 19: Viabilidade de células de <i>C. albicans</i> em expostas a 1% de própolis, por 5, 10 e 20 minutos.....	104
Figura 20: A viabilidade de células de <i>C. albicans</i> em expostas a própolis.	106
Figura 21: Os mutantes para o gene <i>IPF4847</i> são mais tolerantes a própolis que a linhagem selvagem correspondente.....	107
Figura 22: Os mutantes para os <i>cdc35Δ</i> , <i>ras1Δ</i> , são mais tolerantes a própolis que a linhagem selvagem correspondente e também que o mutante <i>pde2Δ</i>	108
Figura 23: Ensaio de diluição para as 3 formas morfológicas de <i>C. albicans</i>	112
Figura 24: Ensaio para verificar a influência na filamentação de <i>C. albicans</i>	113
Figura 25: A própolis afeta a filamentação em <i>C. albicans</i>	115

LISTA DE TABELAS

Tabela 1: Apresentação quantitativa de compostos do Extrato Padronizado de Própolis obtidos a partir de três diferentes lotes (média $\pm$ DP) (n = 3)	62
Tabela 2: Classes específicas <i>gene ontology</i> (GO) derivadas da rede de interação física proteína-proteína (PPPI) observadas no <i>cluster 6</i>	75
Tabela 3: Classes específicas <i>gene ontology</i> (GO) derivadas da rede de interação física proteína-proteína (PPPI) de sub-redes de proteínas <i>unclustered</i>	76
Tabela 4: Genes selecionados que foram mais expressos ($\log_2 \geq 1.0$) durante a exposição de <i>S. cerevisiae</i> a própolis (para uma lista completa de genes mais expressos, ver Anexo C).	90
Tabela 5: Genes selecionados que foram menos expressos ($\log_2 \geq 1.0$) durante a exposição de <i>S. cerevisiae</i> a própolis (para uma lista completa de genes mais expressos, ver Anexo C).	93
Tabela 6: Principais classes de ontologias gênicas observadas nos clusters derivados da rede PPPI fusionada	97

LISTA DE ABREVIATURAS, SIGLAS E SÍMBOLOS

CCCP	<i>Carbonyl cyanide m-chlorophenylhydrazone</i>
cDNA	DNA complementar
CT	<i>Cycle Threshold</i>
DAPI	4,6-diamino-2-fenilindol
DIC	<i>Differential interference contrast</i>
GFP	<i>Green fluorescent protein</i>
H ₂ DCFDA	2', 7'- dichlorodihydrofluorescein diacetate
Kb	Kilo base
Mb	Mega base
MDR	Resistência a múltiplas drogas
mRNA	RNA mensageiro
p/v	Peso/volume
PBS	Tampão salina fosfato
PCR	Reação em cadeia da polimerase
PI	Iodeto de Propídio
PPPI	Interação Física Proteína-Proteína
ROS	Espécies reativas de oxigênio
RT PCR	Reação em cadeia da polimerase em tempo real
SBF	Soro bovino fetal
v/v	Volume/volume
kDa	Kilodalton
wt	<i>Wild type</i>

SUMÁRIO

1. INTRODUÇÃO	18
1.1. Própolis	19
1.2. <i>Saccharomyces cerevisiae</i> e <i>Candida albicans</i>	22
1.2.1. <i>Saccharomyces cerevisiae</i>	22
1.2.2. <i>Candida albicans</i>	24
1.3. A Morte Celular	28
1.3.1 A morte celular e a mitocôndria.....	32
1.4. A morte celular em fungos.....	34
2. OBJETIVOS.....	39
3. MATERIAIS E MÉTODOS	41
3.1. Linhagens, meios e métodos de cultura	42
3.2. Preparação do extrato padronizado de própolis	42
3.3. Screening das coleções de <i>S. cerevisiae</i> e de <i>C. albicans</i>	44
3.3.1. Screening da coleção de <i>S. cerevisiae</i> das linhagens haplóides de genes não essenciais deletados para a sensibilidade a própolis.....	44
3.3.2. Screening com alta eficiência de bibliotecas de <i>C. albicans</i> para fatores de transcrição e quinases para sensibilidade à própolis	44
3.4. Extração de RNA e reação de PCR em tempo real para <i>S. cerevisiae</i>	45
3.5. Microscopia óptica e confocal	45
3.5.2. Ensaio do TUNEL (Terminal Deoxynucleotidyl Transferase-mediated dUTP Nick End Labeling).....	48
3.5.3. Ensaio de Anexina e Iodeto de Propídio (PI)	49
3.6. Estudos de fracionamento mitocondrial	50
3.7. O consumo de oxigênio.....	51
3.8. Projeto de rede de interação física proteína-proteína (PPPI) e análise topológica	51
3.8.1 Análise <i>Gene ontology</i> (GO)	52
3.9. Microarranjo de DNA (Microarray).....	52
3.9.1. Preparo das amostras.....	52
3.9.2. Isolamento do RNA e reação de PCR em tempo real	53
3.9.3. Hibridação do microarray	54
3.9.3.1. Marcação e hibridização dos RNAs e geração das imagens de microarranjo.....	54
3.9.3.2. Análise das imagens e dados obtidos dos microarranjos	55

3.9.4. Projeto e análise topológica global da rede de interação física proteína-proteína (PPPI)	56
3.9.4.1. Centralidades da rede e análises topológicas locais	57
3.9.4.2. Análise Gene ontology (GO).....	57
4. RESULTADOS E DISCUSSÃO	58
Módulo 4.1: Caracterização molecular da morte celular induzida por própolis em <i>Saccharomyces cerevisiae</i>	59
4.1. RESULTADOS	60
4.1.1. A própolis induz morte celular em <i>S. cerevisiae</i>	60
4.1.2. O citocrome c é liberado durante a morte celular induzida por própolis.....	68
4.1.3. Avaliação da biblioteca de deleção de levedura para sensibilidade à própolis	71
4.1.4. Linhagens “Petite” são menos sensíveis a própolis	77
4.1.5. Um papel para o vacúolo e a H ⁺ -ATPase vacuolar na sensibilidade a própolis	79
4.1.6. DISCUSSÃO	83
Módulo 4.2: Análise de sistemas de <i>Saccharomyces cerevisiae</i> exposta a própolis	87
4.2. RESULTADOS E DISCUSSÃO.....	88
4.2.1. A análise da hibridação por microarray.....	88
4.2.2. A validação da análise da hibridação por microarray.....	94
4.2.3. Análise por biologia de sistemas para a exposição a própolis	96
Módulo 4.3: A identificação dos mecanismos de letalidade da própolis em <i>Candida albicans</i>	101
4.3. RESULTADOS E DISCUSSÃO.....	102
4.3.1. A própolis induz morte celular em <i>C. albicans</i>	102
4.3.2. o Envolvimento das caspases e da via metabólica Ras na morte celular induzida por própolis em <i>C. albicans</i>	107
4.3.3. A avaliação de bibliotecas de deleções de <i>C. albicans</i> para sensibilidade à própolis	109
4.3.4. O papel da própolis na transição dimórfica de <i>C. albicans</i>	111
4.3.4.1. A própolis inibe os três morfotipos de <i>C. albicans</i>	111
4.3.4.2. A própolis inibe a transição dimórfica de levedura para hifa em <i>C. albicans</i>	113
5. CONCLUSÕES	116
6. REFERÊNCIAS BIBLIOGRÁFICAS	118
ANEXOS	140

1. Introdução

1.1. Própolis

A palavra própolis é derivada do grego *pro-*, em defesa, e *polis-*, cidade ou comunidade, isto é, em defesa da comunidade. É uma mistura complexa, formada por material resinoso, maleável e balsâmico coletado pelas abelhas dos ramos, flores, pólen, brotos e exudatos de árvores. Na colméia, as abelhas adicionam secreções próprias (salivares e enzimas) a esta mistura (LUSTOSA *et al.*, 2008; PEREIRA *et al.*, 2002; FRANCO *et al.*, 2000). Para a produção de própolis, as abelhas usam materiais resultantes de uma variedade de processos botânicos em diferentes partes das plantas, como substâncias ativamente secretadas por elas, bem como exudatos de feridas: materiais lipofílicos nas folhas e germes, látex, resinas, entre outros (LUSTOSA *et al.*, 2008; BANKOVA, 2005b; CAPASSO & CASTALDO, 2002). Tem sido sugerido que a própolis é usada como vedante nas colméias para bloquear aberturas desnecessárias e evitar a invasão por fungos, bactérias e insetos (CHEN, *et al.*, 2004), além de ser utilizada no preparo de locais assépticos para postura da abelha rainha, na mumificação de insetos invasores e para reparar e fortalecer os favos de mel (NUNES *et al.*, 2009; BANKOVA *et al.*, 2000).

A própolis é um dos muitos produtos naturais que vem sendo utilizado durante séculos pela humanidade (LUSTOSA *et al.*, 2008; VARGAS *et al.*, 2004), uma vez que o uso de extratos de própolis na medicina popular data de 300 a.C. (DA SILVA *et al.*, 2006). O termo própolis já era descrito no século XVI na França, e a Farmacopéia de Londres do século XVII lista a própolis como droga oficial (LUSTOSA *et al.*, 2008; CAPASSO & CASTALDO, 2002; PEREIRA *et al.*, 2002; CAO *et al.*, 2004). Esta substância resinosa é utilizada na medicina popular em muitas regiões do mundo (MARUYAMA, *et al.*, 2009). Os egípcios conheciam as propriedades anti-putrefativas da própolis e empregavam para embalsamar cadáveres. (CAPASSO & CASTALDO, 2002). Na África do Sul, na guerra ao final do século XIX, foi amplamente utilizada devido às suas propriedades cicatrizantes; e na segunda guerra mundial foi empregada em várias clínicas soviéticas (LUSTOSA *et al.*, 2008; PEREIRA *et al.*, 2002). Na ex-União Soviética e em países do leste da Europa, a própolis começou a ser apreciada como meio para tratamento de problemas de saúde nos anos de 1950 e 1960. Na metade dos anos 80, a própolis tornou-se um importante produto na medicina alternativa e complementar. O Japão é

o principal importador de própolis, com preferência pela própolis brasileira (LUSTOSA *et al*, 2008; SALATINO *et al.*, 2005), a qual é muito utilizada por eles como suplemento para a saúde (URUSHISAKI *et al*, 2011). No Brasil, o interesse pela própolis aconteceu somente na década de 80 com o trabalho pioneiro de Ernesto Ulrich Breyer, demonstrando em seu livro, "Abelhas e saúde", as propriedades terapêuticas da própolis e a sua utilização como antibiótico natural (LIMA, 2006). Tem sido relatado que a própolis, devido à sua composição complexa, possui várias atividades biológicas, tais como as atividades antibacteriana (KORU *et al*, 2007; PARK *et al*, 2002), antiviral (VYNOGRAD *et al*, 2000), antiprotozoária (MENNA-BARRETO *et al*, 2009), antiinflamatória (RAMOS e MIRANDA, 2007), antioxidante (SIMÕES *et al*, 2004; RUSSO *et al*, 2002), cicatrizante (BERRETTA *et al*, 2012), imunomodulatória (SFORCIN, 2007), além de possuir propriedades antitumorais (KIMOTO *et al*, 1998). Recentemente, há várias evidências mostrando que a própolis pode ser usada para tratar infecções fúngicas de *Candida* (AGUERO *et al*, 2010; DALKEN-DOTA *et al*, 2010; OTA *et al*, 2001; SALOMÃO *et al*, 2008; SIQUEIRA *et al*, 2009). Assim, a própolis tem atraído a atenção devido à variedade de suas propriedades biológicas e terapêuticas, porém, relatos sobre a sua atividade e os mecanismos de ação relacionados a ela em leveduras/linhagens dermatófitas são raros (SILICI, *et al*, 2005).

A própolis possui composição complexa e variada, dependendo basicamente das fontes de plantas disponíveis para as abelhas. Assim, amostras brasileiras apresentam diferenças notáveis na sua composição química quando comparadas com amostras das zonas temperadas. Além disso, diferenças também são encontradas entre amostras tropicais dependendo da flora do local de coleta (PARK, *et al*, 2002). A atividade antimicrobiana da própolis Européia tem sido associada com a presença de flavonóides e derivados do ácido caféico (HEGAZI, *et al*, 2000). No caso de amostras tropicais, os principais compostos bioativos são ácidos fenólicos e derivados prenilados (PRYTZYK, *et al*, 2003; SALOMÃO, *et al*, 2008). Em relação à composição da própolis brasileira, esta também varia em função da região de origem, mas de modo geral, encontramos compostos flavonóides, como quercetina, aromadendrina, pinocembrina, compostos fenólicos, com destaque para os derivados prenilados como artepelin C, baccharina e drupanin, além de ácidos como p-cumárico, caféico, ferúlico, benzóico, dentre outros (LEITÃO *et al*, 2004). No Brasil, a planta mais importante como fonte de própolis verde é a *Baccharis dracunculifolia*

D. C. (Asteraceae) (KUMAZAWA et al., 2004). Mais de 200 constituintes da própolis foram identificados; contudo, a atividade biológica destas substâncias ainda não é bem compreendida (BANKOVA, *et al*, 1995; CHEN, *et al*, 2004). A própolis contém cera, flavonóides e compostos fenólicos como ácido cinâmico e seus derivados, e também vários aldeídos, cetonas, aminoácidos (GREENAWAY, *et al*, 1991).

O éster fenetil do ácido cafêico (CAPE), componente característico da própolis européia, foi um dos componentes isolados e identificados que apresentou atividade antitumoral e antioxidante (CHEN, *et al*, 2004. Na concentração de 10 μ M, ele bloqueia completamente a produção de espécies reativas de oxigênio (ROS) em neutrófilos humanos e sistemas xantina/xantina oxidase (SUD'INA, *et al*, 1993), todavia, não há traços de CAPE na própolis verde brasileira. Chen *et al* (2004) isolaram e caracterizaram um tipo de prenilflavona, propolina C, da própolis de Taiwan e, foi verificado que esta substância é um poderoso indutor de apoptose em células de melanoma humano e um significativo inibidor da xantina oxidase. Urushisaki *et al* (2011) demonstraram que os ácidos cafeoilquínicos presentes na própolis verde brasileira apresentam atividade contra o vírus da influenza. Estes componentes podem restaurar a viabilidade das células de rim canino "Madin-Darby" (MDCK – Madin- Darby canine kidney cells) que foram infectadas com o vírus da gripe influenza, de maneira dose dependente. Além disso, pela análise combinada utilizando ensaio de viabilidade celular e PCR em tempo real, eles sugerem ser muito possível que estas substâncias não possuam efeito direto sobre o vírus ou interajam com componentes virais, mas que este produto natural pode ser utilizado como profilaxia ou como agente anti-influenza moderado.

Há uma necessidade urgente em descobrir novos medicamentos antifúngicos. Existem, atualmente, cinco classes diferentes de drogas antifúngicas classificadas com base em seus mecanismos de ação (KONTOYIANNIS & LEWIS, 2002). A maioria dessas drogas afeta a membrana celular ou a integridade da parede celular através da inibição da síntese de ergosterol ou da síntese de β -glucana/quitina, respectivamente. Uma complicação adicional para este espectro limitado de drogas é a aquisição de resistência por parte dos fungos patogênicos observados nos casos clínicos (GULSHAN & MOYE-ROWLEY, 2007; KANAFANI & PERFECT, 2008; MORSCHHÄUSER, 2010). Enquanto a anfotericina B tem atividade fungicida, os azólicos são geralmente os agentes fungistáticos para *Candida* spp e agentes fungicidas para *Aspergillus* spp (MANAVATHU *et al*, 1998). Equinocandinas

(caspofungina, micofungina e anidulafungina) exibem atividade fungicida contra *Candida* spp e atividade fungistática contra *Aspergillus* spp (GULLO, 2009).

Assim, as recentes descobertas sobre os possíveis mecanismos de ação da própolis, descritas neste trabalho, associadas às vantagens relacionadas à segurança e a ausência de efeitos adversos importantes (SENEDESE *et al*, 2011), proporcionam uma importante oportunidade para a utilização da própolis na geração de um medicamento que apresente um mecanismo de ação distinto em relação aos demais antifúngicos clássicos, somada às outras atividades conhecidas da própolis, como o aumento da ativação da imunidade (SFORCIN, 2007), e que podem contribuir para a maior resposta do paciente ao tratamento.

1.2. *Saccharomyces cerevisiae* e *Candida albicans*

1.2.1. *Saccharomyces cerevisiae*

S. cerevisiae foi introduzida por volta de 1960 como um sistema experimental para a biologia molecular. Já em 1980, a levedura foi utilizada para produzir vacina contra hepatite B. Em 1996, *S. cerevisiae* foi o primeiro organismo eucariótico que obteve sua sequência genômica completamente estabelecida (GOFFEAU, *et al*, 1996). Nos anos seguintes, a levedura se tornou uma referência útil para estudos comparados com sequência gênicas de humanos, animais, plantas e de vários organismos unicelulares. Além disso, a facilidade de manipulação genética em leveduras abriu a possibilidade de dissecar funcionalmente produtos de genes de outros eucariotos no sistema de levedura. Na era “pós-genômica”, a levedura estava novamente na vanguarda no campo da genômica funcional (FELDMANN, 2005). Nos últimos anos, todos os recursos do genoma que foram desenvolvidos para *S. cerevisiae* (www.yeastgenome.org), promoveram uma excelente plataforma genômica para o desenvolvimento de drogas e a identificação de seus alvos (BHARUCHA & KUMAR, 2007; HOON *et al*, 2008; MAERE *et al*, 2005; WUSTER & MADAN BABU, 2008).

Está bem estabelecido que a levedura é um sistema ideal em que a arquitetura celular e mecanismos celulares fundamentais podem ser investigados com sucesso. Entre todos os organismos eucarióticos modelos, *S. cerevisiae* combina várias vantagens. É um organismo unicelular que, ao contrário dos

eucariotos mais complexos, pode ser cultivado em meios de cultura definidos, dando o controle completo ao investigador em relação aos parâmetros ambientais. Ela pode ser manipulada pelas técnicas de genética clássica e várias funções têm sido estudadas em grande detalhe por abordagens bioquímicas (FELDMANN, 2005; STRATHERN *et al*, 1981; BROACH *et al*, 1991; GUTHRIE & FINK, 1991). Na verdade, uma grande variedade de exemplos fornece evidências de que as funções celulares substanciais são altamente conservadas da levedura aos mamíferos, e que genes correspondentes podem, muitas vezes, se complementarem funcionalmente (FELDMANN, 2005).

O genoma de *S. cerevisiae* é dividido em 16 cromossomos que variam em tamanho entre 250 kb e > 2500 kb. O ciclo de vida de *S. cerevisiae* normalmente alterna entre as fases diplóide e haplóide. Ambas ploidias podem existir como culturas estáveis. Em cepas heterotáticas, células haplóides são de dois “mating types”, a e α . O “mating” das células a e α resultam em diplóides a/ α , que são incapazes de cruzar, mas podem sofrer meiose. Os quatro produtos haplóides resultantes da meiose de uma célula diplóide estão contidos dentro da parede da célula-mãe (o asco). A digestão dos ascos e separação dos esporos por micromanipulação fornece os quatro produtos meióticos haplóides. A análise dos padrões de segregação dos diferentes marcadores heterozigóticos entre os quatro esporos constitui análise de tétrades e revela a ligação entre dois genes (ou entre um gene e seu centrômero) (MORTIMER & SCHILD, 1991). No conjunto, a distância genética em levedura parece ser notavelmente proporcional à distância física, com uma média global de 3 kb/cM (FELDMANN, 2005).

Em uma levedura, podem-se observar as seguintes estruturas celulares: parede celular, periplasma, membrana plasmática, invaginação da membrana, cicatriz de brotamento, citoplasma, núcleo, mitocôndria, retículo endoplasmático, sistema de Golgi, vesículas secretoras, vacúolo, peroxissomo. As células de *S. cerevisiae* possuem geralmente formato elipsóide, sendo que o maior diâmetro das células varia entre 5 e 10 μm e o menor diâmetro varia entre 1 a 7 μm . A média dos volumes celulares é de 29 ou 55 μm^3 para as células haplóides e diplóides, respectivamente, e o tamanho celular aumenta de acordo com a maturação da célula (ZINSER & DAUM, 1995; FELDMANN, 2005). As colônias de *S. cerevisiae* são de coloração branca ou creme, de aspecto cremoso, possuem reprodução vegetativa por brotamento e podem possuir pseudo-hifas simples; possuem ascos

persistentes contendo quatro ou mais ascósporos, redondos ou ovais (BARNETT *et al*, 2000).

Como já descrito, *S. cerevisiae* tem sido utilizada com sucesso como um modelo para complexos processos fisiológicos em células de metazoários. Resultados fundamentais foram atingidos nos campos de envelhecimento, transporte vesicular e divisão celular (MADEO *et al*, 2004; CARMONA-GUTIERREZ *et al*, 2010). Esta levedura também está sendo muito útil para solucionar questões complexas de bioenergética em mitocôndrias saudáveis. Recentemente, resultados fundamentais foram alcançados na pesquisa da função das mitocôndrias durante a execução da apoptose em leveduras (EISENBERG *et al*, 2007).

1.2.2. *Candida albicans*

A *C. albicans* é uma espécie de fungo que normalmente reside como comensais no trato gastrointestinal dos seres humanos e outros animais de sangue quente. O seu genoma diplóide estimado é de aproximadamente 30 Mb, contendo 8 pares de cromossomos homólogos (<http://www.broadinstituit.org>). É o fungo patogênico mais comum em humanos e é responsável por causar uma variedade de infecções de pele e tecidos moles em pessoas saudáveis e doenças mais virulentas, invasivas, e disseminadas em pacientes com o sistema imunológico comprometido. O fato de este microorganismo conseguir permanecer em hospedeiros saudáveis e também causar um espectro de doenças em hospedeiros imunocomprometidos são questões de interesse biológico significativo, bem como de importância clínica e econômica (NOBLE & JOHNSON, 2007).

Candida spp são organismos tipicamente comensais, presentes em aproximadamente 50% da população. Entretanto, por volta de 80% das mulheres sofrem de infecções clínicas de *Candida* e em torno de 5% destas infecções podem ser recorrentes, com algumas delas tornado-se resistente à terapia antifúngica (ODDS, *et al*, 2004; RAMSDALE, 2008). Em indivíduos imunocomprometidos, *C. albicans* pode produzir suaves, mas irritantes infecções superficiais da mucosa oral e vaginal. Em pacientes severamente imunocomprometidos, este fungo pode produzir uma infecção sistêmica disseminada que se não tratada efetivamente está associada com uma alta incidência de mortalidade (ODDS, *et al*, 2004).

C. albicans também é a quarta infecção hospitalar mais comum e é alta a incidência da resistência a tradicionais terapias antifúngicas (YANO *et al*, 2011; KRCMERY, 2000; LAMAGNI, *et al*, 2001; RAMSDALE, 2008). Vários agentes antifúngicos estão disponíveis para tratar infecções por este microorganismo; contudo, o sucesso dos tratamentos antifúngicos depende do estado do paciente, do tipo de agente antifúngico utilizado, e também da resposta biológica do patógeno fúngico. Tem sido demonstrado em muitas ocasiões que *C. albicans* pode desenvolver mecanismos específicos de resistência a partir da exposição a drogas antifúngicas (SHAPIRO *et al*, 2011). Um dos mais bem documentados mecanismos de resistência aos agentes antifúngicos azólicos é o aumento da expressão de genes que codificam transportadores que conferem resistências a múltiplas drogas (MDR) com diferentes mecanismos de ação (SANGLARD *et al* 1997; SANGLARD *et al*, 1995; WHITE, 1997; COSTE *et al*, 2004; SHAPIRO *et al*, 2011). O aumento da expressão destes genes leva a um maior efluxo de azóis e, portanto, resulta na diminuição do acúmulo da droga e na redução da inibição do seu alvo, codificado pelo gene *ERG11*. Pelo menos duas famílias de transportadores MDR, a família de transportadores ABC (“ATP-binding cassette”) e a superfamília MFS (“major facilitator superfamily”), estão envolvidas na resistência aos agentes antifúngicos dos azóis. Cdr1p e Cdr2p (“*Candida* drug resistance”), da família de transportadores ABC, e CaMdr1p (“*C. albicans* multidrug resistance 1”), da família de transportadores MFS, são os principais mediadores de resistência a azóis em *C. albicans* (SHAPIRO *et al*, 2011; SANGLARD *et al*, 1995; COSTE *et al*, 2004).

Além disso, a superfície da célula apresenta dois papéis fundamentais na sobrevivência dos microrganismos: proteção e interação. O papel protetor preserva a integridade da célula diante de ataques ambientais. O papel de interação inclui a detecção de sinais externos e também a ligação a outras células ou superfícies externas (WALKER *et al.*, 2008). *C. albicans* é protegido por uma parede celular constituída por polímeros de açúcares como base, cujo constituinte principal é o β -glucano, um polímero de glicose que confere forma. A quitina fornece a rigidez da parede, enquanto que as manoproteínas que revestem a superfície servem para interagir com o ambiente externo. A parede celular está ligada a vários processos biológicos conhecidos em *C. albicans*, incluindo o crescimento, a morfogênese, o “mating”, e patogenicidade (CHAUHAN *et al.*, 2004).

Com relação à morfologia de *C. albicans*, uma característica marcante é a sua capacidade de crescer ou como uma levedura de brotamento unicelular ou nas formas filamentosas de pseudohifas e hifas (SUDBERY, 2011; SUDBERY *et al*, 2004; ODDS, 1998). As pseudohifas são morfologicamente distinguíveis de hifas, uma vez que elas apresentam constrictões nos locais de septação e são mais largas que as hifas. Por outro lado, as hifas formam longos filamentos semelhantes a tubos, com lados completamente paralelos e sem constrictões no local da septação. As hifas e as pseudohifas também possuem diferenças fundamentais na organização do ciclo celular e nos mecanismos de crescimento polarizado (SUDBERY, 2011). Em *C. albicans*, a orientação do tubo germinativo nos estados leveduriforme e pseudohifa é determinada pelo polarissoma (complexo de proteínas envolvidas na orientação do crescimento do tubo germinativo). Na formação do tubo germinativo da hifa verdadeira, a orientação é determinada tanto pelo polarissoma como pelo *Spitzenkorper* (estrutura responsável pelo crescimento e desenvolvimento polarizado, encontrada apenas em hifas verdadeiras) (WHITEWAY & BACHEWICH, 2007; CRAMPIN *et al*, 2005). Os clamidoconídios (clamidosporos) são estruturas de resistência encontradas em *C. albicans*, formadas quando o fungo se encontra em um local onde não existem todos os nutrientes necessários para o seu desenvolvimento, como por exemplo, meios de cultura feitos à base de arroz e milho, como o industrializado corn meal (WHITEWAY & BACHEWICH, 2007). A plasticidade morfológica de *C. albicans* é um determinante da virulência, visto que a forma de hifas, por exemplo, tem papel fundamental no processo de infecção. Durante as infecções da mucosa, as hifas invadem as células epiteliais e endoteliais e causam danos, provavelmente através da liberação de enzimas hidrolíticas (ZHU & FILLER, 2010). A *C. albicans* é perfeitamente adaptada ao crescimento em seu hospedeiro humano e forma hifas sob uma variedade de condições ambientais que refletem a diversidade de microambientes que ela encontra no hospedeiro (SUDBERY, 2011). Assim, as hifas se formam em resposta à presença de soro, pH neutro, 5% de CO₂ (a pressão parcial de CO₂ no sangue) e N-acetil-D-glucosamina (GlcNAc) (MARDON *et al*, 1969; SUDBERY, 2011).

As vias que operam a transdução de sinais indutores de hifa, assim como os fatores de transcrição necessários, foram revisados recentemente em detalhes por Shapiro *et al* (2011). A expressão de genes hifa-específicos é regulada negativamente por um complexo constituído pelo co-repressor transcrricional geral

Tup1 em associação com o Nrg1 ou o regulador de crescimento filamentosos tipo Rox1p (Rfg1). As células que não possuem qualquer um desses repressores crescem constitutivamente como pseudohifas longas, e a expressão de genes hifa-específicos não é inibida (KADOSH & JOHNSON, 2005; MURAD *et al*, 2001; SUDBERY, 2011). A regulação positiva da expressão de genes hifa-específicos é realizada por vários fatores de transcrição, incluindo Efg1, CHP1, Cph2, Tec1, Flo8, Czf1, Rim101 e Ndt80 (SHAPIRO *et al*, 2011; SUDBERY, 2011; STOLDT *et al*, 1997; LANE *et al*, 2001; CAO *et al*, 2006; DAVIS *et al*, 2000; SELLAM *et al*, 2010). O Efg1 é necessário para a formação de hifas em resposta a soro, pH neutro, CO₂ e GlcNAc em meios líquidos, e em meios sólidos como o meio “Spider”. O Cph1 e a sua via de ativação “upstream” são necessários apenas para a formação de hifas em meio sólido “Spider”, mas não em meios líquidos. Consequentemente, sugere-se que Efg1 seja o principal regulador da formação de hifas na maioria das condições. O Efg1 e o Cph1 são ativados por distintas vias de sinalização “upstream”. No caso de Efg1, a via metabólica é baseada no AMP cíclico, enquanto que no caso de Cph1 ela depende da via metabólica de sinalização de uma proteína quinase ativada por mitógeno (MAPK). O Ras1 (GTPase) regula as vias metabólicas de cAMP e MAPK (STOLDT *et al*, 1997; SUDBERY, 2011; LEBERER *et al*, 2001).

A transição de levedura para hifa de *C. albicans* está ligada a uma série de propriedades: adesão a células epiteliais e endoteliais; invasão primária e intercelular via endocitose induzida e penetração ativa; aquisição específica de ferro a partir de fontes intracelulares do hospedeiro; escape de fagócitos e evasão imune. Assim, o dimorfismo é essencial para a patogenicidade, tanto a nível superficial quanto sistêmico. O potencial patogênico de formação de hifa é devido à morfologia do fungo e também à expressão de um programa transcricional hifa-específico (JACOBSEN *et al*, 2012).

Além disso, o sistema imunológico é capaz de discriminar entre levedura e crescimento na forma de hifas. Este reconhecimento específico de morfologia de *C. albicans*, juntamente com a detecção da carga fúngica e dos danos causados pela invasão das hifas pode ser a chave do hospedeiro para a discriminação entre colonização e infecção. Logo, a inibição da transição de levedura para hifa ou modulação da resposta imune associada com o dimorfismo pode ser uma opção muito atraente para o controle das infecções de *C. albicans* no futuro (JACOBSEN *et al*, 2012; WHITEWAY & BACHEWICH, 2007).

1.3. A Morte Celular

A morte celular pode ser definida como sendo o estado em que as células se encontram totalmente incapazes de realizar qualquer função, seja ela de controle homeostático, mobilidade, síntese, excreção, reprodução ou outro (ANDERSON *et al.*, 1982). É um processo em que uma célula perde a capacidade de manter a homeostasia e com isso, ocorrem alterações na integridade da membrana plasmática, aumento do volume celular e perda irreversível de suas funções. Esse processo ocorre a partir de uma rede de mecanismos elementares que agem em série ou em paralelo (GOLSTEIN e KROEMER, 2005; KUMAR *et al.*, 2010). A morte celular foi, por muito tempo, definida um processo de caráter puramente degenerativo, ocorrendo frente a situações de infecção, lesão celular e ausência de substratos e fatores de crescimento. Entretanto, nem todos os eventos de morte celular apresentam tais finalidades e motivações. Existem diferentes tipos de morte celular que podem ser classificados levando-se em consideração os aspectos funcionais (morte acidental ou programada, fisiológica ou patológica), a morfologia celular (morte apoptótica, necrótica ou autofágica) e a ativação de enzimas específicas (morte com ou sem envolvimento de caspases, nucleases, dentre outras) (GALLUZZI *et al.*, 2007a).

De acordo com a Comissão de Nomenclatura de Morte Celular (CNMC), a classificação oficial das modalidades de morte é definida em termos puramente morfológicos e com poucas referências a alterações bioquímicas. Dessa forma, os tipos de morte celular se resumem a apoptose, necrose e autofagia, sendo que cada um pode ocorrer a partir da ativação de uma série de mecanismos diferentes (GOLSTEIN e KROEMER, 2005; KROEMER *et al.*, 2005). Em algumas situações, a exposição da célula a condições extremas pode provocar lesões reversíveis ou fatais. As lesões fatais provocam morte celular tipo necrose (KROEMER *et al.*, 2005, KUMAR *et al.*, 2010). **A necrose** foi primeiramente descrita por Walker e colaboradores em 1988 e foi considerada por muito tempo como uma forma de morte caracterizada pelo descontrole celular (GALLUZZI *et al.*, 2007). Atualmente, a necrose é considerada como uma morte celular acidental, não descontrolada, uma vez que estudos recentes demonstram a existência de vias de transdução de sinais envolvidas no controle desse processo apesar da ausência de marcadores bioquímicos específicos (GALLUZZI *et al.*, 2007a; EISENBERG *et al.*, 2010).

A necrose compreende um conjunto de alterações morfológicas devido à ação progressiva de enzimas na célula, geralmente causada por uma lesão exógena irreversível ou um estímulo patológico. As células necróticas apresentam aumento no volume celular, intumescimento das organelas, culminando na ruptura da membrana plasmática, com extravasamento do conteúdo celular e, também, algumas características da apoptose, como condensação da cromatina (picnose) e a fragmentação da membrana nuclear (cariorrexe). Muitas vezes, a necrose está associada a diversas patologias e processos inflamatórios locais (GALLUZZI et al., 2007; EISENBERG et al., 2010; KUMAR et al., 2010).

Embora a apoptose seja reconhecida principalmente por seus papéis no desenvolvimento de eucariotos superiores, ela não é restrita somente a metazoários, podendo ocorrer na maioria dos sistemas vivos, incluindo plantas, fungos e bactérias (CAO, *et al*, 2009; RAMSDALE, 2008; SHARON, *et al*, 2009). A apoptose já foi descrita na maioria das linhagens filogenéticas, incluindo eubactérias, protistas, fungos, plantas e animais. A ampla ocorrência da morte celular programada entre os seres vivos aponta uma origem remota desse processo, indicando que o controle da morte celular pode implicar em vantagem seletiva na evolução de um organismo ou uma população (RAMSDALE, 2008; SEVERIN et al., 2008). Ao contrário da necrose, **a apoptose**, também conhecida por morte celular programada, é um processo no qual a execução celular ocorre sem a ruptura da membrana plasmática, envolvendo a expressão e a repressão de genes específicos, sob o preciso controle molecular, com a ativação de um programa de morte intracelular levando a célula a um “suicídio” de uma maneira controlada (ALBERTS et al., 2007; EISENBERG et al., 2010). A apoptose é caracterizada por mudanças morfológicas e processos bioquímicos que são, em grande parte, conservados através da evolução (SHIRTLIFF, *et al*, 2009; GALLUZZI, *et al*, 2008). A morte celular apoptótica é um processo vital em organismos multicelulares, que normalmente ocorre durante o desenvolvimento e está associada com a manutenção da homeostasia celular, eliminação de danos das células, resposta a agentes infecciosos, envelhecimento e diferenciação, assim como respostas adaptativas das células a estresses bióticos e abióticos (DANIAL; KORSMEYER, 2004; GREEN, 2005; SHARON, *et al*, 2009).

As redes apoptóticas são compostas de centenas de proteínas que são estreitamente reguladas por complexas cascatas de sinalização em múltiplos pontos. As apoptoses inapropriadas devido a interferências nestas redes

regulatórias podem ter múltiplos efeitos e estão associadas a muitas malignidades humanas, como doenças neurodegenerativas, autoimunes e câncer (ELMORE, 2007). Muitas das principais proteínas apoptóticas têm sido identificadas; entretanto, em muitos casos, os mecanismos moleculares destas proteínas são parcialmente compreendidos (SHARON, *et al*, 2009). Seguindo a exposição a uma variedade de estímulos, as células iniciam o processo de apoptose de uma maneira estritamente regulada (GALLUZZI *et al.*, 2008; MOVASSAGH, *et al.*, 2008). Dentre essas mudanças, enquadram-se: a exposição da fosfatidilserina na superfície celular, cariorrexe, a quebra do DNA e a picnose, com a formação de um anel típico no interior do envelope nuclear. As células formam corpos apoptóticos envolvidos por membrana plasmática e com as organelas citoplasmáticas. O brotamento de corpos apoptóticos resulta em uma progressiva condensação celular que são endocitadas pelas células fagocíticas (MADEO *et al.*, 1997; GALLUZZI *et al.*, 2007, 2008). As características centrais das respostas da morte celular programada em mamíferos são definidas por mudanças morfológicas e bioquímicas, as quais são mediadas por vias metabólicas de morte extrínseca ou intrínseca. Na via intrínseca, os sinais de morte induzem a liberação de proteínas mitocondriais, resultando na amplificação de uma cascata de caspases (LIU, *et al*, 1996). As caspases são proteases que atuam em diferentes substratos no citoplasma e acarretam várias modificações morfológicas importantes na célula apoptótica (DEGTEREV, *et al*, 2003). Na via metabólica extrínseca, sinais mediados por receptores de morte da superfamília do receptor TNF ativam a cascata de caspases diretamente. As vias de morte independente de caspases, também podem ser iniciadas em resposta a estresses, como por exemplo, após a exposição de células a ROS, que envolvem a translocação do fator de indução de apoptose (AIF) da mitocôndria para o núcleo. Além disso, a apoptose está normalmente associada com a fragmentação do núcleo e à degradação de DNA, que podem estar relacionadas a atividade de diferentes nucleases (ENARI, *et al*, 1998; RAMSDALE, 2008), seguida pela formação de um anel atípico no interior do envelope nuclear (MADEO, *et al*, 1997; GALLUZZI, *et al*, 2008). A mitocôndria exerce um papel crucial na execução da morte celular apoptótica, sendo esta organela alvo de alterações morfológica (fragmentação da rede tubular) e bioquímica (geração de espécies de radicais de oxigênio, EROs). Além disso, várias proteínas reguladoras e executoras da morte celular são liberadas dessa organela (FRÖLICH *et al.*, 2007).

A autofagia é considerada uma via alternativa de morte celular programada na qual ocorre o processo de autodigestão celular (OKADA e MAK, 2004). A autofagia pode constituir uma defesa celular contra o estresse agudo, em particular a privação de nutrientes ou fatores de crescimento, sendo um processo conservado em diversas espécies. Através da autodigestão de algumas organelas e proteínas, a célula consegue adaptar seu metabolismo a falta de nutrientes (MAIURI et al., 2007; LEVINE e KROEMER, 2009). Em todas as células encontra-se um nível basal do processo autofágico, que é necessário e importante para a manutenção da homeostase celular (MIZUSHIMA et al., 2008). Desta forma, a autofagia é uma forma de reciclagem celular que também promove a geração de ATP e controla o dano celular removendo proteínas e organelas não-funcionais (GLICK et al., 2010). Entretanto, células que sofrem autofagia excessiva morrem por uma via alternativa que leva a uma morfologia celular diferente tanto da morte por apoptose quanto por necrose. Essas células destroem grande parte do citosol e das suas organelas causando atrofia celular com colapso das funções vitais, levando a morte (MAIURI et al., 2007). O marco na morfologia da autofagia é a vacuolização do citoplasma, sendo que a maior parte das proteínas ativadas durante a autofagia está envolvida na formação das vesículas. No início desse processo de morte celular, uma dupla membrana de origem desconhecida - provavelmente derivada da bicamada lipídica do retículo endoplasmático, do complexo de Golgi e/ou dos endossomos - forma uma estrutura chamada fagóforo. Essa membrana se expande, capturando porções do citoplasma (organelas e proteínas) para a formação de vesículas, chamadas autofagossomos (GLICK et al., 2010; RUBINSZTEIN et al., 2007). Após a fusão dessas vesículas com lisossomos, forma-se uma vesícula ainda maior, o autofagolisossomo, onde se dá a degradação do conteúdo citoplasmático capturado pelas proteases ácidas (MAIURI et al., 2007). Em *S. cerevisiae*, mutantes defeituosos para autofagia (ATG, “autophagy-related genes”) apresentam viabilidade reduzida durante a privação de nitrogênio, uma vez que este processo tem sido implicado em uma variedade de funções celulares essenciais e possui um papel importante na manutenção da função mitocondrial devido a participação na síntese de proteínas essenciais. A disfunção mitocondrial é a maior causa da morte celular induzida por privação de nutrientes em mutantes defeituosos para autofagia (SUZUKI, et al, 2011; YORIMITSU & KLIONSKY, 2005)

Apesar de a CNMC definir três tipos de morte celular bem distintos morfológicamente, acredita-se que durante a evolução, um mecanismo ancestral de morte celular, provavelmente a necrose, pode ter sido complementada com elementos de outros tipos de morte celular programada como a apoptose e/ou a autofagia, sendo que ambas apresentam vantagem seletiva contribuindo para a eliminação de uma célula sem acarretar prejuízos para as células vizinhas. Esse acúmulo de mecanismos de morte celular resulta na possibilidade de ativação de diferentes vias efetoras frente a diferentes insultos celulares (GOLDSTEIN e KROEMER, 2005). Assim, observa-se que células de um mesmo tecido ou uma mesma população podem apresentar-se em estágios distintos de morte celular, sendo que esse processo não ocorre de forma sincronizada em todas as células.

1.3.1 A morte celular e a mitocôndria

Na ativação da morte celular programada tipo apoptose pela via intrínseca, uma das principais contribuições da mitocôndria é a liberação para o citosol de proteínas normalmente confinadas no espaço intermembranas. A partir da permeabilização da membrana externa, essas proteínas deixam a mitocôndria e exercem sua função pró-apoptótica no citosol ou no núcleo da célula (ROUCOU et al., 2000; KROEMER et al., 2007; PEREIRA et al., 2008). Simultaneamente à permeabilização da membrana externa, observa-se a fissão da mitocôndria que deixa a forma tubular, sofrendo uma notável fragmentação. Essa alteração na morfologia mitocondrial facilita a liberação dos fatores apoptogênicos (BRÖKER et al., 2005; CHENG et al., 2008).

A citocromo c foi a primeira proteína mitocondrial apoptogênica identificada e a sua translocação do espaço intermembranas para o citoplasma foi observada em vários modelos de estudo da apoptose. Após a ativação da apoptose pela via intrínseca, essa enzima é liberada para o citosol onde interage com o fator ativador de proteases pró-apoptóticas (Apaf-1) e forma o apoptossomo, um grande complexo heptamérico que recruta alostericamente a caspase 9 e, conseqüentemente, inicia a cascata de caspases (GALLUZZI et al., 2008; ALBERTS et al., 2007). Os citocromos são proteínas com característica absorção da luz visível. A citocromo c é uma pequena proteína mitocondrial, codificada pelo DNA nuclear, que possui entre 12-14 kDa, e apresenta o grupo prostético heme c e o íon ferro ligado covalentemente a

um resíduo de cisteína (NELSON e COX, 2006). Durante a respiração oxidativa convencional, o citocromo *c* atua como carreador dos elétrons. Depois de receber um elétron do complexo III, o íon ferro dessa proteína sofre redução e se move até o complexo IV, onde doa esse elétron para o centro de cobre do complexo IV. A citocromo *c* está envolvida em processos de redução/oxidação celular e pode existir na sua forma oxidada ou reduzida, dependendo do ambiente oxidativo da mitocôndria, sendo reduzida pelo complexo III e rapidamente oxidada pela citocromo *c* oxidase e, assim, quando a cadeia respiratória está ativa, não há acúmulo de nenhuma das duas formas dessa proteína (NELSON e COX, 2006; BORUTAITE, 2010).

Sobre o papel do citocromo *c* na morte celular em fungos, Ludovico e colaboradores (2006) demonstraram que a citocromo *c* é liberada das células de *S. cerevisiae* na apoptose causada por ácido acético. Em mutantes inativos de Cyt1p, esse estímulo apoptótico não causou a morte celular, demonstrando que essa proteína é essencial para o processo. Como não há relatos de proteína similares a Apaf-1 em leveduras, acredita-se que a liberação da citocromo *c* em leveduras ative a apoptose por vias diferentes das descritas para eucariotos superiores, como por exemplo, pela ativação da pró-caspase Yca1p e a formação de uma estrutura tipo-apoptossomo (GIANNATTASIO et al., 2008; BORUTAITE et al., 2010).

Outra proteína mitocondrial que parece atuar potencialmente no processo de morte celular é a endonuclease G (EndoG). Trata-se de uma nuclease mitocondrial que é normalmente confinada ao espaço mitocondrial intermembranas (LI et al., 2001; PARRISH et al., 2001; BUTTNER et al., 2007). A EndoG pertence a um grupo conservado de DNA-RNA-nucleases não específicas com domínio $\beta\beta\alpha$ -metal-ion-finger (SCHAFER et al., 2004). Essa proteína está localizada na mitocôndria em condições fisiológicas e sua função primária está relacionada com a manutenção da sobrevivência celular e consiste na quebra de fitas duplas de DNA para o início da recombinação e reparo, em especial do DNA mitocondrial (mtDNA). Huang e colaboradores (2006) propuseram a participação de EndoG na replicação mitocondrial, através da geração de oligonucleotídeos de RNA iniciadores da síntese de DNA na matriz da mitocôndria. Em mamíferos, a depleção dessa proteína reduz a proliferação celular e favorece a permanência de células na fase G2 do ciclo celular. A deleção de EndoG já foi associada com tumores em células tetraplóides (ZASSEHAUS e DENNIGER, 1993; HUANG et al., 2006; BÜTTNER et al., 2007a).

Durante o processo de apoptose, a EndoG pode ser liberada e subsequentemente translocada para o núcleo onde é responsável pela quebra do DNA genômico (LI et al., 2001; PARRISH et al. 2001; HUANG et al., 2006; ISHIHARA e SHIMAMOTO, 2006; BÜTTNER et al., 2007; RAMSDALE, 2008, LI et al., 2010). Apesar do conhecimento sobre a localização sub-celular e a função endonucleásica de EndoG, o mecanismo de translocação mitocôndria-núcleo e a função exata dessa proteína na sobrevivência e na morte celular ainda não foram completamente elucidados. A EndoG participa também da fragmentação do DNA nuclear em processos específicos de resposta ao estresse oxidativo (LI et al., 2010; ISHIHARA e SHIMAMOTO, 2006).

As tentativas de definir a função exata da EndoG na sobrevivência e na morte celular produziram resultados controversos. Existem pontos que permanecem contraditórios sobre a atuação da endonuclease G. Alguns estudos afirmam que essa proteína regula a apoptose independentemente da liberação de outras proteínas mitocondriais, como o AIF, e de caspases, enquanto outros sugerem a necessidade desses fatores na morte celular mediada por EndoG (BURHANS e WEINBERGER, 2007). Estudos abordando a EndoG em fungos se limitam às leveduras *S. cerevisiae* e *S. pombe*. Em *S. cerevisiae* foi identificada uma proteína homóloga a EndoG, a Nuc1p, que de forma análoga, desencadeia a morte celular quando deixa a mitocôndria. A Nuc1p induz a apoptose independentemente de caspases e AIF. O mecanismo de ativação de apoptose pela Nuc1p parece estar relacionado ao poro de transição de permeabilidade, a carioferina (Kap123p) e a histona H2B, sugerindo uma via na qual ocorre a abertura do poro mitocondrial com a liberação dessa proteína, a sua importação nuclear e a associação com a cromatina (BÜTTNER et al., 2007). *S. pombe* tem uma proteína homóloga a EndoG, Pnu1p, que é produzida como um precursor com uma sequência de localização mitocondrial, durante a importação para mitocôndria a sequência é clivada e a proteína se torna funcional. A atividade extramitocondrial dessa proteína está relacionada com a degradação do DNA nuclear na morte celular (ODA et al., 2007).

1.4. A morte celular em fungos

Alguns fatores relacionados com a apoptose já foram descritos em fungos, uma vez que células fúngicas expostas a condições que levam a morte exibem

diversos marcadores característicos de apoptose, incluindo a rápida exposição da fosfatidilserina na membrana celular, a marginalização da cromatina no núcleo, a fragmentação nuclear, a degradação do DNA (LUDOVICO *et al*, 2001) e a liberação do citocromo c da mitocôndria para o citoplasma (MANON, *et al*, 1997). A caracterização de ROS atuando como reguladores centrais da apoptose em leveduras, confirmou uma grande similaridade com os processos dos metazoários (MADEO, *et al*, 2004).

A maioria dos estudos de morte celular em fungos está focada em *S. cerevisiae*, *A. nidulans* e *N. crassa* (FIGUEIREDO *et al*, 2011; MADEO, *et al*, 1997; RAMSDALE, 2008). Cheng e colaboradores (2003) demonstraram que em *A. nidulans*, os esfingolípídeos são essenciais para a progressão do ciclo celular e a manutenção da polaridade das células. Já Leiter e colaboradores (2005) observaram que o tratamento das hifas de *A. nidulans* com a proteína antifúngica PAF induziu a hiperpolarização da membrana plasmática, a exposição de fosfatidilserinas na superfície e fragmentação do DNA, o acúmulo intracelular de ROS e a desintegração da mitocôndria, caracterizando um processo de morte celular tipo-apoptose.

Diversos estudos, em diferentes modelos experimentais, dão apoio experimental ao papel do estresse oxidativo no desencadeamento da morte celular. Em fungos, várias drogas são usadas para provocar esse tipo de estresse, dentre elas, o farnesol (um isoprenóide não-esteróide produzido pela defosforilação do farnesil fosfato, um metabólito da via biossintética do colesterol). Semighini e colaboradores (2006) demonstraram que o farnesol causou diminuição de crescimento e presença de características morfológicas típicas de apoptose, incluindo externalização de fosfatidilserina e fragmentação e condensação nuclear. O mecanismo de indução a apoptose em *A. nidulans* requer a função mitocondrial, a produção de espécies reativas de oxigênio e o complexo heterotrimérico da proteína G, além da função da poli-(ADP-ribose) polimerase (PARP) (SEMIGHINI *et al*, 2006). Em um estudo similar, Savoldi e colaboradores (2008) investigaram as vias metabólicas de *A. nidulans* que são influenciadas em presença de farnesol a partir da análise do perfil de transcrição deste fungo filamentoso. Os resultados demonstraram mudanças na expressão de vários genes codificadores de proteínas mitocondriais, incluindo a oxirredutase mitocondrial tipo-AIF em *A. nidulans*. Os mutantes inativos de *aifA* em *A. nidulans* exibiram uma maior sensibilidade a apoptose induzida por essa droga, que pode ser pelo menos em parte devido ao

estresse oxidativo (SAVOLDI et al., 2008). Além disso, foi verificado por COLABARDINI e colaboradores (2010) que a overexpressão da *pkcA* em *A. nidulans* (que codifica a proteína quinase C) durante a exposição a farnesol causa aumento da morte celular; e também, a overexpressão deste gene aumenta tanto a acumulação de RNAm quanto a atividade de metacaspases (COLABARDINI et al, 2010). Já no trabalho de FIGUEIREDO e colaboradores (2011) foi identificado o gene homólogo para a endonuclease G em *A. nidulans* (*nucA*^{ENDO^G}) e foi sugerido que NucA está relacionada apenas com a manutenção da integridade do mtDNA e não está envolvida na morte celular (FIGUEIREDO et al, 2011).

Em *N. crassa*, o tratamento de células com fitoesfingosina (PHS) induz morte celular programada com o envolvimento de proteínas mitocondriais (VIDEIRA et al, 2009). Além disso, foi verificado por CARNEIRO et al (2012) que o genoma de *N. crassa* possui 3 ORFs (“*open reading frames*”) homólogas aos genes que codificam o AIF humano e as proteínas AMID (“*apoptosis-inducing factor-homologous mitochondrion-associated inducer of death*”), que são flavoproteínas com atividades oxidorreduzases envolvidas na apoptose independente de caspase. Este grupo demonstrou que enquanto o AIF exibe resistência aumentada a PHS, o mutante para AMD é mais sensível (CARNEIRO et al, 2012).

Em leveduras, análises funcionais em genes revelaram algumas similaridades, ao nível molecular, entre a apoptose em fungos e em eucariotos superiores, identificando homólogos de proteases tipo caspases, o AIF e a serino protease Htr2A/Omi (WISSING, et al, 2004; RAMSDALE, 2008). Diversos estudos têm mostrado a conexão existente entre a morte celular tipo-apoptose em fungos e importantes processos biológicos como desenvolvimento, envelhecimento, resposta a estresse e patogênese. O papel emergente da apoptose como um regulador chave para o desenvolvimento fúngico, sugere que pode ser possível desenvolver novos recursos para o controle de infecções fúngicas através da manipulação da apoptose (SHARON, et al, 2009). Contudo, existem poucos estudos investigando a relação entre drogas antifúngicas e morte celular (para uma revisão, ver RAMSDALE, 2008).

S. cerevisiae tem sido utilizada com sucesso como modelo experimental para complexos processos fisiológicos de células de metazoários (MADEO, et al, 2004; CARMONA-GUTIERREZ et al, 2010). Esta levedura foi o primeiro organismo eucarioto que obteve sua seqüência genômica completamente elucidada (GOFFEAU, et al, 1996) e, este fato tem permitido o desenvolvimento de muitas

ferramentas novas para a análise dos componentes moleculares das células e suas interações (SCHERENS, *et al*, 2004). Estas ferramentas incluem três coleções de mutantes que têm contribuído para a identificação dos componentes fisiológicos básicos e elucidação das vias metabólicas de *S. cerevisiae* (WINZELER, *et al*, 1999; Saccharomyces Genome Deletion Project).

A maquinaria do núcleo apoptótico é bastante conservada neste organismo e vários ortólogos de proteínas apoptóticas de mamíferos em levedura foram identificados (CARMONA-GUTIERREZ *et al*, 2010). Além disso, várias vias metabólicas de morte celular têm sido caracterizados e ensaios de apoptose e/ou morte celular por necrose estão sendo usados atualmente (CARMONA-GUTIERREZ *et al*, 2010). Resultados importantes foram atingidos relacionados a envelhecimento, transporte vesicular ou divisão celular (MADEO, *et al*, 2004). Como já mencionado, a maquinaria apoptótica tem sido relatada para *S. cerevisiae*; em particular, vários tratamentos rigorosos induzem a liberação do citocromo *c*, a fragmentação mitocondrial e também do DNA nuclear (SEVERIN, *et al*, 2008). Homólogos de proteínas pró-apoptóticas foram encontradas nesta levedura, incluindo AIF, um homólogo do HtrA e também para o gene da metacaspase *Yca1*, o qual é o único gene para estas proteínas encontrado em *S. cerevisiae* (UREN, *et al*, 2000; MADEO, *et al*, 2002). Segundo VACHOVA, *et al* (2007), células $\Delta yca1$ mostram elevada resistência a peróxido de hidrogênio, envelhecimento, estresse hiperosmótico e outras condições de estresse excessivo.

Várias substâncias têm sido usadas para induzir apoptose em células de levedura do tipo selvagem; algumas das quais atuam na via metabólica mitocondrial (MADEO, *et al*, 2004). Enquanto a maioria dos mecanismos “upstream” permanece ainda desconhecida, a indução de apoptose por ácido acético, amiodarona e fator α (“mating-type pheromone”) tem ligação clara com a hiperpolarização do potencial de membrana mitocondrial, redução da atividade da citocromo *c* oxidase (COX), formação de ROS mitocondrial, liberação do citocromo *c* e, eventualmente, perda do potencial de membrana mitocondrial e morte celular (EISENBERG, *et al*, 2007).

C. albicans, como descrito anteriormente, é um fungo polimórfico, capaz de alterar sua morfologia entre as formas levedura, hifa e pseudohifa e, estas transições são o papel central para sua patogênese e formação de biofilme (SHIRTLIFF, *et al*, 2009). Este fungo tem se tornado um modelo na genética molecular para estudo da patogenicidade, virulência e desenvolvimento fúngico.

Segundo PHILLIPS, *et al*, (2006), a resposta apoptótica em *C. albicans* pode ser desencadeada quando há a exposição a H₂O₂, acidificação intracelular, ou baixas doses do antifúngico anfotericina B. Entretanto, os mecanismos pelos quais *C. albicans* sofre apoptose ainda não são claros. A via metabólica Ras parece estar envolvida na sinalização apoptótica neste fungo. Mutantes que possuem a sinalização Ras-cAMP-PKA bloqueada, suprimem ou atrasam a resposta apoptótica, enquanto mutações que estimulam esta sinalização, apresentam a taxa de entrada em apoptose acelerada.

Além disso, não há relatos, para *C. albicans*, de metacaspases envolvidas na apoptose. Entretanto, tem sido demonstrado que o gene *IPF4847* (orf19.5995), que é um gene de *C. albicans* com homologia estrutural para a metacaspase YCA1 de *S. cerevisiae*, quando deletado, atenua a morte celular induzida por estresse oxidativo (CAO, *et al* 2009). Este gene também é homólogo para a caspase em mamíferos MCA1 e, foi demonstrado que o farnesol, um precursor para a síntese de esteróis na via metabólica isoprenóide/esterol, recentemente descrito como uma molécula “quorum-sensing” secretada por *C. albicans* capaz de impedir a conversão de levedura para hifa (HORNBY, *et al*, 2001), promove a apoptose via indução de caspases, produção de ROS e rompimento da integridade da mitocôndria (SHIRTLIFF, *et al*, 2009).

Evidências para a morte celular programada em fungos patogênicos têm sido discutidas juntamente com uma avaliação da capacidade de existir agentes antifúngicos que promovam a apoptose e outras formas de morte celular, a fim de aumentar a possibilidade de manipulação das via metabólicas destes fungos para se obter terapias antifúngicas mais efetivas (RAMSDALE, 2008; CAO, *et al*, 2009). Ademais, a identidade e o mecanismo de ação dos compostos químicos presentes na própolis não são conhecidos e, identificar os alvos celulares de quaisquer compostos bioativos é uma tarefa muito difícil. Porém, nos últimos anos, todos os recursos do genoma que foram desenvolvidos para *S. cerevisiae* (www.yeastgenome.org) proporcionaram uma excelente plataforma genômica para o desenvolvimento de drogas e identificação de alvos de drogas (BHARUCHA & KUMAR, 2007; HOON *et al*, 2008; MAERE *et al*, 2005; WUSTER & MADAN BABU, 2008). Além disso, todas essas ferramentas genéticas tornaram possíveis estudar e construir redes de interação genética para identificação específica de alvos de drogas (COSTANZO *et al*, 2010; BOONE *et al*, 2007).

2. Objetivos

A própolis tem atraído à atenção devido à variedade de suas propriedades biológicas e terapêuticas. Ela é constituída por uma mistura complexa de substâncias. Muitos constituintes da própolis foram identificados, contudo, a atividade biológica destas substâncias ainda não é bem compreendida e estudos sobre a sua atividade em leveduras/linhagens dermatófitas são raros. Além disso, evidências para a morte celular programada em fungos patogênicos têm sido relatadas, juntamente com uma avaliação da capacidade de agentes antifúngicos de promover apoptose e outras formas de morte celular. Neste contexto, a utilização da levedura *S. cerevisiae* como modelo experimental tem apresentado sucesso. Ela foi o primeiro organismo eucarioto que obteve sua seqüência genômica completamente elucidada e este fato tem permitido o desenvolvimento de muitas ferramentas novas, dentre elas três coleções de mutantes, de grande aplicação na análise dos componentes moleculares das células e suas interações.

Dessa forma, o presente projeto tem como propósito avaliar diferentes efeitos causados pela própolis no organismo modelo *S. cerevisiae* e no organismo comensal eventualmente patogênico humano *C. albicans*, visando ampliar os conhecimentos acerca das vias metabólicas de morte celular em fungos, abrindo assim diversas perspectivas para sua utilização como uma terapia antifúngica mais efetiva. Logo, os principais objetivos deste trabalho são:

- (i) A utilização de *S. cerevisiae* como organismo modelo para estudos de genética, biologia celular e genômica com o objetivo de compreender como a própolis atua sobre fungos ao nível celular:
 - a. a avaliação de como esta substância pode afetar a sobrevivência celular de *S. cerevisiae*, avaliando diferentes determinantes genéticos envolvidos na apoptose e/ou necrose nesse organismo;
 - b. a utilização da biblioteca de deleção de genes não-essenciais de *S. cerevisiae* para investigar possíveis alvos celulares da própolis; e
 - c. a análise do perfil transcricional de *S. cerevisiae* exposta a própolis.
- (ii) a avaliação de como este composto pode afetar a sobrevivência celular de *C. albicans*, avaliando diferentes determinantes genéticos envolvidos na morte celular desse organismo:
 - a. a utilização de bibliotecas de deleção de *C. albicans* para investigar possíveis alvos celulares da própolis; e
 - b. a avaliação do papel da própolis na transição dimórfica de *C. albicans*.

3. Materiais e Métodos

3.1. Linhagens, meios e métodos de cultura

As cepas de *S. cerevisiae* utilizadas foram: BY4742 (*MAT α* ; *his3 Δ 1*; *leu2 Δ 0*; *lys2 Δ 0*; *ura3 Δ 0*) (BRACHMANN *et al.*, 1998), BY4741A e BY4742 *rho*-/*0* (ambos *MAT α* ; *his3 Δ 1*; *leu2 Δ 0*; *lys2 Δ 0*; *ura3 Δ 0*; *rho*-/*0*) (neste estudo), YJL208C/*NUC1* (*MAT α* ; *his3 Δ 1*; *leu2 Δ 0*; *met15 Δ 0*; *ura3 Δ 0*; *GFP::nuc1*) (Invitrogen, USA) YJR048W/*CYC1* (*MAT α* ; *his3 Δ 1*; *leu2 Δ 0*; *met15 Δ 0*; *ura3 Δ 0*; *GFP::cyc1*) (Invitrogen, USA), e Δ YCA1 (*Mat α* ; *his3 Δ 1*; *leu Δ 0*; *met15 Δ 0*; *ura3 Δ 0*; *yca1::kanMX4*) (MADEO *et al.*, 2002). As cepas de *C. albicans* foram: SC5314 (*wild type*) (GILLUM *et al.*, 1984), CAI4 (*ura3::imm434/ura3::imm434 iro1/iro1::imm434*) (FONZI & IRWIN, 1993), BWP17(*ura3::imm434/ura3::imm434 iro1/iro1::imm434 his1::hisG/his1::hisG arg4/arg4*) (WILSON *et al.*, 1999), DAY286 (*ura3:: λ imm434 his1::hisG pARG4::URA3::arg4::hisG, ura3:: λ imm434 his1::hisG arg4::hisG*) (DAVIS *et al.*, 2002).

Os meios usados foram: meio completo YPD ágar (2% p/v glicose, 1% p/v extrato de levedura, 2% p/v peptona e 2% p/v ágar), YPGal ágar (2% p/v galactose, 1% p/v extrato de levedura, 2% p/v peptona e 2% p/v ágar), meio mínimo SC ágar (0,7% p/v “Yeast Nitrogen Base” (DIFCO); 2% p/v glicose; 0,1g/l L-leucina; 0,1g/l L-lisina; 0,1g/l L-triptofano; 0,05g/l L-histidina; 2% p/v ágar) e meios YPD, YPGal, SC líquidos com a mesma composição (mas sem ágar). A linhagem de Rosella foi construída pela transformação do plasmídeo c-Rosella (ROSADO *et al.*, 2008) na cepa BY4742. Os transformantes foram selecionados em meio mínimo SC ágar.

3.2. Preparação do extrato padronizado de própolis

O Extrato Padronizado de Própolis (EPP) - (EPP-AF ®), e os três lotes (L. 14004/10, 14401/10 e 010/08), foram produzidos pela Empresa Apis Flora (Ribeirão Preto / SP - Brasil). A padronização do produto é feita para garantir qualitativa e quantitativamente sua composição química em todos os lotes, uma vez que as concentrações das substâncias presentes na própolis variam de região para região. Para isto, os extratos foram padronizados utilizando uma mistura de própolis composta por matérias primas provenientes de diversos locais do Brasil (número de patente PI 0405483-0, publicado na Revista de Propriedade Industrial n ° 1.778 de

01 de fevereiro de 2005), como nos Estados de Minas Gerais, São Paulo, Paraná, Santa Catarina e Rio Grande do Sul . (BERRETTA *et al.* 2005).

A própolis (mistura de matérias-primas) foi mantida em um freezer por 12 horas, reduzida a um pó fino em um triturador, e tamanho de suas partículas foi normalizado usando uma peneira de malha 42. Foi então extraído utilizando solução hidroalcoólica (7:3), com maceração dinâmica seguida pelo processo de percolação e, finalmente por uma filtração. O EPP obtido apresenta 11% p/v de resíduo seco e composição química padronizada qualitativa e quantitativamente por RP-HPLC.

Os extratos de própolis EPP-AF[®] foram caracterizados através da análise dos padrões: Ácido caféico (Fluka, L. 43706045), Ácido *p*-cumárico (Fluka, L.3250759), Ácido cinâmico (Fluka, L.21907066), Isosakuranetina (ChromaDex),Artepin C (Wako, L. 016.19131), além dos reagentes grau HPLC: Água purificada (Milli-Q), Metanol grau HPLC (J. T. Baker, L.9093-68), Ácido fórmico PA (Vetec, L.0804789) e Metanol PA (Cinética, L.16245).

A análise quantitativa do extrato de própolis foi realizada em cromatografia líquida de alta eficiência (CLAE), utilizando um aparelho Shimadzu equipado com um controlador CBM-20A, uma bomba LC-20AT quaternária, um detector de arranjo de diodos SPD-M 20A, e um software Shimadzu LC, versão 1.21 SP1. Utilizou-se uma coluna ShimadzuShim-Pack CLC-ODS (M) (4,6 mm x 250 mm, diâmetro de partícula de 5mM, diâmetro dos poros de 100 Å). A fase móvel consistiu de um gradiente de metanol e água acidificada com ácido fórmico (0,1% v/ v), variando de 20% a 95%, com uma corrida de 77 minutos, a uma vazão de 0,8 mL / min. A detecção foi fixada em 275 nm, conforme Berretta *et al.* (2012).

Preparo da amostra: Para um balão volumétrico de 10 mL, adicionou-se, anotando a massa, uma quantidade de extrato padronizado de própolis referente ao teor de flavonóides de cada amostra, a fim de que o resultado ficasse dentro da curva analítica. Adicionou-se 5 mL de metanol (grau HPLC) e 100 µL de padrão interno (solução a 2000 µg/mL). Colocou-se em banho de ultrassom por 10 minutos. Completou-se o volume com água acidificada com ácido fórmico pH = 2,70, homogeneizando-se cuidadosamente. Filtrou-se em filtro de 0,45 µm, diretamente para um vial de 1,5 mL. Cada lote foi analisado em triplicata (Tabela 1).

3.3. Screening das coleções de *S. cerevisiae* e de *C. albicans*

3.3.1. Screening da coleção de *S. cerevisiae* das linhagens haplóides de genes não essenciais deletados para a sensibilidade a própolis

Para o screening da biblioteca de deleção haplóide, criada no Projeto de Deleção do Genoma de *Saccharomyces* (WINZELER, *et al*, 1999; *Saccharomyces* Genome Deletion Project; http://www-sequence.stanford.edu/group/yeast_deletion_project/), um conjunto de aproximadamente 4.800 mutantes deletados foram inoculados a partir da cultura - estoque, a qual estava em placas de 96 poços a -80°C, e crescidos a 30°C em meio YPD contendo 200 µg/ml of Geneticin (G418; Sigma, St Louis, MO) e então estocados a 4 °C. Para determinar quais mutantes eram sensíveis a própolis, as células foram replicadas em placas de Petri contendo YPD e 0,68% de etanol ou extrato alcoólico de própolis (*Apis Flora*®, Ribeirão Preto, SP) a at 0,25 % and 0,50 % (v/v). As placas foram incubadas a 30°C por 5 dias.

3.3.2. Screening com alta eficiência de bibliotecas de *C. albicans* para fatores de transcrição e quinases para sensibilidade à própolis

Neste ensaio, foram utilizadas 15 bibliotecas de deleção fornecidas, em placas de 96 poços, pelo Dr. Kevin Mc Cluskey do Fungal Genetics Stock Center (www.fgsc.net; Mc CLUSKEY & PLAMANN, 2010) [(CJN TF Homozygous mutants; 2nd set TF Homozygous mutants 1; 2nd set TF Homozygous mutants 2; VB GKO Plate 1; VB GKO Plate 2; VB GKO Plate 3; VB GKO Plate 4; VB GKO Plate 5; Dominique Sanglard TF mutants; SFY Homozygous mutants plate 1; CWP Homozygous mutants plate 1; CWP Homozygous mutants plate 2; Misc Homozygous mutants plate 1; Kinase from A. Mitchell plate K1 e plate K2) (<http://www.fgsc.net/candida/FGSCcandidaresources.htm>)]. Estas placas foram replicadas em novas placas de 96 poços, e crescidas a 30°C em meio YPD líquido. Após isso, para verificar a sensibilidade ao extrato de própolis, foram escaneadas em placas YPD sólido contendo com 5,45% etanol (como um controle), que é a concentração de etanol presente em 1% do extrato alcoólico de própolis ou própolis (0,500 e 1%),

durante 5 dias a 30°C. Cerca de 800 cepas diferentes foram envolvidas nesta triagem.

3.4. Extração de RNA e reação de PCR em tempo real para *S. cerevisiae*

Células de BY4742 em fase de crescimento exponencial (9 horas) foram coletadas, e a uma suspensão de 10^7 céls/ml em YPD líquido foi adicionado 0.125% (v/v) de extrato alcoólico de própolis. O tratamento foi realizado por 5 e 10 minutos a 30 ° C, com agitação mecânica (200 rpm). Em seguida, as células de levedura foram lavadas com água destilada e o RNA foi isolado. Para o isolamento do RNA total, as células foram rompidas por vortexação com pérolas de vidro, e o RNA total foi extraído com o reagente Trizol (Invitrogen, EUA). Dez microgramas do RNA de cada tratamento foram então fracionados em 2,2 M de formaldeído, 1,2% p/v em gel de agarose, corados com brometo de etídio, e então visualizados com luz ultravioleta.

A presença das bandas de RNA ribossomais 25S e 18S intactas foi utilizada como um critério para avaliar a integridade do RNA. O tratamento com DNase I livre de RNase para os experimentos de PCR em tempo real foi realizado como descrito por (SEMIGHINI *et al*, 2002). Todas as reações de PCR foram realizadas utilizando o Sistema ABI 7500 Fast Real-Time PCR (Applied Biosystems, EUA) e Taq-Man TM Universal PCR Master Mix Kit (Applied Biosystems, EUA).

As reações e os cálculos foram realizados de acordo com Semighini *et al* (2002). Os primers e sondas fluorescentes Lux TM (Invitrogen, EUA) utilizadas neste trabalho foram: ATG8_263RL (5'-CGGCTATATGGCAGACATCAACG C[FAM]G-3') e ATG8_263RL/235FU (5'-CCCTGAGAAGGCCATCTTCATTT-3') para YBL078C/ATG8; ATG14_892FL (5'-CGGACAACAAGATGAAGTGTAGG TC[FAM]G-3') e ATG14_892FL/910RUa (5'-AACCATCGAGTACAACATCCC AAA-3') para YBR128C/ATG14; ACT1_366FL (CGGCTGGTTCTGGTATGTGT AAAGC[FAM]G) e ACT1_366FL/385RU (ATGGGAAGACAGCACGAGGAG) para YFL039C/ACT1.

3.5. Microscopia óptica e confocal

Após a incubação em condições adequadas para cada experimento, as lâminas foram montadas e visualizadas em um microscópio de fluorescência. Para a microscopia das cepas YJL208C/NUC1, YJR048W/CYC1 e Rosella, as células na

fase de crescimento estacionária (16 horas) foram coletadas e ressuspensas (10^7 céls/ ml) em meio líquido SC para YJL208C e YJR048W e meio líquido YPD para Rosella, contendo 0,125 % (v/v) de extrato alcoólico de própolis; o tratamento foi realizado por 15, 10 e 90 minutos, respectivamente, a 30 ° C, com agitação mecânica (200 rpm). O controle positivo para a Rosella foi feito crescendo-se as células por 16 horas a 30°C em meio YPD e transferindo as para meio com privação de nutrientes [0,17 % (p/v) de extrato de levedura; 0,1% (p/v) de KH_2PO_4 ; 0,01% (p/v) de CaCl_2 ; 0,0005% (p/v) de FeCl_3 ; 0,07% (p/v) de MgCl_2 ; 0,05% (p/v) de NaCl; 2% (p/v) de glicose] para induzir a autofagia. Após 4 horas de incubação a 30°C, adicionou-se o protonóforo “carbonyl cyanide *m*-chlorophenylhydrazone” (CCCP, a 10 μM) no tampão pH 7,5 por 10 minutos (ROSADO *et al*, 2008).

Para o controle positivo da apoptose na microscopia com NUC1, utilizou-se 0,4mM de H_2O_2 por 20 horas (BÜTTNER *et al*, 2007). As células foram lavadas com tampão fosfato salina (PBS) e incubados por 5 minutos em uma solução contendo 100 ng/ml de DAPI (4,6-diamino-2-fenilindol), a fim de marcar os núcleos (Sigma Chemical, St. Louis). Após a incubação com o corante, elas foram lavadas com tampão PBS por 10 minutos a temperatura ambiente e, em seguida, as lâminas foram montadas. As lâminas foram visualizadas com um Carl Zeiss (Jena, Alemanha) utilizando lente objetiva de imersão em óleo com aumento de 100x (CE Plano Neofluar, NA 1.3).

Com o intuito de se verificar o acúmulo de ROS, as células foram coradas com H_2DCFDA (2',7'-dichlorodihydrofluorescein diacetate). As células de levedura (1×10^7 células/ml) foram cultivadas em tubos de 50ml contendo 10 ml de YPD por 9 ou 16 horas a 30°C, e expostas a 0,125% de própolis por 5 ou 10 minutos. As células foram lavadas com meio YPD e incubadas com tampão PBS pré-aquecido contendo 10 mM H_2DCFDA (Invitrogen Molecular Probes) por 30 minutos, no escuro, a 37°C. As lâminas foram montadas e as imagens foram analisadas por microscopia de fluorescência a temperatura ambiente. Utilizou-se microscópio Carl Zeiss Observer. Z 1 (Jena, Alemanha) e lente objetiva de imersão em óleo com aumento de 100x (46 HE YFP, excitação em 500/20nm e emissão em 535/30nm). O DIC (differential interference contrast) para as imagens em campo claro e imagens fluorescentes foram capturadas com uma câmera AxioCam (Carl Zeiss), processadas utilizando o software AxioVision versão 4.8 e salvos como arquivos TIFF. O tratamento posterior foi realizado usando o Adobe Photoshop 7.0 (Adobe

Systems Incorporated, CA). No ensaio do TUNEL (Terminal Deoxynucleotidyl Transferase-mediated dUTP Nick End Labeling) para *S. cerevisiae*, as lâminas foram visualizadas em um microscópio confocal. As imagens foram analisadas utilizando o laser Leica TCS SP5 do microscópio confocal (Leica Microsystems, Heidelberg, Alemanha) (Laboratório de microscopia confocal, FMRP, Universidade de São Paulo, São Paulo, Brasil) e lente objetiva de imersão em água com aumento de 63x. As imagens foram capturadas e processadas usando o software Leica AF LAS (Leica Microsystems) e tratamentos adicionais foram realizados com o Adobe Photoshop 7.0. Para *C. albicans*, utilizou-se o microscópio Carl Zeiss Observer. Z 1 em nosso laboratório.

Para a microscopia utilizando-se Mitotracker® (reagente que cora mitocôndrias), as células foram fixadas com 500 µL de formaldeído (J. T. Baker) diluído para 3.7% em PBS, por 15 minutos a temperatura ambiente. Após a fixação, as células foram lavadas com PBS e coradas com 250 nM do reagente Mitotracker® (Invitrogen) diluído em PBS, por 10 minutos a temperatura ambiente. Após isso, lavou-se com PBS e, para corar o núcleo das células, elas foram incubadas com 250ng de DAPI diluído em 42,5% de glicerol (em PBS), por 10 min a temperatura ambiente. Finalmente, as células coradas foram lavadas com PBS, montadas e visualizadas no microscópio de fluorescência Carl Zeiss Observer. Z 1 (Jena, Germany) usando lente objetiva de imersão em óleo 100x e os filtros; filter set 38 HE, comprimento de onda de excitação, 470/40 nm; comprimento de onda de emissão, 525/50 nm para o GFP (verde), filter set 49 comprimento de onda de excitação, 365 nm; comprimento de onda de emissão, 445/50 nm para o DAPI e filter set 63 HE comprimento de onda de excitação, 572/25 nm; comprimento de onda de emissão, 629/62 nm para o Mitotracker® (vermelho).

3.5.1. A indução de morte celular e determinação da viabilidade

Células de BY4742 ou SC5314 foram incubadas em meio líquido YPD por 16 horas a 30°C. As células cultivadas foram centrifugadas e inoculadas em 50 ml de meio líquido YPD fresco, em frascos de 250 ml, a uma concentração final de 5×10^5 células/ml. As células foram então cultivadas por 9 horas (fase exponencial; 2×10^7 células/ml) e 16 horas (fase estacionária; 3×10^8 células/ml). As células em fases exponencial e estacionária foram coletadas e inoculadas novamente em 50 ml de

meio líquido YPD fresco, em frascos de 250 ml, a uma concentração final de 1×10^7 células/ml, e expostas a 0,68% de etanol (o que corresponde a 60% do etanol presente em 0,125% do extrato alcoólico de própolis) ou 0,125% (v/v) de extrato alcoólico de própolis para *S. cerevisiae* por 5, 10 e 20 minutos; e a 0,68% e 5,45% de etanol ou 0,125% (v/v) de própolis por 1h, e também com 0,5 e 1% por 20 min para *C. albicans*. O tratamento foi realizado a 30°C, com agitação mecânica (200 rpm). Em seguida, as células foram lavadas com tampão PBS para remover a própolis e os procedimentos para a detecção de morte celular foram iniciados. A viabilidade celular foi determinada pelo plaqueamento das células em concentrações apropriadas e contando-se o número de colônias crescidas.

Para o experimento cujo resultado está mostrado na figura 20, em particular, foi feito um pré inóculo de 16 horas (fase estacionária de crescimento) da cepa SC5314. Após este período, 1×10^6 células/mL foram inoculadas em 30 mL de meio YPD líquido em erlenmeyers de 125mL. O tratamento foi feito com 0,125% de extrato alcoólico de própolis a 11% e para o controle negativo, foi adicionado 0,68% de etanol. Os erlenmeyers foram mantidos em agitação de 180 rpm, a 30 °C, por 24, 48 e 72 horas. Após estes períodos de tratamento, o meio de cultura foi removido por centrifugação em falcons de 50mL e ressuspendeu-se o *pellet* de células em 1mL de meio de cultura fresco. Deste volume, foram feitas diluições seriadas de 10 vezes e plaqueou-se em placas de *Petri* contendo meio YPD sólido de forma a conter 100 colônias por placa, a fim de se verificar a viabilidade das células. Incubou-se a 30°C, por 16 horas. Com o volume restante (0,5mL), analisou-se a morfologia das células no microscópio Carl Zeiss Observer. Z 1 (Jena, Alemanha). Além disso, adicionou-se o corante Iodeto de propídio (PI), a fim de se verificar a ocorrência de necrose.

3.5.2. Ensaio do TUNEL (Terminal Deoxynucleotidyl Transferase-mediated dUTP Nick End Labeling)

As quebras no DNA foram demonstradas pelo TUNEL com o kit "In Situ Cell Death Detection Kit, Fluorescein", da Roche. Este procedimento foi realizado segundo MADEO *et al* (1997) e RIBEIRO *et al* (2006), com algumas modificações. As células de levedura foram fixadas em 3,7% (v/v) de formaldeído por 30 minutos a temperatura ambiente, lavadas três vezes com PBS e as paredes celulares foram

digeridas com 15U/ml de Lyticase (*Arthrobacter luteus* 100KU bruto - 362 unidades/mg; Sigma) a 37°C, por 60 minutos. Aplicou-se 10µl da suspensão celular a uma lamínula de microscópio e deixou-se secar durante 30 minutos a 37 °C. As lamínulas foram lavadas com PBS, incubadas em solução de permeabilização [0,1% (v/v) de Triton X-100 e 0,1% (p/v) de citrato de sódio] por 2 minutos no gelo e lavadas novamente com PBS. Para o controle positivo, a endonuclease DNase I (Sigma) foi aplicada às células nas lamínulas de microscopia. As lamínulas foram colocadas em um recipiente umidificado por 1 hora a 37°C; em seguida, lavadas duas vezes com PBS e posteriormente incubadas com 10µL da mistura reacional do TUNEL, durante 60 minutos, a 37 °C. Elas foram lavadas três vezes com PBS e incubadas por 5 minutos em uma solução com 100 ng/ml de DAPI (Sigma Chemical, St. Louis). Após a incubação com o corante, elas foram lavadas com tampão PBS por 10 minutos em temperatura ambiente e, em seguida, montadas e visualizadas no microscópio.

3.5.3. Ensaio de Anexina e Iodeto de Propídio (PI)

A exposição de fosfatidilserina foi detectada pela anexina-V-fluos (Roche, EUA), conforme descrito por MADEO, *et al* (2004), com algumas modificações. As células foram coletadas e lavadas com tampão de sorbitol (1,2M de sorbitol; 0,5mM de MgCl₂, 35mM de K₂HPO₄, pH 6,8). As paredes celulares foram digeridas com 15U de lyticase (Sigma) em tampão sorbitol por cerca de 30 minutos, a 37 ° C. As células foram então lavadas duas vezes com tampão de ligação (10mM de HEPES/NaOH, pH 7,4; 140mM de NaCl, 2,5mM CaCl₂ contendo 1,2M de sorbitol). Para 38 µL da suspensão celular em tampão de ligação/sorbitol, foram adicionados 2 µL de anexina V (Roche, EUA) e 2 µL da solução de trabalho de PI (50µg/mL); incubou-se por 15 minutos a temperatura ambiente.

As células foram então lavadas e ressuspensas em tampão de ligação/sorbitol e, por fim, montadas com a suspensão celular. As configurações do microscópio e aquisições de imagens foram as descritas acima, usando microscopia de epifluorescência.

Para a avaliação quantitativa da coloração anexina V-PI, pelo menos 100 células de leveduras foram contadas por amostra. Para o controle positivo da

apoptose, as células foram tratadas com 80mM de ácido acético, pH 3,0 por 200 minutos (LUDOVICO *et al*, 2001) e para o controlo positivo de necrose, as células foram fixadas com solução fixadora (formaldeído 3,7%; 50mM tampão fosfato de sódio pH 7,0; 0,2% Triton X-100), por 30 minutos, a temperatura ambiente.

3.6. Estudos de fracionamento mitocondrial

Para a extração de proteínas mitocondriais, as mitocôndrias foram isoladas de células de *S. cerevisiae* com base no procedimento descrito por ATTARDI & CHOMYN (1995); GLICK & SCHATZ (1991). Células cultivadas por 16 horas em meio líquido YPGal (a 30°C), expostas ou não a 0,125% de própolis, por 10 min, a 30 °C foram coletadas a 3500 x g, por 5 min e então lavadas uma vez com 1,2M de sorbitol.

Em seguida, as células lavadas foram ressuspensas em tampão de digestão a uma concentração de 10 gramas de células (peso úmido) em 50 ml de tampão (1,2M de sorbitol; 0,06M de fosfato, pH 7,5; 1mM de EDTA; 15mM de β -mercaptoetanol e 50 mg de Zymolyase 20T (MP Biomedicals) e incubadas a 37°C durante 2 horas para serem convertidas em esferoplastos.

Esferoplastos foram lavados com 1,2M de sorbitol e, em seguida, gentilmente homogeneizados em tampão de ressuspensão (0,6M de sorbitol; 20mM de HEPES, pH 7,4 e 0,5mM de PMSF), utilizando um homogeneizador Elvehjem- Potter. O homogeneizado foi centrifugado a 2.500 x g, 5 min a 4°C. O sobrenadante foi novamente centrifugado a 2.500 x g, 5 min a 4°C por mais duas vezes e depois a 12.000 x g por 10 minutos para sedimentar as mitocôndrias.

O sobrenadante (porção citoplasmática) foi coletado e a porção sedimentada ("pellet" mitocondrial) foi ressuspensa gentilmente com o tampão de ressuspensão e lavadas por mais quatro vezes a 2000 x g, 5 min a 4°C. O "pellet" mitocondrial final foi ressuspensa suavemente em 0,6M de sorbitol, 20mM de HEPES. A concentração de proteínas foi determinada pelo ensaio de Biureto (LOWRY *et al*, 1951), utilizando BSA como padrão. As amostras foram então analisadas em gel de eletroforese de poliacrilamida - dodecilsulfato de sódio (SDS-PAGE).

As amostras foram preparadas para SDS-PAGE pela adição de tampão de amostra 1x (62,5mM de Tris-HCl, pH 6,8; 2% de SDS; 10 % de glicerol; 5 % de β -mercaptoetanol e 5 % de azul de bromofenol, e aquecidas a 100°C por 3 minutos.

Para cada amostra, 20µg do total de proteínas foram colocados em cada poço de um gel 10% SDS-PAGE. Após a separação das proteínas, o gel transferido para uma membrana de nitrocelulose (0,2 µm; Bio-Rad) e então bloqueado com 5% leite em pó em tampão TBS/T (10mM de Tris-HCl; 150mM de NaCl, pH8.0 e 0,05% de Tween 20), a membrana foi marcada com o anticorpo de coelho anti-citocromo c (CNAT - contra citocromo c nativo de levedura, Sigma) na diluição de 1:200 em tampão TBS/T por 1 h a temperatura ambiente. Após isso, foi lavada quatro vezes por 5 minutos cada com tampão TBS / T e então incubada com uma diluição de 1:5000 do anticorpo de cabra anti-IgG de coelho, marcado com peroxidase (KPL, EUA) por 1 hora. Depois de ser lavada, as bandas foram reveladas com o uso do sistema de detecção SuperSignal ULTRA chemiluminescence detection system (Pierce) e pelo uso do Hyperfilm ECL (Amersham Biosciences).

3.7. O consumo de oxigênio

O consumo de oxigênio mitocondrial e de esferoplastos foi monitorado por um computador acoplado com um eletrodo do tipo Clark, a 25 ° C com 1mM de NADH como substrato na presença de mitocôndrias na concentração de 100 µg/ml de proteínas. As mitocôndrias foram isoladas a partir de células de *S. cerevisiae* crescidas em YPGal. A própolis foi adicionada a uma concentração final de 0,125%. A fim de bloquear a respiração - citocromo c oxidase, 1mM de KCN foi adicionado no final (BARROS *et al*, 2004).

3.8. Projeto de rede de interação física proteína-proteína (PPPI) e análise topológica

Os dados de sobrevivência obtidos a partir de cepas de *S. cerevisiae* deletadas e submetidas ao tratamento com própolis foram usados para obter informações sobre como os genes deletados e seus produtos interagem no contexto das interações físicas proteína-proteína (redes PPPI). Nesse sentido, o escaneamento e o projeto de rede de interação de redes PPPI foram realizados usando o software Cytoscape, versão 2.6.3 (SHANNON *et al*, 2003). Para isto, foram utilizados os dados PPPI de *S. cerevisiae* disponíveis no banco de dados STRING

8.3 (<http://string.embl.de>) usando os seguintes parâmetros: todos os métodos de previsão ativa habilitados, exceto mineração de texto; mais que 50 interações; score de confiança médio (0.400); e profundidade de rede igual a 1 com a adição de novos nós até à saturação da rede. As redes PPPI obtidas nesta primeira triagem foram então combinadas em uma única rede PPPI, empregando a função de união do “Cytoscape core plugin Merge Networks” (Tabela 2). Esta rede principal PPPI foi então analisada com o software “Molecular Complex Detection” (MCODE) (Bader & Hogue, 2003), disponível em <http://www.cytoscape.org/plugins2.php>, a fim de detectar sub-redes ou um conjunto de proteínas que poderiam representar diferentes processos biológicos. Os parâmetros usados para o MCODE, a fim de gerar os clusters foram: loops incluídos; degree cutoff of 2; exclusão de nós únicos conectados do cluster (haircut option enable); expansão do cluster por um “neighbor shell” permitida (fluff option enable); node density cutoff de 0.1; node score cutoff de 0.2; *k*-core de 2; e profundidade máxima da rede de 100.

3.8.1 Análise *Gene ontology* (GO)

A análise de agrupamento *Gene ontology* (GO) foi realizada utilizando o software *Biological Network Gene Ontology* (BiNGO) (MAERE *et al*, 2005), disponível em http://chianti.ucsd.edu/cyto_web/plugins/index.php. O grau de enriquecimento funcional para um dado grupo e categoria foi avaliado quantitativamente (valor de *p*) através da distribuição hipergeométrica (RIVALS *et al*, 2007), e um teste de correção múltiplo foi aplicado utilizando a taxa de falsa descoberta (FDR) (BENJAMINI & HOCHBERG, 1995), totalmente implementada no software BiNGO. As categorias de processos biológicos super-representados foram gerados após a correção FDR, com um nível de significância de 0,05.

3.9. Microarranjo de DNA (Microarray)

3.9.1. Preparo das amostras

O ensaio foi realizado com cepa BY4742 de *S. cerevisiae*. As células de levedura foram cultivadas por 9 horas (fase de crescimento exponencial), em 50 mL de YPD líquido, a 30 ° C com agitação mecânica (200 rpm). Em seguida, as células

($\sim 2 \times 10^7$ células ml^{-1}) foram expostas a 0,125% de própolis por 5 ou 10 minutos. Para o controle do experimento usando 0,125% de própolis como tratamento, foi utilizado 0,68% de etanol (o que corresponde a 60% de etanol presente no extrato alcoólico de própolis para a concentração de 0,125%). Dois experimentos independentes foram realizados para o array. A viabilidade celular foi determinada pelo plaqueamento das concentrações apropriadas das células e pela contagem do número de colônias em comparação com controles.

3.9.2. Isolamento do RNA e reação de PCR em tempo real

Para o isolamento do RNA total, as células foram rompidas por vortexação com pérolas de vidro, e o RNA total foi extraído com o reagente Trizol (Invitrogen, EUA). Dez microgramas do RNA de cada tratamento foram então fracionados em 2,2 M de formaldeído, 1,2% p/v em gel de agarose, corados com brometo de etídio, e então visualizados com luz ultravioleta. A presença das bandas de RNA ribossomais 25S e 18S intactas foi utilizada como um critério para avaliar a integridade do RNA. Depois disso, o RNA total foi purificado pelo kit RNeasy® Mini Kit (Qiagen) e as amostras purificadas foram quantificadas no NanoDrop® 2000 - Thermo Scientific (Uniscience). Para validação do microarray, foram escolhidos nove genes diferentes a partir da análise do ensaio, cujo mRNA apresentou abundância aumentada ou diminuída quando *S. cerevisiae* é exposta a própolis. Foram desenhados primers e sondas fluorescentes Lux™ (Invitrogen, EUA), e a PCR em tempo real foi utilizada para quantificar a expressão destes genes em uma nova réplica biológica para cada um dos mRNA isolados dos tratamentos por 5 e 10 minutos, com 0,125% de própolis, e comparados com o controle correspondente exposto a 0,68% de etanol. A reação de PCR em tempo real foi realizada como descrito anteriormente no item 3.4, e os primers e sondas fluorescentes utilizadas foram: *PDR15_4412RL* (5'-cggtaGGCATTGGTGGTGGATA CcG-3') e *PDR15_4412RL/4388FU* (5'-GCTACTTGAGTGACCCGTCTGC-3'); *SNQ2_377FL* (5'-cggatTGAGCAAGGCATCCATATCcG-3') e *SNQ2_377FL/418RU* (5'-CTTCTAGGGCACTCGCATCCAC-3'); *VMA21_184RL* (5'-cgcgTTGTGATCTTC AGTATCCTCGcG-3') e *VMA21_184RL/144FU* (5'-TGGAGGTTTAGCTGCT GCAATG-3'); *VMA7_221RL* (5'-cggaaCCACTCTAGCTCTTATGTTTTCCcG-3') e *VMA7_221RL/208FU* (5'-ACGATATTGCCATCCTTCTAATCA-3'); *CKS1_231RL* (5'-

cggcCCAAAGACTGTGTGATGCcG-3') e *CKS1_231RL/213FU* (5'-GGGACCCTGCGTATATTAACAGAA-3'); *SIR1_1690RL* (5'-cggatAGGAACACAAAAGTGCATCcG-3') e *SIR1_1690RL/1643FU* (5'-CCTACGAGGTGCGGACTTCAA-3'); *RLF2_1426FL* (5'-cggcACTTACAACAACTCAAGGCcG-3') e *RLF2_1426FL/1455RU* (5'-TTTGGGTTCTTTGAATGGATCG-3'); *TIM10_228RL* (5'-cggtgTTTTCTGCATGTTTTCACcG-3') e *TIM10_228RL/169FU* (5'-TCCGAGGGTGAGCTGAATAAGA-3'); *MIA40_1036FL* (5'-cgggAGACTGTTTCAGAAAATACCCcG-3') e *MIA40_1036FL/1059RU* (5'-GGTTCCTCATCATCAGATGTTTCC-3').

3.9.3. Hibridação do microarray

3.9.3.1. Marcação e hibridização dos RNAs e geração das imagens de microarranjo

Para análise de expressão gênica foi utilizado o microarray comercialmente disponível da Agilent, o qual contém todo o genoma *S. cerevisiae* [Yeast (V2) Gene Expression Microarray, modelo 8x15K]. Os slides de microarray contêm 15.208 sondas para *S. cerevisiae* (cepa BY4742). As amostras de RNA foram extraídas e purificadas. A marcação do cRNA foi realizada de acordo com o protocolo padrão descrito pela Agilent, usando "Two-Color Microarray-Based Gene Expression Analysis" (Agilent Technologies, USA). Resumidamente, para a síntese e marcação do cRNA, 5 µg do RNA total foi utilizado, o qual foi incubado com os controles internos do kit, na diluição adequada (RNA Spike A ou B). Após isso, foi realizada a síntese de cDNA, incubando-se as amostras na presença do primer T7 promotor. A desnaturação do primer foi realizada a 65°C por 10 minutos. Posteriormente, foi adicionado à reação o cDNA Master Mix (5X First Strand Buffer, DTT 0,1 M, dNTP mix 10 mM, MMLV-RT e RNaseOut). Esta mistura foi incubada a 40 °C por 2 horas para síntese de cDNA, sendo posteriormente colocada a 70 °C por 15 minutos. A seguir, foi realizada a síntese de cRNA, adicionando-se às amostras o mix do fabricante (Agilent™ Transcription Master Mix, contendo 4X Transcription Buffer, DTT 0,1 M, NTP mix, 50% PEG, RNaseOut, pirofosfatase inorgânica e T7 RNA polymerase). À esta mistura foram adicionados os marcadores Cy-3 ou Cy-5 nas amostras. Para a síntese de cRNA marcado, as reações foram incubadas a 40 °C por 2 horas. O cRNA marcado foi finalmente purificado utilizando-se o kit RNeasy®

Mini Kit (QIAGEN) e quantificados através do NanoDrop® 2000 Thermo Scientific (Uniscience). Após a marcação, 300 ng de cRNAs marcados com Cy3 e Cy5 (atividade específica > 8,0 pmol Cy3-Cy5/μg cRNA) foram fragmentados a 60 ° C, por 30 minutos, em um volume de reação de 25 μl contendo tampão de fragmentação 1x (Agilent) e agente de bloqueio 2x (Agilent), seguindo as instruções do fabricante. A fragmentação foi interrompida adicionando-se o tampão de hibridização (2X GE Hybridization Buffer HI-RPM). Finalmente, 50 μL de cada amostra foi colocada nas lâminas contendo os microarranjos de oligonucleotídeos. As lâminas foram então condicionadas em forno de hibridização (Agilent G2545A Hybridization Oven) a 65 °C por 17 horas. Após a hibridização, os microarrays foram lavados sequencialmente: 1 minuto a temperatura ambiente com o GE Wash Buffer 1 (Agilent) e 1 minuto a 37 ° C com o GE Wash Buffer 2 (Agilent); após isso, lavou-se por 10 segundos com Acetonitrila (Agilent), seguido por 30 segundos de lavagem com Stabilization and Drying Solution (Agilent). As lâminas foram imediatamente submetidas à detecção fluorescente no aparelho GenePix 4000B microarray scanner (Molecular Devices, USA) com simultâneo escaneamento dos canais de Cy3 e Cy5 em uma resolução de 5 μm. O laser foi fixado em 100% e o ganho PMT foi ajustado automaticamente para cada slide utilizando o programa GenePix Pro (Molecular Device), de acordo com a intensidade do sinal de cada array, pelo uso do Feature Extraction Software.

3.9.3.2. Análise das imagens e dados obtidos dos microarranjos

As imagens geradas através do escaneamento das lâminas de microarranjo foram analisadas utilizando-se o software Agilent Feature Extraction (FE), versão 9.5.3.1 (Agilent Technologies, USA), que faz uso do algoritmo de normalização por Lowess, obtendo-se a subtração do *background* e a normalização dos valores de intensidade da fluorescência de Cy-3 e Cy-5. Os dados normalizados pelo FE foram convertidos em novo formato de arquivo utilizando-se o software Express Converter (versão 2.1, TM4 platform disponível em <http://www.tm4.org/utilities.html>), que converte os arquivos Agilent em arquivos com extensão .mev (de *multiple experiment viewer*), arquivos esses compatíveis com a plataforma TM4 para análise de expressão gênica. Os novos arquivos, agora com extensão .mev, foram carregados no software MIDAS (TM4 platform), onde foi calculada a média dos

dados obtidos a partir da replicata de genes em cada microarranjo e também das hibridações dye-swap e de duas duplicatas biológicas para cada tratamento, utilizando as ferramentas “flip dye consistency cheking” e “in slides replicates analysis”, implementadas no software MIDAS. Finalmente, os dados completamente normalizados foram analisados através do software TIGR MeV (TM4 platform, Multi Experiment Viewer, disponível em <http://www.tigr.org/software/microarray.shtml>), por meio do qual os genes diferencialmente expressos foram estatisticamente identificados, utilizando-se teste t ($p > 0.01$). Os genes diferencialmente expressos foram considerados significantes quando a razão do $\log_2$ da expressão (Cy3/Cy5) foi diferente de zero, que indica a ausência de modulação da expressão gênica.

3.9.4. Projeto e análise topológica global da rede de interação física proteína-proteína (PPPI)

Os dados do transcriptoma gerados a partir de cepa de levedura BY4742, submetida ao tratamento com própolis, foram utilizados para obter informações sobre como os genes sub ou super-expressos e seus produtos interagem no contexto das interações físicas de proteína - proteína (redes PPPI) em *S. cerevisiae*. Neste sentido, o screening da mineração dos dados e o projeto de redes PPPI para os genes induzidos e reprimidos foram realizados utilizando o software Cytoscape, versão 2.6.3 (SHANNON *et al*, 2003). Para isto, foram utilizados os dados PPPI de *S. cerevisiae* disponíveis no banco de dados *Saccharomyces* Genome Database (<http://www.yeastgenome.org>). As redes PPPI dos genes induzidos e reprimidos, obtidas do primeiro screening, foram combinadas em uma única rede PPPI empregando a função de união do “Cytoscape core plugin Merge Networks”. A rede fundida foi então analisada com o software “Molecular Complex Detection” (MCODE) (Bader & Hogue, 2003), um Cytoscape plug-in disponível em <http://www.cytoscape.org/plugins2.php>, a fim de detectar clusters de proteínas que poderiam representar diferentes processos biológicos. Os parâmetros usados para o MCODE, a fim de gerar os clusters foram: loops incluídos; degree cutoff of 2; exclusão de nós únicos conectados do cluster (haircut option enable); expansão do cluster por um “neighbor shell” permitida (fluff option enable); node density cutoff de 0.1; node score cutoff de 0.2; k -core de 2; e profundidade máxima da rede de 100. O grau de sobreposição dos dados entre as redes PPI dos genes induzidos e reprimidos foi

obtido a partir de uma área proporcional pela análise do diagrama de Venn, disponível em <http://bioinforx.com/free/bxarrays/overlap.php>.

3.9.4.1. Centralidades da rede e análises topológicas locais

As duas principais centralidades de rede (node degree and betweenness) foram computadas a partir da rede fundida e dos clusters utilizando o Cytoscape plugin CentiScaPe 1.0 (Scardoni *et al.*, 2009). As topologias locais da rede, definidas como bottlenecks, foram obtidas a partir do “threshold” gerado por cada centralidade, calculado pelo CentiScape 1.0. Neste sentido, os “bottlenecks” foram definidos como nós com um valor acima do “threshold” calculado para o “node degree” e “betweenness”.

3.9.4.2. Análise Gene ontology (GO)

A análise de agrupamento *Gene ontology* (GO) foi realizada utilizando o software *Biological Network Gene Ontology Ontology* (BiNGO) (MAERE *et al.*, 2005), um Cytoscape plugin, disponível em http://chianti.ucsd.edu/cyto_web/plugins/index.php, da mesma forma como descrito anteriormente no item 3.8.1.

4. Resultados e Discussão

Módulo 4.1: Caracterização molecular da morte celular induzida por própolis em *Saccharomyces cerevisiae*.

Artigo: “Molecular Characterization of Propolis Induced Cell Death in *Saccharomyces cerevisiae*. de Castro, P. A.; Savoldi, M.; Bonatto, D.; Barros, M. H.; Goldman, M. H. S. ; Berretta, A. A. ; Goldman, G. H. *Eukaryotic Cell*, Mar. 2011, p. 398–411”.

4.1. RESULTADOS

4.1.1. A própolis induz morte celular em *S. cerevisiae*

A própolis é um produto complexo derivado de resinas vegetais e saliva de abelha. Existem diversos compostos químicos presentes neste produto natural que poderiam ser responsáveis por suas propriedades antibióticas. Sendo um produto de origem natural, possui inúmeros componentes potencialmente ativos, que podem variar de acordo com a origem botânica, variação climática, estação do ano, entre outros. Uma questão comumente colocada para quem trabalha com esse tipo de produto é a reprodutibilidade lote-a-lote. Assim, para caracterizar quimicamente a própolis empregada no presente estudo, a mesma foi avaliada por HPLC frente aos padrões previamente descritos na literatura (BERRETA, *et al* 2012) para este tipo de própolis, como os ácidos: cafêico, p-cumárico, cinâmico, o flavonóide isosakuranetina e o derivado prenilado e marcador da própolis verde, artepillin C (Figura 1 e Tabela 1).

Além da caracterização realizada, três lotes distintos foram avaliados, visando verificar a reprodutibilidade lote-a-lote do extrato padronizado de própolis EPP AF[®] fornecido pela empresa Apis Flora, fato que foi constatado após a avaliação da correlação linear dos dados obtidos na tabela 1 para cada lote. O tratamento estatístico foi realizado pelo método ANOVA one-way com nível de significância de 0,05.

Figura 1: Perfil cromatográfico de (a) Extrato Padronizado de Própolis (EPP) Lote 1440110; (b) EPP Lote 1400410 e (c) EPP Lote 010/08. O perfil cromatográfico inclui os compostos identificados como: 1. ácido Cafêico (em torno de 15 min); 2. ácido p-cumárico (em torno de 20 min); 3. ácido cinâmico (em torno de 36 min), 4. isosakuretin (em torno de 48 min) e Artepillin C (em torno de 61 min). Os cromatogramas foram plotados a 275 nm, usando RP-HPLC, coluna C18 (Shim-pack, CLC-ODS (M), 25 cm x 4,6) e gradiente de eluição com metanol e água acidificada (ácido fórmico pH=2,7).

Tabela 1: Apresentação quantitativa de compostos do Extrato Padronizado de Própolis obtidos a partir de três diferentes lotes (média $\pm$ DP) (n = 3)

Extrato Padronizado de Própolis (EPP-AF®)			
Compostos (mg/g)	Lote 14401/10	Lote 14004/10	Lote 010/08
Ácido caféico	0,349 $\pm$ 0,002	0,293 $\pm$ 0,003	0,345 $\pm$ 0,011
Ácido p-cumárico	1,470 $\pm$ 0,006	1,511 $\pm$ 0,020	1,712 $\pm$ 0,069
Ácido cinâmico	0,167 $\pm$ 0,001	0,158 $\pm$ 0,003	0,167 $\pm$ 0,006
Isosakuranetina	2,328 $\pm$ 0,009	2,304 $\pm$ 0,031	2,871 $\pm$ 0,103
Artepin C	8,272 $\pm$ 0,013	7,080 $\pm$ 0,135	6,621 $\pm$ 0,306

O trabalho de caracterização dos lotes do extrato de própolis foi realizado nos laboratórios da companhia Apis Flora Indl coml Ltda. Em seguida, em nosso laboratório, foi testado o crescimento de *S. cerevisiae* na presença de quatro destes compostos químicos (ácidos caféico, p-cumárico, cinâmico e isosakuranetin). Para isto, foram aplicados 1, 2, 5, 10, 25, e 100 μ g/ml de cada composto e também de uma mistura contendo 100 μ g/ml de cada um deles nas culturas contendo 10^4 células em meio líquido YPD. Este ensaio foi realizado em placas de microtitulação de 96 poços, atingindo um volume final de 200uL de cultura mais os compostos. Nenhum desses tratamentos foi capaz de inibir o crescimento de *S. cerevisiae* (BY4742), *Candida albicans* (CAI4) ou *C. parapsilosis* (ATCC 2201) (dados não mostrados). Assim, levando em consideração o fato de que os efeitos da morte celular da própolis poderiam ser devido à combinação de compostos químicos, ou seja, um efeito sinérgico dos demais componentes e de diferentes concentrações, decidiu-se investigar esses efeitos realizando os experimentos com extratos alcoólicos de própolis.

Uma primeira série de experimentos foi feita para verificar a concentração de trabalho de própolis adequada, e também como a viabilidade celular era afetada por ela. Uma vez que a própolis é dissolvida em etanol (60% de extrato alcoólico), o tratamento controle foi inoculado em 2,73% de etanol, visto que 0,5% do extrato

alcoólico de própolis apresenta uma concentração final de cerca de 2,73% de etanol (Figura 2A). Diluições seriadas de dez vezes de células de *S. cerevisiae* sugeriram que 0,125% de própolis seria uma escolha adequada como uma concentração sub-inibitória (Figura 2A). Todos os controles nos experimentos posteriores, utilizando 0,125% de própolis como tratamento possuíam como controle 0,68% de etanol (o que corresponde a 60% de etanol presente em 0,125% do extrato alcoólico de própolis). Não foi possível realizar a determinação da concentração inibitória mínima por microdiluições porque a coloração da própolis é escura, o que interfere nas determinações espectrofotométricas (dados não mostrados).

Os efeitos inibitórios da própolis sobre *S. cerevisiae* não são afetados pelo crescimento em baixas concentrações de oxigênio, como a 1% de O₂ (dados não mostrados). Também foi observado que *C. albicans* foi mais suscetível a concentrações de própolis entre 0,50 e 0,75% (Figura 2A). Além disso, as células de levedura nas fases inicial de crescimento (6 horas; $1,2 \times 10^6$ células/ml) e estacionária (16 horas, 3×10^8 células/ml) são mais resistentes a 0,125% de extrato de própolis, enquanto que na fase exponencial (9 horas, 2×10^7 células/ml) são muito mais sensíveis (Figura 2B).

Numa próxima etapa, investigou-se qual o principal mecanismo de morte celular em células nas fases exponencial e estacionária de crescimento, 9 e 16 horas de crescimento, respectivamente, observando-se o número de células que são marcadas por iodeto de propídio (PI) ou Anexina e Tunel (indicações preliminares de necrose ou apoptose, respectivamente). A apoptose primária é caracterizada por um aumento do número de células Anexina V positivas [A (+)] e PI negativas [PI (-)], enquanto nas células em apoptose tardia (levando à necrose secundária) ocorre um aumento do número de células A (+) e PI(+) positivas. A necrose primária é caracterizada por um aumento do número de células Anexina V negativas (-) e células PI(+). Observou-se cerca de 15% das células A + e PI + no controle negativo (0,68% de EtOH), enquanto 100% das células A (+) e PI (+) foram verificadas no controle positivo (ácido acético, pH 3,0, durante 200 minutos). Após 5 minutos de exposição a 0,125% de própolis, as células cultivadas por 9 e 16 horas apresentaram um aumento do número de células A (+) [cerca de 30% A (+), PI (+) e 80% A (+)], sugerindo ocorrência de apoptose primária (Figuras 2C-D). Após 10 minutos de exposição a 0,125% de própolis, as células cultivadas por 9 e 16 horas apresentaram um número maior de células A (+), PI (+) (cerca de 80% para ambas

as classes), sugerindo a apoptose tardia, levando à necrose secundária (Figura 2C-D). No entanto, coloração positiva para o Tunel manteve-se constante tanto para 5 quanto para 10 minutos de exposição a 0,125% de própolis (cerca de 80%; Figura 3).

A liberação nuclear de HMGB1 (proteína de alta mobilidade do grupo box-1 em mamíferos) é uma característica definidora da necrose em células de mamíferos (SCAFFIDI *et al*, 2002). Uma vez que pode ocorrer vazamento em células apoptóticas durante o cultivo prolongado, resultando assim em células PI positivas, determinou-se a translocação núcleo-citosol do Nhp6Ap (homólogo em levedura do HMGB1) em células cultivadas por 16 horas e expostas a 0,125% de própolis por 5 e 10 minutos. A microscopia de fluorescência mostrou a translocação núcleo-citosol do Nhp6Ap::GFP em 20 e 30% das células após a exposição por 5 e 10 minutos a 0,125% de própolis, respectivamente (Figura 2E). Em conjunto, estes resultados indicam que a própolis é capaz de induzir uma resposta de morte celular apoptótica, mas o aumento da exposição a própolis prevê um aumento correspondente na resposta a necrose.

Figura 2: A própolis induz apoptose e necrose tardia em *S. cerevisiae*. (A) Ensaio de diluição para *S. cerevisiae* e *C. albicans* em meio YPD complementado com diferentes concentrações de própolis. As placas foram incubadas a 30°C, por 72 horas. (B) Viabilidade de células de levedura em diferentes fases do ciclo de crescimento e expostas a própolis (0,125% por 5 ou 10 minutos). A viabilidade celular foi determinada pelo plaqueamento de concentrações apropriadas das células e pela contagem do número de colônias. Os números são referentes à média $\pm$ o desvio padrão de três experimentos independentes. (C) Ensaio de Anexina e PI para células de levedura crescidas por 16 horas e expostas a 0,125% de própolis. (D) Número total de células positivas para Anexina e PI para leveduras expostas a 0,125% de própolis. Barras: 5 μ m. Os números são referentes à média $\pm$ o desvio padrão de três experimentos independentes. Em cada experimento, 100 células foram avaliadas para a coloração de Anexina e PI. (E) Translocação Núcleo-citosol de Nhp6Ap::GFP após a exposição ou não das células de *S. cerevisiae* a 0,125% de própolis por 10 minutos. Barras: 5 μ m.

Figura 3: Ensaio do TUNEL (*Terminal Deoxynucleotidyl Transferase-mediated dUTP Nick End Labeling*). As quebras no DNA foram demonstradas pelo TUNEL com o kit “*In Situ Cell Death Detection Kit, Fluorescein*”, da Roche. A coloração positiva para o Tunel permaneceu constante para ambos 5 e 10 minutos de exposição a 0,125% de própolis (em torno de 80%). Barras: 5 μ m.

Há um acúmulo de Espécies Reativas de Oxigênio (ROS) produzidas na mitocôndria durante a apoptose em levedura (EISENBERG *et al*, 2010; OSIEWACZ *et al*, 2006; PERRONE *et al*, 2008). Para investigar a produção de ROS durante a morte celular induzida por própolis em *S. cerevisiae*, foi utilizado 2',7'-dichlorodihydrofluorescein diacetate (H_2DCFDA), um indicador de ROS célula – permeável, o qual não é fluorescente até que os grupos acetato sejam removidos por esterases intracelulares e ocorra a oxidação dentro da célula.

Embora este indicador possa ser oxidado por diferentes ROS, ele é usado principalmente para detectar superóxidos. As células de levedura cultivadas por 9 ou 16 horas, expostas ou não a 0,125% de própolis por 5 e 10 minutos, foram incubadas com H_2DCFDA e examinadas por microscopia de fluorescência (Figura 4). A fluorescência verde dependente de DCFDA foi detectada em cerca de 10% das células com 9 ou 16 horas de crescimento e não expostas à própolis. O tratamento com 0,125% de própolis

das células cultivadas por 9 horas, por 5 e 10 minutos, resultou em 70 a 80% de fluorescência, enquanto o tratamento das células cultivadas por 16 horas, por 5 e 10 minutos, produziram 40 a 60% de fluorescência (Figura 4). Estes resultados indicam que há um grande acúmulo de ROS durante a morte celular induzida pela própolis.

Figura 4: A própolis induz a produção de ROS. Células de leveduras foram crescidas por 9 ou 16 horas a 30°C em YPD, coradas com H₂DCFDA por 30 minutos a 37°C, e posteriormente, analisadas por microscopia de fluorescência. Barras: 5 µm.

As caspases são membros de uma família conhecida como cisteína proteases, que estão ativamente envolvidas na morte celular em eucariotos (para uma revisão, ver LI & YUAN, 2008). Embora *S. cerevisiae* não tenha caspases em seu genoma, uma proteína tipo caspase chamada *YCA1*, que se encaixa na categoria do tipo I das metacaspases, foi identificada e caracterizada (MADEO *et al*, 2002; MAZZONI & FALCONE, 2008; UREN *et al*, 2000). Tem sido demonstrado que *YCA1* está envolvida em vários estímulos de morte celular (para uma revisão, ver MAZZONI & FALCONE, 2008). Quando células de levedura cultivadas por 16 horas foram expostas a 0,125% de própolis por 5, 10 e 20 minutos, houve uma redução na viabilidade celular de 40, 70 e 95% na linhagem selvagem, enquanto que na *YCA1Δ*, esta redução foi cerca de 10, 50 e 80%, respectivamente (Figura 5). Estes resultados indicam que o gene *YCA1* é importante para a morte celular mediada por própolis.

Figura 5: O mutante da metacaspase *YCA1* é mais tolerante a própolis que a linhagem selvagem correspondente. Viabilidade das células de levedura da cepa selvagem e do mutante *YCA1*, crescidas por 16 horas e expostas a 0,125% de própolis ($p < 0.05$), foi determinada pelo plaqueamento de concentrações apropriadas das células e pela contagem do número de colônias. Os resultados são referentes à média $\pm$ o desvio padrão de três experimentos independentes. O tratamento estatístico foi realizado pelo método ANOVA one way, com pós teste Newman Keuls.

4.1.2. O citocromo c é liberado durante a morte celular induzida por própolis

Neste trabalho, também foi investigado se outros componentes da maquinaria de morte celular são ativados durante a morte celular induzida por própolis em *S. cerevisiae*. Concentramos nossa atenção em dois desses determinantes putativos: *CYC1* e *NUC1*. Em células de mamíferos, pelo menos duas grandes vias apoptóticas foram descritas: (i) a via intrínseca que necessita do envolvimento das mitocôndrias, e (ii) a via extrínseca, onde as mitocôndrias são ignoradas e as caspases são ativadas (para uma revisão, ver ELMORE, 2007). Uma das características da via intrínseca é a liberação de fatores apoptogênicos como citocromo c para o citosol e a consequente montagem organizada, por esta proteína, do complexo de alto peso molecular, o apoptossomo mitocondrial, que ativa caspases (ELMORE, 2007). Em células de *S. cerevisiae* sob processo de apoptose induzida por ácido acético, foi observada a translocação de citocromo c para o citosol (LUDOVICO *et al*, 2002). Foram realizados estudos de fracionamento celular com o objetivo de esclarecer se citocromo c foi translocado para o citoplasma durante a exposição de células de levedura, cultivadas

por 16 horas, a 0,125% de própolis por 10 minutos (Figura 6). Em células de levedura não expostas à própolis, todo citocromo *c* permaneceu localizado nos extratos de proteínas mitocondriais, enquanto que, em consequência da adição de própolis, houve uma significativa presença de citocromo *c* nos extratos de proteínas do citoplasma (Figura 6, painel direito). Existem quantidades correspondentes de proteína em cada fração do extrato celular, como mostrado pela coloração de Coomassie (Figura 6, painel esquerdo).

Figura 6: Exposição de células de levedura a própolis induz a liberação do citocromo *c* para o citoplasma. As células de levedura foram crescidas por 16 horas a 30°C e expostas ou não a 0,125% de própolis por 10 min a 30°C. As células foram coletadas, fracionadas (mitocôndrias e citoplasma), conforme descrito em Materiais e Métodos. As proteínas foram separadas em gel de poliacrilamida (painel a esquerda – gel corado com corante Coomassie) e estavam na mesma concentração das proteínas que foram transferidas para a membrana de nitrocelulose (painel a direita). Esta membrana foi marcada com anticorpo de coelho anti-citocromo *c*.

A EndoG tem sido descrita como uma endonuclease mitocondrial que digere o DNA e o RNA (BÜTTNER *et al*, 2007; PARRISH *et al*, 2001). Após a indução da apoptose, a translocação de endoG em mamíferos e o seu homólogo NUC1 em *S. cerevisiae* para o núcleo coincide com a fragmentação do DNA em larga escala (BÜTTNER *et al*, 2007; PARRISH *et al*, 2001; LI & YUAN, 2008). Não foi observado aumento da sensibilidade das células de levedura Δ NUC1 a própolis (dados não mostrados). Numa próxima etapa, examinou-se se Nuc1p seria translocado para o núcleo quando a célula de levedura fosse exposta a própolis (Figura 7). Assim, em células de levedura cultivadas por 16 horas e não expostas a própolis, o

Nuc1p::GFP mostrou uma distribuição mitocondrial difusa com morfologia filamentosa, como confirmado pela coloração com MitoTracker (Figura 7, a primeira linha). Em células de leveduras expostas a peróxido de hidrogênio, parte da Nuc1p::GFP foi translocada para o núcleo (ver coloração com DAPI), como demonstrado por BÜTTNER *et al*, 2007, mas também acumulado nas mitocôndrias, como indicado pela coloração com MitoTracker (Figura 7, segunda linha). Em contraste, quando as leveduras foram expostas a 0,125% de própolis por 15 minutos, cerca de 100% das células de levedura mostraram uma distribuição da Nuc1p::GFP tipo pontilhada, sugerindo fragmentação mitocondrial, como confirmada também por MitoTracker (Figura 7, terceira linha) . Além disso, a Nuc1p::GFP não translocou para o núcleo, como mostrado pela coloração com DAPI (Figura 7, terceira linha). Estes resultados indicam fortemente que o citocromo c, mas não Nuc1p, está envolvido na morte celular mediada própolis em *S. cerevisiae*.

Figura 7: Nuc1p não está envolvido na sensibilidade à própolis. Células de leveduras Nuc1::GFP foram crescidas por 16 horas a 30°, expostas ou não a própolis (0,125% por 15 minutos) e então visualizadas no microscópio de fluorescência. Um controle positivo foi feito pela incubação das células crescidas por 16 horas com 0,4 mM de H₂O₂, por 20 horas a 30 °C. Barras: 5 µm.

4.1.3. Avaliação da biblioteca de deleção de levedura para sensibilidade à própolis

Para elucidar as funções de genes que possam ser necessárias para a sensibilidade da própolis em eucariotos, a coleção completa de linhagens haplóides deletadas de *S. cerevisiae* foi avaliada em placas YPD contendo 2,73% etanol (como um controle, que é a concentração de etanol presente em 0,5% do extrato alcoólico de própolis), 0,250 e 0,500% de própolis, durante 5 dias a 37°C. Cerca de 4.800 cepas diferentes foram envolvidas nesta triagem, e depois de três “screenings” de confirmação, 138 cepas deletadas foram identificadas com diferentes graus de sensibilidade a própolis quando comparadas com a cepa do tipo selvagem correspondente (Anexo A). Os dados de sobrevivência, obtidos a partir de linhagens deletadas de levedura, levam ao questionamento de como a ausência de determinadas proteínas nas leveduras sensíveis a própolis afetam diferentes processos biológicos que levam à morte celular. Nesse sentido, foi iniciada a busca por proteínas potenciais e/ou mecanismos e seus respectivos processos biológicos que são afetados pela exposição à própolis.

Para atingir esse objetivo, diferentes redes PPPI usando dados de deleção em leveduras foram obtidas do banco de dados STRING. Proteínas compartilhadas e sub-redes presentes na rede principal PPPI (Figura 8) foram identificadas usando o *Cytoscape-associated plugin MCODE* e submetidas a uma análise *Gene Ontology* (GO), a fim de obter informações sobre a natureza e o número de sub-redes pertencentes à rede e seus respectivos processos biológicos associados. Os resultados obtidos das análises MCODE e GO mostraram que a rede final PPPI (Figura 8A) contém 553 nós e 3651 conectores e é composta por seis *clusters* fortemente interligados, cada um compreendendo diferentes processos biológicos. As análises GO revelaram que esses processos biológicos podem ser classificados em: (i) tradução/energia proveniente da oxidação de compostos orgânicos/manutenção do genoma mitocondrial, (ii) a transcrição/organização cromossômica e biogênese/fase G1 do ciclo celular mitótico, (iii) processo metabólico do ácido carboxílico/proteína alvo para o peroxissome/ metabolismo respiratório, (iv) estabelecimento e/ou manutenção da arquitetura da cromatina/desacetilação de histona, (v) processo catabólico do RNA, e (vi) mecanismos de divisão celular. Como esperado, as proteínas que não podem ser classificadas em qualquer um desses *clusters* também foram identificadas na rede (*unclustered proteins subnetwork*; Figuras 8A-B).

Figura 8: Sistemas biológicos para a sensibilidade à própolis em *S. cerevisiae*. (A) Rede de interação física proteína-proteína (PPPI) obtida de dados de deleção em levedura, onde a ausência de determinados genes aumenta a sensibilidade das células para o tratamento com própolis. (B) Distribuição de proteínas (número de nós) associadas à sensibilidade induzida por própolis, como observado nos dados de leveduras deletadas em diferentes *clusters* e *unclustered proteins* da rede principal PPPI. *Cluster 1* = tradução/energia proveniente da oxidação de compostos orgânicos/manutenção do genoma mitocondrial; *cluster 2* = transcrição/organização cromossômica e biogênese/fase G1 do ciclo celular mitótico; *cluster 3* = processo metabólico do ácido carboxílico/proteína alvo para o peroxissome/ metabolismo respiratório; *cluster 4* = estabelecimento e/ou manutenção da arquitetura da cromatina/desacetilação de histona; *cluster 5* = processo catabólico do RNA; *cluster 6* = mecanismos de divisão celular; e *unclustered nodes*.

A análise da distribuição das proteínas cuja deleção aumenta a sensibilidade das linhagens de levedura a própolis indicaram que ambos *cluster 6* e *unclustered protein subnetworks* contêm 61,73% das proteínas totais observadas no ensaio de deleção (Figura 8B; Anexo B). Os outros 38,27% das proteínas ficaram distribuídas entre *cluster 1-5* (Figura 8B; Anexo B), mostrando que o *cluster 6* e *unclustered protein subnetworks* parecem ser importantes para as respostas pleiotrópicas observadas quando as células de levedura são submetidas ao tratamento com própolis. Quando as proteínas da sub-rede do *cluster 6* (Figura 9; Anexo B) foram submetidas à análise GO, os dados indicaram que os principais processos biológicos encontrados dentro desta sub-rede estavam associados à divisão celular e a reprodução, que inclui o estabelecimento da polaridade celular, ciclo celular e brotamento (Tabela 2).

A.

B.

Figura 9: (A) Sub-redes de proteínas (cluster 6) associadas à divisão celular, estabelecimento da polaridade celular, cadeia de transporte de elétrons e acidificação vacuolar. (B) Sub-redes de proteínas *Unclustered* associadas à macroautofagia, proteína alvo para vacúolo, resposta celular à privação de nutrientes e estresse, transporte vacuolar, comunicação celular e regulação da polimerização protéica.

Tabela 2: Classes específicas *gene ontology* (GO) derivadas da rede de interação física proteína-proteína (PPPI) observadas no *cluster* 6.

Processo Biológico	Classificação GO	Valor de p^a	valor corrigido de p^b	K^c	f^d
Divisão celular	51301	2.95×10^{-14}	2.56×10^{-11}	36	323
Brotamento celular	7114	1.31×10^{-13}	5.68×10^{-11}	19	82
Reprodução assexual	19954	4.16×10^{-13}	1.20×10^{-10}	19	87
Reprodução de um organismo unicelular	32505	6.45×10^{-13}	1.40×10^{-10}	19	89
Estabelecimento da polaridade celular	30468	1.23×10^{-12}	1.67×10^{-10}	20	104
Transporte de elétrons mitocondrial, ubiquinol para citocromo c	6122	2.79×10^{-11}	2.11×10^{-9}	8	11
Citocinese	16288	2.91×10^{-11}	2.11×10^{-9}	19	109
Ciclo celular	7049	2.13×10^{-9}	1.05×10^{-7}	40	566
Cadeia de transporte de elétrons	22900	8.55×10^{-7}	1.61×10^{-5}	8	31
Redução do pH	45851	2.63×10^{-5}	3.08×10^{-4}	6	24
Acidificação vacuolar	7035	2.63×10^{-5}	3.08×10^{-4}	6	24
Regulação negativa da transcrição do promotor da RNA polymerase II pela glicose	433	3.30×10^{-4}	2.63×10^{-3}	3	6

^aValor de p calculado pela distribuição hipergeométrica de uma classe ontológica visualizada na rede de interação.

^bValores calculados, baseados nos valores de p obtidos depois que FDR foi aplicada.

^cNúmero total de proteínas encontradas na rede que pertencem a uma ontologia do gene.

^dNúmero total de proteínas que pertencem a uma ontologia do gene específica.

Além disso, as proteínas da sub-rede do *cluster* 6 parecem ser importantes para a cadeia de transporte de elétrons mitocondrial, a acidificação vacuolar e a regulação negativa da transcrição do promotor da RNA polimerase II (Tabela 2). Curiosamente, a análise GO das unclustered proteins subnetwork (Figura 9B; Anexo B) mostrou que essas proteínas são importantes para a regulação da macroautofagia associada com a proteína alvo para vacúolo e a resposta celular à privação de nutrientes, entre outros processos (Tabela 3).

Tabela 3: Classes específicas *gene ontology* (GO) derivadas da rede de interação física proteína-proteína (PPPI) de sub-redes de proteínas *unclustered*.

Processo Biológico	Classificação GO	Valor de p^a	Valor corrigido de p^b	k^c	f^d
Macroautofagia	34262	7.44×10^{-14}	5.44×10^{-11}	11	23
Via metabólica CVT	32258	9.36×10^{-12}	3.43×10^{-9}	9	18
Proteína alvo para vacúolo	6623	7.37×10^{-11}	1.80×10^{-8}	13	62
Resposta celular à privação de nutrientes	9267	3.03×10^{-10}	5.54×10^{-8}	11	44
Resposta celular ao estresse	33554	1.06×10^{-9}	5.09×10^{-7}	11	49
Resposta celular aos níveis de nutrientes	31669	4.86×10^{-9}	5.09×10^{-7}	11	56
Resposta à privação de nutrientes	42594	1.80×10^{-8}	1.46×10^{-6}	11	63
Transporte vacuolar	7034	5.05×10^{-8}	3.69×10^{-6}	14	122
Comunicação celular	7154	1.44×10^{-6}	6.21×10^{-5}	20	317
Regulação da polimerização protéica	32271	1.76×10^{-6}	6.79×10^{-5}	4	6

^aValor de p calculado pela distribuição hipergeométrica de uma classe ontológica visualizada na rede de interação.

^bValores calculados, baseados nos valores de p obtidos depois que FDR foi aplicada.

^cNúmero total de proteínas encontradas na rede que pertencem a uma ontologia do gene.

^dNúmero total de proteínas que pertencem a uma ontologia do gene específica.

Nas próximas duas subseção mostra-se a validação de dados sobre o papel desempenhado pelos *clusters* relacionados à cadeia de transporte mitocondrial de elétrons e a acidificação vacuolar.

4.1.4. Linhagens “Petite” são menos sensíveis a própolis

Recentemente foi demonstrado que durante a necrose em *S. cerevisiae* há uma inibição da função mitocondrial com a correspondente redução da produção de ATP (para uma revisão, ver EISENBERG *et al*, 2010). Ao longo deste trabalho, há vários indícios de que a própolis induz a uma necrose secundária como mecanismo de morte celular em *S. cerevisiae*.

Além disso, como foi demonstrado na seção anterior, quando vários genes relacionados com a energia proveniente da oxidação de compostos orgânicos/manutenção do genoma mitocondrial/cadeia de transporte de elétrons mitocondrial são deletados, há um aumento na sensibilidade à própolis em *S. cerevisiae*. Assim, decidiu-se investigar uma possível correlação entre a sensibilidade de própolis, mitocôndria e a morte celular por necrose em *S. cerevisiae*, isolando linhagens “petite” desta levedura e verificando a sua sensibilidade à própolis. Para isolar células BY4742 ρ^0/ρ^- , a linhagem respiratória - competente BY4742 (BRACHMANN *et al*, 1998) foi mutageneizada com brometo de etídio e as colônias resultantes com deficiência respiratória “petite” foram cruzadas com linhagens teste ρ^0/ρ^- (TZAGOLOFF *et al*, 1975) para confirmar a incapacidade respiratória das “petite” isoladas.

Duas destas cepas “petite”, BY4741A ρ^-/ρ^0 e BY4742 ρ^-/ρ^0 , e a linhagem selvagem correspondente foram expostas a 0,125% de própolis por 20 minutos e determinou-se a viabilidade celular (Figura 10). As cepas “petite” apresentaram a viabilidade dramaticamente aumentada quando comparadas à cepa selvagem (comparar 95 e 75 % das linhagens BY4741A ρ^-/ρ^0 e BY4742 ρ^-/ρ^0 , respectivamente, contra aproximadamente 10 % na linhagem selvagem, como porcentagem de sobrevivência; Figura 10A). Assim, quando foi feito o fracionamento das proteínas celulares destas linhagens “petite”, o citocromo *c* não foi observado nos seus extratos de proteína citoplasmática nas células expostas à própolis, mas apenas em extratos de proteínas mitocondriais (Figura 10B). O efeito da própolis em mitocôndrias isoladas da cepa de levedura W303 foi medido através do monitoramento da taxa de consumo de oxigênio como consequência da atividade da NADH oxidase mitocondrial. Na presença de própolis, a atividade da NADH oxidase é 1/10 do que o observado antes da adição deste composto, indicando o seu efeito tóxico sobre a respiração (Figura 10C). Estes resultados sugerem fortemente que a sensibilidade a própolis de *S. cerevisiae* é dependente da função mitocondrial.

Figura 10: Mutantes “Petite” são mais tolerantes a própolis. (A) A viabilidade das células de levedura da cepa selvagem e dos mutantes “petite”, crescidas por 16 horas e expostas a 0,125% de própolis foi determinada pelo plaqueamento de concentrações apropriadas das células e contagem do número de colônias crescidas. Os resultados são referentes à média $\pm$ o desvio padrão de três experimentos independentes (* $p < 0.05$). O tratamento estatístico foi realizado pelo método ANOVA one way, com pós teste Newman Keuls. (B) Células do mutante “petite” expostas a própolis não induzem a liberação do citocromo c para o citoplasma. As células de levedura foram crescidas por 16 horas a 30°C e expostas ou não a 0,125% de própolis por 10 min a 30°C. Após isso, elas foram coletadas e fracionadas (mitocôndria e citoplasma), como descrito em Materiais e Métodos. As proteínas foram separadas em gel de poliácridamida e coradas com Coomassie (painel esquerdo); elas estavam na mesma concentração das proteínas que foram transferidas para a membrana de nitrocelulose (painel a direita). Esta membrana foi marcada com anticorpo de coelho anti-citocromo c. (C) Atividade da NADH oxidase na levedura selvagem em presença de própolis. Mitocôndrias (100 μ g de proteína) da linhagem selvagem W303 foram utilizadas para o monitoramento da taxa de consumo de oxigênio como consequência da atividade da NADH oxidase mitocondrial. As setas indicam a adição de própolis (concentração final de 0,125%) ou etanol e KCN. Estes são traços representativos de um experimento, que foram reproduzidos com mitocôndrias isoladas em três dias diferentes.

4.1.5. Um papel para o vacúolo e a H⁺-ATPase vacuolar na sensibilidade a própolis

Outro conjunto de genes observados como importantes para a sensibilidade a própolis estava relacionado à acidificação vacuolar e à proteínas ligadas a macroautofagia associada com a proteína alvo para o vacúolo e a resposta celular à privação de nutrientes (Tabela 3). A acidificação de organelas está envolvida na seleção de proteínas nas vias biossintéticas e endocíticas, a ativação proteolítica de precursores de zimogênio e transporte transmembrana de conteúdo viral e toxinas (para uma revisão, ver KANE, 2006). Em fungos, o vacúolo tipo lisossomo também desempenha um papel no armazenamento de blocos de construção metabólica, na homeostase do cálcio e no controle osmótico. A acidificação vacuolar é essencial para todas estas funções. A ATPase vacuolar (V-ATPase) é uma enzima essencial para catalisar todos esses processos. A sua principal função na célula eucariótica é o transporte *ATP-driven* de prótons do citosol em organelas ácidas (KANE, 2006). Quando vários genes (*VPH1*, *VMA3,4,5,11,22*, *RAV1* e *SOP4*) envolvidos na montagem da V-ATPase de levedura foram deletados, as linhagens correspondentes das leveduras deletadas tornaram-se mais sensíveis à própolis (Figura 11A).

Recentemente, Rosado *et al* (2008) desenvolveram um método para monitorar autofagia utilizando Rosella, um biossensor composto de uma proteína fluorescente vermelha *fast-maturing* pH estável fusionada a uma proteína variante fluorescente verde pH sensível. O seu modo de ação baseia-se em diferenças de pH entre os diferentes compartimentos celulares e o vacúolo. A Rosella é um biossensor geneticamente codificado com dupla emissão de cor de um produto da fusão da proteína fluorescente vermelha *fast-maturing* insensível ao pH, DsRed (BEVIS & GLICK 2002), com uma proteína variante da GFP sensível ao pH (MIESENBOCK *et al*, 1998). A lógica deste ensaio depende da diferença entre o pH do vacúolo (ou lisossomo, pH de aproximadamente 6,2 ou 4,8, respectivamente) e de outros compartimentos da célula (pH > 7,0). Em células de leveduras cultivadas por 16 horas, tanto a fluorescência vermelha quanto a verde foram distribuídas uniformemente por todo o citoplasma, mas ausente no vacúolo (Figura 11B, primeira linha). As células foram então transferidas para o meio de cultura com a privação de nutrientes, neste caso, sem nitrogênio, a fim de induzir a autofagia. Após 4 horas de incubação, foi adicionado em tampão pH 7,5 o protonóforo CCCP, por 10 minutos. A

elevação do pH é verificada devido à ação do CCCP no colapso dos gradientes de pH através das membranas (Rosado et al., 2008). Desta maneira, para este controle positivo, observou-se acúmulo da fluorescência vermelha e verde no vacúolo em aproximadamente 40% das células (Figura 11B, segunda linha).

Em seguida, as células foram expostas a 0,125% de própolis e, após 90 minutos de incubação, verificou-se o acúmulo de um fenótipo sutil de fluorescência vermelha, mas não verde no vacúolo em aproximadamente 60% de células da linhagem selvagem (Figura 11B, segundo e linhas de terceiros).

Figura 11: Própolis induz acidificação vacuolar em *S. cerevisiae*. (A) Ensaio de diluição em meio YPD suplementado com diferentes concentrações de própolis. Células de levedura foram crescidas por 96 horas, a 30°C. (B) Rosella expressa no citosol da levedura migra para o vacúolo após exposição a própolis. Imagens em DIC e com fluorescência são mostradas para células do tipo selvagem sob condições normais de crescimento (primeira linha), após a privação de nitrogênio por 4 horas e adição de CCCP (segunda linha) e expostas a própolis (0,125%, por 90 minutos, a 30°C) (terceira linha). A posição do vacúolo nas imagens com DIC está delimitada por uma linha tracejada branca. Os vacúolos vazios são destacados por setas brancas; os vacúolos preenchidos são destacados pelas setas amarelas. Barra: 5 µm.

A autofagia pode estar envolvida no turnover de proteínas de vida longa e em organelas [por exemplo, autofagia mitocondrial (mitophagy) e do retículo endoplasmático (reticulophagy)] (MAIURI *et al*, 2007). Autofagia é também um processo importante durante a privação de nutrientes, ou seja, pelo catabolismo de macromoléculas, a autofagia gera substratos metabólicos (MAIURI *et al*, 2007). A autofagia tem sido seguida por vários métodos bioquímicos e morfológicos (para revisões, ver KLIONSKY *et al*, 2007; KLIONSKY *et al*, 2008). A expressão de GFP-Atg8p em levedura pode ser usada para acompanhar a localização ou a acumulação da estrutura pré-autofagossomal, autofagossomal e corpos autofágicos.

Observou-se que cepas com os genes *ATG1* [uma proteína serina/treonina quinase necessária para a formação de vesículas na autofagia e na via metabólica citoplasma-ao-vacúolo (CVT)], *ATG10* (uma enzima conservada conjugada, tipo E2 que medeia a formação do conjugado Atg12p-Atg5p, que é um passo crítico na autofagia), e *ATG20* [um membro da família nexina necessária para a via citoplasma-ao-vacúolo (CVT) e para a endossomal] deletados são mais sensíveis à própolis (Figura 12A) (para revisões sobre os genes de levedura envolvidos na autofagia, ver KANKI & KLIONSKY, 2010; KOURTIS e TAVERNARAKIS, 2009). O acúmulo de mRNA de *ATG14* e *ATG8* e o monitoramento de GFP::Atg8 foram utilizados como ensaios para verificar se a própolis induz um aumento da autofagia. A própolis induziu o acúmulo de mRNA de *ATG8* e *ATG14* cerca de 4 a 14 vezes, e duas vezes, respectivamente (5 e 10 minutos de exposição à 0,125% de própolis, Figura 12B). Além disso, a GFP::Atg8 é translocada para estruturas semelhantes a vacúolos após a exposição a 0,125 % de própolis por 20 minutos (Figure 12C).

Figura 12: Própolis induz autofagia em *S. cerevisiae*. (A) Ensaio de diluição em meio YPD com diferentes concentrações de própolis. Células de levedura foram crescidas por 96 horas a 30°C. (B) Células da linhagem selvagem foram crescidas por 9 horas em meio líquido YPD, a 30°C e expostas ou não a 0,125% de própolis por 5 ou 10 minutos. O RNA foi extraído e real-time RT-PCR foi o método utilizado para quantificar a acumulação de mRNA. A quantidade medida de mRNA em cada uma das amostras foi normalizada usando os valores de CT (*Cycle Threshold*), obtidos para as amplificações de actina (*ACT1*) na mesma placa. A quantificação relativa da expressão gênica de *ATG8* e *ATG14* e também de *ACT1* foi determinada por uma curva padrão (valores de CT plotados contra o logaritmo do número de cópias de DNA). Os resultados são a média $\pm$ o desvio padrão de quatro séries de experimentos. (C) Linhagem de levedura GFP::Atg8 exposta ou não a própolis (0,125%, por 20 minutos, a 30°C). Barra: 5 μ m.

Em conjunto, estes dados indicam que a acidificação vacuolar e a autofagia são importantes para a morte causada por própolis em leveduras.

4.1.6. DISCUSSÃO

Os três principais tipos de própolis em todo o mundo apresentam diferentes composições: (1) a CAPE (feniletil éster do ácido caféico), presente na própolis Européia, Extremo Oriente e na Nova Zelândia, (2) a artepillin C (ARC), encontrada na própolis verde brasileira e (3) a própolis Brasileira vermelha. Recentemente, foi realizada uma triagem de 1.266 compostos com atividades farmacêuticas conhecidas e identificaram-se 15 que prolongaram a sobrevivência de nematóides infectados com *C. albicans* e que também inibiram a filamentação *in vivo* de *C. albicans* (BREGER *et al*, 2007). Um destes compostos, CAPE, apresentou atividade antifúngica em um modelo de candidíase em camundongos (BREGER *et al*, 2007). A CAPE está presente em um tipo de própolis conhecido como *Poplar-type*, enquanto que a própolis verde Brasileira é um *Baccharis-type*. Ainda não foi possível identificar vestígios de CAPE na própolis verde Brasileira (dados não apresentados; SHIMAZAWA *et al*, 2005; TEIXEIRA *et al*, 2005). Assim, foram investigados alvos moleculares envolvidos na letalidade celular causada pela própolis Brasileira verde. Não foi verificado qualquer efeito da morte celular quando a levedura *S. cerevisiae* foi exposta a altas concentrações dos quatro principais compostos que estão presentes na própolis verde; por isso, decidiu-se verificar os efeitos da própolis na letalidade celular de *S. cerevisiae* usando extratos alcoólicos de própolis.

A morte celular em *S. cerevisiae* é frequentemente demonstrada por marcadores indicativos de apoptose e/ou necrose, tais como a externalização de fosfatidilserina para a porção externa da membrana plasmática, condensação da cromatina, a geração de espécies reativas de oxigênio, o acúmulo de iodeto de propídio, a localização de HMGB1/Nhp6p, a detecção de rupturas da membrana plasmática e a completa desintegração das estruturas subcelulares (CARMONA-GUTIERREZ *et al*, 2010; EISENBERG *et al*, 2010). Além disso, o acúmulo de ROS, a fragmentação mitocondrial, a liberação de citocromo c, a ativação da metacaspase Yca1p, o fator indutor de apoptose Aif1p, a endonuclease G Nuc1p, a serina protease OMI, perturbações do citoesqueleto e modificações epigenéticas da cromatina também são observados (BÜTTNER *et al.*, 2007). A necrose também tem

sido descrita durante o envelhecimento cronológico da levedura (EISENBERG *et al*, 2009). Verificou-se que os marcadores de apoptose e necrose são induzidos por própolis, mas o aumento do tempo de exposição à própolis intensifica o número de células com marcadores para a necrose, como PI e translocação nucleo-citosol do Nhp6Ap::GFP. Os dados obtidos nesta fase do trabalho indicam um duplo papel para o tratamento com a própolis: como um agente que induz apoptose e necrose secundária. Estes efeitos são parcialmente mediados pela metacaspase YCA1, uma vez que o mutante $\Delta YCA1$ é mais resistente a própolis. Em *S. cerevisiae*, a superexpressão de YCA1 causa morte celular enquanto que a deleção deste gene protege contra a morte devido à espécies reativas de oxigênio ou envelhecimento cronológico (MADEO *et al*, 2002). Além disso, observou-se também que o Nuc1p não é translocado para o núcleo quando há indução de morte celular pela própolis. Curiosamente Büttner, *et al* (2007) observaram que a morte por apoptose mediada por Nuc1p não requer YCA1. Além disso, marcadores simultâneos para apoptose e necrose também foram observados durante a indução de morte celular por própolis em *T. cruzi* (MENNA-BARRETO, *et al* 2009).

Existem várias condições em que ROS mitocondriais têm sido associados à apoptose em levedura (para revisões, ver EISENBERG *et al*, 2010; OTA *et al*, 2001; PERRONE *et al*, 2008). A própolis pode induzir a formação de ROS e é mais letal quando *S. cerevisiae* cresce na presença de glicerol e etanol como fontes de carbono (dados não mostrados), sugerindo que a respiração aumenta a letalidade da própolis. Verificou-se que a indução de morte celular pela própolis estimula a liberação do citocromo c e que também células ρ^0 são mais tolerantes a própolis. O citocromo c é uma proteína mitocondrial com uma função na cadeia respiratória e uma função adicional como um ativador da caspase-9 da via intrínseca de apoptose de mamíferos (KUSHNAREVA & NEWMAYER, 2010). O tratamento de células de levedura com ácido acético leva à liberação de citocromo c (LUDOVICO, *et al*, 2006). O rompimento do citocromo c impede parcialmente a morte celular induzida por ácido acético e, conseqüentemente, as células ρ^0 mostram resistência contra a morte celular induzida por este ácido (LUDOVICO, *et al*, 2006). No entanto, não é muito clara se a liberação do citocromo c pode levar à formação em *S. cerevisiae* de uma estrutura semelhante ao apoptossomo e a ativação das caspases nos mamíferos. Em contraste, Sripriya *et al*. (2009) observaram a perda do potencial de

membrana mitocondrial e ausência de liberação de citocromo *c* durante a morte celular por necrose em *S. cerevisiae* induzida pela expressão de um *proteinaceous elicitor hairpin* (Pss) de *Pseudomonas syringae*. Contudo, a deficiência de DNA mitocondrial funcional, e a conseqüente incapacidade de respirar dos mutantes "*petite*" conferiram resistência à morte.

A triagem da coleção completa de linhagens haplóides deletadas de *S. cerevisiae* foi feita com o objetivo de compreender como a própolis, um produto de origem natural com composição química complexa, afeta o metabolismo celular. Essa estratégia já foi usada extensivamente para a compreensão de como outros compostos afetam o metabolismo das células de *S. cerevisiae* (BLACKBURN & AVERY, 2003; CHAN *et al*, 2000; DESMOUCELLES *et al*, 2002; HILLENMEYER *et al*, 2008; SCHERENS & GOFFEAU, 2004; ZEWAİL *et al*, 2003). Desta maneira, através da aplicação de biologia de sistemas, observou-se que a maioria das proteínas, cuja a deleção aumenta a sensibilidade de linhagens de leveduras a própolis, estão envolvidas em mecanismos de divisão celular, cadeia de transporte de elétrons mitocondrial, a acidificação vacuolar, a regulação da macroautofagia associada com o transporte de proteínas para o vacúolo, a resposta celular a privação de nutrientes e a regulação negativa da transcrição do promotor da RNA polimerase II.

A cadeia de transporte de elétrons mitocondrial e as vias de acidificação vacuolar foram investigadas e validadas em detalhes. Mostrou-se que a própolis induz a acidificação vacuolar e a translocação de Atg8p para os vacúolos, uma das marcas da autofagia. Em *S. cerevisiae*, o vacúolo é muito importante para manter a homeostase celular compreendendo a regulação do pH intracelular e a degradação, principalmente durante a limitação de nutrientes, de proteínas e organelas por autofagia (para revisões, ver ARIÑO *et al*, 2010; INOUE & KLIONSKY, 2010; LYNCH-DAY & KLIONSKY, 2010). A morte celular induzida por ácido acético é aumentada em mutantes de *S. cerevisiae* que apresentam o gene *VPS* deletado (os genes *VPS* estão envolvidos na fusão vacuolar homotípica, triagem de proteína vacuolar e são essenciais para a função vacuolar normal) (SCHAUER *et al*, 2009). Tem-se observado que o pH intracelular foi acidificado em células mutantes *VPS* após tratamento com ácido acético (SCHAUER *et al*, 2009). É possível que a perturbação do controle homeostático do pH possa provocar a necrose pela

liberação de proteases pró-necrótica, as quais iriam encontrar um pH ótimo para sua atividade enzimática no citosol acidificado (EISENBERG *et al*, 2010).

Assim, procurou-se enfatizar nesta etapa do trabalho, a importância de *S. cerevisiae* como sistema-modelo para entender, a nível molecular, o mecanismo de ação de um composto de origem natural. Há vários estudos descrevendo os mais diversificados efeitos da própolis sobre os diferentes sistemas biológicos, como a quimioprevenção do câncer (SZLISZKA & KROL, 2010), agentes cariostáticos em estreptococos (LIBÉRIO *et al*, 2009), ação imunomoduladora e a produção de linfócitos e anticorpos (SFORCIN, 2007). Este estudo é o primeiro a investigar, pelo uso da genômica funcional, como a própolis influencia o metabolismo de um organismo. Isto fornece novas possibilidades para a compreensão e a validação da própolis como agente terapêutico contra infecções fúngicas.

Módulo 4.2: Análise de sistemas de *Saccharomyces cerevisiae* exposta a própolis

Artigo: “Systems analysis of *Saccharomyces cerevisiae* exposed to propolis. De Castro, P. A.; Savoldi, M.; Bonatto, D.; Malavazi, I.; Goldman, M.H.S.; Berretta, A. A; Goldman, G. H. BMC Complementary and Alternative Medicine, submitted, dec, 2011.

4.2. RESULTADOS E DISCUSSÃO

4.2.1. A análise da hibridação por microarray

Anteriormente a este trabalho havia apenas um estudo na literatura relatando o perfil transcricional de células eucarióticas expostas a própolis (Gruber & Holtz, 2010), no qual a mesma e outros oito compostos com propriedades antioxidantes conhecidas, foram aplicados por 24 horas para fibroblastos humanos normais de epitélio e queratinócitos. Os autores identificaram 205 genes importantes para a pele; destes, cinco (ATP citrato sintase, aquaporina 3, citocromo c oxidase 1, óxido nítrico sintase 3, e hidroxilase 3) parecem “upregulated” e apenas um (receptor de progesterona) está “downregulated” para ambas as linhagens celulares.

Como foi mostrado anteriormente, *S. cerevisiae* foi utilizada como um sistema modelo eucariótico para identificar genes que são importantes para a morte celular mediada por própolis, uma vez que quando as células desta levedura, na fase de crescimento exponencial, foram expostas a 0,125% de própolis por 5 e 10 minutos, demonstrou-se que há uma sobrevivência de 24,1 e 6,3%, respectivamente. Para ampliar nosso conhecimento e adquirir uma visão sobre quais vias metabólicas são moduladas durante a exposição de *S. cerevisiae* a própolis, determinou-se o seu perfil transcricional. Assim, o RNA total extraído da cultura de células da levedura, na fase exponencial de crescimento, expostas a própolis, foi usado para amplificar cRNAs marcados com fluorescência para hibridações de microarray.

A expressão do RNAm da cepa BY4742 de *S. cerevisiae*, crescida por 9 horas e expostas a 0,125% de própolis por 5 e 10 minutos, foi comparada com células de levedura expostas por 5 e 10 minutos a 0,68% de etanol (o que corresponde a 60% de etanol presente no extrato alcoólico de própolis para a concentração de 0,125%). Nestes experimentos, o objetivo principal era se concentrar em genes que têm a expressão de mRNA aumentada ou diminuída. O conjunto de dados completo foi depositado no “Gene Expression Omnibus” (GEO) do “National Center of Biotechnology Information” (NCBI), com o número GSE33971 (<http://www.ncbi.nlm.nih.gov/projects/geo/query/acc.cgi?acc=GSE33971>).

Foram observados 339 genes modulados em pelo menos um “timepoint” quando comparado com o tempo de referência (p -value 0,01). Estes genes estão envolvidos com uma variedade de processos celulares com expressão de mRNA

aumentada ou diminuída (Anexo C mostra os genes com log ratios ≥ 1 (136 genes) ou ≤ 1 (203 genes), respectivamente. Foi utilizada a análise da Ontologia Gênica (GO) com o objetivo de classificar a quais processos biológicos os genes modulados estão associados (Anexo C). A análise da GO dos genes com aumento da abundância de mRNA mostrou um enriquecimento para (i) transporte de proteínas e transporte (23,1%), (ii) tradução (12,8%), (iii) mitocôndrias (8,3%), (iv) processo metabólico, e (v) processo metabólico do DNA e transcrição (7,3%) (Figura 13).

Figura 13: Análise da Ontologia Gênica dos genes com abundância de RNAm aumentada no ensaio de hibridação de microarray de *S. cerevisiae* exposta a própolis.

Neste grupo de genes, foram observados vários genes que codificam transportadores envolvidos na resistência a múltiplas drogas (MDR) (para revisões ver ERNST *et al*, 2010; SHAHI & MOYE-ROWLEY, 2009; SÁ-CORREIA *et al*, 2009; SIPOS & KUCHLER, 2006), sugerindo que a própolis pode ativar, no âmbito transcricional, o complexo conjunto de genes responsáveis pela MDR em *S. cerevisiae* (Tabela 4).

Tabela 4: Genes selecionados que foram mais expressos ($\log_2 \geq 1.0$) durante a exposição de *S. cerevisiae* a própolis (para uma lista completa de genes mais expressos, ver Anexo C).

FLR1	Multidrug transporter of the MFS, involved in efflux of fluconazole, diazaborine, benomyl, methotrexate, and other drugs
PDR15	ATP binding cassette (ABC) transporter, multidrug transporter and general stress response factor implicated in cellular detoxification
AZR1	Transporter of the MFS, involved in resistance to azole drugs such as ketoconazole and fluconazole
YOR1	ATP-binding cassette (ABC) transporter, multidrug transporter mediates export of many different organic anions including oligomycin
PDR10	ATP-binding cassette (ABC) transporter, multidrug transporter involved in the pleiotropic drug resistance network
PDR12	ATP-binding cassette (ABC) transporter, weak-acid-inducible multidrug transporter required for weak organic acid resistance
ENA1	P-type ATPase sodium pump, involved in Na ⁺ and Li ⁺ efflux to allow salt tolerance
ENA2	P-type ATPase sodium pump, involved in Na ⁺ efflux to allow salt tolerance; likely not involved in Li ⁺ efflux
VMA21	Membrane protein that is required for vacuolar H ⁺ -ATPase (V-ATPase) function, although not an actual component of the V-ATPase
TPO1	Polyamine transporter (MFS) that recognizes spermine, putrescine, and spermidine
TPO4	Polyamine transport protein (MFS) recognizes spermine, putrescine, and spermidine
ATG22	Vacuolar integral membrane protein required for efflux of amino acids during autophagic body breakdown in the vacuole
VMA7	Subunit F of the eight-subunit V1 peripheral membrane domain of vacuolar H ⁺ -ATPase (V-ATPase)
PKR1	V-ATPase assembly factor, functions with other V-ATPase assembly factors in the ER to efficiently assemble the V-ATPase
GRX4	Hydroperoxide and superoxide-radical responsive glutathione-dependent oxidoreductase
GTT3	Protein of unknown function with a possible role in glutathione metabolism
TSA2	Stress inducible cytoplasmic thioredoxin peroxidase; cooperates with Tsa1p in the removal of reactive oxygen
GTT2	Glutathione S-transferase capable of homodimerization; functional overlap with Gtt2p, Grx1p, and Grx2p
GND2	6-phosphogluconate dehydrogenase, catalyzes an NADPH regenerating reaction in the pentose phosphate pathway
TRX1	Cytoplasmic thioredoxin isoenzyme of the thioredoxin system which protects cells against oxidative and reductive stress

Além disso, também foram verificados vários genes que codificam proteínas importantes para a montagem da ATPase vacuolar e na via endocítica (Tabela 4 e Anexo C). Nenhum dos mutantes de deleção para estes genes que codificam transportadores mostrados no Anexo C foram observados como sendo mais sensíveis à própolis, exceto para *TPO1*, indicando uma resposta transcricional redundante destes genes a própolis. No entanto, quando vários genes (*VPH1*, *VMA3,4,5,11,22*, *RAV1* e *SOP4*) envolvidos na montagem da V-ATPase em levedura foram deletados, as correspondentes cepas mutantes tornaram-se mais sensíveis à própolis, como já descrito no módulo 4.1.

Como visto anteriormente (módulo 4.1) pelo screening de uma biblioteca de deleção de levedura, observou-se que a maioria das proteínas, cuja deleção aumenta a sensibilidade de cepas a própolis, estão envolvidas nos mecanismos de divisão celular, na cadeia de transporte de elétrons mitocondrial, na acidificação vacuolar, na regulação da macroautofagia associada com proteína alvo de vacúolo, e na resposta celular à privação de nutrientes e regulação negativa da transcrição do promotor da RNA polimerase II. Foi também mostrado que a própolis induz acidificação vacuolar e translocação de Atg8p para os vacúolos, um dos marcadores de autofagia. Em *S. cerevisiae*, o vacúolo é muito importante para manter a homeostase celular, compreendendo a regulação do pH intracelular e degradação, principalmente durante a limitação de nutrientes, de proteínas e organelas pela autofagia (para revisões, ver [ARIÑO *et al*, 2010; INOUE & KLIONSKY, 2010; LYNCH-DAY & KLIONSKY, 2010]).

A morte celular induzida por ácido acético é aumentada em mutantes deletados *VPS* de *S. cerevisiae* (os genes *VPS* estão envolvidos na fusão homotípica do vacúolo, classificação de proteína vacuolar e são essenciais para a função vacuolar normal) (EISENBERG *et al*, 2009). Foi verificado que o pH intracelular é acidificado em mutantes *VPS* após tratamento com ácido acético (EISENBERG *et al*, 2009). É possível que a perturbação do controle homeostático do pH possa provocar necrose pela liberação de proteases pró-necróticas, as quais iriam encontrar um pH ideal para a suas atividades enzimáticas no citosol acidificado (OTA *et al*, 2001).

Nesta análise do microarray, observaram-se vários genes que codificam proteínas importantes para a desintoxicação de ROS em *S. cerevisiae*, como *GRX*, *GTT2*, *GTT3*, *TSA2*, *GND2* e *TRX1* com aumento na acumulação de RNAm quando

S. cerevisiae é exposta a própolis (Tabela 4 e Anexo C). Curiosamente, há um aumento no acúmulo de RNAm de vários genes que codificam proteínas envolvidas na geração de ATP na mitocôndria, como *ATP17*, *ATP18*, *ATP19*, *ATP20*, *ATP21* e *COX8* (Anexo C). Existem várias condições em que a produção de ROS pelas mitocôndrias têm sido associada à apoptose em leveduras (para revisões, ver [OTA *et al*, 2001; PERRONE *et al*, 2008; EISENBERG *et al*, 2010; TESTE *et al*, 2009]). A própolis pode induzir a formação de ROS e é mais letal quando *S. cerevisiae* cresce na presença de glicerol e etanol como fontes de carbono (módulo 4.1), sugerindo que a respiração aumenta a letalidade da própolis.

A análise da GO dos genes com diminuição da abundância de RNAm mostrou um enriquecimento para (i) o processo metabólico do DNA e transcrição (25,7%), (ii) processamento de RNA (22,8%), e (iii) processo metabólico (14%) (Figura 14).

Figura 14: Análise da Ontologia Gênica dos genes com abundância de RNAm diminuída no ensaio de hibridação de microarray de *S. cerevisiae* exposta a própolis.

Existem vários genes relacionados com o cinetocoro, distribuição cromossômica e silenciamento da cromatina, que tem diminuição do acúmulo de RNAm de *S. cerevisiae* quando ocorre exposição à própolis (Tabela 5). A abundância de RNAm reduzida destes genes sugere que a própolis está ativando controles de “checkpoint” transcricionais envolvidos nas fases S- e M- importantes para a replicação de DNA e segregação cromossômica adequadas.

Tabela 5: Genes selecionados que foram menos expressos ($\log_2 \geq 1.0$) durante a exposição de *S. cerevisiae* a própolis (para uma lista completa de genes mais expressos, ver Anexo C).

SMC4	Subunit of the condensin complex; reorganizes chromosomes during cell division
CM10	Essential chromatin-associated protein involved in the initiation of DNA replication
CIN8	Kinesin motor protein involved in mitotic spindle assembly and chromosome segregation
IBD2	Component of the BUB2-dependent spindle checkpoint pathway, interacts with Bfa1p and functions upstream of Bub2p and Bfa1p
SPC110	Inner plaque spindle pole body (SPB) component, ortholog of human kendrin; involved in connecting nuclear microtubules to SPB
BRN1	Subunit of the condensin complex; required for chromosome condensation and for clustering of tRNA genes at the nucleolus
CEP3	Essential kinetochore protein, component of the CBF3 complex that binds the CDEIII region of the centromere
SLK19	Kinetochore-associated protein required for normal segregation of chromosomes in meiosis and mitosis
NSL1	Component of the MIND kinetochore complex which joins kinetochore subunits contacting DNA to those contacting microtubules
SPC24	Component of the evolutionarily conserved kinetochore-associated Ndc80 complex
MAM1	Monopolin, kinetochore associated protein involved in chromosome attachment to meiotic spindle
RLF2	Largest subunit (p90) of the Chromatin Assembly Complex (CAF-1) and Msi1p that assembles newly synthesized histones
HOS2	Histone deacetylase required for gene activation via specific deacetylation of lysines in H3 and H4 histone tails
ESA1	Catalytic subunit of the histone acetyltransferase complex (NuA4) that acetylates four conserved internal lysines of histone H4
LEO1	Component of the Paf1 complex, which associates with RNA polymerase II and is involved in histone methylation
SPT21	Protein required for normal transcription at several loci including HTA2-HTB2 and HHF2-HHT2
RFM1	Specificity factor that directs the Hst1p histone deacetylase to some of the promoters regulated by Sum1p
SGF29	Probable subunit of SAGA histone acetyltransferase complex
IES2	Protein that associates with the INO80 chromatin remodeling complex under low-salt conditions
IFH1	IFH1 exhibits genetic interactions with FHL1, overexpression interferes with silencing at telomeres and HM loc
NGG1	component of transcriptional adaptor and histone acetyltransferase complexes, the ADA, the SAGA, and the SLIK complexes

4.2.2. A validação da análise da hibridação por microarray

Para validar alguns dos nossos resultados, foram escolhidos nove genes diferentes a partir da análise do microarray, cujo RNAm apresentou abundância aumentada ou diminuída quando *S. cerevisiae* é exposta a própolis. Para isto, foram desenhadas sondas Lux fluorescentes e utilizada a análise de PCR em tempo real para quantificar a expressão de uma nova réplica biológica do RNAm isolado do tratamento por 5 e 10 minutos de exposição a 0,125% de própolis comparada com os controles correspondentes tratados com o 0,68% etanol.

O gene normalizador (constitutivo) utilizado, *TAF10*, codifica uma subunidade (145 kDa) de TFIID e complexos SAGA, envolvidos na iniciação da transcrição pela RNA polimerase II e na modificação da cromatina. Recentemente, este gene foi mostrado como um gene de referência adequado para análise da expressão gênica quantitativa por PCR em tempo real (TESTE *et al*, 2009). Além disso, *TAF10* sofreu modulação gênica em nossos experimentos de hibridação de microarrays (dados não mostrados). Assim, a quantidade medida de mRNA de um gene específico, em cada uma das amostras tratadas, foi normalizada utilizando os valores C_T obtidos para as amplificações de mRNA do *TAF10* nas corridas executadas na mesma placa.

Os resultados foram expressos como o número de vezes em que os genes apresentam abundância aumentada ou diminuída quando as leveduras foram expostas a própolis em relação ao tratamento controle com etanol (Figura 15). Foi avaliada a abundância de mRNA de *PDR15* (YDR406W), o qual codifica um transportador envolvido na resistência múltipla a drogas; *SNQ2* (YDR011W), que codifica um transportador ABC de membrana plasmática [“ATP-binding cassette (ABC) transporter”], um transportador de resistência múltipla a drogas envolvido na resistência a várias drogas e também na resistência a espécies de oxigênio singlete; *VMA7* (YGR020C) e *VMA21* (YGR105W) que codificam a subunidade F do domínio periférico de membrana V1 da H⁺-ATPase vacuolar que possui 8 subunidades (V-ATPase) e a proteína integral de membrana que é necessária para função da H⁺-ATPase vacuolar, respectivamente; *RLF2* (YPR018W) que codifica a maior subunidade (p90) do “Chromatin Assembly Complex” (CAF-1) com Cac2p e Msi1p que reúne as histonas recém-sintetizadas no DNA replicado; e *TIM10* (YHR005C-A) que codifica a proteína essencial do espaço intermembrana mitocondrial, e forma um

complexo com Tim9p (TIM10 complexo) que fornece as proteínas hidrofóbicas ao complexo TIM22 para a inserção na membrana interna.

Todos os genes apresentaram a abundância de mRNA aumentada em nossas hibridações de microarray, exceto para *RLF2*, que obteve diminuição do acúmulo de mRNA. Todos os cinco genes que mostraram aumento (*PDR15*, *TIM10*, *SNQ2*, *VMA7* e *VMA21*) e diminuição (*RLF2*) na análise de hibridação de microarray, mostraram aumento e diminuição correspondentes nos experimentos de PCR em tempo real (Figura 15A-F).

Figura 15: PCR em tempo real para os genes selecionados da análise de hibridação de microarray. *S. cerevisiae* foi crescida por 9 horas, em meio líquido YPD a 30°C, e depois, as células ($\sim 2 \times 10^7$ células ml^{-1}) foram transferidas para meio líquido fresco e expostas a 0,125% de própolis por 5 ou 10 minutos. A quantificação relativa da expressão dos genes *RLF2* (A), *PDR15* (B), *TIM10* (C), *SNQ2* (D), *VMA7* (E), *VMA21* (F), e *TAF10* foi determinada por uma curva padrão (valores de C_T plotados contra p logaritmo do número de cópias de DNA). Os resultados são referente à média $\pm$ o desvio padrão de 4 experimentos independentes.

Desta maneira, pode-se dizer que esta abordagem de hibridação de microarrays é capaz de fornecer informações sobre a modulação da expressão gênica em *S. cerevisiae* com um nível bastante elevado de confiança.

4.2.3. Análise por biologia de sistemas para a exposição a própolis

Os dados obtidos do transcriptoma para a exposição da cepa de *S. cerevisiae*, submetida aos tratamentos com própolis descritos neste trabalho, induzem ao questionamento de como os genes sub-expressos ou super-expressos afetam diferentes processos biológicos que são alterados durante a exposição a própolis. Neste sentido, iniciou-se uma busca por potenciais proteínas e/ou mecanismos e seus processos biológicos associados que são afetados pelas condições de tratamento. Para isto, redes PPPI usando os dados do transcriptoma deste trabalho foram obtidas do banco de dados Saccharomyces Genome Database (SGD) (Anexo D). Sub-redes (“clusters”) presentes na rede PPPI oriunda da união da rede dos genes induzidos com a dos genes reprimidos (Tabela 6 e Anexo D) foram identificadas e recuperadas usando o plugin MCODE associado ao Cytoscape e, juntamente com a rede PPPI fusionada, foram submetidas à análise da Ontologia Gênica (GO), a fim de obter informações sobre a natureza dos processos biológicos associados a cada cluster.

Tabela 6: Principais classes de ontologias gênicas observadas nos clusters derivados da rede PPPI fusionada

<i>Biological process</i>	<i>GO identification no.</i>	<i>P value^a</i>	<i>Corrected P value^b</i>	<i>k^c</i>	<i>f^d</i>
Translation	6416	2.07×10^{-45}	2.93×10^{-43}	45	499
Cellular protein metabolic process	44267	4.00×10^{-21}	9.02×10^{-20}	45	1702
Transcription from RNA polymerase II promoter	6366	3.49×10^{-11}	3.43×10^{-9}	9	162
Mitochondrial transport	6839	1.74×10^{-10}	3.88×10^{-9}	5	67
Histone modification	16570	1.14×10^{-9}	2.23×10^{-8}	7	92
Ribosome assembly	42255	1.82×10^{-9}	2.58×10^{-8}	9	65
Establishment and/or maintenance of chromatin architecture	6325	2.45×10^{-9}	4.37×10^{-8}	9	260
G1 phase of mitotic cell cycle	80	1.05×10^{-8}	1.28×10^{-7}	5	32
Transcription	6350	1.15×10^{-8}	1.33×10^{-7}	11	598
Regulation of RNA metabolic process	51252	1.53×10^{-8}	1.57×10^{-7}	11	614
Mitochondrion organization and biogenesis	7005	2.69×10^{-8}	2.91×10^{-7}	5	179
Chromosome organization and biogenesis	7001	1.04×10^{-7}	8.86×10^{-7}	9	398
DNA double-strand break processing	729	1.77×10^{-6}	5.67×10^{-5}	2	5
Negative regulation of RNA metabolic process	51253	2.49×10^{-6}	8.00×10^{-5}	4	157
Negative regulation of transcription	16481	3.12×10^{-6}	8.00×10^{-5}	4	166
Protein transport	15031	3.78×10^{-6}	2.11×10^{-5}	5	481
Heterochromatin formation	31507	3.74×10^{-5}	2.99×10^{-4}	3	92
Chromatin silencing	6342	3.74×10^{-5}	2.99×10^{-4}	3	92
pH Reduction	45851	4.88×10^{-5}	6.68×10^{-4}	2	24
Vacuolar acidification	7035	4.88×10^{-5}	6.68×10^{-4}	2	24
Gene silencing	16458	5.40×10^{-5}	3.84×10^{-4}	3	104
Regulation of pH	6885	8.21×10^{-5}	8.62×10^{-4}	2	31
Organic acid transport	15849	2.06×10^{-4}	1.41×10^{-2}	3	65
ER to Golgi vesicle-mediated transport	6888	6.12×10^{-4}	2.57×10^{-2}	2	84
Cell cycle	7049	9.16×10^{-4}	3.51×10^{-3}	3	566
Ethanol biosynthetic process during fermentation	43458	6.52×10^{-3}	1.76×10^{-2}	1	2
Regulation of cell cycle	74	7.11×10^{-3}	1.57×10^{-2}	2	160
Telomere maintenance via recombination	722	9.76×10^{-3}	2.53×10^{-2}	1	19
Response to stress	6950	1.26×10^{-2}	2.96×10^{-2}	6	632
Telomere maintenance via recombination	722	9.76×10^{-3}	2.53×10^{-2}	1	19

^a *P* values were calculated by the hypergeometric distribution of one ontology class visualized in the network.

^b Calculated values based on *P* values obtained after FDR was applied.

^c Total number of proteins found in the network which belong to a gene ontology.

^d Total number of proteins that belong to a specific gene ontology.

Para a análise, duas diferentes redes PPPI, usando dados do transcriptoma da levedura foram obtidas a partir do Saccharomyces Genome Database (SGD): uma associada aos genes reprimidos e outra associada aos genes induzidos. A rede PPPI associada aos genes induzidos, obtida do SGD, contém 1.226 nós e 2.854 conectores, enquanto que a rede PPPI associada aos genes reprimidos contém 1.412 nós e 2.782 conectores (Anexo E). Ambas as redes foram analisadas, a fim de verificar a sobreposição dos dados pelo Diagrama de Venn (Area-Proportional Venn Diagram) (Figura 16). A rede PPPI associada aos genes reprimidos contém 919 genes únicos, enquanto a rede PPPI associada aos genes induzidos contém 733 genes únicos; e existem 493 genes que se sobrepõem. Embora o grau de sobreposição da rede não seja elevado, decidiu-se unir as duas redes em um único gráfico, contendo 2158 nós e 5655 conectores (Anexo E).

Induced Network	Repressed network	Combined	Overlap	Not Overlap	Induced network Only	Repressed network Only
1226	1412	2145	493	1652	733	919

Figura 16: Análise, por Diagrama de Venn, do grau de sobreposição dos nós das redes PPPI para os genes induzidos e reprimidos na análise do microarray.

As análises GO da rede fusionada e dos seus clusters indicam a participação de importantes processos biológicos, tais como: Cluster 1: tradução, biogênese ribossomal e regulação negativa de processamento de RNAm; Cluster 2: transporte (ácido orgânico, colina, betaína, etanolamina); Cluster 3: transporte, organização e biogênese mitocondriais, e transporte de proteínas; Cluster 4: regulação da transcrição do promotor da RNA polimerase II, estabelecimento e/ou manutenção da

arquitetura da cromatina, regulamentação do processo metabólico do RNA, transcrição/organização e biogênese dos cromossomos, fase G1 do ciclo celular mitótico, modificação de histona; Cluster 5: processo catabólico do DNA, reparo da quebra da dupla fita, recombinação meiótica, fase M do ciclo celular meiótico, manutenção dos telômeros através de recombinação; Cluster 6: transporte mediado por vesícula do Retículo Endoplasmático para o Golgi; Cluster 7: regulação negativa da transcrição, regulação negativa do processo metabólico do RNA, formação da heterocromatina, silenciamento da cromatina, silenciamento gênico, estabelecimento e / ou manutenção da arquitetura da cromatina, regulação do ciclo celular; Cluster 8: processo biossintético do RNA, transcrição do promotor da RNA polimerase II; Cluster 9: regulação de pH, redução do pH, acidificação vacuolar; Cluster 10: modificação das histonas, transcrição dos promotores da RNA polimerase I, II e III, o estabelecimento e/ou manutenção da arquitetura da cromatina, organização cromossômica e biogênese, processo de biossíntese do etanol durante a fermentação, resposta ao estresse; Cluster 11: organização cromossômica e biogênese, reparo do DNA, ciclo celular mitótico, acetilação das histonas (Anexo D).

Com base em dados obtidos a partir desta análise inicial utilizando biologia de sistemas, decidimos obter mais informações sobre os principais nós envolvidos na informação da rede pesquisando pelas centralidades da rede. As centralidades de rede permitem-nos identificar os nós (e consequentemente os processos biológicos) que passam uma posição relevante na arquitetura geral da rede (BORGATTI, 2005). As centralidades foram recentemente aplicadas para compreender a estrutura e propriedades de complexos biológicos, redes tecnológicas e de infra-estrutura (ESTRADA, 2006; ESTRADA & HATANO, 2010). Muitos dos nós em uma dada rede que apresentam valores elevados de centralidade são pontos importantes de vulnerabilidade, indicando que qualquer ataque a esses nós poderia introduzir fortes perturbações na rede. O grau do nó ("node degree") representa a medida mais simples de centralidade em uma determinada rede, correspondendo ao número de nós adjacentes a um determinado nó, onde adjacente significa diretamente ligado (SCARDONI *et al*, 2009). O grau do nó representa a "popularidade" de um determinado nó, e nós altamente conectados em uma rede são denominados de "hubs". Uma outra medida é o "betweenness", que indica em qual extensão um nó específico está entre todos os outros nós dentro da rede (NEWMAN, 2005). De maneira geral, "betweenness" mostra a influência de um nó sobre a propagação de

informações através da rede. Por outro lado, “bottleneck” é um dado topológico local definido como todos os nós com valores altos de “betweenness” e diferentes graus de nós, indicando que eles são pontos centrais que controlam a comunicação entre os outros nós dentro da rede (GIRVAN & NEWMAN, 2002; YU *et al*, 2007). As medidas de “betweenness” e grau do nó nos permitem definir os nós “bottleneck”. Assim, nós “bottleneck” correspondem às proteínas altamente centrais, que conectam vários complexos ou são membros periféricos de complexos centrais, sendo pontos importantes de comunicação entre dois complexos (YU *et al*, 2007). A maioria dos nós “bottleneck” tende a ser proteínas essenciais em uma rede (YU *et al*, 2007).

A análise de centralidade da rede PPPI fusionada indicou a presença de 419 nós “bottleneck”; 99 desses nós “bottleneck” foram induzidos no array e 92 foram reprimidos (Anexo F planilha 1). A análise dos 11 clusters mostrou a presença de importantes nós “bottleneck” nos clusters (Anexo F planilha 2). A partir da análise GO para os nós “bottleneck”, foram observados nós com participação nos processos biológicos como processo metabólico, o transporte de proteína e transporte, organização de componentes celulares e biogênese, processo metabólico do DNA e transcrição, processamento de RNA, tradução, resposta ao estresse, ciclo celular (para genes induzidos no microarray); e processo metabólico, transporte de proteínas e transporte, processo metabólico do DNA e transcrição, processamento de RNA, tradução, transdução de sinal, ciclo celular (para genes reprimidos no microarray). Estes dados sublinham a importância destas vias metabólicas na resposta de *S. cerevisiae* a própolis.

Módulo 4.3: A identificação dos mecanismos de letalidade da própolis em *Candida albicans*

4.3. RESULTADOS E DISCUSSÃO

4.3.1. A própolis induz morte celular em *C. albicans*

Como verificado anteriormente no módulo 4.1 (Figura 2A), *C. albicans* foi mais suscetível a concentrações de própolis entre 0,50 e 0,75%. Este ensaio constituiu-se do primeiro de uma série de experimentos feitos para verificar a concentração de trabalho adequada de própolis, e também como a viabilidade celular era afetada por ela. Numa próxima etapa, foram feitas novas diluições seriadas de dez vezes de células de *C. albicans* para as duas cepas selvagens CAI4 e SC5314, nas concentrações de 0,250; 0,500; 0,750 e 1% de extrato alcoólico de própolis, utilizando como controle 5,45% de etanol (o que corresponde a 60% de etanol presente em 1% do extrato alcoólico de própolis). Conforme verificado na Figura 17 A, ambas as cepas de *C. albicans* foram mais sensíveis à concentração de 1%, sendo que a cepa CAI4 apresentou-se sutilmente mais sensível que a cepa SC5314.

Posteriormente, testamos também a sensibilidade à própolis de outras cepas de *C. albicans* (3153A e 529L) que são consideradas hipervirulentas no modelo de candidíase vaginal animal (YANO e FIDEL, 2011; RAHMAN *et al*, 2007). A cepa 3153A apresenta maior virulência no modelo de candidíase vaginal animal do que a cepa SC5314 (resultados não-apresentados). Com este ensaio para avaliar a sensibilidade das duas novas linhagens, pôde-se verificar, comparando-se com SC5314, que a cepa 3153A mostrou-se um pouco mais resistente até a concentração de 0,5% de própolis; contudo, acima desta concentração, ambas as cepas são sensíveis. Já para a cepa 529L, observou-se sensibilidade a partir da concentração de 0,125% (Figura 17B). Vale ressaltar que a cepa 529L apresenta um tamanho celular menor comparado às outras cepas, por isso, na diluição de 10^2 células do controle negativo, as mesmas não são muito visíveis na figura.

Figura 17: Ensaio de diluição para cepas de *C. albicans* em meio YPD complementado com diferentes concentrações de própolis. As placas foram incubadas a 30°C, por 72 horas. (A) CAI4 e SC5714 e (B) SC5314, 3153^a e 529L.

Além disso, assim como para *S. cerevisiae*, investigou-se a viabilidade de *C. albicans* em diferentes estágios da curva de crescimento. Para isto, as células da cepa SC5314 (1×10^7 células/mL⁻¹), nos estágios estabelecidos, foram expostas a própolis (0,125; 0,250; 0,500 e 1%) por 20 minutos, e determinou-se a viabilidade celular pelo plaqueamento de concentrações apropriadas das células e pela contagem do número de colônias em relação ao controle negativo, tratado com etanol (5,45%), o qual obteve 100% de sobrevivência em todas as fases da curva de crescimento. Pode-se observar que as células na fase estacionária (16 horas) são mais resistentes ao extrato de própolis em relação às outras fases da curva de crescimento (Figura 18).

Figura 18: Viabilidade de células de *C. albicans* em diferentes fases do ciclo de crescimento e expostas a própolis (0,125; 0,250; 0,500 e 1%), por 20 minutos. A viabilidade celular foi determinada pelo plaqueamento de concentrações apropriadas das células e pela contagem do número de colônias. Os números são referentes à média $\pm$ o desvio padrão de três experimentos independentes.

Posteriormente, outro ensaio de viabilidade foi feito, agora com tempos distintos de exposição à própolis, com um inóculo de SC5314 na fase estacionária (16 horas) (1×10^7 células/mL⁻¹) para apenas 1% de própolis. Verificou-se uma redução na porcentagem de sobrevivência conforme o aumento do período de incubação. Para 20 min de exposição a 1% de própolis, a redução na viabilidade foi de 89,8% (Figura 19).

Figura 19: Viabilidade de células de *C. albicans* em expostas a 1% de própolis, por 5, 10 e 20 minutos. A viabilidade celular foi determinada pelo plaqueamento de concentrações apropriadas das células e pela contagem do número de colônias. Os números são referentes à média $\pm$ o desvio padrão de três experimentos independentes. O tratamento estatístico foi realizado pelo método ANOVA one way, com pós teste Dunnet e $p < 0,01$.

Numa próxima etapa, investigou-se a morte celular em células observando-se o número de células que são marcadas por PI (indicação preliminar de necrose). Para este experimento 1×10^6 células/mL foram tratadas com 0,125% de extrato alcoólico de própolis a 11% ou 0,68% de etanol, a 30 °C por 24, 48 e 72 horas. Após estes períodos de tratamento, verificou-se a viabilidade das células e também a morfologia celular no microscópio. Além disso, adicionou-se o corante PI, a fim de se verificar a ocorrência de necrose. O resultado pode ser observado na Figura 20, onde se sugere a ocorrência de morte celular por necrose, uma vez que a porcentagem de células marcadas com PI apresenta significativa correspondência com a porcentagem de morte das células verificada no ensaio de viabilidade. Ainda vale ressaltar que a morfologia celular das células tratadas com própolis encontra-se alterada para uma significativa porcentagem de células, como pode ser observado nas fotos do DIC. Tal anormalidade morfológica está sendo estudada em nosso laboratório.

Foi escolhida esta concentração de própolis (0,125%), pois não turva muito o meio de cultura e não torna a remoção da própolis extremamente difícil como as outras duas concentrações (0,5% e 1%), facilitando assim a microscopia. Mesmo assim, adicionamos um ponto de 24 horas com 1% de própolis, contudo, a presença de vestígios de própolis é bastante intensa, como pode ser verificada na microscopia correspondente da figura 20. Nela, as setas amarelas apontam a célula de *C. albicans*.

Figura 20: A viabilidade de células de *C. albicans* em expostas a própolis. A viabilidade celular foi determinada pelo plaqueamento de concentrações apropriadas das células e pela contagem do número de colônias. Para o controle considerou-se sobrevivência de 100%; assim, a porcentagem de sobrevivência dos tratamentos foi calculada em relação ao controle. Os números são referentes à média $\pm$ o desvio padrão de três experimentos independentes. Além disso, verificou-se que a própolis induz necrose em *C. albicans*. Ensaio de PI para células de SC5314 expostas a própolis. Os números são referentes à média $\pm$ o desvio padrão de três experimentos independentes. Em cada experimento, 100 células foram avaliadas para a coloração de PI. Barras: 5 μ m.

A microscopia utilizando o TUNEL, a fim de verificar possível morte por apoptose, também foi feita para as mesmas condições do ensaio de necrose (24, 48 e 72 horas de tratamento com 0,125% de própolis), e também por períodos menores como 20 min, 1h e 2h nas concentrações de 0,125% e 0,5%. Contudo, não se observou marcação com este reagente para as células tratadas com própolis (dados não mostrados).

4.3.2. O envolvimento das caspases e da via metabólica Ras na morte celular induzida por própolis em *C. albicans*

Como sugerido no módulo 4.1 (Figura 5), o gene *YCA1*, o qual codifica uma proteína tipo caspase, é importante para a morte celular mediada por própolis em *S. cerevisiae*. Embora não existam relatos de metacaspases envolvidas na apoptose para *C. albicans*, tem sido demonstrado que o gene *IPF4847* (orf19.5995), que é um gene de *C. albicans* com homologia estrutural para a metacaspase *YCA1* de *S. cerevisiae*, quando deletado, resulta na resistência ao estresse oxidativo, sugerindo que este gene seja uma metacaspase responsável pela morte celular programada induzida por estresse oxidativo neste fungo (CAO, *et al* 2009; AERTS *et al*, 2009). Desta maneira, utilizando-se as linhagens de *C. albicans*: BWP17 (*wt*), L1374-BWP17 *mca1* Δ/Δ (*mca1* Δ ::*FRT*/*mca1* Δ ::*FRT*) e L1428-BWP17 *mca1* Δ/Δ -*MCA1* (*mca1* Δ ::*FRT*/*mca1* Δ ::*MCA1*-*SAT1*-*FLIP*) (AERTS *et al*, 2009), foi verificada a viabilidade à própolis. Quando as células de *C. albicans*, cultivadas por 16 horas, foram expostas a 1% de própolis por 5, 10 e 20 minutos, houve uma redução na viabilidade celular de 30, 70 e 94%, respectivamente, na linhagem selvagem BWP17, enquanto que nos mutantes, esta redução foi de cerca de 36, 38 e 72% para o heterozigoto (L1428) e de 45, 46 e 90% para o homozigoto (L1374) (Figura 21). Estes resultados indicam que o gene *IPF4847* é importante para a morte celular mediada por própolis.

Figura 21: Os mutantes para o gene *IPF4847* são mais tolerantes a própolis que a linhagem selvagem correspondente. A viabilidade das células de *C. albicans* para a cepa selvagem e para os mutantes, crescidas por 16 horas e expostas a 1% de própolis (* $p < 0.05$), foi determinada pelo plaqueamento de concentrações apropriadas das células e pela contagem do número de colônias. Os resultados são referentes à média $\pm$ o desvio padrão de três experimentos independentes. O tratamento estatístico foi realizado pelo método ANOVA one way, com pós teste Newman Keuls.

Além disso, segundo PHILLIPS *et al*, (2006), a resposta apoptótica em *C. albicans* pode ser desencadeada quando há a exposição a H_2O_2 , acidificação intracelular, ou baixas doses do antifúngico anfotericina B. Entretanto, os mecanismos pelos quais *C. albicans* sofre apoptose ainda não são claros. A via metabólica Ras parece estar envolvida na sinalização apoptótica neste fungo induzida por ácido acético e peróxido de hidrogênio. A ativação artificial de Ras acelera a taxa na qual as células se tornam apoptóticas, mas a ativação de Ras por si só não é suficiente para matar as células. Mutantes que possuem a sinalização Ras-cAMP-PKA (*ras1Δ*, *cdc35Δ*, *tpk1Δ* e *tpk2Δ*) bloqueada, suprimem ou atrasam a resposta apoptótica, enquanto mutações que estimulam a sinalização *RAS^{val13}* e *pde2Δ*, apresentam a taxa de entrada em apoptose acelerada. Sendo assim, foi verificada a viabilidade frente ao extrato alcoólico de própolis para os seguintes mutantes de *C. albicans* *cdc35Δ*, *ras1Δ*, *tpk1Δ*, *tpk2Δ* e *pde2Δ* (obtidas do laboratório do Dr. Mark Ramsdale) em comparação com a linhagem selvagem correspondente. Pôde-se observar que, com exceção dos mutantes *cdc35Δ* e *ras1Δ*, os quais apresentaram certa redução na morte celular em comparação com o selvagem, os mutantes para os genes *tpk1Δ* e *tpk2Δ* apresentaram maior sensibilidade à própolis. Tal fato confirma a hipótese de que a perturbação da sinalização de Ras inibe a taxa de entrada em apoptose, mas não evita que isso aconteça se o estímulo for suficientemente forte (PHILLIPS *et al*, 2006). Além disso, para o mutante *pde2Δ*, a porcentagem de morte celular foi maior em comparação com o selvagem. (Figura 22).

Figura 22: Os mutantes para os *cdc35Δ*, *ras1Δ*, são mais tolerantes a própolis que a linhagem selvagem correspondente e também que o mutante *pde2Δ*. A viabilidade das células de *C. albicans* para a cepa selvagem e para os mutantes, crescidas por 16 horas e expostas a 1% de própolis (* $p < 0.05$), foi determinada pelo plaqueamento de concentrações apropriadas das células e pela contagem do número de colônias. Os resultados são referentes à média $\pm$ o desvio padrão de três experimentos independentes. O tratamento estatístico foi realizado pelo método ANOVA one way, com pós teste Newman Keuls.

4.3.3. A avaliação de bibliotecas de deleções de *C. albicans* para sensibilidade à própolis

Para tentar elucidar algumas funções de genes que possam ser necessárias para a sensibilidade à própolis do fungo patogênico *C. albicans*, 15 bibliotecas de deleção de *C. albicans* fornecidas pelo Fungal Genetics Stock Center (Mc CLUSKEY & PLAMANN, 2010) [(CJN Transcription Factor (TF) Homozygous mutants; 2nd set TF Homozygous mutants 1; 2nd set TF Homozygous mutants 2; VB GKO Plate 1; VB GKO Plate 2; VB GKO Plate 3; VB GKO Plate 4; VB GKO Plate 5; Dominique Sanglard TF mutants; SFY TF Homozygous mutants plate 1; Cell Wall Protein Homozygous mutants plate 1; Cell Wall Protein Homozygous mutants plate 2; Miscellaneous Homozygous mutants plate 1; Kinase from A. Mitchell plate K1 e plate K2) (<http://www.fgsc.net/candida/FGSCcandidaresources.htm>)] foram escaneadas em placas YPD contendo 5,45% etanol (como um controle), que é a concentração de etanol presente em 1% do extrato alcoólico de própolis ou própolis (0,500 e 1%), durante 5 dias a 30°C.

Cerca de 800 mutantes diferentes foram envolvidos nesta triagem, e depois de três “screenings” para confirmação, 51 foram identificados apresentando maior sensibilidade a própolis quando comparadas com as cepas do tipo selvagem correspondente. Tais mutantes provenientes das bibliotecas são homozigotos para os genes deletados. A partir de uma análise feita no banco de dados *kinase.com*, com o intuito de categorizar as proteínas quinases encontradas no screening, foram encontrados quatro grandes grupos de proteínas quinases: STE (homólogas em levedura das quinases *sterile 7*, *sterile 11*, *sterile 20*); CAMK (proteína quinase dependente de cálcio/calmodulina); AGC (que contém as famílias relacionadas à PKA, PKG e PKC) e CMGC (que contém as famílias CDK, MAPK, GSK3, CLK) (www.cellsignal.com/reference/kinase). Além disso, 9 dos mutantes sensíveis (*VPS34*, *HOG1*, *SUV3*, *MDS3*, *PTP3*, *YHB1*, *CDC10*, *ACE2*, *FKH2*) a própolis apresentam virulência em camundongos reduzida ou ausente em diferentes modelos de virulência (Anexo G). As proteínas quinases desempenham papéis importantes na sinalização e resposta a alterações no ambiente externo; e a capacidade de *C. albicans* rapidamente detectar e responder a mudanças em seu ambiente é fundamental para sua sobrevivência no hospedeiro humano (BLANKENSHIP *et al.*, 2010).

Posteriormente, foi iniciada a pesquisa por proteínas potenciais e/ou mecanismos e seus respectivos processos biológicos que são afetados pela exposição à própolis. Para este objetivo, foram utilizadas duas abordagens: inicialmente, analisamos os mutantes de *C. albicans* através do banco de dados de *Candida* (*Candida* Genome Database), a fim de identificar possíveis processos biológicos de cada gene, assim como seus ortólogos em *S. cerevisiae* (Anexo G). Os mutantes para sensibilidade a própolis estão relacionados principalmente a genes envolvidos no reparo da quebra de dupla fita, remodelagem da cromatina, na fusão nuclear durante “mating”, na regulação positiva da transcrição, na virulência em infecção sistêmica em camundongos, na regulação da morfogênese, nas vias metabólicas de transdução de sinal para o crescimento e formação de hifas, na transição de pseudohifa para hifa, na formação de clamidoconídios e na citocinese, na formação do anel de septina dentro do tubo germinativo, na biogênese da parede celular, em resposta a drogas, na sensibilidade a caspofungina e peróxido de hidrogênio, no transporte vesicular normal, como componente da via metabólica de sinalização TOR [(“target of rapamycin” - quinase essencial conservada ao longo da evolução eucariótica, com papel fundamental na morfogênese (ZACCHI *et al*, 2010)], no transporte do complexo de Golgi para vacúolo, separação celular, aderência, formação de biofilme, e resposta a estresses.

Num segundo momento, levando-se em conta a riqueza de dados para o organismo modelo *Saccharomyces cerevisiae*, que serve como uma forte base de comparação, decidiu-se a partir dos ortólogos encontrados em *S. cerevisiae*, construir uma rede PPPI na levedura utilizando dados de deleção obtidos do banco de dados STRING. Proteínas compartilhadas e sub-redes presentes na rede principal PPPI foram identificadas usando o *Cytoscape-associated plugin MCODE* e submetidas a uma análise de *Gene Ontology* (GO), a fim de obter informações sobre a natureza e o número de sub-redes pertencentes à rede e seus respectivos processos biológicos associados. Os resultados obtidos das análises MCODE e GO mostraram que a rede final contém 309 nós e 3191 conectores e é composta por oito *clusters*. As análises GO revelaram que os *clusters* 1, 2, 3 e 4 possuem processos biológicos em comum, os quais podem ser classificados em: processo catabólico de proteína, regulação do processo metabólico de RNA, transporte e localização, transporte organização e biogênese

nuclear, regulação da transcrição do promotor da RNA polimerase I, II e III, regulação do ciclo celular, regulação da transcrição, estabelecimento e/ou manutenção da arquitetura da cromatina, modificação e remodelamento da cromatina, organização e biogênese, reparo do DNA, resposta a estresse, modificação de histona, processo metabólico do DNA. Já os clusters 5, 6 e 7 se sobressaem com os seguintes processos biológicos: crescimento filamentosos e de pseudohifa, transporte vesicular ao longo de filamentos de actina, resposta a feromônio, estabelecimento da orientação e localização do fuso mitótico, ativação da cascata MAPK (Mitogen-Activated Protein Kinase), transdução de sinal durante conjugação com fissão nuclear, regulação da conjugação, organização e biogênese do citoesqueleto, morfogênese celular, brotamento celular, organização e montagem do anel de septina, processo metabólico da parede celular. No cluster 8, estão em evidência os processos: Transporte vacuolar, resposta estímulos químicos, transporte do complexo de Golgi para o vacúolo (Anexo H). Vários genes observados em nosso “screening”, tais como *kis1*, *hst7*, *gin4*, *ptp3*, entre outros, estão envolvidos na transição dimórfica em *C. albicans* (BLANKENSHIP *et al.*, 2010; SHAPIRO *et al.*, 2011). Levando-se em consideração que no patógeno oportunista *C. albicans* a transição morfológica levedura-hifa desempenha uma papel crítico na capacidade deste fungo colonizar como comensal e causar doença como um patógeno (ZACCHI *et al.*, 2010), e que na análise do screening das bibliotecas, este processo de transição foi bastante evidenciado, decidiu-se verificar se a própolis exerce algum papel na inibição do processo de filamentação. Assim, na próxima subseção, foi realizado um primeiro experimento com o intuito de avaliar a sensibilidade de *C. albicans* em relação à própolis nas suas 3 formas morfológicas: levedura, pseudohifa e hifa, para numa segunda etapa estudar sua atividade na transição dimórfica.

4.3.4. O papel da própolis na transição dimórfica de *C. albicans*

4.3.4.1. A própolis inibe os três morfotipos de *C. albicans*

Inicialmente, com o intuito de avaliar a sensibilidade de *C. albicans* em relação à própolis nas suas 3 formas morfológicas, foi feito um inóculo, em meio YPD líquido com agitação, da cepa SC5314 por 16h a 30°C. Após este período,

as células foram contadas e, 2×10^7 células/ ml foram incubadas por 4 horas, também com agitação, em meio YPD líquido a 30°C (para manter a forma de levedura), em YPD líquido a 37°C (para induzir a forma de pseudohifa), e em meio YPD líquido acrescido Soro Fetal Bovino (SFB) (20%) a 37°C. Posteriormente, para o controle, foi adicionado 5,45% de etanol; em relação à própolis, utilizou-se a concentração de 1% por 20 min e 1 hora. A fim de verificar a eficiência da transição, retirou-se uma alíquota da cultura para verificar no microscópio, observando que a transição para hifa foi em torno de 95%, enquanto que para pseudohifa, 60% (Figura 23). Após os tratamentos, foram feitas diluições seriadas de dez vezes e gotejadas em placa contendo YPD sólido. Neste ensaio, observou-se que a própolis inibe o crescimento da levedura, pseudohifa e hifa em *C. albicans* (Figura 23).

Figura 23: Ensaio de diluição para as 3 formas morfológicas de *C. albicans*. Após a indução das formas de pseudohifa e hifa, as células foram tratadas com 1% de própolis por 20 min e 1 hora. Diluições seriadas de dez vezes foram feitas e gotejadas em placas contendo YPD sólido. As placas foram incubadas a 30°C, por 72 horas. A fim de verificar a eficiência da transição, retirou-se uma alíquota da cultura para verificar no microscópio. Barras de 5 μ m.

4.3.4.2. A própolis inibe a transição dimórfica de levedura para hifa em *C. albicans*

Para analisar o efeito do extrato alcoólico de própolis na filamentação de *C. albicans*, a linhagem SC5314 foi crescida por 16 horas em YPD líquido, a 30 °C, com agitação. As células foram lavadas com água estéril para remoção do meio. Em torno de 2×10^7 cells/mL foram adicionadas a 1mL meio RPMI1640 (para induzir a filamentação) contendo 0,05; 0,1 e 0,2% de própolis. As amostras foram incubadas por 90 min a 37°C sob agitação (100 rpm). Após este período, os tubos foram centrifugados para remover o sobrenadante e as células lavadas com PBS. A viabilidade das células foi checada pelo plaqueamento de diluições seriadas em placas de YPD sólido e, não houve indução significativa de morte celular para as concentrações utilizadas. Para verificar a influência na filamentação, as células foram visualizadas no microscópio e, com o intuito de quantificar as formas filamentosas, pelo menos 100 células por amostra foram contadas e a porcentagem de filamentação foi calculada através da contagem do número de tubos germinativos pelo número total de células contadas (Figura 24).

Figura 24: Ensaio para verificar a influência na filamentação de *C. albicans*. A filamentação da linhagem SC5314 foi induzida em meio RPMI1640, concomitantemente com o tratamento com a própolis. A influência na filamentação foi avaliada analisando-se as células no microscópio e, com o intuito de quantificar as formas filamentosas, pelo menos 100 células por amostra foram contadas e a porcentagem de filamentação foi calculada através da contagem do número de células filamentosas pelo número total de células contadas.

Com este ensaio pôde-se verificar que a própolis afeta a filamentação em *C. albicans*. Desta maneira, o uso da própolis para o tratamento clínico da candidíase

pode ter grande aplicação, principalmente por afetar um mecanismo importante para a patogenicidade deste fungo. Além disso, observou-se, por microscopia de fluorescência utilizando o filtro GFP (verde), que todas as células de *C. albicans* tratadas com própolis possuíam fluorescência verde distribuída por toda a célula e também agrupada em formatos característicos de mitocôndrias. Sendo assim, subsequentemente à quantificação das formas filamentosas, decidiu-se corar as mitocôndrias com o corante Mitotracker® e visualizadas no microscópio de fluorescência (Figura 25).

Com esta microscopia, pôde-se realmente comprovar que compostos fluorescentes da própolis, como por exemplo alguns compostos fenólicos, estão distribuídos por toda o citoplasma e também nas mitocôndrias. Esta microscopia foi repetida crescendo as células de *C. albicans* em meio YPD (sem indução da filamentação), e tratando-as com 0,1 e 0,2% de própolis e, esta fluorescência verde foi verificada em toda a célula. Porém, tal fato não ocorre com as células de *S. cerevisiae*, tratadas com as mesmas concentrações de própolis, uma vez que possivelmente *C. albicans* seja mais permeável que *S. cerevisiae* a estes compostos presentes na própolis ou existam outros compostos em *C. albicans* que quando reagem com a própolis fluorescem. Além disso, segundo WALKER *et al.* (2008), a parede celular de *C. albicans* é altamente conectada a um número muito maior de vias de sinalização do que encontrado para *S. cerevisiae*. Logo, a rede de sinalização da parede celular desse fungo comensal/patogênico pode refletir um balanço de diversas entradas, que são preparadas para promover a modificação e suporte de adaptação, o que pode também ser um dos motivos para a maior permeabilidade a propolis nas células de *C. albicans* em comparação com *S. cerevisiae*.

Vale ainda ressaltar que os mecanismos e vias regulatórias que governam a dinâmica da parede celular em *C. albicans* são uma área crítica de compreensão, fazendo da parede celular um atraente alvo terapêutico, uma vez que a parede celular está ligada a vários processos biológicos conhecidos em *C. albicans*, incluindo o crescimento, a morfogênese, o “mating”, e a patogenicidade (CHAUHAN *et al.*, 2002).

Figura 25: A própolis afeta a filamentação em *C. albicans* e tais células tratadas com própolis apresentam fluorescência distribuída pelo citoplasma e mitocôndrias. A filamentação da linhagem SC5314 foi induzida em meio RPMI1640, concomitantemente com o tratamento com a própolis. A influência na filamentação foi avaliada analisando-se as células no microscópio. As células foram coradas com o corante de mitocôndrias Mitotracker® e também com o corante de núcleo DAPI. Barras: 10 μ m.

5. Conclusões

A própolis é um produto complexo derivado de resinas vegetais e saliva de abelha. Há diversos compostos químicos presentes neste produto natural que podem potencialmente, ser responsáveis por suas propriedades antibióticas. No entanto, levando em consideração o fato de que os efeitos de morte celular de própolis poderiam ser devido a uma grande combinação de compostos químicos e concentrações, decidimos investigar os efeitos de morte celular de própolis, concentrando nossos experimentos no extrato alcoólico de própolis. Em resumo, nossos dados indicam que a própolis afeta várias vias metabólicas na célula eucariótica. Contudo, as vias mais importantes, em *S. cerevisiae*, estão relacionadas ao estresse oxidativo, a cadeia de transporte de elétrons mitocondrial, a acidificação vacuolar, regulação da macroautofagia associadas com proteína alvo para vacúolo, a resposta à privação de nutrientes, a regulação negativa da transcrição do promotor da RNA polimerase II. Em *C. albicans*, na transição dimórfica e na inibição dos três morfotipos (levedura, hifa e pseudohifa), além de se observar morte celular por necrose e também do gene homólogo da metacaspase YCA1 em *S. cerevisiae*. Assim, este trabalho enfatiza a importância de *S. cerevisiae* como um sistema modelo para compreender a nível molecular o mecanismo de ação de um composto natural. Além disso, é um dos primeiros a investigar sistematicamente, pelo uso da genômica funcional, como a própolis influencia na modulação da abundância de RNAm de um organismo. Isto fornece novas possibilidades para a compreensão e validação da própolis como agente terapêutico alternativo contra infecções fúngicas, uma vez que pode-se obter um medicamento que se torne uma alternativa terapêutica ainda inexplorada, com grande potencial terapêutico e que, talvez, considerando seu mecanismo de ação múltiplo, possa ser mais difícil de causar resistência microbiana.

6. Referências Bibliográficas

- AGUERO, M.B.; GONZALEZ, M.; LIMA, B.; SVETAZ, L.; SÁNCHEZ, M.; ZACCHINO, S.; FERESIN, G.E.; SCHMEDA-HIRSCHMANN, G.; PALERMO, J.; WUNDERLIN, D. AND TAPIA, A. Argentinean propolis from *Zuccagnia punctata* Cav. (Caesalpinieae) exudates: phytochemical characterization and antifungal activity. **J. Agric. Food Chem.** 58: 194-201, 2010.
- ALBERTS, B.; JOHNSON A.; LEWIS, J.; RAFF, M.; ROBERTS, K.; WALTER P.; In: **The Molecular Biology of the Cell.**, 5th edition, 1268p, Garland Science, 2007.
- ARIÑO, J.; RAMOS, J. AND SYCHROVÁ, H. Alkali metal cation transport and homeostasis in yeasts. **Microbiol. Mol. Biol. Rev.** 74: 95-120, 2010.
- ATTARDI, G.M., AND CHOMYN, A. Mitochondrial Biogenesis and Genetics. **Methods in Enzymology** 260: 406-417. San Diego: Academic Press, 1995.
- BADER, G.D., AND HOGUE, C.W. An automated method for finding molecular complexes in large protein interaction networks. **BMC Bioinformatics** 4: 2, 2003.
- BANKOVA, V. Chemical diversity of propolis and the problem of standardization. **Journal of Ethnopharmacology**, v.100, n.1/2, p.114-117, 2005.
- BANKOVA, V.; CHRISTOV, R.; KUJUMGIEV, A.; MARCUCCI, M.C.; POPOV, S. Chemical composition and antibacterial activity of Brazilian propolis. **Z Nat Sect C Biosci.**, v. 50, p.167–72, 1995.
- BANKOVA, V.S.; CASTRO, S.L. DE; MARCUCI, M.C. Propolis: recent advances in chemistry and plant origin. **Apidologie**, v.31, p.3-15, 2000.
- BARNETT, J. A.; PAYNE, R. W. e YARROW, D. Yeasts: characteristics and identification. 3ed. Cambridge, U.K.; New York, NY, USA: **Cambridge University Press**, ix 1139 p. p., 2000.
- BASNET P, MATSUNO T, NEIDLEIN RZ: Potent free radical scavenging activity of propol isolated from Brazilian propolis. **Z. Naturforsch.** C 52:828–833, 1997.
- BENJAMINI, Y., AND HOCHBERG, Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing. **J. R. Stat. Soc. B** 57: 289-300, 1995.
- BERRETTA-SILVA, A.A., MEDA, A.C., FERREIRA, M.E.T. – PI 0405483-0. Processo de obtenção do extrato padronizado de própolis, extrato assim obtido, suas formulações, produtos e usos. Revista de Propriedade Industrial n. 1778m de 01/02/2005.

- BERRETTA, A. A.; NASCIMENTO, A. P.; BUENO, P. C. P.; VAZ M. M. O. L. L.; MARCHETTI J. M. Propolis Standardized Extract (EPP-AF®), an Innovative Chemically and Biologically Reproducible Pharmaceutical Compound for Treating Wounds **International Journal of Biological Sciences**, 8(4):512-521, 2012.
- BARROS M.H, BANDY B, TAHARA E.B, KOWALTOWSKI A.J. Higher respiratory activity decreases mitochondrial reactive oxygen release and increases life span in *Saccharomyces cerevisiae*. **J Biol Chem**. v.26;279(48):49883-8, 2004.
- BEVIS, B.J., AND GLICK, B.S. Rapidly maturing variants of the *Discosoma* red fluorescent protein (DsRed). **Nature Biotech** 20: 83-87, 2002.
- BHARUCHA, N. AND KUMAR, A. Yeast Genomics and Drug Target Identification. **Combinatorial Chemistry & High Throughput Screening** 10: 618-634, 2007.
- BLACKBURN, A.S. AND AVERY, S.V. Genome-wide screening of *Saccharomyces cerevisiae* to identify genes required for antibiotic insusceptibility of eukaryotes. *Antimicrobial Agents and Chemotherapy* **47**: 676-681, 2003.
- BLANKENSHIP JR, FANNING S, HAMAKER JJ, MITCHELL AP (2010) An Extensive Circuitry for Cell Wall Regulation in *Candida albicans*. **PLoS Pathog** 6(2):e1000752.
- BOONE, C., H. BUSSEY, AND ANDREWS, B.J. Exploring genetic interactions and networks with yeast. **Nature Reviews Genetics** 8: 437-449, 2007.
- BORGATTI SP: Centrality and network flow. **Soc. Networks** 2005, 27: 55-71.
- BORUTAITE, V. Mitochondria as Decision-Makers in Cell Death. **Environ Mol Mutagen.**, v.51, p.406-416, 2010.
- BRACHMANN, C.B.; DAVIES, A; COST, G.J.; CAPUTO, E.; LI, J.; HIETER, P.; AND BOEKE, J.D. Designer deletion strains derived from *Saccharomyces cerevisiae* S288C: a useful set of strains and plasmids for PCR-mediated gene disruption and other applications. **Yeast** 14:115-132, 1998.
- BRADFORD, M. M. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. **Anal. Biochem.**, v. 72, p.248-254, 1976.

- BREGER, J., B.B. FUCHS, G. APERIS, T.I. MOY, F.M. AUSUBEL, AND E. MYLONAKIS. Antifungal chemical compounds identified using a *C. elegans* pathogenicity assay. **PLoS Pathogens**, 3: e18, 2007.
- BROACH, J.R. PRINGLE, J.R. AND JONES, E.W. The Molecular and Cellular Biology of the Yeast *Saccharomyces*. **Cold Spring Harbor Laboratory Press**, Cold Spring Harbor, New York, 1991.
- BRÖKER, L. E.; KRUYT, F. A. E., GIACCONE, G. Cell Death Independent of Caspases: a Review. **Clin Cancer Res.**, v.11, n.9, p.3155-3162, 2005.
- BÜTTNER, S., T. EISENERG, D. CARMONA-GUTIERREZ, D. RULI, H. KNAUER, C. RUCKENSTUHL, C. SIGRIST, S. WISSING, M. KOLLROSER, K-U. FRÖLICH, S. SIGRIST, AND F. MADEO. Endonuclease G regulates budding yeast life and death. **Mol. Cell.**, v.25: p.233-246, 2007.
- BURHANS, W.; M. WEINBERGER. Yeast endonuclease G: complex matters of death, and of life. **Mol. Cell.**, v.25, p.323-325, 2007.
- CAO YH, WANG Y, YUAN Q. Analysis of flavonoids and phenolic acid in propolis by capillary electrophoresis. **Chromatographia** 59: 135-140, 2004.
- CAO, F., *et al.* The Flo8 transcription factor is essential for hyphal development and virulence in *Candida albicans*. **Mol. Biol. Cell** 17:295–307, 2006.
- CAO, Y.Y.; HUANG, S.; DAI, B., ZHU, Z.; LU, H.; DONG, L.; WANG, Y.; GAO, P.; CHA, Y.; JIANG, Y. *Candida albicans* cells lacking CaMCA1-encoded metacaspase show resistance to oxidative stress-induced death and change in energy metabolism Fungal. **Genetics and Biology**, v. 46, p. 183–189, 2009.
- CAPASSO F, CASTALDO S. Propolis, an old remedy used in modern medicine. **Fitoterapia**. 73: S1-6, 2002.
- CARMONA-GUTIERREZ, D., T. EISENBERG, S. BÜTTNER, C. MEISINGER, G. KROEMER, AND F. MADEO. Apoptosis in yeast: triggers, pathways, subroutines. **Cell Death and Differentiation** 17: 763-773, 2010.
- CARNEIRO, P.; DUARTE, M; VIDEIRA, A. Characterization of apoptosis-related oxidoreductases from *Neurospora crassa*. **PLoS One.**, 7(3):e34270, 2012.

- CHAN, T-F.; CARVALHO, J.; RILES, L. AND ZHENG, X.F.S. A chemical genomics approach toward understanding the global functions of the target of rapamycin protein (TOR). **Proc. Natl. Acad. Sci. USA** 97: 13227–13232, 2000.
- CHAUHAN N, LI D, SINGH P, CALDERONE R, KRUPPA M (2002) The cell wall of CHEN, C.; WU, C.; LIN, J. Propolin C from propolis induces apoptosis through activating caspases, Bid and cytochrome c release in human melanoma cells. **Biochemical Pharmacology**, v.67, p. 53–66, 2004.
- CHEN, C. N., WU, C. L. LIN, J. K. Propolin C from propolis induces apoptosis through activating caspases, Bid and cytochrome c release in human melanoma cells. **Biochemical Pharmacology**, v. 67, p. 53-66, 2004.
- CHENG, W. C.; LEACH, K. M.; HARDWICK, J. M. Mitochondrial death pathways in yeast and mammalian cells. **Bioch. et Bioph. Acta.**, v.1783, p.1272–1279, 2008.
- COLABARDINI, A. C.; DE CASTRO, P. A.; GOUVÊA, P. F.; SAVOLDI, M.; MALAVAZI, I.; GOLDMAN, M. H. S.; GOLDMAN, G. H. Involvement of the *Aspergillus nidulans* protein kinase C with farnesol tolerance is related to the unfolded protein response. **Mol Microbiol.**, v.78, n.5, p.1259-79, 2010.
- COSTANZO, M., A. BARYSHNIKOVA, J. BELLAY, Y. KIM, E.D. SPEAR, C.S. SEVIER, H. DING, J. L. KOH, K. TOUFIGHI, S. MOSTAFAVI, J. PRINZ, R.P. ST ONGE, B. VANDERSLUIS, T. MAKHNEVYCH, F.J. VIZEACOMAR, S. ALIZADEH, S. BAHR, R.L. BROST, Y. CHEN, M. COKOL, R. DESHPANDE, Z. LI, Z.Y. LIN, W. LIANG, M. MARBACK, J. PAW, B.J. SAN LUIS, E. SHUTERIQUI, A.H. TONG, N. VAN DYK, I.M. WALLACE, J.A. WHITNEY, M.T. WEIRAUCH, G. ZHONG, H. ZHU, W.A. HOURS, M. BRUDNO, S. RAGIBIZADEH, B. PAPP, C. PÁL, F.P. ROTH, G. GIAEVER, C. NISLOW, O.G. TROYANSKAYA, H. BUSSEY, G.D. BADER, A.C. GINGRAS, Q.D. MORRIS, P.M. KIM, C.A. KAISER, C.L. MYERS, B.J. ANDREWS, AND C. BOONE. The genetic landscape of a cell. **Science** 327: 425-431, 2010.
- COSTE, A. T., M. KARABABA, F. ISCHER, J. BILLE, AND D. SANGLARD. TAC1, transcriptional activator of CDR genes, is a new transcription factor involved in the regulation of *Candida albicans* ABC transporters CDR1 and CDR2. **Eukaryot. Cell**, 3:1639–1652, 2004.
- CRAMPIN H, FINLEY K, GERAMID-NEJAD M, COURT H, GALE C, BERMAN J ET AL. *Candida albicans* hyphae have a Spitzenkörper that is distinct from the polarisome found in yeast and pseudohyphae. **J Cell Sci.**, 118(13): 2935-2947, 2005.

- DA SILVA J.F.M., SOUZA M.C., MATTA S.R, ANDRADE M.R, VIDAL F.V.N. Correlation analysis between phenolic levels of Brazilian propolis extracts and their antimicrobial and antioxidant activities. **Food Chem** 99: 431-435, 2006.
- DALKEN-DOTA, K.F.; FARIA, M.G.; BRUSCHI, M.L.; PELLOSO, S.M.; LOPES-CONSOLARO, M.E. AND SVIDZINSKI, T.I. Antifungal activity of propolis extract against yeasts isolated from vaginal exudates. **J. Altern. Complement. Med.** 16: 285-290, 2010.
- DANIAL ,N.N.; KORSMEYER ,S.J. Cell death: critical control points. **Cell.**, v. 2, p. 205-19, 2004.
- DAVIS DA, BRUNO VM, LOZA L, FILLER SG, MITCHELL AP. *Candida albicans* Mds3p, a conserved regulator of pH responses and virulence identified through insertional mutagenesis. **Genetics**, v. 162, p. 1573–1581, 2002.
- DAVIS, D., J. E. EDWARDS, JR., A. P. MITCHELL, AND A. S. IBRAHIM. *Candida albicans* RIM101 pH response pathway is required for host-pathogen interactions. **Infect. Immun.**, 68:5953–5959, 2000.
- DEGTEREV, A.; BOYCE, M.; YUAN, J. A decade of caspases., **Oncogene**, v.22, p.8543–8567, 2003.
- DESMOUCELLES, C., B. PINSON, C. SAINT-MARC, AND B. DAIGNAN-FORNIER. Screening the Yeast “Disruptome” for Mutants Affecting Resistance to the Immunosuppressive Drug, Mycophenolic Acid. **J. Biol.Chem.** 277: 27036–27044, 2002.
- EISENBERG T, CARMONA-GUTIERREZ D, BÜTTNER S, TAVERNARAKIS N, MADEO F: Necrosis in yeast. **Apoptosis** 2010, 15: 257-268.
- EISENBERG, T., H. KNAUER, A. SCHAUER, S. BÜTTNER, C. RUCKENSTUHL, D. CARMONA-GUTIERREZ, J. RING, S. SCHROEDER, C. MAGNES, L. ANTONACCI, H. FUSSI, L. DESZCZ, R. HARTL, E. SCHRAML, A. CRIOLLO, E. MEGALOU, D. WEISKOPF, P. LAUN, G. HEEREN, M. BREITENBACH, B. GRUBECK-LOEBENSTEIN, E. HERKER, B. FAHRENKROG, K.U. FRÖHLICH, F. SINER, N. TAVERNARAKIS, N. MINOIS, G. KROEMER, AND F. MADEO. Induction of autophagy by spermidine promotes longevity. **Nat. Cell Biol.** 11: 1305-1314, 2009.
- EISENBERG, T.; BÜTTNER, S.; KROEMER; G.; MADEO, F. The mitochondrial pathway in yeast apoptosis. **Apoptosis**., v.12(5), p.1011-23, 2007.

- EISENBERG, T.; CARMONA-GUTIERREZ, D.; BÜTTNER, S.; TAVERNARAKIS, N.; MADEO, F. Necrosis in yeast. **Apoptosis**, v.15, p.257–268, 2010.
- ELMORE, S. Apoptosis: a review of programmed cell death. **Toxicol Pathol.** v. 35, p. 495–516, 2007.
- ENARI, M.; SAKAHIRA, H.; YOKOYAMA, H.; OKAWA, K.; IWAMATSU, A.; NAGATA, S. Caspase-activated DNase that degrades DNA during apoptosis, and its inhibitor ICAD. **Nature**. v. 391, p. 43–50, 1998.
- ERNST R, KUEPPERS P, STINDT J, KUCHLER K, SCHMITT L: Multidrug efflux pumps: substrate selection in ATP-binding cassette multidrug efflux pumps--first come, first served? **FEBS J.** 2010, 277(3):540-9.
- ESTRADA E, HATANO NA: Vibrational approach to node centrality and vulnerability in complex networks. **Phys. A.** 2010, 389: 3648-3660.
- ESTRADA E: Virtual identification of essential proteins within the protein interaction network of yeast. **Proteomics** 2006, 6: 35-40.
- FEDOR, F.; SEVERIN, A; MARGARITA, V.; MEER, B.; EKATERINA, A.; SMIRNOVA, A.; DMITRY, A.; KNORRE, A.; VLADIMIR, P.;SKULACHEV. Natural causes of programmed death of yeast *Saccharomyces cerevisiae* Biochimica. **Biophysica Acta** v. 1783, p. 1350–1353, 2008.
- FELDMANN, H. Yeast molecular biology: a short compendium on basic features and novel aspects, 2005. Disponível em: <http://biochemie.web.med.unimuenchen.de/Yeast_Biol/>. Acesso em: 12 dez. 2011.
- FONZI WA, IRWIN MY. Isogenic strain construction and gene mapping in *Candida albicans*. **Genetics**. Jul;134(3):717-28, 1993.
- FRANCO S.L, BRUSCH M.L, MOURA L.P.P, BUENO J.H.P. Avaliação Farmacognóstica da própolis da região de Maringá. **Rev Bras Farmacogn** 9: 1-10, 2000.
- FRÖHLICH, K. U.; FUSSI, H.; RUCKENSTUHL, C. Yeast apoptosis-From genes to pathways. **Semin Cancer Biol.**, v.17, p. 112–121, 2007.
- GALLUZZI, L.; JOZA, N.; TASDEMIR, E.; MAIURI, M. C.; HENGARTNER, M.; ABRAMS, J. M.; TAVERNARAKIS, N.; PENNINGER, J.; MADEO, F.; KROEMER, G. No death without life: vital functions of apoptotic effectors. **Cell Death Differ.**, v. 15, p. 1113-1123, 2008.

- GALLUZZI, L.; ZAMZAMI, N.; ROUGE, T. L. M.; LEMAIRE, C.; BRENNER, C.; KROEMER, G. Methods for the assessment of mitochondrial membrane permeabilization in apoptosis. **Apoptosis**, v.12, p.803–813, 2007.
- GIANNATTASIO S, ATLANTE A, ANTONACCI L, GUARAGNELLA N, LATTANZIO P, PASSARELLA S, MARRA E. Cytochrome *c* is released from coupled mitochondria of yeast en route to acetic acid-induced programmed cell death and can work as an electron donor and a ROS scavenger. **FEBS Lett.**, v.582, n.10, p.1519-25, 2008.
- GILLUM AM, TSAY EY, KIRSCH DR. Isolation of the *Candida albicans* gene for orotidine-5'-phosphate decarboxylase by complementation of *S. cerevisiae* *ura3* and *E. coli* *pyrF* mutations. **Mol Gen Genet.**198(1):179-82, 1984.
- GIRVAN M, NEWMAN ME: Community structure in social and biological networks. **Proc. Natl. Acad. Sci. U S A.** 99: 7821-7826, 2002.
- GLICK, B., AND SCHATZ, G. Import of proteins into mitochondria. **Annu. Rev. Genet.** 25: 21-44, 1991.
- GLICK, D.; BARTH, S.; MACLEOD, K. F. Autophagy: cellular and molecular mechanisms. **J Pathol.**, v.221, n.1, p.3–12, 2010.
- GOFFEAU. A.; BARRELL, B.G.; BUSSEY, H.; DAVIS, R.W.; DUJON, B.; FELDMANN, H.; GALIBERT, F.; HOHEISEL, J.D.; JACQ, C.; JOHNSTON, M.; LOUIS, E.J.; MEWES, H.W.; MURAKAMI, Y.; PHILIPPSSEN, P.; TETTELIN, H.; OLIVER, S.G. Life with 6000 genes. **Science.** v. 274(5287), p. 546, 563-7, 1996.
- GOLSTEIN, P.; KROEMER, G.Redundant cell death mechanisms as relics and backups. **Cell Death Differ.**, v.12, p.1490–1496, 2005.
- GREEN, D.R. Apoptotic pathways: ten minutes to dead. **Cell.**, v. 121, p. 671–674, 2005.
- GREENAWAY, W.; SCAYSBROOK, T.; WHATLEY, F.R. The composition and plant origins of propolis. **Bee World.**, v.71, p.107–18, 1991.
- GRUBER JV, HOLTZ R: Examining the genomic influence of skin antioxidants in vitro. **Mediators Inflamm.** 2010, pii: 230450.
- GULLO A. Invasive fungal infections: the challenge continues. **Drugs.** 69 Suppl 1: 65-73, 2009.

- GULSHAN, K., AND MOYE-ROWLEY, W.C. Multidrug Resistance in Fungi. **Eukaryotic Cell** 6: 1933–1942, 2007.
- GUTHRIE, C. AND FINK, G.R. Guide to Yeast Genetics and Molecular Biology, Methods in Enzymology, Vol. 169, **Academic Press**, San Diego, 1991.
- HEGAZI, A.G.; ABD-EL-HADY, F.K; ABD-ALLAH, F.A. Chemical composition and antimicrobial activity of European propolis. **Z Naturforsch**, v. 55C, p.70–5, 2000.
- HILLENMEYER, M.E.; FUNG, E.; WILDENHAIN, J.; PIERCE, S.E.; HOON, S.; LEE, W.; PROCTOR, M.; ST.ONGE, R.P.; TYERS, M.; KOLLER, D.; ALTMAN, R.B.; DAVIS, R.W.; NISLOW, C. AND GIAEVER, G. The chemical genomic portrait of yeast: uncovering a phenotype for all genes. **Science** 320: 362-365, 2008.
- HOON, S.; R.P. ST.ONGE, G. GIAEVER, AND C. NISLOW. Yeast chemical genomics and drug discovery: an update. Trends in Pharmacological Sciences 29: 499-504, 2008.
- HORNBY, J.M.; JENSEN, E.C.; LISEC, A.D.; TASTO, J.J.; JAHNKE, B.; SHOEMAKER, R.; DUSSAULT, P.; NICKERSON, K.W. Quorum sensing in the dimorphic fungus *Candida albicans* is mediated by farnesol. **Appl Environ Microb**, v. 67, p. 2982–2992, 2001.
- HUANG, K-J., C-C. KU, AND I. R. LEHMAN. Endonuclease G: a role for the enzyme in recombination and cellular proliferation. **Proc. Nat. Acad. Sci. USA.**, v.103, p.8995-9000, 2006.
- INOUE, Y., AND KLIONSKY, D. J. Regulation of macroautophagy in *Saccharomyces cerevisiae*. **Semin. Cell Dev. Biol.** 21: 664-70, 2010.
- ISHIHARA, Y.; SHIMAMOTO, N. Involvement of Endonuclease G in Nucleosomal DNA Fragmentation under Sustained Endogenous Oxidative Stress. **J. Biol. Chem.** v.281, n.10, p.6726-33, 2006.
- ITO J, CHANG FR, WANG HK, PARK YK, IKEGAKI M, KILGORE N, LEE KH: Anti-AIDS agents. 48.(1) Anti-HIV activity of moronic acid derivatives and the new melliforone-related triterpenoid isolated from Brazilian propolis. **J. Nat. Prod.** 2001, 64:1278–1281.
- JACOBSEN, I.D, WILSON D., WÄCHTLER B., BRUNKE S., NAGLIK J.R, HUBE B. *Candida albicans* dimorphism as a therapeutic target. **Expert Rev Anti Infect Ther.** Jan;10(1):85-93, 2012.

- KADOSH, D.; A. D. JOHNSON. Induction of the *Candida albicans* filamentous growth program by relief of transcriptional repression: a genome-wide analysis. **Mol. Biol. Cell**, 16:2903–2912., 2005.
- KANAFANI, Z. AND PERFECT, J. Antimicrobial resistance: resistance to antifungal agents: mechanisms and clinical impact. **Clin. Infect. Dis.** 46: 120–128, 2008.
- KANE, P. The where, when, and how of organelle acidification by the yeast vacuolar H⁺-ATPase. *Microbiol. Mol. Biol. Rev.* 70: 177-191, 2006.
- KANKI, T., AND KLIONSKY, D.J. The molecular mechanism of mitochondria autophagy in yeast. **Mol. Microbiol.** 75: 795-800, 2010.
- KIMOTO T, ARAI S, KOHGUCHI M, NOMURA Y, MICALLEF MJ, KURIMOTO M, MITO K: Apoptosis and suppression of tumor growth by artemisinin C extracted from Brazilian propolis. **Cancer Detect. Prev.** 1998, 22:506–515.
- KLIONSKY, D.J., H. ABELIOVICH, P. AGOSTINIS, D.K. AGRAWAL, G. ALIEV, D.S. ASKEW, M. BABA, E.H. BAEHRECKE, B.A. BAHR, A. BALLABIO, AND B.A. BAMBER. Guidelines for the use and interpretation of assays for monitoring autophagy in higher eukaryotes. **Autophagy** 4: 151–175, 2008.
- KLIONSKY, D.J.; CUERVO, A.M. AND SEGLIN, P.O. Methods for monitoring autophagy from yeast to human. **Autophagy** 3: 181-206, 2007.
- KONTOYIANNIS, D.P. AND LEWIS, R.E. Antifungal drug resistance of pathogenic fungi. **Lancet** 359: 1135–1144, 2002.
- KORU, O; TOKSOY F, ACIKEL CH, TUNCA YM, BAYSALLAR M, USKUDAR GUCLU A, AKCA E, OZKOK TUYLU A, SORKUN K, TANYUKSEL M, SALIH B: Oral and dental bacteriology and infection. In vitro antimicrobial activity of propolis samples from different geographical origins against certain oral pathogens. **Anaerobe**, 13:140–145. 2007.
- KOURTIS, N.; TAVERNARAKIS, N. Autophagy and cell death in model organisms. **Cell Death and Differentiation** 16: 21–30, 2009.
- KRCMERV, Y.; FRIC, M.; PISARCIKOVA, M.; HUTTOVA, M.; FILKA, J.; KRALINSKY, K.; HUPKOVA, H.; HANZEN; J.; TRUPJ. L.; LISKOVA, M. Fungemia in neonates: report of 80 cases from seven university hospitals. **Pediatrics**., v. 105, p. 913–914, 2000.

- KRCMERY V Jr. Is there in vivo-in vitro correlation between antifungal susceptibility, species of *Candida* spp. and clinical outcome? **Int J Antimicrob Agents.**, 16(4):537-9.,2000.
- KROEMER, G., RUSTIN, P., BURCELIN, R., AND PENNINGER, J.M. Targeted deletion of AIF decreases mitochondrial oxidative phosphorylation and protects from obesity and diabetes. **Cell.**, v.131, p.476-491, 2007.
- KROEMER, G.; EL-DEIRY, W. S.; GOLSTEIN, P.; PETER, M. E.; VAUX, D.; VANDENABEELE, P.; ZHIVOTOVSKY, B.; BLAGOSKLONNY, M. V.; MALORNI, W. KNIGHT, R. A.; PIACENTINI, M.; NAGATA, S.; MELINO, G. Classification of cell death: recommendations of the Nomenclature Committee on Cell Death. **Cell Death Differ.**, v.12, p.1463–1467, 2005.
- KUMAR,V.; ABBAS, A. K.; FAUSTO, N; ASTER, J. C. In: Robbins e Cotran: Patologia: Bases Patológicas das Doenças. Tradução: B. Alencar. **Editora: Elsevier Ltda.**, Rio de Janeiro, 8ª edição, 1480p., 2010.
- KUMAZAWA, S., HAMASAKA, T. E NAKAYAMA, T. Antioxidant activity of propolis of various geographic origins. **Food Chemistry**, v. 84, p. 329-339, 2004.
- KUSHNAREVA, Y., AND D.D. NEWMAYER. Bioenergetics and cell death. *Ann. N. Y. Acad. Sci.* 1201:50-57, 2010.
- LAMAGNI, T.L.; EVANS, B.G.; SHIGEMATSU, M.; JOHNSON, E.M. Emerging trends in the epidemiology of invasive mycoses in England and Wales (1990–9). **Epidemiol. Infect.**, v.126, p. 397–414, 2001.
- LANE, S., C. BIRSE, S. ZHOU, R. MATSON, AND H. LIU. DNA array studies demonstrate convergent regulation of virulence factors by Cph1, Cph2, and Efg1 in *Candida albicans*. **J. Biol. Chem.** 276:48988–48996, 2001.
- LEBERER, E., *et al.* Ras links cellular morphogenesis to virulence by regulation of the MAP kinase and cAMP signalling pathways in the pathogenic fungus *Candida albicans*. **Mol. Microbiol.**, 42:673–687, 2001.
- LEITÃO, D.P. *et al.* Comparative evaluation of in-vitro effects of Brazilian green própolis and *Baccharis dracunculifolia* extracts on cariogenic factors of *Streptococcus mutans*. **Biol. Pharm. Bull.** 27(11):1834-9, 2004.
- LEITER, E.; SZAPPANOS, H.; OBERPARLEITER, C.; KAISERER, L.; CSERNOCH, L.; PUSZTAHELYI, T.; EMRI, T.; PÓCSI, I.; SALVENMOSER, W.; MARX, F. Antifungal Protein PAF Severely Affects the Integrity of the Plasma Membrane of *Aspergillus nidulans* and Induces an Apoptosis-Like Phenotype. **Antimicrob Agents Chemother.** V.49, n.6, p. 2445–2453, 2005.

- LEVINE B, KROEMER G. Autophagy in aging, disease and death: the true identity of a cell death impostor. **Cell Death Differ.** v.16, n.1, p.1-2, 2009.
- LI, J. AND YUAN, J. Caspases in apoptosis and beyond. **Oncogene** 27: 6194-6206, 2008.
- LI, J.; ZHOU, J.; LI, Y.; QIN, D.; LI, P. Mitochondrial fission controls DNA fragmentation by regulating endonuclease G. **Free Radic Biol Med.**, v.49, p.622–631, 2010.
- LI, L.Y., X. LUO, AND X. WANG. Endonuclease G is an apoptotic DNase when released from mitochondria. **Nature.**, v.412, p.95-99, 2001.
- LIBÉRIO, S.A., A.L. PEREIRA, M.J. ARAÚJO, R.P. DUTRA, F.R. NASCIMENTO, V. MONTEIRO-NETO, M.N. RIBEIRO, A.G. GONÇALVES, AND R.N. GUERRA. The potential use of propolis as a cariostatic agent and its actions on mutans group streptococci. **J. Ethnopharmacol.** 125: 1-9, 2009.
- LIMA M.G. A produção de própolis no Brasil. São João da Boa Vista: **São Sebastião Editora e Gráfica**, 2006.
- LIU, X.; KIM, C.N.; YANG J.; JEMMERSON R.; WANG, X. Induction of apoptotic program in cell-free extracts: requirement for dATP and cytochrome c. **Cell** v. 86, p. 147, 1996.
- LOWRY, O.H., N.J. ROSEBROUGH, A. LEWIS FARR, AND R.J. RANDALL. Protein measurement with the Folin phenol reagent. **J. Biol. Chem.** 193: 265-275, 1951.
- LUDOVICO, P., F. RODRIGUES, A. ALMEIDA, M.T. SILVA, A. BARRIENTOS, AND M. CÔRTE-REAL. Cytochrome c release and mitochondria involvement in programmed cell death induced by acetic acid in *Saccharomyces cerevisiae*. **Mol. Biol. Cell** 13: 2598-2606, 2002.
- LUDOVICO, P.; RODRIGUES, F.; ALMEIDA, A.; SILVA, M. T.; BARRIENTOS, A. AND CORTE-REAL, M. Cytochrome c release and mitochondria involvement in programmed cell death induced by acetic acid in *Saccharomyces cerevisiae*. **Mol. Biol. Cell** 13: 2598-2606, 2006.
- LUDOVICO, P.; SOUSA, M. J.; SILVA, M. T.; LEO, C.; CORTE-REAL, M. *Saccharomyces cerevisiae* commits to a programmed cell death process in response to acetic acid. **Microbiology.** v. 147, p. 2409–2415, 2001.

- LUSTOSA S. R; GALINDO A. B; NUNES L.C.C; RANDAU K.P; ROLIM NETO P.J. Própolis: atualizações sobre a química e a farmacologia. **Rev Bras Farmacogn** 18: 447-454, 2008.
- LYNCH-DAY, M.A., AND KLIONSKY, D.J. The Cvt pathway as a model for selective autophagy. **FEBS Lett.** 584: 1359-1366, 2010.
- MADEO, F.; FRÖHLICH, E.; FRÖHLICH, K.-U. A Yeast Mutant Showing Diagnostic Markers of Early and Late Apoptosis., **The Journal of Cell Biology**, v. 139(3), p. 729–734, 1997.
- MADEO, F.; FRÖLICH, E.; LIGR, M.; GREY, M.; SIGRIST, S.J.; WOLF, D.H. AND FRÖLICH, K.-U. Oxygen stress: a regulator of apoptosis in yeast. **J. Cell Biol.** 145: 757-767, 1999.
- MADEO, F.; HERKER, E.; WISSING, S.; JUNGWIRTH, H.; EISENBERG, T.; FRÖHLICH, K.-U. Apoptosis in yeast. **Current Opinion in Microbiology**, v. 7, p.655–660, 2004.
- MADEO, F.; HERKER; MALDENER, E. C.; WISSING, S.; LACHELT, S.; HERLAN, M.; FEHR, M.; LAUBER, K.; SIGRIST, S.J.; WESSELBORG, S.; FROHLICH, K.U. A caspase related protease regulates apoptosis in yeast, **Mol. Cell**, v.9, p. 911–917, 2002.
- MAERE, S.; HEYMANS, K. AND KUIPER, M. BiNGO: a Cytoscape plugin to assess overrepresentation of Gene Ontology categories in biological networks. **Bioinformatics** 21: 3448-3449, 2005.
- MAIURI, M.C., E. ZALCKVAR, A. KIMCHI, AND G. KROEMER. Self-eating and self-killing: crosstalk between autophagy and apoptosis. **Nature Rev. Mol. Cell Biol.** 8: 741-752, 2007.
- MANAVATHU, E.K.; CULRIGHT, J.L. AND CHANDRASEKAR, C.H. P.H. Organism-dependent fungicidal activities of propolis. **Antimicrobial Agents and Chemotherapy** 42: 3018-3021, 1998.
- MANON, S; CHAUDHURI, B; GUÉRIN, M. Release of cytochrome c and decrease of cytochrome c oxidase in Bax-expressing yeast cells, and prevention of these effects by coexpression of Bcl-xL. **FEBS Lett.** v. 415(1), p. 29-32, 1997.
- MARDON D., BALIST! E., PHIUPS W.: Control of dimorphism in a biochemical variant of *Candida albicans*. **J. Bacteriol.**, 100, 701- 707, 1969.

- MATSUNO T, MATSUMOTO Y, SAITO N, MORIKAWA J: Isolation and characterization of cytotoxic diterpenoid isomers from propolis. **Z. Naturforsch** 1997, C 52:702–704.
- MAZZONI, C., AND FALCONE, C. Caspase-dependent apoptosis in yeast. **Biochim. Biophys. Acta.** 1783: 1320-1327, 2008.
- MENNA-BARRETO, R.F.S., SALOMÃO K., DANTAS A.P., SANTA-RITA R.M., SOARES M.J., BARBOSA H.S., DE CASTRO D S.L. Different cell death pathways induced by drugs in *Trypanosoma cruzi*: An ultrastructural study. **Micron** 40: 157–168, 2009.
- MIESENBOCK, G., D.A. DE ANGELIS, AND J.E. ROTHMAN. Visualizing secretion and synaptic transmission with pH-sensitive Green fluorescent proteins. **Nature** 394: 192-195, 1998.
- MIZUSHIMA, N.; LEVINE, B.; CUERVO, A. M.; KLIONSKY, D. J.; Autophagy fights disease through cellular self-digestion. **Nature.**, v.451, n.7182, p.1069-75, 2008.
- MORSCHHÄUSER, J. Regulation of multidrug resistance in pathogenic fungi. **Fungal Genetics and Biology** 47: 94-106, 2010.
- MORTIMER, R.K. AND SCHILD, D. in: J.R. Broach, J.R. Pringle and E.W. Jones, eds. The Molecular and Cellular Biology of the Yeast *Saccharomyces*. **Cold Spring Harbor Laboratory Press**, Cold Spring Harbor, New York, pp. 11-26, 1991.
- MOVASSAGH, M.; ROGER, S.; FOO, S.-W. Simplified apoptotic cascades. **Heart Fail Rev.**, v.13, p. 111–119, 2008.
- MURAD, A. M., *et al.* NRG1 represses yeast-hypha morphogenesis and hypha-specific gene expression in *Candida albicans*. **EMBO J.** 20:4742–4752, 2001.
- NELSON, D. L.; COX, M. M. In: **Lehninger Princípios de Bioquímica**. Tradução: Wilson Roberto Navega. Editora: Savier, 4ª edição, São Paulo, 2006.
- NEWMAN MEJ: A measure of betweenness centrality based on random walks. **Soc. Networks** 2005, 27: 39-54.
- NOBLE SM, JOHNSON AD. Genetics of *Candida albicans*, a diploid human fungal pathogen. **Annu Rev Genet.** 41:193-211, 2007.

- NUNES, L. C. C.; GALINDO, A. B.; DEUS, A. S. O.; RUFINO, D. A.; RANDAU, K. P.; XAVIER, H. S.; CITÓ, G. L.; NETO, R. J. P. Variabilidade sazonal dos constituintes da própolis vermelha e bioatividade em *Artermia salina*. **Revista Brasileira de Farmacognosia**, 19(2B): 524-529, 2009.
- ODA, K.; KAWASAKI, N.; FUKUYAMA, M.; IKEDA, S. Ectopic Expression of Mitochondria Endonuclease Pnu1p from *Schizosaccharomyces pombe* Induces Cell Death of the Yeast. **J Biochem Mol Biol.**, v.40, n.6, p.1095-1099, 2007.
- ODDS, F. C., *et al.* Evaluation of possible correlation between antifungal susceptibilities of filamentous fungi in vitro and antifungal treatment outcomes in animal infection models. **Antimicrob. Agents Chemother.** 42: 282–288, 1998.
- ODDS, F.C.; BROWN, A.J.P.; GOW, N.A.R.; *Candida albicans* genome sequence: a platform for genomics in the absence of genetics. *Genome Biol.* v. 5, p. 230–235, 2004.
- OKADA H, MAK TW. Pathways of apoptotic and nonapoptotic death in tumour cells. **Nat Rev Cancer.**, v.4, p.592-603, 2004.
- OSIEWACZ, H.D., AND SCHECKHUBER, C.Q. Impact of ROS on ageing of two fungal model systems: *Saccharomyces cerevisiae* and *Podospora anserine*. **Free Radic. Res.** 40: 1350-1358, 2006.
- OTA, C.; UNTERKIRCHER, C.; FANTINATO, V. AND SHIMIZU, M. T. Antifungal activity of propolis on different species of *Candida*. **Mycoses** 44: 375-378, 2001.
- PARK, Y.K; ALENCAR, S.M; AGUIAR; C.L. Botanical origin and chemical composition of Brazilian propolis. **J Agric Food Chem** v. 50, p. 2502–6, 2002.
- PARRISH, J., L. LI, K. KLOTZ, D. LEDWICH, X. WANG, AND D. XUE. Mitochondrial endonuclease G is important for apoptosis in *C. Elegans*. **Nature.**, v.412, p.90-94, 2001.
- PEREIRA A. S; SEIXAS F.R.M.S.; NETO F.R.A. Propolis: 100 anos de pesquisa e suas perspectivas futuras. **Quim Nova**, 25: 321-326, 2002.
- PEREIRA, C.; SILVA, R. D.; SARAIVA, L.; JOHANSSON, B.; SOUSA, M. J.; CÔRTE-REAL, M. Mitochondria-dependent apoptosis in yeast. **Biochim Biophys Acta.**, v.1783, p.1286–1302, 2008.

- PERRONE, G.G., S.X. TAN, AND I.W. DAWES. Reactive oxygen species and yeast apoptosis. **Biochim. Biophys. Acta** 1783: 1354-1368, 2008.
- PHILLIPS, A. J.; CROWE, J. D.; RAMSDALE, M. Ras pathway signaling accelerates programmed cell death in the pathogenic fungus *Candida albicans*. **PNAS** v. 103, 2006.
- PIMENTEL FIGUEIREDO, B. C.; DE CASTRO, P. A.; DINAMARCO, T. M.; GOLDMAN, M. H. S.; GOLDMAN, G. H. The *Aspergillus nidulans* *nucA*^{ENDOG} homologue is not involved in cell death. **Eukaryotic Cell**, 10(2), 2011.
- PRYTZYK, E; DANTA,S A.P; SALOMÃO, K; PEREIRA, A.S; BANKOVA, V.S; DE CASTRO, S.L. Flavonoids and trypanocidal activity of Bulgarian propolis. **J Ethnopharmacol.**, v. 88, p. 189–93, 2003.
- RAHMAN D.; MISTRY M.; THAVARAJ, S.; CHALLACOMBE, S. J.; NAGLIK J. R. Murine model of concurrent oral and vaginal *Candida albicans* colonization to study epithelial host–pathogen interactions. **Microbes Infect.**, v. 9(5): 615–622, 2007.
- RAMOS A.F.N.; MIRANDA J.L. Própolis: a review of its anti-inflamamtory and healing actions. **Journal of Venomous Animal and Toxins Including Tropical Diseases**, 13, 697–710., 2007.
- RAMSDALE, M. Programmed cell death in pathogenic fungi. **Biochimica Biophys. Acta** 1783: 1369-1380, 2008.
- RIBEIRO, G. F.; CÔRTE-REAL M., AND JOHANSSON, B. Characterization of DNA Damage in Yeast Apoptosis Induced by Hydrogen Peroxide, Acetic Acid, and Hyperosmotic Shock. **Molecular Biology of the Cell** 17: 4584–4591, 2006.
- RIVALS, I.; PERSONNAZ, L.; TAING, L. AND POTIER, M.C. Enrichment or depletion of a GO category within a class of genes: which test? **Bioinformatics** 23: 401-407, 2007.
- ROSADO, C. J., D. MIJALJICA, I. HATZINISIRIOU, M. PRESCOTT, AND R.J. DEVENISH. A fluorescent pH-biosensor for reporting vacuolar turnover of cytosol and organelles in yeast. **Autophagy** 4: 205-213, 2008.
- ROUCOU, X.; PRESCOTT, M.; DEVENISH, R. J.; NAGLEY, P. A cytochrome c-GFP fusion is not released from mitochondria into the cytoplasm upon expression of Bax in yeast cells., **FEBS Lett.**, v. 471, p. 235-239, 2000.

- RUBINSZTEIN, D. C.; GESTWICKI, J.E.; MURPHY, L. O.; KLIONSKY, D. J. Potential therapeutic applications of autophagy. **Nat Rev Drug Discov.**, v.6, n.4, p.304-312, 2007.
- RUSSO, A.; LONGO, R.; VANELLA, A. Antioxidant activity of própolis: role of caffeic acid phenethyl ester and galangin. *Fitoterapia*, 73: S21-S29, (Supl. 1), 2002.
- SACCHAROMYCES GENOME DELETION PROJECT (http://www-sequence.stanford.edu/group/yeast_deletion_project/).
- SÁ-CORREIA I, DOS SANTOS SC, TEIXEIRA MC, CABRITO TR, MIRA NP: Drug:H⁺ antiporters in chemical stress response in yeast. **Trends Microbiol.** 2009, 17(1):22-31.
- SALATINO, A.; T EIXEIRA, E.W.; N EGRI, G.; MESSAGE , D. Origin and Chemical Variation of Brazilian Propolis. **Evidencebased Complementary and Alternative Medicine**, v.2, n.1, p.33-38, 2005.
- SALOMÃO, K.; PEREIRA, P. R. S. ;CAMPOS, L. C.; M. BORBA, C. CABELLO; P. H. MARCUCCI; M. C. DE CASTRO S. L. Brazilian Propolis: Correlation Between Chemical Composition and Antimicrobial Activity **eCAM**, v. 5(3), p.317–324, 2008.
- SAMBROOK, J.; RUSSELL, D. W. Molecular cloning: a laboratory manual. **3rd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY**, 2001.
- SANGLARD, D., ET AL. Mechanisms of resistance to azole antifungal agents in *Candida albicans* isolates from AIDS patients involve specific multidrug transporters. **Antimicrob. Agents Chemother.**, 39:2378–2386, 1995.
- SANGLARD, D., F. ISCHER, M. MONOD, AND J. BILLE. Cloning of *Candida albicans* genes conferring resistance to azole antifungal agents: characterization of CDR2, a new multidrug ABC transporter gene. **Microbiology** 143(Pt. 2):405–416, 1997.
- SAVOLDI, M.; MALAVAZI, I.; SORIANI, F. M.; CAPELLARO, J. L.; KITAMOTO, K.; FERREIRA, M. E. S.; GOLDMAN, M. H. S.; GOLDMAN, G. H. Farnesol induces the transcriptional accumulation of the *Aspergillus nidulans* Apoptosis-Inducing Factor (AIF)-like mitochondrial oxidoreductase. **Mol Microbiol.**, v.70, n.1, p.44–59, 2008.
- SCAFFIDI, P.; MISTELI, T. AND BIANCHI, M.E. Release of chromatin protein HMGB1 by necrotic cells triggers inflammation. **Nature** 418: 191-195, 2002.

- SCARDONI G, PETTERLINI M, LAUDANNA C: Analyzing biological network parameters with CentiScaPe. **Bioinformatics** 2009, 25: 2857-2859.
- SCHÄFER P, SCHOLZ SR, GIMADUTDINOW O, CYMERMAN IA, BUJNICKI JM, RUIZ-CARRILLO A, PINGOUD A, MEISS G. Structural and Functional Characterization of Mitochondrial EndoG, a Sugar Non-specific Nuclease which Plays an Important Role During Apoptosis. **J. Mol. Biol.**, v.338, p.217–228, 2004.
- SCHAUER, A.; KNAUER, H.; RUCKENSTUHL, C.; FUSSI, H.; DURCHSCHLAG, M.; POTOČNIK, U. AND FRÖHLICH, K-U. Vacuolar functions determine the mode of cell death. **Biochim. Biophys. Acta** 1793: 540-545, 2009.
- SCHERENS. B.; GOFFEAU, A. The uses of genome-wide yeast mutant collections. **Genome Biology**, v. 5, p. 229, 2004.
- SELLAM, A., *et al.* Role of transcription factor CaNdt80p in cell separation, hyphal growth, and virulence in *Candida albicans*. **Eukaryot. Cell**, 9:634–644, 2010.
- SEMIGHINI, C. P.; HORNBY, J. M.; DUMITRU, R.; NICKERSON, K.W.; HARRIS, S.D. Farnesol-induced apoptosis in *Aspergillus nidulans* reveals a possible mechanism for antagonistic interactions between fungi. **Mol Microbiol.**, v.59, n.3, p.753–764, 2006.
- SEMIGHINI, C.P.; MARINS, M.; GOLDMAN, M.H.S. AND GOLDMAN, G.H. Quantitative analysis of the relative transcript levels of ABC transporter *Atr* genes in *Aspergillus nidulans* by Real-Time Reverse Transcription-PCR assay. **Appl. Environ. Microbiol.** 68: 1351-1357, 2002.
- SENEDESE JM, RODRIGUES AR, FURTADO MA, FAUSTINO VD, BERRETTA AA, MARCHETTI JM, TAVARES DC. Assessment of the mutagenic activity of extracts of brazilian propolis in topical pharmaceutical formulations on Mammalian cells in vitro and in vivo. **Evid Based Complement Alternat Med.** 2011:315701, 2011.
- SEVERIN, F. F.; MEER, M. V.; SMIRNOVA, E. A.; KNORRE, D. A.; SKULACHEV, V. P.; Natural causes of programmed death of yeast *Saccharomyces cerevisiae*. **Biochim Biophys Acta.**, v.1783, p.1350–1353, 2008.
- SFORCIN, J.M. Propolis and the immune system: a review. **J. Ethnopharmacol.** 113: 1-14, 2007.
- SHAHI P, MOYE-ROWLEY WS: Coordinate control of lipid composition and drug transport activities is required for normal multidrug resistance in fungi. **Biochim Biophys Acta.** 2009, 1794(5):852-9.

- SHANNON P, A. MARKIEL, O. OZIER, N.S. BALIGA, J.T. WANG, D. RAMAGE, N. AMIN, B. SCHWIKOWSKI, AND T. IDEKER. Cytoscape: a software environment for integrated models of biomolecular interaction networks. **Genome Res.**13: 2498-2504, 2003.
- SHAPIRO R. S.; ROBBINS, NICOLE; COWEN, L. E. Regulatory Circuitry Governing Fungal Development, Drug Resistance, and Disease. **Microbiol. Mol. Biol. Rev.** 75(2):213, 2011.
- SHARON, A.; FINKELSTEIN, N.; SHLEZINGER; I.; HATAM. Fungal apoptosis: function, genes and gene function. **FEMS Microbiol Rev** v. 33, p. 833–854, 2009.
- SHIMAZAWA, M.; CHIKAMATSU, S.; MORIMOTO, N.; MISHIMA, S.; NAGAI, H. AND HARA, H. Neuroprotection by Brazilian green propolis against *in vitro* and *in vivo* ischemic neuronal damage. **eCAM** 2: 201–207, 2005.
- SHIRTLIFF, M; KROM, B.P; MEIJERING, R.A; PETERS, B.M; ZHU, J; SCHEPER, M.A; HARRIS, M.L; JABRA-RIZK, M.A. Farnesol-induced apoptosis in *Candida albicans*. Antimicrobial agents and chemotherapy. **Epub**, v. 53(6), p. :2392-401, 2009.
- SILICI, S.; N. KOÇ, A.; AYANGID, L; ÇANKAYA, S. Antifungal Activities of Propolis Collected by Different Races of Honeybees Against Yeasts Isolated From Patients With Superficial Mycoses. **J Pharmacol Sci**, v. 99, p. 39 – 44, 2005.
- SIMÕES, L.M.C.; GREGÓRIO, L.E.; DA SILVA FILHO, A.A.; DE SOUZA, M.L.; AZZOLINI, A.E.C. S; BASTOS, J.K.; LUCISANO-VALIM, Y.M. Effect of Brazilian green propolis on the production of reactive oxygen species by stimulated neutrophils. **J. Ethnopharmacol.**, 94, 59-65, 2004.
- SIPOS G, KUCHLER K: Fungal ATP-binding cassette (ABC) transporters in drug resistance & detoxification. **Curr Drug Targets**. 2006, 7(4):471-81.
- SIQUEIRA, A.B.S., B.S. GOMES, I. CAMBUIM, R. MAIA, S. ABREU, C.M. SOUZA-MOTTA, L.A. DE QUEIROZ, AND A.L.F. PORTO. *Trichophyton* species susceptibility to green and red propolis from Brazil. **Letters in Applied Microbiology** 48: 90–96, 2009.
- SRIPRIYA, P.; VEDANTAM, L. V .AND PODILE, A. R.. Involvement of mitochondria and metacaspase elevation in hairpin Pss-induced cell death of *Saccharomyces cerevisiae*. **J. Cell Biochem**. 107: 1150-1159, 2009.
- STOLDT, V. R., A. SONNEBORN, C. E. LEUKER, AND J. F. ERNST. Efg1p, an essential regulator of morphogenesis of the human pathogen *Candida albicans*, is a member of a conserved class of bHLH proteins regulating morphogenetic processes in fungi. **EMBO J**. 16:1982–1991, 1997.

- STRATHERN, J.N., JONES, E.W. AND BROACH, J.R. The Molecular and Cellular Biology of the Yeast *Saccharomyces*. Cold **Spring Harbor Laboratory Press**, Cold Spring Harbor, New York, 1981.
- SUD'INA, G.F; MIRZOEVA, O.K; PUSHKAREVA, M.A; KORSHUNOVA, G.A; SUMBATYAN, N.V; VARFOLOMEEV, S.D. Caffeic acid phenethyl ester as a lipoxygenase inhibitor with antioxidant properties. **FEBS Lett**, v. 329, p. 21-4, 1993.
- SUDBERY, P.E. Growth of *Candida albicans* hyphae. **Nat Rev Microbiol.**, 16;9(10):737-48, 2011.
- SUDBERY, P., N. GOW, AND J. BERMAN. The distinct morphogenic states of *Candida albicans*. **Trends Microbiol.**, 12:317–324, 2004.
- SUZUKI SW, ONODERA J, OHSUMI Y. Starvation induced cell death in autophagy-defective yeast mutants is caused by mitochondria dysfunction. **PLoS One**, 6: e17412. 2011.
- SZLISZKA, E., AND KROL W. The role of dietary polyphenols in tumor necrosis factor-related apoptosis inducing ligand (TRAIL)-induced apoptosis for cancer chemoprevention. **Eur. J. Cancer Prev.** (*in press*), 2010.
- TEIXEIRA, E.W.; NEGRI, G.; MEIRA, R.M.S.A.; MESSAGE, D. AND SALATINO, A. Plant origin of green propolis: bee behavior, plant anatomy and chemistry. **eCAM** 2: 85–92, 2005.
- TESTE MA, DUQUENNE M, FRANÇOIS JM, PARROU JL: Validation of reference genes for quantitative expression analysis by real-time RT-PCR in *Saccharomyces cerevisiae*. **BMC Mol Biol.** 2009, 10():99.
- TZAGOLOFF, A.; AKAI, A.; NEEDLEMAN, R.B. AND ZULCH, G. Assembly of the mitochondrial membrane system. Cytoplasmic mutants of *Saccharomyces cerevisiae* with lesions in enzymes of the respiratory chain and in the mitochondrial ATPase. **J. Biol. Chem.** 250: 8236-8242, 1975.
- UREN, A.G.; O'ROURKE, K.; ARAVIND, L.A.; PISABARRO, M.T.; SESHAGIRI, S.; KOONIN, E.V.; DIXIT, V.M. Identification of paracaspases and metacaspases: two ancient families of caspase-like proteins, one of which plays a key role in MALT lymphoma. **Mol. Cell**, v. 6, p. 961, 2000.
- URUSHISAKI T, TAKEMURA T, TAZAWA S, FUKUOKA M, HOSOKAWA-MUTO J, ARAKI Y, KUWATA K. Caffeoylquinic acids are major constituents with potent anti-influenza effects in brazilian green propolis water extract. **Evid Based Complement Alternat Med.** 2011:254914, 2011.

- VACHOVA, L.; PALKOVA, Z. Caspases in yeast apoptosis-like death: facts and artifacts, **FEMS Yeast Res.** v. 7, p. 12–21, 2007.
- VARGAS A.C, LOGUERCIO A.P, WITT N.M, DA COSTA M.M, SÁ E SILVA M, VIANA L.R. Atividade antimicrobiana “in vitro” de extrato alcoólico de própolis. **Ciência Rural** 34: 159-163, 2004.
- VIDEIRA A, KASUGA T, TIAN C, LEMOS C, CASTRO A, GLASS NL. Transcriptional analysis of the response of *Neurospora crassa* to phytosphingosine reveals links to mitochondrial function. **Microbiology.**, 155(Pt 9):3134-41, 2009.
- VYNOGRAD N, VYNOGRAD I, SOSNOWSKI Z: A comparative multicenter study of the efficacy of propolis, acyclovir and placebo in the treatment of genital herpes (HSV). **Phytomedicine** 2000, 7:1–6.
- WALKER LA, MUNRO CA, DE BRUIJN I, LENARDON MD, MCKINNON A, et al. Stimulation of Chitin Synthesis Rescues *Candida albicans* from Echinocandins. **PLoS Pathog** v. 4(4): e1000040., 2008.
- WHITE, T. C., M. A. PFALLER, M. G. RINALDI, J. SMITH, AND S. W. REDDING. Stable azole drug resistance associated with a substrain of *Candida albicans* from an HIV-infected patient. **Oral Dis.**, 3 (Suppl. 1):S102–S109, 1997.
- WHITEWAY M, BACHEWICH C. Morphogenesis in *Candida albicans*. **Annu Rev Microbiol**, 61: 529-553, 2007.
- WILSON RB, DAVIS D, MITCHELL AP. Rapid hypothesis testing with *Candida albicans* through gene disruption with short homology regions. **J Bacteriol.** Mar;181(6):1868-7, 1999.
- WINZELER, E. A.; SHOEMAKER, D. D.; ASTROMOFF, A.; LIANG, H.; ANDERSON, K.; ANDRE, B.; BANGHAM, R.; BENITO, R.; BOEKE, J. D.; BUSSEY, H.; CHU, A. M.; CONNELLY, C.; DAVIS, K.; DIETRICH, F.; DOW SALLY, W.; EL BAKKOURY, M.; FOURY, F.; FRIEND, S.H.; GENTALLEN, E.; GIAEVER, G.; HEGEMANN, J. H.; JONES, T.; LAUB, M; LIAO, H.; LIEBUNDGUTH, N.; LOCKHART, D. J.; LUCAU-D, A.; LUSSIER, M.; M'RABET, N.; MENARD, P.; MITTMANN, M.; CHAI, REBISCHUNG; C., REVUELTA J.L.; RILES I.; ROBERTS C. J.; ROSS-MACDONALD, P.; SCHERENS, B.; SNYDER, M.; SOOKHAI-MAHADEO, S.; STORMS, R. K.; VÉRONNEAU, S.; VOET, M.; VOLCKAERT, G.; WARD, T. R.; WYSOCKI, R.; YEN, G. S.; YU, K.; ZIMMERMANN, K.; PHILIPPSSEN, P.; JOHNSTON, M.; DAVIS, R. Functional Characterization of the *S. cerevisiae* Genome by Gene Deletion and Parallel Analysis. **Science**, v. 285, p. 901–906, 1999.

- WISSING, S.; LUDOVICO, P.; HERKER, E.; BUTTNER, S.; ENGELHARDT, S.M.; DECKER, T.; LINK, A.; PROKSCH, A.; RODRIGUES, F.; CORTE-REAL, M.; FROHLICH, K.U.; MANNS, J.; CANDE, C.; SIGRIST, S.J.; KROEMER, G.; MADEO, F. An AIF orthologue regulates apoptosis in yeast. **J. Cell Biol.** v. 166, p. 969–974, 2004.
- WUSTER, A. AND MADAN BABU M. Chemogenomics and biotechnology. Trends in Biotechnology 26: 252-258, 2008.
- YANO J, NOVERR MC, FIDEL PL JR. Cytokines in the host response to Candida vaginitis: Identifying a role for non-classical immune mediators, S100 alarmins. **Cytokine**, 58(1):118-28, 2012.
- YORIMITSU T, KLIONSKY DJ. Autophagy: molecular machinery for self-eating. **Cell Death Differ.**, 12 Suppl 2:1542-52, 2005.
- YU. H; KIM P.M; SPRECHER E; TRIFONOV V; GERSTEIN M: The importance of bottlenecks in protein networks: correlation with gene essentiality and expression dynamics. **PLoS Comput. Biol.** 3: e59. 2007.
- ZASSENHAUS, H. P.; DENNIGER, G. Analysis of the role of the NUC1 endo/exonuclease in yeast mitochondrial DNA recombination. **Curr Genet.**, 25: 142-149, 1993.
- ZEWAIL, A.; XIE, M.W.; XING, Y.; LIN, L.; ZHANG, P.F.; ZOU, W.; SAXE, J.P. AND HUANG. J. Novel functions of the phosphatidylinositol metabolic pathway discovered by a chemical genomics screen with wortmannin. **Proc. Natl. Acad. Sci. USA** 100: 3345–3350, 2003.
- ZHU, W., AND S. G. FILLER. Interactions of Candida albicans with epithelial cells. **Cell. Microbiol.** v. 12:273–282, 2010.
- ZINSER, E. AND DAUM, G. Isolation and biochemical characterization of organelles from the yeast *S. cerevisiae*. **Yeast**, v.11 (1995) 493-536.

Anexos

Os anexos **A, B, C, D, E, F, G, H** constituem-se de tabelas extensas e, por este motivo, foram gravados no CD presente na contracapa da dissertação.

Os anexos **I, J, L, M, N, O** são artigos publicados durante o período no qual desenvolvi este trabalho de mestrado, incluindo dois artigos referentes a esta dissertação (um já publicado “**I**” e outro submetido “**J**”) e mais quatro artigos, dos quais participei como co-autora.

O Anexo **P** corresponde ao “**SBBQ Award**”, prêmio para o melhor pôster apresentado durante o XL Encontro da Sociedade Brasileira de Bioquímica e Biologia Molecular (SBBq) (30 de abril a 3 de maio de 2011, em Foz do Iguaçu, Paraná, Brasil). O pôster que recebeu este prêmio é intitulado: “Molecular Characterization of Propolis Induced Cell Death in *Saccharomyces cerevisiae*, de Castro, P. A.; Savoldi, M.; Bonatto, D.; De Barros, M. H.; Goldman, M. H. S.; Berreta, A. A.; Goldman, G. H.”

7.1. Anexo I. Artigo publicado: Molecular Characterization of Propolis Induced Cell Death in *Saccharomyces cerevisiae*. De Castro, P. A.; Savoldi, M.; Bonatto, D.; Barros, M. H.; Goldman, M. H. S. ; Berretta, A. A. ; Goldman, G. H. Eukaryotic Cell, Mar. 2011, p. 398–411.

Molecular Characterization of Propolis-Induced Cell Death in *Saccharomyces cerevisiae*^{▽†}

Patrícia Alves de Castro,¹ Marcela Savoldi,¹ Diego Bonatto,⁴ Mário Henrique Barros,² Maria Helena S. Goldman,³ Andresa A. Berretta,⁶ and Gustavo Henrique Goldman^{1,5*}

Faculdade de Ciências Farmacêuticas de Ribeirão Preto,¹ Departamento de Microbiologia, Instituto de Ciências Biomédicas,² and Faculdade de Filosofia, Ciências e Letras de Ribeirão Preto,³ Universidade de São Paulo, São Paulo, Brazil; Centro de Biotecnologia da UFRGS, Universidade Federal do Rio Grande do Sul, Porto Alegre, Brazil⁴; Laboratório Nacional de Ciência e Tecnologia do Bioetanol, Caixa Postal 6170, 13083-970 Campinas, São Paulo, Brazil⁵; and Apis Flora Industrial e Comercial, Ribeirão Preto, São Paulo, Brazil⁶

Received 7 October 2010/Accepted 21 December 2010

Propolis, a natural product of plant resins, is used by the bees to seal holes in their honeycombs and protect the hive entrance. However, propolis has also been used in folk medicine for centuries. Here, we apply the power of *Saccharomyces cerevisiae* as a model organism for studies of genetics, cell biology, and genomics to determine how propolis affects fungi at the cellular level. Propolis is able to induce an apoptosis cell death response. However, increased exposure to propolis provides a corresponding increase in the necrosis response. We showed that cytochrome *c* but not endonuclease G (Nuc1p) is involved in propolis-mediated cell death in *S. cerevisiae*. We also observed that the metacaspase *YCA1* gene is important for propolis-mediated cell death. To elucidate the gene functions that may be required for propolis sensitivity in eukaryotes, the full collection of about 4,800 haploid *S. cerevisiae* deletion strains was screened for propolis sensitivity. We were able to identify 138 deletion strains that have different degrees of propolis sensitivity compared to the corresponding wild-type strains. Systems biology revealed enrichment for genes involved in the mitochondrial electron transport chain, vacuolar acidification, negative regulation of transcription from RNA polymerase II promoter, regulation of macroautophagy associated with protein targeting to vacuoles, and cellular response to starvation. Validation studies indicated that propolis sensitivity is dependent on the mitochondrial function and that vacuolar acidification and autophagy are important for yeast cell death caused by propolis.

Propolis is a natural product of plant resins collected by honeybees (*Apis mellifera*) from various plant sources. It is used by the bees to seal holes in their honeycombs and protect the hive entrance (22, 24, 82). Propolis has been used in folk medicine for centuries. Its chemical composition is quite complex since more than 300 compounds, such as polyphenols, phenolic aldehydes, sesquiterpene quinones, coumarins, amino acids, steroids, and inorganic compounds, have been identified in propolis samples. Propolis has cytotoxic (53), anti-herpes virus (83), antitumor (34), radical scavenging (4), antimicrobial (38, 60), antiprotozoan (55), and anti-HIV (30) activity and suppressive effects of dioxin toxicity (60). More recently, evidence has shown that propolis can be used to treat *Candida* fungal infections (1, 17, 59, 68, 76).

There is an urgent need to discover novel antifungal drugs. There are currently five different classes of antifungal drugs classified based on their mechanisms of action (37). Most of these drugs affect the cell membrane or cell wall integrity through inhibition of either ergosterol synthesis or β -glucan/chitin synthesis, respectively. A further complication to this

limited spectrum of drugs is the acquisition of resistance by the pathogenic fungi observed in clinical settings (26, 31, 57). While amphotericin B has fungicidal activity, the azoles are generally fungistatic agents for *Candida* spp. and fungicidal agents for *Aspergillus* spp. (52). The echinocandins (caspofungin, micafungin, and anidulafungin) exhibit fungicidal activity against *Candida* spp. and fungistatic activity against *Aspergillus* spp. (25). There are very few studies investigating the relationship between antifungal drugs and cell death (for a review, see reference 63). Recently, *Saccharomyces cerevisiae* has become an excellent model system to characterize cell death (for a review, see reference 14). The apoptotic core machinery is quite conserved in this organism, and multiple yeast orthologues of mammalian apoptotic proteins have been identified (14). In addition, several cell death pathways have been characterized, and assays for apoptotic and/or necrotic cell death are being currently used (14).

The identity and the mechanism of action of the chemical compounds present in the propolis are not known. To identify the cellular targets of any bioactive compounds is a very difficult task. In the last years, however, all the genome resources that were developed for *S. cerevisiae* (www.yeastgenome.org) provided an excellent genomic platform for drug development and drug target identification (7, 28, 50, 85). In addition, all these genetic tools have made it possible to study and construct genetic interaction networks for identification of specific drug targets (9, 16). Here, we apply the power of *S. cerevisiae* as a model organism for studies of genetics, cell biology, and genomics in order to understand how propolis affects fungi at

* Corresponding author. Mailing address: Departamento de Ciências Farmacêuticas, Faculdade de Ciências Farmacêuticas de Ribeirão Preto, Universidade de São Paulo, Av. do Café S/N, CEP 14040-903, Ribeirão Preto, São Paulo, Brazil. Phone and fax: 55 16 36024280. E-mail: ggoldman@usp.br.

† Supplemental material for this article may be found at <http://ec.asm.org/>.

[▽] Published ahead of print on 30 December 2010.

the cellular level. First, we evaluate how propolis can affect *S. cerevisiae* cell survival by assessing several genetic determinants involved in apoptosis and/or necrosis in this organism. As a complementary step, we have used the yeast nonessential gene deletion library to investigate possible cell targets for propolis.

MATERIALS AND METHODS

Strains, media, and culture methods. The following *S. cerevisiae* strains were used: BY4742 (*MAT α his3 Δ 1 leu2 Δ 0 lys2 Δ 0 ura3 Δ 0*) (10), BY4741A and BY4742 *rho⁻* (both *MAT α his3 Δ 1 leu2 Δ 0 lys2 Δ 0 ura3 Δ 0 [*rho⁻*]) (this study), YJL208C/*NUC1* (*MAT α his3 Δ 1 leu2 Δ 0 met15 Δ 0 ura3 Δ 0 GFP::nuc1*) (Invitrogen), YJR048W/*CYC1* (*MAT α his3 Δ 1 leu2 Δ 0 met15 Δ 0 ura3 Δ 0 GFP::cyc1*) (Invitrogen), and Δ YCA1 (*MAT α his3 Δ 1 leu Δ 0 met15 Δ 0 ura3 Δ 0 yca1::kanMX4*) (49). The following media were used: the complete medium YPD agar (2% [wt/vol] glucose, 1% [wt/vol] yeast extract, 2% [wt/vol] peptone, and 2% [wt/vol] agar), YPGal agar (2% [wt/vol] galactose, 1% [wt/vol] yeast extract, 2% [wt/vol] peptone, and 2% [wt/vol] agar), minimal medium SC agar (0.7% [wt/vol] yeast nitrogen base [DIFCO], 2% [wt/vol] glucose, 0.1g/liter L-leucine, 0.1g/liter L-lysine, 0.1g/liter L-tryptophan, 0.05g/liter L-histidine, 2% [wt/vol] agar), and YPD, YPGal, and SC liquid media with the same composition as above but without agar. Rosella strains were constructed by the transformation of c-Rosella plasmid (67) into the BY4742 strain. Transformants were selected in SC agar minimal medium.*

Preparation of the propolis standardized extract. Three batches of propolis standardized extract (PSE), 14004/10, 14401/10, and 010/08, were produced by Apis Flora Company (EPP-AF; Ribeirão Preto, SP, Brazil). The extracts were standardized using a propolis blend composed of raw material obtained from several sites of Brazil (A. A. Beretta-Silva, A. C. Meda, and M. E. T. Ferreira, 1 February 2005, Brazilian patent application PI 0405483), such as the states of Minas Gerais, São Paulo, Paraná, Santa Catarina, and Rio Grande do Sul. Propolis (blend of raw material) was kept in a freezer for 12 h and ground to a fine powder in a blender, and its particle size was standardized using a 42-mesh sieve. It was then extracted using hydroalcoholic solution (7:3), with dynamic maceration followed by a percolation process and finally by filtration. The PSE obtained represents 11% (wt/vol) of dry residue and chemical composition standardized qualitatively and quantitatively by reverse-phase high-pressure liquid chromatography (RP-HPLC) by analyzing the following compounds: caffeic, *p*-coumaric, and cinnamic acids; isosakuranetin; and artemillin C (ARC) (see Table S1 in the supplemental material).

Screening of the yeast deletion library. For the screening of the yeast deletion library (84; http://www-sequence.stanford.edu/group/yeast_deletion_project/) a set of approximately 4,800 haploid deletions in nonessential genes of *S. cerevisiae* were screened for propolis sensitivity. Mutants were inoculated from stock cultures in 96-well master plates at -80°C and were grown at 30°C in YPD medium containing 200 $\mu\text{g/ml}$ of Geneticin (G418; Sigma, St. Louis, MO) and then stored at 4°C . To determine which mutants were sensitive to propolis, the cells were spotted on YPD plates plus 0.68% ethanol or alcoholic propolis extract (Apis Flora, Ribeirão Preto, SP, Brazil) at 0.25% and 0.50% (vol/vol). Plates were incubated at 30°C for 5 days.

RNA isolation and real-time PCR. Exponential (9 h) growth phase BY4742 cells were harvested and suspended (10^7 cells ml^{-1}) in YPD liquid medium containing 0.125% (vol/vol) alcoholic propolis extract. The treatment was carried out for 5 and 10 min at 30°C with mechanical shaking (200 rpm). Then, yeast cells were washed with distilled water, and RNA was isolated. For total RNA isolation, the yeast cells were disrupted by vortexing with glass beads, and total RNA was extracted with Trizol reagent (Invitrogen). Ten micrograms of RNA from each treatment was then fractionated in 2.2 M formaldehyde–1.2% (wt/vol) agarose gel, stained with ethidium bromide, and then visualized with UV light. The presence of intact 25S and 18S rRNA bands was used as a criterion to assess the integrity of the RNA. RNase-free DNase I treatment for the real-time PCR experiments was carried out as previously described by Semighini et al. (72).

All the PCRs were performed using an ABI 7500 Fast Real-Time PCR System (Applied Biosystems) and TaqMan Universal PCR Master Mix kit (Applied Biosystems). The reactions and calculations were performed according to Semighini et al. (72). The primers and Lux fluorescent probes (Invitrogen) used in this work were the following: ATG8_263RL (5'-CGGCTATATGGCAGACATCA CGC-FAM-G-3') and ATG8_263RL/235FU (5'-CCCTGAGAAGGCCATCT TCATT-3') for YBL078C/ATG8; ATG14_892FL (5'-CGGACAACAAGATG AAGTGTAGGTC-FAM-G-3') and ATG14_892FL/910RUa (5'-AACCATCG AGTACAACATCCAAA-3') for YBR128C/ATG14; and ACT1_366FL (CGG

CTGGTTCTGGTATGTGTAAAGC-FAM-G) and ACT1_366FL/385RU (AT GGAAGACAGCAGGAG) for YFL039C/ACT1.

Optical and confocal microscopy. After incubation under the appropriate conditions for each experiment, slides were mounted and visualized in a fluorescence microscope. For YJL208C/*NUC1*, YJR048W/*CYC1*, and Rosella strain microscopy, stationary (16 h)-growth-phase cells were harvested and suspended (10^7 cells ml^{-1}) in SC liquid medium for YJL208C and YJR048W and in YPD liquid medium for Rosella containing 0.125% (vol/vol) alcoholic propolis extract; the treatments were carried out for 15, 10, and 90 min, respectively, at 30°C with mechanical shaking (200 rpm). A positive control for Rosella staining was performed by growing cells for 16 h at 30°C in YPD medium and transferring them to starvation medium (0.17% [wt/vol] yeast extract, 0.1% [wt/vol] KH_2PO_4 , 0.01% [wt/vol] CaCl_2 , 0.0005% [wt/vol] FeCl_3 , 0.07% [wt/vol] MgCl_2 , 0.05% [wt/vol] NaCl, 2% [wt/vol] glucose) to induce autophagy. After 4 h of incubation at 30°C , protonophore carbonyl cyanide *m*-chlorophenylhydrazone (CCCP; at 10 μM) was added in pH 7.5 buffer for 10 min. Incubation of starved cells before imaging in pH 7.5 buffer containing the protonophore CCCP resulted in an increase in green fluorescence emission in the vacuole (67). For positive control to apoptosis in the *NUC1* microscopy, we used 0.4 mM H_2O_2 for 20 h (13). Cells were washed with phosphate-buffered saline (PBS) and incubated for 5 min in a solution supplemented with 100 ng/ml of 4',6'-diamidino-2-phenylindole (DAPI; Sigma Chemical, St. Louis, MO). After incubation with the dye, cells were washed with PBS buffer for 10 min at room temperature, and then the slides were mounted. Slides were viewed with a Carl Zeiss (Jena, Germany) microscope using a 100 $\times$ magnification oil immersion objective lens (EC Plan-Neofluar; numerical aperture, 1.3). In order to verify the accumulation of reactive oxygen species (ROS), cells were stained with 2',7'-dichlorodihydrofluorescein diacetate (H_2DCFDA). Yeast cells (10^7 cells ml^{-1}) were grown in 50-ml tubes in 10 ml of YPD medium for 9 or 16 h at 30°C and exposed to 0.125% propolis for 5 or 10 min. Cells were washed with YPD medium and incubated with prewarmed PBS buffer containing 10 μM H_2DCFDA (Invitrogen, Molecular Probes) for 30 min in the dark at 37°C . Slides were mounted, and images were analyzed by fluorescence microscopy at room temperature. Slides were viewed with a Carl Zeiss (Jena, Germany) microscope using a 100 $\times$ oil immersion objective lens (filter set 46 HE YFP; excitation wavelength, 500/20 nm; emission wavelength, 535/30 nm). Phase-contrast for the bright-field images and fluorescent images were captured with an AxioCam camera (Carl Zeiss), processed using the AxioVision software, version 3.1, and saved as TIFF files. Further processing was performed using Adobe Photoshop, version 7.0 (Adobe Systems Incorporated, CA).

In terminal deoxynucleotidyltransferase-mediated dUTP-biotin nick end labeling (TUNEL) assays, slides were visualized in a confocal microscope. Images were analyzed using a Leica TCS SP5 spectral laser scanning confocal microscope (Leica Microsystems, Heidelberg, Germany) at the Laboratory of Confocal Microscopy, Faculty of Medicine of Ribeirão Preto (FMRP), University of São Paulo, São Paulo, Brazil, using a 63 $\times$ water immersion objective lens, with laser line 488 nm for green fluorescent protein (GFP) and 405 nm for DAPI. Images were captured by direct acquisition, with the Z step ranging from 0.5 to 2 μm with the Leica LAS AF software (Leica Microsystems), and additional processing was carried out using Adobe Photoshop, version 7.0.

Induction of cell death. YPD liquid medium was inoculated with BY4742 cells for 16 h at 30°C . Grown cells were centrifuged and inoculated in 50 ml of fresh YPD liquid medium in 250-ml flasks to a final concentration of 5×10^5 cells ml^{-1} . Cells were then grown for 9 h (exponential; 2.0×10^7 cells ml^{-1}) and 16 h (stationary; 3.0×10^8 cells ml^{-1}). The cells in exponential and stationary phases were harvested and inoculated again in 50 ml of fresh YPD liquid medium in 250-ml flasks to a final concentration of 10^7 cells ml^{-1} and exposed to either 0.68% ethanol or 0.125% (vol/vol) alcoholic propolis extract. The treatment was carried out for 5, 10, and 20 min at 30°C with mechanical shaking (200 rpm). Then, the cells were washed with phosphate-buffered saline (PBS) to remove propolis, and the procedures were initiated for cell death detection. Cell viability was determined by plating appropriate cell concentrations and counting the number of grown colonies.

TUNEL assay. DNA strand breaks were demonstrated by TUNEL with an In Situ Cell Death Detection Kit, Fluorescein, from Roche. This procedure was according to Madeo et al. and Ribeiro et al. (47, 65) with some modifications. Yeast cells were fixed with 3.7% (vol/vol) formaldehyde for 30 min at room temperature and washed three times with PBS, and cell walls were digested with 15 U/ml lyticase, from *Arthrobacter luteus* (crude 100,000 U; 362 units/mg; Sigma) at 37°C for 60 min. Ten microliters of the cell suspensions was applied to a microscope slide and allowed to dry for 30 min at 37°C . The slides were rinsed with PBS, incubated in permeabilization solution (0.1% [vol/vol] Triton X-100 and 0.1% [wt/wt] sodium citrate) for 2 min on ice, and rinsed twice with PBS. For

TABLE 1. Specific gene ontology classes derived from PPPI network observed in cluster 6

Biological process	GO identification no.	<i>P</i> value ^a	Corrected <i>P</i> value ^b	<i>k</i> ^c	<i>f</i> ^d
Cell division	51301	2.95×10^{-14}	2.56×10^{-11}	36	323
Cell budding	7114	1.31×10^{-13}	5.68×10^{-11}	19	82
Asexual reproduction	19954	4.16×10^{-13}	1.20×10^{-10}	19	87
Reproduction of a single-celled organism	32505	6.45×10^{-13}	1.40×10^{-10}	19	89
Establishment of cell polarity	30468	1.23×10^{-12}	1.67×10^{-10}	20	104
Mitochondrial electron transport, ubiquinol to cytochrome ^c	6122	2.79×10^{-11}	2.11×10^{-9}	8	11
Cytokinesis	16288	2.91×10^{-11}	2.11×10^{-9}	19	109
Cell cycle	7049	2.13×10^{-9}	1.05×10^{-7}	40	566
Electron transport chain	22900	8.55×10^{-7}	1.61×10^{-5}	8	31
pH reduction	45851	2.63×10^{-5}	3.08×10^{-4}	6	24
Vacuolar acidification	7035	2.63×10^{-5}	3.08×10^{-4}	6	24
Negative regulation of transcription from RNA polymerase II promoter by glucose	433	3.30×10^{-4}	2.63×10^{-3}	3	6

^a *P* values were calculated by the hypergeometric distribution of one ontology class visualized in the network.

^b Calculated values based on *P* values obtained after FDR was applied.

^c Total number of proteins found in the network which belong to a gene ontology.

^d Total number of proteins that belong to a specific gene ontology.

a positive control the endonuclease DNase I (Sigma) was applied to the cells on microscope slides. The slides were placed in a humidified box for 1 h at 37°C and then washed twice in PBS. Slides were subsequently incubated with 10 μ l of TUNEL reaction mixture, containing terminal deoxynucleotidyltransferase and fluorescein isothiocyanate (FITC)-dUTP, for 60 min at 37°C. They were rinsed three times with PBS and incubated for 5 min in a solution supplemented with 100 ng/ml of DAPI (Sigma Chemical, St. Louis, MO). After incubation with the dye, they were washed with PBS buffer for 10 min at room temperature and then rinsed in distilled water, mounted, and visualized in a confocal microscope.

Annexin and PI assays. Phosphatidylserine exposure was detected by an annexin-V-Fluor staining kit (Roche), as described by Madeo et al. (48) with some modifications. Cells were harvested and washed with sorbitol buffer (1.2 M sorbitol, 0.5 mM MgCl₂, 35 mM K₂HPO₄, pH 6.8). Cell walls were digested with 15 U of lyticase (Sigma) in sorbitol buffer for about 30 min at 37°C. Cells were then washed twice with binding buffer (10 mM HEPES/NaOH, pH 7.4, 140 mM NaCl, 2.5 mM CaCl₂) containing 1.2 M sorbitol (binding-sorbitol buffer). To 38- μ l cell suspensions in binding-sorbitol buffer, 2 μ l of annexin V (Roche) and 2 μ l of a propidium iodide (PI) working solution (50 μ g/ml) were added, and the mixture was incubated for 15 min at room temperature. The cells were then washed and resuspended in binding-sorbitol buffer. Finally, the slides were mounted with the cell suspensions. Microscope settings and image acquisitions were as described above using epifluorescence microscopy. For quantitative assessment of annexin V-PI staining, at least 100 yeast cells were counted per sample. For the apoptosis positive control, the cells were treated with 80 mM acetic acid, pH 3.0, for 200 min (44), and for the necrosis positive control, the cells were fixed with fixative solution (3.7% formaldehyde, 50 mM sodium phosphate buffer, pH 7.0, 0.2% Triton X-100) for 30 min at room temperature.

Mitochondrial fractionation studies. For the extraction of mitochondrial proteins, mitochondria were isolated from *S. cerevisiae* cells based on the a previously described procedure (23, 64). Cells grown for 16 h in YPGal liquid (at 30°C), either untreated or exposed to 0.125% propolis for 10 min at 30°C, were harvested at $3,500 \times g$ for 5 min and were washed once with 1.2 M sorbitol. Then, washed cells were suspended in digestion buffer at a concentration of 10 g of cells (wet weight) in 50 ml of buffer (1.2 M sorbitol, 0.06 M phosphate, pH 7.5, 1 mM EDTA, 15 mM β -mercaptoethanol, and 50 mg Zymolyase 20T [MP Biomedicals]) and incubated at 37°C for 2 h to be converted to spheroplasts. Spheroplasts were washed with 1.2 M sorbitol and then gently homogenized in a resuspension buffer (0.6 M sorbitol, 20 mM HEPES, pH 7.4, and 0.5 mM phenylmethylsulfonyl fluoride [PMSF]) using an Elvehjem-Potter homogenizer. The homogenate was centrifuged at $2,500 \times g$ for 5 min at 4°C. The supernatant was centrifuged again at $2,500 \times g$ for 5 min at 4°C two additional times, and then it was centrifuged at $12,000 \times g$ for 10 min to sediment mitochondria. The supernatant (cytoplasmic portion) was collected, and the mitochondrial pellet was resuspended gently with resuspension buffer and washed four additional times at $2,000 \times g$ for 5 min at 4°C. The final mitochondrial pellet was resuspended gently in 0.6 M sorbitol–20 mM HEPES. Protein concentration was determined by Biuret assay (43) using bovine serum albumin (BSA) as a standard.

Samples were then analyzed by 10% sodium dodecyl sulfate-polyacrylamide

gel electrophoresis (SDS-PAGE). Samples were prepared for SDS-PAGE by the addition of 1 $\times$ sample buffer (62.5 mM Tris-HCl, pH 6.8, 2% SDS, 10% glycerol, 5% β -mercaptoethanol, and 5% bromophenol blue) and heating at 100°C for 3 min. Twenty micrograms of total protein from each sample was loaded into each lane of a 10% SDS-PAGE gel. After separation of the proteins, the gel was blotted onto a pure nitrocellulose membrane (0.2- μ m pore size; Bio-Rad), and after being blocked in 5% dried milk in TBS-T buffer (10 mM Tris-HCl, 150 mM NaCl, pH 8.0, and 0.05% Tween 20), the membrane was probed with the antibody rabbit anti-cytochrome *c* (CNAT against native cytochrome *c* from yeast; Sigma) at a 1:200 dilution in TBS-T buffer for 1 h at room temperature. The membrane was washed four times for 5 min each with TBS-T buffer and then incubated with a 1:5,000 dilution of goat anti-rabbit IgG peroxidase-labeled antibody (KPL) for 1 h. After being washed, the blot was developed by use of the SuperSignal Ultra chemiluminescence detection system (Pierce) and recorded by the use of Hyperfilm ECL (Amersham Biosciences).

Oxygen consumption. Mitochondrial and spheroplast oxygen consumption was monitored on a computer-interfaced Clark-type electrode at 25°C with 1 mM NADH as a substrate in the presence of mitochondria at 100 μ g/ml protein concentration. Mitochondria were isolated from *S. cerevisiae* cells grown in YPGal medium. Propolis was added to a 0.125% final concentration. In order to block cytochrome *c* oxidase respiration, 1 mM KCN was added at the end.

PPPI network design and topological analysis. The survival data gathered from yeast deletion strains submitted to propolis treatment was used to obtain information about how the deleted genes and their products interact in the context of physical protein-protein interaction (PPPI) networks in *S. cerevisiae*. To this end, data mining screening and network design of PPPI networks were performed using Cytoscape software, version 2.6.3 (74). For this purpose, we used the PPPI data of *S. cerevisiae* available in the STRING database, version 8.3 (<http://string.embl.de>), using the following parameters: active prediction methods all enabled except text mining; no more than 50 interactions; medium confidence score (0.400); and network depth equal to 1 with addition of new nodes until saturation of the network. The PPPI networks obtained from this first screening were then combined in a unique PPPI network by employing the union function of the Cytoscape core plug-in Merge Networks (Table 1). This major PPPI network was then analyzed with molecular complex detection (MCODE) software (3), a Cytoscape plug-in (<http://www.cytoscape.org/plugins/mcode>) in order to detect subnetworks or clusters of proteins that could represent distinct biologic processes. The parameters used for MCODE to generate the subnetworks were as follows: loops included; degree cutoff of 2; deletion of single connected nodes from cluster (haircut option enable); expansion of cluster by one neighbor shell allowed (fluff option enable); node density cutoff of 0.1; node score cutoff of 0.2; *k*-core of 2; and maximum depth of network of 100.

GO analysis. Gene ontology (GO) clustering analysis was performed using Biological Network Gene Ontology (BiNGO) (50) software, a Cytoscape plug-in available at http://chianti.ucsd.edu/cyto_web/plugins/index.php. The degree of functional enrichment for a given cluster and category was quantitatively assessed (*P* value) by hypergeometric distribution (66), and a multiple test correction was applied using the false-discovery rate (FDR) (5) algorithm, fully imple-

mented in BiNGO software. Overrepresented biological process categories were generated after FDR correction, with a significance level of 0.05.

RESULTS

Propolis induces cell death in *S. cerevisiae*. Propolis is a complex product derived from plant resins and bee saliva. There are several chemical compounds present in this natural product that could potentially be responsible for its antibiotic properties. As a first step to understand whether individual components of propolis could play such role, we fractionated propolis extracts by HPLC and purified some of the most abundant chemical compounds (see Fig. S1 and Table S1 in the supplemental material). We have obtained similar fingerprints for three different batches of propolis (see Fig. S2). We tested *S. cerevisiae* growth in the presence of four of these chemical compounds (caffeic, *p*-coumaric, and cinnamic acids and isosakuranetin). We applied 1, 2, 5, 10, 25, 50, and 100 $\mu\text{g/ml}$ of each compound and a mix of 100 $\mu\text{g/ml}$ of each in microtiter plate liquid cultures. None of these treatments was able to inhibit the growth of *S. cerevisiae* BY4742, *Candida albicans* CAI4, or *Candida parapsilosis* ATCC 2201 (data not shown). Thus, taking into consideration the fact that the cell death effects of propolis could be due to a combination of chemical compounds and concentrations, we decided to investigate the cell death effects of propolis by concentrating our experiments on alcoholic extracts of propolis.

The first series of experiments was related to verifying a working propolis concentration and determining how yeast viability was affected. Since propolis is dissolved in ethanol (60% alcoholic extract), the control treatment was grown on 2.73% ethanol because 0.5% propolis has a final concentration of about 2.73% ethanol (Fig. 1A). Tenfold dilutions of *S. cerevisiae* cells suggested that 0.125% propolis could be an adequate choice as a subinhibitory concentration (Fig. 1A). All the controls for further experiments using 0.125% propolis as a treatment have 0.68% ethanol (what corresponds to 60% alcoholic extract for 0.125% propolis). We were not able to perform the determination of MICs by using microdilutions because the color of the propolis is dark, which interferes in the spectrophotometer determinations (data not shown). The inhibitory effects of propolis on *S. cerevisiae* are not affected by growth on lower oxygen concentrations, such as 1% O_2 (data not shown). We also observed that *C. albicans* was more susceptible to concentrations of propolis between 0.50 and 0.75% (Fig. 1A). However, *C. parapsilosis* and *Aspergillus fumigatus* growth was not affected by the same concentrations of propolis (data not shown). Yeast cells in the initial mid-exponential (6 h; 1.2×10^6 cells ml^{-1}) and stationary (16 h; 3.0×10^8 cells ml^{-1}) phases are more resistant to 0.125% propolis extracts while cells in the exponential phase (9 h; 2.0×10^7 cells ml^{-1}) are much more sensitive (Fig. 1B). In the next step, we investigated what was the main mechanism of cell death in the cells in the exponential and stationary phases, i.e., 9 and 16 h of growth, respectively, by observing the number of cells which are propidium iodide (PI) positive (PI^+) or annexin and TUNEL positive (preliminary indications of necrosis or apoptosis, respectively).

Early apoptosis is characterized by an increased number of annexin V-positive (A^+) cells and PI negative (PI^-) cells while

in late apoptosis (leading to secondary necrosis), there is an increase in the number of A^+ and PI^+ cells. The primary necrosis is characterized by an increased number of annexin V-negative (A^-) cells and PI^+ cells. We have observed about 15% A^+ and PI^+ cells in the negative control (0.68% ethanol [EtOH]) while 100% A^+ and PI^+ cells in the positive control (acetic acid, pH 3.0, for 200 min). Upon 5 min of exposure to 0.125% propolis, cells grown for 9 and 16 h have an increased number of A^+ cells (about 30% both A^+ and PI^+ and 80% A^+), suggesting early apoptosis (Fig. 1B to D). Upon 10 min of exposure to 0.125% propolis, cells grown for 9 and 16 h have an increased number of A^+ and PI^+ (about 80% for both classes), suggesting late apoptosis, leading to secondary necrosis (Fig. 1B and C). However, the TUNEL-positive staining remained constant for both 5 and 10 min of exposure to 0.125% propolis (about 80%) (see Fig. S3 in the supplemental material).

Nuclear release of mammalian high-mobility group box-1 protein (HMGB1) is a defining feature of necrosis in mammalian cells (69). Since apoptotic cells can become leaky during further cultivation, resulting in PI positives, we determined the nucleo-cytosolic translocation of the yeast HMGB1 homologue Nhp6Ap in cells grown for 16 h exposed to 0.125% propolis for 5 and 10 min. Fluorescence microscopy showed the nucleo-cytosolic translocation of the GFP-tagged Nhp6Ap (Nhp6Ap-GFP) in 20 and 30% of the cells after exposure for 5 and 10 min to 0.125% propolis, respectively (Fig. 1E). Taken together, these results indicate that propolis is able to induce an apoptosis cell death response but that increased exposure to propolis provides a corresponding increase in the necrosis response.

There is an accumulation of mitochondrion-produced reactive oxygen species (ROS) during yeast apoptosis (for reviews, see references 19, 58, and 62). To investigate the production of ROS during propolis-induced cell death in *S. cerevisiae*, we used 2',7'-dichlorodihydrofluorescein diacetate (H_2DCFDA), a cell-permeable ROS indicator that is nonfluorescent until acetate groups are removed (H_2DCF) by intracellular esterases and oxidation occurs within the cell. Although H_2DCF can be oxidized by different ROS, it is mainly used to detect superoxides. Yeast cells grown for 9 or 16 h were either untreated or exposed to 0.125% propolis for 5 and 10 min, incubated with H_2DCFDA , and examined by fluorescence microscopy (Fig. 2). The DCFDA-dependent green fluorescence was detected in about 10% of the 9- or 16 h-grown cells not exposed to propolis. Treatment of 9-h-grown cells with 0.125% propolis for 5 and 10 min yielded 70 to 80% fluorescence while treatment of 16-h-grown cells treated for 5 and 10 min produced 40 to 60% fluorescence (Fig. 2). These results indicate that there is a great accumulation of ROS during cell death induced by propolis.

Caspases are members of a family known as cysteine proteases that are actively involved in cell death in eukaryotes (for a review, see reference 41). Although *S. cerevisiae* does not have caspases in its genome, a caspase-like protein named YCA1, fitting into the type I category of metacaspases, was identified and characterized previously (49, 54, 81). It has been shown that YCA1 is involved in several cell death stimuli (for a review, see reference 54). When yeast cells grown for 16 h were exposed to 0.125% propolis for 5, 10, and 20 min, there

FIG. 1. Propolis induces apoptosis and late necrosis in *S. cerevisiae*. (A) Spot dilution assays for *S. cerevisiae* and *C. albicans* in YPD medium supplemented with different concentrations of propolis. The plates were incubated at 30°C for 72 h. (B) Viability of yeast cells in different growth cycle phases exposed to 0.125% propolis for 5 or 10 min. Cell viability was determined by plating appropriate cell concentrations and counting the number of colonies. The numbers are the average $\pm$ standard deviation of three independent experiments. (C) Annexin and PI assays for 16-h yeast cells exposed to 0.125% propolis. (D) Total numbers for annexin- and PI-positive cells for yeast cells exposed to 0.125% propolis. Bar, 5 μ m. The numbers are the average $\pm$ standard deviation of three independent experiments. In each experiment, 100 cells were assessed for annexin and PI staining. Positive controls for annexin and PI were exposing the cells to acetic acid, pH 3.0, for 200 min and fixing them with fixation solution as described in the Materials and Methods section. (E) Nucleo-cytosolic translocation of Nhp6Ap-GFP in untreated *S. cerevisiae* cells or cells exposed to 0.125% propolis for 10 min. Bar, 5 μ m.

was a reduction in the cell viability of 40, 70, and 95% in the wild-type strain while in the *YCA1* Δ strain, this reduction was about 10, 50, and 80% (Fig. 3). These results strongly indicate that the *YCA1* gene is important for propolis-mediated cell death.

Cytochrome *c* is released during propolis-induced cell death. We have also investigated if other components of the cell death machinery are being activated during propolis-induced cell death in *S. cerevisiae*. We concentrated our attention on two of these putative determinants: *CYC1* and *NUC1*.

FIG. 2. Propolis induces the production of ROS. Yeast cells were grown for 9 or 16 h at 30°C in YPD medium and stained with H_2DCFDA for 30 min at 37°C before fluorescent microscopy analysis. Bar, 5 μm .

In mammalian cells, at least two major apoptotic pathways have been described: (i) the intrinsic pathway that needs the involvement of the mitochondria, and (ii) the extrinsic pathway where mitochondria are bypassed and caspases are activated (for a review, see reference 21). One of the hallmarks of the intrinsic pathway is the release of apoptogenic factors such as cytochrome *c* to the cytosol and the consequent assembly organized by this protein of the high-molecular-weight complex, the mitochondrial apoptosome, that activates caspases (for a review, see reference 21). In *S. cerevisiae* cells undergoing an

apoptotic process induced by acetic acid, translocation of cytochrome *c* to the cytosol was observed (45). We performed cell fractionation studies to determine if cytochrome *c* was translocated to the cytoplasm during exposure of yeast cells grown for 16-h (16-h yeast cells) to 0.125% propolis for 10 min (Fig. 4). When the yeast cells were not exposed to propolis, all the cytochrome *c* was located in the mitochondrial protein extracts while upon propolis addition, there was a significant presence of cytochrome *c* in the cytoplasm protein extracts (Fig. 4, right panel). There are corresponding protein amounts in each cell fraction extract, as shown by Coomassie staining (Fig. 4, left panel).

Endonuclease G (EndoG) has been described as a mitochondrial endonuclease that digests both DNA and RNA (12, 13, 61). Upon apoptosis induction, translocation of mammalian EndoG and its *S. cerevisiae* homologue *NUC1* to the nucleus coincides with large-scale DNA fragmentation (12, 13, 41, 61). We have not observed increased sensitivity of *NUC1* Δ yeast cells to propolis (data not shown). Next, we examined if Nuc1p would translocate to the nucleus when yeast cells were exposed to propolis (Fig. 5). When 16-h yeast cells were not exposed to propolis, the Nuc1p-GFP showed a diffuse mitochondrion filament-like distribution, as confirmed by MitoTracker staining (Fig. 5, first row). In yeast cells exposed to hydrogen peroxide, part of the Nuc1p-GFP was translocated to the nucleus (DAPI staining), as previously shown by Büttner et al. (13), but it also accumulated in the mitochondria as indicated by MitoTracker staining (Fig. 5, second row). In contrast, when yeast cells were exposed to 0.125% propolis for 15 min, about 100% of the yeast cells showed a Nuc1p-GFP dot-like distribution, suggesting mitochondrial fragmentation, as again confirmed by MitoTracker staining (Fig. 5, third row). Moreover, the Nuc1p-GFP did not translocate to the nuclei, as shown by DAPI staining (Fig. 5, third row). These results strongly indicate that cytochrome *c*, but not Nuc1p, is involved in propolis-mediated cell death in *S. cerevisiae*.

FIG. 3. Metacaspase *YCA1* mutant is more tolerant to propolis than the corresponding wild-type strain. Viability of 16-h yeast wild-type and *YCA1* mutant strains treated with 0.125% propolis (*, $P < 0.05$) was determined by plating appropriate cell concentrations and counting the number of colonies. The results are the means $\pm$ standard deviation of three independent experiments.

FIG. 4. Exposure of yeast cells to propolis induces cytochrome *c* release into the cytoplasm. Yeast cells were grown for 16 h at 30°C and left untreated or exposed to 0.125% propolis for 10 min at 30°C. The cells were harvested and fractionated (mitochondria and cytoplasm) as described in the Materials and Methods section, and proteins were run on a polyacrylamide gel. At left is a Coomassie-stained gel with the same amount of proteins that were transferred to a nitrocellulose filter (right panel). This membrane was probed with the antibody rabbit anti-cytochrome *c*.

High-throughput screen of the yeast deletion library for propolis sensitivity. To elucidate the gene functions that may be required for propolis sensitivity in eukaryotes, the full collection of haploid *S. cerevisiae* deletion strains was screened for growth on YPD plates supplemented with 2.73% ethanol (as a

control; 2.73% is the ethanol concentration used to dissolve 0.500% propolis) and 0.250 and 0.500% propolis for 5 days at 37°C. About approximately 4,800 different strains were involved in this screening, and after three confirming screenings, we were able to identify 138 deletion strains that have different

FIG. 5. Nuc1p is not involved in propolis sensitivity. Yeast Nuc1-GFP cells were grown for 16 h at 30° and left untreated or exposed to 0.125% propolis for 15 min, and cells were visualized by a fluorescence microscope. A positive control was made by incubating 16-h-old yeast cells with 0.4 mM H₂O₂ for 20 h at 30°C. Bar, 5 μm. DIC, differential interference contrast. Arrows indicate the nuclei.

FIG. 6. System biology for the propolis sensitivity in *S. cerevisiae*. (A) A physical protein-protein interaction (PPPI) network obtained from yeast deletion data whose absence increases the sensitivity of yeast cells to propolis treatment. (B) Distribution of proteins (number of nodes) associated with propolis-induced sensitivity as observed from yeast deletion data in the different clusters and unclustered proteins of the major PPPI network. Cluster 1, translation/energy derivation by oxidation of organic compounds/mitochondrial genome maintenance; cluster 2, transcription/chromosome organization and biogenesis/ G_1 phase of mitotic cell cycle; cluster 3, carboxylic acid metabolic process/protein targeting to peroxisome/respiration metabolism; cluster 4, establishment and/or maintenance of chromatin architecture/histone deacetylation; cluster 5, RNA catabolic process; cluster 6, cell division mechanisms; and unclustered nodes.

degrees of propolis sensitivity compared to the corresponding wild-type strain (see Table S2 in the supplemental material). The survival data obtained from yeast deletion strains prompted us to ask how the absence of the determined proteins of propolis-sensitive yeast strains affects different biological processes that lead to cell death. In this sense, a search for potential proteins and/or mechanisms and their associated biological processes that are affected by propolis exposure was initiated. To achieve this goal, different PPPI networks using yeast deletion data were retrieved from the STRING database. Shared proteins and subnetworks present in the major PPPI network (Fig. 6) were identified and retrieved using the Cytoscape-associated plug-in MCODE and subjected to a Gene Ontology (GO) analysis in order to obtain information about the nature and number of subnetworks belonging to the network and their associated biological processes. Results obtained from MCODE and GO analysis showed that the final PPPI network (Fig. 6A) contains 553 nodes and 3,651 connectors and is composed of six heavily interconnected clusters, each comprising different biological processes. GO analyses revealed that these biological processes can be classified into the following categories: (i) translation/energy derivation by oxidation of organic compounds/mitochondrial genome maintenance, (ii) transcription/chromosome organization and biogenesis/ G_1 phase of mitotic cell cycle, (iii) carboxylic acid metabolic process/protein targeting to peroxisome/respiration metabolism, (iv) establishment and/or maintenance of chromatin architecture/histone deacetylation, (v) RNA catabolic process, and (vi) cell division mechanisms. As expected, proteins that could not be classified into any cluster

were also identified in the network (unclustered protein subnetworks) (Fig. 6A and B).

The distribution analysis of the proteins whose deletion increases the sensitivity of yeast strains to propolis indicated that both cluster 6 and unclustered protein subnetworks contain 61.73% of the total proteins observed in the deletion assay (Fig. 6B; see also Table S3 in the supplemental material). The other 38.27% proteins were found distributed among clusters 1 to 5 (Fig. 6B; see also Table S3), showing that cluster 6 and the unclustered protein subnetworks appear to be important for the pleiotropic responses observed when yeast cells are submitted to propolis treatment. When the proteins of the cluster 6 subnetwork (Fig. 7; see also Table S3) were subjected to GO analysis, the data indicated that the major biological processes found within this subnetwork were associated with cell division and reproduction, which includes the establishment of cell polarity, cell cycle, and budding (Table 1). Moreover, the proteins of the cluster 6 subnetwork appear to be important to the mitochondrial electron transport chain, vacuolar acidification, and negative regulation of transcription from RNA polymerase II promoter (Table 1). Interestingly, the GO analysis of the unclustered protein subnetworks (Fig. 7B; see also Table S3) showed that these proteins are important for the regulation of macroautophagy associated with protein targeting to vacuoles and the cellular response to starvation, among other processes (Table 2).

In the next two sections, we show validation data about the role played by clusters related to the mitochondrial electron transport chain and vacuolar acidification.

FIG. 7. (A) Subnetwork of proteins (cluster 6) associated with cell division, establishment of cell polarity, electron transport chain, and vacuolar acidification. (B) Unclustered protein subnetworks associated with macroautophagy, protein targeting to vacuoles, cellular response to starvation and stress, vacuolar transport, cell communication, and regulation of protein polymerization.

Petite strains are less sensitive to propolis. It has recently been shown that during necrosis in *S. cerevisiae*, there is an inhibition of mitochondrial function with a corresponding reduction of ATP production (for a review, see reference 19). In this paper, there are several indications that propolis induces a secondary necrosis cell death mechanism in *S. cerevisiae*. In addition, as demonstrated in the previous section, we found that when several genes related to energy derivation by oxidation of organic compounds, mitochondrial genome maintenance, and the mitochondrial electron transport chain are deleted, there is an increase in *S. cerevisiae* propolis sensitivity. We decided to investigate a possible correlation between propolis sensitivity, mitochondria and *S. cerevisiae* necrotic cell death by isolating petite strains of *S. cerevisiae* and checking their sensitivity to propolis. In order to isolate BY4742 [*rho*⁰]/[*rho*⁻] cells, the respiratory-competent strain BY4742 (10) was mutagenized with ethidium bromide, and the resultant petite respiratory-deficient colonies were crossed to [*rho*⁰]/[*rho*⁻] tester strains (80) to confirm the nature of the isolated petite

mutant respiratory incapacity. Two of these petite strains, BY4741A [*rho*⁰]/[*rho*⁻] and BY4742 [*rho*⁻]/[*rho*⁰], and the corresponding wild-type strain were exposed to 0.125% propolis for 20 min, and cell viability was determined (Fig. 8). The petite strains showed dramatic increased viability compared to the wild-type strain (compare 95 and 75% viability of BY4741A [*rho*⁻]/[*rho*⁰] and BY4742 [*rho*⁻]/[*rho*⁰] strains, respectively, against about 10% viability in the wild-type strain) (Fig. 8A). Accordingly, when cell protein extracts were fractionated, cytochrome *c* could not be observed in cytoplasmic protein extracts of petite cells exposed to propolis but only in mitochondrial protein extracts (Fig. 8B). We measured the effect of propolis on mitochondria isolated from yeast strain W303 by monitoring the oxygen consumption rate as a consequence of mitochondria NADH oxidase activity. In the presence of propolis the NADH oxidase activity is 1/10 of that observed before the addition of this compound, indicating its toxic effect on respiration (Fig. 8C). These results strongly suggest that *S. cerevisiae*

TABLE 2. Specific GO classes derived from PPPI of unclustered protein subnetworks

Biological process	GO identification no.	<i>P</i> value ^a	Corrected <i>P</i> value ^b	<i>k</i> ^c	<i>f</i> ^d
Macroautophagy	34262	7.44 × 10 ⁻¹⁴	5.44 × 10 ⁻¹¹	11	23
CVT pathway	32258	9.36 × 10 ⁻¹²	3.43 × 10 ⁻⁹	9	18
Protein targeting to vacuole	6623	7.37 × 10 ⁻¹¹	1.80 × 10 ⁻⁸	13	62
Cellular response to starvation	9267	3.03 × 10 ⁻¹⁰	5.54 × 10 ⁻⁸	11	44
Cellular response to stress	33554	1.06 × 10 ⁻⁹	5.09 × 10 ⁻⁷	11	49
Cellular response to nutrient levels	31669	4.86 × 10 ⁻⁹	5.09 × 10 ⁻⁷	11	56
Response to starvation	42594	1.80 × 10 ⁻⁸	1.46 × 10 ⁻⁶	11	63
Vacuolar transport	7034	5.05 × 10 ⁻⁸	3.69 × 10 ⁻⁶	14	122
Cell communication	7154	1.44 × 10 ⁻⁶	6.21 × 10 ⁻⁵	20	317
Regulation of protein polymerization	32271	1.76 × 10 ⁻⁶	6.79 × 10 ⁻⁵	4	6

^a *P* values calculated by the hypergeometric distribution of one ontology class visualized in the network.
^b Calculated values based on *P* values obtained after FDR was applied.
^c Total number of proteins found in the network which belong to a gene ontology.
^d Total number of proteins that belong to a specific gene ontology.

FIG. 8. Petite mutants are more tolerant to propolis. (A) Viability of 16-h yeast wild-type and petite mutant strains to 0.125% propolis (*, $P < 0.05$) was determined by plating appropriated cell concentrations and counting the number of colonies. The results are the means $\pm$ standard deviations of three independent experiments. (B) Exposure of yeast petite mutant cells to propolis does not induce cytochrome *c* release into the cytoplasm. Yeast cells were grown for 16 h at 30°C and left untreated or exposed to 0.125% propolis for 10 min at 30°C. The cells were harvested and fractionated (mitochondria and cytoplasm) as described in the Materials and Methods section, and proteins were run on a polyacrylamide gel. At left is a Coomassie-stained gel with the same amount of proteins that were transferred to a nitrocellulose filter (right panel). This membrane was probed with the antibody rabbit anti-cytochrome *c*. (C) NADH oxidase activity in wild-type yeast in the presence of propolis. Mitochondria (100 μ g of protein) of the wild-type W303 strain were assayed for NADH-oxidase with a Clark electrode as described in the Materials and Methods section. Arrows indicate the addition of propolis (final concentration of 0.125%) or the identical volume of ethanol and KCN. These are representative traces of one experiment, which were reproduced with mitochondria isolated on three different days.

propolis sensitivity is dependent on the mitochondrial function.

A role for the vacuole and the vacuolar H^+ -ATPase in propolis sensitivity. Another set of genes important for propolis sensitivity observed in our screening were involved in vacuolar acidification and macroautophagy associated with protein targeting to vacuoles and cellular response to starvation (Table 2). Organelle acidification is implicated in protein sorting in the biosynthetic and endocytic pathways, proteolytic activation of zymogen precursors, and transmembrane transport of viral contents and toxins (for a review, see reference 32). In fungi, the lysosome-like vacuole also plays a role in storage of metabolic building blocks, calcium homeostasis, and osmotic control. Vacuolar acidification is essential for all these functions. The vacuolar proton-translocating ATPase (V-ATPase) is an essential enzyme to catalyze all these processes. Their primary role in eukaryotic cell is ATP-driven transport of protons from the cytosol into acidic organelles (32). When several genes (*VPH1*, *VMA3*, *VMA4*, *VMA5*, *VMA11*, *VMA22*,

RAV1, and *SOP4*) involved in the assembly of the yeast V-ATPase were deleted, the corresponding yeast deletion strains became more sensitive to propolis (Fig. 9A). Recently, Rosado et al. (67) developed a method for monitoring autophagy using Rosella, a biosensor comprised of a fast-maturing pH-stable red fluorescent protein fused to a pH-sensitive green fluorescent protein variant. Its mode of action relies upon differences in pH between different cellular compartments and the vacuole. Rosella is a genetically encoded dual-color-emission biosensor, a product of a fusion of a relatively pH-insensitive fast-maturing variant of the red fluorescent protein, DsRed (6) with a pH-sensitive variant of GFP (56). The rationale of this assay relies on the pH difference between the vacuole (or lysosome; approximately pH 6.2 or 4.8, respectively) and other compartments of the cell (pH >7.0). The reporter is a dual-color emission biosensor comprising a relatively pH-stable DsRed and a pH-sensitive GFP variant (67). In yeast cells grown for 16 h, both red and green fluorescence were evenly distributed throughout the cytosol and absent from the vacuole

FIG. 9. Propolis induces vacuolar acidification in *S. cerevisiae*. (A) Spot dilution assays for YPD medium supplemented with different concentrations of propolis. Yeast cells were grown for 96 h at 30°C. (B) Rosella expressed in the yeast cytosol is delivered to the vacuole under cell exposure to propolis. DIC and fluorescence images are shown for wild-type cells under growing conditions (first row), after nitrogen starvation for 4 h (second row), and exposed to 0.125% propolis for 90 min at 30°C. The position of the vacuole in the DIC images is delineated by a dashed white line. Empty vacuoles are highlighted by white arrows. Filled vacuoles are highlighted by yellow arrows. Bar, 5 μ m.

(Fig. 9B, first row). Cells were then transferred to starvation medium (without nitrogen) to induce autophagy. After a 4-h incubation, red fluorescence, but not green, was observed to accumulate in the vacuole of approximately 40% of the cells (Fig. 9B, second row). Then, the cells were exposed to 0.125% propolis, and after a 90-min incubation, a subtle phenotype of red fluorescence, but not green, was observed to accumulate in the vacuole of approximately 60% of wild-type cells (Fig. 9B, second and third rows).

Autophagy can be involved in the turnover of long-lived proteins and whole organelles (for example, mitochondria in mitophagy and the endoplasmic reticulum in reticulophagy) (51). Autophagy is also an important process during starvation; i.e., by the catabolism of macromolecules autophagy generates metabolic substrates (51). Autophagy has been followed by various biochemical and morphological methods (for reviews, see references 35 and 36). Expression of GFP-Atg8p in yeast can be used to follow the localization or accumulation of the preautophagosomal structure, autophagosomes, and autophagic bodies. We have observed that *ATG1* (encoding a protein serine/threonine kinase required for vesicle formation in autophagy and the cytoplasm-to-vacuole targeting [CVT] pathway), *ATG10* (encoding a conserved E2-like conjugating en-

zyme that mediates formation of the Atg12p-Atg5p conjugate, which is a critical step in autophagy), and *ATG20* (encoding a nexin family member required for the CVT pathway and for endosomal sorting) deletion strains are more sensitive to propolis (for reviews about yeast genes involved in autophagy, see references 33 and 39) (Fig. 10A). We have used mRNA accumulation of *ATG14* and *ATG8* and monitoring GFP-Atg8 as assays to verify if propolis induces an increase in autophagy. Propolis induced *ATG8* and *ATG14* mRNA accumulation about 4 times and 14 times, respectively (5- and 10-min exposure to 0.125% propolis) (Fig. 10B). The GFP-Atg8 translocated to vacuolar-like structures after exposure to 0.125% propolis for 20 min (Fig. 10C).

Taken together, these data strongly indicate that vacuolar acidification and autophagy are important for yeast cell death caused by propolis.

DISCUSSION

The three main types of propolis around the world have different compositions: (i) caffeic acid phenethyl ester (CAPE)-based propolis in Europe, the Far East and New Zealand; (ii) artemillin C (ARC)-based Brazilian green propolis;

FIG. 10. Propolis induces autophagy in *S. cerevisiae*. (A) Spot dilution assays for YPD medium supplemented with different concentrations of propolis. Yeast cells were grown for 96 h at 30°C. (B) Yeast wild-type cells were grown for 9 h in liquid YPD medium at 30°C and left untreated or exposed to 0.125% propolis for 5 or 10 min. Total RNA was extracted, and real-time reverse transcription-PCR (RT-PCR) was used to quantify mRNA accumulation. The measured quantity of the mRNA in each of the treated samples was normalized using the threshold cycle (C_T) values obtained for the actin (*ACT1*) mRNA amplifications run in the same plate. The relative quantitation of *ATG8*, *ATG14*, and *ACT1* gene expression was determined by a standard curve (i.e., C_T values plotted against logarithm of the DNA copy number). The results are the means $\pm$ standard deviations of four sets of experiments. (C) Cells of yeast strains harboring GFP-Atg8 either untreated or exposed to 0.125% propolis for 20 min at 30°C. Bar, 5 μ m.

and (iii) Brazilian red propolis. Recently, a screen of 1,266 compounds with known pharmaceutical activities was performed, and 15 compounds that prolonged survival of *C. albicans*-infected nematodes and inhibited *in vivo* filamentation of *C. albicans* were identified (11). One of these compounds, CAPE, exhibited antifungal activity in a murine model of candidiasis (11). CAPE is a poplar-type propolis while Brazilian green propolis is a *Baccharis* type. We along with other investigators have been unable to identify any traces of CAPE in the Brazilian green propolis (75, 79; also data not shown). Here, we have investigated molecular targets involved in cell lethality caused by the Brazilian green propolis. We were not able to see any cell death effect when *S. cerevisiae* was exposed to high concentrations of four of the major compounds that are present in the green propolis. Thus, we decided to verify the effects of propolis on *S. cerevisiae* cell lethality by using alco-

holic propolis extracts. Cell death in *S. cerevisiae* is frequently followed by indicative apoptotic and/or necrotic markers such as externalization of phosphatidylserine to the outer leaflet of the plasma membrane, chromatin condensation, the generation of reactive oxygen species, propidium iodide accumulation, HMGB1/Nhp6p localization, detection of plasma membrane ruptures, and complete disintegration of subcellular structures (14, 19). In addition, ROS accumulation, mitochondrial fragmentation, cytochrome *c* release, the metacaspase Yca1p, the apoptosis-inducing factor Aif1p, the endonuclease G Nuc1p, the serine protease OMI, cytoskeleton perturbations, and chromatin epigenetic modification have also been observed (79). Necrosis has also been described during yeast chronological aging (20).

We have noticed that apoptosis and necrosis markers are induced by propolis, but increased time exposure to propolis intensifies the number of cells with necrotic markers, such as PI and nucleo-cytosolic translocation of the Nhp6Ap-GFP. Our data indicate a dual role for propolis treatment as an agent that induces apoptosis and secondary necrosis. These effects are partially mediated by *YCA1* metacaspase since the *YCA1* Δ mutant is more resistant to propolis. In *S. cerevisiae* overexpression of *YCA1* caused cell death while *YCA1* deletion protected against cell death caused by reactive oxygen species or chronological aging (49). Additionally, we have also observed that Nuc1p does not translocate to the nucleus upon propolis cell death induction. Interestingly, Büttner et al. (13) observed that apoptotic death mediated by Nuc1p does not require *YCA1*. More interestingly, simultaneous markers for apoptosis and necrosis have also been noticed during propolis cell death induction in *Trypanosoma cruzi* (55).

There are several conditions where mitochondria-produced ROS have been associated with yeast apoptosis (for reviews, see references 19, 59, and 62). Propolis can induce ROS formation, and it is more lethal when *S. cerevisiae* grows in the presence of glycerol and ethanol as carbon sources (data not shown), suggesting that respiration increases propolis lethality. Actually, we were able to demonstrate that propolis can inhibit respiration in *S. cerevisiae*. We have observed that propolis cell death induction stimulates the cytochrome *c* release and that also [*rho*⁰] cells are more tolerant to propolis. Cytochrome *c* is a mitochondrial protein with a function in the respiratory chain and an additional function as an activator of caspase-9 in the intrinsic pathway of mammalian apoptosis (40). Treatment of yeast cells with acetic acid leads to mitochondrial cytochrome *c* release (45). Disruption of cytochrome *c* partially prevents acetic acid-induced cell death, and accordingly [*rho*⁰] cells show resistance against acetic acid-induced cell death (45). However, it is not very clear, as it is in mammals, if cytochrome *c* release can lead to the formation of an apoptosome-like structure and activation of caspases. In contrast, Sriprya et al. (77) have observed loss of mitochondrial membrane potential and absence of cytochrome *c* release during necrotic cell death of *S. cerevisiae* induced by expression of a proteinaceous elicitor hairpin (Pss) from *Pseudomonas syringae*. However, the deficiency of functional mitochondrial DNA, and consequent inability to respire from petite mutants, conferred resistance to death.

We have applied the power of genomics by screening the full collection of haploid *S. cerevisiae* deletion strains to compre-

hend how propolis, a complex phytotherapeutic compound, affects cell metabolism. This strategy has already been extensively used for understanding how other compounds affect *S. cerevisiae* cell metabolism (8, 15, 18, 27, 71, 86). By applying systems biology, we observed that most of the proteins whose deletion increases the sensitivity of yeast strains to propolis are involved in cell division mechanisms, the mitochondrial electron transport chain, vacuolar acidification, regulation of macroautophagy associated with protein targeting to vacuoles, cellular response to starvation, and negative regulation of transcription from the RNA polymerase II promoter. We have investigated and validated in more detail the mitochondrial electron transport chain and vacuolar acidification pathways. We have shown that propolis induces vacuolar acidification and translocation of Atg8p to the vacuoles, one of the hallmarks of autophagy. In *S. cerevisiae*, the vacuole is very important for maintaining cellular homeostasis comprising the regulation of intracellular pH and degradation mainly during nutrient limitation of proteins and organelles by autophagy (for reviews, see references 2, 29, and 46). Cell death induced by acetic acid is increased in *S. cerevisiae* *VPS* gene deletion mutants (*VPS* genes are involved in homotypic vacuole fusion and vacuolar protein sorting and are essential for normal vacuolar function) (70). It has been observed that the intracellular pH was acidified in *VPS* mutant cells upon treatment with acetic acid (70). It is possible that the disturbance of the homeostatic pH control may trigger necrosis by release of pro-necrotic proteases, which would find an optimal pH for their enzymatic activity in the acidified cytosol (19).

Our work emphasizes the importance of *S. cerevisiae* as a model system to understand at a molecular level the mechanism of action of a phytotherapeutic compound. There is a multitude of studies describing the most diverse effects of propolis on different biological systems, ranging from cancer chemoprevention (78) to cariostatic activity on streptococci (42) to an influence on immunomodulatory action and lymphocytes and antibody production (73). Our study is the first one that investigates systematically by using functional genomics how propolis influences the metabolism of an organism. This opens new possibilities for understanding and validating propolis as an alternative therapeutic agent against fungal infections.

ACKNOWLEDGMENTS

This research was supported by the Fundação de Amparo à Pesquisa do Estado de São Paulo and Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Brazil.

We thank R. J. Devenish for providing us with the yeast vectors containing Rosella, the four anonymous reviewers for their suggestions, and Rogelio Lopes Brandão for helping us with the yeast deletion library. We also thank D. J. Klionsky for providing the ATG8::GFP construct.

REFERENCES

1. Agüero, M. B., et al. 2010. Argentinean propolis from *Zuccagnia punctata* Cav. (Caesalpinieae) exudates: phytochemical characterization and antifungal activity. *J. Agric. Food Chem.* **58**:194–201.
2. Ariño, J., J. Ramos, and H. Sychrová. 2010. Alkali metal cation transport and homeostasis in yeasts. *Microbiol. Mol. Biol. Rev.* **74**:95–120.
3. Bader, G. D., and C. W. Hogue. 2003. An automated method for finding molecular complexes in large protein interaction networks. *BMC Bioinformatics* **4**:2.
4. Basnet, P., T. Matsuno, and R. Z. Neidlein. 1997. Potent free radical scavenging activity of propolis isolated from Brazilian propolis. *Z. Naturforsch. C* **52**:828–833.
5. Benjamini, Y., and Y. Hochberg. 1995. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J. R. Stat. Soc. B* **57**:289–300.
6. Bevis, B. J., and B. S. Glick. 2002. Rapidly maturing variants of the *Disco-soma* red fluorescent protein (DsRed). *Nat. Biotechnol.* **20**:83–87.
7. Bharucha, N., and A. Kumar. 2007. Yeast genomics and drug target identification. *Comb. Chem. High Throughput Screen.* **10**:618–634.
8. Blackburn, A. S., and S. V. Avery. 2003. Genome-wide screening of *Saccharomyces cerevisiae* to identify genes required for antibiotic insusceptibility of eukaryotes. *Antimicrob. Agents Chemother.* **47**:676–681.
9. Boone, C., H. Bussey, and B. J. Andrews. 2007. Exploring genetic interactions and networks with yeast. *Nat. Rev. Genet.* **8**:437–449.
10. Brachmann, C. B., et al. 1998. Designer deletion strains derived from *Saccharomyces cerevisiae* S288C: a useful set of strains and plasmids for PCR-mediated gene disruption and other applications. *Yeast* **14**:115–132.
11. Breger, J., et al. 2007. Antifungal chemical compounds identified using a *C. elegans* pathogenicity assay. *PLoS Pathog.* **3**:e18.
12. Burhans, W., and M. Weinberger. 2007. Yeast endonuclease G: complex matters of death, and of life. *Mol. Cell* **25**:323–325.
13. Büttner, S., et al. 2007. Endonuclease G regulates budding yeast life and death. *Mol. Cell* **25**:233–246.
14. Carmona-Gutierrez, D., et al. 2010. Apoptosis in yeast: triggers, pathways, subroutines. *Cell Death Differ.* **17**:763–773.
15. Chan, T.-F., J. Carvalho, L. Riles, and X. F. S. Zheng. 2000. A chemical genomics approach toward understanding the global functions of the target of rapamycin protein (TOR). *Proc. Natl. Acad. Sci. U. S. A.* **97**:13227–13232.
16. Costanzo, M., et al. 2010. The genetic landscape of a cell. *Science* **327**:425–431.
17. Dalben-Dota, K. F., et al. 2010. Antifungal activity of propolis extract against yeasts isolated from vaginal exudates. *J. Altern. Complement. Med.* **16**:285–290.
18. Desmoucelles, C., B. Pinson, C. Saint-Marc, and B. Daignan-Fornier. 2002. Screening the yeast “disruptome” for mutants affecting resistance to the immunosuppressive drug, mycophenolic acid. *J. Biol. Chem.* **277**:27036–27044.
19. Eisenberg, T., D. Carmona-Gutierrez, S. Büttner, N. Tavernarakis, and F. Madeo. 2010. Necrosis in yeast. *Apoptosis* **15**:257–268.
20. Eisenberg, T., et al. 2009. Induction of autophagy by spermidine promotes longevity. *Nat. Cell Biol.* **11**:1305–1314.
21. Elmore, S. 2007. Apoptosis: a review of programmed cell death. *Toxicol. Pathol.* **35**:45–516.
22. Ghisalberti, E. L. 1979. Propolis: a review. *Bee World* **60**:59–84.
23. Glick, B., and G. Schatz. 1991. Import of proteins into mitochondria. *Annu. Rev. Genet.* **25**:21–44.
24. Greenaway, W., T. Scaysbrook, and F. R. Whatley. 1990. The composition and plant origins of propolis. *Bee World* **71**:107–118.
25. Gullo, A. 2009. Invasive fungal infections: the challenge continues. *Drugs* **69**(Suppl. 1):65–73.
26. Gulshan, K., and W. C. Moye-Rowley. 2007. Multidrug resistance in fungi. *Eukaryot. Cell* **6**:1933–1942.
27. Hillenmeyer, M. E., et al. 2008. The chemical genomic portrait of yeast: uncovering a phenotype for all genes. *Science* **320**:362–365.
28. Hoon, S., R. P. St. Onge, G. Giaever, and C. Nislow. 2008. Yeast chemical genomics and drug discovery: an update. *Trends Pharmacol. Sci.* **29**:499–504.
29. Inoue, Y., and D. J. Klionsky. 2010. Regulation of macroautophagy in *Saccharomyces cerevisiae*. *Semin. Cell Dev. Biol.* **21**:664–670.
30. Ito, J., et al. 2001. Anti-AIDS agents. 48.(1) Anti-HIV activity of moronic acid derivatives and the new melliflorone-related triterpenoid isolated from Brazilian propolis. *J. Nat. Prod.* **64**:1278–1281.
31. Kanafani, Z., and J. Perfect. 2008. Antimicrobial resistance: resistance to antifungal agents: mechanisms and clinical impact. *Clin. Infect. Dis.* **46**:120–128.
32. Kane, P. 2006. The where, when, and how of organelle acidification by the yeast vacuolar H⁺-ATPase. *Microbiol. Mol. Biol. Rev.* **70**:177–191.
33. Kanki, T., and D. J. Klionsky. 2010. The molecular mechanism of mitochondrial autophagy in yeast. *Mol. Microbiol.* **75**:795–800.
34. Kimoto, T., et al. 1998. Apoptosis and suppression of tumor growth by artemisinin C extracted from Brazilian propolis. *Cancer Detect. Prev.* **22**:506–515.
35. Klionsky, D. J., A. M. Cuervo, and P. O. Seglen. 2007. Methods for monitoring autophagy from yeast to human. *Autophagy* **3**:181–206.
36. Klionsky, D. J., et al. 2008. Guidelines for the use and interpretation of assays for monitoring autophagy in higher eukaryotes. *Autophagy* **4**:151–175.
37. Kontoyiannis, D. P., and R. E. Lewis. 2002. Antifungal drug resistance of pathogenic fungi. *Lancet* **359**:1135–1144.
38. Korua, O., et al. 2007. Oral and dental bacteriology and infection. *In vitro* antimicrobial activity of propolis samples from different geographical origins against certain oral pathogens. *Anaerobe* **13**:140–145.
39. Kourtis, N., and N. Tavernarakis. 2009. Autophagy and cell death in model organisms. *Cell Death Differ.* **16**:21–30.

40. Kushnareva, Y., and D. D. Newmeyer. 2010. Bioenergetics and cell death. *Ann. N. Y. Acad. Sci.* **1201**:50–57.
41. Li, J., and J. Yuan. 2008. Caspases in apoptosis and beyond. *Oncogene* **27**:6194–6206.
42. Libério, S. A., et al. 2009. The potential use of propolis as a cariostatic agent and its actions on mutans group streptococci. *J. Ethnopharmacol.* **125**:1–9.
43. Lowry, O. H., N. J. Rosebrough, A. Lewis Farr, and R. J. Randall. 1951. Protein measurement with the folin phenol reagent. *J. Biol. Chem.* **193**:265–275.
44. Ludovico, P., M. J. Sousa, M. T. Silva, C. Leão, M. Corte-Real. 2001. *Saccharomyces cerevisiae* commits to a programmed cell death process in response to acetic acid. *Microbiology* **147**:2409–2415.
45. Ludovico, P., et al. 2002. Cytochrome *c* release and mitochondria involvement in programmed cell death induced by acetic acid in *Saccharomyces cerevisiae*. *Mol. Biol. Cell* **13**:2598–2606.
46. Lynch-Day, M. A., and D. J. Klionsky. 2010. The Cvt pathway as a model for selective autophagy. *FEBS Lett.* **584**:1359–1366.
47. Madeo, F., E. Fröhlich, and K.-U. Fröhlich. 1997. A yeast mutant showing diagnostic markers of early and late apoptosis. *J. Cell Biol.* **139**:729–734.
48. Madeo, F., et al. 1999. Oxygen stress: a regulator of apoptosis in yeast. *J. Cell Biol.* **145**:757–767.
49. Madeo, F., et al. 2002. A caspase-related protease regulates apoptosis in yeast. *Mol. Cell* **9**:911–917.
50. Maere, S., K. Heymans, and M. Kuiper. 2005. BiNGO: a Cytoscape plugin to assess overrepresentation of Gene Ontology categories in biological networks. *Bioinformatics* **21**:3448–3449.
51. Maiuri, M. C., E. Zalckvar, A. Kimchi, and G. Kroemer. 2007. Self-eating and self-killing: crosstalk between autophagy and apoptosis. *Nat. Rev. Mol. Cell Biol.* **8**:741–752.
52. Manavathu, E. K., J. L. Culright, and P. H. Chandrasekar. 1998. Organism-dependent fungicidal activities of propolis. *Antimicrob. Agents Chemother.* **42**:3018–3021.
53. Matsuno, T., Y. Matsumoto, N. Saito, and J. Morikawa. 1997. Isolation and characterization of cytotoxic diterpenoid isomers from propolis. *Z. Naturforsch. C* **52**:702–704.
54. Mazzoni, C., and C. Falcione. 2008. Caspase-dependent apoptosis in yeast. *Biochim. Biophys. Acta* **1783**:1320–1327.
55. Menna-Barreto, R. F. S., et al. 2009. Different cell death pathways induced by drugs in *Trypanosoma cruzi*: an ultrastructural study. *Micron* **40**:157–168.
56. Miesenböck, G., D. A. De Angelis, and J. E. Rothman. 1998. Visualizing secretion and synaptic transmission with pH-sensitive green fluorescent proteins. *Nature* **394**:192–195.
57. Morschhäuser, J. 2010. Regulation of multidrug resistance in pathogenic fungi. *Fungal Genet. Biol.* **47**:94–106.
58. Osiewacz, H. D., and C. Q. Scheckhuber. 2006. Impact of ROS on ageing of two fungal model systems: *Saccharomyces cerevisiae* and *Podospora anserina*. *Free Radic. Res.* **40**:1350–1358.
59. Ota, C., C. Unterkircher, V. Fantinato, and M. T. Shimizu. 2001. Antifungal activity of propolis on different species of *Candida*. *Mycoses* **44**:375–378.
60. Park, Y. K., et al. 2005. Suppressive effects of ethanolic extracts from propolis and its main botanical origin on dioxin toxicity. *J. Agric. Food Chem.* **53**:10306–10309.
61. Parrish, J., et al. 2001. Mitochondrial endonuclease G is important for apoptosis in *C. elegans*. *Nature* **412**:90–94.
62. Perrone, G. G., S. X. Tan, and I. W. Dawes. 2008. Reactive oxygen species and yeast apoptosis. *Biochim. Biophys. Acta* **1783**:1354–1368.
63. Ramsdale, M. 2008. Programmed cell death in pathogenic fungi. *Biochim. Biophys. Acta* **1783**:1369–1380.
64. Reers, M., et al. 1995. Mitochondrial membrane potential monitored by JC-1 dye. *Methods Enzymol.* **260**:406–417.
65. Ribeiro, G. F., M. Corte-Real, and B. Johansson. 2006. Characterization of DNA damage in yeast apoptosis induced by hydrogen peroxide, acetic acid, and hyperosmotic shock. *Mol. Biol. Cell* **17**:4584–4591.
66. Rivals, I., L. Personnaz, L. Taing, and M. C. Potier. 2007. Enrichment or depletion of a GO category within a class of genes: which test? *Bioinformatics* **23**:401–407.
67. Rosado, C. J., D. Mijaljica, I. Hatzinisiriou, M. Prescott, and R. J. Devenish. 2008. Rosella: a fluorescent pH-biosensor for reporting vacuolar turnover of cytosol and organelles in yeast. *Autophagy* **4**:205–213.
68. Salomão, K., et al. 2008. Brazilian propolis: correlation between chemical composition and antimicrobial activity. *Evid. Based Complement. Alternat. Med.* **5**:317–324.
69. Scaffidi, P., T. Misteli, and M. E. Bianchi. 2002. Release of chromatin protein HMGB1 by necrotic cells triggers inflammation. *Nature* **418**:191–195.
70. Schauer, A., et al. 2009. Vacuolar functions determine the mode of cell death. *Biochim. Biophys. Acta* **1793**:540–545.
71. Scherens, B., and A. Goffeau. 2004. The uses of genome-wide yeast mutant collections. *Genome Biol.* **5**:229.
72. Semighini, C. P., M. Marins, M. H. S. Goldman, and G. H. Goldman. 2002. Quantitative analysis of the relative transcript levels of ABC transporter *Atr* genes in *Aspergillus nidulans* by real-time reverse transcription-PCR assay. *Appl. Environ. Microbiol.* **68**:1351–1357.
73. Sforzin, J. M. 2007. Propolis and the immune system: a review. *J. Ethnopharmacol.* **113**:1–14.
74. Shannon, P., et al. 2003. Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome Res.* **13**:2498–2504.
75. Shimazawa, M., et al. 2005. Neuroprotection by Brazilian green propolis against *in vitro* and *in vivo* ischemic neuronal damage. *Evid. Based Complement. Alternat. Med.* **2**:201–207.
76. Siqueira, A. B. S., et al. 2009. *Trichophyton* species susceptibility to green and red propolis from Brazil. *Lett. Appl. Microbiol.* **48**:90–96.
77. Sripriya, P., L. V. Vedantam, and A. R. Podile. 2009. Involvement of mitochondria and metacaspase elevation in hairpin Pss-induced cell death of *Saccharomyces cerevisiae*. *J. Cell Biochem.* **107**:1150–1159.
78. Szliszka, E., and W. Krol. 2011. The role of dietary polyphenols in tumor necrosis factor-related apoptosis inducing ligand (TRAIL)-induced apoptosis for cancer chemoprevention. *Eur. J. Cancer Prev.* **20**:63–69.
79. Teixeira, E. W., G. Negri, R. M. S. A. Meira, D. Message, and A. Salatino. 2005. Plant origin of green propolis: bee behavior, plant anatomy and chemistry. *Evid. Based Complement. Alternat. Med.* **2**:85–92.
80. Tzagoloff, A., A. Akai, R. B. Needleman, and G. Zulch. 1975. Assembly of the mitochondrial membrane system. Cytoplasmic mutants of *Saccharomyces cerevisiae* with lesions in enzymes of the respiratory chain and in the mitochondrial ATPase. *J. Biol. Chem.* **250**:8236–8242.
81. Uren, A. G., et al. 2000. Identification of paracaspases and metacaspases: two ancient families of caspase-like proteins, one of which plays a key role in MALT lymphoma. *Mol. Cell* **6**:961–967.
82. Viuda-Martos, M., Y. Ruiz-Navajas, J. Fernandez-López, and J. A. Pérez-Álvarez. 2008. Functional properties of honey, propolis, and royal jelly. *J. Food Sci.* **73**:R117–R124.
83. Vynograd, N., I. Vynograd, and Z. Sosnowski. 2000. A comparative multi-center study of the efficacy of propolis, acyclovir and placebo in the treatment of genital herpes (HSV). *Phytotherapy* **7**:1–6.
84. Winzler, E. A., et al. 1999. Functional characterization of the *S. cerevisiae* genome by gene deletion and parallel analysis. *Science* **285**:901–906.
85. Wuster, A., and M. Madan Babu. 2008. Chemogenomics and biotechnology. *Trends Biotechnol.* **26**:252–258.
86. Zewail, A., et al. 2003. Novel functions of the phosphatidylinositol metabolic pathway discovered by a chemical genomics screen with wortmannin. *Proc. Natl. Acad. Sci. U. S. A.* **100**:3345–3350.

7.2. Anexo J. Artigo submetido: Systems analysis of *Saccharomyces cerevisiae* exposed to propolis. De Castro, P. A.; Savoldi, M.; Bonatto, D.; Malavazi, I.; Goldman, M.H.S.; Berretta, A. A; Goldman, G. H. BMC Complementary and Alternative Medicine, submitted, Dec. 2011.

Systems analysis of *Saccharomyces cerevisiae* exposed to propolis

Patrícia Alves de Castro¹, Marcela Savoldi¹, Diego Bonatto², Iran Malavazi³, Maria Helena S. Goldman⁴, Andresa A. Berretta⁵, and Gustavo Henrique Goldman^{1,6}

¹Faculdade de Ciências Farmacêuticas de Ribeirão Preto, and ⁴Faculdade de Filosofia, Ciências e Letras de Ribeirão Preto, Universidade de São Paulo, São Paulo, Brazil; ²Centro de Biotecnologia da UFRGS, Universidade Federal do Rio Grande do Sul; ³Departamento de Genética e Evolução, Centro de Ciências Biológicas e da Saúde (CCBS), Universidade Federal de São Carlos, Brazil; ⁶Laboratório Nacional de Ciência e Tecnologia do Bioetanol – CTBE, Caixa Postal 6170, 13083-970 Campinas, São Paulo, Brazil; and [‡]Apis Flora Industrial e Comercial, Ribeirão Preto, São Paulo, Brazil.

Running head: Transcriptional profiling of propolis

Corresponding author: Dr. Gustavo H. Goldman

Departamento de Ciências Farmacêuticas

Faculdade de Ciências Farmacêuticas de Ribeirão
Preto, Universidade de São Paulo

Av. do Café S/N, CEP 14040-903, Ribeirão Preto,
São Paulo, Brazil, Phone/Fax: 55-16-36024280/81

e-mail address: ggoldman@usp.br

Abstract

Background:

Propolis is a natural product of plant resins collected by honeybees (*Apis mellifera*) from various plant sources. Our previous studies indicated that propolis sensitivity is dependent on the mitochondrial function and that vacuolar acidification and autophagy are important for yeast cell death caused by propolis. Here, we try to extend our understanding about the mechanism of action of propolis by applying systems analysis of the transcriptional profiling of *S. cerevisiae* exposed to propolis.

Methods:

We have used transcriptional profiling of *S. cerevisiae* exposed to propolis. We validated our findings by using real-time PCR of selected genes. Systems biology analysis was applied aiming to understand which pathways are modulated by propolis in *S. cerevisiae*.

Results:

We were able to observe 339 genes modulated in at least one timepoint when compared to the reference time (*t*-test, *p*-value 0.01). Gene Ontology (GO) analysis of the genes with increased mRNA abundance showed an enrichment for (i) protein transport and transport (23.1 %), (ii) translation (12.8 %), (iii) mitochondria (8.3 %), (iv) metabolic process, and (v) DNA metabolic process and transcription (7.3 %). GO analysis of the genes with decreased mRNA abundance showed an enrichment for (i) DNA metabolic process and transcription (25.7 %), (ii) RNA processing (22.8 %), and (ii) metabolic process (14 %). Real-time RT-PCR analysis of selected genes showed our microarray hybridization approach is capable of providing information about *S. cerevisiae* gene expression modulation with a considerably high level of confidence. Finally, a physical protein-protein (PPPI) network design and global topological analysis stressed the importance of these pathways in response of *S. cerevisiae* to propolis.

Conclusions:

In summary, our data indicate that propolis is largely affecting several pathways in the eukaryotic cell. However, the most prominent pathways are related to oxidative stress, mitochondrial electron transport chain, vacuolar acidification, regulation of macroautophagy associated with protein target to vacuole, cellular response to starvation, and negative regulation of transcription from RNA polymerase II promoter. Our work emphasizes again the importance of *S. cerevisiae* as a model system to understand at molecular level the mechanism of action of a apitherapeutic compound. Our study is the first one that investigates systematically by using functional genomics how propolis influences and modulates the mRNA abundance of an organism.

Background

Propolis is a natural product of plant resins collected by honeybees (*Apis mellifera*) from various plant sources. It is used by the bees to seal holes in their honeycombs and protect the hive entrance [1-3]. Propolis has been used in folk medicine for centuries. Its chemical composition is quite complex since more than 300 compounds, such as polyphenols, phenolic aldehydes, sesquiterpene quinones, coumarins, amino acids, steroids, and inorganic compounds, have been identified in propolis samples. Propolis has cytotoxic [4], anti-herpes virus [5], antitumor [6], radical scavenging [7], antimicrobial [8,9], antiprotozoan [10], and anti-HIV [11] activity and suppressive effects of dioxin toxicity [9]. More recently, evidence has shown that propolis can be used to treat *Candida* fungal infections [12-16].

Recently, we applied the power of *Saccharomyces cerevisiae* as a model organism for studies of genetics, cell biology, and genomics to determine how propolis affects eukaryotic cells at the cellular level [17]. Propolis is able to induce an apoptosis cell death response; however, increased exposure to propolis provides a corresponding increase in the necrosis response. We showed that cytochrome *c* but not endonuclease G (Nuc1p) is involved in propolis-mediated cell death in *S. cerevisiae*. We also observed that the metacaspase *YCA1* gene is important for propolis-mediated cell death. We screened the full collection of about 4,800 haploid *S. cerevisiae* non-essential deletion mutants for propolis sensitivity, and we were able to identify 138 deletion strains that have different degrees of propolis sensitivity compared to the corresponding wild-type strains. Our studies indicated that propolis sensitivity is dependent on the mitochondrial function and that vacuolar acidification and autophagy are important for yeast cell death caused by propolis. Here, we try to extend our understanding about the mechanism of action of propolis and the genes that have their mRNA abundance modulated by this phytotherapeutic compound by applying systems analysis of the transcriptional profiling of *S. cerevisiae* exposed to propolis.

Methods

Propolis Standardized Extract

Propolis Standardized Extract - (EPP-AF[®]) (Batch 010/08) were produced and kindly provided by Apis Flora Company (Ribeirão Preto/SP – Brazil). The extract was standardized using a propolis blend composed by raw material obtained from several sites of Brazil (Patent number PI 0405483-0, published in Revista de Propriedade Industrial n. 1778 from 01/02/2005). Propolis (blend of raw material) was kept in a freezer for 12 h, ground to a fine powder in a blender. It was then extracted using hydroalcoholic solution (7:3), with dynamic maceration followed by percolation process and finally by a filtration. Propolis extract obtained presents 11% w/v of dry matter and chemical composition standardized qualitatively and quantitatively by RP-HPLC into compounds: caffeic, p-coumaric and cinnamic acids, aromadendrin, isosakuranetin and artemillin C.

Yeast strain, media and culture methods

The assays were carried out with *S. cerevisiae* yeast strain BY4742 (*MAT α* ; *his3 Δ 1*; *leu2 Δ 0*; *lys2 Δ 0*; *ura3 Δ 0*) [18]. The culture medium used was complete medium YPD (2% w/v glucose, 1% w/v yeast extract, 2% w/v peptone). For the microarray assay, the yeast cells were grown for 9 hours (mid-exponential phase) in 50mL of liquid YPD at 30°C with mechanical shaking (200rpm). After this, the cells ($\sim 2 \times 10^7$ cells ml⁻¹) were exposed to propolis 0.125 % for 5 or 10 minutes. The control for the experiment using propolis 0.125 % as a treatment has 0.68 % ethanol (what corresponds to 60 % alcoholic extract for 0.125 % propolis). Two independent experiments were performed to each array using two different biological samples and dye swap analysis. Cell viability was determined by plating appropriate cell concentrations and counting the number of colonies in comparison to untreated controls.

RNA isolation and real-time PCR

For total RNA isolation, the yeast cells were disrupted by vortexing with glass beads and total RNA was extracted with Trizol reagent (Invitrogen, USA). Ten micrograms of RNA from each treatment were then fractionated in 2.2 M formaldehyde, 1.2 % w/v agarose gel, stained with ethidium bromide, and then visualized with UV-light. The presence of intact 25S and 18S ribosomal RNA bands was used as a criterion to assess the integrity of the RNA. RNase free DNase treatment was carried out as previously described [19]. After this, the total RNA was purification by RNeasy[®] Mini Kit (Qiagen) and the purified samples were measured in the NanoDrop[®] 2000 (Thermo Scientific).

Microarray hybridization

For gene expression analysis commercially-available Agilent whole genome *S. cerevisiae* microarray [Yeast (V2) Gene Expression Microarray, 8x15K] was used. The microarray slides contain 15,208 probes for *S.cerevisiae* (BY4742 strain). The RNA samples obtained under the conditions above described were subjected to cRNA fluorescent labeling. cRNA labeling was performed according to the standard protocol described by Agilent using Two-Color Microarray-Based Gene Expression Analysis (Agilent Technologies, USA). Briefly, for cRNA synthesis and labeling 5 µg of total RNA were used. After labeling, 300 ng of Cy3 and Cy5-labelled cRNAs (specific activity > 8.0 pmol Cy3-Cy5/µg cRNA) was fragmented at 60 °C for 30 minutes in a reaction volume of 25 µl containing 1x Agilent fragmentation buffer and 2x Agilent blocking agent following the manufacturer's instructions. On completion of the fragmentation reaction, 25 µl of 2x Agilent hybridization buffer was added to the fragmentation mixture and hybridized to the *S. cerevisiae* microarrays slides for 17 hours at 65 °C in an Agilent G2545A Hybridization Oven and on Agilent Rotator Rack. After hybridization, microarrays were sequentially washed: 1 minute at room temperature with GE Wash Buffer 1 (Agilent) and 1 minute with 37 °C GE Wash buffer 2 (Agilent), then a 10 seconds Acetonitrile Wash (Agilent) followed by a

30 seconds Stabilization and Drying Solution wash (Agilent). Slides were immediately subjected to fluorescent detection using fluorescent detection with a GenePix 4000B (Molecular Devices, USA) with simultaneously scanning the Cy3 and Cy5 channels at a resolution of 5 μm . Laser was set at 100% and PMT gain was adjusted automatically for each slide using the program GenePix Pro (Molecular Device) according to the signal intensity of each array. Merged Cy3 and Cy5 TIFF images generated by the GenePix Pro were used to analysis in the Agilent Feature Extraction software (version 9.5.3.1, Agilent) using Linear Lowess algorithm to obtain background subtracted and normalized intensity values. The dye-normalized values generated in the Feature Extraction data files were used to upload the software Express Converter (version 2.1, TM4 available at <http://www.tm4.org/utilities.html>) which conveniently converts the Agilent file format to mev (multi experiment view) file format compatible to the TM4 softwares for microarray analysis (available at <http://www.tm4.org/>). The mev files were then uploaded in the MIDAS software where the resulting data were averaged from replicated genes on each array, from dye-swap hybridizations for each experiment and from two biological replicates using the tools “flip dye consistency cheking” and “in slides replicates analysis” implemented in MIDAS software. The mev files generated were then loaded in MEV software (MultiExperiment Viewer) where differentially expressed genes were identified using one-class t-test ($p > 0.01$). Significantly different genes were those whose mean $\log_2$ expression ratio over all included samples was statistically different from 0 which indicates the absence of gene modulation.

Physical protein-protein (PPPI) network design and global topological analysis

The transcriptomic data gathered from yeast BY4742 strain submitted to propolis treatment was used to obtain information about how the under- and overexpressed genes and their products interact in the context of physical protein-protein interactions (PPPI networks) in *S. cerevisiae*. In this sense, the data mining screening and network design of repressed or induced genes-associated PPPI networks was performed using Cytoscape software, version 2.6.3 [20]. For this purpose, we used the PPPI data of *S. cerevisiae* available in

the *Saccharomyces* Genome Database (<http://www.yeastgenome.org>). The induced and repressed PPPI networks obtained from this first screening were then combined in a unique PPPI network by employing the union function of the Cytoscape core plugin Merge Networks. The union PPPI network was then analyzed with molecular complex detection (MCODE) software [21], a Cytoscape plug-in (at <http://www.cytoscape.org/plugins2.php>) in order to detect clusters of proteins that could represent distinct biologic processes. The parameters used for MCODE to generate the subnetworks were as follows: loops included; degree cutoff of 2; deletion of single connected nodes from cluster (haircut option enabled); expansion of cluster by one neighbor shell allowed (fluff option enable); node density cutoff of 0.1; node score cutoff of 0.2; *k*-core of 2; and maximum depth of network of 100. The degree of data overlapping between induced- and repressed-associated PPPI networks was obtained from an area-proportional Venn diagram analysis, available at <http://bioinforx.com/free/bxarrays/overlap.php>.

Network centralities and local topological analyses

Two major network centralities (node degree and betweenness) were computed from the merged network and clusters using the Cytoscape plugin CentiScaPe 1.0 [22]. The local topology of the network, defined as bottlenecks, was obtained from the threshold generated by each centrality calculated by CentiScape 1.0. In this sense, bottlenecks were defined as nodes with a value above the threshold calculated for node degree and betweenness.

Gene ontology analysis

Gene ontology (GO) clustering analysis was performed using Biological Network Gene Ontology (BiNGO) [23] software, a Cytoscape plugin available at http://chianti.ucsd.edu/cyto_web/plugins/index.php. The degree of functional enrichment for a given cluster and category was quantitatively assessed (*p* value) by hypergeometric distribution [24] and a multiple test correction was applied using the false discovery rate (FDR) [25] algorithm, fully implemented in

BiNGO software. Overrepresented biological process categories were generated after FDR correction, with a significance level of 0.05.

Results and Discussion

Microarray hybridization analysis

To our knowledge, previous to our work there is only a single study in the literature reporting transcriptional profiling for eukaryotic cells exposed to propolis [26]. In this study, propolis was applied for 24 hours to normal human dermal fibroblast and keratinocytes. These authors were able to identify 205 genes important for skin and only 5 (ATP citrate synthase, aquaporin 3, cytochrome c oxidase 1, nitric oxide synthase 3, and hydroxylase 3) and 1 (progesterone receptor) that appear to be up- and downregulated in both cell lines, respectively. We have been using *S. cerevisiae* as an eukaryotic model system to identify genes that are important for propolis-mediated cell death. As previously shown, when *S. cerevisiae* exponential cells are exposed to propolis 0.125 % for 5 and 10 minutes, there is a decreased survival of 24.1 and 6.3 %, respectively [17]. To gain an insight on which pathways are modulated during *S. cerevisiae* exposure to propolis, we determined its transcriptional profile. Total RNA extracted from these cultures was used to generate fluorescent-labeled cRNAs for competitive microarray hybridizations. All the controls for further experiments using propolis 0.125 % as a treatment have 0.68 % ethanol (what corresponds to 60 % alcoholic extract for 0.125 % propolis). We have compared the mRNA expression of the *S. cerevisiae* BY4742 strain grown for 9 hours and exposed to 0.125 % propolis for 5 and 10 minutes with yeast cells exposed to 5 and 10 minutes 0.68 % ethanol. In these experiments, the main aim was to focus on genes that have increased or decreased mRNA expression. The full dataset was deposited in the Gene Expression Omnibus (GEO) from the National Center of Biotechnology Information (NCBI) with the number GSE33971

(<http://www.ncbi.nlm.nih.gov/projects/geo/query/acc.cgi?acc=GSE33971>). We were able to observe 339 genes modulated in at least one timepoint when compared to the reference time (p -value 0.01). These genes are involved with a

variety of cellular processes with increased or decreased mRNA expression [Supplementary Table 1S shows the genes with log ratios ≥ 1 (136 genes) or ≤ -1 (203 genes), respectively]. We used Gene Ontology (GO) analysis aiming to classify to which biological processes the modulated genes are associated (Table S1). GO analysis of the genes with increased mRNA abundance showed an enrichment for (i) protein transport and transport (23.1 %), (ii) translation (12.8 %), (iii) mitochondria (8.3 %), (iv) metabolic process, and (v) DNA metabolic process and transcription (7.3 %) (Figure 1A). In this group of genes, we have observed several genes encoding transporters reported as involved in multidrug resistance (MDR; for reviews see [27-30], suggesting propolis can activate at the transcriptional level the complex set of genes responsible for MDR in *S. cerevisiae* (Table 1). In addition, we have also observed several genes encoding proteins important for the assembly of the vacuolar ATPase and the endocytic pathway (Table 1 and Supplementary Table S1). None of the deletion mutants for these transporter encoding genes shown in Table S1 were observed as more sensitive to propolis, except for *TPO1* [17], indicating a redundant transcriptional response of these genes to propolis. However, when several genes (*VPH1*, *VMA3,4,5,11,22*, *RAV1* and *SOP4*) involved in the assembly of the yeast V-ATPase were deleted, the corresponding yeast deletion strains became more sensitive to propolis [17].

Earlier, by screening a yeast deletion library, we observed that most of the proteins whose deletion increases the sensitivity of yeast strains to propolis are involved in cell division mechanisms, mitochondrial electron transport chain, vacuolar acidification, regulation of macroautophagy associated with protein target to vacuole, cellular response to starvation, and negative regulation of transcription from RNA polymerase II promoter [17]. We have shown that propolis induces vacuolar acidification and translocation of Atg8p to the vacuoles, one of the hallmarks of autophagy [17]. In *S. cerevisiae*, the vacuole is very important for keeping cellular homeostasis comprising the regulation of intracellular pH and degradation mainly during nutrient limitation of proteins and organelles by autophagy (for reviews, see [31- 33]). Cell death induced by acetic acid is increased in *S. cerevisiae* *VPS* gene deletion mutants (*VPS* genes are involved in homotypic vacuole fusion, vacuolar protein sorting and are

essential for normal vacuolar function) [34]. It has been observed that the intracellular pH was acidified in *VPS* mutant cells upon treatment with acetic acid [34]. It is possible the disturbance of the homeostatic pH control may trigger necrosis by release of pro-necrotic proteases, which would find an optimal pH for their enzymatic activity in the acidified cytosol [35]. We have observed several genes encoding proteins important for ROS detoxification in *S. cerevisiae*, such as *GRX*, *GTT2*, *GTT3*, *TSA2*, *GND2*, and *TRX1* with increased mRNA accumulation when *S. cerevisiae* is exposed to propolis (Table 1 and Supplementary Table S1). Interestingly, there is an increased mRNA accumulation of quite a lot of genes encoding proteins involved in the generation of ATP into the mitochondria, such as *ATP17*, *ATP18*, *ATP19*, *ATP20*, *ATP21*, and *COX8* (Supplementary Table S1). There are several conditions where mitochondria-produced ROS have been associated to yeast apoptosis (for reviews, see [35-38]). Propolis can induce ROS formation and it is more lethal when *S. cerevisiae* grows in the presence of glycerol and ethanol as carbon sources [17], suggesting that respiration increases propolis lethality.

GO analysis of the genes with decreased mRNA abundance showed an enrichment for (i) DNA metabolic process and transcription (25.7 %), (ii) RNA processing (22.8 %), and (iii) metabolic process (14 %) (Figure 1B). There are several genes related to kinetochore and chromosome distribution and chromatin silencing that have decreased mRNA accumulation when *S. cerevisiae* is exposed to propolis (Table 2). The reduced mRNA abundance of these genes suggested propolis is activating transcriptional checkpoint controls involved in the S- and M-phases important for DNA replication and proper chromosome segregation. It is not possible to infer at this point if this is a direct or indirect effect of propolis on the cell cycle progression.

Validation of the microarray hybridization analysis

To validate some of our findings, we have chosen nine different genes from our microarray analysis whose mRNA has either increased or decreased abundance when *S. cerevisiae* is exposed to propolis. We designed Lux fluorescent probes and used real-time RT-PCR analysis to quantify their

expression in a new biological replicate of the mRNA isolated from 5 and 10 minutes exposure to propolis and compared them with the corresponding 5 and 10 minutes exposure to 0.68 % ethanol. We have used as a normalizer control, *TAF10*, a gene encoding a subunit (145 kDa) of TFIID and SAGA complexes, involved in RNA polymerase II transcription initiation and in chromatin modification. Recently, this gene was shown as an appropriate reference gene for quantitative gene expression analysis by real-time RT-PCR [39]. In addition, *TAF10* was not shown as modulated in our microarray hybridization experiments (data not shown). Thus, the measured quantity of a specific gene mRNA in each of the treated samples was normalized using the C_T values obtained for the *TAF10* mRNA amplifications run in the same plate. The results were expressed as the number of times the genes have increased or decreased abundance when the yeast strains were exposed to propolis compared to the ethanol treatment (Figure 2). We evaluated the mRNA abundance of *PDR15* (YDR406W) encoding a transporter involved in multidrug resistance; *SNQ2* (YDR011W) encoding a plasma membrane ATP-binding cassette (ABC) transporter, multidrug transporter involved in multidrug resistance and resistance to singlet oxygen species; *VMA7* (YGR020C) and *VMA21* (YGR105W) encoding the subunit F of the eight-subunit V1 peripheral membrane domain of vacuolar H⁺-ATPase (V-ATPase) and Integral membrane protein that is required for vacuolar H⁺-ATPase (V-ATPase) function, respectively; *RLF2* (YPR018W) encoding the largest subunit (p90) of the Chromatin Assembly Complex (CAF-1) with Cac2p and Msi1p that assembles newly synthesized histones onto recently replicated DNA; and *TIM10* (YHR005C-A) encoding an essential protein of the mitochondrial intermembrane space, that forms a complex with Tim9p (TIM10 complex) and delivers hydrophobic proteins to the TIM22 complex for insertion into the inner membrane. All the genes had increased abundance in our microarray hybridizations, except for *RLF2* that had decreased mRNA abundance. All five genes that showed increased mRNA abundance (*PDR15*, *TIM10*, *SNQ2*, *VMA7*, and *VMA21*) and decreased mRNA abundance (*RLF2*) in the microarray hybridization analysis showed corresponding increase and decreased in the real-time PCR experiments (Figure 2A-F). Thus, it seems that our microarray

hybridization approach is capable of providing information about *S. cerevisiae* gene expression modulation with a considerably high level of confidence.

Systems analysis for propolis exposure

The transcriptomics data obtained submitted to the treatment conditions described in this work prompt us to ask how the underexpressed or overexpressed genes affect different biological processes that are altered during the exposure to propolis. In this sense, a search for potential proteins and/or mechanisms and their associated biological processes that are affected by the conditions treatment was initiated. To achieve this goal, the union of induced and repressed PPPI networks using the yeast transcriptomics data obtained in this work were retrieved from *Saccharomyces* Genome Database (SGD) (Table 3 and Supplementary Table S2). Subnetworks present in the union PPPI network (Table 2) were identified and retrieved using the Cytoscape-associated plugin MCODE and subjected to a Gene Ontology (GO) analysis in order to obtain information about the nature and number of subgraphs belonging to the network and their associated biological processes.

For analysis, two different PPPI networks using yeast transcriptomics data were retrieved from *Saccharomyces* Genome Database (SGD): one associated to repressed genes (repressed genes-associated PPPI network) and one associated to induced genes (induced genes-associated network). The induced genes-associated PPPI network obtained from SGD contains 1,226 nodes and 2,854 connectors while the repressed-associated gene PPPI network contains 1,412 nodes and 2,782 connectors (Supplementary Table S3). Both induced- and repressed-genes associated PPPI networks were analyzed to see the data overlapping and Area-Proportional Venn Diagram (Figure 3). The repressed-genes associated PPPI network contains 919 unique genes, while the induced-genes associated network contains 733 unique genes, and there are 493 overlapping genes. Although the degree of network overlapping is not elevated, we decided to merge both networks in a unique graph, containing 2158 nodes and 5655 connectors (Supplementary Table S3).

GO analysis of union network and its clusters indicate the participation of important biological processes, like: Cluster 1: Translation, ribosome biogenesis and negative regulation of mRNA processing; Cluster 2: Transport (organic acid, choline, betaine, ethanolamine); Cluster 3: mitochondrion transport, mitochondrion organization and biogenesis, protein transport; Cluster 4: regulation of transcription from RNA polymerase II promoter, establishment and/or maintenance of chromatin architecture, regulation of RNA metabolic process, transcription/chromosome organization and biogenesis, G1 phase of mitotic cell cycle, histone modification; Cluster 5: DNA catabolic process, double-strand break repair, meiotic recombination, M phase of meiotic cell cycle, telomere maintenance via recombination; Cluster 6: ER to Golgi vesicle-mediated transport; Cluster 7: negative regulation of transcription, negative regulation of RNA metabolic process, heterochromatin formation, chromatin silencing, gene silencing, establishment and/or maintenance of chromatin architecture, regulation of cell cycle; Cluster 8: RNA biosynthetic process, transcription from RNA polymerase II promoter; Cluster 9: regulation of pH, pH reduction, vacuolar acidification; Cluster 10: histone modification, transcription from RNA polymerase I, II and III promoter, establishment and/or maintenance of chromatin architecture, chromosome organization and biogenesis, ethanol biosynthetic process during fermentation, response to stress; Cluster 11: chromosome organization and biogenesis, DNA repair, mitotic cell cycle, histone acetylation (Supplementary S2).

Based on data gathered from this initial systems biology analysis, we decided to get more information about the major nodes involved in the information of the network using network centralities. Network centralities allow us to identify nodes (and the consequent biological processes) that have a relevant position in the overall network architecture [40]. Centralities have been recently applied to quantify the centrality and prestige of actors in social networks [40] and to understand the structure and properties of complex biological, technological and infrastructural networks [41, 42]. Many of the nodes in a given network that show elevated values of centrality are important points of vulnerability, indicating that any attack to these nodes could introduce strong perturbations in the network. Node degree represents the simplest

centrality measure in a given network, corresponding to the number of nodes adjacent to a given node, where adjacent means directly connected [22]. The node degree represents the “popularity” of a given node, and highly connected nodes in a network are termed hubs. Next, betweenness is a measure that indicates to what extent a specific node is between all other nodes within the network [43]. In a general sense, betweenness show the influence of a node over the spread of information throughout the network. On the other hand, bottleneck is a local topologic data that is defined as all nodes with high betweenness values and different nodes degrees, indicating that those nodes are central points that control the communication between other nodes within the network [44, 45]. The measures of betweenness and node degree allow us to define the bottleneck nodes. Bottleneck nodes correspond to highly central proteins that connect several complexes or are peripheral members of central complexes, being important communication points between two complexes [45]. Mostly of bottleneck nodes tend to be essential proteins in a network [45].

The centrality analysis of union network indicated the presence of 419 bottleneck nodes; 99 of these bottleneck nodes were induced in array and 92 were repressed in array (Supplementary Table S4, plan 1). The analysis of the 11 clusters showed the presence of important bottleneck nodes in the clusters (Supplementary Table S4, plan 2). By applying GO analysis to the nodes bottleneck, we observed nodes participating in biological processes as metabolic process, protein transport and transport, cellular component organization and biogenesis, DNA metabolic process and transcription, RNA processing, translation, response to stress, cell cycle (for induced genes in the microarray); and metabolic process; protein transport and transport; DNA metabolic process and transcription; RNA processing; translation; signal transduction; cell cycle (for repressed genes in the microarray). These data stress the importance of these pathways in response of *S. cerevisiae* to propolis.

Conclusions

Propolis is a complex product derived from plant resins and bee's saliva. There are several chemical compounds present in this natural product that could potentially be responsible for its antibiotic properties. However, taking into consideration the fact that the cell death effects of propolis could be due to a great combination of chemical compounds and concentrations, we decided to investigate the cell death effects of propolis by concentrating our experiments on alcoholic extracts of propolis. In summary, our data indicates propolis is largely affecting several pathways in the eukaryotic cell. However, the most prominent pathways are related to oxidative stress, mitochondrial electron transport chain, vacuolar acidification, regulation of macroautophagy associated with protein target to vacuole, cellular response to starvation, and negative regulation of transcription from RNA polymerase II promoter. Our work emphasizes again the importance of *S. cerevisiae* as a model system to understand at molecular level the mechanism of action of an apitherapeutic compound. Our study is the first one that investigates systematically by using functional genomics how propolis influences and modulated the mRNA abundance of an organism. This opens new possibilities for understanding and validating propolis as an alternative therapeutic agent.

Acknowledgements

This research was supported by the Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP), and Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), and Financiadora de Estudos e Projetos (FINEP), Brazil.

Author details

¹Faculdade de Ciências Farmacêuticas de Ribeirão Preto, and ⁴Faculdade de Filosofia, Ciências e Letras de Ribeirão Preto, Universidade de São Paulo, São Paulo, Brazil; ²Centro de Biotecnologia da UFRGS, Universidade Federal do Rio Grande do Sul; ³Departamento de Genética e Evolução, Centro de Ciências Biológicas e da Saúde (CCBS), Universidade Federal de São Carlos, Brazil; ⁶Laboratório Nacional de Ciência e Tecnologia do Bioetanol – CTBE,

Caixa Postal 6170, 13083-970 Campinas, São Paulo, Brazil; and [‡]Apis Flora Industrial e Comercial, Ribeirão Preto, São Paulo, Brazil.

Authors' contributions

PAC and MS performed most of the experiments. PAC, DB, AB, IM, MHSG and GHG performed data analysis. GHG wrote the manuscript; conceived, designed and coordinated this study; and is the principal investigator of this work. All authors read and approved final manuscript.

Competing interests

The authors declare that they have no competing interests.

Figure Legends

Figure 1 – GO analysis of the genes observed as modulated in the microarray hybridization analysis of *S. cerevisiae* exposed to propolis. Genes with increased (A) and decreased (B) mRNA abundance.

Figure 2 - Real-time RT-PCR for selected genes from the microarray hybridization analysis. *S. cerevisiae* was grown for 9 hours in liquid YPD at 30 °C and after this, the cells ($\sim 2 \times 10^7$ cells ml⁻¹) were transferred to fresh liquid YPD and exposed to propolis 0.125 % for 5 or 10 minutes. The relative quantitation of *RLF2* (A), *PDR15* (B), *TIM10* (C), *SNQ2* (D), *VMA7* (E), *VMA21*(F), and *TAF10* gene expression was determined by a standard curve (i.e., C_T–values plotted against logarithm of the DNA copy number). The results are the means $\pm$ standard deviation of four sets of experiments.

Figure 3 – An area proportional Venn analysis of node overlapping degree of repressed- and induced-associated gene PPPI networks.

References

1. Ghisalberti EL: **Propolis: a review**. *Bee World* 1979, **60**:59–84.
2. Greenaway W, Scaysbrook T, Whatley FR: **The composition and plant origins of propolis**. *Bee World* 1990, **71**:107–118.
3. Viuda-Martos M, Ruiz-Navajas Y, Fernandez-Lo´pez J, Pe´rez- A´ lvarez J A: **Functional properties of honey, propolis, and royal jelly**. *J. Food Sci.* 2008, **73**:R117–R124.
4. Matsuno T, Matsumoto Y, Saito N, Morikawa J: **Isolation and characterization of cytotoxic diterpenoid isomers from propolis**. *Z. Naturforsch* 1997, **C 52**:702–704.
5. Vynograd N, Vynograd I, Sosnowski Z: **A comparative multicenter study of the efficacy of propolis, acyclovir and placebo in the treatment of genital herpes (HSV)**. *Phytomedicine* 2000, **7**:1–6.
6. Kimoto T, Arai S, Kohguchi M, Nomura Y, Micallef MJ, Kurimoto M, Mito K: **Apoptosis and suppression of tumor growth by artepillin C extracted from Brazilian propolis**. *Cancer Detect. Prev.* 1998, **22**:506–515.
7. Basnet P, Matsuno T, Neidlein RZ: **Potent free radical scavenging activity of propol isolated from Brazilian propolis**. *Z. Naturforsch* 1997, **C 52**:828–833.
8. Koru O, Toksoy F, Acikel CH, Tunca YM, Baysallar M, Uskudar Guclu A, Akca E, Ozkok Tuylu A, Sorkun K, Tanyuksel M, Salih B: **Oral and dental bacteriology and infection. In vitro antimicrobial activity of propolis samples from different geographical origins against certain oral pathogens**. *Anaerobe* 2007, **13**:140–145.

9. Park YK, Fukuda I, Ashida H, Nishiumi S, Yoshida K, Daugsch A, Sato HH, Pastore GM: **Suppressive effects of ethanolic extracts from propolis and its main botanical origin on dioxin toxicity.** *J. Agric. Food Chem* 2005, **53**:10306–10309.

10. Menna-Barreto RF, Salomão K, Dantas AP, Santa-Rita RM, Soares MJ, Barbosa HS, de Castro SL: **Different cell death pathways induced by drugs in *Trypanosoma cruzi*: an ultrastructural study.** *Mícron*. 2009, **40**:157–168.

11. Ito J, Chang FR, Wang HK, Park YK, Ikegaki M, Kilgore N, Lee KH: **Anti-AIDS agents. 48.(1) Anti-HIV activity of moronic acid derivatives and the new melliforone-related triterpenoid isolated from Brazilian propolis.** *J. Nat. Prod.* 2001, **64**:1278–1281.

12. Agüero MB, Gonzalez M, Lima B, Svetaz L, Sánchez M, Zacchino S, Feresin GE, Schmeda-Hirschmann G, Palermo J, Wunderlin D, Tapia A: **Argentinean propolis from *Zuccagnia punctata* Cav. (Caesalpinieae) exudates: phytochemical characterization and antifungal activity.** *J. Agric. Food Chem.* 2010, **58**:194–201.

13. Dalben-Dota KF, Faria MG, Bruschi ML, Pelloso SM, Lopes-Consolaro ME, Svidzinski TI: **Antifungal activity of propolis extract against yeasts isolated from vaginal exudates.** *J. Altern. Complement. Med.* 2010, **16**:285–290.

14. Osiewacz HD, Scheckhuber CQ: **Impact of ROS on ageing of two fungal model systems: *Saccharomyces cerevisiae* and *Podospora anserina*.** *Free Radic. Res.* 2006, **40**:1350–1358.

15. Salomão K, Pereira PR, Campos LC, Borba CM, Cabello PH, Marcucci MC, de Castro SL: **Brazilian propolis: correlation between chemical composition and antimicrobial activity.** *Evid. Based Complement. Alternat. Med.* 2008, **5**:317–324.

16. Siqueira AB, Gomes BS, Cambuim I, Maia R, Abreu S, Souza-Motta CM, de Queiroz LA, Porto AL: **Trichophyton species susceptibility to green and red propolis from Brazil.** *Lett. Appl. Microbiol.* 2009, **48**:90–96.

17. de Castro PA, Savoldi M, Bonatto D, Barros MH, Goldman MH, Berretta AA, Goldman GH: **Molecular characterization of propolis-induced cell death in *Saccharomyces cerevisiae*.** *Eukaryot Cell.* 2011, **10(3)**:398-411.

18. Brachmann CB, Davies A, Cost GJ, Caputo E, Li J, Hieter P, Boeke JD: **Designer deletion strains derived from *Saccharomyces cerevisiae* S288C: a useful set of strains and plasmids for PCR mediated gene disruption and other applications.** *Yeast* 1998, **14**:115–132.

19. Semighini CP, Marins M, Goldman MHS, Goldman GH: **Quantitative analysis of the relative transcript levels of ABC transporter Atr genes in *Aspergillus nidulans* by real-time reverse transcription-PCR assay.** *Appl. Environ. Microbiol.* 2002, **68**:1351–1357.

20. Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N, Schwikowski B, Ideker T: **Cytoscape: a software environment for integrated models of biomolecular interaction networks.** *Genome Res.* 2003, **13**:2498-2504.

21. Bader GD, Hogue CW: **An automated method for finding molecular complexes in large protein interaction networks.** *BMC Bioinformatics* 2003, **4**:2.

22. Scardoni G, Petterlini M, Laudanna C: **Analyzing biological network parameters with CentiScaPe.** *Bioinformatics* 2009, **25**: 2857-2859.

23. Maere S, Heymans K, Kuiper M: **BiNGO: a Cytoscape plugin to assess overrepresentation of gene ontology categories in biological networks.** *Bioinformatics* 2005, **21**: 3448-3449.
24. Rivals I, Personnaz L, Taing L, Potier M-C: **Enrichment or depletion of a GO category within a class of genes: which test?** *Bioinformatics* 2007, **23**: 401-407.
25. Benjamini Y, Hochberg Y: **Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing.** *J. R. Stat. Soc. Series B Stat. Methodol.* 1995, **57**: 289-300.
26. Gruber JV, Holtz R: **Examining the genomic influence of skin antioxidants in vitro.** *Mediators Inflamm.* 2010, pii: 230450.
27. Ernst R, Kueppers P, Stindt J, Kuchler K, Schmitt L: **Multidrug efflux pumps: substrate selection in ATP-binding cassette multidrug efflux pumps--first come, first served?** *FEBS J.* 2010, **277(3)**:540-9.
28. Shahi P, Moye-Rowley WS: **Coordinate control of lipid composition and drug transport activities is required for normal multidrug resistance in fungi.** *Biochim Biophys Acta.* 2009, **1794(5)**:852-9.
29. Sá-Correia I, dos Santos SC, Teixeira MC, Cabrito TR, Mira NP: **Drug:H⁺ antiporters in chemical stress response in yeast.** *Trends Microbiol.* 2009, **17(1)**:22-31.
30. Sipos G, Kuchler K: **Fungal ATP-binding cassette (ABC) transporters in drug resistance & detoxification.** *Curr Drug Targets.* 2006, **7(4)**:471-81.
31. Ariño J, Ramos J, Sychrová H: **Alkali metal cation transport and homeostasis in yeasts.** *Microbiol. Mol. Biol.* 2010, **74**: 95-120.

32. Inoue Y, Klionsky DJ: **Regulation of macroautophagy in *Saccharomyces cerevisiae***. *Semin. Cell Dev. Biol.* 2010, **21**: 664-70.
33. Lynch-Day, MA, Klionsky DJ: **The Cvt pathway as a model for selective autophagy**. *FEBS Lett.* 2010, **584**: 1359-1366.
33. Schauer A, Knauer H, Ruckenstuhl C, Fussi H, Durchschlag M, Potocnik U, Fröhlich K-U: **Vacuolar functions determine the mode of cell death**. *Biochim. Biophys. Acta.* 2009, **1793**: 540-545.
34. Eisenberg T, Knauer H, Schauer A, Büttner S, Ruckenstuhl C, Carmona-Gutierrez D, Ring J, Schroeder S, Magnes C, Antonacci L, Fussi H, Deszcz L, Hartl R, Schraml E, Criollo A, Megalou E, Weiskopf D, Laun P, Heeren G, Breitenbach M, Grubeck-Loebenstien B, Herker E, Fahrenkrog B, Fröhlich KU, Sinner F, Tavernarakis N, Minois N, Kroemer G, Madeo F: **Induction of autophagy by spermidine promotes longevity**. *Nat. Cell Biol.* 2009, **11**: 1305-1314.
35. Ota C, Unterkircher C, Fantinato V, Shimizu MT: **Antifungal activity of propolis on different species of *Candida***. *Mycoses* 2001, **44**: 375-378.
36. Perrone GG, Tan SX, Dawes IW: **Reactive oxygen species and yeast apoptosis**. *Biochim. Biophys. Acta* 2008, **1783**: 1354-1368
37. Eisenberg T, Carmona-Gutierrez D, Büttner S, Tavernarakis N, Madeo F: **Necrosis in yeast**. *Apoptosis* 2010, **15**: 257-268.
38. Teste MA, Duquenne M, François JM, Parrou JL: **Validation of reference genes for quantitative expression analysis by real-time RT-PCR in *Saccharomyces cerevisiae***. *BMC Mol Biol.* 2009, **10**():99.
39. Borgatti SP: **Centrality and network flow**. *Soc. Networks* 2005, **27**: 55-71.

40. Estrada E: **Virtual identification of essential proteins within the protein interaction network of yeast.** *Proteomics* 2006, **6**: 35-40.
41. Estrada E, Hatano NA: **Vibrational approach to node centrality and vulnerability in complex networks.** *Phys. A.* 2010, **389**: 3648-3660.
42. Newman MEJ: **A measure of betweenness centrality based on random walks.** *Soc. Networks* 2005, **27**: 39-54.
43. Girvan M, Newman ME: **Community structure in social and biological networks.** *Proc. Natl. Acad. Sci. U S A* 2002, **99**: 7821-7826.
44. Yu H, Kim PM, Sprecher E, Trifonov V, Gerstein M: **The importance of bottlenecks in protein networks: correlation with gene essentiality and expression dynamics.** *PLoS Comput. Biol.* 2007, **3**: e59.

Table 1 – Selected genes more expressed ($\log_2 \geq 1.0$) during *S. cerevisiae* exposure to propolis (for a complete list of the genes more expressed, see Supplementary Table S1).

FLR1	Multidrug transporter of the MFS, involved in efflux of fluconazole, diazaborine, benomyl, methotrexate, and other drugs
PDR15	ATP binding cassette (ABC) transporter, multidrug transporter and general stress response factor implicated in cellular detoxification
AZR1	Transporter of the MFS, involved in resistance to azole drugs such as ketoconazole and fluconazole
YOR1	ATP-binding cassette (ABC) transporter, multidrug transporter mediates export of many different organic anions including oligomycin
PDR10	ATP-binding cassette (ABC) transporter, multidrug transporter involved in the pleiotropic drug resistance network
PDR12	ATP-binding cassette (ABC) transporter, weak-acid-inducible multidrug transporter required for weak organic acid resistance
ENA1	P-type ATPase sodium pump, involved in Na ⁺ and Li ⁺ efflux to allow salt tolerance
ENA2	P-type ATPase sodium pump, involved in Na ⁺ efflux to allow salt tolerance; likely not involved in Li ⁺ efflux
VMA21	Membrane protein that is required for vacuolar H ⁺ -ATPase (V-ATPase) function, although not an actual component of the V-ATPase
TPO1	Polyamine transporter (MFS) that recognizes spermine, putrescine, and spermidine
TPO4	Polyamine transport protein (MFS) recognizes spermine, putrescine, and spermidine
ATG22	Vacuolar integral membrane protein required for efflux of amino acids during autophagic body breakdown in the vacuole
VMA7	Subunit F of the eight-subunit V1 peripheral membrane domain of vacuolar H ⁺ -ATPase (V-ATPase)
PKR1	V-ATPase assembly factor, functions with other V-ATPase assembly factors in the ER to efficiently assemble the V-ATPase
GRX4	Hydroperoxide and superoxide-radical responsive glutathione-dependent oxidoreductase
GTT3	Protein of unknown function with a possible role in glutathione metabolism
TSA2	Stress inducible cytoplasmic thioredoxin peroxidase; cooperates with Tsa1p in the removal of reactive oxygen
GTT2	Glutathione S-transferase capable of homodimerization; functional overlap with Gtt2p, Grx1p, and Grx2p
GND2	6-phosphogluconate dehydrogenase, catalyzes an NADPH regenerating reaction in the pentose phosphate pathway
TRX1	Cytoplasmic thioredoxin isoenzyme of the thioredoxin system which protects cells against oxidative and reductive stress

Table 2 – Selected genes less expressed ($\log_2 \leq -1.0$) during *S. cerevisiae* exposure to propolis (for a complete list of the genes less expressed, see Supplementary Table S1).

SMC4	Subunit of the condensin complex; reorganizes chromosomes during cell division
CM10	Essential chromatin-associated protein involved in the initiation of DNA replication
CIN8	Kinesin motor protein involved in mitotic spindle assembly and chromosome segregation
IBD2	Component of the BUB2-dependent spindle checkpoint pathway, interacts with Bfa1p and functions upstream of Bub2p and Bfa1p
SPC110	Inner plaque spindle pole body (SPB) component, ortholog of human kendrin; involved in connecting nuclear microtubules to SPB
BRN1	Subunit of the condensin complex; required for chromosome condensation and for clustering of tRNA genes at the nucleolus
CEP3	Essential kinetochore protein, component of the CBF3 complex that binds the CDEIII region of the centromere
SLK19	Kinetochore-associated protein required for normal segregation of chromosomes in meiosis and mitosis
NSL1	Component of the MIND kinetochore complex which joins kinetochore subunits contacting DNA to those contacting microtubules
SPC24	Component of the evolutionarily conserved kinetochore-associated Ndc80 complex
MAM1	Monopolin, kinetochore associated protein involved in chromosome attachment to meiotic spindle
RLF2	Largest subunit (p90) of the Chromatin Assembly Complex (CAF-1) and Msi1p that assembles newly synthesized histones
HOS2	Histone deacetylase required for gene activation via specific deacetylation of lysines in H3 and H4 histone tails
ESA1	Catalytic subunit of the histone acetyltransferase complex (NuA4) that acetylates four conserved internal lysines of histone H4
LEO1	Component of the Paf1 complex, which associates with RNA polymerase II and is involved in histone methylation
SPT21	Protein required for normal transcription at several loci including HTA2-HTB2 and HHF2-HHT2
RFM1	Specificity factor that directs the Hst1p histone deacetylase to some of the promoters regulated by Sum1p
SGF29	Probable subunit of SAGA histone acetyltransferase complex
IES2	Protein that associates with the INO80 chromatin remodeling complex under low-salt conditions
IFH1	IFH1 exhibits genetic interactions with FHL1, overexpression interferes with silencing at telomeres and HM loc
NGG1	component of transcriptional adaptor and histone acetyltransferase complexes, the ADA, the SAGA, and the SLIK complexes

TABLE 3 - Main specific gene ontology classes derived from PPPI network observed in clusters

<i>Biological process</i>	<i>GO</i>		<i>Corrected P value^b</i>	<i>k^c</i>	<i>f^d</i>
	<i>identification</i>	<i>P value^a</i>			
	<i>no.</i>				
Translation	6416	2.07×10^{-45}	2.93×10^{-43}	45	499
Cellular protein metabolic process	44267	4.00×10^{-21}	9.02×10^{-20}	45	170
Transcription from RNA polymerase II promoter	6366	3.49×10^{-11}	3.43×10^{-9}	9	162 ²
Mitochondrial transport	6839	1.74×10^{-10}	3.88×10^{-9}	5	67
Histone modification	16570	1.14×10^{-9}	2.23×10^{-8}	7	92
Ribosome assembly	42255	1.82×10^{-9}	2.58×10^{-8}	9	65
Establishment and/or maintenance of chromatin architecture	6325	2.45×10^{-9}	4.37×10^{-8}	9	260
G1 phase of mitotic cell cycle	80	1.05×10^{-8}	1.28×10^{-7}	5	32
Transcription	6350	1.15×10^{-8}	1.33×10^{-7}	11	598
Regulation of RNA metabolic process	51252	1.53×10^{-8}	1.57×10^{-7}	11	614
Mitochondrion organization and biogenesis	7005	2.69×10^{-8}	2.91×10^{-7}	5	179
Chromosome organization and biogenesis	7001	1.04×10^{-7}	8.86×10^{-7}	9	398
DNA double-strand break processing	729	1.77×10^{-6}	5.67×10^{-5}	2	5
Negative regulation of RNA metabolic process	51253	2.49×10^{-6}	8.00×10^{-5}	4	157
Negative regulation of transcription	16481	3.12×10^{-6}	8.00×10^{-5}	4	166

Protein transport	15031	3.78×10^{-6}	2.11×10^{-5}	5	481
Heterochromatin formation	31507	3.74×10^{-5}	2.99×10^{-4}	3	92
Chromatin silencing	6342	3.74×10^{-5}	2.99×10^{-4}	3	92
pH Reduction	45851	4.88×10^{-5}	6.68×10^{-4}	2	24
Vacuolar acidification	7035	4.88×10^{-5}	6.68×10^{-4}	2	24
Gene silencing	16458	5.40×10^{-5}	3.84×10^{-4}	3	104
Regulation of pH	6885	8.21×10^{-5}	8.62×10^{-4}	2	31
Organic acid transport	15849	2.06×10^{-4}	1.41×10^{-2}	3	65
ER to Golgi vesicle-mediated transport	6888	6.12×10^{-4}	2.57×10^{-2}	2	84
Cell cycle	7049	9.16×10^{-4}	3.51×10^{-3}	3	566
Ethanol biosynthetic process during fermentation	43458	6.52×10^{-3}	1.76×10^{-2}	1	2
Regulation of cell cycle	74	7.11×10^{-3}	1.57×10^{-2}	2	160
Telomere maintenance via recombination	722	9.76×10^{-3}	2.53×10^{-2}	1	19
Response to stress	6950	1.26×10^{-2}	2.96×10^{-2}	6	632
Telomere maintenance via recombination	722	9.76×10^{-3}	2.53×10^{-2}	1	19

^a *P* values were calculated by the hypergeometric distribution of one ontology class visualized in the network.

^b Calculated values based on *P* values obtained after FDR was applied.

^c Total number of proteins found in the network which belong to a gene ontology.

^d Total number of proteins that belong to a specific gene ontology.

A.

B.

Figure 1

A.

B.

C.

D.

E.

F.

Induced Network	Repressed network	Combined	Overlap	Not Overlap	Induced network Only	Repressed network Only
1226	1412	2145	493	1652	733	919

Figure 3

Additional files provided with this submission:

Additional file 1: Supplementary Table S1.xls, 106K

<http://www.biomedcentral.com/imedia/4774560964946544/supp1.xls>

Additional file 2: Supplementary Table S2.ods, 98K

<http://www.biomedcentral.com/imedia/2071216190649465/supp2.ods>

Additional file 3: Supplementary Table S3.xls, 191K

<http://www.biomedcentral.com/imedia/3697035196494649/supp3.xls>

Additional file 4: Supplementary Table S4.xls, 172K

<http://www.biomedcentral.com/imedia/1921029501649465/supp4.xls>

7.3. Anexo L. Artigo publicado: Involvement of the *Aspergillus nidulans* protein kinase C with farnesol tolerance is related to the unfolded protein response. Colabardini, A. C.; de Castro, P. A.; de Gouvêa, P. F.; Savoldi, M.; Malavazi, I.; Goldman, M. H. S.; Goldman, G. H. Molecular Microbiology, Dec. 2010, 78: 1259–1279.

Involvement of the *Aspergillus nidulans* protein kinase C with farnesol tolerance is related to the unfolded protein response

Ana Cristina Colabardini,¹ Patrícia Alves de Castro,¹ Paula Fagundes de Gouvêa,¹ Marcela Savoldi,¹ Iran Malavazi,³ Maria Helena S. Goldman² and Gustavo Henrique Goldman^{1,4*}

¹Faculdade de Ciências Farmacêuticas de Ribeirão Preto and ²Faculdade de Filosofia, Ciências e Letras de Ribeirão Preto, Universidade de São Paulo, São Paulo, Brazil.

³Departamento de Genética e Evolução, Centro de Ciências Biológicas e da Saúde (CCBS), Universidade Federal de São Carlos, Brazil.

⁴Laboratório Nacional de Ciência e Tecnologia do Bioetanol – CTBE, Caixa Postal 6170, 13083-970 Campinas, São Paulo, Brazil.

Summary

Previously, we demonstrated that the *Aspergillus nidulans* *calC2* mutation in protein kinase C *pkcA* was able to confer tolerance to farnesol (FOH), an isoprenoid that has been shown to inhibit proliferation and induce apoptosis. Here, we investigate in more detail the role played by *A. nidulans* *pkcA* in FOH tolerance. We demonstrate that *pkcA* overexpression during FOH exposure causes increased cell death. FOH is also able to activate several markers of endoplasmic reticulum (ER) stress and the unfolded protein response (UPR). Our results suggest an intense cross-talk between PkcA and the UPR during FOH-induced cell death. Furthermore, the overexpression of *pkcA* increases both mRNA accumulation and metacaspases activity, and there is a genetic interaction between PkcA and the caspase-like protein CasA. Mutant analyses imply that MAP kinases are involved in the signal transduction in response to the effects caused by FOH.

Introduction

Protein kinase C (PKC) is a conserved enzyme among eukaryotes. Although mammalian cells have at least 10

isoforms, fungi only have one or two PKCs (Levin, 2005; Shea and Poeta, 2006; Rhone and Poeta, 2009). *Saccharomyces cerevisiae* has only one PKC (*PKC1*) that has a serine/threonine domain, regulatory domains (C1, C2 and HR1), and a pseudosubstrate sequence. The *PKC1* C1 and HR1 domains interact with the small GTPase Rho1 (Nonaka *et al.*, 1995; Schmitz *et al.*, 2002). *PKC1* is involved in the regulation of cell construction via the activation of a MAP kinase signalling cascade in response to cell wall damage (Kamada *et al.*, 1995). Bck1p, the first component in the cell wall integrity (CWI) MAPK signalling cascade, activates Slp2p that phosphorylates and activates the transcription factor Rlm1p, which regulates the expression of genes whose products are important in cell wall biosynthesis (Jung and Levin, 1999). In the filamentous fungus *Aspergillus nidulans*, there are two putative PKC-encoding genes, *pkcA* (AN0106.3) and *pkcB* (AN5973.3) (Herrmann *et al.*, 2006; Ichinomiya *et al.*, 2007; Ronen *et al.*, 2007; Teepe *et al.*, 2007). Although sequence similarity of PkcB with other PKCs is limited to the catalytic region in the carboxy-terminus, PkcA has all the characteristic features of fungal PKCs, including an extended N-terminal regulatory domain containing C1, C2 and HR1 domains (Herrmann *et al.*, 2006). Knockdown of *pkcA* expression leads to reduced growth, conidiation and penicillin production (Herrmann *et al.*, 2006).

Recently, Teepe *et al.* (2007) shown that an *A. nidulans* *calC2* mutation that confers hypersensitivity to calcofluor white along with other drug sensitivities (indicating a defect in CWI and polar growth) was complemented by *pkcA*. The *pkcA* allele of the *calC2* strain contains a mutation predicted to introduce a charged arginine residue in place of neutral glycine at a conserved site located immediately beside the C1B regulatory domain. Ronen *et al.* (2007) and Ichinomiya *et al.* (2007) generated a conditional promoter mutant of *pkcA* and observe that its repression resulted in increased conidial swelling, decreased rates of hyphal growth, changes in the ultrastructure of the cell wall and increased sensitivity to anti-fungal agents.

Farnesol (FOH), a non-sterol isoprenoid produced by dephosphorylation of farnesyl pyrophosphate, has been

Accepted 15 September, 2010. *For correspondence. E-mail ggoldman@usp.br; Tel./Fax (+55) 16 36024280/4281/4311.

shown to inhibit proliferation and induce apoptosis in neoplastic cell lines and also to be effective in chemoprevention and chemotherapy in several *in vivo* cancer models (Adany *et al.*, 1994; Ohizumi *et al.*, 1995; Miquel *et al.*, 1996; He *et al.*, 1997; Hudes *et al.*, 2000; Burke *et al.*, 1997; Wiseman *et al.*, 2007; Joo and Jetten, 2010). In fungi, FOH plays a role as an extracellular quorum-sensing molecule in the dimorphic fungus *Candida albicans* (Nickerson *et al.*, 2006; Langford *et al.*, 2009). FOH inhibits the yeast to mycelium dimorphic transition and induces *C. albicans* to grow as actively budding yeasts (Hornby *et al.*, 2001). In *S. cerevisiae*, FOH blocks growth by raising the concentration of mitochondrial reactive oxygen species (ROS) (Machida *et al.*, 1998; 1999). FOH has no effect on *A. nidulans* hyphal morphogenesis, but it triggers morphological features characteristic of apoptosis through ROS accumulation (Semighini *et al.*, 2006; Savoldi *et al.*, 2008). In mammalian cells, FOH-mediated cell death can be reduced by the addition of phorbol esters (Voziyan *et al.*, 1995; Taylor *et al.*, 2005). Phorbol esters are non-metabolizable structural mimics of diacylglycerol that bind to and activate proteins containing C1 domains, such as PKCs (Yang and Kazanietz, 2003). There is an increase of FOH sensitivity when PKC inhibitors are added, suggesting a role for PKCs as the C1 domain-containing proteins that regulate FOH sensitivity (Voziyan *et al.*, 1995; Taylor *et al.*, 2005; Fairn *et al.*, 2007). Phosphatidylcholine (PC), phosphatidic acid (PA) and diacylglycerol (DAG) were able to prevent induction of apoptosis by FOH in mammalian cells (Taylor *et al.*, 2005). *S. cerevisiae* FOH-induced growth inhibition could be effectively prevented by the coaddition of a DAG analogue (Machida *et al.*, 1999). In *C. albicans*, FOH-growth defect was also relieved by addition of a DAG analogue, implicating phosphatidylinositol signalling in the delay (Uppuluri *et al.*, 2007). However, a mutant strain deleted for *PKC1* responded to FOH and DAG analogue similar to wild-type, suggesting the PKC is not the target of the DAG analogue. Fairn *et al.* (2007) provided a more substantial evidence of the involvement of the *PKC1* pathway in FOH sensitivity. These authors showed that Pkc1 relocalized to the mitochondria upon FOH addition and inactivation of the non-essential and non-redundant member of the Pkc1 signalling pathway, *BCK1*, resulted in FOH sensitivity. Furthermore, expression of activated alleles of *PKC1*, *BCK1* and *MKK1* increased tolerance to FOH and hydrogen peroxide. Savoldi *et al.* (2008) reported that *A. nidulans* PkcA *calC2* mutation was able to confer FOH tolerance. In contrast to *S. cerevisiae*, *A. nidulans* PkcA::GFP did not relocalize to the mitochondria upon exposure of germlings to FOH, but instead the fluorescence was completely relocalized to the entire cytoplasm. Here, we investigate in more detail the role played by *A. nidulans* *pkcA* in FOH tolerance. We demonstrate that

pkcA overexpression during FOH exposure causes increased cell death. FOH is also able to activate several markers of endoplasmic reticulum (ER) stress and the unfolded protein response (UPR). Our results suggest an intense cross-talk between PkcA, CWI and UPR during FOH-induced cell death.

Results

Overexpression of *pkcA* increases FOH sensitivity

We have previously shown that a point mutation (*calC2*) in *pkcA* was able to confer FOH tolerance (Savoldi *et al.*, 2008). We considered the possibility that PkcA overexpression would increase the FOH sensitivity. Thus, we constructed an *alcA::pkcA* mutant by replacing the endogenous *pkcA* promoter by the *alcA* promoter and verified its survival to FOH when PkcA is overexpressed. The *alcA* promoter is repressed by glucose, derepressed by glycerol and induced to high levels by ethanol or L-threonine (Flippin *et al.*, 2002). We were able to select a transformant that when grown in the presence of glycerol 2% as single carbon source for 16 h and transferred to glycerol 2% + ethanol 2% for 6 h at 37°C, the *pkcA* mRNA accumulates about 10-fold (Fig. 1A). When the *alcA::pkcA* mutant strain was grown in MM + glycerol 2% + ethanol 2% for 16 h at 37°C, the PkcA activity was about three times higher than in the wild-type strain (Fig. 1B). As previously reported by other groups (Ichinomiya *et al.*, 2007; Ronen *et al.*, 2007), the *alcA* repression by growing the *alcA::pkcA* mutant strain in the presence of 4% glucose, decreased dramatically the colony diameter (Fig. 1C), indicating once more that *pkcA* is an essential gene. The *pkcA* overexpression increased the susceptibility to FOH (Table 1 and Fig. 5D). When PkcA was overexpressed and the *alcA::pkcA* mutant strain was exposed to FOH, the PkcA activity increased only about 20% (compared with the *alcA::pkcA* mutant not exposed to FOH; Fig. 1B). There are no differences in the survival of both strains in the absence of FOH.

Along this manuscript, we evaluated *A. nidulans* cell death by using three different methods: nuclear condensation, germling viability and 10-fold dilutions of a starting suspension of conidia. Thus, to verify if the *alcA::pkcA* mutant is more sensitive to cell death caused by other stressing agents, we overexpressed PkcA and exposed conidia of this strain to several concentrations/dosages of amphotericin, itraconazole, caspofungin, camptothecin, hydroxyurea, 4-nitroquinoline oxide, sirolimus, calcium, menadione, hydrogen peroxide and paraquat. Overexpression of *pkcA* increased the survival to caspofungin and posaconazole but not for Congo red (A.C. Colabardini, unpublished results). The wild-type and the *alcA::pkcA* mutant strains showed comparable levels of susceptibility

Fig. 1. Overexpression of *pkcA* increases the susceptibility to FOH.

A. Wild-type and *alcA::pkcA* strains were grown for 16 h in MM + glycerol 2% at 37°C and transferred for MM + glycerol 2% + ethanol 2% and grown for further 6 h. The relative quantification of *pkcA* and tubulin gene expression was determined by a standard curve (i.e. C_T – values plotted against logarithm of the DNA copy number). The results are the means $\pm$ standard deviation of four sets of experiments.

B. Protein kinase C activity of the wild-type and *alcA::pkcA* strains exposed or not to FOH 50 μ M. The results are the means $\pm$ standard deviation of three sets of experiments. Statistical differences were determined by one-way analysis of variance (ANOVA) followed, when significant, by Newman–Keuls multiple comparison test, using GraphPad Prism statistical software (GraphPad Software, version 3, 2003).

C. Wild-type and *alcA::pkcA* strains were grown for 72 h at 37°C in either MM + glycerol 2% or MM + glucose 4%.

D. Wild-type strain was grown for 16 h in YG at 37°C and transferred for fresh YG with either 0, 10, 50 or 100 μ M FOH and grown for further 2 h. The relative quantification of *pkcA* and tubulin gene expression was determined by a standard curve (i.e. C_T – values plotted against logarithm of the DNA copy number). The results are the means $\pm$ standard deviation of four sets of experiments.

E. Activation of intracellular caspases and PI labelling in germlings grown for 16 h in MM + 2% glycerol + 100 mM threonine exposed or not to FOH, indicated by green or red fluorescence respectively. Bars, 5 μ m. This figure is available in colour online at wileyonlinelibrary.com.

to all these agents (data not shown). The *calC2* mutant that displays increased FOH tolerance also did not show any growth difference with the wild-type strain, except when exposed to FOH (see Table 1 and Fig. 5D). We also observed that both mRNA accumulation and PKC activity is increased by FOH (Fig. 1B and D).

A Poly-Caspases FLICA apoptosis detection kit showed the presence of green fluorescence in 80% of the FOH-exposed wild-type germlings ($n = 100$), indicating the activation of intracellular caspases, whereas no fluorescence was observed in the unexposed germlings (Fig. 1E, upper panel). Consistently, when the levels of *pkcA* mRNA are increased, there is green fluorescence even in the absence of FOH (Fig. 1E, lower panel). We examined a possible necrotic role in the FOH-cell death by staining protoplasts with propidium iodide (PI). When the wild-type

was grown in the presence of glycerol + 100 mM threonine and not exposed to FOH, approximately 10% of the germlings were labelled with PI (Fig. 1E); when the wild-type was exposed to FOH 50 μ M for 30 min, 30% of the germlings were labelled with PI (Fig. 1E). The *alcA::pkcA* mutant grown in the presence of glycerol + 100 mM threonine and not exposed to FOH showed 15% of the germlings labelled with PI (Fig. 1E); when the *alcA::pkcA* mutant was exposed to FOH 50 μ M for 30 min, 30% of the germlings were labelled with PI (Fig. 1E). All the germlings that showed positive labelling for FLICA also were positive for PI, and vice versa. We were not able to perform TUNEL and Annexin assays for the *alcA::pkcA* mutant because it is not possible to produce protoplasts from this mutant even when it is grown in the presence of glycerol (data not shown). These results indicate that FOH induces

E

Fig. 1. cont.

Table 1. Non-condensed nuclei (%) and viability (%) of wild-type and mutant strains exposed to different FOH concentrations.

Strains	Non-condensed nuclei (%)		Viability (%)	
	20 μ M FOH	50 μ M FOH	20 μ M FOH	50 μ M FOH
Wild-type (A89)	97.5 $\pm$ 0.2	31.0 $\pm$ 5.6	73.0 $\pm$ 2.1	31 $\pm$ 3.6
$\Delta mpkC$	20.5 $\pm$ 6.0	10.5 $\pm$ 4.9	36.0 $\pm$ 3.4	12.0 $\pm$ 3.2
$\Delta hogA$	95.0 $\pm$ 0.0	2.5 $\pm$ 0.7	11.0 $\pm$ 1.2	2.0 $\pm$ 0.8
$\Delta mpkA$	100.0 $\pm$ 0.0	79.0 $\pm$ 1.7	94.0 $\pm$ 6.9	68.5 $\pm$ 2.6
Wild-type (BPU1)	99.3 $\pm$ 0.1	53.0 $\pm$ 3.5	85.0 $\pm$ 4.5	53.5 $\pm$ 5.4
$\Delta attA$	100 $\pm$ 0.0	40.3 $\pm$ 2.7	71.0 $\pm$ 6.8	40.0 $\pm$ 4.5
$\Delta napA$	99.7 $\pm$ 0.1	54.3 $\pm$ 4.2	79.0 $\pm$ 5.8	54.2 $\pm$ 2.9
Wild-type (CLK3)	100.0 $\pm$ 0.0	29.3 $\pm$ 1.1	71.0 $\pm$ 4.9	31.0 $\pm$ 2.5
$\Delta srrA$	100.0 $\pm$ 0.0	34.6 $\pm$ 4.5	72.0 $\pm$ 5.1	36.2 $\pm$ 3.6
$\Delta sskA$	98.0 $\pm$ 2.0	34.0 $\pm$ 6.0	64.0 $\pm$ 3.0	34.5 $\pm$ 4.1
$\Delta srrA \Delta sskA$	98.7 $\pm$ 0.2	43.7 $\pm$ 4.8	81.0 $\pm$ 1.5	42.5 $\pm$ 6.7
Wild-type (GR5)	98.7 $\pm$ 0.2	22.0 $\pm$ 5.3	93.0 $\pm$ 1.9	61.0 $\pm$ 2.9
$calC2$	99.0 $\pm$ 0.0	99.0 $\pm$ 0.0	98.0 $\pm$ 0.0	97.0 $\pm$ 1.0
$alcA::pkcA$	23.0 $\pm$ 0.8	11.7 $\pm$ 5.1	65.0 $\pm$ 3.7	49.0 $\pm$ 4.1
$alcA::pkcA \Delta casA$	83.0 $\pm$ 0.9	50.0 $\pm$ 2.9	90.0 $\pm$ 7.3	67.0 $\pm$ 1.9
$\Delta casA$	99.0 $\pm$ 0.1	78.0 $\pm$ 1.9	95.0 $\pm$ 1.7	71.0 $\pm$ 2.5
$\Delta casB$	69.0 $\pm$ 2.8	7.0 $\pm$ 4.0	56.0 $\pm$ 4.9	23.0 $\pm$ 1.5
$\Delta casA \Delta casB$	92.0 $\pm$ 0.1	72.0 $\pm$ 1.2	84.0 $\pm$ 3.5	58.0 $\pm$ 1.6

caspase-like proteases and necrosis in *A. nidulans*; however, increased PkcA expression does not enhance the necrotic process but activates caspase-like proteases. In addition, increased levels of PkcA expression enhance the *A. nidulans* FOH-induced cell death and influence of PkcA on cell death seems to be specifically related to FOH.

FOH promotes the UPR

Recent studies have revealed that induction of ER stress and the subsequent activation of the UPR play a critical role in the induction of apoptosis by FOH in lung carcinoma cells (for a review, see Joo and Jetten, 2010). Based on this evidence, we raised two hypotheses: (i) FOH is affecting ER stress and activating the UPR in *A. nidulans* and (ii) the specific effect of PkcA on FOH-induced cell death could be due to a synergistic combination of PkcA activity and FOH on ER stress and activation of UPR. To start addressing these hypotheses, we overexpressed PkcA in the presence of agents that disrupt endoplasmic reticulum homeostasis (ER), such as 2-deoxy-D-glucose (2-DG) and dithiothreitol (DTT). These two agents induce ER stress by impairing the N-linked glycosylation that promotes folding of nascent polypeptide chains in the ER and unfolding proteins directly by reducing disulphide bonds respectively (Helenius and Aebi, 2004; Back *et al.*, 2005). The overexpression of PkcA increases the susceptibility of $alcA::pkcA$ mutant strain to these agents (Fig. 2A and B).

The second hypothesis suggests that *A. nidulans* PkcA is influencing the UPR. The accumulation of aberrant folded proteins in the ER activates the bifunctional trans-

membrane kinase/endoribonuclease Ire1 (Korennykh *et al.*, 2009). Ire1 excises an intron from yeast *HAC1* cytoplasmic precursor mRNA *HAC1u* (uninduced), removing it to generate the induced form of the *HAC1* mRNA, *HAC1i* (induced) (Cox and Walter, 1996). This creates a frame-shift in the mRNA, allowing for the translation of a transcription factor that moves to the nucleus and regulates the expression of UPR target genes (Travers *et al.*, 2000; Kimata *et al.*, 2006). Recently, a homologue of *HAC1* was identified in *A. fumigatus* (Richie *et al.*, 2009). The *A. fumigatus* $\Delta hacA$ mutant was unable to activate the UPR in response to ER stress and was hypersensitive to agents that disrupt ER homeostasis or the cell wall. We identified an *A. nidulans* *HAC1* homologue, *hacA* (AN9397.4), that displays about 75% identity and 83% similarity with *A. fumigatus* *hacA* (7e-155) and about 40% identity and 56% similarity with *S. cerevisiae* *HAC1* (2e-11). In *S. cerevisiae*, the Ire1p-mediated splicing of a 252-nucleotide (nt) unconventional intron from *HAC1* mRNA relieves the transcript from a translational block (Kawahara *et al.*, 1997; Rueggsegger *et al.*, 2001). In contrast, in mammalian cells, the *HAC1* homologue *XBP1* is activated in a similar way by the splicing of a 26 nt unconventional intron (Lee *et al.*, 2002), whereas *Trichoderma reesei* *hac1* and *A. nidulans*, *A. niger* and *A. fumigatus* *hacA* are activated by the splicing of a 20 nt unconventional intron (Saloheimo *et al.*, 2003; Mulder *et al.*, 2004; Richie *et al.*, 2009). We developed an RT-PCR assay to distinguish the increased accumulation of *A. nidulans* *hacAⁱ* (Fig. 2C and D). First, we exposed *A. nidulans* to 1 mM DDT for 60 min and verified a 6.0 and 3.5 times increase of the mRNA accumulation of *hacAⁱ* in the wild-type and *calC2* mutant strains respectively (Fig. 2D).

Fig. 2. *hacA'* mRNA expression is increased in the presence of FOH.

A. Wild-type and *alca::pkcA* strains were grown for 96 h at 37°C in liquid MM + glycerol 2% + 100 mM threonine in the presence of DTT 0.5 or 1.0 mM.

B. Wild-type and *alca::pkcA* strains were grown in MM + 2% glycerol + threonine 100 mM in the presence of increasing concentrations of 2-DG.

C. DNA sequences that border the intron processing of the *A. nidulans hacA* gene. Arrows indicate the processing of the 20 bp intron from *hacA^u* (uninduced) to the *hacA'* (induced) mRNA. PCR amplification of the *hacA* and *tubC* (encoding β -tubulin) cDNAs when *A. nidulans* wild-type and *calC2* mutant strains were exposed to DTT (D) and FOH (E). This figure is available in colour online at [wileyonlinelibrary.com](http://www.wileyonlinelibrary.com).

When the wild-type strain was exposed to increasing concentrations of FOH, there was a *hacAⁱ* mRNA accumulation of about 0.5, 2.5 and 3.5 times at 10, 50 and 100 μ M FOH respectively. In contrast, the *calC2* mutant strain did not show any difference in the *hacAⁱ* mRNA accumulation when exposed to FOH (Fig. 2E). By overexpressing *pkcA* in the absence of FOH, we were able to see a three-fold increase in the *hacAⁱ* mRNA accumulation (Fig. 2E). The *hacA* levels of *alcA::pkcA* and its corresponding wild-type are lower than the wild-type exposed or not to FOH

because in the former the strains were grown in MM supplemented with 2% glycerol plus threonine 100 mM while the later was grown in complete medium. We also investigated if FOH led to induction of the UPR as judged by the transcriptional induction of *A. nidulans* homologues of *A. niger* genes known to be affected by ER stress, *bipA*, *pdiA* and *pdiB* (Guillemette *et al.*, 2007). These genes showed diverse degrees of increased mRNA accumulation when *A. nidulans* was exposed to different FOH concentrations (*A. nidulans* homologues AN2062.4,

Fig. 2. *cont.*

AN7436.4 and AN0248.4; Fig. 3A). FOH also enhances the mRNA accumulation of other genes involved in the UPR, such as the transcription factor Hsf1, and the chaperones Hsp90 and Hsp104 (*A. nidulans* homologues AN8035.4, AN8269.4 and AN0858.4 respectively; Fig. 3B).

Taken together, these data suggest FOH is activating the UPR and that PkcA plays a role in the activation of this process in *A. nidulans*.

Metacaspases are induced by FOH

The caspases are members of a family, or structurally related group, known as cysteine proteases (for a review

see Mazzoni and Falcone, 2008). A cell death stimulus activates a cascade of caspases that ultimately cleaves substrates essential for cell death (Donepudi and Grutter, 2002; Degterev *et al.*, 2003; Lavrik *et al.*, 2005). Richie *et al.* (2007) showed that *A. fumigatus* metacaspases CasA and CasB play a role under conditions of ER stress. Thus, we decide to investigate a possible connection among *A. nidulans* PkcA, FOH-induced cell death and metacaspases. In *A. nidulans*, we have identified two putative caspase-like proteins, fitting into the type I category of metacaspases, and named them *casA* (AN2503.3) and *casB* (AN5712.3). The *A. nidulans* CasA and CasB have about 64% identity and 76% similarity (*e*-value: 2×10^{-30}), and *S. cerevisiae* Yca1p metacaspase

Fig. 3. Several genes involved in the unfolded protein response have increased mRNA accumulation when *A. nidulans* is exposed to FOH. Wild-type strain was grown for 16 h in YG at 37°C and transferred for fresh YG with either 0, 10, 50 or 100 μM FOH and grown for further 2 h. The relative quantification of (A) *bipA* (AN2062.4), *pdiA* (AN7436.4) and *pdiB* (AN0248.4), and (B) *hsf1* (AN8035.4), *hsp90* (AN8269.4) and *hsp104* (AN0858.4) and tubulin gene expression was determined by a standard curve (i.e. C_T – values plotted against logarithm of the DNA copy number). The results are the means $\pm$ standard deviation of four sets of experiments.

has great similarity to both CasA (57% identity, 70% similarity, e -value: $8e-21$) and CasB (58% identity, 71% similarity, e -value: $1e-101$). When the wild-type strain is exposed to FOH, there is an increased accumulation of *casA* mRNA of 27.0 and 137.5 times, and *casB* mRNA of 2.2, 8.0 and times at 10 and 50 μM FOH respectively (Fig. 4A). PkcA overexpression increases the level of mRNA accumulation of both *casA* and *casB* mRNA by 2, and 4.5 times, respectively, even in the absence of FOH (Fig. 4B). To gain more insight on *A. nidulans* metacaspase function, we constructed deletion strains for both *casA* and *casB* genes; we also constructed a double mutant strain by crossing both $\Delta casA$ and $\Delta casB$ deletion mutants. None of these mutants displayed either growth reduction or any other apparent phenotype when compared with the corresponding wild-type strain.

We also investigated if necrosis and other caspase-like proteases would be present upon FOH-cell death of the metacaspase mutants by staining with PI and FLICA. When the wild-type and $\Delta casA$ mutant were grown in complete medium and not exposed to FOH, approximately 5% and 0% of the germlings showed PI and FLICA labelling respectively (Fig. S1). Germlings of the $\Delta casB$ and $\Delta casA \Delta casB$ mutants not exposed to FOH showed about 13–15% PI and 15% FLICA labelling (Fig. S1). Upon exposure to FOH 50 μM for 30 min, the wild-type showed germlings with 15% PI and 15% FLICA labelling (Fig. S1). The $\Delta casB$ and $\Delta casA \Delta casB$ mutants exposed to FOH displayed about 70% PI and 100% FLICA labelling (Fig. S1). Interestingly, when the $\Delta casA$ mutant was exposed to 50 μM FOH for 30 min, it showed germlings with 50% PI and 60% FLICA labelling (Fig. S1). We have not observed any PI minus FLICA positive cells. Curiously, these results suggest that the activation of necrosis and caspase-like proteases by FOH in *A. nidulans* is dependent to metacaspases since the absence of at least one of the metacaspase genes increases PI and FLICA labelling. However, *casB* seems to play a more important role in this activation because in its absence the levels of activation are even higher.

We investigated in the detail the viability of germlings of *A. nidulans* $\Delta casA$, $\Delta casB$ and $\Delta casA \Delta casB$ mutants comparatively to the wild-type strain. To verify if these mutants are more sensitive to cell death caused by other oxidative stressing agents, we exposed conidia of these strains to several concentrations/dosages of the agents mentioned earlier in the section *Overexpression of pkcA increases FOH sensitivity*. The wild-type and the $\Delta casA$ mutant strains showed comparable levels of susceptibility to all these agents (data not shown). However, when the mutant strains were exposed to ER disrupting agents, such as 2-DG and DTT, they showed differential susceptibility (Fig. 5A and B). The $\Delta casA$ and $\Delta casA \Delta casB$ strains were as resistant to DTT as the wild-type strain

Fig. 4. Poly-Caspases FLICA showed the presence of increased green fluorescence when PkcA mRNA expression is increased. A. Wild-type strain was grown for 16 h in YG at 37°C and transferred for fresh YG with either 0, 10, 50 or 100 μM FOH and grown for further 2 h. The relative quantification of *casA-B* and tubulin gene expression was determined by a standard curve (i.e. C_T – values plotted against logarithm of the DNA copy number). The results are the means $\pm$ standard deviation of four sets of experiments.

B. Wild-type and *alcA::pkcA* strains were grown for 16 h in MM + glycerol 2% at 37°C and transferred for MM + glycerol 2% + threonine 100 mM and grown for further 6 h. The relative quantification of *casA-B* and tubulin gene expression was determined by a standard curve (i.e. C_T – values plotted against logarithm of the DNA copy number). The results are the means $\pm$ standard deviation of four sets of experiments.

while $\Delta casB$ was more sensitive than the wild-type strain (Fig. 5A). In contrast, $\Delta casA$ was as resistant to 2-DG as the wild-type while $\Delta casB$ and $\Delta casA \Delta casB$ were more sensitive than the wild-type strain (Fig. 5B). Exposing the germ tubes to 20 or 50 μM FOH for 2 h revealed that $\Delta casA$ was significantly more tolerant to FOH than the wild-type, while $\Delta casB$ and $\Delta casA \Delta casB$ mutant strains were less tolerant to FOH than the wild-type and $\Delta casA$ mutant strain (Table 1 and Fig. 5C). We constructed by sexual crosses a double mutant *alcA::pkcA* $\Delta casA$ and evaluated its sensitivity to FOH. We overexpressed the *pkcA* gene in this strain and compared its tolerance to FOH with the *alcA::pkcA* and $\Delta casA$ mutant strains (Table 1 and Fig. 5D). The absence of *casA* confers increased tolerance to FOH when *pkcA* is overexpressed (Table 1 and Fig. 5D). The complemented metacaspase mutants showed the same phenotypes of the wild-type strain (Table 1; Figs S2 and S4).

The results demonstrated FOH-induced upregulation of both *casA-B* genes and that FOH-induced cell death may

be dependent on metacaspases. Furthermore, the overexpression of *pkcA* increases both mRNA accumulation and metacaspases activity, and there is a genetic interaction between PkcA and CasA.

FOH susceptibility is mediated via PKC/MAPK

Mitochondria and ROS participate in FOH-induced apoptosis in *A. nidulans* (Semighini *et al.*, 2006; Savoldi *et al.*, 2008). Two-component systems (Histidine-to-Aspartate phosphorelay) are signal transduction mechanisms involved in several stimuli, including oxidative stress responses (Wuichet *et al.*, 2010; Bahn, 2008). *A. nidulans* has a histidine kinase (NikA) and two response regulators (SskA and SrrA) implicated in oxidative stress responses (Hagiwara *et al.*, 2007; Vargas-Pérez *et al.*, 2007). In *A. nidulans*, the HogA/MAPK kinase cascade plays important roles downstream of the NikA-SskA phosphorelay (Moye-Rowley, 2003; Furukawa *et al.*, 2005; Ikner and Shiozaki, 2005). In this species, it is

Fig. 5. *for cont.*

Fig. 5. *A. nidulans* metacaspases CasA and CasB are involved in FOH-induced cell death.

A and B. (A) Wild-type, $\Delta casA$, $\Delta casB$, $\Delta casA \Delta casB$ mutant strains were grown for 96 h at 37°C in YG medium in the presence of DTT 0.5 or 1.0 mM, or (B) MM + 2-DG 0.3 or 0.5 mM.

C and D. Aliquots (5 μ l) of 10-fold dilutions derived from a starting suspension of 1.0×10^8 conidia ml^{-1} of the indicated strain were spotted on YG-agar plates supplemented with the 0, 0.5 or 1.0 mM FOH. The plates were incubated at 37°C for 48 h.

known that NapA and AtfA are transcription factors activated by the MAPK cascade that are involved in stress responsive transcriptional regulation to oxidative stress (Asano *et al.*, 2007; Hagiwara *et al.*, 2008; Balázs *et al.*,

2010). Thus, we investigated the involvement of SskA-SrrA, AtfA^{YAP1} (the *YAP1* homologue), and NapA signal transduction pathway on FOH tolerance by verifying their viability after exposure to FOH (Table 1 and Fig. 6).

Fig. 6. FOH-susceptibility of mutants from the MAP kinase pathway. Aliquots (5 μ l) of 10-fold dilutions derived from a starting suspension of 1.0×10^8 conidia ml^{-1} of the corresponding wild-type, $\Delta sskA$, $\Delta srrA$, $\Delta sskA \Delta srrA$ (A), $\Delta napA$ and $\Delta atfA$ (B), and $\Delta mpkA$, $\Delta mpkC$ and $\Delta hogA$ (C) strains were spotted on YG-agar plates supplemented with the 0, 0.5 or 1.0 mM FOH. The plates were incubated at 37°C for 48 h.

When exposed to lower concentrations of FOH (20 and 50 μ M), the *ssrA*, *sskA*, *ssrA sskA*, *napA* and *atf* deletion mutants are as sensitive to FOH as the wild-type strain (Table 1). However, when exposed to higher concentrations of FOH (0.5 and 1.0 mM), the *ssrA*, *sskA*, *ssrA sskA* and *atf* deletion mutants are more sensitive to FOH than the wild-type strain (Fig. 6A and B). There are four MAPK kinases in *A. nidulans*: MpkA-C and SakA/HogA (for a review see May, 2008). We evaluate the FOH sensitivity of deletion mutants for three of these kinases, MpkA, MpkC and HogA (Table 1 and Fig. 6C). Interestingly, the Δ *mpkA* mutant shows increased tolerance while Δ *mpkC* and Δ *hogA* showed increased sensitivity to FOH (Table 1 and Fig. 6C). These results suggest SsrA, SskA, AtfA and MAP kinases are involved in the signal transduction in response to the effects caused by FOH.

FOH activates the CWI pathway

In *S. cerevisiae* PKC1 is involved in the regulation of cell construction via the reactivation of *MPK1* and activation of the transcription factor Rlm1p, which regulates the expression of genes whose products are important in cell wall biosynthesis (Jung and Levin, 1999). Fujioka *et al.* (2007) demonstrated that *A. nidulans* *rlmA* and *mpkA* are functional orthologues of *S. cerevisiae* *RLM1* and *MPK1*. Considering that Δ *mpkA* mutant is more resistant to FOH, we speculated if FOH is activating the CWI pathway. Therefore, we tested the mRNA accumulation of *agsA* (AN5885.3; encoding one of the *A. nidulans* α -1,3-glucan synthases), *rlmA* (AN2984.3) and *mpkA* (AN5666.3) genes when *A. nidulans* is exposed to different concentrations of FOH (Fig. 7A). All three genes showed increased mRNA accumulation upon *A. nidulans* exposure to different FOH concentrations. We also tested if the mRNA levels of these genes could be controlled directly or indirectly by PkcA by overexpressing PkcA even in the absence of FOH. Again all three genes showed increased mRNA accumulation when *A. nidulans* PkcA was overexpressed (Fig. 7B). Previously, it was shown that cell wall stress results in the phosphorylation of *A. nidulans* MpkA (Fujioka *et al.*, 2007). We checked if FOH can activate the MpkA phosphorylation by exposing *A. nidulans* cultures to different FOH concentrations (Fig. 7C). Addition of Congo red induced phosphorylation of *A. nidulans* MpkA (Fig. 7C, first row). Interestingly, increasing FOH concentrations also induced *A. nidulans* MpkA phosphorylation (Fig. 7C, second row). MpkA was also phosphorylated when *A. nidulans* was exposed to DTT (Fig. 7C, third row). These results strongly indicate that FOH is able to activate the CWI pathway and that there is a possible cross-talk between CWI and UPR pathways.

Discussion

In mammalian cells, the PKC family consists of 10 related serine/threonine kinases, which are important controllers of cell proliferation, survival and cell death (Reyland, 2009). The *PKC1* gene was originally identified in *S. cerevisiae* (Levin, 2005) and is a close homologue of the human PKC gene. In *A. nidulans*, the PKC/MAPK pathway is involved in the CWI, polarized growth and morphogenesis (Ronen *et al.*, 2007; Teepe *et al.*, 2007), utilization of polyalcohol sugars (Reyes *et al.*, 2006), osmotic and oxidative stresses (Davenport *et al.*, 1995; Han and Prade, 2002; Kawasaki *et al.*, 2002; Furukawa *et al.*, 2005), sexual development and spore viability (Kawasaki *et al.*, 2002; Ichinomiya *et al.*, 2007), and penicillin production (Herrmann *et al.*, 2006). Here, we showed a completely novel feature of the *A. nidulans* PKC/MAPK pathway related to FOH and the UPR. Interestingly, Binder *et al.* (2009) have recently reported that the antifungal protein PAF also interferes with PKC/MAPK pathway.

The PKC/MAPK pathway is required for transducing the response to FOH in mammalian cells, *S. cerevisiae*, *C. albicans* and *A. nidulans* (Voziyan *et al.*, 1995; Yang and Kazanietz, 2003; Taylor *et al.*, 2005; Fairn *et al.*, 2007; Uppuluri *et al.*, 2007; Savoldi *et al.*, 2008; Román *et al.*, 2009). However, it is not very clear how PKC/MAPK could mediate this response. We observed that a mutation in the *pkcA* (*calC2*), which confers sensitivity to calcofluor and SDS, conferred FOH tolerance. By investigating a conditional mutant of *pkcA* and other MAP kinase mutants, we identified a relationship between the CWI, UPR and FOH. Actually, there are several pieces of evidence in the literature showing a connection among these pathways. Mammalian PKC θ is required for autophagy in response to stress in the ER (Sakaki and Kaufman, 2008; Sakaki *et al.*, 2008), while PKC δ communicates ER stress to the mitochondria (Qi and Mochly-Rosen, 2008). In *S. cerevisiae*, ER stress has been shown to trigger signalling through the CWI/MAPK signalling cascade (Bonilla and Cunningham, 2003; Chen *et al.*, 2005; Nita-Lazar and Lennarz, 2005; Pal *et al.*, 2007; Wright *et al.*, 2007; Shaner *et al.*, 2008). More recently, strong evidences consolidated the hypothesis that cellular responses to ER and cell wall stress are co-ordinated to buffer the cell against these two unrelated cellular stresses (Krysan, 2009; Scrimale *et al.*, 2009). Another interesting connection between ER stress and MAP kinases was shown by Bicknell *et al.* (2010). These authors demonstrated that *S. cerevisiae* Hog1 MAP kinase becomes phosphorylated during the late stage of ER stress. *S. cerevisiae* Hog1 co-ordinates a multifaceted response to persistent ER stress activating autophagy by enhancing the stability of Atg8, a critical autophagy protein. Furthermore, the *A. fumigatus* Δ *hacA* mutant was unable to activate the UPR

Fig. 7. FOH activates the cell wall integrity (CWI) pathway.

A. Wild-type strain was grown for 16 h in YG at 37°C and transferred for fresh YG with either 0, 10, 50 or 100 μM FOH and grown for further 2 h. The relative quantification of *agsA*, *rlmA* and *mpkA* and tubulin gene expression was determined by a standard curve (i.e. C_T – values plotted against logarithm of the DNA copy number). The results are the means $\pm$ standard deviation of four sets of experiments.

B. Wild-type and *alcA::pkcA* strains were grown for 16 h in MM + glycerol 2% at 37°C and transferred for either MM + glucose 4% (G), or MM + glycerol 2% (Gly), or MM + glycerol 2% + ethanol 2% (GE) and grown for further 6 h. The relative quantification of *agsA*, *rlmA* and *mpkA* and tubulin gene expression was determined by a standard curve (i.e. C_T – values plotted against logarithm of the DNA copy number). The results are the means $\pm$ standard deviation of four sets of experiments.

C. FOH activates the CWI pathway. A. *nidulans* overnight cultures grown at 37°C were exposed to the indicated concentrations of the indicated agents. Protein extracts were analysed by Western blotting using antibodies directed against phosphorylated MpkA or β -tubulin (TubC) for loading control.

in response to ER stress and was hypersensitive to agents that disrupt ER homeostasis or the cell wall (Richie *et al.*, 2009). Mutations of *HAC1* homologue in *Candida albicans* also lead to increased susceptibility to the clinically used, cell wall-directed antifungal caspofungin (Wimalasena *et al.*, 2008). Our results strongly suggest

that FOH is inducing the UPR in *A. nidulans* and this response is mediated via PKC/MAPK pathway.

FOH-induced apoptosis in human lung carcinoma cells analysed by microarray analysis revealed that many genes implicated in ER stress were highly upregulated, suggesting that FOH-induced apoptosis in these cells is coupled to

Fig. 7. cont.

the activation of an ER stress response pathway and the UPR (Joo *et al.*, 2007). The precise mechanism how FOH induces ER stress has not been elucidated yet (Joo and Jetten, 2010). Actually, the UPR is not the single target for FOH either in mammalian cells or fungi (Miquel *et al.*, 1996; Semighini *et al.*, 2006; Savoldi *et al.*, 2008; Langford *et al.*, 2009; Joo and Jetten, 2010). In mammalian cells, these include regulation of 3-hydroxy-3-methylglutaryl-CoA reductase and CTP:phosphocholine cytidyltransferase α , rate-limiting enzymes in the mevalonate pathway and phosphatidylcholine biosynthesis, respectively, the generation of ROS, and the activation of the NF- κ B signalling pathway and a number of NF- κ B target genes (Joo and Jetten, 2010). Previously, we investigate which pathways are influenced through FOH by examining the transcriptional profile of *A. nidulans* exposed to this isoprenoid (Savoldi *et al.*, 2008). We observed decreased mRNA abundance of several genes involved in RNA processing and modification, transcription, translation, ribosomal structure and biogenesis, amino acid transport and metabolism, and ergosterol biosynthesis. We also observed increased mRNA expression of genes encoding a number of mitochondrial proteins and characterized in detail one of them, the *atfA*, encoding the apoptosis-inducing factor (AIF)-like mitochondrial oxidoreductase. However, surprisingly we have not observed any gene in the CWI and UPR with increased mRNA accumulation, except for chaperone-encoding genes (Savoldi *et al.*, 2008). This could be due to the low resolution of the microarray slides used for these experiments (*A. nidulans* oligonucleotide slides version 1 for microarray hybridizations; for details see http://pfgrc.jcvi.org/index.php/microarray/array_description/aspergillus_nidulans/version1.html).

Interestingly, *A. nidulans* PkcA overexpression activates the UPR in the absence of FOH. This suggests that PkcA activity already induces some major stresses on the cells independently of FOH. FOH exposure has an additive lethal effect what strongly indicates there is a connection among PkcA, CWI and UPR in *A. nidulans*. It is well known that FOH increases ROS production in *A. nidulans* (Semighini *et al.*, 2006; Savoldi *et al.*, 2008; Dinamarco *et al.*, 2010). We have not observed increased ROS susceptibility when PkcA is overexpressed (data not shown). However, *pkcA* mRNA accumulation is increased in the presence of oxidative stress (Fig. S3). Gerik *et al.* (2008) observed that besides its role in the activation of CWI, *Cryptococcus neoformans* PKC1 is essential for defence against both oxidative and nitrosative stresses. Vilella *et al.* (2005) also showed the involvement of Pkc1 in oxidative stress but could not detect a role for this protein in the nitrosative stress response. Thus it is possible that the connection among PkcA and UPR could be mediated through ROS formation upon FOH exposure, by not only

activating PkcA activity but also the UPR. The environment of the ER is oxidizing, supporting the formation of intra- and interchain disulphide bonds that serve to stabilize the folding and assembly of nascent proteins (Shimizu and Hendershot, 2009). Each disulphide bond that forms during oxidative folding should produce a single ROS (Shimizu and Hendershot, 2009). Increased ROS production through FOH will create an intense oxidized environment in the ER with increased production of mis-folded proteins, activating the UPR.

Recently, Dichtl *et al.* (2010) provided evidence that FOH is able to interfere with *A. fumigatus* CWI signalling. These authors showed that FOH inhibits the activation of the CWI pathway and weakens the cell wall, leading to the exit of *AfRho1* and *AfRho3* from the hyphal tip. They also observed a displacement of the actin-binding protein tropomyosin from the apical tip. Furthermore, they showed that $\Delta mkk2$ and $\Delta mpkA$ mutants were highly susceptible to FOH. We have also seen that *A. nidulans* $\Delta mpkC$ and $\Delta hogA$ mutants are FOH-sensitive while $\Delta mpkA$ is resistant to FOH. These results suggest a different mechanism of regulation for FOH-cell death induced in *A. fumigatus* and *A. nidulans*. However, in both organisms the signal transduction for FOH-cell death is mediated via MAP kinases. The *A. nidulans* YAP1 homologue, $\Delta atfA$ is also more susceptible to FOH, suggesting that the FOH response could be mediated through PKC/MAPK/YAP1 pathway. There is also MpkA phosphorylation and increased mRNA accumulation of *A. nidulans* genes involved in the CWI pathway, such as *agsA*, *rlmA* and *mpkA*, both upon FOH exposure and PkcA overexpression. However, we have not addressed here more specifically how FOH is activating the *A. nidulans* CWI pathway.

Caspases are members of a family known as cysteine proteases that are actively involved in cell death in eukaryotes (for a review see Li and Yuan, 2008). Although *S. cerevisiae* does not have caspases in its genome, a caspase-like protein named YCA1, fitting into the type I category of metacaspases, was identified and characterized (Uren *et al.*, 2000; Madeo *et al.*, 2002; Mazzoni and Falcone, 2008). Previously, Richie *et al.* (2007) have identified *A. fumigatus* metacaspases CasA and CasB. These authors have not observed any difference in sensitivity to a range of pro-apoptotic stimuli, but the double *casA casB* deletion mutant showed a growth disadvantage in the presence of agents that disrupt ER, such as 2-DG and DTT. ER stress caused by an increase of unfolded or misfolded proteins activates apoptotic pathways (Momoi, 2004; Rasheva and Domingos, 2009). A subset of caspases, some of them located in the ER, are activated during ER stress (Momoi, 2004; Rasheva and Domingos, 2009). These results suggested a protective role for *A. fumigatus* metacaspases during ER stress rather than a

death-inducing function (Richie *et al.*, 2007). Accordingly, we have also not observed any growth differences among the *A. nidulans* wild-type, single and double *casA-B* deletion mutants for several apoptotic stimuli, except for FOH and ER-disrupting agents. Curiously, the observed effects in the presence of these agents for $\Delta casA$ and $\Delta casB$ were absolutely contrasting, i.e. $\Delta casA$ mutation confers increased survival to them while $\Delta casB$ mutation decreased its viability. These results suggest *A. nidulans* CasA and CasB have different functions. The decreased susceptibility of *A. nidulans* $\Delta casA$ mutant to ER-disrupting agents and FOH is not due to a protective role but rather to an active cell death role for CasA. Thus, it is tempting to speculate that its paralogue, CasB, would keep a function as a protecting agent against ER stress while CasA could have evolved a different function as a cell death effector. Our findings reveal that FOH induces a caspase-mediated process of apoptosis in *A. nidulans*. The same caspase dependency of FOH-induced cell death was recently observed for *C. albicans* (Shirtliff *et al.*, 2009). An interesting and novel aspect of our work was to show a genetic interaction between PkcA and metacaspases.

Experimental procedures

Strains, media and culture methods

Aspergillus nidulans strains used are described in Table S1. The media used were of two basic types, i.e. complete and minimal. The complete media comprised the following three variants: YAG (2% w/v glucose, 0.5% w/v yeast extract, 2% w/v agar, trace elements), YUU [YAG supplemented with 1.2 g l⁻¹ (each) of uracil and uridine], and liquid YG or YG + UU medium with the same composition (but without agar). The minimal media were a modified minimal medium (MM; 1% w/v glucose, original high-nitrate salts, trace elements, 2% w/v agar, pH 6.5). Expression of *pkcA* gene under the control of *alcA* promoter was regulated by carbon source: repression on glucose 4% w/v, derepression on glycerol, and induction on ethanol or threonine. Therefore, MM + Glycerol and MM + Ethanol were identical to MM, except that glycerol (2% v/v) and/or ethanol (2% v/v) or threonine (100 mM) were used, respectively, in place of glucose as the sole carbon source. Trace elements, vitamins, and nitrate salts were included as described by Kafer (1977). Strains were grown at 37°C unless indicated otherwise.

For the FOH viability assay (Noventa-Jordão *et al.*, 1999), 1×10^6 conidia ml⁻¹ were incubated in YG + UU for 5 h at 30°C in a reciprocal shaker (250 r.p.m.). After this period, FOH 50 µM was added to the germlings and they were allowed to grow for additional 120 min, respectively, at the same conditions. Conidia were conveniently diluted and plated in YAG + UU plates. Plates were incubated at 37°C for 48 h. Viability was determined as the percentage of colonies on treated plates compared with untreated controls. For the viability assays using 10-fold dilutions of a starting suspension of conidia, conidia were collected from 5-day-old cul-

tures grown on YAG or YUU agar plates. The amount of conidia was calculated using a counting chamber. We spotted 10⁸, 10⁷, 10⁶ and 10⁵ conidia in a volume of 5 µl on YUU agar plates that were incubated at 37°C for 48 h.

In vivo detection of caspase activation

Activated caspases in *A. nidulans* germlings were detected microscopically using a Poly-Caspases FLICA apoptosis detection kit (Immunochemistry Technologies, LLC) according to the manufacturer's recommendation. Briefly, 16-hour-old *A. nidulans* germlings were exposed to FOH 50 µM for 0 and 30 min at 30°C. Germlings with intracellular active caspases fluoresce green, whereas non-apoptotic cells appear unstained. Thirty microlitres of FLICA reagent was dropped on coverslips with adherent germlings and incubated at 37°C for 1 h in the dark. Then, they were briefly rinsed with PBS buffer (140 mM NaCl, 2 mM KCl, 10 mM NaHPO₄, 1.8 mM KH₂PO₄, pH 7.4), and the slides were treated with a 10 mg ml⁻¹ solution of RNase A for 1 h at 37°C and stained with a solution containing 1.25 µg of PI per millilitre. For the positive control with PI, coverslips containing germlings were fixed for 3 min at room temperature in a solution containing 3.7% formaldehyde, 0.2% Triton X-100, and 50 mM phosphate buffer, pH 7 (Bhabhra *et al.*, 2004). The slides were visualized with an epifluorescence microscope, using a Carls Zeiss (Germany) microscope using 100× magnification oil immersion objective lens (EC Plan-Neofluar, NA 1.3) equipped with a 100W HBO mercury lamp epifluorescence module. Phase contrast for the brightfield images and fluorescent images were captured with a AxioCan camera (Carl Zeiss), processed using the AxioVision software version 3.1 and saved as tiff files. Further processing was performed using Adobe Photoshop 7.0 (Adobe Systems Incorporated, CA, USA).

Molecular techniques

Standard genetic techniques for *A. nidulans* were used for all strain constructions and transformations (Kafer *et al.*, 1977). DNA manipulations were according to Sambrook and Russell (2001). All PCR reactions were performed using Platinum Taq DNA Polymerase High Fidelity (Invitrogen).

For the DNA-mediated transformation, the deletion cassettes were constructed by 'in vivo' recombination in *S. cerevisiae* as previously described by Colot *et al.* (2006). Briefly, about 1.5 kb regions on either side of the ORFs were selected for primer design. For every construction, the primers were named as 5F and 5R in that the 5F primer contains a short homologue sequence to the MCS of the plasmid pRS426. Both primers were used to amplify the 5'-UTR flanking region of the targeted ORF. Likewise, the primers 3F and 3R were used to amplify the 3'-UTR ORF flanking region and the 3R primer also contains a short homologue sequence to the MCS of the plasmid pRS426 (small letters indicated in Table S2). Both fragments 5- and 3-UTR were PCR-amplified from genomic DNA using as templates the A4 strain for *A. nidulans* cassettes. The *pyrG* used in the *A. nidulans* cassettes for generating the strain $\Delta casB$ was used as marker for auxotrophy and was amplified from

pCDA21 plasmid (Chaverocche *et al.*, 2000), and *pyroA* for Δ casA was amplified from *A. fumigatus* Af293 (for details see Fig. S4). For the construction of *alcA::pkcA*, 1000 bp of the *pkcA* encoding region was cloned downstream to the *alcA* promoter into the pMCB17apx vector. This construction was further transformed in *A. nidulans* to replace the endogenous *pkcA* promoter yielding the strain *alcA::pkcA*.

Deletion cassettes generation was achieved by transforming each fragment for each construction along with the plasmid pRS426 BamHI/EcoRI cut in the in *S. cerevisiae* strain SC94721 by the lithium acetate method (Schiestl and Gietz, 1989). The DNA of the yeast transformants was extracted by the method described by Goldman *et al.* (2003), dialysed and transformed by electroporation in *Escherichia coli* strain DH10B to rescue the pRS426 plasmid harbouring the cassettes. The cassettes were PCR-amplified from these plasmids and used for transformation of *A. nidulans* according to the procedure of Osmani *et al.* (1987). Transformants were scored for their ability to grow on minimal medium. PCR or Southern blot analyses were used throughout of the manuscript to demonstrate that the transformation cassettes had integrated homologously at the targeted *A. nidulans* loci (Fig. S4).

RNA isolation, real-time PCR reactions and UPR PCR assay

For total RNA isolation, the germings were disrupted by grinding in liquid nitrogen with pestle and mortar and total RNA was extracted with Trizol reagent (Invitrogen, USA). Ten micrograms of RNA from each treatment was then fractionated in 2.2 M formaldehyde, 1.2% w/v agarose gel, stained with ethidium bromide, and then visualized with UV light. The presence of intact 25S and 18S ribosomal RNA bands was used as a criterion to assess the integrity of the RNA. RNase-free DNase I treatment for the real-time PCR experiments was carried out as previously described (Semighini *et al.*, 2002).

All the PCR reactions were performed using an ABI 7500 Fast Real-Time PCR System (Applied Biosystems, USA) and Taq-Man Universal PCR Master Mix kit (Applied Biosystems, USA). The reactions and calculations were performed according to Semighini *et al.* (2002). The primers and Lux fluorescent probes (Invitrogen, USA) used in this work are described in Table S2.

The *hacA* mRNA splicing was assayed using a modification of a reported protocol (Bicknell *et al.*, 2007). Overnight cultures were grown and treated with FOH 0, 10, 50 or 100 μ M for 2 h or DTT 1 mM for 60 min. The cells were frozen and harvested, and total RNA was isolated using Trizol. Twenty micrograms of total RNA was treated with DNase, purified using a RNA easy kit (Qiagen), and cDNA was generated using the SuperScript III First Strand Synthesis system (Invitrogen) and oligo(dT) primers according to the manufacturer's protocol. The *hacA* uninduced (*hac^u* – 100 bp long) and *hacA* induced (*hacⁱ* – 80 bp long) cDNAs were amplified by PCR using the primers *hacAF* and *hacAR* (described in Table S2), with the following PCR conditions: 94°C, 1 min; 25 cycles of 94°C, 30 s; 60°C, 30 s; 72°C, 30 s; 72°C, 10 min. Amplicons were analysed by electrophoresis on 5% polyacrilamid plus 0.5% agarose gels. The cDNA loading was normalized by

tubC (encoding β -tubulin) PCR amplification under the same conditions by using primers *tubC* 918F and *tubC* 1075R (Table S2). The images generated were subjected to densitometric analysis of pixel intensity using the ImageJ software (<http://rsbweb.nih.gov/ij/index.html>).

PKC assay and Western blotting analysis

Protein assays were performed by initially growing conidia from the wild-type and mutant strains in a reciprocal shaker at 37°C for 16 h in liquid MM 2% glycerol medium supplemented with threonine 100 mM. Each sample was harvested by filtration through a Whatman filter number 1, washed thoroughly with sterile water, quickly frozen in liquid nitrogen, and disrupted by grinding. Total protein was extracted at 4°C with extraction buffer [25 mM Tris-HCl (pH 7.4), 0.5 mM EDTA, 0.5 mM EGTA, 0.05% Triton X-100, 10 mM β -mercaptoethanol, 1 μ g ml⁻¹ leupeptin, 1 μ g ml⁻¹ aprotinin, 0.5 mM of PMSF]. The lysate was centrifuged for 30 min at 4°C, 14 000 *g* in a microcentrifuge. The supernatant was chromatographed on a 1 ml DEAE-cellulose anion-exchange chromatography column that has been pre-equilibrated in extraction buffer. The column was washed with 5 ml of extraction buffer, and the PKC-containing fraction was eluted using 1 ml of extraction buffer containing 200 mM NaCl. Protein concentration was determined using a modified Bradford assay (Bio-Rad, Hercules, CA, USA). The enzymatic assay was performed according to SignaTECT PKC Assay System protocol (Promega).

Conidia (5×10^7 ml⁻¹) were grown in appropriated liquid medium using a reciprocal shaker at 37°C for 16 h, treated with FOH 0, 10, 50 or 100 μ M for 2 h, or DTT 1 mM for 0, 15, 30, or 60 min, or Congo red 1 mM for 0, 20 or 60 min, harvested, frozen and disrupted by grinding. Protein extracts were obtained using HB buffer [25 mM Tris-HCl (pH 7.4), 0.5 mM EDTA, 0.5 mM EGTA, 0.05% Triton X-100, 10 mM β -mercaptoethanol and mini-complete protease inhibitors (Roche Diagnostics, Indianapolis, IN, USA)] at 4°C, and centrifuged for 30 min at 4°C, 14 000 *g* in a microcentrifuge. Extracts were normalized by protein concentration (Bradford assay; Bio-Rad), fractionated by SDS-PAGE electrophoresis (12% gel), and transferred to nitrocellulose (Hybond C⁺) membranes.

The MpkA phosphorylation was assessed by using anti-phosphop44/42 MAPK (Erk1/2) (Thr202/Tyr204) antibodies (1:1500 dilution, Cell Signalling Technology, USA). Primary antibodies were detected by using a rabbit peroxidase (HRP)-conjugated second antibody (Kirkegaard and Perry laboratories), 1:5000 diluted in TBS-T for 1 h at room temperature shaking, and ECL reagents (GE) for chemiluminescent detection. Protein concentrations were normalized by using Mouse monoclonal (TU-06) antibody to β -tubulin (Abcam), diluted 1:5000 em TBS-T plus skimmed milk powder 5%, incubated with agitation for 16 h at 4°C. β -tubulin was detected by using a mouse peroxidase (HRP)-conjugated second antibody (Pierce), 1:3000 diluted in TBS-T, incubated with agitation for 1 h at room temperature, and Super signal West Pico chemiluminescent substrate (Pierce) was used for chemiluminescent detection.

Acknowledgements

This research was supported by the Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP), Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Brazil, John Simon Guggenheim Memorial Foundation, USA. We also thank Drs Jesus Aguirre, K. Abe, and T. Mizuno for supplying us with *A. nidulans* mutant strains, Dr Robert Cramer Jr. for critical reading of the manuscript, and the four anonymous reviewers for their suggestions.

References

- Asano, Y., Hagiwara, D., Yamashino, T., and Mizuno, T. (2007) Characterization of the bZip-type transcription factor NapA with reference to oxidative stress response in *Aspergillus nidulans*. *Biosci Biotechnol Biochem* **71**: 1800–1803.
- Adany, I., Yazlovitskaya, E.M., Haug, J.S., Voziyan, P.A., and Melnykovich, G. (1994) Differences in sensitivity to farnesol toxicity between neoplastically and non-neoplastically-derived cells in culture. *Cancer Lett* **79**: 175–179.
- Back, S.H., Schroder, M., Lee, K., Zhang, K., and Kaufman, R.J. (2005) ER stress signaling by regulated splicing: IRE1/HAC1/XBP1. *Methods* **35**: 395–416.
- Bahn, Y.S. (2008) Master and commander in fungal pathogens: the two-component system and the HOG signaling pathway. *Eukaryot Cell* **7**: 2017–2036.
- Balázs, A., Pócsi, I., Hamari, Z., Leiter, E., Emri, T., Miskei, M., et al. (2010) AtfA bZIP-type transcription factor regulates oxidative and osmotic stress responses in *Aspergillus nidulans*. *Mol Genet Genomics* **283**: 289–303.
- Bhabhra, R., Miley, M.D., Mylonakis, E., Boettner, D., Fortwendel, J., Panepinto, J.C., et al. (2004) Disruption of the *Aspergillus fumigatus* gene encoding nucleolar protein CgrA impairs thermotolerant growth and reduces virulence. *Infect Immun* **72**: 4731–4740.
- Bicknell, A.A., Babour, A., Federovitch, C.M., and Niwa, M. (2007) A novel function in cytokinesis reveals a house-keeping function for the unfolded protein response. *J Cell Biol* **177**: 1017–1027.
- Bicknell, A.A., Tourtellotte, J., and Niwa, M. (2010) Late phase of the endoplasmic reticulum stress response pathway is regulated by Hog1 MAP kinase. *J Biol Chem* **285**: 17545–17555.
- Binder, U., Oberparleiter, C., Meyer, V., and Marx, F. (2009) The antifungal protein PAF interferes with PKC/MPK and cAMP/PKA signalling of *Aspergillus nidulans*. *Mol Microbiol* **75**: 294–307.
- Bonilla, M., and Cunningham, K.W. (2003) Mitogen-activated protein kinase stimulation of Ca(2+) signaling is required for survival of endoplasmic reticulum stress in yeast. *Mol Biol Cell* **14**: 4296–4305.
- Burke, Y.D., Stark, M.J., Roach, S.L., Sen, S.E., and Crowell, P.L. (1997) Inhibition of pancreatic cancer growth by the dietary isoprenoids farnesol and geraniol. *Lipids* **32**: 151–156.
- Chaverroche, M.K., Ghigo J.M., and d'Enfert, C. (2000) A rapid method for efficient gene replacement in the filamentous fungus *Aspergillus nidulans*. *Nucleic Acids Res* **28**: E97–E104.
- Chen, Y., Feldman, D.E., Deng, C., Brown, J.A., De Giacomo, A.F., Gaw, A.F., et al. (2005) Identification of mitogen-activated protein kinase signaling pathways that confer resistance to endoplasmic reticulum stress in *Saccharomyces cerevisiae*. *Mol Cancer Res* **3**: 669–677.
- Colot, H.V., Park, G., Turner, G.E., Ringelberg, C., Crew, C.M., Litvinkova, L., et al. (2006) A high-throughput gene knockout procedure for *Neurospora* reveals functions for multiple transcription factors. *Proc Natl Acad Sci USA* **103**: 10352–10357.
- Cox, J.S., and Walter, P. (1996) A novel mechanism for regulating activity of a transcription factor that controls the unfolded protein response. *Cell* **87**: 391–404.
- Davenport, K.R., Sohaskey, M., Kamada, Y., Levin, D.E., and Gustin, M.C. (1995) A second osmosensing signal transduction pathway in yeast. Hypotonic shock activates the PKC1 protein kinase-regulated cell integrity pathway. *J Biol Chem* **270**: 30157–30161.
- Degterev, A., Boyce, M., and Yuan, J. (2003) A decade of caspases. *Oncogene* **22**: 8543–8567.
- Dichtl, K., Ebel, F., Dirr, F., Routier, F.H., Heesemann, J., and Wagener, J. (2010) Farnesol misplaces tip-localized Rho proteins and inhibits cell wall integrity signaling in *Aspergillus fumigatus*. *Mol Microbiol* **76**: 1191–1204.
- Dinamarco, T.M., Figueiredo, B.C.P., Savoldi, M., Malavazi, I., Soriani, F.M., Uyemura, S.A., et al. (2010) The roles played by *Aspergillus nidulans* apoptosis-inducing factor (Aif)-like mitochondrial oxidoreductase (AifA) and NADH-ubiquinone oxidoreductases (NdeA-B and NdiA) in farnesol resistance. *Fungal Genet Biol* (in press).
- Donepudi, M., and Grutter, M.G. (2002) Structure and zymogen activation of caspases. *Biophys Chem* **101–102**: 145–153.
- Fairn, G.D., MacDonald, K., and McMaster, C.R. (2007) A chemogenomic screen in *Saccharomyces cerevisiae* uncovers a primary role for the mitochondria in farnesol toxicity and its regulation by the Pkc1 pathway. *J Biol Chem* **282**: 4868–4874.
- Flippin, M., Kocalkowska, J., and Felenbok, B. (2002) Characteristics of physiological inducers of the ethanol utilization (alc) pathway in *Aspergillus nidulans*. *Biochem J* **15**: 25–31.
- Fujioka, T., Mizutani, O., Furukawa, K., Sato, N., Yoshimi, A., Yamagata, Y., et al. (2007) MpkA-dependent and -independent cell wall integrity signaling in *Aspergillus nidulans*. *Eukaryot Cell* **6**: 1497–1510.
- Furukawa, K., Hoshi, Y., Maeda, T., Nakajima, T., and Abe, K. (2005) *Aspergillus nidulans* HOG pathway is activated only by two-component signalling pathway in response to osmotic stress. *Mol Microbiol* **56**: 1246–1261.
- Gerik, K.J., Bhimreddy, S.R., Ryerse, J.S., Specht, C.A., and Lodge, J.K. (2008) PKC1 is essential for protection against both oxidative and nitrosative stresses, cell integrity, and normal manifestation of virulence factors in the pathogenic fungus *Cryptococcus neoformans*. *Eukaryot Cell* **7**: 1685–1698.
- Goldman, G.H., dos Reis Marques, E., Duarte Ribeiro, D.C., de Souza Bernardes, L.A., Quiapin, A.C., Vitorelli, P.M., et al. (2003) Expressed sequence tag analysis of the human pathogen *Paracoccidioides brasiliensis* yeast phase: identification of putative homologues of *Candida*

- albicans* virulence and pathogenicity genes. *Eukaryot Cell* **2**: 34–48.
- Guillemette, T., van Peij, N.N., Goosen, T., Lanthaler, K., Robson, G.D., Hondel, C.A., *et al.* (2007) Genomic analysis of the secretion stress response in the enzyme-producing cell factory *Aspergillus niger*. *BMC Genomics* **8**: 158.
- Hagiwara, D., Asano, Y., Marui, J., Furukawa, K., Kanamaru, K., Kato, M., *et al.* (2007) The SskA and SrrA response regulators are implicated in oxidative stress responses of hyphae and asexual spores in the phosphorelay signaling network of *Aspergillus nidulans*. *Biosci Biotechnol Biochem* **71**: 1003–1014.
- Hagiwara, D., Asano, Y., Yamashino, T., and Mizuno, T. (2008) Characterization of bZip-type transcription factor AtfA with reference to stress responses of conidia of *Aspergillus nidulans*. *Biosci Biotechnol Biochem* **72**: 2756–2760.
- Han, K.H., and Prade, R.A. (2002) Osmotic stress-coupled maintenance of polar growth in *Aspergillus nidulans*. *Mol Microbiol* **43**: 1065–1078.
- He, L., Mo, H., Hadisusilo, S., Oureshi, A.A., and Elson, C.E. (1997) Isoprenoids suppress the growth of murine B16 melanomas in vitro and in vivo. *J Nutr* **127**: 668–674.
- Helenius, A., and Aebi, M. (2004) Roles of N-linked glycans in the endoplasmic reticulum. *Annu Rev Biochem* **73**: 1019–1049.
- Herrmann, M., Spröte, P., and Brakhage, A.A. (2006) Protein kinase C (PkcA) of *Aspergillus nidulans* is involved in penicillin production. *Appl Environ Microbiol* **72**: 2957–2970.
- Hornby, J.M., Jensen, E.C., Lisec, A.D., Tasto, J.J., Jahnke, B., Shoemaker, R., *et al.* (2001) Quorum sensing in the dimorphic fungus *Candida albicans* is mediated by farnesol. *Appl Environ Microbiol* **67**: 2982–2992.
- Hudes, G.R., Szarka, C.E., Adams, A., Ranganathan, S., McCauley, R.A., Weiner, L.M., *et al.* (2000) Phase I pharmacokinetic trial of perillyl alcohol (NSC 641066) in patients with refractory solid malignancies. *Clin Cancer Res* **6**: 3071–3080.
- Ichinomiya, M., Uchida, H., Koshi, Y., Ohta, A., and Horiuchi, H. (2007) A protein kinase C-encoding gene, *pkcA*, is essential to the viability of the filamentous fungus *Aspergillus nidulans*. *Biosci Biotechnol Biochem* **71**: 2787–2799.
- Ikner, A., and Shiozaki, K. (2005) Yeast signaling pathways in the oxidative stress response. *Mutat Res* **569**: 13–27.
- Joo, J.H., Liao, G., Collins, J.B., Grissom, S.F., and Jetten, A.M. (2007) Farnesol-induced apoptosis in human lung carcinoma cells is coupled to the endoplasmic reticulum stress response. *Cancer Res* **67**: 7929–7936.
- Joo, J.H., and Jetten, A.M. (2010) Molecular mechanisms involved in farnesol-induced apoptosis. *Cancer Lett* **287**: 123–135.
- Jung, U.S., and Levin, D.E. (1999) Genome-wide analysis of gene expression regulated by the yeast cell wall integrity signalling pathway. *Mol Microbiol* **34**: 1049–1057.
- Kamada, Y., Jung, U.S., Piotrowski, J., and Levin, D.E. (1995) The protein kinase C-activated MAP kinase pathway of *Saccharomyces cerevisiae* mediates a novel aspect of the heat shock response. *Genes Dev* **9**: 1559–1571.
- Kafer, E. (1977) Meiotic and mitotic recombination in *Aspergillus* and its chromosomal aberrations. *Adv Genet* **19**: 33–131.
- Kawahara, T., Yanagi, H., Yura, T., and Mori, K. (1997) Endoplasmic reticulum stress-induced mRNA splicing permits synthesis of transcription factor Hac1p/Ern4p that activates the unfolded protein response. *Mol Biol Cell* **8**: 1845–1862.
- Kawasaki, L., Sánchez, O., Shiozaki, K., and Aguirre, J. (2002) SakA MAP kinase is involved in stress signal transduction, sexual development and spore viability in *Aspergillus nidulans*. *Mol Microbiol* **45**: 1153–1163.
- Kimata, Y., Ishiwata-Kimata, Y., Yamada, S., and Kohno, K. (2006) Yeast unfolded protein response pathway regulates expression of genes for anti-oxidative stress and for cell surface proteins. *Genes Cells* **11**: 59–69.
- Korennykh, A.V., Egea, P.F., Korostelev, A.A., Finer-Moore, J., Zhang, C., Shokat, K.M., *et al.* (2009) The unfolded protein response signals through high-order assembly of Ire1. *Nature* **457**: 687–693.
- Krysan, D.J. (2009) The cell wall and endoplasmic reticulum stress responses are coordinately regulated in *Saccharomyces cerevisiae*. *Commun Integr Biol* **2**: 233–235.
- Langford, M.L., Atkin, A.L., and Nickerson, K.W. (2009) Cellular interactions of farnesol, a quorum-sensing molecule produced by *Candida albicans*. *Future Microbiol* **4**: 1353–1362.
- Lavrik, I.N., Golks, A., and Krammer, P.H. (2005) Caspases: pharmacological manipulation of cell death. *J Clin Invest* **115**: 2665–2672.
- Lee, K., Tirasophon, W., Shen, X., Michalak, M., Prywes, R., Okada, T., *et al.* (2002) IRE1-mediated unconventional mRNA splicing and S2P-mediated ATF6 cleavage merge to regulate XBP1 in signaling the unfolded protein response. *Genes Dev* **16**: 452–466.
- Levin, D.E. (2005) Cell wall integrity signaling in *Saccharomyces cerevisiae*. *Microbiol Mol Biol Rev* **69**: 262–291.
- Li, J., and Yuan, J. (2008) Caspases in apoptosis and beyond. *Oncogene* **27**: 6194–6206.
- Machida, K., Tanaka, T., Fujita, K., and Taniguchi, M. (1998) Farnesol-induced generation of reactive oxygen species via indirect inhibition of the mitochondrial electron transport chain in the yeast *Saccharomyces cerevisiae*. *J Bacteriol* **180**: 4460–4465.
- Machida, K., Tanaka, T., Yano, Y., Otani, S., and Taniguchi, M. (1999) Farnesol-induced growth inhibition in *Saccharomyces cerevisiae* by a cell cycle mechanism. *Microbiology* **145**: 293–299.
- Madeo, F., Herker, E., Maldener, C., Wissing, S., Lachelt, S., Herlan, M., *et al.* (2002) A caspase-related protease regulates apoptosis in yeast. *Mol Cell* **9**: 911–917.
- May, G. (2008) Mitogen-activated protein kinase pathways in *Aspergilli*. In *The Aspergilli. Genomics, Medical Aspects, Biotechnology and Research Methods*. Mycology 26. Goldman, G.H., and Osmani, S.A. (eds). Boca Raton, FL: CRC Press, pp. 121–127.
- Mazzoni, C., and Falcone, C. (2008) Caspase-dependent apoptosis in yeast. *Biochim Biophys Acta* **1783**: 1320–1327.
- Miquel, K., Pradines, A., and Favre, G. (1996) Farnesol and geranylgeraniol induce actin cytoskeleton disorganization and apoptosis in A549 lung adenocarcinoma cells. *Biochem Biophys Res Commun* **225**: 869–876.

- Momoi, T. (2004) Caspases involved in ER stress-mediated cell death. *J Chem Neuroanat* **28**: 101–105.
- Moye-Rowley, W.S. (2003) Regulation of the transcriptional response to oxidative stress in fungi: similarities and differences. *Eukaryot Cell* **2**: 381–389.
- Mulder, H.J., Saloheimo, M., Penttilä, M., and Madrid, S.M. (2004) The transcription factor HACA mediates the unfolded protein response in *Aspergillus niger*, and up-regulates its own transcription. *Mol Genet Genomics* **271**: 130–140.
- Nickerson, K.W., Atkin, A.L., and Hornby, J.M. (2006) Quorum sensing in dimorphic fungi: farnesol and beyond. *Appl Environ Microbiol* **72**: 3805–3813.
- Nita-Lazar, M., and Lennarz, W.J. (2005) Pkc1p modifies CPY degradation in the ERAD pathway. *Biochem Biophys Res Commun* **332**: 357–361.
- Nonaka, H., Tanaka, K., Hirano, H., Fujiwara, T., Kohno, H., Umikawa, M., et al. (1995) A downstream target of RHO1 small GTP-binding protein is PKC1, a homolog of protein kinase C, which leads to activation of the MAP kinase cascade in *Saccharomyces cerevisiae*. *EMBO J* **14**: 5931–5938.
- Noventa-Jordão, M.A., Nascimento, A.M., Goldman, M.H., Terenzi, H.F., and Goldman, G.H. (1999) Molecular characterization of ubiquitin genes from *Aspergillus nidulans*: mRNA expression on different stress and growth conditions. *Biochim Biophys Acta* **1490**: 237–244.
- Ohizumi, H., Masuda, Y., Nakajo, S., Sakai, I., Ohsawa, S., and Nakaya, K. (1995) Geranylgeraniol is a potent inducer of apoptosis in tumor cells. *J Biochem* **117**: 11–13.
- Osmani, S.A., May, G.S., and Morris, N.R. (1987) Regulation of the mRNA levels of *nimA*, a gene required for the G2-M transition in *Aspergillus nidulans*. *J Cell Biol* **104**: 1495–1504.
- Pal, B., Chan, N.C., Helfenbaum, L., Tan, K., Tansey, W.P., and Gething, M.J. (2007) SCFCdc4-mediated degradation of the Hac1p transcription factor regulates the unfolded protein response in *Saccharomyces cerevisiae*. *Mol Biol Cell* **18**: 426–440.
- Qi, X., and Mochly-Rosen, D. (2008) The PKC δ -Abl complex communicates ER stress to the mitochondria – an essential step in subsequent apoptosis. *J Cell Sci* **121**: 804–813.
- Rasheva, V.I., and Domingos, P.M. (2009) Cellular responses to endoplasmic reticulum stress and apoptosis. *Apoptosis* **14**: 996–1007.
- Reyes, G., Romans, A., Nguyen, C.K., and May, G.S. (2006) Novel mitogen-activated protein kinase MpkC of *Aspergillus fumigatus* is required for utilization of polyalcohol sugars. *Eukaryot Cell* **5**: 1934–1940.
- Reyland, M.E. (2009) Protein kinase C isoforms: multi-functional regulators of cell life and death. *Front Biosci* **14**: 2386–2399.
- Rhome, R., and Del Poeta, M. (2009) Lipid signaling in pathogenic fungi. *Annu Rev Microbiol* **63**: 119–131.
- Richie, D.L., Miley, M.D., Bhabhra, R., Robson, G.D., Rhodes, J.C., and Askew, D.S. (2007) The *Aspergillus fumigatus* metacaspases CasA and CasB facilitate growth under conditions of endoplasmic reticulum stress. *Mol Microbiol* **63**: 591–604.
- Richie, D.L., Hartl, L., Amanianda, V., Winters, M.S., Fuller, K.K., Miley, M.D., et al. (2009) A role for the unfolded protein response (UPR) in virulence and antifungal susceptibility in *Aspergillus fumigatus*. *PLoS Pathog* **5**: e1000258.
- Román, E., Alonso-Monge, R., Gong, Q., Li, D., Calderone, R., and Pla, J. (2009) The Cek1 MAPK is a short-lived protein regulated by quorum sensing in the fungal pathogen *Candida albicans*. *FEMS Yeast Res* **9**: 942–955.
- Ronen, R., Sharon, H., Levdansky, E., Romano, J., Shadkhan, Y., and Osherov, N. (2007) The *Aspergillus nidulans* *pkcA* gene is involved in polarized growth, morphogenesis and maintenance of cell wall integrity. *Curr Genet* **51**: 321–329.
- Rueggsegger, U., Leber, J.H., and Walter, P. (2001) Block of HAC1 mRNA translation by long-range base pairing is released by cytoplasmic splicing upon induction of the unfolded protein response. *Cell* **107**: 103–114.
- Sakaki, K., and Kaufman, R.J. (2008) Regulation of ER stress-induced macroautophagy by protein kinase C. *Autophagy* **4**: 841–843.
- Sakaki, K., Wu, J., and Kaufman, R.J. (2008) Protein kinase C θ is required for autophagy in response to stress in the endoplasmic reticulum. *J Biol Chem* **283**: 15370–15380.
- Saloheimo, M., Valkonen, M., and Penttilä, M. (2003) Activation mechanisms of the HAC1-mediated unfolded protein response in filamentous fungi. *Mol Microbiol* **47**: 1149–1161.
- Sambrook, J., and Russell, D.W. (2001) *Molecular Cloning: A Laboratory Manual*, 3rd edn. New York: Cold Spring Harbor Laboratory Press.
- Savoldi, M., Malavazi, I., Soriani, F.M., Capellaro, J.L., Goldman, M.H.S., Ferreira, M.E., and Goldman, G.H. (2008) Farnesol induces the transcriptional accumulation of the *Aspergillus nidulans* apoptosis inducing factor (AIF)-like mitochondrial oxidoreductase. *Mol Microbiol* **70**: 44–59.
- Schiestl, R.H., and Gietz, R.D. (1989) High efficiency transformation of intact yeast cells using single stranded nucleic acids as a carrier. *Curr Genet* **16**: 339–346.
- Schmitz, H.P., Lorberg, A., and Heinisch, J.J. (2002) Regulation of yeast protein kinase C activity by interaction with the small GTPase Rho1p through its amino-terminal HR1 domain. *Mol Microbiol* **44**: 829–840.
- Scrimale, T., Didone, L., Bentley, K.L., and Krysan, D.J. (2009) The unfolded protein response is induced by the cell wall integrity mitogen-activated protein kinase signaling cascade and is required for cell wall integrity in *Saccharomyces cerevisiae*. *Mol Biol Cell* **20**: 164–175.
- Semighini, C.P., Marins, M., Goldman, M.H.S., and Goldman, G.H. (2002) Quantitative analysis of the relative transcript levels of ABC transporter *Atr* genes in *Aspergillus nidulans* by Real-Time Reverse Transcription-PCR assay. *Appl Environ Microbiol* **68**: 1351–1357.
- Semighini, C.P., Hornby, J.M., Dumitru, R., Nickerson, K.W., and Harris, S.D. (2006) Farnesol-induced apoptosis in *Aspergillus nidulans* reveals a possible mechanism for antagonistic interactions between fungi. *Mol Microbiol* **59**: 753–764.
- Shaner, L., Gibney, P.A., and Morano, K.A. (2008) The Hsp110 protein chaperone Sse1 is required for yeast cell wall integrity and morphogenesis. *Curr Genet* **54**: 1–11.
- Shea, J.M., and Del Poeta, M. (2006) Lipid signaling in pathogenic fungi. *Curr Opin Microbiol* **9**: 352–358.

- Shimizu, Y., and Hendershot, L.M. (2009) Oxidative folding: cellular strategies for dealing with the resultant equimolar production of reactive oxygen species. *Antioxid Redox Signal* **11**: 2317–2331.
- Shirliff, M.E., Krom, B.P., Meijering, R.A.M., Peters, B.M., Zhu, J., Scheper, M.A., *et al.* (2009) Farnesol-induced apoptosis in *Candida albicans*. *Antimicrob Agents Chemother* **53**: 2392–2401.
- Taylor, M.M., MacDonald, K., Morris, A.J., and McMaster, C.R. (2005) Enhanced apoptosis through farnesol inhibition of phospholipase D signal transduction. *FEBS J* **272**: 5056–5063.
- Teepe, A.G., Loprete, D.M., He, Z., Hoggard, T.A., and Hill, T.W. (2007) The protein kinase C orthologue PkcA plays a role in cell wall integrity and polarized growth in *Aspergillus nidulans*. *Fungal Genet Biol* **44**: 554–562.
- Travers, K.J., Patil, C.K., Wodicka, L., Lockhart, D.J., Weissman, J.S., and Walter, P. (2000) Functional and genomic analyses reveal an essential coordination between the unfolded protein response and ER-associated degradation. *Cell* **101**: 249–258.
- Uppuluri, P., Mekala, S., and Chaffin, W.L. (2007) Farnesol-mediated inhibition of *Candida albicans* yeast growth and rescue by a diacylglycerol analogue. *Yeast* **24**: 681–693.
- Uren, A.G., O'Rourke, K., Aravind, L.A., Pisabarro, M.T., Seshagiri, S., Koonin, E.V., *et al.* (2000) Identification of paracaspases and metacaspases: two ancient families of caspase-like proteins, one of which plays a key role in MALT lymphoma. *Mol Cell* **6**: 961–967.
- Vargas-Pérez, I., Sánchez, O., Kawasaki, L., Georgellis, D., and Aguirre, J. (2007) Response regulators SrrA and SskA are central components of a phosphorelay system involved in stress signal transduction and asexual sporulation in *Aspergillus nidulans*. *Eukaryot Cell* **6**: 1570–1583.
- Vilella, F., Herrero, E., Torres, J., and Torre-Ruiz, M.A. (2005) Pkc1 and the upstream elements of the cell integrity pathway in *Saccharomyces cerevisiae*, Rom2 and Mtl1, are required for cellular responses to oxidative stress. *J Biol Chem* **280**: 9149–9159.
- Vozian, P.A., Haug, J.S., and Melnykovich, G. (1995) Mechanism of farnesol cytotoxicity: further evidence for the role of PKC-dependent signal transduction in farnesol-induced apoptotic cell death. *Biochem Biophys Res Commun* **212**: 479–486.
- Wimalasena, T.T., Enjalbert, B., Guillemette, T., Plumridge, A., Budge, S., Yin, Z., *et al.* (2008) Impact of the unfolded protein response upon genome-wide expression patterns, and the role of Hac1 in the polarized growth, of *Candida albicans*. *Fungal Genet Biol* **45**: 1235–1247.
- Wiseman, D.A., Werner, S.R., and Crowell, P.L. (2007) Cell cycle arrest by the isoprenoids perillyl alcohol, geraniol, and farnesol is mediated by p21 (Cip1) and p27 (Kip1) adenocarcinoma cells. *J Pharmacol Exp Ther* **320**: 1163–1170.
- Wright, C.M., Fewell, S.W., Sullivan, M.L., Pipas, J.M., Watkins, S.C., and Brodsky, J.L. (2007) The Hsp40 molecular chaperone Ydj1p, along with the protein kinase C pathway, affects cell-wall integrity in the yeast *Saccharomyces cerevisiae*. *Genetics* **175**: 1649–1664.
- Wuichet, K., Cantwell, B.J., and Zhulin, I.B. (2010) Evolution and phyletic distribution of two-component signal transduction systems. *Curr Opin Microbiol* **13**: 219–225.
- Yang, C., and Kazanietz, M.G. (2003) Divergence and complexities in DAG signaling: looking beyond PKC. *Trends Pharmacol Sci* **24**: 602–608.

Supporting information

Additional supporting information may be found in the online version of this article.

Please note: Wiley-Blackwell are not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.

7.4. Anexo M. Artigo publicado: The *Aspergillus nidulans* *nucA*^{ENDO} homologue is not involved in cell death. Pimentel Figueiredo, B. C.; de Castro, P. A.; Dinamarco, T. M.; Goldman, M. H. S.; Goldman, G. H.. *Eukaryotic Cell*, Fev. 2011, 10(2):276-83.

The *Aspergillus nidulans* *nucA*^{EndoG} Homologue Is Not Involved in Cell Death[∇]

Bárbara de Castro Pimentel Figueiredo,¹ Patrícia Alves de Castro,¹ Taísa Magnani Dinamarco,¹ Maria Helena S. Goldman,² and Gustavo Henrique Goldman^{1,3*}

Faculdade de Ciências Farmacêuticas de Ribeirão Preto¹ and Faculdade de Filosofia, Ciências e Letras de Ribeirão Preto,² Universidade de São Paulo, São Paulo, Brazil, and Laboratório Nacional de Ciência e Tecnologia do Bioetanol, CTBE, Caixa Postal 6170, 13083-970 Campinas, São Paulo, Brazil³

Received 14 September 2010/Accepted 8 November 2010

Upon apoptosis induction, translocation of mammalian mitochondrial endonuclease G (EndoG) to the nucleus coincides with large-scale DNA fragmentation. Here, we describe for the first time a homologue of EndoG in filamentous fungi by investigating if the *Aspergillus nidulans* homologue of the EndoG gene, named *nucA*^{EndoG}, is being activated during farnesol-induced cell death. Our results suggest that NucA is not involved in cell death, but it plays a role in the DNA-damaging response in *A. nidulans*.

The mechanisms involved in cell death in filamentous fungi are not well known (for reviews, see references 9, 17, and 24). We have been using *Aspergillus nidulans* and the isoprenoid farnesol (FOH), which inhibits proliferation and induces apoptosis, as a model system and cell death stimulus, respectively, aiming to understand by which means filamentous fungi are driven toward cell death (3, 5, 20, 23). In mammalian cells, at least two major apoptotic pathways have been depicted: (i) the intrinsic pathway, which needs the involvement of the mitochondria, and (ii) the extrinsic pathway, where mitochondria are bypassed and caspases are activated (for a review, see reference 6). One of the hallmarks of the intrinsic pathway is the release of apoptogenic factors, such as cytochrome *c*, to the cytosol and the consequent assembly, organized by this protein, of the high-molecular-weight complex, the mitochondrial apoptosome, which activates caspases (for a review, see reference 6). In *Saccharomyces cerevisiae* cells undergoing an apoptotic process induced by acetic acid, translocation of Cyc1p to the cytosol was observed (13). Another factor of the intrinsic pathway is endonuclease G (EndoG), which has been described as a mitochondrial endonuclease that digests both DNA and RNA (1, 2). Upon apoptosis induction, translocation of mammalian EndoG and its *S. cerevisiae* homologue *NUC1* to the nucleus coincides with large-scale DNA fragmentation (2, 12, 16). Here, we describe for the first time a homologue of EndoG in filamentous fungi by investigating if the *A. nidulans* EndoG homologue, named *nucA*^{EndoG}, is being activated during FOH-induced cell death.

We identified AN2185.2 as the putative homologue of *S. cerevisiae* *NUC1* (53% identity, 68% similarity, E value of 2e−97). *In silico* analysis of NucA (Fig. 1A) shows a putative mitochondrial target sequence (MTS) located at amino acids 1 to 28 (as predicted by MITOPROT [http://ihg.gsf.de/ihg

/mitoprot.html], PrediSi [http://www.predisi.de/home.html], and Predotar [http://urgi.versailles.inra.fr/predotar/predotar.html]) and a putative nuclear localization signal located at amino acids 308 to 314 (as predicted by NucPred [http://www.sbc.su.se/~maccallr/nucpred/] but not confirmed by NetNES 1.1 [http://www.cbs.dtu.dk/services/NetNES-1.1/index.php] or by PredictNLS Server [http://cubic.bioc.columbia.edu/services/predictNLS/]). A DNA/RNA nonspecific endonuclease active site (PS01070; http://www.expasy.org/prosite/), typical of a family of bacterial and eukaryotic endonucleases that cleave double-stranded and single-stranded nucleic acids and require a divalent ion such as magnesium for their activity, is located at amino acids 156 to 164 (Fig. 1A).

For total RNA isolation, RNase-free DNase I treatment for the real-time PCR experiments and the reactions and calculations were performed according to the method described in reference 22. Primers and Lux probes for the normalizer β -tubulin (*tubC*) are 5′-GCAGAATGTCTCGTCCGAATG-3′ and 5′-CACTTTATGCCGTCGCCGAAAG[FAM]G-3′ (where FAM is 6-carboxyfluorescein), while for *nucA*^{EndoG} the primers and Lux probes are 5′-cggatTGGTGGCCTCTGTTCATATCc[FAM]G-3′ and 5′-GCCTCTCTTCCGTGCCAAACT-3′. When the wild-type strain is exposed to FOH, there is an increased accumulation of *nucA*^{EndoG} mRNA of 3.1-, 40.0-, and 10.8-fold at 10, 50, and 100 μ M FOH, respectively (Fig. 1B). There is a low *nucA*^{EndoG} mRNA accumulation when *A. nidulans* germlings were exposed to oxidative stressing agents, such as hydrogen peroxide and paraquat (about 2.0- and 3.5-fold at 100 mM H₂O₂ and 5 mM paraquat, both for 60 min, respectively) (Fig. 1C), and DNA-damaging agents (about 2- to 3-fold for hydroxyurea [HU], methyl methanesulfonate [MMS], and bleomycin [BLEO]) (Fig. 1D).

Standard genetic techniques for *A. nidulans* were used for all strain constructions and transformations (11). DNA manipulations were as described in reference 19. All PCRs were performed using Platinum *Taq* DNA Polymerase High Fidelity (Invitrogen). For the DNA-mediated transformation, the deletion and monomeric red fluorescent protein (mRFP) fusion cassettes were constructed by *in vivo* recombination in *S. cerevisiae* as previously described in reference

* Corresponding author. Mailing address: Departamento de Ciências Farmacêuticas, Faculdade de Ciências Farmacêuticas de Ribeirão Preto, Universidade de São Paulo, Av. do Café S/N, CEP 14040-903, Ribeirão Preto, São Paulo, Brazil. Phone and fax: 55-16-36024280/81. E-mail: ggoldman@usp.br.

[∇] Published ahead of print on 3 December 2010.

FIG. 1. *In silico* analysis of NucA (A) and expression levels of the *nucA*^{EndoG} gene (B to D). The wild-type strain was grown for 16 h in YG at 37°C and transferred to fresh YG with FOH (0, 10, 50, or 100 μM FOH for 2 h) (B), oxidative stressing agents (C), or DNA-damaging agents (25 μM camptothecin [CPT], 0.003% methyl methanesulfonate [MMS], 0.6 μg/ml bleomycin [BLEO], or 0.25 μg/ml 4-nitroquinoline oxide [4-NQO] for 30, 60, 120, and 240 min) (D). The relative quantitation of *nucA*^{EndoG} and tubulin gene expression was determined by a standard curve (i.e., threshold cycle [*C_T*] values plotted against logarithm of the DNA copy number). The results are the means ± standard deviations from four sets of experiments. The values represent the fold change in gene expression compared to that of the wild-type control grown without any drug (represented absolutely as 1.00).

4. Briefly, for the gene deletion, about 1.5-kb regions on either side of the open reading frames (ORFs) were selected for primer design (5'-gtaacgccagggttttccagtcacgacgGTTGATG ATATGGAACCTCGTGCTTGTGCG-3', 5'-GACCCAACAA CCATGATACCATCGTTGCTGTGGATTGAAAAAATG GAGAA-3', 5'-CTGTCGATCATGTGGATGCTACGGCT ACTGTACATCTATATCATACCTATCTGT-3', and 5'-gctg ataacaatttcacacaggaacacgCTAGGCCAGTGTCCGAAC TA CAAACC-3' [lowercase letters indicate homology with the pRS426 vector region]). The NucA::mRFP cassette was constructed by using the primers 5'-gtaacgccagggttttccagtcacgacg GTCATCGACCCAGCATCCCC-3', 5'-GCTCCAGCGCCT GCACAGCTCCAGCCTTCTTCTTCGCATTGTTAAACT CC-3', 5'-GACCCAACAACCATGATACCATTGCTGTTG CCAGGTGAGG-3', and 5'-GGACTGGTGCAGGCGCTG GAGC-3'. The CycA::mRFP cassette was constructed by using the primers 5'-acgccagggttttccagtcacgacgTTGTTATTACGG GGAGGGCTACTCCGG-3', 5'-GCTCCAGCGCCTGCACC AGCTCCAGAGGCCCGGCGATCCC-3', 5'-CTGTCGATC ATGTGGATGCTGTGACAGGCGGGTACTGTAACATT ACACC-3', and 5'-gctgataacaatttcacacaggaacAGCAGCGG TGGCGATTACTTGTGTTTC-3'. Fragments of 5' and 3' untranslated regions of each gene were PCR amplified from genomic DNA using the A4 strain as the template for *A. nidulans* cassettes. The *pyroA* gene used in the *A. nidulans* cassette for generating the Δ *nucA*, *alcA::nucA*, NucA::mRFP, and CycA::mRFP strains was amplified from *A. fumigatus*

Af293. Cassette generation was achieved by transforming each fragment for each construction along with the plasmid pRS426 BamHI/EcoRI cut in the in *S. cerevisiae* strain SC9721 by the lithium acetate method (21). The DNA of the yeast transformants was extracted by the method described in reference 8, dialyzed, and transformed by electroporation in *Escherichia coli* strain DH10B to rescue the pRS426 plasmid harboring the cassettes. The cassettes were PCR amplified from these plasmids and used for transformation of *Aspergilli* according to the procedure described in reference 15. Transformants were scored for their ability to grow on minimal medium. PCR or Southern blot analyses were used throughout this study to demonstrate that the transformation cassettes had integrated homologously at the targeted *A. nidulans* loci.

We constructed a deletion mutant for *nucA*^{EndoG} (Fig. 2A). We investigated the effects of overexpressing *nucA*^{EndoG} in *A. nidulans* by constructing a conditional *alcA* mutant of the *nucA*^{EndoG} gene (Fig. 2B), using the primers 5' CCCGGTAC CGCTAATTAAC 3', 5' CTGTCGATCATGTGGATGCTA AAAGCTGATTGTGATAGTTC 3', 5' AGTTAATTAG CGGTACCGGGATGTCCAAGGCCACTGTTGCTG 3', and 5' GCGGATAACAATTTCACACAGGAAACAGCCTAGGC CAGTGTCCGAAC TACAAACC 3'. The *alcA* promoter is repressed by glucose, derepressed by glycerol, and induced to high levels by ethanol or L-threonine (7). The *nucA*^{EndoG} deletion strain has no apparent growth or conidiation defects and did not show increased susceptibility to FOH (Fig. 2C, left). Overexpres-

FIG. 2. Phenotypes of the $\Delta nucA$ deletion strain. (A) Schematic illustration of the $nucA^{EndoG}$ deletion strategy. Genomic DNA from both wild-type and $\Delta nucA$ strains was isolated and cleaved with the enzyme *EcoRV*; a 1.5-kb DNA fragment was used as a hybridization probe. This fragment recognizes a single DNA band (about 2.8 kb) in the wild-type strain and also a single DNA band (about 5.5 kb) in the $\Delta nucA$ mutant, as shown in the Southern blot analysis. (B) Expression of $nucA^{EndoG}$ in the $alcA::nucA$ strain. The $alcA::nucA$ strain was grown for 16 h at 37°C in 2% minimal medium (MM) plus glycerol. After this period, mycelia were transferred to MM plus different carbon sources and grown for another 6 h at 37°C. The relative $nucA^{EndoG}$ quantitation was performed as described for Fig. 1. (C) FOH susceptibility of $\Delta nucA$ and $alcA::nucA$ mutants. Aliquots (5 μ l) of 10-fold dilutions derived from a starting suspension of 1.0×10^8 conidia/ml of the corresponding wild-type, $\Delta nucA$, and $alcA::nucA$ strains were spotted on YG agar plates supplemented with 0, 0.25, 0.5, or 1.0 mM FOH. The plates were incubated at 37°C for 48 h. (D) Growth phenotypes of the wild-type and $nucA^{EndoG}$ deletion mutant strains grown either in complete medium (YG) supplemented with uridine and uracyl (YUU) or in YUU plus 0.35 μ g/ml of 4-NQO for 48 h at 37°C. (E) Quiescent and germinating conidiospores from different strains were exposed to UV light, and viability was scored after exposure to this DNA-damaging agent. Viability was determined as the percentage of colonies on treated plates compared to that on untreated controls. The results are the means $\pm$ standard deviations from four independent experiments. Statistical differences were determined by one-way analysis of variance (ANOVA) followed, when significant, by the Newman-Keuls multiple comparison test, using Sigma Stat statistical software (Jandel Scientific). *, $P < 0.01$.

FIG. 3. Genetic interactions with the *A. nidulans* $\Delta nucA$ mutant. (A) Growth phenotypes of the wild-type, $\Delta nucA$, $\Delta atrA$, and $\Delta nucA \Delta atrA$ strains grown in YUU, YUU plus MMS (0.0075%), or YUU plus 4-NQO (0.25 $\mu\text{g/ml}$). (B) Growth phenotypes of the wild-type, $\Delta nucA$, $\Delta atrA$, and $\Delta nucA \Delta atrA$ strains grown in YUU, YUU plus MMS (0.005%), YUU plus 4-NQO (0.25 $\mu\text{g/ml}$), or YUU plus menadione (0.1 mM).

sion of the *nucA*^{EndoG} gene (about 7-fold at 10 μM FOH; see Fig. 2B) did not increase the susceptibility to FOH (Fig. 2C, right). To verify if the *nucA*^{EndoG} deletion and overexpression mutants are more sensitive to cell death caused by other stressing agents, we exposed conidia of these strains to several concentrations/dosages of amphotericin, itraconazole, caspofungin, camptothecin, hydroxyurea, sirolimus, calcium, menadione, hydrogen peroxide, and paraquat. We were not able to observe any other phenotype

of susceptibility to these agents. However, the *nucA*^{EndoG} deletion mutant is more sensitive to 4-nitroquinoline oxide (Fig. 2D), and its quiescent and germinating conidia are slightly more sensitive to UV light (Fig. 2E).

ATM and ATR are paralogous phosphoinositide 3-kinase-related kinases (PIKKs) that orchestrate the DNA damage response in eukaryotic cells (14, 25). Aiming to see possible interactions between NucA and other elements of the DNA

damage response, we constructed $\Delta nucA$ double mutants with $\Delta atmA^{ATM}$ and $\Delta uvsB^{ATR}$, the *A. nidulans* ATM and ATR gene homologues. We have tested comparatively their growth in the presence of several DNA-damaging agents. The $\Delta nucA \Delta atmA^{ATM}$ double mutant was more sensitive to 0.0075% methyl methanesulfonate and 0.25 $\mu\text{g/ml}$ 4-nitroquinoline oxide (Fig. 3A). The $\Delta nucA$ and $\Delta uvsB^{ATR}$ double mutants were more sensitive to 0.005% MMS, 0.05 $\mu\text{g/ml}$ 4-NQO, and 0.1 mM menadione (Fig. 3B). These results indicate that NucA is genetically interacting with these two genes, aiming to repair DNA damage caused by these genotoxins. Taken together, all these results suggest that NucA does not seem to be involved in cell death, but it plays a role in the DNA-damaging response in *A. nidulans*.

We also constructed a NucA::mRFP strain that has the same degree of FOH sensitivity as the wild strain (data not shown) and exposed it to FOH. NucA::mRFP accumulation was detected along the cytoplasm but not in the nuclei when NucA::mRFP germlings were exposed or not to FOH (Fig. 4A). The same results were observed when NucA::mRFP was exposed to different concentrations of 4-NQO (data not shown). Since our microscopic localization studies did not allow us to assess the subcellular location of NucA::mRFP, we decided to perform cell fractionation studies. For the extraction of mitochondrial proteins, mitochondria were isolated from *A. nidulans* mycelium based on the method described in reference 26. Germlings grown for 16 h (at 30°C) exposed or not to 500 μM FOH for 2 h at 30°C were ground with liquid nitrogen and resuspended in 50 ml ice-cold isolation buffer (0.4 M mannitol, 0.1% [wt/vol] bovine serum albumin [BSA], 1 mM EGTA, 15 mM β -mercaptoethanol, and 0.05 mM phenylmethylsulfonyl fluoride [PMSF] in 25 mM morpholinepropanesulfonic acid [MOPS] buffer, pH 7.8). Mitochondria were enriched as follows. Cell debris was spun down three times at $3,000 \times g$ for 5 min, and then a centrifugation step at $18,000 \times g$ for 30 min followed. The mitochondrial fraction was carefully resuspended in 3 ml of resuspension buffer (0.4 M mannitol, 1 mM EGTA, 0.2 mM PMSF, and 10 mM $\text{KH}_2\text{PO}_4\text{-KOH}$, pH 7.2) and homogenized with two strokes in an Elvehjem-Potter homogenizer. This resuspension was layered on top of a three-step Percoll gradient (10 ml of 14% [vol/vol], 10 ml of 26% [vol/vol], and 5 ml of 45% [vol/vol] Percoll in 0.4 M mannitol, 1 mM EGTA, and 10 mM $\text{KH}_2\text{PO}_4\text{-KOH}$, pH 7.2). After centrifugation for 60 min at $47,000 \times g$, mitochondria were isolated from the 26%/45% interphase. To remove Percoll, purified mitochondria were centrifuged twice in resuspension buffer for 30 min at $18,000 \times g$. Both the cytoplasm and the purified organelles were finally resuspended in resuspension buffer.

Each sample assayed for the fluorescence intensity for monomeric red fluorescent protein (mRFP) (excitation, 543 nm; emission, scan from 400 to 700 nm, peaking at 550 nm) was detected by using an F4500 Hitachi fluorescence spectrophotometer. First, we used an *A. nidulans* protein located in the mitochondria, CycA::mRFP (*cycA* gene, AN6246.3, encodes cytochrome *c*), as a positive control for our cell fractionation studies. As expected, CycA::mRFP is located mainly in the mitochondria when CycA::mRFP germlings were exposed or not to FOH, although it showed about twice as much fluorescence when exposed to FOH (Fig. 4B). Interestingly, there is

an increase of the *cycA* mRNA accumulation when wild-type germlings were exposed to FOH (about 3.5, 13.3, and 7.0 times more mRNA accumulation when germlings were exposed to 10, 50, and 100 μM FOH, respectively). Cell fractionation studies showed that NucA::mRFP localizes to the mitochondria when NucA::mRFP germlings were exposed or not to FOH (Fig. 4B). Accordingly, there is an induction of about 2.5-fold in the mRFP fluorescence in the mitochondria when NucA::mRFP germlings were exposed to FOH (Fig. 4B). A cytoplasmic marker (mRFP::KipB) was used as a positive control for a protein specifically located in the cytoplasm (Fig. 4B). The *kipB* gene is not induced by FOH and encodes an *A. nidulans* kinesin that localizes to cytoplasmic astral and mitotic microtubules in a discontinuous pattern, and spots of red fluorescent protein move along microtubules toward the plus ends (18). As expected, mRFP::KipB is located predominantly in the cytoplasm (Fig. 4B). Interestingly, our cell fractionation studies suggested that CycA::mRFP does not translocate from the mitochondria to the cytoplasm when germlings were exposed to FOH. The NucA mitochondrial localization was validated by colocalization studies of NucA::mRFP with the mitochondria by using the mitochondrion-specific dye MitoTracker green (Fig. 4C). In the absence of FOH, NucA::mRFP colocalizes to mitochondrial filament-like structures, while when germlings were exposed to 100 μM FOH for either 30 or 120 min, the NucA::mRFP colocalizes to mitochondrial fragment-like structures. Taken together, these results strongly suggest that NucA localizes to the mitochondria exposed or not to FOH.

Aiming to confirm the subcellular localization of cytochrome *c*, another set of cell fractionation studies was performed with the wild-type strain, and the samples were then analyzed by 15% sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE). Samples were prepared for SDS-PAGE by the addition of $1\times$ sample buffer (62.5 mM Tris-HCl [pH 6.8], 2% SDS, 10% glycerol, 5% β -mercaptoethanol, and 5% bromophenol blue) and heating at 100°C for 3 min. Twenty micrograms of total protein from each sample was loaded into each lane of a 15% SDS-PAGE gel. After separation of the proteins, the gel was blotted onto a pure nitrocellulose membrane (0.2 μm ; Bio-Rad), and after being blocked in 5% dried milk in TBS-T buffer (10 mM Tris-HCl, 150 mM NaCl [pH 8.0], and 0.05% Tween 20), the membrane was probed with the rabbit anti-cyc1 antibody (CNAT; against native cytochrome *c* from yeast; Sigma) at a 1:200 dilution in TBS-T buffer for 1 h at room temperature. The membrane was washed four times for 5 min each with TBS-T buffer and then incubated with a 1:5,000 dilution of goat anti-rabbit IgG peroxidase-labeled (KPL) antibody for 1 h. After being washed, the blot was developed by use of the SuperSignal Ultra chemiluminescence detection system (Pierce) and recorded by the use of Hyperfilm ECL (Amersham Biosciences). The Western blot analysis showed that *A. nidulans* CycA is not translocated to the cytoplasm upon exposure to FOH (Fig. 4D). The images generated were subjected to densitometric analysis of pixel intensity using ImageJ software (<http://rsbweb.nih.gov/ij/index.html>). By normalizing the protein loading, we were able to see that there is about 2.3 times more protein upon FOH exposure (Fig. 4D). *A. nidulans* NucA localizes predominantly to the mitochondria, but in contrast to its human and yeast homologues, it does not

FIG. 4. NucA::mRFP does not translocate to the nucleus upon exposure of germlings to FOH. (A) Germlings of NucA::mRFP were grown in liquid MM for 18 h at 30°C. The 18-h germlings were either not exposed (top, control) or exposed (bottom) to 100 μ M FOH for 1 h at 30°C. After the treatment, the germlings were fixed, washed, and DAPI (4',6-diamidino-2-phenylindole) stained. Samples were analyzed by laser scanning confocal microscopy. Images were captured by direct acquisition. Bars, 5 μ m. (B) Cell fractionation studies of the CycA::mRFP and NucA::mRFP strains exposed or not to 500 μ M FOH. Graphs are representative of results from at least three independent assays. (C) NucA::mRFP localizes to the mitochondria. This experiment was performed as described in the legend to Fig. 4A. Mitochondrial staining of live cells was performed using MitoTracker green (Invitrogen Molecular Probes). (D) Protein extracts from the wild-type strain (cell fractionation protein assays) were analyzed by Western blotting using antibodies directed against *S. cerevisiae* Cyc1p. Coomassie staining of these protein fractions was used as a loading control by using ImageJ software (<http://rsbweb.nih.gov/ij/index.html>).

FIG. 4—Continued.

migrate to the nucleus upon FOH-induced cell death. The *A. nidulans* *nucA*^{EndoG} deletion mutant is sensitive neither to FOH nor to several other cell death inducers. Actually, we investigated a possible role for NucA in DNA damage by constructing double mutants with the two major kinases that participate in signaling during DNA damage, ATM and ATR. There is a synergistic interaction between *A. nidulans* $\Delta atmA$, $\Delta atrA$, and $\Delta nucA$ deletion mutants, suggesting that these genes collaborate in parallel pathways for DNA repair. Thus, it is possible that NucA plays a role in mitochondrial recombination and proliferation, as it was previously observed for mammalian cells (10).

During the fractionation studies, we used CycA::mRFP (*cycA* encodes cytochrome *c*) as a positive control for mitochondrial localization. Our results showed that CycA::mRFP is located predominantly in the mitochondria, even when CycA::mRFP germlings were exposed to FOH. Interestingly,

during *A. nidulans* FOH-induced cell death, not only is the cytochrome *c* not released to the cytoplasm, but caspases are activated (3). This could reflect a particular feature of the FOH-induced cell death or a more general observation related to cell death in filamentous fungi linked to the fact that regulation and expression of these genes involved in apoptosis could have undergone a rewiring process in mammals during their evolution. However, in *S. cerevisiae* cells undergoing an apoptotic process induced by acetic acid, translocation of CytC to the cytosol and reactive oxygen species (ROS) production were observed (13). The behavior of *A. nidulans* CycA in response to different cell death stimuli remains to be investigated.

In summary, our study shows that the behaviors of two genetic determinants of the intrinsic pathway of cell death in higher eukaryotes, EndoG and cytochrome *c*, are different in FOH-induced cell death in the filamentous fungus *A. nidulans*.

This opens the interesting possibility that cell death in filamentous fungi is activated by distinct (and probably novel) pathways from animals, opening exciting new avenues for research into new targets for the control of fungal proliferation.

This research was supported by the Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP), Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Brazil.

We thank the two anonymous reviewers for their suggestions.

REFERENCES

- Burhans, W., and M. Weinberger. 2007. Yeast endonuclease G: complex matters of death, and of life. *Mol. Cell* **25**:323–325.
- Büttner, S., et al. 2007. Endonuclease G regulates budding yeast life and death. *Mol. Cell* **25**:233–246.
- Colabardini, A. C., et al. 2010. Involvement of the *Aspergillus nidulans* protein kinase C with farnesol tolerance is related to the unfolded protein response. *Mol. Microbiol.* **78**:1259–1279.
- Colot, H. V., et al. 2006. A high-throughput gene knockout procedure for *Neurospora* reveals functions for multiple transcription factors. *Proc. Natl. Acad. Sci. U. S. A.* **103**:10352–10357.
- Dinamarco, T. M., et al. 2010. The roles played by *Aspergillus nidulans* apoptosis-inducing factor (AIF)-like mitochondrial oxidoreductase (AifA) and NADH-ubiquinone oxidoreductases (NdeA-B and NdiA) in farnesol resistance. *Fungal Genet. Biol.* **47**:1055–1069.
- Elmore, S. 2007. Apoptosis: a review of programmed cell death. *Toxicol. Pathol.* **35**:45–516.
- Flipphi, M., J. Kocalkowska, and B. Felenbok. 2002. Characteristics of physiological inducers of the ethanol utilization (alc) pathway in *Aspergillus nidulans*. *Biochem. J.* **15**:25–31.
- Goldman, G. H., et al. 2003. Expressed sequence tag analysis of the human pathogen *Paracoccidioides brasiliensis* yeast phase: identification of putative homologues of *Candida albicans* virulence and pathogenicity genes. *Eukaryot. Cell* **2**:34–48.
- Hamann, A., D. Brust, and H. D. Osiewacz. 2008. Apoptosis pathways in fungal growth, development and ageing. *Trends Microbiol.* **16**:276–283.
- Huang, K.-J., C.-C. Ku, and I. R. Lehman. 2006. Endonuclease G: a role for the enzyme in recombination and cellular proliferation. *Proc. Natl. Acad. Sci. U. S. A.* **103**:8995–9000.
- Kafer, E. 1977. Meiotic and mitotic recombination in *Aspergillus* and its chromosomal aberrations. *Adv. Genet.* **19**:33–131.
- Li, L. Y., X. Luo, and X. Wang. 2001. Endonuclease G is an apoptotic DNase when released from mitochondria. *Nature* **412**:95–99.
- Ludovico, P., et al. 2002. Cytochrome *c* release and mitochondria involvement in programmed cell death induced by acetic acid in *Saccharomyces cerevisiae*. *Mol. Biol. Cell* **13**:2598–2606.
- McKinnon, P. J. 2004. ATM and ataxia telangiectasia. *EMBO Rep.* **5**:772–776.
- Osmari, S. A., G. S. May, and N. R. Morris. 1987. Regulation of the mRNA levels of *nimA*, a gene required for the G2-M transition in *Aspergillus nidulans*. *J. Cell Biol.* **104**:1495–1504.
- Parrish, J., et al. 2001. Mitochondrial endonuclease G is important for apoptosis in *C. elegans*. *Nature* **412**:90–94.
- Ramsdale, M. 2008. Programmed cell death in pathogenic fungi. *Biochim. Biophys. Acta* **1783**:1369–1380.
- Rischitor, P., S. Konzack, and R. Fischer. 2004. The Kip3-like kinesin KIPB moves along microtubules and determines spindle position during synchronized mitoses in hyphae of *Aspergillus nidulans*. *Eukaryot. Cell* **3**:632–645.
- Sambrook, J., and D. W. Russell. 2001. Molecular cloning: a laboratory manual, 3rd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Savoldi, M., et al. 2008. Farnesol induces the transcriptional accumulation of the *Aspergillus nidulans* apoptosis-inducing factor (AIF)-like mitochondrial oxidoreductase. *Mol. Microbiol.* **70**:44–59.
- Schiestl, R. H., and R. D. Gietz. 1989. High efficiency transformation of intact yeast cells using single stranded nucleic acids as a carrier. *Curr. Genet.* **16**:339–346.
- Semighini, C. P., M. Marins, M. H. S. Goldman, and G. H. Goldman. 2002. Quantitative analysis of the relative transcript levels of ABC transporter *Atr* genes in *Aspergillus nidulans* by real-time reverse transcription-PCR assay. *Appl. Environ. Microbiol.* **68**:1351–1357.
- Semighini, C. P., M. Savoldi, G. H. Goldman, and S. D. Harris. 2006. Functional characterization of the putative *Aspergillus nidulans* poly(ADP-ribose) polymerase homolog PrpA. *Genetics* **173**:87–98.
- Sharon, A., A. Finkelstein, N. Shlezingers, and I. Hatam. 2009. Fungal apoptosis: function, genes and gene function. *FEMS Microbiol. Rev.* **33**:833–854.
- Shiloh, Y., and M. B. Kastan. 2001. ATM: genome stability, neuronal development, and cancer cross paths. *Adv. Cancer Res.* **83**:209–254.
- Vödisch, M., et al. 2009. Two-dimensional proteome reference maps for the human pathogenic filamentous fungus *Aspergillus fumigatus*. *Proteomics* **9**:1407–1415.

7.5. Anexo N. Artigo publicado: Molecular characterization of the putative transcription factor SebA involved in virulence in *Aspergillus fumigatus*. Dinamarco, T. M.; Almeida, R. S.; de Castro, P. A.; Reis, T. F; Ramalho, L. N. Z.; Savoldi, M.; Goldman, M. H. S.; Goldman, G. H.. *Eukaryotic Cell*, Apr. 2012, 11(4):518-31.

Molecular Characterization of the Putative Transcription Factor SebA Involved in Virulence in *Aspergillus fumigatus*

Taísa Magnani Dinamarco, Ricardo S. Almeida, Patrícia Alves de Castro, Neil Andrew Brown, Thaila Fernanda dos Reis, Leandra Naira Zambelli Ramalho, Marcela Savoldi, Maria Helena S. Goldman and Gustavo Henrique Goldman
Eukaryotic Cell 2012, 11(4):518. DOI: 10.1128/EC.00016-12.
Published Ahead of Print 17 February 2012.

Updated information and services can be found at:
<http://ec.asm.org/content/11/4/518>

SUPPLEMENTAL MATERIAL

These include:

<http://ec.asm.org/content/suppl/2012/03/26/11.4.518.DC1.html>

REFERENCES

This article cites 66 articles, 26 of which can be accessed free at: <http://ec.asm.org/content/11/4/518#ref-list-1>

CONTENT ALERTS

Receive: RSS Feeds, eTOCs, free email alerts (when new articles cite this article), [more»](#)

Information about commercial reprint orders: <http://journals.asm.org/site/misc/reprints.xhtml>
To subscribe to to another ASM Journal go to: <http://journals.asm.org/site/subscriptions/>

Molecular Characterization of the Putative Transcription Factor SebA Involved in Virulence in *Aspergillus fumigatus*

Taísa Magnani Dinamarco,^b Ricardo S. Almeida,^b Patrícia Alves de Castro,^b Neil Andrew Brown,^b Thaila Fernanda dos Reis,^b Leandra Naira Zambelli Ramalho,^d Marcela Savoldi,^b Maria Helena S. Goldman,^c and Gustavo Henrique Goldman^{a,b}

Laboratório Nacional de Ciência e Tecnologia do Bioetanol—CTBE, São Paulo, Brazil,^a and Faculdade de Ciências Farmacêuticas de Ribeirão Preto,^b Faculdade de Filosofia, Ciências e Letras de Ribeirão Preto,^c and Faculdade de Medicina de Ribeirão Preto,^d Universidade de São Paulo, São Paulo, Brazil

Aspergillus fumigatus is a major opportunistic pathogen and allergen of mammals. Nutrient sensing and acquisition mechanisms, as well as the capability to cope with different stressing conditions, are essential for *A. fumigatus* virulence and survival in the mammalian host. This study characterized the *A. fumigatus* SebA transcription factor, which is the putative homologue of the factor encoded by *Trichoderma atroviride* *seb1*. The Δ *sebA* mutant demonstrated reduced growth in the presence of paraquat, hydrogen peroxide, CaCl₂, and poor nutritional conditions, while viability associated with *sebA* was also affected by heat shock exposure. Accordingly, SebA::GFP (SebA::green fluorescent protein) was shown to accumulate in the nucleus upon exposure to oxidative stress and heat shock conditions. In addition, genes involved in either the oxidative stress or heat shock response had reduced transcription in the Δ *sebA* mutant. The *A. fumigatus* Δ *sebA* strain was attenuated in virulence in a murine model of invasive pulmonary aspergillosis. Furthermore, killing of the Δ *sebA* mutant by murine alveolar macrophages was increased compared to killing of the wild-type strain. *A. fumigatus* SebA plays a complex role, contributing to several stress tolerance pathways and growth under poor nutritional conditions, and seems to be integrated into different stress responses.

Aspergillus fumigatus is a major opportunistic pathogen and allergen of mammals that is readily isolated from human habitats and vegetable compost heaps (4, 9, 17, 25, 33). *A. fumigatus* produces prodigious amounts of conidia; as a consequence, the human respiratory tract is constantly exposed to the fungus, which is increasingly linked to severe asthma and sinusitis (3, 11, 15, 37). The incidence of invasive infection in immunocompromised individuals can be as high as 50%, often with a higher mortality rate (7, 15). Despite the great medical and ecological importance of this pathogen, the basic biology of the organism is poorly understood, although the completed *A. fumigatus* genome sequence (40) offers new opportunities to probe the genetic basis of *A. fumigatus* biology and physiology. Sensing and nutrient acquisition mechanisms are essential for *A. fumigatus* virulence (2, 61). Fungi sense and respond to the immediate environment via signal transduction cascades, which transduce external information and trigger appropriate responses (53). In opportunistic fungi, such as *A. fumigatus*, transcription factors (TFs) represent an important genetic component for the establishment of infection by activating or repressing different mechanisms that regulate virulence and pathogenicity. Recently, null mutants for 11 *A. fumigatus* TFs, including AcuM, the zinc cluster transcription factor, which has been shown to be essential for virulence in murine models of hematogenously disseminated and invasive pulmonary aspergillosis, were constructed and characterized (35). Transcriptional profiling suggests that AcuM suppresses *sreA* and induces *hapX* to stimulate the expression of genes involved in both reductive iron assimilation and siderophore-mediated iron uptake. Another TF important for virulence and pathogenicity in *A. fumigatus* is CrzA, which is modulated by calcineurin (54, 55). In addition to the identification of TFs, the characterization of the genetic mechanisms that regulate the establishment of *A. fumigatus* infection and disease progression represents an important challenge.

In the present study, the *A. fumigatus* homologue of the TF *seb1* (here named *sebA*) was characterized. The *Trichoderma atroviride*

TF *seb1* has been shown to be involved in the osmotic stress response (43, 51). *T. atroviride* *seb1* encodes a C2H2 zinc finger protein that is ~60% identical, in the zinc finger region, to the *Saccharomyces cerevisiae* TFs Com2p (a protein closely related to Msn2), Msn2p, and Msn4p. *S. cerevisiae* Com2p is a protein of unknown function, while Msn2 and Msn4 participate in multiple stress responses (10, 20, 22, 29). The *Candida albicans* Com2p homologue, Mnl1p, regulates the weak acid stress responses (46). Here, *A. fumigatus* SebA was demonstrated to be involved in tolerance of oxidative and heat shock stresses, in addition to adaptation to poor nutritional conditions. This role as a TF involved in stress responses has a dramatic impact on *A. fumigatus* virulence and pathogenicity, since a Δ *sebA* strain showed attenuated virulence in a murine model of invasive pulmonary aspergillosis. Interestingly, the Δ *sebA* mutant strain was also increasingly killed by murine alveolar macrophages (AM) compared to the wild-type strain.

MATERIALS AND METHODS

Strains and culture conditions. The *A. fumigatus* strains used in this study were CEA17 (*pyrG* mutant) (for the transformation assays), CEA17-80 (as the wild-type strain), Δ *calA* and Δ *crzA* (55), Δ *sebA* (Δ *sebA*::*pyrG*), and Δ *sebA*::*sebA*⁺. Three variants of a complete medium were used: YAG (2% glucose, 0.5% yeast extract, 2% agar, trace elements), YUU (YAG supplemented with 1.2 g/liter of both uracil and uridine), and liquid YG medium (consisting of the same compositions but without agar). A modified minimal medium (MM; 1% glucose, the original high

Received 13 January 2012 Accepted 10 February 2012

Published ahead of print 17 February 2012

Address correspondence to Gustavo H. Goldman, ggoldman@usp.br.

Supplemental material for this article may be found at <http://ec.asm.org/>.

Copyright © 2012, American Society for Microbiology. All Rights Reserved.

doi:10.1128/EC.00016-12

levels of nitrate salts, trace elements, 2% agar, pH 6.5) was also used. The trace elements, vitamins, and nitrate salts in the media are described in reference 31. Additionally, 10% fetal bovine serum (FBS; Gibco) was used as a medium. For the analysis of growth on lung tissue, the methodology described in reference 21 was followed. Briefly, dissected mouse lungs were placed on the surface of a 1% agarose plate in sterile distilled water. The lung tissue was inoculated with 2,000 conidia in 5 μ l of sterile water and grown for 2 days at 37°C. Growth under starvation conditions was determined according to reference 45. Briefly, ~1,000 conidia were spread onto the surface of a YG plate and incubated for 16 h at 37°C. Hyphal plugs containing individual colonies were obtained using a sterile 1-ml tip and transferred onto the center of a 1% agarose plate, and radial growth was determined after 3 days of incubation at 37°C.

Staining and microscopy. SebA::GFP (SebA::green fluorescent protein) conidiospores were grown on coverslips in 4 ml of YG medium for 16 h at 30°C. After incubation, subsets of coverslips with adherent germ-lings were left untreated, treated with paraquat (10 mM), H₂O₂ (2 mM), and CaCl₂ (200 mM), or submitted to heat shock (30°C to 37°C and 30°C to 44°C). Subsequently, the coverslips were rinsed with phosphate-buffered saline (PBS; 140 mM NaCl, 2 mM KCl, 10 mM NaHPO₄, 1.8 mM KH₂PO₄, pH 7.4) and incubated for 3 min in a solution with Hoechst stain (Life Technologies) (12 μ g/ml). After incubation with the dye, the coverslips were washed with PBS and mounted for examination. Slides were visualized on an Observer Z1 fluorescence microscope using a 100 $\times$ objective oil immersion lens (for GFP, filter set 38—high efficiency [HE], excitation wavelength of 450 to 490 nm, and emission wavelength of 525/50 nm; for Hoechst stain, filter set 49, excitation wavelength of 365 nm, and emission wavelength of 420 to 470 nm). DIC (differential interference contrast) images and fluorescent images were captured with an AxioCam camera (Carl Zeiss) and processed using AxioVision software (version 4.8).

Construction of *A. fumigatus* sebA disruption mutant and SebA::GFP. A gene replacement cassette was constructed by *in vivo* recombination in *S. cerevisiae* as previously described in reference 14. Briefly, ~2.0-kb regions on either side of the *sebA* open reading frame (ORF) (Afu4g09080) were selected for primer design. For construction, the primers (named 5F and 5R) were used to amplify the flanking 5' untranslated region (5'-UTR) of the targeted ORF. Likewise, primers 3F and 3R were used to amplify the 3'-UTR ORF flanking region. The primers 5F and 3R also contained a short sequence homologous to the multiple-cloning site (MCS) of the pRS426 plasmid. Both fragments (the 5'-UTR and 3'-UTR) were amplified from genomic DNA (gDNA) by PCR using *A. fumigatus* gDNA as a template. The *A. fumigatus* pyrG selectable marker for uridine or uracil prototrophy was used for generating the Δ sebA strain.

Deletion cassettes were generated by transforming each fragment along with the cut plasmid, pRS426 BamHI-EcoRI, into *S. cerevisiae* SC94721 via the lithium acetate method (14, 49). The DNA of the *S. cerevisiae* transformants was extracted as described in reference 23, dialyzed, and transformed by electroporation in *Escherichia coli* DH10B to rescue the pRS426 plasmid harboring the deletion cassette. The cassette was PCR amplified from the plasmids and used for *A. fumigatus* transformation. Southern blot analysis was used to demonstrate that the cassettes had integrated homologously at the targeted *A. fumigatus* locus. To complement the Δ sebA strain, a region of ~2.0 kb flanking the *sebA* gene was amplified using gDNA from CEA10 as a template. Protoplasts were cotransformed with the 6.8-kb PCR product and the hygromycin B resistance vector (pHAT- α) and then plated on media containing hygromycin B (150 μ g/ml).

To generate the SebA::GFP strain, the *sebA* ORF was cloned in frame with the green fluorescent protein (GFP) gene. The construct links GFP to the C terminus of *sebA* and is separated by four additional codons that, after translation, produce a four-amino-acid linker (glycine-threonine-arginine-glycine) (59). The *S. cerevisiae* *in vivo* recombination system was used for production of the transformation cassette, and then the fragments were PCR amplified. First, the *sebA* ORF and its 5'-UTR flanking

region were amplified from gDNA of the wild-type strain by the use of the primers *sebA* pRS426 5f and *sebA* SPACER GFP Rev. The stop codon of the *sebA* gene was omitted in this construction. The GFP ORF was amplified from the pMCB17apx plasmid (provided by Vladimir P. Efimov) by the use of the primers Spacer GFP Fwd and GFP VE3' AF. The last fragment was PCR amplified from the *sebA* deletion cassette and contained the selective marker *pyrG*. The primers used for this PCR amplification were GFP *pyrG* Fwd and *pyrG* Rev. The amplification of the 3'-UTR was done with the *Afu sebA* 3f and *Afu sebA* 3r primers. *S. cerevisiae* *in vivo* recombination was performed as previously described. The PCR-amplified cassette was transformed into the *A. fumigatus* wild-type strain.

Heat and oxidative stress treatments. The heat shock viability assay (41) was carried out with wild-type and Δ sebA strains. Conidia (1×10^6 ml⁻¹) were incubated in YG for 5 h at 30°C in a shaker (250 rpm). After this period, the germ-lings were exposed to 37°C and 48°C for 15, 30, and 60 min. The untreated control group was maintained at 30°C. Conidia were collected, diluted, and plated on YAG plates. Plates were incubated at 37°C for 48 h. Viability was determined as the percentage of colonies on treated plates compared with untreated controls.

For real-time analysis, heat shock treatment was done according to reference 40. For temperature-shift experiments, conidia (5×10^6 ml) from the wild-type strain were incubated in YG medium for 17 h at 30°C. Cultures were then transferred to 37°C or 48°C for continued growth, for 15, 30, 60, and 120 min. Total RNA extracted from samples before (0 min) and after (15, 30, 60, and 120 min) the two temperature shifts (30°C to 37°C and 30°C to 48°C) were used to profile the *sebA* gene expression.

The *sebA* gene expression was also analyzed under different osmotic and oxidative stress conditions. Cultures containing 10^7 wild-type or Δ sebA conidia were grown overnight in YG media and then treated with either 2 mM H₂O₂ or 10 mM paraquat for 15 and 30 min to assess oxidative stress tolerance and 1.2 M KCl or 1.2 M sorbitol for 30, 60, 120, and 180 min to assess osmotic stress tolerance.

RNA extraction and real-time PCRs. After the different conditional treatments, mycelia were harvested by filtration, washed twice with water, and immediately frozen in liquid nitrogen. Total RNA was isolated from germ-lings disrupted by grinding in liquid nitrogen with a pestle and mortar and extracted using TRIzol reagent (Invitrogen). RNA (10 μ g) from each treatment was fractionated on a 2.2 M formaldehyde–1.2% (wt/vol) agarose gel, stained with ethidium bromide, and visualized under UV light. The presence of intact 25S and 18S rRNA bands was used to assess RNA integrity. RNase-free DNase I treatment was carried out as previously described (52). A 20- μ g volume of total RNA was treated with DNase and purified using an RNeasy kit (Qiagen). cDNA was synthesized using a SuperScript III First Strand Synthesis system (Invitrogen) and oligo(dT) primers according to the manufacturer's protocol.

All PCRs were performed using an ABI 7500 Fast real-time PCR system (Applied Biosystems) and a TaqMan Universal PCR Master Mix kit (Applied Biosystems). The reactions and calculations were performed according to reference 52. The primers and Lux fluorescent probes (Invitrogen) used are described in Table S1 in the supplemental material.

Murine model of pulmonary aspergillosis. Outbreed female mice (BALB/c strain; body weight, 20 to 22 g) were housed in vented cages containing 5 animals. Mice were immunosuppressed with cyclophosphamide at 150 mg/kg of body weight administered intraperitoneally on days -4, -1, and 2 prior to and postinfection. Hydrocortisonacetate (200 mg/kg) was injected subcutaneously on day -3 (modified from reference 39). *A. fumigatus* spores used for inoculation were grown on *Aspergillus* complete medium for 2 days prior to infection. Conidia were freshly harvested in PBS and filtered through a Miracloth (Calbiochem). Conidial suspensions were spun for 5 min at $3,000 \times g$, washed three times with PBS, counted using a hemocytometer, and resuspended at a concentration of 2.5×10^6 conidia/ml. Viable counts of the administered inocula were determined, following serial dilution, by plating on *Aspergillus* complete medium, and the conidia were grown at 37°C. Mice were anesthetized by halothane inhalation and infected by intranasal instillation of

FIG 1 Growth phenotypes of the $\Delta sebA$ mutant strain. The wild-type, the $\Delta sebA::sebA^+$, and the $\Delta sebA$ mutant strains were grown for 72 h in YAG medium (A), YAG medium plus paraquat at different concentrations (B), or YAG medium or YAG medium plus paraquat at 4 mM (C) or for 48 h at 37°C in YAG medium plus different concentrations of H_2O_2 (D).

5.0×10^4 conidia in 20 μ l of PBS. As a negative control, a group of 5 mice received PBS only. Mice were weighed every 24 h from the day of infection and visually inspected twice daily. In the majority of cases, the endpoint for survival experimentation was identified when a 20% reduction in body weight was recorded, at which time the mice were sacrificed. The statistical significance of comparative survival values was calculated using log rank analysis and the Prism statistical analysis package. Additionally, at 3 days postinfection, 2 mice per strain were sacrificed and the lungs were removed, fixed, and processed for histological analysis.

Lung histopathology and fungal burden. After sacrifice, the lungs were removed and fixed for 24 h in 3.7% formaldehyde–PBS. Samples were washed several times in 70% alcohol before dehydration in a series of alcohol solutions of increasing concentrations. Finally, the samples were diafanized in xylol and embedded in paraffin. For each sample, sequential 5- μ m-thick sections were collected on glass slides and stained with Gomori methenamine silver (GMS) or hematoxylin and eosin (HE) stain following standard protocols (25). Briefly, sections were deparaffinized, oxidized with 4% chromic acid, stained with methenamine silver solution, and counterstained with picric acid. For HE staining, sections were deparaffinized and stained first with hematoxylin and then with eosin. All stained slides were immediately washed, preserved with mounting medium, and sealed with a coverslip. Microscopic analyses were done using an Axioplan 2 imaging microscope (Zeiss) at the stated magnifications under bright-field conditions.

To investigate fungal burden in murine lungs, mice were immunosuppressed with cyclophosphamide at 150 mg/kg of body weight administered intraperitoneally on days -4 and -1 , while hydrocortisonacetate was injected subcutaneously at 200 mg/kg on day -3 . Five mice per group (wild-type, $\Delta sebA$, and $\Delta sebA::sebA$ strains and PBS control) were intranasally inoculated with 1×10^6 conidia/20 μ l of suspension. A higher

inoculum, in comparison to the survival experiments, was used to increase fungal DNA detection. Animals were sacrificed 72 h postinfection, and both lungs were harvested and immediately frozen in liquid nitrogen. Samples were homogenized with glass beads by performing a vortex procedure for 10 min, and DNA was extracted via the phenol-chloroform method. DNA quantity and quality were assessed using a NanoDrop 2000 spectrophotometer (Thermo Scientific). At least 500 μ g of total DNA from each sample was used for quantitative real-time PCRs. A primer and a Lux probe (Invitrogen) were used to amplify the 18S rRNA region of *A. fumigatus* (primer, 5'-CTTAAATAGCCCGTCCGCATT-3'; probe, 5'-CATCACAGACCTGTTATTGCCG-3') and an intronic region of mouse GAPDH (glyceraldehyde-3-phosphate dehydrogenase) (primer, 5'-CGA GGGACTTGGAGGACACAG-3'; probe, 5'-GGGCAAGGCTAAAGGTC AGCG-3'). Six-point standard curves were calculated using serial dilutions of gDNA from all the *A. fumigatus* strains used and the noninfected mouse lung. Fungal and mouse DNA quantities were obtained from the threshold cycle (C_T) values from an appropriate standard curve. Fungal burden was determined as the ratio between picograms of fungal and micrograms of mouse DNA.

Phagocytosis index. To estimate the percentage of phagocytosed conidia, alveolar macrophages were harvested from lungs of BALB/c mice (8 to 10 weeks old) by using an intravenous catheter (Angiocath; Becton Dickinson) (18 gauge, 1.88 in. long). Six mice per experiment were sacrificed; the trachea was exposed and the catheter attached. A 1-ml volume of RPMI 1640 medium (Sigma-Aldrich) was introduced and subsequently removed from the lungs three times. Macrophages from all six animals were mixed and centrifuged at 4,000 rpm for 5 min. The supernatant was removed, and the pellet was washed with 5 ml of RPMI 1640 medium and resuspended in 1 ml of RPMI 1640–10% fetal calf serum (FCS) (Gibco). Cells were counted in a hemocytometer. The phagocytic assay was per-

FIG 2 Growth phenotypes of the $\Delta sebA$ mutant strain. (A and B) The wild-type, the $\Delta sebA::sebA^+$, and the $\Delta sebA$ mutant strains were grown for 72 h in either MM or FBS (10%) medium at different temperatures (A) or in FBS (10%) liquid medium from 24 to 96 h at 37°C (B) (data represent the average results of three experiment repetitions $\pm$ standard deviations). (C) *SebA* is important for growth on lung tissue. Equal numbers of conidia were inoculated on dissected mouse lung tissue that was plated on 1% agarose in sterile distilled water and incubated at 37°C for 48 h. (D) *SebA* is required under nutrient starvation conditions. Agar plugs containing hyphae from 16 h of growth on YG plates were transferred to starvation medium (1% agarose in sterile distilled water), and colony diameters were measured after 3 days at 37°C. Values represent the averages of the results of nine biological experiment repetitions $\pm$ standard deviations (*, $P < 0.0001$).

formed according to Mech and coworkers (38) with some changes. Briefly, in a 24-well plate containing one 15-mm-diameter coverslip per well, 5×10^4 macrophages were incubated with 1 ml of RPMI 1640–10% FCS (assay medium) at 37°C with 5% CO_2 for 1 h. Afterward, the wells were washed with 1 ml of assay medium to remove nonadherent cells. In each well, 1 ml of assay medium containing 2.5×10^5 conidia (5 conidia/macrophage) was added. Duplicate wells were assayed for each strain. The samples were incubated at 37°C with 5% CO_2 for 80 min, after which the supernatant was removed and 500 μl of 3.7% formaldehyde–PBS was added. After 15 min, the samples were washed with 1 ml of ultrapure water and incubated for 20 min with 495 μl of water–5 μl of calcofluor (10 mg/ml). The samples were washed and mounted on slides with 50% glycerol. A Zeiss Observer.Z1 fluorescence microscope was used to assess the percentage of phagocytosed spores. Macrophage cells were not permeable; hence, conidia inside the macrophages were not stained, while conidia outside the macrophages were stained, by calcofluor. At least 100 spores were counted per sample, and a phagocytosis index was calculated. The experiments were repeated in triplicate.

Conidial killing by alveolar macrophages. To assess conidial killing by alveolar macrophages, the phagocytic cells were obtained as described above. In a 96-well plate, 5×10^4 macrophages were added with 200 μl of RPMI 1640–10% FCS per well and incubated at 37°C with 5% CO_2 for 1 h. Afterwards, 2.5×10^5 conidia (5 conidia/macrophage) were added and incubated at 37°C with 5% CO_2 for 4 h. As a positive control, wells containing medium and spores without macrophages were used. Triplicate wells were assayed for each strain with and without macrophages. After incubation, the plate was centrifuged at 3,500 rpm for 10 min, supernatant was removed without disturbing the pellet, and 100 μl of 1% Triton X-100 was added. After 10 min at room temperature, samples were removed from the plate and serially diluted in sterile water. The dilutions were plated on *A. fumigatus* complete medium (38) and incubated at 37°C for 2 days. The percentage of conidial killing was calculated by comparing

CFU numbers from samples incubated with macrophages to CFU numbers from those incubated without macrophages. The experiments were repeated three times.

RESULTS

There is a single *Seb1* homologue in *A. fumigatus*. A BLASTp search of the *A. fumigatus* genome database (<http://www.aspgd.org>) performed using *T. atroviride seb1* as a query revealed a single open reading frame with significant similarity. The potential homologue, Afu4g09080 (here named *sebA*), is predicted to be a 618-amino-acid zinc finger C2H2 protein with homology to the *T. atroviride* *Seb1* (from residue 477 to residue 616; *e*-value, 1.2×10^{-55} ; 70.2% similarity and 61.7% identity). *SebA* shares identity, only in the zinc finger domain, with the *S. cerevisiae* TFs Msn2p (from residue 483 to residue 544; *e*-value, 7.2×10^{-17} ; 72% similarity and 54% identity), Msn4p (from residue 437 to residue 544; *e*-value, 1.3×10^{-15} ; 57% similarity and 38% identity), and Com2p (from residue 375 to residue 440; *e*-value, 4.2×10^{-20} ; 72% similarity and 68% identity). Identity with these *S. cerevisiae* homologues is mainly located in the two zinc finger C2H2-type domains (PROSITE entry PS50157; <http://www.expasy.org>) located at the region spanning residue 493 to residue 548 (FVCSLCSRRFRFRQEHLKRHYRSLHTQDKPFECHECGKKFSRSDNLAQHARTHGGGS).

Subsequently, the *A. fumigatus* *SebA* null mutant ($\Delta sebA$) was generated using an *in vivo* *S. cerevisiae* fusion-based approach (see Materials and Methods). Replacement of the *sebA* gene in several transformants was verified by Southern blot analysis; one of the transformants was used for further phenotypic characterization (see Fig. S1 in the supplemental material). The $\Delta sebA$ phenotype

FIG 3 SebA::GFP accumulates in the nuclei upon oxidative stress. The SebA::GFP strain was grown for 16 h at 30°C in YG medium and incubated 1 h more at 30°C (A) or transferred to either H_2O_2 (2 mM) for 15, 30, and 60 min at 30°C (B) or paraquat (10 mM) for 15, 30, and 60 min (C). Bars, 5 μm .

was complemented with *sebA* from the wild-type strain, producing the $\Delta\text{sebA}::\text{sebA}^+$ strain (see Fig. S1 in the supplemental material). All strains were able to grow on uracil, uridine, and hygromycin. The ΔsebA mutant showed radial growth comparable to that of the wild-type and complemented strains, except at lower temperatures (such as 25°C and 30°C), when the radial growth was reduced (Fig. 1A). The ΔsebA mutant was more sensitive to oxidative stress factors, such as paraquat and H_2O_2 (at 37°C), than the wild-type and complemented strains, as shown by the reduction in growth in liquid complete medium (Fig. 1B and D) and solid complete medium (Fig. 1C). No growth reduction was observed when the ΔsebA mutant was grown under osmotic stress conditions, such as in the presence of increasing concentrations of KCl, NaCl, or sorbitol (data not shown). The reduction in radial growth was greater when the ΔsebA mutant was grown on minimal medium (at 37°C) and even greater at different temperatures on solid FBS (10%) medium (Fig. 2A). Growth of the wild-type, $\Delta\text{sebA}::\text{sebA}^+$, and ΔsebA mutant strains in FBS (10%) liquid medium from 24 to 96 h was evaluated (Fig. 2B). The ΔsebA mutant demonstrated a dramatic (between 60 and 70%) reduction in dry mycelial weight compared to the wild-type and complemented strains (Fig. 2B). Growth characteristics on YAG medium and on dissected mouse lung tissue were also compared. As previously shown (Fig. 1), the wild-type and ΔsebA and $\Delta\text{sebA}::\text{sebA}^+$ mutant strains had the same radial diameter after 48 h of growth on YAG medium (Fig. 2C); however, the ΔsebA mutant strain had reduced

growth on lung tissue after 48 h (Fig. 2C). Growth of the ΔsebA strain was also impaired on lung tissue, *in vivo*, as shown by the histopathology of infected lungs (see Fig. 9B) and fungal burden quantifications (see Fig. 9C). To verify whether SebA is important for growth under nutrient-limiting conditions, the wild-type, $\Delta\text{sebA}::\text{sebA}^+$, and ΔsebA strains were grown under starvation conditions as described in reference 45. Both the wild-type and $\Delta\text{sebA}::\text{sebA}^+$ strains were able to produce restricted growth in the absence of extracellular nutrients, while the ΔsebA strain had significantly reduced growth under starvation conditions (Fig. 2D). The ΔsebA mutant showed no differential susceptibility to antifungal agents, such as amphotericin, azoles, and caspofungin, in E-tests or susceptibility to cell wall-targeting agents, such as Congo red (data not shown).

To assess the subcellular location of SebA, we constructed a SebA::GFP strain. This strain behaves exactly like the wild-type strain (data not shown). When grown on YG for 16 h at 30°C, it showed a weak and diffuse fluorescent signal in the cytosol (Fig. 3A). In contrast, when this strain was transferred to YG–2 mM H_2O_2 (Fig. 3B) or YG–10 mM paraquat (Fig. 3C) for 15, 30, or 60 min, SebA::GFP progressively accumulated in the nucleus, as shown by Hoechst costaining (Fig. 3B and C).

Taken together, these results suggest that SebA is important for the oxidative stress response and plays a role during growth under poor nutrient conditions.

The transcriptional basis for *A. fumigatus* ΔsebA sensitivity to oxidative stress. The *A. fumigatus* ΔsebA strain was shown to be

FIG 4 The transcriptional basis for *A. fumigatus* Δ *sebA* sensitivity to hydrogen peroxide. The wild-type, Δ *sebA*, and Δ *sebA*::*sebA*⁺ strains were grown for 16 h in YG at 37°C and transferred to fresh YG medium with either 0 or 2 mM H₂O₂ and grown for a further 15 or 30 min. The relative quantitations of *Afyap1* (A), *Afatf1* (B), *Afccp1* (C), *Afcac1* (D), *Afcac2* (E), *Afsod1* (F), and *Afsod2* (G) and tubulin gene expression were determined by a standard curve (i.e., C_T values plotted against a logarithm of the DNA copy number). Data shown represent the means ($\pm$ standard deviations) of the results of four biological replicate experiments (*, $P < 0.001$).

more sensitive to paraquat and H₂O₂ (Fig. 1B). Therefore, a transcriptional approach was used to characterize the transcriptional basis of the oxidative stress sensitivity of the Δ *sebA* mutant. Several *A. fumigatus* homologues involved in oxidative stress tolerance were identified, Lux fluorescent probes were designed for these genes, and then gene expression in response to H₂O₂ and paraquat was determined in the wild-type, Δ *sebA*::*sebA*⁺, and Δ *sebA* strains via real-time reverse transcription-PCR (RT-PCR) (Fig. 4 and 5). In the wild-type strain, *sebA* transcription did not increase in the presence of H₂O₂ (2 mM) or paraquat (10 mM) for 15 and 30 min or when the strain was exposed to increasing concentrations of KCl and sorbitol (data not shown). The transcriptional response of several *A. fumigatus* genes involved in the oxidative stress response (<http://www.aspgd.org>), including *Afyap1* and *Afatf1* (Afu6g09930 and Afu3g11330), which encode TFs required for resistance to oxidative stress (26, 34, 65), *Afccp1* (Afu4g09110), which encodes a cytochrome *c* peroxidase (1, 34,

57), *Afcac1* (Afu3g02270), which encodes a catalase (57), *Afcac2* (Afu8g01670), which encodes a bifunctional catalase-peroxidase (34), *afsod1* (Afu5g09240), which encodes a Cu/Zn superoxide dismutase (30, 32, 34, 42, 62), and *Afsod2* (Afu4g11580), which encodes a manganese-superoxide dismutase (32, 57), was assessed via real-time RT-PCR.

When the wild-type and the complemented strains were exposed to no or 2 mM H₂O₂ for 15 or 30 min, all the investigated genes showed elevated transcription (Fig. 4). The transcriptional response of the Δ *sebA* strain was lower than that of the wild-type and complemented strains (Fig. 4), except for *Afsod1* and *Afsod2*, which had transcription similar to or higher than that seen with the wild-type strain (Fig. 4F to G). The wild-type, the Δ *sebA*::*sebA*⁺, and the Δ *sebA* strains were also exposed to 10 mM paraquat for 15 and 30 min, and gene expression was determined (Fig. 5). Again, all the genes showed increased transcription in the wild-type and the Δ *sebA*::*sebA*⁺ strains when exposed to paraquat (Fig.

FIG 5 The transcriptional basis for *A. fumigatus* $\Delta sebA$ sensitivity to paraquat. The wild-type, $\Delta sebA$, and $\Delta sebA::sebA^+$ strains were grown for 16 h in YG at 37°C and transferred to fresh YG medium with either 0 or 10 mM paraquat and grown for a further 15 or 30 min. The relative quantifications of *Afyap1* (A), *Afatf1* (B), *Afcp1* (C), *Afcac1* (D), *Afcac2* (E), *Afsod1* (F), and *Afsod2* (G) and tubulin gene expression were determined by a standard curve (i.e., C_T values plotted against a logarithm of the DNA copy number). Data shown represent the means ($\pm$ standard deviations) of the results of four biological replicate experiments (*, $P < 0.001$).

5), while the transcriptional response of the $\Delta sebA$ strain, for the majority of genes, was lower than in the wild-type and complemented strains (Fig. 5), except for *Afyap1*, *Afatf1*, and *Afsod1*, which did not show any increase in transcription compared to the control (Fig. 5A, B, and F). Interestingly, in the absence of either H_2O_2 or paraquat, the $\Delta sebA$ strain showed higher transcription than the wild-type and complemented strains for the majority of the genes, except *Afcac1* and *Afsod2* (Fig. 4 and 5).

Collectively, these data suggest that SebA contributes to the transcriptional regulation of several genes related to *A. fumigatus* oxidative stress adaptation.

***A. fumigatus* sebA is important for heat shock recovery.** The fact that SebA was involved in the oxidative stress response and provided greater fitness under poor nutrient conditions suggested that SebA could be involved in other kinds of stress responses. Despite our not seeing any growth defects for the $\Delta sebA$ mutant at 37 and 44°C (Fig. 1), the involvement of *sebA* in heat adaptation was evaluated by exposing the wild-type, the $\Delta sebA::sebA^+$, and

the $\Delta sebA$ strains to heat shock and measuring the *sebA* gene expression (Fig. 6A). Transcription of *sebA* increased in the wild-type strain after exposure to 37°C and 48°C (approximately 6.5- and 4.0-fold after 30 and 60 min and 2.0-fold after 60 and 120 min; Fig. 6A). Accordingly, when all three strains were grown for 5 h at 30°C and then exposed to 37°C for 60 min, there were decreases of 40%, 35%, and 70% in the viability of the wild-type, $\Delta sebA::sebA^+$, and $\Delta sebA$ strains, respectively (Fig. 6B). In addition, when the strains were grown for 5 h at 30°C and exposed to 48°C for 60 min, there were decreases of 60%, 55%, and 80% in the viability of the wild-type, the $\Delta sebA::sebA^+$, and the $\Delta sebA$ strains, respectively (Fig. 6B). Both heat shock treatments induced the SebA::GFP protein to accumulate in the nucleus (Fig. 6C to E). However, SebA::GFP was seen only after 60 min of heat shock (Fig. 6C to E).

To evaluate if SebA is able to control directly or indirectly the transcriptional response of *A. fumigatus* to heat shock, several homologues recognized as being responsible for the heat shock response in *S. cerevisiae* (<http://www.yeastgenome.org>), including

FIG 6 The *A. fumigatus sebA* gene is involved in the heat shock response. The wild-type, *ΔsebA::sebA⁺*, and *ΔsebA* strains were grown for 16 h in YG at 30°C and transferred to fresh YG medium for 15, 30, 60, or 120 min at either 37 or 48°C. (A) Relative quantitations of *sebA* and tubulin gene expression were determined by a standard curve (i.e., C_T values plotted against a logarithm of the DNA copy number). Data shown represent the means ($\pm$ standard deviations) of the results of four biological replicate experiments (*, $P < 0.001$). (B) Conidia (1×10^6 /ml) were incubated in YG for 5 h at 30°C in a shaker (250 rpm). After this period, the germinated conidia were allowed to grow for an additional 60 min at either 37 or 48°C. Germinated conidia were diluted and plated in YAG plates. Plates were incubated at 37°C for 48 h. Viability was determined as the percentage of colonies on treated plates compared to untreated controls (data represent averages of the results of three experiment repetitions $\pm$ standard deviations; *, $P < 0.001$). (C to E) SebA::GFP accumulates in the nuclei upon heat shock stress. The SebA::GFP strain was grown for 16 h at 30°C in YG medium and then incubated 1 h at 30°C (C) or submitted to heat shock (30°C to 37°C and 30°C to 48°C) (D and E). Bars, 5 μm.

the *HSP1* homologue *Afhf1* (Afu5g01900), which encodes a TF that regulates the response to heat shock, the *HSP30* homologue *Afhsp30* (Afu6g06470), the *HSP90* homologue *Afhsp90* (Afu5g04170), and the *SSP1* homologue *Afssp1* (Af8g03930), which encodes a chaperone that is also a member of the *HSP70* protein family, were assessed via real-time RT-PCR. All these genes, except *Af2g09960*, have been demonstrated to have increased transcription during heat shock in *A. fumigatus* (1, 19). In this study, all these genes showed increased expression in the wild-type and *ΔsebA::sebA⁺* strains upon heat shock, for at least a single

time point, during one or both heat shock transitions (Fig. 7; see also Fig. S2 in the supplemental material). In contrast, much more variable behavior was observed for the *ΔsebA* mutant strain. Some of the genes had a smaller increase in expression (such as *Afhf1*; Fig. 7A) or a much greater increase in expression (such as *Afhsp30* and *Afhsp90*; Fig. 7B and C) or even reduced expression (such as *Afssc1* and *Afssp1*; Fig. 7D and E). Taken together, these data suggest that SebA may also participate in the modulation of the transcriptional regulation of several genes involved in the *A. fumigatus* heat shock response.

FIG 7 The transcriptional basis for *A. fumigatus* $\Delta seba$ heat shock response. The wild-type and $\Delta seba$ strains were grown for 16 h in YG at 30°C and transferred to fresh YG medium for 120 min at 30°C or 15, 30, 60, or 120 min at either 37 or 48°C. Relative quantitations of *Afh1* (A), *Afh30* (B), *Afh90* (C), *Afsc1* (D), and *Afsp1* (E) and tubulin gene expression were determined by a standard curve (i.e., C_T values plotted against a logarithm of the DNA copy number). Data shown represent the means ($\pm$ standard deviations) of the results of four biological replicate experiments (*, $P < 0.001$). The results for the $\Delta seba::seba^+$ strain are shown in Fig. S2 in the supplemental material.

The *A. fumigatus* $\Delta seba$ mutant strain is more sensitive to calcium. In an attempt to further identify which biochemical pathways *SebA* is involved in, and/or regulates, the calcineurin/CrzA pathway was evaluated to determine if it is required for the

action of the *SebA* mechanism. First, the $\Delta seba$ strain was shown to be more sensitive to $CaCl_2$ than the corresponding wild-type and $\Delta seba::seba^+$ strains (Fig. 8A). Interestingly, $\Delta seba$ was as sensitive to $CaCl_2$ as the $\Delta crzA$ mutant (Fig. 8B). The transcrip-

FIG 8 The *A. fumigatus* $\Delta sebA$ is more sensitive to calcium. (A and B) Growth phenotypes of the wild-type strain and the $\Delta sebA::sebA^+$ and $\Delta sebA$ mutant strains (A) or the $\Delta crzA::crzA^+$ and $\Delta crzA$ mutant strains (B) grown for 72 h in YAG medium supplemented with 500 mM CaCl₂ at 37°C. (C) The wild-type, $\Delta calA$, and $\Delta crzA$ strains were grown for 16 h in YG at 30°C and transferred to fresh YG medium plus 200 mM CaCl₂ for either 10 or 30 min. Relative quantifications of *sebA* and tubulin gene expression were determined by a standard curve (i.e., C_T values plotted against a logarithm of the DNA copy number). Data shown represent the means ($\pm$ standard deviations) of the results of four biological replicate experiments.

tional dependence of *sebA* on calcium was evaluated by real-time RT-PCR. The expression of *sebA* in the wild-type, $\Delta calA$, and $\Delta crzA$ mutant strains, in response to a short pulse of CaCl₂ (200 mM) for 10 and 30 min (Fig. 8C), was determined. In the wild type, *sebA* expression increased 2-fold following exposure to CaCl₂ (Fig. 8C). The increases were approximately 3- and 8-fold in the $\Delta crzA$ mutant strain and 2- and 3-fold in the $\Delta calA$ mutant strain for CaCl₂ exposures of 10 and 30 min, respectively (Fig. 8C). Collectively, these results suggest that SebA is involved in the calcium metabolism and that it possibly interacts with the calcineurin-CrzA pathway.

The *A. fumigatus* $\Delta sebA$ mutant strain has decreased virulence in a low-dose murine infection. To assess the role of SebA in pathogenicity, the *A. fumigatus* deletion mutant was assessed in a neutropenic murine model for invasive pulmonary aspergillosis, comparing virulence of the *A. fumigatus* $\Delta sebA$ mutant strain ($n = 10$) to the virulence of the wild-type strain ($n = 10$) and the complemented strain ($n = 10$) (Fig. 9A). Wild-type infection resulted in a mortality rate of over 90% at 5 days postinfection, while infection with the $\Delta sebA$ strain resulted in a significantly reduced mortality rate of approximately 50% after 9 to 15 days postinfection ($P < 0.005$). With respect to the kinetics of infection, mice succumbing to infection with the $\Delta sebA$ mutant had a median survival of 7 days, compared to 3 days for mice infected with the wild-type strain. To directly link the observed attenuation of infection with the loss of the *sebA* gene, an independent strain resulting from a single ectopic reintegration of the wild-type *sebA* gene into the $\Delta sebA$ strain was assessed (see Fig. S1 in the supplemental material). This reintegration restored full virulence to the $\Delta sebA$ mutant (Fig. 9A). Histological examinations were performed to further elucidate the basis of the attenuated virulence of the $\Delta sebA$ strain. Infected tissues were collected during early infection with the aim of identifying differences in growth rate, tissue invasion, and the inflammatory response between the two strains. At 72 h postinfection, the lungs of mice infected with the

wild-type isolate contained multiple foci of invasive hyphal growth, representing penetration of the pulmonary epithelium into the major airways (Fig. 9B) and pockets of branched invading mycelia originating from the alveoli (Fig. 9B). In contrast, infection resulting from $\Delta sebA$ inoculations was typified by contained inflammatory infiltrates in bronchioles (Fig. 9B), some of which contained fungal elements in the form of poorly germinated or ungerminated spores. Fungal burden was determined by real-time PCR and showed that the $\Delta sebA$ mutant strain did not grow as well as the wild-type or the complemented $\Delta sebA$ strain (Fig. 9C; $P < 0.0001$). Taken together, these data strongly indicate that SebA plays a role in *A. fumigatus* virulence.

Alveolar macrophages (AM) play an essential role in clearing *A. fumigatus* conidia from the lung (44). Approximately 50% of *A. fumigatus* wild-type, $\Delta sebA$, and $\Delta sebA::sebA^+$ conidia were internalized after 80 min of incubation with AM (Fig. 10A). After a 4 h incubation, *in vitro* killing of resting conidia reached about 40%, 70%, and 40% for the *A. fumigatus* wild-type, $\Delta sebA$, and $\Delta sebA::sebA^+$ strains, respectively (Fig. 10B). These data suggest that there are no differences in the uptake of $\Delta sebA$ conidia by AM, while the $\Delta sebA$ strain is more sensitive to AM killing.

DISCUSSION

The first member of the SebA/Seb1 family in filamentous fungi was isolated from *T. atroviride* and named the *seb1* (stress response element binding) gene because the zinc finger domain specifically recognizes the 5'-AGGGG-3' sequence (43). In *S. cerevisiae*, this motif is recognized and bound by Msn2p/Msn4p (48). The *T. atroviride* *seb1* gene encodes a C2H2 zinc finger protein that is approximately 63% identical to the *S. cerevisiae* proteins Msn2p-Msn4p and Com2p in their zinc finger regions. Msn2p and Msn4p are activated under several different sets of stress conditions, causing their rapid accumulation in the nucleus (6, 8, 10, 13, 18, 22, 24, 28, 36, 47, 50, 60). However, Seb1 may not be the direct Msn2p-Msn4p homologue, since *seb1* overexpression in *S. cerevisiae* was

FIG 9 *A. fumigatus sebA* contributes to virulence in neutropenic mice. (A) Comparative analysis of wild-type, $\Delta sebA$, and $\Delta sebA::sebA^+$ strains in a neutropenic murine model of pulmonary aspergillosis. Mice in groups of 10 per strain were infected intranasally with a 20- μ l suspension of conidiospores at a dose of 5.0×10^4 . (B) Histological analysis of the infected murine lung 72 h after infection with the wild-type strain reveals invasion of the murine lung epithelium. (C) Fungal burden was determined by real-time RT-PCR based on the 18S rRNA gene of *A. fumigatus*. Real-time PCR was used to quantify infection. The measured quantity of the 18S ribosomal DNA (rDNA) in each of the treated samples was normalized using the C_T values obtained for an intronic region of the mouse GAPDH DNA amplifications. Data shown represent the means ($\pm$ standard deviations) of the results determined with lungs from five mice for each treatment.

unable to complement the $\Delta msn2/msn4$ mutation (43). Transcript levels of *seb1* increased under conditions of osmotic stress (sorbitol, NaCl) but not under other stress conditions (cadmium sulfate, pH, membrane disturbance). A *T. atroviride* $\Delta seb1$ strain showed strongly reduced growth on solid medium but grew normally in liquid medium. In liquid medium, growth of the disrupted strain was significantly more inhibited than that of the wild-type strain by the presence of 1 M sorbitol or NaCl (43). Seb1 is involved in, but is not essential for, the osmotic stress response (51). This study identified and characterized a putative *A. fumigatus seb1* homologue, *sebA*. In contrast to the protein encoded by *T. atroviride seb1*, *A. fumigatus* SebA appears to play no role in osmotic stress tolerance. In contrast, SebA was involved in the response to heat shock, oxidative stress, and poor nutrient conditions, such as reduced growth when protein represented the only carbon source and on FBS media. *A. fumigatus* SebA was shown to contribute to the transcriptional modulation and regulation of several genes involved in oxidative stress and heat shock adaptation. However, only two of the genes assessed, *AfuccpA* (Afu4g09119) and *Afucat2* (Afu8g01670), at positions -236 and -588 bp (relative to the ATG start codon), respectively, possessed the stress response element (STRE) motif. These data suggest that SebA activates target genes directly via recognition of a promoter sequence(s) different from that of the STRE motif. It is also possible that SebA interacts with other transcription factors, such as AfHsf1 and AfYap1, modulating their activation. Additional work is necessary to address these hypotheses. However, it is clear from comparisons of the gene functions associated with *A. fumigatus* SebA to those associated with *T. atroviride* Seb1 that the SebA and Seb1 roles are differently wired in different filamentous fungi.

There is a great deal of speculation about which factors are responsible for efficient *A. fumigatus* virulence and pathogenicity (for reviews, see references 15, 37, and 61). Thermotolerance is the most important genetic trait for fungal survival during mammalian and avian infection and within compost (1, 19, 40). Thus, the genetic determinants involved in heat and stress resistance might also contribute to the virulence of this opportunistic pathogen. Few *A. fumigatus* genes that have been shown to be involved in thermotolerance were not linked to virulence (5, 12, 66). Thus, thermotolerance is a polygenic trait, and the most direct approach to identify genes involved in this genetic program is the isolation of transcription factors that activate signal transduction pathways related to thermal adaptation. As expected, most of the analyses have shown that heat shock proteins are very important for the thermotolerance, but in addition, several proteins involved in oxidative stress response, signal transduction, transcription, translation, and carbohydrate or nitrogen metabolism were observed (1, 19, 40). In *S. cerevisiae*, the heat shock response is mostly controlled by Hsf1, Msn2/Msn4, and Hac1, and targets for these transcription factors have already been identified (22, 27, 63–66). Upon heat shock, the *A. fumigatus* Hsf1 homologue has increased transcription (1, 22). Albrecht et al. searched for putative *A. fumigatus* Hsf1 binding signatures and found 17 genes with a potential heat shock element in the promoter region (1). These genes encode not only proteins that are involved directly in the heat shock response (such as chaperones) but also enzymes involved in the oxidative stress response, signal transduction, and carbohydrate and nitrogen metabolism, among others. In *S. cerevisiae*, there is a link between the heat shock response, oxygen respiration, the formation of reactive oxygen intermediates, and the ac-

FIG 10 Killing of $\Delta sebA$ mutant strain is increased by AM phagocytosis. (A) Percentages of *A. fumigatus* conidia taken up by the AM. Data shown represent the average results of three experiment repetitions $\pm$ standard deviations. (B) Percentages of conidia killed by the AM after 4 h of incubation *in vitro*. Data shown represent the average results of three experiment repetitions $\pm$ standard deviations (*, $P < 0.001$).

tivation of the oxidative stress response (56). *A. fumigatus* SebA has a very complex role, contributing to several pathways of stress tolerance and growth under poor nutrient conditions. The *A. fumigatus* SebA seems to be one of the components that integrates responses to different stresses, such as heat shock and oxidative stress, during growth into the mammalian host. Not only does the $\Delta sebA$ mutant have decreased viability under oxidative and heat shock stress conditions, but also several genes that are important for coping with these stresses have heterogeneous transcription compared to those of the wild-type strain. The influence of the Ca^{+2} -calcineurin-CrzA pathway on *sebA* is evidence of this complex regulation. The $\Delta sebA$ mutant is more sensitive to calcium and *sebA* has increased transcription upon calcium exposure in the $\Delta crzA$ mutant strain. It is possible that CrzA controls the SebA levels in the nucleus; this kind of regulation has already been observed for *S. cerevisiae* Msn2/Msn4 (58). It is also possible that there is a connection between *A. fumigatus* SebA and calcineurin under poor nutrient conditions, considering the fact that the *A. fumigatus* $\Delta calA$ mutant cannot grow in FBS (16). However, the increased expression of SebA in the $\Delta calA$ and $\Delta crzA$ mutants does not affect oxidative stress, heat shock stress, and conidial viability in these mutant strains.

Consistently, the multiple defects of the $\Delta sebA$ mutant affected the ability of the pathogen to cope with different stresses and nutritional conditions, which resulted in the attenuated virulence seen in the neutropenic murine model of invasive pulmonary aspergillosis and the decreased viability during AM phagocytosis. The attenuated virulence of the $\Delta sebA$ could be due to a global reduction in its fitness, while the decreased viability after AM phagocytosis could be due to the increased sensitivity of the mutants to reactive oxygen and nitrogen species present in the AM phagolysosomal compartments. However, there are no differences between the mutant and the wild-type strains during their internalization by the AM, which could suggest that the *sebA* null mutation does not have a considerable impact on the chemical composition of the conidia surface. The reduced ability of the $\Delta sebA$ mutant to survive within the mammalian environment may be the result not only of the decreased capacity to deal with stress conditions but also of the incapacity to metabolize nutrients. We have observed a reduction in growth of this mutant in

poor nutrient media, such as FBS and murine lung. Nevertheless, there was no reduction in proteolytic activity in the $\Delta sebA$ mutant strain (data not shown). Thus, the combination of these phenotypic defects associated with fungal growth and response to stressing conditions led to a significant decrease in its ability to establish disease even in an immunocompromised host.

It remains to be demonstrated which signal transduction pathways and genetic programs are being affected by SebA. However, the current work demonstrated that a transcription factor could have a dramatic impact on virulence and survival in the mammalian host. It is tempting to speculate how SebA homologues would affect virulence and pathogenicity in other plant and mammalian fungal pathogens.

ACKNOWLEDGMENTS

This research was supported by the Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP) and Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Brazil.

We thank the anonymous reviewers for their interesting comments and suggestions.

REFERENCES

- Albrecht D, Guthke R, Brakhage AA, Kniemeyer O. 2010. Integrative analysis of the heat shock response in *Aspergillus fumigatus*. BMC Genomics 11:32.
- Bahn Y-S, et al. 2007. Sensing the environment: lessons from fungi. Nat. Rev. Microbiol. 5:57–69.
- Balloy V, Chignard M. 2009. The innate immune response to *Aspergillus fumigatus*. Microbes Infect. 11:919–927.
- Beffa T, et al. 1998. Mycological control and surveillance of biological waste and compost. Med. Mycol. 36(Suppl. 1):137–145.
- Bhambra R, Askew DS. 2005. Thermotolerance and virulence of *Aspergillus fumigatus*: role of the fungal nucleolus. Med. Mycol. 43(Suppl. 1): S87–S93.
- Boy-Marcotte E, et al. 2006. The transcriptional activation region of Msn2p, in *Saccharomyces cerevisiae*, is regulated by stress but is insensitive to the cAMP signalling pathway. Mol. Genet. Genomics 275:277–287.
- Brakhage AA. 2005. Systemic fungal infections caused by *Aspergillus* species: epidemiology, infection process and virulence determinants. Curr. Drug Targets 6:875–886.
- Cameroni E, Hulo N, Roosen J, Winderickx J, De Virgilio C. 2004. The novel yeast PAS kinase Rim 15 orchestrates G0-associated antioxidant defense mechanisms. Cell Cycle 3:462–468.
- Casadevall A, Pirofski LA. 1999. Host-pathogen interactions: redefining

- the basic concepts of virulence and pathogenicity. *Infect. Immun.* 67:3703–3713.
10. Causton HC, et al. 2001. Remodeling of yeast genome expression in response to environmental changes. *Mol. Biol. Cell* 12:323–337.
 11. Chai LY, Vonk AG, Kullberg BJ, Netea MG. 2011. Immune response to *Aspergillus fumigatus* in compromised hosts: from bedside to bench. *Future Microbiol.* 6:73–83.
 12. Chang YC, Tsai HF, Karos M, Kwon-Chung KJ. 2004. THTA, a thermotolerance gene of *Aspergillus fumigatus*. *Fungal Genet. Biol.* 41:888–896.
 13. Choo Y, Klug A. 1994. Selection of DNA binding sites for zinc fingers using rationally randomized DNA reveals coded interactions. *Proc. Natl. Acad. Sci. U. S. A.* 91:11168–11172.
 14. Colot HH, et al. 2006. A high-throughput gene knockout procedure for *Neurospora* reveals functions for multiple transcription factors. *Proc. Natl. Acad. Sci. U. S. A.* 103:10352–10357.
 15. Dagenais TR, Keller NP. 2009. Pathogenesis of *Aspergillus fumigatus* in invasive aspergillosis. *Clin. Microbiol. Rev.* 22:447–465.
 16. da Silva Ferreira M, et al. 2007. Functional characterization of the *Aspergillus fumigatus* calcineurin. *Fungal Genet. Biol.* 44:219–230.
 17. Denning DW. 1998. Invasive aspergillosis. *Clin. Infect. Dis.* 26:781–803.
 18. DeRisi JL, Iyer VR, Brown PO. 1997. Exploring the metabolic and genetic control of gene expression on a genomic scale. *Science* 278:680–686.
 19. Do JH, Yamaguchi R, Miyano S. 2009. Exploring temporal transcription regulation structure of *Aspergillus fumigatus* in heat shock by state space model. *BMC Genomics* 10:306.
 20. Estruch F, Carlson M. 1993. Two homologous zinc finger genes identified by multicopy suppression in a SNF1 protein kinase mutant of *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* 13:3872–3881.
 21. Feng X, et al. 2011. HacA-independent functions of the ER stress sensor IreA synergize with the canonical UPR to influence virulence traits in *Aspergillus fumigatus*. *PLoS Pathog.* 7:e1002330.
 22. Gasch AP, et al. 2000. Genomic expression programs in the response of yeast cells to environmental changes. *Mol. Biol. Cell* 11:4241–4257.
 23. Goldman GH, et al. 2003. Expressed sequence tag analysis of the human pathogen *Paracoccidioides brasiliensis* yeast phase: identification of putative homologues of *Candida albicans* virulence and pathogenicity genes. *Eukaryot. Cell* 2:34–48.
 24. Görner W, et al. 1998. Nuclear localisation of the C2H2 zinc finger protein Msn2p is regulated by stress and protein kinase A activity. *Genes Dev.* 12:586–597.
 25. Greenberger PA. 2002. Allergic bronchopulmonary aspergillosis. *J. Allergy Clin. Immunol.* 110:685–692.
 26. Hagiwara D, Asano Y, Yamashino T, Mizuno T. 2008. Characterization of bZip-type transcription factor AtfA with reference to stress responses of conidia of *Aspergillus nidulans*. *Biosci. Biotechnol. Biochem.* 72:2756–2760.
 27. Hahn J-S, Hu Z, Thiele DJ, Iyer VR. 2004. Genome-wide analysis of the biology of stress responses through heat shock transcription factor. *Mol. Cell. Biol.* 24:5249–5256.
 28. Hasan R, et al. 2002. The control of the yeast H₂O₂ response by the Msn2/4 transcription factors. *Mol. Microbiol.* 45:233–241.
 29. Hohmann S. 2002. Osmotic stress signaling and osmoadaptation in yeasts. *Microbiol. Mol. Biol. Rev.* 66:300–372.
 30. Holdom MD, Lechenne B, Hay RJ, Hamilton AJ, Monod M. 2000. Production and characterization of recombinant *Aspergillus fumigatus* Cu,Zn superoxide dismutase and its recognition by immune human sera. *J. Clin. Microbiol.* 38:558–562.
 31. Käfer E. 1977. Meiotic and mitotic recombination in *Aspergillus* and its chromosomal aberrations. *Adv. Genet.* 19:33–131.
 32. Lambou K, Lamarre C, Beau R, Dufour N, Latge JP. 2010. Functional analysis of the superoxide dismutase family in *Aspergillus fumigatus*. *Mol. Microbiol.* 75:910–923.
 33. Latgé JP. 1999. *Aspergillus fumigatus* and aspergillosis. *Clin. Microbiol. Rev.* 12:310–350.
 34. Lessing F, et al. 2007. The *Aspergillus fumigatus* transcriptional regulator AfYap1 represents the major regulator for defense against reactive oxygen intermediates but is dispensable for pathogenicity in an intranasal mouse infection model. *Eukaryot. Cell* 6:2290–2302.
 35. Liu H, et al. 2010. *Aspergillus fumigatus* AcuM regulates both iron acquisition and gluconeogenesis. *Mol. Microbiol.* 78:1038–1054.
 36. Martínez-Pastor MT, et al. 1996. The *Saccharomyces cerevisiae* zinc finger proteins Msn2p and Msn4p are required for transcriptional induction through the stress-response element. *EMBO J.* 15:2227–2235.
 37. McCormick A, Loeffler J, Ebel F. 2010. *Aspergillus fumigatus*: contours of an opportunistic human pathogen. *Cell Microbiol.* 12:1535–1543.
 38. Mech F, Thywissen FA, Guthke R, Brakhage AA, Figge MT. 2011. Automated image analysis of the host-pathogen interaction between phagocytes and *Aspergillus fumigatus*. *PLoS One* 6:e19591.
 39. Mota Júnior AO, et al. 2008. Molecular characterization of the *Aspergillus fumigatus* NCS-1 homologue, NcsA. *Mol. Genet. Genomics* 280:483–495.
 40. Nierman W, et al. 2005. Genomic sequence of the pathogenic and allergenic filamentous fungus *Aspergillus fumigatus*. *Nature* 438:1151–1156.
 41. Noventa-Jordão MA, do Nascimento AM, Goldman MH, Terenzi HF, Goldman GH. 2000. Molecular characterization of ubiquitin genes from *Aspergillus nidulans*: mRNA expression on different stress and growth conditions. *Biochim. Biophys. Acta* 1490:237–244.
 42. Oberegger H, Zadra I, Schoeser M, Haas H. 2000. Iron starvation leads to increased expression of Cu/Zn-superoxide dismutase in *Aspergillus*. *FEBS Lett.* 485:113–116.
 43. Peterbauer CK, Litscher D, Kubicek CP. 2002. The *Trichoderma atroviride* seb1 (stress response element binding) gene encodes an AGGGG binding protein which is involved in the response to high osmolarity stress. *Mol. Genet. Genomics* 268:223–231.
 44. Philippe B, et al. 2003. Killing of *Aspergillus fumigatus* by alveolar macrophages is mediated by reactive oxidant intermediates. *Infect. Immun.* 71:3034–3042.
 45. Powers-Fletcher MV, et al. 2011. Impact of the lectin chaperone calnexin on the stress response, virulence and proteolytic secretome of the fungal pathogen *Aspergillus fumigatus*. *PLoS One* 12:e28865.
 46. Ramsdale M, et al. 2008. MNL1 regulates weak acid-induced stress responses of the fungal pathogen *Candida albicans*. *Mol. Biol. Cell* 19:4393–4403.
 47. Rep M, Krantz M, Thevelein JM, Hohmann S. 2000. The transcriptional response of *Saccharomyces cerevisiae* to osmotic shock. Hot1p and Msn2p/Msn4p are required for the induction of subsets of high osmolarity glycerol pathway-dependent genes. *J. Biol. Chem.* 275:8290–8300.
 48. Ruis H, Schüller C. 1995. Stress signalling in yeast. *Bioessays* 17:959–965.
 49. Schiestl RH, Gietz RD. 1989. High efficiency transformation of intact yeast cells using single stranded nucleic acids as a carrier. *Curr. Genet.* 16:339–346.
 50. Schüller C, et al. 2004. Global phenotypic analysis and transcriptional profiling defines the weak acid stress response regulon in *Saccharomyces cerevisiae*. *Mol. Biol. Cell* 15:706–720.
 51. Seidl V, Seiboth B, Karaffa L, Kubicek CP. 2004. The fungal STRE-element-binding protein Seb1 is involved but not essential for glycerol dehydrogenase (gld1) gene expression and glycerol accumulation in *Trichoderma atroviride* during osmotic stress. *Fungal Genet. Biol.* 41:1132–1140.
 52. Semighini CP, Marins M, Goldman MHS, Goldman GH. 2002. Quantitative analysis of the relative transcript levels of ABC transporter *Atr* genes in *Aspergillus nidulans* by real-time reverse transcription-PCR assay. *Appl. Environ. Microbiol.* 68:1351–1357.
 53. Shapiro RS, Robbins N, Cowen LE. 2011. Regulatory circuitry governing fungal development, drug resistance, and disease. *Microbiol. Mol. Biol. Rev.* 75:213–267.
 54. Soriani FM, et al. 2010. Identification of possible targets of the *Aspergillus fumigatus* CRZ1 homologue, CrzA. *BMC Microbiol.* 10:12.
 55. Soriani FM, et al. 2008. Functional characterization of the *Aspergillus fumigatus* CRZ1 homologue, CrzA. *Mol. Microbiol.* 67:1274–1291.
 56. Sugiyama K, Izawa S, Inoue Y. 2000. The Yap1p-dependent induction of glutathione synthesis in heat shock response of *Saccharomyces cerevisiae*. *J. Biol. Chem.* 275:15535–15540.
 57. Sugui JA, et al. 2008. Genes differentially expressed in conidia and hyphae of *Aspergillus fumigatus* upon exposure to human neutrophils. *PLoS One* 3:e2655.
 58. Takatsume Y, et al. 2010. Calcineurin/Crz1 destabilizes Msn2 and Msn4 in the nucleus in response to Ca(2+) in *Saccharomyces cerevisiae*. *Biochem. J.* 427:275–287.
 59. Teepe AG, Loprete DM, He Z, Hoggard TA, Hill TW. 2007. The protein kinase C orthologue PkcA plays a role in cell wall integrity and polarized growth in *Aspergillus nidulans*. *Fungal Genet. Biol.* 44:554–562.
 60. Teixeira MC, et al. 2006. The YEASTRACT database: a tool for the analysis of transcription regulatory associations in *Saccharomyces cerevisiae*. *Nucleic Acids Res.* 34:D446–D451.

61. Tekaia F, Latgé JP. 2005. *Aspergillus fumigatus*: saprophyte or pathogen? Curr. Opin. Microbiol. 8:385–392.
62. Teutschbein J, et al. 2010. Proteome profiling and functional classification of intracellular proteins from conidia of the human-pathogenic mold *Aspergillus fumigatus*. J. Proteome Res. 9:3427–3442.
63. Wu WS, Li WH. 2008. Identifying gene regulatory modules of heat shock response in yeast. BMC Genomics 9:439.
64. Yamamoto A, Mizukami Y, Sakurai H. 2005. Identification of a novel class of target genes and a novel type of binding sequence of heat shock transcription factor in *Saccharomyces cerevisiae*. J. Biol. Chem. 280:11911–11919.
65. Yamamoto N, Maeda Y, Ikeda A, Sakurai H. 2008. Regulation of thermotolerance by stress-induced transcription factors in *Saccharomyces cerevisiae*. Eukaryot. Cell 7:783–790.
66. Zhou H, et al. 2007. O-Mannosyltransferase 1 in *Aspergillus fumigatus* (AfPmt1p) is crucial for cell wall integrity and conidium morphology, especially at an elevated temperature. Eukaryot. Cell 6:2260–2268.

7.6. Anexo O. Artigo publicado: Sugarwin: A sugarcane insect-induced gene with antipathogenic activity. Mol Plant Microbe Interact. Medeiros A. H.; Franco F. P.; Matos J. L.; de Castro P. A.; Santos-Silva L. K.; Henrique-Silva F.; Goldman G. H.; Moura D. S.; Silva-Filho M. C. Mol Plant Microbe Interact., May. 2012, 25(5): 613-24.

Sugarwin: A Sugarcane Insect-Induced Gene with Antipathogenic Activity

Ane H. Medeiros,¹ Flávia P. Franco,¹ Juliana L. Matos,¹ Patrícia A. de Castro,² Ludier K. Santos-Silva,³ Flávio Henrique-Silva,³ Gustavo H. Goldman,^{2,4} Daniel S. Moura,⁵ and Marcio C. Silva-Filho¹

¹Departamento de Genética, Escola Superior de Agricultura Luiz de Queiroz, Universidade de São Paulo, C.P. 83, 13418-900 Piracicaba, SP, Brazil; ²Departamento de Ciências Farmacêuticas, Faculdade de Ciências Farmacêuticas, Universidade de São Paulo, Ribeirão Preto, SP, Brazil; ³Departamento de Genética e Evolução, Universidade Federal de São Carlos, São Carlos, SP, Brazil; ⁴Laboratório Nacional de Ciência e Tecnologia do Bioetanol (CTBE), Campinas 13083-970, Brazil; ⁵Departamento de Ciências Biológicas, Escola Superior de Agricultura Luiz de Queiroz, Universidade de São Paulo, C.P. 9, 13418-900 Piracicaba, SP, Brazil

Submitted 29 September 2011. Accepted 11 January 2012.

In sugarcane fields, colonization of the stalk by opportunistic fungi usually occurs after the caterpillar *Diatraea saccharalis* attacks the sugarcane plant. Plants respond to insect attack by inducing and accumulating a large set of defense proteins. Two homologues of a barley wound-inducible protein (BARWIN), sugarcane wound-inducible proteins SUGARWIN1 and SUGARWIN2, have been identified in sugarcane by an in silico analysis. Antifungal properties have been described for a number of BARWIN homologues. We report that a SUGARWIN::green fluorescent protein fusion protein is located in the endoplasmic reticulum and in the extracellular space of sugarcane plants. The induction of *sugarwin* transcripts occurs in response to mechanical wounding, *D. saccharalis* damage, and methyl jasmonate treatment. The accumulation of transcripts is late induced and is restricted to the site of the wound. Although the transcripts of *sugarwin* genes were strongly increased following insect attack, the protein itself did not show any effect on insect development; rather, it altered fungal morphology, leading to the apoptosis of the germlings. These results suggest that, in the course of evolution, *sugarwin*-encoding genes were recruited by sugarcane due to their antipathogenic activity. We rationalize that sugarcane is able to induce *sugarwin* gene expression in response to *D. saccharalis* feeding as a concerted plant response to the anticipated invasion by the fungi that typically penetrate the plant stalk after insect damage.

Plant responses to herbivores and pathogens are complex and modulate the expression of a large number of genes, many of which are believed to play roles in defense (Agrawal et al. 2003). To effectively combat invasion by microbial pathogens and herbivorous insects, plants have evolved sophisticated defensive strategies to perceive the attack by pathogens and insects and to translate this perception into an appropriate defensive response (Dangl and Jones 2001).

Current address for A. H. Medeiros: Universidade Federal de São Carlos, campus Araras, Piracicaba, SP, Brazil.

Corresponding author: M. C. Silva-Filho; Telephone: +55-19 34294442; Fax: +55-19 34336706; E-mail: mcsilva@usp.br

*The e-Xtra logo stands for “electronic extra” and indicates that two supplementary tables and one supplementary figure are published online.

Both herbivore feeding and mechanical damage induce systemic responses that are rapidly propagated throughout the plant and responses that are restricted to the wound site (Kessler and Baldwin 2002). Many plants respond to herbivore attacks by activating defense genes in leaves whose products inhibit the digestive proteases of herbivores and reduce the nutritional quality of the ingested proteins, making the attackers ill (Ryan 1990). The study of the wound response provoked by herbivores began with the seminal work on inducible proteinase inhibitors of tomato plants (Ryan 1968). Since then, several genes that encode wound-inducible proteins (WIN) have been identified in various plant species (Christopher et al. 2004; Ryan 1990; Schilmiller and Howe 2005).

During a search for sugarcane homologues of plant defense genes in the sugarcane expressed-sequence tag (EST) database SUCEST (Vettore et al. 2001), Falco and associates (2001) identified two sugarcane homologues of a protein named barley wound-inducible (BARWIN). BARWIN is a wound- and pathogen-inducible protein that can be isolated from barley seed and leaves (Hejgaard et al. 1992; Svensson et al. 1992). Although the BARWIN domain can be found in association with a chitin-binding domain typical of lectins (Broekaert et al. 1990), a number of proteins have been isolated from plants that show only the BARWIN domain: tobacco PR4 (Friedrich et al. 1991), tomato P2 (Linthorst et al. 1991), WHEATWINs (Caruso et al. 1999), WjAMP-1 from *Wasabia japonica* (Kiba et al. 2003), ZmPR4 from maize (Bravo et al. 2003), OsPR4 (Agrawal et al. 2003) and OsPR4b (Zhu et al. 2006) from rice, and LrPR4 from *Lycoris radiata* (Li et al. 2010). The *barwin* homologues of several plant species have been associated with the plant responses to fungal infection, such as *Erysiphe graminis* f. sp. *hordei* in barley (Hejgaard et al. 1992), *F. culmorum* in wheat (Caruso et al. 1999), *Magnaporthe grisea* in rice (Agrawal et al. 2003) and *L. radiata* (Li et al. 2010), *F. moniliforme* (presently renamed *F. verticillioides*) in maize (Bravo et al. 2003), and *Cladosporium fulvum* in tobacco (Linthorst et al. 1991). Antifungal activity has been described for BARWIN-like proteins from barley (Hejgaard et al. 1992) and wasabi (Kiba et al. 2003) (which showed activity against both fungi and bacteria) and for recombinant BARWIN-like proteins from wheat (Caruso et al. 2001), maize (Bravo et al. 2003), and rice (Zhu et al. 2006).

In the tropics, sugarcane is an important cash crop cultivated for its stalks, which accumulate high levels of sucrose. This plant contributes to 60% of the raw sugar produced worldwide,

with the remaining 40% coming from the sugar beet. In some countries, a significant part of the sugarcane biomass is used for the production of bioethanol (Grivet and Arruda 2002). In Brazil, the sugarcane borer *Diatraea saccharalis* (F.) (Lepidoptera: Pyralidae) is the major pest of sugarcane and is responsible for both direct and indirect damage. The indirect damage is caused by microorganisms that gain access to the plant through the openings created by *D. saccharalis* (Ogunwolu et al. 1991). Phytopathogens, such as *Colletotrichum falcatum* (Went) and *F. verticillioides* (Sacc.) Nirenberg, are widespread in areas of sugarcane borer infestation; together with insects, these fungi form the borer-rot complex.

We report the molecular cloning, subcellular localization, and expression analysis of two sugarcane homologues of *barwin*, designated herein as sugarcane wound-inducible genes *sugarwin1* and *sugarwin2*. *Sugarwin*-green fluorescent protein (SUGARWIN::GFP) fusions transiently expressed in onion epidermal cells were detected in the endoplasmic reticulum, transvacuolar strands, and apoplast, thus indicating that SUGARWIN are secreted proteins. Gene expression analyses showed that the induction of *sugarwin*-encoding genes occurs in response to wounding, attack by *D. saccharalis*, and the application of methyl jasmonate (MJ). The accumulation of *sugarwin* transcripts is late induced and is restricted to the site of the wound or attack. A recombinant SUGARWIN protein added to the diet of *D. saccharalis* showed no effect on caterpillar development, whereas the same protein incorporated into the growth medium of *F. verticillioides* altered the fungal morphology and caused cell apoptosis. Although our data do not rule out the possibility of *sugarwin* genes being part of a non-specific defensive compound storm, we suggest that *sugarwin* genes may be part of a concerted strategy of plant defense against opportunistic pathogens.

RESULTS

Isolation of cDNAs encoding two new BARWIN domain-containing proteins.

Two cDNAs from the sugarcane EST database (Vettore et al. 2001, 2003) were identified: *sugarwin1* (GenBank accession CA145787) and *sugarwin2* (GenBank accession CA138924). The deduced proteins from the isolated cDNAs contain at least two regions: a signal peptide of 32 and 27 amino acids for SUGARWIN1 and SUGARWIN2, respectively, and the BARWIN domain containing 120 and 121 amino acids, respectively. Further analysis revealed that the probable N-terminal signal peptide cleavage site is located between amino acid Q (32 for SUGARWIN1 and 27 for SUGARWIN2) and Q (Bendtsen et al. 2004). Multiple amino acid sequence alignments of the BARWIN domain from the deduced SUGARWIN1 and SUGARWIN2 proteins revealed a high level of similarities to other HEVEIN-like and BARWIN-like proteins from mono- and dicotyledonous plants (Fig. 1). The structures and domains of these proteins were scrutinized with InterProScan (Quevillon et al. 2005) and revealed, only in the dicotyledonous species, a C-terminal chitin-binding domain, located after the signal peptide, which was followed by an N-terminal BARWIN domain. As with other monocotyledonous species, the SUGARWIN isoforms do not contain the chitin-binding domain, and the proteins fall into a different phylogenetic group (Fig. 2). SUGARWIN1 and SUGARWIN2 showed 73.9% identity and 80.7% similarity, whereas the level of SUGARWIN1 similarity to the BARWIN domain from monocot and dicot plants ranged from 80.7 to 93.3%, with identity values as high as 88.2%. The identity of SUGARWIN2 to other BARWIN-like proteins was approximately 75%, with similarity values as high as 81%.

SUGARWIN proteins are secreted through the default secretory pathway.

The amino-terminal regions of SUGARWIN1 and SUGARWIN2 were highly hydrophobic and displayed features similar to signal peptides for translocation to the endoplasmic reticulum (ER). Plasmolized onion epidermal cells transiently expressing SUGARWIN1::GFP and SUGARWIN2::GFP fusion proteins are shown in Figure 3A to D. These cells showed green fluorescence in the ER, transvacuolar strands (Fig. 3C), and in apoplast. These results indicate that the chimeric proteins travel from the ER to the cell exterior, following the default secretory pathway. GFP5-ER (mgfp5-ER) (Haseloff et al. 1997) was used as a control for GFP signal retention in the ER (Fig. 3E and F).

Wounding, herbivory, and MJ treatments differentially induce *sugarwin* genes.

For the monitoring of the gene expression response after wounding, the plant tissue was separated into two parts that were analyzed separately: the "wounded region" refers to the 2.5-cm region where the damage occurred, and the "unwounded region" represents the upper part of the plant, above the damaged region.

Sugarwin1 and *sugarwin2* transcripts began to accumulate 6 h following *D. saccharalis* attack and continued to accumulate until the last observation, 24 h later. Transcripts of the two *sugarwin* isoforms were highly abundant at the wounded region (Fig. 4A and C) and were expressed at low levels at the unwounded region (Fig. 4B and D). For example, *sugarwin1* was 5-fold more highly expressed at the wound site, whereas *sugarwin2*, after 6 h of feeding, was 53 times more highly expressed at the wound site than in the unwounded plant parts and, after 12 h of feeding, this transcript was 192-fold more highly expressed at the wound site. The transcripts of *sugarwin2* were much more abundant than those of *sugarwin1*. At 24 h after challenge by *D. saccharalis*, the transcript accumulation of both *sugarwin1* and *sugarwin2* increased 15- and 300-fold at the site of herbivore damage, compared with the control plants at the beginning of the experiment. In the undamaged tissue region, the *sugarwin1* and *sugarwin2* transcripts presented a threefold increase (Fig. 4B and D). The transcript accumulation reached its peak for both *sugarwin1* and *sugarwin2* after 12 h of continuous mechanical wounding at the wounded region, while no transcripts accumulated at the unwounded region.

The transcript levels of the sugarcane homologue of *Bowman-Birk proteinase inhibitor* (BBPI), which was used as a positive control for the mechanical wound and insect treatments, were highly induced by *D. saccharalis* feeding. After 6 h of insect damage at the wounded region, BBPI transcript accumulation was as high as 84-fold and continued to increase after 24 h of damage, reaching levels as high as 325-fold induction. In the unwounded plant parts, BBPI transcript induction was even higher than in the wounded plant parts. In contrast, mechanical wounding resulted in higher BBPI transcript accumulation at the site of the wound (Fig. 4E) than at the unwounded parts of the plant (Fig. 4F), and the levels of induction were much lower than those observed after insect wounding. At the site of mechanical wounding, the highest levels of mRNA accumulation were found 12 h after wounding was initiated and subsequently decreased.

The effects of MJ and methyl salicylate (MS) were evaluated by spraying both methyl esters onto sugarcane plants. Only MJ induced the transcription of *sugarwin*- and BBPI-encoding genes (Fig. 5A, C, and E; note the differences in scale). MS did not show any effect on *sugarwin*-encoding genes or BBPI (Fig. 5B, D, and F).

Fig. 1. Amino acid sequence alignment of chitin-binding and barley wound-inducible protein (BARWIN) domain homologues on mono- and dicotyledonous plants: *Arabidopsis* (GenBank accession P43082), rubber tree (AAO63572), grape (AAC33732), tobacco 1 (AAB29959), tobacco 2 (CAA41437), potato (P09762), sugarcane wound-inducible protein (SUGARWIN)2 (CA138924), wheat (CAA06856), barley (P28814), rice (AAL11444), maize (NP001105464), and SUGARWIN1 (CA145787). The alignment was performed on Muscle (Edgar 2004). The bar graph indicates the residues conservation similarity. The background residue coloring were based on RASMOL colors.

SUGARWIN protein does not affect *D. saccharalis* development.

To test the effect of SUGARWIN proteins on *D. saccharalis* development, both isoforms of SUGARWIN were produced in *Pichia pastoris*. Recombinant SUGARWIN1 and SUGARWIN2 proteins with the addition of a histidine tag have expected molecular masses of 14.2 and 16.4 kDa, respectively. In total, 151.2 mg and 1.18 g of recombinant protein per liter of culture was obtained for SUGARWIN1 and SUGARWIN2, respectively. The SUGARWIN proteins were added to the *D. saccharalis* diet at a final concentration of 420 µg/ml (30 and 26.3 µM for SUGARWIN1 and SUGARWIN2, respectively), and the development of neonate larvae was evaluated after 14 days. No significant difference was found between the control diet (phosphate-buffered saline [PBS]) and the diet with SUGARWIN proteins. By comparison, the positive control, in which soybean proteinase inhibitor was incorporated into the caterpillars' diet, resulted in larvae that weighed as little as 0.0012 g (Table 1). Larval survival was 95% on the control diet (PBS) whereas, on the SUGARWIN1 and SUGARWIN2 diets, it was 100%. In the diet containing soybean proteinase inhibitor, only 30% larval survival was observed (Table 1).

The effect of recombinant SUGARWIN2 on the mycelial morphology and cell death of *F. verticillioides*.

To test the hypothesis that SUGARWIN proteins are able to affect fungal growth, we exposed *F. verticillioides* 8-h-old germlings to increasing concentrations of SUGARWIN2 (20, 40, 80, and 160 µM) for 16 h at 25°C. The SUGARWIN2 protein was used in our experiments because its gene expression in sugarcane plants is nearly 30-fold higher than that of SUGARWIN1. The viability test shows that SUGARWIN2 treatment at concentrations of 80 and 160 µM causes germling cell death (Fig. 6A). To further investigate the mechanism causing cell death, the terminal deoxynucleotidyl transferase dUTP nick-mediated end labeling (TUNEL) assay was used to determine whether SUGARWIN2 induces DNA fragmentation. This assay uses terminal deoxynucleotidyltransferase to label 3'-OH DNA termini with FITC-conjugated dUTP, which can be directly visualized by fluorescence microscopy. After

16 h of treatment with 160 µM SUGARWIN2, the majority of the hyphae (approximately 80%) showed TUNEL-positive staining (Fig. 6B, lower panels), whereas the untreated control hyphae showed no staining (Fig. 6B, upper panels). To confirm the effect of SUGARWIN2, differential interference contrast (DIC) and fluorescence microscopy with Calcofluor White analyses were performed. Again, *F. verticillioides* 8-h-old germlings were exposed to 160 µM SUGARWIN2 for 2 to 16 h. The control germlings exposed to PBS showed normal development and clear fluorescent staining of the septa. The germlings exposed to SUGARWIN2 showed dramatic morphological changes (Fig. 7). Approximately 2, 5, 30, 40, and 80% of these germlings showed abnormalities (such as increased vacuolization, multiple points of fractures in the hyphae, and extensive leaking of intracellular material) (Fig. 7, arrows) when exposed to 160 µM SUGARWIN2 for 2, 4, 6, 8, and 16 h, respectively. Taken together, these results strongly suggest that SUGARWIN2 is affecting the morphogenesis and viability of *F. verticillioides*, leading to cell death.

DISCUSSION

Plants are constantly faced with the risk of being attacked by herbivorous insects and pathogenic fungi. Because both insects and fungi are ubiquitous in terrestrial systems, they may encounter each other on the same individual host or may exploit a host that has been altered by previous attack (Denooij et al. 1992). Plant-inducible defenses involve a broad range of proteins and other molecules, the synthesis of which is spatially and temporally controlled (Karban and Baldwin 1997; Walling 2000; Wu and Baldwin 2010). These defense mechanisms appear to be tightly regulated, allowing economy in times of peace and presenting a moving defense horizon to an attacker (Reymond et al. 2004).

In this work, two homologues of the wound- and pathogen-inducible *barwin* gene have been identified in sugarcane. SUGARWIN1 and SUGARWIN2 showed high similarity to the BARWIN domains of monocot and dicot plants (Fig. 1). The BARWIN association with the chitin-binding domain exhibits an intricate and recurrent pattern of evolution. The dicotyledonous plants can be divided into two phylogenetic

Fig. 2. Maximum likelihood tree of chitin-binding and barley wound-inducible protein (BARWIN) domain sequences. The tree was inferred using the amino acid maximum likelihood feature of PhyML (Guindon and Gascuel 2003). Clades support was given by the approximate likelihood ratio test (aLRT) based on the nonparametric Shimodaira-Hasegawa-like (SH-like) procedure (Anisimova and Gascuel 2006). Scale at bottom of figure indicates the branch length associated with 0.08 substitutions per position.

Fig. 3. Confocal laser scanning micrographs of proteins fused to green fluorescent protein (GFP) and transiently expressed in onion epidermal cells. **A** and **B**, sugarcane wound-inducible protein 1 (SUGARWIN1); **C** and **D**, SUGARWIN2; **E** and **F**, the synthetic construct modified GFP protein (mgfp5-ER). Fluorescence and differential interference contrast microscopy were performed using a Zeiss Axiovert 100 M (Carl Zeiss, Thornwood, NY, U.S.A.).

groups: those that have the hevein domain associated with the BARWIN domain and those that solely have the BARWIN domain (Agrawal et al. 2003; Bravo et al. 2003; Zhu et al. 2006). The monocot homologues of BARWIN, including sugarcane, do not have the BARWIN domain associated with the hevein domain (Fig. 2).

Primary structure analyses of the deduced proteins from both sugarcane BARWIN homologues indicated that they are synthesized as preproteins, with an N-terminal signal peptide that is involved in extracellular secretion. Indeed, SUGARWIN signal peptides fused to GFP and transiently expressed in onion epidermal cells showed that both proteins were targeted to the extracellular space through the secretory pathway (Fig. 3A to D). The predicted signal peptide cleavage site, between the first alanine and glutamine residues, and the signal peptide size (32 and 27 amino acids), shows conservation among other BARWIN-like proteins (Agrawal et al. 2003; Caruso et al. 2001; Friedrich et al. 1991; Li et al. 2010; Linthorst et al. 1991; Zhu et al. 2006). The subcellular localization of the SUGARWIN proteins substantiates their antipathogenic function. Phytopathogenic fungi and bacteria are slow-growing organisms that must colonize the extracellular space before penetrating directly or indirectly into the cell lumen. The presence of an antipathogenic protein in the extracellular space would impair microorganism growth and colonization. Thus, this plant strategy appears to be directed toward microorganisms.

The transcript levels of the *sugarwin* genes accumulated in response to herbivory by *D. saccharalis* (Fig. 4). To our knowledge, this is the first report of the induction of a *barwin*-like gene by herbivory. The *sugarwin* genes show a pattern of expression typical of tomato late-wound-induced genes (Ryan

2000). Expression studies have shown a late-inducible pattern of expression for other BARWIN-like proteins: WjAMP mRNAs are significantly increased 24 and 48 h after inoculation with either *Botrytis cinerea* or *Alternaria alternata* (Kiba et al. 2003). Additionally, wounding and inoculation with *F. moniliforme* also induces the expression of ZmPR4 after 24 h (Bravo et al. 2003). Other monocot plants, such as wheat (Bertini et al. 2006), rice (Agrawal et al. 2003), and barley (Hejgaard et al. 1992), also show induced expression of BARWIN-like proteins after inoculation with distinct pathogenic fungi.

It has been demonstrated that the induction of signaling defenses by one type of intruder may also provide protection (or increased susceptibility) against the attack of another organism. For instance, in *Arabidopsis*, the spectrum of the herbivore *Pieris rapae*-induced resistance has been investigated against different pathogens (De Vos et al. 2006); feeding by the herbivore has been shown to significantly reduce the disease caused by the bacterial pathogens *Pseudomonas syringae* pv. *tomato* and *Xanthomonas campestris* pv. *armoraciae* in caterpillar-damaged tissue, although it was not effective against the necrotrophic fungus *A. brassicicola*. Another study used different *Arabidopsis* mutants with impaired jasmonic acid (JA)- and salicylic acid (SA)-dependent defense responses to investigate the impact of pathogen-induced defense responses on herbivore performance (Cui et al. 2002). In the tomato, wounding by *Helicoverpa zea* has been shown to induce resistance to the bacterium *P. syringae* pv. *tomato* (Stout et al. 1999).

Transcripts of *sugarwin1* and *sugarwin2* were induced at the site of mechanical or insect wounding, whereas they were expressed at very low levels in systemic tissues (Fig. 4). The expression of *sugarwin* also diminished when the mechanical

Fig. 4. Expression levels of gene sugarcane wound-inducible 1 (*sugarwin1*) in the A, wounded and B, unwounded regions of sugarcane; *sugarwin2* in the C, wounded and D, unwounded regions; and the Bowman-Birk proteinase inhibitor in the E, wounded and F, unwounded regions. Expression was quantified by real-time quantitative polymerase chain reaction in response to *Diatraea saccharalis* feeding (black bars) and to mechanical wounding (gray bars) with respect to control expression (white bars). The x-axis indicates the time of analysis (hours after treatment) and the y-axis indicates the fold change in gene expression. Values are the means ($\pm$ standard error) of the transcripts from three replicates, normalized to the transcript abundance of glyceraldehyde 3-phosphate dehydrogenase. Regulation of expression was calculated with the REST 2008 software (Pfaffl et al. 2002). Asterisks above the bars represent significant differences in comparison with the control at 0 h.

wounding was halted (after 12 h), indicating that the level of expression is dependent on wound infliction. When the plants were exposed to herbivore attack, the transcript levels accumulated over time, likely due to the constant wounding inflicted on the plant. As a positive control for the systemic response, we analyzed the gene expression of the local and systemic inducible inhibitor BBPI. Herbivore attack induced the accumulation of *BBPI* transcripts at the highest levels in systemic unwounded tissues (Fig. 4E and F). It has been shown for a number of plant families that wounding in the leaves results in the synthesis of defense proteins both at the site of the wound and in the distal systemic leaves (Green and Ryan 1972; Moura and Ryan 2001; Wu and Baldwin 2010). In this study, the high expression level of *BBPI* observed in the unwounded plant tissues could result in increased protection against chewing insects, because their rate of damage and tissue destruction is very swift. Thus, *sugarwin1* and *sugarwin2* transcript accumulation exclusively at the wound site could be related to their roles in plant defense. *ZmPR4*, through in situ RNA hybridization experiments, has been shown to have mRNA accumulation in the same tissues of fungal-infected germinating embryos (Bravo et al. 2003). The authors suggested that the expression of the *ZmPR4* gene could be part of the defense response of germinating maize seed against fungal pathogens. This observation could also be extended to the *sugarwin* genes.

It has often been observed that, in cases of the sugarcane epiphytotic known as red rot, cultures of microorganisms isolated from the stalk show a blend of organisms that cause rot and wilt (Abbott and Hughes 1961; Khanna and Rafay 1953; Singh et al. 1977). The potential sources of inoculum include the seed cane, or sett, because sugarcane is a vegetatively propagated crop; the soil; or contaminated agricultural implements (Upadhyay et al. 2009). These fungi enter the sugarcane stalk

through the root primordia, leaf scars, and bud scales; through wounds caused by cultural operations; and through injuries caused by the sugarcane borer (McKaig 1936; Mian et al. 1969; Upadhyay et al. 2009). Indeed, in sugarcane, attacks by the sugarcane borer, *D. saccharalis*, are often followed by fungi that cause rotting, such as *C. falcatum* and *F. verticillioides* (McKaig 1936; Mian et al. 1969; Ogunwolu et al. 1991). Other examples of insect–fungi association can be found between *Fusarium* spp. from sugarcane and the stalk borer, *Eldana saccharina* (Lepidoptera: Pyralidae) (McFarlane et al. 2009) or between *F. verticillioides* and the European corn-borer *Ostrinia nubilalis* (Gatch and Munkvold 2002).

The wounds created by insects permit fungi to invade plants and establish infection sites within the more vulnerable internal tissue (McFarlane et al. 2009). In our experiments, induction of *sugarwin* genes in unwounded tissue was marginal; therefore, we hypothesize that, if SUGARWIN has antipathogenic activity, it is conceivable that increased levels of *sugarwin* transcripts at the site of wounds and, therefore, of SUG-

Table 1. Effect of sugarcane wound-inducible proteins SUGARWIN1 and SUGARWIN2 on larval survival and larval weight of *Diatraea saccharalis* reared for 14 days on corn flour-based diet

Treatment	Larval survival (%)	Larvae weight (g) ^y
Control diet (PBS) ^z	95	0.016 ± 0.006 a
Soybean proteinase inhibitor	30	0.001 ± 0.001 b
SUGARWIN1	100	0.015 ± 0.006 a
SUGARWIN2	100	0.014 ± 0.006 a

^y Mean values followed by the same letter on the column are not significantly different, according to Tukey test at 5%.

^z PBS = phosphate-buffered saline.

Fig. 5. Expression levels of gene sugarcane wound-inducible 1 (*sugarwin1*) after A, methyl jasmonate (MJ) and B, methyl salicylate (MS) exposure; *sugarwin2* after C, MJ and D, MS exposure; and the Bowman-Birk proteinase inhibitor after E, MJ and F, MS exposure. Expression levels were quantified by real-time quantitative polymerase chain reaction. The x-axis indicates the time of analysis (hours after treatment) and the y-axis indicates the fold change in gene expression. Values are the means (± standard error) of the transcripts from three replicates, normalized to the transcript abundance of glyceraldehyde 3-phosphate dehydrogenase. The regulation of expression was calculated with the REST 2008 software (Pfaffl et al. 2002). Asterisks above the bars represent significant differences in comparison with the control at 0 h.

Fig. 6. Effects of sugarcane wound-inducible protein 2 (SUGARWIN2) on cell death in *Fusarium verticillioides*. **A**, Viability test of *F. verticillioides* germlings. *F. verticillioides* germlings were treated with different concentrations of SUGARWIN2 (20, 40, 80, and 160 μ M) for 16 h. Later, the samples were transferred to new plates containing a solid potato dextrose medium and incubated at 25°C for another 36 h. **B**, The terminal deoxynucleotidyl transferase dUTP nick-mediated end labeling (TUNEL) assay for SUGARWIN-treated *F. verticillioides* germlings. *F. verticillioides* germlings grown in the absence of SUGARWIN2 were either exposed to phosphate-buffered saline (PBS) (upper panel) or grown in the presence of 160 μ M SUGARWIN2 (lower panels) for 16 h at 25°C. Germlings were then fixed and double-stained with 4',6-diamidino-2-phenylindole (DAPI) and TUNEL. Bars represent 5 μ m.

ARWIN proteins, could decrease the likelihood of colonization by fungi. Altogether, the late-inducible pattern of expression, the exclusive expression of *sugarwin* mRNA at the wound site, and the extracellular localization of the mRNA support the antipathogenic activity of SUGARWIN proteins.

Plant defenses against pathogens and insects are differentially regulated by cross-communicating signaling pathways in which SA, JA, and ethylene play key roles (Duby et al. 2001; Wu and Baldwin 2009). In general, it can be stated that JA signaling is crucial for protection against herbivore attack (Lukaszewicz et al. 1998; Scott et al. 1999) and appears to mobilize antimicrobial defenses that are predominantly effective against necrotrophic pathogens, whereas the SA-mediated defense response is effective against biotrophic fungi, bacteria, and viruses (Pompermayer et al. 2003). *Barwin*-like genes have also been reported as being induced by treatment with hormone elicitors. ZmPR4 mRNA accumulates after MJ treatment (Bravo et al. 2003). In tobacco, PR4 is induced 4 days after spraying plants with SA (Linthorst et al. 1991). In rice, OsPR4 is induced by MJ and abscissic acid (Agrawal et al. 2003). The wheat *wPR4ef* gene is induced by JA and SA treatment (Howe et al. 1996). In sugarcane, the transcription of the *sugarwin* genes was induced after treatment with MJ but not with MS (Fig. 5A and B), substantiating the pattern of an insect- and wound-inducible gene rather than the pattern suggested by its antipathogenic activity.

We tested a concentration of recombinant proteins in the caterpillar diet that was greater than the endogenous levels found in other plant species (Kiba et al. 2003; Svensson et al. 1992), and

Fig. 7. Effects of sugarcane wound-inducible protein 2 (SUGARWIN2) on the hyphal morphology of *Fusarium verticillioides*. *F. verticillioides* germlings grown in the absence of SUGARWIN2 with 16 h of exposure to phosphate-buffered saline (PBS) at 25°C were used as a control for *F. verticillioides* germlings grown in the presence of 160 μ M SUGARWIN2 for 16 h at 25°C. CFW = Calcofluor White. Bars represent 5 μ m.

this treatment had no effect on *D. saccharalis* life history parameters (larval survival and larval weight). Although our minimal concentration of SUGARWIN protein that showed an effect on fungal growth is greater than the concentration used in previous experiments with other BARWIN proteins (Caruso et al. 2001; Bravo et al. 2003; Zhu et al. 2006), the SUGARWIN proteins may have experienced a decrease in their activity due to the added histidine tag. However, we demonstrated major alterations in the morphology of *F. verticillioides* hyphae when SUGARWIN2 recombinant protein was incorporated into the liquid medium. The hyphae exhibited visible abnormalities, such as a twisted appearance, and a TUNEL assay showed that the protein induced DNA fragmentation (Figs. 6 and 7).

Previous in vitro recombinant protein studies have attributed antifungal properties to BARWIN-like proteins (Bravo et al. 2003; Caruso et al. 2001; Zhu et al. 2006). Indeed, in our study, we showed an effect on fungal growth and hyphal biology which is likely to be part of the plant defense against biotrophic stressors. SUGARWIN proteins are involved in fungal cell wall disruption, leading to the leakage of intracellular material and eventual fungal cell death. The high level of similarity of SUGARWIN proteins to other BARWIN-like proteins with antipathogenic activities also suggests that the antifungal activity is conserved among proteins containing the BARWIN domain. Plants respond to insects and pathogens by means of a myriad of specific and nonspecific compounds. Usually, nonspecific responses are constitutive, and induced responses show different levels of specificity. Our data do not exclude *sugarwin* genes from being part of a set of nonspecific responses; however, their antipathogenic activity and the lack of an effect against the caterpillars support the hypothesis that these genes may be expressed as part of an orchestrated response against pathogens that is triggered by herbivores.

MATERIALS AND METHODS

Subcellular localization of SUGARWIN1 and SUGARWIN2.

The server SignalP 3.0 (Bendtsen et al. 2004) was used to predict the peptide cleavage site of SUGARWIN1 and SUGARWIN2. The primers contained restriction sites for cloning into the vector EN50PMA4-BK-GFP (Duby et al. 2001). The plasmid pBIN 35S-mGFP5 (Haseloff et al. 1997) was used as an ER control. The identities of the constructs were confirmed by nucleotide sequencing. The constructs pBIN 35S-mGFP5, SUGARWIN1::GFP, and SUGARWIN2::GFP were delivered into onion epidermal cells using tungsten particle bombardment, as previously reported (Lukaszewicz et al. 1998). The tungsten particles were accelerated using a PDS-1000/He Biolistic Particle Delivery System (Bio-Rad, Munich) with rupture discs of 1,100 psi under an Hg pressure of 28 in. After bombardment, the cells were allowed to recover for 18 to 22 h at 22°C in continuous light. The bombarded peels were bathed in 20 mM MOPS (pH 7.0) buffer for 4 h, according to Scott and associates (1999), and plasmolyzed for 10 min with 0.5 M mannitol prior to visualization. The fluorescence and DIC microscopy were performed using a Zeiss Axiovert 100 M equipped with $\times 25$ and $\times 40$ water and $\times 100$ oil-immersion lenses (Carl Zeiss, Thornwood, NY, U.S.A.).

Plant and insect materials and treatments.

The sugarcane plantlets and *D. saccharalis* larvae were provided by the Centro de Tecnologia Canavieira, Piracicaba, SP, Brazil. The sugarcane one-eyed seed sets ('SP80-3280') were planted in 200-ml plastic cups containing a commercial planting mix (Plantmax, Eucatex, São Paulo, Brazil) and cultivated in a greenhouse for 20 days under natural light condi-

tions, a temperature of $30 \pm 5^\circ\text{C}$, and $40 \pm 10\%$ relative humidity. The *D. saccharalis* larvae were reared on an artificial diet (Pompermayer et al. 2003) and maintained at $25 \pm 4^\circ\text{C}$ and $60\% \pm 10\%$ relative humidity with a photophase of 14 h.

To monitor *sugarwin1* and *sugarwin2* transcript accumulation in response to insect attack, third- to fourth-instar larvae were starved for 24 h prior to being transferred individually to the leaf roll of a sugarcane plantlet. The leaf roll is the preferred entrance site of *D. saccharalis*. The control plants were left unattacked. For the mechanical wounding, the leaf roll was pierced with fine tweezers every hour for 12 h. Plant material was collected at 0, 6, 12, and 24 h and immediately frozen in liquid nitrogen. Three replicates of four plantlets were used for each time point.

For evaluation of the sugarcane response to the phytohormones MJ and MS, 20-day-old sugarcane plantlets were sprayed with 0.2 mM MJ (Bedoukian Research Inc., Danbury, CT, U.S.A.) and 5 mM MS (Sigma-Aldrich, St. Louis) in 0.02% ethanol diluted with 0.125% Triton X-100 (Sigma-Aldrich). The control plants were sprayed with the same solution without the phytohormones. The control and treated plants were incubated separately in hermetically sealed plastic boxes, each containing an open jar of 10 N NaOH (CO_2 trap) (Howe et al. 1996). Plant material was collected and immediately frozen in liquid nitrogen at 0, 4, 8, and 12 h after spraying with MJ and at 0, 6, 48, and 72 h after spraying with MS. Three replicates of four plantlets were used for each time point. The MJ and MS experiments were repeated two times each.

RNA isolation and cDNA synthesis.

The total RNA of the sugarcane tissue was isolated with TRIZOL reagent (Invitrogen, Carlsbad, CA, U.S.A.), according to the manufacturer's instructions, followed by DNA removal by treatment with 2 units of RNase-free DNase I (Fermentas, Vilnius, Lithuania) for 20 min at 37°C . To remove any traces of DNA or DNase that might interfere with the RNA quantification, the RNA was re-extracted with TRIZOL reagent. The total RNA was quantified, and the quality was checked by spectrophotometer and agarose gel electrophoresis. First-strand synthesis was performed in a total volume of 20 μl , using 2 μg of total RNA, 2.5 μM poly (dT) primer, and 1 μl of reverse transcriptase, according to the manufacturer's instructions (Improm-II Reverse Transcriptase; Promega Corp., Madison, WI, U.S.A.).

Real-time polymerase chain reaction (PCR) gene expression and data analysis.

Real-time quantitative (q)PCR was performed using the Rotor Gene 3000 machine (Corbett Life Science, Sydney, Australia) and SYBR Green I chemistry (Platinum SYBR Green qPCR SuperMix-UDG; Invitrogen). Gene-specific primers were designed using OligoPerfect (Invitrogen), and the primer specificity was confirmed by analysis on a 2% agarose gel, by melting curve analysis, and by nucleotide sequencing followed by BlastX confirmation (Altschul et al. 1997). Glyceraldehyde 3-phosphate dehydrogenase (*GAPDH*) and β -actin were used as endogenous controls and have shown the same pattern of regulation; therefore, *GAPDH* was chosen as the expression reference gene. *GAPDH* and rRNA 25S are considered good reference genes for measuring the relative expression of other genes in sugarcane (Iskandar et al. 2004). The GenBank accessions used to design the primers for *BBPI*, *sugarwin1*, *sugarwin2*, and *GAPDH* were AY093805, CA145787, CA138924, and CA301409, respectively. The primer sequences used for the relative quantification can be found in Supplementary Table S2. The relative quantification was performed using a correction for primer efficiency (Pfaffl 2001).

Heterologous expression of SUGARWIN1 and SUGARWIN2 in *Pichia pastoris*.

The coding regions for SUGARWIN1 and SUGARWIN2, without the signal peptide, were amplified by PCR from the clones described above. The PCR products were digested with the restriction enzyme *Xho*I and cloned into the pPICZαA vector (Invitrogen), which had been digested with the same enzyme. Next, the constructs were transformed into *Escherichia coli* DH5α and selected in Luria-Bertani low-salt medium (1% tryptone, 0.5% yeast extract, and 0.5% NaCl, pH 7.0) supplemented with Zeocin at 25 µg/ml. The recombinant clones were selected by PCR and verified by DNA sequencing. The pPICZα-SUG1 and pPICZα-SUG2 constructs were linearized with *Pme*I, and 300 ng was used to transform *Pichia pastoris* KM71H competent cells, which were prepared as described by Cregg and James (2007). The competent KM71H cells were electroporated using a Gene Pulser II (Bio-Rad) with the following conditions: 1,500 V, 25 µF, and 200 Ω. The transformants were selected by plating in 1% yeast extract, 2% peptone, 2% dextrose, 1 M sorbitol, and 2% agar medium containing zeocin at 100 µg/ml. Several transformants of the SUG1 and SUG2 constructs were confirmed by PCR and screened for expression in a 2-ml scale using 24-deep well plates (Whatman, Maidstone, U.K.) as described by Boettner and associates (2002). Samples (15 µl) of the supernatant of each culture were analyzed by sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE), and the clones with the highest level of expression were used for recombinant protein production. A single colony of *P. pastoris* containing the *sugarwin1* or the *sugarwin2* construct was used to inoculate 10 ml 1% yeast extract, 2% peptone, 100 mM potassium phosphate buffer (pH 7.0), 1.34% YNB, 4×10^{-5} % biotin, and 1% glycerol (BMGY) that was grown at 30°C until an optical density (OD) at 600 nm of approximately 5 was reached. This culture was used to inoculate 500 ml of BMGY that was grown to an OD of 4 to 5. The cells were harvested by centrifugation at $1,500 \times g$ for 5 min, resuspended in 100 ml of BMGY with 0.5% methanol instead of glycerol, and incubated at 26°C for SUGARWIN1 and 30°C for SUGARWIN2. To induce gene expression, methanol was added to each sample every 24 h to maintain a final concentration of 0.75%. After 96 h, the cells were harvested by centrifugation at $1,500 \times g$ for 5 min, and the supernatant was passed through a membrane filter of 0.45 µm (Millipore, Bedford, MA, U.S.A.). The recombinant proteins in the supernatant were purified by affinity chromatography using Ni-NTA-agarose (Qiagen) pre-equilibrated with purification buffer (10 mM Tris-HCl, pH 8.0; 50 mM NaH₂PO₄; and 100 mM NaCl). After binding, the proteins were eluted with two-column volumes of purification buffer containing increased imidazole concentrations (10, 25, 50, 75, 100, and 250 mM). The fractions containing the SUGARWIN1 and SUGARWIN2 proteins were dialyzed in PBS (137 mM NaCl, 2.7 mM KCl, 10 mM Na₂PO₄, and 2 mM KH₂PO₄; pH 7.4) and sterilized with a 0.22-µm filter (Millipore). The protein concentrations were determined using the BCA protein assay kit (Pierce). The purification of both recombinant proteins was monitored using SDS-PAGE (Supplementary Fig. S1).

The effect of recombinant SUGARWIN on the development of *D. saccharalis*.

Newly hatched larvae were transferred to individual petri dishes containing a corn-based artificial diet (Parra and Mihsfeldt 1992) supplemented with 30% (vol/vol) of a solution containing recombinant SUGARWIN1 and SUGARWIN2 (in a final concentration of 420 µg/ml). The negative control was the same diet supplemented with 30% (vol/vol) of PBS buffer. The diet supplemented with 5% (wt/vol) of semipurified soybean

proteinase inhibitor served as positive control (Pompermyer et al. 2003) (250 µM). The proteins were added to the diet when the temperature reached 40°C and were thoroughly mixed before pouring into individual petri dishes. The diet was allowed to rest for 30 min at room temperature before neonate transfer. *D. saccharalis* was reared with a photophase of 14 h at $25 \pm 4^\circ\text{C}$ and $60 \pm 10\%$ relative humidity. Fourteen days after transferring the larvae to the diet, larval survival and mean larval weight were evaluated. The four treatments—negative control (PBS), positive control (soybean proteinase inhibitor), SUGARWIN1 and SUGARWIN2—were arranged in a completely randomized design with an uneven number of repetitions. In total, 240 caterpillars, each one representing a single repetition, were used in the experiments, which were replicated at least twice.

Viability test on *F. verticillioides*.

Initially, conidia of *F. verticillioides* were treated with SUGARWIN2. Approximately 10 conidia of *F. verticillioides* were inoculated into the wells of a 96-well plate containing liquid PD (potato dextrose) medium. After 8 h of incubation at 25°C, SUGARWIN2 was added to each well at a final concentration of 20, 40, 80, or 160 µM. PBS was used as a negative control. The treatments were performed in triplicate. After 16 h of treatment, all of the material was transferred to a 24-well plate containing solid PD medium. The plates were incubated at 25°C for another 36 h.

The effect of recombinant SUGARWIN2 on the mycelial morphology of *F. verticillioides*.

Approximately 1.5×10^4 conidia of *F. verticillioides* were inoculated into wells with coverslips of a 24-well plate containing liquid PD medium. After 8 h of incubation at 25°C, SUGARWIN2 was added to each well at a final concentration of 160 µM. The plates were incubated at 25°C for 16 h. The treatments were performed in triplicate. The morphological analysis was performed at 2, 4, 6, 8, and 16 h. PBS was used as a negative control.

Microscopy of *F. verticillioides*.

The slides were mounted and the images were analyzed by fluorescence microscopy at room temperature. The slides were viewed with a Carl Zeiss (Jena, Germany) microscope using a $\times 100$ magnification oil-immersion objective lens using a 43HE DsRed filter (excitation 550/25 nm and emission 605/70) for staining with TUNEL and a 49DAPI filter (excitation 365 nm and emission 445/50) for staining with 4',6-diamidino-2-phenylindole (DAPI) and Calcofluor. Phase contrast for the brightfield and fluorescent images was captured with an Axio-Cam camera (Carl Zeiss), processed using the AxioVision software version 3.1, and saved as TIFF files. Further processing was performed using Adobe Photoshop 7.0 (Adobe Systems Incorporated, San Jose, CA, U.S.A.).

Calcofluor assay on *F. verticillioides*.

After treatment with SUGARWIN2, slides of *F. verticillioides* were mounted with the addition of 2 µl of a solution of Fluorescent Brightener 28 (Calcofluor White M2R) (Sigma-Aldrich) at a concentration of 1 mg/ml in each sample, which remained for 10 min in the dark (Bertini et al. 2009). The images were acquired using a Carl Zeiss microscope using a $\times 100$ objective lens, brightfield imaging, and 49DAPI filter for staining with Calcofluor.

TUNEL assay on *F. verticillioides*.

DNA strand breaks were demonstrated by a TUNEL assay using an In Situ Cell Death Detection Kit, TMR red (Roche

Diagnostics GmbH, Mannheim, Germany). After 16 h of treatment with SUGARWIN2 at 25°C, as previously described, the supernatants containing the hyphae were transferred to microtubes and washed with PBS. The TUNEL assay was then performed as described by Madeo and associates (1997) for the yeast *Saccharomyces cerevisiae*, with modifications. The hyphae were first fixed with fixation solution (3.7% formaldehyde, 0.2% Tween 80, and 50 mM $\text{N}_2\text{H}_2\text{PO}_4$) for 30 min at room temperature and then washed in PBS. The hyphae were incubated in permeabilization solution (0.1% Triton X-100 and 0.1% sodium citrate) for 10 min on ice, followed by a further washing with PBS. The hyphae were incubated with a TUNEL reaction solution for 1 h at 37°C, followed by further washing with PBS. The cells were then subjected to nuclear staining for 3 min with DAPI (Sigma-Aldrich) at 0.1 µg/ml, a fluorescent stain that binds strongly to A-T rich regions in the DNA, producing a bright blue nucleus. The images were acquired using a Carl Zeiss microscope with a ×100 objective lens, brightfield imaging, and 49DAPI and 43HE DsRed filters for staining with DAPI and TUNEL, respectively.

ACKNOWLEDGMENTS

We thank S. M. Chabregas and M. C. Falco of the Centro de Tecnologia Canavieira for providing the sugarcane plantlets, *D. saccharalis* larvae, and fungal isolates; E. Calderan for enormous assistance with insect and fungal experiments; A. C. Santiago for technical assistance; J. Haseloff for providing the plasmid pBIN 35S-mGFP5; N. G. Zério for skillful assistance in the *D. saccharalis* rearing assays; A. Figueira for the real-time PCR facilities; and M. M. Brandão for the phylogenetic analyses. This work was supported by the Fundação de Amparo a Pesquisa do Estado de São Paulo (FAPESP) grant 08/52067-3, the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) grant 474542/2010-6, and the Instituto Nacional de Ciência e Tecnologia do Bioetanol grant 574002/2008-1 and FAPESP 08/57908-6. M. C. Silva-Filho, G. H. Goldman, and F. F. Henrique-Silva are also research fellows of CNPq.

LITERATURE CITED

- Abbott, E. V., and Hughes, C. G. 1961. Red rot. Pages 263-282 in: Sugarcane Diseases of the World. J. P. Martin, E. V. Abbott, and C. G. Hughes, eds. Elsevier, Amsterdam.
- Agrawal, G. K., Jwa, N. S., Han, K. S., Agrawal, V. P., and Rakwal, R. 2003. Isolation of a novel rice PR4 type gene whose mRNA expression is modulated by blast pathogen attack and signaling components. *Plant Physiol. Biochem.* 41:81-90.
- Altschul, S. F., Madden, T. L., Schaffer, A. A., Zhang, J., Zhang, Z., Miller, W., and Lipman, D. J. 1997. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res.* 25:3389-3402.
- Anisimova, M., and Gascuel, O. 2006. Approximate likelihood-ratio test for branches: A fast, accurate, and powerful alternative. *Syst. Biol.* 55:539-552.
- Bendtsen, J. D., Nielsen, H., von Heijne, G., and Brunak, S. 2004. Improved prediction of signal peptides: SignalP 3.0. *J. Mol. Biol.* 340:783-795.
- Bertini, L., Cascone, A., Tucci, M., D'Amore, R., Di Bernardino, I., Buonocore, V., Caporale, C., and Caruso, C. 2006. Molecular and functional analysis of new members of the wheat PR4 gene family. *Biol. Chem.* 387:1101-1111.
- Bertini, L., Caporale, C., Testa, M., Proietti, S., and Caruso, C. 2009. Structural basis of the antifungal activity of wheat PR4 proteins. *FEBS (Feb. Eur. Biochem. Soc.) Lett.* 583:2865-2871.
- Boettner, M., Prinz, B., Holz, C., Stahl, U., and Lang, C. 2002. High-throughput screening for expression of heterologous proteins in the yeast *Pichia pastoris*. *J. Biotechnol.* 99:51-62.
- Bravo, J. M., Campo, S., Murillo, I., Coca, M., and Segundo, B. S. 2003. Fungus- and wound-induced accumulation of mRNA containing a class II chitinase of the pathogenesis-related protein 4 (PR-4) family of maize. *Plant Mol. Biol.* 52:745-759.
- Broekaert, I., Lee, H. I., Kush, A., Chua, N. H., and Raikhel, N. 1990. Wound-induced accumulation of mRNA containing a hevein sequence in laticifers of rubber tree (*Hevea brasiliensis*). *Proc. Natl. Acad. Sci. U.S.A.* 87:7633-7637.
- Caruso, C., Bertini, L., Tucci, M., Caporale, C., Leonardi, L., Saccardo, F., Bressan, R. A., Veronese, P., and Buonocore, V. 1999. Isolation and characterisation of wheat cDNA clones encoding PR4 proteins. *DNA Seq.* 10:301-307.
- Caruso, C., Bertini, L., Tucci, M., Caporale, C., Nobile, M., Leonardi, L., and Buonocore, V. 2001. Recombinant wheat antifungal PR4 proteins expressed in *Escherichia coli*. *Protein Expr. Purif.* 23:380-388.
- Christopher, M. E., Miranda, M., Major, I. T., and Constabel, C. P. 2004. Gene expression profiling of systemically wound-induced defenses in hybrid poplar. *Planta* 219:936-947.
- Cregg, J. M. 2007. *Methods in Molecular Biology*.
- Cui, J., Jander, G., Racki, L. R., Kim, P. D., Pierce, N. E., and Ausubel, F. M. 2002. Signals involved in *Arabidopsis* resistance to *Trichoplusia ni* caterpillars induced by virulent and avirulent strains of the phytopathogen *Pseudomonas syringae*. *Plant Physiol.* 129:551-564.
- Dangl, J. L., and Jones, J. D. 2001. Plant pathogens and integrated defence responses to infection. *Nature* 411:826-833.
- De Noij, M. P., Biere, A., and Linders, E. G. A. 1992. Pest and pathogen interactions through host predisposition. Pages 143-160 in: *Pests and Pathogens: Plant Responses to Foliar Attack*. P. G. Ayres, ed. BIOS Scientific, Oxford.
- De Vos, M., Van Zaanen, W., Koornneef, A., Korzelius, J. P., Dicke, M., Van Loon, L. C., and Pieterse, C. M. 2006. Herbivore-induced resistance against microbial pathogens in *Arabidopsis*. *Plant Physiol.* 142:352-363.
- Duby, G., Degand, H., and Boutry, M. 2001. Structure requirement and identification of a cryptic cleavage site in the mitochondrial processing of a plant F1-ATPase beta-subunit presequence. *FEBS (Feb. Eur. Biochem. Soc.) Lett.* 505:409-413.
- Edgar, R. C. 2004. MUSCLE: Multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res.* 32:1792-1797.
- Falco, M. C., Marbach, P. A. S., Pompermayer, P., Lopes, F. C. C., and Silva-Filho, M. C. 2001. Mechanisms of sugarcane response to herbivory. *Genet. Mol. Biol.* 24:113-122.
- Friedrich, L., Moyer, M., Ward, E., and Ryals, J. 1991. Pathogenesis-related protein-4 is structurally homologous to the carboxy-terminal domains of hevein, win-1 and win-2. *Mol. Gen. Genet.* 230:113-119.
- Gatch, E. W., and Munkvold, G. P. 2002. Fungal species composition in maize stalks in relation to European corn borer injury and transgenic insect protection. *Plant Dis.* 86:1156-1162.
- Green, T. R., and Ryan, C. A. 1972. Wound-induced proteinase inhibitor in plant leaves: a possible defense mechanism against insects. *Science* 175:776-777.
- Grivet, L., and Arruda, P. 2002. Sugarcane genomics: depicting the complex genome of an important tropical crop. *Curr. Opin. Plant Biol.* 5:122-127.
- Guindon, S., and Gascuel, O. 2003. A simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood. *Syst. Biol.* 52:696-704.
- Haseloff, J., Siemering, K. R., Prasher, D. C., and Hodge, S. 1997. Removal of a cryptic intron and subcellular localization of green fluorescent protein are required to mark transgenic *Arabidopsis* plants brightly. *Proc. Natl. Acad. Sci. U.S.A.* 94:2122-2127.
- Hejgaard, J., Jacobsen, S., Bjorn, S. E., and Kragh, K. M. 1992. Antifungal activity of chitin-binding PR-4 type proteins from barley grain and stressed leaf. *FEBS (Feb. Eur. Biochem. Soc.) Lett.* 307:389-392.
- Howe, G. A., Lightner, J., Browse, J., and Ryan, C. A. 1996. An octadecanoid pathway mutant (JL5) of tomato is compromised in signaling for defense against insect attack. *Plant Cell* 8:2067-2077.
- Iskandar, H. M., Simpson, R. S., Casu, R. E., Bonnett, G. D., MacLean, D. J., and Manners, J. M. 2004. Comparison of reference genes for quantitative real-time polymerase chain reaction analysis of gene expression in sugarcane. *Plant Mol. Biol. Rep.* 22:325-337.
- Karban, R., and Baldwin, I. T. 1997. *Induced Responses to Herbivory*. Chicago University Press, Chicago.
- Kessler, A., and Baldwin, I. T. 2002. Plant responses to insect herbivory: the emerging molecular analysis. *Annu. Rev. Plant Biol.* 53:299-328.
- Khanna, K. L., and Rafay, S. A. 1953. Annual report of the scheme for the investigation and control of wilt disease of sugar cane for the year ending 3rd May 1953. Central Sugarcane Research Station, Bihar, India.
- Kiba, A., Saitoh, H., Nishihara, M., Omiya, K., and Yamamura, S. 2003. C-terminal domain of a hevein-like protein from *Wasabia japonica* has potent antimicrobial activity. *Plant Cell Physiol.* 44:296-303.
- Li, X., Xia, B., Jiang, Y., Wu, Q., Wang, C., He, L., Peng, F., and Wang, R. 2010. A new pathogenesis-related protein, LrPR4, from *Lycoris radiata*, and its antifungal activity against *Magnaporthe grisea*. *Mol. Biol. Rep.* 37:995-1001.
- Linthorst, H. J. M., Danhash, N., Brederode, F. T., Vankan, J. A. L., Dewit, P. J. G. M., and Bol, J. F. 1991. Tobacco and tomato PR proteins homologous to win and pro-hevein lack the hevein domain. *Mol. Plant-Microbe Interact.* 4:586-592.

- Lukaszewicz, M., Jerouville, B., and Boutry, M. 1998. Signs of translational regulation within the transcript leader of a plant plasma membrane H⁺-ATPase gene. *Plant J.* 14:413-423.
- Madeo, F., Frohlich, E., and Frohlich, K.U. 1997. A yeast mutant showing diagnostic markers of early and late apoptosis. *J. Cell Biol.* 139:729-734.
- McFarlane, S. A., Govender, P., and Rutherford, R. S. 2009. Interactions between *Fusarium* species from sugarcane and the stalk borer, *Eldana saccharina* (Lepidoptera: Pyralidae). *Ann. Appl. Biol.* 155:349-359.
- McKaig, N. 1936. Chemical composition of juice from Louisiana sugarcane injured by the sugarcane borer and the red rot disease. *J. Agric. Res.* 52:0017-0025.
- Mian, A. A., Coates, J., and Cordiner, J. B. 1969. Binary gaseous diffusion coefficients of N₂-HCl, a-HBr and N₂-HBr systems as a function of temperature. *Can. J. Chem. Eng.* 47:499.
- Moura, D. S., and Ryan, C. A. 2001. Wound-inducible proteinase inhibitors in pepper. Differential regulation upon wounding, systemin, and methyl jasmonate. *Plant Physiol.* 126:289-298.
- Ogunwolu, E. O., Reagan, T. E., Flynn, J. L., and Hensley, S. D. 1991. Effects of *Diatraea saccharalis* (F) (Lepidoptera, Pyralidae) damage and stalk rot fungi on sugarcane yield in Louisiana. *Crop Prot.* 10:57-61.
- Parra, J. R. P., and Mihsfeldt, L. H. 1992. Comparison of artificial diets for rearing the sugarcane borer. Pages 195-209 in: *Advances in Insect Rearing for Research and Pest Management* T. A. N. Leppla, ed. Westview Press, Boulder, CO, U.S.A.
- Pfaffl, M. W. 2001. A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Res.* 29:e45. Published online.
- Pfaffl, M. W., Horgan, G. W., and Dempfle, L. 2002. Relative expression software tool (REST (c)) for group-wise comparison and statistical analysis of relative expression results in real-time PCR. *Nucleic Acids Res.* 30:e36. Published online.
- Pompermayer, P., Falco, M. C., Parra, J. R. P., and Silva-Filho, M. C. 2003. Coupling diet quality and Bowman-Birk and Kunitz-type soybean proteinase inhibitor effectiveness to *Diatraea saccharalis* development and mortality. *Entomol. Exp. Appl.* 109:217-224.
- Reymond, P., Bodenhausen, N., Van Poecke, R. M., Krishnamurthy, V., Dicke, M., and Farmer, E. E. 2004. A conserved transcript pattern in response to a specialist and a generalist herbivore. *Plant Cell* 16:3132-3147.
- Quevillon, E., Silventoinen, V., Pillai, S., Harte, N., Mulder, N., Apweiler, R., and Lopez, R. 2005. InterProScan: protein domains identifier. *Nucleic Acids Res.* 33:W116-W120.
- Ryan, C. A. 1968. An inducible protein in potato and tomato leaflets. *Plant Physiol.* 43:1880-1881.
- Ryan, C. A. 1990. Protease Inhibitors in plants—genes for improving defenses against insects and pathogens. *Annu. Rev. Phytopathol.* 28:425-449.
- Ryan, C. A. 2000. The systemin signaling pathway: differential activation of plant defensive genes. *BBA-Protein Struct. M* 1477:112-121.
- Schilmiller, A. L., and Howe, G. A. 2005. Systemic signaling in the wound response. *Curr. Opin. Plant Biol.* 8:369-377.
- Scott, A., Wyatt, S., Tsou, P. L., Robertson, D., and Allen, N. S. 1999. Model system for plant cell biology: GFP imaging in living onion epidermal cells. *Biotechniques* 26:1125, 1128-1132.
- Singh, K., Budhraj, T. R., and Agnihotri, V. P. 1977. Survival of *Colletotrichum falcatum* in soil, its portals of entry and role of inoculum density in causing infection. *Int. Sugar J.* 79:43-44.
- Stout, M. J., Fidantsef, A. L., Duffey, S. S., and Bostock, R. M. 1999. Signal interactions in pathogen and insect attack: systemic plant-mediated interactions between pathogens and herbivores of the tomato, *Lycopersicon esculentum*. *Physiol. Mol. Plant Pathol.* 54:115-130.
- Svensson, B., Svendsen, I., Hojrup, P., Roepstorff, P., Ludvigsen, S., and Poulsen, F. M. 1992. Primary structure of barwin: a barley seed protein closely related to the C-terminal domain of proteins encoded by wound-induced plant genes. *Biochemistry* 31:8767-8770.
- Upadhyay, R. K., Mukerji, K. G., Chamola, B. P., and Dueby, O. P. 2009. *Integrated Pest and Disease Management*. APH Publishing Corporation, New Delhi, India.
- Vettore, A. L., da Silva, F. R., Kemper, E. L., and Arruda, P. 2001. The libraries that made SUCEST. *Genet. Mol. Biol.* 24:1-7.
- Vettore, A. L., da Silva, F. R., Kemper, E. L., Souza, G. M., da Silva, A. M., Ferro, M. L., Henrique-Silva, F., Gigliotti, E. A., Lemos, M. V., Coutinho, L. L., Nobrega, M. P., Carrer, H., Franca, S. C., Bacci Junior, M., Goldman, M. H., Gomes, S. L., Nunes, L. R., Camargo, L. E., Siqueira, W. J., Van Sluys, M. A., Thiemann, O. H., Kuramae, E. E., Santelli, R. V., Marino, C. L., Targon, M. L., Ferro, J. A., Silveira, H. C., Marini, D. C., Lemos, E. G., Monteiro-Vitorello, C. B., Tambor, J. H., Carraro, D. M., Roberto, P. G., Martins, V. G., Goldman, G. H., de Oliveira, R. C., Truffi, D., Colombo, C. A., Rossi, M., de Araujo, P. G., Sculaccio, S. A., Angella, A., Lima, M. M., de Rosa Junior, V. E., Siviero, F., Coscrato, V. E., Machado, M. A., Grivet, L., Di Mauro, S. M., Nobrega, F. G., Menck, C. F., Braga, M. D., Telles, G. P., Cara, F. A., Pedrosa, G., Meidanis, J., and Arruda, P. 2003. Analysis and functional annotation of an expressed sequence tag collection for tropical crop sugarcane. *Genome Res.* 13:2725-2735.
- Walling, L. L. 2000. The myriad plant responses to herbivores. *J. Plant Growth Regul.* 19:195-216.
- Wu, J., Baldwin, I. T. 2009. Herbivory-induced signalling in plants: perception and action. *Plant Cell Environ.* 32:1161-1174.
- Wu, J. Q., and Baldwin, I. T. 2010. New insights into plant responses to the attack from insect herbivores. *Annu. Rev. Genet.* 44:1-24.
- Zhu, T., Song, F., and Zheng, Z. 2006. Molecular characterization of the rice pathogenesis-related protein, OsPR-4b, and its antifungal activity against *Rhizoctonia solani*. *J. Phytopathol.* 154:378-384.

7.7. Anexo P. Prêmio SBBq: “SBBQ Award” para o pôster “Molecular Characterization of Propolis Induced Cell Death in *Saccharomyces cerevisiae*, de Castro, P. A.; Savoldi, M.; Bonatto, D.; De Barros, M. H.; Goldman, M. H. S.; Berreta, A. A.; Goldman, G. H.”

Sociedade Brasileira de Bioquímica e
Biologia Molecular - SBBq

CERTIFICATE

We hereby certify that the abstract entitled
Molecular Characterization of Propolis Induced Cell Death in
Saccharomyces cerevisiae. (H 28)

Castro, P.A. (USP-FCFRP), Savoldi, M. (USP-FCFRP), Bonatto, D. (UCS), Barros, M.H. (USP-
ICB II), Goldman, M.H.S. (USP-FFCLRP), Berretta, A.A. (Apis Flora Industrial e Comercial),
Goldman, G.H. (USP-FCFRP)

received the "**SBBQ AWARD**" for the best poster presented during the XL Annual
Meeting of the Brazilian Society for Biochemistry and Molecular Biology (SBBq),
April 30th to May 3rd, 2011, in the Convention Center of the Rafain Palace Hotel in
Foz do Iguaçu, Paraná, Brazil.

May 3rd, 2011

Helena C.F. Oliveira
Acting President of SBBq