UNIVERSIDADE DE SÃO PAULO

FACULDADE DE MEDICINA DE RIBEIRÃO PRETO

DEPARTAMENTO DE BIOLOGIA CELULAR E MOLECULAR E BIOAGENTES
PATOGÊNICOS

PROGRAMA DE PÓS-GRADUAÇÃO EM BIOLOGIA CELULAR E MOLECULAR

**Suscetibilidade de macrófagos alveolares murinos à
infecção por *Coxiella burnetii* fase II *in vitro***

TALITA DUARTE FERNANDES

RIBEIRÃO PRETO – SP

2018

TALITA DUARTE FERNANDES

Suscetibilidade de macrófagos alveolares murinos à infecção por *Coxiella burnetii* fase II *in vitro*

Versão original

Tese apresentada à Faculdade de Medicina de
Ribeirão Preto, Universidade de São Paulo, para
obtenção do título de Doutor em Ciências.

Área de Concentração: Biologia Celular e
Molecular

Orientador: Prof. Dr. Dario Simões Zamboni

RIBEIRÃO PRETO – SP

2018

Autorizo a reprodução e divulgação total ou parcial deste trabalho, por qualquer meio convencional ou eletrônico, para fins de estudo e pesquisa, desde que citada a fonte.

FICHA CATALOGRÁFICA

Preparada pela Biblioteca Central do Campus Administrativo de Ribeirão Preto / USP

FERNANDES, Talita Duarte

Suscetibilidade de macrófagos alveolares murinos à infecção por *Coxiella burnetii* fase II *in vitro*. Ribeirão Preto, 2018.

161 p. : il.; 30 cm

Tese de Doutorado apresentada à Faculdade de Medicina de Ribeirão Preto/USP. Área de Concentração: Biologia Celular e Molecular

Orientador: ZAMBONI, Dario Simões.

1. *Coxiella burnetii*. 2. Macrófagos murinos. 3. Permissividade. 4. Polarização. 5. Modelo.

FOLHA DE APROVAÇÃO

Nome: FERNANDES, Talita Duarte

Título: Suscetibilidade de macrófagos alveolares murinos à infecção por *Coxiella burnetii* fase II *in vitro*.

Tese apresentada à Faculdade de Medicina de
Ribeirão Preto, Universidade de São Paulo, para
obtenção do título de Doutor em Ciências.

Aprovada em: _____

Banca Examinadora

Prof. Dr. _____ Instituição: _____

Julgamento: _____ Assinatura: _____

Prof. Dr. _____ Instituição: _____

Julgamento: _____ Assinatura: _____

Prof. Dr. _____ Instituição: _____

Julgamento: _____ Assinatura: _____

Prof. Dr. _____ Instituição: _____

Julgamento: _____ Assinatura: _____

Prof. Dr. _____ Instituição: _____

Julgamento: _____ Assinatura: _____

*Aos meus pais, Miguel e Adriana, pela confiança
inabalável em mim e pelo apoio constante em todos
os meus projetos.*

“Science means constantly walking a tightrope between blind faith and curiosity; between expertise and creativity; between bias and openness; between experience and epiphany, between ambition and passion; and between arrogance and conviction – in short, between an old today and a new tomorrow”.
(Heinrich Rohrer)

AGRADECIMENTOS

Aos meus pais, Miguel e Adriana, pelo amor incondicional e pelo apoio em todas as etapas da minha vida. Por sempre estarem ao meu lado, me incentivando e torcendo pelo meu sucesso em tudo aquilo que eu me atrevo a fazer. Obrigada por serem o porto seguro de todas as minhas jornadas! Ao meu irmão, Tiago, pelo convívio diário e cada vez mais próximo, por aguentar o meu humor peculiar e por estar ao meu lado o tempo todo, incluindo as várias madrugadas de trabalho. Vocês três são a minha base e a minha vida. Amo vocês!

A todos os meus familiares, minhas avós amadas, meus tios e tias, primos e primas... Pelos encontros em família, pelas disputas de jogos de tabuleiro, pelos churrascos e almoços de domingo e pela presença constante em minha vida. Obrigada por sempre acreditarem e torcerem por mim!

Ao Professor Dario Simões Zamboni, por ter acreditado em mim e por ter aceitado continuar esse processo de orientação, agora durante o doutorado. Obrigada por todo o apoio, desde a época da iniciação científica, e por demonstrar sempre que fazer ciência de qualidade é um desafio grande, porém extremamente recompensador. Sou grata pela confiança que depositou em mim durante essa trajetória, especialmente durante esse período de mudanças que vivi nos últimos tempos. Obrigada por ter contribuído de forma imprescindível para minha formação enquanto cientista e pesquisadora.

À Prof^a Hayley Newton (*University of Melbourne at the Peter Doherty Institute for Infection and Immunity*, Austrália), pela colaboração com o meu projeto, através de ótimas discussões científicas e por ter gentilmente cedido a *Coxiella burnetii* expressando *mCherry*, que foi fundamental para o desenvolvimento do meu artigo. Muito obrigada!

Aos membros da banca examinadora, por aceitarem meu convite e por se disponibilizarem para avaliação deste trabalho.

Aos colegas do Laboratório de Patogenicidade Microbiana e Imunidade Inata, tanto aqueles que passaram pelo laboratório e seguiram novos caminhos quanto aqueles que ainda estão presentes, meu muito obrigado por serem a melhor equipe com a qual alguém poderia desejar trabalhar. Obrigada por sempre estarem presentes e dispostos a ajudar quando precisei! Vocês tornaram esse processo muito mais divertido.

À Maira, nossa técnica querida, por toda a ajuda durante todos esses anos! Obrigada por sua dedicação ao Laboratório e por sempre contribuir para fazer tudo funcionar direitinho. Nossa vida seria muito mais complicada sem você! Agradeço ainda à Victória e Letícia, ajudarem a manter o laboratório organizado e tornar nosso trabalho muito mais fácil.

Um agradecimento especial ao Djalma, Lili e Ju Ribeiro, não só pelos anos de convivência no laboratório, mas também pelo auxílio fundamental na produção do meu artigo. Suas contribuições foram essenciais para o sucesso da publicação. À Lari, que esteve comigo durante toda a minha iniciação científica e me ensinou a amar e respeitar essa bactéria incrível que é a *Coxiella*, muito obrigada por ter sido meu modelo de cientista logo que entrei nesse mundo de pesquisas. Tenho muito orgulho de ter o seu nome no meu artigo e tenho orgulho de poder dizer que fui sua “cria”.

À Cati, Nati e Alexandre, obrigada por serem alguns dos melhores amigos que eu fiz no laboratório! Obrigada por todas as risadas, todo o apoio, toda a ajuda e toda a cumplicidade durante todos esses anos. A rotina pesada ficava muito mais fácil com vocês, e sentirei muita falta de todos! À Dani, minha amiga amada, muito obrigada por tudo que você faz e fez por mim dentro e fora do laboratório. Obrigada pela amizade, pelas viagens, pelas conversas, pelos treinos, por estar sempre por perto para me ouvir quando precisei, e principalmente por ser sempre sincera e falar aquelas verdades que eu posso não querer ouvir, mas que sei que são pro meu bem. Obrigada por muitas vezes acreditar em mim mais do que eu mesma, por me incentivar e por mostrar que eu capaz. Quando eu crescer, quero ser igual você!

Aos funcionários do biotério, pela paciência e pelo cuidado com os animais, fundamentais para realização desse trabalho. Um agradecimento especial ao Rubilan, que além de tudo sempre arrumava um tempo pra comentar comigo as últimas impressões sobre as partidas de futebol americano. À Maria Helena, pelo cuidado e ajuda no preparo dos materiais. À Tuca, Roberta e José pelo auxílio com as microscopias.

À Gabriela, secretária do Programa de Pós-graduação, pela paciência infinita e por sempre me atender com um sorriso no rosto toda vez que eu chegava desesperada com dúvidas sobre alguma coisa. Obrigada por sempre ser tão prestativa e por sempre conseguir me acalmar, Gabi!

A todos os funcionários, docentes e colegas do Programa de Pós Graduação em Biologia Celular e Molecular da FMRP e do Departamento de Biologia Celular e Molecular e Bioagentes Patogênicos, por contribuírem para criação de um espaço estimulante para discussão de ciência e por ajudarem no meu crescimento pessoal e profissional durante todos esses anos.

Às amigadas de longa data, que estão ao meu lado em todos os momentos importantes e que fazem a minha vida mais feliz: Bia, Lo, Dani, Bá, Lê e Bru. Obrigada por serem as melhores amigas que alguém poderia ter. Obrigada pelos momentos de descontração, pelo

apoio mútuo, pelo carinho de sempre e por serem aquelas com quem eu sei que sempre posso contar pra tudo que precisar. Amo vocês, meninas! À minha querida família francana, em especial à Marizinha e Cinthia, por esse amor que existe desde pequena e que ultrapassa a distância. Meu coração está sempre com vocês!

Aos colegas de trabalho que chegaram há não tanto tempo na minha vida, mas que compartilham as vitórias e as dificuldades dessa nova profissão que escolhi seguir, muito obrigada por participarem da construção dessa nova etapa da minha vida. Um agradecimento especial aos colegas peritos, Jesus, Bruna, André e Thatiany, à Marcella, auxiliar de autópsia mais fofa do mundo, e principalmente à Débora, minha médica legista preferida, que em tão pouco tempo de convivência se tornou uma das minhas melhores amigas. Amiguinha, obrigada por ser tão parecida comigo e por me entender tão bem! Obrigada por me incentivar nas horas difíceis, por sempre me fazer rir das coisas mais absurdas e por me permitir ser quem eu sou sem me julgar (muito).

Agradeço ainda à Universidade de São Paulo, por oferecer as condições necessárias para que eu pudesse crescer pessoal e profissionalmente, por meio de sua estrutura, funcionários e docentes. A todas as agências de fomento que auxiliam financeiramente o laboratório (CRID, INCTV, FAEPA, CNPq, FAPESP, TDR/WHO, PEW), meu muito obrigado pelo apoio essencial durante o desenvolvimento desse projeto. Um agradecimento especial à Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) pelo suporte financeiro concedido durante os anos iniciais do desenvolvimento desse trabalho (Código de Financiamento 001)..

Por fim, deixo ainda meu muito obrigado para aqueles que não mais estão por aqui, mas que sempre torceram por mim e que com certeza estariam orgulhosos: Vô Wilson, Vô Miguel, Tia Tê e Bisa. Saudades sempre...

RESUMO

FERNANDES, Talita Duarte. **Suscetibilidade de macrófagos alveolares murinos à infecção por *Coxiella burnetii* fase II *in vitro***. 2018. 161 f. Tese (Doutorado Direto em Biologia Celular e Molecular) – Faculdade de Medicina de Ribeirão Preto, Universidade de São Paulo, Ribeirão Preto, 2018.

Coxiella burnetii é a bactéria intracelular causadora da Febre Q, capaz de subverter funções celulares e evadir o reconhecimento do sistema imune da célula hospedeira permitindo o estabelecimento do seu nicho replicativo nas células-alvo: macrófagos e monócitos. É um patógeno altamente virulento, sendo necessário poucos organismos para desencadear a doença, e é considerado como um potencial agente de bioterrorismo da categoria B. Entre os danos econômicos causados por *C. burnetii* destaca-se a infecção de animais de gado, seu reservatório natural, pois causa aborto espontâneo dos filhotes e, consequentemente, prejuízo aos produtores. Seres humanos também adquirem a infecção, por inalação de partículas contaminadas. Hospedeiros imunocompetentes são capazes de restringir a infecção por *C. burnetii* apesar dos diversos mecanismos de evasão da resposta imune do hospedeiro, como interação com vias de sinalização celular e utilização de efetores bacterianos. No entanto, em hospedeiros não competentes a infecção pode evoluir para casos de Febre Q crônica e levá-los à morte. Modelos murinos são frequentemente utilizados para entender as interações patógeno-hospedeiro nas infecções por essa bactéria. A diferença na suscetibilidade de macrófagos de diferentes linhagens murinas e de diferentes tipos celulares à infecção por *C. burnetii* ainda é pouco conhecida. No entanto, sabe-se que *C. burnetii* fase II sucumbe a macrófagos derivados da medula óssea (BMDMs) de camundongos C57BL/6, enquanto células das linhagens BALB/c e A/J são suscetíveis à infecção. Nesse contexto, considerando a relevância biomédica de *C. burnetii*, faz-se necessário a determinação de um modelo relevante para melhor compreender as relações patógeno-hospedeiro nas infecções por essa bactéria. Neste trabalho, caracterizamos um novo modelo de estudo com macrófagos primários que permite avaliar a infecção com *C. burnetii* fase II *in vitro*, além de elucidar os mecanismos relacionados à suscetibilidade das células à bactéria. Por meio de quantificação do DNA genômico bacteriano por qPCR verificamos que macrófagos alveolares (AMs) murinos são altamente suscetíveis à replicação da bactéria, mesmo no fundo gênico restritivo C57BL/6. Caracterizamos a replicação da bactéria em AMs por microscopia de fluorescência e microscopia eletrônica de transmissão e demonstramos que essa replicação ocorre dentro dos vacúolos parasitóforos típicos. Pela análise de expressão gênica por RT-PCR e

fenotipagem de marcadores de superfície por FACS identificamos que a alta suscetibilidade dos AMs se deve a uma polarização dessas células para um padrão M2. Por fim, validamos a relevância desse modelo pela análise da suscetibilidade de AMs murinos deficientes para NOS2, IFN- γ e IL-4 como prova de princípio. Verificamos que a suscetibilidade de AMs é comparável à de células Vero e células THP-1, modelos celulares imortalizados descritos como altamente suscetíveis à replicação de *C. burnetii*, e que AMs podem ser utilizados para estudos utilizando células derivadas de camundongos com fundo genético C57BL/6. Dessa forma, nosso trabalho caracterizou os AMs murinos como um relevante modelo celular primário altamente suscetível para o estudo das interações patógeno-hospedeiro frente à infecção *in vitro* por *C. burnetii* fase II, além de elucidar os mecanismos por trás dessa suscetibilidade.

Palavras-chave: *Coxiella burnetii*. Macrófagos murinos. Permissividade. Polarização. Modelo.

ABSTRACT

FERNANDES, Talita Duarte. **Susceptibility of murine alveolar macrophages to infection by *Coxiella burnetii* phase II in vitro**. 2018. 161 f. Tese (Doutorado Direto em Biologia Celular e Molecular) – Faculdade de Medicina de Ribeirão Preto, Universidade de São Paulo, Ribeirão Preto, 2018.

Coxiella burnetii is the intracellular bacterium that causes Q fever, capable of subverting cellular functions and evading recognition by the host cell immune system, thus allowing the establishment of its replicative niche in its target cells: macrophages and monocytes. It's a highly virulent pathogen, with few organisms being sufficient to cause the disease, and considered a potential class B bioterrorism agent. Among the economic damages caused by *C. burnetii* are infection of cattle animals, its natural reservoir, that leads to spontaneous abortion of the offspring and financial loss to the producers. Human beings also acquire the infection, through inhalation of contaminated air particles. Immunocompetent hosts are able to restrict infection by *C. burnetii* despite the several evasion mechanisms from the host immune system, which includes interaction with cell signaling pathways and utilization of bacterial effectors. However, in non-competent hosts the infection can evolve to chronic Q Fever and lead to death. Murine models are frequently used to understand the host-pathogen interactions during infections by this bacterium. The difference in the susceptibility among macrophages from different murine strains, as well as different cell types, still poorly known. However, it is known that *C. burnetii* phase II succumbs to bone-marrow derived macrophages (BMDMs) from C57BL/6 mice, whereas cells from BALB/c and A/J strains are susceptible to infection. In this context, considering the biomedical relevance of *C. burnetii*, it's necessary to establish a relevant study model to better understand the host-pathogen relations in infections by *C. burnetii*. In this work, we characterized a new study model with primary macrophages to assess the infection by *C. burnetii* phase II in vitro and elucidated the mechanisms underlying the susceptibility to *C. burnetii*. By using qPCR for quantification of bacterial genomic DNA, we saw that murine alveolar macrophages (AMs) are highly susceptible to *C. burnetii* replication, even in the usually restrictive C57BL/6 background. We characterized the bacterium replication in murine AMs by fluorescence microscopy and transmission electron microscopy, showing that this replication occurred inside the typical *C. burnetii*-containing vacuole. We also identified, through the gene expression by RT-PCR and the phenotypic profile of surface markers by FACS, that the increased susceptibility of murine AMs were due to a polarization of these cells into a M2 profile. To finish, we validated the relevance of this

model through the analysis of the susceptibility of murine AMs deficient to NOS2, IFN- γ and IL-4 as a proof of principle. We verified that the susceptibility of AMs is comparable to Vero cells and THP-1 cells, immortalized cell models described as highly susceptible to *C. burnetii* replication, and that AMs can be used to studies using cells derived from mice with the C57BL/6 background. In this way, our work characterized murine AMs as a relevant primary cell model highly susceptible to studies among the host-pathogen interactions in infections with *C. burnetii* phase II in vitro, and we also elucidate the mechanisms underlying this susceptibility.

Keywords: *Coxiella burnetii*. Murine macrophages. Permissiveness. Polarization. Model.

LISTA DE FIGURAS

Figura 1 - Representação do ciclo de vida bifásico de <i>C. burnetii</i>	20
Figura 2 - Processo de desenvolvimento e maturação do vacúolo contendo <i>C. burnetii</i>	25
Figura 3 - Representação do modelo de polarização de macrófagos com os estados comuns para humanos e camundongos.....	32
Figura 4 - Representação dos modelos de plasticidade e ontogenia na determinação do estado de polarização dos macrófagos.....	35
Figura 5 - Macrófagos murinos derivados de medula óssea da linhagem A/J são mais suscetíveis à multiplicação de <i>C. burnetii</i> fase II do que macrófagos murinos derivados de medula óssea da linhagem C57BL/6.....	49
Figura 6 - Macrófagos alveolares murinos são permissivos à replicação de <i>C. burnetii</i> fase II.....	52
Figura 7 – Localização intracelular de <i>C. burnetii</i> fase II em macrófagos alveolares murinos visualizada por microscopia confocal de fluorescência.....	55
Figura 8 – Localização intracelular de <i>C. burnetii</i> fase II em macrófagos alveolares murinos visualizada por microscopia eletrônica de transmissão.....	56
Figura 9 - Macrófagos alveolares murinos exibem níveis mais elevados de expressão de genes relacionados com o padrão de polarização M2.....	60
Figura 10 - Macrófagos alveolares murinos exibem um fenótipo de polarização M2.....	61
Figura 11 - Macrófagos alveolares murinos possibilitam a utilização de macrófagos deficientes no fundo gênico C57BL/6 para avaliação da resposta imune inata contra <i>C. burnetii</i> fase II.....	64
Figura 12 - Macrófagos alveolares murinos são permissivos à replicação de <i>C. burnetii</i> fase II <i>in vitro</i> de forma semelhante às células Vero e células THP-1.....	66

LISTA DE ABREVIATURAS, SIGLAS, UNIDADES E SÍMBOLOS

AM: *Alveolar macrophage*

Ank: *Ankyrin*

AKT/PKB: *Protein kinase B*

ANOVA: *Análise de variância*

BAL: *Bronchoalveolar lavage*

BCG: *Bacillus Calmette Guerin*

Bcl2: *B-cell lymphoma 2*

Beclin1: *Bcl-2 interacting coiled-coil protein*

BMDMs: *Bone marrow-derived macrophages*

BSA: *Bovine serum albumin*

***C. burnetii*:** *Coxiella burnetii*

Caspase: *Cysteine-dependent aspartate-specific protease*

CCV: *Coxiella-containing vacuole*

CD206: *receptor de manose*

CD369: *dectina-1*

CETEA: *Comitê de Ética e Experimentação Animal*

CO₂: *Dióxido de carbono*

COX2: *Cyclooxygenase 2*

CR3: *Complement receptor 3*

CT: *Cycle Threshold*

DAMPs: *Damage-associated molecular pattern*

DAPI: *4',6'-diamindino-2-phenylindole dilactate*

DIC: *Differential interference contrast*

DMEM: *Dulbecco's modified Eagle's medium*

DNA: *Deoxyribonucleic acid*

cDNA: *Complementary DNA*

Dot: *Defective organelle trafficking*

EDTA: *Ethylene Diamine Tetracetic Acid*

EEA1: *Early endosome antigen 1*

Erk1: *Extracellular signal-Regulated Kinase 1*

Erk2: *Extracellular signal-Regulated Kinase 2*

FACS: *Fluorescence-activated cell sorting*

FIZZ1: *Found in inflammatory zone 1*

FSC: *Forward scatter*

FWD: *Primer forward*

g: *Grama*

Gy: *Gray*

h: *Horas*

IcaA: *Inhibition of Caspase-1 Activation*

Icm: *Intracellular multiplication*

IFN: *Interferon*

IL: *Interleukin*

IL-1R: *Interleukin-1 receptor*

iNOS: *Inducible nitric oxide synthase*

IRS-PCR: *Infrequent restriction site polymerase chain reaction*

kHZ: quilo-Hertz

LAMP-1: *Lysosomal-membrane-associated glicoprotein -1*

LAMP-2: *Lysosomal-membrane-associated glicoprotein -2*

LC3: *Light-chain 3*

LCCM: *L929 Cell Conditioned Medium*

LCV: *Large cell variant*

log: logaritmo na base 10

LPS: *Lipopolysaccharide*

LRV: *Large replicative vacuole*

M: molar

MAMPs: *Microbe-associated molecular patterns*

MAPK: *Mitogen Activated Protein Kinases*

MOI: *Multiplicity of Infection*

M-CSF: *Macrophage colony-stimulating factor*

mg: Miligrama

MHC: *Major histocompatibility complex*

min: minutos

MLVA: *Multiple locus VNTR analyses*

mL: mililitros

mM: Milimolar

MOI: *Multiplicity of infection*

mRNA: RNA mensageiro

MST: *Multispacer sequence typing*

NADPH: *Nicotinamide adenine dinucleotide phosphate*

NF-κB: *Nuclear Factor Kappa B*

ng: *Nanograma*

NLR: *Nucleotide-binding domain leucine-rich repeat-containing protein*

NLRP: *NLR family, Pyrin domain-containing*

nm: *Nanômetro*

NMI: *Nine Mile I*

NMII: *Nine Mile II*

NO: *Nitric oxide*

NOS2: *Nitric oxide synthase 2*

PARP: *Poly ADP-ribose processing*

PBS: *Phosphate-Buffered Saline*

PCR: *Polimerase Chain Reaction*

PFA: *paraformaldehyde*

PMA: *Phorbol 12-myristate 13-acetate*

Pro-IL: *Pro-interleukin*

PRRs: *Pattern Recognition Receptors*

qPCR: *Quantitative Polimerase Chain Reaction*

REV: *Primer reverse*

RFLP: *Restriction fragments length polymorphisms*

RGD: *arginine-glycine-aspartic acid domain*

Rho: *Ras homologous*

RIF: rifampicina

RNA: *Ribonucleic acid*

RNS: *Reactive nitrogen species*

ROS: *Reactive Oxygen Species*

rpm: rotação por minuto

RPMI: *Roswell Park Memorial Institute medium*

rRNA: RNA ribossômico

RT-PCR: Reverse transcriptase *Polimerase Chain Reaction*

SBF: Soro Bovino Fetal

SCV: *Small cell variant*

SNARE: *Soluble NSF-attachment receptor*

SNP: *Single nucleotide polymorphisms*

SSC: *Side scatter*

STAT: *Signal Transducer and Activator of Transcription*

TGF: *Tumor growth factor*

TH1: *Type 1 T helper*

TH2: *Type 2 T helper*

THP-1: *Tohoku Hospital Pediatrics-1*

TLR: *Toll-Like Receptor*

TNF: *Tumor Necrosis Factor*

T4SS: *Type IV Secretion System*

WT: *wild-type*

YM1/Chil3: *chitinase-like protein 3*

µg: Micrograma

µL: Microlitros

µM: Micromolar

µm: Micrômetro

°C: Graus Celsius

SUMÁRIO

1. INTRODUÇÃO.....	14
1.1. <i>Coxiella burnetii</i> – Aspectos Gerais.....	14
1.2. O vaúcolo parasitóforo de <i>Coxiella burnetii</i> e sua interação com vias do hospedeiro.....	21
1.3. Ativação e subversão da resposta imune por <i>Coxiella burnetii</i>	25
1.4. Polarização de macrófagos e monócitos em resposta à <i>Coxiella burnetii</i>	30
1.5. Modelos de estudo das interações patógeno-hospedeiro com infecção por <i>Coxiella burnetii</i>	37
2. OBJETIVOS.....	39
2.1. Objetivo Geral.....	39
2.2. Objetivos Específicos.....	39
3. MATERIAIS E MÉTODOS.....	40
3.1. Animais.....	40
3.2. Obtenção e preparo de <i>C. burnetii</i> para infecção.....	40
3.3. Cultura de células de linhagem de mamíferos.....	40
3.4. Preparo de macrófagos derivados de medula óssea (BMDMs).....	41
3.5. Preparo de macrófagos alveolares (AMs).....	41
3.6. Infecção com <i>Coxiella burnetii</i>	42
3.7. Microscopia de Fluorescência.....	42
3.8. Microscopia Eletrônica de Transmissão.....	43
3.9. Quantificação de <i>Coxiella burnetii</i> <i>in vitro</i>	44
3.10. Análises de RT-PCR.....	44
3.11. Citometria de fluxo para análises fenotípicas.....	45
3.12. Declaração Ética.....	46
3.13. Análises Estatísticas.....	46
4. RESULTADOS.....	47
4.1. Macrófagos derivados de medula óssea de camundongos A/J são mais suscetíveis à infecção por <i>Coxiella burnetii</i> fase II do que aqueles derivados de camundongos C57BL/6.....	47
4.2. Macrófagos alveolares derivados de camundongos C57BL/6 são permissivos à replicação de <i>C. burnetii</i> fase II.....	50

4.3. Macrófagos alveolares murinos suportam a replicação de grandes quantidades de bactéria intracelular nos vacúolos contendo <i>C. burnetii</i> fase II.....	53
4.4. Macrófagos alveolares murinos mostram uma polarização mais pronunciada em células do tipo M2 do que macrófagos murinos derivados de medula óssea.....	57
4.5. Macrófagos alveolares murinos deficientes em NOS2 e IFN γ são mais suscetíveis à replicação de <i>C. burnetii</i> fase II, enquanto células deficientes em interleucina-4 (IL-4) são mais restritivas à replicação da bactéria.....	62
4.6. A permissividade dos macrófagos alveolares murinos à infecção por <i>C. burnetii</i> fase II é comparável a aquelas de linhagens celulares imortalizadas altamente suscetíveis.....	65
5. DISCUSSÃO.....	67
6. SUMÁRIO.....	76
7. CONCLUSÃO.....	78
8. REFERÊNCIAS BIBLIOGRÁFICAS.....	79
9. ANEXO I – TRABALHO DE PRIMEIRA AUTORIA.....	92
9.1. Anexo 1 - Artigo publicado (FERNANDES, Talita D et al. <i>Infection and Immunity</i> , 2016).....	92
10. ANEXO II – TRABALHOS DE COLABORAÇÃO.....	103
10.1. Anexo 2 - Artigo publicado (CUNHA, Larissa D; RIBEIRO, Juliana M; FERNANDES, Talita D; MASSIS, Liliana M; KHOO, Chen A; MOFFATT, Jennifer H; NEWTON, Hayley J; ROY, Craig R; ZAMBONI, Dario S. <i>Nature Communications</i> , 2015).....	103
10.2. Anexo 3 - Artigo publicado (MASCARENHAS, Danielle PA; CERQUEIRA, Daiane M; PEREIRA, Marcelo SF; CASTANHEIRA, Fernanda VS; FERNANDES, Talita D; MANIN, Grazielle Z; CUNHA, Larissa D; ZAMBONI, Dario S. <i>Pathogens</i> , 2017).....	126

1. INTRODUÇÃO

1.1. *Coxiella burnetii* – Aspectos Gerais

Coxiella burnetii é uma gama-proteobactéria patogênica gram-negativa intracelular não flagelada que sobrevive no interior de um vacúolo parasitóforo denominado CCV (do inglês, *Coxiella-Containing Vacuole*) ou LRV (do inglês, *Large Replicative Vacuole*) o qual apresenta características fagolisossomais que o tornam único (HACKSTADT; WILLIAMS 1981; WEISBURG et al., 1989; VOTH; HEINZEN 2007; CAPO; MEGE 2012; GHIGO; COLOMBO; HEINZEN, 2012; ANDERSON et al., 2013). Essa bactéria é o agente causador de uma infecção zoonótica denominada Febre Q, que afeta principalmente animais de pastoreio e gado. No entanto, seu reservatório natural é extenso e inclui artrópodes como carrapatos, além de aves, répteis e mamíferos selvagens e domésticos (MAURIN; RAOULT 1999; RAOULT et al., 2005; HECHERY, 2012; ANDERSON, et al., 2013; ELDIN et al., 2017). A maioria dos animais não desenvolve sintomas aparentes da doença, exceto cabras e ovelhas, as quais apresentam alta proliferação bacteriana no trato reprodutivo feminino, frequentemente levando a abortos tardios e perdas econômicas consideráveis para os produtores (HECHERY, 2012).

O primeiro surto envolvendo essa bactéria até então desconhecida ocorreu em 1935 em Queensland, Austrália. Nesta ocasião, alguns trabalhadores de um abatedouro em Brisbane foram infectados e começaram a apresentar sintomas de uma doença febril (DERRICK, 1937). Em 1937, Derrick injetou o sangue dos pacientes infectados em porquinhos-da-índia e conseguiu transferir com sucesso a doença, entretanto sem conseguir visualizar seu agente causador. Amostras de tecidos infectados foram então enviadas para o pesquisador Frank Macfarlane Burnet que, juntamente com Mavis Freeman, reproduziram as reações febris características nos porquinhos-da-índia. Eles observaram que os tecidos infectados apresentavam organismos dificilmente filtráveis que aparentavam possuir uma natureza “*rickettsial*” (BURNET; FREEMAN, 1937; CAPO; MEGE 2012). Testes realizados com o soro de pacientes que tiveram Febre Q e suspensões de tecidos clarificados de cobaias infectadas mostraram a ocorrência de aglutinação. Esse fato ocorria quando era utilizado o soro de pacientes que apresentavam tifo por *Rickettsia* (BURNET; FREEMAN, 1937; CAPO; MEGE 2012).

Quase simultaneamente, descobriu-se um agente desconhecido em Hamilton, Montana, nos Estados Unidos, durante estudos de campo sobre a ecologia da febre maculosa das montanhas rochosas. Essa descoberta envolveu a utilização de carrapatos *Dermacentor andersoni* coletados em Nine Mile, Montana, os quais foram alimentados com porquinhos-da Índia. Um desses animais desenvolveu uma doença febril com características divergentes daquelas comumente apresentadas pela febre maculosa das montanhas rochosas (DAVIS; COX 1938; CAPO; MEGE 2012). Chamado de “agente *Nine Mile*”, esse organismo podia ser transferido serialmente em cobaias, era filtrável e não apresentava crescimento em meio de cultura axênico, o que levou os pesquisadores Davis e Cox (1938) a questionarem sua “natureza bacteriana”. A ligação entre o agente *Nine Mile* e o causador da Febre Q foi descoberta mais tarde. Um paciente infectou-se acidentalmente em laboratório pelo agente *Nine Mile* e desenvolveu uma febre com de sintomatologia muito semelhante à da Febre Q. Estudos complementares de proteção cruzada desenvolvidos posteriormente demonstraram que ambos os agentes eram o mesmo patógeno (DYER, 1938; DYER, 1939; CAPO; MEGE 2012).

Por não se comportar como uma *Rickettsia* típica, esse organismo foi inicialmente colocado em um novo gênero na família *Rickettsiaceae*, o gênero “*Coxiella*” e espécie “*burnetii*” em homenagem a Cox e Burnet, respectivamente. Mais tarde, através de estudos com taxonomia molecular envolvendo análises da sequência do gene codificante para 16S rRNA, *C. burnetii* foi reclassificada para o filo *Proteobacteria*, classe γ -*Proteobacteria*, ordem *Legionellales* e família *Coxiellaceae*, sendo próxima de *Rickettsiella grylli*, *Legionella pneumophila* e *Francisella tularensis* (WEISBURG et al., 1989; SAUNDERS et al., 1993; ROUX et al., 1997; WAAG; THOMPSON, 2005; RAOULT et al., 2009; CAPO; HECHERY, 2012; MEGE, 2012; VAN SCHAIK; SAMUEL, 2012; ELDIN et al., 2017). De fato, o genoma de *C. burnetii* contém uma porcentagem das bases nitrogenadas guanossina e citosina mais próxima do que é encontrado em membros da ordem *Legionellales* do que membros da ordem *Rickettsiales*, corroborando sua reclassificação na nova ordem (HECHERY, 2012).

Quando analisada apenas a sequência do gene 16S rRNA, *C. burnetii* apresenta uma homogeneidade genética considerável (STEIN; RAOULT, 1993). Todavia, comparações mais amplas de seu genoma por meio da utilização de técnicas como análises de polimorfismos de fragmentos de restrição com diferentes comprimentos (do inglês, *Restriction Fragment Length Polymorphism* - RFLP) e análises de comparação de *microarray* revelaram

heterogeneidade considerável no padrão de bandas e nos padrões de hibridização. Essas análises indicaram a existência de diversidade genética entre isolados (HENDRIX et al., 1991; BEARE et al., 2006; HECHERY, 2012). De fato, a combinação dessas técnicas de análise resultou na categorização inicial de trinta e dois isolados bacterianos em seis grupos genômicos distintos (I-VI) (HENDRIX et al., 1991).

A diversidade genética de *C. burnetii* é também evidenciada pela presença de moléculas de lipopolissacarídeo (LPS) antigênicas e estruturalmente únicas entre diferentes isolados. Foram descritos três quimiotipos de LPS que são associados com grupos genômicos específicos, atuando como biomarcadores únicos de *C. burnetii* (TOMAN; SKULTETY; IHNATKO, 2009; HECHERY, 2012; SCHAIK; SAMUEL, 2012). Com o desenvolvimento de métodos de cultivo de *C. burnetii* em meio axênico) tornou-se possível analisar um número cada vez maior de isolados bacterianos e até mesmo traçar suas origens evolutivas a partir de diferentes técnicas de genotipagem e sequenciamento, como: a) genotipagem de múltiplos espaçadores (do inglês, MST – *Multispacer Sequence Typing*); b) análise da variação do número de repetições em *tandem* em múltiplos *locus* (do inglês, MLVA – *Multiple Locus VNTR Analysis*); c) reação em cadeia da polimerase (do inglês, PCR – *Polymerase Chain Reaction*) baseada em sítios de restrição pouco frequentes (do inglês, IRS-PCR – *Infrequent Restriction Site-PCR*) e d) sequenciamento de polimorfismos de nucleotídeos únicos (do inglês, SNP – *Single Nucleotide Polymorphisms*) (GLAZUNOVA et al., 2005; ARRICAUBOUVEREY et al., 2006; OMSLAND et al., 2009; HORNSTRA et al., 2011; OMSLAND et al., 2011; MCLAUGHLIN et al., 2017).

Apesar de ser uma doença de caráter zoonótico, a infecção de seres humanos por *C. burnetii* é recorrente. Ela ocorre via aerossol, por meio da inalação de partículas de ar contaminadas, ou através do contato com fluidos e subprodutos animais infectados, frequentemente provenientes de ruminantes. Os principais alvos celulares primários da infecção são monócitos e macrófagos, especialmente macrófagos alveolares (HACKSTADT; WILLIAMS, 1981; KHAVIN; TABIBZADEH, 1988; MAURIN; RAOULT 1999; SHANNON; HEINZEN 2008; CAPO; MEGE 2012). Nos humanos, a infecção com *C. burnetii* pode resultar em soroconversão assintomática ou no desenvolvimento da Febre Q sintomática. Em sua fase aguda, apresenta manifestações clínicas variadas com características semelhantes a de uma gripe, incluindo pneumonia, febre, dores de cabeça, calafrios, fadiga e hepatite granulomatosa (RAOULT; MARRIE; MEGE, 2005; CAPO; MEGE 2012; HECHERY, 2012). A falta de sintomas específicos na fase aguda da infecção, as variações

nos períodos de incubação da bactéria e as dificuldades do emprego de metodologias de diagnóstico específicas para detecção de *C. burnetii* em muitos hospitais sugerem que a mesma é, provavelmente, subdiagnosticada. Mais da metade dos pacientes infectados não exibe nenhum sintoma, sendo a infecção aguda autolimitante. Ainda assim, um estudo feito no Brasil por Costa e colaboradores (2006) sugere que cerca de 4,5% dos casos de febres do tipo *influenza*, febres de origem desconhecida e pneumonias atípicas sejam causados por *C. burnetii*.

Apesar da maior parte dos casos de infecção ser assintomática e se resolver sem tratamento específico com antibióticos, cerca de 2-5% desses casos ou de casos de Febre Q aguda evoluem para a Febre Q crônica, forma mais séria da doença. Suas manifestações clínicas mais sérias incluem endocardite, apresentando fibrose, calcificação, inflamação leve e vascularização, frequentemente levando o paciente à hospitalização. Na ausência do tratamento adequado, pode resultar na morte do paciente (RAOULT, 1990; BACA; LI; KUMAR, 1994; MAURIN; RAOULT, 1999; RAOULT; MARRIE; MEGE, 2005; CAPO; MEGE 2012; HECHERY, 2012).

Dados epidemiológicos mostram que gênero masculino, idade superior a 15 anos, imunocomprometimento e doenças cardíacas valvulares pré-existentes compreendem fatores de risco importantes para o desenvolvimento da Febre Q (CAPO; MEGE 2012; HECHERY, 2012). Além disso, mesmo em pacientes que apresentam aparente cura, a *C. burnetii* persiste em seus organismos por um longo período de tempo, impactando na saúde a longo prazo. Mais de um terço dos pacientes continuam a apresentar estado de saúde prejudicado mesmo após 24 meses do diagnóstico da infecção (HARRIS et al., 2000; VAN DER VELDEN et al., 2015).

Análises epidemiológicas mostraram que a infecção com *C. burnetii* apresenta distribuição mundial, com exceção da Nova Zelândia (MAURIN; RAOULT, 1999; RAOULT; MARRIE; MEGE, 2005; MARRIE, 2010; ELDIN et al., 2017). Além disso, por ser uma doença zoonótica, a epidemiologia humana da Febre Q reflete a circulação da bactéria nos reservatórios animais. No mundo, situações endêmicas ou surtos são observados dependendo da área geográfica. Comumente, áreas de endemicidade como França, Espanha e Estados Unidos apresentam casos esporádicos de Febre Q. Esses casos são observados após a realização de atividades de alto risco por seres humanos, como trabalhos em fazenda, matadouros e áreas rurais em geral. Pequenos surtos também podem ocorrer após exposição a fontes comuns de infecção, principalmente em ambiente familiar (ELDIN et al., 2017).

Existem áreas de hiperendemicidade, como a África, nas quais a transmissão da doença é facilitada devido a uma alta concentração de lares muito próximos a grandes populações de ruminantes domésticos. O consumo de leite cru não pasteurizado nessas áreas também pode atuar como fonte de contaminação humana por *C. burnetii*. Um importante surto de larga escala foi observado na Holanda entre os anos 2007 e 2010, onde mais de 4.000 novos casos foram noticiados. Esse surto apresentou correlação com o aumento da importação de cabras no país entre os anos 1985 e 2009 para fortalecimento das atividades de criação de laticínios provenientes desses animais. A introdução de animais contaminados com *C. burnetii* em regiões próximas de áreas urbanas pode ter favorecido o desenvolvimento do surto (DELSING; KULLBERG; BLEEKER-ROVERS, 2010; SCHNEEBERGER et al., 2012).

Além de sua distribuição mundial, *C. burnetii* apresenta alta estabilidade no ambiente e é altamente virulenta, sendo necessário uma dose infecciosa muito baixa (<10 organismos) para causar a doença. À partir das características apresentadas e da capacidade de aerosolização do patógeno, *C. burnetii* foi classificada pelo Centro para Controle e Prevenção de Doenças dos Estados Unidos como agente potencial de bioterrorismo da categoria B (MARRIE, 2004; ELDIN et al., 2017). Com isso, reforça-se a importância de se estudar essa bactéria e todos os aspectos envolvidos na infecção por esse patógeno.

A alta estabilidade de *C. burnetii* no ambiente deve-se a particularidades existentes em seu ciclo de vida de caráter bifásico. Ele envolve a participação de duas variantes celulares denominadas de variante celular pequena – SCV (*Small Cell Variant*) e variante celular grande – LCV (*Large Cell Variant*). Essas distintas formas celulares estão diretamente relacionadas com os diferentes ambientes que a bactéria enfrenta, sendo eles o meio extracelular (para SCV) e o meio intracelular (principalmente para LCV). Quando no meio externo, *C. burnetii* encontra-se na forma de cocabacilo com cerca de 0,2-0,5 µm de comprimento. Quando vista ao microscópio, a SCV apresenta-se altamente elétron-densa devido ao estado extremamente condensado de sua cromatina. Essa condensação relaciona-se com o fato de que esse morfotipo celular é metabolicamente inativo e se assemelha a uma forma de resistência, tendo sido inicialmente confundido com um esporo. A SCV, considerada a forma infectiva da bactéria, é resistente à dissecação, altas temperaturas, choques osmóticos, luz ultravioleta e agentes químicos desinfetantes, permitindo que essa forma celular permaneça infecciosa no ambiente por meses (MCCAUL; WILLIAMS, 1981; WILLIAMS, 1991; COLEMAN et al., 2004; TISSOT-DUPONT et al., 2004; MINNICK;

RAGHAVAN, 2012; VAN SCHAIK; SAMUEL 2012; SANDOZ et al., 2014; ELDIN et al., 2017).

Ao entrar em contato com seu hospedeiro, a SCV é fagocitada, inicia-se uma série de mudanças conformacionais e metabólicas que acompanham desde a maturação do endossomo até o estabelecimento do vacúolo parasitóforo contendo *C. burnetii*. Essas mudanças são acompanhadas de uma transição da SCV para a LCV, forma replicativa bacteriana com cerca de 1 µm de comprimento. A duração desse processo é de aproximadamente dois dias e caracteriza a fase *lag* do crescimento bacteriano, em que não há replicação (COLEMAN et al., 2004). A transição entre a variante SCV, essencialmente dormente, para a LCV, pleiomórfica e metabolicamente ativa, ocorre em resposta à acidificação do CCV e, provavelmente, a outros sinais como disponibilidade de nutrientes na célula hospedeira e ações enzimáticas (HACKSTADT; WILLIAMS, 1981; SESHADRI; HENDRIX; SAMUEL, 1999; MINNICK; RAGHAVAN 2012; VAN SCHAIK; SAMUEL, 2012; SANDOZ et al., 2014). ScyA é uma proteína de ligação ao DNA de caráter altamente básico que, provavelmente, desempenha um papel na condensação da cromatina. A perda da sua atividade interfere com a manutenção do estado metabolicamente inativo da variante SCV e é um dos marcos que sinalizam a transição da SCV para LCV (COLEMAN et al., 2004; VAN SCHAIK; SAMUEL, 2012). Uma vez que essa transição é finalizada inicia-se a fase de crescimento exponencial da bactéria, denominada fase *log*. Nessa etapa, as LCV irão replicar por aproximadamente quatro dias até atingir a fase *log* tardia ou o início da fase estacionária. Nesse momento, observa-se o reaparecimento da proteína ScyA e uma nova transição da forma LCV para SCV de *C. burnetii*, com a lise das células infectadas e a liberação de novas SCV que infectarão outras células. Com isso, finaliza-se um ciclo de vida bifásico dessa bactéria e inicia-se o próximo (Figura 1) (COLEMAN et al., 2004; MINNICK; RAGHAVAN 2012; VAN SCHAIK; SAMUEL, 2012; SANDOZ et al., 2014).

Figura 1 – Representação do ciclo de vida bifásico de *C. burnetii*. Após a internalização da SCV de *C. burnetii* pela célula hospedeira, ocorre o desenvolvimento de um fagossomo, seguido do recrutamento e fusão de lisossomos (círculos amarelos) para esse compartimento. O vacúolo parasitóforo sofre acidificação, levando à transição da SCV para LCV, finalizando a fase *lag* do ciclo com duração de dois dias. Posteriormente, ocorre a replicação exponencial das LCVs, caracterizando a fase *log* do ciclo, com duração aproximada de quatro dias. No período tardio dessa fase inicia-se novamente a transição da LCV para SCV, iniciando a fase estacionária do ciclo, até a lise da célula hospedeira e liberação das SCVs que infectarão outras células. Adaptado de: Minninck; Raghavan (2012).

1.2. O vacúolo parasitóforo de *Coxiella burnetii* e sua interação com vias do hospedeiro

Para a sobrevivência e replicação de *C. burnetii* no interior da célula hospedeira é essencial que ela estabeleça seu vacúolo parasitóforo e o mantenha durante todo o seu desenvolvimento. Além disso, macrófagos e monócitos são as principais células-alvo da infecção por *C. burnetii* e desempenham papéis importantes em processos biológicos como processamento e apresentação de antígenos, produção de citocinas e eliminação de patógenos. Portanto, é fundamental que a bactéria evite ou subverta suas respostas (AMARA; BECHAH; MEGE, 2012). Ambos os processos dependem da participação ativa da bactéria, em parte por meio da utilização de seu sistema de secreção tipo IVB, homólogo ao sistema de secreção Dot/Icm da bactéria *Legionella pneumophila* (SESHADRI et al., 2003; ZAMBONI et al., 2003; ZUSMAN; YERUSHALMI; SEGAL, 2003). Esse sistema de secreção é montado em resposta à acidificação do ambiente de uma organela endocítica tardia em que a bactéria se encontra. Ele facilita a translocação de mais de 130 proteínas efetoras no citoplasma da célula hospedeira que, conjuntamente, colaboram para modulação do vacúolo contendo *C. burnetii* e para a modulação das repostas imunes do hospedeiro, fornecendo um ambiente favorável à replicação bacteriana (HOWE et al., 2003; NEWTON; MCDONOUGH; ROY, 2013; MOFFATT; NEWTON P.; NEWTON H.J., 2015).

O processo de estabelecimento do vacúolo parasitóforo contendo *C. burnetii* começa com sua internalização pela célula hospedeira. Após a descoberta do sistema de secreção tipo IV (T4SS) nessas bactérias sugeriu-se a utilização de mecanismos ativos para indução da fagocitose após ligação inicial à célula hospedeira. No entanto, estudos realizados com bactérias mutantes para T4SS demonstraram que não houve interferência na habilidade dessas bactérias de invadir células fagocíticas e não fagocíticas (BEARE et al., 2011). Esses dados corroboram a hipótese de que o processo de entrada de *C. burnetii* nas células hospedeiras ocorre, na realidade, de forma passiva, envolvendo o remodelamento de filamentos de actina em um mecanismo de zíper (BACA; KLASSEN; ARAGON, 1993; TUJULIN et al., 1998; BEARE et al., 2011; CAREY et al., 2011; VAN SCHAIK et al., 2013). De fato, o tratamento das células hospedeiras com agentes despolimerizantes de actina, como citocalasina D, inibem a internalização do patógeno, ressaltando a importância dos filamentos de actina para entrada da bactéria (BACA; KLASSEN; ARAGON, 1993).

Após a entrada na célula hospedeira, a bactéria trafega passivamente pela via endocítica, participando de diferentes etapas de maturação, antes de atingir o ambiente tipicamente acidificado e hidrolítico dos lisossomos. Diferentemente de outros patógenos que

são comumente degradados neste ambiente, *C. burnetii* é capaz de sobreviver e continuar seu ciclo de desenvolvimento (HACKSTADT; WILLIAMS, 1981; HEINZEN et al., 1996; VOTH; HEINZEN, 2007).

O início do processo de maturação do fagossomo contendo *C. burnetii* envolve fissão e fusão com vários endossomos iniciais, apresentando diferentes marcadores de membrana e alterações de pH do lúmen. Após cinco minutos de internalização bacteriana é possível detectar marcadores como o Antígeno-1 de Endossomos Iniciais (EEA1, do inglês *Early Endosome Antigen-1*) e a GTPase Rab5, além de alteração na acidificação do lúmen para pH 5,4, aproximadamente. Diferentemente do que ocorre para a maioria dos patógenos, foi observado que *C. burnetii* interage com a via de autofagia. Essa interação foi identificada a partir da presença da proteína de cadeia leve 3 associada a microtúbulo (LC3, do inglês *Light-Chain 3*), marcadora de autofagia, na membrana fagossomal (BERON et al., 2002; ROMANO et al., 2007; VAN SCHAIK et al., 2013).

Os fagossomos iniciais continuam trafegando pela via endocítica até completar 60 minutos, aproximadamente, após a entrada na célula hospedeira. A partir desse momento, há a substituição gradual dos marcadores de endossomos iniciais por marcadores de endossomos tardios, como a GTPase Rab7, a proteína 1 associada à membrana lisossomal (LAMP-1, do inglês *Lysosomal Membrane-Associated Protein-1*). Os marcadores LAMP-2 e ATPase vacuolar também estão presentes e são responsáveis por reduzir o pH do vacúolo para valores próximos a 5, através do bombeamento de prótons no fagossomo em maturação (SCOTT; BOTELHO; GRINSTEIN, 2003; HENRY et al., 2004; ROMANO et al., 2007; GHIGO et al., 2009; GHIGO; COLOMBO; HEINZEN, 2012). Posteriormente ocorre a fusão dos fagossomos com lisossomos, formando os chamados fagolisossomos. Nesse evento, hidrolases lisossomais e enzimas como fosfatase ácida e catepsina D são recrutadas em um processo que é ativamente atrasado pela bactéria. Essa modulação da maturação final do vacúolo por *C. burnetii* pode estar relacionada com novas interações com as vias autofágicas da célula. Desse modo, vesículas autofágicas cheias de nutrientes e membranas provenientes de suas cargas degradadas se fundem ao vacúolo em maturação, favorecendo assim a posterior replicação do patógeno (HOWE et al., 1993; HOWE; MALLAVIA, 2000; GUTIERREZ et al., 2005; ROMANO et al., 2007; VOTH; HEINZEN, 2007; VAN SCHAIK; SAMUEL, 2012; VAN SCHAIK et al., 2013).

Em seguida, há um aumento drástico no volume do vacúolo bacteriano, resultando no deslocamento das organelas da célula hospedeira para a periferia, ocupando quase todo o

volume citoplasmático disponível. Essa expansão ocorre por meio de fusão homotípica entre múltiplos vacúolos de menor tamanho contendo *C. burnetii*, um processo que requer a participação de proteínas do hospedeiro. Entre elas, encontram-se as proteínas que mediam diretamente eventos de fusão de vesículas (SNAREs, do inglês *Soluble NSF-attachment receptor*), bem como a participação de efetores bacterianos translocados pelo sistema de secreção. Dentre as proteínas SNAREs potencialmente envolvidas no processo de fusão homotípica, foram identificadas VAMP-3, VAMP-7 e VAMP-8, bem como a proteína Syntaxin-7, a qual se assemelha a uma proteína SNARE (ITAKURA; KISHI-ITAKURA; MIZUSHIMA, 2012; CAMPOY; MANSILLA; COLOMBO, 2013; HAMASAKI et al., 2013; MCDONOUGH et al., 2013; NEWTON et al., 2014; KOHLER; ROY, 2015; PECHSTEIN; SCHULZE-LUEHRMANN; LÜHRMANN, 2018). Com relação aos efetores bacterianos envolvidos nesse processo, CvpB, também conhecida como Cig2, é essencial para fusão homotípica dos vacúolos. Foi demonstrado que a interferência nesse efector através do uso de transposons resultou em múltiplos vacúolos contendo *C. burnetii* por célula sem capacidade de fusão, impossibilitando a formação do vacúolo único característico (NEWTON; MCDONOUGH; ROY, 2013; NEWTON et al., 2014; WEBER; FARIS, 2018). A expansão do vacúolo continua por meio de eventos de fusão heterotípica com vesículas lisossomais, autofágicas, endocíticas e exocíticas (HOWE et al., 2003a; HOWE et al., 2003b; CAMPOY; ZOPPINO; COLOMBO, 2011).

A manutenção do vacúolo parasitóforo depende da participação do citoesqueleto de actina, da produção de proteínas pela bactéria e da ativação de proteínas quinase do hospedeiro como proteína quinase da cadeia leve de miosina, proteína quinase A e proteína quinase C (AGUILERA et al., 2009; HUSSAIN et al., 2010; VAN SCHAIK et al., 2013). Além disso, o acúmulo de RAB1B e RhoGTPase na membrana do vacúolo demonstra a interação com o retículo endoplasmático e as vias secretórias iniciais, sugerindo o fornecimento de membrana para o vacúolo em expansão (CAMPOY; ZOPPINO, COLOMBO, 2011; VAN SCHAIK; SAMUEL, 2012). Efetores bacterianos como CvpA e Cig57 estão envolvidos com a subversão do tráfego de vesículas mediado por clatrina, facilitando a biogênese do vacúolo fusogênico (LARSON et al., 2013; LATOMANSKI et al., 2016; PECHSTEIN; SCHULZE-LUEHRMANN; LÜHRMANN, 2018).

A membrana do vacúolo parasitóforo contendo *C. burnetii* apresenta enriquecimento de colesterol, fato atípico para membranas de compartimentos com características lisossomais. A infecção com essa bactéria induz o aumento na expressão de genes envolvidos

com a produção de colesterol, além da bactéria expressar uma enzima estero redutase que pode afetar o metabolismo de colesterol do hospedeiro (COLEMAN et al., 2004; HOWE; HEINZEN, 2006; GILK; BEARE; HEINZEN, 2010; KOHLER; ROY, 2015). Essas interações bacterianas com diferentes vias do hospedeiro requerem que a produção de proteínas por *C. burnetii* esteja intacta, sugerindo a possível participação de vários outros efetores bacterianos nesses processos (ZAMBONI et al., 2003; CAMPOY; ZOPPINO, COLOMBO, 2011).

C. burnetii é capaz de intervir nas vias apoptóticas da célula hospedeira, inibindo a apoptose celular. Isso se dá pela interferência em diferentes etapas da via de apoptose, por exemplo: a) antagonização do processamento de caspase-3, caspase-9 e PARP (*Poly ADP-Ribose Processing*); b) alteração dos níveis de expressão de mais de 30 genes pró-sobrevivência, e c) ativação sustentada do fator de transcrição NF- κ B (VOTH; HOWE; HEINZEN, 2007a; VOTH; HOWE; HEINZEN, 2007b; HUSSAIN; VOTH, 2012). Outra interação utilizada pela bactéria com as vias de apoptose inclui o recrutamento do complexo anti-apoptótico formado por Beclin-1/Bcl-2 para a membrana do vacúolo parasitóforo, possivelmente por meio de prevenção da liberação de citocromo C das mitocôndrias, ação conhecida de Bcl-2 (YANG et al., 1997; LUHRMANN; ROY, 2007). Esse complexo Beclin-1/Bcl-2 também participa da regulação da autofagia. A modulação positiva de vias de sinalização pró-sobrevivência mediadas por Akt e Erk1/2 pela bactéria, com a ativação sustentada dessas proteínas, contribui para inibição da apoptose (GUTIERREZ et al., 2005; LUHRMANN; ROY, 2007; ROMANO et al., 2007; VOTH et al., 2007; VOTH; HEINZEN, 2009; VAZQUEZ, COLOMBO, 2010; VAN SCHAİK; SAMUEL, 2012).

Luhrmann e colaboradores (2010) demonstraram que AnkG é um dos efetores bacterianos envolvidos na indução do estado anti-apoptótico de células infectadas com *C. burnetii*, por meio de sua interação com a proteína p32 do hospedeiro. O efector bacteriano CaeB localiza-se na membrana da mitocôndria e previne a liberação de proteínas pró-apoptóticas através da inibição da permeabilização dessa membrana, enquanto CaeA, outra proteína efectora bacteriana, apresenta localização nuclear e inibe a apoptose por mecanismos ainda não esclarecidos (LUHRMANN et al., 2010; KLINGENBECK et al., 2013; VAN SCHAİK et al., 2013). Em conjunto, esses fatores demonstram que o vacúolo parasitóforo maduro contendo *C. burnetii* metabolicamente ativa não é estático e que a bactéria é capaz de modular e interagir ativamente com diversas vias do hospedeiro, garantindo dessa forma a manutenção de seu nicho replicativo (Figura 2) (HEINZEN et al., 1996; HOWE; HEINZEN,

2006; CAMPOY; ZOPINO; COLOMBO, 2011; VAN SCHAİK et al., 2013; KOHLER; ROY, 2015)

Figura 2 - Processo de desenvolvimento e maturação do vacúolo contendo *C. burnetii*. Após a internalização da SCV de *C. burnetii* pela célula hospedeira, em um processo que envolve a remodelação de filamentos de actina e interação com os receptores $\alpha_v\beta_3$ integrina, ocorre a formação de um fagossomo que trafega pela via endocítica. Durante esse tráfego ocorre a interação com endossomos e lisossomos, marcada pela aquisição de marcadores como LC3, RAB5, EEA1, LAMP1 e RAB7, juntamente com uma diminuição de pH, até a maturação final do vacúolo contendo *C. burnetii* que apresenta características fagolisossomais. Adaptado de: Van Schaik e colaboradores (2013).

1.3. Ativação e subversão da resposta imune por *Coxiella burnetii*

Além de interagir com diferentes vias e componentes das células hospedeiras a fim de assegurar o estabelecimento e manutenção de seu vacúolo replicativo, *C. burnetii* precisa garantir que o sistema imune dos hospedeiros não seja capaz de detectá-la. Com isso, é possível montar uma resposta efetiva contra o patógeno. Além do papel do sistema de

secreção do tipo IVB e das interações da bactéria com vias de autofagia e apoptose, *C. burnetii* é capaz de subverter a atividade microbicida de monócitos e macrófagos, principais células alvo da infecção por esse patógeno. Essa subversão ocorre por meio de diversos mecanismos, os quais podem ser diretos ou indiretos. Como mecanismo direto tem-se a ação dos efetores bacterianos, por exemplo. Já como mecanismos indiretos têm-se as alterações no estado de polarização dos macrófagos ou a interferência com citocinas regulatórias (AMARA; BECHAH; MEGE, 2012).

Uma das estratégias empregadas por essa bactéria consiste na subversão da fagocitose, processo que apresenta diferenças entre a infecção com a forma virulenta ou avirulenta de *C. burnetii* (CAPO et al., 1999). A diversidade genética e o potencial de virulência de *C. burnetii* reside principalmente nas diversas formas de LPS expressos. Bactérias do isolado RSA493, derivadas da cepa Nine Mile, são chamadas de “fase I” ou virulentas quando apresentam LPS do tipo liso (com uma cadeia de polissacarídeos com o antígeno-O completo e presente, bem como resíduos frequentes dos açúcares virenose e diidroxistrepnose) (NARASAKI; TOMAN, 2012). Já a variante Nine Mile “fase II”, conhecida como a forma avirulenta para mamíferos, é obtida por meio de passagens seriais da forma virulenta em hospedeiros não imunocompetentes. Com isso, ocorrem modificações irreversíveis na estrutura do LPS, com perda do antígeno-O, podendo ou não apresentar deleções genômicas adicionais. Como resultado, o LPS das bactérias fase II apresenta um fenótipo rugoso. Diferentemente, o lipídeo A presente no LPS das formas virulentas e avirulentas de *C. burnetii* apresenta as mesmas espécies iônicas e perfis de fragmentação iguais, sugerindo uma estrutura similar entre eles (VISHWANATH; HACKSTADT, 1988; TOMAN; SKULTETY, 1996; MAURIN; RAOULT, 1999; ZAMBONI et al., 2004; TOMAN; SULTETY; IHNATKO, 2009; NARASAKI; TOMAN, 2012; VAN SCHAIK et al., 2013; ELDIN et al., 2017).

A subversão da fagocitose por *C. burnetii* difere entre a fase I e II, pois os receptores envolvidos nesse processo variam de acordo com o tipo de LPS apresentado (WILLIAMS; PEACOCK; MCCAUL, 1981; HACKSTADT, 1988; AMARA; BECHAH; MEGE, 2012; VAN SCHAIK et al., 2013; LARSON et al., 2016). A fagocitose das formas avirulentas da bactéria depende da participação de receptores de integrina $\alpha V\beta 3$ e também de receptores CR3 (do inglês, *Complement Receptor 3*). Dessa forma, ocorre uma alta taxa de internalização, o que pode interferir com a sobrevivência da bactéria. A maior exposição de proteínas e a hidrofobicidade aumentada desse tipo de LPS devido à falta do antígeno-O provavelmente facilitam as interações entre a membrana plasmática do hospedeiro e da

bactéria, o que explicaria essa maior facilidade para fagocitose da fase II. Já as formas virulentas de *C. burnetii* desacoplam o receptor CR3 do processo, interagindo apenas com as integrinas $\alpha V\beta 3$ e levando a uma taxa baixa de fagocitose e sobrevivência intracelular. É possível que a interação de ambas as formas da bactéria com as $\alpha V\beta 3$ integrinas ocorra por meio de proteínas associadas a membranas que apresentam o domínio de ligação a integrinas composto por arginina-glicina-ácido aspártico (domínio RGD) ou proteínas com domínios anquirina (CAPO et al., 1999; SESHADRI et al., 2003; VAN SCHAIK et al., 2013; CAPO; MEGE, 2012; ELDIN et al., 2017). É importante ressaltar que no caso da fagocitose das formas virulentas ocorre também a indução de protusões na membrana para restringir ainda mais o acesso aos receptores CR3. Esse processo está relacionado com o forte remodelamento do citoesqueleto de actina, além da ativação de vias de proteínas tirosina quinase, produção deficiente de citocinas como IL-12 e TNF, e inibição da maturação de células dendríticas (BACA; KLASSEN; ARAGON, 1993; MECONI et al., 1998; CAPO et al., 1999; DELLACASAGRANDE et al., 2000; MECONI et al., 2001; SHANNON; HOWE; HEINZEN, 2005; CAPO; MEGE, 2012; VAN SCHAIK; SAMUEL, 2012; VAN SCHAIK et al., 2013; LARSON et al., 2016; ELDIN et al., 2017).

O remodelamento do citoesqueleto de actina induzido pela infecção de macrófagos com *C. burnetii* fase I também conta com a participação de receptores do tipo *Toll* (TLRs, do inglês *Toll-Like Receptors*). Esse processo ocorre mesmo com a bactéria sendo silenciosa e altamente adaptada à evasão do sistema imune, incluindo evasão do reconhecimento pelo sistema complemento e do reconhecimento e desencadeamento de resposta imune mediada pelos TLRs (VISHWANATH; HACKSTADT; 1988; ZAMBONI, 2004; ZAMBONI et al., 2004; VAN SCHAIK et al., 2013). Os TLRs são uma classe de receptores da imunidade inata denominados PRRs (do inglês, *Pattern Recognition Receptors*), identificados pela primeira vez em *Drosophila melanogaster* devido ao seu papel no controle do desenvolvimento dorsoventral desse inseto (LEMAITRE et al., 1996). Apresentam uma estrutura básica composta por um domínio N-terminal extracelular de reconhecimento rico em repetições de leucina (do inglês, *Leucine-Rich Repeat – LRR*), um domínio transmembrana e um domínio C-terminal intracelular de sinalização (MEDZHITOV; PRESTON-HULBURT; JANEWAY, 1997; MEDZHITOV, 2001). Essas proteínas reconhecem, de forma geral, padrões moleculares associados a micróbios (MAMPs, do inglês *Microbe-Associated Molecular Patterns*) e padrões moleculares associados a danos (DAMPs, do inglês *Damage-Associated Molecular Patterns*) provenientes de células que estão morrendo ou que apresentam algum

dano (JANEWAY, 1989; MEDZHITOV; JANEWAY, 1997; SATOH; AKIRA, 2016). Até o momento, foram identificados 13 receptores do tipo TLR em camundongos, sendo que 10 desses receptores são compartilhados com seres humanos (ROCK et al., 1998). Esses receptores localizam-se na membrana plasmática das células, como é o caso de TLR1, TLR2, TLR4, TLR5 e TLR6, ou em endossomos, como ocorre com TLR3, TLR4, TLR7, TLR8, TLR9 e TLR13 (O'NEILL et al., 2013). No caso dos TLRs endossomais, o domínio N-terminal de reconhecimento encontra-se no interior do endossomo (CREAGH; O'NEILL, 2006).

Os receptores TLRs reconhecem moléculas presentes em protozoários, fungos, bactérias e vírus. Entre os TLRs capazes de reconhecer proteínas bacterianas encontram-se TLR2, TLR4 e TLR5. O TLR2 reconhece lipoproteínas, glicolipídios, peptidoglicanas, ácidos lipoteicóicos, zimosam e manana derivadas de bactérias Gram-positivas. A proteína flagelina presente nos flagelos bacterianos ativa TLR5. Além de reconhecer LPS presente na parede celular de bactérias Gram-negativas, TLR4 também é ativado por proteínas do envelope do vírus sincicial, fosfolipídios glicoinositol de tripanossomos e proteínas *heat shock* 60 e 70. (AKIRA; TAKEDA, 2004; MEDZHITOV; JANEWAY, 2000; SATOH; AKIRA, 2016). De modo geral, a sinalização via TLRs tem como consequência a indução da produção de quimiocinas, citocinas pró-inflamatórias e da transcrição de genes envolvidos com a produção de proteínas que atuam na resposta imune inata como caspase-1, caspase-11 e NLRP3 (do inglês, *NLR family of PYRIN domain containing Protein 3*). Além de induzir a produção de citocinas pró-inflamatórias como interferons (IFNs) do tipo I, fator de necrose tumoral (TNF) e interleucina(IL)-12, a sinalização via TLRs participa da produção de formas precursoras das citocinas pró-inflamatórias pró-IL-1 β e pró-IL-18. Esse importante processo prepara as células para a ativação de outras vias do sistema imune inato, como por exemplo a formação de plataformas moleculares intracelulares conhecidas como inflamassomas em resposta ao reconhecimento de patógenos (KAWAI; AKIRA, 2011; LATZ; XIAO; STUTZ, 2013).

Como foi dito anteriormente, a entrada de *C. burnetii* fase I nas células e o remodelamento de actina que ocorre durante esse processo são dependentes de TLRs, principalmente TLR4. Macrófagos deficientes para TLR4 apresentam entrada bacteriana prejudicada e produção reduzida de citocinas pró-inflamatórias. No entanto, ainda dispõem de formação defectiva e transiente de granulomas, uma das formas conhecidas de resposta imune sistêmica mediada por células que atuam na tentativa de controlar a infecção por *C. burnetii*.

(HONSTETTRE et al., 2004). Após a descoberta da estrutura tetra-acilada do lipídio-A de *C. burnetii* verificou-se que o mesmo atua como antagonista da sinalização via TLR4 (ZAMBONI et al., 2004; CUNHA; ZAMBONI, 2012). Este mecanismo de evasão da resposta imune que ocorre tanto em bactérias fase I quanto fase II já foi observado também em outras bactérias patogênicas, como *Yersinia pestis* (TELEPNEV et al., 2009; VAN SCHAIK et al., 2013). Esse antagonismo é um processo complexo e parece envolver, ao menos no caso das bactérias fase I, o desencadeamento prejudicado da ativação de vias de MAPK em macrófagos. Este mecanismo é importante para evitar a conversão de fagolisossomos contendo a bactéria em compartimentos degradativos contendo cathepsina D e, assim, favorecer a sobrevivência da bactéria (BARRY et al., 2012).

Outro importante receptor da família dos TLRs que está envolvido na resposta imune à *C. burnetii* é o TLR2. As formas avirulentas da bactéria podem ser reconhecidas por TLR2, induzindo a ativação de respostas imunes do tipo I, com produção de IL-12 e TNF. Essas citocinas contribuem para o controle da infecção em macrófagos murinos. Macrófagos deficientes para TLR2 apresentam permissividade aumentada para a replicação bacteriana em relação a macrófagos selvagens ou deficientes para TLR4, corroborando o importante papel do receptor TLR2 na restrição do crescimento de *C. burnetii*. No entanto, o mesmo não ocorre para a forma virulenta da bactéria. No caso da fase I, o LPS do tipo liso se organiza criando uma estrutura em forma de escudo que impede o reconhecimento de ligantes na superfície bacteriana via TLR2 (ZAMBONI et al., 2004; SHANNON; HOWE; HEINZEN, 2005).

C. burnetii é uma bactéria intracelular com sistema de secreção do tipo IVB ativo, com capacidade de estabelecer um nicho replicativo e multiplicar no interior de macrófagos. Por essa razão, é razoável especular que *C. burnetii* possa ainda subverter respostas imunes que incluam o reconhecimento por outra classe de PRRs da imunidade inata, como os receptores citoplasmáticos do tipo *Nod* (NLRs, do inglês *Nucleotide-binding domain leucine-rich repeat-containing protein receptors*). Esses receptores também são capazes de reconhecer MAMPs e DAMPs e montam resposta por meio da ativação dos inflamassomas, que são grandes complexos proteicos que processam e ativam proteínas caspases e citocinas pró-inflamatórias (TING; WILLINGHAM; BERGSTRALH, 2008; GUO; CALLAWAY; TING, 2015). Já foi descrito por nosso grupo de pesquisa a participação do efector bacteriano IcaA (do inglês, *Inhibition of caspase Activation*), na inibição da ativação de caspase-11 e suas funções, incluindo formação de poros e ativação não-canônica do inflamassoma (CUNHA et al., 2015).

No entanto, ainda são necessárias mais investigações do possível papel dos inflamassomas na resposta à infecção por *C. burnetii*.

O desenvolvimento e aprimoramento do cultivo da bactéria em meio axênico permitiu o avanço das metodologias de manipulação genética de *C. burnetii* (OMSLAND et al., 2009; OMSLAND et al., 2011). Dessa forma, possibilitou-se a identificação de novos efetores bacterianos envolvidos em processos como biogênese e manutenção do vacúolo parasitóforo e subversão da resposta imune do hospedeiro. A identificação do efector IcaA e a caracterização do seu papel na subversão da resposta imune do hospedeiro por nosso grupo de pesquisa, por exemplo, só foi possível graças à utilização do cultivo em meio axênico para criação da bactéria mutante para IcaA, reforçando a importância do desenvolvimento dessas técnicas. Futuramente, essas técnicas auxiliarão ainda mais no entendimento das relações patógeno-hospedeiro envolvendo infecções com essa bactéria (CAREY et al., 2011; NEWTON et al., 2014; KOHLER et al., 2016; LATOMANSKI et al., 2016; CRABILL et al., 2018).

1.4. Polarização de macrófagos e monócitos em resposta à *Coxiella burnetii*

Um aspecto importante envolvendo a resposta imune do hospedeiro à infecção por *C. burnetii* diz respeito ao estado de polarização induzido pela bactéria em monócitos e macrófagos, suas principais células-alvo. Ambos os tipos celulares participam de diversos processos nos organismos como desenvolvimento e homeostase de tecidos, reparação tecidual, fagocitose de patógenos e células mortas, inflamação e outros aspectos da resposta imune (MANTOVANI et al., 2002; MANTOVANI et al., 2004; WYNN; CHAWLA; POLLARD, 2013). A habilidade dessas células em desempenhar tantas funções distintas reside na grande heterogeneidade e plasticidade que apresentam. De forma geral, diz-se que a polarização dos macrófagos e monócitos refere-se a um estado de ativação especializado adquirido em um determinado local dentro de uma janela temporal específica, em resposta a sinais endógenos e exógenos. Esses sinais podem ser citocinas, produtos microbianos, sinais de perigo, sinais homeostáticos, fatores de crescimento, entre outros. Depois de ativadas, as células desempenham funções especializadas (MANTOVANI; SICA; LOCATI, 2005; MARTINEZ; GORDON, 2014; GLASS; NATOLI, 2016; MURRAY, 2017).

O conceito de ativação clássica dos macrófagos surgiu a partir dos trabalhos sobre resistência celular à infecção realizados por Mackaness (1962). Nesses trabalhos foi demonstrado aumento da atividade microbicida de macrófagos dependente de antígenos contra *Listeria* spp. e BCG (do inglês, *Bacillus Calmette-Guerin*) após exposição secundária

aos patógenos. Esse aumento foi posteriormente relacionado à produção de IFN- γ e respostas do tipo Th1 pelas células imunes ativadas pelos antígenos, sendo estendido para propriedades antitumorais e citotóxicas (NATHAN et al., 1983; PACE et al., 1983; CELADA et al., 1984). Posteriormente, outros pesquisadores identificaram macrófagos que apresentavam respostas relacionadas com o padrão Th2 da resposta imune, diferentemente do fenótipo de ativação proposto no modelo clássico de Mackaness (1962). Como resposta padrão Th2, os macrófagos apresentaram secreção reduzida de citocinas pró-inflamatórias, proteção extracelular de parasitas e alta expressão de MHC-II (STEIN et al., 1992; DOYLE et al., 1994). Considerando observações, juntamente com a descoberta de uma capacidade de resposta diferencial a estímulos clássicos como IFN- γ , LPS ou ambos e diferenças metabólicas entre macrófagos ativados em camundongos com backgrounds Th1 ou Th2, Mills e colaboradores (2000) criaram os termos “M1” e “M2” para se referir a macrófagos ativados de forma “clássica” ou “alternativa”, respectivamente. Ao macrófago naïve, diferenciado, mas não ativado, foi designado o termo “M0” (MILLS et al., 2000; MARTINEZ; GORDON, 2014; MURRAY, 2017).

Durante muitos anos essa classificação dualística foi mantida para se referir aos estados de ativação e polarização dos macrófagos. No entanto, o avanço dos estudos na área e o aprimoramento das técnicas de análise mostraram a existência de outros fenótipos de polarização dos macrófagos que não se enquadravam perfeitamente no modelo existente, como é o caso de IL-10, TGF- β e glicocorticoides, reforçando a necessidade de revisá-lo. De fato, a polarização em si é afetada por múltiplos fatores que incluem, por exemplo, a origem do macrófago e o microambiente tecidual e circulante, os quais por si só apresentam uma heterogeneidade grande e dependem inclusive do momento analisado. Sendo assim, a questão da polarização dos macrófagos deve ser vista como multidimensional (GINHOUX et al., 2015; XUE et al., 2014; MURRAY, 2017). Esses fenótipos distintos foram então integrados e subclassificados entre os dois estados funcionais polarizados que já existiam, sendo a classificação dependente do efeito que causavam em marcadores de macrófagos selecionados. Dessa forma, manteve-se o estado M1 e subdividiu-se o M2 nos estados: a) M2a, induzido por IL-4 e IL-13; b) M2b, induzido por imunocomplexos, receptores Fc, agonistas de TLRs e agonistas de IL-1R; c) M2c, induzido por IL-10 e glicocorticoides e comum para humanos e camundongos (Figura 3); e d) M2d, induzido por agonistas de TLRs via receptor de adenosina, relacionado a macrófagos associados a tumores e exclusivo para camundongos

(MANTOVANI et al., 2004; DULUC et al., 2007; MARTINEZ et al., 2008; MARTINEZ; GORDON, 2014; MURRAY, 2017; SHAPOURI-MOGHADDAM et al., 2018).

Figura 3 - Representação do modelo de polarização de macrófagos com os estados comuns para humanos e camundongos. Macrófagos apresentam fenótipos distintos de polarização dependendo do estímulo recebido, os quais podem induzir um estado M1 (induzido por IFN γ + LPS ou TNF) ou um estado M2. O estado M2 subdivide-se em M2a (induzido por IL-4 e IL-13), M2b (induzido por imunocomplexos + ligantes de TLR/IL-1R) e M2c (induzido por IL-10 e glicocorticóides). Em camundongos, ocorre ainda a polarização em um estado M2d (induzido por agonistas de TLRs via receptor de adenosina), não representado na imagem acima. Cada estado de polarização vai desencadear ações efetoras específicas, refletindo assim a plasticidade e a heterogeneidade observada entre macrófagos. Adaptado de: Mantovani e colaboradores (2004).

De forma geral, macrófagos classicamente ativados, ou M1, participam funcionalmente de respostas pró-inflamatórias, anti-tumorais e antimicrobianas, participando da eliminação de patógenos por meio da ativação do sistema de NADPH oxidase (do inglês, *Nicotinamide Adenine Dinucleotide PHosphate*) e geração subsequente de espécies reativas de oxigênio (ROS), além da geração de espécies reativas de nitrogênio (RNS). Devido ao alto nível de ROS que produzem, esses macrófagos direcionam a indução de danos teciduais e prejudicam a regeneração de tecidos, ocasionalmente levando ao desenvolvimento de quadros de inflamação crônica. A indução desse estado de ativação ocorre tipicamente por citocinas Th1, como IFN- γ e TNF- α , ou pelo reconhecimento de LPS bacteriano, levando à produção e

secreção de altos níveis de citocinas pró-inflamatórias como TNF- α , IL-1 α , IL-1 β , IL-6, IL-12, IL-23 e ciclooxigenase-2 (COX-2) e baixos níveis de IL-10. Além disso, a expressão elevada de genes como *Nos2*, *Tnf*, *Ifn* e *Tlr2* são considerados alguns dos marcadores de polarização M1 (MANTOVANI et al., 2004; BISWAS et al., 2012; LOCATI; MANTOVANI; SICA, 2013; MURRAY et al., 2014; MURRAY, 2017; SHAPOURI-MOGHADDAM et al., 2018).

Os macrófagos alternativamente ativados, ou M2, apresentam de forma geral respostas padrão Th2 com características anti-inflamatórias, desempenhando funções relacionadas com a manutenção da homeostase, apresentando alta capacidade de fagocitose, propriedades pró-fibróticas e pró-angiogênicas, promovendo reparo celular e cicatrização, imunoregulação, remodelamento tecidual, beneficiando a formação e progressão de tumores e a eliminação e encapsulamento de parasitas. A indução do estado M2 se dá principalmente pela ação das citocinas IL-4 e IL-13 via ativação de STAT6 por meio do receptor IL-4R α . No entanto, as citocinas IL-10, IL-33 e IL-21 também podem induzir esse estado de polarização por meio de outras vias. Macrófagos M2 produzem quantidades pequenas de IL-12 e quantidades altas de IL-10 e TGF- β , além da expressão elevada de genes como *Arginase1*, *Fizz1*, *Ym1*, *Il-4*, receptores de manose são considerados alguns dos marcadores desse padrão de polarização (GORDON, 2003; MANTOVANI et al., 2013; MURRAY, 2017; SHAPOURI-MOGHADDAM 2018).

Ainda não está bem estabelecido qual o fator mais importante para a determinação do estado de polarização e, conseqüentemente, para as funções especializadas desempenhadas por macrófagos: a ontogenia ou a plasticidade, representada pelo efeito do microambiente tecidual. Essa questão é de extrema importância, visto que pode auxiliar no entendimento de como as células se comportam em estados de homeostase ou não, como no caso de inflamações e infecções por patógenos, e assim buscar estratégias de intervenção. Está bem elucidado que macrófagos possuem duas fontes de origem, sendo elas os progenitores embrionários e a medula óssea. Ambos apresentam capacidade de proliferação sob ação de estímulos apropriados e desempenham funções específicas, essenciais para garantia da manutenção da homeostase do organismo (LAVIN et al., 2015; PERDIGUERO; GEISSMANN, 2015; GINHOUX et al., 2016; MURRAY, 2017; BONNARDEL; GUILLIAMS, 2018).

Os tecidos apresentam uma população de macrófagos com origem mista, sendo que nesses locais é possível observar a presença de macrófagos residentes, geralmente

terminalmente diferenciados, e também de precursores recrutados. Sendo assim, cientistas têm investigado se a ativação e polarização desses macrófagos dependem de sua origem ou do microambiente tecidual em que estão inseridos. Caso dependam da origem, as células responderiam de forma diferente caso tenham derivado de precursores embrionários ou da medula óssea mesmo frente aos mesmos estímulos do microambiente tecidual. No outro caso, independentemente da origem, o que determinaria o estado de polarização seria o microambiente tecidual (incluindo aqui citocinas, quimiocinas, fatores de sobrevivência e outros) em um determinado espaço e tempo, ou seja, sua grande plasticidade (Figura 4) (MURRAY, 2017; BONNARDEL; GUILLIAMS, 2018). Apesar de ambas as hipóteses apresentarem evidências que as suportam, há uma terceira hipótese que as combina. Nela, o microambiente tecidual em um determinado local dentro de uma janela temporal específica seria essencial para direcionar os macrófagos para um determinado estado de polarização, no qual a amplitude da resposta apresentada variaria dependendo da origem daquela célula.

Figura 4 - Representação dos modelos de plasticidade e ontogenia na determinação do estado de polarização dos macrófagos. Segundo o modelo da plasticidade, a população de macrófagos em um determinado local seria relativamente homogênea, pois os estímulos fornecidos pelo microambiente seriam prevalentes para definição do estado de polarização, independentemente da origem apresentada. Nesse modelo, após o término dos estímulos (por exemplo, a resolução de um cenário de inflamação) a população de macrófagos retornaria a um estado homogêneo de macrófagos residentes, com perfil de expressão gênica idêntico e sem grandes alterações epigenéticas. Já segundo o modelo que considera a ontogenia como fator prevalente, macrófagos residentes que apresentam uma mesma origem teriam padrões similares de expressão gênica entre si, com prevalência de diferenças epigenéticas próprias à cada origem, resultando em uma população heterogênea de células. Frente a um estímulo como um cenário de inflamação, os macrófagos responderiam de forma diferente de acordo com sua origem. Após o término do estímulo, os macrófagos restantes apresentariam diferentes perfis de expressão gênica e acumulariam novas diferenças epigenéticas, resultando cada vez mais em populações heterogêneas. Adaptado de Bonnardel e Williams (2018).

No contexto das infecções por *C. burnetii*, foi observado que o padrão de polarização induzido varia de acordo com o tipo celular infectado (macrófago ou monócito). Para os monócitos, a infecção com *C. burnetii* resulta em polarização no padrão M1, desencadeando uma forte resposta pró-inflamatória e microbicida do tipo Th1, com produção de grandes quantidades de IFN- γ . De fato, IFN- γ parece ser fundamental na resposta microbicida à bactéria, visto que macrófagos deficientes para o gene de IFN- γ mostram-se mais suscetíveis à replicação (ANDOH et al., 2007; DELLACASAGRANDE et al., 1999). A atividade microbicida induzida por IFN- γ é independente de oxigênio e, em modelos murinos, está relacionada com a produção de RNS (YOSHII et al., 1999; ZAMBONI; RABINOVITCH, 2003; BRENNAN et al., 2004; ZAMBONI; RABINOVITCH, 2004). Dentre as respostas microbicidas induzidas por IFN- γ e independentes de oxigênio encontram-se a capacidade de alterar o pH dos fagossomos contendo *C. burnetii* através da interferência com processo de fusão entre os fagossomos e os lisossomos, a habilidade de controlar o metabolismo de ferro por meio de redução na quantidade de receptores de transferina (levando à redução da carga bacteriana), e a promoção da apoptose das células infectadas de forma dependente de TNF (DELLACASAGRANDE et al., 1999; HOWE; MALLAVIA, 1999; GHIGO et al., 2002; CAPO; MEGE, 2012).

Diferentemente, em macrófagos foi observado que *C. burnetii* induz um programa atípico de polarização em um estado M2. O mesmo é caracterizado pela expressão elevada de genes que codificam para IL-10, IL-1ra, TGF- β , Arginase-1 e receptores de manose, bem como diminuição dos níveis de expressão dos genes que codificam para TNF- α e TLR2, marcadores típicos de padrão M1 (BENOIT et al., 2008). Por seu forte papel imunomodulador, essa produção elevada de IL-10 induzida pela bactéria compreende um dos principais mecanismos relacionados com a persistência da infecção, levando à evolução da doença para cenários de infecção crônica (GHIGO et al., 2004; AMARA; BECHAH; MEGE, 2012; MEGHARI et al., 2006; CAPO; MEGE, 2012; ELDIN, 2017). Dessa forma, o tipo de padrão de polarização adquirido em resposta à infecção por *C. burnetii* é um processo fundamental para o desencadeamento de respostas, podendo tanto levar ao controle da infecção quanto à persistência da bactéria no hospedeiro e desenvolvimento da forma crônica da doença. Sendo assim, a interferência com o padrão de polarização dos macrófagos pode compreender um alvo terapêutico para o controle das infecções por *C. burnetii*.

1.5. Modelos de estudo das interações patógeno-hospedeiro com infecção por *Coxiella burnetii*

Apesar da diferença nos tipos de LPS, o tráfego intracelular e a cinética de crescimento das formas virulentas e avirulentas de *C. burnetii* em macrófagos humanos é indistinguível. Além disso, macrófagos humanos são capazes de restringir igualmente a multiplicação tanto da fase I quanto da fase II, portanto ambas as variantes podem ser utilizadas em estudos para avaliação das relações patógeno-hospedeiro. Em adição, há um clone da cepa *Nine Mile* fase II que apresenta uma grande deleção cromossomal que o torna incapaz de retornar ao estado completo de virulência, tornando-o apropriado para uso em condições de biossegurança nível 2 (AMANO; WILLIAMS, 1984; HACKSTADT, 1996; HOWE et al., 2010; GRAHAM et al., 2013; LARSON et al., 2016).

Embora *C. burnetii* seja altamente adaptada à sobrevivência nas células do hospedeiro, subvertendo e evadindo as respostas imunes por meio da ação de seus efetores bacterianos e manipulação de diferentes vias de sinalização, existem hospedeiros imunocompetentes que conseguem controlar sua multiplicação (ZAMBONI, 2004). Portanto, existem respostas efetivas contra *C. burnetii*. Como demonstrado anteriormente, macrófagos de origens distintas diferem quanto à suscetibilidade à replicação intracelular de *C. burnetii*. Macrófagos derivados de medula óssea e macrófagos peritoneais primários se mostraram intrinsecamente mais restritivos (ZAMBONI et al., 2002; ZAMBONI; RABINOVITCH, 2003). Essas observações sugerem que a suscetibilidade dos macrófagos à infecção por *C. burnetii* possa estar relacionada com a origem celular, mais do que com o microambiente em que se encontram. Além disso, macrófagos derivados da medula óssea de camundongos da linhagem C57BL/6 são capazes de controlar o crescimento da bactéria fase II, enquanto macrófagos derivados da medula óssea de camundongos das linhagens A/J e BALB/c são mais suscetíveis, demonstrando que diferentes linhagens de camundongos também apresentam diferenças na suscetibilidade à bactéria (ZAMBONI, 2004; ZAMBONI; RABINOVITCH, 2004). O mesmo foi observado para estudos envolvendo a infecção de camundongos das linhagens C57BL/6 e A/J frente à infecção por *C. burnetii* fase I, com os primeiros se mostrando mais restritivos, levantando a possibilidade de que a suscetibilidade à *C. burnetii* esteja relacionada com alguma característica inerente ao fundo gênico do camundongo (SCOTT; WILLIAMS; STEPHENSON, 1987). No entanto, ao analisar macrófagos peritoneais provenientes de camundongos BALB/c, verificou-se que os mesmos apresentaram

a capacidade de controlar parcialmente a infecção por *C. burnetii* fase II (ZAMBONI et al., 2002).

Além de macrófagos peritoneais e macrófagos derivados de medula óssea, macrófagos alveolares compreendem outro importante modelo de infecção com *C. burnetii* utilizado para estudos sobre a interação entre a bactéria e o hospedeiro. Os macrófagos alveolares têm sido considerados por muito tempo como sítio primário da infecção por esse patógeno (KHAVIN; TABIBZADEH, 1988). Estudos realizados com macrófagos alveolares isolados de tecidos *post-mortem* de seres humanos apresentaram capacidade de suportar a replicação de diferentes patótipos de *C. burnetii* (GRAHAM et al., 2013). Interessantemente, macrófagos alveolares derivados de macacos demonstraram forte atividade anti-apoptótica associada com infecções por *C. burnetii* (VOTH; HOWE; HEINZEN, 2007). Já em modelos murinos, foi demonstrado que macrófagos alveolares produzem óxido nítrico em resposta à infecção (YOSHIIE et al., 1999) e são suscetíveis à replicação do patógeno (CALVERLEY et al., 2012). Nesse contexto, estudos visando o entendimento das relações patógeno-hospedeiro e dos mecanismos pelos quais diferentes fundos gênicos e tipos celulares distintos conseguem controlar o crescimento intracelular de *C. burnetii* permanecem de extrema relevância e continuam a suscitar grande interesse científico. Para isso, é necessário o desenvolvimento de novos modelos de estudo apropriados como ferramenta fundamental para os avanços nessa área.

2. OBJETIVOS

2.1. Objetivo Geral

Avaliar a suscetibilidade de macrófagos alveolares murinos à infecção por *C. burnetii* fase II *in vitro* a fim de estabelecer um modelo de célula primária que seja fisiologicamente relevante para estudos de interação patógeno-hospedeiro com essa bactéria.

2.2. Objetivos Específicos

- Avaliar a resposta de macrófagos derivados de medula óssea e de macrófagos alveolares à infecção por *C. burnetii*.
- Analisar a suscetibilidade de macrófagos alveolares à infecção por *C. burnetii* através de técnicas de microscopia.
- Identificar os possíveis mecanismos envolvidos na suscetibilidade de macrófagos alveolares à infecção por *C. burnetii*.
- Analisar a suscetibilidade à infecção por *C. burnetii* de macrófagos alveolares deficientes em componentes importantes para resposta imune contra esse patógeno.
- Comparar a suscetibilidade de macrófagos alveolares murinos à infecção por *C. burnetii* com outros modelos celulares suscetíveis já descritos.

3. MATERIAIS E MÉTODOS

3.1. Animais

Foram utilizados camundongos adultos (seis a oito semanas) das linhagens A/J, C57BL/6, bem como nocautes para *Nos2*, *Ifn γ* e *Il-4* com fundo genético de camundongos C57BL/6. Os animais foram mantidos no Biotério do Departamento de Biologia Celular e Molecular e Bioagentes Patogênicos e no Biotério do Departamento de Imunologia e Bioquímica. Foram providenciadas água e comida à vontade para todos os animais.

3.2. Obtenção e preparo de *C. burnetii* para infecção

Inóculos de *Coxiella burnetii* da linhagem Nine Mile fase II (RSA493) foram preparados a partir de monocamadas de células Vero infectadas, conforme descrito por Zamboni; Mortara e Rabinovitch (2001). Culturas confluentes de células Vero foram irradiadas com 10 Gy 60Co para bloquear a divisão celular e mantidas a 37°C e 5% de CO₂ em meio DMEM (do inglês, *Dulbecco's Modified Eagle's Medium*) (Gibco – Thermo Fischer) com 10% de soro fetal bovino (SBF) (Gibco – Thermo Fischer) e 1mM L-glutamina (Sigma-Aldrich). As células Vero foram infectadas com *Coxiella* e mantidas nas condições acima por mais seis dias. Para o preparo dos inóculos, as células Vero infectadas foram lisadas por homogeneização em água destilada estéril. O lisado celular foi centrifugado a 1.810 x g por 10 minutos a 4°C. O sobrenadante foi descartado e o pellet ressuscitado em PBS (do inglês, *Phosphate-Buffered Saline*) (Gibco – Thermo Fischer) estéril gelado. Após nova centrifugação a 12.857 x g por 30 minutos a 4°C, o sobrenadante foi novamente descartado e o pellet ressuscitado em RPMI 1640 + HEPES (Gibco – Thermo Fischer) com 10% SBF, submetido à sonicação a 35 kHz durante três minutos, aliquoteado e mantido a temperatura de -80°C. Para a realização dos experimentos, as alíquotas foram descongeladas em banho a 37°C e submetidas à sonicação a 35 kHz por três minutos.

3.3. Cultura de células de linhagem de mamíferos

Células THP-1 foram cultivadas em meio RPMI 1640 + HEPES suplementado com 10% SBF, e células Vero foram mantidas em meio DMEM com 10% SBF e 1mM L-

glutamina a 37°C em 5% CO₂. Para experimentos com células THP-1, a diferenciação em células do tipo macrófago foram mediadas pela adição de 200nM de forbol 12-miristato 13-acetato (PMA) na cultura, como descrito por Voth; Howe e Heinzen (2007). Após doze-quatorze horas, o meio foi substituído por meio sem PMA para a realização das infecções. Para infecção das células Vero, as células foram irradiadas com 10 Gy 60Co e mantidas durante doze-quatorze horas a 37°C em 5% CO₂ antes da infecção.

3.4. Preparo de macrófagos derivados de medula óssea (BMDMs)

Macrófagos derivados de medula óssea foram diferenciados in vitro a partir de células tronco extraídas do fêmur de camundongos adultos (seis a oito semanas) das linhagens citadas anteriormente, segundo protocolo descrito por Marim e colaboradores (2010). Brevemente, células da medula óssea foram coletadas dos fêmures e diferenciadas por sete dias em estufa a 37°C, 5% CO₂ em meio RPMI 1640 + HEPES suplementado com 20% de SBF, 30% de meio condicionado por células L929 (LCCM), fonte de M-CSF (fator importante para diferenciação de células progenitoras em macrófagos), e 100U/mL de penicilina/estreptomicina (Sigma-Aldrich). Após a diferenciação, os macrófagos foram coletados pela lavagem das monocamadas com PBS gelado, centrifugados a 200 x g por cinco minutos a 4°C e o sobrenadante descartado. O pellet foi ressuspensionado em meio RPMI 1640 + HEPES contendo 10% SBF e 5% de LCCM e as células foram quantificadas com auxílio de uma câmara de Neubauer. Os macrófagos foram plaqueados a uma densidade de 10⁵ células/poço em placas tratadas de 24 poços um dia antes da infecção e mantidos em cultura em meio RPMI 1640 + HEPES, contendo 10% SBF e 5% de LCCM.

3.5. Preparo de macrófagos alveolares (AMs)

Macrófagos alveolares foram obtidos de camundongos adultos (seis a oito semanas) através de lavagem brônquio-alveolar (BAL). Os animais foram eutanasiados por dose excessiva de anestésico com injeção intravenosa de cetamina e xilazina a 50 mg/kg e 10 mg/kg, respectivamente. Posteriormente, a traqueia dos camundongos foi perfurada com cânula estéril (BD AngioCath) e os pulmões foram lavados quatro vezes consecutivas com 1 ml de estéril contendo 5mM de EDTA (do inglês, *Ethylenediamine Tetraacetic Acid*) a 37°C e o material coletado foi depositado em um tubo cônico tipo falcon estéril. As células obtidas

foram transferidas para um novo falcon e lavadas duas vezes em PBS estéril, com centrifugações a 200 x g durante 10 minutos a 4°C. O sobrenadante foi descartado e o pellet ressuspensionado em meio RPMI 1640 + HEPES com 10% SBF, 1mM L-glutamina e 100U/ml penicilina/estreptomicina. As células foram quantificadas com auxílio de uma câmara de Neubauer e plaqueadas na densidade de 10⁵ células/poço em placas de 24 poços. Os macrófagos foram mantidos em cultura em meio RPMI 1640 + HEPES com 10% SBF, 1mM L-glutamina e 100U/ml penicilina/estreptomicina, a uma temperatura de 37°C e 5% CO₂. O meio de cultura foi substituído duas vezes antes da infecção por meio sem antibiótico. Meio fresco foi adicionado às culturas após seis dias de infecção.

3.6. Infecção com *Coxiella burnetii*

Para infecção das culturas celulares, alíquotas de *C. burnetii* foram descongeladas em banho a 37°C e submetidas à sonicação a 35 kHz por três minutos para dissolução de aglomerados de bactéria. O meio presente nas placas com as células em cultivo foi aspirado e substituído por meio fresco antes da infecção. As bactérias foram homogeneizadas com pipeta e inoculadas diretamente nas placas com as células em cultura, com multiplicidade de infecção (MOI) de 3 ou 100. As placas foram centrifugadas a 200 x g por cinco minutos a 25°C e colocadas em estufa a 37°C e 5% de CO₂. Após um período de infecção de quatro horas o meio de cultura foi aspirado, os poços lavados com PBS estéril morno e o PBS substituído novamente por meio de cultura fresco. Como um controle experimental interno para replicação de *C. burnetii*, um grupo de células infectadas foi tratado com 20ug/ml de rifampicina (RIF) para inibir a tradução proteica de *C. burnetii* (BRENNAN; SAMUEL, 2003).

3.7. Microscopia de Fluorescência

AMs de camundongos C57BL/6 foram plaqueados em uma densidade de 10⁵ células por poço, em uma placa de 24 poços contendo lamínulas de vidro de 13mm. As células foram infectadas com *C. burnetii* selvagem ou *C. burnetii* expressando *mCherry* em um MOI de 3, como descrito anteriormente (CUNHA et al., 2015). Após quatro horas, três dias, seis dias, nove dias ou 13 dias de infecção, as células foram fixadas com 3,5% de PFA (do inglês,

Paraformaldehyde) contendo 7,5% de sacarose por 20 minutos e então lavadas duas vezes com PBS em temperatura ambiente. Para os ensaios com *C. burnetii* selvagem, as células foram então permeabilizadas por dois minutos com metanol 100% gelado, seguido por mais duas lavagens com PBS. A autofluorescência celular foi removida por meio de incubação com glicina 0.1M diluída em PBS por 10 minutos, e então as amostras foram bloqueadas por 30 minutos em temperatura ambiente utilizando PBS contendo 1% de albumina de soro bovino (BSA – *Sigma-Aldrich*) e 5% de soro de cabra. As células foram incubadas com os anticorpos primários anti- *C. burnetii* (diluição 1:100) feito em coelho e anti-LAMP-1 (diluição 1:2) feito em rato, diluídos em PBS contendo 1% de BSA por uma hora, posteriormente foram lavadas duas vezes com PBS em temperatura ambiente e então incubadas por 30 minutos em PBS com 1% BSA contendo os anticorpos secundários conjugados com o fluoróforo Alexa Fluor 488 e Alexa Fluor 594 (*Molecular probes – Thermo Fisher Scientific*) em uma diluição de 1:2000. Após três lavagens com PBS a temperatura ambiente, as células foram montadas utilizando *Prolong Gold Antifade Reagent* contendo DAPI (*Molecular probes – Thermo Fisher Scientific*). Para os experimentos utilizando a *C. burnetii* expressando *mCherry*, as células foram lavadas duas vezes com PBS em temperatura ambiente após a fixação e então montadas diretamente com *Prolong Gold Antifade Reagent* contendo DAPI. Imagens de microscopia multifóton foram adquiridas utilizando um microscópio *LSM 780 Zeiss AxioObserver*, equipado com uma lente objetiva de imersão à óleo de 63X de aumento e analisadas utilizando o programa *ImageJ*.

3.8. Microscopia Eletrônica de Transmissão

AMs foram infectados por três ou seis dias, lavados uma vez com PBS, fixados por uma hora em temperatura ambiente com fixador de Karnovsky modificado (2% formaldeído, 2,5% glutaraldeído em 0,1M de cacodilato de sódio, pH 7.4), soltos e então processados para e analisados por microscopia eletrônica de transmissão, conforme descrito por Zamboni e colaboradores (2002).

3.9. Quantificação de *Coxiella burnetii* in vitro

BMDMs, AMs, células THP-1 e células Vero irradiadas, infectadas com *C. burnetii*, foram mantidas em cultura a 37°C, em uma atmosfera contendo 5% de CO₂, por até 12 dias. Após seis dias de infecção, as células foram suplementadas com um adicional de 500ul de meio RPMI 1640 + HEPES contendo 10% de SFB e 5% de LCCM para BMDMs, meio RPMI 1640 + HEPES com 10% SFB para AMs e células THP-1 e meio DMEM com 10% SFB e 1mM L-glutamina para células Vero irradiadas. Em diferentes tempos após a infecção, células foram lisadas com 1ml de água deionizada e os lisados celulares foram combinados com o sobrenadante das culturas de cada poço, garantindo assim que toda bactéria contida no poço fosse contada. O material foi então centrifugado a 14.000 x g por 15 minutos em uma temperatura de 4°C para a deposição das bactérias presentes nas amostras. O DNA de *C. burnetii* foi extraído utilizando o kit *Illustra™ Bacteria GenomicPrep* (GE Healthcare) e a presença de material genético foi confirmada por espectrometria (absorbância a 260nm). O par de iniciadores dotA-for e dotA-rev (CAREY et al., 2011) foi utilizado para reação de PCR quantitativo (qPCR), um método confiável para quantificação da replicação bacteriana, como demonstrado por Brennan e Samuel (2003) para análises envolvendo *C. burnetii*. Para realização da reação, foi utilizado o kit *Platinum® SYBR® Green qPCR SuperMix-UDG with ROX* (Invitrogen) e as reações de termociclagem foram montadas de acordo com as instruções do fabricante. As amostras foram lidas em um termociclador *ABI Prism 7000*.

3.10. Análises de RT-PCR

Para análises de RT-PCR foi feita a extração de mRNA de culturas de 10⁶ AMs e BMDMs derivados de camundongos C57BL/6, infectados ou não por 12 horas com *C. burnetii* em um MOI de 3, utilizando o reagente TRIzol. Purificação de RNA foi feita utilizando o minikit *Illustra™ RNAspin* (GE Healthcare), de acordo com as instruções do fabricante, e o material genético foi quantificado usando o espectrofotômetro *NanoDrop* (Thermo Scientific). A transcriptase reversa *ImProm-II* (Promega) foi utilizada para obtenção de cDNA. Os ensaios de PCR quantitativa em tempo real foram realizados usando kit *Platinum® SYBR® Green qPCR SuperMix-UDG* com carboxi-X-rodamina (Invitrogen). Reações de termociclagem foram feitas de acordo com as instruções do fabricante, e as amostras foram lidas em um sistema de PCR em tempo real *StepOnePlus* (Applied

Biosystems). A expressão dos genes alvos foi calculada utilizando os valores do ciclo médio do limite (*CT*) para cada amostra, os quais foram normalizados com relação a aqueles dos genes *housekeeping*. As sequências dos iniciadores usados para a reação de PCR de transcrição reversa (RT-PCR) são as seguintes: *Gapdh*-fwd, 5'-CATCACCATCTTCCAGGAGCG-3'; *Gapdh*-rev, 5'-ACGGACACATTGGGGGTAGG-3'; *Nos2*-fwd, 5'-CGAAACGCTTCSCCTTCCAA-3'; *Nos2*-rev, 5'-TGAGCCTATATTGCTGTGGCT-3'; *Ifn*-fwd, 5'-GCATCTTGGCTTTGCAGCT-3'; *Ifn*-rev, 5'-CCTTTTGGCCCTTGCTGTTG-3'; *Tnf*-fwd, 5'-TGTGCTCAGAGCTTTCAACAA-3'; *Tnf*-rev, 5'-CTTGATGGTGCATGAGA-3'; *Tlr2*-fwd, 5'-AAGTCTCCGGAATTATCAGTCC-3'; *Tlr2*-rev, 5'-TGATGGATGTCGCGGAT-3'; *Il4*-fwd, 5'-AAGAGCATCATGCAAATGGA-3'; *Il4*-rev, 5'-TTAAAGCATGGTGGCTCAGTAC-3'; *Arginase1*-fwd, 5'-GTTCCCAGATGTACCAGGATTC-3'; *Arginase1*-rev, 5'-CGATGTCTTTGGCAGATATGC-3'; *Fizz1*-fwd, 5'-TGTGGCTTTGCCTGTGGAT-3'; *Fizz1*-rev, 5'-TCTTAGGACAGTTGGCAGCAG-3'; *Ym1*-fwd, 5'-GGGTTGGTTATGACAATGTCAG-3'; e *Ym1*-rev, 5'-TGAAGTCATCCATGTCCAGG-3'.

3.11. Citometria de fluxo para análises fenotípicas

Suspensões únicas de AMs e BMDMs de camundongos C57BL/6 foram marcadas com corantes do kit de viabilidade celular *LIVE/DEAD fixable blue dead cell stain kit* (*Invitrogen*) em PBS para excluir células mortas da análise. As células foram então coradas com anticorpos conjugados a fluorocromos contra os marcadores de superfície Cd11b (clone M1/70), F4/80 (clone BM8), CD206 (clone C068C2) e CD86 (clone GL-1), adquiridos da BD Biosciences (San Diego, CA), e dectina-1/CLEC7 (CD369), obtido da *BioLegend*, em PBS contendo 1% de SFB por 30 minutos em 4°C e então lavadas. A marcação foi realizada na presença de 5µg/ml de *Fc Block* (*BD Biosciences*). Aquisição de células foi realizada no citômetro de fluxo *BD FACSCanto II*, utilizando o programa *FACSDiva* (*BD Biosciences*). Para cada amostra, pelo menos 100.000 eventos foram coletados. Os dados foram analisados com o programa *FlowJo* (*TreeStar*).

3.12 – Declaração ética

Os cuidados com os animais neste trabalho foram realizados segundo as diretrizes institucionais de ética em experimentação animal e foram aprovadas pela Comissão de Ética em Experimentação Animal da Faculdade de Medicina de Ribeirão Preto (CETEA – Número de protocolos: 097/2010 e 002/2015). Para a eutanásia, os camundongos foram tratados com cetamina e xilazina (50 mg/kg e 10 mg/kg, respectivamente) por via intravenosa.

3.13. Análises Estatísticas

As análises estatísticas foram feitas utilizando-se o programa de análises *GraphPad Prism* (versão 5). Os resultados foram analisados por aplicação do teste estatístico para análise de variância (ANOVA), seguido de pós-teste Bonferroni ou pós-teste *t* de Student. As diferenças foram consideradas estatisticamente significativas se o valor *P* foi $< 0,05$.

4. RESULTADOS

4.1. Macrófagos derivados de medula óssea de camundongos A/J são mais suscetíveis à infecção por *Coxiella burnetii* fase II do que aqueles derivados de camundongos C57BL/6

C. burnetii apresenta diversos mecanismos de subversão da resposta imune do hospedeiro e, ainda assim, sabe-se que existem hospedeiros imunocompetentes capazes de controlar a replicação da bactéria. Entender de que forma o patógeno interage e manipula as respostas de seu hospedeiro é de grande utilidade para o desenvolvimento de novas estratégias eficientes de combate à infecção por essa bactéria, além de potencialmente elucidar novos mecanismos relacionados com a interação patógeno-hospedeiro e a suscetibilidade a *C. burnetii*. Nesse contexto, é fundamental a determinação e utilização de um modelo de estudo adequado.

Sabe-se que vários tipos de macrófagos murinos têm sido utilizados nos estudos que buscam entender as interações patógeno-hospedeiro que acontecem durante a infecção por *C. burnetii*. Diferenças de suscetibilidade entre macrófagos de linhagens murinas já foram observadas, com macrófagos das linhagens C57BL/6 sendo mais resistentes à infecção do que macrófagos derivados de camundongos A/J e BALB/c (ZAMBONI, 2004; ZAMBONI; RABINOVITCH, 2004). Essa diferença de suscetibilidade descrita por Zamboni (2004) foi observada através da análise quantitativa da formação de vacúolos de *C. burnetii*, uma das metodologias empregadas em estudos de suscetibilidade às infecções por essa bactéria. Dessa forma, nosso primeiro objetivo foi confirmar essa diferença de suscetibilidade descrita na literatura entre macrófagos derivados de camundongos das linhagens C57BL/6 e A/J quando infectados com *C. burnetii* fase II. Em adição, optamos por realizar a análise da suscetibilidade através da quantificação do DNA genômico bacteriano por PCR quantitativo (qPCR), uma metodologia confiável para análises envolvendo *C. burnetii*, como demonstrado por Brennan e Samuel (2003).

Para avaliação da suscetibilidade, BMDMs de camundongos C57BL/6 e A/J foram extraídos e diferenciados segundo protocolo descrito por Marim e colaboradores (2010). Posteriormente, esses macrófagos foram infectados com *C. burnetii* fase II utilizando-se uma proporção de 100 bactérias para cada célula (MOI, do inglês *Multiplicity Of Infection* de 100) por um período de 12 dias, sendo que a cada três dias um grupo de poços era lisado e

combinado com seus sobrenadantes para que toda bactéria ali presente fosse coletada. Como controle da entrada da bactéria nas células, poços foram lavados após quatro horas de infecção e lisados, com o material sendo coletado. Um grupo de células infectadas foi ainda tratado com 20µg/ml de rifampicina (RIF), um inibidor da tradução protéica bacteriana, como controle experimental interno da replicação de *C. burnetii* (BRENNAN; SAMUEL, 2003). Em seguida, o DNA genômico das amostras coletadas foi extraído, purificado e quantificado por qPCR.

Assim como descrito na literatura, BMDMs derivados de camundongos C57BL/6 mostraram-se restritivos à multiplicação de *C. burnetii*, controlando a replicação da bactéria e não permitindo um aumento relativo da quantidade de DNA genômico ao longo dos doze dias de infecção. Já BMDMs derivados de camundongos A/J mostraram-se suscetíveis, permitindo por volta de 1,5 *log* de crescimento bacteriano nesse mesmo período (Figura 5A). Essa diferença de suscetibilidade observada não foi decorrente de uma entrada diferencial da bactéria nos macrófagos, como pôde ser observado quatro horas após a infecção (5A, excerto em destaque). Para descartar uma possível influência da alta dose infecciosa utilizada (MOI 100) nas diferenças de suscetibilidade observadas entre os BMDMs, repetimos o experimento utilizando um MOI mais baixo, que se assemelha ao que ocorre em infecções naturais (MOI 3). Observamos que BMDMs derivados de camundongos A/J são mais suscetíveis à replicação bacteriana do que BMDMs da linhagem C57BL/6. Entretanto, diferentemente do que foi observado com MOI mais alto, em infecções realizadas com MOI 3 verificamos que BMDMs C57BL/6 permitiram um pequeno crescimento bacteriano ao longo dos 12 dias de infecção (Figura 5B). Em ambos os casos, o controle experimental interno com rifampicina funcionou, garantindo que de fato o que estava sendo quantificado era a replicação de *C. burnetii*. Esses dados confirmam as diferenças de suscetibilidade descritas na literatura entre as linhagens C57BL/6 e A/J frente à infecção por *C. burnetii* fase II, estendendo-as para BMDMs, além de indicar que MOIs mais baixos são mais adequados para esses experimentos. Em adição, os dados obtidos demonstram ainda a relevância do qPCR como ferramenta para análise de suscetibilidade à essa bactéria.

Figura 5 - Macrófagos murinos derivados de medula óssea da linhagem A/J são mais suscetíveis à multiplicação de *C. burnetii* fase II do que macrófagos murinos derivados de medula óssea da linhagem C57BL/6. BMDMs das linhagens C57BL/6 e A/J foram infectados com *C. burnetii* fase II em MOI 100 (A) ou 3 (B) e a multiplicação bacteriana foi avaliada por quantificação do DNA genômico por qPCR após 0 (4 horas após infecção), 3, 6, 9 e 12 dias de infecção. O eixo y representa a diferença logarítmica na base 10 entre a quantificação genômica no tempo X e a quantificação no tempo 0 ($\log T(X) - \log T(0)$). Células tratadas com o inibidor da tradução protéica bacteriana rifampicina (RIF) foram utilizadas como controle negativo da replicação bacteriana. Em destaque, os excertos mostram a quantidade de DNA genômico de *C. burnetii* 4 horas após a infecção. Barras de erro representam desvio padrão de triplicatas experimentais, e (*) indica $p < 0,05$; (*two-way* ANOVA). Os dados são representativos de três réplicas experimentais distintas. Adaptado de: FERNANDES et al (2016).

4.2. Macrófagos alveolares derivados de camundongos C57BL/6 são permissivos à replicação de *C. burnetii* fase II

Após a confirmação de que BMDMs derivados de camundongos C57BL/6 são restritivos à replicação de *C. burnetii* fase II, nosso próximo objetivo foi identificar um modelo de célula primária murina que permitisse a multiplicação da bactéria *in vitro*, uma vez que a maioria dos camundongos deficientes para diferentes genes foram gerados no fundo gênico C57BL/6. Levando em consideração o fato de que a infecção por *C. burnetii* ocorre via aerossol, infectando primariamente monócitos e macrófagos, com os macrófagos alveolares consistindo na célula-alvo primária da bactéria (VOTH; HOWE; HEINZEN, 2007; AMARA; BECHAH; MEGE, 2012), decidimos avaliar se macrófagos alveolares derivados de camundongos C57BL/6 seriam permissivos à replicação de *C. burnetii* fase II *in vitro*. Para isso, macrófagos alveolares (AMs) foram obtidos de camundongos C57BL/6 através de lavagem brônquio-alveolar e plaqueados em meio de cultura contendo antibiótico. No dia seguinte as células foram lavadas, o meio substituído por meio sem antibiótico e então foi feita a infecção com *C. burnetii* fase II em MOI 3. BMDMs derivados de camundongos C57BL/6 também foram utilizados para comparação da suscetibilidade desses dois tipos celulares à replicação bacteriana. Para ambas as células a infecção transcorreu por 12 dias e, novamente, as células foram lisadas e combinadas com seus respectivos sobrenadantes a cada três dias.

Após a extração, purificação e quantificação do DNA genômico bacteriano por qPCR, observamos que surpreendentemente AMs derivados de camundongos C57BL/6 foram altamente permissivos à replicação de *C. burnetii* fase II, permitindo um crescimento de quase três *logs* ao longo dos doze dias de infecção, em contraste ao que foi observado anteriormente para BMDMs (Figura 6A). Novamente não houve diferença na entrada da bactéria nesses dois tipos celulares, como observado pela quantificação de DNA genômico após quatro horas de infecção (Figura 6A, excerto em destaque). Em seguida, verificamos se a suscetibilidade observada em AMs da linhagem C57BL/6 também ocorreria em AMs derivados de camundongos A/J, uma vez que essa linhagem já se mostrou permissiva à replicação de *C. burnetii* fase II para infecções em BMDMs. Verificamos que AMs de ambas as linhagens analisadas apresentaram-se igualmente suscetíveis à replicação de *C. burnetii* (Figura 6B), sugerindo que mecanismos genéticos presentes em BMDMs que estão envolvidos na resistência natural desse tipo celular à infecção por *C. burnetii* podem estar ausentes ou ser menos eficientes em AMs. Nesse contexto, AMs mostraram-se como um interessante modelo

de macrófagos primários murinos altamente permissivos à replicação intracelular de *C. burnetii* fase II *in vitro*. Considerando a possível relevância desse modelo para estudos sobre as interações patógeno-hospedeiro durante as infecções com *C. burnetii*, decidimos caracterizar a replicação dessa bactéria em AMs.

Figura 6 – Macrófagos alveolares murinos são permissivos à replicação de *C. burnetii* fase II. BMDMs e AMs da linhagem C57BL/6 (A) e AMs das linhagens C57BL/6 e A/J (B) foram infectados com *C. burnetii* fase II com MOI 3. Após 0 (4 horas após infecção), 3, 6, 9, 12 e 15 dias de infecção foi avaliada a multiplicação bacteriana por quantificação do DNA genômico por qPCR. O eixo y representa a diferença logarítmica na base 10 entre a quantificação genômica no tempo X e a quantificação no tempo 0 ($\log T(X) - \log T(0)$). Células tratadas com o inibidor da tradução protéica bacteriana rifampicina (RIF) foram utilizadas como controle negativo da replicação bacteriana. Em destaque, os excertos mostram a quantidade de DNA genômico de *C. burnetii* 4 horas após a infecção. Barras de erro representam desvio padrão de triplicatas experimentais, e (*) indica $p < 0,05$; (two-way ANOVA). Os dados são representativos de três réplicas experimentais distintas. Adaptado de: FERNANDES et al (2016).

4.3. Macrófagos alveolares murinos suportam a replicação de grandes quantidades de bactéria intracelular nos vacúolos contendo *C. burnetii* fase II

Como observado anteriormente, AMs murinos mostraram-se um modelo celular altamente permissivo à replicação de *C. burnetii* fase II *in vitro*, mesmo no fundo gênico geralmente restritivo de camundongos C57BL/6. Para caracterização da replicação da bactéria nesse modelo optamos por utilizar técnicas de microscopia como ferramenta de estudo, como já padronizado anteriormente (ZAMBONI; MORTARA; RABINOVITCH, 2001). AMs derivados de camundongos C57BL/6 foram infectados com *C. burnetii* fase II em um MOI 3 por quatro horas, três dias e seis dias, fixadas e a quantidade de bactéria intracelular foi avaliada por meio de microscopia de fluorescência multifóton. Para isso, as células foram marcadas com anticorpo primário anti-*C. burnetii* e anticorpo primário anti-LAMP-1, a fim de acompanhar o aumento da quantidade de bactéria nas células e visualizar se as mesmas estariam replicando dentro dos vacúolos parasitóforos característicos, os quais se tornam decorados com esse marcador lisossomal (GHIGO et al., 2009). Observamos que ao longo dos primeiros seis dias de infecção ocorre um aumento na quantidade de *C. burnetii*, visualizado pelo aumento da marcação em vermelho dentro dos AMs, sugerindo que a bactéria está replicando nessas células. Em adição, uma associação com a marcação de LAMP-1, em verde, sugere que essa replicação está de fato ocorrendo dentro dos típicos vacúolos contendo *C. burnetii* (Figura 7A).

Apesar do aumento na quantidade de *C. burnetii* dentro dos AMs e a associação com LAMP-1 ter sido constatada, os processos envolvidos na preparação das amostras e marcação com anticorpos dos alvos de interesse para visualização em microscópio de fluorescência levaram à perda daquelas células que apresentam grandes e espaçosos vacúolos parasitóforos, já que essas células facilmente se soltam do fundo das lamínulas e são perdidas durante os processos de lavagem. Dessa forma, visando analisar a presença dos grandes vacúolos típicos que são desenvolvidos durante a infecção com *C. burnetii* em AMs, optamos por utilizar na infecção uma cepa de *C. burnetii* fase II expressando *mCherry* (BEARE et al., 2011). A análise dos AMs por microscopia de fluorescência multifóton mostrou novamente um aumento significativo na quantidade de bactéria ao longo do curso de 12 dias de infecção, representada pelo aumento da intensidade da marcação em vermelho proveniente das *C. burnetii* expressando *mCherry* (Figura 7B). Após seis dias de infecção já era possível observar que várias células continham grande quantidade de bactéria em vacúolos espaçosos, com essas bactérias tomando completamente o espaço celular após nove dias de infecção,

como ilustrado com a sobreposição das imagens de fluorescência e contraste diferencial de interferência (Figura 7C).

Além da microscopia de fluorescência, utilizamos a microscopia eletrônica de transmissão para caracterizar a replicação intracelular de *C. burnetii* fase II nos AMs (Figura 8). Novamente, AMs derivados de camundongos C57BL/6 foram infectados com *C. burnetii* fase II *in vitro* utilizando MOI 3, com as amostras sendo fixadas e preparadas para microscopia eletrônica de transmissão após três e seis dias, como descrito previamente por Zamboni e colaboradores (2002). Três dias após a infecção, as células mostravam grandes vacúolos espaçosos contendo em seu interior várias variantes celulares grandes de *C. burnetii* (Figura 8A). Após seis dias de infecção foi possível observar um aumento considerável na quantidade de bactérias dentro dos vacúolos (Figura 8B), demonstrando novamente que AMs são permissivos à replicação de *C. burnetii* fase II *in vitro* e podem ser utilizados como um modelo relevante de macrófagos primários para estudar infecções com essa bactéria.

Figura 7 – Localização intracelular de *C. burnetii* fase II em macrófagos alveolares murinos visualizada por microscopia confocal de fluorescência. AMs de camundongos C57BL/6 foram infectados com *C. burnetii* fase II selvagem (MOI 3) por 4 horas, 3 e 6 dias (A) ou infectadas com *C. burnetii* fase II expressando *mCherry* (MOI 3) por 4 horas, 3, 6, 9 e 12 dias (B). Após os tempos indicados, as células foram fixadas, processadas e então examinadas utilizando microscopia de fluorescência multifóton. (C) AMs infectados por 9 dias com *C. burnetii* expressando *mCherry*. As imagens foram adquiridas por microscopia de multifóton utilizando objetiva de óleo de imersão de aumento de 63X e analisadas utilizando o software *ImageJ*. DIC, do inglês *Differential Interference Contrast*. Verde, proteína lisossomal associada à membrana LAMP-1; vermelho, *C. burnetii*; ciano, DNA corado com DAPI. Barras de escala, 5µm. Adaptado de: FERNANDES et al (2016).

Figura 8 – Localização intracelular de *C. burnetii* fase II em macrófagos alveolares murinos visualizada por microscopia eletrônica de transmissão. AMs derivados de camundongos C57BL/6 foram infectados com *C. burnetii* fase II (MOI 3), fixados e processados para microscopia eletrônica de transmissão. Células foram fixadas após 3 (A) e 6 dias (B) de infecção. Os painéis da direita são imagens aumentadas das regiões destacadas nos painéis à esquerda. Setas, formas variantes celulares grandes de *C. burnetii*. Barras de escala, 1μm. Adaptado de: FERNANDES et al (2016).

4.4. Macrófagos alveolares murinos mostram uma polarização mais pronunciada em células do tipo M2 do que macrófagos murinos derivados de medula óssea

Após demonstrar por técnicas de microscopia que AMs murinos são de fato permissivos à replicação de *C. burnetii* fase II *in vitro*, nosso próximo objetivo foi buscar entender quais os mecanismos por trás da grande suscetibilidade apresentada por esse modelo. Sabe-se que a replicação e patogênese de *C. burnetii* estão relacionadas com o padrão de polarização induzido em macrófagos e monócitos, sendo que de forma geral a indução de um padrão M1 resulta em maior atividade pró-inflamatória, com desenvolvimento de funções microbidas que levam à eliminação do patógeno, enquanto que a indução de um padrão M2 reflete em ações com um perfil mais anti-inflamatório, permitindo assim a replicação do patógeno (MURRAY et al., 2017; SHAPOURI-MOGHADDAM et al., 2018). Em 2008, Benoit e colaboradores demonstraram que a infecção de macrófagos derivados de monócitos humanos por *C. burnetii* resultava na indução de um programa atípico de polarização em um estado M2, levando à persistência da bactéria nas células (BENOIT et al., 2008). Esse estado de polarização atípico era acompanhado pelo aumento na expressão de genes relacionados com o padrão M2, como Arginase-1, IL-10, TGF- β e receptores de manose, e pela diminuição na expressão de genes considerados marcadores clássicos do padrão M1, como TLR2 e TNF. Dessa forma, resolvemos investigar qual padrão de polarização estaria sendo induzido no modelo de AMs murinos.

Visando determinar o padrão de polarização apresentado por AMs murinos, avaliamos inicialmente a expressão de genes considerados como marcadores dos padrões M1 e M2 em AMs e comparamos com a expressão em BMDMs murinos, por meio de RT-PCR (do inglês, *Reverse Transcription-PCR*). Para isso, mRNA de culturas de 10^6 AMs e BMDMs de camundongos C57BL/6, infectados ou não por 12 horas com *C. burnetii* fase II (MOI 3), foram extraídos, purificados e quantificados. Após a realização de reações de termociclagem e leitura em um sistema de PCR em tempo real, a expressão dos genes de interesse foi calculada utilizando os valores do ciclo médio do limite (CT) para cada amostra, os quais foram normalizados com relação aos valores de CT dos genes *housekeeping*. Os genes *Nos2*, *Tnfa*, *Ifng* e *Tlr2* foram selecionados como marcadores representantes do padrão de polarização M1, enquanto que os genes *Arginase1*, *Fizz1* e *Ym1* foram selecionados como marcadores representantes do padrão de polarização M2. Para as análises em questão e normalização dos cálculos de CT, o gene *Gapdh* foi selecionado como representante do gene *housekeeping*.

Analisando inicialmente a expressão gênica em AMs e BMDMs murinos não infectados, observamos que AMs apresentam uma expressão consideravelmente reduzida de *Nos2*, *Tnfa*, *Ifng* e *Tlr2*, genes associados a um fenótipo de polarização padrão M1. Em contrapartida, a expressão dos genes *Arginase1*, *Fizz1* e *Ym1*, genes associados com um fenótipo de polarização padrão M2, mostraram-se aumentadas. Comparativamente, BMDMs murinos não infectados apresentaram um perfil inverso ao observado em AMs, com expressão aumentada dos genes associados ao padrão M1 e expressão reduzida de *Arginase1* e *Ym1* (Figura 9). Após a infecção com *C. burnetii* fase II por 12 horas, novamente AMs mostraram uma polarização padrão M2 pronunciada quando comparada com aquela apresentada por BMDMs. Essa forte polarização em um perfil M2 por AMs infectados foi representada pelo aumento na expressão de *Il4*, *Arginase1*, *Fizz1* e *Ym1* e redução na expressão de *Nos2*, *Tnfa*, *Ifng* e *Tlr2* em comparação com a expressão desses mesmos genes em BMDMs infectados (Figura 9). Conjuntamente, esses dados são consistentes com o perfil altamente suscetível dos AMs murinos observado frente à infecção *in vitro* por *C. burnetii* fase II, uma vez que essas células polarizariam para um padrão M2 mais tolerogênico.

Em concordância com os resultados obtidos, sabe-se que macrófagos com perfil de polarização M1 desencadeiam a produção de TNF- α , uma citocina pró-inflamatória já descrita como sendo importante para restrição da replicação de *C. burnetii* em macrófagos (BRADLEY et al., 2016). A restrição ao crescimento bacteriano observada em BMDMs murinos da linhagem C57BL/6 pode estar em parte relacionada justamente com essa polarização em um perfil mais M1, como observado pelo aumento na expressão do gene *Tnfa* após a infecção por *C. burnetii*, o que não ocorreria para AMs. Para continuar testando a hipótese de que a grande permissividade de AMs murinos à replicação de *C. burnetii* fase II seria decorrente de um estado de polarização padrão M2 nessas células, decidimos avaliar a expressão de alguns marcadores de superfície conhecidos classicamente relacionados com macrófagos polarizados nesse padrão, sendo eles o receptor de manose (CD206) e dectina-1 (CLEC7A) (MARTINEZ et al., 2006). Para isso, foram realizados experimentos de FACS (do inglês, *Fluorescence-Activated Cell Sorting*) em suspensões únicas de AMs e BMDMs não infectados ou infectados com *C. burnetii* fase II (MOI 3) por 48 horas, inicialmente marcadas com corantes do kit de viabilidade celular *LIVE/DEAD fixable blue dead cell stain kit* para exclusão das células mortas na análise. Aquelas que se mostraram viáveis foram posteriormente coradas com anticorpos conjugados a fluorocromos contra os marcadores de superfície Cd11b, F4/80, CD206, CD86 e dectina-1/CLEC7A, lavadas e então adquiridas em

citômetro de fluxo. Para cada amostra, pelo menos 100.000 eventos foram coletados. Os dados foram em seguida analisados com o programa *FlowJo* (*TreeStar*).

Para as análises de fenotipagem dos marcadores de superfície citados anteriormente, foram consideradas apenas as células vivas, duplo-positivas para Cd11b e F4/80 (Figuras 10A e 10B). Mesmo sem o estímulo infeccioso, AMs murinos mostraram expressão aumentada do receptor de manose (CD206) e dectina-1 (CLEC7A), moléculas associadas com o padrão de polarização M2, em comparação com a expressão dessas mesmas moléculas em BMDMs murinos (Figura 10C). Após a infecção com *C. burnetii* fase II, o mesmo cenário foi observado com relação ao padrão de expressão das moléculas associadas ao perfil de polarização M2 em AMs e BMDMs (Figura 10D). Adicionalmente, avaliamos os níveis de expressão da molécula co-estimulatória CD86, os quais se apresentaram semelhantes em AMs e BMDMs, infectados ou não (Figuras 10C e 10D, respectivamente). Coletivamente, esses dados são consistentes com a hipótese de que AMs apresentam maior suscetibilidade devido ao seu estado de polarização em um padrão M2, característica essa que já foi anteriormente relacionada com permissividade à replicação de *C. burnetii* (BENOIT; DESNUES; MEGE, 2008).

Figura 9 – Macrófagos alveolares murinos exibem níveis mais elevados de expressão de genes relacionados com o padrão de polarização M2. BMDMs e AMs derivados de camundongos C57BL/6 foram ou não infectados com *C. burnetii* fase II (MOI 3) por 12 horas. O RNA total das células foi extraído, e a reação de PCR de transcrição reversa (RT-PCR) foi realizada para detecção dos genes *Nos2*, *Tnfa*, *Ifng*, *Tlr2*, *Il4*, *Arginase1*, *Fizz1* e *Ym1*, genes considerados marcadores dos padrões de polarização M1 e M2. Os dados mostrados representam a média ± o desvio padrão de um experimento representativo de dois realizados. *, $P < 0.05$ comparando BMDMs e AMs; #, $P < 0.05$ comparando células infectadas com células não infectadas do respectivo grupo (*two-way* ANOVA). Adaptado de: FERNANDES et al (2016).

Figura 10 – Macrófagos alveolares murinos exibem um fenótipo de polarização M2. Estratégia de *gates* utilizada e frequência de BMDMs (A) e AMs (B) murinos derivados de camundongos C57BL/6 não infectados. FSC, do inglês *Forward Scatter*; SSC, do inglês *Side Scatter*. Fenotipagem das moléculas de superfície de BMDMs e AMs murinos não infectados (C) ou infectados por 48 horas (D) com *C. burnetii* fase II para a molécula co-estimulatória CD86 e para os marcadores do padrão de polarização M2 CD206 (receptor de manose) e dectina-1, conforme analisado por citometria de fluxo. Foram selecionadas células vivas Cd11b⁺ F4/80⁺. Adaptado de: FERNANDES et al (2016).

4.5. Macrófagos alveolares murinos deficientes em NOS2 e IFN γ são mais suscetíveis à replicação de *C. burnetii* fase II, enquanto células deficientes em interleucina-4 (IL-4) são mais restritivas à replicação da bactéria

Demonstramos anteriormente, por meio de técnicas de microscopia e quantificação de DNA genômico por qPCR, que AMs murinos são altamente suscetíveis à replicação de *C. burnetii* fase II *in vitro*. Determinamos, ainda, que essa permissividade aumentada ocorre devido ao fato de que essas células apresentam uma polarização padrão M2. Em conjunto, esses achados apontam para AMs como sendo um modelo relevante para estudos envolvendo as interações patógeno-hospedeiro em macrófagos primários frente as infecções *in vitro* com *C. burnetii* fase II. Sendo assim, buscamos validar esse modelo de estudo proposto.

Sabe-se que apesar da bactéria apresentar grande capacidade de evadir e subverter a resposta imune do hospedeiro, respostas efetivas ainda ocorrem. Um dos mecanismos importantes envolvidos na restrição do crescimento de *C. burnetii* se dá pela ação de espécies reativas de nitrogênio, com o óxido nítrico (NO) desempenhando um papel fundamental (YOSHII et al., 1999; HOWE et al., 2002; ZAMBONI; RABINOVITCH, 2003; BRENNAN et al., 2004; ZAMBONI; RABINOVITCH, 2004). Dessa forma, para fortalecer nossa proposta da relevância da utilização de AMs murinos como modelo de estudo das interações de *C. burnetii* com macrófagos murinos primários, nosso próximo passo foi avaliar a replicação da bactéria em AMs murinos deficientes para NOS2 (do inglês, *Nitric Oxide Synthase 2*; ou iNOS, do inglês *inducible Nitric Oxide Synthase*). Para isso, AMs murinos da linhagem C57BL/6, selvagens (WT) ou deficientes para NOS2 (*Nos2*^{-/-}), foram obtidos e infectados (MOI 3) com *C. burnetii* fase II *in vitro* por 12 dias, com as amostras sendo lisadas e coletadas em intervalos de três dias. Novamente um grupo contendo rifampicina foi utilizado como controle experimental interno. Após purificação das amostras e quantificação por qPCR, assim como esperado e em conformidade com o que já estava descrito na literatura, AMs de camundongos *Nos2*^{-/-} se mostraram mais permissivos à replicação de *C. burnetii* fase II do que AMs de camundongos WT (Figura 11A), reforçando a importância de NO para o controle da replicação por essa bactéria.

Em seguida, avaliamos também a replicação bacteriana em AMs derivados de camundongos deficientes para IFN- γ (*Ifng*^{-/-}), visto que essa molécula é essencial para produção de NO e, assim como NO, desempenha um papel microbicida fundamental nas infecções por *C. burnetii* (TURCO; THOMPSON; WINKLER, 1984; DELLACASAGRANDE et al., 1999; DINAKAR et al., 1999; GHIGO et al., 2002; AMARA;

BECHAH; MEGE, 2012). Observamos que ao longo de 12 dias de infecção AMs derivados de camundongos *Ifng*^{-/-} são significativamente mais permissivos à replicação de *C. burnetii* fase II do que AMs derivados de camundongos WT (Figura 11B). Coletivamente, esses resultados suportam o uso de AMs murinos como um modelo relevante para investigações envolvendo infecções com *C. burnetii* fase II em macrófagos primários de camundongos C57BL/6, a linhagem mais comumente utilizada para geração de camundongos transgênicos, uma ferramenta fundamental para estudos sobre interações patógeno-hospedeiro.

Posteriormente, para compreender melhor o efeito da polarização M2 de AMs murinos na alta suscetibilidade dessas células à infecção por *C. burnetii* fase II, infectamos AMs derivados de camundongos selvagens ou deficientes para IL-4 (*Il4*^{-/-}) e avaliamos a replicação bacteriana. Cabe ressaltar que IL-4 é um dos principais estímulos clássicos para indução da polarização no padrão M2, juntamente com IL-13, com ambos ativando STAT6 via receptor IL-4Rα (MANTOVANI et al., 2013; MURRAY, 2017; SHAPOURI-MOGHADDAM 2018). Vimos que na ausência de IL-4 os AMs tornam-se mais restritivos à replicação de *C. burnetii* (Figura 11C), o que pode ser explicado pelo fato de que sem IL-4 há um prejuízo na polarização dessas células para um padrão M2, mais suscetível ao crescimento bacteriano, corroborando os resultados obtidos até então. Importante reforçar que a ausência de NOS2, IFN-γ e IL-4 não interferiram na internalização da bactéria pelos AMs, como pode ser observado na quantificação do DNA genômico de *C. burnetii* após quatro horas de infecção (Figuras 11A-C, excertos em destaque). Em conjunto, esses dados suportam a hipótese de que a polarização em um padrão M1 ou M2 está por trás da suscetibilidade aumentada de AMs murinos à infecção por *C. burnetii* fase II em comparação com BMDMs. Além disso, é preciso destacar que, independentemente dos mecanismos que possam estar associados à resistência apresentada pelos BMDMs, ainda assim esses dados suportam o uso de AMs murinos como um importante e relevante modelo celular para análises da replicação de *C. burnetii* em macrófagos primários.

Figura 11 – Macrófagos alveolares murinos possibilitam a utilização de macrófagos deficientes no fundo genético C57BL/6 para avaliação da resposta imune inata contra *C. burnetii* fase II. AMs primários derivados de camundongos C57BL/6 WT e *Nos2*^{-/-} (A), *Ifng*^{-/-} (B) e *Il4*^{-/-} (C) foram infectados com *C. burnetii* fase II (MOI 3) e a multiplicação bacteriana foi avaliada por quantificação do DNA genômico por qPCR após 0 (4 horas após infecção), 3, 6, 9 e 12 dias de infecção. O eixo y representa a diferença logarítmica na base 10 entre a quantificação genômica no tempo X e a quantificação no tempo 0 ($\log T(X) - \log T(0)$). Células tratadas com o inibidor da tradução protéica bacteriana rifampicina (RIF) foram utilizadas como controle negativo da replicação bacteriana. Em destaque, os excertos mostram a quantidade de DNA genômico de *C. burnetii* 4 horas após a infecção. Barras de erro representam desvio padrão de triplicatas experimentais, e (*) indica $p < 0,05$; (*two-way* ANOVA). Os dados são representativos de três réplicas experimentais distintas. Adaptado de: FERNANDES et al (2016).

4.6. A permissividade dos macrófagos alveolares murinos à infecção por *C. burnetii* fase II é comparável a aquelas de linhagens celulares imortalizadas altamente suscetíveis

Conforme demonstramos anteriormente, nossos resultados mostraram que AMs murinos são permissivos à replicação de *C. burnetii* fase II *in vitro*. Por fim, para ressaltar a relevância e aplicabilidade desse modelo murino primário de infecção para estudos com *C. burnetii*, optamos por comparar a multiplicação dessa bactéria em AMs derivados de camundongos C57BL/6 com linhagens celulares imortalizadas altamente suscetíveis. Utilizamos células Vero, uma linhagem celular derivada de rins de macaco verde, e células THP-1, uma linhagem celular monocítica humana. Para esses experimentos as células Vero foram previamente irradiadas, a fim de garantir que apenas mais um ciclo de replicação ocorreria após o plaqueamento das mesmas. Desta forma, a quantidade de bactéria não seria diluída pela replicação das células. As células THP-1 foram previamente diferenciadas durante a noite em células do tipo macrófago por meio da adição de 200nM de forbol 12-miristato 13-acetato (PMA) na cultura, como descrito anteriormente (VOTH; HOWE; HEINZEN, 2007). Os AMs derivados de camundongos C57BL/6 foram obtidos e plaqueados como já descrito anteriormente. No dia seguinte, todas as células tiveram os meios de cultura trocados por meio fresco antes da infecção com *C. burnetii* fase II (MOI 3).

Tanto células Vero quanto células THP-1 já foram descritas como sendo altamente permissivas à replicação de *C. burnetii* (ZAMBONI; MORTARA; RABINOVITCH, 2001; GHIGO et al., 2002; HOWE et al., 2010), o que foi também observado por nós ao longo de 12 dias de infecção. Após seis dias de infecção com *C. burnetii* fase II em um MOI 3, AMs murinos se mostraram tão suscetíveis quanto as células Vero e as células THP-1 (Figura 12). Já nos tempos mais tardios da infecção, a replicação bacteriana observada em células Vero e células THP-1 foi mais robusta do que a apresentada por AMs murinos. Apesar disso, a taxa comparável de replicação de *C. burnetii* fase II durante os primeiros seis dias de infecção reforçam a demonstração de que AMs murinos são, de fato, altamente permissivos à replicação dessa bactéria e podem ser usados como um modelo celular primário do hospedeiro altamente relevante para investigações acerca da interação de macrófagos com *C. burnetii* fase II em cultura.

Figura 12 – Macrófagos alveolares murinos são permissivos à replicação de *C. burnetii* fase II *in vitro* de forma semelhante às células Vero e células THP-1. AMs derivados de camundongos C57BL/6, células Vero irradiadas e células THP-1 (previamente diferenciadas *overnight* com 200nM de PMA) foram infectadas com *C. burnetii* fase II (MOI 3) e a multiplicação bacteriana foi avaliada por quantificação do DNA genômico por qPCR após 0 (4 horas após infecção), 3, 6, 9 e 12 dias de infecção. O eixo y representa a diferença logarítmica na base 10 entre a quantificação genômica no tempo X e a quantificação no tempo 0 ($\log T(X) - \log T(0)$). Células tratadas com o inibidor da tradução protéica bacteriana rifampicina (RIF) foram utilizadas como controle negativo da replicação bacteriana. Em destaque, os excertos mostram a quantidade de DNA genômico de *C. burnetii* 4 horas após a infecção. Barras de erro representam desvio padrão de triplicatas experimentais, e (*) indica $p < 0,05$; (*two-way* ANOVA). Os dados são representativos de três réplicas experimentais distintas. Adaptado de: FERNANDES et al (2016).

5. DISCUSSÃO

Apesar do grande interesse biomédico de *C. burnetii* e do fato de que essa bactéria é extremamente virulenta e altamente adaptada à evasão da resposta imune, os mecanismos pelos quais os macrófagos e monócitos, células-alvo da infecção, reconhecem *C. burnetii*, bem como as interações patógeno-hospedeiro que acontecem durante a infecção, ainda precisam ser melhor compreendidos (AMARA; BECHAH; MEGE, 2012). O aprimoramento das técnicas de cultivo axênico da bactéria abriu caminho para manipulação genética mais eficiente de *C. burnetii*, possibilitando a geração de um número cada vez maior de bactérias mutantes (OMSLAND et al., 2009; OMSLAND et al., 2011). Como consequência, novos efetores bacterianos e suas funções têm sido descobertos, ajudando a compreender de que forma *C. burnetii* interage com seus hospedeiros (CUNHA et al., 2015; QIU; LUO, 2017; WALLQVIST et al., 2017; CRABILL et al., 2018; WEBER; FARIS, 2018). Entretanto, o entendimento das relações patógeno-hospedeiro não depende apenas do desenvolvimento de técnicas de manipulação do patógeno. É fundamental, também, determinar modelos de estudo apropriados. Nesse contexto, a falta de um modelo relevante de célula primária, principalmente um que permita a utilização dos diversos camundongos geneticamente modificados disponíveis no fundo gênico restritivo C57BL/6, é um problema fundamental que precisa ser endereçado. Isso nos levou, nesse trabalho, a buscar qual seria o modelo adequado de célula primária que pudesse ser utilizado nos estudos com *C. burnetii* fase II.

Já foi descrito na literatura que alguns hospedeiros conseguem restringir o crescimento de *C. burnetii*, com camundongos da linhagem C57BL/6 sendo resistentes à infecção, enquanto que camundongos das linhagens BALB/c e A/J são suscetíveis (ZAMBONI, 2004; ZAMBONI; RABINOVITCH, 2004). Os mecanismos existentes por trás dessa diferença de suscetibilidade entre as linhagens não são conhecidos, mas uma hipótese possível é que sejam decorrentes de características inerentes ao próprio fundo gênico dos camundongos. Há, ainda, diferenças de suscetibilidade dependendo do tipo celular analisado, com macrófagos peritoneais primários e macrófagos derivados de medula óssea apresentando-se mais intrinsecamente restritivos à bactéria, assim como monócitos, os quais apresentam um perfil de polarização do tipo M1 e são mais restritivos à replicação de *C. burnetii* (ZAMBONI et al., 2002; ZAMBONI; RABINOVITCH, 2003; BENOIT et al., 2008; CAPO; MEGE, 2012; ELDIN, 2017). Nesse caso, é possível que a diferença de suscetibilidade entre os tipos celulares seja reflexo de origens embrionárias distintas, ou mesmo de um microambiente tecidual diferente a qual cada célula é exposta.

A escolha de macrófagos como modelo celular para os estudos da interação patógeno-hospedeiro nas infecções com *C. burnetii* levou em consideração o fato de que essa é uma das principais células-alvo na infecção pela bactéria, além de considerar a plasticidade e relevância dos macrófagos na participação de diferentes funções nos organismos, tais como desenvolvimento e homeostase de tecidos, reparação tecidual, fagocitose de patógenos, produção de citocinas e outros aspectos da resposta imune (MANTOVANI et al., 2002; MANTOVANI et al., 2004; AMARA; BECHAH; MEGE, 2012; WYNN; CHAWLA; POLLARD, 2013). Para estudos com infecções por *C. burnetii*, macrófagos peritoneais murinos e macrófagos murinos derivados de medula óssea (BMDMs) são muito utilizados (ZAMBONI et al., 2002; ZAMBONI; RABINOVITCH, 2003; ZAMBONI, 2004; ZAMBONI; RABINOVITCH, 2004), com os últimos sendo de particular relevância visto que é possível congelar os BMDMs por longos períodos de tempo e ainda assim manter sua capacidade de diferenciação e funções de forma comparável a BMDMs frescos (MARIM et al., 2010). Além deles, os macrófagos alveolares compreendem outro modelo importante de estudo, uma vez que a infecção por *C. burnetii* se dá via aerossol. Dados da literatura mostram a utilização de macrófagos alveolares isolados de tecidos *post-mortem* de seres humanos, macrófagos alveolares derivados de macacos e macrófagos alveolares murinos para avaliação das interações patógeno-hospedeiro envolvendo infecções com *C. burnetii* (KHAVIN; TABIBZADEH, 1988; YOSHIIE et al., 1999; VOTH; HOWE; HEINZEN, 2007; CALVERLEY et al., 2012; GRAHAM et al., 2013; CUNHA et al., 2015; SOLTYSIAK; VAN SCHAIK; SAMUEL, 2015). Outros modelos incluem: macrófagos derivados de monócitos humanos, células da linhagem monocítica humana THP-1 e células Vero (derivadas de rim de macaco verde) (BURTON et al., 1978; DELLACASAGRANDE et al., 1999; ZAMBONI; MORTARA; RABINOVITCH, 2001; GHIGO et al., 2002; BENOIT et al., 2008; SHANNON; HEINZEN, 2008; HOWE et al., 2010). É importante ressaltar que, apesar da grande variedade de modelos de estudo com *C. burnetii*, muitos dos trabalhos focam apenas na suscetibilidade ao patógeno e não trazem evidências sobre quais os mecanismos estariam por trás desses fenômenos. Justamente por isso, nesse trabalho buscamos não apenas caracterizar um modelo relevante, mas também entender quais são os mecanismos por trás da suscetibilidade à *C. burnetii*.

Além da escolha do modelo de estudo, é importante ainda optar por um método eficiente para análise da suscetibilidade das células à infecção por *C. burnetii*. Tradicionalmente, a suscetibilidade a essa bactéria foi por muito tempo avaliada através da

estimativa do número de vacúolos parasitóforos nas células infectadas, bem como pela quantificação da intensidade de fluorescência em células infectadas marcadas com DAPI com o auxílio de microscópio de fluorescência confocal, o qual está diretamente correlacionado com a multiplicação do patógeno (ZAMBONI et al., 2001; ZAMBONI et al., 2002). Entretanto, essa metodologia pode incorrer em imprecisões dependendo da experiência e critérios utilizados por cada observador. Por exemplo, já foi descrito que ao menos para macrófagos alveolares humanos *C. burnetii* forma mais frequentemente múltiplos vacúolos pequenos ao invés de um grande vacúolo fusogênico (GRAHAM et al., 2013). Dessa forma, para que não houvesse uma diferença entre os observadores quanto à análise de suscetibilidade pelo critério “número de vacúolos”, seria necessário uma padronização prévia quanto à contagem de vacúolos de tamanhos e/ou graus de fusogenicidade distintos. Sendo assim, para evitar esses problemas, nesse trabalho optamos pela quantificação do DNA genômico bacteriano por PCR quantitativo (qPCR) como metodologia de análise da suscetibilidade dos macrófagos à infecção por *C. burnetii*. Esse é um método mais sensível e reprodutível, que não depende dos critérios de observação do experimentador e que já foi descrito como sendo eficiente para estudos com essa bactéria (BRENNAN; SAMUEL, 2003; KLEE et al., 2006a; KLEE et al., 2006b).

É importante ressaltar, ainda, que para a realização deste trabalho optamos por utilizar a forma avirulenta para seres humanos de *C. burnetii* derivada da cepa Nine Mile, conhecida como *C. burnetii* fase II ou NMII (do inglês, *Nine Mile II*). Essa cepa da bactéria, a qual se diferencia das formas virulentas devido ao tipo de LPS que apresenta, possui um clone que contém uma grande deleção cromossomal que resultou na impossibilidade de retornar a um estado completo de virulência, o que a torna apropriada para realização de estudos em condições de biossegurança nível 2. É importante salientar também que, apesar de apresentar essa deleção, essa bactéria não apresenta diferenças quanto ao tráfego intracelular, a cinética de replicação e a capacidade de restrição da sua multiplicação por parte dos hospedeiros quando em comparação com formas virulentas de *C. burnetii* da cepa Nine Mile, chamadas de *C. burnetii* fase I ou NMI (do inglês, *Nine Mile I*). (AMANO; WILLIAMS, 1984; HACKSTADT, 1996; HOWE et al., 2010; GRAHAM et al., 2013; LARSON et al., 2016).

Considerando então a importância biomédica de *C. burnetii* e a necessidade de uma melhor compreensão dos mecanismos envolvidos nas diferenças de suscetibilidade e no controle da infecção com esse patógeno por macrófagos, nosso trabalho buscou determinar um modelo primário de macrófagos que fosse adequado e relevante e que permitisse avaliar as

interações patógeno-hospedeiro nas infecções por *C. burnetii*. Para isso, optamos inicialmente por utilizar macrófagos murinos derivados de medula óssea em fundo genético C57BL/6 como modelo celular, tendo em vista a facilidade de obtenção dessas células e o grande número de camundongos geneticamente modificados disponíveis (MARIM et al., 2010). Foi escolhida a bactéria *C. burnetii* cepa NMII que apresenta a deleção genômica como modelo de infecção, pela possibilidade de se realizar os experimentos sem necessidade de uma estrutura de biossegurança com maior complexidade do que a disponível em nosso laboratório. Por fim, determinamos a utilização de qPCR como método principal para análise da suscetibilidade, considerando sua eficácia, sensibilidade e reprodutibilidade (BRENNAN; SAMUEL, 2003; KLEE et al., 2006a; KLEE et al., 2006b).

Primeiramente, testamos a suscetibilidade de BMDMs de camundongos C57BL/6 frente à infecção por *C. burnetii* fase II e comparamos com BMDMs da linhagem A/J. Em concordância com dados da literatura, vimos que as células em fundo genético C57BL/6 controlaram a replicação da bactéria, enquanto que aquelas no fundo genético A/J foram permissíveis ao crescimento bacteriano, evidenciado pelo aumento na quantificação do DNA genômico de *C. burnetii* ao longo dos 12 dias de infecção (Figura 5). Uma vez que a maioria dos camundongos geneticamente modificados encontram-se no fundo genético C57BL/6 e este se mostrou restritivo à replicação de *C. burnetii*, buscamos outro modelo de macrófagos primários que permitissem o crescimento bacteriano. Demonstramos que AMs são altamente suscetíveis à replicação de *C. burnetii* fase II, independente do fundo genético do camundongo (Figura 6), o que nos sugeriu que esse poderia ser um modelo relevante para se estudar as interações patógeno-hospedeiro nas infecções com essa bactéria. Vale ressaltar que trabalhos anteriores já haviam descrito que macrófagos alveolares de primatas e macrófagos alveolares humanos são permissivos à replicação de *C. burnetii*, assim como observamos em nosso modelo murino (KHAVIN; TABIBZADEH, 1988; YOSHIE et al., 1999; VOTH; HOWE; HEINZEN, 2007; CALVERLEY et al., 2012; GRAHAM et al., 2013; SOLTYSIAK; VAN SCHAIK; SAMUEL, 2015). Isso sugere que a suscetibilidade à infecção possa estar mais relacionada com o tipo de macrófago em si do que com a espécie ou linhagem analisada, ou seja, que essa suscetibilidade aumentada seja uma característica inerente ao macrófago alveolar. Apesar desses achados, os trabalhos anteriores não se aprofundaram na busca pelos mecanismos que explicassem essa permissividade aumentada apresentada pelos AMs, um ponto fundamental para compreensão da resposta à infecção por *C. burnetii*. Isso nos levou

não apenas a buscar e caracterizar AMs como um modelo de estudos relevante, mas ir mais a fundo na busca dos mecanismos que explicassem a suscetibilidade ao patógeno.

Por meio do uso de microscopia de fluorescência multifóton e microscopia eletrônica de transmissão, fornecemos evidências visuais de que *C. burnetii* fase II prontamente replica no interior de AMs (Figuras 7 e 8). Diferentemente da utilização de microscopia de fluorescência como metodologia para quantificar a suscetibilidade ao patógeno, como descrito por Zamboni e colaboradores (2001, 2002), nossa utilização desse método se deu apenas de forma qualitativa como ferramenta para demonstrar a multiplicação de *C. burnetii*. Mostramos ao longo dos 12 dias de infecção através do aumento na intensidade da fluorescência decorrente da marcação da bactéria corada com anticorpos, assim como da bactéria expressando *mCherry*, que macrófagos alveolares murinos são altamente permissíveis à *C. burnetii* de uma forma tal que a bactéria passa a ocupar completamente todo o espaço celular após nove dias de infecção (Figura 7C). O aumento na intensidade da fluorescência proveniente da bactéria expressando *mCherry* demonstra ainda que a bactéria se encontra viva e metabolicamente ativa, reforçando a viabilidade de AMs como modelo de estudo. Além disso, demonstramos através da marcação da proteína LAMP-1 e da visualização da célula infectada por microscopia eletrônica de transmissão que a bactéria replica no interior de seu vacúolo característico. A visualização de grandes quantidades da forma variante celular grande de *C. burnetii* no interior dos vacúolos também corrobora a hipótese de que a bactéria está replicando no interior dos AMs, visto que essa é a forma metabolicamente ativa e replicante do patógeno (COLEMAN et al., 2004).

Com base em nossas observações indicando que AMs murinos são altamente permissivos à replicação de *C. burnetii* fase II *in vitro* em comparação com BMDMs murinos, os quais se mostram resistentes à infecção, e visando elucidar quais os mecanismos por trás dessa suscetibilidade aumentada, hipotetizamos que essa diferença de suscetibilidade poderia ser decorrente da aquisição de padrões distintos de polarização entre esses tipos celulares. Dados da literatura destacam a importância do estado de polarização de macrófagos e monócitos após o encontro com *C. burnetii* para o desenvolvimento de respostas que podem levar ao controle da infecção ou à sobrevivência da bactéria nos hospedeiros (AMARA; BECHAH; MEGE, 2012). Em conjunto com esses dados, evidências recentes indicam que AMs são originados a partir de monócitos fetais de vida longa que migram para os pulmões e lá permanecem como macrófagos alveolares residentes, sendo mantidos através de auto-renovação, muito mais do que uma manutenção envolvendo a substituição das células mortas

por novos monócitos circulantes derivados de células da medula óssea (GUILLIAMS et al., 2013). Como resultado, o microambiente tecidual ao qual essas células estariam expostas seria muito diferente do microambiente em que se encontrariam os BMDMs, de forma que diferenças epigenéticas e transcriptômicas seriam impressas nessas células resultando em um padrão de resposta fenotípico distinto entre elas, de acordo com o que observamos (GORDON et al., 2014; GOSELIN et al., 2014; LAVIN et al., 2014).

Para testarmos a hipótese de que diferenças no estado de polarização dos macrófagos seriam o mecanismo responsável pela alta permissividade dos AMs murinos à infecção por *C. burnetii* fase II, avaliamos a expressão de genes comumente associados com padrões de polarização do tipo M1 ou M2 por meio de RT-PCR. De forma geral, macrófagos que apresentam polarização em um perfil M1 produzem grandes quantidades de citocinas pró-inflamatórias e são considerados microbicidas, enquanto que macrófagos polarizados em um perfil M2 são geralmente anti-inflamatórios e permitem a sobrevivência de patógenos (MANTOVANI et al., 2004; MANTOVANI et al., 2013; MURRAY, 2017; SHAPOURI-MOGHADDAM 2018). Nossos resultados mostraram que AMs murinos apresentam níveis mais elevados de expressão de genes relacionados com a polarização no padrão M2, como Arginase-1, Fizz1, Ym1 e IL-4, enquanto que em BMDMs a expressão de genes relacionados com o padrão M1, como NOS2, IFN- γ , TNF- α e TLR2, é mais elevada. É interessante ressaltar que essa diferença entre AMs e BMDMs foi observada mesmo sem a infecção por *C. burnetii*, e a infecção com esse patógeno apenas acentuou e reforçou o que fora obtido anteriormente (Figura 9). Considerando a existência de dois modelos, sendo um deles o da ontogenia e o outro o da plasticidade (BONNARDEL. GUILLIAMS, 2018) isso fortaleceria a hipótese de que para a determinação do estado de polarização do macrófago e, conseqüentemente, para o desempenho de suas funções especializadas, a ontogenia seria prevalente sobre a plasticidade. Em outras palavras, o fato de que AMs e BMDMs murinos submetidos às mesmas condições *in vitro* apresentam um perfil de expressão gênica que se relaciona com padrões distintos de polarização sugere que, para a determinação do estado de polarização do macrófago, de onde a célula veio seria mais importante do que o microambiente ao qual ela estaria sujeita.

Outra metodologia utilizada nesse trabalho para confirmar a hipótese de que AMs apresentam um perfil de polarização do tipo M2 que os tornam mais permissivos à replicação de *C. burnetii* fase II *in vitro* foi a fenotipagem dos marcadores de superfície dessas células por meio de citometria de fluxo. Confirmando os dados obtidos com RT-PCR, AMs mostraram expressão aumentada de moléculas de superfície consideradas como marca

registrada do perfil de polarização M2, sendo elas o receptor de manose (CD206) e o receptor delectina-1, quando comparados com BMDMs (Figura 10). Nossos resultados vão de acordo com o que já foi descrito na literatura para macrófagos alveolares humanos, nos quais *C. burnetii* induz a ativação de um perfil atípico de polarização M2 (BENOIT et al., 2008). Outros trabalhos também relataram essa permissividade à replicação de *C. burnetii* em macrófagos alveolares primários humanos e macrófagos residentes murinos, entretanto não buscaram determinar qual seria o mecanismo responsável por essa suscetibilidade (CALVERLEY et al., 2012; GRAHAM et al., 2013). É possível que para esses trabalhos, assim como observado no nosso trabalho e por Benoit e colaboradores (2008), essa permissividade seja resultado de um perfil de polarização do tipo M2. Vale destacar que, na realidade, os padrões de polarização M1 e M2 não são únicos, e sim representam dois extremos de um contínuo de estados funcionais polarizados. O padrão M2 subdivide-se em M2a, M2b, M2c e, para camundongos, existe ainda o estado M2d. Cada um deles é induzido frente a estímulos distintos e, como resultado, apresentam funções especializadas particulares. Considerando o padrão de expressão de genes e moléculas de superfície obtido em nossos resultados para AMs murinos frente à infecção por *C. burnetii* fase II *in vitro*, é possível sugerir que nesse caso ocorre uma polarização no perfil M2a ou M2c. Os níveis elevados de expressão de Arginase-1 e receptor de manose, assim como o desempenho de funções relacionadas com o perfil de resposta mais do tipo II e mais anti-inflamatório suportam essa hipótese. Em conjunto, essas características apresentadas pelos AMs possibilitam a sobrevivência do patógeno (MANTOVANI et al., 2004). Estudos avaliando uma maior quantidade de marcadores associados com padrões de polarização poderão determinar a qual subgrupo específico do padrão M2 se enquadram os AMs murinos após infecção com *C. burnetii* fase II.

Corroboramos ainda a hipótese de que a alta permissividade dos AMs murinos observada em nossos experimentos era decorrente de uma polarização em perfil M2 por meio da infecção de células derivadas de camundongos deficientes para IL-4. Já está bem estabelecido na literatura que IL-4 consiste em uma das mais importantes citocinas para a indução do estado de polarização M2, juntamente com IL-13, via ativação de STAT6 pelo receptor IL-4R α (GORDON, 2003; MANTOVANI et al., 2013; MURRAY, 2017; SHAPOURI-MOGHADDAM 2018). Sendo assim, era esperado que AMs derivados de camundongos *Il-4*^{-/-} apresentassem um prejuízo na polarização em perfil M2 quando em comparação com as mesmas células derivadas de camundongos selvagens. Nossos resultados

confirmaram essa hipótese, com AMs derivados de camundongos *Il-4^{-/-}* se mostrando significativamente mais restritivos à replicação de *C. burnetii* fase II do que AMs derivados de camundongos WT (Figura 11C).

Para fortalecer a relevância de AMs murinos como um novo modelo primário de macrófagos para estudos com infecções *in vitro* por *C. burnetii* fase II, como proposto nesse trabalho, infectamos AMs derivados de camundongos *Ifng^{-/-}* e *Nos2^{-/-}* como prova de princípio. Diversos trabalhos na literatura mostram que a produção de óxido nítrico e espécies reativas intermediárias de nitrogênio são de extrema importância na resposta imune contra patógenos respiratórios, com NO apresentando efeitos citotóxicos em níveis elevados (NATHAN; SHILOH, 2000). Além disso, já está bem estabelecido que NO participa da restrição da replicação intracelular de *C. burnetii* (HOWE et al., 2002; ZAMBONI; RABINOVITCH, 2003; BRENNAN et al., 2004; ZAMBONI, RABINOVITCH, 2004; HILL; SAMUEL, 2011), sendo assim a ausência dessa molécula resultaria em uma maior permissividade à multiplicação da bactéria. Isso foi de fato observado em nossos estudos (Figura 11A). A ausência de IFN- γ também deveria levar a um aumento na suscetibilidade à *C. burnetii*, visto que essa molécula não só participa da indução de um estado de polarização no perfil M1, mas também é fundamental para a produção de NO. Como esperado, AMs derivados de camundongos *Ifng^{-/-}* permitiram a replicação de maiores quantidades da bactéria (Figura 11B). Nesse trabalho, demonstramos também que AMs derivados de camundongos C57BL/6 suportam a replicação intracelular de *C. burnetii* e que essa replicação ocorre de maneira comparável a aquela observada em linhagens celulares imortalizadas que são sabidamente altamente suscetíveis à *C. burnetii*, como células Vero e células THP-1 (Figura 12) (BURTON et al., 1978; DELLACASAGRANDE et al., 1999; ZAMBONI; MORTARA; RABINOVITCH, 2001; GHIGO et al., 2002; BENOIT et al., 2008; SHANNON; HEINZEN, 2008; HOWE et al., 2010). Em conjunto, esses resultados validaram a utilização de AMs murinos como um modelo celular relevante para os estudos *in vitro* com a infecção por *C. burnetii* fase II.

Por fim, vale ressaltar que em oposição ao que observamos em nossos experimentos e ao que já estava descrito na literatura, Cockrell e colaboradores (2017) relataram um crescimento de quase três logs de *C. burnetii* fase II em BMDMs derivados de camundongos no fundo gênico restritivo C57BL/6. Esses pesquisadores explicam as diferenças observadas com base na fonte de M-CSF utilizada em ambos os estudos: enquanto nosso grupo e os demais trabalhos da literatura utilizaram meio condicionado de células L929 como a fonte de

M-CSF, nos experimentos publicados por Cockrell e colaboradores (2017) foi utilizado M-CSF recombinante murino. Ainda segundo os autores, o meio condicionado de células L929 conteria componentes não caracterizados que poderiam interferir no estado de ativação do macrófago, tais como quantidades variáveis de interferons do tipo I, o que refletiria diretamente na permissividade à infecção uma vez que o estado de polarização adotado pelos macrófagos é um mecanismo essencial da resposta imune à *C. burnetii* e interferons do tipo I são fortes polarizadores do padrão M1 (MANTOVANI et al., 2004; AMARA; BECHAH; MEGE, 2012; BISWAS et al., 2012; LOCATI; MANTOVANI; SICA, 2013; MURRAY et al., 2014; MURRAY, 2017; SHAPOURI-MOGHADDAM et al., 2018).

Os resultados obtidos por Cockrell e colaboradores (2017) não invalidam nossos achados de que AMs apresentam um perfil de polarização do tipo M2 que os tornam altamente permissivos à replicação de *C. burnetii* fase II *in vitro*. Entretanto, eles nos levam a pensar que talvez BMDMs também possam ser de fato utilizados como outra opção de modelo primário de macrófagos murinos para estudos de suscetibilidade a essa bactéria, desde que para isso seja utilizada uma fonte de M-CSF que não contenha outros componentes que possam potencialmente alterar o estado de polarização dos macrófagos. Por fim, nosso trabalho apresentou e caracterizou macrófagos alveolares murinos como um modelo relevante de células primárias altamente suscetíveis à replicação de *C. burnetii* fase II *in vitro* que pode ser empregado em estudos visando um melhor entendimento entre as relações patógeno-hospedeiro que acontecem nas infecções por essa bactéria.

6. SUMÁRIO

- Macrófagos derivados de medula óssea (BMDMs) de camundongos C57BL/6 são mais restritivos à replicação de *C. burnetii* fase II do que aqueles derivados de camundongos A/J.
- Macrófagos alveolares (AMs) no fundo gênico restritivo C57BL/6 são permissivos à replicação de *C. burnetii* fase II, em contraste com o que ocorre para BMDMs dessa mesma linhagem.
- AMs derivados de camundongos C57BL/6 são tão suscetíveis à replicação de *C. burnetii* fase II quanto AMs derivados de camundongos A/J.
- A replicação de *C. burnetii* fase II em AMs ocorre dentro dos vacúolos parasitóforos característicos e pode ser observada por meio de microscopia confocal de fluorescência e microscopia eletrônica de transmissão.
- AMs murinos apresentam um perfil de polarização do tipo M2, enquanto para BMDMs é observado um perfil de polarização do tipo M1.
- AMs murinos apresentam maior expressão de genes relacionados com o padrão M2 de polarização do que BMDMs murinos, tanto em células não infectadas quanto em células infectadas com *C. burnetii* fase II.
- A fenotipagem de AMs murinos mostra uma maior expressão de moléculas de superfície relacionadas com o padrão M2 de polarização do que a observada em BMDMs murinos, tanto em células não infectadas quanto em células infectadas com *C. burnetii* fase II.
- AMs em fundo gênico C57BL/6 deficientes para NOS2 e IFN- γ são mais suscetíveis à replicação de *C. burnetii* fase II.
- AMs em fundo gênico C57BL/6 deficientes para IL-4 são mais restritivos à replicação de *C. burnetii* fase II.
- A polarização em um padrão M1 ou M2 está relacionada à alta permissividade de AMs murinos à infecção por *C. burnetii* fase II em comparação com BMDMs murinos.
- A suscetibilidade aumentada de AMs murinos à infecção por *C. burnetii* fase II é comparável com aquela observada para células Vero e células THP-1.

- AMs murinos constituem um modelo primário de células suscetíveis altamente relevante para investigações acerca das interações patógeno-hospedeiro envolvendo a infecção com *C. burnetii* fase II *in vitro*.

7.CONCLUSÃO

Os dados obtidos permitem concluir que macrófagos alveolares murinos compreendem um modelo celular primário altamente permissivo à replicação da bactéria *C. burnetii* fase II e que essa suscetibilidade aumentada deve-se à indução de um estado de polarização do tipo M2. Dessa forma, nossos achados indicam que os macrófagos alveolares murinos podem ser utilizados como um relevante modelo de estudos das interações patógeno-hospedeiro em células primárias frente à infecção *in vitro* por *C. burnetii* fase II, permitindo inclusive a utilização de células derivadas de camundongos geneticamente modificados no fundo gênico C57BL/6.

8. REFERÊNCIAS BIBLIOGRÁFICAS¹

- AGUILERA, M. et al. Actin dynamics and Rho GTPases regulate the size and formation of parasitophorous vacuoles containing *Coxiella burnetii*. **Infection and Immunity**, v. 77, p. 4609-4620, 2009.
- AKIRA, S.; TAKEDA, K. Toll-like receptor signalling. **Nature Reviews Immunology**, v. 4, n. 7, p. 499-511, 2004.
- AMANO, K.; WILLIAMS, J. C. Chemical and immunological characterization of lipopolysaccharides from Phase I and Phase II *Coxiella burnetii*. **Journal of Bacteriology**, v. 160, p. 994-1002, 1984.
- AMARA, A. B.; BECHAH, Y.; MEGE, J. L. Immune Response and *Coxiella burnetii* Invasion. **Advances in Experimental Biology and Medicine**, v. 984, p. 287-298, 2012.
- ANDERSON, A. et al. Diagnosis and management of Q fever – United States, recommendations from CDC and the Q Fever Working group. **Morbidity and Mortality Weekly Report. Recommendations and Reports**, v. 62, p. 1-30, 2013.
- ANDOH, M. et al. T cells are essential for bacterial clearance, and gamma interferon, tumor necrosis factor alpha, and B cells are crucial for disease development in *Coxiella burnetii* infection in mice. **Infection and Immunity**, v. 75, p. 3245-3255, 2007.
- ARRICAU-BOUVERY, N. et al. Molecular characterization of *Coxiella burnetii* isolates by infrequent restriction site-PCR and MLVA typing. **BMC Microbiology**, v. 6, p. 38, 2006.
- BACA, O. G.; KLASSEN, D. A.; ARAGON, A. S. Entry of *Coxiella burnetii* into host cells. **Acta Virologica**, v. 37, p. 143-155, 1993.
- BARRY, A. O. et al. Impaired stimulation of p38alpha-MAPK/Vps41-HOPS by LPS from pathogenic *Coxiella burnetii* prevents trafficking to microbicidal phagolysosomes. **Cell Host Microbe**, v. 12, p. 751-763.
- BEARE, P. A. et al. Dot/Icm type IVB secretion system requirements for *Coxiella burnetii* growth in human macrophages. **mBio**, v. 2, p. e00175-11, 2011.
- BENOIT, M. et al. *Coxiella burnetii*, the agent of Q fever, stimulates an atypical M2 activation program in human macrophages. **European Journal of Immunology**, v. 38, p. 1065-1070, 2008.
- BENOIT, M. et al. The uptake of apoptotic cells drives *Coxiella burnetii* replication and macrophage polarization: a model for Q fever endocarditis. **PLoS Pathogens**, v. 4, n. 5, p. e1000066, 2008.
- BENOIT, M.; DESNUES, B.; MEGE, J. L. Macrophage polarization in bacterial infections. **Journal of Immunology**, v. 181, p. 3733-3739, 2008.

¹¹ ¹De acordo com a Associação Brasileira de Normas Técnicas (ABNT NBR 6023)

BERON, W. et al. *Coxiella burnetii* localizes in a Rab7-labeled compartment with autophagic characteristics. **Infection and Immunity**, v. 70, p. 5816-5821, 2002.

BISWAS, S. K. et al. Macrophage polarization and plasticity in health and disease. **Immunologic Research**, n. 53, v. 1-3, p. 11-24, 2012.

BONNARDEL, J.; GUILLIAMS, M. Developmental control of macrophage function. **Current Opinion in Immunology**, v. 50, p. 64-74, 2018.

BRADLEY, W. P. et al. Primary role for Toll-like receptor-drive tumor necrosis factor rather than cytosolic immune detection in restricting *Coxiella burnetii* phase II replication within mouse macrophages **Infection and Immunity**, v. 84, p. 998-1015, 2016.

BRENNAN, R. E. et al. Both inducible nitric oxide synthase and NADPH oxidase contribute to the control of virulent phase I *Coxiella burnetii* infections. **Infection and Immunity**, v. 72, p. 6666-6675, 2004.

BRENNAN, R. E.; SAMUEL, J. E. Evaluation of *Coxiella burnetii* Antibiotic Susceptibilities by Real-Time PCR Assay. **Journal of Clinical Microbiology**, v. 41, n. 5, p. 1869-1874, 2003.

BURNET, F. M.; FREEMAN, M. Experimental studies on the virus of "Q" fever. **Reviews of Infectious Diseases**, v. 5, n. 4, p. 800-8, 1983.

BURTON, P. R. et al. Some ultrastructural effects of persistent infections by the rickettsia *Coxiella burnetii* in mouse L cells and green monkey kidney (Vero) cells. **Infection and Immunity**, v. 21, n. 2, p. 556-66, 1978.

CALVERLET, M. et al. Resident alveolar macrophages are susceptible to and permissive of *Coxiella burnetii* infection. **PLoS One**, v. 7, p. e51941, 2012.

CAMPOY, E. M.; MANSILLA, M. E.; COLOMBO, M. I. Endocytic SNAREs are involved in optimal *Coxiella burnetii* vacuole development. **Cellular Microbiology**, v. 15, p. 922e41, 2013.

CAMPOY, E. M.; ZOPPINO, F. C. M.; COLOMBO, M. I. The early secretory pathway contributes to the growth of the *Coxiella* replicative niche. **Infection and Immunity**, v. 79, n. 1, p. 402-13, 2011.

CAPO, C. et al. Subversion of monocyte functions by *Coxiella burnetii* impairment of the cross-talk between $\alpha\beta3$ integrin and CR3. **Journal of Immunology**, v. 163, p. 6078-6085, 1999.

CAPO, C.; MEGE, J. L. Role of Innate and Adaptive Immunity in the Control of Q Fever. **Advances in Experimental Biology and Medicine**, v. 984, p.273-286, 2012.

CAREY, K. L. et al. The *Coxiella burnetii* Dot/Icm system delivers a unique repertoire of type IV effectors into host cells and is required for intracellular replication. **PLoS Pathogens**, v. 7, p. e1002056.

CELADA, A. et al. Evidence for a gamma-interferon receptor that regulates macrophage tumoricidal activity. **Journal of Experimental Medicine**, v. 160, p. 55-74, 1984.

CHISTIAKOV, D. A.; BOBRYSEV, Y. V.; OREKHOV, A. N. Changes in transcriptome of macrophages in atherosclerosis. **Journal of Cellular and Molecular Medicine**, v. 19, n. 6, p. 1163-73, 2015.

COCKRELL, D. C. et al. Robust growth of avirulent phase II *Coxiella burnetii* in bone marrow-derived murine macrophages. **PLoS ONE**, v. 12, n. 3, p. e0173528, 2017.

COLEMAN, S. A. et al. Temporal analysis of *Coxiella burnetii* morphological differentiation. **Journal of Bacteriology**, v. 186, p. 7344-7352, 2004.

COSTA, P. S.; BRIGATTE, M. E.; GRECO, D. B. Questing one Brazilian query: reporting 16 cases of Q fever from Minas Gerais, Brazil. **Revista do Instituto de Medicina Tropical de São Paulo**, v. 48, n. 1, p.5-9, 2006.

CRABILL, E. et al. Dot/Icm-Translocated Proteins Important for Biogenesis of the *Coxiella burnetii*-Containing Vacuole Identified by Screening of an Effector Mutant Sublibrary. **Infection and Immunity**, v. 88, n. 4, p. e00758-17, 2018.

CREAGH, E. M.; O'NEILL, L. A. TLRs, NLRs and RLRs: a trinity of pathogen sensors that co-operate in innate immunity. **Trends in Immunology**, v. 27, n. 8, p. 352-7, 2006.

CUNHA, L. D. et al. Inhibition of inflammasome activation by *Coxiella burnetii* type IV secretion system effector IcaA. **Nature Communications**, v. 6, p. 10205, 2015.

CUNHA, L. D.; ZAMBONI, D. S. Subversion of inflammasome activation and pyroptosis by pathogenic bacteria. **Frontiers in Cellular and Infection Microbiology**, v. 3, p. 76, 2013.

DELLACASAGRANDE, J. et al. IFN- γ mediated control of *Coxiella burnetii* survival in monocytes: the role of cell apoptosis and TNF. **Journal of Immunology**, v. 150, n. 162, p. 2259-2264, 1999.

DERRICK, E. H. "Q" fever, a new fever entity: clinical features, diagnosis and laboratory investigation. **Reviews of Infectious Diseases**, v. 5, n. 4, p. 790-800, 1983.

DINAKAR, C. et al. Differential regulation of human blood monocyte and alveolar macrophage inflammatory cytokine production by nitric oxide. **Annals of Ashtma Immunology**, v. 82, p. 217-222, 1999.

DOROTHÉE, D. et al. Tumor-associated leukemia inhibitory factor and IL-6 skew monocyte differentiation into tumor-associated macrophage-like cells. **Blood**, v. 110, p. 4319-4330, 2007.

DOYLE, A. G. et al. Interleukin-13 alters the activation state of murine macrophages in vitro: comparison with interleukin-4 and interferon-gamma. **European Journal of Immunology**, v. 24, p. 1441-5, 1994.

DYER, R. E. A filter passing infectious agent isolated from ticks IV Human infection. **Public Health Reports**, v. 53, p. 2277-2282, 1938.

DYER, R. E. Similarity of Australian "Q" fever and a disease caused by na infectious agent isolated from ticks in Montana. **Public Health Reports**, v. 54, p. 1229-1237, 1939.

ELDIN, C. et al. From Q fever to *Coxiella burnetii* infection: a paradigm change. **Clinical Microbiology Reviews**, v. 30, p. 115-190, 2017.

ENSERINK, M. Questions abound in Q-fever explosion in Netherlands. **Science**, v. 327, n. 5963, p. 266-8, 2010.

FERNANDES, T. D. et al. Murine Alveolar Macrophages Are Highly Susceptible to Replication of *Coxiella burnetii* Phase II In Vitro. **Infection and Immunity**, v. 84, n. 9, p. 2439-48, 2016.

GHIGO, E. et al. *Coxiella burnetii* survival in THP-1 monocytes involves the impairment of phagosome maturation: IFN- γ mediates its restoration and bacterial killing. **Journal of Immunology**, v. 69, p. 4488-4495, 2002.

GHIGO, E. et al. Link between impaired maturation of paghosomes and defective *Coxiella burnetii* killing in patients with chronic Q fever. **Journal of Infectious Diseases**, n. 190, p. 1767-1772, 2004.

GHIGO, E. et al. Intracellular life of *Coxiella burnetii* in macrophages. **Annals of New York Academy of Science**, v. 66, p. 55-66, 2009.

GHIGO, E.; COLOMBO, M. I.; HEINZEN, R. A. The *Coxiella burnetii* Parasitophorous Vacuole. **Advances in Experimental Biology and Medicine**, v. 984, p. 141-169, 2012.

GILK, S. D. Role of Lipids in *Coxiella burnetii* Infection. **Advances in Experimental Biology and Medicine**, v. 984, p. 198-213, 2012.

GILK, S. D.; BEARE, P. A.; HEINZEN, R. A. *Coxiella burnetii* expresses a functional Delta24 sterol reductase. **Journal of Bacteriology**, v. 192, p. 6154e9, 2010.

GINHOUX, F. et al. New insights into the multidimensional concept of macrophage ontogeny, activation and function. **Nature Immunology**, v. 17, n. 1, p. 34-40, 2016.

GLASS, C. K.; NATOLI, G. Molecular control of activation and priming in macrophages. **Nature Immunology**, v. 17, n. 1, p. 26-33, 2016.

GLAZUNOVA, O. et al. *Coxiella burnetii* Genotyping. **Emergent Infectious Diseases**, v. 11, n. 8, p. 1211-7, 2005.

GRAHAM J. G. et al. Virulent *Coxiella burnetii* pathotypes productively infect primary human alveolar macrophages. **Celular Microbiology**, v. 15, p. 1012-1025, 2013.

GORDON, S. Alternative activation of macrophages. **Nature Reviews Immunology**, v. 3, p. 23-35, 2003.

GORDON, S. et al. Macrophage heterogeneity in tissues: phenotypic diversity and functions. **Immunological Reviews**, v. 262, p. 36-55, 2014.

GOSSELIN, D. et al. Environment drives selection and function of enhancers controlling tissue-specific macrophage identities. **Cell**, v. 159, p. 1327-1340.

GUILLIAMS, M. et al. Alveolar macrophages develop from fetal monocytes that differentiate into longlived cells in the first week of life via GM-CSF. **Journal of Experimental Biology**, v. 210, p. 1977-1992, 2013.

GUO, H.; CALLAWAY, J. B.; TING, J. P. Inflammasomes: mechanism of action, role in disease, and therapeutics. **Nature Medicine**, v. 21, n. 7, p. 677-87, 2015.

GUTIERREZ, M. G. et al. Autophagy induction favours the generation and maturation of the *Coxiella*-replicative vacuoles. **Cellular Microbiology**, v. 7, n. 7, p. 981-993, 2005.

HACKSTADT, T. Biosafety concerns and *Coxiella burnetii*. **Trends in Microbiology**, v. 4, p. 341-342, 1996.

HACKSTADT, T. Steric hindrance of antibody binding to surface proteins of *Coxiella burnetii* by Phase II lipopolysaccharide. **Infection and Immunity**, v. 56, p. 802-807, 1998.

HAMASAKI, M. et al. Autophagosomes form at ER-mitochondria contact sites. **Nature**, v. 495, p. 389-393, 2013.

HARRIS, R. J. et al. Long-term persistence of *Coxiella burnetii* in the host after primary Q fever. **Epidemiology and Infection**, v. 124, p. 543-549, 2000.

HECHEMY, K. E. History and prospects of *Coxiella burnetii* research. **Advances in Experimental Biology and Medicine**, v. 984, p. 1-11, 2012.

HEINZEN, R. A. et al. Differential interaction with endocytic and exocytic pathways distinguish parasitophorous vacuoles of *Coxiella burnetii* and *Chlamydia trachomatis*. **Infection and Immunity**, v. 64, n. 3, p. 796-809, 1996.

HENRY, R. M. et al. The uniformity of phagosome maturation in macrophages. **Journal of Cellular Microbiology**, v. 164, p. 185-194, 2004.

HILL, J.; SAMUEL, J. E. *Coxiella burnetii* acid phosphatase inhibits the release of reactive oxygen intermediates in polymorphonuclear leukocytes. **Infection and Immunity**, v. 79, p. 414-420, 2011.

HONSTETTRE, A. et al. Lipopolysaccharide from *Coxiella burnetii* is involved in bacterial phagocytosis, filamentous actin reorganization, and inflammatory responses through Toll-like receptor 4. **Journal of Immunology**, v. 172, p. 3695-3703, 2004.

HORNSTRA, H. M. et al. Rapid Typing of *Coxiella burnetii*. **PLoS ONE**, v. 6, n. 11, p. e26201, 2011.

HOWE, D. et al. Maturation of the *Coxiella burnetii* parasitophorous vacuole requires bacterial protein synthesis but not replication. **Cellular Microbiology**, v. 5, n. 7, p. 469-480, 1993.

HOWE, D. et al. Nitric oxide inhibits *Coxiella burnetii* replication and parasitophorous vacuole maturation. **Infection and Immunity**, v. 70, p. 5140-5147, 2002.

HOWE, D. et al. Fusogenicity of the *Coxiella burnetii* parasitophorous vacuole. **Annals of the New York Academy of Sciences**, v. 990, p. 556-562, 2003a.

HOWE, D. et al. Maturation of the *Coxiella burnetii* parasitophorous vacuole requires bacterial protein synthesis but not replication. **Cellular Microbiology**, v. 5, p. 469e80, 2003b.

HOWE, D. et al. *Coxiella burnetii* phase I and phase II variants replicate with similar kinetics in degradative phagolysosome-like compartments of human macrophages. **Infection and Immunity**, v. 78, p. 3465-3474, 2010.

HOWE, D.; MALLAVIA, L. P. *Coxiella burnetii* infection increases transferrin receptors on J774A.1 cells. **Infection and Immunity**, v. 67, p. 3236-3241, 1999.

HOWE, D.; HEINZEN, R. A. *Coxiella burnetii* inhabits a cholesterol-rich vacuole and influences cellular cholesterol metabolism. **Cellular Microbiology**, v. 8, p. 496e507, 2006.

HUSSAIN, S. K. et al. Host kinase activity is required for *Coxiella burnetii* parasitophorous vacuole formation. **Frontiers in Microbiology**, v. 1, p. 137, 2010.

HUSSAIN, S. K.; VOTH, D. E. *Coxiella* Subversion of Intracellular Host Signaling. **Advances in Experimental Biology and Medicine**, v. 984, p. 131-140, 2012.

IHNATKO, R.; SHAW, E.; TOMAN, R. Proteome of *Coxiella burnetii*. **Advances in Experimental Biology and Medicine**, v. 984, p. 105-130, 2012.

ITAKURA, E.; KISHI-ITAKURA, C.; MIZUSHIMA, N. The hairpin-type tail-anchored SNARE syntaxin 17 targets autophagosomes for fusion with endosomes/lysosomes. **Cell**, v. 151, p. 1256e69, 2012.

KAWAI, T.; AKIRA, S. The role of pattern-recognition receptors in innate immunity: update on Toll-like receptors. **Nature Immunology**, v. 11, n. 5, p. 373-84, 2010.

KHAVKIN, T.; TABIBZADEH, S. S. Histologic, immunofluorescence, and electron microscopy study of infectious process in mouse lung after intranasal challenge with *Coxiella burnetii*. **Infection and Immunity**, v. 56, p. 1792-1799, 1988.

KLEE, S. R. et al. Highly sensitive real-time PCR for specific detection and quantification of *Coxiella burnetii*. **BMC Microbiology**, v. 6, p.2, 2006a.

KLEE, S. R. et al. Evaluation of a real-time PCR assay to detect *Coxiella burnetii*. **Annals of the New York Academy of Science**, v. 1078, p. 563-5, 2006b.

KLINGENBECK, L. et al. The *Coxiella burnetii* type IV secretion system substrate CaeB inhibits intrinsic apoptosis at the mitochondrial level. **Cellular Microbiology**, v. 15, p. 675-687, 2013.

KOHLER, L. J. et al. Effector protein Cig2 decreases host tolerance of infection by directing constitutive fusion of autophagosomes with the *Coxiella*-containing vacuole. **mBio**, v. 7, p. e01127-16, 2016.

LARSON, C. L. et al. *Coxiella burnetii* effector protein subverts clathrin-mediated vesicular trafficking for pathogen vacuole biogenesis. **Proceedings of the National Academy of Sciences of the United States of America**, v. 110, p. E4770-4779, 2013.

LARSON, C. L. et al. Right on Q: genetics begin to unravel *Coxiella burnetii* host cell interaction. **Future Microbiology**, v. 11, p. 919-39, 2016.

LATOMANSKI, E. A. et al. The effector Cig57 hijacks FCHO-mediated vesicular trafficking to facilitate intracellular replication of *Coxiella burnetii*. **PLoS Pathogens**, v. 12, p. e1006101, 2016.

LATZ, E.; XIAO, T. S.; STUTZ, A. Activation and regulation of the inflammasomes. **Nature Reviews Immunology**, v. 13, n. 6, p. 397-411, 2013.

LAVIN, Y. et al. Tissue-resident macrophage enhancer landscapes are shaped by the local microenvironment. **Cell**, v. 159, p. 1312-1326, 2014.

LAVIN, Y. et al. Regulation of macrophage development and function in peripheral tissues. **Nature Reviews Immunology**, v. 15, p. 731-44, 2015.

LEMAITRE, B. et al. The dorsoventral regulatory gene cassette *spätzle*/Toll/cactus controls the potent antifungal response in *Drosophila* adults. **Cell**, v. 86, n. 6, p. 973-83, 1996.

LOCATI, M.; MANTOVANI, A.; SICA, A. Macrophage activation and polarization as an adaptive component of innate immunity. **Advances in Immunology**, v. 120, p. 163-184, 2013.

LÜHRMANN, A. et al. Inhibition of pathogen-induced apoptosis by a *Coxiella burnetii* type IV effector protein. **Proceedings of the National Academy of Sciences of the United States of America**, v. 107, p. 18997-19001, 2010.

LÜHRMANN, A.; ROY, C. R. *Coxiella burnetii* inhibits activation of host cell apoptosis through a mechanism that involves preventing Cytochrome c release from mitochondria. **Infection and Immunity**, v. 75, p. 5282-5289, 2007.

MACKANESS, G. B. Cellular resistance to infection. **Journal of Experimental Medicine**, v. 116, p. 381-406, 1962.

MANTOVANI, A. et al. Macrophage polarization: Tumor-associated macrophages as a paradigm for polarizes M2 mononuclear phagocytes. **Trends in Immunology**, v. 23, p. 549-555, 2002.

MANTOVANI, A. et al. The chemokine system in diverse forms of macrophage activation and polarization. **Trends in Immunology**, v. 35, p. 677-86, 2004.

MANTOVANI, A. et al. Macrophage plasticity and polarization in tissue repair and remodeling. **The Journal of Pathology**, v. 229, n. 2, p. 176-185, 2013.

MANTOVANI, A.; SICA, A.; LOCATI, M. Macrophage Polarization Comes of Age. **Immunity**, v. 23, n. 4, p. 344-6, 2005.

MARIM, F. M. et al. A method for generation of bone marrow-derived macrophages from cryopreserved mouse bone marrow cells. **PLoS One**, v. 5, n. 12, p. e15263, 2010.

MARTINEZ, F. O. et al. Transcriptional profiling of the human monocyte-to-macrophage differentiation and polarization: new molecules and patterns of gene expression.. **Journal of Immunology**, v. 177, p. 7303-7311, 2006.

MARTINEZ, F. O. et al. Macrophage activation and polarization. **Frontiers in Bioscience: A Journal and Virtual Library**, v. 13, p. 453-561, 2008.

MARTINEZ, F. O.; GORDON, S. The M1 and M2 paradigm of macrophage activation: time for reassessment. **F1000Prime Reports**, p. 6-13, eCollection, 2014.

MASCARENHAS, D. P. A. et al. Inhibition of caspase-1 or gasdermin-D enable caspase-8 activation in the Naip5/NLRC4/ASC inflammasome. **PLoS Pathogens**, v. 13, n. 8, p. e1006502, 2017.

McDONOUGH, J. A. et al. Host pathways importante for *Coxiella burnetii* infection revealed by genomewide RNA interference screening. **mBio**, v. 4, p. e00606e12, 2013.

McDONOUGH, J. A.; NEWTON, H. J.; ROY, C. R. *Coxiella burnetii* Secretion Systems. **Advances in Experimental Biology and Medicine**, v. 984, p. 170-197, 2012.

McLAUGHLIN, H. P. et al. Phylogenetic inference of *Coxiella burnetii* by 16S rRNA gene sequencing. **PLoS ONE**, v. 12, n. 12, p. e0189910, 2017.

MECONI, S. et al. *Coxiella burnetii* induces reorganization of the actin cytoskeletal in human monocytes. **Infection and Immunity**, v. 66, p. 5527-5533, 1998.

MEDZHITOV, R. Toll-like receptors and innate immunity. **Nature Reviews Immunology**, v. 1, p. 135-45, 2001.

MEDZHITOV, R.; JANEWAY, C. A. Jr. Innate Immunity: the virtues of a nonclonal system of recognition. **Cell**, v. 91, n. 3, p. 295-298, 1997.

MEDZHITOV, R.; JANEWAY, C. A. Jr. The Toll receptor family and microbial recognition. **Trends in Microbiology**, v. 8, n. 10, p. 452-456, 2000.

MEDZHITOV, R.; PRESTON-HULBURT, P.; JANEWAY, C. A. Jr. A human homologue of the *Drosophila* Toll protein signals activation of adaptive immunity. **Nature**, v. 388, n. 6640, p. 394-7, 1997.

MEGHARI, S. et al. Deficient transendothelial migration of leukocytes in Q fever: the role played by interleukin-10. **Journal of Infectious Diseases**, v. 194, p. 365-369, 2006.

MEHRAJ, V. et al. Monocyte responses in the context of Q fever: from a static polarizes model to a kinetic model of activation. **Journal of Infectious Diseases**, v. 208, n. 6, p. 942-51, 2013.

MERTENS, K.; SAMUEL, J. E. Defense mechanisms against oxidative stress in *Coxiella burnetii*: adaptation to a unique intracellular niche. **Advances in Experimental Biology and Medicine**, v. 984, p. 39-63, 2012.

MILLS, C. D. et al. M-1/M-2 macrophages and the Th1/Th2 paradigm. **Journal of Immunology**, v. 164, p. 6166-73, 2000.

MINNICK, M. F.; RAGHAVAN, R. Developmental Biology of *Coxiella burnetii*. **Advances in Experimental Biology and Medicine**, v. 984, p. 231-248, 2012.

MOFFATT, J.; NEWTON, P.; NEWTON, H. J. *Coxiella burnetii*: turning hostility into a home. **Cellular Microbiology**, v. 17, n. 5, p. 621-631, 2015.

MURRAY, P. J. Macrophage polarization. **Annual Review of Physiology**, v. 79, p. 541-566, 2017.

MURRAY, P. J. et al. Macrophage activation and polarization: Nomenclature and experimental guidelines. **Immunity**, v. 41, n. 1, p. 14-20, 2014.

NARASAKI, C. T.; TOMAN, R. Lipopolysaccharide of *Coxiella burnetii*. **Advances in Experimental Biology and Medicine**, v. 984, p. 64-90, 2012.

NATHAN, C. F. et al. Identification of interferon-gamma as the lymphokine that activates human macrophage oxidative metabolism and antimicrobial activity. **Journal of Experimental Medicine**, v. 158, p. 670-89, 1983.

NATHAN, C.; SHILOH, M. U. Reactive oxygen and nitrogen intermediates in the relationship between mammalian hosts and microbial pathogens. **Proceedings in National Academy of Sciences of the United States of America**, v. 97, p. 8841-8848, 2000.

NEWTON, H. J. et al. A screen of *Coxiella burnetii* mutants reveals important roles for Dot/Icm effectors and host autophagy in vacuole biogenesis. **PLoS Pathogens**, v. 10, p. e1004286, 2014.

NEWTON, H. J.; McDONOUGH, J. A.; ROY, C. R. Effector protein translocation by the *Coxiella burnetii* Dot/Icm type IV secretion system requires endocytic maturation of the pathogen-occupied vacuole. **PLoS ONE**, v. 8, p. e54566, 2013.

O'NEILL, L. A.; GOLENBOCK, D.; BOWIE, A. G. The history of Toll-like receptors – redefining innate immunity. **Nature Reviews Immunology**, v. 13, n. 6, p. 453-60, 2013.

OMSLAND, A. et al. Host cell-free growth of the Q fever bacterium *Coxiella burnetii*. **Proceedings of the National Academy of Sciences of the United States of America**, v. 106, p. 4430-4434, 2009.

OMSLAND, A. et al. Isolation from animal tissue and genetic transformation of *Coxiella burnetii* are facilitated by an improved axenic growth medium. **Applied and Environmental Microbiology**, v. 77, p. 3720-3725, 2011.

PACE, J. L. et al. Macrophage activation: priming activity from a T cell hybridoma is attributable to interferon-gamma. **Proceedings of the National Academy of Sciences of the United States of America**, v. 80, p. 3782-6, 1983.

PECHSTEIN, J.; SCHULZE-LUEHRMANN, J.; LÜHRMANN, A. *Coxiella burnetii* as a useful tool to investigate bacterial-friendly host cell compartments. **International Journal of Medical Microbiology**, v. 308, p. 77-83, 2018.

PERDIGUERO, E. G.; GEISSMAN, F. The development and maintenance of resident macrophages. **Nature Immunology**, v. 17, p. 2-8, 2015.

QIU, J.; LUO, Z.Q. *Legionella* and *Coxiella* effectors: strength in diversity and activity. **Nature Reviews Microbiology**, v. 15, n. 10, p. 591-605, 2017.

RAOULT, D. Host factors in the severity of Q fever. **Annals of the New York Academy of Sciences**, v. 590, p. 33-38, 1990.

ROCK, F. L. et al. A Family of human receptors structurally related to *Drosophila* Toll. **Proceedings of the National Academy of Sciences of the United States of America**, v. 95, n. 2, p. 588-93, 1998.

ROMANO, P. S. et al. The autophagic pathway is actively modulated by phase II *Coxiella burnetii* to efficiently replicate in the host cell. **Cellular Microbiology**, v. 9, p. 891-909, 2007.

ROUX, V. et al. Reassessment of the taxonomic position of *Rickettsiella grylli*. **International Journal of Systematic Bacteriology**, v. 47, p. 1255-1257, 1997.

SANDOZ, K. M. et al. Developmental transitions of *Coxiella burnetii* grown in axenic media. **Journal of Microbiological Methods**, v. 96, p. 104-110, 2014.

SATOH, T.; AKIRA, S. Toll-like receptor signaling and its inducible proteins. **Microbiology Spectrum**, v. 4, n. 6, p. MCHD-0040-2016, 2016.

SCOTT, C. C.; BOTELHO, R. J.; GRINSTEIN, S. Phagosome maturation: a few bugs in the system. **Journal of Membrane Biology**, v. 193, p. 137-152, 2003.

SCOTT, G. H.; WILLIAMS, J. C.; STEPHENSON, E. H. Animal models in Q fever: pathological responses of inbred mice to phase I *Coxiella burnetii*. **Journal of General Microbiology**, n. 133, p. 691-700, 1987.

SHANNON, J. G.; HEINZEN, R. A. Infection of human monocyte-derived macrophages with *Coxiella burnetii*. **Methods in Molecular Biology**, v. 431, p. 189-200, 2008.

SHANNON, J. G.; HOWE, D.; HEINZEN, R. A. Virulent *Coxiella burnetii* does not activate human dendritic cells: role of lipopolysaccharide as a shielding molecule. **Proceedings of the National Academy of Sciences of the United States of America**, v. 102, p. 8722-8727, 2005.

SESHADRI, R. et al. Complete genome sequence of the Q-fever pathogen *Coxiella burnetii*. **Proceedings of the National Academy of Sciences of the United States of America**, v. 100, p. 5455-5460, 2003.

SOLTYSIAK, K. A.; VAN SCHAIK, E. J.; SAMUEL, J. E. Surfactant Protein D Binds to *Coxiella burnetii* and Results in a Decrease in Interactions with Murine Alveolar Macrophages. **PLoS One**, v.10, n. 9, p. e0136699, 2015.

STEIN, M. et al. Interleukin-4 potently enhances murine macrophage mannose receptor activity: a marker of alternative immunologic macrophage activation. **Journal of Experimental Medicine**, v. 176, p. 287-92, 1992.

TELEPNEV, M. V. et al. Tetraacylated lipopolysaccharide of *Yersinia pestis* can inhibit multiple Toll-like receptor mediated signaling pathways in human dendritic cells. **Journal of Infected Diseases**, v. 200, p. 1694-1702, 2009.

TING, J. P.; WILLINGHAM, S. B.; BERGSTRALH, D. T. NLRs at the intersection of cell death and immunity. **Nature Reviews Immunology**, v. 8, n. 5, p. 372-9, 2008.

TOMAN, R. et al. (Eds) *Coxiella burnetii*: Recent Advances and New Perspectives in Research of the Q Fever Bacterium. **Advances in Experimental Medicine and Biology**, v. 984, 2012.

TOMAN, R.; SKULTETY, L. Structural study on a lipopolysaccharide from *Coxiella burnetii* strain Nine Mile in avirulent phase II. **Carbohydrates Research**, v. 283, p. 175-185, 1996.

TOMAN, R.; SKULTETY, L.; IHNATKO, R. *Coxiella burnetii* glycomics and proteomics – tools for linking structure to function. **Annals of the New York Academy of Sciences**, v. 1166, p. 67-78, 2009.

TUJULIN, E. et al. Effect of endocytosis inhibitors on *Coxiella burnetii* interaction with host cells. **Acta Virologica**, v. 42, p. 125-131, 1998.

TURCO, J.; THOMPSON, H. A.; WINKLER, H. H. Interferon-gamma inhibits growth of *Coxiella burnetii* in mouse fibroblasts. **Infection and Immunity**, v. 45, p. 781-783, 1984.

- VAN SCHAİK, E. et al. Molecular pathogenesis of the obligate intracellular bacterium *Coxiella burnetii*. **Nature Reviews Microbiology**, v. 11, p. 561-573, 2013.
- VAN SCHAİK, E. J.; SAMUEL, J. E. Phylogenetic diversity, virulence and comparative genomics. **Advances in Experimental Biology and Medicine**, v. 984, p. 13-38, 2012.
- VAZQUEZ, C. L.; COLOMBO, M. I. *Coxiella burnetii* modulates Beclin-1 and Bcl-2, preventing host cell apoptosis to generate a persistente bacterial infection. **Cell Death and Differentiation**, v. 17, n. 3, p. 421-438, 2010.
- VISHWANATH, S.; HACKSTADT, T. Lipopolysaccharide phase variation determines the complement-mediated serum susceptibility of *Coxiella burnetii*. **Infection and Immunity**, v. 56, p. 40-44, 1988.
- VOTH, D. E. et al. *Coxiella burnetii* inhibits apoptosis in human THP-1 cells and monkey primary alveolar macrophages. **Infection and Immunity**, v. 75, n. 9, p. 4263-4271, 2007.
- VOTH, D. E.; HEINZEN, R. A. Lounging in a lysosome: the intracellular lifestyle of *Coxiella burnetii*. **Cellular Microbiology**, v. 9, p. 829-840, 2007.
- VOTH, D. E.; HEINZEN, R. A. Sustained activation of Akt and Erk1/2 is required for *Coxiella burnetii* antiapoptotic activity. **Infection and Immunity**, v. 77, n. 1, p. 205-213, 2009.
- VOTH, D. E.; HOWE, D.; HEINZEN, R. A. *Coxiella burnetii* inhibits apoptosis and modulates cell survival signaling in macrophages. **FASEB Microbial Pathogenesis Meeting**, 2007a.
- VOTH, D. E.; HOWE, D.; HEINZEN, R. A. *Coxiella burnetii* inhibits apoptosis in human THP-1 cells and monkey primary alveolar macrophages. **Infection and Immunity**, v. 75, p. 4263-4271, 2007b.
- WAAG, D.; THOMPSON, H. A. Pathogenesis of and immunity to *Coxiella burnetii*. In: Linder, L. E.; Lebeda, F. K.; Korch, G. W. (Eds). Biological weapons defense: infectious diseases and counter bioterrorism. **Humana Press, Totowa**, 2005.
- WALLQVIST, A. et al. Mechanisms of action of *Coxiella burnetii* effectors inferred from host-pathogen protein interactions. **PLoS One**, v. 12, n. 11, p. e0188071, 2017.
- WEBER, M. M.; FARIS, R. Subversion of the Endocytic and Secretory Pathways by Bacterial Effector Proteins. **Frontiers in Cell and Developmental Biology**, v. 6, p. 1, 2018.
- WEISBURG, W. G. et al. Phylogenetic diversity of the *Rickettsiae*. **Journal of Bacteriology**, v. 171, p. 4202-4206, 1989.
- WILLIAMS, J. C.; PEACOCK, M. G.; McCAUL, T. F. Immunological and biological characterization of *Coxiella burnetii* Phases I and II, separated from host components. **Infection and Immunity**, v. 32, p. 840-851, 1981.

WYNN, T. A.; CHAWLA, A.; POLLARD, J. W. Macrophage biology in development, homeostasis and disease. **Nature**, v. 496, n. 7446, p. 445-455, 2013.

YANG, J. et al. Prevention of apoptosis by Bcl-2: release of cytochrome c from mitochondria blocked. **Science**, v. 275, p. 1129-1132, 1997.

YOSHIIE, K. et al. Induced production of nitric oxide and sensitivity of alveolar macrophages derived from mice with different sensitivity to *Coxiella burnetii*. **Acta Virologica**, v. 43, p. 273-278, 1999.

ZAMBONI, D. S. Genetic control of natural resistance of mouse macrophages to *Coxiella burnetii* infection in vitro: macrophages from restrictive strains control parasitophorous vacuole maturation. **Infection and Immunity**, v. 72, p. 2395-2399, 2004.

ZAMBONI, D. S. et al. Mouse resident peritoneal macrophages partially control in vitro infection with *Coxiella burnetii* phase II. **Microbes and Infection**, v. 4, p. 591-598, 2002.

ZAMBONI, D. S. et al. *Coxiella burnetii* express type IV secretion system protein that function similarly to components of the *Legionella pneumophila* Dot/Icm system. **Molecular Microbiology**, v. 49, p. 965-976, 2003.

ZAMBONI, D. S. et al. Stimulation of toll-like receptor 2 by *Coxiella burnetii* is required for macrophage production of pro-inflammatory cytokines and resistance to infection. **Journal of Biological Chemistry**, v. 279, p. 54405-54415, 2004.

ZAMBONI, D. S.; RABINOVITCH, M. Nitric oxide partially controls *Coxiella burnetii* phase II infection in mouse primary macrophages. **Infection and Immunity**, v. 71, n. 3, p. 1225-33, 2003.

ZAMBONI, D. S.; RABINOVITCH, M. Phagocytosis of apoptotic cells increases the susceptibility of macrophages to infection with *Coxiella burnetii* phase II through down-modulation of nitric oxide production. **Infection and Immunity**, v. 75, p. 2075-2080, 2004.

ZAMBONI, D. S.; MORTARA, R. A.; RABINOVITCH, M. Infection of Vero cells with *Coxiella burnetii* phase II: relative intracellular bacterial load and distribution estimated by confocal laser scanning microscopy and morphometry. **Journal of Microbiological Methods**, v. 43, p. 22-232, 2001.

ZHANG, G. et al. Mechanisms of vaccine-induced protective immunity against *Coxiella burnetii* infection in BALB/c mice. **Journal of Immunology**, v. 179, p. 8372-8380, 2007.

ZHANG, G.; ZHANG, Y.; SAMUEL, J. E. Components of Protective Immunity. **Advances in Experimental Biology and Medicine**, v. 984, p. 91-104, 2012.

ZUSMAN, T. et al. The response regulator PmrA is a major regulator of the icm/dot type IV secretion system in *Legionella pneumophila* and *Coxiella burnetii*. **Molecular Microbiology**, v. 63, p. 1508-1523, 2007.

9. ANEXO I – TRABALHO DE PRIMEIRA AUTORIA

9.1. Anexo 1 - Artigo publicado (FERNANDES, Talita D et al. *Infection and Immunity*, 2016)

“Murine Alveolar Macrophages Are Highly Susceptible to Replication of *Coxiella burnetii* Phase II *In Vitro*”

- Publicado em: setembro de 2016

Murine Alveolar Macrophages Are Highly Susceptible to Replication of *Coxiella burnetii* Phase II *In Vitro*

Talita D. Fernandes,^a Larissa D. Cunha,^{a*} Juliana M. Ribeiro,^a Liliana M. Massis,^a Djalma S. Lima-Junior,^a Hayley J. Newton,^b Dario S. Zamboni^a

Department of Cell Biology, School of Medicine of Ribeirão Preto, University of São Paulo, FMRP/USP, Ribeirão Preto, São Paulo, Brazil^a; Department of Microbiology and Immunology, University of Melbourne at the Peter Doherty Institute for Infection and Immunity, Melbourne, Victoria, Australia^b

Coxiella burnetii is a Gram-negative bacterium that causes Q fever in humans. Q fever is an atypical pneumonia transmitted through inhalation of contaminated aerosols. In mammalian lungs, *C. burnetii* infects and replicates in several cell types, including alveolar macrophages (AMs). The innate immunity and signaling pathways operating during infection are still poorly understood, in part because of the lack of relevant host cell models for infection *in vitro*. In the study described here, we investigated and characterized the infection of primary murine AMs by *C. burnetii* phase II *in vitro*. Our data reveal that AMs show a pronounced M2 polarization and are highly permissive to *C. burnetii* multiplication *in vitro*. Murine AMs present an increased susceptibility to infection in comparison to primary bone marrow-derived macrophages. AMs support more than 2 logs of bacterial replication during 12 days of infection in culture, similar to highly susceptible host cells, such as Vero and THP-1 cells. As a proof of principle that AMs are useful for investigation of *C. burnetii* replication, we performed experiments with AMs from *Nos2*^{-/-} or *Ifng*^{-/-} mice. In the absence of gamma interferon and nitric oxide synthase 2 (NOS2), AMs were significantly more permissive than wild-type cells. In contrast, AMs from *Il4*^{-/-} mice were more restrictive to *C. burnetii* replication, supporting the importance of M2 polarization for the permissiveness of AMs to *C. burnetii* replication. Collectively, our data account for understanding the high susceptibility of alveolar macrophages to bacterial replication and support the use of AMs as a relevant model of *C. burnetii* growth in primary macrophages.

The intracellular bacterial pathogen *Coxiella burnetii* is the causative agent of the zoonotic infection termed Q fever. Human infection with *C. burnetii* can lead to asymptomatic seroconversion or symptomatic Q fever, which often presents as an acute febrile illness. The nonspecific symptoms associated with acute infection indicate that it is likely significantly underdiagnosed. Acute infection is often self-limited, but in a minority of cases it can progress to a serious chronic infection that predominantly manifests as life-threatening endocarditis. Importantly, recent epidemiological data demonstrated the significant long-term health impact of Q fever, reporting that more than one in three patients continued to suffer from an impaired health status 24 months postdiagnosis (1).

Humans can become infected through the inhalation of contaminated aerosols, often from close contact with infected ruminants. Within the alveolar space, *C. burnetii* can infect a variety of cell types, including alveolar macrophages (AMs) (2). Essential to the capacity of *C. burnetii* to cause disease is the ability of the pathogen to replicate inside host cells within a unique lysosome-derived vacuole. The infectious, or small-cell variant, form of the bacterium enters the host cell and is passively trafficked through the endocytic pathway before reaching the hydrolytic and acidic confines of the lysosome. These conditions trigger the metabolic activation of *C. burnetii* and stimulate the conversion to a large-cell, replicative variant (3). This environment also triggers the active bacteria to assemble the Dot/Icm type IV secretion system that facilitates the translocation of over 130 effector proteins into the host cytoplasm (4). Collectively, the action of these effectors modulates the *Coxiella*-containing vacuole (CCV) to provide an environment conducive to replication (5). This intracellular life cycle holds true for both virulent (phase I) *C. burnetii* and avirulent (phase II) *C. burnetii* bacteria that have a lipopolysaccharide

structure altered through genetic mutations that occur during serial passage in a nonimmunocompetent host (6).

It is clear that while development of the CCV is similar in many cell types, some cells, including primary peritoneal macrophages and bone marrow-derived macrophages (BMDMs), are intrinsically more restrictive to the intracellular replication of *C. burnetii* (7, 8). Furthermore, BMDMs from different inbred mouse strains vary in their ability to restrict *C. burnetii* infection, with A/J and BALB/c mice being more susceptible to *C. burnetii* phase II than many other mouse strains, including the C57BL/6 strain (9). The mechanisms through which different genetic backgrounds and cell types can control *C. burnetii* intracellular growth remain an area of interesting scientific pursuit.

AMs have long been considered the primary site of *C. burnetii* infection (2), and AMs from monkeys and humans have recently been used to explore the host-pathogen interactions that take place during *C. burnetii* infection. Cynomolgus monkey AMs were used to demonstrate the potent antiapoptotic activity asso-

Received 13 May 2016 Returned for modification 1 June 2016

Accepted 7 June 2016

Accepted manuscript posted online 13 June 2016

Citation: Fernandes TD, Cunha LD, Ribeiro JM, Massis LM, Lima-Junior DS, Newton HJ, Zamboni DS. 2016. Murine alveolar macrophages are highly susceptible to replication of *Coxiella burnetii* phase II *in vitro*. Infect Immun 84:2439–2448. doi:10.1128/IAI.00411-16.

Editor: C. R. Roy, Yale University School of Medicine

Address correspondence to Dario S. Zamboni, dszamboni@fmrp.usp.br.

* Present address: Larissa D. Cunha, Department of Immunology, St. Jude Children's Research Hospital, Memphis, Tennessee, USA.

Copyright © 2016, American Society for Microbiology. All Rights Reserved.

ciated with *C. burnetii* infection, although the replication dynamics of *C. burnetii* in this cellular model were not explored (10). More recently, human AMs, extracted from postmortem lung tissue samples, were shown to support the replication of different pathotypes of *C. burnetii* (11). In addition, this study demonstrated that both virulent (phase I) and avirulent (phase II) *C. burnetii* bacteria are able to infect human AMs. Interestingly, it was also observed that, within human AMs, *C. burnetii* more frequently forms multiple smaller CCVs rather than the large fusogenic vacuole observed in other cellular models of infection (11). Within a murine model of infection, it has been demonstrated that AMs are susceptible to *C. burnetii* infection (12), and an early study demonstrated that nitric oxide (NO) is produced by murine AMs in response to infection (13).

In the study described here, we investigated and characterized the infection of primary murine AMs by *C. burnetii* phase II *in vitro*. Our data reveal that AMs are permissive to *C. burnetii* multiplication. The AMs showed a pronounced M2 polarization and showed an increased susceptibility to infection in comparison to murine BMDMs. Finally, we investigated the contribution of signaling molecules to the susceptibility of AMs to *C. burnetii*, demonstrating the utility of AMs for investigation of the host restriction of *C. burnetii* infection in relevant host cells.

MATERIALS AND METHODS

Preparation of *C. burnetii* for infection. *Coxiella burnetii* phase II Nine Mile strain (RSA439) bacteria were prepared from infected Vero cell monolayers as previously described (14). Confluent cultures of Vero cells were irradiated with 10 Gy ^{60}Co to block cell division and maintained at 37°C in 5% CO_2 in Dulbecco modified Eagle medium (DMEM; Gibco) with 10% fetal bovine serum (FBS; Gibco) and 1 mM L-glutamine (Sigma-Aldrich). Infection was carried out, and the cells were maintained under the above-described conditions for an additional 6 days. For the preparation of the *Coxiella* inoculum, infected Vero cells were lysed by homogenization in sterile distilled water. Cell lysates were clarified by centrifugation at $1,810 \times g$ for 10 min at 4°C. The supernatant was extracted and centrifuged at $12,857 \times g$ for 30 min at 4°C. The supernatant was discarded, and the pellet was resuspended in RPMI 1640 medium with 10% FBS. Bacterial aliquots were kept at -80°C . Prior to infection, aliquots were thawed at 37°C and subjected to sonication at 35 KHz for 3 min.

Mouse strains. A/J, C57BL/6, *Nos2*^{-/-}, *Ifng*^{-/-}, and *Il4*^{-/-} mice (all in the C57BL/6 mouse genetic background) were bred in the Animal Facilities of the Medical School of Ribeirão Preto (FMRP/USP). All animals were provided food and water *ad libitum*, and the temperature was controlled at 25°C. Mice were cared for according to the institutional guidelines on ethics in animal experiments (approved by the Comissão de Ética em Experimentação Animal da Faculdade de Medicina de Ribeirão Preto/CETEA, protocol number 097/2010).

Culture of mammalian cell lines. THP-1 cells were cultured in RPMI 1640 medium (Sigma-Aldrich) with 10% FBS, and Vero cells were maintained in DMEM (Gibco) with 10% FBS (Gibco) and 1 mM L-glutamine (Sigma-Aldrich) at 37°C in 5% CO_2 . For assays with THP-1 cells, differentiation into macrophage-like cells was mediated by the addition of 200 nM phorbol 12-myristate 13-acetate (PMA) to the culture overnight as described previously (10), and the medium was replaced with PMA-free medium prior to the infection. For infection of Vero cells, the cells were irradiated with 10 Gy ^{60}Co and maintained overnight at 37°C in 5% CO_2 prior to the infection.

Bone marrow-derived macrophages. BMDMs were obtained and differentiated according to the protocol described previously (15). Briefly, bone marrow cells were harvested from femurs and differentiated with RPMI 1640 medium (Sigma-Aldrich) containing 20% FBS (Invitrogen) and 30% L-929 cell conditioned medium (LCCM) at 37°C in 5% CO_2

(15). Macrophages were harvested, counted, and seeded in 24-well plates at a density of 10^5 cells/well 1 day prior to infection and kept in RPMI 1640 medium with 10% FBS and 5% LCCM.

Alveolar macrophages. AMs were obtained from adult mice (age, 6 to 8 weeks) via bronchoalveolar lavage (BAL). The animals were euthanized, and the trachea was pierced with a sterile cannula (BD AngioCath). The lungs were washed four consecutive times with 1.0 ml of warm sterile phosphate-buffered saline (PBS) containing 5 mM EDTA, and the material was collected. Cells were transferred to a new sterile tube, washed twice in sterile PBS, and resuspended in RPMI 1640 medium with 10% FBS, 1 mM L-glutamine, and 100 U/ml penicillin-streptomycin. Cells were seeded in 24-well plates at a density of 10^5 cells/well. Macrophages were maintained in culture in RPMI 1640 medium with 10% FBS, 1 mM L-glutamine, and 100 U/ml penicillin-streptomycin at 37°C in 5% CO_2 . Culture medium was replaced two times before infection in antibiotic-free medium. Fresh media were added to the macrophage cultures at day 6 postinfection.

***C. burnetii* infection of cells.** Frozen stocks of *C. burnetii* were thawed at 37°C and treated in an ultrasound bath for 3 min to dissolve bacterial clumps. Tissue culture medium was aspirated from the cells and replaced with fresh medium before infection. Bacteria were homogenized with a pipette and applied directly to the cells at a multiplicity of infection (MOI) of 3 (except for the experiment whose results are presented in Fig. 1A, in which an MOI of 100 was used). The plates were centrifuged at $200 \times g$ for 5 min at 25°C and placed at 37°C in 5% CO_2 . After 4 h, the culture medium was aspirated, the wells were washed with warm sterile PBS, and fresh culture medium was added. As an internal experimental control of *C. burnetii* replication, infected cells were treated with 20 $\mu\text{g}/\text{ml}$ rifampin (RIF) to inhibit *C. burnetii* protein translation (16).

Fluorescence microscopy. AMs from C57BL/6 mice were seeded at a density of 10^5 cells per well in a 24-well plate containing 13-mm glass coverslips. Cells were infected with wild-type *C. burnetii* bacteria or *C. burnetii* bacteria expressing mCherry at an MOI of 3 as previously described (17). After 4 h, 3 days, 6 days, 9 days, or 12 days of infection, cells were fixed with 3.5% paraformaldehyde containing 7.5% sucrose for 20 min and then washed twice with room temperature PBS. For the assays with wild-type *C. burnetii* bacteria, the cells were then permeabilized for 2 min with ice-cold 100% methanol, followed by two more washes with PBS. Cellular autofluorescence was removed by incubation with 0.1 M glycine in PBS for 10 min, and then the samples were blocked for 30 min in PBS containing 1% bovine serum albumin (BSA; Sigma-Aldrich) and 5% goat serum at room temperature. The cells were incubated with rabbit anti-*C. burnetii* (1:100) and rat anti-LAMP-1 (1:2) primary antibodies diluted in 1% BSA in PBS for 1 h, washed twice with room temperature PBS, and then incubated for 30 min in 1% BSA in PBS containing Alexa Fluor 488- and Alexa Fluor 594-conjugated secondary antibodies (Invitrogen) at a concentration of 1:2,000. After three washes with room temperature PBS, cells were mounted with Prolong Gold antifade reagent containing DAPI (4',6'-diamidino-2-phenylindole dilactate; Invitrogen). For the assays using mCherry-expressing *C. burnetii*, the cells were washed twice with room temperature PBS after the fixation and then directly mounted with Prolong Gold antifade reagent containing DAPI.

Multiphoton microscopy images were acquired using an LSM 780 Zeiss AxioObserver microscope equipped with a 63 $\times$ oil immersion objective and analyzed using ImageJ software.

Transmission electron microscopy. AMs were infected for 3 or 6 days, washed once with PBS, fixed for 1 h at room temperature with modified Karnovsky's fixative (2% formaldehyde, 2.5% glutaraldehyde in 0.1 M sodium cacodylate, pH 7.4), dislodged by scraping, and then processed for and analyzed by transmission electron microscopy as previously described (7).

Quantification of *C. burnetii*. BMDMs, AMs, THP-1 cells, and irradiated Vero cells infected with *C. burnetii* were maintained in culture for up to 12 days. After 6 days of infection, cells were supplemented with an additional 500 μl of RPMI 1640 medium with 10% FBS and 5% LCCM for

BMDMs, RPMI 1640 medium with 10% FBS for AMs and THP-1 cells, and DMEM (Gibco) with 10% FBS (Gibco) and 1 mM L-glutamine (Sigma-Aldrich) for irradiated Vero cells. At different times postinfection, cells were lysed with 1.0 ml of deionized water and cell lysates were combined with the culture supernatant from the respective wells. The material was centrifuged at $14,000 \times g$ for 15 min at 4°C for deposition of the bacteria present in the samples. The DNA from *C. burnetii* was extracted using a Bacteria GenomicPrep Illustra kit (GE Healthcare), and the presence of genetic material was confirmed by spectrophotometry (determination of the absorbance at 260 nm). The primer pair dotA-for and dotA-rev (18) was used for quantitative PCR (qPCR), a reliable method for quantification of bacterial replication, as demonstrated by Brennan and Samuel (16) for analyses involving *C. burnetii*. The reaction was performed using a Platinum SYBR green qPCR Supermix-UDG with carboxy-X-rhodamine (ROX) kit (Invitrogen), and thermocycling reactions were performed according to the manufacturer's instructions. The samples were read on an ABI Prism 7000 thermocycler.

RT-PCR analysis. mRNA from cultures of 10^6 AMs and BMDMs derived from C57BL/6 mice noninfected or infected for 12 h with *C. burnetii* at an MOI of 3 were extracted with the TRIzol reagent. RNA purification was done using an Illustra RNAspin minikit (GE Healthcare), according to the manufacturer's instructions, and genetic material was quantified using a NanoDrop spectrophotometer (Thermo Scientific). ImProm-II reverse transcriptase (Promega) was used to obtain cDNA. The real-time quantitative PCR assays were performed using the Platinum SYBR Green qPCR Supermix-UDG with carboxy-X-rhodamine kit (Invitrogen). Thermocycling reactions were performed according to the manufacturer's instructions, and the samples were read on a StepOnePlus real-time PCR system (Applied Biosystems). The expression of the target genes was calculated using the mean threshold cycle (C_T) values of each sample, which were normalized to those for the housekeeping genes. The sequences of the primers used for reverse transcription-PCR (RT-PCR) are as follows: *Gapdh*-fwd, 5'-CATCACCATCTTCCAGGAGCG-3'; *Gapdh*-rev, 5'-ACGGACACATTGGGGGTAGG-3'; *Nos2*-fwd, 5'-CGAAACGC TTCTCTTCCAA-3'; *Nos2*-rev, 5'-TGAGCCTATATTGCTGTGGCT-3'; *Irfn*-fwd, 5'-GCATCTTGGCTTTGCAGCT-3'; *Irfn*-rev, 5'-CCTTTTTCG CCTTGCTGTG-3'; *Tnf*-fwd, 5'-TGTGCTCAGAGCTTCAACAA-3'; *Tnf*-rev, 5'-CTTGATGGTGCATGAGA-3'; *Thr2*-fwd, 5'-AAGTCTCC GGAATTATCAGTCC-3'; *Thr2*-rev, 5'-TGATGGATGTCGCGGAT-3'; *Il4*-fwd, 5'-AAGAGCATCATGCAATGGA-3'; *Il4*-rev, 5'-TTAAAGCA TGGTGGCTCAGTAC-3'; *Arginase1*-fwd, 5'-GTTCCAGATGTACCA GGATTC-3'; *Arginase1*-rev, 5'-CGATGCTTTTGGCAGATATGC-3'; *Fizz1*-fwd, 5'-TGTGGCTTTGCTGTGGAT-3'; *Fizz1*-rev, 5'-TCTTAG GACAGTTGGCAGCAG-3'; *Ym1*-fwd, 5'-GGGTTGGTTATGACAA TGTCAG-3'; and *Ym1*-rev, 5'-TGAAGTCATCCATGTCCAGG-3'.

Flow cytometry for phenotypic analysis. Single suspensions of AMs and BMDMs from C57BL/6 mice were stained with stains from the LIVE/DEAD fixable blue dead cell stain kit (Invitrogen) in PBS to exclude dead cells. Cells were stained with fluorochrome-conjugated antibodies against surface markers CD11b (clone M1/70), F4/80 (clone BM8), CD206 (clone C068C2), and CD86 (clone GL-1), purchased from BD Biosciences (San Diego, CA), and dextrin-1/CLEC7A (CD369), obtained from BioLegend, in PBS containing 1% FBS for 30 min at 4°C and then washed. Staining was performed in the presence of 5 μ g/ml of Fc Block (BD Biosciences). Cell acquisition was performed on a BD FACSCanto II flow cytometer using FACSDiva software (BD Biosciences). For each sample, at least 100,000 events were collected. Data were analyzed with FlowJo software (TreeStar).

Statistical analyses. Statistical analyses were performed using GraphPad Prism (version 5) software. The results were analyzed by a statistical test for analysis of variance (ANOVA) followed by the Bonferroni posttest or Student's *t* test. Differences were considered statistically significant when the *P* value was <0.05 .

RESULTS

Alveolar macrophages from C57BL/6 mice are permissive to replication of *C. burnetii*. We have previously shown that BMDMs from A/J and BALB/c are less restrictive to the replication of *C. burnetii* phase II than cells from mice with the C57BL/6 mouse genetic background. This was determined by assessing the formation of the large CCV and by the determination of the number of *C. burnetii* focus-forming units (9). To further confirm these data, we quantified the bacterial genomic DNA by quantitative PCR (qPCR) as a tool to assess the replication of *C. burnetii* in BMDMs as previously described (16). In agreement with previously published data (9), we found that BMDMs from C57BL/6 mice are highly restrictive to *C. burnetii* phase II, whereas BMDMs from A/J mice support some intracellular replication of *C. burnetii* phase II from days 3 to 12 postinfection. This was observed in experiments using high and low infection doses (MOIs, 100 and 3, respectively) (Fig. 1A and B, respectively). We continued the macrophage infection studies using an MOI of 3. Because most of the gene-deficient mice generated thus far were constructed in the C57BL/6 mouse background, we aimed to identify a primary cell that supported the replication of *C. burnetii* phase II in culture. It is known that *C. burnetii* infects macrophages and monocytes, with AMs considered the primary target cell of infection (10, 12, 19–22). Thus, we evaluated whether murine AMs from the C57BL/6 mouse strain support the replication of *C. burnetii* phase II in culture. Strikingly, when we assessed bacterial replication using qPCR, we found that, as opposed to BMDMs, the AMs were highly permissive to *C. burnetii* phase II replication (Fig. 1C). Of note, the internalization of bacteria by these different cells was similar, as measured by determination of the number of *C. burnetii* genomes associated with the cells after 4 h of infection (Fig. 1A to D, insets). As a negative control for bacterial replication, rifampin treatment was included to kill the *C. burnetii* bacteria in the macrophages. Further, when we compared the *C. burnetii* replication in AMs from C57BL/6 and A/J mice, we found that AMs from both strains were equally permissive for *C. burnetii* replication (Fig. 1D). These data suggest that the genetic mechanisms controlling the natural resistance of BMDMs to *C. burnetii* may be less effective or absent in AMs. Importantly, AMs emerge as a very interesting model of the primary macrophage that supports the intracellular replication of *C. burnetii* phase II in culture. As such, we decided to further characterize *C. burnetii* replication in AMs.

Alveolar macrophages accumulate large amounts of intracellular bacteria within the Coxiella-containing vacuole. Given the robust replication of *C. burnetii* phase II in murine AMs (Fig. 1C and D), we aimed to assess bacterial replication in these macrophages by microscopy. AMs obtained from C57BL/6 mice were infected, and intracellular bacteria were visualized by immunofluorescence and transmission electron microscopy. Figure 2 illustrates the susceptibility of AMs over the first 6 days of *C. burnetii* phase II infection in culture. We detected an increased concentration of intracellular *C. burnetii*, stained in red, during the course of infection, suggesting that the bacteria can readily replicate in AMs. Further, the association of the bacteria with the lysosomal membrane marker LAMP-1 confirms that the bacteria replicate in the well-characterized lysosome-derived parasitophorous vacuole. During the procedures to stain the bacteria and LAMP-1, the AMs containing large spacious vacuoles were lost because these cells easily detach from the coverslips during the antibody washings.

FIG 1 Alveolar macrophages are permissive to *C. burnetii* replication. BMDMs from C57BL/6 and A/J mice (A and B), BMDMs and AMs from C57BL/6 mice (C), or AMs from C57BL/6 and A/J mice (D) were infected with *C. burnetii* phase II (MOIs, 100 [A] and 3 [B to D]), and bacterial multiplication was evaluated by quantification of the genomic DNA by qPCR after 0 (4 h after infection), 3, 6, 9, 12 or 15 days (d) of infection. The y axis represents the difference in the base 10 logarithm between the genomic quantity X at time t [$\log X(t)$] and the quantity X at time zero (T_0) [$\log X(0)$]. Cells treated with RIF were used as a negative control of bacterial replication. (Insets) Amount of *C. burnetii* genomic DNA present in cells after 4 h of infection. The data shown are the means $\pm$ standard errors of one representative experiment out of three performed. *, $P < 0.05$ (two-way ANOVA).

FIG 2 *C. burnetii* infection in alveolar macrophages assessed by fluorescence microscopy. AMs from C57BL/6 mice were infected with wild-type *C. burnetii* phase II (MOI, 3) for 4 h, 3 days, 6 days, 9 days, and 12 days (A) or infected with mCherry-expressing *C. burnetii* phase II for 4 h, 3 days, 6 days, 9 days, and 12 days (B). After the indicated times, cells were fixed, processed, and then examined using multiphoton fluorescence microscopy. (C) AM infected for 9 days with mCherry-expressing *C. burnetii*. DIC, differential interference contrast. Green, lysosomal membrane-associated protein LAMP-1; red, *C. burnetii*; cyan, DAPI-stained DNA. Bars, 5 μ m.

Thus, to assess the presence of typical large vacuoles in infected AMs, we used a strain of *C. burnetii* expressing mCherry (23). By examining the infected AMs using two-photon laser scanning fluorescence microscopy, we found a significant increase in bacterial replication in AMs from days 0 (4 h of infection) to 12 after infection (Fig. 2A and B). At day 6 of infection, we observed several cells containing many *C. burnetii* bacteria in large vacuoles (Fig. 2B). At day 9 of infection, the AMs were filled with *C. burnetii* bacteria, as illustrated by a merge image showing both differential interference contrast and fluorescence images (Fig. 2C). Intracellular bacteria were also observed by transmission electron microscopy (Fig. 3). At 3 days after infection, AMs showed large CCVs containing high numbers of large-cell variants of *C. burnetii* (Fig. 3A). At day 6 after infection, we detected vacuoles harboring a large amount of *C. burnetii* bacteria (Fig. 3B). Taken together, the results confirm that AMs support *C. burnetii* replication in culture and can be used as a useful model of the primary macrophage to study *C. burnetii* phase II replication *in vitro*.

Alveolar macrophages show a more pronounced polarization into M2 cells than bone marrow-derived macrophages. To

FIG 3 *C. burnetii* infection in alveolar macrophages assessed by transmission electron microscopy. AMs were infected with *C. burnetii* phase II (MOI, 3), fixed, and processed for transmission electron microscopy. Cells were fixed after 3 days (top) and 6 days (bottom) of infection. The panels on the right are enlarged images of the boxed regions in the panels on the left. Arrows, large-cell variant forms of *C. burnetii*. Bars, 1 μm.

assess the mechanisms underlying the permissiveness of AMs to *C. burnetii*, we evaluated M1/M2 polarization, which has previously been shown to be important for *C. burnetii* replication and pathogenesis (24, 25). Initially, we evaluated the expression of genes known to be hallmarks of M1 and M2 polarization in AMs and BMDMs. By measuring gene expression in noninfected BMDMs and AMs, we found that the latter show a significantly reduced expression of *Nos2*, *Tnfa*, *Ifng*, and *Tlr2* (genes associated with an M1 polarization phenotype) and an increased expression of *Arginase1*, *Fizz1*, and *Ym1*, which are associated with an M2 profile (Fig. 4). The AMs also showed a pronounced M2 polarization compared to that of BMDMs in response to *C. burnetii* infection. We detected reduced expression of *Nos2*, *Tnfa*, *Ifng*, and *Tlr2* and increased expression of *Il4*, *Arginase1*, *Fizz1*, and *Ym1* in AMs infected with *C. burnetii* compared to their expression in infected BMDMs (Fig. 4). These data are consistent with AMs being more susceptible because they polarize toward an M2 phenotype. Accordingly, M1 macrophages effectively trigger the production of tumor necrosis factor alpha (TNF-α), a cytokine known to facilitate the restriction of *C. burnetii* replication in macrophages (26). To further test this hypothesis, we evaluated the expression of surface markers known to be hallmarks of M2 macrophages, including mannose receptor (CD206) and dectin-1 (27). To evaluate protein expression, we performed fluorescence-activated cell sorting experiments in which we gated live macrophages and evaluated the expression of the cell surface markers CD206, dectin-1, and CD86 on cells double positive for CD11b and F4/80 (Fig. 5A

and B). Using specific antibodies, we found that AMs showed increased expression of M2-associated molecules, such as CD206 (mannose receptor) and dectin-1 (CLEC7A), compared to BMDMs. This was evident under the uninfected condition (Fig. 5C) and in response to infection (Fig. 5D). Moreover, we measured the expression of the costimulatory molecule CD86, which was similarly expressed in AMs and BMDMs (Fig. 5C and D). These data are consistent with the hypothesis that AMs are more susceptible because of an M2 polarized phenotype, a characteristic that has previously been associated with permissiveness to *C. burnetii* replication (reviewed in reference 28).

Alveolar macrophages deficient in NOS2 and IFN-γ are more susceptible to *C. burnetii* replication, whereas cells deficient in interleukin-4 (IL-4) are more restrictive to *C. burnetii* replication. It is well established that nitric oxide (NO) is involved in the restriction of *C. burnetii* replication (8, 9, 29, 30). Thus, we assessed bacterial replication in AMs deficient in nitric oxide synthase 2 (NOS2; or inducible nitric oxide synthase [iNOS]) as a proof of principle that AMs can be used as a model to investigate the interactions of *C. burnetii* with primary mouse macrophages in the C57BL/6 mouse genetic background. In agreement with previous reports, we found that AMs from *Nos2*^{-/-} mice are more permissive to *C. burnetii* phase II replication than AMs from wild-type C57BL/6 mice (Fig. 6A). We also evaluated bacterial replication in AMs from *Ifng*^{-/-} mice because gamma interferon (IFN-γ) is known to be essential for NO production (31). Moreover, IFN-γ is well described to be involved in the immune re-

FIG 4 Alveolar macrophages exhibit higher levels of expression of genes related to the M2 phenotype. BMDMs and AMs from C57BL/6 mice were left uninfected or infected for 12 h with *C. burnetii* at an MOI of 3. Total RNA was extracted, and reverse transcription-PCR (RT-PCR) was performed for detection of the genes *Nos2*, *Tnfa*, *Ifng*, *Tlr2*, *Il4*, *Arginase1*, *Fizz1*, and *Ym1*. The data shown are the means $\pm$ standard errors from one representative experiment out of two performed. *, $P < 0.05$ comparing BMDMs and AMs; #, $P < 0.05$ comparing infected cells with the uninfected cells from the respective group (two-way ANOVA).

sponse against *C. burnetii* (19, 32–40). We found that AMs from *Ifng*^{-/-} mice are significantly more permissive to *C. burnetii* phase II replication than AMs from wild-type C57BL/6 mice (Fig. 6B). Collectively, these data support the use of AMs as a model of primary macrophages for investigations involving *C. burnetii* infections in primary macrophages from the C57BL/6 mouse, the strain mostly commonly used for the generation of transgenic mice. To further understand the effect of the M2 polarization of AMs that results in the increased susceptibility to *C. burnetii*, we performed experiments using AMs from mice deficient in IL-4, which is required for M2 differentiation (reviewed in reference 41). We found that AMs from *Il4*^{-/-} mice are less permissive to *C. burnetii* replication than cells from C57BL/6 mice (Fig. 6C). The internalization of bacteria did not differ in AMs from C57BL/6, *Nos2*^{-/-}, *Ifng*^{-/-}, and *Il4*^{-/-} mice, as measured by the number of *C. burnetii* bacteria in cultures infected for 4 h (Fig. 6A to C, insets). Collectively, these data support the hypothesis that the M1/M2 polarization underlies the increased susceptibility of AMs in comparison to BMDMs. Importantly, regardless of the mechanisms associated with the resistance of the BMDMs, these data support the use of AMs as an important and relevant host cell for evaluation of *C. burnetii* replication in primary macrophages.

The permissiveness of mouse alveolar macrophages is comparable to that of highly susceptible immortalized cell lines. Our

results demonstrated that murine AMs are permissive for *C. burnetii* phase II replication *in vitro*. Thus, we decided to compare *C. burnetii* phase II replication in C57BL/6 mouse AMs with that in highly susceptible host cells, such as Vero cells (a green monkey kidney cell line) and cells of a monocytic cell line (THP-1) differentiated into macrophage-like cells. Both of these cell types are known to be very susceptible to *C. burnetii* replication (6, 14, 34). Figure 7 shows that at 6 days postinfection, murine AMs are as susceptible to *C. burnetii* phase II infection as Vero and THP-1 cells. At later time points after infection, the Vero and THP-1 cell lines supported bacterial replication more robust than that in primary AMs. Nonetheless, the comparable level of replication during the first 6 days of infection reinforces the demonstration that murine AMs are indeed highly permissive for *C. burnetii* replication and can be used as a relevant host cell model for investigations involving the macrophage interaction with *C. burnetii* phase II in culture.

DISCUSSION

C. burnetii is a unique intracellular bacterial pathogen capable of replicating to high numbers within a lysosome-derived vacuole. Humans can contract Q fever through the inhalation of contaminated aerosols and the subsequent intracellular replication of *C. burnetii* within AMs. As such, studying the interaction between

FIG 5 Alveolar macrophages exhibit an M2 phenotype. (A and B) Gating strategy and frequency of noninfected BMDMs (A) and AMs (B). FSC, forward scatter; SSC, side scatter. (C and D) Surface phenotyping of BMDMs and AMs infected (for 24 h) (C) or not infected (D) with *C. burnetii* for CD86 and the M2 markers CD206 and dectin-1, as analyzed by flow cytometry. Cells were gated on live CD11b⁺ F4/80⁺ cells.

this specific macrophage lineage and the pathogen provides an important, disease-relevant, *in vitro* model to understand the specific host-pathogen interaction that takes place during *C. burnetii* infection. Previous research has demonstrated that BMDMs from different mouse strains vary in their permissiveness to *C. burnetii* replication (42). BMDMs from A/J and BALB/c mice support the intracellular replication of *C. burnetii*, while BMDMs from

C57BL/6 mice restrict bacterial replication. Here, we recapitulate these results and demonstrate that AMs are significantly more permissive to *C. burnetii* replication than BMDMs. This finding is most striking with C57BL/6 mice, in which BMDMs restrict the intracellular growth of *C. burnetii*, yet the bacteria readily replicate in AMs. Indeed, despite a remarkable difference in the replication of *C. burnetii* in BMDMs of C57BL/6 and A/J mice, there is no

FIG 6 AMs support the assessment of innate immunity against *C. burnetii* phase II using knockout mice in the C57BL/6 mouse genetic background. Primary AMs from wild-type C57BL/6 mice and *Nos2*^{-/-} (A), *Ifng*^{-/-} (B), and *Il4*^{-/-} (C) mice were infected with *C. burnetii* phase II (MOI, 3), and bacterial multiplication was evaluated by quantification of the genomic DNA by qPCR after 0 (4 h after infection), 3, 6, 9, and 12 days of infection. The y axis represents the difference in the base 10 logarithm between the genomic quantity X at time t [logX(t)] and the quantity X at time zero (T₀) [logX(0)]. Cells treated with RIF were used as a negative control of bacterial replication. (Insets) Amount of *C. burnetii* genomic DNA present in cells after 4 h of infection. The data shown are the means ± standard errors from one representative experiment out of three independent experiments performed. *, *P* < 0.05 (two-way ANOVA).

significant difference in the ability of *C. burnetii* to replicate in AMs from these two inbred mouse strains. This suggests that the genetic basis for C57BL/6 mouse BMDM restriction of *C. burnetii* infection reflects phenotypic features that may be specific for this cell type or that operate at a lower efficiency in AMs.

AMs are critical lung resident cells that, in healthy lungs, constitute the vast majority of the cellular content within alveoli (43). As such these cells are intricately essential for lung homeostasis and immune surveillance (44). Recent evidence indicates that AMs originate from fetal monocytes that are long living and are maintained by self-renewal rather than circulating bone marrow-derived monocytes (45). This tissue microenvironment has a significant epigenetic and transcriptomic impact on resident macro-

FIG 7 Alveolar macrophages are permissive to *C. burnetii* phase II replication, similar to Vero and THP-1 cells, *in vitro*. C57BL/6 AMs, irradiated Vero cells, and THP-1 cells (previously differentiated overnight with 200 nM PMA) were infected with *C. burnetii* phase II (MOI, 3), and bacterial multiplication was evaluated by quantification of the genomic DNA by qPCR after 0 (4 h after infection), 3, 6, 9, and 12 days of infection. The y axis represents the difference in the base 10 logarithm between the genomic quantity X at time t [logX(t)] and the quantity X at time zero (T₀) [logX(0)]. Cells treated with RIF were used as a negative control of bacterial replication. (Insets) Amount of *C. burnetii* genomic DNA present in cells after 4 h of infection. The data shown are the means ± standard deviations from one representative experiment out of three independent experiments performed. *, *P* < 0.05 (two-way ANOVA).

phages, rendering AMs phenotypically distinct from BMDMs (46–48). In light of our results, one can suggest that these phenotypic differences between BMDMs and AMs from C57BL/6 mice mediate a significant difference in the capacity to restrict the intracellular replication of *C. burnetii*. Interestingly, our data comparing M1/M2 polarization in BMDMs and AMs suggest that the M2 profile of AMs may underlie their increased susceptibility to infection. This hypothesis is supported by experiments using cells from *Il4*^{-/-} mice. In the absence of IL-4, the AMs became significantly more resistant to infection, a feature that is consistent with the essential role of IL-4 in the induction of M2 polarization (41). It is likely that during coevolution with mammalian hosts, *C. burnetii* has evolved strategies to target and subvert specific responses in alveolar macrophages, a process that may be facilitated by the anti-inflammatory M2 characteristics of this cell type.

In support of our data showing the increased replication of *C. burnetii* in AMs, it was previously reported that tissue-resident AMs are susceptible to *C. burnetii* replication *in vivo* (12). Moreover, human AMs were shown to support *C. burnetii* replication in culture (11). In addition, a system that allows infection of *ex vivo* human lung tissue was recently reported. It was found that within human lung tissue, *C. burnetii* preferentially replicated in human AMs (49). In this context, our data provide a definitive demonstration that mouse AMs can be used as a relevant host cell for investigations of the interactions of *C. burnetii* with host cells *in vitro*. C57BL/6 mouse AMs support the intracellular replication of *C. burnetii*, and the bacteria replicate in a manner that is comparable to that in widely used immortalized cell lines that are highly susceptible to *C. burnetii* replication. Over the first 6 days after infection, we found that *C. burnetii* phase II replicates similarly in AMs, THP-1 cells, and Vero cells. At days 9 and 12, we detected higher bacterial loads in Vero and THP-1 cells, which are possibly caused by the saturation of the host cells available for bacterial replication. This hypothesis is supported by the experiments performed with AMs infected with mCherry-expressing *C. burnetii*. After 9 days of infection, we visualized AMs filled with bacteria, which were contained in large vacuoles that occupy the entire

macrophage cytoplasm (Fig. 2B and C). We did not detect a reduction of bacterial loads from days 9 to 12, a feature that does not support the restriction of bacterial replication in the AMs at later stages of infection.

C57BL/6 mouse AMs can serve as an important host cell for the investigation of host factors that aid or restrict the intracellular replication of *C. burnetii*. Here we have confirmed the importance of both nitric oxide and IFN- γ in restricting the ability of *C. burnetii* to replicate. Significantly higher levels of *C. burnetii* were recovered from both *Nos2*^{-/-} and *Ifng*^{-/-} mouse AMs than from wild-type C57BL/6 mouse AMs. Production of nitric oxide and its reactive nitrogen intermediates is an important cellular response that acts against respiratory pathogens (50). At high levels, nitric oxide has a cytotoxic effect and can participate in the control of intracellular pathogen replication. Indeed, several studies have demonstrated that nitric oxide has the capacity to restrict the intracellular replication of *C. burnetii* (8, 9, 29, 30, 51). Our data support these findings, demonstrating that AMs from C57BL/6 mice lacking *Nos2* support higher levels of *C. burnetii* replication than wild-type cells. Similarly, the absence of IFN- γ , which is critical for nitric oxide production, is known to be important to host restriction of *C. burnetii* replication. Importantly, by using AMs from *Nos2*^{-/-} and *Ifng*^{-/-} mice, we validated the use of AMs as a relevant host cell for the *in vitro* infection setting. Our characterization of this model of infection demonstrates that studying the interaction between *C. burnetii* and AMs will allow intricate investigation of both host and bacterial factors that are important for pathogenesis and host resistance during Q fever.

ACKNOWLEDGMENTS

We are grateful to Maira C. Nakamura and Victoria Maria dos Santos for technical assistance. We also thank Maria D. S. Ferreira, José A. Maulin, and Roberta R. Rosales for technical assistance in the institutional facilities for electron microscopy (M. D. S. Ferreira and J. A. Maulin) and multiphoton microscopy (R. R. Rosales).

We declare no conflict of interest in relation to this work.

This work was supported by grants from the Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP; grants 2013/08216-2, 2014/50268-2, and 2014/04684-4), Conselho Nacional do Desenvolvimento Científico e Tecnológico (CNPq), Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), Fundação de Amparo ao Ensino, Pesquisa e Assistência do Hospital das Clínicas da FMRP/USP (FAEPA), and the Australian National Health and Medical Research Council (NHMRC; grants 1062383 and 1063646 to H.J.N.). T.D.F. and L.M.M. are supported by a fellowship from CAPES. L.D.C., J.M.R., and D.S.L. are supported by fellowships from FAPESP. H.J.N. is a visiting professor from the Science without Borders Program (CNPq; grant 401577/2014-7). D.S.Z. is a research fellow from CNPq, Brazil.

The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

REFERENCES

- van Loenhout JA, Wielders CC, Morroy G, Cox MJ, van der Hoek W, Hautvast JL, Paget WJ, van der Velden J. 2015. Severely impaired health status of non-notified Q fever patients leads to an underestimation of the true burden of disease. *Epidemiol Infect* 143:2580–2587. <http://dx.doi.org/10.1017/S0950268814003689>.
- Khavkin T, Tabibzadeh SS. 1988. Histologic, immunofluorescence, and electron microscopic study of infectious process in mouse lung after intranasal challenge with *Coxiella burnetii*. *Infect Immun* 56:1792–1799.
- Seshadri R, Hendrix LR, Samuel JE. 1999. Differential expression of translational elements by life cycle variants of *Coxiella burnetii*. *Infect Immun* 67:6026–6033.
- Newton HJ, McDonough JA, Roy CR. 2013. Effector protein translocation by the *Coxiella burnetii* Dot/Icm type IV secretion system requires endocytic maturation of the pathogen-occupied vacuole. *PLoS One* 8:e54566. <http://dx.doi.org/10.1371/journal.pone.0054566>.
- Moffatt JH, Newton P, Newton HJ. 2015. *Coxiella burnetii*: turning hostility into a home. *Cell Microbiol* 17:621–631. <http://dx.doi.org/10.1111/cmi.12432>.
- Howe D, Shannon JG, Winfree S, Dorward DW, Heinzen RA. 2010. *Coxiella burnetii* phase I and II variants replicate with similar kinetics in degradative phagolysosome-like compartments of human macrophages. *Infect Immun* 78:3465–3474. <http://dx.doi.org/10.1128/IAI.00406-10>.
- Zamboni DS, Mortara RA, Freymuller E, Rabinovitch M. 2002. Mouse resident peritoneal macrophages partially control *in vitro* infection with *Coxiella burnetii* phase II. *Microbes Infect* 4:591–598. [http://dx.doi.org/10.1016/S1286-4579\(02\)01577-0](http://dx.doi.org/10.1016/S1286-4579(02)01577-0).
- Zamboni DS, Rabinovitch M. 2003. Nitric oxide partially controls *Coxiella burnetii* phase II infection in mouse primary macrophages. *Infect Immun* 71:1225–1233. <http://dx.doi.org/10.1128/IAI.71.3.1225-1233.2003>.
- Zamboni DS, Rabinovitch M. 2004. Phagocytosis of apoptotic cells increases the susceptibility of macrophages to infection with *Coxiella burnetii* phase II through down-modulation of nitric oxide production. *Infect Immun* 72:2075–2080. <http://dx.doi.org/10.1128/IAI.72.4.2075-2080.2004>.
- Voth DE, Howe D, Heinzen RA. 2007. *Coxiella burnetii* inhibits apoptosis in human THP-1 cells and monkey primary alveolar macrophages. *Infect Immun* 75:4263–4271. <http://dx.doi.org/10.1128/IAI.00594-07>.
- Graham JG, MacDonald LJ, Hussain SK, Sharma UM, Kurten RC, Voth DE. 2013. Virulent *Coxiella burnetii* pathotypes productively infect primary human alveolar macrophages. *Cell Microbiol* 15:1012–1025. <http://dx.doi.org/10.1111/cmi.12096>.
- Calverley M, Erickson S, Read AJ, Harmsen AG. 2012. Resident alveolar macrophages are susceptible to and permissive of *Coxiella burnetii* infection. *PLoS One* 7:e51941. <http://dx.doi.org/10.1371/journal.pone.0051941>.
- Yoshiie K, Matayoshi S, Fujimura T, Maeno N, Oda H. 1999. Induced production of nitric oxide and sensitivity of alveolar macrophages derived from mice with different sensitivity to *Coxiella burnetii*. *Acta Virol* 43:273–278.
- Zamboni DS, Mortara RA, Rabinovitch M. 2001. Infection of Vero cells with *Coxiella burnetii* phase II: relative intracellular bacterial load and distribution estimated by confocal laser scanning microscopy and morphometry. *J Microbiol Methods* 43:223–232. [http://dx.doi.org/10.1016/S0167-7012\(00\)00223-2](http://dx.doi.org/10.1016/S0167-7012(00)00223-2).
- Marim FM, Silveira TN, Lima DS, Jr, Zamboni DS. 2010. A method for generation of bone marrow-derived macrophages from cryopreserved mouse bone marrow cells. *PLoS One* 5:e15263. <http://dx.doi.org/10.1371/journal.pone.0015263>.
- Brennan RE, Samuel JE. 2003. Evaluation of *Coxiella burnetii* antibiotic susceptibilities by real-time PCR assay. *J Clin Microbiol* 41:1869–1874. <http://dx.doi.org/10.1128/JCM.41.5.1869-1874.2003>.
- Cunha LD, Ribeiro JM, Fernandes TD, Massis LM, Khoo CA, Moffatt JH, Newton HJ, Roy CR, Zamboni DS. 2015. Inhibition of inflammasome activation by *Coxiella burnetii* type IV secretion system effector IcaA. *Nat Commun* 6:10205. <http://dx.doi.org/10.1038/ncomms10205>.
- Carey KL, Newton HJ, Luhrmann A, Roy CR. 2011. The *Coxiella burnetii* Dot/Icm system delivers a unique repertoire of type IV effectors into host cells and is required for intracellular replication. *PLoS Pathog* 7:e1002056. <http://dx.doi.org/10.1371/journal.ppat.1002056>.
- Amara AB, Bechah Y, Mege JL. 2012. Immune response and *Coxiella burnetii* invasion. *Adv Exp Med Biol* 984:287–298. http://dx.doi.org/10.1007/978-94-007-4315-1_15.
- Hackstadt T, Williams JC. 1981. Biochemical stratagem for obligate parasitism of eukaryotic cells by *Coxiella burnetii*. *Proc Natl Acad Sci U S A* 78:3240–3244. <http://dx.doi.org/10.1073/pnas.78.5.3240>.
- Maurin M, Raoult D. 1999. Q fever. *Clin Microbiol Rev* 12:518–553.
- Shannon JG, Heinzen RA. 2008. Infection of human monocyte-derived macrophages with *Coxiella burnetii*. *Methods Mol Biol* 431:189–200.
- Beare PA, Gilk SD, Larson CL, Hill J, Stead CM, Omsland A, Cockrell DC, Howe D, Voth DE, Heinzen RA. 2011. Dot/Icm type IVB secretion system requirements for *Coxiella burnetii* growth in human macrophages. *mBio* 2:e00175-11. <http://dx.doi.org/10.1128/mBio.00175-11>.
- Benoit M, Barbarat B, Bernard A, Olive D, Mege JL. 2008. *Coxiella burnetii*, the agent of Q fever, stimulates an atypical M2 activation pro-

- gram in human macrophages. *Eur J Immunol* 38:1065–1070. <http://dx.doi.org/10.1002/eji.200738067>.
25. Mehraj V, Textoris J, Ben Amara A, Ghigo E, Raoult D, Capo C, Mege JL. 2013. Monocyte responses in the context of Q fever: from a static polarized model to a kinetic model of activation. *J Infect Dis* 208:942–951. <http://dx.doi.org/10.1093/infdis/jit266>.
 26. Bradley WP, Boyer MA, Nguyen HT, Birdwell LD, Yu J, Ribeiro JM, Weiss SR, Zamboni DS, Roy CR, Shin S. 2016. Primary role for Toll-like receptor-driven tumor necrosis factor rather than cytosolic immune detection in restricting *Coxiella burnetii* phase II replication within mouse macrophages. *Infect Immun* 84:998–1015. <http://dx.doi.org/10.1128/IAI.01536-15>.
 27. Martinez FO, Gordon S, Locati M, Mantovani A. 2006. Transcriptional profiling of the human monocyte-to-macrophage differentiation and polarization: new molecules and patterns of gene expression. *J Immunol* 177:7303–7311. <http://dx.doi.org/10.4049/jimmunol.177.10.7303>.
 28. Benoit M, Desnues B, Mege JL. 2008. Macrophage polarization in bacterial infections. *J Immunol* 181:3733–3739. <http://dx.doi.org/10.4049/jimmunol.181.6.3733>.
 29. Brennan RE, Russell K, Zhang G, Samuel JE. 2004. Both inducible nitric oxide synthase and NADPH oxidase contribute to the control of virulent phase I *Coxiella burnetii* infections. *Infect Immun* 72:6666–6675. <http://dx.doi.org/10.1128/IAI.72.11.6666-6675.2004>.
 30. Howe D, Barrows LF, Lindstrom NM, Heinzen RA. 2002. Nitric oxide inhibits *Coxiella burnetii* replication and parasitophorous vacuole maturation. *Infect Immun* 70:5140–5147. <http://dx.doi.org/10.1128/IAI.70.9.5140-5147.2002>.
 31. Dinakar C, Malur A, Raychaudhuri B, Buhrow LT, Melton AL, Kavuru MS, Thomassen MJ. 1999. Differential regulation of human blood monocyte and alveolar macrophage inflammatory cytokine production by nitric oxide. *Ann Allergy Asthma Immunol* 82:217–222. [http://dx.doi.org/10.1016/S1081-1206\(10\)62600-2](http://dx.doi.org/10.1016/S1081-1206(10)62600-2).
 32. Turco J, Thompson HA, Winkler HH. 1984. Interferon-gamma inhibits growth of *Coxiella burnetii* in mouse fibroblasts. *Infect Immun* 45:781–783.
 33. Dellacasagrande J, Capo C, Raoult D, Mege JL. 1999. IFN-gamma-mediated control of *Coxiella burnetii* survival in monocytes: the role of cell apoptosis and TNF. *J Immunol* 162:2259–2265.
 34. Ghigo E, Capo C, Tung CH, Raoult D, Gorvel JP, Mege JL. 2002. *Coxiella burnetii* survival in THP-1 monocytes involves the impairment of phagosome maturation: IFN-gamma mediates its restoration and bacterial killing. *J Immunol* 169:4488–4495. <http://dx.doi.org/10.4049/jimmunol.169.8.4488>.
 35. Andoh M, Zhang G, Russell-Lodrigue KE, Shive HR, Weeks BR, Samuel JE. 2007. T cells are essential for bacterial clearance, and gamma interferon, tumor necrosis factor alpha, and B cells are crucial for disease development in *Coxiella burnetii* infection in mice. *Infect Immun* 75:3245–3255. <http://dx.doi.org/10.1128/IAI.01767-06>.
 36. Ochoa-Reparaz J, Sentissi J, Trunkle T, Riccardi C, Pascual DW. 2007. Attenuated *Coxiella burnetii* phase II causes a febrile response in gamma interferon knockout and Toll-like receptor 2 knockout mice and protects against reinfection. *Infect Immun* 75:5845–5858. <http://dx.doi.org/10.1128/IAI.00901-07>.
 37. Xiong X, Meng Y, Wang X, Qi Y, Li J, Duan C, Wen B. 2012. Mice immunized with bone marrow-derived dendritic cells stimulated with recombinant *Coxiella burnetii* Com1 and Mip demonstrate enhanced bacterial clearance in association with a Th1 immune response. *Vaccine* 30:6809–6815. <http://dx.doi.org/10.1016/j.vaccine.2012.09.017>.
 38. Schoffelen T, Sprong T, Bleeker-Rovers CP, Wegdam-Blans MC, Ammerdorffer A, Pronk MJ, Soethoudt YE, van Kasteren ME, Herremans T, Bijlmer HA, Netea MG, van der Meer JW, Joosten LA, van Deuren M. 2014. A combination of interferon-gamma and interleukin-2 production by *Coxiella burnetii*-stimulated circulating cells discriminates between chronic Q fever and past Q fever. *Clin Microbiol Infect* 20:642–650. <http://dx.doi.org/10.1111/1469-0691.12423>.
 39. Schoffelen T, Self JS, Fitzpatrick KA, Netea MG, van Deuren M, Joosten LA, Kersh GJ. 2015. Early cytokine and antibody responses against *Coxiella burnetii* in aerosol infection of BALB/c mice. *Diagn Microbiol Infect Dis* 81:234–239. <http://dx.doi.org/10.1016/j.diagmicrobio.2014.12.008>.
 40. Schoffelen T, Wegdam-Blans MC, Ammerdorffer A, Pronk MJ, Soethoudt YE, Netea MG, van der Meer JW, Bleeker-Rovers CP, van Deuren M. 2015. Specific in vitro interferon-gamma and IL-2 production as biomarkers during treatment of chronic Q fever. *Front Microbiol* 6:93. <http://dx.doi.org/10.3389/fmicb.2015.00093>.
 41. Gordon S. 2003. Alternative activation of macrophages. *Nat Rev Immunol* 3:23–35. <http://dx.doi.org/10.1038/nri978>.
 42. Zamboni DS. 2004. Genetic control of natural resistance of mouse macrophages to *Coxiella burnetii* infection in vitro: macrophages from restrictive strains control parasitophorous vacuole maturation. *Infect Immun* 72:2395–2399. <http://dx.doi.org/10.1128/IAI.72.4.2395-2399.2004>.
 43. Kopf M, Schneider C, Nobs SP. 2015. The development and function of lung-resident macrophages and dendritic cells. *Nat Immunol* 16:36–44. <http://dx.doi.org/10.1038/ni.3052>.
 44. Morales-Nebreda L, Misharin AV, Perlman H, Budinger GR. 2015. The heterogeneity of lung macrophages in the susceptibility to disease. *Eur Respir Rev* 24:505–509. <http://dx.doi.org/10.1183/16000617.0031-2015>.
 45. Williams M, De Kleer I, Henri S, Post S, Vanhoutte L, De Priekx S, Deswarte K, Malissen B, Hammad H, Lambrecht BN. 2013. Alveolar macrophages develop from fetal monocytes that differentiate into long-lived cells in the first week of life via GM-CSF. *J Exp Med* 210:1977–1992. <http://dx.doi.org/10.1084/jem.20131199>.
 46. Gosselin D, Link VM, Romanoski CE, Fonseca GJ, Eichenfield DZ, Spann NJ, Stender JD, Chun HB, Garner H, Geissmann F, Glass CK. 2014. Environment drives selection and function of enhancers controlling tissue-specific macrophage identities. *Cell* 159:1327–1340. <http://dx.doi.org/10.1016/j.cell.2014.11.023>.
 47. Lavin Y, Winter D, Blecher-Gonen R, David E, Keren-Shaul H, Merad M, Jung S, Amit I. 2014. Tissue-resident macrophage enhancer landscapes are shaped by the local microenvironment. *Cell* 159:1312–1326. <http://dx.doi.org/10.1016/j.cell.2014.11.018>.
 48. Gordon S, Pluddemann A, Martinez Estrada F. 2014. Macrophage heterogeneity in tissues: phenotypic diversity and functions. *Immunol Rev* 262:36–55. <http://dx.doi.org/10.1111/imr.12223>.
 49. Graham JG, Winchell CG, Kurten RC, Voth DE. 2016. Development of an ex vivo tissue platform to study the human lung response to *Coxiella burnetii*. *Infect Immun* 84:1438–1445. <http://dx.doi.org/10.1128/IAI.00012-16>.
 50. Nathan C, Shiloh MU. 2000. Reactive oxygen and nitrogen intermediates in the relationship between mammalian hosts and microbial pathogens. *Proc Natl Acad Sci U S A* 97:8841–8848. <http://dx.doi.org/10.1073/pnas.97.16.8841>.
 51. Hill J, Samuel JE. 2011. *Coxiella burnetii* acid phosphatase inhibits the release of reactive oxygen intermediates in polymorphonuclear leukocytes. *Infect Immun* 79:414–420. <http://dx.doi.org/10.1128/IAI.01011-10>.

10. ANEXO II – TRABALHOS DE COLABORAÇÃO*

10.1. Anexo 2 - Artigo publicado (CUNHA, Larissa D; RIBEIRO, Juliana M; FERNANDES, Talita D; MASSIS, Liliana M; KHOO, Chen A; MOFFATT, Jennifer H; NEWTON, Hayley J; ROY, Craig R; ZAMBONI, Dario S. *Nature Communications*, 2015)

“Inhibition of inflammasome activation by *Coxiella burnetii* type IV secretion system effector IcaA”

- Publicado em: dezembro de 2015

ARTICLE

Received 3 Sep 2015 | Accepted 13 Nov 2015 | Published 21 Dec 2015

DOI: 10.1038/ncomms10205

OPEN

Inhibition of inflammasome activation by *Coxiella burnetii* type IV secretion system effector IcaA

Larissa D. Cunha^{1,†}, Juliana M. Ribeiro¹, Talita D. Fernandes¹, Liliana M. Massis¹, Chen Ai Khoo², Jennifer H. Moffatt², Hayley J. Newton², Craig R. Roy³ & Dario S. Zamboni¹

Coxiella burnetii is a highly infectious bacterium that promotes its own replication in macrophages by inhibiting several host cell responses. Here, we show that *C. burnetii* inhibits caspase-1 activation in primary mouse macrophages. By using co-infection experiments, we determine that the infection of macrophages with *C. burnetii* inhibits the caspase-11-mediated non-canonical activation of the NLRP3 inflammasome induced by subsequent infection with *Escherichia coli* or *Legionella pneumophila*. Genetic screening using flagellin mutants of *L. pneumophila* as a surrogate host, reveals a novel *C. burnetii* gene (IcaA) involved in the inhibition of caspase activation. Expression of IcaA in *L. pneumophila* inhibited the caspase-11 activation in macrophages. Moreover, *icaA*⁻ mutants of *C. burnetii* failed to suppress the caspase-11-mediated inflammasome activation induced by *L. pneumophila*. Our data reveal IcaA as a novel *C. burnetii* effector protein that is secreted by the Dot/Icm type IV secretion system and interferes with the caspase-11-induced, non-canonical activation of the inflammasome.

¹Department of Cell Biology, Medical School of Ribeirão Preto, University of São Paulo (FMRP/USP), Ribeirão Preto, São Paulo 14049-900, Brazil.

²Department of Microbiology and Immunology, University of Melbourne at the Peter Doherty Institute for Infection and Immunity, Melbourne, Victoria 3000, Australia. ³Department of Microbial Pathogenesis, Yale University School of Medicine, New Haven, Connecticut 06536, USA. † Present address: Department

of Immunology, St. Jude Children's Research Hospital, Memphis, Tennessee 38105, USA. Correspondence and requests for materials should be addressed to D.S.Z. (email: dszamboni@fmrp.usp.br).

Innate immune cells have developed systems to signal and mount appropriate responses to pathogenic microbes by recognizing conserved molecules present in microbes (pathogen-associated molecular patterns, PAMPs)¹. For intracellular pathogens, cytosolic nucleotide binding domain and leucine rich repeat containing proteins (NLR) play a pivotal role in recognizing and mounting effective immune responses. Caspase-1 activation mediated by NLR leads to interleukin-1 β (IL-1 β), IL-18, IL-1 α release, membrane permeabilization and a specific form of inflammatory cell death called pyroptosis². The activation of caspase-1 by NLR occurs in a molecular platform known as an inflammasome³. Notably, the adaptor protein apoptosis-associated speck-like protein containing a carboxy-terminal CARD (ASC) is involved in the structure of different inflammasomes². Several NLR, including NLRC4, NLRP1, NLRP3, NLRP6, NLRP12, neuronal apoptosis inhibitory protein 2 (NAIP2), NAIP5, NAIP6 and the cytosolic non-NLR receptor absent in melanoma 2 (AIM2), have been shown to activate the inflammasome in response to molecules derived from the different classes of pathogens. In particular, the specialized secretion systems of pathogenic Gram-negative bacteria have been shown to be a direct PAMP for inflammasome activation, and to be involved in processes that lead to delivery of specific PAMPs into the host cell cytoplasm^{4–8}. In the case of the intracellular Gram-negative bacterium *Legionella pneumophila*, caspase-1 activation is readily triggered in response to the recognition of bacterial flagellin by NAIP5 and NLRC4, a process that requires the expression of a functional type IV secretion system (T4SS) called defect in organelle trafficking/intracellular multiplication (Dot/Icm)^{9–11}. Most recently, caspase-11 has emerged as an important regulator of inflammasome activation in response to the lipopolysaccharide (LPS) of Gram-negative bacterial species that either express a secretion system or escape to the host cell cytosol^{12–18}. Caspase-11 mediates NLRP3-dependent caspase-1 activation (called the non-canonical activation of the NLRP3 inflammasome), but also induces a caspase-1-independent membrane permeabilization that leads to pyroptosis and the release of inflammatory cytokines, including IL-1 α .

Given the presence of a large repertoire of intracellular pattern-recognition receptors mediating inflammasome activation and pyroptotic cell death, the bona fide intracellular pathogens have developed efficient strategies to manipulate host cell functions and subvert host surveillance (reviewed in Cunha *et al.*¹⁹). In this sense, the expression of specialized secretion systems that mediate the injection of pathogen effector proteins directly into host cell cytosol constitutes a widespread strategy used by intracellular bacterial pathogens to manipulate host cell functions^{20–22}. For intracellular pathogens that target innate immune cells and are associated with severe and highly threatening chronic diseases, such as *Chlamydia trachomatis*, *Coxiella burnetii* and *Mycobacterium tuberculosis*, it is predictable that these pathogens encode unique defence mechanisms to subvert inflammasome activation. In fact, factors that inhibit inflammasome activation have been identified in *M. tuberculosis* and in *C. trachomatis*, although the molecular mechanisms of inflammasome inhibition and host cell targets are unclear^{23–25}.

The intracellular bacterial pathogen *C. burnetii* is the causative agent of an acute pneumonia-like disease called Q fever, which may lead to endocarditis or hepatitis²⁶. The bacteria replicate within a phagolysosomal vacuole and infect primarily alveolar macrophages (AMs)²⁷. *C. burnetii* is highly specialized to subvert host cell functions, including the avoidance of TLRs recognition, the inhibition of apoptosis and the modulation of diverse vesicle traffic pathways^{28–39}. However, immunocompetent individuals can effectively control bacterial multiplication, thus emphasizing

that effective immune responses are critical for host resistance. *C. burnetii* is evolutionarily close to *L. pneumophila*, belonging to the order of Legionellales. Importantly, these pathogens share a homologous Dot/Icm type IV secretion system⁴⁰. Historically, the genetic manipulation of *C. burnetii* has not been easily achieved, and as such, *L. pneumophila* has been used as a surrogate host to characterize and identify the *C. burnetii* Dot/Icm secretion system^{41,42}. *C. burnetii* encodes diverse putative effector proteins in its genome, some of them with sequence motifs predicted to mediate eukaryotic protein–protein interactions such as ankyrin repeat-containing and leucine-rich repetition-containing domains^{40,43}. More than 100 putative *C. burnetii* effectors identified by genetic and bioinformatic whole-genome screens have been shown to be secreted by the Dot/Icm system of *L. pneumophila*^{44–50}. Despite this massive identification of the *C. burnetii* effectors, the molecular function of very few effector proteins has been defined. Because *C. burnetii* is a highly virulent intracellular pathogen and very few cells (<5 organisms) are sufficient to induce disease in a healthy human being, we investigated whether inflammasome activation occurs in response to *C. burnetii* infection. We found that *C. burnetii* inhibits the non-canonical activation of the NLRP3 inflammasome, which occurs in a process dependent on caspase-11 and independent of NLRC4 and bacterial flagellin. To search for the mechanism underlying the inhibition of the inflammasome by the bacterium, we screened potential effectors using *L. pneumophila* as a surrogate host and identified a *C. burnetii* effector protein, denoted herein as IcaA (*Inhibition of caspase activation*), that interferes with the caspase-11-mediated, non-canonical activation of the inflammasome. *C. burnetii* mutants for *icaA* fail to inhibit caspase-11 activation induced by *flaA* mutants of *L. pneumophila*. These findings unveil the process of inflammasome inhibition by *C. burnetii* and therefore contribute to our understanding of the molecular mechanisms underlying the virulence strategies used by this highly threatening obligate intracellular pathogen.

Results

C. burnetii infection does not stimulate inflammasome activation.

To investigate whether *C. burnetii* induces caspase-1 activation, we infected murine bone marrow-derived macrophages (BMDMs) for 24 h and assessed the processing of caspase-1 and pro-IL-1 β by western blot. We found that the infection of wild-type C57BL/6 BMDMs (BL/6 BMDMs) with *C. burnetii* does not induce the cleavage of caspase-1 p20 subunit (Fig. 1a). As a control, we used *L. pneumophila*, which readily triggers the processing of these proteins in infected BMDMs (Fig. 1a). To further evaluate caspase-1 activation in response to *C. burnetii*, we assessed caspase-1 activation in BMDMs using the FAM-YVAD-FMK (FLICA), a previously described fluorescent probe that specifically binds the active form of caspase-1 while inhibiting its autolytic processing¹¹. We found that in contrast to *L. pneumophila* infection, *C. burnetii* infection does not induce an increase in the percentage of cells stained with FLICA, as measured by flow cytometry (Fig. 1b,c). As a control, we used BMDMs from a *Casp1/11*^{−/−} mouse to indicate that this probe is specific to caspase-1. Finally, we measured the secretion of IL-1 β as readout for caspase-1 activation. We found that as opposed to *L. pneumophila* infection, *C. burnetii* infection does not induce the secretion of IL-1 β (Fig. 1a,d). Altogether, these data show that *C. burnetii* does not trigger caspase-1 activation in BMDMs.

C. burnetii inhibits NLRP3 activation induced by caspase-11.

Although macrophages use multiple sensors to detect

Figure 1 | Caspase-1 is not activated in macrophages infected with *C. burnetii*. Bone marrow-derived macrophages (BMDMs) remained noninfected (NI) or were infected with wild-type *L. pneumophila* for 9 h (WT Lp, multiplicity of infection (MOI) 10) or *C. burnetii* for 24 h (Cb, MOI 30). **(a)** Immunoblot showing processed p20 subunit of caspase-1 (caspase-1 p20), unprocessed caspase-1 (pro-caspase-1), p17 subunit of mature IL-1β (IL-1β p17) and pro-IL-1β, as determined in the supernatant (SN) and in the cell extract (CE). **(b)** Flow analysis of FLICA staining in BL/6 BMDM noninfected (green curve), infected with Cb (red curve) or infected with WT Lp (blue curve) and *Casp1/11*^{-/-} BMDMs infected with WT Lp (grey curve). **(c)** Quantification of FLICA-positive cells in **(b)**. **(d)** IL-1β secretion in infected BMDMs was determined by ELISA. Data in **(c,d)** are expressed as average ± s.e.m. of triplicate wells and significance was calculated with *t*-test. **P* < 0.05 compared with NI cells. Data are representative of at least three independent experiments. Full blots are presented in the Supplementary Fig. 10.

Gram-negative bacteria, *C. burnetii* can evade immune responses and interfere with different cellular process to establish a replicative niche in these cells. Therefore, we investigated whether the absence of caspase-1 activation results from the active process of inhibition by *C. burnetii*. Previous work has demonstrated that *C. burnetii* and *L. pneumophila* establish distinct and functional replicative vacuoles in co-infected BMDMs⁵¹. We thus performed co-infection studies to evaluate whether *C. burnetii* can actively inhibit the inflammasome activation triggered by *L. pneumophila*. We found that pre-infection of BMDMs with *C. burnetii* for 24 h resulted in reduced caspase-1 activation and pro-IL-1β cleavage in response to a second infection with *L. pneumophila* (Fig. 2a). Notably, pre-infection with *C. burnetii* induced a significantly higher expression of pro-casp-1 and pro-IL-1β, than single infections with *L. pneumophila*, possibly because of the presence of *C. burnetii* PAMPs during the 24-h pre-stimulation. To guarantee that the cells were stimulated similarly during the co-infection experiments, we pre-infected BMDMs with live or paraformaldehyde (PFA)-fixed *C. burnetii* before infections with *L. pneumophila*. We found that PFA-inactivated *C. burnetii* induced a similar expression of pro-casp-1 as the live bacteria, but only live *C. burnetii* inhibited caspase-1 activation in response to *L. pneumophila* (Supplementary Fig. 1a). These data indicate that the inhibition of caspase-1 activation is an actively induced process that requires *C. burnetii* metabolic activity. Caspase-1 activation in response to *L. pneumophila* occurs via at least two distinct pathways, one dependent on and another independent of bacterial flagellin⁵². To evaluate whether *C. burnetii* inhibits the flagellin-dependent or -independent pathway triggered by *L. pneumophila* infection, we performed co-infections of *C. burnetii* with wild-type (WT Lp) or flagellin-deficient (*flaA*⁻) *L. pneumophila*. We found that *C. burnetii* pre-infection inhibits caspase-1 activation and IL-1β processing in response to either WT Lp or *flaA*⁻ *L. pneumophila* (Fig. 2a). Strikingly, the co-infections performed with *flaA*⁻ *L. pneumophila* revealed that the flagellin-independent pathway

for caspase-1 activation was abolished when the BMDMs were pre-infected with *C. burnetii* (Fig. 2a). These data indicate that *C. burnetii* effectively inhibits the flagellin-independent pathway for caspase-1 activation. Similar experiments performed with PFA-fixed *C. burnetii* indicated that live *C. burnetii* is required for the inhibition of caspase-1 activation in response to *flaA*⁻ *L. pneumophila* (Supplementary Fig. 1b).

These findings were further confirmed by the assessment of endogenous caspase-1 activation with the FLICA assay. BMDMs pre-infected with *C. burnetii* showed a reduction in caspase-1 activation in response to infection with both the WT Lp and *flaA*⁻ *L. pneumophila* (Fig. 2b–d). As demonstrated by western blot, when co-infections were performed using *flaA*⁻ *L. pneumophila*, we found that BMDMs do not trigger caspase-1 activation (Fig. 2c,d). Pre-infections with *C. burnetii* did not reduce the internalization of *L. pneumophila* in BMDMs (Supplementary Fig. 2), confirming that this phenomenon is not caused by differences in *L. pneumophila* internalization after a previous exposure to *C. burnetii*. The *C. burnetii*-mediated inhibition of caspase-1 activation induced by *L. pneumophila* was also assessed by measuring the secretion of active IL-1β by enzyme-linked immunosorbent assay (ELISA). We found that pre-infection with live *C. burnetii* inhibited the secretion of IL-1β in response to infection with WT Lp, and abolished the secretion of IL-1β in response to infection with *flaA*⁻ *L. pneumophila* (Fig. 2e).

We investigated whether *C. burnetii* is able to inhibit caspase-1 activation in response to other stimuli. Thus, BMDMs pre-infected or not with *C. burnetii* were stimulated with LPS + ATP or with LPS + dsDNA to trigger the activation of the inflammasomes dependent on NLRP3 and AIM2, respectively. We found that *C. burnetii* fails to inhibit caspase-1 activation in response to ATP (Supplementary Fig. 3a,c) and dsDNA (Supplementary Fig. 3b,d). These results argue that *C. burnetii* does not directly inhibit caspase-1 activation but may interfere with upstream events in the signalling cascade of the inflammasome

Figure 2 | *C. burnetii* inhibits the caspase-11-mediated non-canonical activation of the NLRP3 inflammasome induced by *L. pneumophila*. Bone marrow-derived macrophages (BMDMs) were left noninfected (NI) or were infected with *C. burnetii* (Cb, MOI 30) for 24 h and further infected with wild-type *L. pneumophila* (WT Lp, MOI 10) or *flaA⁻* (MOI 10) for 9 h. Co-infected BMDMs are indicated (Cb + WT Lp or Cb + *flaA⁻*). **(a)** Immunoblot showing levels of processed p20 subunit of caspase-1 (caspase-1 p20), unprocessed caspase-1 (pro-caspase-1), p17 subunit of mature IL-1β (IL-1β p17) and pro-IL-1β, as determined in supernatant (SN) and cell extract (CE). **(b,c)** Flow analysis of FLICA staining in BL/6 BMDM noninfected (grey curve), infected with Cb (green curve), infected with *L. pneumophila* (blue curve) or co-infected (red curve). **(d)** Quantification of FLICA-positive cells in **(b,c)**. **(e)** IL-1β secretion in infected BMDMs was determined by ELISA. BMDMs infected with live *C. burnetii* (live Cb) or with PFA-fixed were used. **(f)** Caspase-1 cleavage and IL-1β secretion were evaluated by immunoblot as in **(a)**. Data in **(d,e)** are expressed as average ± s.e.m. of triplicate wells and significance was calculated with ANOVA. **P* < 0.05. Data are representative of three **(a,f)** and two **(b-e)** independent experiments.

triggered by *L. pneumophila*. It was recently demonstrated that *L. pneumophila* triggers the non-canonical activation of the NLRP3 inflammasome in a process independent of flagellin and dependent on caspase-11 (ref. 13). Thus, we used BMDMs from *Nlrp3^{-/-}*, *Nlrp3^{-/-}*, *Asc^{-/-}* or *Casp11^{-/-}* mice to evaluate the requirement of these proteins for the *C. burnetii*-mediated inhibition of caspase-1 activation in response to *L. pneumophila*. We found that the inhibition of caspase-1 processing occurs in *Nlrp3^{-/-}* BMDMs, a feature that is consistent with the findings that *C. burnetii* inhibits the flagellin-independent pathway for caspase-1 activation (Fig. 2f). In the absence of ASC, we detected no caspase-1 processing. In *Nlrp3^{-/-}* and *Casp11^{-/-}* BMDMs, we found that *C. burnetii* fails to inhibit caspase-1 cleavage and the processing of pro-IL-1β (Fig. 2f). This feature is consistent with the hypothesis that *C. burnetii* inhibits the non-canonical activation of the NLRP3 inflammasome, which occurs through caspase-11 and culminates in the activation of the NLRP3/ASC/caspase-1 platform, leading to caspase-1 cleavage and IL-1β production⁵³. It was previously demonstrated that Gram-negative bacteria, such as *E. coli*, or the combination of LPS plus the B subunit of cholera toxin

(CTB) triggers the non-canonical activation of the inflammasome^{18,53}. Thus, to test whether *C. burnetii* inhibits the non-canonical activation of the inflammasome, we pre-infected BMDMs with *C. burnetii*, stimulated with either LPS + CTB or *E. coli* and measured caspase-1 activation. We found that *C. burnetii* pre-infection severely inhibited caspase-1 activation and pro-IL-1β processing in response to *E. coli* (Fig. 3a). According to previous reports, the *E. coli*-induced activation of caspase-1 and pro-IL-1β cleavage was dependent on NLRP3 and ASC (Fig. 3a). Supporting the inhibitory activity of *C. burnetii* in the non-canonical inflammasome activation, we found that the release of IL-1β and IL-1α in response to *E. coli* or LPS + CTB is reduced in BMDMs pre-infected with *C. burnetii* (Fig. 3b–e). Next, we used a pore formation assay that reflects a caspase-11 activity that operates independently of NLRC4 inflammasome and upstream of the non-canonical activation of the NLRP3 inflammasome^{13,54}. By co-infecting cells with *C. burnetii* and *E. coli*, we found that *C. burnetii* is able to inhibit the caspase-11-mediated pore formation induced by *E. coli* (Fig. 3f–h). In this experiment, we also included BMDMs from a mouse deficient in caspase-1. This mouse, herein called

Figure 3 | *C. burnetii* inhibits caspase-11-dependent pore formation and NLRP3 activation in response to *E. coli* and cholera toxin B. Bone marrow-derived macrophages (BMDMs) generated from wild-type C57BL/6 (BL/6), *Nlrp3*^{-/-}, *Asc*^{-/-} or *Casp1/11*^{-/-} mice were left noninfected (NI) or were infected with *C. burnetii* (Cb, MOI 30) for 24 h, stimulated with LPS (0.5 μg ml⁻¹) for 4 h and infected with *E. coli* at MOI 20 (Ec or Cb + Ec) or treated with CTB at 20 μg ml⁻¹ (CTB or Cb + CTB). **(a)** Immunoblot showing levels of processed p20 subunit of caspase-1 (caspase-1 p20), unprocessed caspase-1 (pro-caspase-1) and p17 subunit of mature IL-1β (IL-1β p17), as determined in supernatant (SN) and cell extract (CE) after 12 h of infection with *E. coli*. **(b)** IL-1β and **(c)** IL-1α were determined by ELISA from cell supernatants 16 h after *E. coli* infection. **(d)** IL-1β and **(e)** IL-1α were determined by ELISA from cell supernatants 16 h after CTB stimulation. **(f–i)** Pore formation was assessed fluorometrically in real time by the uptake of propidium iodide (RFUs, relative fluorescence units). **(b–e)** Data are expressed as the average ± s.e.m. of triplicate wells and significance was calculated with ANOVA. *P < 0.05. **(a–i)** Data are representative of two independent experiments.

Casp1^{-/-}/*Casp11*^{Tg}, is the *Casp1/11*^{-/-} expressing a transgene encoding a functional copy of the caspase-11 allele, as described previously⁵³. By using *Casp1*^{-/-}/*Casp11*^{Tg} BMDMs, we confirmed that *C. burnetii* is able to inhibit caspase-11-mediated pore formation induced by *E. coli* in the absence of caspase-1 (Fig. 3i). Next, we tested whether the inhibition of caspase-11 also operates in primary mouse AMs, which are permissive for *C. burnetii* replication, and are more similar to the relevant cells that harbour *C. burnetii* during acute Q fever

(Supplementary Fig. 4a). By using AMs, we found that co-infections with live *C. burnetii* but not with heat-killed bacteria inhibit the caspase-11-dependent pore formation induced by *E. coli* in primary AMs similarly to BMDMs (Supplementary Fig. 4b,c).

A genetic screen using *L. pneumophila* identifies *C. burnetii* IcaA. We used *L. pneumophila* as a surrogate host to screen the specific *C. burnetii* effector proteins involved in the inhibition of

non-canonical inflammasome activation, and we constructed a library of *flaA*⁻ mutants of *L. pneumophila* expressing *C. burnetii* effectors that have been confirmed to be secreted via the *L. pneumophila* type IV secretion system^{44,45}. We obtained *L. pneumophila* clones with a confirmed expression of 24 *C. burnetii* effectors (Supplementary Fig. 5a,b). *L. pneumophila* mutants expressing *C. burnetii* genes were screened by western blot for the induction of caspase-1 cleavage in infected BMDMs. This screening identified a strain that induced neither caspase-1 activation nor IL-1 β maturation, which was the clone of *L. pneumophila* that produces the effector protein annotated as CBU1823 (Supplementary Fig. 5c–e). Although biochemical activities displayed by the CBU1823 protein are yet unknown, this protein has been shown to be secreted into host cells through the *L. pneumophila* Dot/Icm system^{44,45}. In addition, a recent report demonstrated that CBU1823 is translocated into BMDMs during *C. burnetii* infection in a process dependent on the *C. burnetii* Dot/Icm system⁵⁵. Because we identified CBU1823 as a *C. burnetii* Dot/Icm effector protein that inhibits caspase activation, we named this protein IcaA. IcaA shows no known protein domain and has no significant homology to any other eukaryotic or bacterial protein to hint at its mechanisms of function.

We further tested whether the expression of IcaA in different strains of *flaA*⁻ *L. pneumophila* displayed reduced caspase-1 cleavage in BMDMs. We found that the inhibition of caspase-1 cleavage by IcaA occurred in both JR32 and Lp01 backgrounds (Supplementary Fig. 6a). Inhibition also occurred when we used the strain Lp02, which is a thymidine auxotroph and does not replicate intracellularly in the absence of thymidine supplementation (Supplementary Fig. 6a). Hence, IcaA-mediated caspase-1 inhibition is independent of bacterial replication. Next, we investigated whether the expression of IcaA influenced *L. pneumophila* fitness. We compared *flaA*⁻ mutants of *L. pneumophila* expressing IcaA (*flaA*⁻/pIcaA) with those encoding the empty vector (*flaA*⁻/pVec) in assays that measured replication in axenic media, internalization in BMDMs and the induction of cytokines. We found that the expression of IcaA does not affect the bacterial replication in liquid ACES-buffered yeast extract (AYE) axenic media, bacterial internalization in BMDMs or the induction of IL-12 in BMDMs (Supplementary Fig. 6b–d). Moreover, the expression of IcaA or the unrelated *C. burnetii* effector protein AnkH in *flaA*⁻ *L. pneumophila* did not interfere with internalization or the early bacterial replication in BMDMs at 9 h of infection compared with the bacteria encoding the empty vector (Supplementary Fig. 6e,f). The bacteria encoding pIcaA, but not those encoding the pVec, showed a reduced activation of caspase-1 and secretion of IL-1 β (Supplementary Fig. 6a,g).

Our studies, using pre-infections with live *C. burnetii*, supported the findings that *C. burnetii* inhibits the non-canonical activation of the inflammasome. Therefore, we used the *flaA*⁻ *L. pneumophila*-carrying pIcaA (or pVec) to evaluate whether IcaA is involved in the inhibition of the caspase-11-mediated, non-canonical activation of the NLRP3 inflammasome. Initially, we investigated whether caspase-11 is required for the IcaA-mediated inhibition of caspase-1 activation. We found that whereas *flaA*⁻ mutants encoding pVec trigger caspase-1 activation in BL/6 BMDMs, the mutants expressing IcaA fail to do so. Importantly, the IcaA-mediated inhibition of caspase-1 activation is not observed in BMDMs from mice of a 129S6/SvEv genetic background (herein called 129), a strain that fails to express functional caspase-11 (Fig. 4a,b). To certify that the lack of caspase-1 inhibition in 129 mouse strains was due to the absence of caspase-11, we transduced 129 BMDMs with a retrovirus-encoding caspase-11 (or GFP as a control). The

transduced cells expressed a significant amount of p43 and p38 isoforms of pro-caspase-11 (Fig. 4b). By infecting the transduced cells, we found that *flaA*⁻ *L. pneumophila* expressing pIcaA inhibits the caspase-1 cleavage in 129 BMDMs expressing caspase-11 but not in cells transduced with retrovirus-encoding GFP (Fig. 4c). In agreement with these findings, we found that *flaA*⁻/pIcaA mutants do not inhibit caspase-1 cleavage or IL-1 β secretion in LPS-primed *Casp11*^{-/-} BMDMs in the BL/6 genetic background (Fig. 4d,e). In contrast, in the BL/6 BMDMs, IcaA expression partially inhibited caspase-1 activation and IL-1 β secretion induced by *flaA*⁻ *L. pneumophila*, which are convenient proxy measures for caspase-11 activation (Fig. 4d,e). Supporting the role of caspase-11 in the flagellin-independent pathway for caspase-1 activation, we found that BMDMs from *Casp11*^{-/-} mice show a reduced activation of caspase-1 and secretion of IL-1 β compared with BL/6 BMDMs (Fig. 4d,e).

IcaA inhibits caspase-11-mediated inflammasome activation.

Our data reported thus far indicate that IcaA is involved in the inhibition of a caspase-11-dependent, non-canonical activation of inflammasomes. Therefore, we tested whether IcaA interferes with a caspase-11-dependent process that is upstream of caspase-1. We have previously reported that caspase-11 is involved in the formation of pores in the macrophage membrane that are independent of NLRP3, ASC and caspase-1 (ref. 13). Therefore, we investigated whether *flaA*⁻ mutants expressing IcaA interfere with caspase-11-mediated pore formation in BMDMs. By analysing the uptake of cell impermeable dye ethidium bromide, we found a partial reduction in pore formation in BMDMs infected with *flaA*⁻/pIcaA bacteria in comparison to those infected with *flaA*⁻/pVec (Fig. 5a,b). Next, we performed analyses using the less toxic dye propidium iodide (PI) to evaluate the kinetics of pore formation in real time. By using this approach, we confirmed that IcaA, but not the unrelated *C. burnetii* effector AnkH, partially inhibits the caspase-11-dependent pore formation observed in response to *flaA*⁻ *L. pneumophila* (Fig. 5c). Pore formation was not detected in *Casp11*^{-/-} or *Casp11*^{+/+} BMDMs, emphasizing that the flagellin-independent pore formation is caspase-11-dependent (Fig. 5d,e). Notably, the IcaA-mediated inhibition of pore formation occurred normally in BMDMs from *Nlrp3*^{-/-}, *Asc*^{-/-} and *Casp1*^{-/-}/*Casp11*^{Tg} mice (Fig. 5f–h). These data further confirm that IcaA interferes with caspase-11-mediated responses but not in the downstream components of the NLRP3 inflammasome, such as NLRP3, ASC and caspase-1. To further evaluate whether the IcaA-mediated inhibition of pore formation is specific to caspase-11, we expressed IcaA in flagellated *L. pneumophila*, which is known to trigger pore formation by a process that is Nlrp4- and caspase-1-dependent but caspase-11-independent^{10,13,52,54,56,57}. By infecting BMDMs from BL/6, *Casp11*^{-/-} and *Casp11*^{+/+} mice with *flaA*⁺/pIcaA or *flaA*⁺/pVec, we found that the IcaA is not effective in inhibiting the flagellin-mediated pore formation that is dependent on caspase-1 but independent of caspase-11 (Supplementary Fig. 7a). As expected, infections performed with *flaA*⁻/pIcaA inhibited caspase-11-dependent pore formation compared with *flaA*⁻/pVec (Supplementary Fig. 7b). All these data indicate that IcaA is specifically involved in the inhibition of caspase-11-mediated processes, but it does not influence caspase-1-mediated processes that do not require caspase-11. Our assertion that *C. burnetii* inhibits a caspase-11-dependent pathway implies that the bacteria express molecules that trigger this pathway. To evaluate whether *C. burnetii* is able to trigger caspase-11 activation, we extracted *C. burnetii* LPS and tested for the induction of pore formation upon

Figure 4 | IcaA requires a functional caspase-11 to inhibit caspase-1 activation. Bone marrow-derived macrophages (BMDMs) generated from C57BL/6 (BL/6) and 129S6/SvEv (129) mice were left noninfected (NI) or were infected with *flaA*⁻ mutants of *L. pneumophila* that were transformed either with an empty vector (*flaA*⁻/pVec) or the vector encoding IcaA (*flaA*⁻/pIcaA) at MOI 10 for 9 h. **(a)** Immunoblot showing processed p20 subunit of caspase-1 (caspase-1 p20) and unprocessed caspase-1 (pro-caspase-1), as determined in supernatant (SN) and cell extract (CE). **(b)** Immunoblot showing the expression of unprocessed isoforms of pro-caspase-11 (p43 and p38) in uninfected BL/6 and caspase-11-deficient BMDMs derived from 129 mouse strain were transduced with the pMSCV virus encoding GFP (pGFP) or pMSCV encoding full-length caspase-11 (pCasp-11). The expression of unprocessed isoforms of pro-caspase-11 (p43 and p38) is shown by western blot. **(c)** Immunoblot showing caspase-1 p20 and pro-caspase-1 as determined in SN and CE from transduced BMDMs infected for 9 h. **(d)** Immunoblot showing caspase-1 p20 and pro-caspase-1 in SN and CE of BMDMs primed with LPS (0.5 μg ml⁻¹) and infected for 3 h. **(e)** IL-1β was determined by ELISA from SN of BMDMs primed with LPS for 3 h. Data are expressed as the average ± s.e.m. of triplicate wells and significance was calculated with ANOVA. **P* < 0.05. Data are representative of at least two (**b,c**) or three (**a,d,e**) independent experiments.

the transfection of BMDMs. We detected a robust pore formation in BMDMs transfected with *C. burnetii* LPS, which was reduced in *Casp1*^{-/-}/*Casp11*^{Tg} and *Casp11*^{-/-} BMDMs (Supplementary Fig. 8). These data support the hypothesis that *C. burnetii* expresses molecules that trigger pore formation if delivered in the macrophage cytoplasm, a feature that supports the development of effectors, such as IcaA, to inhibit this response.

Our data generated thus far indicate that IcaA is an effector protein encoded by *C. burnetii* that mediates the inhibition of caspase-11 functions such as pore formation, but it is not clear whether IcaA inhibits caspase-11 activation. Thus, we tested whether IcaA-expressing *flaA*⁻ *L. pneumophila* can reduce the activation of caspase-11 in the infected BMDMs. To measure caspase-11 activation directly, we pulled down active caspase-11 from macrophages lysates using biotin-VAD-FMK. Active caspase-11 bound to biotin-VAD was concentrated with streptavidin, and the samples were blotted with anti-caspase-11.

We detected a robust caspase-11 activation when BL/6 BMDMs were infected with *flaA*⁻ *L. pneumophila* (Fig. 5i). Notably, IcaA expression in *L. pneumophila* reduced caspase-11 activation compared with the bacteria expressing the empty vector (Fig. 5i). The IcaA-mediated inhibition of caspase-11 activation was also observed in experiments performed with macrophages from the *Casp1*^{-/-}/*Casp11*^{Tg} mice (Fig. 5i). Notably, the overall expression of caspase-11 in these cells is lower than that of C57BL/6 macrophages, possibly because only one copy of the *Casp11*^{Tg} is present in the genome. Regardless of the reduced expression of caspase-11 in the *Casp1*^{-/-}/*Casp11*^{Tg} BMDMs, these data further confirm our assertion that IcaA is involved in the inhibition of caspase-11 activation despite the presence of caspase-1 (Fig. 5e).

***icaA*⁻ *C. burnetii* fails to inhibit inflammasome activation.** To further evaluate whether the endogenously expressed IcaA in

Figure 5 | IcaA does not inhibit caspase-11 activation directly but inhibits a caspase-11 activation upstream of the non-canonical activation of the NLRP3 inflammasome. (a–i) Bone marrow-derived macrophages (BMDMs) were primed with LPS ($0.5 \mu\text{g ml}^{-1}$) for 3 h, remained noninfected (NI) or were infected with *flaA*⁻ *L. pneumophila* transformed either with an empty vector (*flaA*⁻/pVec), the vector encoding IcaA (*flaA*⁻/plcaA), or the vector encoding AnkH (*flaA*⁻/pAnkH) at MOI 10. (a) Fluorescence micrographs revealing cell permeability, as assessed by quantification of ethidium bromide influx (red). Healthy cells are stained with acridine orange (green). Scale bar, 75 μm . (b) Quantification of the experiment shown in (a). Data are expressed as the average $\pm$ s.e.m. of triplicate wells and significance was calculated with t-test. * $P < 0.05$. (c–h) Pore formation was assessed fluorometrically in real time by the uptake of propidium iodide (RFUs, relative fluorescence units). (i) Immunoblot showing the presence of caspase-11 p45 and p38 in the total cell lysate (input) and pull-down fraction using the agarose-streptavidin fraction of BMDMs treated with biotin-VAD-FMK 15 min before infection. Actin (β -actin) was used as a loading control. Data are representative of three (a–h) and two (i) independent experiments.

C. burnetii is involved in the inhibition of the inflammasome activation, we generated the *icaA*⁻ mutant *C. burnetii* (Supplementary Fig. 9). Initially, we tested whether *icaA* is able to trigger caspase-11-dependent responses in single macrophage infections. Measuring caspase-11-dependent pore formation in cells infected with *C. burnetii*, we found that neither the wild-type *C. burnetii* nor the *icaA*⁻ mutants triggered a caspase-11-dependent pore formation (Fig. 6a). Accordingly, *icaA*⁻ mutants

failed to trigger the caspase-11-mediated non-canonical activation of the NLRP3 inflammasome, as measured by the production of IL-1 β (Fig. 6b). These data suggest that additional *C. burnetii* processes may be involved in the activities related to the inhibition of inflammasome activation by this bacterium. Next, we tested whether IcaA expression is required for the inhibition of non-canonical inflammasome activation in co-infection experiments. In these experiments, *Nlrp3*^{-/-} BMDMs or *flaA*⁻

Figure 6 | *C. burnetii* mutants for *icaA* fail to inhibit the non-canonical activation of the inflammasome in response to *flaA*⁻ *L. pneumophila*. Bone marrow-derived macrophages (BMDMs) were left noninfected (NI) or were infected with heat-killed wild-type (HK Cb) wild-type (WT Cb) or *icaA*⁻ mutant (*icaA*⁻ Cb) of *C. burnetii* for 24 h at MOI 30, and further infected with wild-type *L. pneumophila* (WT Lp, MOI 10) or *flaA*⁻ (MOI 10) for 9 h. Co-infected BMDMs are indicated (Cb + Lp or Cb + *flaA*⁻). (a) Pore formation was assessed fluorometrically in real time by the uptake of propidium iodide (RFUs, relative fluorescence units). (b,d,e) IL-1β secretion in infected BMDMs was determined by ELISA. Data are expressed as the average ± s.e.m. of triplicate wells and significance was calculated with t-test. **P* < 0.05. (c) Immunoblot showing processed p20 subunit of caspase-1 (caspase-1 p20), unprocessed caspase-1 (pro-caspase-1) and p17 subunit of mature IL-1β (IL-1β p17), as determined in supernatant (SN) and cell extract (CE). (a–e) Data are representative of at least two independent experiments.

L. pneumophila were used to bypass the activation of NLRP4 inflammasome. We found that, whereas a single infection with wild-type *C. burnetii* or with the *icaA*⁻ mutants did not trigger caspase-1 activation in BMDMs, the co-infection with *icaA*⁻ *C. burnetii* failed to suppress the caspase-1 activation induced by *flaA*⁻ *L. pneumophila* (Fig. 6c). Accordingly, co-infections performed with wild-type *C. burnetii*, but not with *icaA* mutants, inhibited the caspase-11-mediated non-canonical activation of the inflammasome, as measured by IL-1β production (Fig. 6d,e). These data further confirm that endogenously expressed IcaA in *C. burnetii* is important for the activities related to the inhibition of the caspase-11-mediated non-canonical activation of the inflammasome.

Discussion

It is estimated that as few as five infectious *C. burnetii* organisms are able to establish infection in healthy individuals exposed to the bacteria⁵⁸. To achieve such a high infectious efficiency, *C. burnetii* inhibits a large repertoire of specific host cell responses, including those related to the innate immune activation and induction of host cell death. Although the inhibition of caspase-3-mediated apoptosis by *C. burnetii* has been reported previously^{29,31,32,35}, an evaluation of inflammatory caspases and pyroptosis has never been conducted. Herein, we report that *C. burnetii* fails to induce caspase-1 activation upon the infection of primary mouse macrophages. We found that the bacteria actively inhibit the caspase-11 activation and their functions including pore formation and the non-canonical activation of the NLRP3 inflammasome, but they do not inhibit the components of inflammasomes such as ASC, NLRP3 or caspase-1. By searching for the molecular mechanisms underlying this process, we identified a novel *C. burnetii* gene that encodes an

Figure 7 | Model for IcaA-mediated inhibition of non-canonical activation of the inflammasome. Published data demonstrate that caspase-11 senses the presence of cytosolic LPS from Gram-negative bacteria, including *E. coli* and *L. pneumophila*. Activation of caspase-11 in response to *L. pneumophila* requires the expression of Dot/Icm. The effector protein IcaA is translocated by the *C. burnetii* Dot/Icm and inhibits the caspase-11-mediated pore formation and caspase-11-induced non-canonical activation of the NLRP3 inflammasome.

effector protein translocated by the Dot/Icm type IVB secretion system into host cell cytosol. We named this protein IcaA because it is functionally related to the inhibition of caspase activation. IcaA suppresses the caspase-11-mediated, non-canonical activation of caspase-1 in macrophages. Notably, we have no data to support a direct effect of IcaA in the inhibition of caspase-11 activation. Most likely, IcaA is involved in the inhibition of

certain cellular processes, which leads to caspase-11 activation (Fig. 7). It is also possible that IcaA operates in the vacuolar membranes by minimizing the release of bacterial molecules from the vacuole to the cytoplasm, thereby reducing the release of PAMPs that trigger cytosolic innate immune sensors. This hypothesis is consistent with our observed ability to transfer the *C. burnetii* inhibitory activity to *L. pneumophila* *in trans* when IcaA was expressed in *L. pneumophila*. This was achieved because of the high homology of *C. burnetii* and *L. pneumophila* Dot/Icm⁴⁰. In fact, *L. pneumophila* has been extensively used as an effective surrogate host to investigate *C. burnetii* pathogenesis. Earlier studies demonstrated that the structural core components of the *C. burnetii* Dot/Icm can complement *L. pneumophila* mutants deficient in the respective homologous gene^{41,42}. Further studies benefited from the high homology between the two Dot/Icm and used *L. pneumophila* to actively translocate *C. burnetii* effector proteins^{44–50}. This approach supported the identification of several effector proteins encoded by *C. burnetii*, including the protein encoded by gene CBU1823 (refs 44,45). However, only recently has the secretion of *C. burnetii* effectors via *C. burnetii* Dot/Icm been efficiently detected experimentally⁴⁵. Notably, CBU1823 (IcaA) was one protein that has been used to demonstrate the functional activities of the *C. burnetii* Dot/Icm; therefore, IcaA is a bona fide effector secreted by the *C. burnetii* Dot/Icm system into the host cell cytoplasm⁵⁵. These data demonstrate that IcaA is a bona fide substrate of the *C. burnetii* Dot/Icm, which can be translocated by the *Legionella* Dot/Icm or by the *Coxiella* Dot/Icm^{44,45,55}.

The genetic manipulation of *C. burnetii* has only recently been achieved^{44,59}. Therefore, little information is available regarding the role of specific *C. burnetii* effector proteins. It is believed that the bacterial effectors are determinants in the pathogenesis of this bacterium. A functional Dot/Icm system is required for *C. burnetii* growth in macrophages and in permissive cells, supporting an essential role of translocated effector proteins for bacterial survival^{44,59}. By random transposon mutagenesis, several putative effectors have been shown to be important for bacterial replication⁴⁸. Moreover, the deletion of CvpA, an effector that interacts with components of clathrin-coated endocytic vesicles, impairs the maturation of the parasitophorous vacuole and bacterial growth³⁰. The gene-encoding IcaA studied herein was identified in the Nine Mile strain of *C. burnetii*, which is representative of acute disease isolates. Analysis of the genome of other important *C. burnetii* strains showed that this effector is conserved in the Dugway⁴⁴, a strain isolated from rodents that is weakly pathogenic for guinea pigs. Notably, IcaA is encoded in truncated forms in the human endocarditis isolates K (Q154) and G (Q212) strains⁴⁴. Whether the IcaA is non-functional in K and G strains due to frameshift mutations and whether this reflects differences in the virulence and spreading of the bacteria to other tissues is a subject for future investigation. Notably, caspase-11-induced susceptibility to infection has been demonstrated in murine models of bacterial dissemination using *Salmonella typhimurium*⁶⁰. In this context, it remains to be determined whether the IcaA-mediated inhibition of caspase-11 and pyroptosis regulates bacterial dissemination and accounts for the development of chronic versus acute Q fever. Another matter for further investigation is the effect of *icaA* in inflammasome activation within human macrophages. It was recently demonstrated that Nine Mile phase II can trigger the production of active IL-1 β in primary human AMs⁶¹. Although we have demonstrated that the IcaA-mediated inhibition of caspase-11 activation also occurs in primary mouse AMs, a direct comparison with primary human AMs will be important for the comprehensive assessment of the role of IcaA in the bacterial pathogenesis and host responses during Q fever. Nonetheless, the

fact that primary human macrophages do trigger IL-1 β production⁶¹ supports our findings that *C. burnetii* does encode molecules responsible for the non-canonical activation of the inflammasome.

Regardless of the determination of the role of IcaA and caspase-11 in the outcome of *C. burnetii* infection *in vivo*, our study identifies IcaA as a novel bacterial substrate that participates in the inhibition of non-canonical inflammasome activation. To the best of our knowledge, this is the first bacterial effector that interferes with pyroptosis and the activation of the non-canonical inflammasome activation in macrophages¹⁹. By using epithelial cells, OspC3 was identified as a *Shigella flexneri* effector protein that functions as a direct inhibitor of human caspase-4 (ref. 62). This protein was shown to be important for bacterial pathogenesis using a guinea pig model of *S. flexneri* infection. However, it remains unclear whether OspC3 operates in macrophages and whether it interferes with the non-canonical activation of the inflammasome. Regardless of the effect of OspC3 in human epithelial cells, we propose that the *C. burnetii* effector protein IcaA interferes with the process involved in caspase-11 activation, consequently inhibiting non-canonical inflammasome activation in macrophages, a molecular platform that has recently emerged as major player in innate immune responses to Gram-negative bacteria^{12–18,53}. *C. burnetii* does not inhibit caspase-1 or other inflammasomes such as the NAIP5/NLRC4 or the canonical NLRP3. Because *C. burnetii* does not encode genes for flagellin expression, the bacteria may not have encountered selective pressure to develop molecules that target the activation of the NAIP5/NLRC4 inflammasome. In contrast, *C. burnetii* virulence is associated with the presence of the O-antigen in its LPS containing a tetra-acylated lipid A with long fatty acid chains³⁹. Because it has been recently reported that the non-canonical activation of the inflammasome proceeds through the intracellular recognition of cytosolic LPS^{16,17}, it is possible that the *C. burnetii* LPS triggers this pathway, and the bacterium has evolutionarily evolved effectors, such as IcaA, that facilitate the subversion of this important innate immune pathway. In support of this hypothesis, we found that the delivery of *C. burnetii* LPS into the macrophage cytoplasm triggers this pathway. Notably, tetra-acylated lipid A was reported to be inefficient for activating caspase-11 (ref. 16). It is possible that variations in the *Coxiella* lipid A occur according to the bacterial growth phase. Alternatively, it is possible that another bacterial molecule is responsible for triggering caspase-11-dependent pore formation. Nonetheless, the induction of pores in macrophages transfected with *C. burnetii* LPS indicates that *C. burnetii* can trigger a pathway that is inhibited by IcaA. Collectively, the identification of IcaA and the unravelling of its mechanistic functions effectively contribute to our understanding of the biology and pathogenesis of this highly infectious intracellular pathogen and provides novel molecular structures that can be used in the development of therapies for sepsis and chronic inflammatory diseases.

Methods

Bacterial strains. *C. burnetii* (strain Nine Mile RSA 493 phase II) were collected and purified from the infection of irradiated Vero cell monolayers in DMEM (Sigma-Aldrich) supplemented with 5% fetal bovine serum (FBS) at 37 °C in 5% CO₂ (ref. 63). *Legionella pneumophila*. Wild-type (WT) Lp and isogenic flagellin-deficient mutants (*flaA*[−]) of *L. pneumophila* Lp02 (thymidine auxotrophic derivative from serogroup 1 strain Lp01)⁶⁴ were used for co-infection assays. For the surrogate expression of *C. burnetii* effectors, WT Lp, *flaA*[−] and *dotA*[−] isogenic mutants from Lp01 (ref. 64) and *flaA*[−] isogenic mutants of Lp02 (ref. 64) and JR32 (ref. 65) strains were transformed as described below. *L. pneumophila* was cultivated at 37 °C on MOPS-buffered charcoal-yeast extract (CYE) agar plates (1% (w/v) yeast extract, 1% (w/v) MOPS, 3.3 mM L-cysteine, 0.33 mM Fe(NO₃)₃, 1.5% (w/v) Bacto agar and 0.2% (w/v) activated charcoal, pH 6.9) supplemented with 100 μ g ml^{−1} thymidine, as appropriate. *E. coli* (DH5 α strain) was streaked at 37 °C on Luria-Bertani (LB) agar plates without

antibiotics for colony isolation and cultivated in LB broth at 37 °C under rotation (200 r.p.m.).

Mice. C57BL/6, 129S6/SvEv and C57BL/6-derived *Asc*^{-/-}, *Casp11*^{-/-}, *Casp11*^{11-/-}, *Casp1*^{-/-}/*Casp11*^{Tg}, *Nlr4*^{-/-} and *Nlrp3*^{-/-} mice were bred and maintained under specific pathogen-free conditions in the Animal Facilities of the Medical School Ribeirão Preto (FMRP-USP). *Casp1*^{-/-}/*Casp11*^{Tg} mice are *Casp11*^{11-/-} mice expressing a transgene encoding a functional copy of the caspase-11 allele, as described previously⁵³. Male or female mice with 8–12 weeks old were used. All mouse experiments were conducted according to the guidelines of the institutional committee for animal care at the Comissão de Ética em Experimentação Animal da Faculdade de Medicina de Ribeirão Preto, FMRP-USP.

Plasmid construction and expression of Dot/Icm effectors of *C. burnetii* in *L. pneumophila*. The pSN85/Cm^r plasmid with a N-terminal Flag epitope tag was used to clone Dot/Icm effectors from *C. burnetii* (strain Nine Mile RSA 493 phase II). Full-length genes of effectors annotated as CBU were subcloned from pEC33 constructs⁴⁴ through *Bam*HI and *Pst*I sites. Full-length genes of effectors annotated as Ank were amplified from purified genomic DNA⁴⁴ by PCR with sequence-specific primers containing *Bam*HI and *Sph*I sites or *Xba*I and *Sph*I in the case of AnkB (all primers used in this study are provided in Supplementary Table 1). For AnkJ, genomic DNA from the Dugway isolate was amplified by PCR with sequence-specific primers containing *Bam*HI and *Sph*I sites. Following heat-shock transformation and amplification in *E. coli* (DH5 α strain), the constructs were purified with a NucleoSpin plasmid kit (Macherey-Nagel) and subsequently used to transform *L. pneumophila* by electroporation. Transformed *L. pneumophila* clones were selected and expanded in CYE plates with 10 μ g ml⁻¹ chloramphenicol. Effector expression was induced with the addition of 1 mM isopropyl β -D-1-thiogalactopyranoside (IPTG, Invitrogen) to plates and confirmed by immunoblot using an anti-Flag antibody (Sigma-Aldrich).

Construction of *icaA*⁻ mutants of *C. burnetii*. A PCR product consisting of CBU1823 and 2 kb of DNA flanking either end was amplified from *C. burnetii* (Nine Mile RSA 493 phase II) genomic DNA by using PfuUltra II Fusion HotStart DNA polymerase (Agilent Technologies). The product was ligated into the *Bam*HI and *Sa*II sites of pJC-CAT. The resulting construct, pJC-1823FL, was then used as a template in a PCR. The resulting PCR product was then gel purified and digested with *Not*I-HF (New England Biolabs) and ligated with the kanamycin resistance gene amplified from pJB-Kan. The ligation mix was then transformed into DH5 α . The resulting construct, pJC1823KO, in which the CBU1823 gene was replaced with a kanamycin resistance gene, was then used in the gene knockout experiment⁶⁶. CBU1823 knockout was confirmed by Southern hybridization using probes directly binding to CBU1823 and the kanamycin resistance cassette.

Bone marrow-derived and alveolar macrophages. AMs were obtained by bronchoalveolar lavage with warm PBS containing 5 mM EDTA. AMs were washed twice with PBS, seeded on tissue culture plates with RPMI 1640 media 10% FBS, 2 mM L-glutamine and cultivated at 36 °C, 5% CO₂. *C. burnetii* replication in AMs was assessed by quantitative PCR, as described previously⁶⁷. Bone marrow cells were collected from femurs and differentiated with RPMI 1640 (Sigma-Aldrich), 20% FBS (Invitrogen), 30% L-929 cell-conditioned media (LCCM), 2 mM L-glutamine (Sigma-Aldrich) and 100 U ml⁻¹ penicillin–streptomycin (Sigma-Aldrich) cultivated at 36 °C, 5% CO₂. BMDMs were collected and seeded on tissue culture plates and kept in RPMI 1640 media 10% FBS, 5% LCCM and 2 mM L-glutamine⁶⁸.

Construction of retroviral expression vectors and transduction of BMDMs. Full-length genomic C57BL/6 *Casp11* was amplified by PCR with sequence-specific primers containing *Not*I and *Xho*I sites for cloning into pCDNA3.1/Hygro⁺ vector (Invitrogen). *Casp11* was subsequently subcloned into a pMSCV2.0 murine-specific retroviral vector after the digestion of the purified pCDNA3 construct with *Not*I and *Sa*II. pCL vector system⁶⁹ for packaging retroviruses was co-transfected with pMSCV–*Casp11*–IRES–GFP using polyethylenimine in monolayers of Peak cells (ATCC) maintained in RPMI 1640 with 10% FBS at 36 °C and 5% CO₂. BMDMs in day 3 of differentiation were collected, spun down and resuspended in retrovirus-containing peak cell supernatant supplemented with RPMI 1640 containing 20% FBS and 25% LCCM. Cell transduction was confirmed by fluorescence microscopy. BMDMs were then collected, seeded on tissue culture plates one day before infection and kept in RPMI 1640 media 10% FBS and 5% LCCM.

Infection conditions. *C. burnetii* from frozen stocks was thawed at 37 °C. *L. pneumophila* single colonies were seeded and grown for 2 days in CYE agar plates, collected, diluted in distilled water and quantified by measuring optical density (OD) at 600 nm. *L. pneumophila* mutants expressing *C. burnetii* effectors were infected in the presence of 1 mM IPTG. *E. coli* single colonies were grown in LB broth for 8 h and quantified by measuring OD at 600 nm. *E. coli*-infected

macrophages were washed twice with warm PBS one hour after infection, and the media were replaced with media containing gentamicin (Sigma-Aldrich, 50 μ g ml⁻¹). When necessary, the plates were centrifuged for 15 min at 300 g to ensure the comparable adhesion of the bacteria to the cells.

Extraction of *C. burnetii* LPS. LPS was extracted using a modified hot phenol method, as described previously⁷⁰. Briefly, 10⁶ *C. burnetii* were suspended in 1 ml of 50% phenol (pH 8.8), boiled for 10 min, incubated for 5 min on ice and then centrifuged at 14,000 g for 5 min. The aqueous phase was collected, and the extraction was repeated on the pellet. The aqueous phases from both extractions were pooled and vacuum dried for 17 h. The pellet was dissolved in 100 μ l of ultra-pure distilled H₂O. The LPS content after extraction was estimated by a phenol-sulfuric acid method of carbohydrate quantification. Approximately, 1 μ g ml⁻¹ of *C. burnetii* LPS were used to transfect BMDMs using DOTAP (Roche) according to the manufacturer's instructions.

Determination of *L. pneumophila* number per *Legionella*-containing vacuole (LCV). BMDMs were cultivated in coverslips and fixed with 4% PFA after infection. The coverslips were immunostained with rabbit anti-*Legionella* (1:2,000) and goat anti-rabbit IgG Alexa Fluor 488 (Molecular Probes), mounted in glass slides with ProLong Gold containing DAPI (Molecular Probes). The coverslips were analysed with a fluorescence microscope (Leica, Germany), and the number of bacteria per LCV was estimated visually.

Immunoblot analysis. BMDMs were lysed with a RIPA buffer (10 mM Tris-HCl (pH 7.4), 1 mM EDTA, 150 mM NaCl, 1% Nonidet P-40, 1% (w/v) sodium deoxycholate and 0.1% (w/v) SDS) supplemented with a protease inhibitor cocktail. Precleared lysates and supernatants were boiled with Laemmli buffer, resolved by SDS–polyacrylamide gel electrophoresis and transferred to a 0.22- μ m nitrocellulose membrane (GE Healthcare). Rat anti-caspase-1 p20 monoclonal antibody clone 4B4 (Genentech, 1:500), mouse anti-caspase-1 p10 (Santa Cruz Biotechnology, 1:250), goat anti-IL-1 β p17 subunit (Sigma-Aldrich, 1:250), rat anti-caspase-11 p10 (Abcam, 1:500) and rabbit anti-beta actin (Sigma-Aldrich, 1:5,000) were used for antigen detection. To detect the expression of Flag-fused effectors by *L. pneumophila*, bacterial heavy patches streaked from CYE plates after 2 days of growth in the presence of 1 mM IPTG were boiled in Laemmli buffer, immunoblotting was carried out as described above, and anti-Flag M2 (Sigma-Aldrich, 1:5,000) antibody was used for antigen detection.

Reagents for cell stimulation and cytokine determination. Ultra-pure LPS (*E. coli* K12) and poli I:C were from Invivogen. Tumour-necrosis factor- α was from eBioscience. ATP, nigericin and CTB were from Sigma-Aldrich. Cytokines from supernatants of the BMDM culture were detected by ELISA (OptEIA, BD Biosciences), according to the manufacturer's instructions.

Membrane pore formation assay. Pore formation in BMDMs and AMs was quantified by the permeability to ethidium bromide in damaged cells⁵⁷. Images were acquired with a fluorescence microscope (Leica, Germany), five fields per coverslip. Pore formation was quantified using ImageJ software (NIH). For the kinetics of pore formation, the uptake of PI in infected cells was evaluated¹³. BMDMs were seeded in a black, clear-bottom 96-well plate. Before infection or stimulation, BMDMs were washed with warm PBS, and the media were replenished with RPMI 10% without phenol red, 0.038 g ml⁻¹ NaHCO₃, 6 μ g ml⁻¹ PL. Infected BMDMs were kept at 37 °C, and PI was excited at 538 nm. The fluorescence emission was read at 617 nm at every 5 min using a plate fluorometer (FlexStation 3, Molecular Devices).

FLICA staining and flow cytometry analysis. BMDMs were stained with a caspase-1 FLICA probe (FAM-YVAD-FMK), as recommended by the manufacturer. HEK293 reporter cells were likewise stained with a pan-caspase FLICA probe (carboxyfluorescein-VAD-fluoromethylketone, FAM-VAD-FMK; ImmunoChemistry Technologies). Data from 30,000 events were acquired on a FACSCanto II (BD Biosciences) and were analysed with FlowJo software (Tree Star).

Active caspase-11 pull-down assay. The media of primed BMDMs were replenished with fresh media containing 20 μ M biotin-VAD-FMK (Enzo) 15 min before infection. Infected BMDMs were lysed in RIPA buffer (10 mM Tris-HCl (pH 7.4), 1 mM EDTA, 150 mM NaCl, 1% Nonidet P-40, 1% (w/v) sodium deoxycholate and 0.1% (w/v) SDS) supplemented with a protease inhibitor cocktail (Roche). Cleared lysates were equalized according to total protein content, incubated overnight with streptavidin–agarose beads (Novex) and thoroughly rinsed with RIPA buffer. Bound proteins were eluted by re-suspension in Laemmli sample buffer, boiled for 5 min and separated by SDS–polyacrylamide gel electrophoresis.

ARTICLE

NATURE COMMUNICATIONS | DOI: 10.1038/ncomms10205

Statistical analysis. The data were plotted and analysed using GraphPad Prism 5.0 software. The statistical significance was calculated using Student's *t*-test or analysis of variance (ANOVA). Differences were considered statistically significant when the *P* value was <0.05.

References

- Janeway, Jr C. A. Approaching the asymptote? Evolution and revolution in immunology. *Cold Spring Harb. Symp. Quant. Biol.* **54**(Pt 1): 1–13 (1989).
- Mariathasan, S. & Monack, D. M. Inflammasome adaptors and sensors: intracellular regulators of infection and inflammation. *Nat. Rev. Immunol.* **7**, 31–40 (2007).
- Martinon, F., Burns, K. & Tschopp, J. The inflammasome: a molecular platform triggering activation of inflammatory caspases and processing of proIL- β . *Mol. Cell* **10**, 417–426 (2002).
- Miao, E. A. *et al.* Cytoplasmic flagellin activates caspase-1 and secretion of interleukin 1 β via Ipaf. *Nat. Immunol.* **7**, 569–575 (2006).
- Miao, E. A., Ernst, R. K., Dors, M., Mao, D. P. & Aderem, A. *Pseudomonas aeruginosa* activates caspase 1 through Ipaf. *Proc. Natl Acad. Sci. USA* **105**, 2562–2567 (2008).
- Miao, E. A. *et al.* Innate immune detection of the type III secretion apparatus through the NLRC4 inflammasome. *Proc. Natl Acad. Sci. USA* **107**, 3076–3080 (2010).
- Kofoed, E. M. & Vance, R. E. Innate immune recognition of bacterial ligands by NALPs determines inflammasome specificity. *Nature* **477**, 592–595 (2011).
- Franchi, L. *et al.* Critical role for Ipaf in *Pseudomonas aeruginosa*-induced caspase-1 activation. *Eur. J. Immunol.* **37**, 3030–3039 (2007).
- Amer, A. *et al.* Regulation of *Legionella* phagosome maturation and infection through flagellin and host Ipaf. *J. Biol. Chem.* **281**, 35217–35223 (2006).
- Ren, T., Zamboni, D. S., Roy, C. R., Dietrich, W. F. & Vance, R. E. Flagellin-deficient *Legionella* mutants evade caspase-1- and Naip5-mediated macrophage immunity. *PLoS Pathogens* **2**, e18 (2006).
- Zamboni, D. S. *et al.* The Birc1c cytosolic pattern-recognition receptor contributes to the detection and control of *Legionella pneumophila* infection. *Nat. Immunol.* **7**, 318–325 (2006).
- Aachoui, Y. *et al.* Caspase-11 protects against bacteria that escape the vacuole. *Science* **339**, 975–978 (2013).
- Case, C. L. *et al.* Caspase-11 stimulates rapid flagellin-independent pyroptosis in response to *Legionella pneumophila*. *Proc. Natl Acad. Sci. USA* **110**, 1851–1856 (2013).
- Casson, C. N. *et al.* Caspase-11 activation in response to bacterial secretion systems that access the host cytosol. *PLoS Pathogens* **9**, e1003400 (2013).
- Gurung, P. *et al.* Toll or interleukin-1 receptor (TIR) domain-containing adaptor inducing interferon- β (TRIF)-mediated caspase-11 protease production integrates Toll-like receptor 4 (TLR4) protein- and Nlrp3 inflammasome-mediated host defense against enteropathogens. *J. Biol. Chem.* **287**, 34474–34483 (2012).
- Hagar, J. A., Powell, D. A., Aachoui, Y., Ernst, R. K. & Miao, E. A. Cytoplasmic LPS activates caspase-11: implications in TLR4-independent endotoxin shock. *Science* **341**, 1250–1253 (2013).
- Kayagaki, N. *et al.* Noncanonical inflammasome activation by intracellular LPS independent of TLR4. *Science* **341**, 1246–1249 (2013).
- Rathinam, V. A. *et al.* TRIF licenses caspase-11-dependent NLRP3 inflammasome activation by gram-negative bacteria. *Cell* **150**, 606–619 (2012).
- Cunha, L. D. & Zamboni, D. S. Subversion of inflammasome activation and pyroptosis by pathogenic bacteria. *Front. Cell. Infect. Microbiol.* **3**, 76 (2013).
- Nagai, H. & Roy, C. R. Show me the substrates: modulation of host cell function by type IV secretion systems. *Cell. Microbiol.* **5**, 373–383 (2003).
- Galan, J. E. & Wolf-Watz, H. Protein delivery into eukaryotic cells by type III secretion machines. *Nature* **444**, 567–573 (2006).
- Baxt, L. A., Garza-Mayers, A. C. & Goldberg, M. B. Bacterial subversion of host innate immune pathways. *Science* **340**, 697–701 (2013).
- Jorgensen, I. *et al.* The *Chlamydia* protease CPAF regulates host and bacterial proteins to maintain pathogen vacuole integrity and promote virulence. *Cell Host Microbe* **10**, 21–32 (2011).
- Danelishvili, L., Everman, J. L., McNamara, M. J. & Bermudez, L. E. Inhibition of the plasma-membrane-associated serine protease cathepsin G by *Mycobacterium tuberculosis* Rv3364c suppresses caspase-1 and pyroptosis in macrophages. *Front. Microbiol.* **2**, 281 (2011).
- Shah, S. *et al.* Cutting edge: *Mycobacterium tuberculosis* but not nonvirulent mycobacteria inhibits IFN- β and AIM2 inflammasome-dependent IL-1 β production via its ESX-1 secretion system. *J. Immunol.* **191**, 3514–3518 (2013).
- Raoult, D., Marrie, T. & Mege, J. Natural history and pathophysiology of Q fever. *Lancet Infect. Dis.* **5**, 219–226 (2005).
- van Schaik, E. J., Chen, C., Mertens, K., Weber, M. M. & Samuel, J. E. Molecular pathogenesis of the obligate intracellular bacterium *Coxiella burnetii*. *Nat. Rev. Microbiol.* **11**, 561–573 (2013).
- Campoy, E. M., Zoppino, F. C. & Colombo, M. I. The early secretory pathway contributes to the growth of the *Coxiella*-replicative niche. *Infect. Immun.* **79**, 402–413 (2011).
- Klingenbeck, L., Eckart, R. A., Berens, C. & Luhrmann, A. The *Coxiella burnetii* type IV secretion system substrate Caeb inhibits intrinsic apoptosis at the mitochondrial level. *Cell. Microbiol.* **15**, 675–687 (2012).
- Larson, C. L., Beare, P. A., Howe, D. & Heinzen, R. A. *Coxiella burnetii* effector protein subverts clathrin-mediated vesicular trafficking for pathogen vacuole biogenesis. *Proc. Natl Acad. Sci. USA* **110**, E4770–E4779 (2013).
- Luhrmann, A., Nogueira, C. V., Carey, K. L. & Roy, C. R. Inhibition of pathogen-induced apoptosis by a *Coxiella burnetii* type IV effector protein. *Proc. Natl Acad. Sci. USA* **107**, 18997–19001 (2010).
- Luhrmann, A. & Roy, C. R. *Coxiella burnetii* inhibits activation of host cell apoptosis through a mechanism that involves preventing cytochrome c release from mitochondria. *Infect. Immun.* **75**, 5282–5289 (2007).
- McDonough, J. A. *et al.* Host pathways important for *Coxiella burnetii* infection revealed by genome-wide RNA interference screening. *MBio* **4**, e00606–e00612 (2013).
- Voth, D. E. & Heinzen, R. A. Sustained activation of Akt and Erk1/2 is required for *Coxiella burnetii* antiapoptotic activity. *Infect. Immun.* **77**, 205–213 (2009).
- Voth, D. E., Howe, D. & Heinzen, R. A. *Coxiella burnetii* inhibits apoptosis in human THP-1 cells and monkey primary alveolar macrophages. *Infect. Immun.* **75**, 4263–4271 (2007).
- Vazquez, C. L. & Colombo, M. I. *Coxiella burnetii* modulates Beclin 1 and Bcl-2, preventing host cell apoptosis to generate a persistent bacterial infection. *Cell Death Differ.* **17**, 421–438 (2010).
- Shannon, J. G., Howe, D. & Heinzen, R. A. Virulent *Coxiella burnetii* does not activate human dendritic cells: role of lipopolysaccharide as a shielding molecule. *Proc. Natl Acad. Sci. USA* **102**, 8722–8727 (2005).
- Zamboni, D. S. Genetic control of natural resistance of mouse macrophages to *Coxiella burnetii* infection in vitro: macrophages from restrictive strains control parasitophorous vacuole maturation. *Infect. Immun.* **72**, 2395–2399 (2004).
- Zamboni, D. S. *et al.* Stimulation of toll-like receptor 2 by *Coxiella burnetii* is required for macrophage production of pro-inflammatory cytokines and resistance to infection. *J. Biol. Chem.* **279**, 54405–54415 (2004).
- Seshadri, R. *et al.* Complete genome sequence of the Q-fever pathogen *Coxiella burnetii*. *Proc. Natl Acad. Sci. USA* **100**, 5455–5460 (2003).
- Zamboni, D. S., McGrath, S., Rabinovitch, M. & Roy, C. R. *Coxiella burnetii* express type IV secretion system proteins that function similarly to components of the *Legionella pneumophila* Dot/Icm system. *Mol. Microbiol.* **49**, 965–976 (2003).
- Zusman, T., Yerushalmi, G. & Segal, G. Functional similarities between the icm/dot pathogenesis systems of *Coxiella burnetii* and *Legionella pneumophila*. *Infect. Immun.* **71**, 3714–3723 (2003).
- Beare, P. A. *et al.* Comparative genomics reveal extensive transposon-mediated genomic plasticity and diversity among potential effector proteins within the genus *Coxiella*. *Infect. Immun.* **77**, 642–656 (2009).
- Carey, K. L., Newton, H. J., Luhrmann, A. & Roy, C. R. The *Coxiella burnetii* Dot/Icm system delivers a unique repertoire of type IV effectors into host cells and is required for intracellular replication. *PLoS Pathogens* **7**, e1002056 (2011).
- Chen, C. *et al.* Large-scale identification and translocation of type IV secretion substrates by *Coxiella burnetii*. *Proc. Natl Acad. Sci. USA* **107**, 21755–21760 (2010).
- Voth, D. E. *et al.* The *Coxiella burnetii* cryptic plasmid is enriched in genes encoding type IV secretion system substrates. *J. Bacteriol.* **193**, 1493–1503 (2011).
- Voth, D. E. *et al.* The *Coxiella burnetii* ankyrin repeat domain-containing protein family is heterogeneous, with C-terminal truncations that influence Dot/Icm-mediated secretion. *J. Bacteriol.* **191**, 4232–4242 (2009).
- Weber, M. M. *et al.* Identification of *Coxiella burnetii* type IV secretion substrates required for intracellular replication and *Coxiella*-containing vacuole formation. *J. Bacteriol.* **195**, 3914–3924 (2013).
- Lifshitz, Z. *et al.* Computational modeling and experimental validation of the *Legionella* and *Coxiella* virulence-related type-IVB secretion signal. *Proc. Natl Acad. Sci. USA* **110**, E707–E715 (2013).
- Pan, X., Luhrmann, A., Satoh, A., Laskowski-Arce, M. A. & Roy, C. R. Ankyrin repeat proteins comprise a diverse family of bacterial type IV effectors. *Science* **320**, 1651–1654 (2008).
- Sauer, J. D. *et al.* Specificity of *Legionella pneumophila* and *Coxiella burnetii* vacuoles and versatility of *Legionella pneumophila* revealed by coinfection. *Infect. Immun.* **73**, 4494–4504 (2005).
- Case, C. L., Shin, S. & Roy, C. R. Asc and Ipaf Inflammasomes direct distinct pathways for caspase-1 activation in response to *Legionella pneumophila*. *Infect. Immun.* **77**, 1981–1991 (2009).
- Kayagaki, N. *et al.* Non-canonical inflammasome activation targets caspase-11. *Nature* **479**, 117–121 (2011).

54. Cerqueira, D. M., Pereira, M. S., Silva, A. L., Cunha, L. D. & Zamboni, D. S. Caspase-1 but not caspase-11 is required for NLR4-mediated pyroptosis and restriction of infection by flagellated *Legionella* species in mouse macrophages and *in vivo*. *J. Immunol.* **195**, 2303–2311 (2015).
55. Newton, H. J., McDonough, J. A. & Roy, C. R. Effector protein translocation by the *Coxiella burnetii* Dot/Icm type IV secretion system requires endocytic maturation of the pathogen-occupied vacuole. *PLoS ONE* **8**, e54566 (2013).
56. Molofsky, A. B. *et al.* Cytosolic recognition of flagellin by mouse macrophages restricts *Legionella pneumophila* infection. *J. Exp. Med.* **203**, 1093–1104 (2006).
57. Silveira, T. N. & Zamboni, D. S. Pore formation triggered by *Legionella* spp. is an Nlr4 inflammasome-dependent host cell response that precedes pyroptosis. *Infect. Immun.* **78**, 1403–1413 (2010).
58. Brooke, R. J., Kretschmar, M. E., Mutters, N. T. & Teunis, P. F. Human dose response relation for airborne exposure to *Coxiella burnetii*. *BMC Infect. Dis.* **13**, 488 (2013).
59. Beare, P. A. *et al.* Dot/Icm type IVB secretion system requirements for *Coxiella burnetii* growth in human macrophages. *MBio* **2**, e00175–e00211 (2011).
60. Broz, P. *et al.* Caspase-11 increases susceptibility to *Salmonella* infection in the absence of caspase-1. *Nature* **490**, 288–291 (2012).
61. Graham, J. G. *et al.* Virulent *Coxiella burnetii* pathotypes productively infect primary human alveolar macrophages. *Cell. Microbiol.* **15**, 1012–1025 (2013).
62. Kobayashi, T. *et al.* The *Shigella* OspC3 effector inhibits caspase-4, antagonizes inflammatory cell death, and promotes epithelial infection. *Cell Host Microbe* **13**, 570–583 (2013).
63. Zamboni, D. S., Mortara, R. A. & Rabinovitch, M. Infection of Vero cells with *Coxiella burnetii* phase II: relative intracellular bacterial load and distribution estimated by confocal laser scanning microscopy and morphometry. *J. Microbiol. Methods* **43**, 223–232 (2001).
64. Berger, K. H. & Isberg, R. R. Two distinct defects in intracellular growth complemented by a single genetic locus in *Legionella pneumophila*. *Mol. Microbiol.* **7**, 7–19 (1993).
65. Sadosky, A. B., Wiater, L. A. & Shuman, H. A. Identification of *Legionella pneumophila* genes required for growth within and killing of human macrophages. *Infect. Immun.* **61**, 5361–5373 (1993).
66. Beare, P. A. & Heinzen, R. A. Gene inactivation in *Coxiella burnetii*. *Methods Mol. Biol.* **1197**, 329–345 (2014).
67. Brennan, R. E. & Samuel, J. E. Evaluation of *Coxiella burnetii* antibiotic susceptibilities by real-time PCR assay. *J. Clin. Microbiol.* **41**, 1869–1874 (2003).
68. Marim, F. M., Silveira, T. N., Lima, Jr D. S. & Zamboni, D. S. A method for generation of bone marrow-derived macrophages from cryopreserved mouse bone marrow cells. *PLoS ONE* **5**, e15263 (2010).
69. Naviaux, R. K., Costanzi, E., Haas, M. & Verma, I. M. The pCL vector system: rapid production of helper-free, high-titer, recombinant retroviruses. *J. Virol.* **70**, 5701–5705 (1996).
70. Beare, P. A., Howe, D., Cockrell, D. C. & Heinzen, R. A. Efficient method of cloning the obligate intracellular bacterium *Coxiella burnetii*. *Appl. Environ. Microbiol.* **73**, 4048–4054 (2007).

Acknowledgements

This paper is dedicated to Dr Michel Rabinovitch, whose passion for biology and scientific investigation co-infects his students forever. We thank Maira Nakamura for her technical assistance, to Richard Flavell (Yale, USA) for providing us with *Nlr4*^{−/−} and *Casp1/11*^{−/−} mice and Vishva Dixit (Genentech, USA) for providing us with *Nlrp3*^{−/−}, *Casp1*^{−/−}/*Casp11*^{1g} and *Casp11*^{−/−} and the antibody to caspase-1 p20. This work was supported by grants from Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP, grants 2013/08216-2, 2014/50268-2 and 2014/04684-4), Conselho Nacional do Desenvolvimento Científico e Tecnológico (CNPq and INCTV/CNPq), Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), Núcleo de Apoio à Pesquisa em Doenças Inflamatórias (NAPDIN, grant 11.1.21625.01.0) and the PEW Program in Biomedical Sciences. L.D.C. and J.M.R. are supported by fellowships from FAPESP. L.M.M. and T.D.F. are supported by fellowships from CAPES. H.J.N. is a visiting professor from Science Without Borders Program (CNPq, grant 401577/2014-7). D.S.Z. is a research fellow from CNPq, Brazil.

Author contributions

L.D.C., T.D.F., H.J.N., C.R.R. and D.S.Z. designed the experiments; L.D.C., J.M.R., T.D.F., L.M.M., C.A.K., J.H.M. and H.J.N. performed the experiments; L.D.C., J.M.R., T.D.F. and D.S.Z. processed and analysed the data; L.D.C. and D.S.Z. wrote the manuscript; and all authors commented on the manuscript.

Additional information

Supplementary Information accompanies this paper at <http://www.nature.com/naturecommunications>

Competing financial interests: The authors declare no competing financial interests.

Reprints and permission information is available online at <http://npg.nature.com/reprintsandpermissions/>

How to cite this article: Cunha, L. D. *et al.* Inhibition of inflammasome activation by *Coxiella burnetii* type IV secretion system effector IcaA. *Nat. Commun.* **6**:10205 doi: 10.1038/ncomms10205 (2015).

This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

Supplementary Information

Supplementary Figure 1. Viable *C. burnetii* is required for inhibition caspase-1 activation induced by *L. pneumophila*. Bone marrow-derived macrophages (BMDM) were left uninfected (NI) or were infected with PFA-fixed (PFA Cb) or live *C. burnetii* (Live Cb) at MOI 30 for 24 h and further infected with wild-type *L. pneumophila* (WT Lp, MOI 10) or *flaA*⁻ mutant (*flaA*⁻) for 9 h. Co-infected BMDM are indicated (Cb + WT Lp or Cb + *flaA*⁻). **(a)** Immunoblot showing levels of processed p20 subunit of caspase-1 (caspase-1 p20) and unprocessed caspase-1 (pro-caspase-1), as determined in supernatant (SN) and cell extract (CE). **(b)** Similar in **(a)** for BL/6 BMDM.

Supplementary Figure 2. Co-infection with *C. burnetii* does not interfere with *flaA*- *L. pneumophila* replication in BMDM. Bone marrow-derived macrophages (BMDM) from C57BL/6 mice were left uninfected or were infected with *C. burnetii* at MOI 30 for 24 h and further infected with *flaA*- *L. pneumophila* strain Lp01 (*flaA*⁻, MOI 10) for 9 h. Co-infected BMDM are indicated (*Cb* + *flaA*⁻). An estimation of the frequency of bacteria per *Legionella*-containing vacuole (LCV), as assessed by counting on the fluorescence microscopy, is shown.

Supplementary Figure 3. *C. burnetii* does not interfere with ATP-mediated activation of the canonical NLRP3 inflammasome or with dsDNA-mediated activation of the AIM2 inflammasome. Bone marrow-derived macrophages (BMDM) were left uninfected (NI) or were infected with *C. burnetii* (*Cb*, MOI 30) for 24 h and further stimulated with LPS (1 $\mu\text{g ml}^{-1}$) for 4 h. The cells were then treated with either ATP (5 mM) for 20 min or nigericin (20 mM) for 40 min or were transfected with 1 μg of dsDNA for 6 h. (**a**, **b**) Immunoblot showing levels of processed p20 subunit of caspase-1 (caspase-1 p20) and unprocessed caspase-1 (pro-caspase-1), as determined in supernatant (SN) and cell extract (CE) of BL/6 BMDM. (**c**, **d**) IL-1 β secretion in the supernatant was estimated by ELISA.

Supplementary Figure 4. *C. burnetii* inhibits pore formation in response to *flaA*⁻ *L. pneumophila* in alveolar macrophages. (a) Alveolar macrophages (AM) from C57BL/6 mice were infected with *C. burnetii* (MOI 3), and bacterial growth was assessed by qPCR. Relative bacterial replication was calculated as $\log(Y_{x=n}) - \log(Y_{x=0})$. Control AM were treated with bacterial translation inhibitor rifampicin (RIF, 20 mg mL⁻¹). Data are expressed as the average $\pm$ SEM of triplicate wells and significance was calculated with ANOVA. *, $P < 0.05$. (b,c) AM were left uninfected (NI) or were infected with heat-killed (HK Cb) or live *C. burnetii* (Live Cb) at MOI 30 for 6 h and further infected with *flaA*⁻ *L. pneumophila* (*flaA*⁻, MOI 10) for 3 h. (b) Fluorescence micrographs revealing cell permeability, as assessed by the quantification of EtBr influx (red dots) in acridine orange (green) stained AM. Scale bar corresponds to 75 μ m. (c) Quantification of the experiment shown in (b).

Supplementary Figure 5. Identification of CBU1823 (IcaA) as an effector protein that inhibits a flagellin-independent pathway for caspase-1 activation induced by *L. pneumophila*. Bone marrow-derived macrophages (BMDM) from C57BL/6 mice were infected with *flaA*⁻ *L. pneumophila* expressing each indicated effector (indicated is the CBU number or the protein name) or an empty vector (pVec) for 9 h in the presence of 1 mM IPTG. **(a, b)** Mutants were grown in CYE agar plates for two days, and effector expression was assessed by immunoblot for the Flag epitope tag (expression of N-terminal Flag fusions). **(c, d)** Immunoblot showing levels of processed p10 subunit of caspase-1 (caspase-1 p10) and unprocessed caspase-1 (pro-caspase-1) in supernatant. **(e)** Immunoblot showing levels of p17 subunit of mature IL-1β (IL-1β p17) and pro-IL-1β in supernatant fraction. CBU1823 (IcaA) is highlighted in red.

Supplementary Figure 6. Expression of IcaA by *L. pneumophila* interferes with caspase-1 activation but does not influence the replication in axenic media, macrophage infection or the induction of IL-12p40. *flaA*⁻ or *dotA*⁻ mutants from three different *L. pneumophila* strains: Lp01, thymidine auxotroph Lp02 and JR32, transformed with the empty vector (pVec), pSN85 encoding IcaA (pIcaA) or pSN85 encoding Ankh (pAnkH) were evaluated. (a) Immunoblot showing levels of processed p20 subunit of caspase-1 (caspase-1 p20) and unprocessed caspase-1 (pro-Casp-1), as determined in supernatant (SN) and cell extract (CE) of bone marrow-derived macrophages (BMDM) infected for 9 h. (b) Replication of *flaA*⁻/pVec and *flaA*⁻/pIcaA (strain Lp01) in AYE broth was assessed by optical density (OD) at 600 nm. (c) BL/6 BMDM were infected with *flaA*⁻/pVec and *flaA*⁻/pIcaA (strain Lp01) for 1 h, and bacterial internalization was assessed by fluorescence microscopy using anti-*L. pneumophila*. (d) IL-12p40 secretion in the supernatant of BMDM infected for 9 h with *flaA*⁻/pVec and *flaA*⁻/pIcaA (strain Lp02), as estimated by ELISA. (e, f) Estimation of the frequency of bacteria per *Legionella*-containing vacuole (LCV) of BMDM infected with *flaA*⁻/pVec and *flaA*⁻/pIcaA and *flaA*⁻/pAnkH (strain Lp01), as assessed by counting on fluorescence microscopy. (g) IL-1 β secretion in the supernatant of BMDM infected for 9 h with *flaA*⁻/pVec and *flaA*⁻/pIcaA (strain Lp02), as estimated by ELISA. Data are expressed as the average $\pm$ SEM of triplicate wells and significance was calculated with ANOVA. *, $P < 0.05$.

Supplementary Figure 7. IcaA inhibits caspase-11-dependent pore formation in response to *flaA* *L. pneumophila*. Bone marrow-derived macrophages (BMDM) from C57BL/6 or isogenic mice deficient in caspase-11 or caspase-1/11 were primed with LPS (0.5 mg mL⁻¹) for 3 h. Cells were left uninfected (NI) or were infected with wild-type (**a**) or *flaA* *L. pneumophila* (**b**) transformed with either an empty vector (Lp/pVec) or a vector encoding IcaA (Lp/pIcaA) at MOI 10. Pore formation was assessed fluorometrically in real time by the uptake of propidium iodide (RFUs, relative fluorescence units).

Supplementary Figure 8. *C. burnetii* LPS induces pore formation when transfected in macrophages. Bone marrow-derived macrophages (BMDM) generated from C57BL/6 (**a**), *Casp11*^{-/-} (**b**) or *Casp1/11*^{-/-} (**c**) mice were primed with poly I:C (5 µg mL⁻¹) for 4 h. Cells were transfected with 1 µg mL⁻¹ *C. burnetii* LPS (blue) or with MOCK (black) for up to 12 hrs. Pore formation was assessed fluorometrically in real time by the uptake of propidium iodide (RFUs, relative fluorescence units).

Supplementary Figure 9. Validation of the *icaA* mutant *C. burnetii*. CBU_1823 (*icaA*⁻) knockout was confirmed by southern hybridization using probes to *icaA* and the kanamycin resistance cassette. **(a)** Southern blot with probe against kanamycin resistance cassette. **(b)** Southern blot with probe against *icaA*.

Supplementary Figure 10. Full immunoblots of main figures presented in this study.

Supplementary Table1. List of the primers used in this study

Primer	Sequence 5' to 3'	Purpose
AnkA F	AAg gatccTCTTGCTTAGCTTAATGG	pSN85
AnkA R	AAg catgcTTAAAACAGTCCGGGGCC	pSN85
AnkB F	AA tctagaGTATGTTTAACCAATTGGA	pSN85
AnkB R	GGg catgcTTACATGTGCTTACCCGG	pSN85
AnkF F	AAg gatccTCATGAGACAGCGTGAAA	pSN85
AnkF R	AAg catgcCTACCGCTGGAAGCCGC	pSN85
AnkG F	AAg gatccTCATGAGTAGACGTGAGA	pSN85
AnkG R	AAg catgcTCACCGAGGACTAGACAG	pSN85
AnkH F	AAg gatccTCTTGTCAGAATTAGGAG	pSN85
AnkH R	AAg catgcTTAGATTAACGTGCGCGT	pSN85
AnkI F	AAg gatccTCATGAGAGAATCATCAG	pSN85
AnkI R	AAg catgcCTAAATTCCAAAAGAACC	pSN85
AnkJ F	AAg gatccTCATGGCGGAAATTTACTA	pSN85
AnkJ R	AAg catgcTTACGCAGCGCGCATGGT	pSN85
AnkN F	AAg gatccTCTTGCGAGAGCTTATGA	pSN85
AnkN R	AAg catgcCTATAGACCGTATTGCTG	pSN85
AnkO F	AAg gatccTCATGGAAATAATTTCTT	pSN85
AnkO R	AAg catgcTTAATAACGATTTTTTGTT	pSN85
AnkP F	AAg gatccTCGTGGGACAAAATACAA	pSN85
AnkP R	AAg catgcCTTAAGCTAAGCAAGGGG	pSN85
CBU1823_1 FOR	GACg gatccATTCTAAGGTGGATATGTGGGC	pJC1823-FL
CBU1823_1 REV	GACgtcgacCGAGTTGGTTTTTCATTGCCAAGG	pJC1823-FL
CBU1823_2 FOR	GATgtcgacgcggccgcgcatgcATTTCCCTCCGCAA TAGCTTA	pJC1823-KO
CBU1823_2 REV	GATgcatgcgcggccgcCTTGCCTTTTACCCCCTC GT	pJC1823-KO
Kan FOR	GACgcggccgcAGCTTATGGCTTCGTTTCGCAG	pJC1823-KO
Kan REV	GACgcggccgcTCAGAAGAACTCGTCAAGAAGG CG	pJC1823-KO
CBU1823_SH F	CCATTCAAGAGAGTTCATGTATGCCATCA	Southern hybridization
CBU1823_SH R	ATCTCATCTCCTTCAGCCTGACTT	Southern hybridization
Kan_SH F	GTGGAGAGGCTATTCTGGCTATG	Southern hybridization
Kan_SH R	TTGAGCCTGGCGAACAGTT	Southern hybridization
Cb dotA F1	GCGCAATACGCTCAATCACA	qPCR
Cb dotA R1	CCATGGCCCCAATTCTCTT	qPCR

10.2. Anexo 3 - Artigo publicado (MASCARENHAS, Danielle PA; CERQUEIRA, Daiane M; PEREIRA, Marcelo SF; CASTANHEIRA, Fernanda VS; FERNANDES, Talita D; MANIN, Grazielle Z; CUNHA, Larissa D; ZAMBONI, Dario S. *Pathogens*, 2017)

“Inhibition of caspase-1 or gasdermin-D enable caspase-8 activation in the Naip5/NLRC4/ASC inflammasome”

- Publicado em: agosto de 2017

RESEARCH ARTICLE

Inhibition of caspase-1 or gasdermin-D enable caspase-8 activation in the Naip5/NLRC4/ASC inflammasome

Danielle P. A. Mascarenhas, Daiane M. Cerqueira, Marcelo S. F. Pereira, Fernanda V. S. Castanheira, Talita D. Fernandes, Grazielle Z. Manin, Larissa D. Cunha, Dario S. Zamboni*

Department of Cell Biology, School of Medicine of Ribeirão Preto, University of São Paulo. Ribeirão Preto, Brazil

* dszamboni@fmrp.usp.br

OPEN ACCESS

Citation: Mascarenhas DPA, Cerqueira DM, Pereira MSF, Castanheira FVS, Fernandes TD, Manin GZ, et al. (2017) Inhibition of caspase-1 or gasdermin-D enable caspase-8 activation in the Naip5/NLRC4/ASC inflammasome. PLoS Pathog 13(8): e1006502. <https://doi.org/10.1371/journal.ppat.1006502>

Editor: H. Steven Seifert, Northwestern University Feinberg School of Medicine, UNITED STATES

Received: March 9, 2017

Accepted: June 30, 2017

Published: August 3, 2017

Copyright: © 2017 Mascarenhas et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: This work was supported by grants from Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP, grants 2012/09363-6, 2013/08216-2 and 2014/04684-4), Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) and Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES). DSZ is a

Abstract

Legionella pneumophila is a Gram-negative, flagellated bacterium that survives in phagocytes and causes Legionnaires' disease. Upon infection of mammalian macrophages, cytosolic flagellin triggers the activation of Naip/NLRC4 inflammasome, which culminates in pyroptosis and restriction of bacterial replication. Although NLRC4 and caspase-1 participate in the same inflammasome, *Nlrc4*^{-/-} mice and their macrophages are more permissive to *L. pneumophila* replication compared with *Casp1/11*^{-/-}. This feature supports the existence of a pathway that is NLRC4-dependent and caspase-1/11-independent. Here, we demonstrate that caspase-8 is recruited to the Naip5/NLRC4/ASC inflammasome in response to flagellin-positive bacteria. Accordingly, caspase-8 is activated in *Casp1/11*^{-/-} macrophages in a process dependent on flagellin, Naip5, NLRC4 and ASC. Silencing caspase-8 in *Casp1/11*^{-/-} cells culminated in macrophages that were as susceptible as *Nlrc4*^{-/-} for the restriction of *L. pneumophila* replication. Accordingly, macrophages and mice deficient in *Asc/Casp1/11*^{-/-} were more susceptible than *Casp1/11*^{-/-} and as susceptible as *Nlrc4*^{-/-} for the restriction of infection. Mechanistically, we found that caspase-8 activation triggers gasdermin-D-independent pore formation and cell death. Interestingly, caspase-8 is recruited to the Naip5/NLRC4/ASC inflammasome in wild-type macrophages, but it is only activated when caspase-1 or gasdermin-D is inhibited. Our data suggest that caspase-8 activation in the Naip5/NLRC4/ASC inflammasome enable induction of cell death when caspase-1 or gasdermin-D is suppressed.

Author summary

Legionella pneumophila is the causative agent of Legionnaires' disease, an atypical pneumophila that affects people worldwide. Besides the clinical importance, *L. pneumophila* is a very useful model of pathogenic bacteria for investigation of the interactions of innate immune cells with bacterial pathogens. Studies using *L. pneumophila* demonstrated that Naip5 and NLRC4 activate caspase-1 and this inflammasome is activated by bacterial

research fellow from CNPq, Brazil. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

flagellin. However, macrophages and mice deficient in NLRC4 are more susceptible for *L. pneumophila* replication than those deficient in caspase-1, indicating that the flagellin/Naip5/NLRC4 inflammasome triggers responses that are independent on caspase-1. Here, we used *L. pneumophila* to investigate this novel pathway and found that caspase-8 interacts with NLRC4 in a process that is dependent on ASC and independent of caspase-1 and caspase-11. Although caspase-8 is recruited to the Naip5/NLRC4/ASC inflammasome, it is only activated when caspase-1 or gasdermin-D is inhibited. Our data suggest that caspase-8 activation in the Naip5/NLRC4/ASC inflammasome may favor host responses during infections against pathogens that inhibit components of the pyroptotic cell death including caspase-1 and gasdermin-D.

Introduction

Legionella pneumophila is the causative agent of Legionnaires' disease. It was identified for the first time in 1976, after an atypical pneumonia affected the participants of the American Legion Convention in Philadelphia, United States [1]. After isolation, *L. pneumophila* were characterized as Gram-negative, flagellated, intracellular facultative bacteria [2,3]. The species of *Legionella* were found mainly in freshwater and soil environments, including lakes and irrigation systems [4]. Infection of humans occurs upon inhalation of water droplets derived from these environments containing *Legionella* [5]. After inhalation, *L. pneumophila* can subvert the normal vesicle traffic within alveolar macrophages and form LCV (*Legionella*-containing vacuoles), a process that is mediated by the injection of hundreds of bacterial effectors through a type IV secretion system called Dot/Icm [6–9]. During its evolution, *L. pneumophila* were selected based on their replication in protozoa but not in humans, which are accidental hosts [10]. Consequently, *L. pneumophila* can be recognized by many innate immune receptors in mammalian cells, including proteins from the family of the nucleotide-binding domain and leucine-rich repeat-containing proteins (NLRs). These characteristics make *L. pneumophila* an excellent model for the study of innate immunity, including intracellular signaling pathways and inflammasomes.

The major inflammasome that leads to the restriction of *Legionella* replication in macrophages is Naip5/NLRC4. This pathway was discovered in mouse cells upon observations that macrophages from the A/J mouse strain, but not cells from other mice strains, are susceptible to *L. pneumophila* replication [11]. The resistance was mapped to the *Lgn1* locus, which encodes several copies of *Naip* genes, including *Naip5* (*Birc1e*), which is the gene responsible for resistance [12–16]. Successful lines of investigation culminated in the demonstration that Naip5 recognizes bacterial flagellin and interacts with NLRC4 for caspase-1 activation and the restriction of bacterial replication [17–20]. This platform was named the Naip5/NLRC4 inflammasome and triggers pore formation and pyroptosis, which has been considered one of the most important mechanisms for the restriction of intracellular pathogen replication via inflammasomes [21–24]. Host cell death via pyroptosis eliminates intracellular parasite replication and traps intracellular microbes in pyroptotic cells, facilitating microbial destruction by additional phagocytes [23,25–29]. Pyroptosis occurs concomitantly with the secretion of inflammatory cytokines such as IL-1 β and IL-18, a process that requires the adaptor molecule ASC and the formation of NLRC4/ASC puncta [20,30]. ASC also functions as an adaptor protein for other inflammasomes, including AIM2 and NLRP3, which triggers the processing of caspase-1 and caspase-8 [21,31–34]. Of note, Naip5/NLRC4 appears to be the only inflammasome required for the restriction of *L. pneumophila* replication. Macrophages that are deficient

in NLRP3 or AIM2 can efficiently restrict *L. pneumophila* replication [20,21,23,35]. However, the participation of ASC in the resistance of *L. pneumophila* infection is controversial. In murine macrophages, ASC is dispensable for the induction of pyroptosis and the restriction of bacterial replication [20,21]. By contrast, experiments performed with human monocytes indicate that ASC silencing leads to an increase in bacterial replication [36,37]. Thus, the role of ASC in the restriction of *L. pneumophila* replication is still unclear.

We have previously demonstrated the existence of a pathway that is dependent on flagellin and NLRC4 but independent of caspase-1 [38]. Here, we used macrophages and *Casp1/11*^{-/-} mice to systematically assess this pathway. By searching for additional components that operate in the NLRC4 inflammasome independently of caspase-1/11, we found that caspase-8 interacts with NLRC4 in a process that is dependent on ASC. This pathway effectively accounts for resistance to infection in macrophages and in vivo when caspase-1 is absent. In wild-type cells, caspase-8 is recruited to the Naip5/NLRC4/ASC/caspase-1 inflammasome, but is not activated. Caspase-8 activation in this platform only occurs when caspase-1 or gasdermin-D is inhibited, suggesting that this pathway may be important when pyroptosis is inhibited.

Results

Restriction of *L. pneumophila* replication in BMDMs is flagellin/NLRC4-dependent, ASC-independent and partially caspase-1/11-dependent

We have previously demonstrated that activation of the flagellin/NLRC4 inflammasome triggers caspase-1-dependent and independent responses to restrict *Legionella* replication in macrophages and in mouse lungs [38]. However, the caspase-1-independent mechanisms underlying this pathway are unknown. To further characterize this pathway, we performed growth curves using high and very low multiplicity of infections (MOIs) in bone marrow-derived macrophages (BMDMs). Macrophages were infected with wild-type *L. pneumophila* in the JR32 background (WT Lp) and the isogenic mutants *flaA*⁻ and *fliI*⁻. *FliI* is an ATPase that is required for the secretion of flagellin through the flagellar apparatus [39]. Consequently, *fliI*⁻ mutants express flagellin but are non-motile and non-flagellated, making them an appropriate control for *flaA*⁻ mutants for investigations related to the role of flagellin. We found that BMDMs from C57BL/6 and *Asc*^{-/-} mice fully restrict the replication of WT Lp and *fliI*⁻ bacteria at low and high MOIs. In contrast, *Nlrc4*^{-/-} cells are permissive and *Casp1/11*^{-/-} cells are partially restrictive (S1 Fig). Bacterial mutants for flagellin bypass NLRC4-mediated growth restriction and replicate in all macrophages as previously described [17–19,40]. These data support previous reports showing that ASC is not required for the restriction of *L. pneumophila* replication in the presence of caspase-1/11 [20,21]. In addition, these data further support our previous assertion that flagellin triggers an uncharacterized pathway that is dependent on NLRC4 and independent of caspase-1 and caspase-11 [38]. We decided to use BMDMs from *Casp1/11*^{-/-} mice to further investigate this NLRC4-dependent and caspase-1/11-independent pathway.

Flagellated *L. pneumophila* triggers NLRC4 puncta that associate with caspase-8 in a process that is ASC-dependent

The transduction of BMDMs with a retrovirus encoding NLRC4 fused to GFP allows the visualization of NLRC4 puncta in the cytoplasm of macrophages infected with flagellated bacteria [30]. Here, we used this retroviral system to investigate the formation of the NLRC4 inflammasome in the absence of caspase-1/11. BMDMs from *Casp1/11*^{-/-} mice were transduced with

NLRC4-GFP and infected with WT Lp, *fliI*⁻ and *flaA*⁻ at different MOIs and time points. We found that WT Lp and *fliI* triggered the formation of NLRC4 puncta in the absence of caspase-1/11 (Fig 1A). The formation of NLRC4 puncta was influenced by the MOI and significantly diminished in response to *flaA*⁻ bacteria (Fig 1A). Next, we evaluated the requirement of ASC for the formation of NLRC4 puncta in the absence of caspase-1/11. We constructed a mouse that was deficient in ASC and caspase-1/11 and found that whereas the formation of

Fig 1. *Legionella* triggers the formation of NLRC4/caspase-8 puncta in a process that is dependent on ASC and flagellin and independent of caspase-1/11. Bone marrow-derived macrophages (BMDMs) obtained from *Casp1/11*^{-/-} and *Asc/Casp1/11*^{-/-} mice were transduced with retrovirus encoding NLRC4-GFP. Cells were infected with wild-type *L. pneumophila* (WT Lp), motility-deficient mutants expressing flagellin (*fliI*⁻) or with flagellin-deficient mutants (*flaA*⁻) at a MOI of 1, 3 or 10. (A) After 2, 4 and 8 hours of infection, the cells were fixed, and the percentage of transduced cells containing NLRC4-GFP puncta were determined using an epifluorescence microscope. *, *P* < 0.05 compared with BMDMs infected with *flaA*⁻, Student's *t* test. (B-D) Transduced cells were infected with *fliI*⁻ (MOI 10) and fixed after 8 hours of infection. (B) The percentages of transduced *Casp1/11*^{-/-} and *Asc/Casp1/11*^{-/-} cells containing NLRC4-GFP puncta were determined. (C) The percentages of NLRC4-GFP puncta that colocalized with ASC and caspase-8 were determined. (D) Representative images of a transduced BMDM infected with *fliI*⁻ at a MOI of 10. The cultures were stained with anti-caspase-8 (red) and anti-ASC (purple). The cell nuclei were stained with DAPI (cyan); NLRC4-GFP is shown in green. The images show the colocalization of NLRC4-GFP, ASC and caspase-8 in *Casp1/11*^{-/-} BMDMs. The images were acquired by multiphoton microscopy using a 63x oil immersion objective and analyzed using ImageJ Software. Scale bar, 10μm. Data show the average ± SD of triplicate wells. NI, uninfected. Data are presented for one representative experiment of five (A) and two (B-D) experiments with similar results.

<https://doi.org/10.1371/journal.ppat.1006502.g001>

NLRC4 puncta occurred in the absence of caspase-1/11, ASC was essential for formation of the NLRC4 puncta (Fig 1B). These data are in agreement with previous findings indicating that ASC is critical for the nucleation of several inflammasomes, including AIM2, NLRP3 and NLRC4 [30,32,33,41–46]. Our results confirm that ASC is essential to NLRC4 puncta formation formed in the absence of caspase-1/11. Next, we used this NLRC4-GFP system to identify additional components of the NLRC4 inflammasome that operates in the absence of caspase-1/11. Non-inflammatory caspases have been previously shown to participate in the assembly of inflammasomes and to interact with ASC, including caspase-3, caspase-7 and caspase-8 [32–34,37,44,46–51]. Thus, we transduced *Casp1/11*^{−/−} macrophages with a retrovirus encoding NLRC4-GFP and evaluated the colocalization of NLRC4 with these caspases. In this experiment, we used the pan-caspase inhibitor Z-VAD to block caspase activation and to visualize puncta formation. We did not detect significant numbers of NLRC4 or ASC puncta containing caspase-3 and caspase-7 (S2 Fig). In contrast, caspase-8 and ASC was present in more than 90% of the NLRC4 puncta (Fig 1C and S2 Fig). These data are in agreement with our findings indicating that ASC is required for NLRC4 puncta formation, accordingly, endogenous ASC colocalizes with NLRC4 and caspase-8 in the same puncta (Fig 1D). To evaluate the participation of caspase-8 in the NLRC4 inflammasome, we transduced BMDMs from *Casp1/11*^{−/−} mice with a retrovirus encoding ASC fused to GFP (ASC-GFP) and analyzed ASC puncta colocalization with caspase-8. We found that ASC puncta formed readily after the infection and that this process occurred in response to WT Lp and *fliI* but not *flaA*[−] (Fig 2A). After 8 hours of infection, the formation of ASC puncta was partially dependent on flagellin (Fig 2A). We stained caspase-8 in macrophages transduced with retrovirus encoding ASC-GFP and found that caspase-8 colocalized with ASC puncta in response to infection with flagellated bacteria (Fig 2B). Collectively, these results indicate that flagellin triggers the assembly of an inflammasome composed of NLRC4 and ASC, which colocalizes with caspase-8.

The double-stranded DNA sensor AIM2 is known to recruit ASC to trigger puncta formation in response to infection, leading to caspase-1 activation and IL-1 β and IL-18 release [31,52–55]. The role of AIM2 inflammasome in the recognition of *L. pneumophila* has been demonstrated using *sdhA*[−] deficient bacteria. In the absence of SdhA, bacteria do not maintain vacuole integrity and localize in the macrophage cytoplasm, triggering activation of the AIM2 inflammasome [56,57]. In addition, the AIM2 inflammasome has been shown to trigger caspase-8 activation independently of caspase-1 [33,34,58]. Thus, we investigated whether AIM2 is present in the NLRC4/ASC/caspase-8 inflammasome that is formed in response to flagellin-positive *L. pneumophila*. We stained AIM2 in macrophages transduced with retrovirus encoding NLRC4-GFP and found no AIM2 in the NLRC4 puncta (S3A Fig). Moreover, we generated *Aim2/Casp1/11*^{−/−} mice and found that AIM2 was dispensable for the formation of NLRC4 puncta in response to flagellin-positive bacteria (S3B Fig).

The NLRC4/ASC/caspase-8 inflammasome accounts for flagellin-dependent restriction of *Legionella* replication in macrophages in the absence of caspase-1/11

Our data are consistent with the hypothesis that caspase-8 is a part of the inflammasome composed of NLRC4 and ASC. Thus, we investigated whether caspase-8 is activated during infection. We found that caspase-8 was strongly activated in *Casp1/11*^{−/−} BMDMs in response to *fliI* but not *flaA*[−] bacteria (Fig 3A). In agreement with the requirement of ASC for the assembly of the NLRC4/ASC/caspase-8 inflammasome, we found that caspase-8 activation was abolished in *Asc/Casp1/11*^{−/−} cells (Fig 3A). Caspase-8 activation occurred normally in *Aim2/Casp1/11*^{−/−} cells, indicating that AIM2 was not involved in the activation of caspase-8 through the

flagellin/NLRC4/ASC inflammasome (S4 Fig). We also evaluated caspase-8 activation by western blot analysis by measuring the cleavage of p55 and the production of p18 isoforms. We found that flagellated bacteria triggered caspase-8 activation in *Casp1/11*^{-/-} but not in *Asc/Casp1/11*^{-/-} cells. This phenomenon was evident by the reduction in p55 and increased production of p18 in *Casp1/11*^{-/-} BMDMs infected with *fljI*⁻ but not *flaA*⁻ bacteria (Fig 3B).

Next, we evaluated the participation of caspase-8 in caspase-1/11-independent restriction of *L. pneumophila* replication in macrophages, a process that was dependent on flagellin and NLRC4. Endogenous caspase-8 was silenced in *Casp1/11*^{-/-} BMDMs using two independent retrovirus encoding shRNA to target caspase-8. A non-target sequence was used as a control (NT). By western blotting, we detected reduced caspase-8 expression in *Casp1/11*^{-/-} transduced cells. The shRNA Casp8 Seq1 was more efficient than Seq2 for silencing caspase-8 as determined by western blot (Fig 4A and S5 Fig) and RT-PCR (Fig 4B). Importantly, complete silencing of caspase-8 cannot be achieved because caspase-8 expression is required for macrophage survival [59,60]. Nonetheless, using the described silencing conditions, we did not detect signs of cell death or LDH in the supernatant of the transduced macrophages. To evaluate the efficiency of caspase-8 silencing, we quantified caspase-8-containing puncta formation and caspase-8 activation in macrophages infected with flagellated *L. pneumophila*. We found that the frequency of puncta containing caspase-8 and caspase-8 activation was reduced in caspase-

Fig 2. ASC puncta are induced in response to flagellated bacteria and colocalize with caspase-8. Bone marrow-derived macrophages (BMDMs) from *Casp1/11*^{-/-} mice were transduced with retrovirus encoding ASC-GFP. (A) Cells were infected with wild-type *L. pneumophila* (WT Lp), motility-deficient mutants expressing flagellin (*fljI*⁻) or with flagellin-deficient mutants (*flaA*⁻) at a MOI of 10. After 1 or 8 hours of infection, the cells were fixed, and the number of transduced cells containing ASC-GFP puncta was determined using an epifluorescence microscope. (B) Representative images of a transduced BMDM infected with *fljI*⁻ at a MOI of 10. The cultures were stained with anti-caspase-8 (red), cell nuclei were stained with DAPI (cyan) and ASC-GFP is shown in green. The images show the colocalization between ASC-GFP and caspase-8 in *Casp1/11*^{-/-} BMDMs infected for 8 hours. The images were acquired by multiphoton microscopy with a 63x oil immersion objective and analyzed using ImageJ Software. Scale bar, 10μm. Data show the average ± SD of triplicate wells. *, *P* < 0.05, Student's *t* test. NI, uninfected. Data are presented for one representative experiment of three (A) and two (B) experiments with similar results.

<https://doi.org/10.1371/journal.ppat.1006502.g002>

8-silenced cells (Fig 4C and 4D). Next, we evaluated the effect of caspase-8 for the restriction of *L. pneumophila* replication in *Casp1/11*^{-/-} BMDMs. We found that silencing caspase-8 culminated in increased replication of *fliI*⁺ but not *flaA*⁻ bacteria (Fig 4E and 4F). These data indicated that caspase-8 contributed to the restriction of bacterial replication in a process that was dependent on flagellin, supporting the hypothesis that caspase-8 functionally participates in responses that are NLRC4/ASC-dependent and caspase-1/11-independent.

ASC is required for NLRC4/caspase-8-mediated restriction of *Legionella* replication in the absence of caspase-1/11

Our data indicate that caspase-8 is part of the NLRC4/ASC inflammasome and that ASC is essential for the assembly of this inflammasome. Thus, we reasoned that in the absence of

Fig 3. Caspase-8 is activated in response to flagellated bacteria in a process that is ASC-dependent and caspase-1/11-independent. Bone marrow-derived macrophages (BMDMs) from *Casp1/11*^{-/-} and *Asc/Casp1/11*^{-/-} mice were infected with motility-deficient *L. pneumophila* mutants expressing flagellin (*fliI*⁺) or with flagellin-deficient bacteria (*flaA*⁻) at a MOI of 10 for 8 hours. The activity of caspase-8 was measured using the Caspase-8 Glo Assay (A) or by western blot analysis (B). The pro-caspase-8 p55 and the processed form of caspase-8 p18 are indicated. Data show the average $\pm$ SD of triplicate wells. *, $P < 0.05$, Student's *t* test. RLU, relative luminescence units; NI, uninfected. Data are presented for one representative experiment of three (A) and two (B) experiments with similar results.

<https://doi.org/10.1371/journal.ppat.1006502.g003>

Fig 4. Caspase-8 is important for NLRC4-mediated restriction of *L. pneumophila* replication independently of caspase-1/11. Bone marrow-derived macrophages (BMDMs) generated from *Casp1/11*^{-/-} mice were transduced with a retrovirus encoding shRNA sequences to target caspase-8 (Seq1, Seq2) and a non-target shRNA sequence (NT). (A) The silencing was confirmed by western blot analysis and Real Time qPCR. (A) Cell lysates were separated by SDS-PAGE, blotted and probed with anti-caspase-8 (pro-caspase-8 p55) and anti- α -actin. (B) Quantification of the *casp8* gene expression by Real Time qPCR. *Actin beta* gene was used as a control for normalization of expression levels. The number above the bars indicates the percentage of silencing compared to the NT sequence. (C-D) The transduced *Casp1/11*^{-/-} BMDMs were infected with motility-deficient *L. pneumophila* mutants expressing flagellin (*flil*) at a MOI of 10. After 8 hours the cells were fixed, the number of caspase-8 puncta was quantified using an epifluorescence microscope (C) and the activity of caspase-8 was measured using the Caspase-8 Glo Assay (D). (E-F) The transduced *Casp1/11*^{-/-} BMDMs were infected with *flil*⁻ (E) and *flaA*⁻ (F) at a MOI of 10 to evaluate bacterial replication. The cells were incubated for 24, 48 and 72 hours for CFU determination. Data show the average $\pm$ SD of triplicate wells. * $P < 0.05$, compared with NT. (C-D) Student's *t* test, (E-F) ANOVA. RLU, relative luminescence units; NI, uninfected. Data are presented for one representative experiment of three experiments with similar results.

<https://doi.org/10.1371/journal.ppat.1006502.g004>

ASC, the NLRC4/ASC/caspase-8 inflammasome would not be functional. If this hypothesis is correct, *Asc/Casp1/11*^{-/-} macrophages should be more permissive than *Casp1/11*^{-/-} and as permissive as *Nlr4*^{-/-}. We infected C57BL/6, *Casp1/11*^{-/-}, *Asc/Casp1/11*^{-/-} and *Nlr4*^{-/-} BMDMs

with *fliI*⁻ and *flaA*⁻, and evaluated bacterial replication after 24, 48 and 72 hours. Using flagellin-positive bacteria, we confirmed that C57BL/6 BMDMs were restrictive to bacterial growth, *Nlrc4*^{-/-} were permissive and *Casp1/11*^{-/-} were partially restrictive (Fig 5A). Importantly, *Asc*/

Fig 5. ASC is important for NLRC4/caspase-8-mediated restriction of *L. pneumophila* replication independently of caspase-1/11. (A-D) Bone marrow-derived macrophages (BMDMs) from C57BL/6 (open circles), *Nlrc4*^{-/-} (closed squares), *Casp1/11*^{-/-} (closed triangles) and *Asc/Casp1/11*^{-/-} (open triangles) mice were infected with motility-deficient mutants expressing flagellin (*fliI*, A), with flagellin-deficient mutants (*flaA*, B), *L. gratiana* (C) or with *L. micdadei* (D) at a MOI of 10. The cells were incubated for 24, 48 and 72 hours for CFU determination. Data show the average $\pm$ SD of triplicate wells. *, $P < 0.05$ compared with *Casp1/11*^{-/-} BMDMs. #, $P < 0.05$ compared with C57BL/6 BMDMs, ANOVA. (E-I) BMDMs from *Casp1/11*^{-/-} and *Asc/Casp1/11*^{-/-} mice were transduced with a retrovirus encoding shRNA sequences to target caspase-8 (Seq1, Seq2) and a non-target shRNA sequence (NT). (E) Caspase-8 silencing was confirmed by western blot analysis. Cell lysates were separated by SDS-PAGE, blotted and probed with anti-caspase-8 (pro-caspase-8 p55) and anti- α -actin. (F-I) Transduced *Casp1/11*^{-/-} (F, H) and *Asc/Casp1/11*^{-/-} (G, I) BMDMs were infected with *fliI* (F, G) or *flaA* (H, I) at a MOI of 10 and incubated for 24, 48 and 72 hours for CFU determination. Data show the average $\pm$ SD of triplicate wells. *, $P < 0.05$ compared with NT shRNA, ANOVA. NT, non-target shRNA. Data are presented for one representative experiment of four (A), two (B-D) and one (F-I) experiments performed with similar results.

<https://doi.org/10.1371/journal.ppat.1006502.g005>

Casp1/11^{-/-} cells were highly permissive and phenocopied the *Nlrc4*^{-/-} cells (Fig 5A). As predicted, flagellin mutants bypassed the NLRC4 and replicated in all macrophages evaluated (Fig 5B). These data further confirmed that *Casp1/11*^{-/-} cells were more restrictive than *Nlrc4*^{-/-} macrophages, possibly due to the presence of the NLRC4/ASC/caspase-8 inflammasome. We also used non-pneumophila species to compare bacterial replication in *Nlrc4*^{-/-} and *Asc/Casp1/11*^{-/-} macrophages. Infections performed with *L. gratiana* and *L. micdadei* indicated that *Asc/Casp1/11*^{-/-} macrophages were as susceptible as *Nlrc4*^{-/-}, whereas *Casp1/11*^{-/-} cells were partially restrictive (Fig 5C and 5D). Similar experiments performed with *Aim2/Casp1/11*^{-/-} macrophages did not support the role of AIM2 in the NLRC4/ASC-dependent growth restriction that occurred in the absence of caspase-1/11 (S6 Fig). These data indicate that flagellated species of *Legionellae* trigger NLRC4 responses that are independent of caspase-1/11 but dependent on ASC. To further confirm the participation of caspase-8 in this NLRC4/ASC inflammasome, we silenced caspase-8 in *Casp1/11*^{-/-} and *Asc/Casp1/11*^{-/-} macrophages. We confirmed the silencing by western blot analysis (Fig 5E and S7 Fig) and found that the reduction in caspase-8 expression impaired the restriction of bacterial replication in *Casp1/11*^{-/-} infected with *fliI* (Fig 5F). Inhibition of caspase-8 expression affected neither the replication of *fliI* in *Asc/Casp1/11*^{-/-} cells (Fig 5G) nor the replication of *flaA* in *Casp1/11*^{-/-} and in *Asc/Casp1/11*^{-/-} cells (Fig 5H and 5I). Collectively, these data are consistent with the hypothesis that flagellin activates a response that is dependent on NLRC4, ASC and caspase-8 and occurs in the absence of caspase-1/11.

Casp8^{-/-} mice are embryonic lethal [59], and we were not able to generate *Casp8/11*^{-/-} mice. Since the deletion of ASC impairs the assembly of the NLRC4/ASC/caspase-8 inflammasome and caspase-8 activation, we used *Asc/Casp1/11*^{-/-} mice to assess the role of the NLRC4/ASC/caspase-8 inflammasome in the restriction of *Legionella* replication in vivo. Using flagellin-positive bacteria such as *L. pneumophila*, *L. gratiana* and *L. micdadei*, we demonstrated that *Asc/Casp1/11*^{-/-} mice were highly permissive to bacterial replication and phenocopied infection of *Nlrc4*^{-/-} mice (Fig 6A–6C). *Casp1/11*^{-/-} mice were more permissive than C57BL/6, but they were less permissive than *Nlrc4*^{-/-} mice (Fig 6A–6C). Experiments performed with *fliI* and *flaA* indicated that *Asc/Casp1/11*^{-/-} were more permissive than *Casp1/11*^{-/-} when infected with *fliI* but not *flaA* (Fig 6D). To determine whether AIM2 accounted for the restriction of bacterial replication in the absence of caspase-1/11, we compared infection of *Aim2/Casp1/11*^{-/-} with *Casp1/11*^{-/-}. We found that *Aim2/Casp1/11*^{-/-} and *Casp1/11*^{-/-} supported similar replication levels of *fliI* *L. pneumophila* in the lungs. In contrast, *Asc/Casp1/11*^{-/-} and *Nlrc4*^{-/-} mice were significantly more permissive to bacterial replication (S8 Fig). Collectively, these data indicate that AIM2 is dispensable for the functions of the NLRC4/ASC/caspase-8 inflammasome. This molecular platform is assembled in response to flagellin-positive bacteria and operates to restrict bacterial replication in vitro and in vivo in a process that is independent of both caspase-1 and caspase-11.

Activation of the NLRC4/ASC/caspase-8 inflammasome triggers pore formation and cell death

Activation of caspase-1 inflammasomes induces pyroptosis and contributes to the restriction of infection by flagellated bacteria such as *L. pneumophila*, *Salmonella typhimurium* and *Burkholderia thailandensis* [23]. Accordingly, we have previously demonstrated that *L. pneumophila* trigger pyroptosis in a process mediated by caspase-1 and caspase-11, which are activated in response to flagellin and LPS, respectively [20,24,35,61]. Thus, we investigated whether activation of the NLRC4/ASC/caspase-8 inflammasome could trigger cell death in *Casp1/11*^{-/-} macrophages. We assessed membrane permeabilization fluorometrically in real time via the

Fig 6. ASC is essential for NLRC4-mediated restriction of *L. pneumophila* replication independently of caspase-1/11 in vivo. C57BL/6 (open circles), *Nlrc4*^{-/-} (closed squares), *Casp1/11*^{-/-} (closed triangles) and *Asc/Casp1/11*^{-/-} (open triangles) mice were infected with motility-deficient *L. pneumophila* mutants expressing flagellin (*fliI*, A, D), with flagellin-deficient *L. pneumophila* (*flaA*⁻, D), with *L. gratiana* (B) or with *L. micdadei* (C) at a dose of 10⁵ bacteria/mice. The mice were euthanized at 4, 48 or 72 hours after infection. Dilutions of the lung homogenates were added to charcoal-yeast extract agar plates for colony-forming unit determination. Each dot represents a single animal, and the horizontal lines represent the averages. *, *P* < 0.05, #, indicates *P* < 0.05 compared with C57BL/6, Mann Whitney test. Data are presented for one representative experiment of five (A), one (B-C) and two (D) experiments performed with similar results.

<https://doi.org/10.1371/journal.ppat.1006502.g006>

uptake of propidium iodide. Macrophages were infected with *fliI* or *flaA*⁻, and pore formation was monitored in real time for 6 hours. C57BL/6 macrophages triggered robust pore formation in response to infection with *fliI* and reduced pore formation in response to *flaA*⁻ (Fig 7A–7C). The pore formation observed in C57BL/6 macrophages infected with *flaA*⁻ mutants was dependent on caspase-11 [61] and will not be addressed herein. Importantly, despite the absence of both caspase-1 and caspase-11, we detected significant pore formation in *Casp1/11*^{-/-} cells infected with *fliI* (Fig 7B). This response was not detected in *Casp1/11*^{-/-} cells infected with *flaA*⁻ or in *Asc/Casp1/11*^{-/-} macrophages infected with *fliI* or *flaA*⁻ (Fig 7B and 7C). These data support the hypothesis that NLRC4/ASC/caspase-8 induces pore formation. Experiments

Fig 7. The NLRC4/ASC/caspase-8 inflammasome is important for pore formation and cell death independently of caspase-1/11. (A-D; K-M) Bone marrow-derived macrophages (BMDMs) were generated from C57BL/6, *Casp1/11*^{-/-}, *Aim2/Casp1/11*^{-/-} and *Asc/Casp1/11*^{-/-} mice and infected with motility-deficient *L. pneumophila* mutants expressing flagellin (*flii*) or with flagellin-deficient bacteria (*flaA*) at a MOI of 5. (E-J) BMDMs were transduced with retrovirus encoding shRNA sequences to target caspase-8 (Seq1, Seq2) and a non-target shRNA sequence (NT) and infected with *flii* or *flaA* at a MOI of 5. (A-C; E-J) Pore formation was assessed fluorometrically in real time by the uptake of propidium iodide. RFU (%) represents the percentage of RFU estimated with cells lysed with Triton X-100. (D) LDH release was measured using the CytoTox 96 LDH-release kit. The LDH release (%) represents the percentage of LDH release estimated with cells lysed with Triton X-100. (K-M) BMDMs were treated with 50 μ M of Z-IETD or DMSO for 1 hour and infected with (L) *flii* or (M) *flaA* at a MOI of 5. Pore formation was assessed fluorometrically in real time by the uptake of propidium iodide. Data show the average $\pm$ SD of triplicate wells. *, $P < 0.05$, Student's *t* test. NS, not significant; RFUs, relative fluorescence units; NI, uninfected. Data are presented for one representative experiment of five (A-C), three (E-J) and two (D and K-M) experiments with similar results.

<https://doi.org/10.1371/journal.ppat.1006502.g007>

performed using *Aim2/Casp1/11*^{-/-} macrophages corroborate our previous findings, indicating that AIM2 is not required for the activities of the NLRC4/ASC/caspase-8 inflammasome (Fig 7B and 7C). Pore formation induced in response to caspase-1 and caspase-11 activation

culminates with the induction of macrophage lysis, a process that can be assessed by the presence of LDH in tissue culture supernatants [24,61,62]. Thus, we measured LDH release in the supernatants of macrophages infected with *fliI*⁺ and *flaA*⁺ for 8 hours. By comparing *Casp1/11*^{-/-} and *Asc/Casp1/11*^{-/-} cells, we found that macrophage lysis occurred despite the absence of caspase-1/11 (Fig 7D). Cell death was flagellin-dependent because infections with *fliI*⁺ but not *flaA*⁺ induced LDH release (Fig 7D). The participation of caspase-8 in pore formation and cell death induced in caspase-1/11-deficient macrophages was evident using both MOI 5 (Fig 7) and MOI 10 (S9 Fig). Importantly, cell death was not observed in *Asc/Casp1/11*^{-/-}, a feature that corroborates the pore formation studies and indicates that the NLRC4/ASC/caspase-8 inflammasome triggers pore formation and lysis of infected cells. We also assessed whether the NLRC4/ASC/caspase-8 inflammasome was important for the activation of inflammatory cytokines. We found that whereas C57BL/6 macrophages readily triggered the production of IL-1 β after 24 hours of infection with flagellated bacteria, the *Casp1/11*^{-/-} or *Asc/Casp1/11*^{-/-} deficient cells do not trigger a IL-1 β production (S10A Fig). The production of IL-12p40 by these cells confirmed that all macrophages were primed and could respond to *L. pneumophila* infection (S10B Fig). To evaluate the participation of caspase-8 in cell death induced by the NLRC4/ASC/caspase-8 inflammasome, we silenced endogenous caspase-8 using shRNA. Macrophages that were transduced with retrovirus encoding shRNA did not exhibit pore formation before infection, indicating that transduction itself did not trigger cell death (Fig 7E and 7H). In contrast, pore formation was evident in *Casp1/11*^{-/-} but not in *Asc/Casp1/11*^{-/-} macrophages in response to *fliI*⁺ infection (Fig 7F and 7I). Pore formation induced in *Casp1/11*^{-/-} was diminished in caspase-8-silenced cells (Fig 7F). In support of the role of flagellin for triggering these responses, we did not detect pore formation in cells infected with *flaA*⁺ (Fig 7G and 7J). To further evaluate the participation of caspase-8 in pore formation induced by flagellin, we performed pore formation assays using Z-IETD, a cell permeable peptide that binds irreversibly to the catalytic site of caspase-8 [63–65]. We found that treatment of *Casp1/11*^{-/-} macrophages with DMSO or Z-IETD did not cause pore formation in uninfected cells (Fig 7K). However, Z-IETD treatment reduced the pore formation induced by *fliI*⁺ (Fig 7L) but not by *flaA*⁺ (Fig 7M). Collectively, these data indicate that flagellin-positive bacteria trigger pore formation and cell death-independent of caspase-1/11 via a process that requires ASC and caspase-8.

Naip5 is required for NLRC4/ASC/caspase-8 inflammasome activation in response to flagellated *Legionella*

Our data reveal that the NLRC4/ASC/caspase-8 inflammasome is activated in *Casp1/11*^{-/-} macrophages in response to infection with flagellated *Legionella*. To evaluate if Naip5 is required for activation of this inflammasome, we used shRNA to silence endogenous Naip5. In Naip5 silenced *Casp1/11*^{-/-} macrophages, we detected a reduced activation of caspase-8 in response to WT Lp and *fliI*⁺ bacteria (Fig 8A). Naip5 silencing was confirmed by RT-PCR (Fig 8B). We also tested if Naip5 is important for pore formation induced via caspase-8 in *Casp1/11*^{-/-} macrophages. By evaluating pore formation, we found that Naip5 is important for efficient pore formation in response to WT Lp and *fliI*⁺ bacteria (Fig 8D–8E). As previously reported, no pore formation was detected in response to *flaA*⁺ bacteria (Fig 8F) or in *Asc/Casp1/11*^{-/-} macrophages (Fig 8G–8J). Finally, we tested if Naip5 is important for restriction of *L. pneumophila* replication in *Casp1/11*^{-/-} macrophages. We found that Naip5 is important for restriction of flagellin-positive *L. pneumophila* replication in *Casp1/11*^{-/-} macrophages (Fig 8K). Naip5 silencing did not affect the replication of *flaA*⁺ bacteria in *Casp1/11*^{-/-} macrophages (Fig 8L). As predicted, *Asc/Casp1/11*^{-/-} macrophages were permissive to replication of both *flaA*⁺ and *fliI*⁺ bacteria and Naip5 did not influence this process (Fig 8M and 8N). Taken together, these

Fig 8. Naip5 is required for the functions of the NLRC4/ASC/Caspase-8 inflammasome independently of caspase-1/11. Bone marrow-derived macrophages (BMDMs) generated from *Casp1/11*^{-/-} and *Asc/Casp1/11*^{-/-} mice were transduced with a retrovirus encoding shRNA sequence to target Naip5 and a non-target shRNA sequence (NT). Transduced BMDMs were infected with wild-type *L. pneumophila* (WT Lp), motility-deficient mutants expressing flagellin (*fliI*⁻) or with flagellin-deficient mutants (*flaA*⁻) at a MOI of 10. (A) 8 hours after infection the activity of caspase-8 was measured using the Caspase-8 Glo Assay. (B) Quantification of the *Naip5* gene expression by Real Time qPCR. *Gapdh* gene was used as a control for normalization of expression levels. The number above the bars indicates the percentage of silencing compared to the NT shRNA (open bars). (C–J) Pore formation was assessed fluorometrically in real time by the uptake of propidium iodide. The RFU (%) represents the percentage of RFU estimated with cells lysed with Triton X-100. (K–N) The cells were infected with *fliI*⁻ or *flaA*⁻ for 24, 48 and 72 hours for CFU determination. Data show the average ± SD of triplicate wells. *, *P* < 0.05, Student's *t* test (A) and ANOVA (K–N). NS, not significant; RFU, relative fluorescence units; NI, uninfected. Data are presented for one representative experiment of two (A–J) and three (K–N) experiments with similar results.

<https://doi.org/10.1371/journal.ppat.1006502.g008>

data indicates that Naip5 participate of the NLRC4/ASC inflammasome that trigger caspase-8 activation in the absence of caspase-1/11.

Caspase-8 is recruited to the NLRC4/ASC/caspase-1 inflammasome but it is only activated when caspase-1 or gasdermin-D is suppressed

In the experiments shown thus far, we used *Casp1/11*^{-/-} macrophage as a tool to assess the caspase-8 effects without the interference of caspase-1. However, because caspase-1 is present in natural conditions, we tested if caspase-8 participates in the NLRC4/ASC inflammasome in

the presence of caspase-1. First, we infected C57BL/6 macrophages with flagellin-positive *L. pneumophila* to assess if endogenous caspase-8 colocalizes with the Naip5/NLRC4/ASC inflammasome. In uninfected conditions, we detected no significant puncta formation (Fig 9A). However, in response to *fliI* bacteria, caspase-8 colocalizes with ASC (Fig 9B) and caspase-1 (Fig 9C). We determined that caspase-8 is present in more than 60% puncta containing caspase-1 (Fig 9C). These data indicates that regardless to the presence of caspase-1, the caspase-8 is recruited to the inflammasome during activation.

Next, we evaluated caspase-8 activation in wild-type macrophages. We found that caspase-8 activation does not occur in C57BL/6 macrophages infected with *L. pneumophila*. As expected caspase-8 is readily activated in *Casp1/11*^{-/-}, but not in *Asc/Casp1/11*^{-/-} macrophages infected with *fliI* bacteria (Fig 9D). To ensure similar genetic background, we intercrossed a F1 progeny of *Casp1/11*^{-/-} x C57BL/6 to obtain F2 littermate controls. Infections in macrophages from littermate control mice indicated that caspase-8 activation occur in *Casp1/11*^{-/-}, but not in *Casp1/11*^{+/-} and *Casp1/11*^{+/+} macrophages (Fig 9E). To further test whether the deficiency in caspase-1 or caspase-11 enable caspase-8 activation, we performed experiments using mice single deficient in caspase-1 or caspase-11 as previously described [35,66]. We found that caspase-11 deficiency alone is not sufficient to enable caspase-8 activation in response to *L. pneumophila* infection (Fig 9F). In contrast, caspase-1-deficient cells expressing caspase-11 as a transgene (*Casp1*^{-/-}*Casp11*^{tg}) effectively trigger caspase-8 activation in response to *fliI* *L. pneumophila* (Fig 9F). These data indicate that caspase-1 but not caspase-11 is required to prevent caspase-8 activation.

Caspase-1 activation via the NLRC4 inflammasome is known to trigger activation of gasdermin-D (GSDMD) to induce cell death [67,68]. Thus, we tested if inhibition of GSDMD is sufficient to enable caspase-8 activation in the presence of caspase-1. To achieve this, we inhibited endogenous GSDMD using shRNA and found that despite the presence of caspase-1, caspase-8 is robustly activated when GSDMD is inhibited (Fig 9G). RT-PCR was used to confirm the silencing of the two different sequences of shRNA used (Fig 9H). To further confirm the GSDMD silencing, we performed pore formation assay in C57BL/6 macrophages infected with *L. pneumophila*. We found that GSDMD silencing inhibited the caspase-1-mediated pore formation induced by WT Lp and *fliI* *L. pneumophila* (S11A–S11D Fig). However, GSDMD did not participate of pore formation induced via caspase-8 that occurs in *Casp1/11*^{-/-} macrophages (S11E–S11H Fig). Collectively, these data indicates that despite the presence of caspase-1, caspase-8 activation occur in the Naip5/NLRC4/ASC inflammasome when GSDMD is inhibited.

Discussion

The recognition of *Legionella* flagellin by the Naip5/NLRC4 inflammasome in macrophages is a major mechanism for the restriction of bacterial replication in mouse cells [17–20,22,69,70]. It is well accepted in the field that not all NLRC4 functions require caspase-1 [29,38,71,72]. This conclusion is supported by the observation that *Nlrc4*^{-/-} mice (and their macrophages) are significantly more permissive to *L. pneumophila* replication than *Casp1/11*^{-/-} mice [29,38]. Here, we unraveled this caspase-1/11-independent pathway and characterized an inflammasome composed of Naip5, NLRC4, ASC and caspase-8, which operates in the absence of caspase-1 and caspase-11. This inflammasome effectively participates in the mechanisms involved in the restriction of bacterial replication in macrophages and in vivo. Previous biochemistry studies using yeast two-hybrid screening showed that NLRC4 is ubiquitinated by Sug1, a process that facilitates the activation of caspase-8 [51]. Thus, it is possible that Sug1 is also a component of this NLRC4 inflammasome. In addition, previous studies using the *Salmonella*

Fig 9. Caspase-8 colocalizes with the NLRC4/ASC/Caspase-1 inflammasome, but it is only activated if caspase-1 or gasdermin-D is inhibited. (A-C) Bone marrow-derived macrophages (BMDMs) generated from C57BL/6 mice were infected with motility-deficient *L. pneumophila* mutants expressing flagellin (*flii*) at a MOI of 10 for 8 h. The cultures were stained with anti-caspase-1 (green) (A, C) or anti-ASC (green) (B), anti-caspase-8 (red). Cell nuclei were stained with DAPI (cyan). Images were acquired by multiphoton microscopy with a 63x oil immersion objective and analyzed using ImageJ software. The images are the maximal projection of a z project. Scale bars, 10 μm. (C) The percentage of the caspase-1 puncta containing caspase-8 was determined using an epifluorescence microscope. (D) BMDMs were generated from C57BL/6, *Casp1*^{11-/-} and *Asc/Casp1*^{11-/-} mice and infected with *flii*⁻ or *flii*⁺. (E) BMDMs were generated from *Casp1*^{11-/-}, *Casp1*^{11-/-} and *Casp1*^{11-/-} littermate control mice and infected with WT, *flii*⁻ and *flii*⁺. (F) BMDMs were generated from C57BL/6, *Casp1*^{11tg} and *Casp1*^{11-/-} mice and infected with WT, *flii*⁻ and *flii*⁺. (G-H) BMDMs generated from C57BL/6 and *Casp1*^{11-/-} mice were transduced with a retrovirus encoding shRNA sequences to target Gasdermin D (GSDMD) (Seq1, Seq2) and a non-target shRNA sequence (NT). (G) Transduced cells were infected with WT Lp, *flii*⁻ and *flii*⁺. (D-G) Cells were infected with an MOI of 10 and after 8 hours the activity of caspase-8 was measured using the Caspase-8 Glo Assay. (H) GSDMD silencing was confirmed by Real Time qPCR. *Gapdh* gene was used as a control for normalization of expression levels. The numbers above the bars indicate the percentage of silencing compared to the NT sequence. Data show the average ± SD of triplicate wells. *, P < 0.05, Student's t test in relation to *flii*⁺. RLU, relative luminescence units; NI, non-infected. Data are presented for one representative experiment of two (A-C and E-H) and five (D) experiments with similar results.

<https://doi.org/10.1371/journal.ppat.1006502.g009>

enterica serovar Typhimurium have indicated that both caspase-8 and caspase-1 are recruited to the ASC puncta in response to infection. However, caspase-8 is involved in the synthesis of pro-IL-1 β and is dispensable for *Salmonella*-induced cell death [32]. These data contrast with published data using *L. pneumophila*, which indicate that in the absence of caspase-1/11, no inflammatory cytokines are produced [20,21,25,30,35,66]. Accordingly, our data indicates that this Naip5/NLRC4/ASC/Caspase-8 inflammasome is very inefficient to trigger IL-1 β maturation when caspase-1 is not present.

Importantly, AIM2 is not part of this NLRC4/ASC/caspase-8 inflammasome. AIM2 is well-known to trigger caspase-8 activation via ASC [33,34,58]. Our data unequivocally demonstrate that AIM2 is neither a component of this inflammasome, nor is it required for inflammasome functions. AIM2 did not colocalize with the NLRC4/ASC/caspase-8 puncta, it was dispensable for the activation of caspase-8 in response to flagellated *L. pneumophila* and for the induction of cell death and restriction of *L. pneumophila* replication. The pyrin domain of ASC can bind to the death domain of caspase-8 [46]. Thus, it is possible that the interactions of the ASC pyrin domain with the caspase-8 dead domain are critical for the recruitment of caspase-8 to the complex. Consistent with this hypothesis, our studies unequivocally show that ASC is required for the assembly and function of this NLRC4 inflammasome. Importantly, the characterization of this NLRC4/ASC/caspase-8 inflammasome accounted to clarify a controversy in the field concerning the participation of ASC in the NLRC4 inflammasome. Studies utilizing biochemistry and cells from gene-deficient mice have demonstrated that ASC is dispensable for NLRC4 functions, including pyroptosis and the restriction of *L. pneumophila* replication [20,21,29]. However, ASC is essential for caspase-1 cleavage and the processing of inflammatory cytokines in response to flagellated *L. pneumophila*, a process that is NLRC4-dependent and NLRP3-independent [20,21,25,30]. In addition, ASC participates in the restriction of intracellular replication of *L. pneumophila* under certain circumstances [36,37]. Our studies provide data that help to consolidate these data in a cohesive model. NLRC4 can operate to form an inflammasome in absence of ASC that triggers pore formation and the restriction of bacterial replication [20,21,25,30]. This platform does not form puncta and is ineffective for triggering caspase-1 cleavage and processing inflammatory cytokines. When ASC is present, NLRC4 inflammasome associates with ASC and recruit caspase-1 and caspase-8 to the puncta. This inflammasome is very efficient to cleave caspase-1 and inflammatory cytokines such as IL-1 β . Interestingly, our data indicate that caspase-8 is not activated when caspase-1 is present. However, when caspase-1 is missing or when Gasdermin-D is inhibited, we detected a robust caspase-8 activation. These data suggest that activation of caspase-8 in the Naip5/NLRC4/ASC inflammasome functions as a backup strategy to guarantee cell death when key components of the pyroptotic cell death are inhibited. Interestingly, our data and previously published data indicate that gasdermin-D is dispensable for caspase-8-induced cell death [73]. Therefore, when gasdermin-D is inhibited, caspase-8 engages gasdermin-D-independent cell death. It is possible that caspase-8 induces caspase-3 and caspase-7 to targeted gasdermin-E (also known as DFNA5) to induce pore formation and cell death independent of gasdermin-D [74,75]. This may guarantee appropriate responses to pathogens that inhibit canonical components of pyroptotic cell death such as caspase-1 or gasdermin-D.

Materials and methods

Bacterial culture

The *L. pneumophila* bacteria used were JR32 and isogenic clean deletion mutants for motility (*fliT*) and flagellin (*flaA*) [19,38]. *L. micdadei* (ATCC 33218) and *L. gratiana* (ATCC 49413) were used to generate streptomycin-resistant strains. *RpsL* mutants of *L. micdadei* and *L.*

gratiana were isolated by plating these strains on CYE agar containing 100 µg/ml of streptomycin. All bacteria were grown on buffered charcoal-yeast extract (CYE) agar plates [1% yeast extract, 1% MOPS, 3.3 mM L-cysteine, 0.33 mM Fe(NO₃)₃, 1.5% Bacto agar and 0.2% activated charcoal, pH 6.9] [76].

Macrophages

Bone Marrow derived macrophages (BMDMs) were generated from mice as previously described [77]. Briefly, bone marrow cells were harvested from femurs and differentiated with RPMI 1640 (Gibco) containing 20% fetal bovine serum (FBS—Invitrogen) and 30% L-929 cell-conditioned medium (LCCM), 2 mM L-glutamine (Sigma-Aldrich), 15 mM Hepes (Gibco) and 100 U/ml penicillin-streptomycin (Sigma-Aldrich) at 37°C with 5% CO₂ [77]. BMDMs were seeded at 2 X 10⁵ cells/well in 24-well plates and cultivated in RPMI 1640 medium (Gibco) supplemented with 10% FBS, 5% LCCM, 2 mM L-glutamine and 15 mM Hepes.

In vitro infections and CFU determination

For the in vitro infections, the cultures were infected at a multiplicity of infection of 0.015, 5 or 10 followed by centrifugation for 5 minutes at 300 X g at room temperature and incubation at 37°C in a 5% CO₂ atmosphere. In the colony-forming units (CFU) experiments, cultures infected at a MOI of 10 were washed two times with PBS, and 1 ml of medium was added to each well. For CFU determination, the cultures were lysed in sterile water, and the cell lysates were combined with the cell culture supernatant from the respective wells. Lysates plus supernatants from each well were diluted in water, plated on CYE agar plates, and incubated for 4 days at 37°C for CFU determination as described previously [28,38].

Retroviral transduction and quantification of NLRC4-GFP and ASC-GFP puncta

Murine *Nlrc4* or *Asc* were cloned into the pEGFP (N2) vector (Clontech) using XhoI and BamHI restriction sites as previously described [30]. NLRC4-GFP or ASC-GFP and GFP were cloned into the pMSCV2.2 murine-specific retroviral vector (Clontech). The pCL vector system 51 was used to package the retroviruses in transfected monolayers of Hek Peak cells (ATCC CRL-2828), which were maintained in RPMI with 10% FBS. The supernatant from the Hek Peak cells containing retrovirus was collected three days after transfection, filtered using a 0.45-µm filter and used for BMDM transduction. BMDMs were obtained from *Casp1/11*^{-/-}, *Asc/Casp1/11*^{-/-} and *Aim2/Casp1/11*^{-/-} mice and seeded in differentiation medium. On day 3 of differentiation, the supernatants containing retroviral were added to BMDMs in 20% FBS and 25% LCCM. After differentiation, the BMDMs were seeded at 2 X 10⁵ cells/well in 24-well plates containing 12-mm glass cover slides and cultivated in RPMI 1640 medium supplemented with 10% FBS and 5% LCCM. For the caspase colocalization experiments, the cultures were treated with 20 µM of Z-VAD for 1 hour and infected at a MOI of 1, 3 or 10. After infection, the plates were centrifuged for 5 minutes at 300 X g at room temperature and incubated at 37°C in a 5% CO₂ atmosphere. At 1, 2, 4 and 8 hours after infection, the cells were fixed with 4% paraformaldehyde, permeabilized with 0.05% saponin, stained with DAPI and mounted on glass slides using Prolong Gold Antifade Reagent (Invitrogen). For the colocalization assay, the cells were stained with rat anti-caspase-8 (Enzo—1G12; 1:50), rabbit anti-cleaved caspase-8 (Cell signaling- 8592; 1:800), rabbit anti-cleaved caspase-3 (Cell signaling- 9664; 1:400); rabbit anti-cleaved caspase-7 (Cell signaling-8438; 1:400), anti-ASC (Adipogen- AL177; 1:250), goat anti-ASC (Santa Cruz—sc33958; 1:50), rabbit anti-caspase-1 (Santa Cruz- sc514;

1:500) or anti-AIM2 (Cell signaling- 13095; 1:400), followed by Alexa 594-conjugated goat anti-rabbit secondary Ab (Invitrogen; 1:3000), Alexa 594-conjugated goat anti-rat secondary Ab (Invitrogen; 1:3000) or Alexa 647-conjugated chicken anti-rabbit secondary Ab (Invitrogen; 1:2000) and DAPI and mounted on glass slides using Prolong Gold Antifade Reagent (Invitrogen). The images were processed using LAS AF software (Leica Microsystems) and analyzed under fluorescence using a Leica DMI 4000B inverted microscope with a 100X oil objective. The number of NLRC4-GFP or ASC-GFP puncta in the transduced cells and the colocalization were quantified. Bacteria were not stained. Therefore, the whole cell population was scored. Multiphoton microscopy images were acquired using an LSM 780 Zeiss AxioObserver microscope equipped with a 63X oil immersion objective and analyzed using ImageJ software.

Retroviral silencing of caspase-8, Naip5 and GSDMD

For retroviral silencing of caspase-8, Naip5 and GSDMD (Gasdermin D), Hek Peak cells were transfected with lentiviral vectors encoding a small hairpin RNA (shRNA) targeting caspase-8 [Sigma- Seq1: TRCN0000231279 (Sequence- CCGGTCATCTCACAAGAACTATATTCTCG AGAATATAGTTCTTGTGAGATGATTTTGTG); Seq2: TRCN0000231281 (Sequence-CCGG TCCTGACTGGCGTGAAGTATGCTCGAGCATAGTTCACGCCAGTCAGGATTTTGTG)], Naip5 [Sigma- TRCN0000114742 (Sequence—CCGGCGCTTGATTATCTTCTGGAAACT CGAGTTTCCAGAAGATAATCAAGCGTTTGTG)], GSDMD [Sigma- TRCN0000219619 (Sequence—CCGGGATTGATGAGGAGGAATTAATCTCGAGATTAATTCCTCCTCATC AATCTTTTGTG); TRCN0000219620 (Sequence—CCGGCCTAAGGCTGCAGGTAGAATC CTCGAGGATTCTACCTGCAGCCTTAGGTTTTGTG)] and a negative control vector that included a non-target shRNA sequence (NT). The plates were treated with polyethylenimine (Sigma-Aldrich) (Corning). Transduced cells were maintained in RPMI 1640 medium supplemented with 10% FBS at 37°C and 5% CO₂. Lentiviruses expressing shRNAs were collected, filtered using a 0.45-μm filter and added to the BMDMs. After selection with puromycin, the resistant cells were seeded at 2 X 10⁵ cells/well in 24-well plates and infected with *fliI* or *flaA* for CFU determination. The caspase-8 silencing efficiency was measured by immunoblotting: 1 X 10⁶ cells were lysed in RIPA buffer (10 mM Tris-HCl, pH 7.4, 1 mM EDTA, 150 mM NaCl, 1% Nonidet P-40, 1% deoxycholate, and 0.1% SDS) in the presence of a protease inhibitor cocktail (Roche). The lysates were suspended in 4X Laemmli buffer, boiled for 5 minutes, resolved by 15% SDS PAGE and transferred (Semidry Transfer Cell, Bio-Rad) to 0.22-μm nitrocellulose membranes (GE Healthcare). The membranes were blocked in Tris-buffered saline (TBS) with 0.01% Tween-20 and 5% non-fat dry milk for 1 hour. Rat anti-caspase-8 p55 (Enzo- 1G12; 1:500) and anti-rat peroxidase-conjugated antibody (KPL, 1:3000) were diluted in blocking buffer for the incubations (overnight for anti-caspase-8 and 1 hour for the secondary antibody). The ECL luminol reagent (GE Healthcare) was used for antibody detection. For evaluation of caspase-8 silencing by immunoblots, Image J software were used to estimate the ratio of caspase-8 p55 to α-actin.

Caspase-8 activation

To assess caspase-8 activation, BMDMs were infected with *fliI* or *flaA* for 8 hours, and the activity of caspase-8 was measured using the Caspase-8 Glo Assay (Promega) according to manufacturer's recommendations. To evaluate caspase-8 activation by western blot analysis, 1 X 10⁷ cells were infected at a MOI of 10 and lysed 8 hours after infection in RIPA buffer with protease inhibitor, as previously described. The lysates were suspended in 4X Laemmli buffer, boiled for 5 minutes, resolved by 15% SDS PAGE and transferred to 0.22-μm nitrocellulose

membranes. The membranes were blocked in Tris-buffered saline (TBS) with 0.01% Tween-20 and 5% non-fat dry milk or for 1 hour. The mouse anti-caspase-8 (Enzo- 1G12) and anti-rabbit peroxidase-conjugated antibody (KPL; 1:3000) were diluted in blocking buffer for the incubations (overnight for anti-cleaved caspase-8 and 1 hour for the secondary antibody). ECL luminol reagent (GE Healthcare) was used for antibody detection.

Real Time q-PCR

Total RNA was extracted from 2×10^6 macrophages using total RNA isolation kit (illustra RNAspin, GE Healthcare, UK), according to manufacturer's instructions. After extraction, an aliquot of 2 μ l was used to determine the RNA concentration in NanoDrop (Thermo Fisher Scientific) and 1 μ g of the extracted RNA was used for the cDNA conversion using the iScriptTM cDNA Synthesis kit (BIO-RAD) in a thermal cycler. The cDNA (10 ng) was used for the quantification of the Caspase 8 gene expression (TaqMan Assay: Casp 8—Mm01255716_m1) by real-time PCR using TaqMan Fast Advanced Master Mix, according to the manufacturer's instruction (Applied Biosystems). Actin beta (Actb) gene (TaqMan Assay: Actb-Mm00607939_s1) was used as a control for normalization of expression levels. The quantification of GSDMD and Naip5 were performed using 25 ng of cDNA and 10 μ M of each primer, 1X SYBR Green (Applied Biosystems), and was normalized using the housekeeping gene glyceraldehyde-3-phosphate dehydrogenase (Gapdh). The specificity of the PCR products was assessed by melting curve analysis for all samples. The following primers were used: Gapdh, FWD- AGGTCGGGTG TGAACGGATTG, REV-TGTAGACCATGTAGTTGAGGTCA, GSDMD, FWD-TCATGT GTCAACCTGTCAATCAAGGACAT, REV- CATCGACGACATCAGAGACTTTGAAGGA, NAIP5, FWD- GTTGAGATTGGAGAAGACCTCG, REV-CACACGTGAAAGCAACCATGG. The real-time quantitative reaction was performed in the Viia 7 Real-Time PCR System (Applied Biosystems). The results were analyzed using the $2^{-\Delta\Delta CT}$ method and are expressed in relation to the reference group. The percentage of silencing knockdown was estimated using the $[1-(2^{-\Delta\Delta CT}) \times 100]$ equation [78].

Mice and in vivo infections

Mice used in this study were breed and maintained in institutional animal facilities. Mice used were C57BL/6 (Jax 000664), *Nlrc4*^{-/-} [79], *Casp1*¹¹^{-/-} [80], *Asc*^{-/-} [81], *Casp11*^{-/-} and *Casp1*^{-/-} *Casp11*^{tg} [66] were. Double-deficient mice were generated by intercrossing a F1 progeny of the parental strains. All mice were matched by sex and age (all were at least 8 weeks old at the time of infection) and were in a C57BL/6 mouse genetic background. For the in vivo experiments, approximately 5–7 mice per group were used, as indicated in the figures. For in vivo infections, the mice were anesthetized with ketamine and xylazine (50 mg/kg and 10 mg/kg, respectively) by intraperitoneal injection followed by intranasal inoculation with 40 μ l of phosphate-buffered saline (PBS) containing 1×10^5 bacteria per mouse. For CFU determination, the lungs were harvested and homogenized in 5 ml of sterile water for 30 seconds using a tissue homogenizer (Power Gen 125; Thermo Scientific). Lung homogenates were diluted in sterile water and plated on CYE agar plates for CFU determination as previously described [28,38].

Pore formation assay

Pore formation in BMDMs was quantified based on the permeability to propidium iodide (PI) in damaged cells as previously described [61]. BMDMs were seeded in a black, clear-bottom 96-well plate (1×10^5 cells/well). Before infection, the medium was replaced with 10% RPMI without phenol red, 0.038 g/ml NaHCO₃, 6 μ l/ml PI and anti-*L. pneumophila* (1:1000). Infected BMDMs were maintained at 37 °C, and PI was excited at 538 nm. The fluorescence

emission was read at 617 nm every 5 minutes using a plate fluorometer (SpectraMax i3x, Molecular Devices). Total pore formation was determined by lysing cells with Triton X-100.

Lactate dehydrogenase release assay and ELISA

BMDMs were seeded in 24-well plates (5×10^5 cells/well). Infections were performed in RPMI1640 medium without phenol red, 15 mM HEPES and 2 g/l NaHCO_3 supplemented with 10% FBS. After 8 hours of infection, the supernatants were collected for analysis of lactate dehydrogenase (LDH) release. Total LDH was determined by lysing the cultures with Triton X-100. LDH was quantified using the CytoTox 96 LDH-release kit (Promega). For cytokine determination, enzyme-linked immunosorbent assay (ELISA) were used. BMDMs were seeded into 24-well plates (5×10^5 cells/well) and infected with WT Lp, *fliI*⁻ and *flaA*⁻ (MOI 10) for 24 hours. BMDMs supernatant was assessed using ELISA kits according to manufacturer's recommendations (BD Biosciences).

Ethics statement

The care of the mice was in compliance with the institutional guidelines on ethics in animal experiments; approved by CETEA (Comissão de Ética em Experimentação Animal da Faculdade de Medicina de Ribeirão Preto, approved protocol number 218/2014). CETEA follow the Brazilian national guidelines recommended by CONCEA (Conselho Nacional de Controle em Experimentação o Animal). For euthanasia, the mice were treated with ketamine and xylazine (50 mg/kg and 10 mg/kg, respectively) by intravenous injection.

Statistical analysis

The data were plotted and analyzed using GraphPad Prism 5.0 software. The statistical significance was calculated using the Student's t-test or analysis of variance (ANOVA). Nonparametric test Mann–Whitney U test were used for analysis of in vivo experiments. Differences were considered statistically significant when P was <0.05 , as indicated by an asterisk in the figures.

Supporting information

S1 Fig. Restriction of *L. pneumophila* replication is fully dependent on NLRC4 and flagellin, and partially dependent on caspase-1/11. Bone marrow-derived macrophages (BMDMs) from C57BL/6 (open circles), *Nlrc4*^{-/-} (closed squares), *Casp1/11*^{-/-} (closed triangles) and *Asc*^{-/-} (open inverted triangles) mice were infected with *L. pneumophila* for CFU determination. (A, B) Cells were infected with wild-type *L. pneumophila* (WT Lp). (C, D) Cells were infected with motility-deficient mutants expressing flagellin (*fliI*). (E, F) Cells were infected with flagellin-deficient mutants (*flaA*⁻). BMDMs were infected with 3×10^3 (MOI 0.015) or 2×10^5 (MOI 10) bacteria per well and incubated for 24, 48, 72 and 96 hours for CFU determination. Data show the average $\pm$ SD of triplicate wells. *, $P < 0.05$ compared with *Casp1/11*^{-/-} BMDMs, ANOVA. Data are presented for one representative experiment of two experiments with similar results. (TIF)

S2 Fig. Caspase-8 but not caspase-3 and caspase-7 colocalize with NLRC4-GFP and ASC-GFP puncta. Bone marrow-derived macrophages (BMDMs) generated from *Casp1/11*^{-/-} mice were transduced with retrovirus encoding NLRC4-GFP (A) or ASC-GFP (B) and infected with motility-deficient *L. pneumophila* mutants expressing flagellin (*fliI*) at a MOI of 10 for 8 h. (A–B) The percentage of colocalization of caspase-3, caspase-7 and caspase-8 with NLRC4-GFP and ASC-GFP was determined using an epifluorescence microscope. (C–E) BMDMs generated from *Casp1/11*^{-/-} mice were infected with *fliI* at a MOI of 10 for 8 h. The

cultures were fixed and stained with anti-ASC (green), anti-caspase-3 (red) (C), anti-caspase-7 (red) (D), anti-caspase-8 (red) (E). Cell nuclei were stained with DAPI (cyan). Images were acquired by multiphoton microscopy with a 63x oil immersion objective and analyzed using ImageJ software. The images are the maximal projection of a z project. Scale bar, 10µm. Data show the average ± SD of triplicate wells. *, $P < 0.05$, Student's t test. Data are presented for one representative experiment of two experiments with similar results.

(TIF)

S3 Fig. AIM2 does not colocalize and is not required for NLRC4-GFP puncta formation.

Bone marrow-derived macrophages (BMDMs) generated from *Casp1/11^{-/-}* and *Aim2/Casp1/11^{-/-}* mice were transduced with retrovirus encoding NLRC4-GFP and infected with wild-type *L. pneumophila* (WT) or with motility-deficient *L. pneumophila* mutants expressing flagellin (*fliI*) at a MOI of 10 for 8 h. (A) The cultures were stained with anti-AIM2 (red), the cell nuclei were stained with DAPI (cyan) and the NLRC4-GFP puncta is shown in green. The percentage of colocalization of AIM2 with NLRC4-GFP is shown. Images were acquired by multiphoton microscopy with a 63x oil immersion objective and analyzed using ImageJ software. Scale bar, 10µm. (B) Quantification of the number of transduced cells containing NLRC4-GFP in response to WT or *fliI* infection was estimated in *Casp1/11^{-/-}* and *Aim2/Casp1/11^{-/-}* BMDMs. Data show the average ± SD of triplicate wells. NS, not significant, Student's t test. NI, uninfected. Data are presented for one representative experiment of two experiments with similar results.

(TIF)

S4 Fig. AIM2 is not required for caspase-8 activation in response to flagellated *L. pneumophila*.

Bone marrow-derived macrophages (BMDMs) generated from *Casp1/11^{-/-}* and *Aim2/Casp1/11^{-/-}* mice were infected with motility-deficient *L. pneumophila* mutants expressing flagellin (*fliI*) or with flagellin-deficient bacteria (*flaA⁻*) at a MOI of 10 for 8 hours. The activity of caspase-8 was measured using the Caspase-8 Glo Assay. Data show the average ± SD of triplicate wells. *, $P < 0.05$, Student's t test. NI, uninfected. Data are presented for one representative experiment of two experiments with similar results.

(TIF)

S5 Fig. Caspase-8 quantification in the western blot shown in Fig 4A.

Bone marrow-derived macrophages (BMDMs) generated from *Casp1/11^{-/-}* mice were transduced with a retrovirus encoding shRNA sequences to target caspase-8 (Seq1, Seq2) and a non-target shRNA sequence (NT). The silencing was confirmed by western blot analysis (Fig 4A). Cell lysates were separated by SDS-PAGE, blotted and probed with anti-caspase-8 (pro-caspase-8 p55) and anti-α-actin. Immunoblots were analyzed in Image J software and the caspase-8 p55 to α-actin ratio is shown.

(TIF)

S6 Fig. AIM2 is not required for NLRC4-mediated restriction of *L. pneumophila* replication in macrophages.

Bone marrow-derived macrophages (BMDMs) from C57BL/6, *Nlrc4^{-/-}*, *Casp1/11^{-/-}* and *Aim2/Casp1/11^{-/-}* mice were infected with motility-deficient *L. pneumophila* mutants expressing flagellin (*fliI*) at a MOI of 0.015. The cultures were incubated for 24, 48, 72 and 96 hours after infection for CFU determination. Data show the averages ± SD of triplicate wells. *, $P < 0.05$, compared with *Casp1/11^{-/-}* cells. Student's t test. Data are presented for one representative experiment of three experiments with similar results.

(TIF)

S7 Fig. Caspase-8 quantification in the western blot shown in Fig 5E.

Bone marrow-derived macrophages (BMDMs) generated from *Casp1/11^{-/-}* and *Asc/Casp1/11^{-/-}* mice were transduced with a retrovirus encoding shRNA sequences to target caspase-8 (Seq1, Seq2) and a non-target

shRNA sequence (NT). The silencing was confirmed by western blot analysis (Fig 5E). Cell lysates were separated by SDS-PAGE, blotted and probed with anti-caspase-8 (pro-caspase-8 p55) and anti- α -actin. Immunoblots were analyzed in Image J software and the caspase-8 p55 to α -actin ratio is shown.

(TIF)

S8 Fig. AIM2 is not required for NLRC4-mediated restriction of *L. pneumophila* infection in vivo. C57BL/6 (open circles), *Nlrp4*^{-/-} (closed squares), *Casp1/11*^{-/-} (closed triangles), *Aim2*/*Casp1/11*^{-/-} (open diamond) and *Asc/Casp1/11*^{-/-} (closed triangles) mice were infected intranasally with 1×10^5 motility-deficient *L. pneumophila* mutants expressing flagellin (*fliI*). The mice were euthanatized at 4 and 48 hours after infection. Dilutions of the lung homogenates were added to charcoal-yeast extract agar plates for colony-forming unit determination. Each dot represents a single animal, and the horizontal lines represent averages. *, $P < 0.05$, Student's t test. NS, not significant.

(TIF)

S9 Fig. The NLRC4/ASC/caspase-8 inflammasome is important for pore formation and cell death independently of caspase-1/11. Bone marrow-derived macrophages (BMDMs) were generated from C57BL/6, *Casp1/11*^{-/-} and *Asc/Casp1/11*^{-/-} mice and infected with motility-deficient *L. pneumophila* mutants expressing flagellin (*fliI*) or with flagellin-deficient bacteria (*flaA*) at a MOI of 10. (A-C) Pore formation was assessed fluorometrically in real time by the uptake of propidium iodide. RFU (%) represents the percentage of RFU estimated with cells lysed with Triton X-100. (D) After 8 hours the LDH release was measured using the CytoTox 96 LDH-release kit. The LDH release (%) represents the percentage of LDH released compared with cells lysed with Triton X-100. Data show the average $\pm$ SD of triplicate wells. *, $P < 0.05$, Student's t test. NS, not significant; RFU, relative fluorescence units; NI, uninfected. Data are presented for one representative experiment of five (A-C) and two (D) experiments with similar results.

(TIF)

S10 Fig. IL-1 β is not efficiently produced by *Casp1/11*^{-/-} and *Asc/Casp1/11*^{-/-} macrophages. Bone marrow-derived macrophages (BMDMs) were generated from C57BL/6, *Casp1/11*^{-/-} and *Asc/Casp1/11*^{-/-} mice and infected with wild-type *L. pneumophila* (WT Lp), motility-deficient mutants expressing flagellin (*fliI*) or with flagellin-deficient mutants (*flaA*) at a MOI of 10. The production of IL-1 β (A) and IL-12p40 (B) in the tissue culture supernatants was estimated by ELISA at 24 hours after infection. Data show the average $\pm$ SD of triplicate wells. *, $P < 0.05$, Student's t test. nd, not detected; RFU, relative fluorescence units; NI, uninfected. Data are presented for one representative experiment of two experiments with similar results.

(TIF)

S11 Fig. Gasdermin-D is important for pore formation in C57BL/6 but not in *Casp1/11*^{-/-} macrophages. BMDMs generated from C57BL/6 (A-D) and *Casp1/11*^{-/-} (E-H) mice were transduced with a retrovirus encoding shRNA sequence to target Gasdermin D (GSDMD) (Seq1) and a non-target shRNA sequence (NT). Transduced cells were infected with wild-type *L. pneumophila* (WT Lp) (B and F), motility-deficient mutants expressing flagellin (*fliI*) (C and G) or with flagellin-deficient mutants (*flaA*) (D and H) at a MOI of 10. Pore formation was assessed fluorometrically in real time by the uptake of propidium iodide. The RFU (%) represents the percentage of RFU compared with cells lysed with Triton X-100. Data show the average $\pm$ SD of triplicate wells. RFU, relative fluorescence units; NI, uninfected. Data are presented for one representative experiment of two experiments with similar results.

(TIF)

Acknowledgments

We are grateful to Maira Nakamura, Catarina Horta, Leticia Corsi and Victoria dos Santos for their technical assistance. We also thank Roberta Ribeiro Rosales for technical assistance in the institutional facility of the Multiphoton Microscopy. We thank Richard Flavell (Yale University) for providing the *Casp1/11^{-/-}* mice and Vishva Dixit (Genentech) for providing the *Casp11^{-/-}* and *Casp1^{-/-}Casp11^{tg}* mice.

Author Contributions

Conceptualization: Danielle P. A. Mascarenhas, Dario S. Zamboni.

Data curation: Danielle P. A. Mascarenhas.

Formal analysis: Danielle P. A. Mascarenhas, Daiane M. Cerqueira, Marcelo S. F. Pereira.

Funding acquisition: Dario S. Zamboni.

Investigation: Danielle P. A. Mascarenhas, Daiane M. Cerqueira, Marcelo S. F. Pereira, Fernanda V. S. Castanheira, Talita D. Fernandes, Grazielle Z. Manin, Larissa D. Cunha.

Methodology: Danielle P. A. Mascarenhas, Grazielle Z. Manin, Larissa D. Cunha.

Project administration: Dario S. Zamboni.

Resources: Dario S. Zamboni.

Supervision: Dario S. Zamboni.

Validation: Danielle P. A. Mascarenhas, Daiane M. Cerqueira, Marcelo S. F. Pereira.

Visualization: Danielle P. A. Mascarenhas, Dario S. Zamboni.

Writing – original draft: Danielle P. A. Mascarenhas, Dario S. Zamboni.

Writing – review & editing: Danielle P. A. Mascarenhas, Dario S. Zamboni.

References

1. Fraser DW, Tsai TR, Orenstein W, Parkin WE, Beecham HJ, et al. (1977) Legionnaires' disease: description of an epidemic of pneumonia. *N Engl J Med* 297: 1189–1197. <https://doi.org/10.1056/NEJM197712012972201> PMID: 335244
2. Horwitz MA (1983) The Legionnaires' disease bacterium (*Legionella pneumophila*) inhibits phagosome-lysosome fusion in human monocytes. *J Exp Med* 158: 2108–2126. PMID: 6644240
3. McDade JE, Shepard CC, Fraser DW, Tsai TR, Redus MA, et al. (1977) Legionnaires' disease: isolation of a bacterium and demonstration of its role in other respiratory disease. *N Engl J Med* 297: 1197–1203. <https://doi.org/10.1056/NEJM197712012972202> PMID: 335245
4. Newton HJ, Ang DK, van Driel IR, Hartland EL (2010) Molecular pathogenesis of infections caused by *Legionella pneumophila*. *Clin Microbiol Rev* 23: 274–298. <https://doi.org/10.1128/CMR.00052-09> PMID: 20375353
5. Khodr A, Kay E, Gomez-Valero L, Ginevra C, Doublet P, et al. (2016) Molecular epidemiology, phylogeny and evolution of *Legionella*. *Infect Genet Evol* 43: 108–122. <https://doi.org/10.1016/j.meegid.2016.04.033> PMID: 27180896
6. Berger KH, Isberg RR (1993) Two distinct defects in intracellular growth complemented by a single genetic locus in *Legionella pneumophila*. *Mol Microbiol* 7: 7–19. PMID: 8382332
7. Berger KH, Merriam JJ, Isberg RR (1994) Altered intracellular targeting properties associated with mutations in the *Legionella pneumophila* dotA gene. *Mol Microbiol* 14: 809–822. PMID: 7891566
8. Isberg RR, O'Connor TJ, Heidtman M (2009) The *Legionella pneumophila* replication vacuole: making a cosy niche inside host cells. *Nat Rev Microbiol* 7: 13–24. <https://doi.org/10.1038/nrmicro1967> PMID: 19011659

9. Roy CR, Berger KH, Isberg RR (1998) Legionella pneumophila DotA protein is required for early phagosome trafficking decisions that occur within minutes of bacterial uptake. *Mol Microbiol* 28: 663–674. PMID: [9632267](#)
10. Swanson MS, Hammer BK (2000) Legionella pneumophila pathogenesis: a fateful journey from amoebae to macrophages. *Annu Rev Microbiol* 54: 567–613. <https://doi.org/10.1146/annurev.micro.54.1.567> PMID: [11018138](#)
11. Yamamoto Y, Klein TW, Newton CA, Widen R, Friedman H (1988) Growth of Legionella pneumophila in thioglycolate-elicited peritoneal macrophages from A/J mice. *Infect Immun* 56: 370–375. PMID: [3257460](#)
12. Beckers MC, Yoshida S, Morgan K, Skamene E, Gros P (1995) Natural resistance to infection with Legionella pneumophila: chromosomal localization of the Lgn1 susceptibility gene. *Mamm Genome* 6: 540–545. PMID: [8589525](#)
13. Dietrich WF, Damron DM, Isberg RR, Lander ES, Swanson MS (1995) Lgn1, a gene that determines susceptibility to Legionella pneumophila, maps to mouse chromosome 13. *Genomics* 26: 443–450. PMID: [7607666](#)
14. Diez E, Yaraghi Z, MacKenzie A, Gros P (2000) The neuronal apoptosis inhibitory protein (Naip) is expressed in macrophages and is modulated after phagocytosis and during intracellular infection with Legionella pneumophila. *J Immunol* 164: 1470–1477. PMID: [10640764](#)
15. Diez E, Lee SH, Gauthier S, Yaraghi Z, Tremblay M, et al. (2003) Birc1e is the gene within the Lgn1 locus associated with resistance to Legionella pneumophila. *Nat Genet* 33: 55–60. <https://doi.org/10.1038/ng1065> PMID: [12483212](#)
16. Wright EK, Goodart SA, Growney JD, Hadinoto V, Endrizzi MG, et al. (2003) Naip5 affects host susceptibility to the intracellular pathogen Legionella pneumophila. *Curr Biol* 13: 27–36. PMID: [12526741](#)
17. Amer A, Franchi L, Kanneganti TD, Body-Malapel M, Ozoren N, et al. (2006) Regulation of Legionella phagosome maturation and infection through flagellin and host Ipaf. *J Biol Chem* 281: 35217–35223. <https://doi.org/10.1074/jbc.M604933200> PMID: [16984919](#)
18. Molofsky AB, Byrne BG, Whitfield NN, Madigan CA, Fuse ET, et al. (2006) Cytosolic recognition of flagellin by mouse macrophages restricts Legionella pneumophila infection. *J Exp Med* 203: 1093–1104. <https://doi.org/10.1084/jem.20051659> PMID: [16606669](#)
19. Ren T, Zamboni DS, Roy CR, Dietrich WF, Vance RE (2006) Flagellin-deficient Legionella mutants evade caspase-1- and Naip5-mediated macrophage immunity. *PLoS Pathog* 2: e18. <https://doi.org/10.1371/journal.ppat.0020018> PMID: [16552444](#)
20. Zamboni DS, Kobayashi KS, Kohlsdorf T, Ogura Y, Long EM, et al. (2006) The Birc1e cytosolic pattern-recognition receptor contributes to the detection and control of Legionella pneumophila infection. *Nat Immunol* 7: 318–325. <https://doi.org/10.1038/ni1305> PMID: [16444259](#)
21. Case CL, Shin S, Roy CR (2009) Asc and Ipaf Inflammasomes direct distinct pathways for caspase-1 activation in response to Legionella pneumophila. *Infect Immun* 77: 1981–1991. <https://doi.org/10.1128/IAI.01382-08> PMID: [19237518](#)
22. Lightfield KL, Persson J, Brubaker SW, Witte CE, von Moltke J, et al. (2008) Critical function for Naip5 in inflammasome activation by a conserved carboxy-terminal domain of flagellin. *Nat Immunol* 9: 1171–1178. <https://doi.org/10.1038/ni.1646> PMID: [18724372](#)
23. Miao EA, Leaf IA, Treuting PM, Mao DP, Dors M, et al. (2010) Caspase-1-induced pyroptosis is an innate immune effector mechanism against intracellular bacteria. *Nat Immunol* 11: 1136–1142. <https://doi.org/10.1038/ni.1960> PMID: [21057511](#)
24. Silveira TN, Zamboni DS (2010) Pore formation triggered by Legionella spp. is an Nlr4 inflammasome-dependent host cell response that precedes pyroptosis. *Infect Immun* 78: 1403–1413. <https://doi.org/10.1128/IAI.00905-09> PMID: [20048047](#)
25. Casson CN, Copenhaver AM, Zwack EE, Nguyen HT, Strowig T, et al. (2013) Caspase-11 activation in response to bacterial secretion systems that access the host cytosol. *PLoS Pathog* 9: e1003400. <https://doi.org/10.1371/journal.ppat.1003400> PMID: [23762026](#)
26. Jorgensen I, Lopez JP, Laufer SA, Miao EA (2016) IL-1 β , IL-18, and eicosanoids promote neutrophil recruitment to pore-induced intracellular traps following pyroptosis. *Eur J Immunol*.
27. LaRock CN, Cookson BT (2013) Burning down the house: cellular actions during pyroptosis. *PLoS Pathog* 9: e1003793. <https://doi.org/10.1371/journal.ppat.1003793> PMID: [24367258](#)
28. Mascarenhas DP, Pereira MS, Manin GZ, Hori JI, Zamboni DS (2015) Interleukin 1 receptor-driven neutrophil recruitment accounts to MyD88-dependent pulmonary clearance of legionella pneumophila infection in vivo. *J Infect Dis* 211: 322–330. <https://doi.org/10.1093/infdis/jiu430> PMID: [25104770](#)

29. Pereira MS, Marques GG, Dellama JE, Zamboni DS (2011) The Nlrp4 Inflammasome Contributes to Restriction of Pulmonary Infection by Flagellated *Legionella* spp. that Trigger Pyroptosis. *Front Microbiol* 2: 33. <https://doi.org/10.3389/fmicb.2011.00033> PMID: 21687424
30. Case CL, Roy CR (2011) Asc modulates the function of NLRC4 in response to infection of macrophages by *Legionella pneumophila*. *MBio* 2.
31. Hornung V, Ablasser A, Charrel-Dennis M, Bauernfeind F, Horvath G, et al. (2009) AIM2 recognizes cytosolic dsDNA and forms a caspase-1-activating inflammasome with ASC. *Nature* 458: 514–518. <https://doi.org/10.1038/nature07725> PMID: 19158675
32. Man SM, Tourlomousis P, Hopkins L, Monie TP, Fitzgerald KA, et al. (2013) Salmonella infection induces recruitment of Caspase-8 to the inflammasome to modulate IL-1 β production. *J Immunol* 191: 5239–5246. <https://doi.org/10.4049/jimmunol.1301581> PMID: 24123685
33. Pierini R, Juruj C, Perret M, Jones CL, Mangeot P, et al. (2012) AIM2/ASC triggers caspase-8-dependent apoptosis in Francisella-infected caspase-1-deficient macrophages. *Cell Death Differ* 19: 1709–1721. <https://doi.org/10.1038/cdd.2012.51> PMID: 22555457
34. Sagulenko V, Thygesen SJ, Sester DP, Idris A, Cridland JA, et al. (2013) AIM2 and NLRP3 inflammasomes activate both apoptotic and pyroptotic death pathways via ASC. *Cell Death Differ* 20: 1149–1160. <https://doi.org/10.1038/cdd.2013.37> PMID: 23645208
35. Cerqueira DM, Pereira MS, Silva AL, Cunha LD, Zamboni DS (2015) Caspase-1 but Not Caspase-11 Is Required for NLRC4-Mediated Pyroptosis and Restriction of Infection by Flagellated *Legionella* Species in Mouse Macrophages and In Vivo. *J Immunol* 195: 2303–2311. <https://doi.org/10.4049/jimmunol.1501223> PMID: 26232428
36. Abdelaziz DH, Gavrilin MA, Akhter A, Caution K, Kotrange S, et al. (2011) Apoptosis-associated speck-like protein (ASC) controls *Legionella pneumophila* infection in human monocytes. *J Biol Chem* 286: 3203–3208. <https://doi.org/10.1074/jbc.M110.197681> PMID: 21097506
37. Abdelaziz DH, Gavrilin MA, Akhter A, Caution K, Kotrange S, et al. (2011) Asc-dependent and independent mechanisms contribute to restriction of legionella pneumophila infection in murine macrophages. *Front Microbiol* 2: 18. <https://doi.org/10.3389/fmicb.2011.00018> PMID: 21713115
38. Pereira MS, Morgantetti GF, Massis LM, Horta CV, Hori JI, et al. (2011) Activation of NLRC4 by flagellated bacteria triggers caspase-1-dependent and -independent responses to restrict *Legionella pneumophila* replication in macrophages and in vivo. *J Immunol* 187: 6447–6455. <https://doi.org/10.4049/jimmunol.1003784> PMID: 22079982
39. Merriam JJ, Mathur R, Maxfield-Boumil R, Isberg RR (1997) Analysis of the *Legionella pneumophila* flil gene: intracellular growth of a defined mutant defective for flagellum biosynthesis. *Infect Immun* 65: 2497–2501. PMID: 9169800
40. Franchi L, Amer A, Body-Malapel M, Kanneganti TD, Ozoren N, et al. (2006) Cytosolic flagellin requires Ipaf for activation of caspase-1 and interleukin 1 β in salmonella-infected macrophages. *Nat Immunol* 7: 576–582. <https://doi.org/10.1038/ni1346> PMID: 16648852
41. Stutz A, Horvath GL, Monks BG, Latz E (2013) ASC speck formation as a readout for inflammasome activation. *Methods Mol Biol* 1040: 91–101. https://doi.org/10.1007/978-1-62703-523-1_8 PMID: 23852599
42. Masumoto J, Taniguchi S, Ayukawa K, Sarvotham H, Kishino T, et al. (1999) ASC, a novel 22-kDa protein, aggregates during apoptosis of human promyelocytic leukemia HL-60 cells. *J Biol Chem* 274: 33835–33838. PMID: 10567338
43. Fernandes-Alnemri T, Wu J, Yu JW, Datta P, Miller B, et al. (2007) The pyroptosome: a supramolecular assembly of ASC dimers mediating inflammatory cell death via caspase-1 activation. *Cell Death Differ* 14: 1590–1604. <https://doi.org/10.1038/sj.cdd.4402194> PMID: 17599095
44. Hasegawa M, Imamura R, Kinoshita T, Matsumoto N, Masumoto J, et al. (2005) ASC-mediated NF- κ B activation leading to interleukin-8 production requires caspase-8 and is inhibited by CLARP. *J Biol Chem* 280: 15122–15130. <https://doi.org/10.1074/jbc.M412284200> PMID: 15701651
45. Broz P, von Moltke J, Jones JW, Vance RE, Monack DM (2010) Differential requirement for Caspase-1 autoproteolysis in pathogen-induced cell death and cytokine processing. *Cell Host Microbe* 8: 471–483. <https://doi.org/10.1016/j.chom.2010.11.007> PMID: 21147462
46. Vajjhala PR, Lu A, Brown DL, Pang SW, Sagulenko V, et al. (2015) The Inflammasome Adaptor ASC Induces Pro-caspase-8 Death Effector Domain Filaments. *J Biol Chem* 290: 29217–29230. <https://doi.org/10.1074/jbc.M115.687731> PMID: 26468282
47. Masumoto J, Dowds TA, Schaner P, Chen FF, Ogura Y, et al. (2003) ASC is an activating adaptor for NF- κ B and caspase-8-dependent apoptosis. *Biochem Biophys Res Commun* 303: 69–73. PMID: 12646168

48. Akhter A, Gavrilin MA, Frantz L, Washington S, Ditty C, et al. (2009) Caspase-7 activation by the Nlrc4/Ipafl inflammasome restricts *Legionella pneumophila* infection. *PLoS Pathog* 5: e1000361. <https://doi.org/10.1371/journal.ppat.1000361> PMID: 19343209
49. Man SM, Hopkins LJ, Nugent E, Cox S, Gluck IM, et al. (2014) Inflammasome activation causes dual recruitment of NLRC4 and NLRP3 to the same macromolecular complex. *Proc Natl Acad Sci U S A* 111: 7403–7408. <https://doi.org/10.1073/pnas.1402911111> PMID: 24803432
50. Motani K, Kushiya H, Imamura R, Kinoshita T, Nishiuchi T, et al. (2011) Caspase-1 protein induces apoptosis-associated speck-like protein containing a caspase recruitment domain (ASC)-mediated necrosis independently of its catalytic activity. *J Biol Chem* 286: 33963–33972. <https://doi.org/10.1074/jbc.M111.286823> PMID: 21832064
51. Kumar Y, Radha V, Swarup G (2010) Interaction with Sug1 enables Ipafl ubiquitination leading to caspase 8 activation and cell death. *Biochem J* 427: 91–104. <https://doi.org/10.1042/BJ20091349> PMID: 20085538
52. Muruve DA, Petrilli V, Zaiss AK, White LR, Clark SA, et al. (2008) The inflammasome recognizes cytosolic microbial and host DNA and triggers an innate immune response. *Nature* 452: 103–107. <https://doi.org/10.1038/nature06664> PMID: 18288107
53. Roberts TL, Idris A, Dunn JA, Kelly GM, Burnton CM, et al. (2009) HIN-200 proteins regulate caspase activation in response to foreign cytoplasmic DNA. *Science* 323: 1057–1060. <https://doi.org/10.1126/science.1169841> PMID: 19131592
54. Burckstummer T, Baumann C, Bluml S, Dixit E, Durnberger G, et al. (2009) An orthogonal proteomic-genomic screen identifies AIM2 as a cytoplasmic DNA sensor for the inflammasome. *Nat Immunol* 10: 266–272. <https://doi.org/10.1038/ni.1702> PMID: 19158679
55. Fernandes-Alnemri T, Yu JW, Datta P, Wu J, Alnemri ES (2009) AIM2 activates the inflammasome and cell death in response to cytoplasmic DNA. *Nature* 458: 509–513. <https://doi.org/10.1038/nature07710> PMID: 19158676
56. Aachoui Y, Leaf IA, Hagar JA, Fontana MF, Campos CG, et al. (2013) Caspase-11 protects against bacteria that escape the vacuole. *Science* 339: 975–978. <https://doi.org/10.1126/science.1230751> PMID: 23348507
57. Ge J, Gong YN, Xu Y, Shao F (2012) Preventing bacterial DNA release and absent in melanoma 2 inflammasome activation by a *Legionella* effector functioning in membrane trafficking. *Proc Natl Acad Sci U S A* 109: 6193–6198. <https://doi.org/10.1073/pnas.1117490109> PMID: 22474394
58. Pierini R, Perret M, Djebali S, Juruj C, Michallet MC, et al. (2013) ASC controls IFN- γ levels in an IL-18-dependent manner in caspase-1-deficient mice infected with *Francisella novicida*. *J Immunol* 191: 3847–3857. <https://doi.org/10.4049/jimmunol.1203326> PMID: 23975862
59. Kaiser WJ, Upton JW, Long AB, Livingston-Rosanoff D, Daley-Bauer LP, et al. (2011) RIP3 mediates the embryonic lethality of caspase-8-deficient mice. *Nature* 471: 368–372. <https://doi.org/10.1038/nature09857> PMID: 21368762
60. Varfolomeev EE, Schuchmann M, Luria V, Chiannikulchai N, Beckmann JS, et al. (1998) Targeted disruption of the mouse Caspase 8 gene ablates cell death induction by the TNF receptors, Fas/Apo1, and DR3 and is lethal prenatally. *Immunity* 9: 267–276. PMID: 9729047
61. Case CL, Kohler LJ, Lima JB, Strowig T, de Zoete MR, et al. (2013) Caspase-11 stimulates rapid flagellin-independent pyroptosis in response to *Legionella pneumophila*. *Proc Natl Acad Sci U S A* 110: 1851–1856. <https://doi.org/10.1073/pnas.1211521110> PMID: 23307811
62. Fink SL, Cookson BT (2006) Caspase-1-dependent pore formation during pyroptosis leads to osmotic lysis of infected host macrophages. *Cell Microbiol* 8: 1812–1825. <https://doi.org/10.1111/j.1462-5822.2006.00751.x> PMID: 16824040
63. Talanian RV, Quinlan C, Trautz S, Hackett MC, Mankovich JA, et al. (1997) Substrate specificities of caspase family proteases. *J Biol Chem* 272: 9677–9682. PMID: 9092497
64. Thornberry NA, Rano TA, Peterson EP, Rasper DM, Timkey T, et al. (1997) A combinatorial approach defines specificities of members of the caspase family and granzyme B. Functional relationships established for key mediators of apoptosis. *J Biol Chem* 272: 17907–17911. PMID: 9218414
65. Garcia-Calvo M, Peterson EP, Leiting B, Ruel R, Nicholson DW, et al. (1998) Inhibition of human caspases by peptide-based and macromolecular inhibitors. *J Biol Chem* 273: 32608–32613. PMID: 9829999
66. Kayagaki N, Warming S, Lamkanfi M, Vande Walle L, Louie S, et al. (2011) Non-canonical inflammasome activation targets caspase-11. *Nature* 479: 117–121. <https://doi.org/10.1038/nature10558> PMID: 22002608

67. Kayagaki N, Stowe IB, Lee BL, O'Rourke K, Anderson K, et al. (2015) Caspase-11 cleaves gasdermin D for non-canonical inflammasome signalling. *Nature* 526: 666–671. <https://doi.org/10.1038/nature15541> PMID: 26375259
68. Shi J, Zhao Y, Wang K, Shi X, Wang Y, et al. (2015) Cleavage of GSDMD by inflammatory caspases determines pyroptotic cell death. *Nature* 526: 660–665. <https://doi.org/10.1038/nature15514> PMID: 26375003
69. Kofoed EM, Vance RE (2011) Innate immune recognition of bacterial ligands by NAIPs determines inflammasome specificity. *Nature* 477: 592–595. <https://doi.org/10.1038/nature10394> PMID: 21874021
70. Zhao Y, Yang J, Shi J, Gong YN, Lu Q, et al. (2011) The NLRC4 inflammasome receptors for bacterial flagellin and type III secretion apparatus. *Nature* 477: 596–600. <https://doi.org/10.1038/nature10510> PMID: 21918512
71. Coers J, Vance RE, Fontana MF, Dietrich WF (2007) Restriction of *Legionella pneumophila* growth in macrophages requires the concerted action of cytokine and Naip5/lpafl signalling pathways. *Cell Microbiol* 9: 2344–2357. <https://doi.org/10.1111/j.1462-5822.2007.00963.x> PMID: 17506816
72. Fortier A, Doiron K, Saleh M, Grinstein S, Gros P (2009) Restriction of *Legionella pneumophila* replication in macrophages requires concerted action of the transcriptional regulators Irf1 and Irf8 and nod-like receptors Naip5 and Nlrc4. *Infect Immun* 77: 4794–4805. <https://doi.org/10.1128/IAI.01546-08> PMID: 19720760
73. Rauch I, Deets KA, Ji DX, von Moltke J, Tentorey JL, et al. (2017) NAIP-NLRC4 Inflammasomes Coordinate Intestinal Epithelial Cell Expulsion with Eicosanoid and IL-18 Release via Activation of Caspase-1 and -8. *Immunity* 46: 649–659. <https://doi.org/10.1016/j.immuni.2017.03.016> PMID: 28410991
74. Rogers C, Fernandes-Alnemri T, Mayes L, Alnemri D, Cingolani G, et al. (2017) Cleavage of DFNA5 by caspase-3 during apoptosis mediates progression to secondary necrotic/pyroptotic cell death. *Nat Commun* 8: 14128. <https://doi.org/10.1038/ncomms14128> PMID: 28045099
75. Wang Y, Gao W, Shi X, Ding J, Liu W, et al. (2017) Chemotherapy drugs induce pyroptosis through caspase-3 cleavage of a Gasdermin. *Nature*.
76. Feeley JC, Gibson RJ, Gorman GW, Langford NC, Rasheed JK, et al. (1979) Charcoal-yeast extract agar: primary isolation medium for *Legionella pneumophila*. *J Clin Microbiol* 10: 437–441. PMID: 393713
77. Marim FM, Silveira TN, Lima DS Jr., Zamboni DS (2010) A method for generation of bone marrow-derived macrophages from cryopreserved mouse bone marrow cells. *PLoS One* 5: e15263. <https://doi.org/10.1371/journal.pone.0015263> PMID: 21179419
78. Milstein S, Nguyen M, Meyers R, de Fougères A (2013) Measuring RNAi knockdown using qPCR. *Methods Enzymol* 533: 57–77. <https://doi.org/10.1016/B978-0-12-420067-8.00006-4> PMID: 24182918
79. Lara-Tejero M, Sutterwala FS, Ogura Y, Grant EP, Bertin J, et al. (2006) Role of the caspase-1 inflammasome in *Salmonella typhimurium* pathogenesis. *J Exp Med* 203: 1407–1412. <https://doi.org/10.1084/jem.20060206> PMID: 16717117
80. Kuida K, Lippke JA, Ku G, Harding MW, Livingston DJ, et al. (1995) Altered cytokine export and apoptosis in mice deficient in interleukin-1 beta converting enzyme. *Science* 267: 2000–2003. PMID: 7535475
81. Sutterwala FS, Ogura Y, Szczepanik M, Lara-Tejero M, Lichtenberger GS, et al. (2006) Critical role for NALP3/CIA1/Cryopyrin in innate and adaptive immunity through its regulation of caspase-1. *Immunity* 24: 317–327. <https://doi.org/10.1016/j.immuni.2006.02.004> PMID: 16546100

Supplementary Fig.1 Mascarenhas et al.

Supplementary Fig.2. Mascarenhas et al.

Supplementary Fig.3. Mascarenhas et al.

Supplementary Fig.4. Mascarenhas et al.

Supplementary Fig.5. Mascarenhas et al.

Supplementary Fig.6. Mascarenhas et al.

Supplementary Fig.7. Mascarenhas et al.

Supplementary Fig. 8. Mascarenhas et al.

Supplementary Fig.9 Mascarenhas et al.

Supplementary Fig.10 Mascarenhas et al.

Supplementary Fig.11 Mascarenhas et al.