Sistemática de Cyclanthaceae

Eduardo da Silva Leal

São Paulo
2018

Eduardo da Silva Leal

Sistemática de Cyclanthaceae

Systematics of Cyclanthaceae

São Paulo

2018

Eduardo da Silva Leal

Sistemática de Cyclanthaceae

Systematics of Cyclanthaceae

Tese apresentada ao Instituto de
Biociências da Universidade de São
Paulo, para obtenção do Título de
Doutor em Ciências, na Área de
Botânica.

Orientador: Prof. Dr. Renato de
Mello-Silva

São Paulo

2018

Resumo

Cyclanthaceae possui cerca de 230 espécies. São ervas terrestres, hemiepífitas, lianas ou epífitas distribuídas na região Neotropical, se estendendo do sul do México a Mata Atlântica do sul do Brasil. A família é dividida em duas subfamílias de acordo com a morfologia da inflorescência: Cyclanthoideae com flores estaminadas e pistiladas dispostas em ciclos alternos, incluindo apenas *Cyclanthus*; e Carludovicoideae com uma flor pistilada envolvida por quatro flores estaminadas e compreende o restante dos gêneros. Até o presente, estudos filogenéticos baseados somente em dados morfológicos, e com amostragem a nível genérico, não foram suficientes para clarificar suas relações. Além disso, um dos dois grandes grupos em Carludovicoideae emergiu como parafilético. Neste estudo, usamos dados morfológicos e de sete marcadores moleculares, cinco de cloroplastos e dois nucleares, com uma ampla amostragem à nível específico, para propor uma hipótese filogenética para Cyclanthaceae. Nosso resultado recuperou a família como monofilética e corroborou sua divisão em duas subfamílias. Os gêneros pertencentes a Carludovicoideae emergiram em três clados: (i) o gênero *Carludovica*; (ii) o clado *Asplundia* que contém *Asplundia*, *Dicranopygium* e *Schultesiophytum*; e (iii) o clado *Evodianthus* formado por *Chorigyne*, *Dianthoveus*, *Evodianthus*, *Ludovia*, *Sphaeradenia*, *Stelesstylis* e *Thoracocarpus*. Todos os gêneros previamente circunscritos emergiram como monofiléticos. O grupo *Sphaeradenia*, composto por *Chorigyne*, *Ludovia*, *Sphaeradenia* e *Stelesstylis* emerge como monofilético fortemente sustentado por caracteres morfológicos e moleculares. Para maximizar a informação filogenética, bem como facilitar a identificação, o gênero *Dianthoveus* é sinonimizado em *Evodianthus*. No capítulo 2 uma chave de identificação para todas as espécies de Cyclanthaceae ocorrentes na Mata Atlântica é fornecida. As espécies endêmicas de *Asplundia* são revisadas, resultando em três táxons sinonimizados sob *A. brachypus*, e uma nova espécie, *A. orthostigma*, é descrita. No capítulo 3 é apresentado um índice taxonômico para todas as espécies já descritas de Cyclanthaceae. Nós indicamos 225 nomes aceitos e 60 sinônimos. Quatro lectótipos e dois neótipos são designados para quatro nomes. Para mais quatro nomes, a tipificação é completada, com a designação do segundo passo do lectótipo. Finalmente, doze nomes são considerados como novos sinônimos.

Abstract.

Cyclanthaceae has ca. 230 species. They are terrestrial herbs, hemiephytic, climbing or epiphytic distributed in the Neotropics, ranging from the South Mexico to Southward Atlantic forest. The family is divided into two subfamilies based on the inflorescence morphology: Cyclanthoideae with staminate and pistillate flowers arranged in alternate whorls, includes only *Cyclanthus*; and Carludovicoideae which has a pistillate flower with four surrounding staminate flower, encompass the others genera. Nowadays, phylogenetic studies based on morphological data in generic level were insufficient to clarify its relationship. Furthermore, one of the groups of Carludovicoideae emerged as paraphyletic. In this study we combined morphological and seven molecular markers, five from plastid and two nuclear, in a broad sampling in species level to recover a consistent phylogenetic hypothesis to Cyclanthaceae. We reconstructed a monophyletic Cyclanthaceae with two subfamilies. Genera belonging to Carludovicoideae emerged in three clades: (i) the genus *Carludovica*, (ii) the Asplundia Clade which includes *Asplundia*, *Dicranopygium* and *Schultesiophytum*; and (iii) the Evodianthus Clade comprising *Chorigyne*, *Dianthoveus*, *Evodianthus*, *Ludovia*, *Sphaeradenia*, *Stelestylis* and *Thoracocarpus*. All previous known genera were recovered as monophyletic. The Sphaeradenia Group, which includes *Chorigyne*, *Ludovia*, *Sphaeradenia* and *Stelestylis*, emerged as monophyletic strongly supported by morphological and molecular characters. To maximize the phylogenetic outcome and make the identification easier *Dianthoveus* is synonymized under *Evodianthus*. In the Chapter 2 an identification key for all Cyclanthaceae species in Atlantic forest is provided. The endemic species of *Asplundia* are revisited resulting in three taxa synonymized under *A. brachypus*, and *A. orthostigma* newly described. In the Chapter 3 a taxonomic index for all Cyclanthaceae species is provided. We highlighted 225 accepted names and 60 synonyms heterotypics. Four lectotypes and two neotypes are designated. The typification is complemented to four taxa more with second-step lectotypification. Finally, twelve names are placed as synonyms.

General Introduction

Cyclanthaceae are terrestrial herbs, hemiephytic, climbing or epiphytic plants distributed in the Neotropics, ranging from the South Mexico to Southward Atlantic forest (Harling & al., 1998). They are highly diversified in the biogeographic Choco lowland forest and cloudy forest from tropical Andes, reaching up to 3000 m elevation (Harling, 1958). Cyclanthaceae is recognized by its inflorescence with alternate cycles of staminate and pistillate flowers or a pistillate flower with four surrounding staminate flowers, and by the almost ubiquitous bifid leave. The inflorescence is sustained by an unbranched peduncle supporting bracts subtending the inflorescence in a cluster or loose arrangement. Flowers are arranged in a spiralled or ring continuum in Cyclanthoideae making impossible to recognize an isolated flower. Conversely, Carludovicoideae has actinomorphic or zigomorphic staminate flower with numerous stamens and absent perianth (Sajo & al., 2014) while the pistillate flower has tetramerous perianth arranged in one whorl. Four filiform and long staminoid are opposite to the tepals in this flower. Four stigma can be attached to the ovary summit or elevated by a style.

Cyclanthaceae encompass ca. 225 species distributed into 12 genera: *Asplundia* Harling (97 spp.), *Carludovica* Ruiz & Pav. (4 spp.), *Chorigyne* R. Erikss. (7 spp.), *Cyclanthus* Poit. ex A. Rich. (2 sp.), *Dianthoveus* Hammel & G.J. Wilder (1 sp.), *Dicranopygium* Harling (51 spp.), *Evodianthus* Oersted (1 sp.), *Ludovia* Brongn. (3 spp.), *Schultesiophytum* Harling (1 sp.), *Sphaeradenia* Harling (52 spp.), *Stelestylis* Drude (4 spp.) and *Thoracocarpus* Harling (1 sp.). Cyclanthaceae is represented by 29 species and nine genera in Brazil, specially occurring in Amazonic (21 spp.) and Atlantic (10 spp.) forest with seven endemic species at the latter (Leal, 2018). Amazonian-Atlantic forest disjunction was recorded only for *Evodianthus funifer* and *Thoracocarpus bissectus* whereas *Asplundia gardneri* occurs in the *brejos* and riparian forest in the Central Brazil.

The positioning of Cyclanthaceae in suprafamiliar categories has been subject of changes in successive classifications. Bentham & Hooker (1883) grouped it with Aroideae, Lemnaceae, Pandaneae and Typhaceae, in series Nudiflorae (category corresponding to order). Cronquist (1981, 1988) positioned it alone in order Cyclanthales, in subclass Arecidae, assuming the family's affinities with Araceae, Arecaceae, Lemnaceae and Pandanaceae. Dahlgren & al. (1985) also positioned

Cyclanthaceae alone in Cyclanthales, the sole order in superorder Cyclanthiflorae. More recent classifications, based on molecular phylogenies, positioned Cyclanthaceae in order Pandanales, along with Pandanaceae, Stemonaceae, Triuridaceae and Velloziaceae (APG IV, 2016 and precedents). The order has been well supported in recent phylogenetic studies based on morphological (Rudall & Bateman, 2006) and molecular data (Chase & al. 2000, Caddick & al. 2002, Davis & al. 2004, Mennes & al. 2013). Cyclanthaceae is monophyletic and distinct from the remaining Pandanales by the following synapomorphies: leaves with open sheath and monoic inflorescences, one pistillate flower surrounded by four staminate flowers or alternating whorls of pistillate and staminate flowers.

The first infrafamiliar classification for Cyclanthaceae was proposed by Drude (1881), who recognized two tribes: Carludoviceae and Cyclantheae. He also proposed three sections in tribe Carludoviceae: Anomalae, Bifidae and Palmatae, according to leaf blade shape and division. Later, Harling (1954a, b) made profound taxonomic changes in genus *Carludovica*, splitting it into *Asplundia*, *Dicranopygium* and *Sphaeradenia*. Posteriorly, Harling, in his large monograph about the family (1958), elevated the tribes defined by Drude (1881) to subfamilies, taking into account mainly inflorescence structure. He characterized subfamily Cyclanthoideae, at the time containing only *Cyclanthus bipartitus* Poit. ex A.Rich., by the spadix containing alternating whorls of staminate and pistillate flowers, and subfamily Carludovicoideae with the remaining genera, characterized by groups of four staminate flowers surrounding one pistillate flower. Harling (1958) distributed the genera of Carludovicoideae in two groups not formally classified: a cohesive *Sphaeradenia* group, and *Asplundia* group, both inconsistent and presented as an attempt to position genera.

Eriksson (1994) performed a new morphological analysis of Cyclanthaceae, using sixty-three morphological and anatomical characters. *Sphaeradenia* group, comprehending *Chorigyne*, previously described by Eriksson (1989), *Ludovia*, *Sphaeradenia* and *Stelestylis*, emerged as monophyletic, supported by the synapomorphies plants predominantly epiphytes, distichous phyllotaxy, coriaceous leaf blade, apical or subapical placentation and endosperm with thickened cell walls. On the other hand, *Asplundia* group emerged as paraphyletic and affinities among genera remained uncertain, suggesting the need of analysis with more characters (Eriksson, 1994). Additionally, according to the hypothesis presented by Eriksson

(1994) *Asplundia*, the richest genus, is paraphyletic.

In this perspective, this thesis is the first to reconstruct the phylogeny of Cyclanthaceae with a broad sampling combining morphological and molecular data. Moreover, this thesis aims to contribute to the understanding to the species from Atlantic forest as well as provide a full taxonomic index of Cyclanthaceae. Thus, this study is presented in three chapters as follow:

Chapter 1. Phylogeny of Cyclanthaceae (the Panama-hat family) based on molecular and morphological data. This Chapter provide a phylogenetic reconstruction of the family based on seven molecular markers and 42 morphological and anatomical characters. All known genera were recovered as monophyletic according to Maximum Parsimony and Bayesian analysis.

Chapter 2. Revisiting the Atlantic forest cyclants with a key for the species and a revision of the endemic clade of *Asplundia*. Identification key for seven species from the Atlantic forest is provided. Three names are newly synonymized under *Asplundia brachypus*, and a new species is described.

Chapter 3. Nomenclator botanicus of the Cyclanthaceae. A total of 225 species are currently accepted. Four lectotypes and two neotypes are designated. Second-step lectotypification is given for four names. Twelve names are newly synonymized.

Figura 1. Cladograms showing the affinity between the genera of Cyclanthaceae. The Sphaeradenia group is evidenced by dashed ellipses and the Asplundia group by dashed rectangles. A-Harling (1958), the dashed line indicates the uncertainty of the position of Schultesiophytum. B-Hammel and Wilder (1989) indicating the two groups as monophyletic and C. Eriksson (1994) showing the non-monophyletic Asplundia group and monophyletic group Sphaeradenia, as well as two lineages for the genus *Asplundia*.

Final Considerations

We newly presented a consistent phylogeny of Cyclanthaceae combining molecular and morphological datasets in species level and broad sampling. We recovered a monophyletic Cyclanthaceae using combined and total-evidence trees in both Parsimony and Bayesian analysis. Morphological analysis showed that Cyclanthaceae is widely supported by longitudinal incisions in the blade (except *Ludovia*), basal bulb in the filament base, and connate pistillate flowers. Also the inflorescence morphology is unique in the family.

A new perspective of the phylogenetic relationship of Cyclanthaceae was provided. All genera and the subdivision of Cyclanthoideae and Carludovicoideae were supported, however one of the informal categories of Carludovicoideae was recovered non-monophyletic. The subfamily Carludovicoideae is divided into three lineages: i) *Carludovica*; ii) *Asplundia*, *Dicranopygium* and *Schultesiophytum*, named Asplundia Clade; and iii) *Dianthoveus*, *Evodianthus*, *Thoracocarpus* and the Sphaeradenia group, named Evodianthus Clade. Our results recovered a monophyletic *Evodianthus* including *Dianthoveus* supported by styloid idioblasts in the mesophyll oriented in all directions, terminal inflorescence, bracts of the same size, filament absent or inconspicuous, basal bulbs absent, psilate and inaperturate pollen. Sphaeradenia group was recovered as monophyletic strongly supported by molecular and morphological dataset, however, morphological circumscription of its genera should be reevaluated in future studies.

The morphological circumscription of *Asplundia* from Atlantic forest were reassessed and reduced to three endemic species including the species newly described. The exceptional sampling, a robust phylogeny, and the nomenclator botanicus brings an avenue for future studies on biogeography, diversification and evolution of Cyclanthaceae as well as it can give insights on further systematics studies on the poorly known and rich species genera *Asplundia* e *Dicranopygium*.

Literature cited

APG IV. 2016. An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG IV. *Bot. J. Linn. Soc.* 181: 1--20. <https://doi.org/10.1111/boj.12385>

- Bentham, G. & Hooker, J. D. 1883. *Genera Plantarum*, vol. 3. London: Lovell Reeve & Co.
- Chase, M.W., Stevenson, D.W. & Loconte, H. 1995. Monocot systematics: A combined analyses. Pp. 685--730 in Rudall, P.J., Cribb, P.J., Cutler, D.F. & Humphries, C.J. (eds). *Monocotyledons: Systematics and Evolution*. London: Royal Botanic Gardens, Kew.
- Cronquist, A. 1981. *An integrated system of classification of flowering plants*. New York: Columbia University Press.
- Cronquist, A. 1988. *The evolution and classification of flowering plants. Second edition*. New York: New York Botanical Garden Press.
- Dahlgren, R.M.T., Clifford, H.T., & Yeo, P.F. 1985. *The families of the monocotyledons. Structure, evolution, and taxonomy*. Berlin: Springer-Verlag.
- Drude, O. 1881. Cyclanthaceae. Pp. 226--250 in: Martius, C.F.P & Eichler, A.W. (eds.), *Flora Brasiliensis*, vol. 3 pars 2. Lipzig: Fridrich Fleischer.
- Eriksson, R. 1989. *Chorigyne*, a new genus of the Cyclanthaceae from Central America. *Nordic J. Bot.* 9: 31--45. <https://doi.org/10.1111/j.1756-1051.1989.tb00978.x>
- Eriksson, R. 1994. Phylogeny of the *Cyclanthaceae*. *Pl. Syst. Evol.* 190: 31--47. <https://doi.org/10.1007/BF00937857>
- Hammel, B.E. & Wilder, G.J. 1989. *Dianthoveus*, a new genus of Cyclanthaceae. *Ann. Missouri Bot. Gard.* 76: 112--123. <https://doi.org/10.2307/2399344>
- Harling, G. 1954a. *Sphaeradenia* , a new genus of the Cyclanthaceae. *Acta Horti Berg.* 17: 1--6.
- Harling, G. 1954b. Taxonomical studies in the genus *Carludovica* R. & P. *Acta Horti Berg.* 17: 39--45.
- Harling, G. 1958. Monograph of the Cyclanthaceae. *Acta Horti Berg.* 18: 1--428.
- Harling, G., Wilder, G.J. & Eriksson, R. 1998. Cyclanthaceae. Pp. 202--215 in: Kubitzki, K. (ed.), *The families and genera of vascular plants. Flowering plants Monocotyledons-Lilianaes (except Orchidaceae)*, Vol. 3. Berlin: Springer-Verlag.
- Leal, E.S. 2018. Cyclanthaceae in Flora do Brasil 2020 em construção. Jardim Botânico do Riode Janeiro.

<http://floradobrasil.jbrj.gov.br/reflora/floradobrasil/FB98>. Acesso em: 20 Jul. 2018.

- Rudall, P.J. & Bateman, R.M. 2006. Morphological phylogenetic analysis of Pandanales: Testing contrasting hypotheses of floral evolution. *Syst. Bot.* 31: 223--238. <https://doi.org/10.1600/036364406777585766>
- Sajo, M.G., Lombardi, J.A., Forzza, R.C. & Rudall, P.J. 2014. Comparative anatomy of reproductive structures in Cyclanthaceae (Pandanales). *Int. J. Pl. Sci.* 175: 814--827. <https://doi.org/10.1086/676817>
- Soltis, D., Soltis, P., Endress, P., Chase, M.W., Manchester, S., Judd, W., Majure, L. & Mavrodiev, E. 2017. *Phylogeny and Evolution of the Angiosperms: Revised and Updated Edition*. Chicago: University of Chicago Press.
- Stevenson, D.W. & Loconte, H. 1995. Cladistic analysis of monocot families. Pp. 543--578 in: Rudall, P.J., Cribb, P.J., Cutler, D.F. & Humphries, C.J. (eds.), *Monocotyledons: Systematics and Evolution*. London: Royal Botanic Gardens, Kew.