Universidade de São Paulo
Instituto de Física

Medida de elétrons provenientes do decaimento de
hadrons que contêm quarks pesados em colisões pp a
a $\sqrt{s} = 13$ TeV e elétrons provenientes de hadrons
que contêm quarks *beauty* em colisões Pb-Pb a
 $\sqrt{s_{NN}} = 5.02$ TeV pelo ALICE, no LHC

Camila de Conti

Orientador: Prof. Dr. Marcelo Gameiro Munhoz

Tese de doutorado apresentada ao Instituto de Física
da Universidade de São Paulo, como requisito parcial
para a obtenção do título de Doutor(a) em Ciências.

Banca Examinadora:

Prof. Dr. Marcelo Gameiro Munhoz (IFUSP)

Prof. Dr. Luiz Vitor de Souza Filho (IFSC/USP)

Prof. Dr. Fernando S. Navarra (IFUSP)

Profa. Dra. Sandra S. Padula (IFT/UNESP)

Profa. Dra. Carla Göbel Burlamaqui de Mello (PUC/RJ)

São Paulo

2018

University of São Paulo
Physics Institute

Measurement of electrons from open heavy-flavor
hadron decays in pp at $\sqrt{s} = 13$ TeV and electrons
from beauty-hadron decays in Pb-Pb at $\sqrt{s_{\text{NN}}} = 5.02$
TeV collisions by ALICE at LHC

Camila de Conti

Supervisor: Prof. Dr. Marcelo Gameiro Munhoz

Thesis submitted to the Physics Institute of the University of São Paulo in partial fulfillment of the requirements for the degree of Doctor of Science.

Examining Committee:

Prof. Dr. Marcelo Gameiro Munhoz (IFUSP)

Prof. Dr. Luiz Vitor de Souza Filho (IFSC/USP)

Prof. Dr. Fernando S. Navarra (IFUSP)

Profa. Dra. Sandra S. Padula (IFT/UNESP)

Profa. Dra. Carla Göbel Burlamaqui de Mello (PUC/RJ)

São Paulo

2018

Acknowledgments

I am deeply thankful for the opportunity I had to do this work, and for the so many wonderful things I have learned during these years. My special thanks to Marcelo, for trusting this work to me and for giving me chances to grow. My thanks to all the friends from HEPIC@IFUSP, for being around day-by-day, sharing discussions, difficult questions and life. Many thanks to Silvia for gladly receiving me at GSI twice. My special thanks to Andrea, who was my guide since the beginning of the first analysis and with who I have learned a huge part of what I know about this work. Thanks for all the support, for the strong sincerity, and for the constant trust. Your pragmatism and dedication are inspiring. My special thanks also to Martin, for the so many discussions about the beauty analysis, and for all the valuable clarifications and knowledge he transmitted to me with so much patience and care. My work on beauty is not so complete as the work he did in run 1, but I really did the best I could in the timescale given to me, and every conversation with him was extremely necessary and precious for my accomplishment. Thanks to all the HFE group and to the HF conveners, for making the working environment so pleasant, fruitful, and for giving a lot of support. Thanks to all the friends at GSI, that made me feel at home.

This study was financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES) - Finance Code 001.

*”Oh, how I wish I could shut up like a telescope!
I think I could, if only I knew how to begin.’
For, you see, so many out-of-the-way things had
happened lately, that Alice had begun to think that
very few things indeed were really impossible.”*

Alice in Wonderland

Abstract

In this thesis it is performed the measurement of electrons from heavy-flavor hadron decays, with the use of the ALICE detector, at LHC. Two analysis are performed. The first one consists in the measurement of the invariant cross section of electrons from open heavy-flavor (*charm* or *beauty*) hadron decays in pp collisions at $\sqrt{s} = 13$ TeV, the largest center of mass energy generated by LHC up to now, in the transverse momentum interval $0.5 < p_T < 4$ GeV/ c . The experimental cross-section is compared to the pQCD-based prediction given by FONLL, and it sits on the upper edge of the theoretical uncertainty band over the full momentum range. The ratios between this experimental result and the equivalent measurements performed at lower center of mass energy have the potential to bring constraints to the FONLL calculation. The second analysis consists in the measurement of the nuclear modification factor, R_{AA} , of electrons from beauty-hadron decays in the 10% most central Pb-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, in the interval $2 < p_T < 8$ GeV/ c . Electrons from beauty-hadron decays are extracted from the inclusive electron yield via fits to the impact parameter distribution using templates of electrons from different sources, obtained from Monte Carlo simulations. This measurement allows to probe the energy loss of *beauty* quarks when traversing the Quark-Gluon-Plasma (QGP), a color-deconfined medium produced in relativistic heavy-ion collisions. A suppression of about 40-50% is observed on the R_{AA} for $p_T > 3$ GeV/ c and it is consistent, over the full momentum range and within experimental uncertainties, with the prediction of pQCD-based models that describe a mass-dependent energy loss of the heavy-quarks while traversing the QGP, interacting with its constituents via radiative and collisional processes. This analysis brings significant improvement of the experimental uncertainties with respect to the equivalent measurement performed with the run 1 data and consists in an important step in the study of the mass-dependence of the energy loss of heavy quarks inside the QGP at low p_T .

Key-words: Experimental physics; Quark-Gluon-Plasma; Heavy quarks; Beauty quarks; Heavy-flavor electrons.

Resumo

Nessa tese é feita a medida de elétrons provenientes do decaimento de hadrons que contêm quarks pesados (*charm* ou *beauty*), com o uso dos detectores do experimento ALICE, no LHC. Duas análises são realizadas. A primeira consiste na medida da seção de choque invariante de elétrons provenientes do decaimento de hadrons que contêm quarks *charm* ou *beauty* em colisões pp a $\sqrt{s} = 13$ TeV, a maior energia de centro de massa gerada pelo LHC até então, na região de momento transversal $0.5 < p_T < 4$ GeV/ c . Esse resultado é comparado com a previsão dada pelo FONLL (cálculo baseado em QCD perturbativa) e se situa no limite superior da incerteza teórica, sobre toda a faixa de momento. As razões entre este resultado experimental e medidas equivalentes realizadas em mais baixa energia podem trazer informações importantes para tornar o cálculo teórico mais preciso. A segunda análise consiste na medida do fator de modificação nuclear, R_{AA} , de elétrons provenientes do decaimento de hadrons que contêm quarks *beauty*, em colisões centrais de Pb-Pb a $\sqrt{s_{NN}} = 5.02$ TeV, no intervalo $2 < p_T < 8$ GeV/ c . Elétrons provenientes de hadrons que contêm quarks *beauty* são extraídos do conjunto de elétrons inclusivos a partir do ajuste da distribuição do parâmetro de impacto com templates de elétrons de diferentes fontes, extraídos de simulações de Monte Carlo. Essa medida permite estudar a perda de energia dos quarks *beauty* ao atravessarem o Plasma de Quarks e Glúons (QGP), produzido em colisões entre íons pesados em altas energias. É observada uma supressão de $\approx 40\text{-}50\%$ no R_{AA} medido para $p_T > 3$ GeV/ c e esta é consistente, sobre toda a faixa de p_T , com a previsão de modelos baseados em pQCD, que descrevem a perda de energia dos quarks pesados via processos elásticos e inelásticos com os constituintes do QGP, de maneira dependente da massa. Este resultado possui incertezas experimentais significativamente menores com relação à medida realizada com dados do run 1 e consiste num importante passo no estudo da dependência em massa da perda de energia dos quarks pesados no QGP em baixo p_T .

Palavras-chave: Física experimental; Plasma de quarks e glúons; Quarks pesados; Quarks beauty; Elétrons de heavy-flavor.

Contents

1	Introduction	9
2	Heavy-Ion Collisions to Study Strong Interactions	13
3	Heavy-flavor as a Tool to Study the Quark-Gluon Plasma	21
3.1	Heavy-Flavor Production in the "QCD Vacuum"	24
3.2	Nuclear Modification Factor	27
3.2.1	Heavy-Flavor Production in pA Collisions: Testing Cold Nuclear Matter Effects	29
3.2.2	Heavy-Flavor Production in AA Collisions: Probing the QGP Related Effects	33
3.3	Transport Models of Heavy-Quarks Inside QGP	42
4	Experimental Apparatus	47
5	Measurement of Electrons from Open Heavy-Flavor Hadron Decays in pp Collisions at $\sqrt{s} = 13$ TeV	59
5.1	General Strategy	59
5.2	Data Set and Event Selection	61
5.3	Monte Carlo Samples	63
5.4	Track Selection	63
5.5	Selection of Electron Candidates	65
5.6	Efficiency of Track and PID Cuts	67
5.7	Remaining Hadron Contamination	69

5.8	Non Heavy-Flavor Electron Identification	73
5.8.1	Non Heavy-Flavor Electron Tagging Efficiency	77
5.9	Normalization and FONLL Prediction	83
5.9.1	FONLL	85
5.10	Systematic Uncertainties	86
5.10.1	Systematic Uncertainties Coming from the Track Quality Cuts and PID Cuts	88
5.10.2	Systematic Uncertainties due to the Subtraction of Electrons from non Heavy-Flavor Sources	91
5.10.3	Systematic Uncertainties due to π^0 and η Weights	93
5.10.4	Systematic Uncertainties from Other Sources	94
5.11	Results and Discussion	101
6	Measurement of Electrons from Beauty Hadron Decays in Central Pb- Pb Collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV	107
6.1	General Strategy	107
6.2	Data Set and Event Selection	110
6.3	Monte Carlo Samples	111
6.4	Centrality Selection	111
6.5	Track Selection and its Efficiency	114
6.6	Selection of Electron Candidates in Data and its Efficiency	118
6.7	Selection of Electrons in Monte Carlo: Building the Templates	122
6.8	The Maximum Likelihood Template Fit	126
6.8.1	The Likelihood Function	126
6.8.2	The Maximum Likelihood Estimation	128
6.8.3	Bins Without Counts from the Templates	131
6.9	Uncertainty Estimation	132
6.10	Corrections on the Templates and Systematic Uncertainties	135
6.10.1	Mean and Resolution Correction	135
6.10.2	D meson p_{T} -spectrum Correction	137

6.10.3 Beauty p_T -spectrum Correction	141
6.10.4 Hadron Contamination	146
6.10.5 Correction of the Proportion Between Charmed Baryons and Mesons	148
6.10.6 Other Systematic Uncertainties and Summary	151
6.11 Results and Discussion	153
6.11.1 pp Reference	154
6.11.2 R_{AA} of Electrons from Beauty Hadron Decays in Pb-Pb at $\sqrt{s_{NN}} =$ 5.02 TeV	156
7 Summary	165
A Other Evidences of the QGP Formation	167
A.1 Azimutal Anisotropy	167
A.2 Jet Suppression	169
A.3 Quarkonia Sequential Melting	171
B Pseudo-Rapidity and Rapidity	175
C Charm and Beauty Contribution to Open Heavy-Flavor Electrons Spec- trum	177
D Gamma Conversion in the Detector Material	179
E ITS Quality Cuts	181
F Fit Instability due to DCA Bin Size	183
Referências Bibliográficas	185

Chapter 1

Introduction

The Large Hadron Collider (LHC) is the largest particle accelerator built so far and it represents the big efforts from humanity to understand nature in a very detailed level. LHC belongs to the European Organization for Nuclear Research (CERN) and it is composed by four main experiments dedicated to different physics purposes. A Large Ion Collider Experiment (ALICE) [1] is the one dedicated to study relativistic heavy ion collisions. A phase transition from the color neutral matter to a color deconfined medium is predicted by Quantum Chromodynamics (QCD) [2][3] when the matter is submitted to conditions of high energy density and temperature as the ones created during relativistic heavy-ion collisions [4]. This new state of matter is called Quark-Gluon Plasma (QGP) [2] [3] and experimental evidences of its creation were already observed in the past in heavy-ion collisions performed at the Relativistic Heavy Ion Collider (RHIC) [5] [6] [7] [8]. ALICE has the purpose of investigating the properties of this color-deconfined strongly-interacting medium by studying the debris of heavy-ion collisions performed at the unprecedented energies generated by LHC. In ALICE experiment, proton-proton (pp) and proton-nucleus (pA) collisions are also studied. Although the QGP is not expected to be produced in these collisions, they are an important baseline for the study of medium-induced effects on the particle production in heavy-ion collisions [9]. In addition, results from pp collisions allow to test theoretical predictions of particle production in the absence of any nuclear effect [9] [10] [11] [12] [13].

Hadrons composed of heavy quarks are important observables for the purpose of

studying the QGP properties. Heavy quarks are produced mainly by hard processes that takes place in the first instants of the collision, before the QGP formation, and consequently they experience the full evolution of this medium, interacting with its constituents via inelastic and elastic processes [14].

In this work, two different analysis were performed within the ALICE experiment in the heavy-flavor sector, using data from LHC run 2. The first analysis consists in the measurement of the production cross section of electrons from heavy-flavor hadron decays in pp collisions with a center of mass energy of $\sqrt{s} = 13$ TeV, the highest energy generated by LHC up to now. This measurement allows to test the perturbative QCD (pQCD) predictions in this new energy domain and can bring constrains to the current theoretical understanding of heavy-flavor production in the absence nuclear effects [15]. Within the ALICE experiment, due to its lower magnetic field, this measurement can be extended to lower transverse momentum (p_T) with respect to heavy-flavor measurements from other LHC experiments. In particular, this production cross section compared to the ones measured at lower center of mass energies, mainly at low momentum, can provide important constrains for the gluon PDF [15], that is known with limited precision at low Bjorken- x [16] [2]. The second analysis consists in the measurement of electrons from beauty-hadron decays in Pb-Pb collisions performed at a center of mass energy per nucleon pair of $\sqrt{s_{NN}} = 5.02$ TeV. In this environment, the QGP is produced and it modifies the measured spectrum of electrons from beauty-hadron decays with respect to what is measured in pp collisions. The nuclear modification factor, R_{AA} , is an important quantity to characterize the QGP-induced spectrum modification. The R_{AA} of electrons from beauty-hadron decays is measured and compared to theoretical models, what allows to test the pQCD prediction of mass dependent energy loss of heavy quarks while traversing the QGP and can help to constrain microscopic properties of this medium. Both analysis are performed in the pseudo-rapidity interval $|\eta| < 0.8$ and low p_T regions.

This thesis is organized as the following: in [chapter 2](#) some basic principles of QCD are introduced and the heavy-ion collisions are presented as the experimental method used to produce the QGP in laboratory, making possible its experimental study. In [chapter 3](#) the measurement of heavy-flavor particles is introduced as an important tool to

explore properties of the QGP. Recent measurements of heavy-flavor particles in pp, pA and AA collisions are presented and discussed, together with the current open questions and limitations related to them. In this context, the two analysis of this thesis are motivated. The nuclear modification factor R_{AA} is introduced and its sensitivity to different ingredients is discussed. In [chapter 4](#) the ALICE detector is briefly presented and the specific sub-detectors used in this work are described in more detail. In [chapter 5](#) the measurement of electrons from heavy-flavor hadron decays in pp at $\sqrt{s} = 13$ TeV collisions is described in detail and the invariant cross section of these electrons is compared to a pQCD prediction and discussed. In [chapter 6](#) the measurement of electrons from beauty-hadron decays in Pb-Pb at $\sqrt{s_{NN}} = 5.02$ TeV collisions is presented in detail and discussed. The R_{AA} of electrons from beauty-hadron decays is obtained and compared to pQCD predictions. In [chapter 7](#) the main results and conclusions of the two analysis performed in this thesis are summarized.

Chapter 2

Heavy-Ion Collisions to Study Strong Interactions

Particle physics studies the fundamental constituents of matter and their interactions. The Standard Model is the most up-to-date theory that classify the elementary particles and describe their interactions. This model describes three of the four fundamental interactions of nature: the electromagnetic, weak and strong interactions, not including the gravitational interaction. According to the Standard Model, matter is made up of quarks and leptons. The interaction between particles is carried out by gauge bosons, like the photon (for the electromagnetic interaction) and the gluon (for the strong interaction).

The strong nuclear interaction is described by Quantum Chromodynamics (QCD) [2] [3]. According to QCD, each quark carry one color charge. Unlike the theory of Quantum Electrodynamics (QED), where the electric charge is unique and can manifest itself in positive or negative signs (charge and anti-charge), the color charge of QCD has three different possibilities, named red, green and blue (with the corresponding anti-charges antired, antigreen and antiblue). Also differently from QED, the gauge boson in QCD also carries charge: the color of the gluon is a combination of a color charge and an anti-color charge. This results in eight linearly independent possible color states for the gluons, that are represented by the Gell-Mann matrices [3].

Neither quarks or gluons were never observed as free particles: they are always grouped into color neutral hadrons. In other words, quarks and gluons in the ordi-

nary matter are always arranged in a group of quarks, anti-quarks and gluons whose colors combined result in "white" particles. This particular property of particles that carry color charge is called **confinement**. The confinement is a consequence of the behavior of the coupling constant α_s [17], the QCD parameter that gives the strength of the interaction between two color carrying particles as function of their relative distance. The strong coupling increases proportionally to the distance between the two color carrying particles, and when the distance becomes sufficiently large (and by consequence the color field becomes too strong), the corresponding anti-particle of each of the two particles can be created from the vacuum: in this way, the attempt to isolate one color charge from the other originates a system of two color neutral objects.

Nevertheless, QCD predicts situations in which the confinement of the color charge can "vanish" asymptotically. This occurs when instead of trying to take two color charges apart from each other, they are approximated to each other up to very small distances: this can occur in the limits of very high density and/or temperature of the nuclear matter [3]. Calculations in perturbative QCD can determine the evolution of the strong coupling as a function of the transferred momentum Q of an interaction [17]. The dependence of the strong coupling of the interaction on the energy scale Q involved in the process is a property of QCD referred to as the **running of the coupling**. A first order (LO) perturbative QCD (pQCD) calculation (valid at large Q) gives Equation 2.1:

$$\alpha_s(Q) = \frac{12\pi}{(33 - 2n_f)\ln(\frac{Q^2}{\Lambda_{\text{QCD}}^2})}, \quad (2.1)$$

where:

- $\Lambda_{\text{QCD}} \approx 200$ MeV is the non-perturbative scale of QCD, that indicates where the perturbatively-defined coupling would diverge (when $Q \rightarrow \Lambda_{\text{QCD}}$, $\alpha_s(Q) \rightarrow \infty$).
- $n_f = 6$ is the number of flavors in the theory;

The running of the coupling shown in Equation 2.1 indicates the **asymptotic freedom** of the color charge: the larger the momentum transfer Q involved in the processes, the weaker the strong coupling gets, going asymptotically to zero. In the limit of very

large Q , the quarks and gluons behave as almost free particles, and are able to move along larger distances than the typical size of a nucleon. This is called **deconfinement** [18] of the color charge. This deconfined (although still strongly interacting) medium produced in large Q processes is named **Quark-Gluon Plasma (QGP)**.

The analytical perturbative solution of QCD fails in the regime of large $\alpha_s(Q)$, what prevents it to give a quantitative description of confinement. Lattice QCD [19] is a well-established non-perturbative approach to solve the QCD, formulated on a discrete space-time lattice. Lattice QCD calculations predict that a phase transition from baryonic matter to the QGP happens around temperature $T \approx 170$ MeV. However, these calculations are limited, for numerical reasons, and cannot predict a complete phase diagram for quark matter: instead Lattice QCD covers mainly the region of $\mu_B < T$ [19], where μ_B is baryonic potential (quantity proportional to matter density). So $T \approx 170$ MeV is the phase transition temperature predicted by Lattice QCD for matter at low μ_B .

Despite the fact that the phase diagram of quark matter is not currently well known experimentally or theoretically, a commonly conjectured diagram is considered, as shown at Figure 2.1. The phase diagram of QCD indicates different temperature and matter density regimes in which the baryonic matter can undergo color deconfinement. At low T and low μ_B , the baryonic matter consists of colorless hadrons. When the temperature and/or density of the system is increased, after a certain limit, the QGP is created.

The possibility to study the QGP experimentally allows to test and improve the current theoretical understanding of the strong interaction, and can help to build a more solid knowledge about the QCD phase-diagram. The study of the QGP is also important to understand the evolution of our own universe, since the current cosmological model predicts that the universe was in a QGP state (in the limit of low μ_B and high T) during few microseconds after its creation by the Big Bang [21] [22]. The evolution of the QGP and how it transits to hadronic matter can have an important influence on how the universe is today, and it can help to explain fundamental phenomena, like the matter-antimatter asymmetry [23].

The only experimentally known way of creating the QGP in laboratory is by per-

Figure 2.1: Conjectured phase diagram of nuclear matter, where the hadronic and the QGP phases are expected in different temperature T and baryonic potential μ_B . The regions of the phase space probed by the LHC and RHIC colliders are indicated. The transition between hadronic and QGP phases are indicated by the continuous line where a first order phase transition is expected. For small μ_B a second order phase transition (cross-over) is expected, indicated by dashed lines. A possible critical point is also indicated. Figure taken from [20].

forming collisions of heavy nuclei at very high energies. During the first instants of a heavy-ion collision, the baryonic matter achieves conditions of T and μ_B in which quarks and gluons can undergo deconfinement. In heavy-ion collisions the deconfinement is possible also because the region of space in which the matter is under high energy density is much larger than a typical hadron size, in a way that the quarks and gluons can move through a much larger region of space with respect to what they do in ordinary matter [24].

A heavy-ion program has been created with the purpose of producing the QGP in laboratory, with the aim of studying its properties and exploring the QCD phase diagram. Large colliders have been built, like the Relativistic Heavy Ion Collider (RHIC) at Brookhaven National Laboratory and the Large Hadron Collider (LHC) at CERN,

in which collisions of heavy nuclei, like gold (Au) and lead (Pb), can take place. At RHIC, the Au-Au collisions are generated with center of mass energy per nucleon pair ($\sqrt{s_{NN}}$) ranging from few to hundreds of GeV, while at LHC Pb-Pb collisions can take place with energies up to $\sqrt{s_{NN}} = 5.5$ TeV, allowing the creation of the QGP in different regions of the QCD phase diagram, as indicated in [Figure 2.1](#). In particular, the LHC covers a region of very low μ_B and high temperatures, overlapping the region where lattice QCD can give predictions and covering the conditions that are believed to be the ones of the early universe during the first instants after the Big Bang. The environment corresponding to the other extreme case in which the QGP can be formed, low T and high μ_B regime, is expected to be found inside neutron stars, and is not experimentally accessible up to now.

The temporal evolution of a high energy heavy-ion collision can be thought as a sequence of steps [\[3\]](#) [\[21\]](#) as described below. The existence of the QGP is an intermediate step of this evolution. [Figure 2.2](#) represents the time evolution of a heavy-ion collision in which the QGP is created.

Figure 2.2: Diagram representing the time evolution of a heavy-ion collision in the scenario where the QGP is formed. Indicated in the figure are the instant of thermalization τ_0 , the critical temperature T_c below which the QGP is hadronized, and the temperatures T_{ch} and T_{fo} in which the chemical and kinetic freezeout takes place, respectively.

Two nuclei traveling in opposite directions with extremely high velocities (close to c), collide with each other (at time $t = 0$ and position $z = 0$) and a large fraction of their energy is deposited in the overlap region of the collision.

In the first instants of the collision, the system undergoes the so called **pre-equilibrium** phase. In this phase, hard scatterings between partons from the two incoming nuclei take place and the hard probes, that consists of high momentum particles, heavy-quarks, jets and direct photons, are produced.

The continuous interaction between the constituents of the system makes it evolve towards a local **thermal equilibrium**, that is achieved at a time $t \approx \tau_0$. In the case where the energy density and temperature of the system are high enough and go above the critical values (predicted by Lattice QCD) of $\varepsilon_c \approx 1 \text{ GeV}/fm^3$ and $T_c \approx 170 \text{ GeV}$, respectively, the nuclear matter can undergo **color-deconfinement**, and the QGP is formed. The transition from hadronic matter to a Quark-Gluon plasma can happen via different processes at different regions of the phase space, as indicated in [Figure 2.1](#): a first order transition can take place in the region of high μ_B , in which the two states can coexist and the transition from the hadronic state to QGP happens via the absorption of a latent heat. At low μ_B regions, below the critical point, like the ones probed at LHC energies and the ones existing in the first microseconds of the universe, the phase transition is expected, from Lattice QCD, to be of second order (crossover) [\[22\]](#) [\[3\]](#), that is characterized by a continuous evolution of the matter from one state to the other, with no coexistence of different states.

The pressure of this system relatively to the surrounding vacuum induces its collective expansion. The system expands and flows as a relativistic hydrodynamic fluid [\[25\]](#) [\[21\]](#), and the energy density and temperature of the system goes down with time, up to the point where the the nuclear matter can no longer be deconfined ($T < T_c$). At this point, the **hadronization** starts to occur, and the color charges group itself in many different colorless hadrons.

Once the hadrons are formed, they can still have energy to interact inelastically with each other, so the chemical composition (abundances of the different hadron species) of the system can still change. From the moment the hadrons no longer interact inelasti-

cally, the system undergo the so called **chemical freezout** at a temperature T_{ch} , and its chemical composition stops to evolve.

The system continues to expand, and the hadrons can still interact elastically, exchanging energy and momentum. From the moment the particles are too far away from each other for any kind of interaction and momentum exchange, the system undergo the so called **kinetic freezout** at a temperature T_{fo} .

After this stage, the particles reach the detectors. The chemical composition and momentum distribution measured by the detectors are the experimentally available informations for the study of the all phenomena that took place since the first instants of the collision, and, in particular, for the study of the QGP.

Soft and Hard Probes

The experimental study of the QGP produced in relativistic heavy-ion collisions rely on a broad set of different observables. Many properties of this new state of matter can be revealed by the measurement of **soft probes** [26], that consists on light-flavor and low momentum particles. Soft probes allows to access global properties of the QGP by testing theoretical hydrodynamic description of this medium and its evolution, from which quantities like temperature, viscosity, density and expansion velocity can be predicted. On the other hand, the study of **hard probes** [14], that consists on high p_T particles, heavy-flavor particles, quarkonia and jets, allows to resolve the medium constituents (quarks and gluons) and to probe microscopic properties of the QGP. Transport models, that simulate heavy quarks traversing the QGP and interacting elastically and inelastically with the medium constituents, for example, allow to predict (when confronted to experimental measurements) quantities like mean free path and transport coefficients of the QGP.

This work is dedicated to the study of hard probes, and, in particular, of heavy-flavor. In the next chapter it will be presented the motivation to heavy-flavor studies and how the characterization of the QGP can be performed via the measurement of the nuclear modification factor R_{AA} of heavy-flavor particles compared with theoretical calculations. Other hard probe observables, like quarkonia, jets and elliptic flow v_2 of heavy-flavor

particles are discussed in [Appendix A](#).

Chapter 3

Heavy-flavor as a Tool to Study the Quark-Gluon Plasma

From all particles measured in the final state of a heavy-ion collision, the ones related to heavy quarks, charm (c) and beauty (b), are unique observables for the study of the QGP. Due to the large masses of heavy-quarks ($m_c \approx 1.3 \text{ GeV}/c^2$, $m_b \approx 4.2 \text{ GeV}/c^2$), they are produced mainly via the hard parton scatterings that take place in the first instants of the collision, during the pre-equilibrium phase [14]. Consequently, the heavy quarks are expected to be produced at times $t < \tau_0$, where τ_0 is the timescale of the formation of the QGP (Figure 2.2). Because of their early creation, the heavy quarks interact with the QGP during all its evolution, by elastic [27] and inelastic [28] collisions with the quarks and gluons of the medium, what makes them probes that are sensitive to the microscopic properties of the QGP.

Another important feature of heavy quarks is that, since their main production mechanism is the hard scattering of partons, their production cross section can be calculated by perturbative QCD over their full momentum range [29]. This allows the theoretical control of the production of heavy-flavor particles, what makes them a solid reference for the study of heavy-ion collisions.

In order to probe the interaction of heavy quarks with the QGP, the measurement of heavy-flavor particles must be performed. This can be done in two possible scenarios: or the heavy-flavor hadrons are fully reconstructed via an invariant mass analysis of their

hadronic decay products, or the measurement of leptons coming from the decay of the heavy-flavor hadrons is performed.

The heavy-flavor hadrons decay before reaching the detectors. The branching ratios of the most commonly used hadronic decay channels for a charm hadron reconstruction via an invariant mass analysis vary from ≈ 1 to 10 %, depending on the particle specie, as shown in [Table 3.1](#):

Decay channel	Branching Ratio
$D^0 \rightarrow \pi^- K^+$	$BR \approx 3.93\%$
$D^+ \rightarrow K^- \pi^+ \pi^+$	$BR \approx 9.46\%$
$D^{*+} \rightarrow D^0 (\rightarrow K^- \pi^+) \pi^+$	$BR \approx 2.66\%$
$D_s^+ \rightarrow \Phi (\rightarrow K^- K^+) \pi^+$	$BR \approx 2.27\%$
$\Lambda_c^+ \rightarrow p K^- \pi^+$	$BR \approx 6.35\%$
$\Lambda_c^+ \rightarrow p K_s^0 (\rightarrow \pi^+ \pi^-)$	$BR \approx 1.58\%$

Table 3.1: Hadronic decay channel of some charm hadrons and their respective branching ratios, taken from [\[30\]](#).

Since heavy-flavor hadrons have their decay vertex significantly displaced from the primary vertex with respect to light hadrons (due to the larger decay length of the former), topological cuts [\[31\]](#) on the decay vertex are applied in order to enhance the signal of the heavy-flavor hadrons before the invariant mass analysis.

The beauty hadrons have many hadronic decay channels with large branching ratios [\[30\]](#). Nevertheless, they correspond to decays that produce many bodies in the final state, what makes the reconstruction of the beauty hadrons extremely challenging, mainly in face of the high multiplicity environment of a heavy-ion collision. Currently, the full B meson reconstruction is possible via decay channels which are rare, for instance, via the decay channel:

$$B^+ \rightarrow J/\Psi (\rightarrow l^- + l^+) + K^+,$$

that has a branching ratio of $BR \approx 0.006\%$. Therefore, the reconstruction of heavy-flavor hadrons via the hadronic decay channel has the disadvantage of relying in rare signals,

mainly on the beauty sector, that require a big amount of data for a measurement with a good significance.

The semi-leptonic decay channels, on the other hand, produce abundant amount of leptons from both charm and beauty hadrons decays, since:

$$c,b(\rightarrow c) \text{ hadrons} \rightarrow e + X,$$

$$c,b(\rightarrow c) \text{ hadrons} \rightarrow \mu + X$$

have a branching ratio of $BR \approx 10\%$.

Although the semi-leptonic decay channel produce a neutrino that cannot be detected, it is possible to study heavy quark production via the exclusive measurement of the leptons coming from heavy-flavor hadron decays. The disadvantage of this approach is that the kinematic properties of the decay lepton is not so closely related to the original parton ones, what can cause a smearing of the kinematic dependence of the quantities being measured.

Due to the large abundance of leptons from heavy-flavor sources produced in the collisions, heavy quark production will be studied in this work via the measurement of electrons coming from heavy-flavor hadron decays. In ALICE, electrons are measured at mid-rapidity, while the measurement of muons is restricted to forward rapidities only.

Below, in [section 3.1](#), a brief summary of recent heavy-flavor (hadrons and leptons) measurements in minimum bias (MB) proton-proton (pp) collisions will be presented. These pp collisions are an important baseline for the study of the QGP and also allow to test pQCD predictions of particle production. In [section 3.2](#) it will be presented the nuclear modification factor, an experimental tool used to probe modifications on the particle production in proton-nucleus (pA) and nucleus-nucleus (AA) collisions with respect to MB pp collisions. Recent measurements of the nuclear modification factor of heavy-flavor particles in pA and AA collisions will be discussed. During the discussions in these two sections, the two analysis performed in this thesis will be motivated.

The nuclear modification factor of heavy-flavor particles measured in AA collisions, where the QGP is produced, allows to study properties of this medium when confronted to theoretical predictions. Many phenomenological models that describe a heavy-ion collision and the transport of the heavy quarks within the produced QGP are available.

They are a key tool to constrain properties of the QGP and to understand the mechanisms of interaction between the heavy quarks and the color deconfined medium. These models make use of the theoretically calculable heavy quark cross section as input, and then calculate the propagation of these heavy quarks through a color deconfined medium, followed by the hadronization processes and decay of the produced particles, embracing the full evolution of a heavy-ion collision. The comparison between the production of heavy-flavor particles predicted by the models and experimental measurements in heavy-ion collisions allows to test QCD and the QGP description. By the end of this chapter, in [section 3.3](#), the main characteristics of some of these models will be presented.

3.1 Heavy-Flavor Production in the "QCD Vacuum"

Since heavy quarks are produced mainly via hard parton scatterings, the production cross section of heavy-flavor hadrons or their decay products can be described over the full momentum range by perturbative QCD calculations. The pQCD calculations can be performed at next-to-leading order with next-to-leading-log resummation, like the General-Mass Variable-Flavor-Number Scheme (GM-VFNS, [\[32\]](#) [\[33\]](#)) or at Fixed-Order Next-to-LeadingLog (FONLL, [\[34\]](#) [\[35\]](#)) approaches. These pQCD based calculations describe the production cross section of a heavy-flavor particle as a convolution of three terms: i) the Parton Distribution Function (PDF) of the incoming hadrons, ii) the partonic scattering cross section and iii) the fragmentation function, that models the non-perturbative transition of a heavy quark to a given heavy-flavor hadron. This procedure is called factorization approach [\[36\]](#) [\[37\]](#).

In pQCD calculations, the fragmentation function of a heavy quark to a hadron is tuned in order to reproduce results from e^-e^+ experiments [\[38\]](#) [\[39\]](#) [\[40\]](#), under the assumption that they are universal (do not depend on the collision system). FONLL calculations currently does not provide predictions for the production cross section of baryons, while GM-VFNS does, using the quark to baryon fragmentation function $f(c \rightarrow \Lambda_c)$ taken from [\[40\]](#).

Up to now, a large amount of measurements of heavy-flavor particles in pp collisions

have been performed by all LHC experiments in order to test the pQCD prediction of heavy-flavor production in the QCD vacuum in different center of mass energies.

The D^0 meson p_T -differential cross section was measured by ALICE at mid-rapidity down to $p_T = 0$ in pp collisions at $\sqrt{s} = 7$ TeV. As shown in the left panel of Figure 3.1, the measurement is on the upper edge of the FONLL uncertainty band along most of the p_T range. The experimental uncertainties are much smaller than the theoretical ones, that comes mainly from factorization and renormalization scales. On the other hand, GM-VFNS prediction tends to overestimate the data, as shown in the right panel of Figure 3.1.

Figure 3.1: Production cross section of D^0 mesons in pp collisions at $\sqrt{s} = 7$ TeV measured at mid-rapidity by ALICE experiment. The result is compared to FONLL and GM-VFNS pQCD calculations [31].

The cross section of muons from open heavy-flavor hadron decays (muons from the decay of both charm and beauty hadrons), measured at forward rapidity in pp collisions at $\sqrt{s} = 5$ TeV, also sits on the upper edge of FONLL uncertainty band, as shown in Figure 3.2, from $p_T = 2$ to 20 GeV/c. From this comparison to theory, it is possible to see that the open heavy-flavor muon measurement at low p_T ($\lesssim 4$ GeV/c) probes mainly the charm production, while for larger p_T it probes mainly the beauty production.

Figure 3.2: Production cross section of muons from open heavy-flavor hadron decays measured at forward-rapidity by ALICE experiment in pp collisions at $\sqrt{s} = 5.02$ TeV compared to FONLL prediction (the same measurement performed in $\sqrt{s} = 2.76$ TeV is published in [41]).

The comparison between the measurement of electrons from heavy-flavor hadron decays at mid-rapidity and the FONLL prediction will be presented in [section 5.11](#), together with the discussion of one of the measurements performed in this thesis. In this thesis, electrons from open heavy-flavor hadron decays are measured in pp collisions at $\sqrt{s} = 13$ TeV, that is a new and the largest center of mass energy generated by LHC up to now. With this new measurement, it is possible to test the FONLL prediction in a new energy domain, and also, together with the measurements obtained in the past for lower center of mass energies, it is possible to add further constraints to FONLL uncertainties, as will also be discussed in [section 5.11](#).

Differently from what is observed in the measurement of heavy-flavor mesons and leptons presented above, a large deviation between experimental results and theory is observed in the baryon sector at mid-rapidity. The production cross section of Λ_c^+ baryon is systematically underestimated by the pQCD GM-VFNS calculation, as shown in [Figure 3.3](#) [42]. In this same figure the prediction of the Λ_c^+ cross section by the

POWHEG+PYTHIA6 event generator [43] [44] is shown. In both cases, theory subestimates the experimental results by a factor ≈ 10 . Nevertheless, the baryon production at forward rapidity measured by LHCb, presented in the right panel of Figure 3.3, finds a better agreement with GM-VFNS prediction.

Figure 3.3: Production cross section of Λ_c^+ baryon in pp collisions at $\sqrt{s} = 7$ TeV measured at (left) mid-rapidity by ALICE experiment [42] and (right) forward rapidity by LHCb experiment [11]. The results are compared to pQCD GM-VFNS calculation (both) and to prediction from POWHEG+PYTHIA6 event generator (left).

This lack of understanding about the baryon production is one of the current open big questions in the heavy-flavor production sector, and the current results point to a hint of a violation of the universality of the fragmentation function of heavy quarks to baryons.

3.2 Nuclear Modification Factor

Knowledge about the QGP can be acquired by measuring how the existence of this medium modifies the momentum distribution of the heavy-flavor particles. A quantity

suited for this study is the nuclear modification factor R_{AA} , that compares the production spectrum of a particle in Pb-Pb collisions, where the QGP is expected to be produced, with the production spectrum of the same particle in MB pp collisions, at the same center of mass energy.

$$R_{AA} = \frac{dN_{AA}/dp_T}{\langle N_{\text{coll}} \rangle dN_{\text{pp}}/dp_T} = \frac{dN_{AA}/dp_T}{\langle T_{AA} \rangle d\sigma_{\text{pp}}/dp_T}, \quad (3.1)$$

where dN_{AA}/dp_T is the measured p_T spectrum of the particle in the nucleus-nucleus collision, the dN_{pp}/dp_T ($d\sigma_{\text{pp}}/dp_T$) is the measured p_T spectrum (cross section) in MB pp collision and $\langle N_{\text{coll}} \rangle$ ($\langle T_{AA} \rangle$) is the average number of binary collisions (nuclear overlap function) in a nucleus-nucleus collision at a given centrality¹. Both $\langle N_{\text{coll}} \rangle$ and $\langle T_{AA} \rangle$ are obtained from the Glauber model [45] [46] [47] [48] (see also [section 6.4](#)). The relation between $\langle T_{AA} \rangle$ and $\langle N_{\text{coll}} \rangle$ is given by: $N_{\text{coll}} = \sigma_{\text{NN}}^{\text{inel}} \times T_{AA}$, where $\sigma_{\text{NN}}^{\text{inel}}$ is the inelastic nucleon-nucleon cross section at given center of mass energy $\sqrt{s}$.

The rate of hard processes is expected to scale with the average number of binary nucleon-nucleon collisions that occurs in a nucleus-nucleus collision, $\langle N_{\text{coll}} \rangle$ [45] [49] [50] [51]. This so-called binary scaling of hard processes was proven experimentally for the first time via the measurement of direct photon production in Au-Au collisions at $\sqrt{s_{\text{NN}}} = 200 \text{ GeV}/c$ at RHIC [51]. Because of this property, the measurement of the production spectrum of heavy-flavor particles in heavy-ion collisions works as a type of tomography of the QGP: any deviation of the R_{AA} from unity comes from nuclear effects. Nevertheless, the parton interaction with the QGP is not the only possible nuclear effect that can modify the spectrum produced in Pb-Pb with respect to the one produced in pp: **different initial conditions** can alter the parton production cross section in these two different systems, and also the abundance of particles can be modified by **different hadronization mechanisms** that can take place in the scenario where the QGP is formed. **Final stated effects**, like the interaction between particles in the hadron gas

¹The centrality of a collision is a quantity that indicates the level of overlap between the two colliding nuclei. The larger the level of overlap, the more central is the collision. The centrality is defined according to the multiplicity of particles produced in the collision, as will be explained in [section 6.4](#)

phase, can also be different in pp and Pb-Pb collisions. Schematically, the produced spectra in Pb-Pb collision can be thought as a convolution of these many ingredients:

$$\frac{dN_{AA}}{dp_T} = \underset{\text{parton spectra}}{\text{"vacuum"}} \otimes \underset{\text{effects}}{\text{initial-state}} \otimes \underset{\text{with QGP}}{\text{parton interaction}} \otimes \underset{\text{mechanisms}}{\text{(modified?) hadronization}} \otimes \underset{\text{phase}}{\text{hadronic}} \otimes \underset{\text{kinematics}}{\text{decay}} . \quad (3.2)$$

In order to disentangle the different effects that can cause a $R_{AA} \neq 1$ and quantify what comes only from the interaction of the parton with the QGP and what comes from other effects, it is necessary to perform measurements of different observables, that are sensitive to the different ingredients pointed above, in pp, pA and AA collisions.

Below, a brief summary of some recent results regarding heavy-flavor production at LHC in pA and AA collisions are shown. Results from pA collisions constrain initial state effects on the produced heavy-flavor spectra, while results from AA collisions point to modifications on the produced spectra caused by the interaction of heavy quarks with the QGP and by different hadronization mechanisms.

3.2.1 Heavy-Flavor Production in pA Collisions: Testing Cold Nuclear Matter Effects

In collisions where a nucleus is involved, like pA or AA collisions, the particle production can be modified, relatively to MB pp collisions, by initial state effects due to the presence of the nucleus, called Cold Nuclear Matter (CNM) effects.

It is known that the Parton Distribution Function (PDF) of a free nucleon (not bounded in a nucleus) is different from the PDF of a bounded nucleon (that is called nuclear PDF, or nPDF). The difference between PDFs and nPDFs is estimated via experimental measurements together with theoretical calculations. Experimental data from Deep Inelastic Scatterings (DIS) between leptons and nuclei ($l+A$) and leptons and protons ($l+p$) can be compared with the predictions from different pQCD calculations,

that are based on the assumption of a certain PDF for the partons in the different systems and can be tuned to reproduce these experimental results [16]. In Figure 3.4 is shown the nuclear effect on the PDF of a parton i (valence quark, sea quark and gluons, respectively in each panel) predicted by different model calculations. In this notation, x refers to the Bjorken- x [16] [2], that is the fraction of the momentum of the nucleon carried by a parton, Q^2 is the squared of the transferred momentum in the DIS, and $R_i^{\text{Pb}}(x, Q^2)$ is the ratio between the PDF of a bounded nucleon and the PDF of a free nucleon.

Figure 3.4: Ratio $R_i^{\text{Pb}}(x, Q^2)$ between the PDF of a nucleon bounded to a Pb nucleus and the PDF of a free nucleon, obtained from different model calculations. The modifications are shown for valence quarks (left), sea quarks (center) and gluons (right). Figure taken from [16]

The suppression of the nPDF, indicated by $R_i^{\text{Pb}}(x, Q^2) < 1$, in the region of Bjorken- $x \lesssim 0.1$ is called **nuclear shadowing**. At the LHC energies this is the modification of the nPDF that have more influence on the particle production [52] and it can lead to a reduction of heavy-flavor production rates mainly at low p_T in mid-rapidity [16].

Another CNM effect that can be present in a collision involving a nucleus is the energy loss of partons in the cold nuclear matter. Since a nucleus is an extended object,

an incoming parton with momentum k_T can interact elastically with many partons of the colliding nucleus before the hard scattering. This phenomena can modify the momentum spectrum of the produced heavy quarks, and it is called k_T -broadening.

In pA collisions, nuclear effects are expected to be present while the QGP is not expected to be produced. The analogous of R_{AA} (Equation 3.3) for pA collisions is called R_{pA} and it is a quantity suited to measure how much the CNM effects modify the production of a given particle with respect to pp collisions:

$$R_{pA} = \frac{dN_{pA}/dp_T}{A d\sigma_{pp}/dp_T}, \quad (3.3)$$

where A in the denominator refers to the atomic mass number.

The right panel of Figure 3.5 shows the R_{pPb} of D mesons measured by ALICE at mid-rapidity, from $0 < p_T < 35$ GeV/c. The result is comparable to unity within uncertainties, indicating that all cold nuclear matter effects considered together lead to negligible modifications in the production of the D mesons in p-Pb with respect to pp collisions. Some of the theoretical models deviate from the measurement mainly at very low p_T , so the later can be used to constrain the CNM description by these models. In the right panel of Figure 3.5 it is presented the R_{pPb} of electrons from beauty hadron decays measured at mid-rapidity by ALICE [53]. Also via the electron channel, no modification of the spectrum of the heavy-flavor particles is observed within uncertainties. The theoretical predictions, that consider CNM effects and are essentially at unity, describe the experimental result within uncertainties.

Measurement of p-Pb collisions at forward rapidity performed by the LHCb collaboration can access different Bjorken- x regions of the parton PDFs depending on the symmetry of the collision, named forward (p-Pb) or backward (Pb-p). Given that the Bjorken- x can be written as $x = (p_T/\sqrt{s})e^{-y}$, forward collisions (where the particles are measured at positive rapidity y , which is defined with respect to the direction of the proton beam) give access to very low Bjorken- x [2], where the shadowing of the nPDF is stronger. Measurements of the R_{pPb} of D mesons at LHCb result in values comparable to unity in the backward region, while in the forward region a clear deviation from unity

is observed, as shown in Figure 3.6.

The measurements performed in p-Pb collisions guarantee a good control of the initial conditions of the Pb-Pb measurements at different rapidity regions. The general good agreement with theoretical calculations shows that the cold nuclear matter effects, like

Figure 3.5: Nuclear modification factor R_{pPb} of prompt D mesons (left) and electrons from beauty hadron decays [53] (right), both measured at mid-rapidity by ALICE, together with theoretical predictions that includes CNM effects.

Figure 3.6: Nuclear modification factor R_{pPb} of prompt D^0 meson production in the forward (p-Pb) and backward (Pb-p) collisions measured by LHCb collaboration [54].

the PDF modeling, are reasonably well understood. In particular, the data provided by LHCb can be used to constrain the PDF uncertainties in the model calculations, given the significantly smaller uncertainties of the experimental results compared to the model uncertainties.

3.2.2 Heavy-Flavor Production in AA Collisions: Probing the QGP Related Effects

Once the effects of the initial conditions on the produced particle spectra are studied and quantified in pA collisions, the study of the QGP via the R_{AA} measurement can be performed.

A suppression of high p_T partons is expected in Pb-Pb with respect to pp collisions as a consequence of the energy loss of these partons via collisional (elastic) and radiative (inelastic) interactions with the medium constituents while traversing the QGP.

Quarks are expected to lose less energy than gluons while propagating through the QGP, as a consequence of their smaller color-coupling with the medium [55] [56] [57] [58] [59]. QCD also predicts a mass-dependence of the energy loss: heavy-quarks are expected to lose less energy than light quarks via radiative processes, due to the dead-cone effect [60] [61] [62] [63] [64]. The dead-cone-effect is a fundamental prediction of QCD that defines that a quark with mass m_q and energy E_q has its energy loss by radiative processes suppressed within the angle $\theta \lesssim m_q/E_q$ (defined with respect to the direction of its momentum).

Among the theoretical frameworks that describe the energy loss of a heavy-quark in the medium via both radiative and collisional processes is the model by Djordjevic *et al* [65]. Figure 3.7 shows how this model considers the fraction $\Delta E/E$ of energy loss of quarks charm and beauty via collisional and radiative processes, as function of the quark p_T . For charm quarks and p_T above 10 GeV/ c , the radiative processes become more important than the collisional ones, while for beauty quarks this transition occurs for p_T above 25 GeV/ c . Although the radiative energy loss plays the most important role at high p_T , the model shows that the collisional energy loss is not negligible at LHC

energies. For the collisional energy loss, beauty quarks are expected to loose less energy than charm quarks in the low p_T region, while the fraction of energy loss becomes the same for both at higher p_T . For the radiative energy loss, the beauty quark is expected to loose less energy than the charm quark over a very wide p_T range, as a consequence of the dead-cone-effect.

Figure 3.7: Fractional energy loss $\Delta E/E$ evaluated for collisional and radiative processes for charm and beauty quarks, at a constant temperature $T = 304$ MeV and path length $L = 5$ fm. Figure taken from [9].

Precise measurements of the nuclear modification factor R_{AA} of observables related to both light and heavy-quarks can provide important tests of the flavor-dependence of the in-medium energy loss predicted by QCD. On the heavy-flavor sector, is of particular importance to measure observables related to beauty and charm quarks separately.

In the left panel of Figure 3.8 it is shown a preliminary measurement of the R_{AA} of prompt D mesons by ALICE in Pb-Pb at $\sqrt{s_{NN}} = 5.02$ TeV at mid-rapidity in three different centrality classes, named 0-10%, 30-50%, 60-80% (the class 0-10% corresponds to central collisions and the 60-80% to peripheral collisions, as will be discussed in section 6.4). A strong suppression of high p_T D mesons is observed, and this suppression increases from more peripheral to more central collisions. Comparison with the R_{pA} of D mesons (at Figure 3.5) suggests that this strong suppression is a medium-induced effect, reflecting the interaction between the heavy-quarks with the QGP. In the right

panel of the same figure it is shown the R_{AA} of non-prompt J/ψ (from B meson decay) measured by ATLAS in Pb-Pb at $\sqrt{s_{NN}} = 5.02$ TeV at mid-rapidity in the centrality classes 0-10%, 20-40% and 40-80% [66]. In agreement with the previous measurement, a larger suppression of the heavy-flavor particles is observed in more central collisions, where a larger color-deconfined medium is expected to be produced.

Figure 3.8: (Left) R_{AA} of prompt D mesons measured by ALICE in Pb-Pb at $\sqrt{s_{NN}} = 5.02$ TeV at mid-rapidity in the centrality classes 0-10%, 30-50%, 60-80% and (right) R_{AA} of non-prompt J/ψ measured by ATLAS in Pb-Pb at $\sqrt{s_{NN}} = 5.02$ TeV at mid-rapidity in the centrality classes 0-10%, 20-40% and 40-80% [66].

In order to test the mass dependence of energy loss in the heavy-flavor sector, efforts have been made to probe beauty and charm quarks separately. At LHC run 1, many measurements have been performed. In Figure 3.9 it is shown the R_{AA} of D mesons (that probes the interaction of charm quarks with the QGP) measured by ALICE [67] in comparison with the R_{AA} of non-prompt J/ψ (that come from B meson decays and consequently probe the interaction of the beauty quarks with the QGP) measured by CMS [68], as a function of the mean number of participants in the collision, $\langle N_{part} \rangle$, that is proportional to its centrality. The R_{AA} shown are the mean values obtained within a centrality class and considering the high p_T region of the spectra, where the suppression is expected (see figure caption). These results point to a smaller suppression

for non-prompt J/ψ than for D mesons, mainly for more central events. The Djordjevic model can describe the trend of the observed suppression as function of $\langle N_{part} \rangle$, for both particle species. The difference on the R_{AA} of non-prompt J/ψ and D mesons derives mainly from the quark-mass dependence of the energy loss: the dotted blue lines indicate the predicted R_{AA} for the non-prompt J/ψ in the case the b quark loose energy as c quark. In this scenario, this R_{AA} becomes much closer to the R_{AA} of D mesons.

Figure 3.9: R_{AA} of D mesons measured by ALICE compared with the R_{AA} of non-prompt J/ψ measured by CMS in Pb-Pb at $\sqrt{s} = 2.76$ TeV as function of the mean number of participants in the collision $\langle N_{part} \rangle$, that is proportional to its centrality [67].

The results obtained from ALICE during LHC run 1 on the electronic channel support the results shown previously, for central collisions. In the left panel of Figure 3.10 it is shown the comparison between the R_{AA} of electrons from beauty hadron decays and the R_{AA} of electrons from open heavy-flavor hadron decays, both measured in the 20% most central collisions. A hint of smaller suppression for electrons from beauty hadron decays than for electrons from open heavy flavor hadron decays is observed in the p_T range from 3 to 8 GeV/c, that is the common p_T range of both analysis. The R_{AA} of electrons from

beauty hadron decays exhibits a suppression of about 30-40% for $p_T > 3$ GeV/ c and the trend of this suppression is described reasonably well by theoretical models that includes mass dependent energy loss of heavy quarks inside QGP, as shown in the right panel of Figure 3.10. Nevertheless, this particular measurement have very large uncertainties at very low p_T (from ≈ 1.25 to 2.5 GeV/ c), and although the experimental measurement points to an $R_{AA} > 1$ in this region, the values are still comparable with the theory, that predicts an R_{AA} very close to (or even below) unity in this p_T range.

Figure 3.10: (Left) R_{AA} of electrons from beauty hadron decays (red) compared to the R_{AA} of open heavy-flavor electrons measured by ALICE at mid-rapidity in central Pb-Pb collisions. (Right) R_{AA} of electrons from beauty hadron decays compared to different theoretical predictions [53].

The CMS collaboration have performed recently the reconstruction of B^+ mesons in Pb-Pb collisions for the first time [69], using LHC run 2 data, in a centrality integrated analysis (0-100%). As shown in the left panel of Figure 3.11, the R_{AA} of B^+ mesons and non-prompt J/ψ are in good agreement in the common p_T range within experimental uncertainties. In this same figure it is shown the R_{AA} of D mesons [70] and charged particles. A hint of smaller suppression of beauty hadrons (or beauty hadron decay products, like the non-prompt J/ψ) with respect to charm hadrons is observed at low p_T , supporting the previous results from run 1 shown at Figure 3.10. In general, at low

p_T , it is observed a hint of an hierarchy: $R_{AA}(B^+) > R_{AA}(D) > R_{AA}(\pi^\pm)$, expected from the flavor-dependence of parton energy loss predicted by QCD (note that pions are produced mainly via gluon fragmentation [71] [72], so their production in heavy-ion collisions is sensitive to the energy loss of gluons in the QGP). The right panel of Figure 3.11 shows the R_{AA} of B^+ mesons compared with theoretical models. The B^+ mesons R_{AA} is suppressed of about 50% for $p_T > 7 \text{ GeV}/c$.

Figure 3.11: R_{AA} of D mesons, B mesons, non-prompt J/ψ and charged particles for centrality integrated events measured at mid-rapidity by CMS [70], [69]. On the right, the R_{AA} of B mesons is compared to different theoretical models.

By the comparison between the results from ALICE (Figure 3.10) and CMS (Figure 3.11), a suppression of $\approx 40\%$ is observed on the R_{AA} of both B^+ mesons and electrons from beauty hadron decays at large p_T , what indicates that both observables points to a consistent physics message with respect to the energy loss of beauty quarks. The current large uncertainties of both experimental measurements, nevertheless, do not allow to impose stringent constraints to the different theoretical models. With what regards the minimum p_T range that could be accessed experimentally, the B^+ measurement starts from $p_T = 7 \text{ GeV}/c$, while the non-prompt J/ψ measurement starts from $p_T = 3 \text{ GeV}/c$. ALICE, on the other hand, allows the measurement to be extended to smaller p_T , starting from $p_T = 1.25 \text{ GeV}/c$. As indicated in Figure 3.7, the low p_T region is of particular interest because there the mass dependence of the energy loss processes is

expected to be stronger. In addition, the low p_T region of the R_{AA} is expected to be influenced by effects from different origins. The shadowing of the nuclear PDFs is expected to reduce the production of heavy-flavor hadrons in nuclear collisions with respect to pp collisions at low and intermediate p_T [73] [74]. On the other hand, the coalescence of heavy quarks with light quarks of the medium is expected to increase the heavy-flavor production at low and intermediate p_T [74] (see also discussion in [section 3.2.2](#) below). In addition, it is also known that theoretical models which includes radial flow of the heavy quarks (collisional energy loss of heavy quarks in a fluid-dynamical expanding medium) predicts a bump in the R_{AA} of heavy-flavor particles at low p_T [9]. Because of that, it is of particular importance to have a more precise measurement of the R_{AA} of electrons from beauty hadron decays at very low p_T , in order to test the description of the models that contains these different ingredients.

One of the analysis performed in this work is the measurement of electrons from beauty hadron decays at low p_T using the data from Pb-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, collected during LHC run 2. This analysis will be discussed in detail on [chapter 6](#). With the larger amount of data collected at run 2 (about 3 times more events are analyzed in run 2 with respect to run 1), the total error bars of the run 1 measurement ([Figure 3.10](#)) could be significantly reduced, constraining the R_{AA} p_T -dependence, in particular for $p_T \lesssim 3$ GeV/ c , where the result from run 1 shows poor precision. In addition, the R_{AA} of electrons from both beauty and open heavy-flavor hadron decays measured in run 2 could be extended to larger a p_T ranges with respect to the run 1 measurements, improving the experimental picture of mass-dependent energy loss of heavy-quarks shown in the left panel of [Figure 3.10](#).

As presented above, the production of heavy-flavor particles measured in Pb-Pb collisions at LHC have shown a clear deviation from binary scaling, having an associated R_{AA} that deviates from unity. The suppression of the R_{AA} of heavy-flavor particles is consistent, within experimental uncertainties, with the scenario of energy loss of heavy quarks in a color deconfined medium, although the experimental errors do not allow the discrimination among many different theoretical models, that have slightly different descriptions of the heavy quark propagation through the QGP. The mass dependence of

energy loss of heavy quarks is probed with limited precision. This makes essential, in order to test QCD in a more effective way, to perform measurements using newer and more abundant data samples as soon as they are available, as is the aim of one of the works of this thesis.

The Interplay of Hadronization Mechanisms

The QGP can modify the production of heavy-flavor particles not only by inducing energy loss of the heavy quarks, but also by inducing a different hadronization mechanism. A quark in the vacuum hadronize via fragmentation, that is the QCD-based process of creation of another quarks from the vacuum, by the intense color field that is originated in the attempt to isolate a color-carrying object. This causes the confinement of this color-carrying object into a color neutral hadron. This process creates a color neutral hadron with momentum $p_h = z.p_q$, that is a fraction $z \leq 1$ of the initial quark momentum p_q . When quarks hadronize inside a color deconfined medium, it is considered that another hadronization mechanism can take place, which is called coalescence. This different hadronization mechanism was proposed as an explanation for the anomalous baryon to meson ratio ("proton anomaly") and scaling of the elliptic flow with the number of constituents quarks of the hadrons observed in early results at RHIC in the light flavor sector [75] [76]. Coalescence consists in the recombination of quarks that are inside QGP close to each other in phase space. In this process, the resulting hadron has a momentum that corresponds to the sum of the momentum of the initial quarks (see diagram on [Figure 3.12](#)). The existence of the coalescence as a possible hadronization mechanism inside QGP modifies the momentum spectrum of the produced hadrons. In particular, the coalescence can modify the chemical composition of the produced hadrons, since it favors the baryon production with respect to meson production: since the amount of low momentum quarks in the medium are much larger than the amount of intermediate and high momentum quarks, the production of a hadron with a given momentum p_h is more probable via coalescence of three quarks with very low momentum ($p_q \approx p_h/3$), what creates a baryon, than via the coalescence of a quark anti-quark pair with larger momentum ($p_q \approx p_h/2$), what creates a meson, as sketched

on Figure 3.12.

Figure 3.12: Diagram representing the production of a hadron with momentum $p_T \approx 6$ GeV/c via quark fragmentation, coalescence of a quark anti-quark pair or coalescence of three quarks. The black line represents the momentum distribution of the quarks in the medium. Figure taken from [77].

It is also known that the production of strange hadrons is favored in the coalescence scenario, where quarks can recombine inside a quark and gluon soup rich of strange and anti-strange quarks, that are produced mainly by gluon fusion [78] [3] [79].

The measurement of the production of baryons and strange particles in heavy-ion collisions can, because of the reasons discussed above, be sensitive to the effects of the hadronization mechanisms on the produced particle spectra, allowing to quantify and disentangle these effects from energy loss inside the QGP.

In Figure 3.13 it is shown the R_{AA} of many different particle species measured by ALICE. The R_{AA} of charged particles and D mesons are compared to the ones of strange-D meson and Λ_c baryon. These measurements give a hint of the pattern expected in a scenario where coalescence in QGP takes place: the strange-D meson production is enhanced in comparison with non-strange D meson production, and the Λ_c baryon production is even more enhanced than any of the others, since it not only contains a strange

quark but it is also a baryon. It is important, nevertheless, to point out that the measurement of Λ_c is performed in the centrality range 0-80% while the other measurements are performed at 0-10%. The strange-D mesons and Λ_c baryon are particularly valuable probes to disentangle energy loss and hadronization effects in the heavy-flavor sector. More precise measurements, possible during the upcoming LHC run 3, will allow to better constrain these effects.

Figure 3.13: R_{AA} of charged particles, non-strange D mesons, strange D mesons and Λ_c baryon measured in Pb-Pb at $\sqrt{s_{NN}} = 5.02$ TeV collisions by ALICE.

In the [Appendix A](#) it is introduced briefly some measurements about anisotropic flow, jets and quarkonia, that are also important topics of study that allows to investigate QGP properties, although they are not directly related to the analysis of the present work. The reader are invited to read this topics and also to enjoy the references within in case of interest.

3.3 Transport Models of Heavy-Quarks Inside QGP

In order to predict the heavy-flavor production in heavy ion collisions, theoretical models need to include a series of ingredients. In addition to modeling the heavy-quark

energy loss mechanisms, the models also need to include a description of their initial production (that can include cold nuclear matter effects or not), a description of the space-time evolution of the QGP up to the freeze-out, mechanisms of hadronization (coalescence, fragmentation) of the quarks and also the interaction of the beauty and charm hadrons with the hadronic matter after the freeze-out. Many energy loss models have been developed taking into account different ingredients. Different choices of these ingredients can give a factor two of difference in the prediction of a model [9]. Below, the main ingredients that are considered by different theoretical models are summarized. In general, the models can be subdivided in two groups: the ones that make use of perturbative QCD calculation in the description of the interaction of heavy quarks with the QGP and the ones that use non-perturbative approaches.

A more detailed description of many models can also be found here [9]

pQCD based models:

Djordjeovich *et al.*: pQCD energy loss in a static fireball [65]

Initial production of heavy-quarks: given by FONLL and no CNM effects included;

Quark-medium interactions: Both collisional and radiative energy loss are computed within leading order pQCD. Includes running of the coupling;

Hadronization: Fragmentation of the heavy quarks.

Medium: The medium has finite size and it is described as a thermalized QGP at temperature T . Partons of the medium are not static scattering centers, but dynamical (their recoil is taken into account). The medium does not expand.

BAMPS [80]

Initial production of heavy-quarks: given by Monte Carlo event generator for next-to-leading order (MC@NLO [81]). No CNM effects included.

Quark-medium interactions: Collisional interactions described by pQCD and a K-factor of $K = 4$ to account for radiative energy loss (tuned to match RHIC results).

Transport of the heavy quarks is given by Boltzmann equation. Running of coupling constant is considered.

Hadronization: Fragmentation of the heavy quarks;

Medium: Partons of the medium interacting according to Boltzmann equation (3+1)d;

MC@sHQ+EPOS, Coll+Rad(LPM) [82]

Initial production of heavy-quarks: given by FONLL or MC@NLO;

Quark-medium interactions: Collisional and radiative energy loss are considered. The transport of the heavy quarks is given by Boltzman equation. Running of the coupling constant is considered;

Hadronization: Fragmentation of the heavy quarks and coalescence of heavy quarks with light quarks from the medium;

Medium: (3+1)d ideal fluid-dynamical evolution from EPOS [83] initial conditions.

POWLANG: pQCD embedded in viscous hydrodynamics [84]

Initial production of heavy-quarks: NLO pQCD calculation of initial heavy-quark yields done by POWHEG. CNM effects considered (EPS09 [16] nuclear corrections on the PDFs and k_T -broadening).

Quark-medium interactions: Only collisional energy loss is considered. The transport of heavy quarks is given by Langevin equation. pQCD calculations used for hard-scatterings and Hard Thermal Loop (HTL) for soft interactions.

Hadronization: Fragmentation of the heavy quarks and coalescence of heavy quarks with light quarks from the medium (coalescence just for most recent models);

Medium: The fireball evolution is described by a (2+1)d relativistic viscous hydrodynamics.

Parton-Hadron-String Dynamics (PHSD) [85] [73]

Initial production of heavy-quarks: given by PYTHIA [44] (tuned to fit FONLL predictions for charm quarks). CNM effects included;

Quark-medium interactions: Collisional energy loss computed within pQCD;

Hadronization: Fragmentation of the heavy quarks and coalescence of heavy quarks with light quarks from the medium;

Medium: No hydrodynamics. QGP described by Dynamical Quasi-Particle Model [86];

Non-perturbative approaches:

TAMU [87]

Initial production of heavy-quarks: Given by FONLL;

Quark-medium interactions: Collisional energy loss is calculated in a non-perturbative framework (makes use of lattice QCD potentials);

Hadronization: Fragmentation of the heavy quarks and coalescence of heavy quarks with light quarks from the medium;

Medium: Hydrodynamical evolution;

AdS/CFT [88] [89]

Initial production of heavy-quarks: given by FONLL, without inclusion of CNM effects;

Quark-medium interactions: AdS/CFT drag derivations;

Hadronization: Fragmentation of the heavy quarks;

Medium: Static fireball based on Glauber model [45] [46] [47] [48];

Chapter 4

Experimental Apparatus

The ALICE (A Large Hadron Collider Experiment) is an experiment located in one of the four interaction points of the LHC (Large Hadron Collider). This experiment is optimized to perform measurements of particles in the high multiplicity environment created in Pb-Pb collisions. The main goal of the experiment is to study the properties of the nuclear matter in conditions of very high temperatures and energy densities, probing the phase of color deconfinement predicted by QCD. A sketch of ALICE is presented in [Figure 4.1](#). ALICE can be divided in two parts: detectors at central rapidity (in the so-called central barrel) and detectors at forward rapidity, mainly used for triggering, event plane and centrality determination, and muon detection.

The ALICE central barrel is embedded in a magnetic solenoid that produces a field of $B = 0.5$ T parallel to the beam pipe. The charged particles have their trajectory bended by the magnetic field according to its mass and velocity, what makes possible the momentum determination of charged particles by the tracking detectors inside the central barrel.

A brief description of the ALICE detectors is given below, and a more detailed discussion of the main detectors used in the present work is presented later in this chapter. An overview of the ALICE detectors can be found in reference [\[90\]](#) and a summary of particle detection principles can be found in reference [\[91\]](#).

THE ALICE DETECTOR

Figure 4.1: Sketch of the ALICE detector

Central rapidity detectors:

- Inner Tracking System (ITS) [92]: High granularity silicon detector dedicated to tracking, primary and secondary vertex reconstruction, impact parameter determination and particle identification;
- Time-Projection-Chamber (TPC) [93]: Gas chamber that consists in the main tracking and particle identification detector in ALICE;
- Transition-Radiation-Detector (TRD) [94]: Dedicated to the identification of electrons with large momenta and to trigger on high transverse momentum particles;
- Time-of-Flight (TOF) [95]: Dedicated to the measurement of time-of-flight of particles over a known distance. Provide mass information that allows p/K and K/ π separation at low momentum;

- High-Momentum-Particle-IDentification (HMPID) [96]: Performs hadron discrimination at high transverse momenta ($1 < p_T < 3$ GeV/ c for K and π , $1 < p_T < 5$ GeV/ c for p);
- PHOton Spectrometer (PHOS) [97]: High resolution electromagnetic calorimeter;
- EletroMagnetic Calorimeter (EMCal) [98]: Electromagnetic calorimeter and trigger on high momentum electrons and photons;
- ACORDE (A COsmic Ray DEtector) [99]: Array of plastic scintillator counters placed outside ALICE magnet. Dedicated to cosmic ray measurements;

Forward rapidity detectors:

- T0 [100][101]: Two arrays of Cherenkov counters (T0A and T0C), dedicated to trigger the event and to give the event collision time with high resolution (≈ 25 ps);
- VZERO [102][101]: Two arrays of scintillator counters (V0A and V0C) used as Minimum Bias trigger, centrality trigger in Pb-Pb collisions and for event plane determination;
- Forward-Multiplicity-Detector (FMD) [103][101]: Segmented silicon strip detector dedicated to extend charged particle multiplicity measurements for the forward region and to event plane determination;
- Photon-Multiplicity-Detector (PMD) [104]: Measure the multiplicity and spacial distribution of photons at forward rapidity;
- Zero-Degree Calorimeters (ZDC) [105]: Set of proton (ZP) and neutron (ZN) forward calorimeters dedicated to measure the energy of spectators (that do not participate in the collision) protons and neutrons and to estimate the centrality of Pb-Pb collisions;

- Muon spectrometer [106]: Set of muon tracker, trigger, hadron absorbers that allows muons measurement. The physics goal is to study heavy-flavor and quarkonia production (heavy quark resonances like J/ψ and Υ) via the muon decay channel;

Find below more detailed description of ITS, TPC, TOF and VZERO, that are the main detectors used in the analysis performed in this thesis.

Inner Tracking System

The Inner Tracking System (ITS) [107] [92] is the innermost tracking detector in ALICE. It covers the central rapidity region ($\eta < |0.9|$ for the full detector) and the azimuthal range of $\Delta\phi = 2\pi$.

One of the main tasks of the ITS is to reconstruct the primary vertex of the collision with a good resolution (better than $100 \mu\text{m}$ along z direction) and to reconstruct the secondary vertices of beauty and charm hadron decays, improving the identification of the different heavy-flavor species. The ITS detector allows the reconstruction of the tracks transverse impact parameter to the primary vertex ($d_{0r\phi}$) with a resolution better than $100 \mu\text{m}$, as shown in Figure 4.2 for the run 1 data.

The ITS has a central role in tracking and PID of very low momentum charged particles ($p_T < 200 \text{ MeV}/c$), that can bend in the magnetic field to the point of not reaching the TPC. For higher momentum charged particles, that also reaches the TPC, the matching between the ITS and TPC tracks improves the momentum and angle resolution provided by the TPC alone.

The ITS is composed by six cylindrical layers of silicon detectors, and every two layers is built with a different technology, in order to address the requirements for a good tracking at different distances from the primary vertex.

The inner radius of the ITS is the minimum radius compatible with the beam pipe (that has a radius of $\approx 3 \text{ cm}$), and the outer radius is determined by the matching with the TPC.

The first two layers are located at 4 and 7.2 cm from the interaction vertex, where the charged track density is very high. In order to have a high spacial resolution that allows a good separation between the tracks, these first two layers are built with Silicon

Figure 4.2: Resolution of the transverse impact parameter to the primary vertex given by ITS, as function of the track p_T . The estimation is performed with Pb-Pb $\sqrt{s_{NN}} = 2.76$ TeV data. The experimental result is compared with a Monte Carlo simulation [107].

Pixel Detectors (SPD). The SPD layers have acceptance of $\eta < |1.75|$. The second two layers are located at 15 and 23.9 cm distance from the primary vertex and they are built with Silicon Drift Detectors (SDD), while the third two layers are located at 38.5 and 43.6 cm from the primary vertex and are built with Silicon Strip Detectors (SSD). The SDD and SSD layers have acceptance of $\eta < |0.9|$ and they have an analogical readout that allows particle identification via the measurement of energy loss per unit length (dE/dx) in the material.

A summary of the main characteristics of the ITS detector can be found in [Table 4.1](#).

Time Projection Chamber

The Time-Projection-Chamber (TPC) [93] is the main tracking detector in ALICE. It covers the pseudo-rapidity range of $\eta < |0.9|$, and the azimuthal range of $\Delta\phi = 2\pi$. It has a cylindrical shape of about 5 m width and 5 m diameter. The TPC is used for track reconstruction and particle identification (PID).

Layer	Detector	r(cm)	$ \eta $	$\sigma_{r\varphi}(\mu m)$	$\sigma_z(\mu m)$
1	pixel	4.0	1.75	12	100
2	pixel	7.2	1.75		
3	drift	15.0	0.9	38	25
4	drift	23.9	0.9		
5	strip	38.5	0.9	20	830
6	strip	43.6	0.9		

Table 4.1: Summary of characteristics of ITS detectors. The columns $\sigma_{r\varphi}(\mu m)$ and $\sigma_z(\mu m)$ refer to the tracking resolution in the transverse ($r\varphi$) and longitudinal (z) directions, respectively.

Figure 4.3: Sketch of the TPC showing its inner and outer radius, readout pads in the end-plates and the central electrode, that defines two regions of electric field of opposite directions.

The TPC is filled with a gas mixture of Ne (90%) and CO₂ (10%). A charged particle traversing this gas mixture prints a 3D ionization path in the gas. The static electric field between each TPC end-plate and the central electrode (see [Figure 4.3](#)) makes the electrons created in the ionization to drift towards the TPC end-plates, where a 2D (xy

plane) projection of the particle path is read by the read-out modules in the end-plates. The TPC end-plates contains three different sizes of read-out pads (going from smaller ones in the inner radius to larger ones in the outer radius). In each read-out pad, the signals that are close in space and time are combined into clusters. By knowing the drift time of the electrons in the gas and also the time in which the collision took place (given by T0 detector), the z-coordinate of each cluster of the trajectory can be reconstructed, leading to a 3D path information for charged particles inside the TPC.

For each reconstructed track in the TPC, the amount of charge in each cluster is used to estimate the track energy loss in the corresponding track length. The procedure used to estimate the track mean energy loss per length corresponds to a truncated mean of the values of energy loss per length obtained for each cluster. The mean is obtained using the 50-70% clusters with the lowest deposited charge (Sec. 31.6.5 of [108]). This method improves the particle species separation by making the $\langle dE/dx \rangle$ of different particles species more Gaussian like (reducing the tail of the Landau distribution, typical of the process of energy loss in matter [109]). Figure 4.4 shows an example of $\langle dE/dx \rangle$ distribution of different particle species traveling inside the TPC as a function of the particle momentum. Together is shown the theoretical prediction given by the Bethe-Bloch formula ¹.

In order to select a particle based on its dE/dx in TPC, it is used the quantity named $n\sigma_{\text{TPC}}$, that represents the significance of the deviation between the TPC dE/dx signal and a particle identity hypothesis:

¹ The Bethe-Bloch formula, Equation 4.1, gives the $\langle dE/dx \rangle$ of a given particle propagating through a given material as a function of the particle $\beta = v/c$ and mass m_e and the material properties (see Sec. 30 of [108] for a complete discussion).

Since energy loss is a statistical process, the actual energy loss of a particle will show fluctuations around the average given by the Bethe-Bloch prediction. These fluctuations follow a Landau distribution [91].

$$-\left\langle \frac{dE}{dx} \right\rangle = Kz^2 \frac{Z}{A} \frac{1}{\beta^2} \left[\frac{1}{2} \ln \frac{2m_e c^2 \beta^2 \gamma^2 T_{max}}{I^2} - \beta^2 - \frac{\delta(\beta\gamma)}{2} \right] \quad (4.1)$$

$$n\sigma_{\text{TPC}} = \frac{\langle dE/dx \rangle_{\text{measured}} - \langle dE/dx \rangle_{\text{expected}}}{\sigma_{dE/dx}}, \quad (4.2)$$

where $\langle dE/dx \rangle_{\text{measured}}$ is the mean energy loss measured in TPC for a given track, $\langle dE/dx \rangle_{\text{expected}}$ is the expected energy loss in TPC for a given particle identity hypothesis and $\sigma_{dE/dx}$ is the width of the measured signal distribution.

Practical examples of the use of Equation 4.2 will be given later in the analysis developed in this thesis.

The TPC was designed to have a good performance in a high multiplicity environment in which the particle density per unity of rapidity is of the order of $dN_{\text{ch}}/dy = 8000$. The minimum possible inner radius of the TPC ($r_{\text{in}} \approx 90$ cm) is constrained by the maximum acceptable hit density in this high multiplicity environment. The outer radius ($r_{\text{out}} \approx 250$ cm) is determined by the minimum length required for the track to have a dE/dx resolution better than 10%. The gas mixture guaranties a small drift time ($\lesssim 100$ μs) and the two cluster resolution is about ≈ 1 cm [93].

Figure 4.4: TPC dE/dx for various particle species measured in Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV as a function of the track momentum, superimposed with Bethe-Bloch lines.

Time-Of-Flight

The Time-Of-Flight (TOF) [95] is a cylindrical detector located at the central barrel, with pseudo-rapidity coverage of $\eta < |0.9|$ and full azimuth coverage $\Delta\phi = 2\pi$. The detector is divided in 90 modules, and each module is equipped with many MRCP (Multigap Resistive Plate Chamber) strip detectors, that provides a time resolution of the order of 50 ps.

The TOF is responsible for particle identification by measuring the time it takes for the particle to travel from the primary vertex to its MRCP. The time measurement provided by TOF together with the momentum and length of tracks provided by the tracking detectors allows the calculation of the mass of the particles. With the TOF, particles with different masses can be distinguished, and this is particularly important for the separation between pions, kaons and protons, that have their TPC dE/dx distribution overlapped in a considerable momentum region (see Figure 4.4). The identification of pions and kaons are particularly important for the D meson reconstruction (see Table 3.1).

In the analysis performed in this thesis, dedicated to the measurement of electrons, the TOF is used to eliminate part of the contamination of kaons and protons in the electron candidates sample. In Figure 4.5 is shown the β (v/c) measured by TOF as function of the track momentum, and it is possible to see that up to $p \approx 2 \text{ GeV}/c$ the TOF has an important role in the discrimination between pions, kaons and protons and in the discrimination of kaons and protons from electrons.

VZERO

The VZERO [102] detector is composed by two subsystems, named V0A and V0C, that corresponds to two arrays of plastic scintillator counters. They are located around the beam pipe, at forward rapidity, one at each side of the interaction point: the V0A covers the pseudo-rapidity range of $2.8 < \eta < 5.1$ while the V0C covers $-3.7 < \eta < -1.7$. Both V0A and V0C are segmented in four concentric rings around the beam pipe, as sketched in Figure 4.6, and each ring is divided in 8 sections in the azimuthal direction.

The VZERO plays many roles. It can be used as collision centrality estimator.

Figure 4.5: β of tracks measured by TOF as function of their momentum.

Figure 4.6: Sketch of the front view of a VZERO subsystem and its segmentations.

The sum of the energy deposited in V0A and V0C scales with the number of primary particles produced in the collision, and therefore with the centrality of the colliding nucleus (further discussion about this will be presented in [section 6.4](#)).

One of the most important functions of VZERO is that of a Minimum Bias (MB) trigger. The MB trigger is defined such that an event is recorded only if a signal is present in either of the VZERO arrays or in the SPD. The VZERO also provides the time of arrival of the particles on its surface with a good precision. A VZERO trigger can be used to select beam-beam interactions in detriment of beam-gas interactions, and it explores the fact that the arrival time of particles at each of the two arrays will be

different in these two different situations.

The segmentation of the VZERO, as shown in [Figure 4.6](#), allows the determination of azimuthal particle distribution and event plane Ψ_ϕ .

Chapter 5

Measurement of Electrons from Open Heavy-Flavor Hadron Decays in pp Collisions at $\sqrt{s} = 13$ TeV

5.1 General Strategy

In this chapter, the measurement of the invariant cross section of electrons from open heavy-flavor (*charm* or *beauty*) hadron decays in pp collisions at low momentum is presented. In order to calculate this invariant cross section, $\frac{1}{2\pi p_T} \frac{d^2\sigma}{dp_T dy}$, the ingredients shown in [Equation 5.1](#) are necessary. These ingredients are discussed qualitatively in this section and quantitatively along this chapter.

$$\frac{1}{2\pi p_T} \frac{d^2\sigma}{dp_T dy} = \frac{1}{2} \frac{1}{2\pi p_T^{\text{center}}} \frac{1}{\Delta y \Delta p_T} \frac{N_{\text{raw}}^{e^\pm}(p_T)}{(\varepsilon^{\text{reco}} \times \varepsilon^{\text{ID}})} \frac{\sigma_{V0}}{N_{\text{ev}}^{V0}} \quad (5.1)$$

The pp collisions at $\sqrt{s} = 13$ TeV have a visible cross-section σ_{V0} , that is determined based on the signal these events generate in the VZERO detector.

From all the events generated in the experiment, N_{ev}^{V0} events are analyzed. Those are the ones selected by the minimum bias trigger, that, for consistency with the σ_{V0} , is also based on the signal generated in the VZERO detector. In addition, the analyzed

events need to satisfy some requirements that ensure their quality.

From all the tracks generated in each event, just the ones that pass a quality selection criteria are used. This track selection has an efficiency ϵ^{reco} for electrons, that is estimated using simulations of pp events generated by Monte Carlo.

The raw p_T -spectrum of electrons and positrons from heavy-flavor hadron decays, $N_{\text{raw}}^{e^\pm}$, is measured via a sequence of many steps. For practical purposes, "electrons and positrons" will be referred to as just "electrons" during the whole analysis procedure. In the end, for the calculation of the invariant cross section, as shown in [Equation 5.1](#), a factor $1/2$ is considered in order to make the final result account just for electrons, that consist in half of the electron + positron measured yield. The first step for the calculation of $N_{\text{raw}}^{e^\pm}$ is the selection of electron candidates via PID (Particle IDentification) cuts on the signals generated in TPC and TOF detectors. Cuts are applied in order to select tracks which have a time-of-flight measured by TOF and the energy loss per unity length $\langle dE/dx \rangle$ in the TPC compatible with the electron hypothesis. These PID cuts select electrons with an efficiency ϵ^{PID} , also estimated in this analysis via Monte Carlo simulations. This sample of electron candidates selected by the PID cuts is composed by electrons from heavy-flavor hadron decays, electrons from non heavy-flavor hadron decays and also a hadron contamination. The hadron contamination is estimated via a data-driven method based on the signal generated by electrons and hadrons in the TPC, in each p_T bin. With this method is possible to estimate how many hadrons are selected together with the electrons after the PID cuts on the TOF and TPC signals. Once the hadron contamination is estimated and subtracted, the background of electrons from non heavy-flavor hadron decays, that consists mainly in electrons from light-flavor mesons Dalitz decay and gamma conversion in the detector material, are identified via a data-driven method called *photonic tagging method*. Pairs e^-e^+ from this sources can be tagged in data via their low invariant mass. The efficiency of this method, the *tagging efficiency*, is estimated from Monte Carlo simulations and it is used to correct the yield of electrons from non heavy-flavor sources. The identification of the non heavy-flavor electrons via the data-driven approach used in this work is known to lead to smaller final systematic uncertainties on the final heavy-flavor electron yield with respect to the

analysis in which the non heavy-flavor electrons background is estimated via cocktail simulations¹ and subtracted [13].

After the subtraction of the hadron contamination and of the electrons from non heavy-flavor sources from the electron candidate sample, the yield of electrons from heavy-flavor hadron decays $N_{\text{raw}}^{e^\pm}$ is obtained.

This yield is corrected by the efficiencies of the track selection $\varepsilon^{\text{reco}}$ and of the PID cuts applied in the TPC and TOF signals in order to select electrons, ε^{eID} .

The other factors in Equation 5.1 are geometrical factors, that normalize the yield per unity of rapidity Δy and unity of azimuthal angle $\Delta\phi = 2\pi$, or factors used to remove the p_T bin size (Δp_T) and p_T bin center (p_T^{center}) dependence of the measured spectrum.

Along this chapter, the quantitative determination of each of the ingredients of Equation 5.1 is presented. In the end, the obtained invariant cross section of electrons from heavy-flavor hadron decays compared to FONLL prediction and discussed.

5.2 Data Set and Event Selection

In this analysis, events from two different Minimum Bias (MB) data samples (LHC16k and LHC16l) recorded in pp collisions at $\sqrt{s} = 13$ TeV in 2016 by ALICE are used.

The online MB trigger, used to select events during the data taken, requires at least one hit in the SPD or in either of the two VZERO arrays. An additional selection based on the VZERO is applied during the offline analysis, with the aim of selecting beam-beam interactions in detriment of beam-gas interactions. It is based on the time the created particles reach V0A and V0C with respect to the bunch crossing time in the nominal interaction point.

An amount of ≈ 180 M MB events are available considering both samples together, from which ≈ 160 M were analyzed in this work, that correspond to the events that

¹A cocktail is built using a Monte Carlo hadron-decay generator, that can produce the p_T -spectrum of a certain decay particle using as input the measured p_T -spectrum of the source. In this case, the spectra of non heavy-flavor electrons are built by the MC generator from the measured p_T -spectra of the non heavy-flavor sources, measured experimentally [13].

passed the selection criteria described below.

One of the most important requirements to select an event is a good quality of the primary interaction vertex determination. The cuts applied on the primary vertex, described below, have mainly the purposes of:

- Guarantee that the collision occurred very close to the nominal point, at the center of the detectors along the z direction. This ensures a uniform reconstruction efficiency of charged particles, since it allows the particles emitted on both directions to generate tracks in the detectors with proper length;
- Identify and reject cases where two different events occurred very close to each other in time and were recorded as a single event (pile-up rejection);

Global tracks, that are tracks reconstructed using TPC and ITS tracks combined, are used to find the interaction vertex with high precision. Only events with a primary vertex reconstructed with at least two contributor tracks are selected. In addition, the primary vertex needs to be within 10 cm, in the longitudinal direction, from the nominal interaction point ($|V_z| < 10$ cm).

A pile-up rejection is performed by using two predefined algorithms for multiple vertex finding. One is based on vertexes found with the use of the signal on SPD only, V_{SPD} . The minimum number of contributors for V_{SPD} determination is predefined and multiplicity dependent (varying from 3 to 5). If within one event two vertexes are found by this algorithm and the distance between them is larger than 0.8 cm then the event is rejected.

Additional pile-up rejection based on another algorithm of multiple vertex finding is applied. In this algorithm, the minimum number of tracks for a vertex determination is 5 and the event is rejected if two (or more) vertexes are found and the distance between two vertexes weighted by the convolution of their errors is larger than 15.

Around 85% of the events satisfy these selection criteria.

A summary of the event selection cuts is shown in [Table 5.1](#).

Type	Value
Min. global track contributors	2
$ V_z $	< 10 cm
Min. contributors to V_{SPD}	3 - 5
Distance in z between two V_{SPD}	< 0.8 cm
Min. track contributors to vertexes	5
Weighted dist. in z between two vertexes	< 15

Table 5.1: Event quality cuts used on the pp at $\sqrt{s} = 13$ TeV analysis.

5.3 Monte Carlo Samples

All Monte Carlo (MC) samples used in the analysis have the same detector configuration and beam conditions as the ones found during the data taken, what means that they are *anchored* to the data sample. Two MC samples anchored to each data sample are used in this analysis. General purpose (minimum bias) MC productions are used for the calculation of the tagging efficiency. The other MC samples are special PYTHIA [44] MC simulations which have an enhancement of π^0 and η mesons and also in which a trigger on charm and beauty hadrons decaying semi-electronically is employed. These MC productions are used for the calculation of the track selection and PID cut efficiencies for the heavy-flavor electrons sample ($\varepsilon^{\text{reco}}$ and ε^{eID}). All the event selection criteria applied in the data samples are also applied in the MC samples.

All samples include a full simulation of the detector response performed using GEANT 3.

5.4 Track Selection

Once the events are selected, the tracks of a given event need to pass through some quality requirements. These requirements aim mainly to select a sample of tracks that can represent the kinematics of the particles with good quality, for example: the tracks should be within a region in the detector not affected by border effects, they should have

a minimum number of ionized clusters in order to give a precise information about the momentum and energy loss of the particles in the gas or material, etc.

The track quality cuts applied in this analysis follows the criteria applied in former analysis (for example [13]), and a brief discussion of them can be found below.

Acceptance:

- The analysis is performed using tracks inside the pseudo-rapidity range $|\eta| < 0.8$, to make sure that the tracks do not go through the end-caps of the TPC. For electrons, that have very low mass, the rapidity y and pseudo-rapity η converges to the same value, such that this cut is equivalent to a rapidity cut of $|y| < 0.8$ (see [Appendix B](#));

TPC tracks:

- Tracks which include a significant number of TPC clusters originating from more than one charged particle trajectory were rejected by requiring that the χ^2/ndf (where ndf is the number of degrees of freedom) of the momentum fit in the TPC is smaller than 4;
- The track quality in the TPC is characterized by the number of clusters used for the reconstruction of the track. Only tracks with at least 100 (out of a maximum of 159) clusters used for their reconstruction were accepted in this analysis. The requirement on the number of TPC clusters is used to improve the discrimination between electrons and pions, because electron tracks have, on average, a higher number of clusters;
- Not all clusters used for track reconstruction are used for the energy loss calculation. For example, clusters close to the border of TPC sectors are not considered. Only tracks with at least 80 clusters used for the energy loss calculation were accepted in this analysis;

ITS tracks:

- A minimal number of 3 hits in the ITS is required, and two of them need to be on both SPD layers, in order to minimize the contribution of electrons originated from gamma conversions in the ITS. With this cut, only electrons from gamma conversion in the beam pipe or in the first SPD layer are accepted;
- In order to reject background tracks and non-primary tracks, the maximal distance of closest approach (DCA) of the tracks to the primary vertex is restricted to 1 cm in radial (xy) direction and 2 cm along the beam (z) direction;

Only global tracks, the ones created by matching TPC and ITS tracks, are used in the analysis.

A summary of the track quality cuts discussed above can be found in [Table 5.2](#).

Property	Electron candidate
p_T range (GeV/c)	0.5–4
$ y $	< 0.8
Number of TPC clusters	≥ 100
Number of TPC clusters in dE/dx calculation	≥ 80
$\chi^2/\text{clusters}$ of the momentum fit in the TPC	< 4
DCA_{xy}	$< 1 \text{ cm}$
DCA_z	$< 2 \text{ cm}$
Number of ITS hits	≥ 3
Number of hits in the SPD layers	2 (kBoth)

Table 5.2: Summary of the track selection criteria used in the pp at $\sqrt{s} = 13 \text{ TeV}$ analysis.

5.5 Selection of Electron Candidates

In this analysis, the information from TPC and TOF detectors is used to identify electrons. In [Figure 5.1](#) it is shown the $n\sigma_{\text{TPC}}$, that is the distance, in number of standard

deviations of the electron signal in the TPC (σ_{TPC}), between the $\langle dE/dx \rangle$ of a given track and the expected $\langle dE/dx \rangle$ for electrons (as discussed earlier in Equation 4.2). The electron signal is found around $n\sigma_{\text{TPC}} = 0$. In the left panel, it is possible to see a large contamination from protons, pions and kaons crossing the electron signal. The electron selection starts by rejecting particle tracks based on their time-of-flight, measured by TOF, with respect to the expected time-of-flight for electrons: only tracks with $|n\sigma_{\text{TOF}}| < 3$ are accepted, where $n\sigma_{\text{TOF}}$ is defined analogously to $n\sigma_{\text{TPC}}$. In the p_{T} -region up to about 2 GeV/c, the information from TOF is essential to reduce the hadron contamination on the electron sample, as can be noticed by comparing the two panels of Figure 5.1.

A further refinement of the electron selection can be reached by introducing a rejection of tracks based on the dE/dx of electrons in the TPC. Tracks with a $n\sigma_{\text{TPC}}$ between -1 and 3 are selected as electron candidates. This cut is asymmetric in order to minimize the pion contamination, that have negative values of $n\sigma_{\text{TPC}}$, without compromising the abundance of the electron signal (see right panel of Figure 5.1).

A hadron contamination remains in the electron candidates sample, and the procedure used to identify and remove it from this sample is described in section 5.7.

Figure 5.1: The $n\sigma_{\text{TPC}}$ as a function of the track momentum before (left) and after (right) applying the PID selection with TOF. The TPC PID selection is shown as the red lines on the right panel and all tracks inside this band are used for the further analysis steps.

A summary of the the PID cuts applied on TPC and TOF signals to select electron candidates is shown in Table 6.3

p_T range (GeV/c)	TPC dE/dx cut	TOF compatibility with e hypothesis
0.5–4	$-1 < n_{\sigma}^{\text{TPC}} < 3$	$ n_{\sigma}^{\text{TOF}} < 3$

Table 5.3: Summary of the electron identification criteria used in the pp at $\sqrt{s} = 13$ TeV analysis.

5.6 Efficiency of Track and PID Cuts

Next, it is shown the efficiency of the track and PID cuts, discussed in the previous sections, for the heavy-flavor electron sample. These efficiencies are calculated using the MC event samples with a trigger on heavy-flavor hadrons decaying semi-electronically. This results in large abundance for the heavy-flavor electrons sample and allows a precise determination of these efficiencies.

The track reconstruction efficiency, $\varepsilon^{\text{reco}}$, comprises the fraction of heavy-flavor electrons that passes the track quality cuts shown in [Table 5.2](#). For both numerator and denominator of the efficiency, tracks within $|\eta| < 0.8$ are considered. This efficiency is shown in black in the left panel of [Figure 5.2](#). In this same figure is shown $\varepsilon^{\text{reco}}$ combined with the efficiency of electron identification by the PID cut on TPC (blue), and combined with the efficiencies of the PID cuts on TPC and TOF (red) together. This last efficiency, $\varepsilon^{\text{reco}} \times \varepsilon^{\text{elD}}$, is the final one used to correct the heavy-flavor electron yield that will be obtained in the end of the analysis.

The right panel of [Figure 5.2](#) shows the efficiency of electron identification by the PID cuts, ε^{elD} , disentangled from the track reconstruction efficiency. It is possible to see that the efficiency of the TPC PID cut, $-1 < n_{\sigma\text{TPC}} < 3$, shown in blue, is constant at $\approx 84\%$. The efficiency of TOF PID cut, $|n_{\sigma\text{TOF}}| < 3$, shown in green, reaches 70% at high p_T and falls steeply at low p_T , reflecting the less precise resolution of the electron time-of-flight measured by TOF at low momentum. The efficiency of both PID cuts in TPC and TOF together is shown in red.

Studies were made to guarantee that the TPC PID efficiency, estimated from MC, can be used to correct the data without introducing any bias to the final measurement.

Figure 5.2: (Left) Efficiency of track+PID cuts and (right) only PID cuts as a function of the transverse momentum for electrons from heavy-flavor decays.

In Figure 5.3 it is shown the $n\sigma_{\text{TPC}}$ distribution as function of the track momentum, after the TOF PID cut, in data (left panel) and in MC (right panel) samples. In both cases the electron distribution is fitted by a Gaussian, in many intervals of p , and the mean and sigma of these fits are extracted and indicated in the plots. These parameters of the electron $n\sigma_{\text{TPC}}$ distribution are around their expected values, for both data and MC: the mean is around zero and the sigma is around unity. Consequently, since no discrepancy is observed between data and MC, the efficiency of the TPC PID cut estimated from MC can correct properly the data without introducing any bias. In these plots only the electron distributions are considered for the fits.

Similar studies were performed comparing the time-of-flight signal from TOF in data and in MC. The similarities between these samples ensure that the efficiency of the TOF PID cut estimated from MC can correct the data without introducing biases.

Figure 5.3: $n\sigma_{\text{TPC}}$ distributions, after the TOF cut, for data (left panel) and MC (right panel) together with the values of the mean (black) and sigma (red) extracted from the fit of the electron distribution by a Gaussian.

5.7 Remaining Hadron Contamination

In this section it is discussed the procedure used to remove the hadron contamination that remains in the electron candidate sample selected by the TOF and TPC PID cuts. This remaining hadron contamination is estimated statistically. The procedure is to build a hadron contamination spectrum by assigning to each track a weight which describes the probability of the track to be a hadron. After that, the hadron contamination spectrum can be subtracted from the electron candidate spectrum. The procedure to obtain the weight is described below.

The $n\sigma_{\text{TPC}}$ distribution, after the rejection of part of the hadron contamination by the TOF PID cut, is projected in many momentum slices, like the one shown in [Figure 5.4](#). The signals corresponding to hadrons and electrons are fitted simultaneously by a mathematical function $f(x)$ that is a sum of three parts: one part aims to describe the $n\sigma_{\text{TPC}}$ of electrons ($f_e(x)$), a second part aims to describe the $n\sigma_{\text{TPC}}$ of pions and muons ($f_\pi(x)$), and the third part aims to describe the $n\sigma_{\text{TPC}}$ from kaons and protons ($f_p(x)$):

$$f(x) = f_p(x) + f_\pi(x) + f_e(x). \quad (5.2)$$

As discussed in [section 4](#), the TPC signal corresponds to several measurements of the

energy loss of charged particles in an effective track length and, therefore, should follow the Landau distribution. In order to remove the large Landau-tail of the TPC signal distribution, a truncated mean requirement is applied during the $\langle dE/dx \rangle$ calculation, which modifies the distribution. While, after the truncated mean, the signal of electrons, protons and kaons can be nicely described by a Gaussian, the pion signal is still not completely Gaussian-like. Instead, it can be well represented by a Landau multiplied by an exponential function. So the pieces that compose the Equation 5.2, used to fit the total $n\sigma_{\text{TPC}}$ distribution, are chosen to be:

- $f_p(x) = A_1 e^{-\frac{(x-A_2)^2}{2A_3^2}}$ fits protons and kaons;
- $f_\pi(x) = B_1 e^{B_2 x} \times \text{Landau}(x, C_1, C_2, 0)$ fits pions e muons;
- $f_e(x) = D_1 e^{-\frac{(x-D_2)^2}{2D_3^2}}$ fits electrons;

where A_n , B_n , C_n and D_n are free parameters.

Figure 5.4: Distribution of the $n\sigma_{\text{TPC}}$, in the momentum slice between 2.3 to 2.5 GeV/c, with a simultaneous fit of the electron (green), pion (magenta) and kaon (dashed blue) distributions. The open black markers around unity indicates the ratio between the data and the fit.

Figure 5.4 shows the fit of the $n\sigma_{\text{TPC}}$ signal to the Equation 5.2, in a particular momentum slice. The distributions of electrons (green), pions and muons (magenta) and kaons and protons (dashed blue) are fitted simultaneously. The ratio of the data to the fit (open markers) is around unity, pointing to the good quality of the fit.

The hadron contamination in each of the momentum slices of $n\sigma_{\text{TPC}}$ is evaluated according to Equation 5.3:

$$\text{cont.} = \frac{\int_{x_{\min}}^{x_{\max}} (f_{\pi}(x) + f_{\text{K}}(x)) dx}{\int_{x_{\min}}^{x_{\max}} (\text{data}) dx}, \quad (5.3)$$

where $x_{\min} = -1.0$ and $x_{\max} = 3.0$ correspond to the TPC PID cut, $f_i(x)$ are the fit functions for the respective particle type i , and the denominator is the integral of the data itself.

Figure 5.5 shows the hadron contamination obtained in the different momentum bins, in logarithmic scale. It is parameterized using two different functional forms, an Error function (Erf) and a Landau.

The Error function:

$$f(p) = 0.010113(5) + 0.010114(5) \times \text{Erf}(0.75(7) \cdot p - 3.2(2)) \quad (5.4)$$

describes the hadron contamination best. The Landau function is used to evaluate the systematic uncertainty due to this parametrization (as will be discussed in section 5.10). The parameters of the Landau function, obtained from the fit, are:

$$f(p) = 0.6(1) \times \text{Landau}(p, 6(2), 1.6(7), 0). \quad (5.5)$$

Equation 5.4 is used as the weight which describes the probability of a track with momentum p , $-1 < n\sigma_{\text{TPC}} < 3$ and $|n\sigma_{\text{TOF}}| < 3$ to be a hadron. A p_{T} spectrum of this remaining hadron contamination is obtained by assigning to the tracks of transversal momentum p_{T} (and that passes the PID cuts) this weight, which is calculated according to the track momentum p .

The effect of subtracting the remaining hadron contamination from the electron candidate p_{T} spectrum is shown in Figure 5.6. The hadron contamination reaches up to

about 2% of the yield at $p_T = 4\text{--}4.5$ GeV/ c . At low and intermediate p_T , the hadron contamination accounts for less than 1% of the inclusive electron yield.

After all these selection steps and the removal of the remaining hadron contamination, it is obtained a raw inclusive electron p_T spectrum.

Figure 5.5: The fraction of hadrons that remains together with the electron candidates sample even after the PID cuts on TPC and TOF.

Figure 5.6: Ratio between the inclusive electron spectra after and before the remaining hadron contamination subtraction.

5.8 Non Heavy-Flavor Electron Identification

The raw inclusive electron spectrum, obtained via the PID cuts on TPC and TOF and removal of the remaining hadron contamination, contains heavy-flavor electrons plus background electrons that can come from many sources:

- Dalitz-decays of light neutral mesons (π^0 , η and η');
- Gamma conversion;
- Dielectron decays of light mesons (ρ , ω and ϕ);
- Dielectron decays of heavy quarkonia (J/ψ , Υ , ψ');
- Weak $K \rightarrow e\pi\nu$ (K_{e3}) decays;
- Partonic hard scattering processes, like prompt real and virtual photons conversion;

Figure 5.7 shows the inclusive electron yield measured by ALICE in pp at $\sqrt{s} = 7$ TeV collisions compared to the background of non heavy-flavor electrons, from a cocktail simulation [13]. It is possible to see that the dominating sources of background electrons are Dalitz-decays of light neutral mesons, mainly π^0 and η , and gamma conversions. The dielectron decays of heavy quarkonia (J/ψ) get important towards high p_T , while the contribution from the weak kaon decays (K_{e3}) gets important at very low p_T .

The contribution of electrons from ρ , ω , ϕ and η' via dielectron decays or Dalitz decays, is negligible for practical purposes. The same can be said about the contribution of electrons from direct photon conversion. The bottom panel of this figure shows the ratio between the inclusive electron sample and the non heavy-flavor electron background obtained from the cocktail. From this ratio it is possible to see that the low p_T inclusive electron spectrum is dominated by the background, while the contribution of heavy-flavor electrons becomes important at higher p_T .

In this analysis, the main non heavy-flavor electron background, composed by electrons from Dalitz-decays of light neutral mesons and gamma conversions, is estimated and subtracted from the inclusive electron spectrum via the data-driven method described below. The other sources of background that have secondary importance, namely

Figure 5.7: Non-heavy flavor electron sources in pp at $\sqrt{s} = 7$ TeV together with the measured inclusive electron cross section. In the bottom panel, the ratio between the inclusive electron yield and the background estimated by the cocktail shows an increasing trend, indicating the dominance of heavy-flavor electrons at high p_T [13].

K_{e3} processes and J/ψ decays, will be estimated and subtracted for future publication by the usage of a cocktail simulation. Considering that these contributions depends on the cuts for track selection applied in the analysis and that here are used similar cuts from previous analysis, it is known that the contribution of the K_{e3} to the non heavy-flavor electron background amounts to $\approx 0.5\%$ in the p_T interval 0.5–0.7 GeV/c, decreasing

even more towards larger p_T , and the contribution from J/ψ is about 2% in the p_T interval between 3–4 GeV/ c [13].

The **photonic tagging method** is the data-driven method adopted for the identification of the non heavy-flavor electrons background. This method exploits the fact that the electrons from Dalitz decay of π^0 ($\pi^0 \rightarrow e^-e^+\gamma$, BR = $1.174 \pm 0.035\%$ [108]) and η ($\eta \rightarrow e^-e^+\gamma$, BR = $6.9 \pm 0.4 \times 10^{-3}$ [108]), that are very light mesons, or from gamma conversion ($\gamma \rightarrow e^-e^+$) are created in electron-positron pairs. Consequently, electrons from these sources can be tagged in data directly by the low invariant mass of the pair. These pairs must have invariant mass around the mass of their mother, shown in Table 5.4. Hence, the non heavy-flavor background electrons are reconstructed by pairing electrons (positrons) with tracks identified as positrons (electrons) and the pairs with low mass are tagged and subtracted from the inclusive electron yield as explained below.

mother	e^-e^+ invariant mass (GeV/ c^2)
π^0	≈ 0.13
η	≈ 0.54
gamma	≈ 0
mass cut	< 0.1

Table 5.4: Invariant mass of the main non heavy-flavor electron background.

In order to build the invariant mass distribution, the electron (positron) candidates, which fulfill the strict selection criteria discussed above (shown in Table 5.2), are paired with tracks identified as positrons (electrons), that fulfill the less strict cuts shown in Table 5.5.

A pool of unlike sign (ULS) pairs is built, and it contains pairs from the non heavy-flavor electron sources mentioned above and also a combinatorial background. The combinatorial background is estimated from like-sign (LS) pairs, which are built up by pairing electron (positron) candidates and tracks identified as electrons (positrons) by the cuts in Table 5.5. This invariant mass analysis is performed for each electron p_T -bin.

The invariant mass distribution of unlike-sign and like-sign pairs for one p_T -bin is shown in Figure 5.8.

Property	Associated track
p_T range (GeV/ c)	≥ 0
$ y $	< 0.8
$n\sigma_{\text{TPC}}$	-3.5 to 3.5
Number of TPC clusters in dE/dx calculation	≥ 60
$\chi^2/\text{clusters}$ of the momentum fit in the TPC	< 4
DCA_{xy}	< 1 cm
DCA_z	< 2 cm
Number of ITS hits	≥ 3
Number of hits in the SPD layers	2 (kBoth)

Table 5.5: Summary of the criteria used to select the tracks that are paired to the electron candidates for the background tagging.

Figure 5.8: Invariant mass distributions for the unlike- and like-sign pairs.

For each p_T -bin, the yield of non heavy-flavor electrons is evaluated using [Equation 5.6](#):

$$N_{\text{nHFE}} = \frac{N_{\text{ULS}} - N_{\text{LS}}}{\varepsilon_{\text{tagging}}}, \quad (5.6)$$

where N_{ULS} and N_{LS} are the amount of unlike- and like-sign pairs with an invariant mass smaller than the requirement on the pair mass, respectively, and $\varepsilon_{\text{tagging}}$ is the tagging efficiency (that will be discussed below).

The cut on the invariant mass of the pair is chosen to be $m_{ee} < 0.1 \text{ GeV}/c^2$, as also indicated in [Table 5.4](#). All pairs within this cut are used for the calculation of the yield of non heavy-flavor electrons according to [Equation 5.6](#).

5.8.1 Non Heavy-Flavor Electron Tagging Efficiency

The tagging efficiency $\varepsilon_{\text{tagging}}$ quantifies the probability that a non heavy-flavor electron in a given p_T -bin is measured by the tagging method presented above. For the determination of this efficiency, general purpose (minimum bias) MC simulations are used. The tagging efficiency is calculated according to [Equation 5.7](#) as the ratio of non heavy-flavor electrons for which the partner is found in the pool of associated tracks (N_{found}) divided by all electrons from non heavy-flavor sources (N_{nHFE}), for each electron candidate p_T bin.

$$\varepsilon_{\text{tagging}} = \frac{N_{\text{found}}}{N_{\text{nHFE}}} \quad (5.7)$$

The MC simulation needs to undergo some corrections before the calculation of the tagging efficiency is performed. This is necessary because the p_T spectra slope of the π^0 and η mesons in the MC do not match the corresponding slopes found in data, mainly at low p_T , as will be discussed in more detail below. This difference in slope can bias the

tagging efficiency calculation in MC, since it changes slightly the p_T distribution of the electrons from π^0 and η decays, making this efficiency not representative of the tagging efficiency in data. Therefore the slopes of the π^0 and η spectra in MC are corrected to match the corresponding slopes in data as described below.

Correction of π^0 and η Spectra in MC In the left panel of [Figure 5.9](#) the π^0 spectrum from MC is compared to the charged pion spectrum measured in pp collisions at $\sqrt{s} = 13$ TeV. The measured charged pion spectrum is used instead of the measured π^0 spectrum in this comparison with the MC because it is known that both of these measured spectra have essentially the same shape, and the charged pion spectrum have smaller uncertainties relatively to the π^0 spectrum. Consequently, the use of the charged pion spectrum leads to smaller systematic uncertainties associated to the correction that will be presented here. In the right panel of [Figure 5.9](#) it is shown the comparison between η spectrum from MC and η spectrum obtained from m_T -scaling of the measured charged pions spectrum. The m_T -scaling [\[110\]](#) [\[111\]](#) is a property of some charged hadrons of having the same slope in the transversal mass (m_T) spectra, where $m_T = \sqrt{p_T^2 + m^2}$.

In the selection of the π^0 and η spectra from MC, is not included the ones coming from secondary decays (from feed-down of light meson or heavy-flavor hadrons). In this way the spectrum from MC is comparable to the spectrum measured in data, in the sense that both contain only primary particles. It is possible to see in [Figure 5.9](#) that the difference in slope between the spectra from data and MC occurs mainly at low p_T .

A weight for π^0 and η is calculated as the ratio $data/MC$ of the distributions shown in [Figure 5.9](#). These weights can be seen in [Figure 5.10](#) and they are fitted by a Hagedorn function, $f(x) = p_0/(e^{-(p_1x+p_2x^2)} + (x/p_3))^{p_4}$. The values given by the fitted function is used to weight the p_T -spectrum of electrons coming from π^0 and η in MC according to the p_T of the electron mother.

For the tagging efficiency calculation, the electrons from conversion are chosen just if the γ mother comes from a π^0 or η meson decay. The π^0 decay via the channel $\pi^0 \rightarrow \gamma\gamma$, $BR = 98.823 \pm 0.034\%$ [\[108\]](#), is the main contribution for γ production. For electrons coming from gamma conversion, the p_T of their grandmother is used to evaluate the

correct weight from the fitting function.

Figure 5.9: Spectrum of prompt π^0 (left) and η (right) from the MC simulations reported together with the measured charged pion spectrum and η spectrum (obtained from m_T -scaling), respectively.

Figure 5.10: Weights calculated for π^0 (red) and η (blue) as the ratio between the p_T spectrum of charged pions measured in data (η taken from m_T -scaling) and the π^0 (η) spectrum taken from Monte Carlo, fitted by a Hagedorn function.

With this weighting procedure, the non heavy-flavor electron spectrum that will be used for the calculation of the tagging efficiency has a p_T -shape consistent to data, so the tagging efficiency can be used to properly correct the non heavy-flavor electron spectrum measured in data by the tagging method.

The tagging efficiency calculated from the MC after weighting the electrons coming from π^0 , η or gammas (that comes from π^0 and η) is shown in [Figure 5.11](#) as function of the electron p_T . The top panel refers to the tagging efficiency of electrons from these three sources combined, while the bottom panel shows the tagging efficiency for each electron source separately. The tagging efficiency for electrons coming from η decays is systematically lower than the one from π^0 or gamma conversion because of the larger η mass (see [Table 5.4](#)).

The ratio between the inclusive (Signal (S) + Background (B)) and the non heavy-flavor electron spectrum (Background), obtained by the photonic tagging method described above, is shown in [Figure 5.12](#). This ratio can be thought as $(S + B)/(B) = (S/B) + 1$, where S refers to the signal of heavy-flavor electrons. As expected, for very low p_T the inclusive spectrum is dominated by non heavy-flavor electron background, and the ratio is close to unity. The contribution of heavy-flavor electrons increases significantly with p_T , overcoming the background for $p_T \gtrsim 2.5$ GeV/ c . This ratio shows excellent agreement to what is obtained in pp collisions at lower center of mass energies, where the non heavy-flavor background is estimated with the use of the cocktail, like in pp at $\sqrt{s} = 7$ TeV (shown at [Figure 5.7](#)) and in pp at $\sqrt{s} = 2.76$ TeV (shown in reference [\[112\]](#)).

ALI-PREL-133151

Figure 5.11: Tagging efficiency as function of p_T of the electron candidate calculated for all sources combined (top) and for each source separately (bottom).

Figure 5.12: Ratio between inclusive and non heavy-flavor electron spectra as function of p_T .

5.9 Normalization and FONLL Prediction

The raw yield of heavy-flavor electrons, $N_{\text{raw}}^{e\pm}$, is obtained via the subtraction of the non heavy-flavor electron yield from the inclusive electron yield. In this section, the other ingredients used for the determination of the invariant cross section of heavy-flavor electrons, according to [Equation 5.1](#), are resumed.

Number of Events The number of analyzed events corresponds to the events selected by the MB trigger and that also passed the event selection criteria described in [section 5.2](#). These events necessarily have at least two tracks in the central barrel, since this is the minimum number of tracks required by the event selection criteria for the construction of the primary vertex. The number of analyzed events are referred to as N_{vertex} , since they have a vertex from tracks. But the MB trigger can also accept events that generated just signal on the VZERO arrays and no (or less than two) tracks in the central barrel. These events are not analyzed in this work (since they don't have enough tracks) but they need to be taken into account for the calculation of the number of events considered in the analysis, that is the final number of events used for the determination of the heavy-flavor electrons cross section shown in [Equation 5.1](#), $N_{\text{ev}}^{\text{V0}}$.

$N_{\text{ev}}^{\text{V0}}$ is calculated as a sum of the number of analyzed events N_{vertex} and a fraction of the number of events selected by the MB trigger with no tracks in the central barrel:

$$N_{\text{ev}}^{\text{V0}} = N_{\text{vertex}} + f_{\text{vertex}} \times N_{\text{no vertex}}, \quad (5.8)$$

where:

- N_{vertex} denotes the number of events with a vertex from tracks which have passed the event selection (analyzed events);
- $N_{\text{no vertex}}$ is the total number of events selected by the MB trigger where no vertex from tracks could be found;
- f_{vertex} is the fraction of events with a vertex from tracks, with respect to all events selected by the MB trigger, which have passed the event selection criteria;

Cross Section of pp at $\sqrt{s} = 13$ TeV Collisions The visible cross section of pp at $\sqrt{s} = 13$ TeV collisions, referred as σ_{V0} , is measured via van der Meer (vdM) scans [113], with the use of two different trigger detectors. Events triggered by VZERO detector were found to have a cross section of $\sigma_{V0} = (57.8 \pm 1.2)$ mb and events triggered by T0 detector were found to have a cross section of $\sigma_{T0} = (30.1 \pm 0.6)$ mb [114].

Since the triggers used to select the minimum bias data sample used in this analysis are based in the signal in the VZERO detector, the σ_{V0} must be used for the normalization shown in Equation 5.1. However, the VZERO detector have shown instabilities during the pp cross-section measurement, and in order to avoid the direct use of σ_{V0} , it is used the property that

$$L = \frac{N_{ev}^{V0}}{\sigma_{V0}} = \frac{N_{ev}^{T0}}{\sigma_{T0}}, \quad (5.9)$$

where L is the luminosity of the beam and N_{ev}^{V0} and N_{ev}^{T0} are the number of events collected with the VZERO and T0 trigger, respectively. Then, it is possible to write:

$$\sigma_{V0} = \frac{N_{ev}^{V0}}{N_{ev}^{T0}} \sigma_{T0}. \quad (5.10)$$

The relation given by Equation 5.10 is used as replacement for the value $\sigma_{V0} = (57.8 \pm 1.2)$ mb in the normalization. In the used data sample it is found that $N_{ev}^{V0}/N_{ev}^{T0} = 1.9$.

Efficiencies The raw yield of heavy-flavor electrons is normalized to the efficiencies discussed in section 5.6. $\varepsilon^{\text{reco}}$ corresponds to the efficiency associated to the track selection and ε^{eID} is the efficiency associated to the PID cuts applied to TPC and TOF signals for electron identification.

Other Normalizations The raw yield of electrons from heavy-flavor hadron decays is normalized to unit of azimuthal angle, by dividing it by $\Delta\phi = 2\pi$, and to unity of rapidity, by dividing it by the rapidity range $\Delta y = 1.6$. For electrons, which have usually negligible mass compared to their momentum, the pseudo-rapidity and rapidity converges

to the same value (see [Appendix B](#)). Therefore the rapidity range is determined by the pseudo-rapidity range $\Delta\eta = 2 \times 0.8$.

Each p_T bin is divided by its width, Δp_T , and by its central value, p_T^{center} . A factor $1/2$ is considered so the result is representative of electron ($\frac{e^+ + e^-}{2}$) production only.

5.9.1 FONLL

In order to obtain the FONLL prediction for the production cross section of electrons from open heavy-flavor hadron decays in pp at $\sqrt{s} = 13$ TeV, the cross section of the different heavy-flavor electron species are taken separately from this pQCD calculation:

- electrons from B meson decays ($B \rightarrow e$);
- electrons from non-prompt D mesons (that come from B meson decays) ($B \rightarrow D \rightarrow e$);
- electrons from prompt D mesons ($D \rightarrow e$);

These p_T -differential cross sections are shown in the left panel of [Figure 5.13](#). The dashed lines correspond to the central value of the FONLL prediction and the continuous lines corresponds to the uncertainty range of the theoretical calculation, that comes from quark masses uncertainties, renormalization and factorization scale dependencies and PDF uncertainties. For prompt D mesons, are considered just the more abundant species, the D^0 and D^+ , and the theory already provides the proper branching ratios of these species to electrons ($0.7 D^0 + 0.3 D^+$).

The right panel of [Figure 5.13](#) shows the fraction, with respect to all heavy-flavor electrons, of electrons coming from $B(\rightarrow D)$ and prompt D meson decays in the FONLL prediction as function of p_T . In the low p_T region, the electrons come mainly from prompt D meson decays while for the higher p_T measured in this analysis the contribution of electrons from $B(\rightarrow D)$ decays reaches 50%.

Figure 5.13: The p_T -differential cross-section of electrons coming from the decay of heavy-flavor mesons (left panel) and fraction of electrons from B and D meson decays (right panel), given by the FONLL calculation.

5.10 Systematic Uncertainties

In this section, all the cuts and corrections applied in the analysis of this chapter are submitted to variations. The aim is to check if these cuts and corrections are introducing any systematic bias in the measurement. In such case, this bias needs to be quantified and assigned as systematic uncertainty to the final measurement.

The estimation of systematic uncertainty in the heavy-flavor electron yield is performed by classifying the different cuts and corrections in different categories, where each category represents cuts with a common physical origin. These cuts are varied and the systematic behavior associated to these variations are quantified. The considered categories are:

- Track quality cuts and PID cuts;
- Cuts related to the non heavy-flavor electrons background estimation;
- Weights applied on electrons that have π^0 or η as mother or grandmother;
- Pseudo-rapidity range cut;
- SPD requirement;
- Parametrization of the hadron contamination;

The final systematic uncertainty in the heavy-flavor electron yield is the combination of the uncertainties coming from the different categories. This combination is performed by adding the values in quadrature, under the assumption that the different sources are uncorrelated, which means that the biases found within the different categories are independent of each other.

Systematic uncertainties due to the track matching between TPC and ITS tracks and due to K_{e3} and J/ψ electron contribution are taken from previous studies and will be calculated for this data sample for future publication.

The requirements for the track selection and electron identification are varied in a similar way as in [13].

The strategy used for the estimation of systematic uncertainty for each category can be one of the two:

1. In order to spot any possible systematic trend due to a specific cut variation, the ratio var/ref , between the heavy-flavor electron yield obtained with the varied cut (var) to the yield obtained with default cut (ref), is analyzed. Ratios comparable to unity within statistical uncertainties indicates absence of systematic behaviors. The statistical uncertainties of these ratios are calculated using binomial error propagation, which takes into account that the numerator is a sub-set of the denominator (or vice versa) and, therefore, they are fully correlated.
2. In the case where many cut variations can be applied within the same category, the different estimates can be combined into a single contribution to the final uncertainty by the calculation of an overall spread or a mean. This procedure will be referred to as Root-Mean-Square (RMS) approach. For each bin of p_T of the heavy-flavor electron yield, the distribution $(ref - var)/ref$ is built, and values of the mean and RMS of this distribution are used to estimate the systematic trend due to these cut variations. The RMS approach can also be used to estimate the systematic uncertainty related to ratios from different categories in the case these ratios do not have itself a strong systematic behavior.

Below, each source of systematic uncertainty is discussed separately. A summary of

the final systematic uncertainty found for each source is presented in [Table 5.9](#) in the end of this chapter.

5.10.1 Systematic Uncertainties Coming from the Track Quality Cuts and PID Cuts

The track quality cuts and the PID cuts applied to select electron candidates are shown in [Table 5.6](#) together with the variations on those cuts used for systematic studies.

Source of Uncertainty	Reference	Variations
TPC clusters	≥ 100	$\geq 70, \geq 80, \geq 90, \geq 110,$ ≥ 120
TPC PID clusters	≥ 80	$\geq 50, \geq 60, \geq 70, \geq 90, \geq$ 100
ITS hits	≥ 3	$\geq 2, \geq 4$
$DCA_{xy}[cm]$	< 1	< 2.4
$DCA_z[cm]$	< 2	< 3.2
TOF PID	$\pm 3.0 \sigma$	$\pm 2.0\sigma, \pm 2.5\sigma$
TPC PID lower boundary	-1.0σ	$0.0\sigma, -0.5\sigma, -1.5\sigma$
TPC PID upper boundary	3.0σ	3.5σ

Table 5.6: Track and PID cut variations used for the study of systematic uncertainties.

[Figure 5.14](#) shows the ratios associated to track cut variations and PID cut variations separately. In general, no systematic trend is observed in these ratios. The only exception is the variation of the TPC PID selection $-1.5 < n\sigma_{\text{TPC}} < 3$ (shown in green in the top panel), that exhibits a systematic trend at high p_T . It is understood that this trend reflects the larger amount of hadron contamination that are included by this PID cut compared to the default cut, and that are not properly subtracted (since the parameterization of the hadron contamination obtained with the standard PID cut is used to determine the yields in all cases). The variation of the upper boundary of the TPC PID cut does not account for any considerable change in the spectrum.

Since, despite the specific case discussed above, no specific systematic trend is observed in the ratios related to track quality cuts and PID cuts variations, an RMS approach is used to estimate the systematic uncertainty due to these variations, and both categories are considered together.

Figure 5.14: Ratios of var/ref yields, where the variation is performed on the track selection cuts (top panel) and in the PID cuts (bottom panel).

Figure 5.15 shows, for the $1.5 < p_T < 1.75$ GeV/ c bin, the distribution of the deviation between the reference spectrum and all the varied spectra, normalized by the reference value. Similar distributions are provided for all the p_T intervals of the spectrum, and the mean and RMS of each distribution can be seen in Figure 5.16.

The mean of the distributions, shown in the left panel of Figure 5.16, fluctuates

Figure 5.15: Distribution of the difference between the varied and reference spectra, divided by the reference value, in one bin of p_T .

around zero, as demonstrated by the fit of these values by a zero-degree polynomial. No systematic trend is assigned to the behavior of the mean. The RMS as function of p_T , shown in the right panel of the same figure, have an approximate constant value around 1%. As a conservative estimation, the value of the RMS is attributed as systematic uncertainty for the heavy-flavor electron yield. It is a quantification of the spread of the variations around the reference value. At the same time, the RMS exclude effects related to statistical fluctuations of the variations, since it does not consider the maximum values of the variations by their own.

Figure 5.16: Mean and RMS of the distributions $(ref - var/ref)$, related to the track selection and PID cuts, as function of p_T .

5.10.2 Systematic Uncertainties due to the Subtraction of Electrons from non Heavy-Flavor Sources

The cuts associated to the identification of electrons from non heavy-flavor sources, that are the cuts on the e^+e^- pair invariant mass and on the minimum p_T and number of TPC clusters of the associated particles, are shown in [Table 5.7](#) together with the variations performed for the systematic studies.

Source of Uncertainty	Reference	Variations
Pair mass (GeV/c^2)	< 0.1	$< 0.07, < 0.08, < 0.09, < 0.11, < 0.12, < 0.14$
$p_T^{\min}$ (GeV/c)	0.0	0.1, 0.15, 0.2, 0.3
TPC clusters for associated particle	≥ 60	$\geq 50, \geq 70, \geq 75, \geq 80$

Table 5.7: Variation of cuts related to the identification of electrons from non heavy-flavor sources.

The ratios related to each cut are shown separately in [Figure 5.17](#), and while a small systematic behavior is observed at very low p_T , no particular systematic trend is observed for $p_T > 1.5 \text{ GeV}/c$.

In [Figure 5.18](#), the mean and RMS of the distributions ($ref - var/ref$) in each p_T -bin are shown. All the three cut categories are considered together to build these distributions, since no particular strong trend is observed within each category. The mean of the distributions fluctuates around zero and doesn't exhibit any systematic trend. The RMS, nevertheless, clearly shows a systematic trend as function of p_T . This is expected since at low p_T the inclusive electron spectrum is dominated by the non heavy-flavor electron background (like was shown in [Figure 5.12](#)), what makes this region more sensitive to the cuts related to the background estimation. This systematic trend of the RMS is fitted by three zero-degree polynomial functions, as shown in [Figure 5.18](#), and the parameters of the fits are used as a p_T -dependent systematic uncertainty related to background identification, varying from 2.5% at low p_T to 1% at high p_T .

Figure 5.17: Ratios of Varied/Reference heavy-flavor electron yield. Variations are performed on the mass cut (top panel), minimum p_T of the associated particle (middle panel) and minimum number of clusters in the TPC for the associated particle (bottom panel).

Figure 5.18: Mean and RMS of the distributions $(ref - var/ref)$ related to the non heavy-flavor electron background estimation as function of p_T .

5.10.3 Systematic Uncertainties due to π^0 and η Weights

In order to estimate the systematic uncertainty due to the weights that are applied to correct the slopes of the π^0 and η spectra from MC (as discussed in [subsection 5.8.1](#)), it is observed how the final heavy-flavor electron spectrum changes if reasonable variations on the weights are performed. The variations of the weight are produced by tilting up and down the charged pion spectrum as much as its own systematic uncertainties allows. In [Figure 5.19](#) one can see the fits that are performed on the reference charged pion spectrum (black) and on the tilted ones (green and blue). Two different p_T ranges of the fits are shown in each panel for better visualization of the effect of the tilt at low and high p_T .

The corresponding η spectra for each case (up and down tilt) is obtained by m_T -scaling of the tilted charged pion spectra.

Two modified weights, given by the ratio $data/MC$, are calculated using as *data* these tilted charged pion and η spectra (one for the tilted up and one for the tilted down cases). The systematic uncertainty associated to the use of these different weights are estimated looking at the ratio between the reference heavy-flavor electron yield and the yields obtained by using the tilted weights.

In this case, a clear systematic trend is observed in the ratios, as can be seen in [Figure 5.20](#). This is expected because the weights have an effect on the neutral meson

Figure 5.19: Fits of the charged pion spectra tilted up and down inside its systematic uncertainties together with the fit of the reference charged pion spectrum (black). Two plots are shown in order to highlight the tilt of the spectra at low (left panel) and high (right panel) p_T .

spectra only at low p_T ($p_T < 1.5\text{-}2$ GeV/ c), that is the region where the slopes of the spectra from MC and data differ more (like shown in [Figure 5.9](#) and [Figure 5.10](#)). At higher p_T , the effect of the weight is well known to have a negligible contribution, since it is approximately a constant. The ratio between the varied and the reference spectra, shown in [Figure 5.20](#), is fitted by a forth order degree polynomial, and this function is used to obtain the systematic uncertainty associated to the weight, as function of p_T . This systematic uncertainty ranges from 4% at low p_T to negligible values at higher p_T .

5.10.4 Systematic Uncertainties from Other Sources

The systematic uncertainties due to the remaining cuts applied in the analysis are discussed here. The sources of these systematics are presented in [Table 5.8](#), as well as the variations performed for the systematic studies.

Pseudo-Rapidity Range The analysis is performed using tracks inside $|\eta| < 0.8$, which is close to the maximum acceptance of the detectors used. In order to ensure that the result is not affected by being at the edges of the detectors, the heavy-flavor electron yield inside $|\eta| < 0.5$, $|\eta| < 0.6$ and $|\eta| < 0.7$ is determined and compared

Figure 5.20: Ratio of Varied/Reference heavy-flavor electron yield, where the variation is performed in the weight applied to correct the spectra of electrons coming from π^0 and η decays. The fit used to estimate the systematic uncertainty is shown by the red line.

Source of Uncertainty	Reference	Variations
η	-0.8 to 0.8	-0.5 to 0.5, -0.6 to 0.6, -0.7 to 0.7, -0.8 to 0, 0 to 0.8
SPD	kBoth	kAny, kFirst
Hadron parametrization	Error function	Landau
Track matching		
Remaining cocktail components		
$(J/\psi, K_{e3})$		

Table 5.8: Variation of the remaining cuts

to the reference result. In addition, this yield is also calculated in the asymmetric η regions of $-0.8 < \eta < 0$ and $0 < \eta < 0.8$, in order to check if the event selection is such that really avoids any bias in the particle production towards positive or negative η . The ratios relative to these variations are shown in [Figure 5.21](#), and no systematic uncertainty associated to any of the η -range is found.

Figure 5.21: Ratio of Varied/Reference heavy-flavor electron yield, where the variation is performed on the pseudo-rapidity interval considered.

SPD Requirement The SPD default requirement for the track selection is that tracks must have one hit in each SPD layer (kBoth). For the systematic studies this requirement is lowered to ask for at least one hit in any of the layers (kAny). This adds all electrons from gammas converting in the first layer or in the very first part of the second layer to the electron candidate tracks. In addition, the requirement was also lowered to ask for at least one hit in the first layer (kFirst). As can be seen in the ratios presented in [Figure 5.22](#), these variations on the SPD requirement leads to a systematic increase of the final heavy-flavor electron spectrum of about 3%. When kAny or kFirst requirements are applied, some of the additional electrons from gamma conversion that are included in the inclusive electron sample, with respect to the kBoth case, may end up accounted in the heavy-flavor electron yield, increasing this yield by about 3%.

Hadron Contamination The default parametrization for the hadron contamination is the Error Function, as was presented in [section 5.7](#). For systematic studies, the analysis is repeated with a different parametrization, given by the Landau function. The fit of those two functions to the hadron contamination was already shown in [Figure 5.5](#). The ratio between the final heavy-flavor electron spectrum corrected by these two different

Figure 5.22: Ratio of Varied/Reference heavy-flavor electron yield, where the variation is performed on the SPD requirement.

hadron contamination parameterizations is shown in [Figure 5.23](#). A negligible systematic uncertainty is associated to the different parameterizations, since even at high p_T , where a systematic trend appears, it accounts for less than 1%.

Figure 5.23: Ratio of Varied/Reference heavy-flavor electron yield, where the variation is performed by parameterizing the hadron contamination by a Landau function (while for the reference spectrum it is parametrized by an Error function).

Systematics Uncertainties Taken from Previous Measurements and Summary

- TPC-ITS matching: a single charged particle generates a track in the ITS and in the TPC, and both of these tracks can be matched to each other by offline algorithms, creating global tracks, that are used in the analysis. A mismatch between TPC and ITS tracks coming from the same particle can occur if the particle traverses inactive areas of one of the detectors, what prevents its reconstruction to be performed. This mismatch between TPC and ITS tracks that occurs in data not necessarily is reproduced properly in the MC, and the differences in the matching efficiency between TPC-ITS tracks in data and MC can lead to systematic effects in the final measurement. In this analysis, it is assigned a systematic uncertainty of 3% to the ITS-TPC track matching, taken from the charged hadrons spectra analysis in pp at $\sqrt{s} = 13$ TeV [115].
- Since the electrons coming from K_{e3} processes and J/ψ decays are still not subtracted in this analysis (the subtraction will be done for publication), a systematic uncertainty is assigned to their contribution. The values used as systematic uncertainty are the same ones used in pp $\sqrt{s} = 2.76$ TeV analysis: 3.8% for $0.5 \leq p_T < 0.7$, 1.1% for $0.7 \leq p_T < 0.9$ and 1.0% for $3.0 \leq p_T < 4.0$.

A summary of the systematic uncertainties from each category discussed above can be found in [Table 5.9](#).

In order to calculate the total systematic uncertainty, the different contributions discussed previously are summed in quadrature. At low p_T , the largest contributions to the systematic uncertainty come from the weight given to electrons coming from π^0 and η decays and also from K_{e3} process, while at higher p_T the track matching is the dominant source of systematic uncertainty. The final systematic uncertainties ranges from 4 to 7%, and the values in each p_T -bin can be found in [Table 5.10](#).

Source	Value
Track cuts	1%
Associated particle cuts	2.5% at $(0.5 \leq p_T < 1)$, 1% at $(p_T > 1)$
SPD requirements	3%
η range	negligible
Hadron Contamination	negligible
Weight for π^0 and η	$\approx 4\%$ at low p_T to negligible values at high p_T
Track matching	3%
Remaining cocktail components (J/ψ , K_{e3})	3.8% for $(0.5 \leq p_T < 0.7)$, 1.1% for $(0.7 \leq p_T < 0.9)$, 1.0% for $(3.0 \leq p_T < 4.0)$

Table 5.9: Summary of the systematic uncertainties in the HFE spectrum coming from the uncorrelated sources discussed above.

p_T (GeV/ c)	Systematic uncertainty (%)
0.5 - 0.6	7
0.6 - 0.7	7
0.7 - 0.8	6
0.8 - 0.9	5
0.9 - 1.0	5
1.0 - 1.1	5
1.1 - 1.2	5
1.2 - 1.3	4
1.3 - 1.4	4
1.4 - 1.5	4
1.5 - 1.75	4
1.75 - 2.0	4
2.0 - 2.25	4
2.25 - 2.5	4
2.5 - 2.75	4
2.75 - 3.0	4
3.0 - 3.5	4
3.5 - 4.0	5
4.0 - 4.5	5

Table 5.10: Final systematic uncertainty in each p_T -bin of the HFE spectrum.

5.11 Results and Discussion

Figure 5.24: The p_T -differential cross-section for electrons from heavy-flavor hadron decays using the photonic-electron tagging, compared to the theoretical prediction of this cross-section given by FONLL. On the bottom is shown the ratio between the measured cross section and the mean value given by FONLL together with the model prediction band.

Figure 5.24 shows the invariant p_T -differential cross-section of electrons from open heavy-flavor hadron decays measured in pp collisions at $\sqrt{s} = 13$ TeV by ALICE, from $0.5 < p_T < 4$ GeV/ c and central rapidity, compared to the theoretical prediction given by FONLL. The experimental points, shown in black, have statistical uncertainties (shown as vertical bars) and systematic uncertainties (shown as boxes). An additional uncertainty of 5% in the determination of the heavy-flavor electron cross-section is indicated

in the figure, and it comes from the determination of the visible cross section of the pp collisions, σ_{V0} . It is not included in the experimental data points since it is not related to the heavy-flavor electron yield extraction. The FONLL prediction corresponds to the sum of the production cross section of the three different heavy-flavor electron species described above in [subsection 5.9.1](#): electrons from B mesons, prompt and non-prompt D mesons decays. The central value of the FONLL calculation is presented together with its uncertainty band, defined by the filled red area. In the lower panel of this figure, the ratio between the heavy-flavor electron cross section and the central value of the FONLL prediction is shown together with the theoretical uncertainties relative to unity (obtained by the ratio between the upper and lower limits of the FONLL prediction and its central value).

The current result shows agreement with the upper limit of the FONLL prediction band. This same pattern is observed in many previous measurements in the heavy-flavor sector [\[9\]](#), some of them already presented in [section 3.1](#): measurements of heavy-flavor hadrons (B mesons [\[10\]](#), D mesons [\[31\]](#)), heavy-flavor muons [\[11\]](#) and heavy-flavor electrons in pp collisions at lower center of mass energies [\[112\]](#) [\[13\]](#) also can be described by FONLL and, in particular, are comparable to the upper limit of the theoretical calculation along most part of the p_T -range.

The FONLL prediction includes electrons from decays of heavy-flavor mesons but does not include electrons that come from heavy-flavor baryon decays. The measurement, nevertheless, includes electrons from all open heavy-flavor hadrons, including baryons. Since the fraction of electrons from heavy-flavor baryons relatively to heavy-flavor mesons is small (the Λ_c/D^0 ratio in pp at $\sqrt{s} = 7$ TeV ranges from ≈ 0.6 to 0.5 in the p_T interval of this analysis [\[42\]](#)), this absence of the baryon description in FONLL should not account for all the discrepancy between the measurement and the FONLL central values, that, as shown in the [Figure 5.24](#), is a factor of ≈ 2.2 at low p_T and ≈ 1.6 at large p_T .

The present measurement allows to test, for the first time, the pQCD prediction for the heavy-flavor electron cross section at a new collision energy, the largest collected by LHC up to now. In addition to this, this result is part of a bigger picture: ALICE collaboration has performed measurement of the production cross section of heavy-flavor

electrons in all different center of mass energies of pp collisions collected since run 1, namely $\sqrt{s} = 2.76$, $\sqrt{s} = 5.02$ and $\sqrt{s} = 7$ TeV. This energy scan of the heavy-flavor electron production cross section is shown in Figure 5.25, where the present measurement at $\sqrt{s} = 13$ TeV is included. Each measurement is compared to the FONLL prediction in the corresponding center of mass energy.

Figure 5.25: Heavy-flavor electron p_T -differential production cross section measured at central rapidity by ALICE. The measurements are done for pp collisions at $\sqrt{s} = 2.76$, $\sqrt{s} = 5.02$, $\sqrt{s} = 7$ and $\sqrt{s} = 13$ TeV. In red in shown the FONLL prediction. On the bottom of each panel is shown the ratio between the experimental data and central value of FONLL prediction together with the theoretical uncertainty band.

The FONLL calculation is dominated by large uncertainties, particularly in the charm sector, as can be seen in Figure 5.13. The dominating uncertainties are the ones from renormalization and factorization scales and from the quark mass values. At small p_T ($p_T \approx m_Q$, where m_Q is the heavy-quark mass), the scale uncertainty for charm can reach values of about $\approx 100\%$, and $\approx 50\%$ for beauty [15].

These large uncertainties may compromise the comparison between theory and experimental data. Nevertheless, the ratio between the production cross section measured at different energies is shown to be a good observable to test FONLL predictions since many of the theoretical uncertainties that are correlated and independent of the collision energy are significantly reduced in the ratio. The remaining theoretical uncertainties on the ratio have sensitivity mainly to the uncertainty on the PDF of the partons [15].

Among all LHC experiments, measurements at low p_T can be provided specially by ALICE due to its lower magnetic field ($B = 0.5$ T) with respect to the other experiments ($B = 2$ T for ATLAS, $B = 4$ T for CMS and an integrated magnetic field of 4 Tm for LHCb). With this configuration of magnetic field, the particles produced in the collision bend less in ALICE, and tracks with lower p_T can reach the necessary detectors. Low p_T measurements are important to access partons in the low Bjorken- x regime, since, for a given pseudo-rapidity, the probed parton x follows the relation $x \propto p_T/\sqrt{s}$. Consequently, ratios between heavy-flavor production cross section at different energies measured by ALICE can be used to constrain the gluon PDF, that is currently known with poor theoretical precision at low Bjorken- x .

In Figure 5.26 is presented the ratio between heavy-flavor electron cross sections measured in different energies: $\sqrt{s} = 5.02$ TeV/ $\sqrt{s} = 2.76$ TeV, $\sqrt{s} = 7$ TeV/ $\sqrt{s} = 2.76$ TeV and $\sqrt{s} = 13$ TeV/ $\sqrt{s} = 2.76$ TeV, respectively. These ratios are compared to FONLL ratios. With this observable, the uncertainties from theoretical calculation become comparable to the experimental ones. For the two lower energy ratios (left and middle panels), the FONLL describes the data without any bias, while for the $\sqrt{s} = 13$ TeV/ $\sqrt{s} = 2.76$ TeV ratio the experimental data shows a bias to the upper edge of the theoretical prediction. This behavior might indicate a possible theoretical tension in the description of the PDFs probed in large momentum transfer Q processes accessed in pp at $\sqrt{s} = 13$ collisions.

Considering the ratios of the measurement in pp at $\sqrt{s} = 13$ TeV to the ones performed at lower energies, shown in Figure 5.27, it becomes more clear that the new measurement performed in this work can add constraints to the theoretical calculation

Figure 5.26: Ratio between heavy-flavor electron production cross section measured at different center of mass energies: $\sqrt{s} = 5.02 \text{ TeV} / \sqrt{s} = 2.76 \text{ TeV}$, $\sqrt{s} = 7 \text{ TeV} / \sqrt{s} = 2.76 \text{ TeV}$ and $\sqrt{s} = 13 \text{ TeV} / \sqrt{s} = 2.76 \text{ TeV}$, respectively. The red bands indicate the ratio between FONLL predictions at the different energies.

in the low p_T region, mainly for $p_T < 1 \text{ GeV}/c$, where experimental and theoretical ratios follow different trends. Inputs from the theoretical side are necessary to further understand the constraining power of these ratios.

Figure 5.27: Ratio between heavy-flavor electron production cross section measured at different center of mass energies: $\sqrt{s} = 13 \text{ TeV} / \sqrt{s} = 2.76 \text{ TeV}$, $\sqrt{s} = 13 \text{ TeV} / \sqrt{s} = 5.02 \text{ TeV}$ and $\sqrt{s} = 13 \text{ TeV} / \sqrt{s} = 7 \text{ TeV}$, respectively. The red bands indicate the ratio between FONLL predictions at the different energies.

Chapter 6

Measurement of Electrons from Beauty Hadron Decays in Central Pb-Pb Collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV

6.1 General Strategy

In this chapter, the measurement of the invariant yield of electrons from beauty hadron decays in Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV is described. The nuclear modification factor, R_{AA} , is obtained by the ratio between this measurement to the equivalent invariant yield measured in pp collisions at the same center of mass energy, scaled by the proper Glauber model factor.

The invariant yield of electrons from beauty hadron decays is given by [Equation 6.1](#).

$$\frac{dN_{\text{AA}}}{dp_{\text{T}}} = \frac{1}{2N_{\text{ev}}} \frac{1}{2\pi p_{\text{T}}^{\text{center}}} \frac{1}{\Delta y \Delta p_{\text{T}}} \frac{N_{\text{raw}}^{e^{\pm}}(p_{\text{T}})}{(\varepsilon^{\text{reco}} \times \varepsilon^{\text{eID}})} \quad (6.1)$$

From all the Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV generated during the data taken period, N_{ev} events are analyzed. These events are the ones selected by the minimum

bias trigger, based on the signal in the VZERO detector, and that satisfy requirements that ensure the event quality. In addition, just the events in which the two Pb nuclei collide with each other with a large level of overlap are selected, corresponding to the 10% most central collisions.

From all the tracks generated in each event, just the ones that passed a quality selection criteria are used. This track selection has an efficiency $\varepsilon^{\text{reco}}$, that is estimated using simulations of Pb-Pb events generated by Monte Carlo.

The raw p_T -spectrum of electrons from beauty hadron decays, $N_{\text{raw}}^{e^\pm}(p_T)$, is obtained by a sequence of steps. Firstly, electron candidates are selected via PID cuts on the signals generated in TPC and TOF detectors. These cuts select an electron candidate sample, that consists in tracks which have a time-of-flight measured by TOF and energy loss $\langle dE/dx \rangle$ in the TPC compatible with the electron hypothesis. This sample of electron candidates is composed by electrons from beauty hadron decays, charm hadron decays, electrons from non heavy-flavor hadron decays and also a hadron contamination. The identification of electrons from beauty hadron decays is performed in this work by exploring the different shapes of the DCA (Distance of Closest Approach) distribution of electrons from these different sources, following the same technique used for the measurement of electrons from beauty hadron decays in Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 2.76$ TeV in ALICE [53]. The DCA of an electron is determined by prolonging its trajectory backwards in time until the distance of the closest approach of this trajectory to the primary vertex is found. The left panel of Figure 6.1 illustrates a B and D hadron decaying via their semi-leptonic channel and gives a hint that the decay length of the hadron influences directly the DCA of the electron, such that this quantity can reflect the electron source. Electrons from beauty hadron decays, due to the longer decay length of B hadrons, have a wider DCA distribution when compared to the background electrons, allowing their identification by this method. The analysis procedure consists in taking from Monte Carlo simulations templates of the DCA distribution of electrons from beauty hadron decays and from the background sources. The DCA distribution of the electron candidates in data is fitted by these Monte Carlo templates, in each p_T bin independently. An example of such a fit is illustrated in the right panel of Figure 6.1.

The fit procedure consists in a maximum likelihood fit, and the free parameters of the fit are the amplitude of each template. When summed together with the correct amplitudes (the ones that reflect the abundances of electrons from the different sources in data), the templates can reproduce the measured DCA distribution of inclusive electrons, indicating the good quality of the fits. The yield of electrons from beauty hadron decays in each p_T bin, $N_{\text{raw}}^{e^\pm}(p_T)$, can be determined once the amplitude of the template of electrons from beauty hadron decays in each p_T is known.

Figure 6.1: (Left) Sketch of a B and D hadron decaying via semi-leptonic channel and the backward prolongation of the trajectory of the electron for the determination of its DCA. (Right) Example of a template fit to the electron candidate DCA distribution performed for the analysis of LHC run 1 data [53].

This yield is corrected by the efficiencies of the track selection $\varepsilon^{\text{reco}}$ and of the PID cuts applied in the TPC and TOF signals in order to select electrons, ε^{elD} , also estimated from Monte Carlo simulations.

The other factors in Equation 6.1 are geometrical factors, that normalize the yield per unity of rapidity Δy and unity of azimuthal angle $\Delta\phi = 2\pi$, or factors used to remove the p_T bin size (Δp_T) and p_T bin center (p_T^{center}) dependence of the measured spectrum.

Along this chapter, the quantitative determination of each of the ingredients of Equation 6.1 is presented. In the end, the obtained invariant yield of electrons from beauty

hadron decays is compared to the pp reference and the R_{AA} is calculated. The R_{AA} of electrons from beauty hadron decays in this new collision energy is compared to theoretical models and other experimental results and discussed.

6.2 Data Set and Event Selection

In this analysis, events from a minimum bias data sample (LHC15o) recorded in Pb-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV in 2015 by ALICE are used.

An amount of ≈ 100 M events are available, from which ≈ 9 M were analyzed, corresponding to the ones that passed the event selection criteria described below.

In addition to the online MB trigger and offline event selection based on the VZERO detector, as was done in the pp analysis, in this analysis only the 10% most central events are used, as will be discussed in detail in [section 6.4](#). These events correspond to an almost complete overlap between the colliding nuclei.

The event selection also requires that the reconstructed primary vertex has at least two contributor tracks and that the distance of the primary vertex to the nominal collision point in the longitudinal direction is $|V_z| < 10$ cm.

The pile-up rejection in this case is performed by comparing the primary vertex found with global tracks, V_{trc} , with the primary vertex found using just the signal on SPD, V_{SPD} . The minimum number of track contributors for the determination of V_{SPD} is required to be 1. Correlation cuts between V_{trc} and V_{SPD} are applied. The event is rejected if at least one of the three conditions are satisfied:

- The absolute value of the difference between the z position of the two vertexes is greater than 0.2 cm;
- The absolute value of this distance normalized to the convolution of the errors of the two vertexes is larger than 10;
- The absolute value of this distance normalized to the error of the V_{trc} is larger than 20;

In the [Table 6.1](#) the event quality cuts are summarized.

Type	Value
Min. contributors to V_{trc}	2
Min. contributors to V_{SPD}	1
$ V_{\text{trc}}z $	$< 10 \text{ cm}$
$ V_{\text{trc}}z - V_{\text{SPD}}z $	$< 0.2 \text{ cm}$
$ V_{\text{trc}}z - V_{\text{SPD}}z / \sqrt{(V_{\text{trc}}err)^2 + (V_{\text{SPD}}err)^2}$	< 10
$ V_{\text{trc}}z - V_{\text{SPD}}z / V_{\text{trc}}err$	< 20

Table 6.1: Event quality cuts applied in the Pb-Pb at $\sqrt{s_{\text{NN}}} = 5.02 \text{ TeV}$ analysis.

6.3 Monte Carlo Samples

Three Monte Carlo (MC) samples anchored to the data sample are used in this analysis. One of these samples is a special PYTHIA MC simulation that contains enhancement of π^0 and η mesons and a trigger on heavy-flavor hadrons decaying semi-electronically. This MC sample contains events in the centrality class $0 - 10\%$. From this MC production the DCA templates are extracted and it is also used for the calculation of the track selection and PID cut efficiencies for electrons from beauty hadron decays ($\varepsilon^{\text{reco}}$ and ε^{eID}).

The other two MC samples are general purpose (minimum bias) MC productions and they are used for the studies of the mean and resolution of the DCA templates in MC. One sample contains events in the centrality class $0 - 10\%$ while the other one contains events in all centrality classes. The selection of the centrality class of interest, $0 - 10\%$, is performed within the analysis code.

6.4 Centrality Selection

In a nucleus-nucleus collision, the two nuclei can have different levels of overlap and this influences directly the amount of produced particles: the multiplicity of the collision increases linearly with the overlap region of the nuclei. Experimentally, the overlap between the colliding nuclei are classified according to the *centrality* of the collision.

Head-on collisions are referred to as *central* collisions, producing thousands of particles, while collisions with a very small overlap region are referred to as *peripheral* collisions, and produce much less particles. One can define the centrality of a collision as the percentile of observed events N_{ev} with a charged particle multiplicity above a given threshold ($N_{\text{ch}}^{\text{THR}}$):

$$c = \frac{1}{N_{\text{ev}}} \int_{N_{\text{ch}}^{\text{THR}}}^{\infty} \frac{dN_{\text{ev}}}{dN'_{\text{ch}}} dN'_{\text{ch}}. \quad (6.2)$$

In ALICE, many detectors can be used to estimate the charged particle multiplicity of the events for the centrality determination. In this analysis, the VZERO detector is used (the combined signal of V0A and V0C, as described in [section 4](#)). A distribution of the VZERO signal can be found in [Figure 6.2](#), where the signal amplitude is divided in centrality ranges: the 2.5% most central collisions corresponds to the 2.5% events with larger signal amplitude in the VZERO. This approach allows the determination of events within intermediate centrality classes, like the 10-20% most central events, and so on.

For this analysis, the 10% most central collisions are selected.

It is important to make the connection between the centrality and the geometrical parameters of the collision, that cannot be determined experimentally. Geometrically, the degree of superposition between two colliding nuclei is characterized by the impact parameter b , which is defined as the distance between the center of each nucleus in a plane transverse to the momentum of the nuclei. A collision with a given impact parameter b has an associated number of binary collisions (N_{coll}) and a number of participant nucleons (N_{part}), that is the number of nucleons in a nucleus that undergoes one or more binary collisions. The number of spectator nucleons, those that do not participate in the collision, is given by $N_{\text{spec}} = A - N_{\text{part}}$, where A is the total number of nucleons in one nucleus. The number of binary collisions that occurs in a given centrality class is used for the determination of the nuclear modification factor R_{AA} . The determination of the geometrical parameters is also useful for the comparison between results from collisions of different nuclei.

In order to establish this connection between the centrality classes and the geomet-

rical parameters, a Monte Carlo Glauber [45] [46] [47] [48] approach is used. In this approach, a model of the nucleus is built by sorting the position of the nucleons stochastically according to a given nuclear density function. The two nucleus are made to collide with an impact parameter b , which already defines N_{part} . The interaction of each nucleus is treated as a sequence of independent nucleon-nucleon collisions. Each nucleon travels on a straight-line trajectory and the nucleon-nucleon inelastic cross section, $\sigma_{\text{NN}}^{\text{inel}}$, is assumed to be independent of the number of collisions a nucleon underwent previously. The inelastic nucleon-nucleon cross section $\sigma_{\text{NN}}^{\text{inel}}$ can be either directly taken from measurements in pp collisions, or extracted from interpolations of the available data [47]. The Glauber Monte-Carlo model is then combined with a particle production model, which simulates the particles created in each nucleon-nucleon collision and models it as a negative binomial distribution (NBD), since this distribution is known to describe well the charged particle multiplicity measured in pp collisions [116].

The VZERO signal distribution is simulated through this approach and a fit to the experimental VZERO signal amplitude is performed in order to find the NBD parameters that can better describe the data.

Figure 6.2 shows in red the Monte Carlo Glauber fit to the charged particle multiplicity measured by VZERO. From this fit, it is possible to perform the correspondence between the geometrical parameters like N_{coll} , N_{part} and b to each class of centrality. Mean values of the geometrical quantities are calculated for each centrality class.

For the 10% most central Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV used in this analysis, the Monte-Carlo Glauber model gives $\langle N_{\text{coll}} \rangle = 1560$ and $\langle T_{\text{AA}} \rangle = 23.44 \text{ mbarn}^{-1}$ (calculated as $T_{\text{AA}} = N_{\text{coll}} / \sigma_{\text{NN}}^{\text{inel}}$, where $\sigma_{\text{NN}}^{\text{inel}} = 67.6 \pm 0.6 \text{ mb}$ for $\sqrt{s_{\text{NN}}} = 5.02$ TeV [47]).

Figure 6.2: Experimental VZERO amplitude shown in blue, divided in centrality classes, from the most central (0-2.5%) to the most peripheral (80-90%) events. In red is shown the NBD-Glauber fit to the experimental distribution [47].

6.5 Track Selection and its Efficiency

The track selection criteria applied on the current analysis is summarized in Table 6.2, and most part of the cuts are very similar to the ones applied in the previous analysis, described in detail in section 5.4. Nevertheless, two additional cuts are applied here, which are the ones highlighted in table Table 6.2. A qualitative motivation for these new cuts can be found below.

As discussed earlier in section 5.4, the tracks used in the analysis are required to have two hits on the SPD (*kBoth*). This requirement is made in order to suppress the huge background of electrons from gamma conversion by just accepting the ones that are created in the beam pipe or in the very beginning of the first SPD layer, rejecting the ones that are originated later. The first SPD layer is located at a distance of $R \approx 4$ cm from

Property	Electron candidate
p_T range (GeV/ c)	1.5–8
$ y $	< 0.8
Number of TPC clusters	≥ 100
Number of TPC clusters in dE/dx calculation	≥ 80
$\chi^2/\text{clusters}$ of the momentum fit in the TPC	< 4
DCA_{xy}	< 1 cm
DCA_z	< 2 cm
Number of ITS hits	≥ 4
$\chi^2/\text{clusters of the momentum fit in the ITS}$	< 5
f_{shared} ITS clusters	< 0.3
Number of hits in the SPD layers	2 (kBoth)

Table 6.2: Summary of the track selection criteria used in the Pb-Pb at $\sqrt{s_{\text{NN}}} = 5.02$ TeV analysis.

the nominal point, meaning that the *kBoth* requirement would accept only electrons from gamma conversion created at positions $R \lesssim 4$ cm. Nevertheless, while checking through MC simulation the position R where the electrons from gamma conversion are created (already considering the tracks that passed the *kBoth* requirement) it is possible to see that many of them are created after the first SPD layer, with $R > 4$ cm, like shown in the left plot of [Figure 6.3](#). These electrons are accepted by the *kBoth* requirement because their tracks are wrongly associated with hits (clusters) on the SPD during the track reconstruction, as a result of the high occupancy of the detectors during heavy-ion collisions. These tracks with misassociated hits on the SPD will be referred to as tracks from late gamma conversion. This effect is known to be present in the data and to be well reproduced in MC simulations [\[20\]](#).

It is important to notice that, as a consequence of their larger production radius R , the electrons from late gamma conversion have a broader DCA distribution with respect to the electrons from gammas converting in the beam pipe, as can be seen in the right

panel of Figure 6.3. It is important for the current analysis to reduce the amount of electrons from late gamma conversion, making the final DCA distribution of conversion electrons slimmer than and less competitive with the DCA distribution of electrons from beauty hadron decays.

Figure 6.3: (Left) Distribution of the production radius R of electrons from gamma conversion as taken from MB MC simulations. (Right) DCA distribution of electrons from gamma conversion with large production radius ($R > 3.5$ cm) and with small production radius ($R < 3.5$ cm).

Quality cuts on the ITS tracks were studied in order to investigate how to suppress electrons from late gamma conversion, as it was performed in previous works [20] [117]. Due to the misassociated hits on the SPD, ITS tracks from late gamma conversion will most probably have larger values of χ^2 of the momentum fit as well as a larger fraction of shared clusters with respect to ITS tracks of electrons from other sources.

The distribution of the χ^2/ndf of the momentum fit of ITS tracks is checked in the MB MC simulations for different electron species, as shown in the left panel of Figure 6.4. As expected, tracks of electrons from gamma conversion have a χ^2/ndf distribution with a longer tail, contrasting with the χ^2/ndf distribution of electrons from other sources. The tail should be populated mainly by electrons from late gamma conversion. The right panel of figure Figure 6.4 shows the distribution of the fraction of clusters a given ITS track shares with other tracks, f_{shared} , for different electron species. The electrons from gamma conversion tends to have a large amount of f_{shared} , while tracks of electrons from

other sources are more abundant in the region of low f_{shared} .

Monte Carlo studies show that the selection of tracks with small values of χ^2/ndf and f_{shared} leads to a sample with a reduced amount of electrons from late gamma conversion, making the final template of electrons from gamma conversion slimmer (see [Appendix E](#)). Cuts on χ^2/ndf and f_{shared} were optimized in order to increase the signal of electrons from beauty hadron decays with respect to the total background. The optimal cuts were found to be $\chi^2/ndf < 5$ and $f_{shared} < 0.3$.

Figure 6.4: Distribution of χ^2/ndf (left) and f_{shared} (right) of electrons from different sources as taken from the MB MC simulations. Electrons from late gamma conversion tend to have larger values of χ^2/ndf and f_{shared} than the electrons that convert earlier and other electron sources.

The reconstruction efficiency (ϵ^{reco}) for electrons from beauty hadron decays is the fraction of these electrons that passes the track quality cuts in [Table 6.2](#), as given by the simulation. This efficiency is shown in [Figure 6.5](#) as function of the track p_T . For both numerator and denominator, tracks within $|\eta| < 0.8$ are considered.

Figure 6.5: Reconstruction efficiency for electrons from beauty hadron decays in central Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV.

6.6 Selection of Electron Candidates in Data and its Efficiency

In this analysis, the TPC and the TOF are used as PID detectors to identify the electron candidates.

The electron selection starts by rejecting all tracks that have time-of-flight, measured by TOF, outside 3 sigmas from the expected value for electrons. Considering only tracks with $|n_{\sigma}^{\text{TOF}}| < 3$, a considerable amount of kaons, protons and deuterons that have values of TPC dE/dx crossing the electron TPC dE/dx distribution are removed, guaranteeing an electron candidate sample with less contamination, as can be seen by the comparison of panels (a) and (b) in [Figure 6.6](#);

A further refinement of the electron selection can be reached by introducing a rejection of tracks based on the deviation from the expected dE/dx for electrons in the TPC, see panel (b) of [Figure 6.6](#): tracks with a n_{σ}^{TPC} between -0.1 and 3 are selected as electron candidates, as indicated by the red lines in the figure. The asymmetry of this cut aims to avoid the large contamination of hadrons in the region of negative TPC dE/dx , that is populated mainly by charged pions.

Figure 6.6: $n\sigma_{\text{TPC}}$ as a function of the track momentum before (panel (a)) and after (panel (b)) applying the TOF PID cut. The TPC PID selection is shown as the red lines in panel (b) and all tracks inside this band are used for the analysis.

A summary of the PID cuts used to select the electron candidates in this analysis is shown in [Table 6.3](#).

p_{T} range (GeV/ c)	TPC dE/dx cut	TOF compatibility with e hypothesis
1.5–8	$-0.1 < n_{\sigma}^{\text{TPC}} < 3$	$ n_{\sigma}^{\text{TOF}} < 3$

Table 6.3: Summary of the electron identification criteria used in the Pb-Pb at $\sqrt{s_{\text{NN}}} = 5.02$ TeV analysis.

The efficiency of each PID cut, ε^{PID} , applied in the data sample can be estimated by the usage of the MC simulations, as far as those represents accurately the signal distributions, TPC dE/dx and time-of-flight, found in data.

The similarity between the time-of-flight signal from TOF in data and in MC ensure that the efficiency of the TOF PID cut estimated from MC can correct the data without introducing biases. The efficiency of the TOF PID cut is shown in [Figure 6.7](#) as function of the track p_{T} . It represents the fraction of electrons from beauty hadron decays that passed the TOF PID cut for electrons, as given by the simulation. The TOF PID efficiency reaches very low values at low p_{T} because the resolution of the time-of-flight

of electrons becomes worse as the p_T gets smaller.

The TPC PID efficiency is calculated in this analysis without the usage of the MC, since the mean value of the $n\sigma_{\text{TPC}}$ signal for electrons in data and in MC are slightly displaced with respect to each other. The shape of the electrons $n\sigma_{\text{TPC}}$ distribution is well known to be Gaussian-like, and ideally this Gaussian have center at $n\sigma_{\text{TPC}} = 0$ and width $\sigma = 1$, as happens in the MC simulation. In this data sample it was found that the Gaussian signal has $\sigma = 1$ but it is centered at $n\sigma_{\text{TPC}} = -0.1$, consequence of the imperfect calibration of the TPC signal. Since the TPC PID cut used to select electrons is $-0.1 < n_{\sigma}^{\text{TPC}} < 3$, it selects 50% of the electron Gaussian signal, so the efficiency of the TPC PID cut is taken as 50%, constant with p_T .

Figure 6.7: Efficiency of the TOF PID cut for the selection of electrons from beauty hadron decays, as function of p_T .

The final efficiency, obtained considering efficiencies of track and PID cuts in combination ($\varepsilon^{\text{reco}} \times \varepsilon^{\text{PID}}$), is shown in [Figure 6.8](#) as function of p_T . This efficiency, that amounts in about 14% on its plateau, is used to correct the final yield of electrons from beauty hadron decays.

Figure 6.8: Final efficiency ($\varepsilon^{\text{reco}} \times \varepsilon^{\text{eID}}$) for electrons from beauty hadron decays.

Unfolding of the p_T -spectrum In the Monte Carlo simulations it is possible to check that the true (generated) p_T of a particle is not the same as the p_T of the reconstructed track, created when the particle traverses the detector material or gas. The reconstruction of the track as well as the bremsstrahlung that occurs when the particle interacts with the detector material affect the reconstructed p_T , making it usually smaller than the true p_T , see [Figure 6.9](#). This means that the real p_T spectrum of a given particle is not directly accessible experimentally. Nevertheless, this p_T bin-migration that is embedded in the measured signal can be corrected by the unfolding procedure. In this analysis, the unfolding is performed by considering during the efficiency calculation (discussed above) the generated p_T spectrum from the MC as denominator and the reconstructed p_T spectrum from the MC as numerator. Hence, when correcting the measured electron spectrum by this efficiency it also undo this p_T bin-migration caused by reconstruction and detector effects, in a way that the corrected spectrum is closer to the true distribution generated in the collision. In low p_T analysis, like the ones performed in this thesis, the effect of the unfolding in the final spectrum is about 2%. At larger p_T , where the particles can be more affected by bremsstrahlung, this effect can be larger.

In the pp analysis presented in the previous chapter, the unfolding is performed via the same strategy as in the present analysis.

Figure 6.9: Reconstructed p_T vs generated p_T of electrons from beauty hadron decays in Pb-Pb at $\sqrt{s_{NN}} = 5.02$ TeV, taken from simulations.

6.7 Selection of Electrons in Monte Carlo: Building the Templates

In order to extract, from the DCA distribution of electron candidates, the signal of electrons from beauty hadron decays, DCA templates of electrons from different sources are taken from the MC simulations. The electrons from the MC are grouped in four different categories, each one with a distinct DCA distribution shape that reflects the electron source.

The first group of electrons to be considered is the one that consists in the signal to be measured, electrons from beauty hadrons decays, that will be referred to as **beauty electrons**. Electrons from this group have a wider DCA distribution when compared to background electrons. The wider DCA of the signal reflects the larger decay length of beauty hadrons, that ranges from $c\tau \approx 400 \mu m$ for the Λ_b^0 baryon to $c\tau \approx 500 \mu m$ for the B mesons. Beauty hadrons also can decay into hadrons that contains charm and then those can decay into electrons. This adds a considerable decay length for some of the electrons of this group, making the DCA shape of electrons from beauty hadron decays even wider.

Electrons coming from the decay of hadrons that contains a charm quark (and that does not come from beauty hadrons feeddown) are grouped together, and they will be referred to as **charm electrons**. They have a slimmer DCA distribution compared to beauty electrons. The decay length of charmed hadrons ranges from $c\tau \approx 60 \mu m$ for the Λ_c^+ baryon to $c\tau \approx 300 \mu m$ for $D^\pm$ mesons. Mesons D^0 and $D_s^\pm$ have an intermediate decay length of $\approx 100 \mu m$.

Electrons coming from Dalitz decay of light mesons, that consists mainly of π^0 and η , are considered in another group, that will be referred to as **Dalitz electrons**. Dalitz electrons have a very narrow DCA distribution, since their mothers decay very close to the primary vertex, with a decay length of the order of $c\tau \approx 20 nm$. Because of this, their DCA template is very slim and limited by the detector resolution.

The last group consists of electrons that comes from conversion of gammas in the detector material, the **conversion electrons**. Although these electrons are created much farther away from the primary vertex with respect to the other electrons discussed above (the beam pipe, the first material that the particles can interact with, is situated 3 cm far away from the primary vertex, while the beauty hadrons have decay length of $c\tau \approx 500 \mu m$ (≈ 0.05 cm)), the DCA values of conversion electrons are comparable to the ones from the other sources. This happens because since the gammas do not interact with the magnetic field, its trajectory before the interaction with the material is straight. In addition, the electrons created by gamma conversion has a very small angle with respect to the gamma, and consequently their impact parameter is essentially due to the existence of the magnetic field B (see [Appendix D](#)).

The symmetry in the kinematics of the gamma conversion is something useful to be explored. By convention, the impact parameter has a positive or negative sign depending on whether the primary vertex is on the left or on right side of the track (with respect to the momentum vector). By consequence, the DCA of electrons is a mirrored version of the DCA of positrons: as can be seen in the two panels of [Figure 6.10](#), positrons and electrons have a different preferred sign of DCA, so the peak of the DCA distribution is centered at negative values for electrons and at positive values for positrons. Therefore, the total DCA template for conversion (electrons + positrons) is symmetric with respect to zero.

A very useful tool is to build the templates as the distribution of $DCA \times charge \times B \text{ field sign}$ instead of just DCA, because the multiplication of the DCA by charge and B field sign makes the conversion template significantly asymmetric: in this case, both electrons and positrons distributions gets shifted to the same side, leading to a final (electrons + positrons) asymmetric distribution (see Figure 6.10). While the conversion electrons template becomes asymmetric if the quantity $DCA \times charge \times B \text{ field sign}$ is considered, the other templates remain unchanged. This asymmetry in the conversion electrons template makes more difficult the occurrence of possible interplays between this template and the one from beauty electrons during the fit procedure, even if both of these templates have a broad distribution. The ITS quality cuts discussed in section 6.5 also help to disfavor this interplay, by making the template of conversion electrons slimmer.

Figure 6.10: DCA distributions of electrons and positrons from gamma conversion (left) compared to the $DCA \times charge \times B \text{ field sign}$ distribution (right) of the same particles. An asymmetry is introduced in the final template (electrons + positrons) if the DCA is multiplied by the charge of the track and B field sign.

The templates are all built using the MC that contains a trigger on electrons from heavy-flavor decays. Because of this trigger, the relative abundance of heavy-flavor in this MC is increased with respect to the MB production. Consequently, in the construction of the templates of conversion and Dalitz electrons it is taken care to not include electrons from gammas, π^0 or η that come from the feeddown of heavy-flavor hadrons, or that come from feeddown of kaons from heavy-flavor hadrons, since the contribution from

those feeddowns would distort the shape of those templates.

Figure 6.11 shows the templates of the four groups of electrons. In particular, for the templates of electrons from beauty and charmed hadrons decays, the contribution from baryons and mesons are shown separately, in order to make it visible how the different decay lengths of baryons and mesons discussed above directly influences the DCA shapes of their decay electrons. But, in the end, baryons and mesons are summed together to build the final beauty and charm electrons templates used in the fitting procedure. In Table 6.4 a summary of the decay length of the different electrons sources is presented.

Figure 6.11: DCA templates of electrons from beauty hadron decays, charmed hadron decays, Dalitz decays and gamma conversion, respectively.

Electron source	Decay length (production radius in the case of conversion)
Λ_b^0	$c\tau \approx 400 \mu m$
B	$c\tau \approx 500 \mu m$
Λ_c^+	$c\tau \approx 60 \mu m$
D^0 and $D_s^\pm$	$c\tau \approx 100 \mu m$
$D^\pm$	$c\tau \approx 300 \mu m$
Dalitz	$c\tau \approx 20 nm$
Conversion	$> 3 cm$

Table 6.4: Different sources of electrons and approximate radial position where electrons from these different sources are created.

6.8 The Maximum Likelihood Template Fit

The fit method used in this analysis is the same used for the run 1 analysis [118] [53]: it is a maximum likelihood method based on the Barlow and Beeston approach [119] slightly improved. This approach allows to fit an experimental distribution (in this case, the DCA distribution of measured electrons) with other distributions that have no analytical formula, taken from MC simulations (DCA templates). The template fit is performed independently in each p_T bin, and the DCA distributions follows Poissonian statistics. Below will be discussed the likelihood function of the problem and how its minimization is performed.

6.8.1 The Likelihood Function

Considering the DCA distribution of electrons measured in data, a number d_i of electrons is found for each DCA bin i . For each bin i there is also an unknown expectation value of electrons, the f_i .

Considering the templates taken from MC, the number of entries in the DCA bin i for the source j is named a_{ij} , and the unknown corresponding expectation value is A_{ij} .

The assumption that a weighted sum of the templates can reproduce the data means that there are amplitudes p_j such that the expectation value of the number of electrons in each DCA bin, f_i , consists in the sum of the expectation value of each source, A_{ij} , scaled by the amplitude p_j :

$$f_i = \sum_{j=1}^m p_j \times A_{ji}. \quad (6.3)$$

For each source j there is one associated amplitude p_j . By the knowledge of the amplitude of the beauty template, p_{beauty} , it is easy to determine the amount of electrons from beauty hadron decays in each p_T bin, so this is the parameter to be estimated in this analysis.

The likelihood of having d_i counts in a bin where the expected number of counts is f_i is given by the Poisson distribution:

$$p(d_i, f_i) = e^{-f_i} \frac{f_i^{d_i}}{d_i!}. \quad (6.4)$$

Analogously, the likelihood of having in the template j and bin i the amount a_{ij} when the expected value is A_{ij} is given by:

$$p(a_{ji}, A_{ji}) = e^{-A_{ji}} \frac{A_{ji}^{a_{ji}}}{a_{ji}!}. \quad (6.5)$$

The total likelihood to be considered in this analysis is the the product between the likelihood of the observed d_i ($p(d_i, f_i)$) and the likelihood of the a_{ij} taken from MC ($p(a_{ji}, A_{ji})$), summed over all the DCA bins i and template sources j :

$$L = \sum_{i=1}^n e^{-f_i} \frac{f_i^{d_i}}{d_i!} \times \sum_{i=1}^n \sum_{j=1}^m e^{-A_{ji}} \frac{A_{ji}^{a_{ji}}}{a_{ji}!}. \quad (6.6)$$

As it is common in maximum likelihood problems, the logarithm of the total likelihood is considered:

$$\ln L = \sum_{i=1}^n d_i \ln(f_i) - f_i + \sum_{i=1}^n \sum_{j=1}^m a_{ij} \ln(A_{ij}) - A_{ij}. \quad (6.7)$$

6.8.2 The Maximum Likelihood Estimation

The total likelihood given by Equation 6.7 has a total of $n \times (m + 1)$ free parameters, that corresponds to the $n \times m$ parameters A_{ij} and the m parameters p_j . In order to determine the parameters that maximizes the likelihood, the derivative of $\ln L$ is taken with respect to the free parameters and made equal to zero, giving the two coupled equations:

$$\frac{d \ln L}{dp_j} = \frac{d_i A_{ji}}{f_i} - A_{ji} = 0; \forall i, j, \quad (6.8)$$

$$\frac{d \ln L}{dA_{ji}} = \frac{d_i p_j}{f_i} - p_j + \frac{a_{ji}}{A_{ji}} - 1 = 0; \forall i, j. \quad (6.9)$$

The manipulation of Equation 6.9 leads to:

$$A_{ji} = \frac{a_{ji}}{1 + p_j(1 - \frac{d_i}{f_i})}. \quad (6.10)$$

Equation 6.10 shows that for the determination of A_{ji} , information from both data and Monte Carlo templates are used. Without the data, the maximum likelihood would imply that the expected values A_{ji} would be equal to the a_{ji} taken from MC. The information from the data should make the A_{ji} converge towards the true distribution, and thereby be slightly smoothed with respect to a_{ji} .

The solution of Equation 6.9 with respect to A_{ij} leads to:

$$A_{ji} = -\frac{1}{2} \frac{d_i p_j - f_{i \setminus j}(p_j + 1) + a_{ji} p_j}{p_j(p_j + 1)} + \sqrt{\frac{1}{4} \left(\frac{d_i p_j - f_{i \setminus j}(p_j + 1) + a_{ji} p_j}{p_j(p_j + 1)} \right)^2 + \frac{a_{ji} f_{i \setminus j}}{p_j(p_j + 1)}}, \quad (6.11)$$

where:

$$f_{i \setminus j} = \sum_{k \neq j} p_k A_{ki}. \quad (6.12)$$

The interaction of Equation 6.11 leads to a distribution of the expected values A_{ji} that does not have any systematic deviation from the templates a_{ji} taken from MC,

as can be seen in Figure 6.12, indicating that the procedure was successful. This is an important improvement of this method (already used in run 1 analysis [118] [53]) with respect to the approach proposed by Barlow and Beeston [119], where the A_{ji} is calculated by the interactive update of Equation 6.10 (f_j depends on A_{ji}). By the pure Barlow and Beeston approach, a systematic difference was observed between A_{ji} and a_{ji} , mainly on the tails (this was deeply studied in [118]), while when calculating A_{ji} by Equation 6.11 a good convergence is observed within few interactions (≈ 5).

Figure 6.12: Templates (a_{ji}) for electrons from charm and beauty hadron decays together with the corresponding expectation values A_{ji} found by the maximum likelihood fit. No systematic deviations are found between a_{ji} and A_{ji} , indicating that the procedure was successful.

The solution of Equation 6.8 with respect to p_j gives:

$$p_j = \frac{d_i - \sum_{k \neq j} p_k A_{ki}}{A_{ji}}. \quad (6.13)$$

The template fit starts by setting the initial parameters p_j for every source j . Then, for a specific source j , the A_{ji} is updated according to Equation 6.11. After this, the p_j is updated according to Equation 6.13, that comes from the maximization of the likelihood with respect to p_j , making use of the last A_{ji} calculated. Due to this interactive procedure, the number of free parameters of the problem is reduced to m , the amount of parameters p_j . The A_{ji} are strictly determined by the p_j , as given by Equation 6.11.

The initial values of A_{ji} are set up as a_{ji} , and the initial values of p_j are set up as $p_j = 0.25$. In this analysis, $m = 4$, since four templates are used to fit the data.

Once the numerical computation of the free amplitude parameters p_j have converged to the solution that maximizes the total likelihood, the sum of the templates normalized by the correct amplitudes (shown as the "fit" histogram in Figure 6.13) is shown to reproduce accurately the DCA distribution of inclusive electrons. An example of the result of the maximum likelihood fit for one p_T bin is shown in Figure 6.13. The normalization pull corresponds to the difference, in each p_T -bin, between data and fit, divided by the sum of the errors of the data and fit histograms. The integral of the beauty template gives the amount of beauty electrons in this p_T bin.

Figure 6.13: DCA distribution of electrons from data (black dots) together with the templates of the four electron sources (charm, beauty, Dalitz and conversion) with the amplitudes given by the maximum likelihood fit. The sum of the templates are a good representation of the data.

6.8.3 Bins Without Counts from the Templates

A special care must be given to the DCA bins with a finite count from data, d_i , but zero count from the templates, $a_{ji} = 0, \forall j$. For this case, [Equation 6.9](#) leads to

$$\frac{d_i}{f_i} - 1 = \frac{1}{p_j}; \forall j, \quad (6.14)$$

which does not contain any information about A_{ji} . The maximum of the likelihood in such bins is not associated necessarily with a zero derivative of the likelihood in the A_{ji} phase space, leading to inconsistencies of the currently used method [\[118\]](#). Instead, the maximum of the likelihood in these cases corresponds to the situation in which the counts coming from data is associated to a single source A_{ji} . The determination of the source A_{ji} that maximizes the likelihood can be extremely biased by the amplitude p_j of the templates [\[118\]](#), which has no physical meaning by itself and just reflects the initial abundances of the electrons from different sources in the Monte Carlo (the number p_j can be, for example, very different if an enhanced or Minimum Bias Monte Carlo is used to build a template).

The maximum likelihood fit is performed independently in each DCA bin. This means that if the order of the bins of the DCA distributions was shuffled (consistently for data and all templates), the result of the fit on each DCA bin would remain the same, there is no requirement for local smoothness. Nevertheless, the available prior information about the DCA distributions of the electrons from different sources, dictated by the underlying physical process that originate the electrons, can be used in favor of the fit procedure in this case of DCA bins with no counts from templates.

Bins without counts from the templates will most probably appear on the tails of the DCA distribution. It is known that DCA distributions have a peak centered around zero, and decrease towards larger and smaller DCA values (this is not true just for the conversion template, that is not centered at zero). In particular, the electrons from beauty hadron decays should be the dominating contribution in the tails of the DCA distribution, since their distribution is the wider one. It is not likely that a count in the tail of the DCA distribution comes from the very slim Dalitz template, for example. This means that the counts from data, d_i , in the bins with no count from templates (that

are mainly on the tails) should most probably come from electrons from beauty-hadron decays.

Therefore, for this analysis, all electrons in bins with no counts from the templates are assumed to be electrons from beauty hadron decay, since these have the widest distribution. This assumption can bring a bias to the final result, since, even though most probably, not always this count d_i really comes from electrons from beauty hadron decay. But since this assumption is more realistic than the attribution of d_i to any other template, the bias will be smaller than in any other case. This bias is estimated together with the estimation of the statistical uncertainties, as explained below.

6.9 Uncertainty Estimation

When the measurement of a given quantity is performed within an experiment, an estimation of the statistical uncertainty of this quantity can be performed by repeating the experiment many times. By doing this, a distribution of the measured quantity is obtained and the width of this distribution indicates how much is the statistical fluctuation associated to this measurement while the mean of the distribution indicates its most probable value.

In order to estimate the statistical uncertainty of the yield of electrons from beauty hadron decays measured in this analysis via the template fits discussed above, a "toy model" is built in order to virtually repeat this measurement many times, using a new and independent data set for each fit (toy model data and templates).

The toy model builds the new data set, in each p_T bin, as described below:

- Each original DCA template is normalized to have its maximum at 1;
- A smoothing is applied to these templates, in order to improve the representativeness of bins with few entries;
- **New templates** are sampled from the original ones, by sorting random numbers following the probability distribution given by the scaled and smoothed original templates;

- The scaled templates for the different sources are summed together according to the proportions found in the original measurement;
- A **toy model data** is built by sampling the distribution resultant from the previous step;

The created toy model data and templates have the same number of entries as the corresponding original distributions.

In this toy model approach, the toy model data is built under control, as a sum of a known contribution of electrons from the different sources, as taken from the original measurement. For each p_T bin, it is performed the fit of the toy model data with the toy model templates. The amount of electrons from beauty hadron decays is determined for this virtual experiment.

A number of 1000 template fits was performed with the toy model approach in each p_T bin. Figure 6.14 shows the fraction of electrons from beauty hadron decays (relative to the total amount of electrons) obtained from each of these virtual experiments, for two different p_T bins.

Figure 6.14: Distribution of the fraction of electrons from beauty hadron decays ($b \rightarrow e / (\text{all } e)$) obtained from 1000 interactions of the toy model, for two different p_T bins. The mean of the toy model distribution and the toy model true $b \rightarrow e / (\text{all } e)$ are also indicated.

The RMS of these distributions are taken and divided by the fraction of electrons from beauty hadron decays from the original experiment, in each p_T bin. The resulting

number is the statistical uncertainty, in percentage, of the amount of electrons from beauty hadron decays measured in the original experiment. The values of this statistical uncertainty as a function of p_T are shown in Figure 6.15.

The usage of the toy model approach for the statistical uncertainty estimation in this analysis is important since it allows to take into account the contribution of statistical uncertainty from both data and Monte Carlo templates.

Figure 6.15: Statistical uncertainty (in percentage) obtained in each p_T for the electrons from beauty hadron decays.

In Figure 6.14 it is also indicated the mean of the toy model distribution and the fraction of electrons from beauty hadron decays from the original measurement (the true amount inside the toy model data). The difference between these two quantities indicates the bias associated to the template fits as performed in this work. The bias was found to be very small, smaller than 5% for most of the p_T bins, so no correction of the yield due to this bias was performed.

6.10 Corrections on the Templates and Systematic Uncertainties

A Monte Carlo simulation does not reproduce the particle p_T -spectra or even the relative abundances between different particle species accurately, and this brings the necessity of performing a series of corrections in the Monte Carlo. These corrections can affect directly the different DCA templates used in the analysis and they are necessary in order to make the DCA templates as close as possible to the DCA distributions that would be found in data for each electron category, avoiding the introduction of biases in the template fit procedure. The estimation of the systematics uncertainties in the present analysis is performed by varying these corrections performed in the Monte Carlo within reasonable limits. Below, together with each correction, will be discussed the associated systematic uncertainties.

6.10.1 Mean and Resolution Correction

The mean and resolution of the DCA distributions in data are different from the ones in MC.

The mean of DCA distributions in data is shifted towards negative values for the ALICE data samples collected during LHC run 2. This is understood to be a consequence of a misalignment of some SPD modules during this data taken period. This shift is dependent on the p_T and azimuthal angle (φ) of the track, and also on the polarity of the B field. The shift of the mean is not existent in the MC DCA distributions. A code (named *improver task*) was created with the purpose of mimicking in the MC the DCA mean shift found in data. The code performs the mean correction by summing to the DCA of a single track in the MC the quantity $\langle \text{DCA}(\text{data}) \rangle - \langle \text{DCA}(\text{MC}) \rangle$, where $\langle \text{DCA}(\text{data}) \rangle$ is the mean of the DCA found in data and $\langle \text{DCA}(\text{MC}) \rangle$ is the mean of the DCA found in MC for a particle specie.

The resolution of the DCA distributions is found to be worse in data than in MC. This is not a particular issue from run 2 data sample like the mean shift discussed above, but it is a consequence of the fact that there are more fluctuations in the real

experiment than in the MC simulations. The improver task also takes care of correcting the resolution in MC in order to make it to match the resolution found in data.

In order to check the corrections performed by the improver task, a consistent particle sample needs to be selected both in data and in MC. For this purpose, charged hadrons were selected both in data and in MC by the TPC PID cut $-3 < n\sigma_{\text{TPC}} < -5$. A minimum bias MC simulation was chosen for this study. For each p_T bin, the DCA distributions are fitted by a Gaussian in the region $-rms < \text{DCA} < rms$, that comprehends a very narrow DCA range and has minor contribution from secondary hadrons. An example of the Gaussian fits, performed in one bin of p_T , can be seen in [Figure 6.16](#). The fits are performed in data, in MC without improver task corrections and in MC with the improver task corrections. The mean and resolution of the DCA distributions is taken as the mean and σ of the fitted Gaussians.

Figure 6.16: DCA distribution of charged hadrons in data (black) and in MC before (red) and after (blue) the corrections of the mean and resolution. Also the Gaussian fits performed for the extraction of the mean and resolution are shown.

The comparison between the DCA mean and resolution found in data and in MC with and without the corrections, as a function of p_T , can be seen in [Figure 6.17](#). In particular, the top panels show the mean correction, and the bottom panels show the

resolution correction. Both DCA mean and resolution in the MC are in much better agreement with data after the corrections applied by the improver task.

Figure 6.17: Mean (top) and resolution (bottom) of DCA in data (black) compared to the ones from MC without (red) and with (blue) the corrections performed by the improver task.

6.10.2 D meson p_T -spectrum Correction

The D meson spectrum generated in the enhanced MC simulation, from which the DCA templates are taken, has a very different slope with respect to the spectrum measured in data. A correction in the slope of the p_T spectrum of D mesons in MC is necessary because it can affect directly the width of the DCA distribution of electrons from charm hadron decays.

In [Figure 6.18](#) it is possible to see the comparison between the D meson p_T spectrum from the enhanced MC (green) and the D^0 p_T spectrum measured in Pb-Pb collisions

at 5.02 TeV (blue). The spectrum generated in the MC is much harder than the one measured experimentally. It is also known that the different D meson species measured experimentally have essentially the same slope, and here the D^0 is used as representative of the slope of all D meson species in data.

Each one of these spectra is fitted by a TSallis function:

$$f(x) = \text{par}(0) \times (1 - (1 - \text{par}(1)) \times x/\text{par}(2))^{1/(1-\text{par}(1))}. \quad (6.15)$$

The ratio of both spectra, $data/MC$, is shown in the right panel of [Figure 6.18](#) together with the ratio of the functions that fit each spectrum. The ratio of the functions interpolates well the ratio of the two spectra, and this function gives the correction that is necessary to be applied to the MC in order restore the shape of the D meson spectrum found in data.

The strategy adopted to correct the slope of the D meson spectrum from MC in this analysis is to perform a statistical rejection of D mesons: the probability to accept a D meson is given by the correction function (right panel of [Figure 6.18](#)) normalized to be unity at its higher value. Since the probability is p_T dependent, this stochastic correction will change the slope of the spectrum, as required.

For the correction of the D meson p_T -shape, the usage of a weighting procedure like it is usually performed in many analysis, as described in [section 5.8.1](#), would invalidate the assumption of a Poissonian statistics of the templates. But this is an assumption of the maximum likelihood fit performed in this analysis, as discussed above in [subsection 6.8.1](#). For this reason, the stochastic correction is chosen instead of a weighting procedure.

In order to minimize the loss of data due to the statistical rejection of D mesons, three features are explored:

1. The template fit is performed bin by bin;
2. This correction depends only on the slope of the correction function and not on its absolute value;
3. In most cases, the D mother p_T is larger or equal to the decay electron p_T , like it is shown in the left panel of [Figure 6.19](#);

Figure 6.18: Left: p_T spectrum of the D mother of electrons used in this analysis (green) and p_T spectrum of D^0 mesons measured in data (blue) fitted by the functions shown in the corresponding colors. The red D^0 spectrum and fit are used for systematic uncertainty studies. Right: Ratio between the D^0 measurement and the spectrum of D mesons from MC together with the ratio of the two fits. The red ratio is used for systematic uncertainty studies and it is discussed in the text.

Considering these three features, the correction function can be normalized to be unity at the beginning of each electron p_T bin, like it is shown in the right panel of [Figure 6.19](#). In this way, the correction is performed in each electron p_T bin minimizing the loss of data, since the probability of accepting a D meson always starts at 100% in each electron p_T bin. Since the D mother p_T is usually larger or equal to the electron p_T , just few cases will be not corrected in a p_T dependent way (since the probability of rejecting a D meson is not defined in an electron p_T bin for the D mother that has p_T smaller than the lower edge of p_T bin). These few cases where D mother p_T is smaller than the lower edge of the electron p_T bin are all considered, with probability 100%.

Once done the statistical rejection of D mesons according to the probabilities described above, a cross check of the procedure is performed: for different electron p_T bins, the D mother p_T spectra is compared to the slope of the D^0 spectrum measured in data, as shown in [Figure 6.20](#). This check ensures that the statistical subtraction of D mesons in MC is indeed correcting their slope to match the slope of the data.

The effect of the correction of the D mesons p_T -spectrum in MC on the DCA of

Figure 6.19: Left: Electron p_T shown as function of the D mother p_T . A black line indicates where the electron p_T is the same as the D mother p_T . Right: Probability of accepting D mesons from MC, normalized to be unity at the beginning of each p_T bin of the electron, used to correct the slope of D mesons in MC in each electron p_T bin.

Figure 6.20: Spectra of the D mother of electrons in MC, for many electron p_T bins, after the p_T correction, compared to the slope of the spectra of D^0 measured in data.

electrons from D mesons can be observed in [Figure 6.21](#). After the correction, the DCA distribution of electrons from D meson decay gets visibly narrower.

In order to estimate the systematic effect that this correction can bring to the yield of electrons from beauty hadron decays, the correction is varied within reasonable limits and

Figure 6.21: DCA of electrons from D meson decay before and after correcting the D meson p_T spectrum slope.

the modification of the yield is quantified. The statistical and systematic uncertainties of the D^0 measurement are summed in quadrature and the slope of the D^0 spectrum is changed as much as the combined uncertainties allows. In the left panel of [Figure 6.18](#) it is possible to see in red the modification performed in the D^0 spectrum for the systematics studies: its uncertainty is summed to the central values at low p_T and subtracted from the central values at high p_T , causing a tilt on the slope. In red is also shown the function that fits this modified D^0 spectrum. In the right panel, the red function is the corresponding correction function used for systematic studies.

The ratio of the p_T spectrum of electrons from beauty hadron decays obtained using the modified correction function and the default correction function is shown in [Figure 6.22](#). To the D meson p_T -shape correction it is assigned a systematic effect of 5%, constant in p_T , obtained from this ratio.

6.10.3 Beauty p_T -spectrum Correction

As shown previously, the slope of the p_T spectrum of D mesons influences directly the shape of the DCA distribution of electrons from D meson decays. For an analogous

Figure 6.22: Ratio of spectrum of electrons from beauty hadron decays obtained when correcting the D meson spectrum with the modified correction function and with the default correction function. The lines in green represent the systematic uncertainty assigned to this correction. The dots at unity have the statistical error bars of the spectrum of electrons from beauty hadron decays.

reason, it is needed to guarantee that the spectrum of B mesons taken from MC has the correct slope. There is no available measurement of the B meson p_T spectrum produced in Pb-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV in the p_T range of the current analysis: the motivation of measuring electrons from B hadron decays is exactly to probe, even if indirectly, the B hadron production. So, differently from the case of the D meson p_T -spectrum correction, the B meson correction cannot be performed by comparing the MC and measured p_T -spectrum. The B meson p_T -spectrum correction is performed by considering two different factors that are not taken into account (properly) in the MC. These effects are:

- In the currently used MC, it is observed that the generated B meson spectrum does not follow exactly the pattern predicted by FONLL calculations, like shown in the left panel of [Figure 6.23](#). The FONLL calculations are known to be in a good agreement with heavy-flavor mesons measurements in pp collisions, and by

consequence, the PYTHIA event generator (that produce particles via pp collisions in the MC) should be in good agreement with FONLL. This does not happen in the current MC since the B meson spectrum was produced by the sum of many spectra generated in different p_T ranges (technique used to extend the p_T range of the produced spectrum). For this reason, **the produced B meson p_T spectrum needs to undergo a correction to match the one from FONLL calculation.**

- The MC does not reproduce effects of the QGP on the spectrum: the B meson spectrum taken from MC does not have the p_T dependent suppression induced by the existence of the medium produced in a real collision. **This p_T dependent suppression induced by the medium needs to be added in the B meson spectrum from MC, in order to have a template of DCA of beauty electrons with a realistic shape.** A quantification of this suppression can be given by the R_{AA} of B mesons taken from model predictions, as shown in the right panel of [Figure 6.23](#);

Due to these two factors above mentioned, in the current analysis the function used to correct the B meson spectrum is given by

$$\frac{R_{AA} \times FONLL}{MC}, \quad (6.16)$$

where:

- " R_{AA} " accounts for the parametrization of B meson R_{AA} given by TAMU model for Pb-Pb at 2.76 TeV collisions [87] (shown in red in the right panel of [Figure 6.23](#)). It is important to remember that the R_{AA} from the model consists in the ratio of what would be the B spectrum produced in Pb-Pb over the FONLL B spectrum;
- "FONLL" is the central value of B meson spectrum given by FONLL calculation (shown by the continuous blue line in the left panel of [Figure 6.23](#));
- "MC" is the B spectrum given by the MC simulation (in red in the left panel of [Figure 6.23](#));

The resulting correction function is shown in black in the right panel of [Figure 6.23](#). The correction of the B meson spectrum is performed by subtracting B mesons statistically according to this function, that gives the probability to accept the B mesons from the MC. This correction function is normalized to have its maximum value at one.

Figure 6.23: Left: B meson spectrum from the MC simulation (red) and from FONLL calculation with uncertainties (blue). Right: correction function for the B meson spectrum (black) shown together with TAMU parametrization of the R_{AA} for B mesons in Pb-Pb at 2.76 TeV collisions.

Differently from the case of the charm correction, the B meson p_T -spectrum correction is not performed bin by bin, since the correction function for B mesons is smoother and does not imply in a so large loss of data. Also, the amount of cases in which the B mother p_T is smaller than the electron p_T is not negligible as it is for D mesons, as can be seen in [Figure 6.24](#). This makes a correction bin by bin for the B mesons not viable.

The effect of the correction of the B mesons p_T -spectrum on the DCA of electrons from B mesons decay can be observed in [Figure 6.25](#). As for the D mesons, the electron DCA distribution becomes slimmer after the correction of the mother p_T slope.

The study of the systematic effect of this correction is performed by varying the correction function, [Equation 6.16](#), within reasonable limits. In the left side of [Figure 6.23](#), two variations of the parametrization of the B mesons R_{AA} from TAMU model are shown. These variations are built by adding (or subtracting) to the central value of the R_{AA} parametrization half of the difference between it and unity. In this way, the slope of the R_{AA} is significantly varied. Two different correction functions are calculated

Figure 6.24: Electron p_T shown as function of the B mother p_T .

Figure 6.25: DCA of electrons from B mesons decay before and after correcting the B meson p_T spectrum slope.

according to Equation 6.16 by making use of these two variations of the R_{AA} . In the left panel of Figure 6.26, the central B meson correction function (default, in black) is shown together with the two variations of it, obtained as described above. The ratios between the beauty electrons spectrum obtained using the varied correction functions and the default correction function is shown in the right panel of Figure 6.26. A systematic

effect of about 10% is observed, and it is constant in p_T ;

Figure 6.26: Left: correction functions for the B meson spectrum used for the systematics studies (red) shown together with the one used for the central value (black). Right: Ratios between the beauty electron spectrum obtained with the modified correction functions and the central value.

6.10.4 Hadron Contamination

The template fit performed in this analysis makes use of four DCA templates (from beauty, charm, Dalitz and conversion electrons). Although the electron candidates sample have a hadron contamination, no template of hadrons is used in the fits to take into account the hadron contamination directly. Nevertheless, as can be seen in [Figure 6.27](#), the template of Dalitz electrons is very similar in shape to the DCA distribution of hadrons. In this figure, the hadrons considered are the ones selected by the TPC PID cut $-3 < n\sigma_{\text{TPC}} < -5$, so they consist mainly of charged pions. Charged pions are also the main source of hadron contamination in data for $p_T > 1 \text{ GeV}/c$.

Due to the similarity between the DCA distribution of electrons from Dalitz decay and hadrons, the hadron contamination is expected to be accommodated by the Dalitz template: the resulting amplitude of the Dalitz template will reflect the amount of electrons from Dalitz decay and the hadron contamination within the data sample. The other extreme case can be considered: if the hadron DCA distribution is used instead of the Dalitz template for the fit, then the final amplitude of the hadron template would em-

Figure 6.27: Comparison between Dalitz electron template and hadron DCA distribution.

brace both hadrons and Dalitz electrons. The most accurate situation would consist in an intermediate case, where hadrons and Dalitz electrons within the data sample would be each fitted by its own proper template. However, in practice, even if both of these templates were considered for the fits, the very similar shape they have would make almost impossible to disentangle these two different contributions, making the use of both templates pointless.

The approach of this analysis is to use for the central (default) fit just the Dalitz template to fit both the signal from Dalitz electrons and hadron contamination. For a systematic study, the other extreme case discussed above is considered. The systematic difference between the final result obtained by these two approaches covers any systematic effect caused by the fact that no hadron template is used in the default fit.

In [Figure 6.28](#) is shown the ratio of the beauty electron spectrum obtained in the case where the hadron template is used instead of Dalitz template and the default case, where Dalitz template is used instead of hadrons. A systematic effect of about 10% is observed. The systematic error due to the hadron contamination is applied in the final beauty electron spectrum bin by bin according to the values in the histogram shown in [Figure 6.28](#).

Figure 6.28: Estimation of systematic effect in the beauty electron spectrum due to the hadron contamination in the electron candidate sample.

6.10.5 Correction of the Proportion Between Charmed Baryons and Mesons

The template of electrons from charm hadron decays is built as the sum of the template of electrons from all different charmed baryons and mesons species (mainly Λ_c^+ , D^0 , $D^\pm$, $D_s^\pm$). As discussed above (see table [Table 6.4](#)), these different sources of electrons have different decay lengths, and, in particular, the Λ_c^+ has the shorter decay length, leading to the most narrow electron DCA distribution among the contributions to the charm template. It is particularly important to correct in the MC the proportion between the Λ_c^+ baryon and the mesons, since the MC does not reproduce the correct proportion between baryons and mesons found in data and these sources have very distinct decay lengths.

In order to perform this correction, the measurement of the ratio Λ_c^+/D^0 is taken into account. In [Figure 6.29](#) is shown the measurement of this fraction in Pb-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV. Based on the central value of this measurement, the MC is corrected in order to have a fraction $\Lambda_c^+/D^0 = 1$, for all p_T bins of the analysis. The correction of the MC is performed by scaling the template of electrons from Λ_c^+ by the weight:

$$w = (\Lambda_c^+/\text{D}^0)_{\text{data}} \times \frac{BR(\Lambda_c^+ \rightarrow e)}{BR(\text{D}^0 \rightarrow e)} \times \frac{(\text{nEntries } \text{D}^0 \rightarrow e)}{(\text{nEntries } \Lambda_c^+ \rightarrow e)}, \quad (6.17)$$

where

- $(\Lambda_c^+/\text{D}^0)_{\text{data}}$ is the correct fraction (measured in data);
- $BR(\Lambda_c^+ \rightarrow e)$ is the branching ratio of the decay of Λ_c^+ in a semi-leptonic channel $\rightarrow \text{BR} = 4.5 \pm 1.7$ for $\Lambda_c^+ \rightarrow e + X$;
- $BR(\text{D}^0 \rightarrow e)$ is the branching ratio of the decay of D^0 in a semi-leptonic channel $\rightarrow \text{BR} = 6.49 \pm 0.11$ for $\text{D}^0 \rightarrow e + X$;
- $\text{nEntries } \text{D}^0 (\Lambda_c^+)$ is the number of entries in the template of electrons from $\text{D}^0 (\Lambda_c^+)$;

In order to estimate the systematic uncertainty related to this correction, two extreme values of the fraction are considered: $\Lambda_c^+/\text{D}^0 = 0.5$ and $\Lambda_c^+/\text{D}^0 = 1.5$. As can be seen from [Figure 6.29](#), this variation covers (and even exceeds) the statistical and systematics uncertainties of the measurement.

The ratios of the final yield of electron from beauty hadron decays obtained using the default correction $\Lambda_c^+/\text{D}^0 = 1$ and the variations of the correction ($\Lambda_c^+/\text{D}^0 = 0.5$ and $\Lambda_c^+/\text{D}^0 = 1.5$) are shown in [Figure 6.30](#). A 3% systematic effect due to this correction is assigned to the final beauty electrons yield, constant with p_T .

During the discussion of the D meson p_T -shape correction, it was mentioned that the correction is performed by a statistical rejection of D mesons instead of a weighting procedure. This is done in order to keep the charm template with an integer number of entries in each DCA bin, keeping the poissonian statistics assumed in the fit procedure. Nevertheless, in the correction of the Λ_c^+/D^0 ratio presented here, a weight is applied on the Λ_c^+ template in order to bring its number of entries to the right proportion with respect to the number of entries of the D^0 template, in each p_T bin. With this weight, the entries in each DCA bin of the Λ_c^+ template are no longer integers. Although it violates the poissonian statistics assumed in the analysis, it was observed that the yield of electrons from beauty hadron decays with and without the present correction are very

Figure 6.29: Measurement of the fraction (Λ_c^+/D^0) in Pb-Pb at 5.02 TeV collisions (green) and also in other collision systems.

Figure 6.30: Systematic effect in the beauty electron spectrum due to the (Λ_c^+/D^0) correction.

similar to each other, showing that this correction has a small effect to the final yield and that the analysis result is stable under this weighting procedure. In despite of the drawback of not maintaining the poissonian statistics of this particular template, it was decided, under the mentioned observations, to keep this correction and the associated

systematic uncertainties in this analysis.

6.10.6 Other Systematic Uncertainties and Summary

- TPC PID: the efficiency of the TPC PID cut applied to select electrons is different in data and in MC. As previously discussed, the center of the $n\sigma_{TPC}$ of the electron distribution in data is located at -0.1, while in MC it is located at zero. The efficiency of the PID cut $-0.1 < n\sigma_{TPC} < 3$ is calculated in data (as discussed in [section 6.6](#)) and via MC simulations. The ratio between the spectrum of electrons from beauty hadron decays obtained with the use of the default efficiency correction (obtained from data as 50% under the assumption that the electron $n\sigma_{TPC}$ is Gaussian-like) and the varied efficiency correction (obtained from MC) is shown in [Figure 6.31](#), and they differ of about 7%.

Figure 6.31: Systematic effect due to the difference of the TPC PID in data and MC samples.

Some sources of systematic errors could not be treated in this work due to time constraints. Nevertheless, they are added in the final result and their values are based on other analysis. These sources are:

- TPC-TOF matching: charged particles traversing the TPC ionizes the gas, creating tracks. These tracks need to be prolonged and associated to a signal on TOF, so the time-of-flight of the corresponding particle can be calculated. Due to the limited acceptance of the detectors, some of these charged particles, that generates a track in the TPC, reaches the TOF in an inactive area, and the corresponding time-of-flight cannot be known. This mismatch between TPC and TOF signals, if not reproduced properly in the MC simulations, can lead to systematic effects in the experimental measurement. Studies performed in Pb-Pb at $\sqrt{s_{\text{NN}}} = 5.02$ TeV (for the calculation of the R_{AA} of electrons from open heavy-flavor decays) have shown that the difference between TPC-TOF matching efficiencies in data and MC have a difference of about 2%. This value is included in the present analysis as systematic uncertainty in the final yield of electrons from beauty hadron decays;
- TOF PID: the efficiency of the TOF PID cut for the identification of electrons can be different in data and in MC. This difference was studied using data from Pb-Pb at $\sqrt{s_{\text{NN}}} = 2.76$ TeV [118]. In data, a sample of electrons from neutral meson decays can be selected based on their decay topology [118] and this sample is used for the calculation of this efficiency in data. A difference of about 10% was found between the TOF PID efficiency in data and in MC. This value is used in this analysis as systematic uncertainty due to the TOF PID cut;

A summary of all systematic uncertainties considered in the yield of electrons from beauty hadron decays is presented in the Table 6.5. The B meson correction, hadron contamination and TOF PID cut are the main sources of systematic uncertainties.

The final systematic uncertainty is calculated summing the different contributions in quadrature, and it ranges from 19 to 21% in the different p_{T} -bins of the beauty electron spectrum, as shown in Table 6.6.

Source	Value
B meson correction	10%
D meson correction	5%
Hadron contamination	$\approx 10\%$ for low p_T and $\approx 5\%$ at high p_T
Λ_c/D^0 ratio correction	3%
TPC-TOF matching	2% (from HFE in Pb-Pb at 5.02 TeV analysis)
TPC PID	7%
TOF PID	10% (from run1 $b \rightarrow e$ analysis)

Table 6.5: Summary of the systematic uncertainties on the yield of electrons from beauty hadron decays from the uncorrelated sources discussed above.

p_T (GeV/ c)	Systematic uncertainty (%)
2.0 - 2.5	21
2.5 - 3.0	19
3.0 - 4.0	19
4.0 - 5.0	20
5.0 - 6.0	19
6.0 - 8	19

Table 6.6: Final systematic uncertainty in each p_T -bin of the beauty electrons spectrum.

6.11 Results and Discussion

In this section, the results from the analysis presented in this chapter will be shown and discussed. The invariant yield of electrons from beauty hadron decays measured in Pb-Pb at $\sqrt{s_{NN}} = 5.02$ TeV is shown and compared to the pp reference. The nuclear modification factor R_{AA} is obtained and compared to pQCD-based theoretical predictions and also to the R_{AA} of electrons from open heavy-flavor hadron decays.

Before showing these final results, the analysis method used to obtain the pp reference used in this work is briefly discussed below.

6.11.1 pp Reference

The measurement of electrons from beauty hadron decays was performed for the first time in ALICE using data from pp at $\sqrt{s} = 7$ TeV collisions [12]. The procedure used in this analysis is different from the one presented in the current work. The first step of the analysis in pp at $\sqrt{s} = 7$ TeV is to select a sample of electrons with large DCA values, from the tails of the DCA distribution, that is dominated by electrons from beauty hadron decays. The background on the selected sample is then estimated via cocktail simulation and subtracted.

The applied DCA cut is p_T -dependent, and given by $|d_0| > 64 + 780 \pm \exp(-0.56p_T)$ (with d_0 in μm and p_T in GeV/c). Such a cut is performed in order to account for the p_T -dependence of the resolution of the DCA distributions as well as to maximize the significance for the beauty decay electron spectrum. Applying this cut, the hadron contamination in the electron sample was found to be reduced to less than 3% in all p_T bins. The background components of the selected sample are mainly electrons from Dalitz decays, from gamma conversion and from charm hadron decays.

The p_T -spectrum of the background electrons are taken from Monte Carlo (cocktail) simulations. As was already discussed, the Monte Carlo does not reproduce correctly the p_T spectra of many particles, therefore they need to be corrected. The production cross section of π^0 and η mesons were measured in ALICE in pp collisions at $\sqrt{s} = 7$ TeV [71], and these mesons are the main sources of electrons from Dalitz decays and of gammas that convert in the detector material into e^-e^+ pairs. The measured π^0 and η spectra are compared to the corresponding spectra in the Monte Carlo simulation and the weights used to correct electrons from Dalitz decays and gamma conversion could be estimated. The p_T spectrum of D mesons in MC was corrected based on the shape of the production cross section of D mesons measured in ALICE in pp collisions at $\sqrt{s} = 7$ TeV [120] [121]. The spectrum of B hadrons was not corrected, since the Monte Carlo was shown to reproduce correctly the p_T -shape measured in data.

In the right panel of Figure 6.32 it is shown the p_T -spectra of the inclusive electrons and background sources, after the DCA cut and background p_T -shape correction. In the same figure it is shown the electrons from beauty hadron decays, obtained after

subtracting the background from the inclusive spectrum.

Figure 6.32: (Left) p_T -spectrum of inclusive electrons, background electrons (from charm, Dalitz decay and gamma conversion) and electrons from beauty hadron decays, obtained by the analysis based on the d_0 cut. (Right) Cross section of electrons from beauty hadron decays obtained via the direct measurement based on d_0 cut (black) and via an alternative method (blue).

The corresponding cross section of electron from beauty-hadron decays is shown in the right panel of Figure 6.32. This cross section, obtained from the analysis based on the d_0 cut, is compared with the outcome from a different analysis method shown in blue, in which no d_0 cut is applied. In this analysis, the electrons from beauty hadron decays are obtained by considering the measurement of electrons from open heavy-flavor hadron decays performed by ALICE in pp at $\sqrt{s} = 7$ TeV [13] and by subtracting the charm hadron decay contribution. The spectrum of electrons from charm hadron decay is estimated by considering the measured D meson spectrum and by performing the semi-leptonic decay of those D mesons via PYTHIA simulations. The two results show excellent compatibility within uncertainties, and the method based on the d_0 cut leads to significantly reduced systematics uncertainties in the final cross section.

The calculation of the R_{AA} of electrons from beauty hadron decays performed in this work uses as pp reference the measurement in pp at $\sqrt{s} = 7$ TeV that makes use of the

d_0 cut, discussed above. An energy scaling based on FONLL is performed in order to scale the result from $\sqrt{s} = 7$ TeV to the energy of $\sqrt{s} = 5.02$ TeV.

6.11.2 R_{AA} of Electrons from Beauty Hadron Decays in Pb-Pb at $\sqrt{s_{NN}} = 5.02$ TeV

After the template fits, the amount of electrons from beauty hadron decays in each p_T bin is extracted by integrating the corresponding template over the full DCA range.

The invariant yield of electrons from beauty hadron decays is obtained according to [Equation 6.1](#), with the use of the ingredients obtained along this chapter. The measured yield of electrons from beauty hadron decays, $N_{\text{raw}}^{e^\pm}(p_T)$, is corrected by the track reconstruction efficiency for $b \rightarrow e$ ($\varepsilon_{\text{reco}}$, shown in [Figure 6.5](#)), and by the efficiency of identification of $b \rightarrow e$ by the TOF and TPC PID cuts, ε^{ID} ([Figure 6.7](#) for TOF and 50% for the TPC). In addition, the raw yield of $b \rightarrow e$ is normalized to unit rapidity by dividing it by the rapidity range $\Delta y = 1.6$. For electrons, which have usually negligible mass compared to their momentum, the pseudo-rapidity and rapidity are the same. Therefore the rapidity range is determined by the pseudo-rapidity range of the measurement $\Delta \eta = 2 \times 0.8$. The yield is also normalized to unity of the azimuthal angle ϕ , so it is divided by the $\Delta \phi$ coverage of the used detectors, that is 2π . Each bin of p_T is normalized by its width (Δp_T) and by its central value (p_T^{center}). The yield is also normalized to the number of analyzed events, N_{ev} (discussed in [section 6.2](#)), and divided by 2, to account only for electrons ($\frac{e^+ + e^-}{2}$).

The invariant yield of electrons from beauty hadrons decays measured in this work, using the 10% most central Pb-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, is shown in [Figure 6.33](#) together with the pp reference scaled by the Glauber model factor of the corresponding centrality, $\langle T_{AA} \rangle = 23.44 \text{ mb}^{-1}$. The modification of the p_T -spectrum of electrons from beauty hadron decays measured in Pb-Pb relative to pp at the same center of mass energy is quantified by the R_{AA} , that is obtained by the ratio of these two spectra. In this ratio, the uncertainties from the reference, statistical and systematics, are propagated to the R_{AA} . In this [Figure 6.33](#) it is possible to see that the systematic errors relative to the

signal extraction in the reference (represented by the boxes), are large, comparable to the ones from the measurement of this work, that amounts in about 20% at low p_T . But the reference also has additional systematic errors that do not come from the signal extraction, but from the energy scaling from $\sqrt{s} = 7$ TeV to $\sqrt{s} = 5.02$ TeV (3.5%) and from the $\langle T_{AA} \rangle$ factor (3%), as indicated on the bottom left of the figure. Currently, a measurement of electrons from beauty hadron decays in pp at $\sqrt{s} = 5.02$ TeV is being performed using the same strategy of this work. This new reference will have smaller systematic errors than the reference used in this work, since the energy scaling will not be necessary.

The R_{AA} of electrons from beauty hadron decays in $\sqrt{s_{NN}} = 5.02$ TeV is shown in the left panel of [Figure 6.34](#). This result indicates that electrons from beauty hadron decays produced in the 10% most central Pb-Pb collisions are suppressed by a factor 40-60% with respect to the measurement in pp, for $p_T \gtrsim 3$ GeV/ c . By confronting this result with the R_{pPb} of electrons from beauty hadron decays, shown in the right panel of [Figure 6.34](#), that is compatible with unity in the full p_T range, it is possible to argue that this suppression in the R_{AA} is not due to initial state effects, but a consequence of the interaction of the beauty quarks with the produced medium. It is known that the effect of the energy loss should be visible in the region of larger p_T mainly. At very low p_T , other effects that can modify the produced spectrum compete with the energy loss, so a suppression of the R_{AA} is not necessarily expected. In the obtained result, is visible that the R_{AA} of electrons from beauty hadron decays is not suppressed, within uncertainties, for $p_T < 3$ GeV/ c .

The comparison between the R_{AA} of electrons from beauty hadron decays measured in this work and the one measured in Pb-Pb at $\sqrt{s_{NN}} = 2.76$ TeV collisions [[53](#)] is shown in the left panel of [Figure 6.35](#). The results are compatible with each other within uncertainties. The small discrepancy between them, that points to a slightly larger suppression of the R_{AA} measured at $\sqrt{s_{NN}} = 5.02$ TeV, could be due to the fact that this measurement is performed using the 10% most central Pb-Pb collisions and the result from $\sqrt{s_{NN}} = 2.76$ TeV is obtained for the 20% most central Pb-Pb collisions. This same level of compatibility between run 1 ($\sqrt{s_{NN}} = 2.76$ TeV) and run 2 ($\sqrt{s_{NN}} = 5.02$ TeV)

Figure 6.33: Invariant yield of electrons from beauty hadron decays measured in this work (blue) compared with the pp reference scaled by the $\langle T_{AA} \rangle$ (red).

R_{AA} measurements is observed also for non-strange D mesons [122], and it is supported by theoretical models. The R_{AA} of non-strange D mesons measured in central collisions at $\sqrt{s_{NN}} = 2.76$ TeV by ALICE is also compatible with the R_{AA} of D^0 mesons measured in central Au-Au collisions at $\sqrt{s_{NN}} = 200$ GeV at RHIC [123], as discussed in [124]. The combined effect of a denser medium and harder initial p_T spectra could result in similar values of the R_{AA} measured in different center of mass energies.

As shown in the left panel of Figure 6.35, the analysis performed in this work covers the p_T range from 2 to 8 GeV/c, while the run 1 analysis starts at $p_T = 1.25$ GeV/c. Although the two bins inside the p_T range from 1.25-2 GeV/c were also analyzed in the present work, they are not included in the final result, since their values presented a strong dependence on the DCA bin size of the templates, as explained in Appendix F. The origins of the observed instability are currently under investigation.

An independent analysis is performed by ALICE Collaboration at the same center of mass energy with the purpose of extending the measurement performed in this work to

Figure 6.34: (Left) R_{AA} of electrons from beauty hadron decays measured in the current work and (right) R_{pPb} of electrons from beauty hadron decays at the same center of mass energy [53]. The filled boxes at unity, in both panels, represent the systematic uncertainties not related to the signal extraction (normalization and scaling uncertainties combined in quadrature).

larger p_T regions, by making use of the EMCal triggered events. The high p_T analysis is completely independent of the low p_T one discussed in this chapter, but it is performed with a similar strategy. The main differences are that in the high p_T analysis the EMCal detector is used for the electron identification together with the TPC (TOF is not used) and the non heavy-flavor electron background as well as the hadron contamination are identified via data-driven approaches. Then, the electrons from beauty hadron decays are identified by performing a template fit on the DCA distribution of the *heavy-flavor electrons*, in which just two templates, of electrons from charm and beauty hadron decays, are considered. The high p_T analysis could be extended to low p_T down to 3 GeV/c and the comparison of the obtained R_{AA} with the measurement of the present work shows a complete agreement between the two results within uncertainties in the common p_T range, as shown in the right panel of Figure 6.35. Both comparisons shown in Figure 6.35 ensure the reliability of the result of this work and of the method used.

In the further discussions, the result from this work will be presented together with the complementary result from the high p_T analysis.

Figure 6.35: (Left) Comparison between the R_{AA} of electrons from beauty hadron decays measured in the current work and the one measured in Pb-Pb at $\sqrt{s_{NN}} = 2.76$ TeV [53]. (Right) Comparison between the R_{AA} measured in the current work with the one obtained from the high p_T analysis performed with the usage of EMCal.

For the first time in ALICE the R_{AA} of electrons from beauty hadron decays is obtained in the wide p_T range of 2 to 26 GeV/c (where the range $2 < p_T < 8$ GeV/c comprises the result from this work).

The R_{AA} of electrons from beauty hadron decays and electrons from open heavy-flavor (charm + beauty) hadron decays measured in Pb-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV are compared in Figure 6.36. From this comparison, a hint of less suppression of electrons from beauty hadron decays than of electrons from open heavy-flavor hadron decays is observed for $2 < p_T \lesssim 7$ GeV/c. For $p_T \gtrsim 7$ GeV/c, both R_{AA} show similar suppression. It is known from pp measurements as well as from FONLL calculations that at ≈ 4 GeV/c the open heavy-flavor electrons spectrum have a contribution from beauty and charm hadron decays of $\approx 50\%$ each (see Appendix C and Figure 5.13). As the p_T increases, the beauty contribution overcomes the one of charm hadron decays, becoming dominant. This can explain why at $p_T \gtrsim 7$ GeV/c the suppression of the R_{AA} of electrons from beauty and open heavy-flavor hadron decays is essentially the same.

The R_{AA} of electrons from beauty hadron decays measured at $\sqrt{s_{NN}} = 5.02$ TeV

Figure 6.36: R_{AA} of electrons from beauty hadron decays (red: this work, blue: high p_T analysis), compared to the R_{AA} of electrons from open heavy-flavor hadron decays (gray) in Pb-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV.

could be compared to three available pQCD-based predictions. The PHSD ¹ [85] [73], Djordjevic [65] and MC@sHQ+EPOS2 [82] predictions are fully compatible with the measurement, over the full p_T range accessible by the models, as shown in Figure 6.37. In Table 6.7, some important features of these models, also discussed briefly in section 3.3, are summarized in order to facilitate the present discussion.

As pointed in Table 6.7, the PHSD model describes the energy loss of heavy quarks in the color-deconfined medium via only collisional processes. This is the reason why its prediction is restricted to the low p_T region of the spectrum, where the collisional processes are dominant. The Djordjevic and MC@sHQ+EPOS2 predictions describe,

¹It is important to note that the PHSD references pointed out correspond to a version of this model in which just the energy loss of charm quarks is implemented, and no description of beauty quarks is made. The prediction from PHSD of the R_{AA} of electrons from beauty hadron decays shown in Figure 6.37 is a very recent work, and there is still no published paper that describes the implementation of the beauty quark interaction with the QGP, although all the processes should be analogous to the charm quark description.

within experimental uncertainties, the suppression of about 50% observed in the experimental data at large p_T and they are also fully comparable with each other from $p_T > 5$ GeV/ c . The Djordjevich model does not provide prediction for the very low p_T region of the R_{AA} since it models the hadronization via only fragmentation processes, and the missing coalescence could compromise the prediction of the particle production in Pb-Pb at low p_T , where this process can become important. From this comparison with models, it is possible to affirm that the theoretical description of a mass-dependent energy loss of heavy-quarks inside QGP as predicted by QCD leads to a reasonably good description of the R_{AA} of electrons from beauty hadron decays presented in this work, from $p_T = 2$ to 26 GeV/ c . As shown in the left panel of [Figure 6.35](#), the current result has significantly reduced systematic uncertainties at $p_T \approx 2 - 3$ GeV/ c with respect to what was measured with run 1 data, and it hints that the R_{AA} stays very close to unity in this region. This is in very good agreement with what PHSD and MC@sHQ+EPOS2 predicts, as shown in [Figure 6.37](#).

Other processes can have an important influence in the R_{AA} at very low p_T , in addition to the energy loss, as the shadowing of the nuclear PDFs, that can suppress the low momentum particle production, and coalescence as a possible hadronization mechanism, that can enhance the low momentum particle production in Pb-Pb [\[74\]](#). From [Table 6.7](#) it is possible to see that both PHSD and MC@sHQ+EPOS2 consider these effects. Another important effect is the radial flow, that is a consequence of the hydrodynamical expansion of the medium, and that can enhance the R_{AA} at low p_T [\[9\]](#). From [Table 6.7](#) it is possible to see that the MC@sHQ+EPOS2 is the only of these three models that includes the radial flow. This can be one of the reasons why it predicts a slightly larger R_{AA} at very low p_T with respect to PHSD. In despite of this, within the current experimental resolution, both models agree very well with the R_{AA} measured in this work at very low p_T .

No measurement of v_2 of electrons from beauty hadron decays in Pb-Pb at $\sqrt{s_{NN}} = 5.02$ TeV is available currently, therefore the simultaneous description of v_2 and R_{AA} by theoretical models still cannot be tested.

Figure 6.37: R_{AA} of electrons from beauty hadron decays measured in Pb-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV in central rapidity compared with three pQCD-based model predictions.

	PHSD [85] [73]	Djordjevic [65]	MC@sHQ [82]
Heavy-quark production	PYTHIA tunned on FONLL, PDF shadowing	FONLL, no PDF shadowing	FONLL, PDF shadowing (EPS09 (LO) [16])
Medium modeling	no fl. dyn. evolution	Glauber model nuclear overlap, no fl. dyn. evolution	3+1 d expansion (EPOS model)
Quark-medium interactions	collisional energy loss	rad. + coll. energy loss	rad. + coll. energy loss (transport with Boltzman equation)
Heavy quark hadronization	fragmentation and coalescence	fragmentation	fragmentation and coalescence

Table 6.7: Main features of the three pQCD-based model predictions available for comparison with the measured R_{AA} of electrons from beauty hadron decays in Pb-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV in central rapidity.

Chapter 7

Summary

In this work, two analysis are performed on the heavy-flavor sector using recent data collected by ALICE, at LHC. Both analysis make use of electrons from heavy-flavor hadron decays to probe the production of heavy quarks.

In the first analysis, the invariant cross section of electrons from heavy-flavor hadron decays is measured in pp at $\sqrt{s} = 13$ TeV collisions, at low p_T (from 0.5 to 4 GeV/ c) and central rapidity ($|\eta| < 0.8$). In this analysis, the non heavy-flavor electron background is identified via the data-driven *photonic tagging method*, leading to a high precision heavy-flavor electron cross section. The experimental measurement is compared to the theoretical prediction from FONLL. In this way, this pQCD-based prediction could be tested on pp collisions at the largest center of mass energy generated by LHC up to now. The experimental result sits on the upper edge of the FONLL prediction band, as observed on similar measurements performed at lower center of mass energy and also in the measurements of heavy-flavor mesons. The ratios between the cross sections of electrons from heavy-flavor hadron decays at different center of mass energies could be performed and compared to the corresponding ratios from FONLL. These ratios can bring constrains to further improve the theoretical calculations.

In the second analysis, the production of electrons from beauty hadron decay is measured in Pb-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, for low p_T (from 2 to 8 GeV/ c) and central rapidity ($|\eta| < 0.8$) regimes. This analysis explores the fact that electrons from beauty hadron decays have a broader DCA distribution with respect to electrons from

the background sources. This feature allows the extraction of the beauty contribution, from the DCA distribution of inclusive electrons, via a maximum-likelihood template fit of the different electron sources. The templates are taken from Monte Carlo simulations and they consist in DCA distributions of electrons from Dalitz decay of light mesons, gamma conversion, charm hadron and beauty hadron decays. The QGP is expected to be produced in Pb-Pb collisions, and the modification of the p_T -spectrum of the measured electrons from beauty hadron decays with respect to the measurement in pp collisions at the same center of mass energy is quantified by the nuclear modification factor R_{AA} . A suppression of about 40-50% is observed in the R_{AA} for $p_T > 3$ GeV/ c . This suppression is consistent with the scenario in which the color-deconfined medium is produced in the early stages of Pb-Pb collisions and the heavy quarks interact with this medium losing energy via collisional and radiative processes in a mass-dependent way, following pQCD descriptions over the full measured p_T range. At very low p_T (from ≈ 2 to 3 GeV/ c) this analysis results in significantly reduced final uncertainties with respect to the run 1 measurement and it is consistent with unity within 1σ , following the prediction of models which include shadowing of the nuclear PDF and coalescence as a possible hadronization mechanism. The usage of an upcoming new pp reference will further reduce the final systematic uncertainties of the measured R_{AA} , improving the comparison of this result to models predictions and to other experimental results. The result of this work hints that electrons from beauty hadron decays undergo less suppression with respect to electrons from open heavy-flavor hadron decays in the range $2 < p_T < 6$ GeV/ c . For larger p_T it is observed a similar suppression on both results, that can be understood by the dominance of the beauty contribution to the open heavy-flavor electrons sample in this p_T range.

Both of the analysis performed in this work were approved by ALICE Collaboration as preliminary results and presented in important conferences of the field.

Appendix A

Other Evidences of the QGP Formation

Together with the nuclear modification factor, the quantities presented here are sensitive to different properties of the QGP. Above all, these quantities are observables that were early used to indicate the very existence of the color-deconfined medium.

A.1 Azimutal Anisotropy

Another observable that is sensitive to the dynamics of the early stages of the collision is the azimuthal anisotropy of the produced particles in space and momentum. In a heavy-ion collisions in which the nuclei collide with a non-zero impact parameter (non-central collision), the overlap region has an anisotropic shape (ellipsoid shape), like shown at [Figure A.1](#).

In the case the matter inside this overlap region becomes a strongly interacting macroscopic medium, an anisotropic pressure gradient will be developed as a consequence of the initial anisotropy and of the strong re-interactions among the produced medium constituents. Due to this pressure gradient, this initial anisotropy is converted, during the expansion of the system, into a momentum anisotropy, that leads to an anisotropic particle distribution along the azimuthal angle ϕ [\[125\]](#) [\[26\]](#).

The anisotropic distribution of particles along the azimuthal angle can be character-

Figure A.1: Schematic view of a nucleus-nucleus collision in a plane perpendicular to the beam direction. Are indicated the overlap region (with ellipsoid shape), the spectators (that refers to the nucleons that do not participate in the collision) the event plane of the collision Ψ_{RP} , and the impact parameter b , that is defined as the distance between the centers of the two colliding nucleus.

ized by the coefficients v_n of the Fourier series [126]:

$$v_n = \langle \cos[n(\phi - \Psi_n)] \rangle, \quad (\text{A.1})$$

where Ψ_n is the event plane angle and n refers to the n -th harmonic.

In non-central collision, the largest contribution corresponds to the second harmonic, v_2 , called elliptic flow. It represents the ellipticity of the produced particle distribution.

The large measured elliptic flow of light particles at low momentum measured firstly at RHIC can be described by models that assumes that the bulk of the produced matter underwent a thermodynamical equilibrium in the early stages of the collision (at $\tau \approx 1$ fm/c), evolved following hydrodynamical laws and behaved as a nearly perfect fluid. It is understood that this early thermalization can be achieved only in a color-deconfined strongly interacting medium. In a scenario without this quick thermalization induced by the strong interactions and without hydrodynamical expansion, the large v_2 observed experimentally can not be reproduced by theoretical models [125]. The measurements of v_2 of the bulk matter supports the existence of the QGP and indicates it behaves like a non-viscous fluid.

The elliptic flow of heavy-flavor particles can provide further information about the transport properties of the medium. Figure A.2 shows the measurement of D mesons v_2 performed at central rapidity in ALICE. The non-zero elliptic flow of the D mesons at low momentum ($2 < p_T < 8$ GeV/c) indicates a participation of the charm quark in the collective flow of the QCD medium [127], what implies that also heavy-flavor quarks can undergo a certain degree of thermalization inside the QGP, despite their large mass. The measurement of v_2 of heavy-flavor particles at high momentum is shown to be sensitive to the path-length (that refers to the size of the traversed medium) dependence of energy loss [128]: it is predicted that the energy loss of the quark can give rise to a positive v_2 at high momentum.

Figure A.2: Elliptic flow coefficient v_2 for average non-strange D mesons measured at mid-rapidity by ALICE [129].

A.2 Jet Suppression

High momentum quarks, as well as heavy quarks, are created mainly by the hard scattering of partons from the two incoming nucleon. In this process, a pair of energetic partons is created in a back-to-back configuration. Since these partons carry a lot of energy, each one fragment in a shower of other partons, which undergo hadronization. This process gives origin to two showers of hadrons emitted in opposite directions, separated by the angle $\Delta\phi = \pi$, which can be measured experimentally. A jet is defined as

a group of particles, mostly hadrons, emitted within a narrow cone, which is expected to originate from the fragmentation of a high momentum parton.

Since jets are always produced in pairs, the two-hadron spacial correlation in pp collisions can give insight about the back-to-back jet structure. The azimuthal correlation between a particle from a jet and other particles from the same jet creates an enhancement at $\Delta\phi \approx 0$ (near side peak), while the azimuthal correlation between this particle and the particles from the second jet creates an enhancement at $\Delta\phi \approx \pi$ (away side peak). In nucleus-nucleus collisions, medium related effects can modify the two hadron azimuthal correlation with respect to what is measured in pp. The high momentum parton can lose a large fraction of its energy while interacting with the QGP by elastic and inelastic collisions with the partons of the medium. This energy loss may effectively quench (reduce the intensity of) the outgoing jet. In addition, studies of two particle correlation in nucleus-nucleus collisions can indicate possible medium-induced modification of the fragmentation function [5] [130].

Figure A.3 shows the two-hadron azimuthal correlation $\Delta\phi$ measured in pp, d-Au and Au-Au collisions by STAR [130]. While the same-side correlations remain unchanged in the three collision systems, the away-side correlations are strongly suppressed in Au-Au, vanishing completely. The same does not happen in d-Au collisions, indicating that this strong quenching of the away-side peak is not related to initial conditions, but with energy loss in the dense medium. In the situation where the QGP is formed, the hard hadrons that are measured (constituents of the near-side correlation peak) would arise preferentially from partons scattered outward from the surface region of the collision zone, while the away-side partons traverse a large fraction of the QGP and lose a large fraction of its energy, what leads to a strong or complete quenching of the jet. A sketch of this idea is depicted on Figure A.4.

The integral over the near-side and away-side peak distributions gives the yield that is used to study quantitatively the away-side suppression in nucleus-nucleus relative to pp collisions and other medium induced effects.

Figure A.3: Azimuthal correlation between high momentum hadrons measured by STAR in pp, d-Au and Au-Au collisions [130].

Figure A.4: Representation of the production of a high momentum quark pair in a back-to-back configuration. One of the quarks is emitted in the outward direction, interacting very little with the QGP, while the second one is emitted inwards and loses energy in the dense QCD medium.

A.3 Quarkonia Sequential Melting

Quarkonia consists in the bound states of heavy-quarks, like $c\bar{c}$ (charmonium) and $b\bar{b}$ (bottomonium). The study of quarkonia states in high-energy nucleus-nucleus collisions

has shown to be a sensitive probe of the thermodynamical properties of color-deconfined matter. The QGP is expected to screen the potential that confines the $c\bar{c}$ or $b\bar{b}$ bound states, causing their "melting" (dissociation). The melting temperature depends on the binding energy of the quarkonium state. The ground states, more strongly bounded, are expected to melt at higher temperatures than the less-tightly bounded resonant states. The dissociation happens when the color (Debye) screening radius of the medium falls below their corresponding $Q\bar{Q}$ binding radius [21].

Experimental studies of the J/ψ suppression in different collision energies have found a good description from the theoretical side, which considers the melting of the charmonium states by color screening and the charmonium enhancement due to coalescence as interplaying phenomena [131].

LHC brings unique opportunities to study the dissociation of the bottomonium $b\bar{b}$ bound states as well. According to lattice QCD, the Υ is expected to survive up to temperatures of $4 T_c$, whereas the resonances $\Upsilon(2s)$ survives up to $1.6 T_c$ and $\Upsilon(3s)$ up to $1.2 T_c$. This creates a sequential melting pattern, that is understood to be a signature of the QGP [132]. This sequential melting can be seen at Figure A.5, which shows the signal of Υ , $\Upsilon(2s)$ and $\Upsilon(3s)$ originated from the invariant mass analysis of the decay products $\mu^+\mu^-$, measured in Pb-Pb collisions. In this figure it is also shown the signal of the same quarkonia states from pp collisions (fit of the measurement), where a normalization is performed to make the ground state of both measurements to have the same amplitude. The resonant states are suppressed in central Pb-Pb relative to pp collisions, and the suppression is larger the less bounded the state is [133].

Figure A.5: $\Upsilon(nS)$ states measured in Pb-Pb (data points) compared to the fit of the pp measurement (dashed blue line) at 2.76 TeV by CMS [131].

Appendix B

Pseudo-Rapidity and Rapidity

The **rapidity** is a variable used in relativistic kinematics, defined as

$$y = \frac{1}{2} \ln \frac{E + p_z}{E - p_z} = \frac{1}{2} \ln \frac{1 + p_z/E}{1 - p_z/E} = \tanh^{-1} \left(\frac{p_z}{E} \right), \quad (\text{B.1})$$

and it has the advantage of being additive under longitudinal boost.

The **pseudo-rapidity** η can be derived from the rapidity y as shown below:

$$y = \frac{1}{2} \ln \frac{E + p_z}{E - p_z} = \frac{1}{2} \ln \frac{\sqrt{m^2 + p^2} + p \cdot \cos\theta}{\sqrt{m^2 + p^2} - p \cdot \cos\theta}. \quad (\text{B.2})$$

In the limit $p \gg m$, the mass can be neglected:

$$y = \frac{1}{2} \ln \frac{p + p \cdot \cos\theta}{p - p \cdot \cos\theta} = -\ln \tan(\theta/2) \equiv \eta. \quad (\text{B.3})$$

The pseudo-rapidity η depends only on polar angle of emission θ of the particle, and it is very useful when other properties of the particle (like mass or energy) are not known necessarily. When $\theta = 90$ (in the direction perpendicular to the beams), $\eta = 0$, as can be seen in [Figure B.1](#).

Figure B.1: Pseudo-rapidity η as a function of the polar emission angle θ .

Appendix C

Charm and Beauty Contribution to Open Heavy-Flavor Electrons Spectrum

Measurement of electrons from beauty hadron decays performed using the impact parameter cut, as describe on [subsection 6.11.1](#). [Figure C.1](#) shows the cross section of electrons from beauty hadron decays ($b(\rightarrow c)\rightarrow e$) and also the one of electrons from charm hadron decays ($c\rightarrow e$) compared to FONLL predictions. On the bottom panel the ratio between $b(\rightarrow c)\rightarrow e$ and $c\rightarrow e$ is shown, and is possible to see that the contribution of beauty overcomes the contribution of charm for $p_T \gtrsim 4$ GeV/ c .

Figure C.1: Cross section of electrons from beauty and charm hadron decays performed in pp at $\sqrt{s} = 7$ TeV [12]. On the bottom panel is shown the ratio between electrons from beauty and electrons from charm hadron decays as function of p_T .

Appendix D

Gamma Conversion in the Detector Material

Figure D.1: Sketch of a gamma conversion on the beam pipe. Figure taken from [118].

Figure D.1 shows a sketch of a gamma conversion on the detector material, as described on section 6.7. The gamma have a straight trajectory from the primary vertex (most part of the gammas comes from $\pi^0 \rightarrow 2\gamma$ decays, so they originate essentially on the primary vertex). Electrons from gamma conversion have a typical opening angle of the order of $m_e c^2 / E_e$ [134]. For the energies considered in this work, the initial trajec-

tories of the electrons can be considered parallel to the trajectory of the gamma. The magnetic field B bends the electrons trajectory, and it will have in the end the radius r :

$$r = \frac{p_T}{|q|B}, \quad (\text{D.1})$$

where q is the charge of the particle and p_T its transversal momentum.

According to [Figure D.1](#) is possible to conclude that a typical impact parameter of electrons from gamma conversion will be:

$$d_0 = \sqrt{R^2 + r^2} - r = r \left(\sqrt{1 + \frac{R^2}{r^2}} - 1 \right), \quad (\text{D.2})$$

where R is the radius related to the location of the detector material. For the energies considered here, $R \ll r$, and a Taylor expansion leads to:

$$d_0 \approx \frac{R^2 |q| B}{2p_T}. \quad (\text{D.3})$$

Appendix E

ITS Quality Cuts

Figure E.1 shows the template of electrons from different sources, from MB Monte Carlo simulations, before any ITS quality cut (top left), after the cut on $\chi^2/ndf < 5$ (top right) and after the cuts on $\chi^2/ndf < 5$ and $f_{shared} < 0.3$ (bottom). These cuts reduce the tails of the template of conversion electrons, making the final distribution slimmer than the beauty electron template.

Figure E.1: Template of electrons from different sources, from MB Monte Carlo, before any ITS quality cut (top left), after the cut on the χ^2 (top right) and after the cuts on χ^2 and f_{shared} (bottom).

Appendix F

Fit Instability due to DCA Bin Size

Chronologically, the first DCA distributions used in this work had a DCA bin size of 0.005 cm. As soon as the whole analysis was performed and the R_{AA} of electrons from beauty hadron decays was compared to the result from run 1, an incompatibility between these two results was observed for the p_T bins in the interval 1 to 2 GeV/ c . Nevertheless, this large incompatibility has disappeared when the DCA distributions were built with DCA bin size of 0.0005 cm, the same used in run 1 analysis. From this observation, the DCA bin size of 0.0005 cm was taken as default bin size in this analysis, as in run 1. Also, the default DCA range in which the DCA template fit is performed is $-0.2 < \text{DCA} < 0.2$ cm. Further studies showed that the DCA resolution of the templates are of the order of 0.001 cm, so a reasonable choice was to use a significantly smaller DCA bin size as default.

But in face of this unexpected dependence of the final result (in the p_T interval of 1 to 2 GeV/ c) on the DCA bin size of the distributions, different DCA bin sizes were tested. [Figure F.1](#) shows the ratios between the yield of electrons from beauty hadron decays obtained using different variations of the DCA bin size and of the DCA range used for the fit and the yield obtained with default values. Also cases in which the DCA bin size on the tails ($|\text{DCA}| > 0.1$ cm) is different from the DCA bin size used in the peak of the distribution were tested.

The two bins with $p_T < 2$ GeV/ c show a strong sensitivity to these different ingredients. The usage of a coarser tail seems to have almost the same effect of not considering

at all the tails (a region where the beauty contribution is important) for the fits. In these two first bins, these variations makes the final yield to be easily smaller than the default yield in more than 10%.

Since from $p_T > 2 \text{ GeV}/c$ the final yield is much more stable under these variations, the final results of this analysis is presented from $p_T > 2 \text{ GeV}/c$. Further investigations on this subject are being carried on.

Figure F.1: Effect, on the final yield of beauty electrons, of considering different DCA bin sizes and DCA ranges for the fit, relatively to the default yield. The DCA bin sizes in the legend are given in centimeters.

Bibliography

- [1] K. Aamodt et al. The ALICE experiment at the CERN LHC. **JINST**, v. 3, p. S08002, 2008.
- [2] Edmond Iancu. QCD in heavy ion collisions. In **Proceedings, 2011 European School of High-Energy Physics (ESHEP 2011): Cheile Gradistei, Romania, September 7-20, 2011**, pages 197–266, 2014.
- [3] A. K. Chaudhuri. **A short course on Relativistic Heavy Ion Collisions**. IOPP, 2014.
- [4] Ulrich W. Heinz. The Strongly coupled quark-gluon plasma created at RHIC. **J. Phys.**, v. A42, p. 214003, 2009.
- [5] John Adams et al. Experimental and theoretical challenges in the search for the quark gluon plasma: The STAR Collaboration’s critical assessment of the evidence from RHIC collisions. **Nucl. Phys.**, v. A757, p. 102–183, 2005.
- [6] K. Adcox et al. Formation of dense partonic matter in relativistic nucleus-nucleus collisions at RHIC: Experimental evaluation by the PHENIX collaboration. **Nucl. Phys.**, v. A757, p. 184–283, 2005.
- [7] B. B. Back et al. The PHOBOS perspective on discoveries at RHIC. **Nucl. Phys.**, v. A757, p. 28–101, 2005.
- [8] I. Arsene et al. Quark gluon plasma and color glass condensate at RHIC? The Perspective from the BRAHMS experiment. **Nucl. Phys.**, v. A757, p. 1–27, 2005.

- [9] A. Andronic et al. Heavy-flavour and quarkonium production in the LHC era: from protonâ€‘proton to heavy-ion collisions. **Eur. Phys. J.**, v. C76, n. 3, p. 107, 2016.
- [10] Georges Aad et al. Measurement of the differential cross-section of B^+ meson production in pp collisions at $\sqrt{s} = 7$ TeV at ATLAS. **JHEP**, v. 10, p. 042, 2013.
- [11] R Aaij et al. Prompt charm production in pp collisions at $\sqrt{s}=7$ TeV. **Nucl. Phys.**, v. B871, p. 1–20, 2013.
- [12] Betty Abelev et al. Measurement of electrons from beauty hadron decays in pp collisions at $\sqrt{s} = 7$ TeV. **Phys. Lett.**, v. B721, p. 13–23, 2013. [Erratum: Phys. Lett.B763,507(2016)].
- [13] Betty Abelev et al. Measurement of electrons from semileptonic heavy-flavour hadron decays in pp collisions at $\sqrt{s} = 7$ TeV. **Phys. Rev.**, v. D86, p. 112007, 2012.
- [14] Panagiota Foka and MaÅgorzata Anna Janik. An overview of experimental results from ultra-relativistic heavy-ion collisions at the CERN LHC: Hard probes. **Rev. Phys.**, v. 1, p. 172–194, 2016.
- [15] Matteo Cacciari, Michelangelo L. Mangano, and Paolo Nason. Gluon PDF constraints from the ratio of forward heavy-quark production at the LHC at $\sqrt{S} = 7$ and 13 TeV. **Eur. Phys. J.**, v. C75, n. 12, p. 610, 2015.
- [16] K. J. Eskola, H. Paukkunen, and C. A. Salgado. EPS09: A New Generation of NLO and LO Nuclear Parton Distribution Functions. **JHEP**, v. 04, p. 065, 2009.
- [17] Alexandre Deur, Stanley J. Brodsky, and Guy F. de Teramond. The QCD Running Coupling. **Prog. Part. Nucl. Phys.**, v. 90, p. 1–74, 2016.
- [18] David J. Gross and Frank Wilczek. Ultraviolet behavior of non-abelian gauge theories. **Phys. Rev. Lett.**, v. 30, p. 1343–1346, Jun 1973.

- [19] Claudia Ratti. Lattice QCD and heavy ion collisions: a review of recent progress. **Rept. Prog. Phys.**, v. 81, n. 8, p. 084301, 2018.
- [20] Andrea Dubla, Raimond Snellings, and Alessandro Grelli. Elliptic flow at different collision stages, Apr 2016. Presented 06 Apr 2016.
- [21] Wit Busza, Krishna Rajagopal, and Wilke van der Schee. Heavy Ion Collisions: The Big Picture, and the Big Questions. **Ann. Rev. Nucl. Part. Sci.**, v. 68, p. 339–376, 2018.
- [22] Y. Aoki, G. Endrodi, Z. Fodor, S. D. Katz, and K. K. Szabo. The Order of the quantum chromodynamics transition predicted by the standard model of particle physics. **Nature**, v. 443, p. 675–678, 2006.
- [23] V. Antunes, I. Bediaga, and M. Novello. Gravitational mechanism for baryogenesis in the cosmological QCD phase transition. v., 2016.
- [24] Francesco Becattini. The Quark Gluon Plasma and relativistic heavy ion collisions in the LHC era. **J. Phys. Conf. Ser.**, v. 527, p. 012012, 2014.
- [25] Raimond Snellings. Elliptic Flow: A Brief Review. **New J. Phys.**, v. 13, p. 055008, 2011.
- [26] Panagiota Foka and MaÅgorzata Anna Janik. An overview of experimental results from ultra-relativistic heavy-ion collisions at the CERN LHC: Bulk properties and dynamical evolution. **Rev. Phys.**, v. 1, p. 154–171, 2016.
- [27] Markus H. Thoma and Miklos Gyulassy. Quark Damping and Energy Loss in the High Temperature QCD. **Nucl. Phys.**, v. B351, p. 491–506, 1991.
- [28] Miklos Gyulassy and Michael Plumer. Jet Quenching in Dense Matter. **Phys. Lett.**, v. B243, p. 432–438, 1990.
- [29] Michelangelo L. Mangano. Two lectures on heavy quark production in hadronic collisions. **Proc. Int. Sch. Phys. Fermi**, v. 137, p. 95–137, 1998.

- [30] M. Tanabashi et al. Review of Particle Physics. **Phys. Rev.**, v. D98, n. 3, p. 030001, 2018.
- [31] Shreyasi Acharya et al. Measurement of D-meson production at mid-rapidity in pp collisions at $\sqrt{s} = 7$ TeV. **Eur. Phys. J.**, v. C77, n. 8, p. 550, 2017.
- [32] B. A. Kniehl, G. Kramer, I. Schienbein, and H. Spiesberger. Collinear subtractions in hadroproduction of heavy quarks. **Eur. Phys. J.**, v. C41, p. 199–212, 2005.
- [33] B. A. Kniehl, G. Kramer, I. Schienbein, and H. Spiesberger. Inclusive Charmed-Meson Production at the CERN LHC. **Eur. Phys. J.**, v. C72, p. 2082, 2012.
- [34] Matteo Cacciari, Mario Greco, and Paolo Nason. The P(T) spectrum in heavy flavor hadroproduction. **JHEP**, v. 05, p. 007, 1998.
- [35] Matteo Cacciari, Stefano Frixione, Nicolas Houdeau, Michelangelo L. Mangano, Paolo Nason, and Giovanni Ridolfi. Theoretical predictions for charm and bottom production at the LHC. **JHEP**, v. 10, p. 137, 2012.
- [36] J. F. Owens. Large-momentum-transfer production of direct photons, jets, and particles. **Rev. Mod. Phys.**, v. 59, p. 465–503, Apr 1987.
- [37] John C. Collins, Davison E. Soper, and George F. Sterman. Factorization of Hard Processes in QCD. **Adv. Ser. Direct. High Energy Phys.**, v. 5, p. 1–91, 1989.
- [38] Matteo Cacciari, Paolo Nason, and Carlo Oleari. A Study of heavy flavored meson fragmentation functions in $e^+ e^-$ annihilation. **JHEP**, v. 04, p. 006, 2006.
- [39] S. Kretzer and I. Schienbein. Heavy quark fragmentation in deep inelastic scattering. **Phys. Rev.**, v. D59, p. 054004, 1999.
- [40] Bernd A. Kniehl and Gustav Kramer. Charmed-hadron fragmentation functions from CERN LEP1 revisited. **Phys. Rev.**, v. D74, p. 037502, 2006.

- [41] Diego Stocco. Measurement of heavy-flavour decay muon production at forward rapidity in pp and Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the ALICE experiment. **Nucl. Phys.**, v. A910-911, p. 355–358, 2013.
- [42] Shreyasi Acharya et al. Λ_c^+ production in pp collisions at $\sqrt{s} = 7$ TeV and in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV. **JHEP**, v. 04, p. 108, 2018.
- [43] Stefano Frixione, Paolo Nason, and Giovanni Ridolfi. A Positive-weight next-to-leading-order Monte Carlo for heavy flavour hadroproduction. **JHEP**, v. 09, p. 126, 2007.
- [44] Torbjorn Sjostrand, Stephen Mrenna, and Peter Z. Skands. PYTHIA 6.4 Physics and Manual. **JHEP**, v. 05, p. 026, 2006.
- [45] Michael L. Miller, Klaus Reygers, Stephen J. Sanders, and Peter Steinberg. Glauber modeling in high energy nuclear collisions. **Ann. Rev. Nucl. Part. Sci.**, v. 57, p. 205–243, 2007.
- [46] T. W. Ludlam, A. Pfoh, and A. Shor. HIJET. A MONTE CARLO EVENT GENERATOR FOR P NUCLEUS AND NUCLEUS NUCLEUS COLLISIONS. Em Brookhaven Nat. Lab. Upton - BNL-37196 (REC.FEB.86) 9p, IN *BROOKHAVEN 1985, PROCEEDINGS, EXPERIMENTS FOR A RELATIVISTIC HEAVY ION COLLIDER*, 373-381., 1986.
- [47] Constantin Loizides, Jason Kamin, and David d’Enterria. Improved Monte Carlo Glauber predictions at present and future nuclear colliders. **Phys. Rev.**, v. C97, n. 5, p. 054910, 2018.
- [48] Roy J. Glauber. Quantum Optics and Heavy Ion Physics. **Nucl. Phys.**, v. A774, p. 3–13, 2006.
- [49] Serguei Chatrchyan et al. Measurement of isolated photon production in pp and PbPb collisions at $\sqrt{s_{NN}} = 2.76$ TeV. **Phys. Lett.**, v. B710, p. 256–277, 2012.
- [50] M. M. Aggarwal et al. Observation of direct photons in central 158-A-GeV Pb-208 + Pb-208 collisions. **Phys. Rev. Lett.**, v. 85, p. 3595–3599, 2000.

- [51] S. S. Adler et al. Centrality dependence of direct photon production in $s(NN)^{**}(1/2) = 200\text{-GeV Au} + \text{Au}$ collisions. **Phys. Rev. Lett.**, v. 94, p. 232301, 2005.
- [52] Leonid Frankfurt, Mark Strikman, and Christian Weiss. Small-x physics: From HERA to LHC and beyond. **Ann. Rev. Nucl. Part. Sci.**, v. 55, p. 403–465, 2005.
- [53] Jaroslav Adam et al. Measurement of electrons from beauty-hadron decays in p-Pb collisions at $\sqrt{s_{NN}} = 5.02\text{ TeV}$ and Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76\text{ TeV}$. **JHEP**, v. 07, p. 052, 2017.
- [54] Roel Aaij et al. Study of prompt D^0 meson production in pPb collisions at $\sqrt{s_{NN}} = 5\text{ TeV}$. **JHEP**, v. 10, p. 090, 2017.
- [55] Néstor Armesto, Andrea Dainese, Carlos A. Salgado, and Urs Achim Wiedemann. Testing the color charge and mass dependence of parton energy loss with heavy-to-light ratios at bnl rhic and cern lh. **Phys. Rev. D**, v. 71, p. 054027, Mar 2005.
- [56] Miklos Gyulassy and Xin nian Wang. Multiple collisions and induced gluon bremsstrahlung in qcd. **Nuclear Physics B**, v. 420, n. 3, p. 583 – 614, 1994.
- [57] R. Baier, Yuri L. Dokshitzer, Alfred H. Mueller, S. Peigne, and D. Schiff. Radiative energy loss of high-energy quarks and gluons in a finite volume quark - gluon plasma. **Nucl. Phys.**, v. B483, p. 291–320, 1997.
- [58] B. G. Zakharov. Radiative energy loss of high-energy quarks in finite size nuclear matter and quark - gluon plasma. **JETP Lett.**, v. 65, p. 615–620, 1997.
- [59] Urs Achim Wiedemann. Gluon radiation off hard quarks in a nuclear environment: Opacity expansion. **Nucl. Phys.**, v. B588, p. 303–344, 2000.
- [60] Yuri L. Dokshitzer, Valery A. Khoze, and S. I. Troian. On specific QCD properties of heavy quark fragmentation ('dead cone'). **J. Phys.**, v. G17, p. 1602–1604, 1991.

- [61] Fabio Maltoni, Michele Selvaggi, and Jesse Thaler. Exposing the dead cone effect with jet substructure techniques. **Phys. Rev.**, v. D94, n. 5, p. 054015, 2016.
- [62] Yuri L. Dokshitzer and D. E. Kharzeev. Heavy quark colorimetry of QCD matter. **Phys. Lett.**, v. B519, p. 199–206, 2001.
- [63] Ben-Wei Zhang, Enke Wang, and Xin-Nian Wang. Heavy quark energy loss in a nuclear medium. **Phys. Rev. Lett.**, v. 93, p. 072301, Aug 2004.
- [64] R. Thomas, Burkhard Kampfer, and G. Soff. Gluon emission of heavy quarks: Dead cone effect. **Acta Phys. Hung.**, v. A22, p. 83–91, 2005.
- [65] Magdalena Djordjevic and Marko Djordjevic. LHC jet suppression of light and heavy flavor observables. **Phys. Lett.**, v. B734, p. 286–289, 2014.
- [66] Morad Aaboud et al. Prompt and non-prompt J/ψ and $\psi(2S)$ suppression at high transverse momentum in 5.02 TeV Pb+Pb collisions with the ATLAS experiment. v., 2018.
- [67] Andrea Festanti. Measurement of D-meson production in Pb-Pb collisions at the LHC with ALICE. **J. Phys. Conf. Ser.**, v. 668, n. 1, p. 012054, 2016.
- [68] Vardan Khachatryan et al. Suppression and azimuthal anisotropy of prompt and nonprompt J/ψ production in PbPb collisions at $\sqrt{s_{NN}} = 2.76$ TeV. **Eur. Phys. J.**, v. C77, n. 4, p. 252, 2017.
- [69] Albert M Sirunyan et al. Measurement of the $B^\pm$ Meson Nuclear Modification Factor in Pb-Pb Collisions at $\sqrt{s_{NN}} = 5.02$ TeV. **Phys. Rev. Lett.**, v. 119, n. 15, p. 152301, 2017.
- [70] Albert M Sirunyan et al. Nuclear modification factor of D^0 mesons in PbPb collisions at $\sqrt{s_{NN}} = 5.02$ TeV. **Phys. Lett.**, v. B782, p. 474–496, 2018.
- [71] B. Abelev et al. Neutral pion and η meson production in proton-proton collisions at $\sqrt{s} = 0.9$ TeV and $\sqrt{s} = 7$ TeV. **Phys. Lett.**, v. B717, p. 162–172, 2012.

- [72] Rodolfo Sassot, Pia Zurita, and Marco Stratmann. Inclusive Hadron Production in the CERN-LHC Era. **Phys. Rev.**, v. D82, p. 074011, 2010.
- [73] E. L. Bratkovskaya, T. Song, H. Berrehrah, D. Cabrera, J. M. Torres-Rincon, L. Tolos, and W. Cassing. Heavy flavor in relativistic heavy-ion collisions. **J. Phys. Conf. Ser.**, v. 668, n. 1, p. 012008, 2016.
- [74] Francesco Scardina, Santosh K. Das, Vincenzo Minissale, Salvatore Plumari, and Vincenzo Greco. Estimating the charm quark diffusion coefficient and thermalization time from d meson spectra at energies available at the bnl relativistic heavy ion collider and the cern large hadron collider. **Phys. Rev. C**, v. 96, p. 044905, Oct 2017.
- [75] V. Greco, H. van Hees, and R. Rapp. Dynamics and Hadronization at intermediate transverse momentum at RHIC. **AIP Conf. Proc.**, v. 964, p. 232–237, 2007.
- [76] R. J. Fries, Berndt Muller, C. Nonaka, and S. A. Bass. Hadronization in heavy ion collisions: Recombination and fragmentation of partons. **Phys. Rev. Lett.**, v. 90, p. 202303, 2003.
- [77] Tuva Richert. Baryon enhancement in heavy ion collisions measured by the ALICE experiment. Taken from the link:
<http://agenda.albanova.se/getFile.py/access?contribId=259&resId=250&materialId=slides&confId=3646>.
- [78] Peter Koch, Berndt Müller, and Johann Rafelski. From strangeness enhancement to quark-gluon plasma discovery. **Int. J. Mod. Phys.**, v. A32, n. 31, p. 1730024, 2017.
- [79] P. Koch, Berndt Muller, and Johann Rafelski. Strangeness in Relativistic Heavy Ion Collisions. **Phys. Rept.**, v. 142, p. 167–262, 1986.

- [80] Jan Uphoff, Oliver Fochler, Zhe Xu, and Carsten Greiner. Elliptic Flow and Energy Loss of Heavy Quarks in Ultra-Relativistic heavy Ion Collisions. **Phys. Rev.**, v. C84, p. 024908, 2011.
- [81] Stefano Frixione and Bryan R. Webber. Matching NLO QCD computations and parton shower simulations. **JHEP**, v. 06, p. 029, 2002.
- [82] Marlene Nahrgang, Joerg Aichelin, Pol Bernard Gossiaux, and Klaus Werner. Influence of hadronic bound states above T_c on heavy-quark observables in Pb + Pb collisions at the CERN Large Hadron Collider. **Phys. Rev.**, v. C89, n. 1, p. 014905, 2014.
- [83] K. Werner, Iu. Karpenko, T. Pierog, M. Bleicher, and K. Mikhailov. Event-by-Event Simulation of the Three-Dimensional Hydrodynamic Evolution from Flux Tube Initial Conditions in Ultrarelativistic Heavy Ion Collisions. **Phys. Rev.**, v. C82, p. 044904, 2010.
- [84] M. Monteno, W. M. Alberico, A. Beraudo, A. De Pace, A. Molinari, M. Nardi, and F. Prino. Heavy-flavor dynamics in nucleus-nucleus collisions: from RHIC to LHC. **J. Phys.**, v. G38, p. 124144, 2011.
- [85] Taesoo Song, Hamza Berrehrah, Daniel Cabrera, Juan M. Torres-Rincon, Laura Tolos, Wolfgang Cassing, and Elena Bratkovskaya. Tomography of the Quark-Gluon-Plasma by Charm Quarks. **Phys. Rev.**, v. C92, n. 1, p. 014910, 2015.
- [86] Hamza Berrehrah, Elena Bratkovskaya, Thorsten Steinert, and Wolfgang Cassing. A dynamical quasiparticle approach for the QGP bulk and transport properties. **Int. J. Mod. Phys.**, v. E25, n. 07, p. 1642003, 2016.
- [87] Min He, Rainer J. Fries, and Ralf Rapp. Heavy Flavor at the Large Hadron Collider in a Strong Coupling Approach. **Phys. Lett.**, v. B735, p. 445–450, 2014.
- [88] W. A. Horowitz. Testing pQCD and AdS/CFT Energy Loss at RHIC and LHC. **AIP Conf. Proc.**, v. 1441, p. 889–891, 2012.

- [89] W. A. Horowitz and M. Gyulassy. Heavy quark jet tomography of Pb + Pb at LHC: AdS/CFT drag or pQCD energy loss? **Phys. Lett.**, v. B666, p. 320–323, 2008.
- [90] ALICE Collaboration, F Carminati, P Foka, P Giubellino, A Morsch, G Paic, J-P Revol, K SafarÅk, Y Schutz, and U A Wiedemann. Alice: Physics performance report, volume i. **Journal of Physics G: Nuclear and Particle Physics**, v. 30, n. 11, p. 1517, 2004.
- [91] Werner Riegler. Particle Physics Instrumentation. Em **Proceedings, 6th CERN - Latin-American School of High-Energy Physics (CLASHEP 2011): Natal, Brazil , March 23 - April 5, 2011**, pages 393–404, 2015. [,235(2015)].
- [92] G. Dellacasa et al. ALICE technical design report of the inner tracking system (ITS). v., 1999.
- [93] G. Dellacasa et al. ALICE: Technical design report of the time projection chamber. v., 2000.
- [94] P. Cortese. ALICE transition-radiation detector. v., 2001.
- [95] Francesca Carnesecchi. Performance of the ALICE Time-Of-Flight detector at the LHC. Em **14th Workshop on Resistive Plate Chambers and Related Detectors (RCP2018) Puerto Vallarta, Jalisco State, Mexico, February 19-23, 2018**, 2018.
- [96] G. De Cataldo. The high momentum particle identification detector in ALICE at LHC. **Nucl. Phys. Proc. Suppl.**, v. 177-178, p. 273–275, 2008.
- [97] G. Dellacasa et al. ALICE technical design report of the photon spectrometer (PHOS). v., 1999.
- [98] P. Cortese et al. ALICE electromagnetic calorimeter technical design report. v., 2008.

- [99] A. Fernandez, S. Kartal, and C. Pagliarone. ACORDE a cosmic ray detector for ALICE. Em **Summer School and Conference on New Trends in High-Energy Physics: Experiment, Phenomenology, Theory Yalta, Crimea, Ukraine, September 16-23, 2006**, 2006.
- [100] M. Bondila et al. ALICE T0 detector. **IEEE Trans. Nucl. Sci.**, v. 52, p. 1705–1711, 2005.
- [101] P Cortese et al. ALICE technical design report on forward detectors: FMD, T0 and V0. v., 2004.
- [102] E. Abbas et al. Performance of the ALICE VZERO system. **JINST**, v. 8, p. P10016, 2013.
- [103] Christian Holm Christensen, Jens Jorgen Gaardhoje, Kristjan Gulbrandsen, Borge Svane Nielsen, and Carsten Sogaard. The ALICE Forward Multiplicity Detector. **Int. J. Mod. Phys.**, v. E16, p. 2432–2437, 2007.
- [104] G. Dellacasa et al. ALICE technical design report: Photon multiplicity detector (PMD). v., 1999.
- [105] G. Dellacasa et al. ALICE technical design report of the zero degree calorimeter (ZDC). v., 1999.
- [106] Gines Martinez. The Muon spectrometer of the ALICE experiment. **Nucl. Phys.**, v. A749, p. 313–319, 2005.
- [107] G. Contin. Performance of the present ALICE Inner Tracking System and studies for the upgrade. **JINST**, v. 7, p. C06007, 2012.
- [108] J. Beringer et al. Review of Particle Physics (RPP). **Phys. Rev.**, v. D86, p. 010001, 2012.
- [109] L. Landau. On the energy loss of fast particles by ionization. **J. Phys.(USSR)**, v. 8, p. 201–205, 1944.

- [110] J. Bartke et al. Simplicity of Transverse Spectra Analysis in Terms of Transverse Energy. **Nucl. Phys.**, v. B120, p. 14–22, 1977.
- [111] Richard Witt. $p(t)$ δ systematics and $m(t)$ scaling. Em **Ultra-relativistic nucleus-nucleus collisions. Proceedings, 17th International Conference, Quark Matter 2004, Oakland, USA, January 11-17, 2004**, 2004.
- [112] Betty Bezverkhny Abelev et al. Measurement of electrons from semileptonic heavy-flavor hadron decays in pp collisions at $\sqrt{s} = 2.76$ TeV. **Phys. Rev.**, v. D91, n. 1, p. 012001, 2015.
- [113] Vladislav Balagura. Notes on van der Meer Scan for Absolute Luminosity Measurement. **Nucl. Instrum. Meth.**, v. A654, p. 634–638, 2011.
- [114] ALICE luminosity determination for pp collisions at $\sqrt{s} = 13$ TeV. v., Jun 2016.
- [115] Jaroslav Adam et al. Pseudorapidity and transverse-momentum distributions of charged particles in proton-proton collisions at $\sqrt{s} = 13$ TeV. **Phys. Lett.**, v. B753, p. 319–329, 2016.
- [116] Jan Fiete Grosse-Oetringhaus and Klaus Reygers. Charged-Particle Multiplicity in Proton-Proton Collisions. **J. Phys.**, v. G37, p. 083001, 2010.
- [117] Alberto Caliva and Thomas Peitzmann. Low-mass dielectron measurement in Pb–Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV with ALICE at the LHC. Meting van de productie van elektronparen met lage invariante massa in Pb–Pb botsingen met $\sqrt{s_{NN}} = 2.76$ TeV met ALICE bij de LHC, Mar 2017. Presented 04 Apr 2017.
- [118] Martin Volkl. Electrons from beauty-hadron decays in central Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV. 2016-10-28, PhD Thesis, Combined Faculties of the Natural Sciences and Mathematics of the Ruperto-Carola-University of Heidelberg, Germany.
- [119] Roger J. Barlow and Christine Beeston. Fitting using finite Monte Carlo samples. **Comput. Phys. Commun.**, v. 77, p. 219–228, 1993.

- [120] B. Abelev et al. Measurement of charm production at central rapidity in proton-proton collisions at $\sqrt{s} = 7$ TeV. **JHEP**, v. 01, p. 128, 2012.
- [121] v.
- [122] Shreyasi Acharya et al. Measurement of D^0 , D^+ , D^{*+} and D_s^+ production in Pb-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV. **Submitted to: JHEP**, v., 2018.
- [123] L. Adamczyk et al. Observation of D^0 Meson Nuclear Modifications in Au+Au Collisions at $\sqrt{s_{NN}} = 200$ GeV. **Phys. Rev. Lett.**, v. 113, n. 14, p. 142301, 2014.
- [124] Francesco Prino and Ralf Rapp. Open Heavy Flavor in QCD Matter and in Nuclear Collisions. **J. Phys.**, v. G43, n. 9, p. 093002, 2016.
- [125] Peter F. Kolb and Ulrich W. Heinz. Hydrodynamic description of ultrarelativistic heavy ion collisions. v., p. 634–714, 2003.
- [126] S. Voloshin and Y. Zhang. Flow study in relativistic nuclear collisions by Fourier expansion of Azimuthal particle distributions. **Z. Phys.**, v. C70, p. 665–672, 1996.
- [127] S. Batsouli, S. Kelly, M. Gyulassy, and J. L. Nagle. Does the charm flow at RHIC? **Phys. Lett.**, v. B557, p. 26–32, 2003.
- [128] M. Gyulassy, I. Vitev, and X. N. Wang. High p(T) azimuthal asymmetry in noncentral A+A at RHIC. **Phys. Rev. Lett.**, v. 86, p. 2537–2540, 2001.
- [129] Shreyasi Acharya et al. D -meson azimuthal anisotropy in midcentral Pb-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV. **Phys. Rev. Lett.**, v. 120, n. 10, p. 102301, 2018.
- [130] Xin-Nian Wang. Discovery of jet quenching and beyond. **Nucl. Phys.**, v. A750, p. 98–120, 2005.
- [131] David d’Enterria. Perturbative probes of QCD matter at the Large Hadron Collider. **PoS**, v. QNP2012, p. 018, 2012.

- [132] S. Chatrchyan et al. Indications of suppression of excited Υ states in pb-pb collisions at $\sqrt{s_{\text{NN}}} = 2.76$ TeV. v.
- [133] Vardan Khachatryan et al. Suppression of $\Upsilon(1S)$, $\Upsilon(2S)$ and $\Upsilon(3S)$ production in PbPb collisions at $\sqrt{s_{\text{NN}}} = 2.76$ TeV. **Phys. Lett.**, v. B770, p. 357–379, 2017.
- [134] H. Bethe and W. Heitler. On the Stopping of fast particles and on the creation of positive electrons. **Proc. Roy. Soc. Lond.**, v. A146, p. 83–112, 1934.