UNIVERSIDADE DE SÃO PAULO
INSTITUTO DE GEOCIÊNCIAS

**MECANISMOS DE DEFORMAÇÃO E CRONOLOGIA DE
EVENTOS NA ZONA DE CISALHAMENTO PATOS
(PROVÍNCIA BORBOREMA, NORDESTE DO BRASIL)**

Luís Gustavo Ferreira Viegas

Orientador: Prof. Dr. Carlos José Archanjo

TESE DE DOUTORAMENTO
Programa de Pós-graduação em Geoquímica e Geotectônica

São Paulo
2013

UNIVERSIDADE DE SÃO PAULO
INSTITUTO DE GEOCIÊNCIAS

Luís Gustavo Ferreira Viegas

Orientador: Prof. Dr. Carlos José Archanjo

TESE DE DOUTORAMENTO

COMISSÃO EXAMINADORA

São Paulo
2013

SUMÁRIO

AGRADECIMENTOS	4
RESUMO	7
ABSTRACT	9
CAPÍTULO 1	
INTRODUÇÃO	11
1. Apresentação	11
2. Contextualização	13
<i>2.1. O conceito de Orógenos Quentes</i>	13
<i>2.2. Influência da fusão parcial</i>	14
<i>2.3. Segregação magmática em mesoscala: migmatitos</i>	15
3. Zonas de cisalhamento e fusão parcial na Província Borborema	16
<i>3.1. O sistema de cisalhamento Patos-Seridó</i>	18
<i>3.2. Geometria e cinemática</i>	18
<i>3.3. Geocronologia</i>	21
4. Organização da tese	21
CAPÍTULO 2	
FUNDAMENTAÇÃO TEÓRICA E MÉTODOS DE ESTUDO	24
1. Introdução	24
2. Anisotropia de Suscetibilidade Magnética (ASM)	24
<i>2.1. Elementos de mineralogia magnética</i>	25
<i>2.2. Aplicações da ASM em petrologia estrutural</i>	29
<i>2.2.1. Rochas parcialmente fundidas (migmatitos)</i>	29
<i>2.2.2. Granitoides</i>	30
3. Difração de elétrons retroespalhados (EBSD)	30
<i>3.1. Características do equipamento e procedimentos analíticos</i>	32
<i>3.2. Indexação das figuras de difração</i>	33
4. Geocronologia U-Pb em zircão (SHRIMP)	35
5. Termometria de Ti em Quartzo – método TitaniQ	36

CAPÍTULO 3

ALOJAMENTO DE MAGMAS NA CROSTA INTERMEDIÁRIA PARCIALMENTE FUNDIDA E SUAS CONEXÕES COM O PLUTONISMO GRANÍTICO EM NÍVEIS CRUSTAIS SUPERIORES 38

1. Introdução 38

Timing of HT/LP transpression in the Neoproterozoic Seridó Belt (Borborema Province, Brazil): constraints from U-Pb (SHRIMP) geochronology and implications for the connections between NE Brazil and West Africa 39

CAPÍTULO 4

TRAMAS DE ROCHAS PARCIALMENTE FUNDIDAS E ALOJAMENTO DE MAGMAS EM REGIMES TRANSCORRENTES 53

1. Introdução 53

Fabrics of migmatites and the relationships between partial melting and deformation in high-grade transpressional shear zones: the Espinho Branco anatexite (Borborema Province, NE Brazil) 54

CAPÍTULO 5

CRONOLOGIA DE EVENTOS E LOCALIZAÇÃO DA DEFORMAÇÃO EM ZONAS DE CISALHAMENTO DÚCTEIS HETEROGÊNEAS 66

1. Introdução 66

Microfabrics and zircon U-Pb (SHRIMP) chronology of high- and low-temperature mylonites from the Patos shear zone (Borborema Province, NE Brazil) 67

CAPÍTULO 6

CONCLUSÕES E CONSIDERAÇÕES FINAIS 97

1. Síntese 97

2. Metamorfismo de alto grau e fusão parcial na Faixa Seridó 97

3. Tramas migmatíticas e alojamento de magmas em regimes transcorrentes 98

4. Localização da deformação e reativação na zona de cisalhamento Patos 99

5. Considerações finais 100

CAPÍTULO 7

REFERÊNCIAS

103

AGRADECIMENTOS

Esta tese é fruto de um trabalho que, sem a devida colaboração e participação de diversas pessoas e instituições, jamais teria chegado ao fim. Dessa forma, aqui expresso minha mais sincera gratidão a todos que estiveram envolvidos, direta ou indiretamente, na elaboração desta pesquisa.

Em primeiro lugar, agradeço ao Instituto de Geociências da Universidade de São Paulo (IGc-USP) pela oportunidade de realizar o doutorado em um ambiente estimulante, agradável e com possibilidade de intensas trocas científicas e interpessoais. Devo agradecer também às instituições que financiaram esta pesquisa: FAPESP (processos 2009/17537-1 e 2010/50060-1) e CNPq (processo 200496/2011-5).

Minha família merece um agradecimento especial, por representar o suporte sólido que me deu a liberdade e oportunidade de optar pela trajetória acadêmica. Agradeço ao meu pai por ser um grande conselheiro e mentor em diversas questões práticas de trabalho e de vida, e à minha querida mãe e irmão pelo constante apoio e estímulo ao longo desses anos. Muito obrigado.

Voltando à esfera acadêmica, várias foram as pessoas que participaram ao longo do desenvolvimento desse trabalho. Agradeço aos técnicos do Laboratório de Paleomagnetismo do IAG-USP, Daniele Brandt e Giovanni Moreira, pela enorme ajuda nas medidas de anisotropia magnética. Obrigado também aos técnicos Vasco Loios e Maurício Dias pelo apoio na preparação de amostras para geocronologia. Um agradecimento especial aos funcionários do IGc, Leonardo, Magali, Tadeu, Solange e Erickson, por estarem sempre disponíveis e aptos a ajudar com questões técnicas, burocráticas e financeiras.

Aos professores do IGc com os quais tive a oportunidade de trocar conhecimentos e aprender coisas novas: H. Ulbrich, Renato de Moraes, Marcos Egydio, Ginaldo Campanha, Caetano Juliani e Ricardo Trindade. Ao Prof. Ricardo agradeço pelas discussões sobre dados de ASM, bem como ao Marcos pela análise de dados de microestruturas e mecanismos de deformação.

Agradeço ao Professor Leonardo Lagoeiro por disponibilizar a infraestrutura de seu laboratório de microanálises, me receber durante a visita a Ouro Preto e auxiliar no início da aquisição dos dados de EBSD. Agradeço também aos graduandos e pós-graduandos do MicroLAB: Paola, Gilberto, Fábio e Mônica, pela valiosa ajuda e momentos de descontração na hora do café. Ainda em Ouro Preto, não posso deixar de agradecer à galera da República Kome Keto (www.komeketo.com), que me recebeu em Ouro Preto, e ao Tosado e Broz, que me ajudaram na procura de estada, nos contatos iniciais, etc. Muito obrigado a todos.

À Professora Maria Helena Hollanda, agradeço por participar ativamente na aquisição dos dados geocronológicos contidos nesta tese, esclarecer diversas dúvidas sobre geologia isotópica e, especialmente nos meses finais, pelas valiosas conversas e conselhos profissionais. Muito obrigado.

Aos colegas da Geo, Alejandro, Marta, Alba, Johanna, Donald, Angela Faleiros, Felipe Meira, Eduardo, Cláudia, Soraya, Alexandra, Andréa, Nívea, Maurício, Bruna, Marcelo, PH, Rafael, Fábio, Natali, Laura, Vicente, Angela, Eline, e tantos outros que passaram pelo instituto enquanto estive por aqui. Muito obrigado pelos momentos partilhados de descontração, conversas e risadas no dia-a-dia.

Um agradecimento especial aos dois bons amigos que fiz, John Rico e Camilo Bustamante, pela amizade, companheirismo, parceria nos trabalhos de campo e pelas diversas conversas geológicas. Muito obrigado.

Uma parte importante desta tese foi realizada no exterior e, dessa forma, tive contato com um mundo novo onde as trocas que fiz foram extremamente enriquecedoras, tanto no plano profissional quanto pessoal.

Alors, pour commencer, je veux remercier a Alain Vauchez et Andrea Tommasi, mes directeurs de thèse au l'étranger, pour toute l'aide avec le déménagement, les premiers jours à Montpellier, les diners d'arrivée et départ, la langue, etc. Merci aussi pour être toujours disponible avec les questions concernant la vie en France.

Je tiens aussi que remercier tous les gents au sein du laboratoire Géosciences Montpellier, qu'ont m'aidée avec tout les choses, depuis la thèse, les trucs administratifs, et

même les astuces pour se balader en ville, etc. Je veux dire un grand merci à Bernadette, Françoise, Dave Mainprice, Christel, Benoit, Diane, Olivier, Alain Chauvet, Anne Delplanque, Sylvie, Fabrice Grosbeau, Josianne Tack, Dominique Arnau, Jose Atienza, Christophe Nevado, Doriane et Fabrice Barou. Merci par tout.

Sur le plan personnel, mon expérience n'aurait pas pu être plus riche. J'ai croisé avec des gents super sympas, que seront toujours dans mon cœur, et que j'oublierais jamais. Mes cher(e)s Virginie, Ana Ruiz, Kate, Steve, Roberto, Mikael, Erwin, Antoine, Yannick, Fatna, Stéphanie, Julie, Carole, Greg Fanjat, Ousmane, Greg Ballas, Théo, Marcus, Dino, Sergio, Camile, Damien, Jeremy et Shantanu, je vous remercie vivement pour l'hospitalité, bon humeur et amitié pendant mon année en France. C'était super !

De volta ao Brasil, agradeço aos amigos antigos, sempre presentes, sem os quais as coisas teriam sido certamente menos interessantes. Obrigado à amiga Geane Carolina Cavalcante, por partilhar de um tema de tese similar, pelas trocas científicas, pela ajuda em diversos momentos e pela descontração no ambiente de trabalho. E ao amigo Mauro Barreto, incansável parceiro de cervejas nos finais de semana. Muito obrigado.

Por fim, o último agradecimento vai para o Professor, Orientador e, ao cabo desses três anos e meio de colaboração, bom amigo Carlos Archanjo. Agradeço por ter aceitado a supervisão deste trabalho, ter me oferecido a oportunidade de trabalhar com um assunto interessante em uma área de trabalho mais interessante ainda (Província Borborema), pela constante disponibilidade, desde os contatos iniciais até os momentos finais, pela enorme paciência durante a progressão de minha (nada rápida) curva de aprendizado nos métodos magnéticos, pelos conselhos acadêmicos, pelas discussões geológicas, e por todo o apoio prestado durante o desenvolvimento deste trabalho. Porém, este espaço final é reservado para que eu possa expressar minha honesta e sincera gratidão pela liberdade com a qual pude desenvolver esta tese, pelo constante estímulo que recebi para expressar (e formar) minha filosofia de pensamento geológico, individualizar minha linha de pesquisa e estabelecer meu método de trabalho, e pelo exemplo que tive de como se fazer ciência pela ciência em si. Por tudo isso, Archanjo, muito obrigado.

VIEGAS, L.G.F. **Mecanismos de deformação e cronologia de eventos na zona de cisalhamento Patos (Província Borborema, NE Brasil)**. Tese (doutorado) – Instituto de Geociências, Universidade de São Paulo, São Paulo, 2013.

A zona de cisalhamento Patos consiste em uma mega-estrutura transcorrente E-W que deformou rochas pré-cambrianas durante a orogênese brasileira. As relações entre fusão parcial e deformação, os mecanismos de deformação envolvidos na nucleação da estrutura e a evolução termocronológica da milonitização foram investigados por meio de trabalhos de campo e técnicas multidisciplinares tais como a anisotropia de suscetibilidade magnética (ASM), geocronologia (U-Pb SHRIMP) e orientações preferenciais cristalográficas medidas através de difração de elétrons retroespalhados (EBSD). O domo de Santa Luzia, situado na conexão da zona de cisalhamento com a Faixa Seridó, caracteriza-se pela trama magnética levemente oblíqua à elongação NE do corpo. Idades U-Pb obtidas na borda recristalizada de zircões do migmatito forneceram idades de 575 ± 3.4 Ma. Trama magnética e idade semelhantes foram encontradas no granito Acari, sugerindo a contemporaneidade entre o alojamento dos magmas e a deformação cisalhante. No anatexito Espinho Branco, geometrias complexas observadas em campo não são reproduzidas na trama de ASM, a qual concorda com a cinemática da zona de cisalhamento Patos. Entretanto, sítios com suscetibilidades baixas (< 0.5 mSI) exibem dispersões na trama magnética, as quais contrastam com orientações cristalográficas de biotita e dados de campo. Contudo, a anisotropia de remanência anisterética mantém-se coerente com o campo de deformação a que foi submetido o anatexito, sugerindo que a dispersão das direções principais de ASM relaciona-se à fases hidrotermais tardias que desorganizam a trama magnética. Ortognaisses miloníticos situados na borda norte da zona de cisalhamento são marcados por migração de bordas dos grãos, fraturamento intracristalino e mirmequitização periférica. As tramas de eixos [0001] de quartzo exibem máximos em Y nos milonitos de alto-grau, sugerindo ativação do sistema de deslizamento prismático-*a*. As microestruturas gradam para o estado magmático no contato com o anatexito, sendo comuns tramas randômicas de eixo-*c* de quartzo. Zircões extraídos de leucossomas fornecem uma idade de 566 ± 5 Ma, atribuídos à cristalização do magma. A margem sul da estrutura exhibe uma progressiva redução do tamanho dos grãos e geração de milonitos finos a ultramilonitos sem a presença de fusão parcial. Zircões extraídos de um *augen* granítico forneceram uma idade de intercepto inferior de 545 ± 14 Ma, atribuída ao metamorfismo. Os milonitos da borda sul registram tramas de eixo-*c* de quartzo com

máximos entre Z e Y, sugerindo ativação dos sistemas romboédrico- e basal-*a*. Nos feldspatos, os principais sistemas de deslizamento são (010)[001] e (010)[100] em todos os litotipos, com exceção do sistema (100)[010], observado na transição entre migmatitos e os milonitos de baixo grau. Estimativas de temperatura por meio do método TitaniQ registram temperaturas médias de ~ 490°C nos milonitos da borda sul. O conjunto de resultados permite concluir que: i) a ASM reflete os incrementos finais da deformação regional que afetou os migmatitos, ii) a fusão parcial associada ao cisalhamento dextral do Patos ocorreu em c. 565 Ma, enquanto a milonitização de médio a baixo grau, datada em 545 Ma, caracteriza o retrabalhamento metamórfico ocasionado pela localização da deformação em regime dúctil-rúptil; iii) este retrabalhamento é correlacionável a outras zonas de cisalhamento da Província Borborema, constituindo um evento de caráter regional possivelmente relacionado às colisões tardias na margem oeste do Gondwana e, iv) a zona de cisalhamento Patos foi formada após a colisão dos crátons Oeste Africano e Congo-São Francisco ao escudo da Borborema, acomodando deslocamentos transcorrentes a partir de descontinuidades crustais formadas em estágio pré-colisional.

Palavras-chave: Tectonofísica, migmatito, magmatismo, geologia estrutural, geocronologia, reologia.

VIEGAS, L.G.F. **Deformation mechanisms and chronology of events in the Patos shear zone (Borborema Province, NE Brazil).** Tese (doutorado) – Instituto de Geociências, Universidade de São Paulo, São Paulo, 2013.

The Patos shear zone is an E-W transcurrent structure that deformed Precambrian rocks from the Borborema Province during the Neoproterozoic. The relationships between partial melting and deformation, the nucleation of the shear zone and the thermochronological evolution of mylonites were investigated by means of detailed field work coupled with Anisotropy of Magnetic Susceptibility (AMS), U-Pb SHRIMP geochronology and crystallographic fabrics measured through electron backscatter diffraction (EBSD). In the Santa Luzia dome, located within the Seridó Belt, the magnetic lineations are slightly oblique to the dome NE elongation. Zircon recrystallized rims yield U-Pb ages of 575 ± 3.4 Ma. These data are consistent with ages and magnetic fabrics found in the Acari granite, suggesting that magma emplacement and shearing were coeval in the Seridó belt. In the Espinho Branco anatexite, emplaced within the mylonites of the Patos shear zone, complex field leucosome geometries are not reproduced in the magnetic fabric, which is consistent with the kinematics of the shear zone. However, low susceptibilities (< 0.5 mSI) display a scattering of magnetic orientations and do not correlate with biotite lattice fabrics and field foliations. Nevertheless, anisotropy of anhysteretic remanence fabrics remain parallel to the external strain field, suggesting that AMS dispersions may be related to late percolation of hydrothermal fluids. Mylonitic orthogneisses from the northern border of the Patos shear zone are marked by grain boundary migration, intracrystalline fracturing and myrmekitization at the periphery of grains. Quartz [0001] lattice fabrics form maxima mainly on Y, suggesting activation of prism-*a* slip systems. Towards the contact with migmatites, the microstructures progressively change from solid-state to magmatic, characterized by widespread interstitial quartz and random orientations of quartz *c*-axis fabrics. Zircons from the leucosomes yield an age of 566 ± 5 Ma, which is attributed to magma crystallization. The southern border of the Patos shear zone is marked by progressive grain size reduction and formation of fine-grained mylonites to ultramylonites without traces of partial melting. An age of 545 ± 14 Ma, obtained in zircons from an ultramylonitic *augen* granite, constrains the timing of the low-grade metamorphism. These Southern mylonites show quartz [0001] fabrics with maxima spreading between Z and Y, suggesting activation of prism and basal slip systems. In feldspars, the main activated slip systems are (010)[001] and (010)[100] in

all rock types, except for the (100)[010] slip system, observed in the transition from migmatites to low-grade mylonites. TitaniQ temperature estimates record mean temperatures of $\sim 490^{\circ}\text{C}$ for the southern mylonites. These data allows us to conclude that: i) AMS reflects the final strain increments recorded in migmatites during regional deformation, ii) partial melting and dextral shearing in the Patos shear zone occurred at ~ 565 Ma, while low-grade mylonitisation dated at 545 Ma defines a late reworking at ductile-brittle conditions, iii) this reworking event is observed in other areas of the Borborema Province, defining a regional event possibly associated with late collisions in the western margin of Gondwana, iv) the Patos shear zone was nucleated after the collision between the Congo-São Francisco and West-African Cratons, through accommodation of strike-slip displacements localized in crustal discontinuities previously formed in an early pre-collisional stage.

Keywords: Tectonophysics, migmatite, magmatism, structural geology, geochronology, rheology.

CAPÍTULO 1

INTRODUÇÃO

1. Apresentação

As relações entre magmatismo e deformação em zonas de cisalhamento litosféricas têm sido alvo de diversos estudos há mais de três décadas (Ramsay & Graham 1970; Nicolas et al. 1972, 1977; Atherton 1993; Pitcher 1993). Estudos experimentais demonstram que, em condições de deformação próximas da fácies anfibolito alto, quantidades de líquido magmático (*melt*) em torno de 7% em volume são suficientes para causar uma mudança drástica nas propriedades reológicas dos materiais rochosos, promovendo localização da deformação nos domínios magmáticos (Arzi 1978; Van der Molen & Paterson 1979; Rosenberg & Handy 2005). Por esta razão, diversos autores consideram geologicamente viável uma relação de retroalimentação entre fusão parcial e deformação; o calor advindo da fusão facilita a deformação de alta temperatura, enquanto que estruturas regionais podem servir de conduto para migração e transporte de magmas ao longo da crosta intermediária em direção a níveis crustais superiores (Rutter & Neumann 1995; Brown & Rushmer 1997; Brown 2001).

As características reológicas do magma são influenciadas pela profundidade de alojamento e, conseqüentemente, as interações entre magmas e falhas regionais são afetadas (ou não) pela presença de zonas de cisalhamento em determinadas profundidades (Schofield & D'Lemos 1998). Magmas que interagem com zonas de cisalhamento em níveis crustais intermediários, por exemplo, tendem a desenvolver tramas influenciadas pelo campo deformacional externo em decorrência do longo tempo de cristalização na presença de gradientes geotérmicos elevados (Paterson et al. 1998; Brown & Solar 1998). Alternativamente, magmas que cristalizam em níveis crustais superiores exibem maior probabilidade de preservar as tramas magmáticas relacionadas à forças de corpo (ascensionais, convectivas, etc.; Weinberg et al. 2004).

Terrenos migmatíticos são caracterizados pela associação frequente entre anatexitos-granitos-zonas de cisalhamento em contextos geológicos diversos e diferentes intervalos de tempo (Whitney et al. 2004). Veios e/ou mobilizados de material granítico constituem o produto da fusão parcial e são caracterizados como *leucossomas* (Sawyer 2008). A geometria

dos leucossomas e sua relação com as rochas encaixantes fornece informações importantes sobre os mecanismos de migração e transporte do material magmático na crosta intermediária, fazendo dos terrenos migmatíticos um registro de parte do caminho de migração do magma desde a área-fonte até o local de alojamento (Brown & Rushmer 1997). Entretanto, fatores como deformação, taxa de fusão e diferenças reológicas entre anatexitos e encaixantes podem resultar em distribuições heterogêneas do material parcialmente fundido ao longo destes terrenos, caracterizando geometrias complexas em escala de afloramento. Nestes casos, estudos baseados em análise estrutural clássica podem levar a interpretações ambíguas e/ou não-conclusivas.

No nordeste do Brasil, zonas de cisalhamento são a principal expressão da orogênese brasileira/Pan-africana na Província Borborema (Fig. 1). Estas zonas estão geralmente associadas a uma intensa atividade magmática expressa na forma de plútons graníticos, os quais têm sido alvo de diversos estudos cartográficos-estruturais e geocronológicos com o objetivo de elucidar seus mecanismos de alojamento e a relação com a tectônica regional (Brito Neves et al. 2003). O sistema de cisalhamento Patos-Seridó (Fig. 1), uma das principais estruturas presentes na referida província, caracteriza-se como uma zona mecanicamente conectada onde deslocamentos transcorrentes são transferidos para uma faixa metassedimentar deformada em regime compressivo. As condições metamórficas nos metapelitos da faixa dobrada são de alta temperatura-baixa pressão, com generalizada fusão parcial em gnaisses do embasamento (Corsini et al. 1991).

Milonitos de alta temperatura associados com leucossomas graníticos podem ser observados na borda norte do setor central da zona de cisalhamento Patos, gradando progressivamente para migmatitos na porção interna da estrutura. Os migmatitos exibem percentagens variadas de fusão parcial, podendo ser seguidos desde a zona de cisalhamento até o interior da faixa metassedimentar do Seridó. Embora concordantes com respeito à orientação da trama tectônica regional, as relações entre migmatitos, milonitos e a deformação do sistema Patos-Seridó são ainda pouco conhecidas.

As interações entre anatexitos e a deformação regional em níveis crustais mais profundos podem, por vezes, exercer um controle importante sobre as intrusões mais rasas, particularmente por meio de mecanismos de migração e segregação do magma ao longo de sua trajetória de ascensão, a qual pode ser facilitada pela presença de zonas de cisalhamento. Dessa forma, o estudo de rochas parcialmente fundidas constitui um aspecto relevante para o entendimento das características reológicas da crosta durante eventos orogênicos, além de fornecer informações sobre os processos de transferência de magmas e a formação de

instabilidades crustais marcadas pela acumulação da deformação em zonas de cisalhamento dúcteis.

Esta tese está incluída dentro desta temática e aborda, sob o ponto de vista da geologia estrutural, as relações entre a deformação dúctil e os produtos da fusão parcial, *i.e.*, migmatitos, no interior de zonas de cisalhamento. Por meio de uma abordagem integrada envolvendo técnicas multidisciplinares, pretende-se contribuir para o entendimento dos processos responsáveis pelo fluxo de material crustal em níveis intermediários da crosta durante diferentes estágios evolutivos de um ciclo orogênico (Whitney et al. 2004). Estes estudos auxiliarão na caracterização dos mecanismos de alojamento de rochas parcialmente fundidas, na correlação entre magmatismo e deformação em diferentes níveis crustais, na evolução estrutural de uma zona de cisalhamento gerada na presença de magma, e no estabelecimento de uma cronologia absoluta de eventos magmáticos e deformacionais no contexto da orogênese brasileira.

Fig. 1: Principais zonas de cisalhamento e granitos neoproterozóicos da porção central da Província Borborema.

2. Contextualização

2.1. O conceito de Orógenos Quentes

Temperaturas altas associadas à presença de grandes volumes de magma em segmentos crustais foram primeiramente descritas em terrenos arqueanos (Bouhallier et al. 1995). As anomalias termais e estilos particulares de deformação observados nestes terrenos foram atribuídos ao elevado fluxo de calor registrado no Arqueano. Entretanto, estudos

recentes revelaram que a crosta intermediária abaixo dos Himalaias e da Cordilheira Central dos Andes registra temperaturas elevadas e encontra-se parcialmente fundida (Fig. 2; Nelson et al. 1996). Com base em dados geofísicos, foram detectadas “camadas” de crosta fundida com geotermas que ultrapassam 30°C/km em profundidades relativamente rasas (20-40 km; Unsworth et al. 2005). Estas evidências levaram a um renovado interesse no estudo da crosta intermediária presente nestes *orógenos quentes*.

Embora recentemente caracterizados como tal, orógenos quentes têm sido identificados em diversas partes do mundo ao longo do tempo geológico, englobando o Pré-Cambriano (Fritz et al. 2009), Neoproterozóico (Vauchez et al. 2007) e Fanerozóico (Chardon et al. 2009). Em todos os casos, as principais características são a presença de um gradiente geotérmico elevado e grandes volumes de material magmático alojado na forma de batólitos graníticos e/ou corpos de anatexitos.

Fig. 2: Seção interpretativa da zona de colisão himalaiana evidenciando a natureza parcialmente fundida da crosta intermediária. MFT: Main Frontal Thrust; MCT: Main Central Thrust; STD: South Tibetan Detachment. Adaptado de Nelson et al. 1996.

O padrão deformacional é marcado por uma complexa repartição de *strain* em domínios de localização da deformação, como zonas de cisalhamento, e domínios onde a deformação é distribuída de forma heterogênea, envolvendo componentes gravitacionais e convergentes. A diferenciação entre esses dois padrões se dá a partir da taxa de fusão parcial envolvida na deformação.

2.2. Influência da fusão parcial

Evidências experimentais indicam que a presença de magma reduz de forma drástica a resistência reológica da crosta durante eventos orogênicos. Pequenas quantidades de líquido magmático (~ 7% em volume) são suficientes para causar uma mudança brusca na resistência

das rochas e favorecer a localização da deformação em zonas de cisalhamento onde mecanismos de alta temperatura predominam (Fig. 3; Rosenberg & Handy 2005; Rosenberg & Handy 2000). A fusão parcial tem início nos contatos entre grãos, com a geração de filmes de magma que podem ou não conectar-se entre si, em função da distribuição de ângulos diedrais entre os cristais (Sawyer 2001): ângulos diedrais menores do que 30° favorecem um escape do líquido em decorrência da pressão intragranular na região entre os grãos (Rosenberg 2001). A progressão da fusão leva a uma atividade continuada de escape do líquido e subsequente coleta do mesmo em uma rede interconectada de filmes de magma que atuam como condutos de transporte e posterior migração do material parcialmente fundido. Por outro lado, grandes quantidades de fusão promovem uma localização ineficiente da deformação e resultam em padrões heterogêneos de migração do material magmático, o qual geralmente flui a partir de fluxo lateral (Whitney et al. 2004). Nesses contextos, a distribuição da deformação e o comportamento reológico da crosta são complexos (Vanderhaege 2009).

Fig. 3: Diminuição da resistência da rocha e localização da deformação na presença de fusão parcial (modificado de Rosenberg et al. 2006; Rosenberg & Handy 2000).

2.3. Segregação magmática em mesoscala: migmatitos

Os terrenos migmatíticos são fragmentos preservados da crosta intermediária parcialmente fundida presentes em orógenos antigos e jovens (Mehnert 1968; Sawyer 2008). Assinaturas geoquímicas similares em anateixitos e granitos da Província Superior, no Canadá, sugerem que rochas parcialmente fundidas representam estágios transicionais da

migração e transporte do magma ao longo da crosta intermediária até o alojamento em níveis crustais superiores (Sawyer 1998).

Leucossomas orientados paralelamente à foliação da rocha, sem exibir ruptura da trama planar e com estruturas pré-fusão preservadas definem *metatexitos*. Com o aumento da fusão, a morfologia torna-se difusa, a foliação prévia é gradativamente desagregada e a rocha evolui para se tornar mais homogênea e isotrópica, caracterizando *diatexitos* (Sawyer 2008). O produto final da fusão parcial são *leucogranitos*, frequentemente observados em associação com diatexitos em zonas de cisalhamento de alto grau (Sawyer 2001).

A geometria do leucossoma auxilia na caracterização dos caminhos de migração do magma e, com base em estudos estruturais detalhados, pode-se deduzir os mecanismos de segregação e transporte em zonas deformadas (Blumenfeld & Bouchez 1992; Marchildon & Brown 2003). Porém, a natureza heterogênea da fusão parcial em escala de afloramento pode resultar em morfologias complexas, levando muitas vezes à interpretações ambíguas e/ou não-conclusivas acerca do padrão estrutural em anatexitos.

Nesse sentido, o uso de técnicas de cartografia estrutural de detalhe, como a Anisotropia de Susceptibilidade Magnética (ASM), constitui uma ferramenta útil na caracterização da trama de alojamento de rochas parcialmente fundidas em zonas de cisalhamento. A importância da ASM na análise estrutural de granitos é hoje amplamente reconhecida (Bouchez 1997) e, aplicações mais recentes da técnica em migmatitos forneceram resultados satisfatórios no estudo dos mecanismos de fluxo destas rochas (Ferre et al. 2003).

Esta tese tem como um dos objetivos a caracterização estrutural da trama interna de migmatitos alojados no interior de estruturas dúcteis de alto grau no nordeste do Brasil. Um breve esboço petroestrutural das principais zonas de cisalhamento observadas na região é apresentado a seguir.

3. Zonas de cisalhamento e fusão parcial na Província Borborema

A feição marcante da orogênese Brasileira/Pan-Africana (~ 600 Ma) na Província Borborema consiste em uma rede de zonas de cisalhamento dúcteis associadas a um expressivo plutonismo granítico (Fig. 4; Vauchez et al. 1995). As zonas de cisalhamento Pernambuco e Patos, de direção E-W, representam duas descontinuidades importantes no interior da Província, deformando em cinemática dextral rochas do embasamento, cobertura metassedimentar e plutons graníticos sin- a pós-cinemáticos (Van Schmus et al. 2008).

Rochas parcialmente fundidas estão presentes em ambas as zonas de cisalhamento, em contato com milonitos de alto e baixo grau metamórfico. Os leucossomas podem apresentar morfologias distintas em decorrência de diferentes percentagens de fusão parcial. A continuidade estrutural de leucossomas paralelos à foliação é frequentemente rompida por acumulações métricas de mobilizados quartzo-feldspáticos, caracterizando diatexitos, ou intrusão de *stocks* graníticos por vezes truncando a foliação milonítica.

Fig. 4: Província Borborema, NE Brasil, destacando as principais zonas de cisalhamento (em vermelho).

ZCPO: zona de cisalhamento Pernambuco Oeste; ZCPE: zona de cisalhamento Pernambuco Este.

A zona de cisalhamento Patos, área-alvo deste estudo, estende-se por cerca de 600 km na direção E-W até se conectar com a Faixa Seridó em sua porção central, e tem sido interpretada como uma ramificação de falhas de direção NE observadas na porção ocidental da Província (Corsini et al. 1991; 1996). A região de conexão entre a zona de cisalhamento Patos e a Faixa Seridó exibe uma rotação gradual das tramas tectônicas, em concordância com o eixo de alongação principal da faixa metassedimentar (NE). Esta geometria arqueada é interpretada como um sistema transpressivo conjugado onde deslocamentos transcorrentes são transferidos para uma faixa compressional através de rotação progressiva do eixo de estiramento principal da zona (Corsini et al. 1991).

Acumulações de material parcialmente fundido são observadas em duas localidades principais: i) na área de conexão com a Faixa Seridó, onde ocorre o diatexito de Santa Luzia

(Capítulo 3); e ii) no interior da zona de cisalhamento Patos, na forma do corpo sigmoide denominado migmatito Espinho Branco (Capítulo 4). Os leucossomas exibem morfologias diversas e estruturas de natureza magmática; estruturas decorrentes de deformação no estado-sólido ocorrem de forma localizada. Estas duas ocorrências estão restritas à borda norte da zona de cisalhamento, em contato com milonitos de alta temperatura (Fig. 5).

A margem sul da estrutura é caracterizada por uma faixa de milonitos finos a ultramilonitos, onde evidências de fusão parcial são ausentes (Fig. 6). Estes milonitos desenvolveram-se principalmente a partir de ortognaisses do embasamento e granitos sintectônicos, e suas tramas diferem daquelas observadas nos milonitos de alto grau. Um estudo estrutural detalhado nestes milonitos está apresentado no Capítulo 5.

3.1. O sistema de cisalhamento Patos-Seridó

O Sistema de cisalhamento Patos-Seridó localiza-se no domínio centro-leste da Província Borborema (Fig. 1), onde a zona de cisalhamento transcorrente de Patos representa um limite crustal dividindo duas unidades geotectônicas: o domínio Jaguaribe-Seridó, a norte, e o domínio conhecido como Zona Transversal, a sul (Van Schmus et al. 2008).

A terminação oeste da estrutura conecta-se com zonas de cisalhamento NE-SW formando um duplex no qual lentes de rochas do embasamento alternam-se com metassedimentos e plutons granitoides (Corsini et al. 1996). Em sua terminação leste, a zona de cisalhamento encurva-se para NE até desaparecer sob os sedimentos da costa atlântica. Milonitos de médio a baixo-grau metamórfico são as principais rochas observadas na terminação leste (Archanjo & Fetter 2004; Rodrigues 2008).

3.2. Geometria e cinemática

Os padrões geométricos e cinemáticos ao longo do sistema Patos-Seridó são heterogêneos: de oeste para leste, a zona de cisalhamento passa gradualmente de uma estrutura em duplex para uma zona de curvatura e conexão com a Faixa Seridó. Na sua porção central, as tramas planares são subverticais e estão associadas a uma lineação de estiramento sub-horizontal, sugerindo deslocamentos transcorrentes ao longo da estrutura.

Fig. 5: Desagregação da estrutura planar em decorrência do aumento gradual da quantidade de fusão; diatexito de Santa Luzia (foto superior) e anatexito Espinho Branco (foto inferior).

Fig. 6: Milonitos de alta temperatura (esquerda) associados à fusão parcial e milonitos de média- a baixa temperatura (direita) na zona de cisalhamento Patos.

Uma forte componente de estiramento E-W é observada em seixos conglomeráticos estirados, agregados quartzo-feldspáticos, plútons graníticos alongados e foliações *boudinadas* em migmatitos e milonitos (Corsini et al. 1991). As microestruturas e os indicadores cinemáticos mostram, de forma sistemática, predominância do sentido de cisalhamento dextral em toda a zona.

As principais unidades litológicas observadas no interior da estrutura são (Vauchez et al. 1995; Corsini et al. 1998; Costa 2002): i) um embasamento gnáissico-migmatítico composto por ortognaisses migmatíticos bandados, migmatitos e leucogranitos; ii) metassedimentos de médio- a baixo-grau, compreendendo micaxistos, quartzitos, paragnaisses, anfibolitos, conglomerados e, localmente, mármore; iii) plútons graníticos porfiríticos de composição monzogranítica a sienogranítica. Estas rochas estão milonitizadas em intensidades variáveis e a deformação é comumente associada a processos de fusão parcial (Corsini et al. 1991).

As porções central e leste da zona de cisalhamento Patos são compostas por faixas miloníticas de alta temperatura exibindo uma foliação milonítica subvertical e uma lineação de estiramento sub-horizontal. As microestruturas observadas são principalmente de estado-sólido, com evidências de deformação em fácies anfibolito (Corsini et al. 1991; Vauchez et al. 1995). No setor de conexão entre a zona de cisalhamento Patos e a Faixa Seridó, migmatitos estromáticos ocorrem intercalados com ortognaisses miloníticos. Nas proximidades da cidade de Patos, os diatexitos e leucogranitos formam um corpo alongado de cerca de 25 km de comprimento por 5-8 km de largura, ligeiramente paralelo ao eixo maior da zona de cisalhamento. Este corpo é denominado anatexito Espinho Branco (Capítulo 4; Viegas et al. 2013). Na junção com a Faixa Seridó, os migmatitos se localizam no domo anatético de Santa Luzia (Capítulo 3; Archanjo et al. 2013), o qual exibe uma geometria elíptica e alongada na direção NE-SW, concordante com o eixo de alongamento principal da faixa metassedimentar.

A borda sul da zona de cisalhamento Patos é marcada por um corredor estreito (~ 5 km de largura) de milonitos finos a ultramilonitos deformados em fácies xisto-verde, e sem evidências de fusão parcial (Fig. 6). Estas rochas exibem porfiroclastos de quartzo e feldspatos imersos em uma matriz félsica fina, além de assembleias minerais de baixo grau (clorita, muscovita, epidoto).

3.3. Geocronologia

Os principais dados geocronológicos disponíveis na zona de cisalhamento Patos consistem em termocronologia $^{40}\text{Ar}/^{39}\text{Ar}$, idades de cristalização U/Pb e idades modelo (TDM) Sm/Nd. Corsini et al. (1998) conduziram um estudo termocronológico em cristais de biotita, muscovita e anfibólio de amostras localizadas principalmente na porção central da zona de cisalhamento, além de algumas amostras na zona de conexão com o Seridó. Esses autores obtiveram, para a zona de cisalhamento Patos, temperaturas da ordem de 750 – 800°C e pressões de ~ 0,4 – 0,5 MPa, enquanto que a Faixa Seridó registra temperaturas entre 550 – 600°C. Este metamorfismo de alta temperatura é atribuído ao alojamento de magmas máficos derivados do manto e intrusão de plútons graníticos (Corsini et al. 1998). As taxas de exumação calculadas são baixas (0,1 – 0,15 mm/Ma), assumindo o pico metamórfico em 580 Ma com a colocação de granitos e soerguimento em 500 Ma. Idades similares de resfriamento em grãos de muscovita e biotita, em torno de 500 Ma, sugerem rápidas taxas de soerguimento e subsequente resfriamento (Corsini et al. 1998).

Costa (2002) realizou um estudo detalhado de idades de cristalização a partir do método U/Pb (TIMS), além do cálculo de idades modelo Sm/Nd, em gnaisses e leucossomas localizados nas porções central e leste da zona de cisalhamento Patos. As principais idades modelo (TDM) obtidas variam entre 3.4-2.6 Ga em rocha total, enquanto que os zircões forneceram idades U/Pb no intervalo 2.2-2.0 Ga. Estas idades são consideradas por Costa (2002) como sendo idades de um protólito arqueano-paleoproterozóico, posteriormente retrabalhado durante os eventos da orogênese brasileira. Ainda de acordo com Costa (2002), a zona de cisalhamento Patos teria atuado como uma zona transpressiva de escape tectônico lateral e vertical nos estágios finais da orogênese brasileira, sendo portanto responsável pelo deslocamento de dois domínios diferentes; o domínio Jaguaribe-Seridó, a norte, e o domínio da Zona Transversal ao sul.

4. Organização da tese

A abordagem da temática acima exposta é apresentada sob a forma de artigos científicos, os quais foram publicados e submetidos a periódicos internacionais. Dessa forma, cada capítulo pode ser lido de forma relativamente independente dos demais e, ao final da

tese, as principais informações estão reunidas em um capítulo integrador, o qual discute as conclusões finais alcançadas com a realização do presente trabalho.

O Capítulo 2 contém uma descrição sucinta dos principais métodos utilizados na investigação das relações entre magmatismo e deformação em zonas de cisalhamento dúcteis. Sínteses teóricas sobre anisotropia magnética (ASM), medidas de orientações preferenciais cristalográficas (OPCs) e geocronologia U-Pb em zircão (SHRIMP) estão contidas neste capítulo.

O Capítulo 3 apresenta o artigo científico intitulado “*Timing of HT/LP transpression in the Neoproterozoic Seridó Belt (Borborema Province, Brazil): constraints from U-Pb (SHRIMP) geochronology and implications for the connections between NE Brazil and West Africa*”, publicado no periódico *Gondwana Research* (Archanjo et al. 2013). Com base em estudos de campo, cartografia estrutural da trama magnética (ASM), e determinações isotópicas U-Pb (SHRIMP) em migmatitos e granitos alojados na faixa metassedimentar do Seridó, esta contribuição investiga os mecanismos de alojamento do material parcialmente fundido na crosta intermediária e suas relações com granitos alojados em níveis crustais mais rasos. As informações obtidas permitem tecer correlações entre as contrapartes continentais Americana e Africana dentro do contexto do Gondwana Ocidental.

Os migmatitos alojados no interior da zona de cisalhamento Patos foram alvo de um estudo detalhado de petrologia estrutural, o qual consiste no artigo científico apresentado no Capítulo 4, intitulado “*Fabrics of migmatites and the relationships between partial melting and deformation in high-grade transpressional shear zones: the Espinho Branco anatexite (Borborema Province, NE Brazil)*”. Este manuscrito, publicado no *Journal of Structural Geology* (Viegas et al. 2013), compreende uma análise da influência da deformação cisalhante no desenvolvimento da trama magmática no migmatito Espinho Branco.

A combinação de dados de campo, anisotropia de suscetibilidade e remanência (ASM – ARA) e medidas de orientações preferenciais cristalográficas por meio da técnica de difração de elétrons retroespalhados (EBSD) permite concluir que a geometria complexa de rochas parcialmente fundidas, frequentemente observada na escala de campo, não se reproduz nos dados da trama magnética. Entretanto, a heterogeneidade da fusão parcial pode resultar em dados de ASM por vezes discordantes do padrão global da rocha. Dessa forma, a aplicação de técnicas complementares, como a remanência magnética e análises de OPCs, se faz necessária a fim de se obter dados consistentes sobre os mecanismos de fluxo crustal na crosta intermediária parcialmente fundida.

A informação obtida com a cartografia estrutural dos anatexitos indica que os processos de migração e alojamento de magmas são controlados pela deformação cisalhante. Tendo em vista estas características, um estudo termo-tectônico e isotópico foi realizado na zona de cisalhamento Patos, com o objetivo de investigar seus mecanismos de deformação e sua influência no transporte de magmas, bem como a cronologia do processo magmático-deformacional. Estes dados estão reunidos no Capítulo 5, o qual compreende o artigo científico intitulado “*Microfabrics and zircon U-Pb (SHRIMP) chronology of high- and low-temperature mylonites from the Patos shear zone (Borborema Province, NE Brazil)*”, submetido ao periódico *Precambrian Research* (Viegas et al., submetido).

O Capítulo 6 consiste em uma síntese dos resultados e considerações finais.

CAPÍTULO 2

FUNDAMENTAÇÃO TEÓRICA E MÉTODOS DE ESTUDO

1. Introdução

O desenvolvimento da presente tese compreendeu uma abordagem integrada da geologia estrutural de rochas parcialmente fundidas e a deformação em zonas de cisalhamento, a partir do uso de diversas técnicas, tais como a anisotropia magnética (ASM-ARA), difração de elétrons retroespalhados (EBSD) e datações isotópicas (U-Pb SHRIMP). Os princípios teóricos dos métodos de trabalho aqui utilizados estão descritos de forma sucinta neste capítulo.

2. Anisotropia de Suscetibilidade Magnética (ASM)

Um dado material, ao sofrer influência de um campo magnético indutor (H), adquire uma magnetização (M) que pode ser descrita a partir da seguinte relação:

$$M = kH$$

onde k representa a suscetibilidade magnética. Como as unidades de M e H são definidas em A/m (SI), k é adimensional. Em materiais isotrópicos, M é constante e paralelo à orientação do campo indutor (H). Entretanto, os minerais são naturalmente anisotrópicos, de forma que M e H não são coaxiais. Assim, para uma determinada direção de indução, as componentes de k podem ser representadas a partir da seguinte equação:

$$M_i = k_{ij}H_j \quad (i, j = 1, 2, 3)$$

onde M_i é a magnetização na direção i e H_j é o campo indutor na direção j . k_{ij} é uma matriz simétrica de segunda ordem, chamada tensor de suscetibilidade (Nye 1957). A diagonalização da matriz permite encontrar as magnitudes e direções principais do tensor ($k_1 > k_2 > k_3$). A representação espacial do tensor suscetibilidade pode ser descrita por um elipsoide, de onde podem ser extraídas informações sobre o grau de anisotropia (parâmetro P

$= k_1/k_3$), e forma ou simetria do elipsoide que varia de oblato ($k_1 = k_2 > k_3$), prolato ($k_1 > k_2 = k_3$) a triaxial ou neutro ($k_1 > k_2 > k_3$). Nesse trabalho a simetria do elipsoide foi determinada pelo parâmetro T :

$$T = \left[\frac{2(\ln k_2 - \ln k_3)}{\ln k_1 - \ln k_3} \right] - 1$$

onde a variação de T entre 0 e 1 define elipsoides neutros a oblatos e de 0 a -1 os neutros ou prolatos.

Uma propriedade magnética particular observada nos minerais ferromagnéticos é a magnetização remanente (Fuller 1963; Daly 1967; Jackson 1991), a qual pode ser intrínseca ao material analisado (magnetização remanente natural, MRN), ou induzida a partir de experimentos laboratoriais, como a magnetização remanente isoterma (MRI) e a magnetização remanente anisterética (MRA). Assim como na ASM, o tensor da anisotropia de remanência magnética (ARM) é calculado de forma semelhante, onde:

$$M_{ri} = k_{rij} H_j$$

onde k_{rij} é a suscetibilidade remanente. Da mesma forma que na ASM, a ARM é também representada espacialmente por um elipsóide. Entretanto, uma diferença importante entre os tensores de anisotropia de suscetibilidade e de remanência é que enquanto a ASM fornece as contribuições magnéticas de todos os minerais presentes na rocha, a ARM identifica somente a contribuição dos minerais portadores de remanência (Jackson 1991; Trindade et al. 1999; Trindade et al. 2001).

2.1. Elementos de mineralogia magnética

Os minerais constituintes das rochas possuem propriedades magnéticas particulares que, somadas, compõem a suscetibilidade total da amostra. De acordo com essas propriedades, são verificados três tipos principais de comportamento magnético (Tarling & Hrouda 1993): diamagnetismo, paramagnetismo e ferromagnetismo.

O *diamagnetismo* é um tipo de magnetismo presente em todas as substâncias naturais, tendo sua origem relacionada ao movimento orbital dos elétrons sob a ação de um campo indutor. A suscetibilidade diamagnética é de intensidade muito baixa, negativa (direção

oposta ao campo indutor) e desaparece com a remoção do campo indutor (Fig. 1). A suscetibilidade diamagnética não depende das variações de temperatura e é característica dos minerais como quartzo, feldspato e calcita. Os valores de suscetibilidade diamagnética são em geral muito baixos (entre -13 e -17×10^{-6} SI), sendo, portanto, facilmente obliterados por suscetibilidades de intensidades mais altas.

Fig. 1: Propriedades dia- e paramagnéticas sob ação de um campo indutor (H). a) suscetibilidade diamagnética negativa (kd) em contraste com suscetibilidade paramagnética (kp) positiva; b) suscetibilidade paramagnética varia com a temperatura, enquanto que a suscetibilidade diamagnética é constante. M = magnetização; T = temperatura (extraído de Archanjo 2003).

A presença de íons de ferro nos silicatos dá origem a um tipo de comportamento magnético conhecido como *paramagnetismo*. A suscetibilidade paramagnética é positiva, independe do campo indutor e decresce com o aumento da temperatura. A variação da suscetibilidade paramagnética com a temperatura obedece à lei de Curie:

$$k = (T - \theta)^{-1}$$

onde T é a temperatura e θ é a temperatura de Curie. A anisotropia dos minerais paramagnéticos é controlada pela simetria cristalina (Fig. 2). Em minerais do sistema cúbico, como na granada, a anisotropia é nula, enquanto que em sistemas ortorrômbicos (*e.g.* ortopiroxênio) os eixos cristalográficos a , b e c coincidem com as direções principais de anisotropia de suscetibilidade (k_1 , k_2 e k_3). Por outro lado, no sistema monoclinico (anfibólio e biotita), somente o eixo b é paralelo a uma direção de suscetibilidade (Borradaile & Jackson 2004).

Fig. 2: Direções de suscetibilidade e simetria cristalina em diferentes sistemas cristalinos (modificado de Borradaile & Jackson 2004).

Fig. 3: Relações entre trama cristalina e trama magnética em um granito. A) foliação mesoscópica definida pela orientação de forma da biotita e lineação definida pelo alinhamento de cristais alongados ou “eixo” no qual cristais de biotita rotacionam; b) Polo do plano da foliação da biotita (centro do diagrama; clivagem da biotita; 001), e polo do plano que contém os polos de 001 (triângulo na borda sul do diagrama), definindo a lineação ou “eixo”; c) eixos de ASM correspondentes (k_1 : lineação; k_3 : polo da foliação), correlacionados aos eixos da trama cristalina (extraído de Bouchez 2000).

Silicatos ferromagnesianos, como micas e anfibólios, são os principais responsáveis pela suscetibilidade paramagnética em granitoides (Bouchez 1997). Em casos onde o paramagnetismo domina a suscetibilidade, a anisotropia desses minerais é fundamental para a caracterização da trama magnética, a exemplo do alinhamento preferencial de cristais monoclinicos de biotita paralelos à lineação magnética (Fig. 3).

O *ferromagnetismo senso lato* é caracterizado por uma magnetização natural espontânea gerada pela interação de íons de Fe, além de Mn, Cr e Co (Archanjo 2003). Minerais ferromagnéticos *sensu stricto* são caracterizados por momentos magnéticos

paralelos, apontando uniformemente para a mesma direção. Nos minerais ferrimagnéticos, os momentos são antiparalelos, resultando em uma magnetização residual, e os minerais antiferromagnéticos são marcados por momentos antiparalelos de mesma intensidade, definindo uma magnetização resultante nula (Fig. 4). Acima de uma determinada temperatura, os momentos magnéticos perdem sua organização e tornam-se aleatórios. Nos ferromagnéticos *s.s.*, esta temperatura é conhecida como Temperatura de Curie (T_c), enquanto que nos antiferrimagnéticos a mesma é dita Temperatura de Néel (T_n). Acima dessa temperatura de referência os minerais ferromagnéticos passam a se comportar como paramagnéticos com a suscetibilidade diminuindo com o aumento da temperatura (de acordo com a lei de Curie-Weiss).

Fig. 4: Momentos magnéticos com características paralelas e antiparalelas. Desvios suaves dos momentos magnéticos nos minerais antiferromagnéticos resultam nos clino-antiferromagnéticos (clino-afm) (extraído de Archanjo 2003).

As propriedades ferromagnéticas importantes podem ser definidas no ciclo de histerese. A suscetibilidade ferromagnética é obtida na primeira curva de magnetização sob campo indutor muito baixo (< 100 mT) quando há uma relação linear entre M e H . Com o aumento da intensidade do campo indutor a magnetização alcança a saturação (M_s). A partir do ponto de saturação, a magnetização é constante e máxima; removendo-se o campo indutor o mineral adquire uma magnetização remanente (M_r).

Os principais minerais ferromagnéticos encontrados nas rochas são as Ti-magnetitas e Ti-hematitas. A anisotropia magnetocristalina da Ti-magnetita, definida pela organização do retículo cristalino, é obliterada por uma anisotropia magnetoestática (ou de forma), cuja magnitude é dependente da morfologia e tamanho do grão, sendo maior nos grãos prolatos quando comparada aos cristais oblatos (Banerjee 1991; Borradaile & Henry 1997). Além

disso, o tamanho do cristal está relacionado com a presença de domínios magnéticos no interior do grão, os quais existem para minimizar a energia magnetoestática do cristal (Tarling & Hrouda 1993).

A presença de domínios magnéticos permite dividir os minerais ferromagnéticos em multidomínio (MD) ou monodomínio (SD). Grãos MD possuem magnetização estável e paralela ao eixo longo dos grãos, enquanto que nos grãos SD a direção da magnetização é perpendicular ao alongamento do grão. Dessa forma, os grãos SD exibem uma suscetibilidade máxima perpendicular à direção do eixo longo do grão, o que define uma trama “inversa” (Borradaile 1987; Rochette et al. 1992).

Estudos mostram que mesmo quando grãos de magnetita estão presentes em quantidades muito pequenas, a presença destes óxidos é suficiente para obliterar a suscetibilidade paramagnética e imprimir um caráter ferromagnético à rocha (Dunlop & Ozdemir 1997). Teores da ordem de 0,1% de magnetita, por exemplo, já seriam suficientes para que as propriedades magnéticas da rocha sejam dominadas pelo mineral. Quando esse teor aumenta, a magnetita pode desenvolver uma anisotropia textural ou de distribuição que ocorre quando os grãos estão próximos o suficiente para interagir magneticamente (Fig. 5). Nestes casos, a organização espacial do conjunto de grãos pode aumentar ou diminuir a anisotropia de suscetibilidade, dependendo do arranjo dos grãos (Fuller 1963, Hargraves et al. 1991; Gregoire et al. 1995; Bouchez 2000).

Fig. 5: Interações magnéticas entre grãos de magnetita: a) grão de magnetita individual e elipse de suscetibilidade; b) grãos alinhados paralelamente à direção de maior alongamento do grão promovem aumento na anisotropia; c) alinhamento “lado-a-lado” pode ocasionar diminuição da anisotropia (Gregoire et al. 1995; Bouchez 2000).

2.2. Aplicações da ASM em petrologia estrutural

2.2.1. Rochas parcialmente fundidas (migmatitos)

A primeira aplicação bem sucedida da metodologia de ASM em migmatitos foi realizada por Ferré et al. (2003), sendo posteriormente utilizada por diversos autores (Ferré et

al. 2004; Egydio-Silva et al. 2005; Denèle et al. 2007; Charles et al. 2009; Denèle et al. 2009; Schulmann et al. 2009; Kruckenberg et al. 2010). A análise sistemática das características magnéticas de migmatitos mostra que as relações estruturais complexas observadas em campo podem ser facilmente modeladas e interpretadas com base na trama magnética, a qual, em geral, obedece a um regime simples de fluxo magmático (Kruckenberg et al. 2010). A presença de migmatitos em terrenos granítico-gnáissicos, e o seu consequente estudo por meio da ASM, permite inferir sobre a dinâmica de fluxo da crosta intermediária parcialmente fundida, além de fornecer dados sobre migração e alojamento do material magmático em diferentes níveis crustais (Chardon et al. 2009, Vanderhaege 2009). A integração destes dados com informações sobre alojamento de corpos plutônicos permite reconstruir a trajetória de migração e ascensão do material magmático, constituindo, assim, uma importante contribuição para o entendimento da evolução crustal de orógenos antigos (Whitney et al. 2004).

2.2.2. Granitoides

A aplicação da ASM em granitos é, atualmente, uma técnica amplamente difundida e frequentemente utilizada no estudo estrutural de plútons (Bouchez 2000; Benn 2009). Trabalhos pioneiros detalhando conceitos, métodos e aplicações da ASM em granitos foram feitos por Bouchez et al. (1990), Benn (1994), Gregoire et al. (1995), Borradaile & Henry (1997), entre outros. Na Província Borborema, Archanjo (1993) realizou um amplo estudo das tramas magnéticas de granitos brasileiros alojados na Faixa Seridó, seguido por vários estudos semelhantes em diversas áreas da província (Olivier & Archanjo 1995; Neves 1996; Trindade 1999; Archanjo et al. 2002; Rodrigues 2008, etc.). A notável homogeneidade estrutural de granitos, observada desde a escala de batólitos até a escala de afloramento (Olivier et al. 1997), faz da ASM uma ferramenta essencial no estudo do fluxo magmático em contextos tectônicos variados, além de fornecer informações sobre a reologia de magma quando combinada com investigações microestruturais (Bouchez 2000).

3. Difração de elétrons retroespalhados (EBSD)

As técnicas mais comuns para medida das orientações preferenciais cristalográficas (OPCs) em rochas deformadas eram, até recentemente, a platina universal e o goniômetro de

textura. Embora ainda úteis em casos particulares, estas técnicas possuem limitações, como a identificação de apenas um eixo cristalográfico específico (platina universal) e análise em volumes de amostra, sem relações diretas com a microestrutura (goniômetro de textura). Tais restrições foram superadas com o método de difração de elétrons retroespalhados em um microscópio eletrônico de varredura (MEV-EBSD). Este método fornece uma imagem completa dos padrões de orientação cristalográfica de grãos individuais ou agregados poliminerálicos, de tamanhos variados, e sua correlação com a microestrutura observada em rochas deformadas. O sistema MEV-EBSD constitui atualmente uma ferramenta fundamental para estudos estruturais em zonas de cisalhamento (Adams et al. 1993; Prior et al. 1999; Lloyd 2004).

O princípio da técnica MEV-EBSD está fundamentado na interação entre feixes de elétrons e superfícies cristalográficas sólidas (amostras analisadas). Quando o feixe de elétrons atinge a amostra, os elétrons sofrem espalhamentos em direções diversas e a difração dos mesmos através dos planos do retículo produz dois cones, conhecidos como *cones de Kossel* (Fig. 6). A interação dos elétrons retroespalhados para formar os dois cones é regida pela lei de Bragg:

$$n\lambda = 2d_{hkl} \sin\theta$$

onde n é a ordem de reflexão, d_{hkl} a distância reticular (hkl), θ o ângulo de Bragg e λ o comprimento da onda ligada à tensão de aceleração dos elétrons.

Quando os elétrons retroespalhados são projetados em uma tela de fósforo eles desenvolvem faixas ou linhas denominadas bandas de Kikuchi (Fig. 6). Estas linhas possuem padrão paralelo e são caracterizadas como bandas brilhantes com uma largura distal de cerca de 2θ . Dessa forma, diferentes bandas constituem diferentes padrões de difração de elétrons retroespalhados em uma família de planos reticulares. Os ângulos entre as linhas de Kikuchi são correlacionados aos ângulos entre os planos reticulares, e a intersecção entre bandas caracteriza uma direção cristalográfica. Este padrão, portanto, constitui a essência do método EBSD (Lloyd et al. 1987; Prior et al. 1999; Higgins 2006).

Fig. 6: Formação dos cones de Kossel e linhas de Kikuchi a partir de difração dos elétrons sobre os planos do retículo cristalino (modificado de Randle & Enger 2000; in Morales 2007).

3.1. Características do equipamento e procedimentos analíticos

O sistema MEV-EBSD está ilustrado na Fig. 7. Um feixe de elétrons emitido pelo MEV atinge a amostra posicionada em uma superfície que faz um ângulo de $\sim 20^\circ$ com o feixe. Esta inclinação é adotada de forma a restringir a penetração do feixe de elétrons na amostra, otimizando, assim, o espalhamento dos elétrons na superfície da mesma (Fig. 7).

A tela de fósforo localizada no interior da câmara recebe os elétrons retroespalhados e a interação entre os mesmos gera uma figura de difração. Uma câmera de vídeo captura a imagem e a converte em uma imagem de maior resolução, minimizando ruídos.

Finalmente, o padrão de difração obtido é indexado e comparado com a base de dados minerais dos *softwares* de tratamento das imagens de difração, com o objetivo de obter as informações cristalográficas da amostra analisada (Schmidt & Olesen 1989; Prior et al. 1999; Randle & Enger 2000).

A indexação dos padrões de difração ocorre a partir da identificação, por parte do *software*, das intersecções entre as linhas de Kikuchi, que por sua vez vão caracterizar orientações cristalográficas. Para que esse processo seja feito de forma adequada, o feixe de elétrons incidente é carregado com uma aceleração em torno de 15-20 keV.

Feixes com carga energética superior a este valor padrão podem gerar imagens com menor contraste, uma vez que os elétrons podem penetrar a amostra e aumentar o ruído através de interações inelásticas indesejadas (Prior et al. 1999; Bascou 2002).

Fig. 7: Visão geral do sistema MEV-EBSD (fonte: www.dstu.univ-montp2.fr/TECTONOPHY/EBSD/EBSD-measurements.html).

3.2. Indexação das figuras de difração

A determinação das orientações cristalográficas extraída das linhas de Kikuchi é feita de forma automática a partir da transformada de Hough (1962). Este procedimento consiste em transformar os padrões simples de intersecção de linhas em pontos claros, os quais são facilmente reconhecíveis pelo computador a partir do algoritmo de detecção de bandas (Fig. 8). O algoritmo determina as orientações cristalográficas comparando a figura de difração com um modelo de esfera de difração (Fig. 9). Esta esfera é construída a partir de parâmetros dos planos dos retículos e das posições dos átomos equivalentes a partir de simulação

numérica (Schimdt & Olesen 1989). Os diferentes passos para indexação e identificação das orientações cristalográficas estão resumidos na Fig. 10.

Fig. 8: Transformada de Hough. Os pontos claros definem bandas de Kikuchi (manual channel 4.2, HKL technology; fonte: Lagoeiro 2010; http://www.igc.usp.br/fileadmin/disciplinasweb/d78/EBSD_USP.pdf).

Fig. 9: Esfera de difração construída a partir de simulação numérica de padrões de difração EBSD (modificado de Schimdt & Olesen 1989; in Morales 2007).

Fig. 10: Etapas do ciclo de indexação de uma imagem EBSD (fonte: Lagoeiro 2010; http://www.igc.usp.br/fileadmin/disciplinasweb/d78/EBSD_USP.pdf).

Os procedimentos específicos e condições de trabalho nas quais foram feitas as análises nos milonitos da zona de cisalhamento Patos podem ser encontrados no Capítulo 5.

4. Geocronologia U-Pb em zircão (*SHRIMP*)

O método U-Pb em zircão baseia-se essencialmente no decaimento radioativo de isótopos de ^{235}U e ^{238}U para isótopos de ^{207}Pb e ^{206}Pb , respectivamente. Em condições naturais, o U decai para Pb ao longo do tempo geológico e, dessa forma, a partir dos valores de meia-vida e da constante de decaimento, pode-se inferir o tempo necessário para que um isótopo de U sofra desintegração para um isótopo de Pb.

A determinação de idades geocronológicas a partir do decaimento radioativo no sistema U-Pb se dá por meio das razões isotópicas $^{238}\text{U}/^{206}\text{Pb}$ e $^{235}\text{U}/^{207}\text{Pb}$. O zircão é o mineral mais comumente usado neste tipo de análise por possuir uma estrutura cristalina bastante resistente a perturbações externas, além de exibir teores muito baixos de Pb, o que faz com que praticamente toda quantidade de Pb acumulada no zircão seja proveniente do decaimento radioativo do U (Dickin 1995).

A técnica de microsonda iônica de alta resolução (*Sensitive High Resolution Ion MicroProbe – SHRIMP*) tem sido utilizada para datação pontual de grãos de zircão, fornecendo informações precisas sobre idades de cristalização, bem como possíveis eventos metamórficos registrados no grão (Compston et al. 1984).

Em rochas ígneas e metamórficas, a datação U-Pb por meio do *SHRIMP* tem se mostrado eficiente na obtenção de idades de cristalização magmática e recristalização metamórfica. Como o zircão é um mineral que mantém sua estrutura cristalina estável mesmo em altas temperaturas ($\sim 600^\circ$), a datação dos núcleos dos grãos geralmente fornece informações acerca da idade de cristalização da rocha hospedeira. Por outro lado, bordas recristalizadas podem se desenvolver em eventos posteriores à cristalização do grão, permitindo a determinação da idade de eventos metamórficos, notadamente aqueles de alta temperatura (Dickin 1995).

As datações de zircão por meio do *SHRIMP* envolvem um estudo microestrutural preliminar dos grãos selecionados através da Catodo Luminescência (CL). A técnica permite visualizar a morfologia e a estrutura interna do zircão, além de auxiliar na escolha da região do cristal a ser atingida pelo feixe iônico (Dickin 1995). As características genéticas dos grãos de zircão também são analisadas por meio de CL e são de fundamental importância para a análise geocronológica. Zircões que apresentam zoneamentos oscilatórios normalmente cristalizam a partir de processos magmáticos, enquanto que zoneamentos setoriais ou localizados podem indicar eventos secundários de recristalização (Dickin 1995).

Dessa forma, um dado cristal de zircão que apresente núcleos com zoneamentos oscilatórios circundados por bordas sobrecrescidas pode fornecer idades contrastantes quando analisado nessas áreas distintas. A CL, portanto, é de fundamental importância na escolha da região de análise do grão, de acordo com o objeto de interesse (processos magmáticos, metamórficos, etc.).

5. Termometria de Ti em Quartzo – método TitaniQ

O conteúdo de Ti em cristais de quartzo fornece informações sobre temperaturas de (re)cristalização durante eventos magmáticos e metamórficos, fazendo do referido método um geotermômetro eficiente em estudos tectonotermiais (Wark & Watson 2006). O princípio teórico se baseia na atividade de óxidos de Ti (a_{TiO_2}), a qual é função da quantidade de Ti presente em cristais de quartzo. Experimentos em quartzo sintético na presença de rutilo, fluidos aquosos e silicáticos (*e.g.* magma), em temperaturas entre 600°C e 1000°C e pressão de 1.0 GPa, atestam para uma relação de substituição entre átomos de Ti e átomos de Si na estrutura do quartzo, a qual aumenta exponencialmente com o aumento de temperatura (Wark & Watson 2006). Rochas ígneas e metamórficas frequentemente possuem fases minerais

portadoras de Ti, como rutilo, ilmenita, biotita, etc. Estas fases são fonte de titânio para as trocas catiônicas com silício em cristais de quartzo.

Em milonitos, o método TitaniQ tem sido aplicado na identificação de temperaturas deformacionais a partir da análise de grãos de quartzo recristalizados em diferentes intervalos termais (fácies xisto verde – anfibolito). Kohn & Northrup (2009) demonstraram que, durante milonitização, o conteúdo de Ti pode reequilibrar-se entre porfiroclastos e neoblastos de quartzo; os grãos reliquiares preservariam um maior conteúdo de Ti e, consequentemente, forneceriam temperaturas mais altas, enquanto que os grãos recristalizados mais finos possuiriam conteúdos de Ti menores e registrariam temperaturas mais baixas.

A análise do conteúdo de Ti em quartzo é feita por meio de microsonda eletrônica, quando as temperaturas de cristalização de cristais de quartzo são superiores a 600°C, ou microsonda iônica para detecção de conteúdos de Ti em intervalos de temperatura a partir de 400°C (Wark & Watson 2006). Os cristais são previamente selecionados a partir de padrões de luminescência obtidos em imagens de CL. Fortes intensidades de emissão são características de quantidades altas de Ti, enquanto que emissões luminescentes de baixa intensidade são comuns em grãos recristalizados (Kohn & Northrup 2009). Estas características fazem da CL uma ferramenta útil na investigação de condições de (re)equilíbrio dos conteúdos de Ti durante perturbações termais no sistema (*i.e.* deformação).

No presente estudo, o método TitaniQ foi aplicado especialmente em milonitos de média- a baixa-temperatura (fácies xisto verde alto a anfibolito), com o objetivo de investigar as variações térmicas no interior da zona de cisalhamento Patos. Maiores detalhes sobre os resultados obtidos podem ser encontrados no Capítulo 5.

CAPÍTULO 3

ALOJAMENTO DE MAGMAS NA CROSTA INTERMEDIÁRIA PARCIALMENTE FUNDIDA E SUAS CONEXÕES COM O PLUTONISMO GRANÍTICO EM NÍVEIS CRUSTAIS SUPERIORES

1. Introdução

Este capítulo consiste em um estudo detalhado das relações entre o alojamento de anatexitos e suas relações com intrusões graníticas mais rasas.

A área de estudo localiza-se na Faixa Seridó, compreendendo o diatexito de Santa Luzia e os granitos Acari e Totoró. O migmatito Santa Luzia é caracterizado como um domo elíptico alongado na direção NE-SW, flanqueado por metassedimentos da Faixa Seridó a oeste e *augen* gnaisses do embasamento paleoproterozóico a ESE. É formado principalmente por *schlieren*-diatexitos e leucogranitos, onde são observados diversos xenólitos das rochas encaixantes na forma de blocos assimilados pelo magma granítico. Os migmatitos podem ser seguidos por uma faixa contínua de leucossomas e leucogranitos de direção NE, a qual exhibe conexões com o granito Acari.

Estudos prévios sugerem que a tectônica compressiva da faixa Seridó controla a cinemática de alojamento dos plútons adjacentes (Archanjo et al. 2002). Com efeito, as lineações magnéticas observadas no granito Acari exibem um padrão concordante com a elongação da Faixa Seridó. Entretanto, o plúton Totoró, localizado a norte de Acari, exhibe uma forma sub-circular e discordante do padrão tectônico regional, sugerindo tratar-se de uma intrusão pré-cinemática.

O artigo científico aqui apresentado reúne dados de ASM no diatexito de Santa Luzia e determinações isotópicas U-Pb em zircão (SHRIMP) nos migmatitos e granitos, afim de estabelecer a relação cronológica entre os corpos e investigar os mecanismos de migração do material parcialmente fundido e alojamento dos granitos na Faixa Seridó.

Os resultados obtidos sugerem que extensas áreas da Província Borborema foram metamorfisadas em alta temperatura/baixa pressão durante o Neoproterozóico (~ 575 Ma), e que os eventos de fusão parcial e alojamento de plútons graníticos são contemporâneos à deformação transpressiva observada no sistema de cisalhamento Patos-Seridó.

Timing of the HT/LP transpression in the Neoproterozoic Seridó Belt (Borborema Province, Brazil): Constraints from U–Pb (SHRIMP) geochronology and implications for the connections between NE Brazil and West Africa

Carlos J. Archanjo^{a,*}, Luiz G.F. Viegas^a, Maria Helena B.M. Hollanda^a, Laécio C. Souza^b, Dunyi Liu^c

^a Instituto de Geociências, Universidade de São Paulo, 05508-080 São Paulo, SP, Brazil

^b Departamento de Geologia, CCEI/UFRN, 59000-900 Natal, RN, Brazil

^c Beijing SHRIMP Center, Chinese Academy of Geological Sciences, Beijing 10037, China

ARTICLE INFO

Article history:

Received 23 November 2011

Received in revised form 9 April 2012

Accepted 1 May 2012

Available online 19 May 2012

Handling Editor: A.S. Collins

Keywords:

U–Pb (SHRIMP) geochronology

Granite

AMS

Seridó belt

Neoproterozoic

NE Brazil

ABSTRACT

The Seridó schist belt is a classical domain of the Borborema Province (NE Brazil) recording crustal melting at HT/LP conditions in a tectonic setting dominated by dextral transpressive deformation. A comparable tectonometamorphic setting has been described in the Pan-African Igarra and Ilesha schist belts of south Nigeria (Africa). The HT-transpression of the Seridó has previously been imprecisely dated by using a range of isotopic methods on a suite of syntectonic granites. We provide new U–Pb zircon ages of the Acari granite and Santa Luzia migmatite that constrain the HT/LP metamorphism of the Seridó at c. 575 Ma. Anisotropy of magnetic susceptibility of the Santa Luzia migmatite shows that the magnetic fabric records the NE-trending dextral kinematics similar to those previously described in the Acari pluton. On the other hand, the U–Pb ages of the Totoró pluton provide evidence for an older, mostly mafic to intermediate magmatism intruding the pelitic schist that precedes the HT/LP metamorphism by approximately 15 m.y. In contrast with the Acari pluton and Santa Luzia migmatite, the magnetic fabrics of Totoró show a concentric steep-dipping structure apparently related to magma ascent. Thermobarometric studies in the Seridó, Igarra and Ilesha schist belts indicate that a large domain in the northern Borborema and south of the Nigeria shields was deformed at a high temperature and at relatively shallow depths. Although the tectonic setting for such a post-collision HT event remains an open issue, it allows us nevertheless to refine the across-Atlantic geological correlation between the continents.

© 2012 International Association for Gondwana Research. Published by Elsevier B.V. All rights reserved.

1. Introduction

The break-up of West Gondwana in the Mesozoic led to the rupture of the Brasiliano–Pan-African belts between the South American and African continents to form the Atlantic Ocean. To reassemble West Gondwana in order to define the key geodynamics elements of these continents requires, therefore, a fine adjustment between the rock records on both margins across-Atlantic. However, despite a good fit between the platforms of the central and southern Atlantic, the precise correlation between the Neoproterozoic belts is still unresolved. Part of the problem is the perception that during their break-up and drift, the major continents have not always responded as single rigid blocks, leading to the distortion of the plates. To overcome this problem De Witt et al. (2008) proposed detailing the geology of ten piercing points to improve the correlations across the central and south Atlantic. Four of these piercing points are situated in the Borborema (NE Brazil) and Nigeria shields, and one of them corresponds to the vertical shear zones that deform the dominantly pelitic

Neoproterozoic deposits (schist belts) of the Seridó–Jaguaribe and Igarra–Ifewara–Ilesha domains (Fig. 1; Arthaud et al., 2008; Caby et al., 2008).

A critical issue related to the Borborema and Nigeria shields is the age of the magmatism, metamorphism and associated structures. A suture zone that closed the Pharusian Ocean at c. 600 Ma has been identified along the east margin of the West African craton through gravity anomalies and rock assemblages comprising mafic complex and high-pressure metamorphism (Attoh and Nude, 2008; Caby et al., 2008). Until now, no equivalent rock assemblages have been described on the Brazilian side. Thus, such a suture should be covered by Paleozoic sediments of the Paraíba Basin. Some transcontinental shear zones that match across the Atlantic, such as the Transbrasiliano/Kandi and Senador Pompeu/Ifewara (Fig. 1) resulted from collision processes and late accommodation structures due to the continued convergence of the cratons. In Seridó–Jaguaribe and Igarra–Ifewara domains, the shear zones coeval with the granitic magmatism are associated with a high temperature/low-pressure metamorphism (HT/LP; Lima, 1987; Zeh and Holness, 2003). However, there is growing evidence that some shear zones and associated plutons, dyke swarms and pegmatites (re)activate as late as in the Cambrian (Baumgartner et al., 2006; Archanjo et al., 2009; Hollanda et al., 2010). This implies that an

* Corresponding author.

E-mail address: archan@usp.br (C.J. Archanjo).

Fig. 1. Pre-Mesozoic fit between the northern Borborema Province (NE Brazil) and southern Nigerian shield (Africa) highlighting the schist belts and main vertical shear zones (sz). Inset: Sketch map of the West Gondwana and the study area between the Congo and West African cratons. The geology of south Nigeria after Caby and Boessé (2001).

accurate fit requires matching not only the vertical fault structures across the Atlantic, but also the timing in that tectono-magmatic events affected the rock record.

This study provides new U–Pb (SHRIMP) data for granites and migmatites formed during the HT/LP transpression that deformed the Seridó schist belt (Figs. 1 and 2). It combines zircon geochronology with the study of magmatic fabric using the anisotropy of magnetic susceptibility (AMS). In a pre-drift reconstruction the Seridó matches with the schist belts of the East Nigeria shield (Dada, 2008), particularly with the Igbarra sequence (Omitogun et al., 1991). The Seridó is bounded to the south by the Pátos shear zone that, in Nigeria, probably connects with the tectonic line that separates the East from the West Nigeria basement (Van Schmus et al., 2008). The Seridó is a classical domain of the Borborema Province that has been extensively studied due to its important mineral deposits (Au in quartz-veins, W in skarns and Ta–Nb in pegmatites and semi-precious stones). Although the structure and geochemistry of magmatic rocks have been studied in some detail (Leterrier et al., 1994; Araújo et al., 2001; Archânjo et al., 2002; Hollanda et al. 2003), an accurate chronology of the igneous pulses

and the relationship with deformation remain an open issue. We provide reliable crystallisation ages for zircons of syntectonic plutons as well as for overgrowths of inherited zircons that survived the migmatization of the basement, that allow us to constrain the timing of the HT/LP events. In addition, magnetic fabric data in migmatites indicate that the melting of the basement rocks was coeval with the emplacement of granitic plutons and transpressive deformation recorded along the Seridó-Pátos shear system.

2. Geological setting

2.1. The Seridó belt

The Seridó defines the NE-trending schist belt covered to the north by the sediments of the Mesozoic Potiguar basin and bordered to the south by the E-trending Pátos shear zone (Fig. 2). The belt is flanked by an older Paleoproterozoic basement of 2.1 to 2.4 Ga in age, named the Caicó Complex, which includes local Archean rocks (Dantas et al., 2004; Souza et al., 2008; Hollanda et al., 2011). An unconformity

Fig. 2. Simplified geological map of the Seridó-Jaguaribe domain. SPSZ, Senador Pompeu shear zone.

separates the basement from the succession of mainly clastic Neoproterozoic sedimentary rocks that is known as the Seridó Group. The group comprises a thick metaturbidite deposit above a clastic sequence of quartzite and metaconglomerate. Marble and paragneiss associated with calc-silicate rocks may be found locally below the psamo-pelitic sequence or near the base of the Seridó schist.

The deformation in the Seridó Group is characterised by a main phase that formed upright to inclined, open-to-tight NE-trending folds, associated with a variably dipping schistosity with a common fan-like appearance (Archanjo and Bouchez, 1991; Araújo et al., 2005). This event overprints a gentle-dipping foliation and recumbent folds attributed to a former thrust tectonics. In the core of large synclinal structures, such as around the Cruzeta dam (Fig. 3), greenschist facies conditions are preserved. Towards the central domain of the belt, the temperature increases to form andalusite, cordierite and sillimanite in aluminous schists accompanied by abundant quartz-feldspathic veins, aplitic dykes and granite intrusions. Exposures of the contact between the gneissic basement and cover rocks, such as those to the southwest around Santa Luzia, are deeply migmatized. Thermobarometric data

using garnet-biotite and plagioclase-garnet exchange reactions along E–W traverses crossing Florânia-Currais Novos, Cruzeta-Acari and Jardim do Seridó-Carnaúba dos Dantas (Fig. 3) allowed Lima (1987) to estimate a thermal gradient of 30–40 °C/km. Temperatures below 500 °C and pressures of about 3 kbar are recorded to the west approaching the core of the Cruzeta synform. Flanking the Acari pluton the temperatures increase up to 620 °C at pressures of 4 kbar. Stretched conglomerate clasts and elongated porphyroblasts of cordierite and andalusite are subhorizontal and parallel to oblique relative to main elongation of the belt. Kinematic criteria using C–C'–S planes and asymmetrical porphyroclasts are systematically dextral and include a component of flattening normal to the regional NE-trending foliation (Archanjo and Bouchez, 1991).

The Acari pluton consists of a porphyritic syeno- to monzogranite with biotite and hastingsitic to Fe-edinite hornblende. Lenses of biotite diorite, which sometimes occur as large angular enclaves, usually display features of mixing and mingling with granites. Sheet-like equigranular leucogranite often occupies the periphery of the batholith. The Totoró pluton consist of a monzodiorite to porphyritic quartz

Fig. 3. Geological map of the central domain of the Seridó belt. The foliation trends were outlined by using satellite and radar images. Thermobarometric geographic distribution after Lima (1987) and Souza (1996).

Table 1
Summary of SHRIMP U–Pb zircon data for Acari and Totoró plutons and Santa Luzia leucogranite.

Grain spot #	$^{206}\text{Pb}/^{238}\text{U}$	U [ppm]	Th [ppm]	$^{232}\text{Th}/^{238}\text{U}$	$^{206}\text{Pb}/^{238}\text{U}$ age	$^{207}\text{Pb}/^{235}\text{U}$ age	% disc	$^{207}\text{Pb}/^{206}\text{Pb}^*$	$^{207}\text{Pb}/^{235}\text{U}$	$^{206}\text{Pb}^*/^{238}\text{U}$	$\pm\%$	$\pm\%$	Err cor			
Acari porphyritic granite																
1.1	0.68	559	226	0.42	590	± 15	531	± 57	-11	0.0580	2.6	0.7665	3.7	0.0958	2.6	.7097
2.1	0.58	461	145	0.32	581	± 15	500	± 61	-16	0.0572	2.8	0.7444	3.9	0.0944	2.7	.7008
3.1	0.36	814	461	0.59	583	± 14	641	± 37	9	0.0610	1.7	0.7959	3.1	0.0946	2.6	8.295
4.1	1.06	370	692	1.93	586	± 15	608	± 92	4	0.0601	4.3	0.7892	5.0	0.0952	2.6	.5244
5.1	0.50	610	583	0.99	570	± 14	584	± 46	2	0.0595	2.1	0.7576	3.3	0.0934	2.6	.7717
6.1	0.97	676	272	0.42	569	± 15	540	± 62	-11	0.0583	2.8	0.7797	3.8	0.0970	2.6	6.758
7.1	0.86	351	268	0.79	563	± 14	513	± 77	-10	0.0576	3.5	0.7239	4.4	0.0912	2.6	5.976
8.1	2.09	106	86	0.84	595	± 17	611	± 230	3	0.0602	10.7	0.8026	11.1	0.0967	2.9	2.663
9.1	0.90	326	662	2.10	560	± 15	508	± 87	-10	0.0574	4.0	0.7182	4.8	0.0907	2.8	5.764
10.1	0.75	253	168	0.69	573	± 17	569	± 77	-11	0.0590	3.5	0.7562	4.7	0.0929	3.1	6.591
11.1	0.50	1269	545	0.44	624	± 15	523	± 35	-19	0.0578	1.7	0.8100	3.1	0.1016	2.6	8.256
12.1	3.19	105	109	1.08	659	± 28	170	± 351	-287	0.0495	15.0	0.7347	15.7	0.1077	4.5	2.882
13.1	0.53	797	382	0.50	589	± 15	509	± 54	-16	0.0574	2.4	0.7580	3.6	0.0957	2.6	7.258
14.1	0.30	1013	788	0.80	651	± 16	558	± 32	-17	0.0588	1.5	0.8607	3.0	0.1063	2.6	8.662
15.1	1.02	366	394	1.11	616	± 16	448	± 101	-38	0.0559	4.6	0.7727	5.3	0.1003	2.6	.5014
Acari fine-grained granite																
1.1	0.33	227	185	0.84	575	± 13	513	± 43	-12	0.0575	1.9	0.7404	3.0	0.0933	2.3	.7640
2.1	3.47	47	54	1.18	551	± 18	-221	± 469	349	0.0421	18.6	0.5175	18.9	0.0892	3.3	1.755
3.1	1.24	70	99	1.46	557	± 20	325	± 168	-72	0.0529	7.4	0.6590	8.3	0.0903	3.7	4.495
4.1	1.05	61	77	1.32	586	± 15	439	± 139	-34	0.0557	6.2	0.7296	6.8	0.0951	2.7	3.930
5.1	0.96	95	136	1.47	500	± 14	303	± 155	-65	0.0524	6.8	0.5833	7.4	0.0807	2.8	3.810
6.1	0.69	133	143	1.11	584	± 15	368	± 85	-59	0.0539	3.8	0.7053	4.6	0.0948	2.6	5.690
7.1	1.14	63	73	1.20	566	± 14	418	± 149	-35	0.0552	6.7	0.6978	7.2	0.0918	2.6	3.617
8.1	0.28	202	217	1.11	568	± 13	569	± 45	0	0.0591	2.1	0.7497	3.2	0.0921	2.4	7.639
9.1	3.31	89	130	1.52	586	± 14	652	± 265	10	0.0614	12.4	0.8053	12.6	0.0952	2.6	2.031
10.1	0.24	452	178	0.41	577	± 13	539	± 31	-7	0.0582	1.4	0.7521	2.7	0.0936	2.3	8.499
11.1	0.17	1008	1193	1.22	732	± 16	580	± 14	-30	0.0594	2.0	1.0127	2.4	0.1237	2.3	9.595
12.1	0.39	183	237	1.34	575	± 13	573	± 44	0	0.0592	2.0	0.7604	3.1	0.0932	2.3	7.584
13.1	1.58	79	98	1.28	540	± 13	326	± 133	-66	0.0529	5.9	0.6382	6.4	0.0874	2.6	4.007
14.1	2.27	66	89	1.39	563	± 14	491	± 204	-15	0.0570	9.2	0.7177	9.6	0.0913	2.7	2.776
15.1	3.04	68	76	1.16	587	± 15	12	± 341	-4829	0.0463	14.2	0.6088	14.4	0.0954	2.7	1.849
Totoró gabbro																
1.1	0.70	160	225	1.45	576	± 13	468	± 77	-23	0.0564	3.5	0.7271	4.2	0.0935	2.4	5.626
2.1	2.41	36	18	0.52	586	± 17	501	± 280	-17	0.0572	12.7	0.7512	13.1	0.0952	3.1	2.541
3.1	2.52	68	43	0.64	577	± 16	332	± 305	-74	0.0531	13.5	0.6858	13.8	0.0937	2.9	2.087
4.1	1.40	81	115	1.46	599	± 15	581	± 160	-3	0.0594	7.4	0.7970	7.8	0.0974	2.6	3.317
5.1	1.66	94	131	1.44	575	± 14	400	± 159	-44	0.0547	7.1	0.7036	7.5	0.0933	2.6	3.385
6.1	2.49	110	113	1.06	572	± 15	444	± 212	-29	0.0558	9.5	0.7135	9.9	0.0927	2.7	2.721
7.1	4.98	125	193	1.61	554	± 20	650	± 598	15	0.0613	27.9	0.7587	28.1	0.0898	3.8	1.335
8.1	1.47	64	46	0.74	553	± 12	735	± 208	25	0.0638	9.8	0.7875	10.1	0.0895	2.2	2.236
9.1	1.30	68	67	1.01	579	± 13	324	± 157	-78	0.0529	6.9	0.6849	7.3	0.0939	2.3	3.117
10.1	0.24	301	582	2.00	590	± 9	561	± 36	-5	0.0588	1.6	0.7768	2.3	0.0958	1.6	7.041
11.1	0.46	404	749	1.92	604	± 9	527	± 45	-14	0.0579	2.0	0.7840	2.6	0.0981	1.6	6.122
12.1	0.58	322	635	2.04	596	± 9	519	± 62	-15	0.0577	2.8	0.7703	3.3	0.0968	1.6	4.947
13.1	0.98	142	215	1.56	594	± 10	661	± 91	10	0.0616	4.3	0.8196	4.6	0.0965	1.8	3.886
14.1	0.59	470	1039	2.28	587	± 9	559	± 48	-5	0.0588	2.2	0.7725	2.7	0.0953	1.6	5.884
15.1	0.51	251	405	1.67	596	± 9	690	± 60	14	0.0625	2.8	0.8341	3.3	0.0968	1.7	5.102
16.1	0.88	251	443	1.83	610	± 10	578	± 74	-6	0.0593	3.4	0.8117	3.8	0.0993	1.7	4.482
Totoró diorite																
1.1	1.30	193	236	1.26	594	± 16	379	± 139	-57	0.0542	6.2	0.7215	6.8	0.0965	2.8	4.152
2.1	1.64	252	212	0.87	598	± 21	568	± 129	-5	0.0590	5.9	0.7911	6.9	0.0972	3.6	5.217

3.1	1.48	258	285	1.14	20.4	560	±15	607	±149	8	0.0601	6.9	0.7518	7.4	0.0907	2.7	.3659
4.1	1.16	141	148	1.08	11.5	578	±16	646	±128	10	0.0612	6.0	0.7918	6.7	0.0939	2.9	.4406
5.1	1.33	305	87	0.30	25.8	598	±15	566	±96	–6	0.0590	4.4	0.7898	5.2	0.0972	2.7	.5167
6.1	0.72	296	479	1.67	25.2	603	±15	475	±76	–27	0.0566	3.4	0.7650	4.3	0.0981	2.6	.6070
7.1	1.36	142	130	0.95	11.7	582	±20	726	±108	20	0.0635	5.1	0.8282	6.2	0.0945	3.5	.5702
8.1	0.19	447	790	1.83	39.2	626	±17	626	±50	0	0.0606	2.3	0.8522	3.7	0.1020	2.9	.7766
9.1	0.65	272	433	1.65	24.0	625	±16	631	±67	1	0.0608	3.1	0.8537	4.1	0.1019	2.6	.6449
10.1	0.46	499	62	0.13	42.9	613	±15	576	±40	–6	0.0583	1.9	0.8144	3.2	0.0997	2.6	.8102
11.1	0.74	416	878	2.18	35.7	609	±15	473	±58	–29	0.0565	2.6	0.7722	3.7	0.0991	2.6	.7038
Totoró porphyritic granite																	
1.1	3.47	140	124	0.92	11.8	584	±11	545	±318	–7	0.0584	14.6	0.7634	14.7	0.0948	1.9	.1323
2.1	2.18	158	149	0.98	13.9	615	±11	742	±196	17	0.0640	9.3	0.8829	9.4	0.1000	1.8	.1953
3.1	3.59	135	173	1.32	11.7	597	±9	322	±298	–85	0.0528	13.1	0.7064	13.2	0.0970	1.5	.1154
4.1	2.08	223	396	1.84	19.5	611	±7	581	±147	–5	0.0594	6.8	0.8141	6.9	0.0994	1.1	.1658
5.1	4.04	180	177	1.01	15.4	586	±7	147	±326	–299	0.0490	13.9	0.6426	14.0	0.0952	1.3	.0948
6.1	2.14	230	242	1.08	19.5	594	±6	655	±154	9	0.0614	7.2	0.8180	7.2	0.0966	1.1	.1542
7.1	2.12	258	317	1.27	21.6	587	±6	547	±177	–7	0.0585	8.1	0.7688	8.2	0.0954	1.0	.1265
8.1	0.86	263	334	1.31	21.9	592	±5	733	±70	19	0.0637	3.3	0.8458	3.5	0.0963	1.0	.2791
9.1	5.68	154	175	1.17	8.6	384	±6	910	±311	58	0.0684	15.1	0.5867	15.2	0.0613	1.7	.1087
10.1	1.67	197	232	1.22	16.1	577	±9	630	±192	8	0.0607	8.9	0.7843	9.0	0.0936	1.6	.1716
11.1	1.37	261	342	1.35	20.9	567	±8	534	±129	–6	0.0581	5.9	0.7370	6.1	0.0920	1.5	.2424
12.1	1.07	305	319	1.08	25.6	595	±5	517	±94	–15	0.0577	4.3	0.7683	4.4	0.0968	0.9	.2058
13.1	1.13	496	689	1.43	39.3	563	±4	580	±102	3	0.0594	4.7	0.7466	4.7	0.0912	0.8	.1706
14.1	2.12	188	197	1.09	15.7	587	±6	347	±185	–69	0.0534	8.2	0.7019	8.3	0.0953	1.1	.1363
15.1	2.49	209	426	2.10	16.9	564	±7	196	±293	–188	0.0500	12.6	0.6308	12.7	0.0915	1.2	.0984
16.1	1.52	255	279	1.13	20.5	569	±6	521	±124	–9	0.0578	5.7	0.7353	5.8	0.0923	1.1	.1876
17.1	7.34	46	51	1.16	4.0	577	±24	178	±1051	–224	0.0496	45.1	0.6412	45.3	0.0937	4.4	.0966
18.1	1.32	160	205	1.33	13.8	606	±8	762	±162	20	0.0646	7.7	0.8785	7.8	0.0986	1.3	.1685
19.1	4.20	142	143	1.04	12.1	583	±9	269	±438	–116	0.0516	19.1	0.6735	19.2	0.0946	1.7	.0865
20.1	7.93	109	87	0.83	9.4	572	±13	1036	±419	45	0.0738	20.7	0.9432	20.9	0.0927	2.4	.1136
21.1	3.01	207	176	0.88	17.5	586	7	467	±199	–26	0.0564	9.0	0.7400	9.1	0.0952	1.2	.1290
Santa Luzia leucogranite																	
GSL-1 sample																	
1.1 [r]	0.02	1384	41	0.03	111.6	578	±12	549	±14	–5	0.0585	0.6	0.76	2.3	0.0939	2.2	.9614
2.1 [r]	0.07	1512	42	0.03	119.8	568	±12	527	±14	–8	0.0579	0.7	0.74	2.3	0.0922	2.2	.9587
3.1 [r]	0.09	1217	56	0.05	97.4	574	±12	526	±17	–9	0.0579	0.8	0.74	2.4	0.0931	2.2	.9422
4.1 [r]	0.06	1642	89	0.06	132.5	579	±12	545	±14	–6	0.0584	0.7	0.76	2.4	0.0939	2.3	.9604
5.1 [r]	0.16	1759	74	0.04	139.4	568	±12	510	±19	–11	0.0575	0.9	0.73	2.4	0.0921	2.2	.9306
6.1 [r]	0.04	1527	39	0.03	117.8	554	±11	535	±14	–4	0.0581	0.6	0.72	2.3	0.0897	2.2	.9637
7.1 [r]	0.14	2208	154	0.07	169.0	549	±12	500	±14	–10	0.0572	0.6	0.70	2.4	0.0890	2.3	.9634
8.1 [r]	0.08	1799	54	0.03	142.5	568	±12	544	±13	–4	0.0584	0.6	0.74	2.3	0.0922	2.2	.9648
9.1 [r]	0.07	1604	48	0.03	126.4	566	±12	507	±13	–11	0.0574	0.6	0.73	2.3	0.0917	2.2	.9647
10.1 [r]	0.02	2020	85	0.04	159.1	565	±2	547	±16	–3	0.0584	0.7	0.74	0.8	0.0917	0.4	.4627
11.1 [r]	0.00	1676	66	0.04	133.8	573	±2	547	±17	–5	0.0585	0.8	0.75	0.9	0.0929	0.4	.4668
12.1 [r]	0.00	2110	68	0.03	170.2	579	±2	554	±16	–5	0.0586	0.7	0.76	0.8	0.0939	0.4	.4545
13.1 [r]	0.01	2171	135	0.06	174.8	578	±2	599	±14	4	0.0599	0.6	0.77	0.8	0.0937	0.4	.5627
GSL-2 sample																	
1.1 [c]	0.43	95	49	0.53	32.8	2161	±53	2208	±20	2	0.1385	1.2	7.60	3.1	0.3982	2.9	.9264
2.1 [c]	0.48	63	27	0.44	19.4	1969	±42	2137	±23	8	0.1329	1.3	6.55	2.8	0.3573	2.5	.8861
3.1 [c]	0.48	116	54	0.48	39.4	2136	±47	2204	±21	3	0.1382	1.2	7.48	2.7	0.3929	2.4	.8962
4.1 [c]	0.27	84	34	0.42	29.1	2175	±44	2220	±19	2	0.1384	1.1	7.72	2.8	0.4014	2.5	.9184
5.2 [r]	0.22	1281	10	0.01	103.6	578	±12	544	±22	–6	0.0584	1.0	0.76	2.5	0.0939	2.2	.9110
6.1 [c]	0.42	72	31	0.45	20.8	2098	±46	2198	±21	5	0.1376	1.2	7.30	2.8	0.3846	2.6	.9053
7.1 [c]	0.18	141	73	0.54	49.7	2217	±44	2198	±14	–1	0.1376	0.8	7.79	2.5	0.4104	2.3	.9452
8.1 [c]	0.49	43	26	0.63	15.1	2200	±50	2205	±28	0	0.1382	1.6	7.75	3.1	0.4067	2.7	.8550
8.2 [r]	0.38	987	20	0.02	77.3	561	±25	602	±25	7	0.0600	1.2	0.75	2.2	0.0909	2.2	.8863
9.1 [c]	0.18	157	107	0.70	57.6	2293	±48	2205	±11	–4	0.1382	0.6	8.14	2.6	0.4271	2.5	.9681

monzonite, including small stocks of diorite and gabbro-norite. Mixing and mingling features between mafic and felsic magmas are common. Geochemical Harker's diagrams show lower SiO_2 and higher FeO and MgO contents in Totoró compared to Acari, which essentially indicate the more mafic composition of the former (Nascimento et al., 2000).

Field structures and AMS studies show that the Acari pluton is syntectonic due to the high temperature dextral transpressive deformation of the metapelites (Jardim de Sá et al., 1986; Archanjo et al., 2002). The feldspar megacrysts of the Acari display a well-defined shape fabric parallel to the pluton elongation; however, this locally bends to outline a pluton-scale dextral magmatic shear zone (Fig. 3). The magnetic fabric is well-organised throughout with steep foliations mimetising the C-S surfaces of the shear zone at the centre of the intrusion, while lineations plunge to the SW. In contrast, magnetic foliation in the Totoró shows a concentric pattern with inward down to dip lineations. The contrasting internal structures of the Acari and Totoró lead Archanjo et al. (2002) to suggest that these plutons have different emplacement histories such that the fabric of Totoró traces the magma ascent. Available geochronological data (U/Pb TIMs, Rb/Sr) are restricted to the Acari granite. A single zircon fraction of a diorite provided a U/Pb age of 579 ± 7 Ma (Leterrier et al., 1994) while zircons from the porphyritic granite facies yielded a discordant age of 555 ± 5 Ma (Legrand et al., 1991). Furthermore, monazite from a leucogranite yielded an age of 528 ± 12 Ma (Baumgartner et al., 2006). These results indicate that the diorites are older than the granites but they conflict with field evidences suggesting that the diorites and porphyritic granites coexisted at the magmatic stage. Part of the difficulty in dating the Acari granite by isotope dilution on zircon may be related to the strong isotopic inheritance recorded in the Seridó granites (Van Schmus et al., 2008). Accordingly, we opted to study the Totoró and Acari plutons by using punctual (SHRIMP) analysis where primary igneous zoning of zircons was previously detected by cathodoluminescence images.

2.2. Geochronology of the Totoró and Acari plutons

2.2.1. Sampling and analytical methods

We collected three samples from the Totoró (a gabbro, a diorite and a porphyritic granite) and two samples from the Acari (a coarse porphyritic granite and an equigranular leucogranite; see locations in Fig. 3). All samples were prepared using standard crushing and separating procedures. Zircon grains were hand-picked from the mineral separates mounted in epoxy together with chips of TEMORA 1 reference zircon ($^{206}\text{Pb}/^{238}\text{U} = 416.75 \pm 0.24$ Ma; Black et al., 2003), sectioned approximately in half and polished. Reflected and transmitted light photomicrographs and cathodoluminescence (CL) SEM images were prepared for all grains. The spot areas (about 20–30 μm in diameter) for isotopic analysis were chosen in igneous growth sectors to obtain the zircon crystallisation ages. The analyses of U–Th–Pb were conducted using the SHRIMP II at the Beijing SHRIMP Centre (China), and a summary of the isotopic results is listed in Table 1. The isotopic data were collected in sets of five scans throughout the masses, and the TEMORA standard was measured every third unknown analysis. The abundances of U, Th and Pb, as well as the Pb isotope ratios, were normalised using the SL13 zircon standard ($\text{U} = 238$ ppm). Common Pb was corrected using the measured ^{204}Pb and assuming the appropriate model of Pb composition of Stacey and Kramers (1975). The data reduction was performed using the SQUID/Excel macro (Ludwig, 2000) and the statistical assessments (at the 95% confidence level) were calculated using the geochronological ISOPLOT/Excel 3.0 software of Ludwig (2003). All Concordia age errors were calculated at the 1σ level.

2.2.2. Results

The zircon crystals from the gabbro of Totoró sample are euhedral to subeuhedral showing elongated to stubbier forms. They are 100–250 μm in length and have a variable aspect ratio between 1:1 and 4:1 (typically 3:1). They are mostly colourless and inclusion-free,

but fractures are pervasive in all grains. The CL images show well-developed sector and subordinate concentric compositional zoning. No cores were observed. Eighteen analyses were performed on different grains, but two of them were aborted (#8.1 and #12.1) due to the instability of the primary beam. Eight analyses lie significantly to the higher- or lower-age side of the main data cluster (#4.1, #11.1 to #18.1) which provides a $^{206}\text{Pb}/^{238}\text{U}$ concordia age of 595.3 ± 3.4 Ma (Fig. 4A). Zircons separated from the dioritic sample are transparent, strongly fractured fragments with variable sizes and elongation ratios. Inclusions or cores occur locally. Most grains display magmatic, banded or slightly concentric compositional zoning; a few exhibit convoluted zonation and recrystallised inner areas. Eleven spot analyses were performed, but three (#1.1, #6.1, #11.1) were very discordant and were excluded from the age estimation. The remaining eight analyses are all near or concordant, yielding a concordia age of 597.0 ± 5.7 Ma (Fig. 4B) that is statistically indistinguishable from that of the gabbro sample.

Zircon crystals from the porphyritic granite sample are euhedral to sub-euhedral and clear; they are 100–350 μm in length with elongation ratios ranging from 1:1 to 3:1. Fractures, inherited cores and oscillatory igneous zoning are common features in the grains. Twenty-one analyses were conducted on magmatic sectors from different grains; ten

Fig. 4. U–Pb concordia diagrams for the diorite (A), gabbro-norite (B) and porphyritic granite (C) of the Totoró pluton (see analytical data in Table 1).

analyses were excluded as they were discordant or characterised by large errors in their individual analyses. The remaining eleven spot analyses (#1.1, #3.1, #5.1 to #8.1, #10.1, #12.1, #14.1, #19.1, #21.1) were grouped to define a concordia age of 591.0 ± 3.8 Ma (Fig. 4C), which is considered the best estimate of the crystallisation age of the porphyritic suite.

Two new U–Pb SHRIMP ages are reported here for the Acari pluton. Zircon crystals from the coarse porphyritic granite are euhedral, 150–400 μm in length and with aspect ratio ranging from 1:1 to 4:1. Most grains contain needle-like mineral (apatite?) and melt inclusions observed under transmitted light. The CL images show highly zoned crystals forming concentric patterns. Partially reabsorbed (embayed) and recrystallised sectors are also common features. Fifteen analyses were performed for different grains but only ten (#1.1 to #7.1, #9.1, #10.1, #13.1) define a concordia age of 577.7 ± 4.5 Ma (Fig. 5A). If considering only the absolute age, this result is just slightly older than the concordia age calculated from the eight concordant spot analyses (#1.1, #4.1, #7.1 to #10.1, #12.1, #14.1) of the zircon crystals of the equigranular leucogranite sample, which provided a concordia age of 572.1 ± 4.6 Ma (Fig. 5B). The zircon grains of the leucogranite sample are the same size as those from the coarse porphyritic sample, but the former show broad to poorly defined oscillatory igneous zoning. They are sometimes rimmed by discrete U-rich overgrowths that, however, were not large enough to be analysed.

In summary, the SHRIMP U–Pb absolute age for the sample of the porphyritic granite from the Totoró pluton, although a few million younger in age, cannot be distinguished within the uncertainty limits from those of the gabbro and diorite samples. Furthermore field evidence suggests that diorites and porphyritic granites coexisted at the magmatic stage (cf. Archanjo et al., 2002) indicating that the mafic and felsic magmas of the Totoró pluton crystallised at c. 595 Ma. In contrast, the temporal relationship between the Acari and Totoró plutons is

Fig. 5. U–Pb concordia diagrams for the coarse porphyritic granite (A) and equigranular leucogranite (B) of the Acari pluton.

now clear, the former being younger as shown by its reliable crystallisation ages at c. 575 Ma.

2.3. The anatexite of Santa Luzia

The anatexite of Santa Luzia consists of a dome-like structure that is 80 km² and elongated in the NE–SW direction. It comprises a core of migmatised rocks flanked to the west and to the east, respectively, by quartzite and sillimanite schist of the Seridó Group and the basement complex (Figs. 3 and 6). The structure of the host rocks consists of narrow lateral synclines recorded, respectively, by a keel of the supracrustal sequence and a tight folding of the gneissic banding. The axial plane foliation in the lateral synclines is steep-dipping, while the lineations (mineral and stretching) are subhorizontal. In contrast, the migmatitic core displays magmatic structures. Internally, it varies from a nebulite including relicts of pre-existing rocks or rock structures in diffuse contacts to more homogeneous portions of equigranular leucogranite (Fig. 7A). Pegmatite sheets usually occupy the contact with the host rocks. Angular to oval-shaped enclaves of the basement rocks (banded gneiss, augen gneiss and amphibole) of variable sizes are relatively frequent. Field evidence suggests that some enclaves were partially digested by the melt (Fig. 7B) to leave a

Fig. 6. Santa Luzia migmatite. (A) Schlieren structure defined by biotite-rich films alternating with granitic layers (leucosome) enclosing an apparently rotated older mesosome; note a more homogeneous domain (lower left) consisting of a biotite leucogranite (site 9). (B) Angular enclave of banded gneiss partially digested by the melt (left); the rafts of the banded gneiss in the host nebulite show curved contacts, sometimes looking stirred and disrupted resembling a schollen structure (site 16).

Fig. 7. Simplified geological map of the Santa Luzia migmatite and distribution of AMS sites.

mica-rich residue hosted by the nebulite. The field and structural relations suggest that the Santa Luzia migmatite was emplaced and expanded to form a dome exploiting the contact between the Seridó Group and its basement rocks.

The rocks that form the Santa Luzia dome have traditionally been ascribed in the regional mapping studies to the basement complex of the Seridó belt. However, indisputable basement rocks formed by banded gneiss, granitic orthogneiss and metamorphosed mafic dykes dated in the range of 2.4 to 2.2 Ga occur to the east of the dome (Hollanda et al., 2011). Migmatized rocks associated with leucogranite sheets and pegmatite can be traced northeastward in continuity with the Acari and Totoró plutons, which allowed us to suppose a genetic link between the intrusion of granite batholiths and melting of the underlying basement complex. The structural connections between the migmatites and granite plutons were investigated by comparing the anisotropy of magnetic susceptibility (AMS) of the Santa Luzia dome and those of the Acari and Totoró plutons which had been formerly investigated by the same technique (Archanjo et al., 2002). We include, in addition, a new zircon U–Pb age of a felsic neosome of the migmatite. This study, therefore, aims to investigate the structures and kinematic of the melted fraction of the Santa Luzia dome. We do not explore the ascent mechanisms of partially melted rocks in the crust neither the doming process in migmatitic rocks.

3. Anisotropy of magnetic susceptibility

3.1. Sampling and methods

The locations of the sites sampled for the rock magnetic studies are shown in Fig. 7. At each site we extracted at least three cores of approximately 8 cm in length and 2.5 cm in diameter using a portable gasoline-powered rock drill. The cores were oriented with a magnetic

compass. In the laboratory each core was cut into standard specimens that were 2.2 cm in length, such that the 42 sites provided 308 specimens for AMS measurements. The magnetic properties of the specimens were studied at the *Laboratório de Paleomagnetismo* of the *Universidade de São Paulo*. Low-field AMS was determined in a Kappabridge KLY-4 device and the mean-site parameters (magnitude, directions) of the anisotropy tensor ($K_1 > K_2 > K_3$; see Table 2) were calculated by ANISOFT v. 3.2 software provided by the constructor (AGICO Ltd). The anisotropy of anhysteretic remanence (AAR) was determined in a shielded room using an automatic JR-6 magnetometer and an alternate field tumbling demagnetiser coupled to a device that applies a weak direct (anhysteretic) field at different positions on a specimen. The basic procedure consists of exposing the specimen in cycles of demagnetisation in an AF on the order of 100 mT, followed by anhysteretic remanence acquisition and measurement of the resulting magnetisation. In contrast to AMS which describes the specimen bulk anisotropy, AAR isolates the contribution of remanence-bearing minerals from that of the paramagnetic and/or diamagnetic phases (see Jackson, 1991). To determine the best position scheme we initially computed the AAR ellipsoid by fitting the data for 6 and 12 positions. The results showed that the tensors were well-resolved and similar to each other for both schemes, so the AAR was calculated using the 6 positions protocol.

The temperature dependence of the magnetic susceptibility (k – T) was investigated from -200 °C to 700 °C using a CS-3/CS-L furnace apparatus linked to the KLY-4 Kappabridge. The samples were crushed to a fine sand grain size and the k – T was measured at each 3 °C interval. The coercivity spectrum analysis (Dunlop, 1972) was performed by placing a specimen in a steady magnetic field that was progressively increased. Applied fields from 10 mT up to approximately 2 T were imparted using a pulse magnetiser, and the isothermal remanent magnetisation (IRM) was measured in a Molspin magnetometer. Stepwise k – T and IRM measurements were investigated in 12 different specimens to cover the range of magnetic susceptibility values of the migmatite. At low temperatures, k – T was recorded by warming the sample to room temperature. From room temperature to 700 °C (heating cycle) and back (cooling cycle), k – T was measured in an argon environment to minimise the effects of atmospheric oxygen on the heating/cooling cycle.

3.2. Magnetic susceptibility and anisotropy

The bulk magnetic susceptibility (K) ranges from 0.02 to 55.59×10^{-3} SI (average = 7.01 mSI; SD 7.76). Ferromagnetic susceptibilities ($K > 1$ mSI) occur in 63% of the sites (Fig. 7) and very low susceptibilities, as low as 0.5 mSI, occur in 21% of the sites. Values of $K > 1$ mSI are found in the whole dome irrespective of the geographic location but sites with $K < 0.5$ mSI (#5, 16, 19, 20) occur to the west close to the contact with a pegmatite (Fig. 6). To assist the interpretations of the magnetic fabric we investigated the anisotropy directions of the AMS and AAR of representative sites with high (#10, #6), moderate (#18) and low K values (#19).

3.2.1. High susceptibility and anisotropy degree sites

Specimens with a high susceptibility ($K > 5$ mSI) and anisotropy degree ($P \geq 1.20$) comprise 51% of the sites. The IRM saturate above applied fields of 150 mT indicating the remanence is carried by a soft component (Fig. 8A, B). The k – T curves are very simple. They are characterised by a net Verwey transition at c. -160 °C and a susceptibility that vanishes almost totally above 580 °C. The cooling cycle is practically reversible indicating no new phases were formed on heating and cooling. Therefore, the IRM and k – T for the sites with very high (Fig. 8A) and high (Fig. 8B) susceptibility and anisotropy are typical of a low coercive coarse (Ti-poor) magnetite. The ellipsoids that describe the AMS and AAR are similar in shape and orientation. The anisotropy of

Table 2
AMS of the Santa Luzia anateixite.

Site	N	Km	P	T	K1			K2			K3		
					dec	inc	α_1	dec	inc	α_2	dec	inc	α_3
SL01	7	22.400	1.212	0.372	344	58	27.6	202	26	27	104	17	31.3
SL02	11	9.580	1.137	−0.187	18	24	13.3	271	34	23.9	135	46	23.7
SL03	10	2.870	1.234	0.134	345	36	82.5	158	55	82.5	253	3	15.8
SL04	8	3.310	1.259	−0.199	180	16	11.2	301	61	21.5	83	23	21.5
SL05	7	0.278	1.102	0.148	183	22	27.4	281	18	30.7	46	60	18
SL06	7	8.470	1.386	0.304	199	27	16.6	298	16	17.3	55	58	17.9
SL07	7	11.200	1.212	0.069	193	18	58.8	310	54	58.3	92	30	29.2
SL08	7	4.780	1.222	0.207	188	7	21.2	284	41	32.3	91	48	32.8
SL09	6	0.909	1.121	0.281	338	12	33.2	80	45	54	237	42	50.9
SL10	9	17.300	1.282	0.136	225	40	21.7	135	0	19.5	45	50	15.6
SL11	8	1.400	1.179	0.350	359	1	35.4	268	72	33.3	89	18	18.7
SL12	9	5.100	1.261	−0.019	214	41	28.5	33	49	30.5	124	26	26.5
SL13	8	4.620	1.279	0.075	213	14	15.1	317	45	16.5	110	42	16.5
SL14	8	10.600	1.122	−0.001	195	10	4.8	293	37	12.4	92	51	12.5
SL15	7	1.280	1.182	0.026	8	46	35.4	227	37	34.7	120	20	22.5
SL16	7	0.390	1.061	−0.135	24	5	11.6	219	85	21.3	114	1	22
SL17	6	21.500	1.180	0.072	40	1	7.9	310	18	16	131	72	16
SL18	12	6.720	1.169	0.121	15	5	9.9	276	57	14.4	109	32	14.4
SL19	8	0.176	1.021	−0.175	5	23	22.4	210	65	28.8	99	9	25.6
SL20	7	0.141	1.095	0.147	331	8	44.2	236	32	73.6	73	57	74.1
SL21	8	4.130	1.308	0.161	218	11	29.4	356	75	49.9	126	10	50.5
SL22	7	12.100	1.415	−0.238	212	19	20.7	68	67	24.1	307	12	17.5
SL23	8	8.140	1.213	−0.053	256	52	56	349	2	55.8	80	38	36.7
SL24	7	7.070	1.270	−0.068	199	11	15.1	297	36	15.6	95	52	16.1
SL25	7	3.280	1.191	0.178	181	19	30.2	290	44	53.9	74	40	53.7
SL26	7	4.500	1.367	−0.228	185	34	15.8	18	56	36.7	279	6	36.7
SL27	8	11.600	1.333	−0.052	210	28	22.4	119	2	23.2	26	62	22.2
SL28	7	5.650	1.267	−0.008	178	14	17.9	284	47	23.5	77	39	24.2
SL29	7	2.390	1.173	0.025	5	23	32.2	201	66	32.8	98	6	17.8
SL30	8	14.500	1.315	0.285	28	9	53.4	291	38	53.4	129	51	18.4
SL31	8	6.120	1.270	−0.258	12	15	14.5	205	75	14.3	103	3	10
SL32	6	17.400	1.259	0.118	238	9	24.8	336	41	19.1	137	48	20.4
SL33	7	0.217	1.019	−0.228	38	23	17.1	260	60	18.8	135	18	17
SL34	6	10.000	1.297	0.031	203	4	22.3	325	82	28	112	6	28.6
SL35	7	7.680	1.120	−0.098	36	11	10	178	76	12.3	305	8	10.3
SL36	6	7.240	1.179	0.187	210	6	17.7	318	72	16	118	17	9.1
SL37	7	1.240	1.260	−0.199	32	27	24.6	174	57	26.1	293	16	15.9
SL38	7	1.200	1.094	−0.015	31	12	5.9	273	65	8.1	126	21	8.1
SL39	5	2.290	1.290	−0.297	196	24	39.6	34	65	39.5	289	7	15.3
SL40	5	14.900	1.358	0.021	198	20	19.8	289	4	20.7	28	69	17
SL41	5	9.660	1.203	0.264	4	2	16.4	98	54	18.9	273	36	15.1
SL42	6	4.530	1.204	0.018	196	10	83.3	291	25	83.4	85	63	34.3

N, number of specimens; Km, mean-site susceptibility (10^{-3} SI); P, anisotropy degree; T, shape parameter. K1, K2 and K3 are the orientations (declination, inclination) of the AMS principal directions, and α_1 , α_2 and α_3 are the angular dispersion represented by the maximum semi-angle (degrees) of the confidence cone around the mean direction.

the AAR is higher than that of the AMS and the ellipsoids show oblate shapes ($T > 0$) and the respective principal directions are nearly coaxial.

3.2.2. Moderate to low susceptibility and anisotropy degree sites

Specimens with moderate susceptibility ($1 < K$ (mSI) ≤ 5 ; $1.10 < P < 1.20$) comprise 22% of the sites while those with low susceptibility (K (mSI) < 1 mSI) comprise 27%. Specimens with moderate susceptibility (Fig. 8C) have k - T curves quite similar to the specimens with high susceptibility, indicating that magnetite is the main carrier of the susceptibility and anisotropy. However, IRM does not saturate as shown by the smooth, but progressive increase of the remanence between 0.5 and 2.0 T. Thus, besides a soft component, the IRM analysis indicates the presence of a high coercive (hard) component. The k - T shows that the soft component must correspond to a coarse Ti-poor magnetite, while the hard component is attributed to hematite which does not saturate even for fields above 2 T. Considering the gentle slope of the IRM above 0.5 T and the lack of evidence for Morin (-10 °C) or Néel (670 °C) transitions, we estimate that a small amount of hematite must occur in the sample.

IRM acquisition curves of specimens with $K < 0.5$ mSI show a steep initial slope and reach nearly 90% of their saturation magnetisation by 150 mT. However, the curve continues to climb and does not saturate at 1.4 T (Fig. 8D). The k - T curve at a low temperature shows a poorly

defined Verwey transition, which may be attributed to a very low content of fine magnetite grains and/or its suppression by an oxidising magnetite (Ozdemir et al., 1993). During the heating cycle, the susceptibility decreases between 280 and 320 °C suggesting that the sample contains a magnetic sulphide (pyrrhotite, greigite; $T_C \sim 320$ °C). The curves also show a strong increase in susceptibility between 500 and 550 °C but this is not reproduced on cooling, indicating that new magnetic phases were formed on heating. The presence of magnetite and hematite is therefore only suggested by, respectively, the soft and hard coercive components detected in the IRM curve.

The principal directions of the AMS and AAR share a similar orientation even for the site displaying contrasting ellipsoid shapes (Fig. 8C, D). Like those recorded in sites with high susceptibility, the anisotropy degree is higher in the AAR than the AMS. Variations in the magnitude of the anisotropy directions and, therefore, in the ellipsoid shape (Fig. 8D) can be attributed to the increasing contribution of paramagnetic silicates to the bulk susceptibility as the content of the ferromagnetic fraction decreases down to 0.1% (Tarling and Hrouda, 1993).

3.3. Anisotropy of susceptibility directions

The magnetic fabric, represented in Fig. 9 by the mean-site lineation (K1) and foliation (plane normal to K3), has a relatively uniform

Fig. 8. Isothermal remanent magnetisation (IRM) acquisition and respective susceptibility (k)-temperature (T) dependence for high (A, B), moderate (C) and low (D) magnetic susceptibility for representative specimens of the Santa Luzia dome. The ASM and AAR principal directions ($k_1 \geq k_2 \geq k_3$) of the corresponding sites reveal a good concordance between the fabric elements (N, number of measured specimens). Lower-hemisphere, Schmidt stereonet.

orientation. The magnetic lineation plunges gently to the NNE or SSW and the foliation generally dips to the NW. The magnetic fabric is slightly oblique with regards to the mean NE-trending elongation of the dome, a feature that has been also observed further north in the Acari pluton (Archanjo et al., 2002). We also observe that the magnetic foliation does not reproduce the opposite dip of the contacts in the flanks of the dome, which suggest that the AMS records the latest incremental deformations that have affected the melt.

3.4. Geochronology

We collected a sample from a nebulite located in the centre of the dome (see location on Fig. 6). It is composed of a leucocratic rock containing “gosh” layers of alternate lighter and darker material. The hand sample consists of a homogeneous, coarse-grained and equigranular biotite granitic nebulite. The zircon crystals are euhedral and many

display dipyrnidal terminations. Their lengths are of 50–200 μm with an aspect ratio between 1:1 and 3:1. In transmitted light they show xenocrystic cores and large dark areas that appear to be metamict. Many grains are rimmed by zircon overgrowths. In CL images (Fig. 10) these overgrowths form dark (U-rich), slightly zoned rims measuring about 5–30 μm in width that are probably metamorphic in origin.

Sixteen analyses of the dark high-U rims yielded slightly reverse ages with a weighted mean $^{206}\text{Pb}/^{238}\text{U}$ age of 573.2 ± 3.4 Ma (Fig. 11). Eight analyses of the zircon cores resulted in an upper intercept age of 2203 ± 17 Ma that is consistent with ages we have determined from nearby basement exposures (Hollanda et al., 2011). When plotted together these two groups align quite well along a discordia line with a lower intercept at 575.1 ± 3.4 Ma. This younger age obtained on the U-rich rims is interpreted as the crystallisation age of the granitic neosome resulting from the HT event that melted the basement complex.

Fig. 9. Magnetic fabric (foliation, lineation) of the migmatitic dome. Density (Kamb) contours, lower hemisphere; the grey line correspond to the elongation direction of the dome.

Fig. 10. Cathodoluminescence image of selected zircons from the Santa Luzia migmatite. The grains usually show an inner concentric zoning rimmed by a dark, U-rich overgrowths.

4. Discussion

4.1. Significance of magnetic fabrics in the migmatite

The AMS sampling of the Santa Luzia dome was restricted to the more homogenous granitic portions of the migmatite where a magmatic microstructure can be recognised. Thus, the AMS properties of the samples are analogous to the common granitoids in that their general principles are described in Bouchez (1997). Rock magnetic studies showed that magnetite and minor hematite are the main mineral phases responsible for the magnetic susceptibility and anisotropy. Magnetite can be detected by the steep initial slope of the IRM, the Verwey transition at c. -160°C and the net drop of susceptibility on heating near 580°C . Hematite was inferred by the presence of a highly coercive fraction for samples that do not saturate up to induced fields of 2 T and, in some k - T runs, by the progressive decreasing of k after 570°C until it completely disappears at c. 700°C (not shown in Fig. 8). These magnetic characteristics are consistent with hematite which has a Néel temperature of 670°C . Moreover, some samples of the low susceptibility group may contain traces of a magnetic sulphide as suggested by the small drop in susceptibility on heating at c. 300°C . Alternatively, the decreasing of the susceptibility magnitude at c. 300°C could correspond to mineral alteration during the heating cycle as evidenced the strong increase of k just before the Curie temperature of magnetite.

Evidence for magnetite is also seen in the AAR studies as a fabric induced by an anhysteretic induction is only recorded in the phases that retain a remanence, which excludes dia- and paramagnetic minerals. The results show a very good correlation between the orientations of the principal directions of the AAR and ASM irrespective of the magnitude of the susceptibility. Moreover, at sites with moderate to high susceptibility ($K > 1$ mSI), there is a good correspondence between the symmetry of the AAR and AMS ellipsoids. Differences in the ellipsoid symmetry were found in samples with low susceptibility values. This can be attributed to the increasing importance of paramagnetic silicates in the susceptibility and anisotropy as the susceptibility declines down to 10^{-3} SI. For a very low susceptibility, say less than 0.30 mSI, the contribution to the AMS of paramagnetic silicates becomes dominant and the ferrimagnetic grains, if any are available, occur as submicroscopic inclusions usually hosted in silicates (Rochette, 1987; Borradaile and Werner, 1994). The different grain sizes of magnetite are also evidenced in the thermomagnetic curves. Well-defined Verwey transitions at c. -160°C and net Curie temperatures of c. 580°C are typical of a coarse multidomain Fe-rich grain, while poorly defined Verwey transitions are attributed to fine and/or oxidising grains (Ozdemir et al., 1993). In our samples, however, the finest grains are not small enough to develop fabric inversions produced by single-domain grains as indicated at site SL19 (Fig. 8D) which, despite of differences between the AAR and AMS ellipsoid shapes, shows a good correspondence between their principal directions.

In summary, we conclude that the AMS of Santa Luzia dome is dominated by shape anisotropy of magnetite. Some hematite, magnetic sulphide and biotite contribute locally to the bulk AMS but they do not substantially modify the fabric produced by the magnetite. The resulting magnetic fabric is well-organized in the whole dome, but symptomatically it does not mimic the shape of the dome. This is evident in the magnetic foliation that does not reproduce the dip to the east of the eastern dome margin as well as the mean lineation direction oblique to the dome elongation. These results imply that the: (i) AMS records the latest magmatic deformation events before the full crystallisation of the neosome, and (ii) magnetic fabrics can be interpreted as syntectonic to a dextral HT event in that the lineation and foliation tend to align in the direction of the principal flattening plane of the regional strain ellipsoid (Fig. 12). The magnetic fabric of the Santa Luzia migmatite, therefore, is similar to that recorded in the Acari pluton.

Fig. 11. U–Pb concordia diagram for a granitic nebulite of the Santa Luzia dome detailing the analysis in the zircons cores (upper intercept) and in the U-rich overgrowths around the zircon margins (lower intercept). The zircon overgrowths yields a ^{207}Pb -corrected $^{206}\text{Pb}/^{238}\text{U}$ mean weighted age of 573.2 ± 3.4 Ma.

4.2. Timing of the HT deformational event

The new SHRIMP U–Pb zircon ages of a porphyritic granite and a leucogranite of the Acari pluton and the granitic leucosome of the Santa Luzia migmatite allow us to constrain the timing of the HT deformation of the Seridó belt at c. 575 Ma. This event was responsible for melting the Caicó Complex to form migmatite domes that, like in Santa Luzia, host large enclaves of the Paleoproterozoic basement and zircon xenocrysts rimmed by overgrowths of Neoproterozoic

age. Furthermore, the melts and the HT/LP metamorphism coexisted with a general dextral shear deformation that produced regional fan-like foliations including belt-parallel stretching lineations (Archanjo and Bouchez, 1991; Araújo et al., 2001). SHRIMP U–Pb zircon ages of the porphyritic granite and leucogranite are consistent with the age of the Acari diorite yielded by conventional U–Pb methods (Leterrier et al., 1994). However, the younger U–Pb ages calculated from the zircon and monazite of samples from the same granite are here attributed to the resetting of the isotopic system by a newer thermal episode

Fig. 12. Schematic block-diagram illustrating the relationship between the fabric of the magmatic rocks (dashed pattern) and the structure of the central domain of the Seridó belt. The block containing the Acari and Totoró plutons is at the same topographic level than that of Santa Luzia, being shown uplifted to illustrate the inferred deep structure. It combines E–W shortening and NE-trending dextral strike-slip shearing synchronous with the emplacement of the Acari pluton and Santa Luzia migmatitic dome. The significance of the magnetic fabrics (lineation and foliation) of the Totoró and Acari plutons are discussed in Archanjo et al. (2002). Density (Kamb) contour in the lower hemisphere stereonet.

possibly related to the intrusion of the Seridó pegmatitic province (Araújo et al., 2001; Baumgartner et al., 2006). The HT-metamorphism and melting of the basement dated between 578 and 574 Ma have also been described to the east of the Seridó belt. These ages were recorded by Sm–Nd on garnet-whole rock from a granitic leucosome and on garnet bearing alkaline granites emplaced in an extensional setting (Souza et al., 2006).

SHRIMP U–Pb concordia ages between 597 and 591 Ma were given by, respectively, the mafic and felsic intrusions that constitute the Totoró pluton. At first glance, we could consider the felsic and mafic rocks to have been formed by separate magmatic pulses but field evidence indicates that the diorites and granites coexisted at the magmatic stage. Therefore, if we assume that the pluton was built from a single magmatic event and take into account the analytical errors of the individual analyses, we estimate the crystallisation age of mafic and felsic rocks at c. 595 Ma. By comparing the results from the Totoró with Acari plutons and Santa Luzia migmatite and even considering the errors from the individual analyses of the samples, it becomes evident that the Totoró magmatism is approximately 15 m.y. older than the HT tectonothermal event recorded in the central domain of the Seridó belt. Further evidence for a distinct evolution of the Totoró pluton come from the magnetic fabric as its steep plunging lineation contrasts with that of the Acari pluton and Santa Luzia migmatite (Fig. 12). The geochronological and structural evidence indicate, therefore, that the Totoró type magmatism precedes the HT-transpression of the Seridó belt. An older mafic magmatism has been identified in a few places in the Seridó-Jaguaribe domain, such as a leuconorite dated by the U–Pb method at 599 ± 16 Ma (Dantas, 1997) and the Riachão gabbro at 588 ± 6 Ma (Guimarães et al., 2009). Similar ages have been found for carbonatites and alkaline granites in the Pan-African Dahomeyide (West Africa) and these have been linked to an extensional episode following the collision of the West African craton (Attoh et al., 2007).

Lima et al. (1989) describe pyroxene hornfels in the contact of the Totoró gabbro-norite with the host mica schist and in a 50 m² wide pelitic enclave within the mafic rock. In hornfels-bearing garnet-biotite-cordierite assemblages they analysed the exchange reactions of Fe–Mg to estimate the equilibrium reactions at temperatures in the range of 580–700 °C and pressures of 4–6 kbar. These results do not differ from the thermobarometric conditions calculated in the host rocks of the Acari pluton (Lima, 1987; Souza, 1996), allowing us to propose that the ionic exchange reactions of the Totoró hornfels were reset by the late HT transpressive event at c. 575 Ma. An important consequence of the thermobarometric data is that the HT mineral assemblages equilibrated at relatively shallow depths, possibly after suturing the Nigeria and Borborema shields in West Gondwana at 600 Ma.

4.3. Implications for the connections with the Nigeria shield

The Igarra and Ilesha schist belts of southern Nigeria share a similar structural history as that described in the Seridó-Jaguaribe domain (Arthaud et al., 2008; Dada, 2008). Metamorphic assemblages in the Igarra metapelites comprise andalusite-staurolite-garnet equilibrated at temperatures of 550 ± 50 °C and pressures of c. 3 kbar (Omitogun et al., 1991). In the Ilesha schist belt the metamorphic assemblages also include sillimanite with peak metamorphic conditions calculated at 590 ± 20 °C at c. 5 kbar, followed retrogression to 550 °C and 3 kbar (Zeh and Holness, 2003). Both in Igarra and Ilesha, the HT/LP metamorphism was coeval with N-trending dextral strike-slip shear zones.

In a pre-drift reconstruction a close match between the Igarra-Seridó and Senador Pompeu/Orós-Ifewara geological domains becomes evident. The Orós belt (see Fig. 2) is situated near the dextral Senador Pompeu shear zone. It consists of a rift sequence including an aluminous schist, quartzite lenses and local Ca- to Mg-rich carbonate rocks overlaying acid metavolcanics. The metamorphism varies from greenschist facies to amphibolite facies with andalusite-staurolite-

garnet assemblages to increases up to HT-amphibolite and granulite approaching the Atlantic coast (Arthaud et al., 2008). Barrovian-type metamorphism prevails to the west of the Senador Pompeu shear zone. However, there are important differences between the geology of the Orós and the Neoproterozoic Seridó belt. Firstly, the Orós volcanism occurred in the Late Paleoproterozoic (Sá et al., 1995) and its rift deposits include magnesite layers and possibly gypsum, which is suggestive of an evaporitic environment (Parente et al., 2004). Orós-type sequences have not been described until now in southern Nigeria and, evidently, the search for similar rock associations is critical to establish fine-scale geological correlations across the Atlantic.

5. Conclusion

Large areas situated in the Borborema Province to the north of the Patos shear zone and in south Nigeria comprising the Igarra and Ilesha schist belts were metamorphosed under a high temperature/low-pressure regime. Andalusite and garnet are common index minerals in pelitic rocks on both sides of the Atlantic, and the Seridó schist belt in the Brazilian side include in addition syn-kinematic cordierite-sillimanite assemblages usually forming aureoles around granite plutons. The timing of the emplacement of such plutons and the migmatization of basement rocks are now constrained by the U–Pb (SHRIMP) ages determined in the magmatic zoning sectors of zircons from the Acari granite and in the metamorphic overgrowths rimming older zircons from a felsic neosome of the Santa Luzia migmatite. Coarse porphyritic granites and leucogranites from the Acari samples provided crystallisation ages between 577 Ma and 572, while dark U-rich overgrowths from the Santa Luzia sample yielded a concordant lower intercept age of 575 Ma. However, the errors of the analysis of the zircons from the granite and migmatite overlap (1 σ level) to indicate that the respective ages are indistinguishable from each other. The AMS of the Santa Luzia anatetic dome show a relatively simple organisation in that foliations dip mostly to the SW and lineations align slightly oblique with respect to the dome elongation. A similar fabric pattern reflecting a general dextral NE-trending transpression has been previously described in the Acari pluton. Therefore, the U–Pb (SHRIMP) ages and the AMS results combine to fix the HT/LP transpressional event of the Seridó belt at c. 575 Ma. This thermotectonic episode succeeds an older bimodal magmatism represented in the Seridó belt by the Totoró pluton. The U–Pb zircon data from samples collected from different igneous units (diorite, gabbro-norite and porphyritic granite) yielded reliable ages in the range from 597 to 591 Ma. Assuming that the Totoró pluton was built in a single magmatic pulse, as supported by field evidence of mixing and mingling between dioritic and granitic magmas, we estimate that the crystallisation of the pluton would have occurred at c. 595 Ma. The geological significance of such a pre-tectonic magmatism and the timing of the HT-metamorphism in the Igarra and Ilesha schist belts are unknown. However, similar lithologies and thermobarometric conditions place these schist belts as a first order element in searching for across-Atlantic geological correlations.

Acknowledgements

Carlos J. Archanjó and Luis G. Viegas thank Fundação de Amparo a Pesquisa do Estado de São Paulo (FAPESP) for the financial support (grants 2010/50060-1 and 2009/17537-1). We also wish to thank Sérgio Neves and an anonymous reviewer who provided useful corrections which improved the presentation of our work.

References

- Araújo, M.N.C., Alves da Silva, F.C., Jardim de Sá, E.F., 2001. Pegmatite emplacement in the Seridó belt, Northeastern Brazil: late stage kinematics of the Brasiliano orogen. *Gondwana Research* 1, 75–85.
- Araújo, M.N.C., Vasconcelos, P.M., Alves da Silva, F.C.A., Jardim de Sá, E.F., Sá, J.M., 2005. ⁴⁰Ar/³⁹Ar geochronology of the gold mineralization in Brasiliano strike-slip shear

- zones in the Borborema province, NE Brazil. *Journal of South American Earth Sciences* 19, 445–460. <http://dx.doi.org/10.1016/j.sames.2005.06.009>.
- Archânjo, C.J., Bouchez, J.L., 1991. Le Seridó, une chaîne transpressive dextre au Proterozoïque Supérieur du Nord-Est du Brésil. *Bulletin de la Société Géologique de France* 168, 509–520.
- Archânjo, C.J., Trindade, R.L.F., Bouchez, J.L., Ernesto, M., 2002. Granite fabrics and regional-scale strain partitioning in the Seridó belt (Borborema Province, NE Brazil). *Tectonics* 21 (1). <http://dx.doi.org/10.1029/2000TC001269>.
- Archânjo, C.J., Launeau, P., Hollanda, M.H.B.M., Macedo, J.W.P., Liu, D., 2009. Scattering of magnetic fabrics in the epizonal Cambrian alkaline granite of Meruoca (Ceará State, northeast Brazil). *International Journal of Earth Sciences* 98, 1793–1807.
- Arthaud, M.H., Caby, R., Fuck, R.A., Dantas, E.L., Parente, C.V., 2008. Geology of the Northern Borborema Province, NE Brazil and its Correlation with Nigeria, NW Africa. *West Gondwana: Pre-Cenozoic Correlations Across the South Atlantic Region*. In: Pankhurst, R.J., Trouw, R.A.J., Brito Neves, B.B., De Wit, M.J. (Eds.), Geological Society of London, Special Publications, 294, pp. 49–67. <http://dx.doi.org/10.1144/SP294.5>.
- Attoh, K., Nade, P.M., 2008. Tectonic Significance of Carbonatite and Ultrahigh-pressure Rocks in the Pan-African Dahomeyide Suture Zone, Southeastern Ghana. The boundaries of the West African Craton. In: Ennih, N., Liegeois, J.P. (Eds.), Geological Society of London, Special Publications, 297, pp. 217–231. <http://dx.doi.org/10.1144/SP297.10>.
- Attoh, K., Corfu, F., Nade, P.M., 2007. U–Pb zircon age of deformed carbonatite and alkaline rocks in the Pan-African Dahomeyide suture zone, West Africa. *Precambrian Research* 155, 251–260.
- Baumgartner, R., Romer, R.L., Moritz, R., Sallet, R., Chiaradia, M., 2006. Columbite-tantalite bearing granitic pegmatites from the Seridó belt, northeastern Brazil: genetic constraints from U–Pb dating and Pb isotopes. *The Canadian Mineralogist* 44, 69–86.
- Black, L.P., Kamo, S.L., Allen, C.M., Aleinikoff, J.N., Davis, D.W., Korsch, R.J., Foudolis, C., 2003. TEMORA 1: a new zircon standard for Phanerozoic U–Pb geochronology. *Chemical Geology* 200, 155–170.
- Borradale, G.J., Werner, T., 1994. Magnetic anisotropy of some phyllosilicates. *Tectonophysics* 235, 223–248.
- Bouchez, J.L., 1997. Granite is never isotropic: an introduction to AMS studies of granitic rocks. In: Bouchez, J.L., Hutton, D.H.W., Stephens, W.E. (Eds.), Granite: from segregation of melt to emplacement fabrics. Kluwer, Dordrecht, pp. 95–112.
- Caby, R., Boessé, J.M., 2001. Pan-African nappe system in southwest Nigeria: the Ife-Ilesha schist belt. *Journal of African Earth Sciences* 33, 211–225.
- Caby, R., Buscail, F., Dembélé, D., Diakité, S., Sacko, S., Bal, M., 2008. Neoproterozoic Garnet–glauconophanes and Eclogites: New Insights for Subduction Metamorphism of the Gourma Fold and Thrust Belt (Eastern Mali). The boundaries of the West African Craton. In: Ennih, N., Liegeois, J.P. (Eds.), Geological Society of London, Special Publications, 297, pp. 203–216. <http://dx.doi.org/10.1144/SP297.9>.
- Dada, S.S., 2008. Proterozoic Evolution of the Nigeria–Borborema Province. *West Gondwana: Pre-Cenozoic Correlations Across the South Atlantic Region*. In: Pankhurst, R.J., Trouw, R.A.J., Brito Neves, B.B., De Wit, M.J. (Eds.), Geological Society of London, Special Publications, 294, pp. 121–136. <http://dx.doi.org/10.1144/SP294.7>.
- Dantas, E.L., 1997. Geocronologia U–Pb e Sm–Nd de terrenos arqueanos e paleoproterozoicos do Maciço Caldas Brandão, NE Brasil. Tese de Doutorado, IGCE/UNESP, Rio Claro, SP. 208 pp.
- Dantas, E.L., Van Schmus, W.R., Hackspacher, P.C., Fetter, A.H., Brito Neves, B.B., Cordani, U., Nutman, A.P., Williams, I.S., 2004. The 3.4–3.5 São José de Campestre massif, NE Brazil: remnants of the oldest crust in South America. *Precambrian Research* 130, 113–137.
- De Wit, M., Stankiewicz, J., Reeves, C., 2008. Restoring Pan-African–Brasiliano connections: more Gondwana control, less Tran-Atlantic corruption. *Geological Society of London, Special Publication*, 294, pp. 399–412. <http://dx.doi.org/10.1144/SP294.20>.
- Dunlop, D.J., 1972. Magnetic mineralogy of unheated and heated red sediments by co-circuity spectrum analysis. *Geophysical Journal of the Royal Astronomical Society* 27, 37–55.
- Guimarães, I.P., Araújo, D.B., Silva Filho, A.F., Silva, F.M.V., Armstrong, R., 2009. Idades U–Pb em zircão por SHRIMP do magmatismo máfico e félsico do Complexo Serrinha-Pedro Velho, Província Borborema, NE Brasil. 23 Simpósio de Geologia do Nordeste, Fortaleza, Abstracts, CD-ROM.
- Hollanda, M.H.B.M., Pimentel, M.M., Jardim de Sá, E.F., 2003. Paleoproterozoic subduction-related metasomatic signatures in the lithospheric mantle beneath NE Brazil: inferences from trace element and Sr–Nd–Pb isotopic compositions of Neoproterozoic high-K igneous rocks. *Journal of South American Earth Sciences* 15, 885–900.
- Hollanda, M.H.B.M., Archânjo, C.J., Souza, L.C., Armstrong, R., Vasconcelos, P.M., 2010. Cambrian mafic to felsic magmatism and its connections with transcurrent shear zones of the Borborema Province (NE Brazil): implications for the late assembly of the West Gondwana. *Precambrian Research* 178, 1–14. <http://dx.doi.org/10.1016/j.precamres.2009.12.004>.
- Hollanda, M.H.B.M., Archânjo, C.J., Souza, L.C., Liu, D., Armstrong, R., 2011. Long-lived Paleoproterozoic granitic magmatism in the Seridó–Jaguaribe domain, Borborema Province – NE Brazil. *Journal of South American Earth Sciences* 32, 287–300. <http://dx.doi.org/10.1016/j.sames.2011.02.008>.
- Jackson, M., 1991. Anisotropy of magnetic remanence: a brief review of mineralogical sources, physical origins and geological applications, and comparison with susceptibility anisotropy. *Pure and Applied Geophysics* 136, 1–28.
- Jardim de Sá, E.F., Legrand, J.M., Galindo, A.C., Sá, J.M., Hackspacher, P., 1986. Granitogênese brasileira no Seridó: o maciço de Acari (RN). *Revista Brasileira de Geociências* 16, 95–105.
- Legrand, J.M., Deutsch, S., Souza, L.C., 1991. Datação U–Pb e granitogênese do maciço de Acari (RN). 14 Simpósio de Geologia do Nordeste, Recife, pp. 172–174.
- Leterrier, J., Jardim de Sá, E.F., Bertrand, J.M., Pin, C., 1994. Ages U–Pb sur zircon de granitoides “brasilianos” de la ceinture du Seridó (Province Borborema, NE Brésil). *Comptes Rendus de l’Académie de Science, Paris. série II t. 318*, 1505–1511.
- Lima, E.S., 1987. Evolução termo-barométrica das rochas metapelíticas da região do Seridó, Nordeste Brasileiro. *Revista Brasileira de Geociências* 17, 315–323.
- Lima, E.S., Pessoa, R.J.R., Accioly, A.C.A., 1989. O metamorfismo de contato causado pelo gabronorito de Totoró, Currais Novos, Rio Grande do Norte, Nordeste Brasileiro. *Revista Brasileira de Geociências* 19, 323–329.
- Ludwig, K.R., 2000. SQUID 1.00. A User Manual. Berkeley Geochronology Center, Special Publication, vol. 2. Berkeley, CA, 17 pp.
- Ludwig, K.R., 2003. Isoplot 3.00: A Geochronological Toolkit for Microsoft® Excel® (revised version) Berkeley Geochronological Center, Special Publication, vol. 4. Berkeley, CA, 70 pp.
- Nascimento, M.A.L., Antunes, A.F., Galindo, A.C., Jardim de Sá, E.F., Souza, Z.S., 2000. Geochemical signature of the Brasiliano-age plutonism in the Seridó belt, Northeastern Borborema Province (NE Brazil). *Revista Brasileira de Geociências* 30, 161–164.
- Omitogun, A.A., Caby, R., Debat, P., 1991. Le métamorphisme pan-africain de la partie centrale de la zone mobile du Nigéria (Ceinture schisteuse d’Igarrá s.l.). *Comptes Rendus de l’Académie de Science, Paris t. 313. Série II*, 1417–1423.
- Ozdemir, O., Dunlop, D.J., Moskowitz, B.M., 1993. The effect of oxidation on the Verwey transition in magnetite. *Geophysical Research Letters* 20 (16), 1671–1674.
- Parente, C.V., Ronchi, L.H., Sial, A.N., Guillou, J.G., Arthaud, M.H., Fuzikawa, K., Verissimo, C.U.V., 2004. Geology and geochemistry of paleoproterozoic magnetite deposits (~1.8 Ga), state of Ceará, Northeastern Brazil. *Carbonates and Evaporites* 19, 28–50.
- Rochette, P., 1987. Magnetic susceptibility of the rock matrix related to magnetic fabric studies. *Journal of Structural Geology* 9 (8), 1015–1020.
- Sá, J.M., McReath, I., Leterrier, J., 1995. Petrology, geochemistry and geodynamic setting of Proterozoic igneous suites of the Orós fold belt (Borborema Province, Northeast Brazil). *Journal of South American Earth Sciences* 8 (3/4), 299–314.
- Souza, L.C., 1996. Zoneographie metamorphique, chimie des minéraux, pétrochimie, géochronologie ⁴⁰Ar/³⁹Ar et histoire P–T–t des micaschistes englobant le massif gabbro-granitique d’Acari (Brasiliano), Ceinture mobile du Seridó (NE du Brésil). Thèse de Doctorat, Laboratoire de Géologie et Minéralogie, Université Catholique de Louvain, Belgique, 345 pp.
- Souza, Z.S., Montel, J.M., Gioia, S.M.L.C., Hollanda, M.H.B.M., Nascimento, M.A.L., Jardim de Sá, E.F., Amaro, V.E., Pimental, M., Lardeaux, J.M., Veschambre, M., 2006. Electron microprobe dating of monazite from high-T shear zones in the São José de Campestre Massif, NE Brazil. *Gondwana Research* 9, 441–455. <http://dx.doi.org/10.1016/j.jgr.2005.11.008>.
- Souza, Z.S., Martin, H., Peucat, J.J., Jardim de Sá, E.F., Macedo, M.H.F., 2008. Calc-alkaline magmatism at the Archean–Proterozoic transition: the Caicó Complex basement (NE Brazil). *Journal of Petrology* 48, 2149–2185.
- Stacey, J.S., Kramers, J.D., 1975. Approximation of terrestrial lead isotope evolution by a two stage model. *Earth and Planetary Science Letters* 26, 207–221.
- Tarling, D.H., Hrouda, F., 1993. The Magnetic Anisotropy of Rocks. Chapman & Hall, 217 pp.
- Van Schmus, W.R., Oliveira, E.P., Silva Filho, A.F., Toteu, S.F., Penaye, J., Guimarães, I.P., 2008. Proterozoic Links Between the Borborema Province, NE Brazil, and the Central African Fold Belt. *West Gondwana: Pre-Cenozoic Correlations Across the South Atlantic Region*. In: Pankhurst, R.J., Trouw, R.A.J., Brito Neves, B.B., De Wit, M.J. (Eds.), Geological Society of London, Special Publications, 294, pp. 69–99. <http://dx.doi.org/10.1144/SP294.5>.
- Zeh, A., Holness, M.B., 2003. The effect of reaction overstep on the garnet microstructures in metapelitic rocks of the Ilesha schist belt, SW Nigeria. *Journal of Petrology* 44, 967–994.

CAPÍTULO 4

TRAMAS DE ROCHAS PARCIALMENTE FUNDIDAS E ALOJAMENTO DE MAGMAS EM REGIMES TRANSCORRENTES

1. Introdução

O anatexito Espinho Branco, detalhado no artigo científico contido neste capítulo, consiste no segundo corpo migmatítico aflorante na região de estudo e ilustra as relações entre magmatismo e deformação no interior de zonas de cisalhamento dúcteis. O migmatito exhibe percentagens variáveis de fusão parcial, englobando metatexitos estromáticos, *schlieren*-diatexitos e leucogranitos. Uma foliação magmática pode ser observada nos metatexitos e localmente nos leucogranitos; nos diatexitos, as altas percentagens de fusão promovem desagregação da estrutura prévia e resultam em rochas quartzo-feldspáticas homogêneas contendo blocos das rochas encaixantes assimilados.

As diferentes percentagens de fusão parcial resultam em leucossomas com geometrias complexas, dificultando a coleta de dados estruturais confiáveis na escala de campo. Dessa forma, um estudo de ASM foi realizado com o objetivo de investigar a cinemática do material parcialmente fundido e a influência da deformação cisalhante no alojamento do migmatito. A trama magnética é relativamente simples, em contraste com a geometria complexa dos leucossomas observada em mesoescala. Contudo, diferentes valores de suscetibilidade magnética (k) podem resultar em padrões distintos de orientações magnéticas, os quais não são diretamente visíveis na trama geral de ASM. Estudos complementares de remanência anisterética (ARA) e trama cristalográfica (EBSD) foram realizados a fim de refinar os dados de ASM.

Os resultados obtidos sugerem que, apesar da complexidade mesoscópica, a trama magnética de migmatitos registra os incrementos deformacionais finais aos quais os leucossomas foram submetidos. Entretanto, em locais onde há variações consideráveis na suscetibilidade, a informação da ASM pode ser inconclusiva; estudos complementares de remanência e orientações preferenciais cristalográficas são necessários para obter uma imagem real da trama interna e avaliar o significado da ASM em anatexitos alojados no interior de zonas de cisalhamento dúcteis.

Fabrics of migmatites and the relationships between partial melting and deformation in high-grade transpressional shear zones: The Espinho Branco anatexite (Borborema Province, NE Brazil)

Luís Gustavo F. Viegas^{a,b,*}, Carlos J. Archanjo^a, Alain Vauchez^b

^a Instituto de Geociências, USP, rua do lago 562, 05508-080 São Paulo, SP, Brazil

^b Géosciences Montpellier, Place E. Bataillon, 34095 Montpellier Cedex 5, France

ARTICLE INFO

Article history:

Received 10 July 2012

Received in revised form

13 December 2012

Accepted 17 December 2012

Available online 27 December 2012

Keywords:

Migmatite

AMS

EBS

Borborema Province

Petrofabrics

NE Brazil

ABSTRACT

The Espinho Branco anatexite, located within a transcurrent, high-temperature shear zone in NE Brazil, was the subject of a comprehensive petrostructural study (Anisotropy of Magnetic Susceptibility – AMS, Anisotropy of Anisotropic Remanence – AAR, Electron Backscatter Diffraction – EBSD) to evaluate the compatibility of different fabrics with the kinematics of melt deformation. Magnetite dominates susceptibilities larger than 1 mSI and biotite displays [001] lattice directions consistent with AMS k_3 axes. In contrast, migmatites with a susceptibility lower than 0.5 mSI and no visible mesoscopic foliation provide crystallographic fabrics distinct from AMS and AAR. However, AAR remains consistent with the regional strain field. These results suggest that the correlation of field, AMS and crystallographic fabrics is not always straightforward despite the relatively simple organisation of the magnetic fabric in the anatexite. We conclude that AMS recorded the final stages of the strain field in the migmatite irrespective of its complex mesoscale structures and contrasting crystallographic fabrics.

© 2012 Elsevier Ltd. All rights reserved.

1. Introduction

Migmatites are composite igneous and metamorphic high-grade rocks that record crustal flow processes at the roots of orogens (Whitney et al., 2004). Migmatite petrology gives insights into the composition and differentiation of the middle to upper crust, while migmatite structures register the deformation that is active during orogenesis (Ashworth, 1985).

The petrostructural characteristics of migmatites have been studied for over than thirty years (Mehnert, 1968). Quantitative (Blumenfeld and Bouchez, 1988; Leitch and Weinberg, 2002), and qualitative (Brown and Rushmer, 1997; Weinberg and Mark, 2008) work was extensively employed, encompassing field mapping, melt topology and fabric analysis to characterise the structure and reconstruct the emplacement and strain history and their relationships with large-scale crustal deformation. However, the inherently complex geometry at the outcrop scale constantly renders tectonic interpretations uncertain or ambiguous. Due to both their igneous and metamorphic nature, migmatites behave as

two-phase materials enhancing strain localisation in the liquid (magma) phase and promoting strain hardening in the solid (host rock) phase; textures are commonly divided into solid-state in the metamorphic host and magmatic microstructures in the leucosomes or magma (Vigneresse et al., 1996; Vernon, 2000).

More recently, new methodologies were tested with the aim of shedding some light onto deformation patterns in migmatitic bodies. Ferré et al. (2003) pioneered an AMS study on anatexites and concluded that the apparent structural complexity at the mesoscale masks a simple magmatic flow pattern that can be mapped in detail using magnetic fabrics. This methodology was followed by other workers (Denèle et al., 2007; Charles et al., 2009; Archanjo et al., 2012) and coupled with crystallographic preferred orientation (CPO) measurements of rock-forming minerals through electron backscatter diffraction (Kruckenberg et al., 2010). These tools proved useful in establishing correlations between magnetic and mesoscale structural fabrics.

Migmatites emplaced in high-strain zones usually retain fabrics consistent with the regional strain field (Brown, 1994; Paterson et al., 1998). However, the exact chronology between partial melting and deformation is difficult to establish due to the complex crosscutting relationships between the foliations and the mesoscopic melt pockets (Rutter and Neumann, 1995; Rosenberg and Handy, 2005).

* Corresponding author. Géosciences Montpellier, Place E. Bataillon, 34095 Montpellier Cedex 5, France.

E-mail addresses: lviegas@usp.br, viegas@gm.univ-montp2.fr (L.G.F. Viegas).

This paper presents a comprehensive petrostructural study on migmatites outcropping in the high-grade, transcurrent Patos shear zone located in the Borborema Province, northeast Brazil. This shear zone constitutes a major Neoproterozoic structure regarded as a crustal boundary dividing different tectonic terranes (Vauchez et al., 1995; Van Schmus et al., 2008). The shearing is associated with partial melting, and migmatites display contrasting morphologies, comprising foliation-parallel leucosome veins, randomly oriented nebulites and isotropic leucogranites. Combining field studies with magnetic (AMS, AAR) and crystallographic fabric (EBSD) measurements, we show that migmatite internal structures can be mapped with good consistency throughout the shear zone even when melt pockets accumulate with no visible mesoscopic fabric. Furthermore, our study shows that detailed mapping of the fabric of anatexites emplaced in shear zones is needed before placing constraints on the melting and deformation relationships in hot orogens.

2. Geological setting

The Patos shear zone consists of a ~600 km E-trending strike-slip shear zone that deforms the Precambrian rocks of the Borborema Province (Fig. 1). It forms part of a continental-scale shear system that can be followed from NE Brazil to West Africa (Vauchez et al., 1995; Arthaud et al., 2008; De Witt et al., 2008). Three major structural domains can be defined in the Patos shear zone. The Central Domain, located between the towns of Catingueira and Patos (Fig. 2), comprises E-trending mylonitic gneisses approximately 30 km in width in structural continuity with the N-NE fabric of the Seridó belt. This led Corsini et al. (1991) to argue that the Patos-Seridó structure forms a mechanically coupled system in which transcurrent displacements are transferred to a transpressional belt.

The Western Domain consists of a duplex structure where NE-trending lenses of orthogneisses, metapelites and granitoids are bounded by E-trending mylonites that merge with the western

NE-trending Senador Pompeu shear zone. This suggests that the Borborema shear zones form a crustal-scale branched system (Vauchez et al., 1995; Oliveira, 2008). The Eastern Domain displays a gradual rotation of the mylonitic foliation from E-W to NE-SW, close to the Atlantic coastal deposits. The shear zone foliations are mainly subvertical, and stretching lineations are subhorizontal, which is consistent with a dominant transcurrent dextral motion along the entire zone (Corsini et al., 1991).

Anatexites are frequent in the Central Domain, forming lenses of granitic domains mixed with “unmelted” material. They display folded stromatic, schollen, schlieren and boudinage structures and derive mostly from the melting of the Paleoproterozoic basement. A narrow medium-to-low temperature mylonite belt outlines the southern margin of the shear zone (Fig. 2) and reworks the orthogneisses and migmatites under lower amphibolite-green schist facies conditions. Although they have the same fabric orientation and kinematics, it is not clear whether the high- and low-grade mylonites are coeval or were formed at different reworking episodes.

Available geochronological studies of the Patos mylonites include zircon U/Pb (TIMs) data and Sm/Nd model ages obtained in the Central Domain (Costa, 2002). The zircons show strong isotopic discordance but provide unconstrained upper intercept ages ranging from 2.2 Ga to 2.0 Ga, indicating that a Paleoproterozoic protolith was involved in deformation and migmatization. Whole-rock Sm/Nd model ages range from 3.4 Ga to 2.6 Ga and also suggest the presence of Archean sources. Ar^{40}/Ar^{39} ages range from 540 Ma to 490 Ma and are attributed to final cooling and late exhumation of the shear zone (Monié et al., 1997; Corsini et al., 1998).

3. The Espinho Branco anatexite: field characteristics

The Espinho Branco anatexite, located in the Central Domain, occupies an elliptical area of ~25 km² elongated in the E-W direction. This migmatitic zone can be traced in structural

Fig. 1. The Neoproterozoic Borborema shear zone system in NE Brazil. The box shows the study area located within the Patos shear zone.

Fig. 2. Geological map of the connection zone between the Patos shear zone and the Seridó belt.

continuity with the NE-trending Santa Luzia diatexite (Fig. 2). The mineralogical assemblage comprises K-feldspar, quartz, plagioclase and biotite, with amphibole, zircon, titanite and opaques as accessory minerals. It can be divided into three lithological facies that are mainly distinguished by the degree of partial melting and leucosome geometry (Figs. 3 and 4): i) stromatic metatexites, in which the leucosomes are mostly parallel to the mylonitic foliation (Fig. 4a, b); ii) nebulitic schlieren-diatexites with a higher melt fraction and nearly complete disaggregation of compositional layering (Fig. 4d, e) resulting in an homogeneous quartz-feldspathic rock with a magmatic texture; and iii) biotite-leucogranites closely associated with diatexite (Fig. 4f). The contacts between these units are gradual, with no evidence of overprint, except for the leucogranites that may truncate the foliation of metatexites and diatexites.

Shape-preferred orientations of centimetre-scale K-feldspar grains and biotite flakes up to 2 mm define a mesoscopic steeply dipping magmatic foliation (Fig. 3). This fabric is marked by the preferred orientation of the leucosome in the metatexites (Fig. 4a, b) and by the alignment of biotite flakes in the leucogranites. In the diatexites, the higher melt fraction erases the previous foliation and leaves a faint orientation of metric-size mafic lenses parallel to the E-trending magmatic flow. The leucosome defines an interconnected network in which dextral shear sense indicators can be deduced (Fig. 4c). In the western, central and eastern portions of the migmatite body, the leucosome is locally collected into SSW–NNE and NW–SE mesoscale magmatic shear zones. These structures show both synthetic and antithetic shear senses with respect to the Patos shear zone and act as zones of melt channelling throughout the anatexite (Fig. 3, Fig. 4d).

Fig. 3. Lithological facies, mesoscale structural pattern and cross-sections in the Espinho Branco anatexite. The stereograms correspond to foliations measured in the field. Kamb density contours, lower hemisphere. The geographical coordinates are given in UTM units.

Fig. 4. The field aspects of the Espinho Branco migmatite at increasing melt fractions: a) and b) stromatic metatexites showing parallelism of leucosome veins with the host rock foliation; c) syntectonic melt pockets within leucosome bands; d) complex geometrical patterns in metatexite with increasing melt fraction; e) disaggregation and assimilation of host rocks as xenoliths in diatexites; f) contact between metatexite and leucogranite.

3.1. Microstructures and melt segregation at the grain-scale

Vernon (2000), Rosenberg (2001) and Sawyer (2010) revised the major microstructural criteria for identifying deformation mechanisms in melt-bearing systems. In this study, microstructural observations are focused in the leucosome that forms the direct product of partial melting (Brown, 2001; Sawyer, 2008).

The metatexites show a granular texture consisting of up to 1–2 mm quartz and feldspar grains displaying lobate to straight boundaries (Fig. 5a). The microcracks in K-feldspar may be filled with new quartz crystallised as small “drops” (arrows in Fig. 5b). The coarse quartz grains show undulatory extinction, subgrains and fill embayments in plagioclase. The fine-grained quartz is usually free of substructure. The K-feldspar displays local undulatory extinction and myrmekites at K-feldspar-quartz-plagioclase boundaries. The plagioclase occurs as medium-sized (0.5 mm) subhedral grains in contact with quartz lobes (Fig. 5a).

The diatexites display coarse quartz grains (~200 µm) forming “pools” at quartz-feldspar-plagioclase triple junctions (Fig. 5c, d). The quartz may have lobate boundaries towards K-feldspar (Fig. 5c) and also displays local undulatory extinction. The plagioclase grains are sometimes fractured along the acute faces of their internal zoning, resulting in small fragments surrounded by newly crystallised quartz (Fig. 5d). Such microstructures suggest deformation by fracturing, possibly induced by melt percolation and subsequent quartz crystallisation in dilatant sites within the plagioclase crystals (Brown and Rushmer, 1997; Rosenberg and Handy, 2005).

The leucogranites comprise coarse quartz (~2 mm) grains displaying subgrains and lobate to straight boundaries (Fig. 5e). The quartz may form large-sized (~500 µm) elongated crystals parallel to the foliation and also bordering K-feldspar porphyroclasts (Fig. 5f). These elongated grains show undulatory extinctions and may locally display subgrain walls normal to the grain length. The K-feldspar crystals may display anhedral shapes with microcracks

Fig. 5. The microstructures indicative of partial melting at increasing melt fractions: a) and b) interstitial quartz at grain boundaries or filling microcracks in metatexites (arrows); c) and d) interstitial newly crystallised quartz at triple junctions in diatexites (arrows). Note the plagioclase crystal fractured along the acute angle of internal zoning in Fig. 5d; e) microfractures normal to the foliation plane (oriented WNW-ESE) in leucogranite; f) elongate quartz grains surrounding a fractured K-feldspar porphyroblast. The quartz show subgrains parallel to fractures in the feldspar. All photos are in crossed polarisers. The shear sense is dextral. Abbreviations: Qtz - quartz; kfs - K-feldspar; pl - plagioclase; Bt - biotite; Hb - hornblende.

and local myrmekite exsolutions at the grain boundaries. The plagioclase occurs as coarse subhedral grains with local undulatory extinction and deformation twins.

4. Anisotropy of magnetic susceptibility (AMS) and anhysteretic remanence (AAR)

4.1. Sampling and methods

Magnetic studies were carried out in the granitic leucosome and in the intruding leucogranites of the Espinho Branco anatexite. Because leucosomes do not always show grain shapes in a preferred orientation at the mesoscale, this technique was employed to investigate magma flow directions under partially molten conditions (Bouchez, 2000; Ferré et al., 2003). Oriented cores were collected at 49 sites

(Fig. 6). At each site, at least 3 cores of ~8 cm in length and 2.5 cm in diameter were extracted with a gasoline-powered portable rock drill. The samples were later cut into 2.2-cm-long pieces, yielding approximately 5–7 specimens per site. A total of 331 specimens were available for the magnetic study. Low-field AMS was measured on a KLY-4S Kappabridge (AGICO, 300 A/m, Ac field at 920 Hz). The mean susceptibility directions of the AMS tensor were calculated using Jelinek (1978) statistics, which provide the main directions ($k_1 \geq k_2 \geq k_3$) of the magnetic susceptibility ellipsoid. These data are summarised in Table 1.

Anisotropy of anhysteretic remanence (AAR) is especially suited for the investigation of ferromagnetic fabrics because it isolates the contribution of phases that provide the remanence (Jackson, 1991; Borradaile and Jackson, 2004). AF demagnetisation and anhysteretic remanence acquisition were performed with a LDU-AMU

Fig. 6. The distribution of AMS sampling sites, susceptibility and anisotropy frequency histograms in the Espinho Branco anatexite.

Table 1

The AMS parameters for the Espinho Branco migmatite.

Site	Locations		n	k (mSI)	P	T	k1			k3		
	UTM W	UTM S					Dec	Inc	$\alpha K1$	Dec	Inc	$\alpha K3$
PA01	685,544	9,221,757	10	0.08	1.06	-0.2	21	84	26.3	95	88	57.7
PA02	686,323	9,222,428	6	21.7	1.42	-0.497	282	6	16.3	202	33	19.1
PA03	687,490	9,221,686	9	0.07	1.07	0.457	267	31	62.9	179	87	43.9
PA04	687,229	9,221,291	10	10.7	1.19	0.214	267	19	17.7	354	80	17.9
PA05	686,941	9,219,735	14	5.83	1.24	-0.681	250	26	23.2	207	34	60
PA06	687,151	9,218,987	7	9.14	1.37	-0.095	223	28	16.3	229	28	10.6
PA07	687,641	9,219,970	10	6.07	1.28	-0.434	218	4	17.8	133	36	23.1
PA08	688,370	9,221,455	11	0.02	1.04	-0.203	127	49	26.7	206	80	41.9
PA09	690,435	9,221,544	7	12.6	1.28	-0.245	262	13	16.8	349	78	30.8
PA10	692,569	9,220,995	5	9.95	1.13	-0.074	280	3	32.5	190	84	27.9
PA11	693,606	9,220,046	12	1.85	1.22	0.131	287	11	15.5	16	86	9.1
PA12	694,271	9,219,131	8	6.33	1.41	-0.698	250	21	14.3	176	54	42.5
PA13	695,471	9,219,268	10	10.7	1.14	-0.551	260	20	30.5	177	71	63.6
PA14	695,938	9,220,261	7	4.41	1.57	0.526	113	68	31.5	188	85	23.2
PA15	677,270	9,221,855	7	0.14	1.04	0.087	205	39	37.2	116	88	20.5
PA16	676,203	9,222,121	10	0.62	1.11	0.047	72	8	18.5	1	24	28.5
PA17	679,446	9,220,945	5	0.53	1.09	0.728	34	7	59.1	304	86	15
PA18	679,573	9,219,392	6	5.69	1.38	-0.479	264	7	12.3	188	27	15
PA19	680,098	9,218,627	7	9.58	1.18	0.496	231	33	55.6	184	43	30.9
PA20	683,796	9,218,353	10	3.66	1.07	-0.61	246	29	25	297	42	55.2
PA21	686,116	9,218,631	5	11.8	1.27	-0.662	248	7	6.9	216	9	60
PA22	696,810	9,219,840	7	12.9	1.2	-0.357	279	17	9.9	253	18	30.2
PA23	697,767	9,219,593	7	1.36	1.08	-0.371	241	10	39.8	155	69	36.8
PA24	698,707	9,219,650	5	5.08	1.17	0.125	247	25	18.7	187	43	17.4
PA25	702,831	9,214,359	5	0.46	1.04	0.853	335	35	80.4	315	36	42.5
PA26	702,458	9,215,562	7	0.42	1.08	-0.513	247	37	34.9	167	77	66.2
PA27	698,360	9,218,361	7	20.8	1.14	0.754	273	43	53.6	222	56	20.4
PA28	699,518	9,216,907	6	4.24	1.25	-0.708	230	24	14.1	158	55	50.6
PA29	700,298	9,216,148	5	11.2	1.32	-0.315	62	10	38.4	335	73	26.4
PA30	701,229	9,214,873	6	0.15	1.1	0.383	195	4	60.4	282	57	22.4
PA31	696,935	9,218,060	7	5.75	1.32	0.04	239	26	20.9	222	27	21
PA32	696,018	9,218,199	7	33.3	1.35	-0.24	259	30	30.7	212	40	22.3
PA33	685,123	9,220,631	8	16	1.88	0.025	249	14	19.6	168	61	14.6
PA34	681,598	9,220,934	6	4.19	1.45	0.059	258	13	11.8	199	24	19.7
PA35	680,384	9,220,379	6	42.1	1.48	0.025	253	30	14.4	306	45	20.6
PA36	688,004	9,222,233	7	20.5	1.37	-0.811	318	22	24.8	45	82	29.2
PA37	687,113	9,221,782	8	18.5	1.23	-0.735	279	10	14.3	194	64	57.2
PA38	686,356	9,221,677	6	0.12	1.05	-0.509	265	51	21.5	280	52	60.8
PA39	685,489	9,220,514	5	13	1.19	-0.292	257	19	35.4	169	84	42.2
PA40	683,315	9,220,244	6	15.1	1.49	-0.414	266	27	13.4	205	45	15.7
PA41	682,094	9,220,131	7	2.92	1.24	-0.747	254	24	6.1	328	58	47.1
PA42	681,667	9,218,643	6	8.14	1.45	-0.129	262	7	15.9	174	79	17.6
PA43	690,481	9,219,196	5	0.83	1.05	-0.725	5	2	38.2	92	43	66.5
PA44	691,999	9,218,144	6	20.6	1.17	0.684	222	12	64	151	34	32.4
PA45	693,351	9,217,204	11	15.3	1.29	-0.379	250	4	18.7	163	56	46.8
PA46	694,998	9,216,611	6	2.37	1.22	-0.028	228	10	8.2	151	42	10.2
PA47	700,637	9,217,451	6	9.11	1.3	-0.672	250	16	12.5	7	33	61.7
PA48	696,224	9,215,879	6	0.16	1.06	0.483	247	29	44.4	101	35	21.5
PA49	684,273	9,221,331	6	12.5	1.29	-0.261	267	7	10.3	156	21	24.6

n = number of specimens; k (mSI), mean-site susceptibility; P, anisotropy degree; T, shape parameter, k₁ and k₃, orientations of AMS main directions (declination and inclination); αk_1 and αk_3 are the angular dispersion by the maximum semi-angle (degrees) of the confidence cone around the mean direction.

demagnetiser/magnetiser, and the remanence was measured with a JR6A magnetometer (AGICO) housed in a magnetic field-free room in the Laboratório de Paleomagnetismo, University of São Paulo. The specimens were AF demagnetised at peak AF fields of 100 mT. The AAR parameters (magnitude, orientation) were determined after anhysteretic remanence acquisition along six different orientations with a peak field of 50 mT and a biasing field of 100 μ T (Trindade et al., 2001). Statistics were obtained using the method of Jelinek (1978), implemented in the ANISOFT program package (Hrouda et al., 1990).

Magnetic mineralogy experiments were performed by isothermal remanent magnetisation (IRM) acquisition and by determining the temperature dependence of the magnetic susceptibility (k – T) using a KLY-4S Kappabridge connected to a CS-3 furnace. The acquisition of IRM was obtained through successive increments of a steady-magnetic field (0.001 T–2.0 T) and was measured with a Molspin magnetometer. The stepwise k – T curves were obtained from –192° to 700 °C and performed in an argon environment to prevent sample oxidation.

4.2. Magnetic susceptibility and anisotropy

The bulk susceptibility ($k = 1/3[k_1 + k_2 + k_3]$) ranges from 2.00 to 128.22 mSI, and the anisotropy degree ($P = k_1/k_3$) from 1.01 to 2.91. Low susceptibilities ($k < 0.5$ mSI) are recorded in 30% of the specimens and high susceptibilities ($k > 10$ mSI) are recorded in ~27% of them. A mean susceptibility of 7.9 mSI (SD 10.38) and an anisotropy degree of 1.31 (SD 0.28) are observed throughout the migmatite. The IRM acquisition and the k – T curves, given in Fig. 7,

attest to the contribution of ferromagnetic minerals irrespective of the susceptibility magnitude.

Specimens with moderate to high susceptibilities ($1 < k < 10$ mSI) comprise more than 60% of the sites. The IRM and the k – T curves are very simple. The steep initial IRM slope indicates the presence of a soft fraction that saturates after 200 mT (Fig. 7). The k – T curve shows a well-defined Verwey transition at ~150 °C and a net susceptibility decrease at approximately 590 °C. These parameters indicate the presence of a low-coercive, coarse Ti-poor magnetite. Therefore, the susceptibilities depend essentially on the content of magnetite.

In specimens with susceptibilities lower than 1 mSI (38% of the sites), a steep initial slope in the IRM curve also indicates the presence of a “soft” coercive fraction (Fig. 7b and c), but the remanence does not saturate at high-fields, revealing an additional “hard” coercive fraction. The k – T curves in these rather low-susceptibility specimens display a well-defined Verwey transition (Fig. 7b). The susceptibility remains relatively constant during heating but decreases just below 600 °C and vanishes totally at ~700 °C. These properties point to the presence of both Ti-poor magnetite, which shows a Curie temperature of 580 °C, and haematite that shows a Néel temperature of 680 °C. Haematite must be responsible for the hard coercive fraction detected in the IRM curves. The absence of the Verwey transition in the samples with very weak susceptibility suggests the presence of a fine and oxidised magnetite (Özdemir and Dunlop, 1993; Muxworthy and McClelland, 2000). New ferrimagnetic phases are observed to appear during heating, as indicated by the net susceptibility increase between 450 and 600 °C (Fig. 7b and c).

Fig. 7. The isothermal remanent magnetisation (IRM) acquisition and temperature dependence of the magnetic susceptibility for three selected specimens.

Fig. 8. The AMS-AAR magnetic fabric ellipsoids in the Espinho Branco migmatite. Kamb density contours, lower hemisphere, equal-area projections. Symbols: k_1 = square; k_2 = triangle; k_3 = circle; n = number of specimens.

4.3. Anisotropy of anhysteretic remanence results

The AAR anisotropy degree (P_{AAR}) is usually higher than that of AMS (Fig. 8). Although differences in the shape ratios ($T = [2(\ln k_2 - \ln k_3)/(\ln k_1 - \ln k_3)] - 1$) of the AAR and AMS ellipsoids are recorded, their principal directions remain close to each other, mainly in sites with moderate to high susceptibilities (Fig. 8a). However, in sites having very low susceptibilities, the shape ratios can be quite different, most likely due to the contribution of paramagnetic silicates to AMS. In site PA17 (Fig. 8b) for instance, AMS shows a typical oblate fabric ($T = 0.73$), while AAR displays a prolate shape ($T = -0.32$). In some cases, AMS and AAR are well-defined, but their respective orientations are different (Fig. 8c). Similar results were also found in the magnetic fabrics of the Santa Luzia nebulite (Archanjo et al., 2012) located in the northeast of the Espinho Branco migmatite (Fig. 2).

In summary, AAR indicates that magnetite is the dominant carrier of AMS in the studied migmatite, although biotite may contribute to the anisotropy in specimens having very low susceptibilities. In such sites, magnetite observed under the microscope occurs as fine grains usually hosted in silicates. The fact that the susceptibility and remanence anisotropies are close in orientation discards an inverse AMS due to the very fine single domain magnetite grains (Jackson, 1991).

4.4. Magnetic fabric

The magnetic lineations are well-organised (Fig. 9a), contrasting with the apparently disordered leucosome orientations in the field. The mean-site magnetic lineations (k_1 , Table 1) plunge gently to the SW and WSW, oblique to the migmatite elongation axis (Fig. 9). The magnetic foliation (poles to k_3 , Fig. 9b) defines a NNW-SSE trending girdle with shallow to intermediate dips that rotates around a zone axis formed by the magnetic lineation. To compare the structural interpretations

Fig. 9. The magnetic fabric maps (foliation, lineation) and pole diagrams of the Espinho Branco anatexite. Kamb density contours, lower hemisphere.

between field foliations and magnetic fabrics, crystallographic preferred orientations were determined by electron backscatter diffraction (EBSD).

5. Crystallographic preferred orientations

5.1. Methods

Representative samples were chosen for EBSD analysis of biotite crystals in detail because they are the main mafic silicate and tend to orient themselves conformably with the magmatic flow.

Each sample, cut in the XZ section of the AMS ellipsoid, was carefully polished. The orientation measurements were performed using a JEOL JSM 5600 scanning electron microscope (SEM) equipped with an Oxford Instruments/HKL Nordlys EBSD detector at Géosciences Montpellier. The samples were inserted in the microscope chamber at an angle of 70° relative to the electron beam and at a working distance of 25 mm. An accelerated electron beam of 17 kV and a spot size of 75 µm were used to generate diffraction bands on a phosphorous screen. The acquisition was made automatically on a regular grid with step sizes from 25 to 50 µm depending on grain size. The Kikuchi bands were identified and then indexed using the Channel 5 Program (Oxford Instruments). Finally, the orientation measurements were rotated in the geographical frame.

5.2. Lattice preferred orientation results

The biotite fabrics show a good correlation with the magnetic axes and, to some extent, with the field foliations (Fig. 10). In most

samples, the AMS short axis (k_3) is close to the [001] direction. In addition, the [001] direction may be in an asymmetrical position in relation to the pole of the field foliation (Fig. 10, PA14). The AMS k_1 long axis is observed close to {100} poles and may locally correspond to a maximum in the [001] direction (Fig. 10, PA17). The (010) poles configure NW-SE, E-W and NE-SW trending girdles that may contain the k_1 and k_2 AMS axes, with the pole of the girdle being the AMS k_3 direction (Fig. 10).

6. Discussion

6.1. Deformation mechanisms of the Espinho Branco anatexite

The overall microstructure indicates that deformation occurred in the magmatic state. This is supported by several lines of evidence (Fig. 5): i) interstitial quartz located at feldspar boundaries; ii) microcracks in feldspar often sealed by fine-grained quartz; iii) newly crystallised strain-free quartz; iv) quartz grains as “pools” at the quartz-feldspar triple junctions. The dominance of interstitial quartz with little evidence of solid-state deformation indicates that the external strain field was not imprinted in quartz due to post-deformation crystallisation (Rosenberg, 2001).

The biotite fabrics are marked by [001] concentrations close to Z, which suggest a preferred orientation of platy crystals in the flow plane. These patterns, consistent with biotite fabrics in shear zones, can be attributed to rigid-body rotation of biotite crystals in magmatic flow (Nicolas and Poirier, 1976). This process is consistent with the obliquity of [100] directions with respect to the magmatic foliation, in agreement with the dextral shear sense observed in the Patos shear zone.

Fig. 10. The crystallographic preferred orientations of biotite. The lower hemisphere pole figures, equal area stereographic projection, contours at 1, 1.5, 2, 2.5, 3% etc. by 1% area. The magnetic data were plotted in the stereogram to allow fabric comparisons. Filled symbols: AMS k_1 , k_2 , k_3 axes; empty symbols: AAR k_1 , k_2 , k_3 axes. N = number of measurements (one point per grain). J = fabric strength. The mean field foliation is represented by the dashed line. The pole figures were rotated to the geographical frame with the peripheries of the stereonet (Z and X) corresponding to the vertical N-S and horizontal E-W directions, respectively.

6.2. Magnetic fabrics and relationships with field data

The comparisons between AMS and AAR indicate that, in the low susceptibility ($k < 0.5$ mSI) specimens, the anisotropy of Fe-bearing silicates (possibly associated with small inclusions of iron oxides, such as haematite) can account for the dispersion of the anisotropy directions. However, AAR still confirms the presence of magnetite as fine grains dominating the magnetic anisotropy (Fig. 8).

The AMS lineations are consistent through most of the migmatite body. The local differences in orientation are not reproduced in AAR, which maintains the observed orientation across the whole anatectic domain. These results indicate that magnetite is mainly responsible for the magnetic anisotropy even in samples where the AMS is disorganised. When the magnetite contribution to the bulk magnetic fabric decreases in volume, the AMS directions may become locally dispersed (e.g., at the specimen scale) and inconsistent with the overall orientation pattern. However, these are local perturbations and do not account for a global modification of magnetic directions at the scale of the migmatite body.

Well-constrained field foliations tend to have coaxial AMS-AAR, which is consistent with the kinematics of the shear zone (sample PA14 in Fig. 11). In addition, the magnetic fabric of melt pockets records the same strain field as the rest of the migmatite. However, local inconsistencies are present, as in site PA03, where the AMS k_1 axis is obliquely opposing the AAR k_1 axis (Fig. 11).

When the melt fraction increases gradually and promotes disaggregation of the structural fabric, field foliations may become incompatible with AMS and AAR. In site PA41, the geometrical complexity of the migmatite structure is not reproduced in the magnetic fabric, which tends to organise itself in agreement with the anatectic strain field (Fig. 11).

These observations therefore allow us to conclude that: i) in samples with susceptibilities higher than 1 mSI, AMS may be correlated with the rock structure and the regional shear sense directions; ii) at low susceptibilities ($k < 0.5$ mSI), AMS does not match the field foliation. A possible origin for the AMS discrepancy might be the formation of haematite during late hydrothermal alteration. Haematitisation effects may account for the lowering of bulk susceptibilities, scattering of magnetic fabrics and

Fig. 11. The synthetic view of selected samples with different fabric patterns in the Espinho Branco migmatite. Filled symbols: AMS k_1 , k_2 , k_3 axes; empty symbols: AAR k_1 , k_2 , k_3 axes. The mean field foliation is represented by the dashed line. N = number of measurements; J = fabric strength. See text for discussion.

development of a new preferred orientation of fine grains mimicking the fluid pathways (Trindade et al., 2001; Just et al., 2004; Archanjo et al., 2009). However, it must be stressed that the anomalous AMS directions concern only 18% of the sites. The dominant, well-defined magnetic fabric of the migmatite records the deformational processes that occur during emplacement.

6.3. Relationships between field, magnetic and crystallographic fabrics

In high susceptibility specimens ($k > 1$ mSI), magnetic and lattice fabrics are consistent with the field foliation at places where it can be reasonably mapped (PA14 in Fig. 11). In contrast, in sites where the field foliation is poorly constrained due to the increasing melt fractions (PA41 in Fig. 11), no relationship can be observed between magnetic and crystallographic fabrics and field foliations. However, biotite [001] directions still agree with the AMS k_3 axis.

These observations suggest that: i) the magnetic foliation describes the biotite foliation of the anatexite quite well, while k_1 marks the lineation defined by magnetite grains; ii) the high-susceptibility specimens ($k > 1$ mSI, ~75% of the sites) usually display well-defined magnetic fabrics that correlate with crystallographic fabrics, giving reliable information on magma flow regardless of the observed mesoscopic fabric; iii) a combination of magnetic and crystallographic fabrics is efficient for migmatite fabric determination when foliation is hard to measure in the field.

The low-susceptibility specimens ($k < 0.5$ mSI) are related to sites where no visible structure is present except for mesoscopic shear zones (PA03 and PA16, Fig. 11). At such sites (#17, #25, #30, Table 1), the magnetic axes are either dispersed or clustered but do not necessarily correlate with the strain field. Oblique AMS-AAR directions can develop (Fig. 8c; PA03 in Fig. 11), confirming the presence of sub-fabrics in these sites. These inconsistencies are attributed to late hydrothermal oxidation processes affecting magnetite. In such cases, structural interpretations based on AMS may be misleading.

Hence, we postulate that the correlation between the three fabrics analysed (field, magnetic, crystallographic) provides reliable results when there is a good record of field foliations, as already demonstrated in previous studies (Ferré et al., 2003; Kruckenberg et al., 2010). However, when there is no straightforward correlation between the aforementioned fabrics, AMS alone may not truly represent the magmatic flow, notably in sites with low susceptibilities. Additional AAR measurements and their subsequent comparison with lattice fabrics should be performed to better constrain synkinematic melt deformation patterns in high-grade shear zones.

7. Conclusions

The well-defined AAR subfabrics prove that the low-field AMS of the migmatite is controlled by the anisotropy of magnetite. The well-defined magnetic foliation dips dominantly to the south and the subhorizontal lineation is oblique to the E-trending body elongation, making a map-scale dextral sigmoid. The biotite crystallographic fabrics agree with AMS, notably its [001] axes concentrated in a direction close to the magnetic foliation pole (k_3).

Our results reveal a relatively simple magnetic fabric pattern, despite the structural and compositional complexity of the migmatite. They indicate that AMS is independent of mesoscale heterogeneities and confirm that magnetic fabrics are recorded in the neosome during the latest stages of synmagmatic shear deformation. However, magnetic fabrics may be distinct from the regional strain field at very low susceptibilities (<0.5 mSI), primarily if late oxidising fluids interact with the magnetic phases.

In such cases, comprehensive studies encompassing AAR and CPO should be employed to detect subfabrics and evaluate the significance of AMS data.

Acknowledgements

L. G. Viegas and C. Archanjo thank Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP, grants 2010/50060-1 and 2009/17537-1) and the Brazilian research Council (CNPq, grant 200496/2011-5) for financial support. This paper is part of the first author's PhD thesis. We thank John Rico and Camilo Bustamante for assistance during field trips. Danielle Brandt, Giovanni Moreira (Lab. Paleomagnetismo) and Fabrice Barou (Géosciences Montpellier) are thanked for their invaluable help during the magnetic and EBSD laboratory measurements, respectively. Eric Ferré and Jean Luc Bouchez are thanked for constructive reviews that greatly improved the manuscript.

References

- Archano, C.J., Launeau, P., Hollanda, M.H.B.M., Macedo, J.W.P., Liu, D., 2009. Scattering of magnetic fabrics in the Cambrian alkaline granite of Meruoca (Ceará state, northeastern Brazil). *International Journal of Earth Sciences* 98 (8), 1793–1807.
- Archano, C.J., Viegas, L.G., Hollanda, M.H.B.M., Souza, L.C., Liu, D., 2012. Timing of the HT/LP transpression in the Neoproterozoic Seridó Belt (Borboroma Province, NE Brazil): constraints from U–Pb (SHRIMP) geochronology and implications for the connections between NE Brazil and West Africa. *Gondwana Research*, <http://dx.doi.org/10.1016/j.gr.2012.05.005>.
- Arthaud, M.H., Cabry, R., Fuck, R.A., Dantas, E.L., Parente, C.V., 2008. Geology of the Northern Borboroma Province NE Brazil and Its Correlation with Nigeria, NW Africa. In: Geological Society of London, Special Publication, vol. 294, pp. 49–67.
- Ashworth, J.R., 1985. *Migmatites*. Blackie, Glasgow.
- Blumenfeld, P., Bouchez, J.L., 1988. Shear criteria in granite and migmatite deformed in the magmatic and solid states. *Journal of Structural Geology* 10, 361–372.
- Bouchez, J.L., 2000. Anisotropie de susceptibilité magnétique et fabrique des granites, vol. 330. *Comptes Rendus de l'Académie des Sciences de Paris*, pp. 1–14.
- Borradaile, G.J., Jackson, M., 2004. Anisotropy of magnetic susceptibility (AMS): magnetic petrofabrics of deformed rocks. In: Martin-Hernandez, F., Lünenburg, C.M., Aubourg, C., Jackson, M. (Eds.), *Magnetic Fabric: Methods and Applications*. Geological Society of London, Special Publication, vol. 238, pp. 299–360.
- Brown, M., 1994. The generation, segregation, ascent and emplacement of granite magma: the migmatite- to -crustally-derived granite connection in thickened orogens. *Earth Science Reviews* 36, 83–130.
- Brown, M., 2001. Orogeny, migmatites and leucogranites: a review. *Proceedings of the Indian Academy of Sciences (Earth and Planetary Science)* 110, 313–336.
- Brown, M., Rushmer, T., 1997. The role of deformation in the movement of granite melt: views from the laboratory and the field. In: Holness, M.B. (Ed.), *Deformation-enhanced Fluid Transport in the Earth's Crust and Mantle*. The Mineralogical Society Series 8. Chapman & Hall, London, pp. 111–144.
- Corsini, M., Vauchez, A., Archanjo, C.J., Jardim de Sá, E.F., 1991. Strain transfer at a continental scale from a transcurrent shear zone to a transpressional fold belt: the Patos-Seridó belt system, north-eastern Brazil. *Geology* 19, 586–589.
- Corsini, M., Figueiredo, L.L., Cabry, R., Féraud, G., Ruffet, G., Vauchez, A., 1998. Thermal history of the Pan-African/Brasiliano Borboroma Province of northeast Brazil deduced from $^{40}\text{Ar}/^{39}\text{Ar}$ analysis. *Tectonophysics* 285, 103–117.
- Costa, A.C.D., 2002. *Geologia e geocronologia Sm–Nd e U–Pb na região do Lineamento Patos, limite entre os blocos crustais do domínio Rio Grande do Norte e da Zona Transversal, Província Borboroma*. PhD thesis, State University of São Paulo.
- Charles, N., Faure, M., Chen, Y., 2009. The Montagne Noire migmatitic dome emplacement (French Massif Central): new insights from petrofabric and AMS studies. *Journal of Structural Geology* 31, 1423–1440.
- De Witt, M., Stankiewicz, J., Reeves, C., 2008. Restoring Pan-African-Brasiliano Connections: More Gondwana Control, Less Transatlantic Correlation. In: Geological Society of London, Special Publication, vol. 294, pp. 399–412.
- Denèle, Y., Olivier, P., Gleizes, G., Barbey, P., 2007. The Hospitalet gneiss dome (Pyrenees) revisited: lateral flow during Variscan transpression in the middle crust. *Terra Nova* 19, 445–453.
- Ferré, E., Teyssier, C., Jackson, M., Thill, J.W., Rainey, E.G., 2003. Magnetic susceptibility anisotropy: a new petrofabric tool in migmatites. *Journal of Geophysical Research* 108, 2086. <http://dx.doi.org/10.1029/2002JB001790>.
- Hrouda, F., Jelinek, V., Hruskova, L., 1990. A package of programs for statistical evaluation of magnetic anisotropy data using IBM-PC computers (abstract). *EOS Transactions of the American Geophysical Union*, 1289.
- Jackson, M., 1991. Anisotropy of magnetic remanence: a brief review of mineralogical sources, physical origins and geological applications, and

- comparison with susceptibility anisotropy. *Pure and Applied Geophysics* 136, 1–28.
- Jelinek, V., 1978. Statistical processing of anisotropy of magnetic susceptibility measured on groups of specimens. *Studia Geophysica et Geodetica* 22, 50–62.
- Just, J., Kontny, A., Wall, H., Hirt, A.M., Martín-Hernández, F., 2004. Complex Fabrics: Superposition and Alteration Development of Magnetic Fabrics during Hydrothermal Alteration in the Soultz-sous-Forêts Granite from the EPS-1 Borehole, Upper Rhine Graben. In: Geological Society of London, Special Publication, vol. 238. <http://dx.doi.org/10.1144/GSL.SP.2004.238.01.26>, pp. 509–526.
- Kruckenberg, S.C., Ferré, E., Teyssier, C., Vanderhaeghe, O., Whitney, D.L., Seaton, N.C.A., Skord, J.A., 2010. Viscoplastic flow in migmatites deduced from fabric anisotropy: an example from the Naxos dome, Greece. *Journal of Geophysical Research* 115, B09401. <http://dx.doi.org/10.1029/2009JB007012>.
- Leitch, A.M., Weinberg, R.F., 2002. Modelling granite migration by mesoscale pervasive flow. *Earth and Planetary Science Letters* 200, 131–146.
- Mehnert, K.R., 1968. *Migmatites and the Origin of Granitic Rocks*. Elsevier, Amsterdam, London.
- Monié, P., Caby, R., Arthaud, M.H., 1997. The Neoproterozoic Brasiliano orogeny in northeast Brazil: $^{40}\text{Ar}/^{39}\text{Ar}$ and petrostructural data from Ceará. *Precambrian Research* 81, 241–264.
- Muxworthy, A.R., McClelland, E., 2000. Review of the low-temperature magnetic properties of magnetite from a rock magnetic perspective. *Geophysical Journal International* 140 (1), 101–114. <http://dx.doi.org/10.1046/j.1365-246x.2000.00999.x>.
- Nicolas, A., Poirier, J.P., 1976. *Crystalline Plasticity and Solid State Flow in Metamorphic Rocks*. Wiley & Sons, Ltd, London.
- Oliveira, R.G., 2008. Arcabouço geofísico, isostasia e causas do magmatismo cenozoico da Província Borborema e de sua margem continental (Nordeste do Brasil). PhD thesis, Federal University of Rio Grande do Norte.
- Özdemir, O., Dunlop, D.J., 1993. The effect of oxidation on the Verwey transition in magnetite. *Geophysical Research Letters* 20 (16), 1671–1674.
- Paterson, S.R., Fowler, T.K., Schmidt, K.L., Yoshinobu, A.S., Yuan, E.S., Miller, R.B., 1998. Interpreting magmatic fabric patterns in plutons. *Lithos* 44, 53–82.
- Rosenberg, C., 2001. Deformation of partially molten granite: a review and comparison of experimental and natural case studies. *International Journal of Earth Sciences* 90, 60–76. <http://dx.doi.org/10.1007/s005310000164>.
- Rosenberg, C., Handy, M., 2005. Experimental deformation of partially melted granite revisited: implications for the continental crust. *Journal of Metamorphic Geology* 23, 19–28. <http://dx.doi.org/10.1111/j.1525-1314.2005.00555.x>.
- Rutter, E., Neumann, D.H.K., 1995. Experimental deformation of partially molten Westerly granite under fluid-absent conditions, with implications for the extraction of granitic magmas. *Journal of Geophysical Research* 100 (15), 697–715.
- Sawyer, E.W., 2008. Atlas of Migmatites. In: The Canadian Mineralogist, Special Publication, vol. 9. NRC Research Press, Ottawa, Ontario.
- Sawyer, E.W., 2010. Migmatites formed by water-fluxed partial melting of a leucogranodiorite protolith: microstructures in the residual rocks and source of the fluid. *Lithos* 116, 273–286.
- Trindade, R.F., Bouchez, J.L., Bolle, O., Nedelec, A., Peschler, A., Poitrasson, F., 2001. Secondary fabrics revealed by remanence anisotropy: methodological study and examples from plutonic rocks. *Geophysical Journal International* 147, 310–318.
- Van Schmus, W.R., Oliveira, E.P., Silva Filho, A.F., Toteu, S.F., Penaye, J., Guimarães, I.P., 2008. Proterozoic links between the Borborema Province, NE Brazil, and the Central African Fold Belt. In: Pankhurst, R.J., Trouw, R.A.J., Brito Neves, B.B., De Wit, M.J. (Eds.), *West Gondwana: Pre-Cenozoic Correlations Across the South Atlantic Region*. Geological Society of London, Special Publication, vol. 294, pp. 69–99.
- Vauchez, A., Neves, S.P., Caby, R., Corsini, M.E., Egydio Silva, M., Arthaud, M., Amaro, V.E., 1995. The Borborema shear zone system, NE Brazil. *Journal of South American Earth Sciences* 8, 247–266.
- Vernon, R., 2000. Review of microstructural evidence of magmatic and solid-state flow. *Electronic Geosciences* 5 (2), 1436–2511.
- Vigneresse, J.L., Barbey, P., Cuney, M., 1996. Rheological transitions during partial melting and crystallization with application to felsic magma segregation and transfer. *Journal of Petrology* 37 (6), 1579–1600.
- Weinberg, R.F., Mark, G., 2008. Magma Migration, Folding, and Disaggregation of Migmatites in the Karakoram Shear Zone, Ladakh, NW India. In: Geological Society of America Bulletin, vol. 120. <http://dx.doi.org/10.1130/B262271>, pp. 994–1009.
- Whitney, D., Teyssier, C., Vanderhaeghe, O., 2004. Gneiss Domes and Crustal Flow. In: Geological Society of America Special Papers, vol. 380. <http://dx.doi.org/10.1130/0-8137-2380-9.15>, pp. 15–33.

CAPÍTULO 5

CRONOLOGIA DE EVENTOS E LOCALIZAÇÃO DA DEFORMAÇÃO EM ZONAS DE CISALHAMENTO DÚCTEIS HETEROGÊNEAS

1. Introdução

A zona de cisalhamento Patos é caracterizada como uma estrutura transcorrente composta por milonitos grossos de alta-temperatura (fácies anfibolito) cortados por veios de leucossoma granítico, migmatitos cisalhados (anatexito Espinho Branco) e milonitos finos de média- a baixa-temperatura restritos à borda sul da falha. As relações de contato entre milonitos de alto grau e migmatitos são graduais, com a percentagem de fusão parcial aumentando em direção ao contato entre as duas unidades; os milonitos de baixo grau são observados como uma faixa localizada em contato abrupto com as demais litologias.

Estudos petroestruturais no anatexito Espinho Branco indicam que o alojamento do material parcialmente fundido é controlado pela cinemática da zona de cisalhamento Patos. Com efeito, leucossomas presentes nos milonitos de alta temperatura exibem microestruturas indicativas de deformação em estado sólido. Por outro lado, o anatexito Espinho Branco, alojado no interior da zona de cisalhamento, é predominantemente deformado no estado magmático, com feições de recristalização localizadas (Capítulo 4). Os milonitos da borda sul não exibem evidências de fusão parcial e possuem microestruturas de médio- a baixo-grau metamórfico, associadas a assembleias minerais hidratadas contendo clorita, epidoto, etc.

Este capítulo contém um artigo científico submetido ao periódico *Precambrian Research*, onde são investigadas as microestruturas e tramas cristalográficas dos milonitos de alta- e baixa-temperatura por meio da técnica de difração de elétrons retroespalhados (EBSD). A cronologia dos eventos de alta temperatura é estabelecida a partir de idades U-Pb em zircão (SHRIMP) nos migmatitos e milonitos de alta temperatura, complementada por análises das temperaturas deformacionais na faixa de médio- a baixo-grau obtidas a partir do método TitaniQ.

Os resultados obtidos sugerem que a localização da deformação exerceu um papel importante nos processos de milonitização de alto-grau, fusão parcial e retrabalhamento em baixo-grau observados na zona de cisalhamento Patos.

Microfabrics and zircon U-Pb (SHRIMP) chronology of high- and low-temperature mylonites from the Patos shear zone (Borborema Province, NE Brazil)

Luís Gustavo F. Viegas^{1, 2, *}, Carlos J. Archanjó¹, Maria Helena B. M. Hollanda¹, Alain Vauchez²

1. Instituto de Geociências, USP, Rua do Lago 562, 05508-080 São Paulo, SP, Brazil

2. Géosciences Montpellier, Place E. Bataillon 34095, Montpellier Cedex 5, France

* Corresponding author. Email address: lgviegas@usp.br; present address: Instituto de Geociências, USP, Rua do Lago 562, 05508-080 São Paulo, SP, Brazil.

Phone: +55 11 30 91 42 16 (L. G. Viegas).

Keywords: Mylonite, microstructure, EBSD, Borborema Province, geochronology, NE Brazil

Abstract

Microstructures and U-Pb SHRIMP zircon ages were used to constrain the fabric evolution and timing of mylonitisation in the Patos shear zone (NE Brazil). Melt-bearing HT-mylonites display solid-state fabrics with coarse quartz ribbons and sutured grain boundaries. K-feldspar is often fractured and shows peripheral myrmekite. The textures typically become magmatic at the contacts with anatexites. Zircons from two samples of the neosome show Palaeoproterozoic (~ 2.2 Ga) inherited cores that are enveloped by homogeneously recrystallised Neoproterozoic rims. These zircons define a discordia with lower intercept ages of 558 and 562 Ma but with large analytical errors. A leucogranite with transitional contacts with the host diatexites provides a well-constrained zircon mean age of 566 ± 5 Ma, which is considered to be the best estimate for the peak metamorphic conditions. The transition from the HT-mylonites to the fine-grained mylonites and ultramylonites is marked by progressive grain size reduction that is assisted by recrystallisation. Quartz CPOs display [0001] concentrations between Z and Y. The K-feldspar and plagioclase fabrics record the activity of the (010)[001] and (010)[100] slip systems and, in the lower-temperature tectonites, the (100)[010] slip system. Zircons from an ultramylonitic granite are very discordant but yield a well-defined lower intercept age of 545 ± 14 Ma. These results suggest a diachronic evolution in which an initial HT northern domain deformed by intracrystalline slip and melt-assisted granular flow and was followed by medium- to low temperature solid-state creep at the southern shear zone domain. This later event corresponds with the localisation of deformation along shear corridors, probably during exhumation of the structure.

1. Introduction

The Neoproterozoic Brasiliano/Pan-African orogeny produced an anastomosing, wide network of conjugate shear zones across an area of 200000 km² in Hoggar, Nigeria/Cameroon (Africa) and in the Borborema Province (NE Brazil). This type of

transcontinental fault system is believed to be typical of mid- to upper continental crustal deformation during large strike-slip movements. In the Borborema Province, ductile shear zones are associated with widespread granitic plutonism and synkinematic anatexis (Fig. 1; Vauchez et al. 1995). The interplay between magmatism and deformation resulted in the formation of > 20 km wide high-grade zones in which solid-state fabrics are observed in high-temperature mylonitic orthogneisses, migmatites and granites. Detailed field, structural and geochronological studies (Neves et al. 1996; Neves et al. 2000) have shown that magmatic activity played a major role in the nucleation and localisation of strain in these large shear zones and acted as a weak rheological layer that accommodated strain during strike-slip shearing.

Nevertheless, low-temperature mylonitic belts have been documented in several shear zones in the Borborema Province (Vauchez and Egydio-Silva 1992; Davison et al. 1995; Hollanda et al. 2010). These zones are composed of distinct fine-grained mylonites to ultramylonites that often form narrow zones or, in some cases, such as in the Patos shear zone, define abrupt contacts with high-temperature coarse mylonite belts (Corsini et al. 1991). Structural and geochronological investigations in some shear zones have shown that these low-temperature corridors record late-stage strain localisation.

The Patos shear zone consists primarily of high-temperature mylonites that can be divided into three structural domains. The Central Domain, which includes the study area, is composed of E-trending mylonitic orthogneisses that, further west towards the Western Domain, envelope a duplex structure consisting of NE-trending lenses of orthogneisses, metapelites and granitoids (Corsini et al. 1996). In the Eastern Domain, near the Atlantic coast, the mylonites gradually rotate from E-W to NE-SW. The foliations are mainly subvertical, and stretching lineations are subhorizontal along the entire zone, which is consistent with a dominant transcurrent dextral motion (Corsini et al. 1991). Geophysical studies have shown that the Patos shear zone connects with the NE-trending Senador Pompeu shear zone beneath the Parnaíba basin and forms a branched shear zone system that was active in the Neoproterozoic (Oliveira 2008).

Recent anisotropy of magnetic susceptibility (AMS) studies in the migmatites (Viegas et al. 2013) showed that the partially molten material flows laterally along the strike of the structure, which agrees with the dextral shear sense of the zone. However, despite being the subject of extensive work for more than fifteen years (Archanjo et al. 2008 and references therein), little is known about the strain patterns involved in the shearing, particularly with

respect to the mechanisms of strain localisation and the chronology of the high- and low-grade mylonite belts. This contribution focuses on the microstructures, crystallographic preferred orientations (CPOs) and zircon U-Pb (SHRIMP) analyses to investigate the fabric development and age of mylonitisation in the Patos shear zone. We present the first direct age determinations of synkinematic migmatisation and late fault reactivation using punctual methods. The roles of partial melting and strain localisation are addressed with respect to the evolution of a continental-scale strike-slip shear zone in which relatively short-lived (~30 Ma) tectonic events occurred in a single kinematic framework.

Fig. 1: The Neoproterozoic Borborema shear zone system in NE Brazil. The small box shows the study area, which is located within the Patos shear zone.

2. Geological setting

The Patos shear zone consists of a ~ 600 km long strike-slip shear zone that deforms the Precambrian rocks of the Borborema Province (Fig. 1). In the Central Domain, located at the junction between the Patos shear zone and the Seridó Belt, anatexites occur within a ~ 30 km wide connection zone. They are composed of stromatic metatexites, nebulitic diatexites and partially molten leucogranites (Viegas et al. 2013). Unpublished U-Pb and Sm-Nd data indicate that these anatexites derive from melting of the Palaeoproterozoic basement (Costa 2002). A narrow medium- to low-temperature mylonite belt forms the southern margin of the shear zone (Fig. 2) and reworked the coarse mylonites and migmatites of the northern border

at low temperatures. Recent structural and geochronological studies south of the Patos shear zone have shown that some E-trending transcurrent structures record late-stage strain localisation (Hollanda et al. 2010). This is demonstrated in the Coxixola shear zone, in which A-type granite plutons and dikes were emplaced in a strain field that combined dextral shear and extension. U-Pb (SHRIMP) zircon ages in these sheared granites suggest that the Coxixola fault developed between 550 and 530 Ma (Hollanda et al. 2010).

In the Patos mylonites, zircons dated by U/Pb (TIMs) (Costa 2002; Monié et al. 1997) showed strong disturbance but provided unconstrained upper intercept ages between 2.2 to 2.0 Ga. $^{40}\text{Ar}/^{39}\text{Ar}$ plateau ages in amphibole, muscovite and biotite range from 540 Ma to 490 Ma and were attributed to cooling of the shear zone (Corsini et al. 1998). Similar $^{40}\text{Ar}/^{39}\text{Ar}$ plateau ages were documented in muscovite from the Coxixola mylonites; the ~ 510 Ma age was attributed to the final uplift and exhumation of the shear zone (Hollanda et al. 2010).

Fig. 2: Main geological units of the Patos shear zone at the intersection with the Seridó belt. Red arrows in the anatexites and contoured stereograms represent magnetic lineation measurements and main orientations, respectively. Kamb density contours, lower hemisphere.

3. Rock types and shear zone structures

The Patos mylonites share a common orthogneiss/granite protolith and can be divided into four main rock types based on grain size, mesoscale structure and melt content: i) high-temperature melt-bearing mylonites that are located in the northern (upper) part of the shear zone; ii) sheared anatexites in the central sector; iii) “transitional” mylonites, which are

located immediately south of the migmatites, and; iv) fine-grained mylonites to ultramylonites at the southernmost border (Figs. 2, 3).

The coarse-grained melt-bearing mylonites often contain irregular pods of monzogranitic leucosome (Fig. 3a) and feldspar clasts surrounded by elongated biotite flakes, which usually define a layered/banded structure (Fig. 3a). Centimetric quartz ribbons define an E-W subhorizontal stretching lineation. The interlayered leucosomes show evidence of solid-state deformation. Rotated K-feldspar porphyroclasts and fine-grained recrystallised quartz in pressure shadows at feldspar edges indicate dextral motion. The melt content progressively increases towards the contact with the anatexites, and the microstructure gradually changes from solid-state to magmatic.

The anatexites were mostly deformed in the magmatic state and have local solid-state overprints (Viegas et al. 2013). The nebulites and leucogranites are mainly isotropic with little or no mesoscale deformational features. The metatexites show an E-W preferred orientation of elongated K-feldspar and biotite grains (Fig. 3b). Varying degrees of partial melting result in complex field foliation geometries that are not always compatible with the shear zone's strain field. However, the magnetic and crystallographic fabrics indicate that all of these structures resulted from dextral shear deformation in the Patos shear zone (Viegas et al. 2013).

South of the migmatites, the melt fractions gradually decrease, and the deformed rocks show a “transitional” character between migmatites and mylonites (Fig. 3c). Mafic phases become more common, while the leucosomes become increasingly sheared and less common. Medium-grained biotite-gneisses develop and usually display a preferred orientation of flattened phyllosilicates. Quartz is observed as dextrally-rotated sigmoids and as elongated rods that define a subhorizontal E-W lineation.

The southernmost part of the shear zone is composed of fine-grained ($\sim 20 \mu\text{m}$) mylonites to ultramylonites that are mostly derived from high-temperature mylonites and local granite intrusions (Fig. 2). Mesoscale evidence of partial melting is lacking, and these rocks are mainly composed of fractured K-feldspar porphyroclasts that are immersed in a fine (5-15 μm) felsic matrix (Fig. 3d). Millimetre-scale quartz ribbons define the lineations, and rotated feldspar clasts show a dextral shear sense.

Fig. 3: Field structures of the Patos mylonites: a) melt-bearing orthogneisses with sheared leucosomes and coarse layering; b) stromatic metatexites with sheared leucosomes parallel to the host rock fabric; c) transitional mylonite with increasingly-sheared leucosome bands in the foliation; d) fine-grained mylonite with centimetric quartz ribbons in a felsic matrix.

4. Results

4.1. Microstructural data

The microstructures were investigated by conventional optical microscopy and SEM imagery. The mineralogical composition of the different rock types is constant throughout the zone and is formed of nearly equal proportions of K-feldspar + quartz + plagioclase with biotite as the main Fe-silicate. Accessory ilmenite, magnetite and zircon are common, and minor chlorite, epidote, muscovite and sericite can be observed towards the southern border.

4.1.1. High-temperature melt-bearing mylonites

Quartz ribbons occur as rectangular grains with straight boundaries, subgrains and undulatory extinction (Figs. 4a, b). The subgrain boundaries are often oriented oblique to normal to the foliation. Polycrystalline aggregates of coarse equant grains ($> 350 \mu\text{m}$) display

sutured/serrated boundaries and migrating borders towards the adjacent phases (Fig. 4a). The quartz grains outside the ribbons are smaller ($\sim 100\text{-}200\text{ }\mu\text{m}$) and have embayed/lobate boundaries with small bulges at quartz-quartz contacts (Fig. 4c, d). Internal subgrain boundaries remain oblique to normal to the foliation, and bulging structures are common (Figs. 4c, d). Late microcracks crosscut the coarse grains (Fig. 4d).

K-feldspar forms coarse ($150\text{-}300\text{ }\mu\text{m}$), rectangular to subelliptical porphyroclasts with straight boundaries (Figs. 4c, d). Fine-grained myrmekites mantle coarse anhedral clasts and may be associated with fine recrystallised quartz ($< 15\text{ }\mu\text{m}$) that has been collected in the shear zones at grain boundaries (Figs. 4e, f). Undulatory extinction is common (Figs. 4e, f). Sets of late microcracks may be slightly oblique to the grain elongation (Fig. 4d).

The plagioclase ($\text{An}_{35\text{-}38}$) is recrystallised as medium-sized anhedral crystals with straight boundaries in a local polygonal texture. Some of the boundaries are interfingered with fine quartz grains ($\sim 20\text{-}25\text{ }\mu\text{m}$) and show lobate borders towards the quartz and K-feldspar grains (Figs. 4c, d). Mechanical twins are locally bent, and late microcracks occur locally.

4.1.2. Migmatites

Quartz often occurs as coarse anhedral grains up to 2 mm in size and as smaller crystallised grains $\sim 70\text{-}200\text{ }\mu\text{m}$ in size. The coarse grains have straight boundaries and show undulatory extinction and subgrains. Medium- to fine-grained quartz grains occur widely as an interstitial phase that crystallised at quartz-Kfs-plagioclase boundaries (Figs. 5a, b, c). Fractures in the feldspar are preferential sites for the nucleation of newly crystallised quartz as “films” with bulging and pinning microstructures (Fig. 5b).

The K-feldspar forms coarse inequant porphyroclasts with pervasive crosshatch twinning and undulose to straight boundaries (Fig. 5a, c, d). Fine patches ($\sim 10\text{-}30\text{ }\mu\text{m}$) of albite lamellae crosscut the porphyroclasts (Fig. 5d) oblique to normal to the mylonitic foliation. Dilatant fractures cut the porphyroclasts normal to the foliation and either cut across the entire grain or terminate within the grain. The cracks occur preferentially at cleavage planes or twin boundaries and may be filled by newly-crystallised quartz or fine albite grains. The boundaries between adjacent K-feldspar grains are preferential sites for myrmekite nucleation as small ($<15\text{ }\mu\text{m}$) films (Fig. 5a) or within fine rounded grains (Figs. 5c).

Fig. 4: Microstructures in the melt-bearing mylonites: a) coarse-grained quartz ribbons in the foliation with small bulges at grain boundaries (arrows); b) stretched quartz and feldspar clasts that define an *augen* microstructure; c) and d) interstitial quartz at grain boundaries with local bulges at like-like contacts (arrows); e) and f) phase mixing between fine-grained quartz and myrmekite at K-feldspar edges. All photos are with crossed polarizers. The shear sense is dextral. Abbreviations: Qtz - quartz; Kfs - K-feldspar; plag - plagioclase; Bt - biotite.

The plagioclase is mainly andesitic (An_{35-38}) and ranges from 100 to 300 μm in size (Figs. 5a, b). Mechanical twins are oriented normal to the foliation direction, and straight boundaries are usually overprinted by quartz bulges (Fig. 5b). Microcracks are common and may be locally filled by sericite.

Fig. 5: Microstructures in the migmatites: a) and b) interstitial quartz forming melt films (arrows in Fig. 5a) and as small “drops” (arrows in Fig. 5b) at grain boundaries; c) myrmekite exsolutions at K-feldspar contacts. The arrow points to interstitial quartz at triple junctions; d) microcracks in K-feldspar filled with albite exsolution laths (arrows).

4.1.3. Transitional mylonites

Coarse quartz grains (150-300 μm) have sutured/serrated boundaries, may form foliation-parallel ribbons that surround coarse K-feldspar porphyroclasts (Figs. 6a, b) and have oblique subgrain boundaries and undulatory extinction. The coarse grains may be locally associated with traces of melt that occur as fine crystals ($< 30 \mu\text{m}$) at grain boundaries (arrows in Fig. 6a). The quartz may also be recrystallised as fine (50-100 μm) grains associated with micas (Fig. 6b) as well as polygonal mosaics of finer ($< 45 \mu\text{m}$) grains with straight borders (Fig. 6e); these polycrystalline aggregates define mesoscale ribbons that are parallel to the mylonitic foliation.

The K-feldspar occurs either as rotated porphyroclasts (Figs. 6a, b) or as elongated *augen* (Figs. 6c, d). Fine grains ($< 20 \mu\text{m}$) fill fractures and newly-formed millimetre-thick shear zones that developed both normal and parallel to the foliation (Figs. 6e, f). The fine-grained K-feldspar appears to be strain free and is sometimes associated with fine recrystallised quartz. The microshear zones locally contain small amounts of low-grade hydrous minerals, such as chlorite and epidote.

Fig. 6: Microstructures in the transitional mylonites: a) residual coarse-grained texture with K-feldspar porphyroclasts and interstitial quartz (arrows); b) bimodal grain size observed in the transition from a coarse to a medium-grained texture; c) and d) augen structures developed under increasing strain rates; e) polygonal mosaics of recrystallised quartz grains that define foliation-parallel ribbons; f) intense recrystallisation of quartz and the development of microcracks and micro shear zones in K-feldspar. The arrows point to fine K-feldspar aggregates in the shear zones.

Plagioclase sometimes occurs as coarse porphyroclasts with traces of melt (Fig. 6a) but is completely recrystallised when in association with fine-grained K-feldspar (Figs. 6e, f). The plagioclase crystals are partially replaced by sericite and epidote with progressive grain size reduction.

4.1.4. Low-temperature mylonites

Quartz forms millimetre-scale ($\sim 30 \mu\text{m}$) elongated ribbons and fine ($< 15 \mu\text{m}$) recrystallised grains in the matrix. The ribbons usually have straight boundaries with rectangular subgrain walls that are normal to the foliation (Figs. 7a, d, e, f). Fine aggregates may be recrystallised as ribbons that are oblique to the foliation (Fig. 7e). The quartz in the matrix ($\sim 5 \mu\text{m}$) is completely recrystallised and is associated with recrystallised K-feldspar, plagioclase and biotite. Microfaults displace the matrix material alongside the quartz bands (Fig. 7a).

K-feldspar ($\sim 100\text{-}200 \mu\text{m}$) is immersed in the felsic matrix as anhedral grains or stretched porphyroclasts (Figs. 7b, c, d, e), which show undulatory extinction and are pervasively crosscut by fractures that extend throughout the entire grain. Fractures can occur as sets of oriented and equally-spaced cracks that transect the grains in “domino” configurations (Fig. 7b) and can also displace the porphyroclast with normal offsets that are filled by the matrix material (Fig. 7c). Fine trails of matrix minerals may also surround the porphyroclasts (Figs. 7b, c). The elongated porphyroclasts are strongly stretched in the foliation direction and show pinch-and-swell/boudinage structures with straight to curved boundaries (Fig. 7d). The stretching sometimes results in dilaceration of former porphyroclasts into discontinuous fine ribbons (Fig. 7d).

Plagioclase rarely occurs as porphyroclasts except for medium-grained ($\sim 100 \mu\text{m}$) clasts that are crosscut by sets of equally-spaced fractures that are oriented in a domino arrangement (Fig. 7b). Fine-grained plagioclase ($\sim 5 \mu\text{m}$) dominates the matrix.

4.2. Crystallographic preferred orientations

4.2.1. Methods

Representative samples of the Patos mylonites were chosen for EBSD analysis of quartz and feldspars. Each sample was cut in the XZ section of the strain ellipsoid and was polished to ensure a flat surface. The orientation measurements were performed in a JEOL JSM 5600 scanning electron microscope (SEM) that was equipped with an Oxford Instruments/HKL Nordlys EBSD detector at Géosciences Montpellier. The Channel 5 Program (Oxford Instruments) was used for the data treatment. Samples were inserted into the microscope chamber at an angle of 70° relative to the electron beam and with a working

distance of 25 mm. A 17 kV accelerated electron beam was applied for a spot size of 75 μm . The acquisition step sizes ranged from 25 to 50 μm , and the orientation measurements were rotated into the geographical frame.

Fig. 7: Microstructures in the fine-grained mylonites and ultramylonites: a) microcracks and microfaults displacing quartz-rich foliated bands (arrows); b) sets of microcracks that form a “domino” configuration in K-feldspar and plagioclase porphyroclasts; c) microfaults in K-feldspar with normal offsets filled by the matrix material (arrow); d) highly-stretched K-feldspar clast (arrow) that defines a pinch-and-swell/ribbon microstructure alongside fine quartz ribbons; e) and f) oblique quartz ribbons in contact with a K-feldspar porphyroclast and subgrain walls normal to oblique to the foliation (arrow in Fig. 7f).

4.2.2. Lattice fabric results

The quartz [0001] axis in the melt-bearing mylonites displays well-defined concentrations in the centre of the diagram with symmetrical a -axis maxima in the periphery around Z and X (Fig. 8a). The increase in melt content towards the centre of the shear zone results in weak patterns and random crystallographic fabrics in the anatexites, which are marked by small [0001] axis point maxima close to Z and Y and a -axes that form isolated weak concentrations near Y (Fig. 8b). The transitional mylonites show a dispersion of [0001] axes along the YZ plane with maxima in Y and minor concentrations located between Y and Z. This pattern resembles a faint single-girdle in the NS direction (Fig. 8c). The a -axes are scattered through the periphery of the diagram with maxima in X, which is consistent with the orientation of the stretching lineations observed in the field. The low-temperature mylonites display strong [0001] axis concentrations in Z that spread to intermediary positions between Z and Y, while the a -axes are mainly concentrated in the periphery and form maxima around X (Fig. 8d).

Fig. 8: Crystallographic preferred orientations of quartz. Lower hemisphere pole figures, equal area stereographic projection, contours at 1, 1.5, 2, 2.5, 3% etc. by 1% area. The pole figures were rotated to the geographical frame of reference with the peripheries of the stereonet (Z and X) corresponding to the vertical N-S and horizontal E-W directions, respectively.

The K-feldspar in the melt-bearing mylonites displays maxima of (100) poles that are spread near the stretching lineation direction (X), but they also form girdles that split into two small maxima in the periphery close to X. The (010) poles are mainly concentrated close to Z, and the (001) poles show weak or no crystallographic preferred orientation (Fig. 9a, b). The migmatites display concentrations of K-feldspar (010) poles in Z and maxima of (100) poles close to X, *i.e.*, in the lineation direction (Fig. 9c), which suggests activity of the commonly reported (010)[100] slip system (Schulmann et al. 1996; Martelat et al. 1999; Kruse et al. 2001). In the transitional mylonites, the (100) poles define maxima close to the stretching lineation, and the (010) poles form an asymmetrical girdle over the YZ plane with concentrations around Z (Fig. 10a). Near the contact with the low-temperature mylonites, the (100) poles display weak concentrations over Z, and the (010) poles form small (10-15°) angles with X, which suggests activity of the (100)[010] slip system (Fig. 10b; Ishii et al. 2007). The fine-grained mylonites display evidence of activity on the (100)[001] and (010)[001] slip systems (Fig. 10c; Tullis 1983; Martelat et al. 1999; Menegon et al. 2008).

The plagioclase fabrics have (100) poles that form girdles parallel to the stretching lineation and maxima close to Y, which suggests activity of the (010)[100] and (010)[001] slip systems in the melt-bearing mylonites (Figs. 9a, b). The (010)[100] slip system is also observed in the migmatites as girdles of (100) poles in the lineation direction (Fig. 9c). In the transitional mylonites, the activated slip systems include (010)[100] and (100)[010], which is indicated by the (100) and (010) poles close to X, respectively (Figs. 10a, b). The fabrics in the low-temperature mylonites (Fig. 10c) are marked by (100) poles that are asymmetrically concentrated close to X, which suggests the contribution of the tentative (100)[001] slip system (Kruse et al. 2001).

4.3. Titanium-in-Quartz Thermometry

The Ti content in the recrystallised quartz was measured according to the procedures described by Wark & Watson (2006). The method is based on the activity of TiO_2 (a_{TiO_2}), which can be used as a strain geothermometer in deformed rocks. For rutile-saturated rocks, $a_{\text{TiO}_2} = 1$. The Patos mylonites do not contain rutile, but an a_{TiO_2} of 0.6 - 0.8 for crustal granitic mylonites is widely accepted in the literature (Wark & Watson 2006, Wark et al. 2007). We consider an a_{TiO_2} of 0.7 ± 0.2 for our samples, which is consistent with previous

work on mylonites in similar geological settings (Kohn & Northrup 2009; Menegon et al. 2011).

Fig. 9: Crystallographic preferred orientations of feldspars in the melt-bearing mylonites and migmatites. Lower hemisphere pole figures, equal area stereographic projection, contours at 1, 1.5, 2, 2.5, 3% etc. by 1% area. The pole figures were rotated to the geographical frame of reference with the peripheries of the stereonet (Z and X) corresponding to the vertical N-S and horizontal E-W directions, respectively.

Fig. 10: Crystallographic preferred orientations of feldspars in the transitional and fine-grained mylonites. Lower hemisphere pole figures, equal area stereographic projection, contours at 1, 1.5, 2, 2.5, 3% etc. by 1% area. The pole figures were rotated to the geographical frame of reference with the peripheries of the stereonet (Z and X) corresponding to vertical N-S and horizontal E-W directions, respectively.

The measurements were focused on the transitional and low-temperature mylonites (sample location in Fig. 2) to determine the temperatures during the low-grade mylonitisation along the southern border. For each sample, cross-sections were performed through recrystallised grains to measure the Ti contents using a microprobe at Géosciences Montpellier. At least five cross-sections were performed in the recrystallised grains in each

sample. Anomalously high Ti concentrations at grain boundaries or close to Ti-rich minerals were excluded from the data set. The results are summarised in Table 1.

The average Ti concentrations in the transitional mylonites (sample GPT02) range from 1 to 439 ppm, with a mean value of 8.71 ppm. The temperatures vary from 515°C to 555°C, with a calculated mean value of 534°C. In the low-grade mylonites (sample PT251), the Ti concentrations are lower and range from 1 to 21 ppm, with a mean value of 5.42 ppm. Mean temperatures of ~ 495°C are recorded in the southern sector of the Patos shear zone.

Table 1. Ti contents and average temperatures in Ti-in-Quartz thermometry

Ti (ppm)		T(°C) aTiO ₂ = 0.7
Transitional mylonites		
Average	8.71	534.94
SD	6.4	
N	32	
Low-temperature mylonites		
Average	5.42	495.92
SD	5.08	
N	47	

4.4. U/Pb (SHRIMP) zircon geochronology

4.4.1. Sampling and methods

We collected two samples of the felsic leucosomes (PTS, PAUG), one sample of a leucogranite (PA4) associated with the migmatites, and a sample of a mylonitic *augen*-granite (URT). Samples PTS, PAUG and PA4 come from the Espinho Branco anatexite (Viegas et al. 2013), while sample URT comes from the Urtiga granite, which is located in the southern margin of the Patos shear zone (location in Fig. 2). Samples PTS and PAUG comprise medium-grained K-feldspar, quartz, plagioclase and biotite in a granular texture with feldspar clasts that are surrounded by a faint shape-preferred orientation of biotite flakes. Sample PA4 is an equigranular biotite-leucogranite with irregular to diffuse contacts with diatexite. The

augen mylonite (sample URT) is fine-grained and contains equal proportions of K-feldspar, quartz and plagioclase that define a banded structure.

The zircon grains from all of the samples were handpicked after standard crushing and separation. The clearest euhedral to subhedral grains were selected and mounted in epoxy with chips of TEMORA 1 reference zircon ($^{206}\text{Pb}/^{238}\text{U} = 416.78 \pm 0.33$ Ma; Black et al. 2004), sectioned approximately in half and polished. Reflected and transmitted light photomicrographs and cathodoluminescence (CL) SEM images were obtained for all of the grains. The analyses of the U-Th-Pb elements were conducted at the SHRIMP IIe laboratory of the Centro de Pesquisas Geocronológicas at the University of São Paulo. The isotopic data were collected in sets of five scans throughout the masses, and the TEMORA standard was measured after four unknown analyses. The abundances of U, Th and Pb, and the Pb isotopic ratios, were normalised using the SL13 zircon standard (U = 238 ppm). Common Pb was corrected using the measured ^{204}Pb and assuming the appropriate model of Pb composition from Stacey and Kramers (1975). The data reduction was performed using the SQUID 1.06 Excel macro (Ludwig 2000), and the statistical assessments were calculated using the geochronological ISOPLOT software of Ludwig (2003). The data are reported in Table 2 and are plotted on a Wetherill U-Pb concordia diagram. The uncertainties in the isotopic ratios and the ages are reported at the 1σ level, and the weighted mean ages are reported as 95% confidence limits.

4.4.2. Zircon microstructures

The zircons from the leucosomes and the leucogranite are morphologically similar. The grain sizes range from 50 to 400 μm in length, but the zircons from sample PTS reach 300 μm in length. Axial ratios (long/short dimensions) between 2:1 and 4:1 are typical for all of the grains.

The crystals are euhedral to subhedral and are mostly colourless and inclusion free. CL imaging shows typical inner cores that display discrete fine-scale oscillatory zoning, which is interpreted to reflect magmatic growth (Fig. 11). Most of the cores are surrounded by homogeneous bright luminescent overgrowths (Fig. 11a, b, c). In some of the zircons, these overgrowths transgress the darker luminescent inherited cores (Fig. 11a, b). In other cases, the overgrowths are epitaxial and nearly continuous, and the cores preserve the magmatic prismatic shape (Fig. 11c). We interpret the overgrowths as having been produced

during a high temperature metamorphic event. The analysis of cores with oscillatory zoning and the respective overgrowths showed that they crystallised at distinct time periods.

Table 2
Summary of U-Pb SHRIMP zircon data for the Espinho Branco migmatite and the Urtiga granite

[illegible]

Most of the zircons from sample URT (Fig. 11d) are broken with abundant internal fracturing. However, some preserved crystals display subhedral to euhedral shapes with grain sizes ranging from 50 to 300 μm in length and axial ratios from 1.4:1 to 2.7:1. CL imaging highlights the intense internal fracturing that sometimes forms a radial pattern (Fig. 11d). Bright luminescent inclusions are common and are normally aligned with crystal faces. Some zircons preserve oscillatory zoning, which indicates magmatic growth (Fig. 11d). Overgrowths were not detected. The ion-microprobe analysis was therefore focused on the more homogenous domains as far as possible from the fracture traces. Some of the zircons are porous and are characterised by abundant fine inclusions. These altered zones were not analysed because they probably represent dissolution and precipitation of new phases due to metamorphic fluid percolation (Hay & Dempster 2009).

Spot	% comm			ppm			2022H			2023H			Ln	Ln U	Ln D	Corr 2026	ppm Rad 2026H	Total 2022H	204cent 2026H	207cent 2026H	208cent 2026H	204cent 2026H	204cent 2026H	Dis-const	4ccorr 2022	Total 2026	Total 2027	4ccorr 2026	4ccorr 2027	4ccorr 2028	4ccorr 2029	4ccorr 2030	Bccorr 2026	Bccorr 2027	Bccorr 2028	Bccorr 2029	Bccorr 2030	Bccorr 2031							
	206	207	208	206H	207H	208H	206H	207H	208H	206H	207H	208H																											206H	207H	208H	206H	207H	208H	206H
Sample P44																																													
1.1	0.09	0.64	124	0.15	1.746	-2.099	0.90	1.71	0.671	0.278	30	557.2	94	555.5	93	557.7	94	650	37	531	22	1.0267	4.1	0900	0.004	0004	0.016	11.08	1.7	0.613	1.7	0.736	2.4	0903	1.7	705	11.07	0.021	1.5	0.77	2.3	0904	1.75	0.681	
2	0.05	619	86	0.15	1.626	-2.047	0.91	1.48	0.493	20.1	34	556.7	94	557.0	96	558.5	96	546	73	476	21	1.0240	0.13	0903	0.016	0005	0.016	11.18	0.584	3	0.73	2.8	0902	1.8	149	11.06	0.140	0.2	0.76	2.6	0.905	1.85	1.660		
3	-0.12	675	73	0.15	1.623	-2.334	0.92	1.8	0.455	0.297	36	568.2	96	566.8	98	567.1	98	645	49	627	23	1.0314	3.7	0919	0.017	0920	0.017	10.18	1.8	0.611	2.3	0.78	2.9	0921	1.8	161	11.07	0.18	0.695	2.4	0.75	2.9	0.920	1.8	1.631
3.1	0.44	336	79	0.25	1.710	-1.723	1.09	1.28	0.505	0.295	27.2	662.9	127	649.3	157	657	1687	645	1177	205	126	76	0.0254	24.3	0910	0.027	1033	0.025	10.20	0.792	14.7	11.8	0.143	10.9	0.21	0.36	1.95	1.51	1.34	1.093	2.0	1.601			
4	0.24	151	63	0.43	1.736	-1.762	0.47	2.0	0.452	0.497	28	1916.5	329	1874.4	368	1901.5	343	2177	27	1739	64	1.4	0903	3.5	0981	0.075	3482	0.072	2.20	0.190	1.5	0.649	2.5	3452	2.0	790	28.07	0.2	1.402	1.3	0.73	2.4	0.894	2.0	1.708
4.2	-0.13	591	33	0.06	1.722	-0.164	0.98	1.9	0.449	0.239	49	547.3	100	545.4	102	546.3	101	650	46	680	61	1.0335	8.9	0983	0.071	0984	0.071	11.29	1.9	0.613	2.1	0.679	2.5	0986	1.9	665	11.31	0.092	1.9	0.73	2.7	0.943	1.9	1.620	
5	0.24	359	131	0.38	1.727	-0.930	2.99	1.9	0.222	0.267	25	1680.0	267	1643.3	314	1688.0	301	1999	21	1459	49	1.0774	3.1	0903	0.083	2995	0.081	3.36	1.9	1.229	1.2	0.505	2.3	2977	1.9	855	33.4	1.9	1.274	1.0	0.56	2.2	0.959	1.9	1.692
5.2	0.16	780	99	0.13	1.647	-2.261	0.93	1.28	0.622	0.302	32	571.2	91.6	571.7

Fig. 11: Cathodoluminescence images of selected zircons from the Patos shear zone with core and rim mean isotopic ages. Scale bar is 100 μm in A, B, and D and 50 μm in C.

4.4.3. Results

Twelve zircons from sample PTS were investigated and led to fifteen analyses of cores and their respective zircon overgrowths. The data define a discordia line with an upper intercept age at 2195 ± 13 Ma (Fig. 12a). Four concordant analyses (#2.2, #5.2, #6.2 and #9.2) fix the lower intercept age at 558 ± 21 Ma. The upper intercept is interpreted to date the crystallisation of the protolith, while the lower intercept represents the time of the thermal disturbance that formed the zircon overgrowths (Mezger and Krogstad 1997).

The analyses of the inner zircon cores of sample PAUG align in a discordia. After removing two reverse ages (#5.1, #10.1) and three discordant zircons (#9.2, #6.2b, #13.3), the remaining data define an upper intercept age of 2201 ± 12 Ma (Fig. 12b). Conversely, ten concordant zircon overgrowths cluster to yield a mean age of 552 ± 11 Ma (1σ , MSWD = 0.25). Furthermore, the combined analyses define a lower intercept age of 562 ± 12 Ma (Fig.

12b), which is indistinguishable within the uncertainty limits of the mean age provided by the zircon overgrowths. Therefore, these results indicate that melting of Palaeoproterozoic rocks in the late Ediacaran formed the felsic leucosomes.

Fig. 12: U-Pb concordia diagrams for the Espinho Branco migmatite. Insets in samples PTS and PAUG show rim age results.

Sample PA4 yielded three analyses in the zircon cores that displayed oscillatory zoning (#4.1, #5.1, #9.1) and twenty analyses of the respective overgrowths. The ages of the cores confirm the Palaeoproterozoic inheritance of the zircons and, together with analyses of the overgrowths, yield lower and upper intercept ages of 568 ± 10 Ma and 2147 ± 47 Ma, respectively (Fig. 12c). The data reduction of the overgrowths highlighted two zircons with high ^{204}Pb contents (#8.2, #13.1), two with high Th/U ratios (#15.1, #16.1), one with a high U content (#14.1) and one that is very discordant (#3.1). After discarding these analyses, the remaining zircons cluster on the concordia and provide a mean age of 566 ± 5 Ma (Fig. 12d). This age is indistinguishable within the uncertainty limits of the lower intercept age and is considered to be the best estimate for the crystallisation of sample PA4.

Fourteen zircons were investigated in sample URT. The analyses are very discordant. After discarding one zircon with a high ^{206}Pb content (#11.1), the data define upper and lower intercept ages at 2197 ± 33 Ma and 545 ± 14 Ma, respectively (Fig. 13). The array of zircons that plot in the left half of the discordia, including a nearly concordant zircon (#13.1; Fig. 13b), indicate that the lower intercept age dates the mylonitisation of the southern margin of the Patos shear zone (Mezger & Krogstad 1997).

Fig. 13: U-Pb concordia diagram for the Urtiga ultramylonitic granite.

5. Discussion

5.1. High-temperature deformation mechanisms and age of migmatisation

The sutured/lobate high-energy grain boundaries and subgrain walls that are normal to the coarse ribbons in the quartz, the myrmekite that replaces K-feldspar and the recrystallisation of plagioclase indicate deformation in the melt-bearing mylonites by high-temperature dislocation creep (Passchier & Trouw 1996; Rosenberg & Berger 2001). Melt is observed mainly at interstitial grain boundaries and phase mixing distributions between quartz and feldspars (Fig. 4c, d, e). These structures are evidence of synkinematic melting and enhanced diffusional processes such as grain boundary migration (GBM), as exemplified by serrated quartz boundaries with small bulges at like-like (same phase) contacts (Nicolas & Poirier 1976; Vernon 2000; Rosenberg 2001; Sawyer 2010). Further evidence for high temperature conditions is provided by the activity of the prism $\langle a \rangle$ slip system in quartz ($> 500^\circ\text{C}$; Schmid & Casey 1986) and the activation of the (010)[001] and (010)[100] slip systems in plagioclase (Ji & Mainprice 1988; Egydio-Silva et al. 2002).

The microstructure in the migmatites is mainly magmatic with interstitial melt films at grain boundaries and microcracks in K-feldspar (Fig. 5). In addition, the presence of fine-grained strain-free quartz and the occurrence of myrmekite suggest deformation mainly by granular flow with a limited contribution of plastic strain (Vernon 2000; Rosenberg 2001; Sawyer 2001). The quartz CPOs display no visible preferred orientation (Fig. 8b), but the feldspars show activity of the (010)[100] slip system (Fig. 9c). Taken together, these data indicate that melt-enhanced diffusion played an important part in the deformation of the migmatites and prevented the development of a crystallographic preferred orientation in the quartz due to crystallisation from the melt. On the other hand, the formation of a CPO in the feldspars might be attributed to oriented crystal growth during synmagmatic shearing (Tommasi et al. 1994; Garlick & Gromet 2004; Hasalová et al. 2008).

High temperature conditions are also recorded in zircon crystals from the migmatites, which usually have cores with fine magmatic oscillatory zoning enveloped by late homogeneously recrystallised zircon rims. Anatectic zircon can be formed by several different processes, such as dissolution-precipitation of pre-existing grains, breakdown of Zr-bearing phases (e.g., biotite and monazite) and crystallisation from externally-derived Zr-bearing fluids and melts (Rubatto et al. 2001; Andersson et al. 2002; Flowerdew et al. 2006). Nevertheless, irrespective of (re)crystallisation processes, zircon formation temperatures usually range from 850°C to 600°C and vary with the water content (Harley et al. 2007). The $^{207}\text{Pb}/^{238}\text{U}$ ages of the zircon cores define an upper intercept at ~ 2.200 Ma, which indicates that the migmatite derived from partial melting of an older Palaeoproterozoic basement. In contrast, the homogeneous regions surrounding the inherited cores are Neoproterozoic and correspond to the melting event. The recrystallised zircons yielded lower intercept ages that are marked by nearly concordant analysis in the range of 558 - 562 Ma but have a relatively high analytical error. However, a leucogranite with diffuse to lobate contacts with the host migmatite provided a well-constrained lower intercept at 568 ± 10 Ma (MSWD = 2.2) and a mean age of 566 ± 5 Ma (MSWD = 1.3, 2σ level). Hence, we consider the 565 - 568 Ma age range to be the best estimate for the crystallisation of the leucogranite. Field evidence suggests that the leucogranite was emplaced while the host rock was still molten, thus providing the age of the synkinematic melting at the Patos shear zone.

5.2. Transition from high-temperature to low-temperature mylonites

The progressive grain size reduction at medium- to low-temperatures that is observed in the transitional mylonites indicates that these rocks may mark a transient stage between high- and low-grade microstructures. Polygonal flattened or stretched quartz aggregates in the foliation result from subgrain rotation recrystallisation (Fig. 6e; Passchier & Trouw 1996). In the K-feldspar, elongated clasts, microcracks that crosscut porphyroclasts, often in “domino” arrangements (Fig. 6f), and the widespread occurrence of recrystallised grains as fine trails that fill fractures indicate the contribution of pressure solution and/or grain boundary sliding mechanisms (Tullis 1983; Hasalová et al. 2008; Menegon et al. 2008).

The quartz fabrics suggest the activity of prism $\langle a \rangle$ and rhomb $\langle a+c \rangle$ slip systems, which are typical of high-temperature deformation at increasing strain rates (Stipp et al. 2002). The feldspar CPOs record the activity of both the (010)[100] and (100)[010] slip systems (Figs. 10a, b). The activity of the (100)[010] slip system has been reported in granitic ultramylonites that deformed mainly via microfracturing and solution-precipitation creep in the Ryoke metamorphic belt (Ishii et al. 2007). Based on the lack of homogeneous phase mixing, these authors assumed that grain boundary sliding was not the dominant deformation mechanism responsible for the formation of feldspar CPO. In the transitional mylonites from the Patos shear zone, the fine-grained ($< 10 \mu\text{m}$) recrystallised feldspars that fill fractures may have resulted from dissolution-precipitation processes (Wintsch & Yi 2002; Berger & Stunitz 1996). In addition, phase mixing is observed between quartz and feldspar in these dilatational sites, which suggests the involvement of grain boundary sliding (GBS) at upper greenschist/lower-amphibolite facies conditions ($\sim 500^\circ\text{C}$), which was corroborated by the Ti-in-quartz measurements (section 4.3). The fact that low-grade mineral assemblages are locally observed indicates that hydrothermal fluids may have had a limited influence on the dissolution-precipitation processes at the southern border of the Patos shear zone.

5.3. Low-temperature mylonitisation

The fine-grained mylonites to ultramylonites have a homogeneously-mixed felsic matrix that contains highly-deformed K-feldspar porphyroclasts and quartz-rich bands (Fig. 7). The peak temperatures of $\sim 490^\circ\text{C}$ that were calculated from the Ti contents of quartz indicate that this fine-grained mylonitic zone was deformed at upper greenschist facies conditions. The fine-grained recrystallised quartz grains ($< 15 \mu\text{m}$) contain straight

boundaries and display undulatory extinction, which suggests deformation mainly by dislocation creep (Passchier & Trouw 1996). The K-feldspar porphyroclasts are crosscut by fractures and are sometimes stretched in the foliation direction (Fig. 7d), which may indicate crystal plastic deformation processes such as dislocation creep (FitzGerald & Stunitz 1993); the mixture of K-feldspar with other minerals in the matrix suggests a contribution of grain boundary sliding through homogeneous phase-mixing during high-strain deformation (Tullis 1983).

The quartz CPOs that have concentrations between Z and Y indicate a progressive change from prism to rhomb planes activity at increasing strain conditions (Stipp et al. 2002). The development of feldspar CPOs suggests that recrystallisation was mainly attained by grain boundary sliding that was accommodated by dislocation creep; GBS is especially important in fine grains and can develop CPOs in feldspar grains (Tullis 1983). Moreover, the phase mixing that is widely observed in the matrix material also indicates that high-strain rates were achieved by a combination of crystal plastic processes at low-temperature conditions, probably in the presence of residual hydrothermal fluids (FitzGerald & Stunitz 1993; Ishii et al. 2007; Menegon et al. 2008).

5.3. Strain localisation and regional implications

The high-temperature mylonites of the Patos shear zone and the Seridó metasedimentary belt form a branched system that is characterised by a continuous foliation trend from E-W to NE-SW and by zones of partially molten rocks (Corsini et al. 1991). U-Pb (SHRIMP) zircon ages of ~ 575 Ma in the Acari granite and the Santa Luzia diatexite precisely constrain the syntectonic magmatism and the main high-temperature/low-pressure metamorphism in the Seridó belt (Archanjo et al. 2013). The high-temperature event in the Patos shear zone appears to be younger (565-568 Ma) than that in the Seridó belt. Alternatively, the lower intercept age of ~ 545 Ma recorded in the zircons from sample URT records upper greenschist facies reactivation at the southern border. These data suggest that the high- and low-temperature mylonitic processes are diachronic and occurred at distinct time intervals within the shear zone.

The main HT/LP metamorphism observed in the Seridó belt is coeval with widespread amphibolite facies mylonitisation and the emplacement of partially molten rocks within the Patos-Seridó shear system, which is suggested by their magnetic fabrics that are

consistent with the regional strain field (Santa Luzia and Espinho Branco anatexites, Fig. 2; Archanjo et al. 2013; Viegas et al. 2013). Partial melting induced weakening of the quartz fabrics, while the feldspars show well-defined CPOs due to grain growth during magmatic flow. These patterns indicate the partitioning of deformation into solid-state flow in the HT mylonites and granular flow in the migmatites. The presence of migmatites at the intersection between the Patos shear zone and the Seridó belt suggests that the curvature of the mylonitic fabric from E-W to NE-SW might have been facilitated by the distribution of the melting zones, which would act as weak layers of strain localisation at high temperatures (Rosenberg & Handy 2005).

In contrast, the melt-absent shear localisation along the southern border of the fault occurred at lower temperatures ($\sim 490^\circ\text{C}$). This event partially reworked the structures of the Espinho Branco anatexite and is consistent with the younger age calculated from zircons of a sheared granite (now an *augen*-gneiss) that is hosted in fine-grained mylonites and phyllonites. The quartz CPOs that spread between Z and Y indicate the activity of dislocation creep at high strain rates. On the other hand, the feldspars deformed mainly by solution-precipitation creep and grain boundary sliding, as indicated by recrystallisation, phase mixing, CPOs and the localised presence of low-temperature mineral assemblages that are probably associated with hydrothermal fluid percolation.

These observations suggest that the mylonitisation at the southern margin of the shear zone represents a second-stage reactivation process that is marked by further localisation of deformation within narrow zones. Because the greenschist-facies mylonites form the northern limit of the metasedimentary rocks and variably deformed granites of the Cachoeirinha belt (Archanjo et al. 2008), this corridor acted as a preferential layer that accommodated deformation due to the rheological heterogeneities between the two geological domains.

The tectonic reactivations observed in the Patos shear zone have also been documented in other shear domains of the Borborema Province (e.g., Davison et al. 1995; Neves et al. 2000; Vauchez & Egydio-Silva 1992; Hollanda et al. 2010). For instance, fine mylonites and ultramylonites from the dextral transcurrent Coxixola shear zone (Fig. 1) overprinted older basement structures for hundreds of kilometres in the E-W direction (Hollanda et al. 2010). Related granite plutons and dikes constrain the movement of the Coxixola mylonites between 548 and 533 Ma. Furthermore, late shear reactivation in the Seridó belt appeared to control the emplacement of Li-Be bearing pegmatites and Au mineralisations that are hosted in quartz veins (Baumgartner et al. 2006; Araújo et al. 2005). Available $^{40}\text{Ar}/^{39}\text{Ar}$ biotite and muscovite ages in the Patos, Coxixola and Seridó domains

provide cooling ages between 545 and 500 Ma, which are mainly attributed to differential regional uplift and final exhumation in the Late Cambrian (Corsini et al. 1998; Araújo et al. 2005; Hollanda et al. 2010).

6. Conclusions

Microstructures, CPOs and U/Pb (SHRIMP) zircon ages indicate a diachronic, short-lived (~ 20 Ma) evolution composed of two distinct events in a midcrustal transcurrent shear zone. The northern high-temperature border is coupled with synkinematic anatexis and is associated with the main HT/LP metamorphism that is recorded in the Seridó belt. Such high-temperature conditions enhanced the strain distribution and localised partial melting at the Patos-Seridó boundary. Alternatively, reactivation in the southern sector indicates that melt-absent solid-state creep was active at least until the early Cambrian.

These results suggest that the heterogeneous evolution of the Patos shear zone occurred in a single kinematic strain field where distinct rheological conditions were present at different time intervals and ultimately led to different patterns of deformation within the strike-slip structure. The fact that the second-stage reactivation observed at the southern border correlates with the Coxixola fault suggests that the Cambrian tectono-magmatic activity is of regional extent and occurred in other shear zones across the Borborema Province. These late-stage low-temperature faults developed mainly along previously formed discontinuities at contrasting lithological boundaries, which are usually characterised by deformation at the semi-brittle to ductile transition.

Acknowledgments

L. G. Viegas and C. Archanjo thank Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP, grants 2010/50060-1 and 2009/17537-1) and the Brazilian Research Council (CNPq, grant 200496/2011-5) for financial support. This paper is part of the first author's PhD thesis. We thank John Rico and Camilo Bustamante for assistance during the field trips. Prof. Leonardo Lagoeiro and the students at MICROLAB (DEGEO-UFOP) are thanked for invaluable assistance during the initial EBSD work. Fabrice Barou, Claude Merlet and Bernard Boyer are thanked for their help during EBSD and Ti-in-quartz measurements at Géosciences Montpellier.

References

- Andersson, T., Moller, C., Johansson, L., 2002. Zircon geochronology of migmatite gneisses along the Mylonite Zone (S Sweden): a major Sveconorwegian terrane boundary in the Baltic Shield. *Precambrian Research* 114, 121-147.
- Araújo, M.N.C., Vasconcelos, P.M., Alves da Silva, F.C.A., Jardim de Sá, E.F., Sá, J.M., 2005. $^{40}\text{Ar}/^{39}\text{Ar}$ geochronology of the gold mineralization in Brasiliano strike-slip shear zones in the Borborema province, NE Brazil. *Journal of South American Earth Sciences* 19, 445-460, <http://dx.doi.org/10.1016/j.sames.2005.06.009>.
- Archanjo, C.J., Hollanda, M.H.B.M., Rodrigues, S.W.O., Brito Neves, B.B., Armstrong, R., 2008. Fabrics of pre- and syntectonic granite plutons and chronology of shear zones in the Eastern Borborema, NE Brazil. *Journal of Structural Geology* 30, 310-326.
- Archanjo, C.J., Viegas, L.G., Hollanda, M.H.B.M., Souza, L.C., Liu, D., 2013. Timing of the HT/LP transpression in the Neoproterozoic Seridó Belt (Borborema Province, Brazil): Constraints from U-Pb (SHRIMP) geochronology and implications for the connections between NE Brazil and West Africa. *Gondwana Research* 23, 701-714, <http://dx.doi.org/10.1016/j.gr.2012.05.005>.
- Arthaud, M.H., Caby, R., Fuck, R.A., Dantas, E.L., Parente, C.V., 2008. Geology of the Northern Borborema Province NE Brazil and its Correlation with Nigeria, NW Africa. Geological Society of London, Special Publication 294, 49-67.
- Baumgartner, R., Romer, R.L., Moritz, R., Sallet, R., Chiaradia, M., 2006. Columbite-tantalite bearing granitic pegmatites from the Seridó belt, northeastern Brazil: genetic constraints from U-Pb dating and Pb isotopes. *Canadian Mineralogist* 44, 69-86, doi:10.2113/gscanmin.44.1.69.
- Berger, A., Stunitz, H., 1996. Deformation mechanisms and reactions of hornblende: examples from the Bergell tonalite (Central Alps). *Tectonophysics* 257, 149-174.
- Black, L.P., Kamo, S.L., Allen, M.C., Davis, D.W., Aleinikoff, J.N., Valley, J.W., Mundif, R., Campbell, I.H., Korsch, R.J., Williams, I.S., Foudoulis, C., 2004. Improved $^{206}\text{Pb}/^{238}\text{U}$ microprobe geochronology by monitoring of a trace element related matrix effect; SHRIMP, ID-TIMS, ELA-ICP-MS and oxygen isotope documentation for a series of zircon standards. *Chemical Geology* 205, 115-140.
- Corsini, M., Vauchez, A., Archanjo, C.J., Jardim de Sá, E.F., 1991. Strain transfer at a continental scale from a transcurrent shear zone to a transpressional fold belt: the Patos-Seridó belt system, north-eastern Brazil. *Geology* 19, 586-589.
- Corsini, M., Vauchez, A., Caby, R., 1996. Ductile duplexing at a bend of a continental-scale strike-slip shear zone: example from NE Brazil. *Journal of Structural Geology* 18, 385-394.
- Corsini, M., Figueiredo, L.L., Caby, R., Féraud, G., Ruffet, G., Vauchez, A., 1998. Thermal history of the Pan-African/Brasiliano Borborema Province of northeast Brazil deduced from $^{40}\text{Ar}/^{39}\text{Ar}$ analysis. *Tectonophysics* 285, 103-117.
- Costa, A.C.D., 2002. Geologia e geocronologia Sm-Nd e U-Pb na região do Lineamento Patos, limite entre os blocos crustais do domínio Rio Grande do Norte e da Zona Transversal, Província Borborema. PhD thesis, State University of São Paulo.
- Davison, I., MacCarthy, M., Powell, D., Torres, H., Santos, C.A., 1995. Laminar flow in shear zones: The Pernambuco shear zone, NE-Brazil. *Journal of Structural Geology* 17, 149-161.
- De Witt, M., Stankiewicz, J., Reeves, C., 2008. Restoring Pan-African-Brasiliano connections: more Gondwana control, less Tran-Atlantic corruption. Geological Society of London, Special Publication 294, 399-412.
- Egydio-Silva, M., Vauchez, A., Bascou, J., Hippertt, J., 2002. High-temperature deformation in the Neoproterozoic transpressional Ribeira belt, southeast Brazil. *Tectonophysics* 352, 203-224.

- Fitz Gerald, J.D., Stünitz, H., 1993. Deformation of granitoids at low metamorphic grade. I: reactions and grain size reduction. *Tectonophysics* 221, 269-297.
- Flowerdew, M.J., Millar, I.L., Vaughan, A.P.M., Horstwood, M.S.A., Fanning, C.M., 2006. The source of granitic gneisses and migmatites in the Antarctic Peninsula: a combined U-Pb SHRIMP and laser ablation Hf isotope study of complex zircons. *Contribution to Mineralogy and Petrology* 151, 751-768.
- Garlick, S.R., Gromet, L.P., 2004. Diffusion creep and partial melting in high-temperature mylonitic gneisses, Hope Valley shear zone, New England, Appalachians, USA. *Journal of Metamorphic Geology* 22, 45-62.
- Hasalová, P., Schulmann, K., Lexa, O., Stipská, P., Hrouda, F., Ulrich, S., Aloda, J., Tycova, P., 2008. Origin of migmatites by deformation-enhanced melt infiltration of orthogneiss: a new model based on quantitative microstructural analysis. *Journal of Metamorphic Geology* 26, 29-53.
- Hay, D.C., Dempster, T.J., 2009. Zircon behavior during low-temperature metamorphism. *Journal of Petrology*, 50, 571-589, doi: 10.1093/petrology/egp011.
- Hollanda, M.H.B.M., Archanjo, C.J., Souza, L.C., Armstrong, R., Vasconcelos, P.M., 2010. Cambrian mafic to felsic magmatism and its connections with transcurrent shear zones of the Borborema Province (NE Brazil): implications for the late assembly of the West Gondwana. *Precambrian Research* 178, 1-14. <http://dx.doi.org/10.1016/j.precamres.2009.12.004>.
- Ishii, K., Kanagawa, K., Shigematsu, N., Okudaira, T., 2007. High ductility of K-feldspar and development of granitic banded ultramylonite in the Ryoke metamorphic belt, SW Japan. *Journal of Structural Geology* 29, 1083-1098.
- Ji, S., Mainprice, D., 1988. Natural deformation fabrics of plagioclase: implications for slip systems and seismic anisotropy. *Tectonophysics* 147, 145-163.
- Kohn, M. J., Northrup, C.J., 2009. Taking mylonites' temperatures. *Geology* 37, 47-50, doi:10.1130/G25081A.1.
- Kruse, R., Stünitz, H., Kunze, A., 2001. Dynamic recrystallization processes in plagioclase porphyroclasts. *Journal of Structural Geology* 24, 587-589.
- Ludwig, K.R., 2000. SQUID 1.00: A User Manual, vol. 2. Berkeley Geochronology Center, Special Publication, Berkeley, CA.
- Ludwig, K.R., 2003. Isoplot 3.00: A Geochronological Toolkit for Microsoft Excel® (Revised Version), vol. 4. Berkeley Geochronological Center, Special Publication, Berkeley, CA.
- Martelat, J.E., Schulmann, K., Lardeaux, J.M., Nicollet, C., 1999. Granulite microfabric and deformation mechanisms in southern Madagascar, *Journal of Structural Geology* 21, 671-687, doi:10.1016/S0191-8141(99)00052-8.
- Menegon, L., Pennacchioni, G., Spiess, R., 2008. Dissolution-precipitation creep of K-feldspar in mid-crustal granite mylonites. *Journal of Structural Geology* 30, 565-579.
- Menegon, L., Nasipuri, P., Stünitz, H., Behrens, H., Ravna, E., 2011. Dry and strong quartz during deformation of the lower crust in the presence of melt. *Journal of Geophysical Research* 116, B10410, doi:10.1029/2011JB008371.
- Mezger, K., Krogstad, E.J. 1997. Interpretation of discordant U-Pb zircon ages: an evaluation. *Journal of Metamorphic Geology*, 15: 127-140.
- Monié, P., Caby, R., Arthaud, M.H., 1997. The Neoproterozoic Brasiliano orogeny in northeast Brazil: ⁴⁰Ar/³⁹Ar and petrostructural data from Ceará. *Precambrian Research* 81, 241-264.

- Neves, S.P., Vauchez, A., Archanjo, C.J., 1996. Shear-zone controlled magma emplacement or magma-assisted nucleation of shear zones? Insights from northeast Brazil. *Tectonophysics* 262, 349-365.
- Neves, S.P., Vauchez, A., Féraud, G., 2000. Tectono-thermal evolution, magma emplacement, and shear zone development in the Caruaru area (Borborema Province, NE Brazil). *Precambrian Research* 99, 1–32.
- Nicolas, A., Poirier, J.P., 1976. Crystalline plasticity and solid state flow in metamorphic rocks. Wiley & Sons, Ltd. London.
- Oliveira, R.G., 2008. Arcabouço geofísico, isostasia e causas do magmatismo cenozóico da Província Borborema e de sua margem continental (Nordeste do Brasil). PhD thesis, Federal University of Rio Grande do Norte.
- Passchier, C.W., Trouw, R.A.J., 1996. *Microtectonics*. Springer-Verlag, Heidelberg.
- Rosenberg, C., 2001. Deformation of partially molten granite: a review and comparison of experimental and natural case studies. *International Journal of Earth Sciences* 90, 60-76, doi: 10.1007/s005310000164.
- Rosenberg, C., Berger, A., 2001. Syntectonic melt pathways in granitic gneisses, and melt-induced transitions in deformation mechanisms. *Physics and Chemistry of the Earth* 26, 287–293, doi:10.1016/S1464-1895(01)00058-8.
- Rosenberg, C., Handy, M., 2005. Experimental deformation of partially melted granite revisited: implications for the continental crust. *Journal of Metamorphic Geology* 23, 19-28.
- Rubatto, D., Williams, I.S., Buick, I.S., 2001. Zircon and monazite response to prograde metamorphism in the Reynolds Range, central Australia. *Contributions to Mineralogy and Petrology* 140, 458-468.
- Sawyer, E.W., 2001. Grain-scale and outcrop-scale distribution and movement of melt in crystallizing granite. *Transactions of the Royal Society of Edinburgh: Earth Sciences* 91, 73–85.
- Sawyer, E.W., 2010. Migmatites formed by water-fluxed partial melting of a leucogranodiorite protolith: Microstructures in the residual rocks and source of the fluid. *Lithos* 116, 273-286.
- Schmid, S.M., Casey, M., 1986. Complete fabric analysis of some commonly observed quartz [c]-axis patterns. In: Hobbs, B.E., Heard, H.C. (Eds.), *Mineral and Rock Deformation: Laboratory Studies*. American Geophysical Union, Geophysical Monograph 36, 263–286.
- Schulmann, K., Mlcoch, B., Radek, M., 1996. High-temperature microstructures and rheology of deformed granite, Erzgebirge, Bohemian Massif. *Journal of Structural Geology* 18, 719-733.
- Stacey, J.S., Kramers, J.D., 1975. Approximation of terrestrial lead isotope evolution by a two-stage model. *Earth and Planetary Science Letters* 26, 207-221.
- Stipp, M., Stunitz, H., Heilbronner, R., Schmid, S.M., 2002. The eastern Tonale fault zone: a ‘natural laboratory’ for crystal plastic deformation of quartz over a temperature range from 250 to 700°C. *Journal of Structural Geology* 24, 1861-1884.
- Tommasi, A., Vauchez, A., Fernandes, L.A.D., Porcher, C.C., 1994. Magma-assisted strain localization in a orogen-parallel transcurrent shear zone of southern Brazil. *Tectonics* 13, 421-437.
- Tullis, J., 1983. Deformation of feldspars. In: Ribbe, P.H. (Ed.), *Feldspar mineralogy*. American Reviews in Mineralogy 2, 297-323.
- Vauchez, A., Egydio-Silva, M., 1992. Termination of a continental-scale strike-slip fault in partially melted crust: The West-Pernambuco shear zone, northeast Brazil. *Geology* 20, 1007-1010.
- Vauchez, A., Neves, S.P., Cabby, R., Corsini, M.E., Egydio Silva, M., Arthaud, M., Amaro, V.E., 1995. The Borborema shear zone system, NE Brazil. *Journal of South American Earth Sciences* 8, 247-266.

- Vernon, R., 2000. Review of microstructural evidence of magmatic and solid-state flow. *Electronic Geosciences* 5, 2, 1436-2511.
- Viegas, L.G.F., Archanjo, C., Vauchez, A., 2013. Fabrics of migmatites and the relationships between partial melting and deformation in high-grade transpressional shear zones: the Espinho Branco anatexite (Borborema Province, NE Brazil). *Journal of Structural Geology* 48, 45-56, <http://dx.doi.org/10.1016/j.jsg.2012.12.008>.
- Wark, D.A., Hildreth, W., Spear, F.S., Cherniak, D.J., Watson, E.B., 2007. Pre-eruption recharge of the Bishop magma system. *Geology* 35, 235-238, doi: 10.1130/G23316A.1.
- Wark, D.A., Watson, E.B., 2006. TitaniQ: A titanium-in-quartz geothermometer. *Contributions to Mineralogy and Petrology* 152, 743–754, doi:10.1007/s00410-006-0132-1.
- Wintsch, R.P., Yi, K., 2002. Dissolution and replacement creep: a significant deformation mechanism in mid-crustal rocks. *Journal of Structural Geology* 24, 1179-1193.

CAPÍTULO 6

CONCLUSÕES E CONSIDERAÇÕES FINAIS

1. Síntese

Esta tese compreende o estudo das relações estruturais entre magmatismo e zonas de cisalhamento em níveis crustais intermediários. Com base na integração entre trabalhos de campo e o uso de técnicas multidisciplinares, tais como ASM, ARA, EBSD, U-Pb (SHRIMP) e TitaniQ, foi possível avaliar: i) os mecanismos de alojamento de migmatitos em contextos tectônicos distintos, ii) as relações estruturais entre o fluxo de material parcialmente fundido e o alojamento de granitos, iii) a correlação entre estruturas mesoscópicas e a trama interna em corpos migmatíticos, iv) o desenvolvimento de tramas magmáticas em anatexitos deformados em regimes transcorrentes, v) os processos de nucleação e localização da deformação em zonas de cisalhamento dúcteis, e vi) a cronologia de eventos magmáticos e deformacionais em um contexto de anatexia sincinemática.

As principais conclusões obtidas a partir do presente estudo são apresentadas a seguir.

2. Metamorfismo de alto grau e fusão parcial na Faixa Seridó

O diatexito de Santa Luzia exhibe uma trama magnética bem organizada e consistente com a orientação regional da Faixa Seridó. A diferença entre as foliações de campo, que configuram uma geometria dômica, e a trama magnética marcada por lineações concordantes ao eixo de elongação do domo, refletem a sensibilidade da ASM aos incrementos finais da deformação que afetou o migmatito, possivelmente após o alojamento. Estes dados sugerem que o domo de Santa Luzia foi alojado e deformado em um regime de fluxo viscoplástico, onde a trama interna reflete o padrão tectônico regional.

A idade de 575 Ma obtida nos sobrecrescimentos metamórficos de zircões extraídos do leucossoma posiciona de forma precisa a fusão parcial no sistema de cisalhamento Patos-Seridó, a qual correlaciona-se com a idade de alojamento do pluton de Acari (~ 577 Ma), localizado mais a norte. Adicionalmente, a trama magnética é concordante no domo de Santa Luzia e no granito Acari, sugerindo que a fusão parcial do embasamento e a colocação de granitos porfiríticos estão mecanicamente associados na Faixa Seridó. Por outro lado, as

tramas circulares observadas no granito Totoró indicam que o mesmo é pré-cinemático em relação à deformação transcorrente do Seridó, o que é corroborado pela idade de ~ 595 Ma obtida em dioritos e granitos deste corpo plutônico.

Dessa forma, os dados geocronológicos e estruturais permitem concluir que a atividade magmática observada na região de Santa Luzia teve início antes da deformação registrada na Faixa Seridó, sendo subsequentemente associada a metamorfismo das rochas do embasamento e fusão parcial na crosta intermediária, esta última conectada ao plutonismo granítico observado em níveis crustais mais rasos.

3. Tramas migmatíticas e alojamento de magmas em regimes transcorrentes

A complexidade geométrica observada no anatexito Espinho Branco não é observada na orientação da trama magnética, a qual se mostra concordante ao campo deformacional da zona de cisalhamento Patos. Entretanto, as relações entre trama de campo e trama magnética não são diretas, especialmente em sítios onde as foliações de campo não são confiáveis e os valores de suscetibilidade são muito baixos. Nestes casos, o mapeamento da trama interna através da ASM pode levar a resultados não-conclusivos.

Nas amostras com suscetibilidades moderadas a altas (> 1.0 mSI), a magnetita é o mineral portador do sinal magnético e domina a anisotropia de suscetibilidade. A trama magnética exibe boa compatibilidade com foliações de campo e os eixos $[001]$ de cristais de biotita alinham-se em concordância com as direções de k_3 da ASM. Alternativamente, suscetibilidades baixas (< 0.5 mSI) registram uma dispersão das direções de k_1 e k_3 , evidenciando a presença de sub-tramas que influenciam a trama magnética. Com efeito, as tramas magnética e cristalográfica podem exibir direções contrastantes, e comumente não se correlacionam com geometrias de campo. Contudo, as direções de ARA permanecem consistentes com o padrão regional e registram a orientação geral da trama magnética no migmatito.

Este banco de dados sugere que a geometria complexa de anatexitos alojados em zonas de cisalhamento transcorrentes não é compatível com sua respectiva trama interna, a qual registra os incrementos finais da deformação cisalhante. A trama de ARA confirma o caráter ferromagnético do migmatito, dominado por magnetita, enquanto que as tramas de biotita são concordantes com as direções de k_3 . Entretanto, suscetibilidades muito baixas (< 0.5 mSI) podem indicar a presença de sub-tramas definidas por silicatos ferromagnesianos e fases hidrotermais tardias, *e.g.* hematita, as quais seriam responsáveis pela diminuição dos

valores de suscetibilidade, dispersão da trama magnética e o desenvolvimento de orientações preferenciais secundárias (Trindade et al. 2001; Just et al. 2004; Archanjo et al. 2009).

Dessa forma, pode-se concluir que a ASM é uma ferramenta útil para recuperar a trama interna de migmatitos alojados no interior de zonas de cisalhamento dúcteis, particularmente em rochas com suscetibilidades altas (> 1.0 mSI). Entretanto, baixos valores de suscetibilidade (< 0.5 mSI) podem refletir a presença de sub-tramas caracterizadas pela interação de fases hidrotermais tardias com os minerais magnéticos, as quais promovem dispersão das orientações da ASM. Nesses casos, investigações complementares englobando ARA e OPCs devem ser empregadas com o objetivo de detectar sub-tramas e avaliar o significado dos dados de ASM.

4. Localização da deformação e reativação na zona de cisalhamento Patos

A história deformacional da zona de cisalhamento Patos evoluiu de forma diacrônica, com a borda norte registrando idades mais antigas que o setor sul. A idade de ~ 566 Ma obtida nos leucossomas do migmatito Espinho Branco é ligeiramente mais jovem que a idade do metamorfismo e fusão parcial registrados na Faixa Seridó (~ 575 Ma). Adicionalmente, os milonitos finos a ultramilonitos da borda sul registram idades em torno de 545 Ma. Estes dados sugerem que i) a zona de cisalhamento Patos registrou temperaturas altas por mais tempo durante a evolução do sistema, caracterizadas por uma prolongada atividade da fusão parcial, ou ii) o resfriamento se deu de forma mais rápida na faixa metassedimentar, possivelmente devido ao contraste litológico observado no contato entre *augen*-gnaisses do embasamento, o migmatito Santa Luzia e os metassedimentos da Faixa Seridó.

Dados termocronológicos obtidos pelo método $^{40}\text{Ar}/^{39}\text{Ar}$ em anfibólio e biotita indicam uma taxa de resfriamento lenta, em torno de $3\text{--}4^\circ\text{C}$ por ano, entre 540 e 500 Ma na zona de cisalhamento Patos (Corsini et al. 1998; Monié et al. 1997). Tais características sugerem que a nucleação da estrutura se deu em condições de alta temperatura, envolvendo fusão parcial, e possivelmente associada ao metamorfismo em fácies anfibolito registrado na Faixa Seridó. A presença de anatexitos favoreceu a localização da deformação na porção norte da estrutura, configurando uma zona de instabilidade reológica marcada pela curvatura geométrica e conexão entre a zona de cisalhamento e a faixa metassedimentar. Por outro lado, a borda sul evoluiu na ausência de magma e em condições de médio- a baixo-grau, como demonstrado pelas estimativas de temperatura calculadas pelo método TitaniQ (Capítulo 5). Estas condições metamórficas envolveram geração de milonitos finos a

ultramilonitos a partir de recristalização extensiva e progressiva diminuição do tamanho dos grãos.

A localização da deformação em corredores de cisalhamento ao longo da margem sul da zona de cisalhamento Patos exibe semelhança com outras áreas da Província Borborema, como as zonas de cisalhamento Coxixola e Pernambuco (Hollanda et al. 2010; Neves & Mariano 1999). Nestas, transformações metamórficas de médio- a baixo-grau retrabalham rochas de alta-temperatura gerando faixas miloníticas intensamente deformadas. Estas zonas de cisalhamento possuem idades entre 548 e 533 Ma (Hollanda et al. 2010), concordantes com a idade de 545 Ma obtida nos ultramilonitos da zona de cisalhamento Patos.

Dessa forma, a reativação registrada na zona de cisalhamento Patos constitui um evento regional, ativo em outras áreas da Província Borborema durante a transição Neoproterozóico-Cambriano, e caracterizado pela atuação de mecanismos de deformação que incluem fluência de deslocções e fraturamento intracristalino. Estes mecanismos são observados nos corredores miloníticos relativamente retilíneos e pouco espessos, restritos à borda sul da estrutura, onde o contraste litológico inerente às condições de limite da zona favorece a acumulação da deformação e resulta em microestruturas geradas principalmente na faixa de transição reológica dúctil-rúptil.

5. Considerações finais

i) A ASM constitui uma ferramenta útil no mapeamento da trama interna de migmatitos alojados em sistemas de cisalhamento transpressivos. A trama interna do domo de Santa Luzia revela um alojamento controlado pelo cisalhamento dextral da Faixa Seridó, em concordância com o fluxo magmático em níveis crustais mais rasos, representado pelo granito Acari. Na zona de cisalhamento Patos, a geometria complexa dos leucossomas do migmatito Espinho Branco não é reproduzida em sua trama magnética, a qual se mostra bem organizada e consistente com a cinemática da zona de cisalhamento.

ii) Investigações estruturais por meio de técnicas complementares, como a ARA e tramas cristalográficas, são particularmente úteis quando a trama de ASM exibe variações nos valores de suscetibilidade e direção magnética. Nestes casos, as OPCs fornecem informações importantes sobre as orientações preferenciais de eventuais sub-tramas formadas por fases minerais distintas (silicatos, óxidos), enquanto que a ARA revela o padrão direcional das fases ferromagnéticas. Dada a complexidade da geometria de campo e a heterogeneidade da

percentagem de fusão parcial observada em rochas parcialmente fundidas, a aplicação integrada das referidas técnicas, em conjunto com trabalhos de campo detalhados, se faz fundamental no estudo dos mecanismos de migração e transporte de magmas ao longo da crosta intermediária.

iii) A investigação cronológica do magmatismo na região do Seridó sugere que a fusão parcial é sincrônica ao plutonismo granítico observado em níveis crustais mais rasos. Com efeito, as tramas magnéticas concordantes em migmatitos e granitos corroboram essa hipótese. Análises geoquímicas complementares constituem o próximo passo na investigação geológica destas rochas; tais dados serão importantes na caracterização tipológica do magmatismo e identificação de suas fontes, bem como no refinamento da correlação entre rochas parcialmente fundidas e granitoides na crosta continental intermediária-superior.

iv) A presença de microestruturas e mecanismos de deformação contrastantes nas bordas norte e sul da zona de cisalhamento Patos, em associação com as idades obtidas em zircões de leucossomas na borda norte, milonitos na borda sul e as estimativas de temperatura obtidas por meio do método TitaniQ, sugerem que a nucleação da estrutura se deu em condições de alta temperatura, na presença de fusão parcial, por volta de 565 Ma. Após esse primeiro evento, uma reativação subsequente em 545 Ma retrabalhou os milonitos grossos e migmatitos gerando milonitos finos a ultramilonitos em temperaturas intermediárias a baixas, definindo faixas miloníticas concentradas na borda sul.

v) A localização da deformação de médio- a baixo-grau na porção sul da estrutura se deve ao contraste litológico entre a zona de cisalhamento Patos e a Zona Transversal, a qual é formada por metapelitos e granitos fracamente deformados. Estas diferenças litológicas, não-observáveis na borda norte da estrutura, provavelmente contribuíram para o caráter dúctil-rúptil da reativação observada na borda sul da zona de cisalhamento Patos.

vi) A idade dos eventos de magmatismo, metamorfismo e fusão parcial registrados a norte da zona de cisalhamento Patos (~ 575 Ma), quando comparados com idades dos plutons mais antigos alojados no domínio a sul da estrutura, conhecido como Zona Transversal (~ 630-610 Ma), corroboram a hipótese de a zona de cisalhamento Patos representar um limite crustal dividindo terrenos com evoluções geológicas distintas (Vauchez et al. 1995). Com efeito, o batólito Teixeira, localizado no contato sul da estrutura, exibe tramas obliquas a normais à

orientação do cinturão de cisalhamento, e possui uma idade de cristalização em torno de 590 Ma, associada a um estágio inicial de alojamento de magmas anterior ao desenvolvimento de estruturas transcorrentes (Archanjo et al. 2008; Neves et al. 2006). Os dados obtidos no presente trabalho indicam que a nucleação e desenvolvimento da zona de cisalhamento Patos se deu a partir de 565 Ma. Dessa forma, esta zona de cisalhamento representa uma estrutura pós-colisional, nucleada no intervalo 570-560 Ma, e gerada após a colagem de blocos crustais ao final do Neoproterozóico, possivelmente como resultado da transferência progressiva de esforços colisionais em deslocamentos transcorrentes.

vii) A reativação tardia observada na zona de cisalhamento Patos, datada em 545 Ma, exhibe características similares com a deformação observada em outras zonas de cisalhamento da Província Borborema (e.g. Coxixola, Pernambuco), as quais estão associadas à intrusão de granitoides máficos e félsicos na transição Neoproterozóico-Cambriano. Portanto, pode-se concluir que a deformação de baixo-grau observada na zona de cisalhamento Patos constitui um evento de caráter regional, observado em outras áreas da Província Borborema, e possivelmente associado à colisões tardias na margem oeste do Gondwana (Hollanda et al. 2010).

CAPÍTULO 7

REFERÊNCIAS

- Adams, B.I., Wright, S.I., Kunze, K., 1993. Orientation imaging: the emergence of new microscopy. *Metallurgical Transactions* 24, 819-831.
- Archanjo, C.J., 1993. Fabriques des plutons granitiques et deformation crustale du Nord-Est du Brésil: une etude par anisotropie de susceptibilité magnétique de granites ferromagnétiques. *These de Doctorat*. Université Toulouse III (Paul Sabatier).
- Archanjo, C.J., 2003. Anisotropia de suscetibilidade magnética aplicada às rochas ígneas máficas intrusivas. *Tese de livre docência*. IGc – Universidade de São Paulo. São Paulo.
- Archanjo, C.J., Fetter, A.H. 2004. Emplacement setting of the granite sheeted pluton of Esperança (Brasiliano orogen, Northeastern Brazil). *Precambrian Research* 135, 193-215.
- Archanjo, C.J., Launeau, P., Hollanda, M.H.B.M., Macedo, J.W.P., Liu, D., 2009. Scattering of magnetic fabrics in the epizonal Cambrian alkaline granite of Meruoca (Ceará State, northeast Brazil). *International Journal of Earth Sciences* 98, 1793–1807, doi:10.1007/s00531-008-0342-z.
- Archanjo, C.J., Trindade, R.I.F., Bouchez, J.L., Ernesto, M., 2002. Granite fabrics and regional-scale strain partitioning in the Seridó belt (Borborema Province, NE Brazil). *Tectonics* 21, 1-14, doi: 10.1029/2000TC001269.
- Archanjo, C.J., Viegas, L.G., Hollanda, M.H.B.M., Souza, L.C., Liu, D., 2013. Timing of the HT/LP transpression in the Neoproterozoic Seridó Belt (Borborema Province, Brazil): Constraints from U-Pb (SHRIMP) geochronology and implications for the connections between NE Brazil and West Africa. *Gondwana Research* 23, 701-714, doi: <http://dx.doi.org/10.1016/j.gr.2012.05.005>.
- Arzi, A., 1978. Critical phenomena in the rheology of partially melted rocks. *Tectonophysics* 44, 173-184.
- Ashworth, J.R., 1985. *Migmatites*. Blackie, Glasgow.
- Atherton, M.P., 1993. Granite magmatism. *Journal of the Geological Society* 150, 1009-1023.
- Banerjee, S.K., 1991. Magnetic properties of Ti-Fe oxides. In: Lindsley, D.H. (Ed.), *Oxide minerals: petrologic and magnetic significance*. *Reviews in Mineralogy*. Mineralogical Society of America 25, 107-128.
- Bascou, J.F.L., 2002. Relações entre microestruturas, mecanismos de deformação e propriedades físicas anisotrópicas em rochas de alto grau de metamorfismo: estudo de alguns eclogitos e granulitos. *Tese de doutorado*, IGc – Universidade de São Paulo. São Paulo.
- Benn, K., 1994. Overprinting of magnetic fabrics in granites by small strains: numerical modeling. *Tectonophysics* 233, 153-162.
- Benn, K., 2009. Anisotropy of magnetic susceptibility fabrics in syntectonic plutons as tectonic strain markers: the example of the Canso pluton, Meguma Terrane, Nova Scotia. *Transactions of the Royal Society of Edinburgh* 100, 147–158.
- Bouchez, J.L., 2000. Anisotropie de susceptibilité magnétique et fabrique des granites. *Comptes Rendus de l'académie des Sciences de Paris* 330, 1-14.

- Bouchez, J.L., Gleizes, G., Djouadi, T., Rochette, P., 1990. Microstructure and magnetic susceptibility applied to emplacement kinematics of granites: the example of the Foix pluton (French Pyrenees). *Tectonophysics* 184, 157-171.
- Bouhallier, H., Choukroune, P., Ballèvre, M., 1993. Diapirism, Bulk Homogeneous Shortening and Transcurrent Shearing in the Archaean Dharwar Craton: the Holenarsipur area, southern India. *Precambrian Research* 63, 43-58.
- Borradaile, G.J., 1987. Magnetic susceptibility, petrofabrics and strain. *Tectonophysics* 156, 1-20.
- Borradaile, G.J., Henry, B., 1997. Tectonic applications of magnetic susceptibility and its anisotropy. *Earth Science Reviews* 42, 49-93.
- Borradaile, G.J., Jackson, M., 2004. Anisotropy of magnetic susceptibility (AMS): magnetic petrofabrics of deformed rocks. In: Martin-Hernandez, F., Lunenburg, C. M., Aubourg, C., Jackson, M. (Eds.), 2004. Magnetic fabric: methods and applications. Geological Society of London Special Publications 238, 299-360.
- Blumenfeld, P., Bouchez, J.L., 1988. Shear criteria in granite and migmatite deformed in the magmatic and solid states. *Journal of Structural Geology* 10, 361-372.
- Brito Neves, B.B., Passareli, C.R., Basei, M.A.S., Santos, E.J., 2003. Idades U-Pb em zircão de alguns granitos clássicos da Província Borborema. *Geologia USP Série Científica* 3, 25-38.
- Brown, M., 2001. Orogeny, migmatites and leucogranites: a review. *Proceedings of the Indian Academy of Sciences (Earth and Planetary Science)* 110, 313-336.
- Brown, M., 2007. Crustal melting and melt extraction, ascent and emplacement in orogens: mechanisms and consequences. *Journal of the Geological Society* 164, 709-730.
- Brown, M., Rushmer, T., 1997. The role of deformation in the movement of granite melt: views from the laboratory and the field. In: Holness, M. B. (Ed.), *Deformation-enhanced fluid transport in the Earth's Crust and Mantle*. The Mineralogical Society Series 8, Chapman & Hall, London. 111-144.
- Brown, M., Solar, G.S., 1998. Granite ascent and emplacement during contractional deformation in convergent orogens. *Journal of Structural Geology* 20, 1365-1393.
- Compston, W., Williams, I.S., Meyer, C., 1984. U-Pb geochronology of zircon from lunar breccia 73217 using a sensitive high mass-resolution ion microprobe. *Journal of Geophysical Research* 89, B525-B534.
- Corsini, M., Vauchez, A., Archanjo, C.J., Jardim de Sá, E.F., 1991. Strain transfer at a continental scale from a transcurrent shear zone to a transpressional fold belt: the Patos-Seridó belt system, north-eastern Brazil. *Geology* 19, 586-589.
- Corsini, M., Vauchez, A., Caby, R. 1996. Ductile duplexing at a bend of a continental scale strike-slip shear zone: example from NE Brazil. *Journal of Structural Geology* 18, 385-394.
- Corsini, M., Figueiredo, L.L., Caby, R., Féraud, G., Ruffet, G., Vauchez, A., 1998. Thermal history of the Pan-African/Brasiliano Borborema Province of northeast Brasil deduced from $^{40}\text{Ar}/^{39}\text{Ar}$ analysis. *Tectonophysics* 285, 103-117.
- Costa, A.C.D., 2002. Geologia e geocronologia Sm-Nd e U-Pb na região do Lineamento Patos, limite entre os blocos crustais do domínio Rio Grande do Norte e da Zona Transversal, Província Borborema. PhD thesis, State University of São Paulo.

- Chardon, D., Gapais, D., Cagnard, F., 2009. Flow of ultra-hot orogens: a view from the Precambrian, clues for the Phanerozoic. In: Chardon, D., Rey, P., (Eds.), *Hot Orogens. Special Issue, Tectonophysics* 477, 105-118.
- Charles, N., Faure, M., Chen, Y., 2009. The Montagne Noire migmatitic dome emplacement (French Massif Central): new insights from petrofabric and AMS studies. *Journal of Structural Geology* 31, 1423-1440.
- Daly, L., 1967. Possibilité d'existence dans les roches des plusieurs anisotropies magnétiques superpostes: leur separation. *Comptes Rendus de l'académie des Sciences de Paris* 264, 1377-1380.
- Denèle, Y., Olivier, P., Gleizes, G., Barbey, P., 2007. The Hospitalet gneiss dome (Pyrenees) revisited: lateral flow during Variscan transpression in the middle crust. *Terra Nova* 19, 445-453.
- Denèle, Y., Olivier, P., Gleizes, G., Barbey, P., 2009. Decoupling between the middle and upper crust during transpression-related lateral flow: Variscan evolution of the Aston gneiss dome (Pyrenees, France). In: Chardon, D., Rey, P. (Eds.), *Hot Orogens Special Issue, Tectonophysics* 477, 105-118.
- Dickin, A.P., 1995. *Radiogenic Isotope Geology*. Cambridge University Press.
- Dunlop, J.D., Ozdemir, O., 1997. *Rock magnetism fundamentals and frontiers*. Cambridge University Press.
- Egydio-Silva, M., Vauchez, A., Raposo, M.I.B., Bascou, J., Uhlein, A., 2005. Deformation regime variations in an arcuate transpressional orogen (Ribeira belt, SE Brazil) imaged by anisotropy of magnetic susceptibility in granulites. *Journal of Structural Geology* 27, 1750-1764.
- Ferré, E., Teyssier, C., Jackson, M., Thill, J.W., Rainey, E.S.G., 2003. Magnetic susceptibility anisotropy: a new petrofabric tool in migmatites. *Journal of Geophysical Research* 108, 2086, doi:10.1029/2002JB001790.
- Ferré, E.C., Martin-Hernandez, F., Teyssier, C., Jackson, M., 2004. Paramagnetic and ferromagnetic anisotropy of magnetic susceptibility in migmatites: measurements in high and low fields and kinematic implications. *Geophysical Journal International* 157, 1119-1129 doi: 10.1111/j.1365-246X.2004.02294.x.
- Fuller, M.D., 1963. Magnetic anisotropy and paleomagnetism. *Journal of Geophysical Research* 68, 293-309.
- Fritz, H., Tenczer, V., Hauzenberger, C., Wallbrecher, E., Muhongo, S., 2009. Hot granulite nappes – Tectonic styles and thermal evolution of the Proterozoic granulite belts in East Africa. In: Chardon, D., Rey, P., (eds) 2009, *Hot Orogens, Special Issue, Tectonophysics* 477, 160-173.
- Grégoire V., Saint Blanquat M., Nédélec A., Bouchez J.L., 1995. Shape anisotropy versus magnetic interactions of magnetite grains: experiments and application to AMS in granitic rocks. *Geophysical Research Letters* 22, 2765-2768.
- Hargraves R.B., Johnson D., Chan C.Y., 1991. Distribution anisotropy: the cause of AMS in igneous rocks? *Geophysical Research Letters* 18, 2193-2196.
- Higgins, M.D., 2006. *Quantitative textural measurements in igneous and metamorphic petrology*. Université du Québec à Chicoutimi, Québec, Canada.
- Hough, P.V.C., 1962. A method and means for recognizing complex patterns. U. S. Patent 3 069 654.
- Ishihara, S., 1977. The magnetite-series and ilmenite-series granitic rocks. *Mining geology* 27, 293-305.
- Jackson, M., 1991. Anisotropy of magnetic remanence: a brief review of mineralogical sources, physical origins and geological applications. *Pure and Applied Geophysics* 136, 1-28.
- Just, J., Kontny, A., Wall, H., Hirt, A.M., Martín-Hernández, F., 2004. Complex fabrics: Superposition and alteration development of magnetic fabrics during hydrothermal alteration in the Soultz-sous-Forêts granite

- from the EPS-1 borehole, Upper Rhine Graben. Geological Society of London, Special Publication 238, 509- 526, doi: 10.1144/GSL.SP.2004.238.01.26.
- Kohn, M.J., Northrup, C.J., 2009. Taking mylonites' temperatures. *Geology* 37, 47–50, doi:10.1130/G25081A.1.
- Kruckenberg, S.C., Ferré, E., Teyssier, C., Vanderhaeghe, O., Whitney, D., Seaton, N., Skord, J., 2010. Viscoplastic flow in migmatites deduced from fabric anisotropy: an example from the Naxos dome, Greece. *Journal of Geophysical Research* 115, B09401, doi:10.1029/2009JB007012.
- Lagroix, F., Borradaile, G.J., 2000. Magnetic fabric interpretation complicated by inclusions in mafic silicates. *Tectonophysics* 325, 207-225.
- Lloyd, G.E., 2004. Microstructural evolution in a mylonitic quartz simple shear zone: the significant roles of dauphine twinning and misorientation. In: Alsop, G.I., Holdsworth, R.E., McCaffrey, K.J.W., Hand M. (eds.), *Flow Processes in Faults and Shear zones*. Geological Society of London Special Publication 224, 39-61.
- Lloyd, G.E., Law, R.D., Schmid, S.M. 1987. A spherical electron channeling pattern map for use in quartz petrofabric analysis: correction and verification. *Journal of Structural Geology* 9, 251-253.
- Marchildon, N., Brown, M., 2003. Spatial distribution of melt-bearing structures in anatectic rocks from Southern Brittany, France: Implications for melt transfer from grain- to orogen-scale. *Tectonophysics* 364, 215-235.
- Mehnert, K.R., 1968. *Migmatites and the origin of granitic rocks*. Elsevier, Amsterdam, London.
- van der Molen, I., Paterson, S.R., 1979. Experimental deformation of partially-melted granite. *Contributions to Mineralogy and Petrology* 70, 299-318.
- Morales, L. F. G., 2007. Estudo da orientação cristalográfica em dobras, limites de grãos e anisotropia sísmica em muscovita-quartzo milonitos. Tese de Doutorado. Universidade Federal do Rio Grande do Sul.
- Nelson, K.D., Zhao, W., Brown, L.D., Kuo, J., Che, J., Liu, X., Klemperer, S.L., Makovsky, Y., Meissner, R., Mechie, J., Kind, R., Wenzel, F., Ni, J., Nabelek, J., Chen, L., Tan, H., Wei, W., Jones, A.G., Booker, J., Unsworth, M., Kidd, W.S.F., Hauck, M., Alsdorf, D., Ross, A., Cogan, M., Wu, C., Sandvol, E.A., Edwards, M., 1996. Partially molten middle crust beneath southern Tibet: synthesis of Project INDEPTH results. *Science* 274 (5293), 1684-1688.
- Neves, S.P., Vauchez, A., Archanjo, C.J., 1996. Shear zone controlled magma emplacement or magma-assisted nucleation of shear zones? Insights from northeast Brazil. *Tectonophysics* 262, 349-364.
- Neves, S.P., Mariano, G., Correria, P.B., Silva, J.M.R., 2006. 70 m.y. of synorogenic plutonism in eastern Borborema Province (NE Brazil): temporal and kinematic constraints on the Brasiliano Orogeny. *Geodinamica Acta* 19, 213-236.
- Nicolas, A., Bouchez, J.L., Boudier, F., 1972. Interprétation cinématique des déformations plastiques dans le massif de lherzolites de Lanzo (Alpes piémontaises). *Tectonophysics* 25, 233-260.
- Nicolas, A., Bouchez, J.L., Blaise, J., Poirier, J.P., 1977. Geological aspects of deformation in continental shear zones. *Tectonophysics* 42, 55-73.
- Nye, F.J., 1957. *Physical properties of crystals*. Oxford University Press, New York.

- Olivier, P., Archanjo, C.J., 1995. Magnetic and magmatic structures of the Emas granodioritic pluton (Cachoeirinha belt, NE Brazil): Relationships with Pan-African strike-slip fault systems. *Tectonophysics* 229, 239-250.
- Olivier, P., Saint-Blanquat, M., Gleizes, G., Leblanc, D., 1997. Homogeneity of granite fabrics at the metre and decametre scales. In: Bouchez, J.L. et al. (Eds.), *Granite: from segregation of melt to emplacement fabrics*. Kluwer Academic Publishers, Dordrecht, 113-128.
- Pitcher, W.S., 1993. *The Nature and Origin of Granite*. Chapman & Hall, London.
- Prior, D.J., Boyle, A.P., Brenker, F., Cheadle, M.C., Day, A., Lopez, G., Potts, G.J., Reddy, S., Spiess, R., Timms, N., Trimby, P., Wheeler, J., Zetterstrom, L., 1999. The application of electron backscatter diffraction and orientation contrast imaging in the SEM to textural problems in rocks. *American Mineralogist* 84, 1741-1759.
- Ramsay, J.G. & Graham, R.H., 1970. Strain variation in shear belts. *Canadian Journal of Earth Sciences* 7, 786-813.
- Randle, V., Engler, O., 2000. *Introduction to texture analysis*. Gordon and Breach Science.
- Rochette, P., Jackson, M., Aubourg, C., 1992. Rock magnetism and the interpretation of magnetic susceptibility. *Review Geophysics* 30, 209-226.
- Rodrigues, S.W.O., 2008. *Evolução estrutural brasileira da Província Borborema na região de Campina Grande (PB)*. Tese de doutorado. Universidade de São Paulo.
- Rosenberg, C., 2001. Deformation of partially molten granite: a review and comparison of experimental and natural case studies. *International Journal of Earth Sciences* 90, 60-76, doi: 10.1007/s005310000164.
- Rosenberg, C.L., & Handy, M.R. 2000. Syntectonic melt pathways during simple shearing of a partially molten rock analogue (Norcamphor-Benzamide). *Journal of Geophysical Research* 105(B2), 3135–3149.
- Rosenberg, C., Handy, M., 2005. Experimental deformation of partially melted granite revisited: implications for the continental crust. *Journal of Metamorphic Geology* 23, 19-28, doi: 10.1111/j.1525-1314.2005.00555.x.
- Rosenberg, C., Medvedev, S., Handy, M., 2006. Effects of melting on faulting and continental deformation. In: Handy, M., Hirth, G., Hovius, N. (Eds.), *Tectonic faults: Agents of change on a dynamic Earth*. Dahlem Workshop Reports, MIT Press, Cambridge.
- Rudnick, R.L., 1995. Making continental crust. *Nature* 378, 571-578.
- Rutter, E., Neumann, D.H.K., 1995. Experimental deformation of partially molten Westerly granite under fluid-absent conditions, with implications for the extraction of granitic magmas. *Journal of Geophysical Research* 100, 697-715.
- Sawyer, E.W., 1998. Formation and evolution of granite magmas during crustal reworking: the significance of diatexites. *Journal of Petrology* 39, 1147-1167.
- Sawyer, E.W., 2001. Grain-scale and outcrop-scale distribution and movement of melt in crystallizing granite. *Transactions of the Royal Society of Edinburgh: Earth Sciences* 91, 73–85.
- Sawyer, E.W., 2008. *Atlas of migmatites*. The Canadian Mineralogist, Special Publication 9, NRC Research Press, Ottawa, Ontario.
- Schmidt, N.H., Olesen, N.O., 1989. Computer-aided determination of crystal-lattice orientation from electron channeling patterns in the SEM. *Canadian Mineralogist* 27, 15-22.

- Schofield, D.L., D'Lemos, R.S., 1998. Relationships between syntectonic granite fabrics and regional PTtd paths: an example from Gander-Avalon boundary of NE Newfoundland. *Journal of Structural Geology* 20, 459-471.
- Schullmann, K., Edel, J., Hasalova, P., Cosgrove, J., Zek, J., Lexa, O., 2009. Influence of melt induced mechanical anisotropy on the magnetic fabrics and rheology of deforming migmatites, Central Vosges, France. *Journal of Structural Geology* 31, 1223–1237.
- Tarling, D.H., Hrouda, F., 1993. The magnetic anisotropy of rocks. Chapman & Hall, London.
- Trindade, R.I.F., 1999. Magnetismo de corpos graníticos e a evolução tectônica brasileira da porção ocidental da Faixa Seridó (NE do Brasil). Tese de doutorado. Universidade de São Paulo.
- Trindade, R., Bouchez, J.L., Bolle, O., Nédélec, A., Peschler, A., Poitrasson, F., 2001. Secondary fabrics revealed by remanence anisotropy: methodological study and examples from plutonic rocks. *Geophysical Journal International* 147, 310–318.
- Trindade, R., Raposo, M., Ernesto, M., Siqueira, R., 1999. Magnetic susceptibility and partial anhysteretic remanence anisotropies in the magnetite-bearing granite pluton of Tourão, NE Brazil. *Tectonophysics* 314, 443-468.
- Unsworth, M.J., Jones, A.G., Wei, W., Marquis, G., Gokarn, S.G., Spratt, J.E. (INDEPTH-MT TEAM) 2005. Crustal rheology of the Himalaya and Southern Tibet inferred from magnetotelluric data. *Nature* 438, 78–81.
- Vanderhaeghe, O., 2009. Migmatites, granites and orogeny: Flow modes of partially molten rocks and magmas associated with melt/solid segregation in orogenic belts. In: Chardon, D., Rey, P., (eds), *Hot Orogens*, Special Issue, *Tectonophysics* 477, 119-134.
- Van Schmus, W.R., Oliveira, E.P., Silva Filho, A.F., Toteu, S.F., Penaye, J., Guimarães, I.P., 2008. Proterozoic links between the Borborema Province, NE Brazil, and the Central African Fold Belt. In: Pankhurst, R.J., Trouw, R.A.J., Brito Neves, B.B., De Wit, M.J. (Eds.), *West Gondwana: Pre-Cenozoic Correlations Across the South Atlantic Region*: Geological Society of London, Special Publication 294, 69–99.
- Vaucher, A., Neves, S.P., Caby, R., Corsini, M.E., Egydio Silva, M., Arthaud, M., Amaro, V.E., 1995. The Borborema shear zone system, NE Brazil. *Journal of South American Earth Sciences* 8, 247-266.
- Vaucher, A., Egydio-Silva, M., Babinski, M., Tommasi, A., Uhlein, A., Li, D., 2007. Deformation of a pervasively molten middle-crust: Insights from the neoproterozoic Ribeira-Araçuaí orogen (SE Brazil). *Terra Nova* 19, 278-286, doi: 10.1111/j.1365-3121-2007.00747.
- Viegas, L.G.F., Archanjo, C., Vaucher, A., 2013. Fabrics of migmatites and the relationships between partial melting and deformation in high-grade transpressional shear zones: the Espinho Branco anatexite (Borborema Province, NE Brazil). *Journal of Structural Geology* 48, 45-56, doi: <http://dx.doi.org/10.1016/j.jsg.2012.12.008>.
- Wark, D.A., Watson, E.B., 2006. TitaniQ: A titanium-in-quartz geothermometer. *Contributions to Mineralogy and Petrology* 152, 743–754, doi:10.1007/s00410-006-0132-1.
- Weinberg, R., Sial, A.N., Mariano, G., 2004. Close spatial relationship between plutons and shear zones. *Geology* 32, 377-380.
- Whitney, D., Teyssier, C., Vanderhaeghe, O., 2004. Gneiss domes and crustal flow. *Geological Society of America Special Papers* 380, 15-33, doi: 10.1130/0-8137-2380-9.15.