

- ADAMS, D.B. (1979). Brain mechanisms for offense, defense and submission. *Behv. Brain Sci.*, 2, 201-241.
- AGUADO, L., HALL, G. & HARRINGTON, N. (1998). Illness-induced aversion learning in rats with lesion of the dorsal hippocampus. *Behav. Neurosci.*, 112, 1142-1151.
- ALBINSSON, A., ANDERSSON, G., ANDERSSON, K., VEGA-MATUSZCZYK, J. & LARSSON, K. (1996). The effects of lesions in the mesencephalic raphe systems on male rat sexual behavior and locomotor activity. *Behav. Brain Res.*, 80, 57-63.
- ANDRADE, T.G., SILVA, A.M., SILVA, C.L. & GRAEFF, F.G. (1999). Effect of eletrolytic lesion of the median raphe nucleus on behavioral and physiological measures of stress. *Acta Physiol. Pharmacol. Ther. Latinoam.*, 49 (4), 279-289.
- ANTONIADIS, E.A. & McDONALD, R.J. (1999). Discriminative fear conditioning to context expressed by multiple measures of fear in the rat. *Behav. Brain Res.*, 101, 1-13.
- ANTONIADIS, E.A. & McDONALD, R.J. (2000). Amygdala, hippocampus and discriminative fear conditioning to context. *Behav. Brain Res.*, 108, 1-19.
- ANTONIADIS, E.A. & McDONALD, R.J. (2001). Amygdala, hippocampus, and unconditioned fear. *Exp. Brain Res.*, 138, 200-209.
- AVANZI, V., CASTILHO, V.M., ANDRADE, T.G.C. & BRANDÃO, M.L. (1998). Regulation of contextual conditioning by the median raphe nucleus. *Brain Res.*, 790, 178-184.
- AZMITIA, E.C. & SEGAL, M. (1978). An autoradiographic analysis of the differential ascending projections of the dorsal and median raphe. *J. Comp. Neurol.*, 179, 641-668.
- AZRIN, N.H. (1960). Effects of punishment intensity during variable-interval reinforcement. *J. Exp. Anal. Behav.*, 3, 123-142.
- AZRIN, N.H., ULRICH, R.E., HUTCHINSON, R.R. & NORMAN, D.C. (1964). Effect of shock duration on shock-induced fighting. *J. Exp. Anal. Behav.*, 7, 9-11.
- BANDLER, R. & CARRIVE, P. (1988). Integrated defence reaction elicited by excitatory amino acid microinjection in the midbrain periaqueductal grey region of the unretrained cat. *Brain Res.*, 439, 95-106.
- BEAR, M.F., CONNORS, B.W. & PARADISO, M.A. (2001). *Neuroscience: exploring the brain*. Second edition. Williams & Wilkins, Baltimore – USA.

- BEITZ, A., CLEMENTS, J.R., MULLER, M. & ECKLUND, L. (1986). Differential origin of brainstem serotonergic projections to the midbrain periaqueductal gray and superior colliculus of the rat. *J. Comp. Neurol.*, 250, 498-509.
- BLANCHARD, D.C. & BLANCHARD, R.J. (1972). Innate and conditioned reactions to threat in rats with amygdaloid lesions. *J. Comp. Physiol. Psychol.*, 81 (2), 281-290.
- BLANCHARD, D.C. & BLANCHARD, R.J. (1988). Ethoexperimental approaches to the biology of emotion. *Ann. Rev. Psychol.*, 39, 43-68.
- BLANCHARD, R.J. & BLANCHARD, D.C. (1969). Crouching as an index of fear. *J. Comp. Physiol. Psychol.*, 67 (3), 370-375.
- BLANCHARD, R.J. & BLANCHARD, D.C. (1987). An ethoexperimental approach to the study of fear. *Psychol. Rec.*, 37, 305-316.
- BLANCHARD, R.J., DIELMAN, T.E. & BLANCHARD, D.C. (1968). Postshock crouching: familiarity with the stimulus situation. *Psychonom. Sci.*, 10, 371-372.
- BLANCHARD, R.J., FLANNELLY, K.J. & BLANCHARD, D.C. (1986). Defensive reactions of laboratory and wild *Rattus norvegicus*. *J. Comp. Physiol. Psychol.*, 100, 101-107.
- BOBILLIER, P., PETIJEAN, F., SALVERT, D., LIGIER, M. & SEGUIN, S. (1975). Differential projections of the nucleus raphe dorsalis and nucleus raphe centralis as revealed by autoradiography. *Brain Res.*, 85, 205-210.
- BOBILLIER, P., SEGIN, S., PETIJEAN, F., SALVERY, D., TOURET, M. & JOUVET, M. (1976). The raphe nuclei of the cat brainstem: a topographical atlas of their efferent projections as revealed by autoradiography. *Brain Res.*, 113, 449-486.
- BOLLES, R.C. & COLLIER, A.C. (1976). The effect of predictive cues on freezing in rats. *Animal Learn. Behav.*, 4 (1A), 6-8.
- BOLLES, R.C. (1970). Species- specific defensive reactions and avoidance learning. *Psychol. Rev.*, 71, 32-48.
- BOLLES, R.C. & FANSELOW, M.S. (1980). A perceptual-defensive-recuperative model of fear and pain. *Behav. Brain Sci.*, 3, 291-301.
- BOLLES, R.C. & RILEY, A.L. (1973). Freezing as an avoidance response: another look at the operant-respondent distinction. *Learn. Motiv.*, 4, 268-275.

- BOUTON, M.E. & BOLLES, R.C. (1980). Conditioned fear assessed by freezing and by the suppression of three different baselines. *Animal Learn. Behav.*, 8 (3), 429-434.
- BRADY, J.V. & NAUTA, W.J.H. (1955). Subcortical mechanisms in emotional behavior: the duration of affective changes following septal and habenular lesions in the albino rat. *J.Comp. Physiol. Psychol.*, 48, 412-420.
- BRANDÃO, M.L., ANSELONI, V.Z., PANDOSSIO, J.E., De ARAUJO, J.E. & CASTILHO, V.M. (1999). Neurochemical mechanisms of the defensive behavior in the dorsal midbrain, *Neurosci. Biobehav. Rev.*, 23: 863-875.
- BRILEY, M., CHOPIN, P. & MORET, C. (1990). Effect of serotonergic lesion on “anxious” behaviour measured in the elevated plus-maze test in the rat. *Psychopharmacology*, 101, 187-189.
- CAMPEAU, S. & DAVIS, M. (1995). Involvement of the central nucleus and basolateral complex of the amygdala in fear conditioning measured with fear-potentiated startle in rats trained concurrently with auditory and visual conditioned stimuli. *J. Neurosci.*, 15, 2301-2311.
- CARLI, M. & SAMANIN, R. (1992). 8-Hydroxy-2-(di-n-propylamino) tetralin, a 5-HT_{1A} agonist, impairs performance in a passive avoidance task. *Eu. J. Pharmacol.*, 105, 720-726.
- CARRIVE, P., LEE, J. & SU, A. (2000). Lidocaine blockade of amygdala output in fear-conditioned rats reduces Fos expression in the ventrolateral periaqueductal gray. *Neuroscience*, 95(4), 1071-1080.
- CARRIVE, P., LEUNG, P., HARRIS, J. & PAXINOS, G. (1997). Conditioned fear to context is associated with increased Fos expression in the caudal ventrolateral region of the midbrain periaqueductal gray. *Neuroscience*, 78(1), 165-177.
- CASTILHO, V.M., AVANZI, V. & BRANDÃO, M.L. Antinociception elicited by aversive stimulation of the inferior colliculus, *Pharmacol. Biochem. Behav.*, 62: 425-431, 1999.
- CHAN, K-H, MORELL, J.R., JARRARD, L.E. & DAVIDSON, T.L. (2001). Reconsideration of the role of hippocampus in learned inhibition. *Behav. Brain Res.*, 119, 111-130.
- CHEETHAN, S.C. (1990). Brain 5-HT_{1A} binding sites in depressed suicides. *Psychopharmacology*, 102, 544-548.

- CHOPIN, P. & BRILEY, M. (1987). Animal models of anxiety: the effects of compounds that modify 5-HT neurotransmission. *TIPS.*, 8, 383-388.
- COIMBRA, N.C. & BRANDÃO, M.L. (1997). Effects of 5-HT₂ receptors blockade on fear-induced analgesia elicited by electrical stimulation of the deep layers of the superior colliculus and dorsal periaqueductal gray. *Behav. Brain Res.*, 87, 97-103.
- COIMBRA, N.C., TOMAZ, C. & BRANDÃO, M.L. (1992). Evidence for the involvement of serotonin in the antinociception induced by electrical or chemical stimulation of the mesencephalic tectum. *Behav. Brain Res.*, 50, 77-83.
- CONNOR, H.E. & HIGGINS, G.A. (1990). Cardiovascular effects of 5-HT_{1A} receptor agonists injected into the dorsal raphe nucleus of conscious rats. *Eur. J. Pharmacol.*, 182, 63-72.
- CORCORAN, K.A. & MAREN, S. (2001). Hippocampal inactivation disrupts contextual retrieval of fear memory after extinction. *J. Neurosci.*, 21 (5), 1720-1726.
- DAHLSTRÖM, A. & FÜXE, K. (1964). Evidence for the existence of monoamine neurons in the central nervous system. I. Demonstration of monoamines in the cell bodies of brain stem neurons. *Acta Physiol. Scand.*, 62 (supl. 232), 1-55.
- DAHLSTRÖM, A. & FÜXE, K. (1965). Evidence for the existence of monoamine neurons in the central nervous system. II. Experimentally induced changes in the intraneuronal amine levels of bulbospinal neuron systems. *Acta Physiol. Scand.*, 64 (supl. 247), 1-35.
- DARWIN, C. (1872). The expression of emotion in men and animals. *Philosophical Library*, New York (reprint 1985), pp. 134.
- DAVIS, M. (1992). The role of the amigdala in conditioned fear. Em: Aggleton, J.P. (Ed.), *The amigdala: neurobiological aspects of emotion, memory and mental dysfunction*. New York, Wiley-Liss. pp. 255-305.
- De ALMEIDA, R.M.M., GIOVENARDI, M., CHARCHAT, H. & LUCION, A.B. (1998). 8-OH-DPAT in the median raphe nucleus decreases while in the medial septal area it may increase anxiety in female rats. *Neurosci. Biobehav. Rev.*, 23, 259-264.
- De OCA, B.M., De COLA, J.P. MAREN, S. & FANSELOW, M.S. (1998). Distinct regions of the periaqueductal gray are involved in the acquisition and expression of defensive responses. *J. Neurosci.*, 18, 3426-3432.

- DEAKIN, J.W.F. & GRAEFF, F.G. (1991). 5-HT mechanisms of defense, *J. Psychopharmacol.*, 5, 305-315.
- DEAKIN, J.F.W., GUIMARÃES, F.S. & GRAEFF, F.G. (1994). Testing 5-HT theories of anxiety in normal volunteers. Em: PALOMO, T. & ARCHER, T. *Strategies for studying brain disorders, vol. 1. Depressive, anxiety and drug abuse disorders*. Madrid, Editorial Complutense, pp. 211-238.
- DRATCU, L. & LADER, M. (1993). Ansiedade: conceito, classificação e biologia. *J. Bras. Psiq.*, 42, 19-32.
- DREYER, P.I. & CHURCH, R. (1968). Shock-induced fighting as a function of the intensity and duration of the aversive stimulus. *Psychonom. Sci.*, 10, 271- 272.
- ERSPAMER, V. & ASERO, B. (1952). Identification of enteramine, the specific hormone of enterochromaffin cell system, as 5-hidroxytryptamine. *Nature*. 169, 800-801.
- FALCK, B., HILLARP, N.A., THIEME, G. & TORP, A. (1962). Fluorescence of catecholamines and related compounds condensed with formaldehyde. *J. Histochem. Cytochem.*, 10, 348-354.
- FANSELOW, M.S. (1982). The postshock activity burst, *Animal Learn. Behav.* 4, 448-454.
- FANSELOW, M.S. (1984). What is conditioned fear?, *TINS*, 7, 460-462.
- FANSELOW, M.S., (1986). Associative vs topographical accounts of the immediate shock-freezing deficit in rats: Implication for the response selection rules governing species-specific defensive reactions. *Learn. Motiv.*, 17, 16-39.
- FANSELOW, M.S. (1991). The midbrain periaqueductal gray as coordinator of action in response to fear and anxiety. Em: Depaulis A., Bandler R. (Eds.), *The midbrain periaqueductal gray matter: functional, anatomical and immunohistochemical organization*. New York, Plenum Publishing. pp. 151-173.
- FANSELOW, M.S. (2000). Contextual fear, gestalt memories, and the hippocampus. *Behav. Brain Res.*, 110, 73-81.
- FANSELOW, M.S. & BAACKES, M.P. (1982). Conditioned fear-induced opiate analgesia on the formalin test: evidence for two aversive motivational systems. *Learn. Motiv.*, 13, 200-221.

- FANSELOW, M.S. & BOLLES, R.C. (1979). Triggering of the endorphin analgesic reaction by a cue previously associated with shock: Reserval by naloxone. *Bull. Psychonom. Soc.*, 14 (2), 88-90.
- FANSELOW, M.S. & HELMSTETTER, F.J. (1988). Conditional analgesia, defensive freezing, and benzodiazepines. *Behav. Neurosci.*, 102 (2), 233-243.
- FANSELOW, M.S., De COLA, J.P., De OCA, B.M. & LANDEIRA-FERNADEZ, J. (1995). Ventral and dorsolateral regions of the midbrain periaqueductal gray (PAG) control different stages of defensive behavior: Dorsolateral PAG lesions enhance the defensive freezing produced by massed and immediate shock. *Aggressive Behav.*, 21, 63-67.
- FANSELOW, M.S., LESTER, L.S. & HELMSTETTER, F.J. (1988). Changes in feeding and foraging patterns as an antipredator defensive strategy: A laboratory stimulation using aversive stimulation in a closed economy. *J. Exp. Anal. Behav.*, 50, 361-374.
- FANSELOW, M.S., SIGMUNDI, R.A. & BOLLES, R. C. (1980). Naloxone pretreatment enhances shock-elicited aggression. *Physiol. Psychol.*, 8 (3), 369-371.
- FENDT, M. & FANSELOW, M.S. (1999). The neuroanatomical and neurochemical basis of conditioned fear. *Neurosci. Biobehav. Rev.*, 23, 743-760.
- FILE, S.E. (1992). Behavioural detection of anxiolytic action. Em: Elliot, J.M.; Heal, D.J. & Marsden, C.A. (Orgs), *Experimental Approaches to anxiety and depression*. John Wiley & Sons Ltd, Chichester, 25-44.
- FRANKLAND, P.W., DOCKSTADER, C.L. & McDONALD, R.J. (1998). Discriminative and nondiscriminative contextual fear conditioning potentiate the acoustic startle reflex. *Psychobiology*, 26, 267-264.
- FUXE, K. (1965). Evidence for the existence of monoamine containing neurons in the central nervous system. IV. The distribution of monoamine terminals in the central nervous system. *Acta Physiol. Scan.*, 64 (supl. 247), 37-84.
- GADDUM, J.H. (1953). Antagonism between lysergic acid diethylamide and 5-hydroxytryptamine. *J. Physiol.*, 121 (sup.), p. 15.
- GELLER, I. & BLUM, K. (1970). The effects if 5-HTP on para-chlorophenylalanine (PCPA) attenuation on “conflit” behavior. *Eu. J. Pharmacol.*, 9, 319-324.

- GELLER, I. & SEIFTER, J. (1960). The effects of meprobamate, barbiturates, d-amphetamine and promazine on experimentally induced conflict in the rat. *Psychopharmacology*, 1, 482-492.
- GERSON, S.C. & BALDESSARINI, R.J. (1980). Motor effects of serotonin in the central nervous system. *Life Sci.*, 27, 1435-1451.
- GEWIRTZ, J.C., McNISH, K.A. & DAVIS, M. (2000). Is the hippocampus necessary for contextual fear conditioning?. *Behav. Brain Res.*, 110, 83-95.
- GEYER, M.A., PUERTO, A., DAWSEY, W.J., KNAPP, S., BULLARD, W.P. & MANDELL, A.J. (1976). Histologic and enzymatic studies of the mesolimbic and mesostriatal serotonergic pathways. *Brain Res.*, 106, 241-256.
- GOOSENS, K.A. & MAREN, S. (2001). Contextual and auditory fear conditioning are mediated by lateral, basal, and central amygdaloid nuclei in rats. *Learn. Mem.*, 8 (3), 148-155.
- GOZLAN, H., EI MESTIKAWY, S., PICHAT, L., GLOWINSKI, J. & HAMON, M. (1983). Identification of presynaptic serotonin autoreceptors using a new ligand. *Nature*, 305, 140-142.
- GRAEFF, F.G. (1981). Minor tranquilizers and brain defense system, *Braz. J. Med. Biol. Res.*, 14, 239-265.
- GRAEFF, F.G. (1990). Brain defense systems and anxiety, Em: BURROWS, G.D., ROTH, M. & NOYERS Jr., R. (eds). *Handbook of Anxiety, Vol. 3: The Neurobiology of Anxiety*. Elsevier Science Publishers, Amsterdam, pp. 307-354.
- GRAEFF, F.G. (1994). Neuroanatomy and neurotransmitter regulation of defensive behaviors and related emotions in mammals. *Braz. J. Med. Biol. Res.*, 27, 811-829.
- GRAEFF, F.G. (1997). Serotonergic systems. Em: *Neuropsychiatry of the basal ganglia*, 20 (4), 723-739.
- GRAEFF, F.G. (1999). Ansiedade, Em: GRAEFF, F.G. & BRANDÃO, M.L. (1999). *Neurobiologia das doenças mentais*. Quinta Edição. São Paulo, Lemos Editorial & Gráficos Ltda, 109-144.

- GRAEFF, F.G. & BRANDÃO, M.L. (1999). *Neurobiologia das doenças mentais*. Quinta Edição. São Paulo, Lemos Editorial & Gráficos Ltda.
- GRAEFF, F.G. & GUIMARÃES, F.S. (1999). *Fundamentos de Psicofarmacologia*. Ed. Atheneu.
- GRAEFF, F.G. & SCHOENFELD, R.I. (1970). Tryptaminergic mechanisms in punished and nonpunished behavior. *J.Pharmacol. Experim. Therap.*, 173, 277-283.
- GRAEFF, F.G. & SILVEIRA FILHO, G. (1978). Behavioral inhibition induced by electrical stimulation of the median raphe nucleus of the rat. *Physiol. Behav.*, 71, 477-484.
- GRAEFF, F.G., GUIMARÃES, F.S., ANDRADE, T.G.C. & DEAKIN, J.F.W. (1996). Role of 5-HT in anxiety, stress and depression. *Pharmacol. Biochem. Behav.*, 54, 129-141.
- GRAY, J.A. (1982). *The neuropsychology of anxiety*. New York, Oxford University Press.
- GRAY, J.A. (1987). *The psychology of fear and stress*. Cambridge, Cambridge University Press.
- GRAY, J.A. & McNAUGHTON, N. (2000). *The neuropsychology of anxiety*. Second Edition, New York, Oxford University Press.
- HALL, J.R.S. & WARACZYNSKI, M. (1991). The effect of excitoxin lesions of the lateral hypothalamus on self-stimulation reward. *Brain Res.*, 541, 29-40.
- HALL, M.D., EL MESTIKAWY, S., EMIRIT, M.B., PICHAT, L., HAMON, M. & GOZLAN, H. (1985). [³H]8-hidroxy-2-(di-n-propylamino) tetralin binding to pre- and pos-synaptic 5-hydroxytryptamine sites in various regions of the brain. *J. Neurochem.*, 44, 1685-96.
- HANDLEY, S.L. & McBLANE, J.W. (1993). Serotonin mechanisms in animal models of anxiety. *Braz. J. Med. Biol. Res.*, 26, 1-13.
- HARING, J. H. (1991). Reorganization of the area dentata serotonergic plexus after lesions of the median raphe nucleus. *J. Comp. Neurol.*, 306, 576-584.
- HARING, J.H. & DAVIS, J.N. (1985). Differential distribution of locus coeruleus projections to the hippocampal formation: anatomical and biochemical evidence. *Brain Res.*, 325, 366-369.

- HASHIMOTO, S., INOUE, T. & KOYAMA, T. (1997). Effects of the co-administration of 5-HT_{1A} receptor antagonists with an SSRI in conditioned fear stress-induced freezing behavior. *Pharmacol. Biochem. Behav.*, 58 (2), 471-475.
- HASHIMOTO, S., INOUE, T. & KOYAMA, T. (1999). Effects of conditioned fear stress on serotonin neurotransmission and freezing behavior in rats. *Eur. J. Pharmacol.*, 378, 23-30.
- HELMSTETTER, F.J. & BELLGOWAN, P.S. (1993). Lesions of the amygdala block conditional hypoalgesia on the tail flick test. *Brain Res.*, 612: 253-257.
- HELMSTETTER, F.J. & TERSHNER, S.A. (1994). Lesions of the periaqueductal gray and rostral ventromedial medulla disrupt antinociceptive but not cardiovascular aversive conditional responses. *J. Neurosci.*, 14(11), 7099-7108.
- HELMSTETTER, F.J. (1992). The amygdala is essential for the expression of conditional hypoalgesia. *Behav. Neurosci.*, 106, 518-528.
- HEURING, R.E. & PEROUTKA, S.J. (1987). Characterization of a novel [³H] 5-hydroxytryptamine binding site subtype in bovine brain membranes. *J. Neurosci.*, 7, 894-903.
- HILLEGART, V. (1990). Effects of application of 5-HT and 8-OH-DPAT into the dorsal and median raphe nuclei on motor activity in the rat. *Physiol. Behav.*, 48, 143-148.
- HILLEGART, V. & HJORTH, S. (1989). Median raphe, but not dorsal raphe, application of the 5-HT_{1A} agonist 8-OH-DPAT stimulates rat motor activity. *Eu. J. Pharmacol.*, 160, 303-307.
- HILLEGART, V., AHLENIUS, S. & LARSSON, K. (1991). Region-selective inhibition of male rat sexual behavior and motor performance by localized forebrain 5-HT injections: a comparison with effects produced by 8-OH-DPAT. *Behav. Brain Res.*, 42, 169-180.
- HITCHCOCK, J.M. & DAVIS, M. (1986). Lesions of amygdala, but not the cerebellum or red nucleus, block conditioned fear as measured with the potentiated startle paradigm. *Behav. Neurosci.*, 100, 11-22.
- HJORTH, S. & MAGNUSSON, T. (1988). The 5-HT_{1A} receptor agonist, 8-OH-DPAT, preferentially activates cell body 5-HT autoreceptors in rat brain in vivo. *Arch. Pharmac.* 388, 463-471.

- HOFFER, M.A. (1970). Cardiac and respiratory function during sudden prolonged immobility in wild rodents. *Psychonom. Med.*, 633-647.
- HOLE, K., FUXE, K. & JONHSON, G. (1976). Behavioural effects of 5,7-dihydroxytryptamine lesions of ascending 5-hydroxytryptamine pathways. *Brain Res.*, 107, 385-399.
- HOYER, D., ENGEL, G. & KALKMAN, H.O. (1985). Molecular pharmacology of 5-HT₁ and 5-HT₂ recognition sites in rat and pig membranes: radioligand binding studies with [³H] 5-HT, [³H] 8-OH-DPAT, (-)[¹²⁵I]iodocyanopindolol, [³H] mesulergine and [³H] ketanserin. *Eu. J. Pharmacol.*, 118, 13-23.
- HUMPHREY, P.P.A., HARTIG, P. & HOYER, D. (1993). A proposed new nomenclature for 5-HT receptors. *TIPS*, 14, 233-236.
- INOUE, T. TSUCHIYA, K. & KOYAMA, T. (1996). Serotonergic activation reduces defensive freezing in the conditioned fear paradigm. *Pharmacol. Biochem. Behav.*, 53 (4), 825-831.
- IWATA, J., LeDOUX, J.E., MEELEY, M.P., ARNERIC, S. & REIS, D.J. (1986). Intrinsic neurons in the amygdaloid field projected to by the medial geniculate body mediate emotional responses conditioned to acoustic stimuli. *Brain Res.*, 383, 195-214.
- JACOBS, B.L., WISE, W.D. & TAYLOR, K.M. (1974). Differential behavioral and neurochemical effects following lesions of the dorsal or median raphe nuclei in rats. *Brain Res.*, 79, 353-361.
- KAPP, B.S., FRYSSINGER, R.C., GALLAGHER, M. & HASELTON, J. (1979). Amygdala central nucleus lesions: Effects on heart rate conditioning in the rabbit. *Physiol. Behav.*, 23, 1109-1117.
- KIERMAN, M. & CRANNEY, J. (1992). Excitotoxic lesions of the central nucleus of the amygdala but not of the periaqueductal gray block integrated fear responding as indexed by both freezing responses and augmentation of startle. *Soc. Neurosci. Abstr.*, 18 (653.9), 1566-1567.
- KILPATRICK, G.J., JONES, B.J. & TYERS, M.B. (1987). Identification and distribution of 5-HT₃ receptors in rat brain using radioligand binding. *Nature*, 330, 746-748.
- KIM, J.J. & FANSELOW, M.S. (1992). Modality-specific retrograde amnesia of fear. *Science*, 256, 675-677.

- KIM, J.J., RISON, R.A. & FANSELOW, M.S. (1993). Effects of amygdala, hippocampus, and periaqueductal gray lesions on short- and long-term contextual fear. *Behav. Neurosci.*, 107, 1093-1098.
- KIMURA, D. (1958). Effects of selective hippocampal damage on avoidance behavior in the rat. *Can. J. Psychol.*, 12, 213-218.
- KOHLER, C. & SCHWARCZ, R. (1983). Comparison of ibotenic and kainic neurotoxicity in the rat brain: a histological study. *Neuroscience.*, 8, 819- 835.
- KOSOFSKY, B.E. & MOLLIVER, M.E. (1987). The serotonergic innervation of cerebral cortex: different classes of axon terminals arise from dorsal and median raphe nuclei. *Synapse*, 1, 153-168.
- KRIEGER, J.E. & GRAEFF, F.G. (1985). Defensive behaviour and hypertension induced by glutamate in the midbrain central gray of the rat. *Braz. J. Med. Biol. Res.*, 18, 61-67.
- LeDOUX, J.E. (1994) Emotion Memory and the Brain. *Myst. Mind.*, 68-75.
- LeDOUX, J.E. (1996). A few degrees of separation. Em: LeDoux, J.E. *The Emotional Brain*, New York, pp.139-178.
- LeDOUX, J.E., FARB, C. & RUGGIERO, D.A. (1990a). Topographic organization of neurons in the acoustic thalamus that project to the amygdala. *J. Neurosci.*, 10, 1043-1054.
- LeDOUX, J.E., CICCETTI, P., XAGORARIS, A. & ROMANSKI, L.M. (1990b). The lateral amygdaloid nucleus: Sensory interface of the amygdala in fear conditioning. *J. Neurosci.*, 10, 1062-1069.
- LeDOUX, J.E., IWATA, J., CICCETTI, P. & REIS, D.J. (1988). Different projections of the central amygdaloid nucleus mediate autonomic and behavioral correlates of conditioned fear. *J. Neurosci.*, 8, 2517-2529.
- LeDOUX, J.E., RUGGIERO, D.A., FOREST, R., STORNETTA, R. & REIS, D.J. (1987). Topographic organization of convergent projections to the thalamus from the inferior colliculus and spinal cord in the rat. *J. Comp. Neurol.*, 264, 123-146.
- LENT, R. (2001). *Cem bilhões de neurônios: conceitos fundamentais de Neurociência*. São Paulo: Editora Atheneu.

- LI, X.B., INOUE, T., HASHIMOTO, S. & KOYAMA, T. (2001). Effect of chronic administration of flesinoxam and fluvoxamine on freezing behavior induced by conditioned fear. *Eu. J. Pharmacol.*, 425, 43-50.
- LIEBMAN, J.M., MAYER, D.J. & LIEBSKIND, J.C. (1970). Mesencephalic central gray lesions and fear-motivated behavior in rats. *Brain Res.*, 23, 353-370.
- LISTER, R.G. (1990). Ethologically-based animal models of anxiety disorders. *Pharmacol. Ther.*, 46, 321-340.
- LORENS, S.A. (1978). Some behavioral effects of serotonin depletion depend on method: a comparison of 5,7-dihydroxytryptamine, p-chlorophenylalanina, p-chloramphetamine and electrolytic raphe nucleus. *Ann. N.Y. Acad. Sci.*, 305, 522-555.
- LOVICK, T.A., (1991). Interactions between pathways from the dorsal and ventrolateral periaqueductal gray matter in the rat. Em: Depaulis, A. & Bandler, R. (eds.). *The Midbrain Periaqueductal Gray Matter*, Plenum Press, New York, pp.101-120.
- MAISONNETTE, S.S., KAWASAKI, M.C., COIMBRA, N.C. & BRANDÃO, M.L. (1996). Effects of lesion amygdaloid nuclei and substantia nigra on aversive responses induced by electrical stimulation of the inferior colliculus. *Brain Res. Bull.*, 40 (2), 93-98.
- MAISONNETTE, S.S., MORATO, S. & BRANDÃO, M.L. (1993). Role of resocialization and of 5-HT_{1A} receptor activation on the anxiogenic effects induced by isolation in the elevated plus-maze test. *Physiol. Behav.*, 54, 753-758.
- MAMOUNAS, L.A. & MOLLIVER, M.E. (1988). Evidence for dual serotonergic projections to neocortex: axons from the dorsal and median raphe nuclei are differentially vulnerable to the neurotoxin p-chloroamphetamine (PCA). *Exp. Neurol.*, 102, 23-36.
- MAREN, S. (1999). Neurotoxic basolateral amygdala lesions impair learning and memory but not the performance of conditional fear in rats. *J. Neurosci.*, 19 (19), 8696-8703.
- MAREN, S. & HOLT, W. (2000). The hippocampus and contextual memory retrieval in Pavlovian conditioning. *Behav. Brain Res.*, 110, 97-108.
- MAREN, S., AHARONOV, G. & FANSELOW, M.S. (1997). Neurotoxic lesions of the dorsal hippocampus and Pavlovian fear conditioning in rats. *Behav. Brain Res.*, 88, 261-274.

- MARSDEN, C.A. & CURZON, G. (1976). Studies on the behavioural effects of tryptophan and p-chorophenylalanine. *Neuropharmacology*, 15, 165-170.
- MARTIN, G.R. & HUMPHREY, P.P.A. (1994). Receptors for 5-hidroxytryptamine: current perspectives on classification and nomenclature. *Neuropharmacology*, 33 (3/4), 261-273.
- McKENNA, J.T. & VERTES, R.P. (2001). Collateral projections from the median raphe nucleus to the medial septum and hippocampus. *Brain Res. Bull.*, 54 (6), 619-630.
- McNISH, K.A., GEWIRTZ, J.C. & DAVIS, M. (1997). Evidence of contextual fear after lesions of the hippocampus: A disruption of freezing but not fear-potentiated startle. *J. Neurosci.*, 17, 9253-9360.
- MELIK, E., BABAR-MELIK, E., OZGUNEN, T. & BINOKAY, S. (2000) Median raphe nucleus mediates forming long-term but not short term contextual fear conditioning in rats. *Behav. Brain Res.*, 112, 145-150.
- MELO, L.L., CARDOSO, S.H. & BRANDÃO, M.L. (1992). Antiaversive action of benzodiazepines on escape behavior induced by electrical stimulation of the inferior colliculus. *Physiol. Behav.*, 51, 557-562.
- MENDONÇA NETTO, S. & GUIMARÃES, F.S. (1996). Role of hippocampal 5-HT_{1A} receptors on elevated plus maze exploration after a single restraint experience. *Behav. Brain Res.*, 77, 215-218.
- MOLLIVER, M.E. (1987). Serotonergic neuronal systems: what their anatomic organization tells us about function. *J. Clinical Pharmacol.*, 7, 3-23.
- MONGEAU, R., BLIER, P. & De MONTIGNY, C. (1997). The serotonergic and noradrenergic systems of the hippocampus: their interactions and the effects of antidepressant treatments. *Brain Res. Rev.*, 23, 145-195.
- MORRIS, R.G., ANDERSON, E., LYNCH G.S. & BAUDRY, M. (1986). Selective impairment of learning and blockade of long-term potentiation by an N-methyl-D-aspartate receptor agonist, AP5. *Nature*, 319, 774-776.
- MURUA, V.S. & MOLINA, V.A. (1991). Chronic antidepressant counteracts the conditioned analgesia induced by a context previously associated with inescapable shock. *Psychopharmacology*, 105, 439-441.

- MYERS, R.D. (1981). Serotonin and thermoregulation: old and news views. *J. Physiol.*, 77, 505-513.
- NADER, K., MAJIDISHAD, P., AMORAPANTH, P. & LeDOUX, J.E. (2001). Damage to lateral and central, but not other, amygdaloid nuclei prevents the acquisition of auditory fear conditioning. *Learn. Mem.*, 8 (3), 156-163.
- NIEUWENHUYS, R. (1985). *Chemoarchitecture of the brain*. Berlin, Springer - Verlag. pp. 33-41.
- NUTT, D.J. (1990). The pharmacology of human anxiety. *Pharmacol. Ther.* 47, 233-266.
- OBRIST, P. A., SUTTERER, J. R. & HOWARD, J. L. (1972). Preparatory cardiac changes: A psychobiological approach. Em: BLACK, A. H. e PROKASY, W. F. (eds.). *Classical conditioning II: Current theory and research*. Appleton, New York, pp. 312-340.
- OLTON, D.S., BECKER, J.T. & Handelmann, G.E. (1979). Hippocampus, space, and memory. *Behav. Brain Sci.*, 2, 313-365.
- PADOVAN, C.M. & GUIMARÃES, F.S. (1993). Attenuation of behavioral consequences of immobilization stress by intra-hippocampal microinjection of zimelidine. *Braz. J. Med. Biol. Res.*, 26, 1085-1089.
- PALACIOS, J.M. (1990). Distribution of serotonin receptors. *Ann. N. Y. Acad. Sci.*, 600: 36-52.
- PAPAIOANNOU, V.E. & FOX, J.G. (1993) Efficacy of tribromoethanol anesthesia in mice. *Lab. Animal Sci.*, 43, 189-192.
- PAXINOS, G. & WATSON, C. (1997). *The rat brain in stereotaxic coordinates*. Third Edition. New York, Academic Press.
- PAZOS, A. & PALACIOS, J.M. (1985). Quantitative autoradiographic mapping of serotonin receptors in the rat brain I. Serotonin-1 receptors. *Brain Res.*, 346, 205-230.
- PAZOS, A; PROBST, A. & PALACIOS, J.M. (1987). Serotonin receptors in the human brain - III. Autoradiographic mapping of serotonin - 1 receptors. *Neuroscience*, 21, 97-122.
- PEDIGO, N.W., YAMAMURA, H.I. & NELSON, D.L. (1981) Discrimination of multiple [³H] 5-hydroxytryptamine binding sites by the neuroleptic spiperona in rat brain. *J. Neurochem.*, 36, 220-226.

- PELLOW, S., CHOPIN, P., FILE, S.E. & BRILEY, M. (1985). Validation of open: closed arm entries in an elevated plus-maze as a measure of anxiety in the rat. *J. Neurosci. Meth.*, 14, 149-167.
- PEROUTKA, S.J. & SNYDER, S.H. (1979). Multiple serotonin receptors differential binding of [³H] 5-hydroxytryptamine, [³H] lysergic acid diethylamide and [³H] spiroperidol. *Mol. Pharmacol.*, 16, 687-699.
- PHILLIPS, R.G. & LeDOUX, J.E. (1992). Differential contributions of amygdala and hippocampus to cued and contextual fear conditioning. *Behav. Neurosci.*, 106, 274-285.
- PHILLIPS, R.G. & LeDOUX, J.E. (1994). Lesions of the dorsal hippocampal formation interfere with background but not foreground contextual fear conditioning. *Learn. Mem.*, 1, 34-44.
- PIRES, J.G., SILVA, S.R., RAMAGE, A.G. & FUTURO-NETO, H.A. (1993). Influence of median raphe nucleus lesions on neuroleptic- induced catalepsy and on the anticataleptic effect of buspirone. *Braz. J. Med. Biol. Res.*, 26, 323-326.
- RAPPORT, M.M., GREEN, A.A. & PAGE, I.H. (1948). Serum vasoconstrictor (serotonin). Isolation and characterization. *J. Biol. Chem.*, 176, 1243-1251.
- REDISH, A.D. (2001). The hippocampal debate: are we asking the right questions? *Behav. Brain Res.*, 127, 81-98.
- ROBICHAUD, R.C. & SLEDGE, K.L. (1969). The effects of p-chlorophenylalanine on experimentally induced conflict in the rat. *Life Sci.*, 8, 965-968.
- SANANES, C.B. & DAVIS, M. (1992). N-methyl-D-aspartate lesions of the lateral and basolateral nuclei of the amygdala block fear-potentiated startle and shock sensitization of startle. *Behav. Neurosci.*, 106, 72-80.
- SCHREIBER, R. & De VRY, J.D. (1993). Studies on the neural circuits involved in the discriminative stimulus effects of 5-HT_{1A} receptor agonist in the rat. *J. Pharmacol. Exp. Ther.*, 265, 572-579.
- SHARP, T., BRAMWELL, S.R. & GRAHAME-SMITH, D.G. (1989). 5-HT_{1A} agonists reduce 5-hydroxytryptamine release in the rat hippocampus in vivo as determined by brain microdialysis. *Braz. J. Pharmacol.*, 96, 283-290.

- SINTON, C.M. & FALLON, S.L. (1988). Electrophysiological evidence for a functional differentiation between subtypes of the 5-HT_{1A}receptor. *Eur. J. Pharmac.*, 157, 173-181.
- SOTELO, C., CHOLLEY, B., EL MESTIKAWY, S., GOZLAN, H. & HAMON, M. (1990). Direct immunohistochemical evidence of the existence of 5-HT_{1A}autoreceptors on serotonergic neurons in the midbrain raphe nuclei. *Eur. J. Neurosci.*, 2, 1144-1154.
- SOUBRIÉ, P. (1986). Reconciling the role of central serotonin neurons in human and animal behavior, *Behav. Brain Sci.*, 9, 319-364.
- STEIN, L., WISE, C.D. & BERGER, B.D. (1973). Antianxiety action of benzodiazepines: decrease in activity of serotonin neurons in punishment system. Em: GARATTINI, S., MUSSINI, E. & RANDALL, L.O. (eds). *Benzodiazepines*. Raven Press, New York, pp. 299-326.
- STIEDL, O., MISANE, I., SPIESS, J. & ÖGREN, S.O. (2000). Involvement of the 5-HT_{1A} receptors in classical fear conditioning in C57BL/6J mice. *J. Neurosci.*, 20 (22), 8515-8527.
- SUTHERLAND, R.J. & McDONALD, R.J. (1990) Hippocampus, amygdala, and memory deficits in rats. *Behav. Brain Res.*, 37, 57-79.
- THOMAS, H., FINK, H., SOHR, R. & VOITS, M. (2000). Lesion of the median raphe nucleus: a combined behavioral and microdialysis study in rats. *Pharmacol. Biochem. Behav.*, 65, 15-21.
- TÖRK, J. (1990). Anatomy of the serotonergic system. *Ann. N. Y. Acad. Sci.*, 600, 53-67.
- TÖRK, J. & HORNUNG, J.P. (1990). Raphe nuclei and serotonin containing systems. Em: Paxinos, G (ed.). *The Human Nervous System*. San Diego, C.A. Academic Press, pp. 1001-1022.
- TREIT, D. (1985). Animal models for the study of anti-anxiety agents: a review. *Neurosci. Biobehav. Rev.*, 9, 203-222.
- TWAROG, B.M. & PAGE, I.H. (1953). Serotonin content of some mammalian tissues and urine and a method for its determination. *Am. J. Physiol.*, 175, 157-161.
- VERGÉ, D., DAVAL, G., MARCINKIEWICZ, M., PATEY, A., EL MESTIKAWY, S., GOZLAN, H. & HAMON, M. (1986). Quantitative autoradiography of multiple 5-HT₁

- receptors subtypes in the brain of control or 5,7-dihidroxytryptamine-treated rats. *J. Neurosci.*, 6, 3474-3482.
- VERGÉ, D., DAVAL, G., PATEY, A., GOZLAN, H., EL MESTIKAWY, S. & HAMON, M. (1985). Presynaptic 5-HT autoreceptors on serotonergic cell bodies and/or dendrites but not terminal are of the 5-HT_{1A} subtype. *Eu. J. Pharmacol.*, 113, 463-464.
- VIANNA, D.M.L., LANDEIRA-FERNANDEZ, J. & BRANDÃO, M.L. (2001). Dorsolateral and ventral regions of the periaqueductal gray matter are involved in distinct types of fear. *Neurosci. Biobehav. Rev.*, 25, 711-719.
- VOGEL, J.R., BEER, B. & CLODY, D.E. (1971). A simple and reliable conflit of procedure for testing anti-anxiety agents. *Psychopharmacology*, 21, 1-7.
- WILENSKY, A.E., SCHAFE, G.E. & LeDOUX, J.E. (2000). The amygdala modulates memory consolidation of fear-motivated inhibitory avoidance learning but not classical fear conditioning. *J. Neurosci.*, 20 (18), 7059-7066.
- WIRTSHAFTER, D. & ASIN, K.E. (1982). Evidence that electrolytic median raphe lesions increase locomotion but not exploration. *Physiol. Behav.*, 28, 749-754.
- WIRTSHAFTER, D. & McWILLIAMS, C. (1987). Supression of locomotor activity produced by acute injections of the kainic acid into the median raphe nucleus. *Brain Res.*, 408, 349-352.
- WIRTSHAFTER, D., ASIN, K.E. & KENT, E.W. (1980). An analysis of open field hiperactivity following electrolitic median raphe lesion in the rat. *Soc. Neurosci. Abstr.* 6, 422.
- WIRTSHAFTER, D., TRIFUNOVIC, R. & KREBS, J.C. (1989). Behavioral and biochemical evidence for a functional role of excitatory amino acids in the median raphe nucleus. *Brain Res.*, 482, 225-234.
- WOOLEY, D.W. & SHAW, E.A. (1954a). A biochemical and pharmacological suggestion about certain mental disorders. *Science*, 119, 587-588.
- WOOLEY, D.W. & SHAW, E.A. (1954b). A biochemical and pharmacological suggestion about certain mental disorders. *Proc. Nat. Acad. Sci.*, 40, 228-31.