**UNIVERSIDADE DE SÃO PAULO
INSTITUTO DE QUÍMICA DE SÃO PAULO
QUÍMICA ORGÂNICA E BIOLÓGICA**

MARCELO TAVARES DE OLIVEIRA

**QUANTUM CHEMICAL EXPLORATIONS INTO THE BIOSYNTHESIS OF
PENTACYCLIC TRITERPENE FRIEDELIN**

TESE DE DOUTORADO

**SÃO CARLOS
2019**

MARCELO TAVARES DE OLIVEIRA

**QUANTUM CHEMICAL EXPLORATIONS INTO THE BIOSYNTHESIS OF
PENTACYCLIC TRITERPENE FRIEDELIN**

Tese apresentada ao Instituto de Química
de São Carlos da Universidade de São
Paulo como parte dos requisitos para a
obtenção do título de doutor em ciências

Área de concentração: Química Orgânica e Biológica

Orientador: Prof. Dr. Albérico B. F. da Silva

**SÃO CARLOS
2019**

To Sarah.

Acknowledgments

Firstly, I wish to express my sincere thanks to Prof. Albérico for facilitating “in his very own way” the work described here in the course of the past year and a half.

I thank Prof. Ataulpa Braga (IQ/USP) for his most helpful discussions on methods, especially the tweaks of gaussian.

Prof. Glaucius Oliva (IFSC/USP) is acknowledged for his precious contribution towards the zeitgeist workstation where most computations were carried out.

Mr. Gilmar Bertollo Jr. (IFSC/USP), a very knowledgeable tech guy, for his great assistance with hardware – very appreciated.

I express my gratitude to all colleagues (and a few new friends) as well as all members of the community at large in the chemistry institute (IQSC/USP) and the physics institute (IFSC/USP). To all those I came across over the past few years of postgraduate studies who had a share of contribution to make things easier somehow.

The Coordination for the Improvement of Higher Education Personnel, CAPES (Coordenação de Aperfeiçoamento de Pessoal de Nível Superior) is acknowledged for an institutional studentship.

Last but not least, to my family and friends, who have an immense understanding for my absence during the times of hard work. To you, my greatest thank you and I have a debt.

ABSTRACT

TAVARES DE OLIVEIRA, Marcelo. **Quantum chemical explorations into the biosynthesis of pentacyclic triterpene friedelin.** Thesis (Doctorate in Sciences with emphasis on Biological and Organic Chemistry) – 2019. 105 pages. São Carlos Chemistry Institute, University of São Paulo. São Carlos, 2019.

Terpenes comprise the largest class of natural products. These metabolites have several applications in agriculture, as flavours and fragrances, and medicines. In Nature, terpenes play key roles in chemical communication. The vast diversity in the family is achieved by enzymes working on a limited number of building blocks. Terpene synthases are major enzymes in the process responsible for cyclisations and various rearrangements, which produce numerous scaffolds and configurations from a template structure. After activation, carbocation structures are present throughout the biosynthesis process. We explored by means of quantum chemistry, a series of cyclisations/rearrangements in the pathway towards the pentacyclic triterpene friedelin starting from dammarenyl cation, a key precursor in the biosynthesis of triterpenes in plants. The sequence of transformations represents the longest route in pentacyclic triterpenes. We aimed at investigating the intrinsic reactivity of carbocations in the cascade of reactions. By locating all structures relevant to the mechanisms and determining associated energy barriers, we expand knowledge on the topic by providing an improved and more detailed mechanism, which should open up possibilities for further computational and experimental work. Of particular interest to the sequence, we highlight three structures including two secondary carbocations. In the cyclisation of baccharenyl, a non-classical carbocation very close to the transition state was located. Due to considerable strain, conversion of germanicyl I to germanicyl II requires stabilisation by non-covalent interactions to occur, which should implicate in the participation of the enzyme in pre-organising the substrate to a favourable conformation. Lastly, tertiary carbocation glutinyl proved to be the only structure with the hydroxyl group in ring A in axial position accounting partly for the highest barrier in the route. A pattern in dipole moments increasing from the centre to the edges was observed and implications discussed. Regarding methods, by comparison to BB1K, B3LYP showed very similar mean absolute deviation to mPW1PW91.

Keywords: triterpene, mechanism, DFT, carbocation, friedelin.

RESUMO

TAVARES DE OLIVEIRA, Marcelo. **Estudo Químico Quântico Computacional da Biossíntese do Triterpeno Pentacíclico Friedelina**. Tese (Doutorado em Ciências com ênfase em Química Orgânica e Biológica) – 2019. 105 páginas. Instituto de Química de São Carlos da Universidade de São Paulo. São Carlos, 2019.

Terpenos representam a maior classe de produtos naturais com variadas aplicações na agricultura, medicina, como fragrâncias e aromatizantes. Na natureza, terpenos possuem importante função na comunicação química. Apesar da grande variedade de compostos, as enzimas responsáveis realizam as transformações em apenas alguns poucos substratos. Terpeno sintetases são enzimas importantes no processo sendo responsáveis por ciclizações e diversos rearranjos conduzindo a variados núcleos e configurações a partir de uma estrutura base. Após o terpeno sofrer ativação, carbocátions estão presentes por todo mecanismo. Para esta tese, exploramos uma série de ciclizações e rearranjos envolvidos na biossíntese do triterpeno friedelina utilizando métodos de química quântica. Partiu-se do cátion damarenila, precursor-chave na biossíntese de triterpenos em plantas. Objetivou-se investigar a reatividade intrínseca dos diversos carbocátions na cascata de reações. Todas estruturas relevantes foram localizadas permitindo se determinar as barreiras. Os resultados mostram um mecanismo mais detalhado do que aquele apresentado anteriormente, que permite avançar em novos estudos computacionais e/ou experimentais. Entre as estruturas de carbocátions do mecanismo destacamos alguns, dentre os quais dois são secundários. Na ciclização do cátion bacharenila, um carbocátion não-clássico muito próximo ao estado de transição foi localizado. Para a conversão do cátion germanicila I em germanicila II, inferiu-se a existência de interações intermoleculares envolvidas na estabilização do carbocátion favorecendo uma conformação menos estável necessária à transformação, que sugere a participação da enzima para a conversão. O carbocátion glutinila foi o único a apresentar a hidroxila do anel A na posição axial sendo responsável parcialmente pela maior barreira de toda a sequência. Ainda observamos ainda que o momento de dipolo dos cátions aumenta seguindo um padrão do centro para os extremos da estrutura; implicações são discutidas. Os métodos utilizados no estudo foram comparados e não se observou nenhuma diferença significativa para o desvio absoluto médio entre os funcionais mPW1PW91 e B3LYP.

Palavras-chave: triterpeno, mecanismo, DFT, carbocátion, friedelina

LIST OF FIGURES

Figure 1. Examples Important bioactive natural products	13
Figure 2. Some monoterpenes (C ₁₀) present in turpentine	15
Figure 3. (A) Early synthesised terpenes, (B) diterpenes, phomactins R and T	15
Figure 4. Terpene pheromones from insects	16
Figure 5. Terpenes of various biological functions	17
Figure 6. Triterpenes of pharmacological interest	22
Figure 7. Terpene biosynthesis summarised in 3 major events	24
Figure 8. Ruzicka's isoprene rule applied to cadalane and cedrane scaffolds	24
Figure 9. Pathways leading to isoprenic precursors in terpene biosynthesis	25
Figure 10. Biosynthesis of monoterpene key intermediate GPP, and isomerisation LPP and NPP	26
Figure 11. General scheme for the biosynthesis of acyclic terpene precursors	27
Figure 12. Cyclic monoterpene products from GPP having α -terpinyl cation as intermediate	29
Figure 13. Diversity of domains in terpene synthases	31
Figure 14. Catalytic mechanism of taxadiene synthase	32
Figure 15. Examples of pentacyclic triterpenes of baccharene type	33
Figure 16. Biosynthetic route to sterols and triterpenes from squalene	35
Figure 17. Lanosterol biosynthesis from 2,3-oxidosqualene and residues in interaction	36
Figure 18. Biosynthesis of sesquiterpenes ledol and viridiflorol showing key steps	40
Figure 19. Proposed mechanism for the synthesis of sesquiterpene avermitilol	41
Figure 20. Mechanism for the formation of trefalone A	44
Figure 21. Originally proposed mechanism for pentalenene formation	45
Figure 22. Theoretically proposed mechanisms for the formation of pentalenene	46
Figure 23. Theoretically computed pathways for the biosynthesis of trichodiene	47
Figure 24. Computed transition states for taxadiene cations	50
Figure 25. Highlights in the development of computational organic chemistry	52
Figure 26. Perdew's Jacob's ladder of exchange-correlation functionals	62
Figure 27. (a) Two-dimensional representation of a reaction path	64
Figure 27. (b) Three-dimensional representation of a potential energy surface	64
Figure 28. Putative biosynthetic mechanism for the formation of pentacyclic triterpene friedelin...	68
Figure 29. A benzene in non-covalent interactions with a triterpene carbocation	69

Figure 30. Examples of incorrect structures for friedelin	71
Figure 31. Ring naming in the triterpene pentacyclic core and carbon atom numbering in oleanane..	72
Figure 32. Dammarenyl cation best clusters	73
Figure 33. Transformations from dammarenyl (1) to germanicyl I (4)	74
Figure 34. Conversion of dammarenyl cation 1 into 4 showing cyclisations	75
Figure 35. Nonclassical and classical baccharenyl cations	76
Figure 36. Energy profile for the reaction pathway from dammarenyl (1) to lupanyl cation (4)	77
Figure 37. Computed structure for germanicyl I cation (4) showing elongated distances	78
Figure 38. Sequence of transformations from germanicyl I (4) to oleanyl (6) cation	78
Figure 39. Energy barrier for conformational change in ring E.....	79
Figure 40. Attempts to locate TS_{4-to-5} using a non-covalent interaction	80
Figure 41. Transtion state TS_{4-to-5} in non-covalent interactions	81
Figure 42. Energy profile for the reaction pathway from lupanyl (3) to germanicyl II cation (5)	82
Figure 43. Details of the interaction of TS_{4-to-5} with ammonia and phenol	83
Figure 44. Energy profile for the conversion of germanicyl II cation (6) to oleanyl cation (7)	84
Figure 45. Conversion of oleanyl cation (6) to friedelin cation (13)	85
Figure 46. Conversion of oleanyl cation (6) into campanulyl (10)	86
Figure 47. Conversion of campanulyl (10) into friedelin cation (13)	87
Figure 48. Energy profile for the cyclisations/rearrangements from dammarenyl to friedelin	88
Figure 49. Dipole moments for the series pentacyclic triterpene cations	90

LIST OF TABLES

Table 1. Classification of Terpenes	18
Table 2. Terpenes present in various products and/or of commercial interest	19
Table 3. Synthase classification and domains	30
Table 4. Relative stabilities of carbocations	42
Table 5. Energies barriers for transformations in bisabolyl and cuprenyl carbocations	50
Table 6. Activation energies and deviations for the density functions applied in the study	93

SUMMARY

1	Introduction	11
	Part I – Terpenes as a Major Class of Natural Products and their Biosynthesis	12
1.1.	Terpenes	14
1.1.1.	Biological Relevance	16
1.1.2.	Structural Classification	17
1.1.3.	Occurrence, Applications and Uses	18
1.2.	Triterpenes	21
1.3.	Biosynthesis	23
1.3.1.	Cyclizations and Rearrangements in Terpene Biosynthesis	28
1.3.2.	Biosynthesis of Triterpenes	33
	Part II – Quantum Chemistry Explorations into the Biosynthesis of Terpenes	37
1.4.	Quantum Chemistry in the Biosynthesis of Terpenes	39
2	Methods	51
2.1.	Quantum Mechanis. Schrödinger Equation	53
2.2.	Density Functional Theory	57
2.3.	Exchange Correlation Functionals. Applications to Terpene Biosynthesis	61
2.4.	Potential Energy Surface	63
3	Aims and Scope	66
4	Results and Discussion	70
4.1.	Formation of Rings D and E	74
4.2.	Formation of Oleanyl Cation	78
4.3.	Formation of Friedelin Cation	85
4.4.	Performance of Density Functionals	91
5	Conclusion	93
6	References	96

1 INTRODUCTION

PART I – Terpenes as a major class of Natural Products and their Biosyntheses

Our continuous strive to understand Nature has led to numerous scientific discoveries and technological advances that we now enjoy and take for granted. In various scientific fields including some in the realm of chemistry, efforts are largely represented by our endeavours to master the structure of the molecules of life – carbohydrates, lipids, nucleic acids, proteins, and the classes of secondary metabolites, and their intricate roles in a myriad of processes in biological systems.

From the dawn of our civilisation, mankind has explored natural sources particularly plants for multiple purposes. By experimentation in a trial and error approach, remedies to all sorts of ailments were discovered. Poisonous and hallucinogenic species were also identified along the way (Fig. 1).¹ In the past couple centuries, our research advances have resulted in the identification of over 200,000 natural products from plants and many more considering micro-organisms.² Screening these substances for prospective applications represents a challenge and a continuous effort. Developments of tools for agile separation and identification of compounds in the past decades have largely contributed to automation in the field, and bioscreening is an area which has also experienced considerable progress.

The complexity of secondary metabolites has long captivated chemists. Attracted by unprecedented chemical scaffolds displaying unique biological activities, countless research groups have embarked on projects to produce the chemical syntheses of such compounds. In so doing, new reactions and strategies had to be developed.³ Others went on to explore the biochemistry of secondary metabolites aiming at understanding their biological functions while having the perspective of drug discovery. In that sense, secondary metabolites continue to represent an important source of lead compounds and inspiration to the design of new drugs.^{4,5} They provide privileged structures displaying exquisite pharmacophoric groups developed by the enzymatic machinery through millions of years of evolutionary pressure to serve that organism to a certain purpose from chemical defence to inter- or intraspecies communication. In some instances, there are not many alternatives to natural product drugs as it is the case of morphine and cardiac glycosides. Moreover, the presence of natural products in some drug classes including antiparasitics and anticancer drugs is particularly relevant.

Figure 1. Examples of important bioactive natural products.

On the downside, secondary metabolites are often produced in minute quantities requiring considerable effort to access enough amounts for further chemical and biological studies. Chemical synthesis or hemi-synthesis are options to be considered depending on the economic viability for commercialisation. To that end, biological approaches are on the rise. Identification of the gene(s) that encode production of a given secondary metabolite offers the possibility of insertion into a host organism e.g., *E. coli* for biosynthetic production. Bioengineering enzymes to boost yields represents an attractive solution, which has gained substantial attention in recent years; however, successful examples are currently limited.⁶

Beyond drugs, natural products occupy an important place in our society in numerous applications as agrochemicals, cosmetics, dyes, fibres, oils, perfumes among others.⁷ Therefore, harnessing the chemical space of natural products remains relevant as ever in view of the vastness of new compounds yet to be identified particularly from ecosystems like the

oceans and other less explored corners of the globe representing extreme environments, which we have started to explore only more recently.

1.1 Terpenes

Terpenes comprise the largest class of natural products with more than 50,000 compounds already identified considering a strict definition, and over 80,000 compounds when structurally related carotenoids and steroids are taken into account, which represent a great deal of all natural products known to date.² Therefore, implying it is a very diverse family. Terpenes and their oxidised derivatives terpenoids represent the most ancient class of small molecule natural products on the planet. The triterpene-like skeleton of monoaromatic steroids have been found in fossils of the earliest animals of the Ediacara biota (571 million to 541 million of years ago).⁸ Terpenes have also been found in fossils of plants and micro-organisms.^{9,10}

The term terpene was originally given to a mixture of isomeric hydrocarbons of molecular formula $C_{10}H_{16}$ isolated from turpentine (latin *balsamun terebinthinae*), which is a volatile clear liquid of pungent odour and bitter taste; extracted from the pine tree. The terpene term was coined by the French chemist Jean-Baptiste Dumas in 1866, although the analyses of turpentine oil had been made decades before in 1818. Turpentine contains a range of monoterpenes and monoterpenoids (C_{10}) with distribution depending on the species and several other factors such as geographical location, age of the plant, and season of the year. The structures, nonetheless, provide an initial hint on the chemical diversity within the terpene family (Fig. 2).

The history of terpenes can be traced to early developments in organic chemistry from XIX century. By the early 1800s, techniques were improved to allow extraction of the fragile active ingredients from their natural sources by using organic solvents at ambient temperatures providing direct access to essential oils rich in terpenes, which are key ingredients in perfumery.

Figure 2. Some monoterpenes (C_{10}) present in turpentine.

At the turn of XIX century, structures of several terpenes were determined including camphor (Bredt, 1893), pinene (Wagner, 1894), and citral (Tiemann, 1895). Further advances propelled the production of naturally occurring fragrances of interest to the perfume industry e.g. musk, vanilla and violet by end of the century.⁷ Later on, α -terpineol by Perkin (1904)¹³ and camphor (Perkin, 1904)¹⁴ represented early examples of terpenes targeted in organic synthesis (Fig. 3A). Many more followed and the trend never ceased. For their structural complexity and biological relevance, terpenes continue to attract attention as it is the case of the bioinspired synthesis of diterpenoids phomactins reported just a few months ago.¹⁵ Phomactins has shown notable activity as platelet activating factor receptor (PFAR) antagonists rendering the prospect of application in cancer therapy (Fig. 3B).

Figure 3. (A) Early synthesised terpenes, α -terpineol and camphor. Both accomplished by Perkin in 1904; (B) diterpenes, phomactins R and T, as examples of recently achieved synthesis.

Discovery of chromatographic and spectroscopic techniques after the wars led to an explosion in the chemistry of natural products providing a sustained growth in the number of new compounds each decade and enabling access to complex structures. Nowadays, hyphenated techniques e.g., gas chromatography-mass spectroscopy (GC-MS) or liquid chromatography-nuclear magnetic resonance (LC-NMR) associated with access to large databases of already identified compounds, considerably speed up the process of identification and isolation of novel compounds.

1.1.1 Biological Relevance

While the class of flavonoids are best remembered as pigments in plants and for its photoprotective effects and alkaloids for their neurostimulant and poisonous properties, terpenes bring to mind their aromas and tastes. It is not surprising that terpenes of low molecular weight as hydrocarbons are volatile. The biological and ecological roles of these chemicals have been studied only to a certain extent to a limited number of species.

Many plants use terpenes to attract certain insects for pollination while others produce these metabolites as antifeedants to prevent animals from eating them. Interestingly, many insects in turn metabolize terpenes ingested from their food to growth hormones and pheromones of various kinds – alarm pheromones to warn, trace pheromones to mark and locate food resources, aggregation pheromones to determine an assembly place and sexual pheromones to lure potential sexual partners to copulate (Fig. 4). Pheromones exert their functions on tiny quantities and are environment benign, thus offering an attractive alternative to harmful insecticide.¹⁶

Figure 4. Terpene pheromones from insects.

As structures are diverse so do their functions. Terpenes are also growth regulators (phytohormones) in plants showing an important role in signalling. Among other physiological roles, brassinolide control cell elongation and division in plants. Cholesterol is a steroid essential component of biomebranes and responsible for its fluidity. Phytol, the side chain of the respiratory pigment chlorophyll, represents the most abundant terpenoid (Fig. 5).

Figure 5. Terpenes of various biological functions.

1.1.2 Structure and Classification

From examples shown thus far, we have seen acyclic and cyclic structures but also a number of functionalised cases (alcohols, ethers, ketones, esters, etc) of terpenoids extending the concept of terpenes, which are hydrocarbons by definition. Nonetheless, we should use the term “terpene” throughout in a broader sense to include terpenoids.

The core of terpenes is built by five-carbon (C_5) building blocks according to the biosynthetic pathways that will be discussed soon in detail. Therefore, each distinct class of terpenes originates from the successive addition of a C_5 -unit or multiples extending the chain, that is, monoterpenes (C_{10}), sesquiterpenes (C_{15}), diterpenoids (C_{20}), etc. The simplest of all consisting of a single C_5 isoprene unit (Table 1).

Table 1. Classification of Terpenes

Terpenes	Isoprene units	Carbon atoms
Hemiterpenes	1	5
Monoterpenes	2	10
Sesquiterpenes	3	15
Diterpenes	4	20
Sesterpenes	5	25
Triterpenes	6	30
Carotenoids	8	40
Rubber	>100	>500

Although the majority of terpenes follow this pattern, degradation to isoprene moieties is possible as well as further addition of carbons. That is the case of taiwanianquinoid C₁₉ diterpenes¹⁷ and methylated C₁₁ monoterpenes.¹⁸ Apart from variations in the carbon backbone as promptly illustrate in Fig. 2, the presence of oxygenated functions and multiple bonds increase the structural diversity. Moreover, stereocentres may also be present increasing possibilities exponentially.

1.1.3 Occurrence, Applications and Uses

Terpenes are widely distributed in all kingdoms of life despite the majority of terpenes have been isolated from plant species. As much as they are structurally diverse, many are their roles in primary metabolism. A great number of terpenes have economical value to various industries including the polyterpene known as natural rubber, essentials oils in cosmetics, carotenoids pigments in food, and some have even become important drugs. We next show a range of examples to outline the structural diversity of this class of natural products: from simple achiral monoterpene α -terpineol present in various essential oils to the intricate structure of the anticancer drug taxol (Table 2). The uses demonstrate the presence of these compounds in our daily lives. Some are components of essential oils or produced by chemical synthesis to be sold to many industries.¹⁶

Table 2. Terpenes present in various products and/or of commercial interest.

Food and Drinks

Terpenes in food and drinks are responsible for taste and colour

Rosemary

carene
*sweet, diffuse;
coniferous*

Coriander

geranyl acetate
floral; rosy

linalool
floral, slightly spicy

Bay leaves

α -phellandrene

peppery mints; slightly citrusy

β -phellandrene

Black pepper

sabinene
*spicy, woody
and camphoreous*

caryophyllene
*spicy, woody, sweet
and bitter*

Lemongrass

nerol
fruity, spicy, floral

Hops

humulene
hoppy; bitter

Citrus Fruits

limonene
sweet, citric

Wines

Terpenes are partly responsible for the complex flavour and aroma (bouquet) in finished wines

rotundone
peppery, spicy
found in Shiraz, Cabernet-Sauvignon,
Zinfandel, Merlot

1,8-cineole
*fresh, camphoreous,
balsamic*
Australian Shiraz

(-)-cis-rose oxide
lychee taste
Gewürztraminer

α -terpineol
lime taste; sweet
Muscat

Gin

Terpenes in juniper berry responsible for the distinctive flavour of gin

α -pinene
woody, piney; sweet, citrus

limonene

1,4-terpineol
pine; woody, spicy

β -myrcene
woody, herbaceous

bornyl acetate
woody, camphoreous

Table 2. Terpenes present in various products and/or of commercial interest (cont).

Food and Drinks (cont)

Carotenoids as food colouring

β-carotene

carrots, kale, sweet potato
registered as E160a

lycopene

tomatoes, carrots, pawpaw
E160d

Cosmetics and Perfumery

α(-)-bisabol
present in chamomile

has a sweet floral aroma;
skin healing properties,
anti-irritant, anti-inflammatory

cis(-)-carveol
present in spearmint

spearmint and caraway aroma;
fragrance in cosmetics

patchoulol
from patchouli

woody, earthy, balsamic
perfumery

(-)-α-cedrene
from cedar

woody, cedar, fresh, sweet
perfumery

cadinene
from cade

herbal, woody, smoky
cosmetics for skin treatment

α-ocimene
from basil

terpy, woody, floral nuances
perfumery

(+)-iongifolene
pine species

woody, cedar, fresh, sweet
perfumery

(+)-geranyl linalool
various flowers

floral, rosy
perfumery as fixative for rose
fragrances

Table 2. Terpenes present in various products and/or of commercial interest (cont.).

Drugs

1.2 Triterpenes

Triterpenes (C_{30}) represent the most numerous group of terpenes with over 20,000 compounds derived from the common acyclic precursor squalene, a C_{30} linear polyene originated from isoprene. Triterpenes form a very structurally diverse class which display nearly 200 distinct carbon skeletons. Most members display 6-6-6-5 tetracycles, 6-6-6-6-5 or 6-6-6-6-6 pentacycles. Higher hexacyclic, smaller monocyclic, bicyclic and tricyclic as well as acyclic systems also exist in small numbers.¹⁹ Pentacyclic and tetracyclic triterpenes are more prevalent in higher plants while lower systems such as mono-, bi-, and tricyclic occur more often in ferns and cryptogams (non-flowering plants). In these organisms, triterpenes constitute an important portion of the lipids. Importantly, they are also precursors to steroids in both plants and animals. Furthermore, triterpenes have primary and secondary roles in plants. While the vast majority of triterpenes displays roles as secondary metabolites such as defence, ecological or as part of the adaptation system, phytosterols have an essential role in cell membrane formation.

A number of triterpenes show important pharmacological properties to human health, hence of high interest to the pharmaceutical industry. Triterpenes with well-acknowledged biological activity include saponins, steroids, and sterols. The steroidal saponin diosgenin is an important example as source of human hormones. The phytosteroid sapogenin is extracted from tubers of wild yams; genus *Dioscorea*. The obtained aglycone (sugar-free) diosgenin (Fig. 6) after hydrolysis is a precursor to the commercial synthesis of several hormones including cortisone, pregnenolone, progesterone among other steroids and compounds present in contraceptive pills. The class of cardiac glycosides represents another relevant example. Digoxin is produced by the foxglove plant, *Digitalis lanata*, and used in the treatment of heart conditions including atrial fibrillation and heart failure.

The group of ginsenosides, tetracyclic triterpenoid saponins, are the main pharmacologically active substances present in the roots of *Panax sp.*, which have been used in the far orient for over 2,000 years for various ailments and to promote longevity. Ginsenosides have demonstrated neuroprotective properties which could be useful in treating neurodegenerative diseases such as Alzheimer's and Parkinson's. Some ginsenosides have shown inhibitory effect of cancer cell growth.²⁰ Numerous reports on the chemopreventive effects of triterpenes, mostly pentacyclic triterpenes, have shown biological effects related to potential anticancer activity. Several other biological activities have been described for triterpenes: anti-inflammatory, antidiabetic, antioxidant, anti-infectious, anti-viral, anti-bacterial, immunomodulatory, among others.²¹ Despite the potential and attractive action mechanisms, further studies towards drug development are hampered by availability as extraction from plants usually afford limited amounts.

Figure 6. Triterpenes of pharmacological interest.

1.3 Biosynthesis

Thus far, we have shown structures and applications of terpenes while introducing concepts related to the subject matter, which has laid the foundation for what comes next. Probably some reading the text have wondered how Nature constructs all these elaborate terpene structures from just plain C₅ building blocks. Put differently that is to state how the enzymatic machinery achieve such complexity and diversity from sheer simplicity.

Terpenes are found in nearly all organism in the planet demonstrating the importance of the class to many essential processes of life. To understand the biosynthesis of terpenes represents a challenge and not only of academic interest but also practical to industries aiming at developing efficient biotechnological processes to produce molecules of high-added value.

The biosynthesis of terpenes follows an established modular approach applying a series of enzymes. From simple C₅ building blocks, the whole range of distinct classes of terpenes are built in successive additions of five-carbon units, isopentenyl diphosphate (IPP) and dimethyl diphosphate (DMAPP). Prenyltransferases are responsible for this assembly line producing universal diphosphate substrates within five-carbon increments, which constitutes the basis of the different terpene classes (e.g., monoterpenes, sesquiterpenes, etc). In the second major module, each basic activated substrate is converted to different terpene backbones by the action of terpene synthases or cyclases. In fact, cyclisation represents the module of our greatest interest and, therefore, we will elaborate considerably more on it at this stage.

At the third and final module, further modifications introducing functionalities take place decorating the terpene skeleton. Various enzymes participate in this process such as *O*-acetyltransferases, *O*-methyltransferases, dehydrogenases, cytochrome P450s, among others.²² Differently from the other two initial modules, these transformations adding decorative functionalities are not exclusive to terpenes. Prenylations to assemble the C₅ building blocks and cyclisations followed by eventual rearrangements best represent the terpene biosynthetic code.

To illustrate, we outline the biosynthesis of the diterpene, cancer drug taxol (Fig. 7). Firstly, C₅ isoprenoid units are assembled in a determined manner, which we shall discuss in further detail shortly, to form the C₂₀ diterpene general substrate GGPP (geranyl-geranyl diphosphate). Cyclization and rearrangements follow to generate the taxadiene tricyclic core. Various decorations to the main backbone would eventually lead to the taxol structure as we know it.

Figure 7. Terpene biosynthesis summarised in 3 major events: prenylation, cyclisation and decoration.

Research into the biosynthesis of terpenes started in the early XIX century. A turning point in the efforts towards unravelling the mysteries in the biosynthesis of terpenes arrived at the early 1950s with the seminal contributions from Leopold Ruzicka. By 1953, he discovered while studying cadalenes and cedranes that the five-carbon unit isoprene, which follows the rule of same name previously set, was a common precursor in the construction of terpenes and were assembled by connections in a head-to-tail fashion (Fig. 8).²³ This became later known as the Ruzicka's rule. Back then, it was believed that connections following head-to-head or tail-to-tail were irregular. Today, we understand that these result from rearrangements in the course of terpene cyclisation.

Figure 8. Ruzicka's isoprene rule applied to cadalane and cedrane scaffolds.

Further investigation into the precursors in the biosynthesis of terpenes has shown that two activated molecules, isopentenyl diphosphate (IPP) and dimethylallyl diphosphate (DMAPP), displaying the same isoprene-like backbone participate in the construction of terpenes. DMAPP is produced in the well-known mevalonate pathway from two molecules of acetyl-CoA, which is a process known to take place in animals, cytosol of plants, fungi, archaea and a few bacteria. The deoxyxylulose phosphate pathway is relatively recent (1990s – early 2000s), and starts from pyruvate and the simplest activated sugar, glyceraldehyde 3-phosphate. Ultimately, the isoprene equivalents DMAPP and IPP are produced in a ratio of approximately 1:5 (Fig. 9).²⁴ This pathway occurs in most bacteria, plastid of plants and green algae. Despite following distinct pathways in multiple steps through different intermediates and set of enzymes, both pathways deliver the same C₅-activated isoprenes, DMAPP and IPP.

Figure 9. Schematic representation of pathways leading to isoprenic precursors in terpene biosynthesis.

From these activated isoprene units, all terpenes are formed from various combinations and further modifications to access diversity up to higher terpenes of various biological properties. While IPP is the nucleophilic unit, DMAPP is highly electrophilic. Removal of diphosphate in DMAPP leads to a stabilised allylic cation (Fig. 10a). Enzymes collectively known as prenyltransferases carry out the biosynthesis of linear polyprenyl diphosphates. The backbone of monoterpenes results from the catalysed addition of IPP to DMAPP by geranyl diphosphate synthase (GPP synthase). By stereoselective abstraction of H_R, this addition most

often produces (*E*)-configured geranyldiphosphate (GPP). Nonetheless, isomerisation can readily occur to (*Z*)-configured neryl diphosphate (NPP) via carbocation formation and subsequent diphosphate addition to the tertiary carbon producing linalyl diphosphate (LPP), which is conformationally flexible (Fig. 10b). Alternatively, LPP can be directly accessed by enzymes favouring stereoselective deprotonation of H_S.

Figure 10. (a) Biosynthesis of monoterpene (C₁₀) key intermediate GPP. (b) Isomerisations between GPP, LPP and NPP.

The same chemical rationale applies to the biosynthesis of larger acyclic terpenes. From geranyl diphosphate (GPP, C₁₀) and a further isopentenyl diphosphate (IPP, C₅) unit, farnesyl diphosphate (FPP, C₁₅) is produced; and in the continuation of the elongation process geranyl-geranyl diphosphate (GGPP, C₂₀). Squalene (C₃₀) as the triterpene precursor, however, is formed by dimerization of two FPP (C₁₅) units (Fig. 11). In these cases, the chain length is determined by the size of the hydrophobic pocket in the enzyme, which contains a binding site for IPP and another for the diphosphate substrate.²⁵

Figure 11. General scheme for the biosynthesis of acyclic terpene precursors.

The acyclic activated intermediates function as branch points in terpene metabolism to further modifications particularly cyclisations under the action of terpene synthases (or cyclases) to produce the great diversity of scaffolds in cycles of various sizes and connectivity in monocyclic or polycyclic systems. Interestingly, terpene biosynthesis in plant cells is a compartmentalised process occurring in different locations in the cell. Monoterpenes and diterpenes (as well as tetraterpenes) arise in the plastids making use of the deoxyxylulose phosphate pathway. Sesquiterpenes and triterpenes, however, are produced in the cytosol via the mevalonate pathway. In both compartments, IPP isomerase are present.^{26,27}

At this stage and as we move along, it becomes increasingly patent that carbocations represent a hallmark in terpene catalysis through various reactions (e.g., cyclisations, ring expansions, hydride and alkyl shifts, nucleophile additions, Wagner-Meerwein rearrangements and so forth). These reactive intermediates define reactivity along the cascade. As alkylating agents, they could alkylate a residue in the active site resulting in inactivation of the enzyme. To deal with such reactivity, the enzyme contains mostly a nonpolar active site to manage these intermediate through non-covalent interactions, namely cation- π . It is also likely that carbocations do not occur as discrete species along the reaction pathway when ion pairing (e.g. diphosphate) is considered, for instance. However, this does not preclude eliminations forming double bonds along the backbone or the occasional water quenching the reactive species.^{27,28} It is also observed a preference for transformations resorting to more stable tertiary carbocation.

1.3.1 Cyclizations and Rearrangements in Terpene Biosynthesis

Prenylations provide a few general substrates for further transformations. Leaving late-stage functionalisations (decorations) aside, which contributes to the diversity of terpene structures, cyclisations and associated rearrangements are mostly responsible for the diversity of scaffolds and isomers in terpenes. Terpene cyclisations are regarded as the most complex chemical transformation in Nature.²⁹ Terpenoid cyclases or syntases bring about a range of transformations including cyclisations and rearrangements in which great extensions of the structure undergo modifications in bonding, hybridisation and stereochemistry over the course of the cascade of reactions. Cyclases are enzymes which evolved to deal with carbocations and differentiated to produce the impressive diversity of terpenes. Understanding the pertaining mechanisms could unlock the potential of protein engineering to reprogram a given enzyme to the cyclisation of a desired product of commercial interest.

Similarly to prenyltransferases that catalyse intermolecular electrophilic couplings, cyclases can be viewed as intramolecular analogues. However, the number of cyclases are far greater than the former for the many products it generates. For exerting their function on carbocations as substrates, these families of enzymes share some structural features. They exhibit considerable conservation in the primary sequence and have similar properties such

as the fact that they are acidic enzymes, functionally soluble, native structural size in the range of 35–80 kDa and have the requirement of a divalent metal cation in some cases.³⁰

Monoterpene synthases catalyse formation of acyclic, monocyclic and bicyclic natural products as alcohols, olefins or diphosphate esters (Fig. 12). A typical cyclisation involves ionisation of the prenyl diphosphate substrate followed by intramolecular attack of a double bond to a carbon to yield a cyclic carbocation intermediate. Following steps may result from further additions to the remaining double bonds resulting in an extra ring. Atom or group migrations *e.g.* hydride shifts, methyl migrations or Wagner-Meerwein (WM) rearrangements may follow before termination by water capture or deprotonation by a nucleophile.

Figure 12. Cyclic monoterpene products from GPP having α -terpinyl cation as intermediate (adapted from ref. 31).

Monoterpene synthases have been relatively well studied. These enzymes catalyse the formation of multiple products in given fixed ratios from the electrophilic cyclisation mechanism.³² Several reactive carbocation intermediates generated in the active site over the course of the reaction experience interactions with residues differently influencing their reactivity. Furthermore, intrinsic carbocation reactivity may lead to transformations e.g. deprotonation resulting in stabilisation.

Exceptions aside, terpene synthases or cyclases generally fall into two main categories depending on the chemical strategy adopted for carbocation formation. Class I terpene cyclases utilise a trinuclear metal (Mg^{2+}) cluster to promote the ionisation of an isoprenoid diphosphate substrate furnishing an allylic cation and inorganic pyrophosphate. Class II terpene cyclases, however, apply a distinct approach making use of a general acid e.g. aspartic acid side chain in the active site to protonate a terminal carbon-carbon double bond in the substrate producing a tertiary carbocation or via an epoxide to alcohol. The distribution of both classes varies according to the substrate with class I being predominantly responsible for the biosynthesis of smaller terpenes (Table 3).

Table 3. Synthase classification and domains.

Terpenes	Synthase Class	Domain Architecture
Hemiterpenes (C_5)	I	$\alpha\beta$
Monoterpenes (C_{10})	I	$\alpha\beta$
Sesquiterpenes (C_{15})	I	$\alpha, \alpha\alpha, \alpha\beta, \alpha\beta\gamma$
Diterpenes (C_{20})	I, II	$\alpha, \alpha\alpha, \alpha\beta\gamma$
Sesterpenes (C_{25})	I, II	$\alpha, \alpha\alpha$
Triterpene (C_{30})	II	$\beta\gamma, \alpha\beta\gamma$

The structure of terpene synthase structures typically consists of α , β and γ domains in various possible combinations. The pentalenene synthase, which is responsible for the biosynthesis of sesquiterpene pentalenene, has α and β domains. As a class I cyclase, the active site is located in the middle of a region known as “ α fold”, which is an α helical bundle.

That is also the case of tobacco *epi*-aristolochene synthase, which has an $\alpha\beta$ domain assembly, and taxadiene synthase from the Pacific yew exhibiting a $\alpha\beta\gamma$ domain (Fig. 13). In class I cyclases, a trinuclear metal cluster responsible for activating isoprenoid diphosphate substrate coordinates to an aspartate-rich motif (DDXXD) present in α domain in the active site. A class II synthase as the squalene-hopene cyclase ($\beta\gamma$ domain) initiates catalysis by protonation of the epoxide or a terminal π bond by the central aspartate in the DXDD motif. This sequence is unrelated to that of class I as the active site in class II cyclases is located at the interface of β and γ domains.²⁹

Figure 13. Diversity of domains in terpene synthases (α , β and γ domains are represented in blue, green and yellow, respectively). Metal-binding motif in α domain is shown in red and orange (adapted from ref. 33).

Bifunctional terpene cyclases possessing $\alpha\beta\gamma$ -domain exist with type I and II catalytic activity. These enzymes contain highly conserved DDXXD motifs in the α -domain and DXDD in β -domain. Nonetheless, these enzymes are exclusive to fungi and plant species.

The taxadiene synthase responsible for the production of potent anticancer drug taxol offers an interesting case for further illustration. The enzyme catalyses with structural precision the construction of the unique tricyclic hydrocarbon skeleton of taxol. Initially isolated from taxol-producing Pacific yew, *Taxus brevifolia*, the protein was the first class I cyclase to reveal a fusion of $\alpha\beta\gamma$ domains. The sequence was later cloned and expressed into *E. coli* as an 862-residue sequence, which yielded the desired taxa-4(5),11(2)-diene as major cyclised product. The taxadiene taxa-4(20),11(12)-diene and verticillene were also produced in smaller quantities. Although taxadiene synthase exhibit β and γ domains as other class II terpene cyclases, the $\beta\gamma$ interface is afunctional as the essential acid residue is absent.^{34,35}

The catalysis mechanism of GGPP (geranyl-geranyl diphosphate) cyclization has been extensively studied owing to the profile of taxol as a blockbuster drug in cancer chemotherapy. The binding of the substrate to the active site in α domain is accompanied by coordination to 3 Mg^{2+} . These ions also coordinate to residues in the active site including those of the aspartate-rich motif. Once metal atoms trigger ionisation of GGPP, C1–C14 bond formation, ring closure occurs producing a 14-membered ring with inversion of configuration at C1; and a tertiary carbocation at C15, which is attacked by a double bond in the structure leading to a C10–C15 ring closure that is followed by deprotonation to yield the transient species (1S)-verticillene (Fig. 14). An unusual intramolecular proton transfer from C11 to *re* face at C7 α follows producing another tertiary carbocation, which upon transannular ring closure at C3–C8 (C2–C7 of GGPP) forms rings B and C. Final proton elimination gives targeted taxadiene.^{34,36}

Figure 14. Catalytic mechanism of taxadiene synthase.

1.3.2 Biosynthesis of Triterpenes

We have shown previously (Fig. 11) that two C_{15} farnesyl diphosphate (FPP) units dimerise to produce squalene (C_{30}), which is the key precursor in the biosynthesis of triterpenes. Steroids and triterpenes alike are synthesised via the mevalonate pathway; and class II synthases are responsible for the astounding diversity of triterpenes comprising the largest sub-class of terpenes. These synthases are collectively known as oxidosqualene cyclases (OSC). More than 80 such enzymes have been identified only from plants and about a third of which are triterpene synthases. By and large, triterpene diversity occurs mostly in plants; just on oleanes more than 300 have been isolated.³⁷ As stereodense structures, variations in the stereochemistry of triterpenes are quite common contributing to the diversity (Fig. 15).

Figure 15. Examples of pentacyclic triterpenes of baccharane type.

The structural diversity in triterpenes correlates with the application of distinct biosynthetic strategies for the members of this class across kingdoms of life.³⁸ Following a class II terpene synthase mechanism, substrate requires activation by an acid residue in the active site, which is carried out by a well-conserved aspartic acid residue (D455 in the human enzyme) in the general DXDD motif. All-trans squalene itself is only directly activated in bacteria, which leads to the production of pentacyclic triterpenes of hopene family (Fig. 17). In animals, fungi and plants, squalene is oxidised to 2,3-oxidosqualene by the enzyme squalene epoxidase (or monooxygenase) prior to activation.³⁹

After 2,3-oxidosqualene activation, sterols and triterpenes have their origins from separate pathways following distinct key intermediates: protosteryl cation in the case of former and dammarenyl cation for the latter. The difference between their structures lies on the stereochemistry of just a few centres as a result of different conformations imposed by their enzymes prior to cyclisation. Triterpenes cyclize using a chair-chair-chair (CCC) conformation while sterols follow a chair-boat-chair (CBC) conformation in regard to A–B–C rings. Plants have enzymes capable of folding oxidosqualene substrate in both conformations: CCC-folding enzymes leading to the production of triterpenes and CBC in the case of cucurbitacins and β -sitosterols. Lanosterol synthases, which are responsible for production of ergosterol in fungi and cholesterol in animals, fold substrates into CBC conformation (Fig. 16).

As an example, the biosynthesis of steroid lanosterol, which is an intermediate en-route to cholesterol from 2,3-oxidosqualene, is outlined in further molecular detail (Fig. 17). The human oxidosqualene cyclase is only partly embedded in the membrane. Through a hydrophobic channel leading to the region in the membrane where the protein is anchored, substrate access the interface between $\beta\gamma$ domains where the active site is located. Once in the cavity of the synthase active site, the mechanism starts in a pre-organization step when the substrate is recognised leading to a particular folded conformation of 2,3-oxidosqualene, which is chair-boat-chair in this case. It has been shown that this initial step is critical in predisposing the substrate to follow a particular pathway. Next, the aspartic acid residue D455 protonates the epoxide ring in 2,3-oxidosqualene triggering the ring-forming cascade of cyclisations and a series of carbocation rearrangements. Point mutation of this amino acid implicates in complete loss of function. Additionally, two cysteine residues nearby (Cys456 and Cys533) are believed to enhance Asp455 acidity via hydrogen bonding.

Figure 16. Biosynthetic route to sterols and triterpenes from squalene showing key intermediates, protosteryl and dammarenyl cations, respectively (adapted from ref. 39)

Acid-promoted epoxide ring opening and A-ring formation should happen in a concerted fashion as the following cyclisations (Fig. 17a). These rings A-D interact with several aromatic residues in the active site to stabilize high-energy carbocation intermediates. Once D ring is formed producing the key intermediate protosteryl cation, a series of rearrangements including hydride and methyl transfers follow. Ultimate proton elimination at C9 by His232 or

Tyr503 produce lanosterol (Fig. 17b).²⁹ Side reaction such as termination by deprotonation elsewhere or a molecule of water quenching the carbocation may occur along the way resulting in various sterols.

Figure 17. (a) Lanosterol biosynthesis from 2,3-oxidosqualene showing the sequence of transformations. (b) Residues in the active site displaying lanosterol–oxidosqualene cyclase complex. The side chain of residues Phe696 and His232 should stabilise carbocation at C20 of protosteryl through cation– π interactions (ref. 40).

PART II – Quantum Chemistry Explorations into the Biosynthesis of Terpenes

Among the breakthroughs of science in the 20th century, we highlight developments in quantum chemistry to the scope of this text. As a field directly derived from quantum mechanics, probably one of most important discoveries from last century, quantum chemistry rationalizes the electronic structure of atoms and chemical bonds, which has unprecedentedly enlightened our understanding about chemical structures and transformations. Over the past few decades, advances in theoretical methods and sustained improvements in computational power of personal desktop and high-performance CPUs alike have enabled theoretical/computational quantum chemists to tackle increasingly larger systems.

Currently, we can assess many real-world problems relevant to various chemical transformations including those pertaining to the field of catalysis, for instance, with reasonable levels of theory that would allow sufficient accuracy to compare with experimental results and, more interestingly, to make quantitative predictions. At this intersection, insights from quantum chemistry meet empirical discoveries in the domain of experimental areas of chemistry including biological and organic chemistry providing an exciting playground of opportunities for computational-experimental partnerships aiming to improve our understanding of a given process or reaction mechanism; while nurturing the higher ambition of designing novel and/or more efficient processes to meet the challenges of our times.

Applications of quantum chemistry are abundant spanning from computation of properties for materials to reaction mechanistic studies of various types – basic organic transformations or complex organometallic catalysis, conformational analysis to spectroscopic properties of natural products. It is not surprising, therefore, that the number of publications describing computations and experimental results together have climbed sharply. The situation is such that computations can now be carried out in conjugation to experiments or even ahead of.⁴¹

In the field of natural products, a particular application which has gained quite a reputation over the past decade or so is the computation of ¹H and ¹³C NMR chemical shifts, and coupling constants in some cases. That has been applied successfully to a number of natural products to assist in structure elucidation.⁴² There has also been instances that such calculations were instrumental in the revision of structures, which were originally misassigned.^{43,44}

In this session, however, we shall focus on the discussion of applications of quantum chemistry to the biosynthesis of terpenes, a major class of natural products.

Quantum chemistry have been applied to a number of such studies on natural products or related structures of biological interest for self-contained projects or in conjugation with major experimental work.⁴⁵⁻⁵² To that end, a range of methods have been applied at different levels of theory. These computations provide useful information on the structure of intermediates and transition states along the reaction path and associated energetics. These are critical in some instances as they cannot be easily accessed otherwise. That is the case of information related to transition state structures, which can only be obtained indirectly through experiments. Therefore, application of quantum chemistry to study reactions in general plays a key role in assisting in the task of mechanism elucidation by offering molecular details and the overall reactivity principles.

The biosynthesis of natural products including terpenes in particular relies heavily on the intermediacy of carbocation structures. The structural modifications taking place along a biosynthetic route are promoted by carbocation rearrangements in interaction with amino acid residues, water molecules, co-factors or ions in the interior of the enzyme. This interplay largely controls the stability/reactivity of carbocations and is ultimately responsible for the great diversity of molecular architectures in terpenes. In this context, the structural features and reactivity of carbocations as well as the strategies an enzyme may utilise to achieve its catalytic task has been investigated by means of quantum chemistry in a number of studies. Several are the strategies an enzyme may resort to in promoting unfavourable reactions ranging from the stabilisation of transition states by electrostatic effects to reshaping the substrate to a less stable conformation required for reaction. Nonetheless, quantum chemistry studies on terpene biosynthesis have largely focused on the details related to the intrinsic reactivity of carbocation substrates, which has proven to be critical in terpene biosynthesis.⁵³

In this arena, the earliest contribution, to the best of our knowledge, dates back to 1974 when Gleiter and Mullen reported a model on the squalene cyclisation using semi-empirical methods.⁵⁴ The field remained essentially dormant till late 1990s when Jenson and Jorgensen published a contribution on the structure of important carbocations in steroid biosynthesis applying hartree-fock calculations.⁵⁵ Quantum chemical investigations into terpene biosynthesis only started to gather momentum from a few years later into the

following decade. Since then about 100 papers – virtually all using density functional theory (DFT), have been published on the topic from just a few groups with Dean Tantillo's (University of California, Davis, U.S.) being unarguably the top player and responsible for about half of all contributions.

We now turn to examples illustrating applications of quantum chemistry to the reactivity of carbocations in the biosynthesis of several terpenes showcasing the relevance of these studies in association with experimental work in addressing mechanistic questions at a molecular level. These are not organised in any chronological order or aim at being exhaustive, but rather we focus on introducing major concepts from top-notch works in the area.

1.4 Quantum Chemistry in the Biosynthesis of Terpenes

In 2016, as part of a biochemical study on terpene synthases in the livewort *Marchantia polymorpha*, the authors proposed a 1,4-alkyl shift for a step in the mechanism leading to sesquiterpenes ledol and viridol among other related structures.⁵⁶ While hydride shifts (1,2-, 1,3- and higher) are common among terpene rearrangements⁵⁷, extended alkyl shifts beyond a neighbouring carbon (1,2-alkyl shift) are much more rare due to the higher energy barrier associated. A well-established exception is the 1,3-alkyl shift involved in the interconversion of cyclopropylcarbinyl cations, which has been confirmed by labelling experiments⁵⁸ and validated by quantum chemical computations for their relatively low barrier *ca.* 6-11 kcal/mol.^{59,60}

The 1,4-alkyl rearrangement in question (path a, Fig. 18) was investigated by means of quantum chemistry⁶¹ and compared to previously computed data⁶² for an alternative mechanism (path b, Fig. 18) at the very same level of theory, mPW1PW91/6-31+G(d,p)//B3LYP/6-31+G(d,p). The predicted barrier for the initially proposed path A is considerably higher *ca.* 60 kcal/mol than that from the alternative proposal (path B) computed at 30–40 kcal/mol, which demonstrates that path A is not energetically viable.

An interesting feature of secondary carbocation A is that it is a nonclassical carbocation exhibiting 3-centre, 2-electron delocalisation implicating a nearby C–C bond at 2.03 Å away, which is elongated to 1.70 Å and has reduced C–C–C angle to 81°. This bond assist in the

stabilization of the carbocation via hyperconjugation. In the 1,4-alkyl shift mechanism, this stabilization is compromised, hence, the high activation energy.

Figure 18. Biosynthesis of sesquiterpenes ledol and viridiflorol showing key steps including evaluated paths a and b in the synthesis of carbocation C. Activation energies (E_a) are in kcal/mol as follows: B3LYP/6-31+G(d,p)//B3LYP/6-31+G(d,p) in normal text and mPW1PW91/6-31+G(d,p)//B3LYP/6-31+G(d,p) in brackets. Computed distances (Å) for carbocation A are also shown. Adapted from ref. 61.

Carbocation B (Fig. 18) has been located before in the context of a quantum chemistry study on the biosynthesis of sesquiterpene alcohol avermitilol.⁶² The biosynthetic proposed route contains two secondary carbocations (Fig. 19). One of these was located (carbocation 4) as a minimum in reaction path, the other structure (carbocation 5) was not located as such (Fig. 19).

Figure 19. Proposed mechanism for the synthesis of sesquiterpene avermitilol (adapted from ref. 62). Computed distances in Å.

Similarly, carbocation 4 at C2 (Fig. 19) benefits from stabilisation for being adjacent to a cyclopropenylcarbenyl cation, which shows the C–C bond immediately after the carbocation elongated to 1.61 Å, while the opposite C–C bond is shortened to 1.47 Å, as expected. It is noteworthy that this hyperconjugative effect is not an exclusivity of secondary carbocations. Quite contrary, it is a structural feature commonly observed for its ability to stabilize carbocation structures. Moreover, the carbocation centre at C2 is also in interaction with the C=C π -bond opposite in the ring as it is located close in space (2.44 Å and 2.69 Å) providing additional internal stabilisation (Fig. 19).

On the other hand, carbocation 5 (Fig. 19) does not enjoy any sort of structural feature relevant to stabilisation. Consequently, this carbocation was not located as a minimum. In fact, optimisation attempts of this structure have failed, which could be correlated to the fact that this carbocation may not exist in the absence of stabilizing non-covalent intermolecular interactions in the active site of the enzyme. That prompted the authors to consider a complex with a water molecule, $4 \cdot \text{H}_2\text{O}$, for the sake of a stabilising C–H $\cdots$ O interaction (Fig. 19).

Therefore, a concerted C2,C7-cyclisation and water attack to C6 leading directly to avermitilol ($4\bullet\text{H}^+$) was found with a low energy barrier.⁶²

The vast majority of tertiary carbocations found over cascade reactions in terpene biosynthesis contrasts sharply with very few secondary carbocations located as true minima.⁶³ As shown above, secondary carbocations may be avoided by participation in concerted events.⁶⁴ Oftentimes, secondary carbocations are rather transition state structures than minima in the reaction coordinate.

From the textbook general order of carbocation stability: $\text{CH}_3 < \text{primary} < \text{secondary} < \text{tertiary}$, one would not anticipate the general behaviour of secondary carbocations when compared to a tertiary by the presence of an additional alkyl group. Quantitative appreciation of carbocation relative stabilities reveal that the addition of a methyl group to carbocation CH_3^+ greatly increases stability by 25–40 kcal/mol (Table 4).⁶⁵ A second methyl group to CH_3CH_2^+ further improves thermodynamic stability by *ca.* 20–25 kcal/mol positioning secondary, isopropyl carbocation at the same level of stability of an allylic carbocation. From that, *t*-butyl carbocation gains additional 16 kcal/mol. While the values clearly show the considerably higher stability of tertiary carbocation, which may explain preference for this species in terpene biosynthesis, the case of secondary carbocation requires further elaboration as the reason for not observing this species in terpene reaction paths more often remains unanswered.

Table 4. Relative stabilities of carbocations in gas phase (kcal/mol).⁶⁵

Carbocation	MP4(SDQ)/6-31G(d,p)	MP2/6-31G(d)
CH_3^+	0.0	0.0
C_6H_5^+	–	–26.5
CH_3CH_2^+ (bridged)	–33.8	–40.8
CH_3CH_2^+ (open)	–39.0	–34.4
$\text{CH}_2=\text{CHCH}_2^+$	–54.5	–60.0
$(\text{CH}_3)_2\text{CH}^+$	–58.5	–58.9
$\text{C}_6\text{H}_5\text{CH}_2^+$	–	–74.5
$(\text{CH}_3)_3\text{C}^+$	–	–75.0

Evaluation of the kinetics of secondary carbocations shed light into the issue. Dynamics trajectory calculations on secondary carbocations⁶⁶ have shown that average lifetimes were found in the 35–100 fs range (standard deviations: 10–35 fs), which is in the same timeframe of a C–C σ -bond stretch (35–60 fs) but faster than bond-forming/breaking occurring over rearrangements of various carbocations (*ca.* 150–250 fs).⁶⁷ Therefore, it no surprise that several secondary carbocations are found close to the structure of transition states in the reaction coordinate. The remarkably short lifetimes of these carbocations can be considerably increased by non-covalent interactions,^{66,67} which should be the case in the interior of the active site.

In the context of terpene biosynthesis, the importance of non-covalent interactions have been demonstrated in various instances. We have already mentioned that these interactions, namely cation– π and C–H $\cdots\pi$, should be beneficial in altering the energy surface by stabilizing carbocations in the ground and transition state possibly resulting in lower energy barriers. Nonetheless, these interactions could have other more subtle purposes. The thought-provoking case of an important step responsible for major rearrangements in the biosynthesis of sesquiterpene trefalone A is presented (Fig. 20a).⁶⁸

The key transformation under investigation comprises a cyclisation step to form a cyclobutane ring followed by concerted ring opening and 1,2-hydride shift resulting in ring expansion. To evaluate the influence of a non-covalent interaction to the mechanism, energies were computed at the M06-2X/6-31+G(d,p) level of theory in gas phase and in the presence of a benzene molecule strategically positioned in the reaction centre to form a carbocation– π interaction. Benzene represents a simple model for the side chain of aromatic residues (e.g. Phe, Tyr), which are found in the active site of terpene synthases

Although energies have lowered in the first step of the mechanism, the barrier remained the same as reduction occurred by the same amount on reactant and transition state, which on its own shows that the benzene molecule in interaction had apparently no special effect on the energetics in this case (Fig. 20b). However, it has promoted an alternative mechanism in which ring expansion occurs directly without the intermediacy of cyclobutane. In contrast with the first step, activation energy dropped abruptly from 12.5 kcal/mol in the gas phase to 0.4 kcal/mol in the presence of benzene. That is, the ring opening/1,2-hydride shift step becomes nearly barrierless. More interestingly, the curvature has been altered to remove the deep minimum. In the benzene-free mechanism, the secondary carbocation is

avoided while this carbocation is stabilised by C–H $\cdots\pi$ interactions in the presence of benzene contributing to the strength of the hyperconjugation.

Figure 20. (a) Schematic mechanism for the formation of trefolone A showing key sequence under investigation (dotted-blue box). (b) Potential energy surface of carbocation rearrangement: in gas phase (black) and +C₆H₆ (red). Values in kcal/mol computed at M06-2X/6-31+G(d,p) level of theory. Original numbering from ref. 68 was kept.

Interestingly, most valuable information for an alternative mechanism could be abstracted from the theoretical model just shown. Theozymes as such are advantageous in allowing an expedient, qualitative assessment of the catalytic mechanism, which could guide mutation studies in the enzyme assisting in the search for more efficient catalytic processes.⁶⁹

Next, we discuss the theoretical studies on the biosynthesis of three-fused five-membered ring sesquiterpene pentalenene, which contains four adjacent stereocentres including one of them quaternary. A usually represented mechanism for this terpene formation is shown in Fig. 21.⁷⁰

Figure 21. Originally proposed mechanism for pentalenene formation.

The originally proposed mechanism represents a straightforward set of reactions *e.g.*, cyclisations and 1,2-hydride shifts leading to pentalenene. By means of quantum chemical computations, a couple of alternative mechanisms were identified. The structure of carbocation 1 was located as well as the transition state responsible for its conversion to 2 (Fig. 22). The structure of secondary carbocation 2 was also found under the influence of considerable hyperconjugation effect in a structure which can be viewed as a hybrid between a homoallylic cation and a cyclopropylcarbanyl cation (2, Fig. 22). However, attempts to locate the structure of a minimum corresponding to carbocation 3 failed (Fig. 22). Instead, a transition state structure linking directly carbocation 2 to 7 was identified, which represents the fusion of five-, six- and four-membered rings through the formation of two σ -bonds in a concerted manner (Fig. 22, red). As the next step, a transition state connecting 7 to 5 directly was found accounting for the simultaneous migration of a hydrogen and an alkyl group. Such

transformations are known as a dyotropic rearrangement; and by definition they imply simultaneous migration of two groups past each other.⁷⁰ There are various examples of mechanistic proposals from quantum chemical studies in terpene biosynthesis contemplating dyotropic rearrangements.⁷¹

A second mechanism for the transformation was outlined from the conformation 1' (Fig. 22, structures in teal), which is approximately 4 kcal/mol lower in energy than its conformer 1. Similarly, a transition state structure converting 1' to 2' was also identified. Carbocation 2' showed the same stabilisation pattern as discussed for 2. However, a transition state structure for the conversion of 2' to 3' could be located in this case, and showed comparable barrier (6–8 kcal/mol) to that associated with the rearrangement of 2 to 7 (5–6 kcal/mol). In fact, structures 3' and 7 bear some structural similarities. The transition state associated with the final rearrangement, a hydrogen shift, proved to be an unusual proton sandwich between two alkenes (Fig. 22), which is located on an energy plateau. Carbocation 8 is just 0.8 kcal/mol lower in energy than the transition state responsible for its formation from 3', and 2.0 kcal/mol lower than the transition state for its conversion to 5 at mPW1PW91/6-31+G(d,p)//B3LYP/6-31+G(d,p) level of theory.

This case highlights the possible role of the enzyme in the preorganisation of reactants as quite different mechanisms followed from distinct conformations (1 and 1').

Figure 22. Theoretically proposed mechanisms for the formation of sesquiterpene pentalenene. In above proposal (in purple), a dyotropic rearrangement at the last step is outlined. For the lower proposal (in teal), a proton sandwich appears as the last rearrangement.

Thus far, we have mainly shown examples involving hydride transfer. In the last example, the unusual proton-sandwich in one of the steps was considered as an alternative mechanism. In that case, it was not compared against a hydride alternative. Is there any preference for hydride or proton transfer?

Bearing the question in mind, let us examine the case of yet another sesquiterpene trichodiene produced by fungi and precursor of various antibiotics and toxins.^{72,73} The structure of the key intermediate to this pathway, cuprenyl cation, can be formed in various ways (Fig. 23).⁷² All of them from bisabolyl cation, which is another major intermediate in sesquiterpene biosynthesis. Alkene cyclisation of bisabolyl cation could form secondary carbocation A (Fig. 23, in brown) which after 1,4-hydride transfer would produce cuprenyl cation. However, the structure of a minimum corresponding to carbocation A was not located. Authors would instead consistently find the minimum structure of more stable tertiary carbocation B (Fig. 23, in purple). Attempts to locate a transition state for the formation of A also failed but pointed to the formation of B following a concerted cyclisation and methyl shift. Authors also failed but pointed to the formation of B following a concerted cyclisation and methyl shift.

Figure 23. Theoretically computed pathways for the biosynthesis of sesquiterpene trichodiene (adapted from ref. 73).

From that, a transition state directly connecting B to cuprenyl cation was found through a 1,4-hydride transfer followed by [1,2]-methyl shift. Note that methyl group through sequential [1,2]-hydride shifts returns to its original position. This rearrangement is often referred to as “temporary methyl shift”. Another alternative pathway linking bisaboly and cuprenyl cations involves a 1,5-proton transfer rather than a hydride transfer to form carbocation C. In this case, the C=C π -bond acts as a base capturing the proton. Conformation change in C followed by cyclisation leads to cuprenyl cation.

Barriers associated to these rearrangement were computed at B3LYP/6-31+G(d,p) and mPW1PW91/6-31+G(d,p)//B3LYP/6-31+G(d,p) levels of theory (Table 5). Energies associated with the pathway involving cation B has considerably higher energies (ca. 20–33 kcal/mol) than related to the intermediacy of carbocation C (ca. 0.5–9 kcal/mol). Final cycylisation to cuprenyl from C occur with a very small barrier.

Table 5. Energies barriers for transformations in bisaboly and cuprenyl carbocations

	B3LYP/6-31+G(d,p)	mPW1PW91/6-31+G(d,p)// B3LYP/6-31+G(d,p)
TS_{bisaboly-to-B}	25.9	20.3
TS_{B-to-cuprenyl}	33.4	30.9
TS_{bisaboly-to-Ca*}	8.6	5.4
TS_{Ca-to-Cb}	4.6	4.3
TS_{Cb-to-Cc}	0.3	0.3
TS_{Cc-to-cuprenyl}	0.5	−0.7

* Ca, Cb and Cc refer to conformations on cation C which requires conformation changes prior to cyclisation

From the quantum chemical calculations, there is strong support for the proton-transfer mechanism pathway in trichodiene biosynthesis. To overcome the energetic preference, the enzyme would need to incur considerable interference towards the hydride transfer mechanism.

Our favourite diterpene taxol was also subject to a few computational works.⁷⁴⁻⁷⁷ In a recent study combining molecular dynamics and QM/MM, it was demonstrated that taxadiene synthase is a slow-starting enzyme as a strategy to avoid excess release of heat at the early steps – activation leading to formation of carbocation and cyclisation, when there is a higher tendency to promiscuity.⁷⁷ Terpene synthases produces a pool of products favouring some for various reasons. Promiscuity is, therefore, a recurrent topic and also associate to carbocation reactivity. In the case of taxadiene synthase, for instance, product distribution reveals that the main product taxa-4,11-diene is formed in 93.2% yield and isomer taxa-4(20),11-diene in 4.7%. Side products verticillene is formed in 2.1% yield and cembrene A is obtained in negligible amounts. In the same study, mutation of a tyrosine residue (Y841) to histidine would favour formation of cembrene by placing the basic residue in an advantageous position to deprotonate cembrenyl cation, which would radically alter product ratios.⁷⁷

The topic of proton transfer as we have discussed a few lines above and the various possible mechanisms in terpene biosynthesis brings us to the discussion of proton promiscuity. In regard to the biosynthesis of taxol (Fig. 14, page), we have shown an unusual intramolecular proton transfer (C11-to-C7 α) as part of the proposed mechanism. How this rearrangement is brought about was subject of a quantum chemical study at the mPW1PW91/6-31+G(d,p)//B3LYP/6-31+G(d,p) level of theory.⁷⁴

Labelling studies have ruled out the intermolecular deprotonation/reprotonation possibility.^{34,36} To probe the intramolecular proton transfer hypothesis, a transition state (TS_{1-to-2}) structure representing the direct proton transfer in question was located as well as the cations reactant 1 and product 2. The computed barrier associated with this rearrangement was found at 11.3 kcal/mol, and structures 1 and 2 showed similar energy content with cation 2 only 2.2 kcal/mol higher (Fig. 24). Inspection of the transition state structure revealed that the migrating proton was at a similar distance from the recipient carbon at 2.46 Å away and the other π -double bond (2.20 Å). The structure of a transition (TS_{1-to-3}) responsible for this proton transfer was found and the related product 3; the energy barrier was considerably lower at 5.8 kcal/mol. A similar barrier (5.0 kcal/mol) was obtained for the next transition state leading to the expected product 2.

This novel proposal shows that the proton transfer in two steps is favoured, and from computed barriers their movement inside the structure as in a cage requires little energy.

Figure 24. Computed transition states geometries and energies [mPW1PW91/6-31+G(d,p)//B3LYP/6-31+G(d,p)] in kcal/mol for both pathways leading to the formation of taxadiene cation 2. Selected distances in Å.

As we bring this introduction to a close, a few final words. Enzymes are ingenious biomachines capable of accelerating chemical reactions rates⁷⁸ by up to 10^{20} fold enabling the complexity and diversity of life on the planet. Terpenes synthases are indeed a special bunch for being responsible for about 60% of all natural products and so doing by utilising just a handful of acyclic precursors. The application of quantum chemistry and other computational methods to investigate the biosynthesis of terpenes has opened up the doors to the fascinating world of molecular transformations likely to occur in the interior of these enzymes offering a privileged microscopic perspective of events in unmatched detail. Advances in computational methods and hardware would further speed up the process of turning computations into chemical insight and facilitate tackling increasingly more complex problems with higher accuracy.

2 METHODS

Quantum chemistry has come a long way since the early theoretical developments in quantum mechanics with the seminal contributions from Heisenberg, Schrödinger and Dirac among several others. A number of important advances in the following decades would set the ground for a range of methods that we now well-recognise (Fig. 25). The application of quantum mechanics to organic chemistry, for instance, can be traced back to Hückel's π -model and the molecular orbital theory of the 1930s. Nonetheless, several decades would follow until solutions to multielectron multiatom molecules would be available for the general community. The introduction of faster computers together with the discovery of various computational principles prompted computational chemists to tackle basic organic molecules. The turning point was the release of Gaussian 70, the first general-purpose programme to bring the *ab initio* method to the mainstream. To give an idea of situation back then, a single point HF/STO-3G energy calculation on vinylcyclobutane would be considered a challenging case.⁷⁹ Today, the same calculation takes no time – one spends more to prepare the input than Gaussian 09⁸⁰, which was used in this thesis, does to optimise the geometry of vinylcyclobutane in a consumer-grade desktop.

Figure 25. Highlights in the development of computational organic chemistry since discovery of quantum mechanics in the 1920s.

Over the decades, the computational costs have largely decreased to the point that a mere student as this one writing can make an invest into a decent machine. That is a

continuous trend following Moore's law which projected that computational power or at least the number of transistors would double every two years.⁸¹ Presently, that leaves the community in a very comfortable position but also great responsibility to make the best use of resources and methods; and exercise creativity –, dare to take risks for the sake of advancing our cause. Unfortunately, that is not always the case.

While researchers in computational chemistry are largely free to approach a chemical problem in a variety of ways leaving ample room for creativity, there are assumptions being made routinely based on the method and model, which are intrinsically related to the question being addressed. Beyond good chemical reasoning, one should not be too carried away by the beauty or simplicity of a given answer but reflect on the limitations imposed by methods and the model itself. More importantly, computations benefit from experimental validation.

2.1 Quantum Mechanics. The Schrödinger Equation

The quantum mechanics theory applies to the observation of effects at a microscopic level to which the laws of classical mechanics fail to provide a satisfactory description. All bodies and particles obey the laws of quantum mechanics, while classical mechanics is an adequate approximation for large systems. Therefore, both offer the same answer in the macro world.

Some fundamental concepts emerge from the quantum mechanics theory. One of them concerns the wave-particle duality which states that all particles possess wave and particle properties. Discretisation of magnitudes is another principle, which says that a property can only have discrete or quantised values. That is the case of electronic energies in an atom, for example.

A third concept is the uncertainty principle formulated by Heisenberg,⁸² which postulates on the accuracy that given pairs of physical properties of a particle, such as momentum and position, cannot be determined simultaneously with precision. The uncertainty principle offers a series of probabilities for the location of a particle for its given momentum. These probabilities are designated by the wavefunction (Ψ), which depends on the quantum state at the very moment of the measurement.

To solve problems involving the flow of electrons in a chemical structure such as the ones of interest here in rearrangements with bonds being broken and new ones formed, quantum chemistry offers a quality solution for its representation of electrons.

Calculations in quantum chemistry require solution of the Schrödinger equation (Eq. 2.1), which associates energy with the distribution of electrons in a particular position around the nuclei.

$$H\Psi = E\Psi \quad (2.1)$$

In this equation, a given energy (E) is associated with an electron density expressed as a wave function (Ψ) by means of a Hamiltonian operator (H). In math, an operator is a function that acts on another function. In this case, the Hamilton operator contains terms that associate energy with the relative positions of electrons and the nuclei. When applied to the wave function (Ψ), the Hamilton provides the total energy of a given system. As in classical mechanics, the operator is usually expressed as the sum of the operators for kinetic (T) and potential (V) energies, Eq. 2.2.

$$H = T + V \quad (2.2)$$

There are elaborations to the Schrödinger equation, namely the time-dependent variation or the relativistic formalism. Nevertheless, calculations in the present work makes use the time-independent solution to the Schrödinger equation throughout. We shall focus on other aspects in the continuation of this session.

An exact solution to Schrödinger equation would lead to an accurate description of the system at quantum mechanics level, which would result in high quality prediction of properties. For multi-electron systems, however, that represents a challenge due electron-electron repulsive interactions, and an exact analytical solution is unavailable from the electronic Hamiltonian. Various wavefunctions could be applied to solve the Schrödinger equation (Eq. 2.1). The variational principle, however, states that the ground state is better represented by the wave function with the lowest energy associated.

Hartree offered a classical approach to solve the problem. He suggested that the electronic wave function dependent of the N -electrons coordinates was simplified to the product of dependent functions related to one-electron coordinates (Eq. 2.3).⁸³ The product

of these single electron functions known as the Hartree product would generate what we know as molecular orbitals.

$$\Psi_{ele}(r) = \phi_1(r_1) \phi_2(r_2) \dots \phi_i(r_i) \quad (2.3)$$

Introducing the spin component (ω) into the spatial function brings about the spin-orbitals. Fock and Slater subsequently gave their contribution by adding the anti-symmetry related to the permutation of electronic coordinates in the function, which came as the determinant in Eq. 2.4.⁸⁴ The so-called Slater determinant has the spin-orbitals (ϕ_i) written in relation to spatial coordinates (r_i) and spin (ω_i), where N is the number of electrons in the system. Following Pauli exclusion principle, this function describes the quantum state of a system.⁸⁵

$$\Psi_{ele}(r_1, \omega_1 \dots r_i, \omega_i) = \frac{1}{\sqrt{(N!)}} \begin{vmatrix} \phi_1(r_1, \omega_1) & \phi_2(r_1, \omega_1) & \dots & \phi_i(r_1, \omega_1) \\ \phi_1(r_2, \omega_2) & \phi_2(r_2, \omega_2) & \dots & \phi_i(r_2, \omega_2) \\ \dots & \dots & \dots & \dots \\ \phi_1(r_n, \omega_i) & \phi_2(r_i, \omega_i) & \dots & \phi_i(r_i, \omega_i) \end{vmatrix} \quad (2.4)$$

By applying the variational method and the Slater determinant as a tentative wavefunction, Hartree-Fock equations can be written leading to solutions represented as the best set of $\{\phi_i\}$, which in turn are associated with the lowest energy for a given system.

On perhaps a different tone, the Born-Oppenheimer approximation method offers an elegant solution by computing analysis of the nuclei and electrons separately.⁸⁶ In so doing, it assumes that atomic nuclei are approximately three orders of magnitude heavier than the electrons. Therefore, it is reasonable to solve the time independent Schrödinger equation for an electronic system considering stationary nucleus.

The Schrödinger equation is then decomposed in two parts: one part describing the electronic wavefunction (Ψ_{ele}) considering a fixed geometry for the nucleus; and the other accounting for the nuclear wavefunction (Ψ_{nuc}). For the latter, the energy associated with the electronic wavefunction is responsible for the potential energy (Eq. 2.5). Another important assumption in the Born-Oppenheimer approximation is that the electronic wavefunction depends on the coordinates of the nuclei but not on their momenta.

$$\Psi_{tot}(R, r) = \Psi_{ele}(R, r) \Psi_{nuc}(R) \quad (2.5)$$

where R and r denote nuclear and electronic coordinates, respectively.

To proceed, let us rewrite the Schrödinger equation for a given molecule (Eq. 2.6) in which the total Hamiltonian (H_{tot}) is in full represented by Eq. 2.7

$$H_{tot}\Psi_{tot}(R, r) = E_{tot}\Psi_{tot}(R, r) \quad (2.6)$$

$$H_{tot} = T_{tot} + V_{tot} = (T_{ele} + T_{nuc}) + (V_{ne} + V_{ee} + V_{nn}) \quad (2.7)$$

where T_{ele} and T_{nuc} denote kinetic energy of electrons and nuclei, respectively. V_{ne} is the coulombic attraction between electrons and nuclei, and V_{ee} and V_{nn} represent, respectively, coulombic repulsion between electrons and between nuclei, respectively.

Making the assumptions: nuclei move far more slowly than orbiting electrons, and that these electrons are moving around in the potential of stationary nuclei, it implies that the term related to nuclear kinetic energy can be cancelled out from Eq. 2.7. Moreover, repulsion between nuclei in the system can be considered constant. Hence, we can write the electronic Hamiltonian as follows (Eq. 2.8):

$$H_{ele} = T_{ele} + (V_{ne} + V_{ee} + V_{nn}) = T_{ele} + V \quad (2.8)$$

Once applied (Eq. 2.8) to the electronic wavefunction, solving the equation (Eq. 2.9) result in the electronic state of a given molecule.

$$(T_{ele} + V)\Psi_{ele}(R, r) = U_n(R)\Psi_{ele}(R, r) \quad (2.9)$$

Having worked out a solution to the electronic part of Schrödinger equation, solving the complete equation represents the next step. To that end, application of the complete Hamiltonian (Eq. 2.7) to the nuclear wavefunction is required (Eq. 2.10).

$$(T_{nuc} + T_{ele} + V)\Psi_{nuc}(R) = E_{tot}\Psi_{nuc}(R) \quad (2.10)$$

Bearing in mind that electrons move way faster than the nuclei, it is reasonable to replace the electronic Hamiltonian by its average value, which modifies Eq. 2.10 into:

$$[T_{nuc} + U_n(R)]\Psi_{nuc}(R) = E_{tot}\Psi_{nuc}(R) \quad (2.11)$$

The potential U_n , adiabatic potential, comes from the resolution of the electronic version of Schrödinger equation. This potential applied to the calculation of a number of geometries is known as the potential energy surface (PES)

To sum up, the Born-Oppenheimer approximation assumes that nuclei move on a potential energy surface, which is obtained from the solution of the electronic Schrödinger equation. This approach based on the nuclei moving on a potential energy surface stimulated the development of methods focused on solving the electronic Schrödinger equation, which are generally known as electronic structure calculations. These methods for their part allow computation of energies and properties that are widely used in computational quantum chemistry in numerous applications.

2.2 Density Functional Theory

For delivering good quality results at a reasonable computational cost, density functional theory (DFT) has become the method of choice for daily applications in computational quantum chemistry. Its status is such that DFT protocols are implemented in virtually all programme suites. DFT is most often applied following the Kohn-Sham scheme. According to which, an approximate exchange correlation has to be selected. The quality of a DFT calculation largely depends on the exchange-functional chosen as we shall discuss shortly for the case carbocations. For now, let us get to the foundations of this theory.

The resolution of the electronic Schrödinger equation (Eq. 2.9) for a system of general interest to an organic chemist despite all the assumptions, represents still a complicated problem. It implies the computation of the wavefunction for nuclear coordinates, which represents a function of $3N$ spatial coordinates and N spin variables where N represents the number of electrons. However, there is a way out with a different formulation, which simplifies the issue. Density functional theory (DFT) offers an alternative by using the electron density instead of the wavefunction to calculate electronic properties, therefore, offering the advantage of requiring only three spatial coordinates (x, y, z).

By definition, the electron density $\rho(\vec{r}_1)$ represents the multiple integral across the spin variables for all electrons over all spatial coordinates except one (Eq. 2.12),

$$\rho(\vec{r}_1) = N \int \dots \int |\psi(\vec{x}_1, \vec{x}_2, \dots, \vec{x}_n)|^2 ds_1 d\vec{x}_2 \dots d\vec{x}_N \quad (2.12)$$

where,

$$d\vec{x}_i = ds_i d\vec{r}_i \quad (2.13)$$

The electron density determines the probability of finding any of the N electrons within the volume element $d\vec{r}_1$ but with arbitrary spin while the other $N-1$ electrons have arbitrarily random positions and spin in the state represented by the wavefunction ψ . Therefore, $\rho(\vec{r}_1)$ in strict terms is a probability density rather than the electron density, although in practice it is known as such. Furthermore, the electronic density approach is more appealing for being more computationally efficient and having a physically interpretation.

The origins of the density functional theory as we currently know can be traced back to 1964 when Hohenberg and Kohn published a seminal paper describing two theorems, which are the steppingstones of DFT.⁸⁷

The first Hohenberg-Kohn theorem as they are known offer “proof of existence”. Quoting directly from the original⁸⁷: “the external potential $V_{ext}(\vec{r})$ is (to within a constant) a unique functional of $\rho(\vec{r})$; since, in turn, $V_{ext}(\vec{r})$ fixes $\hat{H}$ we see that the full many-particle ground state is a unique functional of $\rho(\vec{r})$.” That implies the existence of an exclusive one-to-one correspondence between the ground state electron density ρ_o and the external potential $V_{ext}(\vec{r})$. As a consequence, all ground state properties of a given system are defined by its electron density.

As the ground state energy represents a functional of the ground state electron density, the energy of a given system can be written as equation 2.14.

$$\begin{aligned} E[\rho] &= F_{HK}[\rho] + V_{ext}[\rho] \\ &= F_{HK}[\rho] + \int \rho(\vec{r}) v_{ext}(\vec{r}) d\vec{r} \end{aligned} \quad (2.14)$$

where $F_{HK}[\rho]$ is the so-called Hohenberg-Kohn functional, which consists of the kinetic energy functional $T[\rho]$ for the electron and the electron-electron repulsion energy functional $V_{ee}[\rho]$. Therefore,

$$F_{HK}[\rho] = T[\rho] + V_{ee}[\rho] \quad (2.14)$$

From equation 2.14, Schrödinger equation can be solved if we know $F_{HK}[\rho]$. Unfortunately, the exact form of the functional $F_{HK}[\rho]$ is not known. Nonetheless, the functional is independent of the system; it could be applied to whichever.

While the ‘holy grail’ of an exact functional is not available, the quest for expressions as potential quality functionals represents a major area of research in DFT. We will elaborate a bit further on this shortly.

Once more reading from the original, the second theorem states: “The functional $F_{HK}[\rho]$, which provides the ground state energy of the system, provides the lowest energy if and only the input density is truly the ground state density.” That is, the second theorem aims at demonstrating that a certain density corresponds to the desired ground state density, which simply put establishes the variational principle in terms of density (Eq. 2.15).

$$E[\rho_0] = \min_{p \rightarrow N} E[\rho] \quad (2.15)$$

$E[\rho_0]$ functional reaches its minimum in relation to all allowed densities if only the provided density is indeed a ground state density. To obtain the exact energy for a given system, the electron density that minimises the energy is required (Eq. 2.16).

$$\left[\frac{\partial E[\rho]}{\partial \rho} \right] = 0 \quad (2.16)$$

As aforementioned, a major drawback of the density functional theory is that the exact expression of the functional $F_{HK}[\rho]$ is not known. An alternative to overcome this issue is to solve the N -electron problem applying an approximate expression of the exact functional. Aiming at developing approximations for the exact functional $F_{HK}[\rho]$ led to the Thomas-Fermi (TF) and Thomas-Fermi-Dirac (TFD) formulations, which represents the basis for the Kohn-Sham method. Differently from their predecessors who focused on determining accurately the kinetic energy by means of a functional of density, Kohn and Sham concentrated on calculating the kinetic energy exactly.

Kohn-Sham approach aims at obtaining the exact kinetic energy of a non-interacting system with the same density as the real interacting system. Even if both system have the same density, a non-interacting and interacting systems have non-equal energies – that represents the assumption in their approach to simplify the problem. They wrote a Slater

determinant representing the exact kinetic energy of a non-interacting system with the same an interacting system (Eq. 2.17).

$$T_S = -\frac{1}{2} \sum_i^N \langle \phi_i | \nabla^2 | \phi_i \rangle \quad (2.17)$$

All necessary corrections were computed into a new term, E_{xc} . Hence, the expression for $F_{HK}[\rho]$, Eq. 2.18:

$$F[\rho] = T_s[\rho] + J[\rho] + E_{xc}[\rho] \quad (2.18)$$

and $E_{xc}[\rho]$, exchange-correlation energy, is defined by Eq. 2.19:

$$E_{xc}[\rho] = (T[\rho] - T_s[\rho]) + (V_{ee}[\rho] - J[\rho]) \quad (2.19)$$

where the two first terms aim at correcting the difference in kinetic energy between interacting and non-interacting system, and the last two refer to the correction between the classical electron-electron interaction and the real case. The exchange correlation energy is the functional that contains all that is unknown.

Cutting short a long way into further mathematical manipulations, we present Kohn-Sham equations:

$$-\left(\frac{1}{2}\nabla^2 + v_{eff}\right)\phi_i = \varepsilon_i\phi_i \quad (2.20)$$

where the term in parenthesis is h_{KS} , the Kohn-Sham monoelectronic Hamiltonian, and the term v_{eff} is the effective core potential, which can be defined as:

$$v_{eff}(\vec{r}) = \int \frac{\rho(\vec{r}_2)}{r_{12}} d\vec{r}_2 + V_{xc}(\vec{r}_1) - \sum_a^M \frac{Z_a}{r_{1a}} \quad (2.21)$$

In Eq. 2.21, the only unknown term is the potential V_{xc} as it has a correlation to the exchange-correlation E_{xc} . And V_{xc} is defined as a functional derivative of E_{xc} related to the electron density:

$$V_{xc} \equiv \frac{\partial E_{xc}}{\partial \rho} \quad (2.22)$$

Although the Kohn-Sham method is exact in principle, the exact forms of E_{xc} and V_{xc} are not known. Otherwise, the method would compute exact energies. In practice, the Kohn-Sham equations have to be solved iteratively as the effective potential depends on the electron density. From an initial trial electron density, the effective potential is calculated with which Kohn-Sham equation can be solved and give out orbitals ϕ_i . These are then computed resulting in a new electron density. This iterative process is continually repeated until a desired convergence criterion is satisfied.

2.3 Exchange Correlation Functionals: Application to Terpene Biosynthesis

Exchange-correlation functionals are commonly designed following an exchange term, E_x , and a correlation term, E_c , (Eq. 2.23), which are usually calculated separately by different approximations, and then combined into exchange-correlation functionals. Fitting parameters from accurate data – experimental or higher level wavefunction-based methods, may also be used in the process.

$$E_{xc}[\rho] = E_x[\rho] + E_c[\rho] \quad (2.23)$$

Each DF adopts a different formulation following varying levels of empiricism. Some focus on parametrisation exclusive to the rules of quantum mechanics e.g., PBE, TPSS; others have empirical parameters fitted to experimental data e.g. B3LYP, M062X, etc.

Today, computational chemists are confronted with an overwhelming amount of a few hundred density functionals (DFs) approximations to choose from.⁸⁸ Nonetheless, many recently generated DFs fail short in accuracy.⁸⁹ Benchmarking studies offer help by systematically assessing performance of various DFs for a given problem by comparing DFT results to high-level wave mechanics calculations or experimental results. However, in the wide world (or zoo⁹⁰, if you like) of DFs, popularity speaks volumes as very few DFs end up

being applied in most cases. In this context, B3LYP functional is second to none; and to this day very much in use despite mounting evidence about its shortcomings.^{91,92}

Density functional are classified according to various systems. Perdew's approach representing DFs in hierarchical levels as rungs in an analogy to the biblical Jacob's ladder has become popular.⁹³ In his view, the rungs in DFT ascend from the Earth, that is, the Hartree-Fock level of theory towards the chemical accuracy "heaven" (Fig. 26).

Figure 26. (a) Perdew's Jacob's ladder with exchange-correlation functionals represented in hierarchical rungs. (b) Yoram Raanan's artistic view of Jacob's ladder.

In each of the rungs in Perdew's ladder, another classification is represented e.g. LDA, GGA, which refers to the approach taken in constructing these functionals in the class. For instance, density ρ (see detail in teal on each box, Fig. 26) for the local density approximation (LDA), the Laplacian of the density, $(\nabla^2\rho(r))$, and/or the kinetic energy density, τ , for meta-GGA (generalized gradient approximation) functionals and so forth.

As no density functional performs well on all tasks, one needs to choose DFs carefully on a case-by-case basis. Furthermore, depending on necessity, one can go up or down in the ladder of DFs according to necessity, however, there is always a cost-benefit ratio to consider.

Going up the ladder would most likely improve accuracy of results but also represents greater computational cost.

In the context of carbocation reactions in terpene biosynthesis in the absence of the enzyme to assess intrinsic reactivity, various studies have surveyed DFs. Despite drawbacks of B3LYP, it has become the most employed DFs for computing geometries. That is not an exclusivity of terpene carbocations but still a general trend in computational organic chemistry. Results from B3LYP have been compared to MP2 in various instances for their ability to predict energy and geometry of carbocation terpenes.⁹⁴⁻⁹⁷ Most often, results agree except in those cases that carbocation structures reside in a flat surface of potential energy. MP2 method for its turn is known to favour delocalised/bridged carbocations.⁷¹

A series of DFs were assessed for their ability to reproduce energies of C₂–C₄ carbocations by comparison with higher level methods, MP4SDTQ/6-311+G(2d,p)//MP2/6-311+G(2d,p) and CCSD(T)/6-311+G(2d,p)//MP2/6-311+G(2d,p). BB1K/6-31G+(d,p) proved to be the best DF among those examined. Other methods also performed relatively well including B98, BMK, mPWB1K, mPW1PW91, PBEPBE and PBEPBEh. Energies from B3LYP showed biggest deviations, but never exceeded 4 kcal/mol as expected for this functional.⁹⁸ In the case of terpenes, the tendency of B3LYP to underestimate terpene carbocations had been reported previously, and single points at mPW1PW91/6-31+G(d,p)//B3LYP/6-31+G(d,p) level of theory are recommended.⁹⁹ This said approach has been widely applied to a variety of terpenes including several examples from Tantillo's group. It is worth mentioning that B3LYP can identify transition structures when other methods have difficulty or fail to locate.

2.4 Potential Energy Surface

We have previously mentioned that the potential energy (PES) is the adiabatic potential, U_n , obtained in the Born-Oppenheimer approximation solution to the Schrödinger equation (Eq. 2.9) for a given set of fixed nuclear coordinates. Thus, solving the equation for all possible configurations produce a complete PES. That is a feasible task only for a small molecule with 3–4 atoms given that it is $3N-6$ space, where N is the number of atoms. For molecules of general interest in organic chemistry, it is practically impossible. Therefore, the general strategy in computational quantum chemistry is to focus on the relevant parts of the

surface to obtain the important information. From the chemical point of view, the interesting parts are often related to the points of minima and first-order saddles, where the energy is stationary in respect to the nuclear coordinates (Fig. 27).¹⁰⁰

Figure 27. (a) A two-dimensional representation of a reaction path according to the intrinsic reaction coordinate from reactants to products going through the transition state structure.¹⁰⁰ (b) A three-dimensional representation of a potential energy surface (PES) of a general reaction.¹⁰¹

Minima along the surface are potentially observable as stable structures: reactants, products or intermediates, which supply information on the nuclear configuration of a molecule. On the other hand, first-order saddle points or transition states are peaks in the surface; points of maxima that are minimum in all but one direction. We look for these related points of minima and maxima (TSs) in the PES for the steps in a chemical transformation. Analysis of these structures are critical in the study of reaction mechanism. The geometries offer a means of observing the movement of atoms along the transformation and chemical groups in interaction, which by comparison provides the basis for understanding the molecular origins of selectivity.¹⁰² Energy barriers or activation energies ($\Delta G^\ddagger$), which is the difference in free energy between a reactant and associated transition state, offers a quantitative evaluation. Equally important is the reaction kinetics that can be brought into play by considering the Eyring¹⁰³ equation (Eq. 2.24), which connects the activation energy ($\Delta G^\ddagger$) to the reaction rate (k).

$$k = \frac{k_B T}{h} e^{-\frac{\Delta G^\ddagger}{RT}} \quad (2.24)$$

Under the transition state theory framework,^{103,104} molecular systems follow the downhill path represented in an intrinsic reaction coordinate (IRC), which connects two minima through the structure of a transition state (Fig. 27).¹⁰⁵ However, some systems may deviate from the IRC path as it is the case of reactive intermediates, e.g. carbocation rearrangement reactions. These cases are not explained by the transition state theory as they are largely controlled by dynamical effects. Kinetic energy becomes particularly important in the reaction path implicating in structural changes as a result of energy redistribution.¹⁰⁶ There are certain known patterns in which the system diverges from the typical IRC pathway: (1) the presence of a shallow intermediate especially when the preceding TS has a high energy represents one such case; (2) “ambimodal” transition states is another, which are present in reaction paths that show post-transition state bifurcations (PTSB) leading to several minima.^{100,107} These cases are better approached by modern computational methods applying quantum dynamics.^{108,109}

3 AIMS & SCOPE

In the class of terpenes, smaller members e.g. monoterpenes (C₁₀), sesquiterpenes (C₁₅) and diterpenes (C₂₀) have been favoured in quantum chemical studies. There are also a few investigations on sesterpenes, but the sub-class of triterpenes have been under-represented despite being the most numerous.^{3,110-121} Part of the reason for that is probably down to the size and associated complexity. Most computational works on triterpenes have focused on various aspects of the squalene cyclization. Therefore, we initially considered there could be potentially missed opportunities for late stage rearrangements.

By mid-2017, we became aware of the pentacyclic triterpene friedelin from a collaboration of researchers in institutions in the local area.¹²² For the challenging sequence of rearrangements according to the proposed mechanism (Fig. 28), the title triterpene got our attention as a prospective target for quantum chemical studies. It contains a diversity of transformations: cyclisations, ring expansions and hydride/alkyl shifts in various chemical environments. Although mostly composed of tertiary carbocations, there is a couple of secondary carbocations in the initial steps. Besides it represents the most rearranged pentacyclic triterpene produced by an oxidosqualene cyclase.¹²²

Moreover, friedelin has shown various biological activities, which also justifies the study of mechanisms related to its biosynthesis. Among important activities, we highlight: antimicrobial against Gram-positive and -negative bacteria including *Mycobacterium tuberculosis* as well as being able to affect the growth of *Candida sp.*,^{123,124} analgesic, antipyretic, anti-histaminic, gastroprotective and anti-inflammatory properties.¹²⁵⁻¹²⁷

Further literature search on computational studies dedicated to triterpene structures revealed that friedelin – or more precisely friedelane (ketone absent), had already been subject of a quantum chemical investigation of rearrangements in its biosynthesis by the group of Prof. E. J. Corey about a decade ago.¹²⁸ Interestingly, it should be pointed out that the structure of friedelin was characterized by Prof. Corey himself over 50 years earlier(!)¹²⁹

Nonetheless, we found various points that we could improve towards a more robust contribution in a complete study with the due level of detail, including: (a) use of the full structure of friedelin to include the ketone in ring A instead of friedelinane used by Corey and co-workers; (b) expand the route to include structures starting from the common triterpene precursor, cation dammarenyl. Corey and co-workers started from an equivalent to lupanyl cation (Fig. 28); (c) locate transition state structures for all steps, which are lacking altogether in the original; (d) improve the level of theory by applying the larger basis set 6-31G+(d,p)

instead of 6-31G(d), and use other density functionals that have shown better performance in carbocation rearrangement studies.

Figure 28. Putative biosynthetic carbocation cyclization/rearrangement mechanism for the formation of pentacyclic triterpene friedelin.

Each carbocation in the sequence above may be intercepted by a nucleophile such as water, for instance, to produce an alcohol or have a neighbouring proton abstracted to form an alkene triterpene. Carbocation intermediates may also serve in parallel pathways. From the project proposal, we also planned to utilise molecules (e.g., water, ammonia, benzene) as non-covalent interaction donors mimicking residues in the active to assess the effect on the reactivity of carbocations whenever strategically necessary (Fig. 29).

Figure 29. A benzene in non-covalent interactions (cation- π and C-H $\cdots\pi$) with a triterpene carbocation.

In overall, we aimed at investigating the inherent reactivity of carbocations pertaining to the pathways shown above. Towards this end, all density functional computations were run at the gas phase. While all intermediates and transition states were analysed for their geometrical features and energetics for a comprehensive discussion. We also intended to examine the full picture trying to correlate our results to mechanistic proposals and information available in the literature.

4 RESULTS & DISCUSSION

Before disclosing our results, an important note should be made on the structure of friedelin. We have run into several wrong representations of this triterpene in the literature and structures of intermediates found in its biosynthetic pathway, including from highly-cited items reported by major players in the field. That may seem an obvious remark but we could not stress enough the importance of checking various reports especially originals. Natural products are often complex molecules containing many structural nuances – and here the devil is in the detail. Mistakes may appear as a single inversion of configuration to an absurdly wrong structure (Fig. 30). By so doing, one can save computational time and avoid the pains of digressing on alternative mechanisms. We leave this cautionary note based on our own experience in this particular case.

Figure 30. Examples of incorrect structures in the literature for friedelin and carbocation intermediates in friedelin biosynthesis. Differences from correct structure is highlighted in red.

To begin with, we present a naming system for the rings in the pentacyclic core and numbering for the carbons in the structure backbone, which will be used in our discussions throughout. As rearrangements occur and particularly in the early steps of ring formation, numbering may vary slightly (Fig. 31).

Figure 31. Ring naming (A-E) in the triterpene pentacyclic core (left) and carbon atom numbering (1-30) in oleanane.

We started computations by carrying out a conformational search using molecular mechanics on all structures to provide reasonably good quality structures for initial DFT optimisations. The analysis of generated structures could assist in establishing best conformation for the hydroxyl group in ring A, which could possibly be applied to all structures in the pathway. We were also interested in the conformation of cation dammarenyl, which represents the initial structure in the pathway towards friedelin and has a flexible arm as a carbon chain attached to C18.

Conformational search in the gas phase was carried out with OPLS3¹³⁰ force-fields available in Macromodel¹³¹ using standard settings.

For most structures, hydroxyl group in ring A would adopt a conformation in equatorial position facing away from *gem*-dimethyl group present in C4. For the sake of consistency, we adopted this conformation for structures in all DFT optimisations unless any particular structural feature would dictate otherwise at the quantum mechanics level.

Conformational search on dammarenyl cation (**1**) had a 10 kcal/mol cutoff. Structures in this range were grouped and lowest energy conformation of each cluster had its geometry subjected to optimisation at the mPW1PW91/6-31+G(d,p) level of theory (Fig. 32). Conformation of lowest energy in cluster 1 (C1) was used as reference in the friedelin pathway and energy values of structures thereafter are reported in relation to it.

Figure 32. Dammarenyl cation (1) from best clusters (C1-C10). Relative energies (mPW1PW91/6-31+G(d,p) with ZPE corrections) are given in kcal/mol.

Before moving to the reaction results, some computational details are necessary. All calculations were carried out with the Gaussian 09 software package.⁷⁹ Based on literature information, we selected the following density functionals: B3LYP¹³²⁻¹³⁴, mPW1PW91¹³⁵ and BB1K¹³⁶ (for a discussion refer to methods, session 2.4). The double zeta basis set 6-31+G(d,p) was used throughout. An ultrafine integration grid of 99,590 points was requested as well as a tight convergence criteria for geometry optimisations. Frequency calculations were computed at 298.15 K to verify structures as either a minimum or transition state. Reported energies include corrections for zero-point vibration energy. In most cases, it was also possible to confirm transition state structures by running IRC (intrinsic reaction coordinate).

Results are presented in stages according to details in the mechanism and to make presentation more readable according to the discussions.

4.1 Formation of Rings D and E

The first sequence refers to a few transformations leading to the formation of 6-membered rings D and E starting from dammarenyl cation, which is the key precursor in the pathway to triterpenes in plants (Fig. 33).

Figure 33. Sequence of transformations showing formation of 6-6-6-6-6 pentacyclic intermediate cation germanicyl I (4) from dammarenyl (1).

The first step represents the expansion of 5-membered ring D in dammarenyl cation (1) to a 6-membered unit (2). We managed to locate a transition state (TS_{1-to-2}) for this cyclisation with all density functionals (B3LYP, mPW1PW91 and BB1K) selected for this study (Fig. 33). The barrier associated with the cyclization is around 10 kcal/mol depending on the density functional. Distances in the C–C bond under formation (ca. 1.9 Å) and breaking (ca. 2.0 Å) were similar making a triangle where the charge is stabilised in a two-electron, three-atom system through π electron donation from the double bond at one of the sides in the triangle ($d = 1.40$ Å, not shown).

Figure 34. Conversion of dammarenyl cation 1 into 4 showing cyclisations. Computed structures (selected distances in Å) and relative energies (in kcal/mol). Values in black refer to B3LYP, blue (mPW1PW91), and brown (BB1K). Energies in parenthesis are single points based on B3LYP geometry.

Once the cyclization takes place, the secondary carbocation baccharenyl (2) is generated. A structure for this cation was located in interaction with the double bond ($d = 2.45$ and $2.93 \text{ \AA}$) present in the alkyl chain arm attached to C18. Despite our efforts, we were unable to locate this structure with all density functionals, and only information related to B3LYP is presented. Other values of energy for this structure correspond to single points based on B3LYP geometry. That we failed to find this transition state with other density functionals is an indicative that the referred structure is not a true minimum in the reaction coordinate, which is not an uncommon situation in quantum chemical studies of terpene biosynthesis.^{63,71} Indeed, very few minima corresponding to secondary carbocation have been found in such studies. In the present case, the structure 2 is located exceedingly close (ca. $0.05 \text{ \AA}$) to that of the transition state corresponding to the next cyclisation ($\text{TS}_{2\rightarrow 3}$), and within 1 kcal in energy (Fig. 34).

A structure as the one located for baccharenyl cation represents a nonclassical carbocation.¹³⁷ By definition, a nonclassical carbocation cannot be represented by a single Lewis structure due to the delocalization of electrons. The stabilization effect of a double to carbocations have been demonstrated by comparing experimental and quantum chemically calculated ^{13}C NMR values for classical and nonclassical carbocations.¹³⁸

We were able to identify a second structure for baccharenyl cation as a classical structure (Fig. 35). However, as expected, higher in energy ($\Delta E = 3.5 \text{ kcal/mol}$) and, therefore, a non-productive structure for the mechanism under investigation.

Figure 35. Nonclassical (left) and classical (right) baccharenyl cations. Relative energies at the B3LYP/6-31+G(d,p)//B3LYP/6-31+G(d,p) level of theory (in kcal/mol).

Similarly, the transition state accounting for the conversion of baccharenyl to lupanyl ($\text{TS}_{2\rightarrow 3}$) could not be located with all density functionals but only B3LYP. That comes as no

surprise considering that various reports in the literature often locate structures with B3LYP and show single point energies with other functionals, especially mPW1PW91, which has given good performance on energetics for this class of molecules and related transformations.⁶³

Considering the energetics in the sequence 1-to-2-to-3, after the initial barrier of 10 kcal/mol associated with TS_{1-to-2} (Fig. 36), baccharenyl (2) and TS_{2-to-3} structures are just 1-2 kcal lower in energy revealing that once ring expansion takes place, cyclisation to produce the 5-membered should promptly occur leading to lupanyl cation without requiring much of enzyme intervention (if any). Preference for the sequence: cyclisation to produce the five-membered ring cation (3) in a Markovnikov-favoured closure followed by anti-Markovnikov ring expansion (barrier: 6–8 kcal/mol) to form the last six-membered ring in the pentacyclic triterpene core was demonstrated previously by means of computation^{55,97} and experiment.^{139,140}

Figure 36. Energy profile for the reaction pathway from dammarenyl (1) to lupanyl cation (4). Values in black refer to B3LYP, blue (mPW1PW91), and brown (BB1K). Energies in parenthesis are single points based on B3LYP geometry.

Interestingly, a structure for germanicyl I (4) cation, a secondary carbocation, was located as a candidate for minimum in the reaction coordinate. As no double bond is present in this structure to provide stabilization by π electron donation, lupanyl might be described as a classical carbocation. A closer inspection on this structure revealed that two single bonds are elongated – particularly C18–C19 (1.75–1.82 depending on the functionals), revealing strong hyperconjugative interaction in action to stabilise the secondary carbocation (Fig. 37).

Figure 37. Computed structure for germanicyl I cation (4). Elongated distances reveal strong hyperconjugative interaction to stabilize the secondary carbocation. Values in black correspond to B3LYP, blue (mPW1PW91) and brown (BB1K).

Assuming that the density functional BB1K shows best performance according to the literature.⁹⁸ For the sequence shown above (Fig. 36), B3LYP on its own displays a better performance with a smaller deviation from BB1K values than mPW1PW91. The latter is often used in conjugation with B3LYP geometries.

4.2 Formation of Oleanyl Cation

The next sequence refers to the formation of oleanyl cation following a sequence of two hydride transfers from the secondary carbocation germanicyl I (Fig. 38)

Figure 38. Sequence of transformations showing conversion of germanicyl I (4) to oleanyl (6) cation.

Although a short and apparently simple sequence of transformations, we have reasons for choosing to discuss it separately. In various texts in the literature, this sequence is not well represented with hydrides being transferred in a single step.^{39,122,141,142}

Numerous attempts to locate a transition state responsible for the hydride transfer in question systematically failed leading always to one of the minima – reactant or product. A major conformation change to **4** is required to allow hydride transfer from C18 to C19. At the initially identified conformation (Fig. 34), ring E in germanicyl I adopts the most stable chair conformation. However, the hydrogen in C18 as such is positioned perpendicular to the vacant p orbital in the carbocation in C19 and, therefore, not interacting for the hydride transfer to take place. We identified a transition state for the associated ring flipping in C19 (Fig. 39) corresponding to an energy barrier of 4 kcal/mol. Following a major change in conformation, hyperconjugation in the transition state structure is attenuated, which is partly responsible for the increase in energy. The distance C18–C19 decreases from 1.75–1.82 Å to 1.61 Å, and the C–C bond for one of the methyl groups attached to C20 bond takes over previous C20–C21 at similar bond length.

Figure 39. Energy barrier for conformational change in ring E leading to a productive conformation for hydride shift. Values in black correspond to B3LYP and blue (mPW1PW91).

A structure corresponding to a minimum of energy similar in geometry to the conformational transition state also could not be found. Therefore, we decided to take a different approach in the search for the transition state responsible for the hydride transfer from germanicyl I (**4**) to germanicyl II (**5**), TS_{4-to-5}. We aimed at stabilising the carbocation by introducing non-covalent interactions (e.g., cation- π , cation- π , C–H $\cdots\pi$) with an electron-

donating molecule (e.g. H₂O, NH₃, benzene) mimicking conditions in the active site that are likely to have a crucial role for a given transformation. In some cases, we used more than one of these donors including in mixed combinations. A few of our attempts are shown on Figure 40. We focused our efforts on B3LYP density functional for its better track record in locating transition structures in terpene biosynthesis studies including in this study.

Figure 40. Attempts (A1–A6) to locate a transition state structure (TS_{4-to-5}) using a non-covalent interaction.

Our initial effort with a single water molecule (A1) on the top face of the triterpene moiety was unsuccessful and we could not locate a transition state structure. An interesting result was obtained for one of our efforts when two water molecules were placed at the same face (A2). A transition state representing a concerted hydride shift and water attack to the carbocation was obtained (not shown).

Next, we tried two water molecules on opposite faces. A transition state structure for the desired transformation was once more obtained but at different distances ($d = 1.16$; 1.73 Å, Fig. 41) from those obtained in A1. Attempts making use of benzene as a π -donor positioned to interact with the hydrogen in migration (A4 and A5) were unsuccessful partly due to the competition for interaction with various hydrogens in the structure moving benzene away

from a more desired position. Inspired by literature information that tyrosine is an important residue in the formation of the last ring E,¹⁴¹ a phenol as a truncated model for tyrosine was placed in contact to the cation as a weaker, in comparison with water, electron donor for n-cation interactions. In this case, we also swapped water by an ammonia molecule at the top face of the triterpene (Fig. 41).

A3) TS_{4-to-5} • 2H₂O

A6) TS_{4-to-5} • NH₃ + PhOH

Figure 41. TS_{4-to-5} in non-covalent interactions. Selected distances are shown (Å): values in black correspond to B3LYP and blue (mPW1PW91). Dotted lines represent bonds forming and breaking in the transition state, in green cation-n and C-H...n interactions directly to the transition state, shaded lines represent the network of C-H...n interactions in the structure (d<3 Å).

Distances in comparison with water-bearing structure (A3) were exactly the same (d= 1.16; 1.73 Å, Fig. 41) despite different molecules in interaction with the transition state structures, which may indicate that the intrinsic reactivity of the carbocation governs the transformation. We computed the frequency on the triterpene geometry alone and a single imaginary frequency was verified (unoptimised structure). The energy on this structure (9.5

kcal/mol for A6) is similar to that observed for A3 (9.2 kcal/mol), which is close to the conformational transition state (TS_{conf.}) as shown on Fig. 42.

Encouraged by the result, we carried out a meticulous (calcall, stepsize=5) attempt in the quest for TS_{4-to-5} using only the triterpene geometry from optimised A6. Unfortunately, it again converged to the structure of the reactant, cation germanicyl I (4).

Previously, we happened to located the transition state of a hydride transfer (TS_{2-to-3}) in which a secondary carbocation was involved, but not the preceding structure as a true minimum. These structures were quite close in energy and geometry. Now, we find a conformational transition state but not the following intermediate and transition state. The latter was only found in the presence of additional molecules (donors) in interaction with the carbocation. Different molecules in interaction with the transition state in question produced equivalent structures and similar in energy from single points to the isolated triterpene structure (Fig. 42). We also computed the energy of germanicyl I cation (4) in interaction with ammonia and phenol (dubbed 4*). In the presence of the donors, conversion of germanicyl I to germanicyl II cation (4-to-5) occur with a minimum barrier (0.2–0.5 kcal/mol).

Figure 42. Energy profile for the reaction pathway from lupanyl (3) to germanicyl II cation (5). Values in black refer to B3LYP, blue (mPW1PW91), and brown (BB1K).

The donor molecules in interaction with TS_{4*-to-5} (or attempt A6) have a major role in holding the conformation through a network of C–H···n interactions; and stabilising the carbocation through a direct cation–n interaction allowing the hydride transfer to take place following a nearly barrierless transformation despite major conformational changes. There is

considerable tension around the reaction centre in the transition state as the atoms between two 6-membered rings in a decalin-like system are forced into a plane (Fig. 43). Indeed, we have not found the transition state in the reaction path (without non-covalent interactions) but, more importantly, we identified a critical step in the cascade of transformations, which very likely requires enzyme intervention.

Figure 43. Details of the interaction of $TS_{4\text{-to-}5}$ with ammonia and phenol, dubbed $TS_{4^*\text{-to-}5}$. (a) Atoms marked in orange are in the same plane. (b) A zoom into the reaction centre. Highlighted atoms are part of a dihedral (13-18-19-20) that are found close to a plane angle ($D=172^\circ$).

Considering the potential energy landscape where the conversion of germanicyl I (4) to germanicyl II (5) is present (Fig. 42), and the structural features related to the events in the course of the reaction, which transfers a hydride from a secondary to tertiary carbocation, relevant dynamical effects are operating. In the presence or not of non-covalent interactions, the energy barrier is not particularly high while major events occur along the reaction path – conformation change to a high-energy situation, but required for the reaction to occur followed by a very early transition state structure towards an exothermic process. With non-covalent interactions in place, the transformation is virtually barrierless suggesting that once ring expansion occurs in the previous step, hydride transfer should readily take place provided conformational changes occur. Similarly, this was also the case in an earlier step of a cyclisation to form ring E in which the transformation had a secondary carbocation as reactant and a small barrier associated to it. Moreover, there are kinetics issues. Secondary carbocations have very short half-lives in the 35–100 fs range.^{66,67} How short would be the half-life of germanicyl I in the reactive conformation? Considering the tension present in the rings, this carbocation may well be in lower end of the above stated range.

Among non-covalent interactions reported in similar studies, cation– π and C–H $\cdots\pi$ are by far most invoked. In the present case, C–H $\cdots\pi$ operating in a network proved vital in organising atoms into a locked conformation required for the reaction.

On this transformation, a final comment. We have also pursued alternative mechanisms for this step including proton shuttle in various possibilities, stepwise proton transfer, and combinations of more than one transformation into a single step to avoid the intermediacy of the secondary carbocation as it was the case of attempts on the direct conversion of lupanyl (3) to germanicyl II (5). We repeatedly faced issues while exploring any of these avenues, which prevented progression.

Conversion of germanicyl II (5) to oleanyl (6) cation was a less eventful transformation as both reactant and product are tertiary carbocations. An energy barrier of 10 kcal/mol has to be overcome and the reaction is slightly endergonic by 0.7 kcal/mol (Fig. 44). Once more, as mentioned previously, B3LYP has a better agreement with BB1K energy and geometry than mPW1PW91, which has a better reputation in quantum chemical studies of terpene biosynthesis.

Figure 44. Energy profile for the conversion of germanicyl II cation (6) to oleanyl cation (7). Values in black refer to B3LYP, blue (mPW1PW91), and brown (BB1K).

4.3 Formation of Friedelin Cation

The next set of transformations comprise the longest sequence with a few hydride and methyl shifts in various chemical environments but only tertiary carbocations involved (Fig. 45). Nonetheless, the energetics of these transformation may vary considerably.

Figure 45. Sequence of transformations showing conversion of oleanyl cation (6) to friedelin cation (13).

Figures 46 and 47 on next couple pages show all structures in the sequence with energies and selected distances in the transition states for all three density functionals.

Figure 46. Conversion of oleanyl cation (6) into campanulyl (10) showing hydride and methyl transfers. Computed structures (selected distances in Å) and relative energies (in kcal/mol). Values in black refer to B3LYP, blue (mPW1PW91), and brown (BB1K).

For the cascade of reactions under analysis, hydride transfers occur at lower energies with barriers around 3–4 kcal/mol while those associated to methyl transfer typically require 8–10 kcal/mol, except for the methyl transfer in glutinyl cation (11), which displays the barrier

of 15–21 kcal/mol, depending on the density functional. To explain this higher barrier, consideration of hydroxyl group preference for axial position is necessary.

Figure 47. Conversion of campanulyl (10) into friedelin cation (13) showing hydride and methyl transfers. Computed structures (selected distances in Å) and relative energies (in kcal/mol). Values in black refer to B3LYP, blue (mPW1PW91), and brown (BB1K).

As we are heading towards the end of the sequence, an energy profile for the full cascade is shown (Fig. 48) with all energies computed with the different density functionals applied in the study.

Figure 48. Energy profile for sequence of cyclisation/rearrangements from dammarenyl (1) to friedelin (13) cation. Relative energies (in kcal/mol) in black refer to B3LYP, blue (mPW1PW91), and brown (BB1K).

Despite conformational changes throughout the cascade, all structures except for one had the hydroxyl group in ring A at the equatorial position – glutinyl cation (11). Interestingly, the axial hydroxyl functionality in glutinyl (11) cation is close in space to the carbocation at 2.78 Å away in the BB1K/6-31+G(d,p) optimised structure. Moreover, we were unable to identify any particularly elongated C–C bond (> 1.6 Å) in hyperconjugation with the carbocation. A glutinyl cation with the hydroxyl in a nearly-equatorial was optimised and located 8.5 kcal/mol higher in energy than the originally found structure. In this case, the “equatorial” hydroxyl group was 3.6 Å away from the cation demonstrating that the group in axial contributes to the stabilisation of the cation. Therefore, it is likely that part of the energy in the high barrier towards TS_{11-to-12} originates from the conformational change, which disrupts the stabilising n – p electron-donating effect of the hydroxyl group in axial position.

In fact, the transformation glutinyl (11) to friedelanyl (12) cation represents the highest barrier in the cascade of reactions and a critical step to access pentacyclic triterpenes containing a ketone in ring A. Despite that hydroxyl in axial represents the most stable conformation, the transition state, however, has the group in equatorial. Interactions aiming at stabilising the equatorial conformation would lower this barrier facilitating this transformation.

The following and final hydride transfer results in the formation of the cetonic functionality and placing the charge at the oxygen, which would be later removed by a basic residue in the enzyme. This is a highly exergonic transformation by 17 kcal/mol, and the product has approximately the same energy content as the starting dammarenyl cation in this series.

Throughout the cascade, conformations changes occur to the carbocations as a result charge migration. The structure dynamically readapts to better accommodate the sp^2 carbon and stabilise the charge. It would adopt a more linear or an arched conformation; alter the conformation of a particular point like rings D–E, for instance, adopting a *cis*-decaline like conformation instead of more stable *trans* conformation, and so forth. These observations could be discussed on pages on end but without a purpose in mind, it is probably less productive. For the moment, we would be interested in major trends to the overall sequence and specific points would come in the continuation of our studies when a particular question would emerge from a real problem.

Bearing that in mind, we looked for properties in the structures along the sequence which could reveal any relevant information to the reactivity of the carbocations.

While checking dipole moments, we observed that values would vary largely across the series. Initially, we considered it could be an issue with the method. Dipole moments (μ) were then computed with the B2PLYP density functional following a recent recommendation from a dedicated study on the property assessing performance of a collection of 88 density functionals of various types. The benchmark investigation revealed better results for the family of double hybrids.¹⁴³

Dipole moment values in the series varied from 1.4 to 20.2 Debye, which is quite a large amplitude possibly indicating dependency on the property. By ordering dipole moments in the sequence of rearrangements, we noticed that as the charge travels along the carbon backbone, values of dipole moment progressively decrease reaching a minimum ($\mu=1.4$) at the centre of the structure in multiflorenyl cation (8). Moving towards the left end, it increases reaching a maximum with the charge placed in the ketone oxygen of friedelin (13) cation (Fig. 49).

Figure 49. Dipole moments (in Debye) for the series pentacyclic triterpene cations computed at the B2PLYP/6-31+G(d,p) level of theory.

A clear pattern is observed for the values of dipole moments in relation to the position of the positive charge in the pentacyclic scaffold. Accordingly, higher exposure of the charge at the edges of the structure produce considerable values ($\mu = 20.2$ D). In contrast, burying it deep inside the molecule leads to the considerably low value of 1.7 D. The difference should indicate enzyme adaptation by providing charge stabilisation accordingly, that is, amino acid residues in the active site matching the polarity of the triterpene cation throughout the rearrangements. An unfavourable interaction with a mismatching residue in polarity could bring the sequence to a halt. Electrostatics plays a key role in enzyme function¹⁴⁴ and the reported trend on dipole moments could have implications in guiding site directed mutations and development of devices capable of carrying out enzymatic transformations.¹⁴⁵

4.4 Density Functional Performance

We have taken note from earlier steps in the reaction cascade that B3LYP seemed to provide better agreement with BB1K energies than mPW1PW91 density functional. Having shown all transformations with their associated energetics, we revisit this issue by comparing all B3LYP and mPW1PW91 activation energies against BB1K values as reference (Table 6).

Table 6. Activation energies (kcal/mol) and deviations for B3LYP and mPW91PW91 by comparison with BB1K values. Mean absolute deviation (MAD) is also shown.

Reaction	B3LYP	mPW1PW91	BB1K
TS _{1-to-2}	10.3 (0.1)	9.3 (1.1)	10.4
TS _{3-to-4}	6.7 (1.3)	6.2 (1.8)	8.0
TS _{5-to-6}	10.2 (0.2)	8.7 (1.3)	10.0
TS _{6-to-7}	7.3 (1.2)	5.4 (0.7)	6.1
TS _{7-to-8}	9.2 (1.4)	7.8 (0)	7.8
TS _{8-to-9}	4.2 (1.1)	2.7 (0.6)	3.3
TS _{9-to-10}	4.9 (1.7)	3.4 (0.2)	3.2
TS _{10-to-11}	4.3 (1.0)	2.7 (0.6)	3.3
TS _{11-to-12}	20.6 (2.9)	14.8 (2.9)	17.7
TS _{12-to-13}	3.1 (0.2)	1.9 (1.0)	2.9
MAD:	1.1	1.0	

Only reactions that energy and geometry were computed with the same functional (Table 6). For comparison, we considered deviation for each transformation with B3LYP and mPW1PW91 in relation to BB1K (values in parenthesis, Table 6). The statistical parameter mean absolute deviation (MAD) revealed negligible difference: 1.0 kcal/mol for B3LYP and 1.1 kcal/mol for mPW1PW91. The deviation for the highest energy barrier is a representative case. Both functionals showed 2.9 kcal/mol as deviation. B3LYP, as expected, overestimated while mPW1PW91 by the same amount. Nonetheless, our earlier statements regarding initial steps in the sequence still hold.

5 CONCLUSION

A quantum chemical investigation into the cyclisation/rearrangement mechanism of the pentacyclic triterpene friedelin was presented in detail. Starting from the common triterpene precursor in plants, dammarenyl cation, all structures including transition states were located. The cascade of reactions contains a couple of secondary carbocations, which proved challenging structures, but also critical to an understanding of the early steps in the mechanism. Baccharenyl is one such structure, which was found as a nonclassical carbocation remarkably close to the transition state. The other, germanicyl I cation, had its transition state to the next step located only when stabilised by electron-donating molecules, that is, two water molecules or ammonia and phenol, which provided non-covalent interactions (cation- π and a network of C-H $\cdots\pi$) that also acts by holding the unfavourable conformation required for a hydrogen transfer to happen. Such situation should hint on the importance of interactions in the active site for this step. From our attempts, C-H $\cdots\pi$ was most successful. In the context of terpene biosynthesis, the observation is novel and should have further implications to other related mechanisms, and possibly also an addition to the overall understanding about the interplay between substrate and enzyme. The transformation of highest barrier in the sequence (11-to-12) represents another site of potential enzyme intervention and, therefore, manipulation for reactivity modulation. The cation and associate transition state structure have a hydroxyl group at axial and equatorial position, respectively. Strategies aiming at stabilising any such structures would facilitate (or not) access to the cetonic functionality in ring A.

For the general bystander, the sequence of reactions in the mechanism towards friedelin would possibly seem most unappealing – there is no fancy chemistry going on. As mentioned earlier, the devil is in the detail – and to that, it should be added: so does beauty. Going deep into the fine intricacies in the mechanism reveal information which could be of practical implication (one wishes): to improve a given process or even change the course towards another direction and product.

Regarding method, no relevant different was observed between (in)famous B3LYP and mPW1PW91, which is more recommended for energies in terpenes when compared to the generally accepted reference functional BB1K.

On the whole, the goal of improving knowledge on the mechanism for this sequence was achieved to an extent and largely when compared to previous contribution by Corey and co-workers. Nonetheless, it is a model – a proposal among possible solutions to the problem; not quite the truth. It will be interesting to see anything for any use out of it.

6 REFERENCES

1. Stone, T.; Darlington, G. *Pills, Potions, and Poisons: How Medicines and Other Drugs Work*; Oxford University Press: Oxford, 2000; 476 p.
2. Buckingham, J.; Cooper, C. M.; Purchase, R. *Natural Products Desk Reference*; CRC Press, Taylor & Francis: Boca Raton, 2016; p. 235
3. Sorensen, E. J.; Nicolaou, K. C. *Classics in Total Synthesis*; VCH: New York, 1996. 821 p.
4. *Natural Product Chemistry for Drug Discovery*; Buss, A. D.; Butler, M. S. Eds.; RSC: Cambridge, 2010, 440 p.
5. Newman, D. J.; Cragg, G. M. Natural products as sources of new drugs from 1981 to 2014. *J. Nat. Prod.* **2016**, 79(3), 629–661.
6. Wu, M.; Law, B.; Wilkinson, B. Micklefield. Bioengineering natural product biosynthesis pathways for therapeutic applications. *Curr. Opin. Biotech.*, **2012**, 23, 931–940.
7. Nicolaou, K. C.; Montagnon, T. *Molecules That Changed the World: A Brief History of the Art and Science of Synthesis and Its Impact on Society*; Wiley-VCH: Weinheim, 2008.
8. Bobrovskiy, I.; Hope, J. M.; Ivantsov, A.; Nettersheim, B. J.; Hallmann, C.; Brocks, J. J. Ancient steroids establish the Ediacaran fossil *Dickinsonia* as one of the earliest animals. *Science*, **2018**, 361, 1246–1249.
9. Otto, A.; White, J. D.; Simoneti, B. R. Natural product terpenoids in Eocene and Miocene conifer fossils. *Science*, **2002**, 297, 1543–1545.
10. Love, G. D.; Grosjean, E.; Stalvies, C.; Fike, D. A.; Grotzinger, J. P.; Bradley, A. S.; Kelly, A. E. et al. Fossil steroids record the appearance of *Demospongiae* during the Cryogenian period. *Nature*, **2009**, 457, 718–721.
11. Matthaus, B.; Ozcan, M. M.; Juhaimi, F. Oil content, fatty acid composition and tocopherol contents of turpentine (*Pistachia terebinthus* L.). *Z. Arznei-Gewurzpfla.* **2015**, 12(1), 136–140.
12. Artem, R.; Salakhutdinov, N. F. Chemical composition of *Pinus sibirica* (Pinaceae). *Chem. Biodivers.* **2015**, 12(1), 1–53.
13. Perkin, W. H. *J. Chem. Soc.* **1904**, 85, 654–671.
14. Thomas, A. F. In: *The total synthesis of natural products*, ApSimons, J., Ed., John Wiley & Sons: New York, 1973, Vol. 2, p. 149–154.
15. Kuroda, Y.; Nicacio, K. J.; Silva-Jr., I. A.; Leger, P. R.; Chang, S.; Gubiani, J. R.; Deflon, V. M.; Nagashina, N. et al. *Nat. Chem.*, **2018**, 10, 938–945.

16. Breitmaier, E. Terpenes. Flavors, fragrances, pharmaca, pheromones; Wiley-VCH: Weinheim, 2006, 214 p.
17. Lin, W. H.; Fang, J. M.; Cheng, Y. S. Uncommon diterpenes with the skeleton of 6-5-6-fused-rings from *Taiwania-cryptomerioides*. *Phytochem.*, **1995**, 40(3), 871–873.
18. Ignea, C.; Pontini, M.; Motawia, M. S.; Maffei, M. E.; Makris, A. M.; Kampranis, S. C. Synthesis of 11-carbon terpenoids in yeast using protein and metabolic engineering. *Nat. Chem. Bio.*, **2018**, 14(12), 1090–1098.
19. Xu, R.; Fazio, G. C.; Matsuda, S. P. T. *Phytochem.* **2004**, 65, 261–291.
20. Lü, J.-M.; Yao, Q.; Chen, C. Ginseng compounds: and update of their molecular mechanisms and medical applications. *Curr. Vasc. Pharmacol.* **2009**, 7(3), 293–302.
21. Rascón-Valenzuela, L. A.; Torres-Moreno, H.; Velázquez-Contreras, C.; Garibay-Escobar, R. E.; Robles-Zepeda, R. E. 2016. “Chapter 6. Triterpenoids: Synthesis, Uses in Cancer Treatment and other Biological Activities.” In *Advances in Medicine and Biology*. Berhardt, L. V., Ed., U.K.: Nova Science Publishers.
22. Gosh, S. Triterpenes structural diversification by plant cytochrome P450 enzymes. *Front. Plant Sci.*, **2017**, 8, article 1886.
23. Ruzicka, L. The isoprene rule and the biogenesis of terpenic compounds. *Experientia*, **1953**, 9(10), 357–367.
24. Eisenreich, W.; Bacher, A.; Arigoni, D.; Rohdich, F. Biosynthesis of isoprenoids via non-mevalonate pathway. *Cell. Mol. Life. Sci.*, **2004**, 61(12), 1401–1426.
25. Ogura, K.; Koyama, T. Enzymatic aspects of isoprenoid chain elongation. *Chem. Rev.*, **1998**, 98(4), 1263–1276.
26. Nagegowda, D. A. Plant volatile terpenoid metabolism: biosynthetic genes, transcriptional regulation and subcellular compartmentation. *FEBS Lett.*, **2010**, 584, 2695–2973.
27. Jenson, C.; Jorgensen, W. L. Computational investigations of carbenium ion reactions relevant to sterol biosynthesis. *Chem. Rev.*, **1997**, 119(44), 10846–10854.
28. Lesburg, C. A.; Caruthers, J. M.; Paschall, C. M.; Christianson, D. W. Managing and manipulating carbocations in biology: terpenoid cyclase structure and mechanism. *Curr. Opin. Struct. Bio.*, **1998**, 8, 695–703.
29. Christianson, D. W. Structural and chemical biology of terpenoid cyclases. *Chem. Rev.*, **2017**, 117, 11570–11648.
30. Croteau, R.; Cane, D. E. 1985. Monoterpene and sesquiterpene cyclases. In *Methods in enzymology – steroids and isoprenoids*. New York: Academic Press, 383–405.
31. Davis, E. M.; Croteau, R. Cyclization Enzymes in the biosynthesis of monoterpenes, sesquiterpenes, and diterpenes. *Top. Curr. Chem.*, **2000**, 209, 2812–2833.

32. Gambliel, H.; Croteau, R. Biosynthesis of (+/-)- α -pinene and (-)- β -pinene from geranyl pyrophosphate by a soluble enzyme-system from sage *Salvia officinalis*. *J. Biol. Chem.*, **1984**, 259, 740–748.
33. Kösal, M.; Jin, Y.; Coates, R. M.; Croteau, R.; Christianson, D. W. Taxadiene synthase structure and evolution of modular architecture in terpene biosynthesis. *Nature*, **2011**, 460, 116–120.
34. Lin, X; Hezari, M.; Koepp, A. E.; Floss, H. G.; Croteau, R. Mechanism of taxadiene synthase, a diterpene cyclase that catalyses the first step of taxol biosynthesis in Pacific yew. *Biochemistry*, **1996**, 35, 2968–2977.
35. Schrepfer, P.; Buettner, A.; Goerner, C.; Hertel, M.; van Rijn, J.; Wallrapp, F.; Eisenreich, W.; Sieber, V.; Kourist, R.; Brück, T. Identification of amino acid networks governing catalysis in the closed complex of class I terpene synthases. *Proc. Natl. Acad. Sci. U. S. A.*, **2016**, 113, E958–E967.
36. Williams, D. C.; Carroll, B. J.; Jin, Q.; Rithner, C. D.; Lenger, S. R.; Floss, H. G.; Coates, R. M.; Williams, R. M.; Croteau, R. Intermolecular proton transfer in the cyclization of geranylgeranyl diphosphate to the taxadiene precursor to taxol catalyzed by recombinant taxadiene synthase. *Chem. Biol.*, **2000**, 7, 969–977.
37. Connolly, J. D.; Hill, R. A. 1991. Dictionary of Terpenoids. Vol. 2: Di- and higher Terpenoids. London: Chapman & Hall.
38. Xu, R.; Fazio, G. C.; Matsuda, S. P. T. On the origins of triterpenoid skeletal diversity. *Phytochem.*, **2003**, 65, 261–291.
39. Thimmappa, R.; Geisler, K.; Louveau, T.; O'Maille, P.; Osbourn, A. Triterpene biosynthesis in plants. *Annu. Rev. Plant Biol.*, **2014**, 65, 225–257.
40. Thoma, R.; Schulz-Gasch, T.; D'Arcy, B.; Benz, J.; Aebi, J.; Dehmlow, H.; Hennig, M.; Stihle, M.; Ruf, A. Insight into Steroid Scaffold Formation from the Structure of Human Oxidosqualene Cyclase. *Nature* **2004**, 432, 118–122.
41. Nguyen, Q. N. N.; Tantillo, D. J. The many roles of quantum chemical predictions in synthetic organic chemistry. *Chem. Asian J.*, **2014**, 9, 674–680.
42. Nicolaou, K. C.; Frederick, M. O. On the structure of maitotoxin. *Angew. Chemie Int. Ed.*, **2007**, 46, 5278–5282.
43. Lodewky, M. W.; Siebert, M. R.; Tantillo, D. J. Computational prediction of ^1H and ^{13}C chemical shifts: a useful tool for natural product, mechanistic and synthetic organic chemistry. *Chem. Rev.*, **2012**, 112, 1839–1862.
44. Lodewky, M. W.; Soldi, C.; Jones, P. B.; Olmstead, M. M.; Rita, J.; Shaw, J. T.; Tantillo, D. J. The correct structure of aqualide – experimental validation of a theoretically-predicted structural revision. *J. Am. Chem. Soc.*, **2012**, 134, 18550–18553.

45. Howe, G. W.; van der Donk, W. ^{18}O kinetic isotope effect reveal an associative transition state for phosphite dehydrogenase catalyzed phosphoryl transfer. *J. Am. Chem. Soc.*, **2018**, *140*, 17820–17824.
46. Miton, C. M.; Jonas, S.; Fischer, G.; Duarte, F.; Mohamed, M. F.; van Loo, B.; Kintsjes, B.; Kamerlin, S. C. L.; Tokuriki, N.; Hyvönem, M.; Hollfelder, F. Evolutionary repurposing of a sulfatase: a new Michaelis complex leads to efficient transition state charge offset. *Proc. Natl. Acad. Sci. USA*, **2018**, *115*, E7293–E7302.
47. Wang, E. M. J.; Li, P.; Shaik, S.; Davies, G. J.; Walton, P. H.; Rovira, C. QM/MM studies into the H_2O_2 -dependent activity of lytic polysaccharide monooxygenase: evidence for the formation of a caged hydroxyl radical intermediate. *ACS Catal.*, **2018**, *8*, 1346–1351.
48. Iglesias-Fernández, J.; Hancock, S. M.; Lee, S. S.; Khan, M.; Kirkpatrick, J.; Oldham, N. J.; McAuley, K.; Fordham-Skelton, A.; Rovira, C.; Davis, B. G. A front-face 'S_Ni synthase' engineered from a retaining 'double-S_N2' hydrolase. *Nat. Chem. Bio.*, **2017**, *13*, 874–881.
49. Tantillo, D. J. Does nature know best? Pericyclic reactions of the Daphniphyllum alkaloid-forming cation cascade. *Org. Lett.*, **2016**, *18*, 4482–4484.
50. Hotta, K.; Chen, X.; Paton, R. S.; Minami, A.; Li, H.; Swaminathan, K.; Mathews, I. I.; Watanabe, K.; Oikawa, H.; Houk, K. N.; Kim, C. Y. Enzymatic catalysis of anti-Baldwin ring closure in polyether biosynthesis. *Nature*, **2012**, *483*, 355–359.
51. Biarnes, X.; Ardevol, A.; Planas, A.; Rovira, C.; Laio, A.; Parrinello, M. The conformational free energy landscape of α -D-glucopyranose: implications for substrate preactivation in α -glucosidase hydrolases. *J. Am. Chem. Soc.*, **2007**, *129*, 10686–10693.
52. de Visser, S. P.; Ogliaro, F.; Sharma, P. K.; Shaik, S. What factors affect the regioselectivity of oxidation by cytochrome P450? A DFT study of allylic hydroxylation and double bond epoxidation in a model reaction. *J. Am. Chem. Soc.*, **2002**, *124*, 11809–11826.
53. Tantillo, D. J. Importance of inherent substrate reactivity in enzyme-promoted carbocation cyclization/rearrangements. *Angew. Chem. Int. Ed.*, **2017**, *56*, 10040–10045.
54. Gleiter, M.; Mullen, K. Model calculations on the squalene cyclization. *Helv. Chim. Acta*, **1974**, *57*, 823–831.
55. Jenson, C.; Jorgensen, W. J. Computational investigations of carbenium ion reactions relevant to sterol biosynthesis. *J. Am. Chem. Soc.*, **1997**, *119*, 10846–10854.
56. kumar, S.; Kempinski, C.; Zhuang, X.; Norris, A.; Mafu, S.; Zi, J.; Bell, S. A.; Nybo, S. E.; Kinison, S. E.; Jiang, Z.; Goklany, S.; Linscott, K. B.; Chen, X.; Jia, Q.; Brown, S. D.; Bowman, J. L.; Babbitt, P. C.; Peters, R. J.; Chen, F.; Chappel, J. Molecular diversity of terpenes synthases in the livewort *Marchantia polymorpha*. *Plant Cell*, **2016**, *28*, 2632–2650.
57. Hong, Y. J.; Tantillo, D. J. Feasibility of intramolecular proton transfers in terpene biosynthesis – guiding principles. *J. Am. Chem. Soc.*, **2015**, *137*, 4134–4140.

58. Meguro, A.; Motoyoshi, Y.; Teramoto, K.; Ueda, S.; Totsuka, Y.; Ando, Y.; Tomita, T.; Kim, S. Y.; Kimura, T.; Igarashi, M.; Sawa, R.; Shinada, T.; Nishiyama, M.; Kuzuyama, T. An unusual terpene cyclization mechanism involving a carbon-carbon bond rearrangement. *Angew. Chemie Int. Ed.*, **2015**, *54*, 4353–4356.
59. Sato, H.; Teramoto, K.; Masumoto, Y.; Tezuka, N.; Sakai, K.; Ueda, S.; Totsuka, Y.; Shinada, T.; Nishiyama, M.; Wang, C.; Kuzuyama, T.; Uchiyama, M. “Cation-stitching cascade”: exquisite control of terpene cyclization in cyclooctatin biosynthesis. *Sci. Rep.*, **2016**, *5*, 18471.
60. Hong, Y. J.; Tantillo, D. J. The energetic viability of an unexpected skeletal rearrangement in cyclooctatin biosynthesis. *Org. Biomol. Chem.*, **2015**, *13*, 10273–10278.
61. Hong, Y. J.; Tantillo, D. J. Is a 1,4-alkyl shift involved in the biosynthesis of ledol and viridiflorol? *J. Org. Chem.*, **2017**, *82*, 3957–3959.
62. Hong, Y. J.; Tantillo, D. How many secondary carbocations are involved in the biosynthesis of avermitilol. *J. Org. Lett.*, **2011**, *13*, 1294–1297.
63. Tantillo, D. J. The carbocation continuum in terpene biosynthesis – where are the secondary carbocations? *Chem. Soc. Rev.*, **2010**, *39*, 2847–2854.
64. Tantillo, D. J. Recent excursion to the lands between concerted and stepwise: from natural products biosynthesis to reaction design. *J. Phys. Org. Chem.*, **2008**, *21*, 561–570.
65. Apeloig, Y.; Müller, T. 1997. Chapter 2. Theory and calculations. In *Dicoordinated carbocations*. Rapport, Z.; Stang, P. J., Ed., New York: Wiley.
66. Pemberton, R. P.; Tantillo, D. J. Lifetimes of carbocations encountered along reaction coordinates for terpene formation. *Chem. Sci.*, **2014**, 3301–3308.
67. Siebert, M. R.; Zhang, J.; Addepalli, S. V.; Tantillo, D. J.; Hase, W. L. The need for enzymatic steering in abietic acid biosynthesis: gas-phase chemical dynamics simulations of carbocation rearrangements on a bifurcation potential energy surface. *J. Am. Chem. Soc.*, **2011**, *133*, 8335–8343.
68. Hong, Y. J.; Tantillo, D. J. Tension between internal and external modes of stabilization in carbocations relevant to terpene biosynthesis: modulating minima depth via C–H $\cdots\pi$ interactions. *Org. Lett.*, **2015**, *17*, 5388–5391.
69. Tantillo, D. J.; Chen, J.; Houk, K. N. Theozymes and compuzymes: theoretical models for biological catalysis. *Curr. Op. Chem. Biol.*, **1998**, *2*, 743–750.
70. Gutta, P.; Tantillo, D. J. Theoretical studies on farnesyl cation cyclisation: pathways of pentalenene. *J. Am. Chem. Soc.*, **2006**, *128*, 6172–6179.
71. Tantillo, D. J. Biosynthesis via carbocations: theoretical studies on terpene formation. *Nat. Prod. Rep.*, **2011**, *28*, 1035–1053.

72. Hong, Y. J.; Tantillo, D. J. Which is more likely in trichodiene biosynthesis: hydride or proton transfer? *Org. Lett.*, **2006**, *8*, 4601–4604.
73. Hong, Y. J.; Tantillo, D. J. Branching out from the bisaboly cation. Unifying mechanistic pathways to barbatene, bazzanene, chamigrene, chamipinene, cumacrene, cuprenene, dunniene, isobazzanene, iso- γ -bisabolene, laurene, microbiotene, sesquithuejene, sesquisabinene, thujopsene, trichodiene, and widdradiene sesquiterpenes. *J. Am. Chem. Soc.*, **2014**, *136*, 2450–2463.
74. Gutta, P.; and Tantillo, D. J. A promiscuous proton in taxadiene biosynthesis? *Org. Lett.*, **2007**, *9*, 1069–1071.
75. Hong, Y. J.; and Tantillo, D. J. The Taxadiene-Forming Carbocation Cascade. *J. Am. Chem. Soc.*, **2011**, *133*, 18249–1825.
76. Freud, Y.; Ansbacher, T.; Major, D. T. Catalytic Control in the Facile Proton Transfer in Taxadiene Synthase. *ACS Catal.*, **2017**, *7*, 7653–7657.
77. Ansbacher, T.; Freud, Y.; Major, D. T. Slow-Starter Enzymes: Role of Active-Site Architecture in the Catalytic Control of the Biosynthesis of Taxadiene by Taxadiene Synthase. *Biochem.*, **2018**, *57*, 3773–3779.
78. Gao, J.; Ma, S.; Major, D. T.; Nam, K.; Pu, J.; Truhlar, D. G. Mechanism and free energies of enzymatic reactions. *Chem. Rev.*, **2006**, *106*, 3188–3209.
79. Gaussian website. <https://gaussian.com> (accessed Jan 15, 2019).
80. M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, B. Mennucci, G. A. Petersson, H. Nakatsuji, M. Caricato, X. Li, H. P. Hratchian, A. F. Izmaylov, J. Bloino, G. Zheng, J. L. Sonnenberg, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven, J. A. Montgomery, Jr., J. E. Peralta, F. Ogliaro, M. Bearpark, J. J. Heyd, E. Brothers, K. N. Kudin, V. N. Staroverov, R. Kobayashi, J. Normand, K. Raghavachari, A. Rendell, J. C. Burant, S. S. Iyengar, J. Tomasi, M. Cossi, N. Rega, J. M. Millam, M. Klene, J. E. Knox, J. B. Cross, V. Bakken, C. Adamo, J. Jaramillo, R. Gomperts, R. E. Stratmann, O. Yazyev, A. J. Austin, R. Cammi, C. Pomelli, J. W. Ochterski, R. L. Martin, K. Morokuma, V. G. Zakrzewski, G. A. Voth, P. Salvador, J. J. Dannenberg, S. Dapprich, A. D. Daniels, Ö. Farkas, J. B. Foresman, J. V. Ortiz, J. Cioslowski, D. J. Fox, Gaussian, Inc., Wallingford, CT, 2013, Revision D.01.
81. Meeting the man behind Moore's law. <http://news.bbc.co.uk/2/hi/technology/7080646.stm> (accessed Jan 16, 2019).
82. Heisenberg, W. über den anschaulichen inhalt der quantentheoretischen kinematik und mechanic. *Z. Phys. A*, **1927**, *43*, 172–198.
83. Hartree, D. R. The wave mechanics of an atom with a non-Coulomb central field. Part I. Theory and methods. *Math. Proc. Camb Phil. Soc.*, **1928**, *24*, 89–110.
84. Slater, J. C. The theory of complex spectra. *Phys. Rev.*, **1929**, *34*, 1293–1322.

85. Pauli, W. über den zusammenhang des abschlusses der elektronengruppen im atom mit der komplexstruktur der spektren. *Z. Phys. A*, **1925**, 31, 765–783.
86. Born, M.; Oppenheimer, R. Quantum theory of molecules. *Ann. Phys.*, **1927**, 84, 457–484.
87. Hohenberg, P.; Kohn, W. Inhomogeneous electron gas. *Phys. Rev. B*, **1964**, 136, 864–871.
88. Head-Gordon, M.; Mardirossian, N. Thirty years of density functional theory in computational chemistry: an overview and extensive assessment of 200 density functionals. *Mol. Phys.*, **2017**, 115, 2315–2372.
89. Medvedev, M. G.; Bushmarinov, I. S.; Sun, J.; Perdew, J. P.; Lyssenko, K. A. Density functional theory is straying from the path toward the exact functional. *Science*, **2017**, 355, 49–52.
90. Goerigk, L.; Hansen, A.; Bauer, C.; Ehrlich, S.; Naibi, A.; Grime, S. A look at the density functional theory zoo with the advanced GMTKN55 database for general main group thermochemistry, kinetics and noncovalent interactions. *Phys. Chem. Chem. Phys.*, **2017**, 68, 32184–32215.
91. Krüse, H.; Goerigk, L.; Grimme, S. Why the standard B3LYP/6-31G* model chemistry should not be used in DFT calculations of molecular thermochemistry: understanding and correcting the problem. *J. Org. Chem.*, **2012**, 77, 10824–10834.
92. Simón, L.; Goodman, J. M. How reliable are DFT transition structures? Comparison of GGA, hybrid-meta-GGA and meta-GGA functionals. *Org. Biomol. Chem.*, **2011**, 9, 689–700.
93. Perdew, J. P.; Schmidt, K. Jacob's ladder of density functional approximations for the exchange-correlation energy. *AIP Conference Proceedings*, **2001**, 577, 1–20.
94. Vrcek, I. V.; Vrcek, V.; Siehl, H. U. Quantum chemical study of degenerate hydride shifts in acyclic tertiary carbocations. *J. Phys. Chem. A*, **2002**, 106, 1604–1611.
95. Gutta, P.; Tantillo, D. J. Proton sandwiches: nonclassical carbocations with tetracoordinate protons. *Angew. Chem. Int. Ed.*, **2005**, 44, 2719–2723.
96. Bojin, M. D.; Tantillo, D. J. Nonclassical carbocations as C–H hydrogen donors. *J. Phys. Chem. A*, **2006**, 110, 4810–4816.
97. Siebert, M. R.; Tantillo, D. J. Brother versus brother: competitive stabilization of carbocationic centers by flanking cyclopropanes and π -systems. *J. Phys. Org. Chem.*, **2007**, 20, 384–394.
98. Weitman, M.; Major, D. T. Challenges posed to bornyl diphosphate synthase: diverging reaction mechanisms in monoterpenes. *J. Am. Chem. Soc.*, **2010**, 132, 6439–6360.
99. Matsuda, S. P. T.; Wilson, W. K.; Xiong, Q. Mechanistic insights into triterpene synthesis from quantum mechanical calculations. Detection of systematic errors in B3LYP cyclization energies. *Org. Biomol. Chem.*, **2006**, 4, 530–543.
100. Hare, S. R.; Tantillo, D. J. Dynamic behavior of rearranging carbocations – implications for terpene biosynthesis. *Beilstein J. Org. Chem.*, **2016**, 12, 377–390.

101. Carpenter, B. K. Energy disposition in reactive intermediates. *Chem. Rev.*, **2013**, *113*, 7265–7286.
102. Peng, Q.; Durte, F.; Paton, R. S. Computing organic chemistry stereoselectivity – from concepts to quantitative calculations and predictions. *Chem. Soc. Rev.*, **2016**, *45*, 6093–6107.
103. Eyring, H. The activated complex in chemical reactions. *J. Chem. Phys.*, **1935**, *3*, 107–115.
104. Evans, M. G.; Polanyi, M. Some applications of the transition state method to the calculation of reaction velocities, especially in solution. *Trans. Faraday Soc.*, **2006**, *31*, 875–893.
105. Fukui, K. The path of chemical reactions – The IRC approach. *Acc. Chem. Res.*, **1981**, *14*, 363–368.
106. Rehbein, J.; Carpenter, B. K. Do we fully understand what controls chemical selectivity? *Phys. Chem. Chem. Phys.*, **2011**, *13*, 20906–20922.
107. Ess, D. H.; Wheeler, S. E.; Iafe, R. G.; Xu, L.; Çelebi-Ölçüm, N.; Houk, K. N. Bifurcations on potential energy surface. *Angew. Chem. Int. Ed.*, **2010**, *47*, 7592–7601.
108. Hong, Y. J.; Tantillo, D. J. Biosynthetic consequences of multiple sequential post-transition-state bifurcations. *Nat. Chem.*, **2014**, *6*, 104–111.
109. Nyman, G. Computational methods of quantum reaction dynamics. *Int. J. Quantum Chem.*, **2014**, *114*, 1183–1198.
110. Hess, B. A. Computational studies on the cyclization of squalene to the steroids and hopenes. *Org. Biomol. Chem.*, **2017**, *15*, 2133–2145.
111. Purino, M.; Ardiles, A. E.; Callies, O.; Ignacio, A. J.; Bazzocchi, I. L. Montecrinanes A–C: triterpene with an unprecedented rearranged tetracyclic skeleton from *Celastrus vulcanicola*. Insights into triterpenoid biosynthesis based on DFT calculations. *Chem. Eur. J.*, **2016**, *100*, 12771–12800.
112. Chen, N.; Shenglong, W.; Smentek, L.; Hess, B. A.; Wu, R. B. Biosynthetic mechanism of lanosterol: cyclization. *Angew. Chem. Int. Ed.*, **2015**, *54*, 8693–8696.
113. Hess, B. A.; Smentek, L. The concerted nature of the cyclization of squalene oxide to the protosterol cation. *Angew. Chem.*, **2013**, *125*, 11235–11238.
114. Smentek, L.; Hess, B. A. Compelling computational evidence for the concerted cyclization of the ABC rings of hopene from protonated squalene. *J. Am. Chem. Soc.*, **2010**, *132*, 17111–17117.
115. Xiong, Q.; Rocco, F.; Wilson, W. K.; Xu, R.; Ceruti, M.; Matsuda, S. P. T. Structure and reactivity of the dammarenyl cation: configuration transmission in triterpene synthesis. *J. Org. Chem.*, **2005**, *70*, 5362–5375.
116. Hess, B. A.; Smentek, L. Concerted nature of AB ring formation in the enzymatic cyclozation of squalene to hopenes. *Org. Lett.*, **2004**, *6*, 1717–1720.
117. Hess, B. A.; Smentek, L. Density functional study on formation A and B rings in conversion of 2,3-oxidosqualene to lanosterol. *Mol. Phys.*, **2004**, *102*, 1201–1206.
118. Hess, B. A. Formation of the C ring in the lanosterol biosynthesis in the lanosterol biosynthesis from squalene. *Org. Lett.*, **2003**, *5*, 165–167.

119. Hess, B. A. Concomitant C-ring expansion and D-ring formation in lanosterol biosynthesis from squalene without violation of markovnikov's rule. *J. Am. Chem. Soc.*, **2002**, *124*, 10286–10287.
120. Gao, D.; Pan, Y.-K.; Byun, K.; Gao, J. Theoretical evidence for a concerted mechanism of the oxirane cleavage and A-ring formation in oxidosqualene cyclization. *J. Am. Chem. Soc.*, **1998**, *120*, 4045–4046.
121. Jenson, C.; Jorgensen, W. L. Computational investigations of carbenium ion reactions relevant to sterol biosynthesis. *J. Am. Chem. Soc.*, **1997**, *119*, 10849–10854.
122. Souza-Moreira, T.; Alves, T. B.; Pinheiro, K. A.; Felipe, L. G.; de Lima, G. M. A.; Watanabe, T. F.; Barbosa, C. C.; Santos, V. A. F. F. M.; Lopes, N. P.; Valentin, S. R.; Guido, R. V. C.; Furlan, M.; Zanelli, C. F. Friedelin synthase from *Maytenus ilicifolia*: leucine 482 plays an essential role in the production of the most rearranged pentacyclic triterpene. *Sci. Rep.*, **2016**, *6*, 36858.
123. Kuete, V.; Nguemaving, J. R.; Beng, V. P.; Ezebaze, A. G.; Etoa, F. X.; Meyer, M.; Bodo, B.; Nkengfack, A. E. Antimicrobial activity of the methanolic extracts and compounds from *Vismia laurentii* De Wild (Guttiferae). *J. Ethnopharmacol.* **2007**, *109*, 372–379.
124. Mann, A.; Ibrahim, K.; Oyewale, A. O.; Amupitan, J. O.; Fatope, M. O.; Okogun, J. I. *Am. J. Chem.* **2011**, *1*, 52–55.
125. Antonisamy, P.; Duraipandiyar, V.; Igancimuthu, S. Anti-inflammatory, analgesic and antipyretic effects of friedelin isolated from *Azima tetraacantha* Lam. in mouse and rat models. *J. Pharm. Pharmacol.*, **2011**, *63*, 1070–1077.
126. Antonisamy, P.; Duraipandiyar, V.; Aravinthan, A.; Al-Dhabi, N. A.; Choi, K. C.; Kim, J. H. Protective effects of friedelin isolated from *Azima tetraacantha* Lam. against ethanol-induced gastric ulcer in rats and possible underlying mechanisms. *Eur. J. Pharmacol.* **2015**, *750*, 167–175.
127. Sunil, C.; Duraipandiyar, V.; Ignacimuthu, S.; Al-Dhabi, N. A. *Food Chem.* Antioxidant, free radical scavenging and liver protective effects of friedelin isolated from *Azima tetraacantha* Lam. leaves. **2013**, *139*, 860–865.
128. Corey, E. J.; Ursprung, J. J. The structures of the triterpenes friedelin and cerin. *J. Am. Chem. Soc.*, **1956**, *78*, 5041–5051.
129. Laszlo, K.; Chein, R. J.; Corey, E. J. Conformational Energetics of Cationic Backbone Rearrangements in Triterpenoid Biosynthesis Provide an Insight into Enzymatic Control of Product. *J. Am. Chem. Soc.* **2008**, *140*, 9031–9036.
130. Harder, E.; Wolfgang, D.; Maple, J.; Wu, C.; Reboul, M.; Xiang, J. Y.; Wang, L.; Lupyan, D.; Dahlgren, M. K.; Knight, J. L.; Kaus, J. W.; Cerutti, D. S.; Krilov, G.; Jorgensen, W. L.; Abel, R.; Friesner, R. A. OPLS3: A Force Field Providing Broad Coverage of Drug-like Small Molecules and Proteins. *J. Chem. Theory Comp.*, **2015**, *12*, 281–296.
131. Schrödinger Release 2017-1 (2017) Small-Molecule Drug Discovery Suite, Schrödinger, LCC, New York.

132. Becke, A. D. Density-Functional Thermochemistry. 3. The Role of Exact Exchange. *J. Chem. Phys.* **1993**, *98*, 5648–5652.
133. Stephens, P. J.; Devlin, F. J.; Chabalowski, C. F.; Frisch, M. J. Ab Initio Calculation of Vibrational Absorption and Circular Dichroism Spectra Using Density Functional Force Fields. *J. Phys. Chem.* **1994**, *98*, 11623–11627.
134. Barone, V.; Orlandini, L.; Adamo, C. Proton Transfer in Model Hydrogen-Bonded Systems by a Density Functional Approach. *Chem. Phys. Lett.* **1994**, *231*, 295–300.
135. Adamo, C.; Barone, V. Toward Reliable Adiabatic Connection Models Free from Adjustable Parameters. *Chem. Phys. Lett.* **1997**, *274*, 242–250.
136. Zhao, Y.; Lynch, B. J.; Truhlar, D. G. Development and assessment of a new hybrid density functional model for thermochemical kinetics. *J. Phys. Chem. A*, **2004**, *108*, 2715–2719.
137. Aue, D. H. Carbocations. *Wiley Interdiscip. Rev. Comput. Mol. Sci.*, **2011**, *1*, 487–508.
138. Schleyer, P. R. The ab initio energy difference favoring nonclassical over the classical structure of the bicyclo[2.1.0]hexyl cation. Comparison of calculated (IGLO) and experimental ¹³C chemical shifts. *J. Am. Chem. Soc.*, **1988**, *110*, 300–301.
139. Corey, E. J.; Cheng, H.; Baker, C. H.; Matsuda, S. P. T.; Li, D.; Song, X. Methodology for the preparation of pure recombinant *S. cerevisiae* lanosterol synthase using a Baculovirus expression system. Evidence that oxirane cleavage and A-ring formation are concerted in the biosynthesis of lanosterol from 2,3-oxidosqualene. *J. Am. Chem. Soc.*, **1997**, *119*, 1277–1288.
140. Nishizawa, M.; Takenaka, H.; Hayashi, Y. Experimental evidence of the stepwise mechanism of a biomimetic olefin cyclization: trapping of cation intermediates. *J. Am. Chem. Soc.*, **1985**, *107*, 522–523.
141. Kushiro, T.; Shibuya, M.; Masuda, K.; Ebizuka, Y. Mutational studies on triterpene synthases: engineering lupeol synthase into α -amyrin synthase. *J. Am. Chem. Soc.*, **2000**, *122*, 6816–6824.
142. Dev, S., Ed. In: CRC handbook of terpenoids triterpenoids. Volume II: pentacyclic and hexacyclic triterpenoids. Taylor & Francis: Boca Raton, 1989, 624p.
143. Hait, D.; Head-Gordon, M. How accurate is density functional theory at predicting dipole moments? An assessment using a new database of 200 benchmark values. *J. Chem. Theory Comput.*, **2018**, *14*, 1969–1981.
144. Major, D. T. Electrostatic control of chemistry in terpene cyclases. *ACS Catal.*, **2017**, *7*, 5461–5465.