VESÍCULAS

ANA MARIA CARMONA-RIBEIRO

1992

**Texto apresentado ao Instituto de Química da Universidade de
São Paulo para o Concurso de Professor Livre-Docente junto
ao Departamento de Bioquímica**

ÍNDICE:

	Pág.
1. O princípio e o fim	1
2. O meio.....	7
3. Referências.....	17
4. Referência 6.....	21
5. Referência 7.....	30
6. Referência 8.....	42
7. Referência 9.....	49
8. Referência 10.....	58
9. Referência 11.....	65
10. Referência 12.....	71
11. Referência 13.....	80
12. Referência 14.....	87
13. Referência 15.....	94
14. Referência 16.....	101
15. Referência 17.....	132
16. Referência 18.....	164
17. Referência 19.....	181
18. Referência 28.....	199

VESÍCULAS

1. O princípio e o fim.

Em 1980, iniciei meu doutoramento no laboratório do prof. Chaimovich.

A partir do material bibliográfico disponível na época o qual incluía uma abrangente revisão no *Accounts of Chemical Research* por Janos Fendler (1) e de uma rápida conscientização do que seria factível com o material e equipamentos existentes no laboratório, elaborei meu primeiro projeto de pesquisa. Esse projeto se chamou: "Transporte de íons e substâncias neutras através de bicamadas de cloreto de dioctadecildimetilamonio (DODAC)". O título e o projeto agradaram ao Hernan que queria um estudo do transporte de protons através dessas bicamadas (o que eu jamais cheguei a fazer a contento) para começar a avaliar reatividade química em interfaces de vesículas.

Kunitake e colaboradores haviam descrito a formação de lipossomos multilamelares de DODAC por sonicação em banho (2).

Chaimovich e colaboradores haviam preparado pela primeira vez bicamadas de dihexadecilfosfato (DHP) por sonicação com sonda (3).

Fendler e colaboradores haviam descrito de forma aparentemente definitiva e completa a formação e caracterização de vesículas unilamelares pequenas de DODAC também por sonicação com sonda (4).

Os seis primeiros meses de meu doutoramento foram penosos tentando inútilmente incorporar porcentagens significativas de sacarose radioativa em vesículas de DODAC obtidas por sonicação para iniciar os experimentos de "leak-in" ou "leak-out" de substratos radioativos.

O primeiro relatório FAPESP/doutoramento fechou com a impossibilidade de se estudar analiticamente permeação de substâncias através de bicamadas sintéticas usando vesículas tão pequenas quanto as sonicadas de DODAC. Por outro lado, algumas tentativas de reproduzir os dados de Fendler (4) nesse mesmo sistema haviam falhado.

Não conseguira obter resposta osmótica em vesículas sonicadas de DODAC (4) quando usava um soluto não- iônico como a sacarose para gerar gradiente osmótico. Ainda, a reportada estabilidade em presença de sal monovalente (4) era claramente inexistente. A turbidez aumentava a olhos vistos por adição de 10 mM de NaCl à preparação de vesículas sonicadas. Assim sendo decidi consultar o recheado arquivo de separatas sobre lipossomos de fosfolípidos que existia no laboratório prestando especial atenção ao assunto

"Preparation of large phospholipid vesicles".

Achei o "report" do David Deamer & A D Bangham sobre preparação de vesículas grandes de fosfolípides por vaporização de uma solução etérea de fosfolípide em solução aquosa (qual não foi minha emoção ao conhecer pessoalmente o David em Canberra anos depois!) (5). Como o DODAC se mostrou insolúvel em éter, fiz uma solução clorofórmica de DODAC que injetei em solução aquosa mantida a uma temperatura superior à temperatura de ebulição do clorofórmio. Copiei quanto pude o arranjo e as condições experimentais para preparação de vesículas grandes de fosfolípides (poucos perdoam a rolha de cortiça que usei quando obtive as vesículas grandes de DODAC pela primeira vez (6)).

A partir da medida de incorporação de sacarose radioativa usando cromatografia em gel, a qual foi 100 vezes maior que a obtida para as sonicadas (6), sai pulando pelo corredor com uma descoberta nas mãos. De fato, a obtenção das vesículas grandes de DODAC em 3 de fevereiro de 1981 definiu um sistema adequado para estudar permeação e incorporação de substratos em bicamadas artificiais fechadas de anfifílicos sintéticos que vinham sendo propostas como modelos de membrana (6, 7).

Ademais, a comparação das vesículas grandes obtidas por injeção com as pequenas obtidas por sonicação permitiu

definir diversas propriedades de sistemas vesiculares que são dependentes de tamanho como, por exemplo, a capacidade de responder a gradientes osmóticos. Contrariando o trabalho de Fendler (4), a não-responsividade de vesículas sonicadas (pequenas) de DODAC e de DHP a gradientes osmóticos foi descrita (6, 7).

Mais tarde, com estudos de efeitos de NaCl sôbre a estabilidade das vesículas de anfifílicos sintéticos, ficou clara a origem das supostas cinéticas de encolhimento de Fendler por adição de gradientes osmóticos de KCl (3).

As vesículas agregavam e/ou fundiam em presença de sal (8, 9). Assim, sensatamente, determinara o comportamento de osmômetro para as vesículas grandes usando um soluto neutro impermeante (sacarose) como gerador do gradiente osmótico (Fig. 3, referência 6) ao invés de estabelecer gradientes osmóticos usando sal (6, 7).

O novo sistema vem sendo caracterizado de forma bastante completa e com uma certa... paixão (!).

Depois de estudar qualitativa e quantitativamente estrutura, função e interações em vesículas de anfifílicos sintéticos sempre comparando pequenas com grandes e essas com sistemas fosfolipídicos análogos (10- 14) comecei a me preocupar com o destino e o uso de minhas diletas filhas. Queria que se tornassem úteis, ou pelo menos, mais úteis (15- 18). Essa

fase se iniciou em 1987 com um projeto reprovado pelo PADCT.

Esse mesmo projeto, mais tarde modestamente aprovado pelo CNPq (1988), quase ignorado pela FINEP (1988), completamente ignorado pela Fundação Banco do Brasil e vencedor dentre outros tantos analisados por assessores do BID- Recursos Humanos (1990) procura dar um sentido para a existência das vesículas de anfifílicos sintéticos (15- 17).

A idéia central é estudar interações entre as vesículas e superfícies biológicas, minerais ou poliméricas verificando até que ponto vesículas podem modificar propriedades superficiais.

Nessa linha, mostramos que vesículas de anfifílicos sintéticos interagem com microesferas de poliestireno de carga oposta recobrando-as com bicamadas (15). Esse último trabalho tem implicações importantes em colóides e superfícies (17) e em ciências biomédicas e biológicas (18). Pela primeira vez percebi as vesículas de anfifílicos sintéticos caminhando à frente das fosfolipídicas.

Interações entre membranas fosfolipídicas e superfícies é um campo totalmente aberto à investigação. A interação de vesículas pequenas de fosfolípides com microesferas de poliestireno (18) ocorre por um mecanismo completamente diferente daquele mostrado para as vesículas sintéticas

(15). Há predominância da atração hidrofóbica entre a superfície do poliestireno e as cadeias hidrocarbônicas dos fosfolípides sobre a atração eletrostática entre a membrana e o poliestireno de carga oposta (18). Fosfolípides recobrem o poliestireno formando primeiro uma monocamada onde as cabeças polares se voltam para a água e as cadeias hidrocarbônicas se voltam para a superfície da microesfera (18). Depois disso, um aumento da concentração de fosfolípide na mistura, gera recobrimento com uma ou duas bicamadas (18).

Sobre esses alicerces as possibilidades são múltiplas. Interações entre membranas e superfícies são relevantes em design de kits imunológicos, em reconhecimento celular, em tratamento de água poluída com microorganismos, em formulação de fármacos ou defensivos agrícolas, em estabilização ou floculação de dispersões aquosas e em todos os casos de modificação de superfícies por agentes interfaciais.

A par das aplicações, há ainda a possibilidade de novos "insights" quanto aos fatores que determinam essas interações e que se encontram pouco estudados (17).

De um ponto de vista mais físico-químico, a obtenção das partículas de poliestireno recobertas com bicamadas sintéticas (15) possibilita uma verificação sistemática de

teorias elaboradas para partículas coloidais carregadas idealmente esféricas e homodispersas (17).

2. O meio.

A idéia fundamental deste texto é oferecer a estrutura sobre a qual os trabalhos foram pensados e realizados.

De modo geral, eu me percebi repetindo um dado ciclo diversas vezes e isso foi totalmente pessoal. Foi um processo solitário de ter uma idéia central, escolher uma ou varias estratégias para desenvolver essa idéia, ler sobre correlatos significativos, aprender com as experiências na bancada (inclusive as negativas), ler de novo, remoer resultados (com as madrugadas e os despertares repentinos gerando sempre os melhores "insights"), escrever a primeira horrível versão, ver o trabalho voltar recusado por causa da forma (e uma vez, por críticas conceituais severas), reescrever incorporando críticas, reenviar e finalmente ter o trabalho aceito para publicação.

Esses ciclos se originaram em um esqueleto de idéias relativamente simples:

- 1) fazer vesículas grandes de anfifílicos sintéticos por analogia com métodos já descritos para fosfolípides;
- 2) caracterizar as vesículas de anfifílicos sintéticos em

comparação com as fosfolipídicas procurando semelhanças e diferenças;

3) entender a razão por trás das diferenças;

4) encontrar usos para as vesículas de anfifílicos sintéticos;

5) estender esses usos para as vesículas de fosfolípides.

Uma importante diferença entre vesículas de fosfolípides e vesículas de anfifílicos sintéticos é a estabilidade em presença de NaCl. As primeiras são estáveis enquanto que as sintéticas agregam ou fundem por adição de sal (6, 7).

Em 1985 propus um estudo dessa instabilidade usando medidas diretas de interações entre as bicamadas de anfifílicos sintéticos. Nesse mesmo ano, Marra & Israelachvili publicaram um deslumbrante trabalho no *Biochemistry* medindo diretamente interações entre bicamadas fosfolipídicas adsorvidas em superfícies de mica (20) com o "surface force apparatus" (21). Por outro lado, eu estivera acompanhando regularmente a produção científica não apenas de Israelachvili como também de outros cientistas eméritos do Department of Applied Mathematics/ Research School of Physical Sciences/ Australian National University em Canberra. Barry Ninham, Department Head na época, havia sido referee de meu terceiro trabalho tendo se identificado durante o processo de revisão.

Assim, quando escrevi para Israelachvili em 1985 propondo

medir diretamente interações entre bicamadas de anfifílicos sintéticos para descobrir porque bicamadas de DHP agregavam enquanto que bicamadas de DODAC fundiam por adição de sal, recebi 3 cartas de volta: uma do próprio Israelachvili dizendo que estava de partida para a Califórnia, outra do Ninham dizendo que me conhecia (imaginem!) e que poderia inclusive pagar minhas despesas de estadia e outra do Pashley dizendo que ele próprio estava trabalhando ao longo dessa linha medindo interações entre bicamadas de DODAC adsorvidas na mica (uma idéia que se vai- pensei então).

E lá fui eu via Tierra del Fuego e Pólo Sul alcançar a Austrália do outro lado !

Foram dois meses bem vividos cientificamente cruzando com diversas sumidades pelos corredores e vasculhando o fabuloso arquivo de publicações do departamento.

Pashley estava muito ocupado medindo a força que mantém unidas as duas monocamadas que constituem uma bicamada (22) e não tinha tempo para dedicar à outra metade de minha idéia (medir interações entre bicamadas de DHP adsorvidas na mica). Ele e seus colaboradores haviam enviado o trabalho com o DODAC para publicação (23).

Além do mais parecia difícil adsorver uma bicamada negativamente carregada como a de DHP em um substrato também

negativamente carregado como a mica. Por outro lado, fui informada que era impossível aprender a usar o surface force apparatus em apenas dois meses, o tempo de duração de minha visita ao departamento.

A situação estava nesse impasse até o dia do seminário quando contei minha estória para tantas sumidades quantas quisessem ouvir (e a sala estava cheia). Enfatizei a dicotomia FUSÃO/AGREGAÇÃO - quando? porque e então? através de medidas diretas de forças de interação. Na saída eu arranjava um colaborador: o recém-doutor Per Martin Claesson que se dispôs a fazer e de fato fez todas as medidas de forças de interação entre camadas de dihexadecilfosfato adsorvidas na mica (10).

A adsorção foi obtida graças a uma idéia simples: adsorver primeiro uma monocamada positiva de DODAC (com as cabeças voltadas para a mica e as cadeias hidrocarbônicas voltadas para o ar) e a seguir, uma monocamada de DHP usando a técnica de Langmuir-Blodgett (10). A mim coube o trabalho com as monocamadas sendo que descrevi a monocamada insolúvel de DHP na interface água/ar pela primeira vez (10).

Como o Per não gostava do Langmuir-Blodgett australiano que era "feito em casa", todos meus resultados de monocamada foram repetidos pela Kazue Kurihara (cientista japonesa que atualmente integra o grupo do prof. Kunitake e que só vim a

conhecer recentemente em Compiègne na França) em Estocolmo com um aparelho de Langmuir-Blodgett computadorizado (10).

Fiquei especialmente orgulhosa ao saber de Israelachvili (em 1990, no Guarujá) que uma das figuras desse trabalho seria mostrada e discutida na segunda edição de seu livro (página 401, Fig. 18.3, referência 28). Essa figura mostra as forças extraatrativas e extrarepulsivas que ocorrem entre camadas de DHP adsorvidas na mica (10). A pH 9.5, a adsorção de íon sódio hidratado nos grupamentos fosfato dissociados gera uma força extrarepulsiva de hidratação (10). A pH 5.5, foi medida uma força extraatrativa de curto alcance (distância de separação menor que $20 \text{ \AA}$) de natureza ainda pouco clara (talvez pontes de hidrogenio interbicamadas, talvez defeitos estruturais na membrana com exposição de regiões hidrofóbicas) (10).

Ainda é meu plano uma investigação sistemática da ocorrência de fusão a pH ácido em vesículas de DHP para poder distinguir entre as duas possibilidades.

A determinação dos perfis de força de interação em função da distância de separação para verificar se fusão ou agregação poderiam ser previstas a partir dos mesmos não se mostrou uma estratégia muito reveladora. Os usuários do surface force apparatus não mediram esses perfis em concentrações de sal onde os fenômenos ocorrem (19). Não há na literatura

perfis de força de interação em 50 mM de NaCl para bicamadas de DODAC e de DHP (19).

Um "rationale" para a ocorrência de fusão com DODAC e apenas agregação com DHP (9) derivou na verdade dos experimentos de efeito de sal em monocamadas de DODAC ou de DHP na interface água/ar (10). A área por monômero a dada pressão superficial aumenta com a concentração de NaCl para o DHP e diminui, para o DODAC. Assim, o tamanho do agregado (vesícula) tende a diminuir e a aumentar com a concentração de sal, respectivamente (10). Outras razões foram encontradas posteriormente com as determinações de temperaturas de transição de fase gel para a líquido-cristalina em vesículas de DHP (12, 13). O grau de empacotamento da bicamada de DHP é tão alto que fica difícil imaginar fusão ocorrendo nesse sistema (12, 13).

De volta ao Brasil em fins de 1986, cheia de idéias, energia e sonhos depois daquela injeção de 2 meses de Austrália estava sem lenço e sem documento, nada no bolso ou nas mãos. Assim sendo, pensei no factível a partir de quase nada (DHP emprestado, água, sal, espectrofotômetro emprestado e eu mesma...!).

Escrevi um projeto assentado na terra para estudar efeitos de pH sobre estrutura, função e interações em vesículas de dihexadecilfosfato e um projeto voando na estratosfera para

estudar mobilidades eletroforéticas de vesículas de anfifílicos sintéticos e adsorção de bicamadas sintéticas em partículas coloidais (sem aparelho para microeletroforese de partículas, sem partículas, sem light-scattering para determinar tamanho e sem DODAC).

1987 foi o ano dos projetos e de minha contratação pelo IQUSP. Com apoio da FAPESP e do CNPq comecei a estudar efeitos do pH sobre propriedades de membranas de DHP extrapolando o conhecimento das forças de interação intra- e intermonocamadas de DHP (10) para as vesículas em solução. Diversas propriedades vesiculares como tamanho, fluidez da bicamada, permeabilidade à água e estabilidade dependem criticamente da composição do meio de preparação (12- 14). A molécula de DHP é especialmente interessante por permitir variar a carga da membrana apenas variando o pH do meio de preparação. Assim, alguns efeitos de carga sobre as propriedades de membranas de DHP foram descritos (12- 14).

Em 1988, Ohara Augusto e eu devíamos preparar uma aula prática para os alunos da Medicina que tivesse a ver com membranas. Já há algum tempo, Ohara queria demonstrar para os alunos a importância de alguns efeitos deletérios do oxigênio como os que ocorrem em processos de lipoperoxidação (25). Gostei da idéia e propus que fizéssemos isso usando lipossomos de asolecitina já que esta é muito rica em fosfolipídios com cadeias hidrocarbônicas insaturadas (26)

e, por acaso eu sabia que o Joaquim Procopio do ICB havia recentemente publicado um trabalho com "black lipid membranes" feitas de asolecitina (27). Assim, fui ao ICB pedir um pouco de asolecitina emprestada.

Depois de todos os acertos experimentais para mostrar vazamento por causa da lipoperoxidação e de uma parceria que deu certo, Ohara e eu publicamos no Biochemical Education o experimento que vem sendo há alguns anos a aula prática de membranas para os estudantes da Medicina (28).

Para os tempos áridos, outra estratégia providencial, apesar de difícil, foi fazer um trabalho teórico (11). Desde 1985 a pedra fundamental da teoria DLVO (29) não saía de minha cabeça e eu já estivera quantificando a instabilidade das vesículas usando determinações turbidimétricas de velocidades iniciais de floculação em função da concentração de sal (8) como queriam o Derjaguin, o Landau, o Verwey e o Overbeek (30).

Pareceu-me que faltava pouco para poder comparar as estabilidades experimentais com as previstas teoricamente considerando apenas a força de repulsão eletrostática e a força de atração de van der Waals. Copiei as equações que descreviam essas duas forças para partículas coloidais, "chutei" valores de potencial de superfície, ajustei constantes de Hamaker variando valores dentro de uma ordem

de magnitude (constantes variáveis!), usei para as vesículas uma equação de Van der Waals que valia para partículas esféricas massivas (colóides), calculei os perfis de energia de interação em função da distância de separação, integrei de acordo com a equação de Fuchs e mandei para o J. Phys. Chem. ainda recomendando que o Overbeek fosse o revisor.

A resposta veio certa. Todos os erros foram criteriosamente apontados e eu levei mais 3 anos para achar as equações certas, introduzir alguns valores de potencial de superfície que haviam sido obtidos experimentalmente pelo Lukac (31), usar apenas um valor razoável de constante de Hamaker e enfim adaptar o modelo DLVO que fora feito para partículas coloidais para o cálculo de estabilidades teóricas das vesículas (11).

Mesmo assim, o trabalho ainda estava incompleto. A dependência do potencial eletrostático superficial das vesículas grandes com a concentração de sal monovalente ainda não fôra determinada experimentalmente na literatura e portanto, a energia de repulsão eletrostática não podia ser calculada para as vesículas (11, 17).

Com alguma luz surgindo no final do túnel o CNPq aprovou em 1988 o projeto alado que me concedeu o aparelho de mobilidade eletroforética em fins de 1989. Finalmente eu ia poder terminar as medidas de mobilidade que iniciara em 1986

na Austrália, quando ganhei do BID/Recursos Humanos uma viagem para a Inglaterra para poder aterrisar o projeto voador de 1987. Lá, em 1990, encontrei as partículas, o light-scattering e o DODAC que me faltavam. Também, usei e abusei do aparelho de mobilidade eletroforética de forma a resolver a questão dos potenciais de superfície para as vesículas grandes de anfifílicos sintéticos (16). Neste ano da graça de 1992, completei o trabalho teórico calculando o termo eletrostático da energia de interação a partir dos valores de potencial eletrostático obtidos por microeletroforese de vesículas (19). Resultou daí que a estabilidade teóricamente calculada é maior que a experimental, o que sugere a existência de interações extraatrativas entre as vesículas que não se enquadram no arcabouço teórico da DLVO (19). Como gostaria de ter um surface force apparatus para medir diretamente essa interação extraatrativa em 50 mM de NaCl !

3. Referências

- 1 Fendler, J H (1980) *Acc. Chem. Res.* 13 7.
- 2 Kunitake, T, Okahata, Y, Tamaki, K, Kumamaru, F & Takayanagi, M (1977) *Chem. Lett.* 387.
- 3 Mortara, R A, Quina, F H & Chaimovich, H (1978) *Biochem. Biophys. Res. Commun.* 81 1080.
- 4 Kano, K, Romero, A, Djermouni, B, Ache, H & Fendler, J H (1979) *J. Am. Chem. Soc.* 101 4030.
- 5 Deamer, D W & Bangham, A D (1976) *Biochim. Biophys. Acta* 443 629.
- 6 Carmona-Ribeiro, A M & Chaimovich, H (1983) *Biochim. Biophys. Acta* 733 172.
- 7 Carmona-Ribeiro, A M , Yoshida, L S, Sesso, A & Chaimovich, H (1984) *J. Colloid Interface Sci.* 100 433.
- 8 Carmona-Ribeiro, A M , Yoshida, L S & Chaimovich, H (1985) *J. Phys. Chem.* 89 2928.
- 9 Carmona-Ribeiro, A M & Chaimovich, H (1986) *Biophys.*

J. 50 621.

- 10 Claesson, P M, Carmona-Ribeiro, A M & Kurihara, K
(1989) J. Phys. Chem. 93 917.
- 11 Carmona-Ribeiro, A M (1989) J. Phys. Chem. 93 2630.
- 12 Carmona-Ribeiro, A M (1990) J. Colloid Interface
Sci. 139 343.
- 13 Carmona-Ribeiro, A M & Hix, S (1991) J. Phys. Chem.
95 1812.
- 14 Carmona-Ribeiro, A M , Castuma, C E , Sesso, A &
Schreier, S (1991) J. Phys. Chem. 95 5362.
- 15 Carmona-Ribeiro, A M & Midmore, B R (1992) Langmuir
8 (3), 801.
- 16 Carmona-Ribeiro, A M & Midmore, B R (1992) J. Phys. .
Chem. 96 (8), 3542.
- 17 Carmona-Ribeiro, A M (1992) Chem. Soc. Reviews 21
(3), 207.
- 18 Carmona-Ribeiro, A M & Herrington, T M J. Colloid
Interface Sci., no prelo .

- 19 Carmona-Ribeiro, A M J. *Phys. Chem.*, in press.
- 20 Marra, J & Israelachvili, J N (1985) *Biochemistry* 24 4608.
- 21 Israelachvili, J N & Adams, G E (1978) *J. Chem. Soc. Faraday Trans. I* 74, 975.
- 22 Pashley, R M, McGuiggan, P M, Ninham, B W and Evans, D F (1985). *Science* 229, 1088.
- 23 Pashley, R M, McGuiggan, P M, Ninham, B W, Brady, J & Evans, D F (1986) *J. Phys. Chem.* 90 1637.
- 24 Israelachvili, J N (1992) "Intermolecular and Surface Forces" 2nd Edition, Academic Press, London and New York.
- 25 Halliwell, B & Gutteridge, J M (1985) "Free Radicals in Biology and Medicine", Clarendon Press, Oxford.
- 26 Kagawa, Y & Racker, E (1966) *J. Biol. Chem.* 241 2467.
- 27 Fornés, J A & Procopio, J (1987) *J Colloid*

Interface Sci. 117 570.

- 28 Augusto, O & Carmona-Ribeiro, A M (1989)
Biochemical Education 17 (4) 209.
- 29 Reerink, H & Overbeek, J Th G (1954) Disc. Faraday
Soc. 18 74.
- 30 Verwey, E J W & Overbeek, J Th G (1948) "Theory of
Stability of Lyophobic Colloids", Elsevier,
Amsterdam.
- 31 Lukac, S. (1983) J. Phys. Chem. 87 5045.

Referência 6

BBA 71805

PREPARATION AND CHARACTERIZATION OF LARGE DIOCTADECYLDIMETHYLAMMONIUM CHLORIDE LIPOSOMES AND COMPARISON WITH SMALL SONICATED VESICLES

A.M. CARMONA RIBEIRO and H. CHAIMOVICH *

Departamento de Bioquímica, Instituto de Química, USP, C.P. 20.780, São Paulo, SP (Brazil)

(Received February 14th, 1983)

Key words: Liposome preparation; Dioctadecyldimethylammonium chloride; Internal volume; Osmotic behavior

Dioctadecyldimethylammonium chloride (DODAC) unilamellar liposomes with a mean external diameter of 0.5 μm and sharp gel-to-liquid-crystalline phase transition temperatures (T_c) were obtained by chloroform vaporization and compared with small sonicated DODAC vesicles. Sucrose, impermeant through large DODAC liposomes and sonicated vesicles, was used for internal volume determinations. The internal volumes for large DODAC liposomes and sonicated DODAC vesicles were 9.0 ± 1.3 and 0.13 ± 0.2 l/mol, respectively. Ideal osmometer behaviour, towards KCl (0–50 mM) and sucrose, was observed only for large DODAC liposomes. Sonicated DODAC vesicles were osmotically non-responsive towards sucrose and flocculated upon addition of KCl. At temperatures near the T_c , a steep increase in the initial shrinkage rate and a minimum for the total extent of shrinkage were observed for large DODAC liposomes. Large DODAC liposomes are proposed as an adequate synthetic membrane model.

Introduction

Much of our present knowledge of membrane properties has been obtained with models prepared with phospholipids [1]. A variety of bilayer structures, formed by dialkyldimethylammonium halides [2] and other synthetic amphiphiles [3–7], have recently been used to mimic membrane properties, since they furnish unique opportunities to investigate structure-function relationships.

Closed-vesicle systems constitute one of the most explored models of membranes. While several methods for the preparation of phospholipid vesicles [8–11] or liposomes [12,13] have been described, synthetic amphiphile vesicles have been

obtained mainly by sonication [7]. This latter method yields small unilamellar vesicles with a mean external diameter of approx. 300 Å [14,15] and a corresponding small internal aqueous compartment. Thus incorporation of water-soluble substrates and permeability studies in synthetic amphiphile systems have been restricted.

In this report we describe the preparation and characterization of large DODAC liposomes that: (a) possess a mean external diameter of 0.5 μm ; (b) entrap sizeable amounts of sucrose; (c) have a sharp gel to liquid-crystalline phase transition; and (d) exhibit ideal osmometer behaviour over a range of conditions. In addition, we compare the properties of sonicated DODAC vesicles with those of large DODAC liposomes and conclude that this latter system constitutes a model that is more adequate for the investigation of transport properties in membranes.

* To whom correspondence should be addressed.

Abbreviation: DODAC, dioctadecyldimethylammonium chloride.

Materials and Methods

The purification and analysis of dioctadecyltrimethylammonium chloride (DODAC) (obtained from Herga Indústrias Químicas do Brasil) have been described [16]. Sephadex G-25 (fine) was obtained from Sigma (St. Louis, MO) and [^{14}C]sucrose from Schwarz-Mann (Orangeburg, NY, spec. act. 480 mCi/mmol). All other reagents were analytical grade. Deionized water, doubly distilled in glass, was used throughout. DODAC concentration was determined by chloride microtitration [17].

Preparation of small DODAC vesicles. Small DODAC vesicles were prepared by sonication (10 min, 50°C) of 60 mg DODAC in 10 ml water using the microprobe of a Cell Disrupter Virsonic Model 150 operated at a nominal output of 90 W. The dispersion was centrifuged (50 min, $10\,000 \times g$, 22°C) in order to eliminate residual Ti.

Preparation of large DODAC liposomes. The vaporization method of Deamer and Bangham [10] was adapted for DODAC after experimenting with several solvent combinations, temperatures and DODAC concentrations. 1.0 ml DODAC (0.02 M) in CHCl_3 was injected (0.2 ml/min) into 4.0 ml of water maintained at $70.0 \pm 0.5^\circ\text{C}$. The experimental arrangement for injection was that described by Deamer and Bangham [10]. Aliquots (1.0 ml) of the DODAC dispersion were filtered through a Sephadex G-25 column (1.8×14.0 cm) previously saturated with DODAC. DODAC recoveries were higher than 90% in all cases. The CHCl_3 contents [18] of the chloride-containing fractions eluting at the void volume (V_0) were less than 0.5 mol% ($\text{CHCl}_3/\text{DODAC}$).

Electron microscopy. Aliquots of DODAC dispersed by chloroform vaporization and filtered on Sephadex G-25 were mixed with equal volumes of ammonium molybdate (1%). After drying, the sample was examined in a Zeiss EM-9 S2 Electron Microscope operated at 60 kV. Mean external diameter were calculated from histograms obtained from several electron micrographs.

Determination of apparent internal volume (V). Aliquots of DODAC dispersion, prepared in water containing [^{14}C]sucrose, were filtered through a Sephadex G-25 column. The [^{14}C]sucrose entrapped/free ratio (R) was expressed as $R =$

$\text{cpm}(V_0)/\text{cpm}(V_i)$ where $\text{cpm}(V_0)$ is the total radioactivity eluting in the void volume (V_0), and $\text{cpm}(V_i)$, the total radioactivity eluting in the internal volume (V_i). The apparent internal volume (V) was expressed as $V = R/[\text{DODAC}]$, where $[\text{DODAC}]$ is the concentration of the DODAC dispersion applied to the filtration column. The extent of sucrose adsorption to preformed DODAC vesicles or liposomes was determined as follows. Small DODAC vesicles (or large DODAC liposomes) were prepared in water as described. To these preparations was added [^{14}C]sucrose, and the dispersions were filtered through a Sephadex G-25 column. The extent of sucrose adsorption per mol DODAC was expressed as an apparent volume (V_a) (see above).

The true internal volume (V_{in}) was defined as

$$V_{in} = V - V_a$$

Determination of osmotic properties. DODAC dispersions were added to aqueous solutions of KCl or sucrose (or aqueous solutions of KCl or sucrose were added to DODAC dispersions) and the time-dependent scatter-derived absorbance changes were registered in a Cary-14 Spectrophotometer at either 400 or 280 nm. The time lag between mixing and the start of the register was usually less than 6 s. The initial shrinking rate ($v\%$) was defined [19] as:

$$v\% = \frac{1}{A_0} \cdot \frac{\Delta A}{\Delta t} \cdot 100 \quad (1)$$

where $\Delta A/\Delta t$ represents the initial slope of the curve absorbance vs. time and A_0 , the initial absorbance. The reciprocal of the total absorbance change ($1/\Delta A$) was taken to be:

$$\frac{1}{\Delta A} = \frac{1}{(A_f - A_0)} \quad (2)$$

where A_f is the final measured absorbance. The extent of shrinkage (S) was defined as:

$$S = (1/A_0 - 1/A_f) \cdot A_0 \cdot 100 \quad (3)$$

Mean values for these parameters were obtained from at least two independent experiments and errors are expressed as mean square deviations.

Phase transition determinations. The changes of absorbance as a function of temperature were obtained in a Cary-14 Spectrophotometer equipped with a thermostatically controlled cell compartment (Haake E-12 Circulating Bath). Heating or cooling rates were generally 2 K/min. Temperatures were measured inside the cuvette with a copper-constantin thermocouple connected to a millivoltmeter. The response of this measuring system is linear between 0 and 100°C with a relative error of less than 0.01%.

Results

Physical characterization

DODAC dispersions, obtained by chloroform vaporization, eluted at the void volume of a Sep-

hadex G-25 filtration column. The apparent absorptivity (400 nm) of the chloride-containing fractions was $300 \text{ M}^{-1} \cdot \text{cm}^{-1}$.

These latter fractions examined by electron microscopy with both negative and positive staining, exhibited closed vesicular structures (Fig. 1) with a mean external diameter of $0.51 \mu\text{m}$ (range $0.2\text{--}1.0 \mu\text{m}$, Fig. 1C). Thus, the chloroform vaporization-Sephadex filtration preparation procedure yielded large DODAC liposomes which will thereafter be referred to as such.

DODAC liposomes exhibited a gel-to-liquid-crystalline phase transition temperature range narrower than sonicated vesicles, no pretransition being observed in either system (Fig. 2). The mid-points of the phase transition (T_c) for DODAC liposomes were 38.6°C (heating) and 34.8°C (cool-

Fig. 1. Electron micrographs of large DODAC liposomes positively stained with 2% ammonium molybdate, $6630\times$ (A); $24225\times$ (B). The distribution of liposome sizes can be seen in (C).

Fig. 2. Absorbance of 4.96 mM DODAC as a function of temperature. Arrows indicate the direction of temperature variation. (A) Large DODAC liposomes. (B) Sonicated DODAC vesicles.

ing). For the sonicated DODAC vesicles, the t_c values were 37.5°C (heating) and 34.4°C (cooling).

Functional characterization

Sucrose impermeability, entrapment and adsorption. The apparent internal volume (V) of sonicated DODAC vesicles, estimated using

Fig. 3. Time-dependent absorbance changes after addition of: (A) 0.05 M KCl (—) or 0.10 M sucrose (---) to large DODAC liposomes and (B) 0.05 M KCl (—) or sucrose (---) to sonicated DODAC vesicles.

Fig. 4. Tests of osmometer behaviour towards KCl for large DODAC liposomes (O) and sonicated DODAC vesicles (●). The reciprocal of the total absorbance change ($1/\Delta A$) was plotted against the reciprocal of the concentration gradient of KCl ($1/\Delta C$) and the initial shrinking rate ($v\%$) against the concentration gradient of KCl (ΔC). These parameters were calculated from the registered time-dependent absorbance changes obtained as follows: (A) and (B) 0.1 ml sonicated DODAC (4.96 mM) vesicles or large DODAC (4.96 mM) liposomes (prepared in water) were added to KCl solution (0.9 ml); (C) and (D) 0.2 ml large DODAC liposomes (3 mM) (prepared in water) were added to 0.8 ml KCl solutions. The insert in (D) contains an amplified plot of $v\%$ against ΔC for the lowest range of [KCl].

[^{14}C]sucrose as a water-soluble marker, was 0.33 ± 0.20 l/mol ($n = 4$). Vesicles containing [^{14}C]sucrose did not leak this marker when incubated up to 24 h in water at room temperature. The extent of external sucrose adsorption (see

Fig. 5. Osmometer behaviour of large DODAC liposomes towards sucrose. Large DODAC liposomes (0.1 ml, 2.1 mM) were added to 0.9 ml sucrose solutions and time-dependent absorbance changes registered for calculations of $1/\Delta A$ (total absorbance change) or $v\%$ (initial shrinking rate).

Fig. 6. Effect of temperature on the total extent of shrinkage and on the initial shrinking rate for large DODAC liposomes. A suspension of large DODAC liposomes (0.2 ml, 4.96 mM) was added to a solution of KCl (0.8 ml, 30.4 mM).

Materials and Methods) corresponded to a volume (V_s) of 0.20 ± 0.08 l/mol ($n = 6$). This latter value did not change when the unlabelled vesicles were incubated with externally added marker up to 24 h.

For large DODAC liposomes, an apparent internal volume of 9.66 ± 1.30 l/mol ($n = 3$) was determined. Equilibrium dialysis and filtration on Sephadex G-25 columns demonstrated (not shown), that no leak-out of entrapped [^{14}C]sucrose or leak-in of added [^{14}C]sucrose occurred over a period of 48 h. The extent of marker adsorption was 0.64 ± 0.30 l/mol ($n = 14$).

Osmotic properties

The time-dependent absorbance changes of DODAC sonicated vesicles (Fig. 3A) or DODAC liposomes (Fig. 3B) were registered for various osmotic gradients generated with KCl or sucrose. When KCl (0–50 mM) was the impermeant solute, linear dependences of $v\%$ as a function of KCl concentration gradient or of $1/\Delta A$ on the reciprocal of the KCl concentration gradient were obtained only for the large DODAC liposomes (Fig. 4). No changes in absorbance with time were

detected for sonicated DODAC vesicles upon establishing osmotic gradients with sucrose. For DODAC liposomes, linear functions relating $v\%$ and ΔC or $1/\Delta A$ and $1/\Delta C$ were obtained with sucrose osmotic gradients (Fig. 5).

The effect of temperature on the osmometric behaviour of large DODAC liposomes was characterized by a steep increase in the initial shrinkage rate and a minimum for the total extent of shrinkage near the T_c (33–37°C) (Fig. 6).

Discussion

In this section, we show that DODAC liposomes, in contrast to small sonicated DODAC vesicles, are suitable as models for permeability and transport studies through bilayers composed of synthetic amphiphiles.

A fundamental property of a closed membrane model is to act as a barrier towards some substances entrapped in its internal aqueous compartment. Sucrose was shown here to be a suitable impermeant marker for internal volume determinations [20]. The apparent internal volume of DODAC liposomes (9.7 ± 1.3 l/mol), obtained from sucrose entrapment and adsorption data, was comparable to that of large unilamellar vesicles prepared by diethyl ether vaporization of phospholipids [10].

The internal volume of DODAC liposomes can also be calculated from the mean diameter measured by electron microscopy. The mean area per DODAC monomer in the bilayer is approx. $40 \text{ \AA}^2$ [21] and the number of monomers in the inner and outer monolayer of DODAC liposomes can be considered equal. Assuming that DODAC liposomes (0.5 μm mean external diameter) are unilamellar, one calculates an internal volume of 10 l/mol, a value that is in very good agreement with that obtained experimentally by measuring sucrose entrapment.

The absence of typical multilamellar structures in electron microscopy (see, for example, Ref. 2), the similarity of the preparation method with that of large unilamellar phospholipidic vesicles [10] and the close agreement between the calculated and measured internal volumes indicate strongly that DODAC liposomes are unilamellar.

An identical calculation can be made for

DODAC sonicated vesicles. Taking a mean radius of 150 Å [14], an area per monomer of 40 Å² [21] and a 2:1 ratio of monomers in the outer and inner monolayer, one calculates an internal volume of 0.15 l/mol for these vesicles. From our sucrose data, subtracting the volume corresponding to adsorption (V_a) from the apparent internal volume (V), one obtains an upper limit of 0.13 l/mol for the internal volume (V_{in}) of sonicated DODAC vesicles. This value is comparable to the calculated internal volume and identical to that measured by Papahadjopoulos and co-workers [20] for sonicated phosphatidylserine vesicles.

Sonicated DODAC vesicles are osmotically non-responsive towards sucrose concentration gradients. Osmotic inactivity of small phospholipid vesicles has been described [23] specially for vesicles of external diameters smaller than 400 Å [24]. Absence of osmotic response for small sonicated vesicles has been associated with a rigid-sphere behaviour derived from the high radius of curvature of these structures [24,25]. Osmotic non-response in small sonicated vesicles can also be rationalized in terms of the organization of water in the internal aqueous compartment. Taking 0.13 l/mol as the internal volume and a 2:1 ratio for the number of monomers in the outer and inner monolayer, a simple calculation shows that there are only approx. 22 water molecules per DODAC monomer in the internal aqueous compartment of sonicated DODAC vesicles. This latter value is comparable to the number of water molecules undergoing slow exchange with bulk water in phosphatidylserine (Na⁺) bilayers [26]. Thus, the osmotic non-responsiveness could be related not only to the rigid structure of the bilayer but also to the nature of the tightly bound water existing in the small compartment. In other restricted environments such as that of inverted micelles there is firm experimental evidence for motion-restricted water under a variety of conditions [27].

A large part of this work has dealt with permeability of DODAC liposomes and sonicated vesicles towards water. Osmotic properties of some phospholipidic systems have been characterized extensively [28,29] and the demonstration of ideal osmometer behaviour of phosphatidylcholine liposomes was based on the verification that this latter system follows Boyle-Van't Hoff's law for solutes

such as glucose, manitol and salts [28]. In the present work, we have taken advantage of the linear relationship between the final liposome volume after shrinkage and the reciprocal of the final absorbance [19,28,29] to obtain some of the osmometric parameters. Our criteria for osmometric behaviour was also Boyle-Van't Hoff's law, which predicts a linear relationship between $1/\Delta A$ and $1/\Delta C$ and also between $v\%$ and ΔC . DODAC liposomes display the expected linear relationships between 0 and 50 mM KCl (Fig. 6). Above this salt concentration, the liposome preparation was seen to flocculate readily. Correspondingly, the linearity of the osmometric parameters with the concentration, or the reciprocal of the concentration, is not maintained. For sonicated vesicles, ideal osmometric behaviour was not observed under any set of conditions. In fact, visible aggregation and/or flocculation were noticed at most of the concentration gradients used here. The published osmometric behaviour of sonicated DODAC dispersions [14,30] is probably due to aggregation phenomena. The fact that sucrose gradients do not elicit an osmotic response with sonicated vesicles (*vide supra*) further establishes the difficulty of expecting ideal osmometric behaviour in the sonicated DODAC system. The marked difference of behaviour towards salts between the DODAC liposomes and the sonicated vesicles may have several explanations. One should notice, however, that intervesicular interactions are more probable in sonicated vesicles since the total exposed area is larger than that in large liposomes. Moreover, for equal amphiphile concentrations, the encounter probability is larger for the smaller particles. In view of the lack of data concerning surface potential of these systems, it is premature to offer definitive rationalizations that might account for the observed difference.

The pronounced hysteresis obtained for the phase transition temperature curves (Fig. 3) was expected, since the molar volume of surfactant in the liquid crystalline phase is greater than that in the gel phase and a domain of liquid crystalline phase will be under compression, whereas a domain of gel will be under tension [31]. McDonald et al. [32] have indeed shown that hysteresis is more likely to occur for charged lipids and that aqueous dispersions of phosphatidylserine do not

have superimposable heating and cooling curves. It is also well established that if the sharpness of the transition increases, so does the co-operativity [31,33]. Accordingly, the phase transition of large DODAC liposomes was more cooperative than that of sonicated DODAC vesicles (Fig. 2).

Phospholipid liposomes are permeable to water even below the T_c , being osmotically-sensitive structures both above and below the T_c [31]. In the vicinity of the T_c , a markedly reduced change in absorbance after an osmotic shock may occur due to increased permeability to water or to solute [34–38]. Large DODAC liposomes, in particular, behaved near the T_c as dimyristoylphosphatidylcholine liposomes [19], attaining a minimum for their total extent of shrinking and abruptly increasing the initial shrinking rate (Fig. 6). Increased permeability near the T_c to water and KCl is probably due to the formation of statistical pores between the gel and liquid-crystalline domains.

In conclusion, the two most interesting features of this work are: (i) the introduction of a confident criteria for stability of synthetic amphiphile vesicles towards salt, i.e., the determination of salt concentration ranges in which osmometric behaviour occurs; and (ii) the perspective of employing methods available for phospholipids to obtain new surfactant systems in which functional properties of membranes can be systematically checked.

Acknowledgements

This work was supported in part by the Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP), The Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) and Financiadora de Estudos e Projetos (FINEP B-76-81-295-00-00). We gratefully acknowledge Dr. Antonio Sesso and Mr. H. Brandão for the electron micrographs.

References

- Gomperts, B.D. (1977) *The Plasma Membrane: Models for Structure and Function*, 1st Edn., Academic Press, London.
- Kunitake, T., Okahata, Y., Tamaki, K., Kumamaru, F. and Takayanagi, M. (1977) *Chem. Lett.*, 387–390
- Hargreaves, W.R., and Deamer, D.W. (1978) *Biochemistry* 17, 3759–3768
- Mortara, R.A., Quina, F.H. and Chaimovich, H. (1978) *Biochem. Biophys. Res. Commun.* 81, 1080–1086
- Czarniecki, M.F. and Breslow, R. (1979) *J. Am. Chem. Soc.* 101, 3675–3676
- Murakami, Y., Nakano, A. and Fukuya, K. (1980) *J. Am. Chem. Soc.* 102, 4253–4254
- Fendler, J.H. (1982) *Membrane Mimetic Chemistry: Characterization and Applications of Micelles, Microemulsions, Monolayers, Bilayers, Vesicles, Host-Guest Systems and Polyions*, 1st edn., Ch. 6, Wiley-Interscience, New York
- Huang, C. (1969) *Biochemistry* 8, 344–352
- Batzri, S. and Korn, F.D. (1973) *Biochim. Biophys. Acta* 298, 1015–1019
- Deamer, D. and Bangham, A.D. (1976) *Biochim. Biophys. Acta* 443, 629–634
- Barenholz, Y., Amselem, S. and Lichtenberg, D. (1979) *FEBS Lett.* 99, 210–214
- Bangham, A.D. and Horne, R.W. (1964) *J. Mol. Biol.* 8, 660–668
- Kinsky, S.C. (1974) *Methods Enzymol.* 32, 501–513
- Kano, K., Romero, A., Djermouni, B., Ache, H. and Fendler, J.H. (1979) *J. Am. Chem. Soc.* 101, 4030–4037
- Fendler, J.H. (1980) *Acc. Chem. Res.* 13, 7–13
- Cuccovia, I.M., Aleixo, R.M.V., Mortara, R.A., Filho, P.B. Bonilha, J.B.S., Quina, F.H. and Chaimovich, H. (1979) *Tetrahedron Lett.* 3065–3068
- Schales, O. and Schales, S.S. (1941) *J. Biol. Chem.* 140, 879–884
- Snell, F.D. and Snell, C.T. (1937) *Colorimetric Methods of Analysis*, 2nd Edn., Vol. 2, pp. 557–558, D. Van Nostrand Company, Inc., New York
- Blok, M.C., Van Deenen, L.L.M. and De Gier, J. (1976) *Biochim. Biophys. Acta* 433, 1–12
- Papahadjopoulos, D., Nir, S. and Ohki, S. (1972) *Biochim. Biophys. Acta* 266, 561–583
- Herrmann, U. and Fendler, J.H. (1979) *Chem. Phys. Lett.* 64, 270–274
- Schreier, S., Polnaszek, C.F. and Smith, I.C.P. (1978) *Biochim. Biophys. Acta* 515, 375–436
- Johnson, S.M. and Buttress, N. (1973) *Biochim. Biophys. Acta* 307, 20–26
- Lichtenberg, D., Freire, E., Schmidt, C.F., Barenholz, Y., Felgner, P.L. and Thompson, T.E. (1981) *Biochemistry* 20, 3462–3467
- Schullery, S.E., Schmidt, C.F., Felgner, P., Tillack, T.W. and Thompson, T.E. (1980) *Biochemistry* 19, 3919–3923
- Finer, E.G. and Darke, A. (1974) *Chem. Phys. Lipids* 12, 1–16
- El Seoud, O.A., Chinelatto, A.M. and Shimizu, M.R. (1982) *J. Colloid Interface Sci.* 88, 420–427
- Bangham, A.D., De Gier, J. and Greville, G.D. (1967) *Chem. Phys. Lipids* 1, 225–246
- Rendi, R. (1967) *Biochim. Biophys. Acta* 135, 333–346
- Tran, C.D., Klahn, P.L., Romero, A. and Fendler, J.H. (1978) *J. Am. Chem. Soc.* 100, 1622–1624
- Lee, A.G. (1977) *Biochim. Biophys. Acta* 472, 237–281
- McDonald, R.C., Simon, S.A. and Baer, E. (1976) *Biochemistry* 15, 885–891

- 33 Takemoto, H., Inoue, S., Yasunaga, T., Sukigara, M. and Toyoshima, Y. (1981) *J. Phys. Chem.* 85, 1032-1037
- 34 Haest, C.W.M., De Gier, J., Van Es, G.A., Verkleij, A.J. and Van Deenen, L.L.M. (1972) *Biochim. Biophys. Acta* 288, 43-53
- 35 Papahadjopoulos, D., Jacobson, K., Nir, S. and Isac, T. (1973) *Biochim. Biophys. Acta* 311, 330-348
- 36 Inoue, K. (1974) *Biochim. Biophys. Acta* 339, 390-402
- 37 Nichols, P. and Miller, N. (1974) *Biochim. Biophys. Acta* 356, 184-198
- 38 Blok, M.C., Van der Neut-Kok, E.C.M., Van Deenen, L.L.M. and De Gier, J. (1975) *Biochim. Biophys. Acta* 406, 187-196

Referência 7

Permeabilities and Stabilities of Large Dihexadecylphosphate and Dioctadecyldimethylammonium Chloride Vesicles

ANA MARIA CARMONA-RIBEIRO,* LUCIA SATIKO YOSHIDA,*
ANTONIO SESSO,[†] AND HERNAN CHAIMOVICH*¹

**Departamento de Bioquímica, Instituto de Química, USP, São Paulo, Brasil, and*

[†]*Departamento de Patologia, Faculdade de Medicina, USP, São Paulo, Brasil*

Received October 18, 1983; accepted January 9, 1984

Sodium dihexadecylphosphate (DHP) vesicles, with a mean external diameter of 0.27 μm , were obtained by chloroform vaporization followed by gel filtration. The relative volume of the internal aqueous compartment, estimated measuring entrapment of [^{14}C]sucrose, was 13 liters/mol. DHP and dioctadecyldimethyl ammonium chloride (DODAC) large vesicles were stable up to 20 days at room temperature. In contrast, small sonicated DHP and DODAC vesicles were stable for some hours after preparation. Both DHP and DODAC large vesicles were impermeable toward most solutes tested. Large DHP vesicles responded as ideal osmometers toward gradients of sucrose, NaCl, or NaOH. The behavior of large vesicles of synthetic amphiphiles and of phospholipids was found to be analogous in the gel state.

INTRODUCTION

Synthetic amphiphiles such as dioctadecyldimethylammonium chloride (DODAC) or dihexadecylphosphate (DHP) form smectic mesophases when dispersed in aqueous solutions (1–8). Sonication of DODAC or DHP dispersions produces closed unilamellar vesicles with average external diameters of 300 and 500 Å, respectively (3, 7). In spite of the extensive use of these vesicles in biomimetic chemistry (9), little is known about the permeability properties of these bilayers since permeability studies are analytically limited by the size of the internal aqueous compartment. Thus, for a vesicle of 150 Å in radius 10 molecules of the solute yield a concentration of 1 mM in the intravesicular compartment. Larger vesicles are required in order to entrap larger amounts of the starting aqueous phase rendering these systems more useful in biomimetic chemistry and facilitating trans-

port studies through bilayers of synthetic amphiphiles.

We have recently reported the preparation and characterization of large DODAC vesicles with a mean external diameter of 0.5 μm (10) obtained by a modification of the vaporization method used by Deamer and Bangham (11). In this report, we have used this method for the preparation of negatively charged unilamellar vesicles of DHP and extended our studies on transport properties of synthetic amphiphile bilayers. DHP vesicles, prepared by vaporization of chloroform solutions, exhibited a mean external diameter of 0.27 μm , showed ideal osmometer behavior, and were more stable than the corresponding sonicated vesicles. Relative permeability studies of large DODAC and DHP vesicles showed that they are highly impermeable toward a variety of solutes with the exception of water, ethylene glycol, and ammonium acetate. Similar behavior for synthetic amphiphilic vesicles and phospholipidic vesicles in the gel state was demonstrated.

¹ To whom correspondence should be addressed.

MATERIAL AND METHODS

Dihexadecylphosphoric acid and Sephadex were from Sigma (St. Louis, Mo), [^{14}C]sucrose from Schwarz-Mann (Orangeburg, N. Y., specific activity 480 mCi/mole), and Sepharose 4B from Pharmacia (Upsala, Sweden). Sodium dihexadecylphosphate (DHP) was obtained by titration of a methanol solution of dihexadecylphosphoric acid with sodium methoxide, evaporation of the solvent, and drying *in vacuo*. All other reagents were analytical grade. Deionized water, doubly distilled in glass, was used throughout. DHP concentration was measured determining total phosphate (12). Spectral data were obtained in a Cary-14 or Beckmann-M25 spectrophotometers.

Preparation of sonicated DHP vesicles. Sonicated DHP vesicles were obtained as described in (7).

Preparation of large DHP vesicles. The vaporization method of Deamer and Bangham (11) was adapted for DHP after experimenting several DHP concentrations. DHP (1.0 ml) (0.0012–0.004 *M*) in CHCl_3 (40°C) was injected (0.2 ml/min) into 4.0 ml of water thermostatted at $70.0 \pm 0.5^\circ\text{C}$. The experimental set-up for injection was that described by Deamer and Bangham (11). Aliquots of the DHP dispersion were always filtered through a Sephadex G-25 column (1.8 $\times$ 14.0 cm) previously saturated with DHP or through a Sepharose 4-B column (43.5 $\times$ 1 cm). DHP recoveries were higher than 90% in all cases. Preparation of sonicated DODAC vesicles or large DODAC vesicles was as described in (10).

Electron microscopy. Aliquots of DHP dispersed by chloroform vaporization before or after filtering on Sephadex G-25 were mixed with equal volumes of uranyl acetate (2%). After drying, the samples were examined in a Zeiss EM-3 52 electron microscope operated at 60 kV or in a Philips 301 electron microscope. Mean external diameters were calculated from histograms obtained from several electron micrographs.

Determination of internal volume (V_{in}). Aliquots of DHP dispersion prepared in water containing [^{14}C]sucrose were filtered through

a Sephadex G-25 column (1.8 $\times$ 14.0 cm) or through a Sepharose 4-B column (43.5 $\times$ 1 cm). The [^{14}C]sucrose entrapped/free ratio (*R*) was expressed as $R = \text{cpm}(V_0)/\text{cpm}(V_i)$, where $\text{cpm}(V_0)$ is the total radioactivity eluting in the void volume V_0 and $\text{cpm}(V_i)$, the total radioactivity eluting in the internal volume V_i . The internal volume (V_{in}) was expressed as $V_{in} = R/[\text{DHP}]$, where [DHP] is the concentration of the DHP dispersion applied to the filtration column.

Determination of osmometer properties. Large DHP vesicles, prepared in water and filtered through Sephadex G-25, were added to a cuvette containing aqueous solutions of sucrose, NaCl, or NaOH. Time-dependent scatter-derived absorbance changes were registered at 400 nm. The time lag between mixing and the start of the register was usually less than 6 sec. The initial shrinking rate ($v\%$) and the reciprocal of the total absorbance change ($1/\Delta A$) were calculated as in (10) from at least two experiments.

Determination of relative permeabilities. Aliquots of large DHP vesicles or large DODAC vesicles (10) were added to equal volumes of aqueous solution of the solute. Absorbance changes as a function of time were followed at 400 nm at room temperature (22°C). From the registered absorbance, the total extent of shrinkage was calculated as $1/(A_f - A_0)$ where A_f is the final absorbance and A_0 , the initial absorbance. This parameter is considered to be representative of the liposomal final volume after shrinkage (13). Refractive index for all the solutions tested were determined in a Abbe-Atago Refractometer Model 301 (Japan) at 25°C previously calibrated with distilled water.

Determination of stabilities. Scatter-derived absorbances as a function of the wavelength (λ) (14) were registered at different times after the vesicles preparation. For sonicated vesicles, λ varied between 400 and 220 nm and for injected vesicles, between 770 and 400 nm.

RESULTS

Electron microscopy of DHP dispersions obtained by chloroform vaporization showed

FIG. 1. Electron micrographs of large DHP liposomes positively stained with uranyl acetate (5040X) before (A) or after (C) elution through Sephadex G-25. The distributions of liposome sizes before or after gel filtration is shown in (B) and (D), respectively.

closed vesicular structures (Fig. 1A) with a mean external diameter of $0.36 \mu\text{m}$ (range $0.1\text{--}1.4 \mu\text{m}$) (Fig. 1B).

When DHP dispersions were filtered through Sephadex G-25 the phosphate-containing fractions eluted at the void volume. Electron microscopy of these fractions exhibited vesicles (Fig. 1C) with a mean external diameter of $0.27 \mu\text{m}$ (range $0.1\text{--}0.6 \mu\text{m}$) (Fig. 1D), the particles bigger than $0.6 \mu\text{m}$ being retained upon filtration.

DHP vesicles, prepared in solutions of [^{14}C]sucrose, incorporated this water-soluble marker in their internal aqueous compartment (Figs. 2A, C). Upon rechromatography of such labeled preparation the radioactivity was totally recovered in the vesicle peak (Figs. 2C, D). In contrast with the results obtained with DODAC (10), sucrose did not adsorb externally to vesicles prepared in the absence of this marker (Fig. 2B). Using this filtration procedure and [^{14}C]sucrose (10) the internal vol-

ume of DHP vesicles was found to be 13.6 ± 3.0 liters/mole ($n = 7$) (Fig. 2A).

The apparent spectra due to scattering of freshly prepared DODAC or DHP vesicles obtained by sonication changed as a function of time (Fig. 3). However, the shape of the curve relating absorbance (240 nm) with time for sonicated DODAC vesicles (Fig. 3B) is different from that of sonicated DHP vesicles (Fig. 3D).

For large DODAC or DHP vesicles prepared in water, the spectra remain practically the same during ca. 20 days. Taking the values of the absorbance at 460 nm and plotting against time one obtains curves such as those in Figs. 4A and B. Large DHP vesicles yielded linear dependences of absorbance as a function of the inverse square of the wavelength (Fig. 5B). For large DODAC vesicles, similar plots do not extrapolate to zero and are not linear (Fig. 5A).

The time-dependent scatter-derived absor-

FIG. 2. Gel filtration elution profiles of large DHP liposomes prepared in the presence (A and C) or in the absence (B) of [^{14}C]sucrose. In B, the radioactive marker was added to previously formed vesicles. In D, rechromatography of fraction 5 in C (0.4 ml), 2 hr after its elution. A, B, and D show elutions through a Sepharose 4-B column (43.5×1.0 cm) and C, elution through a Sephadex G-25 column (1.8×12.0 cm).

FIG. 3. Spectra of sonicated DODAC vesicles 9.5 mM (A) or of sonicated DHP vesicles 7.0 mM (C) as a function of time. The insets show the absorbance at 240 nm as a function of time for sonicated DODAC (B) or DHP vesicles (D). In (A), the spectra were obtained at 2 (a), 24 (b), 149 (c), 190 (d), and 360 (e) hr. In (B), they were obtained at 1.5 (a), 16.6 (b), 23 (c), and 42 (d) hr.

bance changes of large DHP vesicles were registered for various osmotic gradients generated with sucrose (Fig. 6A), NaOH (Fig. 6B), or NaCl (Fig. 6C). Linear dependences of $v\%$ as a function of concentration gradients or of

$1/\Delta A$ on the reciprocal of the NaCl or NaOH concentration gradient were obtained for restricted ranges of NaCl or NaOH concentrations (0–50 mM) or over all the range of sucrose concentrations (Figs. 6D, E, F, G, H, I).

The values for the total extent of shrinkage ($1/\Delta A$) after submitting large DHP or DODAC vesicles to the same osmotic gradient of various solutes are in Tables I and II.

DISCUSSION

Vesicles prepared with double chain amphiphiles such as DODAC or DHP provide an opportunity to address the question of the relationship(s) between the structure of the monomer and the architecture of the aggregates (15), offer an efficient media for remarkable catalysis of chemical reactions (9, 16) and a defined media for species separation in photochemically produced pairs (9), along with the possibility to quantify ion binding in membrane models, which have been thoroughly analyzed in micellar systems (17). Recently, we demonstrated that large unilamellar DODAC vesicles obtained by vaporization are more interesting than small unilamellar DO-

FIG. 4. Absorbances (460 nm) as a function of time for large DODAC (A) or DHP vesicles (B). In A, DODAC concentrations are 1.2 (○) and 6.3 (●) mM. In B, DHP concentrations are 1.1 (▲), 2.4 (○), and 3.9 (●), mM.

FIG. 5. Absorbance as a function of the reversed square of the wavelength for large DODAC (A) or DHP vesicles (B). In A, curves were obtained at 0 (a) and 240 (b) hr for 1.2 mM DODAC or at 1.2 (c) and 242 (d) hr for 6.3 mM DODAC. In B, curves were obtained at 0.8 (a) and 208 (b) hr for 1.1 mM DHP; at 1.3 (c) and 210 (d) hr for 2.4 mM DHP; at 2.3 (e) and 210 (f) hr for 3.9 mM DHP. The lines were extrapolated below 400 nm.

DAC vesicles as membrane models (10). Here, we show that large DHP vesicles may be obtained by the same general procedure with some modifications and extend the characterization of these large anionic (DHP) or cationic (DODAC) vesicles presenting data of permeabilities and stabilities. With large amphiphilic vesicles, in contrast with the small sonicated vesicles, quantification of substrate passive transport is possible.

The determination of the internal volume (V_{in}) of vesicles requires an impermeant, water-soluble solute, which does not adsorb to the bilayer. Sucrose was such a solute since it does not adsorb to preformed DHP vesicles (Fig. 2B) and once entrapped does not leak out (Fig. 2D). Furthermore, as expected for an impermeant solute, ideal osmometric behavior was obtained over a wide range of concentrations when DHP vesicles were shrunk in the presence of sucrose.

The average value of V_{in} (13.6 liters/mole) and the relatively small standard deviation (± 3 liters/mole, $n = 7$) demonstrated the reproducibility of our preparation procedure.

If the vesicle is unilamellar, and the bilayer thickness is negligible with respect to the radius, it can easily be shown that $V_{in} = (rNaA)/$

6, where r is the radius of the vesicle, N , Avogadro's number, and A , the area occupied by a monomer. Taking $r = 1.35 \times 10^{-5}$ cm (Fig. 1B) and $V_{in} = 13,600 \text{ cm}^3/\text{mole}$ (our mean measured value for the internal volume), the calculated area per monomer is $98 \text{ \AA}^2$, a value which is higher than expected.

Hermann and Fendler (18) obtained sonicated DHP vesicles with an hydrodynamic radius of $592 \text{ \AA}$ and an average molecular weight of 23.6×10^6 . From these data, an area per monomer of $58 \text{ \AA}^2$ for a prolate shape ($a = 2451 \text{ \AA}$, $b = 79 \text{ \AA}$) or a value of $105 \text{ \AA}^2$ for a spherical shape (hydrodynamic radius of $592 \text{ \AA}$) can be calculated. Since hydrodynamic values were considered, the true area per monomer must be smaller. Areas per monomer for noncompressed monolayers of phosphatidic acids are around $50 \text{ \AA}^2$ (19). Sonicated amphiphilic vesicles in the gel state may possess interdigitated hydrocarbonic chains (20), and if interdigitation occurs, the expected area per monomer could be higher than in a classical bilayer.

The overestimation of A can also arise from an underestimation in r (see equation above). Since r was obtained from electron micrographs of stained preparations the measured

FIG. 6. Osmometer behavior of large DHP vesicles toward sucrose, NaCl, and NaOH. Time-dependent absorbance changes were registered at 400 nm after addition of sucrose (A), NaOH (B), or NaCl (C) to large DHP vesicles. Final concentrations of vesicles and solute in the cuvette were, respectively, 0.116 mM P_i; 0.083 M sucrose (A); 0.07 mM P_i; 0.052 M NaOH (B); 0.092 mM P_i; 0.049 M NaCl (C). From the time-dependent absorbance changes the initial shrinking rate ($v\%$) or the reciprocal of the total absorbance ($1/\Delta A$) was calculated and plotted, respectively, against the solute concentration gradient (ΔC) or the reciprocal of the solute concentration gradient ($1/\Delta C$). The osmometer behavior towards sucrose is shown in (D) and (G); toward NaOH, in (E) and (H) and, toward NaCl, in (F) and (I). The insets in (D) or in (H) show the behavior of the parameters along extended solute concentration ranges.

values for r can correspond to that of osmotically shrunk vesicles. In fact, vesicular shrinkage occurs due to osmotic gradients generated by the added electronic stain (uranyl acetate) and can account in part for the overestimated value calculated for A .

Small sonicated phospholipidic vesicles spontaneously fuse to yield large particles below the T_c (21–24). It has been proposed on geometrical grounds that the molecular packing in small unilamellar vesicles is dictated by the surface curvature and is highly asymmetric with respect to the inner and outer halves of the bilayer (25, 26). The extreme surface cur-

vature, intrinsic to small unilamellar vesicles, induces a profound asymmetry in the packing arrangement of the hydrocarbon chains in the two leaflets of the bilayer and causes the two monolayers of the bilayer to exist in markedly different motional states. Recently, it was argued that the restricted range of radii of the minimum size vesicle, which can be produced from phospholipids with hydrocarbon chain lengths ranging from 12 to 18 carbon bonds, suggests that the limiting size is not determined by a geometrical constraint associated with the hydrocarbon chain length of the lipid as suggested by Israelachvili *et al.* (27) but it

TABLE I
Permeabilities of Large DODAC Vesicles
Relative to Sucrose^a

Solute	<i>n</i> ^b	<i>O</i> ^c (mOsm/kg)	1/Δ <i>A</i> ^d	<i>P</i> ^e
NH ₄ Ac	1.3331	97.0	71.1 ± 17.3	10.5
Na Ac	1.3335	77.5	12.7 ± 2.8	1.9
NaCl	1.3335	96.5	11.4 ± 0.2	1.7
KCl	1.3336	102.0	10.3 ± 3.8	1.5
Ethylene glycol	1.3336	104.0	34.4 ± 5.8	5.1
Urea	1.3339	108.0	11.4 ± 1.4	1.7
Glycerol	1.3343	111.4	13.8 ± 3.6	2.0
Erythritol	1.3344	105.0	10.2 ± 2.1	1.5
Dextrose	1.3355	103.5	8.4 ± 1.6	1.2
Sucrose	1.3375	92.5	6.8 ± 0.7	(1.0)

^a Large (0.5 ml) DODAC vesicles (5.61 mM) were added to 0.5 ml of the solution containing the solute to be tested.

^b Refractive index of the solution added to the vesicles.

^c Osmolality of the solution added to the vesicles.

^d Final volume of the vesicles after shrinkage represented by 1/Δ*A* ± standard deviation.

^e Permeability of the vesicles measured as 1/Δ*A*/1/Δ*A*_{sucrose}.

would be the result of the polar group interaction on the vesicle interior (28, 29).

For synthetic amphiphilic vesicles obtained by sonication, stability over months was suggested (3–6). In the present work, we show that DODAC and DHP sonicated vesicles are unstable (Fig. 3). In contrast, large DODAC or DHP vesicles remained unaltered over weeks (Fig. 4). Our turbidity measurements indicated that sonicated vesicles form larger structures spontaneously. It is not possible, however, in terms of these turbidimetric data to decide if the larger structures result from vesicle aggregation or from fusion. However, we must observe that different intervesicular interactions seem to account for the observed absorbance changes along time in Figs. 4B and D.

Figure 3B presents data for DODAC sonicated vesicles, which differ from that of DHP sonicated vesicles (Fig. 3D). This difference can arise from continuous aggregation for sonicated DODAC vesicles, and rapid fusion,

yielding larger stable vesicles, for sonicated DHP vesicles.

A good fitting of experimental data was obtained for large DHP vesicles (Fig. 5A) in accordance with the Jöbst approximation (30) for scattering by particles having dimensions comparable to those of visible light and spherical shape, the intensity of scattered light being proportional to the inverse square of the wavelength. This is a strong evidence that large DHP vesicles are perfect spheres and that the variation of refractive index for these vesicles with the vesicles concentration (*dn/dC*), does not depend on wavelength (31). In contrast, large DODAC vesicles exhibit *dn/dC* varying with the incident wavelength and may have a different shape.

In phospholipid systems, osmotic behavior was characterized for liposomes (13, 32, 33) and osmotic nonresponsiveness was reported for sonicated vesicles (34). Recently, we have characterized the osmometer behavior of large unilamellar DODAC vesicles in comparison with the osmotic insensitivity of small soni-

TABLE II
Permeabilities of Large DHP Vesicles
Relative to Sucrose^a

Solute	<i>n</i> ^b	<i>O</i> ^c (mOsm/kg)	1/Δ <i>A</i> ^d	<i>P</i> ^e
NH ₄ Ac	1.3331	100.5	83.3	2.3
Na Ac	1.3335	79.0	31.8 ± 0.1	0.9
NaCl	1.3334	81.0	44.6 ± 4.1	1.2
KCl	1.3335	84.0	43.6 ± 2.7	1.2
Ethylene glycol	1.3335	82.0	87.1 ± 5.3	2.4
Urea	1.3334	62.0	39.2 ± 1.1	1.1
Glycerol	1.3340	84.0	48.1 ± 6.5	1.3
Ascorbic Acid	1.3340	40.0	40.8 ± 1.2	1.1
Erythritol	1.3342	68.0	36.7 ± 4.7	1.0
Dextrose	1.3349	76.0	41.7 ± 2.5	1.2
Ammonium molybdate	1.3354	61.0	39.2 ± 1.1	1.1
Sucrose	1.3373	71.0	36.4 ± 0.9	(1.0)

^a Large (0.6 ml) DHP vesicles (0.18 mM) were added to 0.6 ml of the solution containing the solute to be tested.

^b Refractive index of the solution added to the vesicles.

^c Osmolality of the solution added to the vesicles.

^d Final volume of the vesicles after shrinkage presented by 1/Δ*A* ± standard deviation.

^e Permeability of the vesicles measured as 1/Δ*A*/1/Δ*A*_{sucrose}.

cated DODAC vesicles (10). In this latter work we proposed that the osmometric behavior of a vesicular system could be used as a criteria for the stability of the system toward salts. Using this criteria it can be concluded that DHP vesicles are stable toward NaOH and NaCl in the 0–50 mM range and toward sucrose over the concentration range used (Fig. 6). Ideal osmometer behavior suggests that large DHP vesicles are nonpermeant toward NaOH. Isotope-exchange studies on liposomes reveal that the replacement of zwitterionic phosphatidylcholine by negatively charged lipids decreases the anionic and increases the cationic permeability (35, 36). OH^-/H^+ were suggested to be permeant across phospholipidic vesicles in the liquid crystalline state (37–41), or across phosphatidic acid bilayers in the gel state (41) but nonpermeant across phospholipidic vesicles below the T_c (40, 42). Our result, demonstrating nonpermeability of NaOH through large DHP vesicles in the gel state, agree with those presented by Keana *et al.* (42) or Trandinh *et al.* (40) for systems below the T_c .

In the concentration region where DHP vesicles behave as ideal osmometers, the initial shrinking rates obtained with NaOH and NaCl are practically identical (Fig. 6). However, when the DHP system behaves as osmotically nonideal, that is in the region where the system aggregates and/or flocculates, the observed rates of absorbance change are greater for NaOH than for the NaCl. Since aggregation depends on vesicle-vesicle interactions, and the DHP vesicles are charged, the most efficient salt is that which minimizes the extent of the electrostatic repulsion. The effective dielectric constants of the NaOH solutions are smaller than those of NaCl of identical concentrations (43). Since the thickness of the diffuse double layer is proportional to the square root of the effective dielectric constant of the media (44, 45), it follows that this thickness is smaller for large DHP vesicles in NaOH. Consequently the surface potential profile will decay faster for vesicles in solutions of NaOH.

The final volume of the liposomal population after shrinking is proportional to the inverse of the final absorbance of the shrunk or swelled dispersion (13). We took advantage of this property in order to measure permeabilities of large DODAC liposomes or of large DHP vesicles relative to sucrose (Tables I and II). The greater the final volume of the vesicle after shrinkage due to a solute gradient, the higher the permeability of the solute. The relation between the final volume after shrinkage due to a solute gradient and the final volume after shrinkage due to an equivalent sucrose gradient represents a relative permeability value. Most solutes were nonpermeant, with the exception of ethylene glycol and ammonium acetate. Nonionics in liquid-crystalline phospholipid bilayers can permeate by dissolving and diffusing across by following the fluctuations (kinks) in molecular packing (46, 47). For charged species, this pathway is unfavorable, since it involves a displacement from a medium of high to a medium of low dielectric constant (48, 49). The diffusion of such permeants would then depend on the occurrence of statistical pores, whose interior is lined by the polar headgroups of the phospholipidic molecules. Such pores of short duration might result from thermal fluctuations which could be identified as a fast lateral movement of the phospholipids molecules on the surface of the membrane (50), and a slow flip-flop reorientation between the two sides of the bilayer (51, 52). Such effects would be inhibited by cholesterol which tends to make the membranes more rigid above the T_c and to produce a decrease in the permeability coefficients of phospholipid vesicles to Na^+ , K^+ , Cl^- , and glucose (53). Below the gel to liquid-crystalline phase transition temperature, the rate of water permeation through lipid bilayers is strongly reduced and only a limited number of solutes are able to permeate through those solid bilayers (33). For example, bilayers of 16:0/16:0 phosphatidylcholine at 30°C ($T_c = 41.5^\circ\text{C}$) are impermeable even for thiourea, being permeable for formamide, ethylene glycol, and 1,3-propanediol (54). These data

closely parallel our results in Tables I and II, which indicated that large DODAC or DHP vesicles in the gel state are impermeable even for urea and permeable only for water, ethylene glycol, and ammonium acetate. The observed permeability toward NH_4Ac is probably due to NH_3 and HAc permeation, since all the ionic solutes tested were impermeant (Tables I and II) and large phospholipidic vesicles are rather permeable toward NH_3 or HAc in the neutral form (55, 56).

Thus, close analogy of behavior was found for phospholipid and the correspondent synthetic amphiphile system in the gel state.

ACKNOWLEDGMENTS

This work was supported in part by the Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP), the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), and Financiadora de Estudos e Projetos (FINEP).

We gratefully acknowledge Dr. Iolanda Midea Cuccovia for her help and criticism of this manuscript and Mr. Helio Brandão for assistance with the electron micrographs.

REFERENCES

1. Tanaki, K., Kumamuru, F., and Takayanagi, T., *Chem. Lett.*, 387 (1977).
2. Kunitake, T., and Okahata, Y., *J. Amer. Chem. Soc.* **99**, 3860 (1977).
3. Deguchi, K., and Mino, J., *J. Colloid Interface Sci.* **65**, 155 (1978).
4. Tran, C. D., Klahn, P. L., Romero, A., and Fendler, J. H., *J. Amer. Chem. Soc.* **100**, 1672 (1978).
5. Kano, K., Romero, A., Djermouni, B., Ache, H., and Fendler, J. H., *J. Amer. Chem. Soc.* **101**, 4030 (1979).
6. Lim, Y. Y., and Fendler, J. H., *J. Amer. Chem. Soc.* **101**, 4023 (1979).
7. Mortara, R. A., Quina, F. H., and Chaimovich, H., *Biochim. Biophys. Res. Commun.* **81**, 1080 (1978).
8. Paradies, H. H., *Biochim. Biophys. Res. Commun.* **101**, 1096 (1981).
9. Fendler, J. H., *Acc. Chem. Res.* **13**, 7 (1980).
10. Carmona-Ribeiro, A. M., and Chaimovich, H., *Biochim. Biophys. Acta* **733**, 172 (1983).
11. Deamer, D., and Bangham, A. D., *Biochim. Biophys. Acta* **443**, 629 (1976).
12. Rouser, G., Fleischer, S., and Yamamoto, A., *Lipids* **5**, 494 (1970).
13. Bangham, A. D., De Gier, J., and Greville, G. D., *Chem. Phys. Lipids* **1**, 225 (1967).
14. Chong, C. S., and Colbow, K., *Biochim. Biophys. Acta* **436**, 260 (1976).
15. Murakami, Y., Nakano, A., and Fukuya, K., *J. Amer. Chem. Soc.* **102**, 4253 (1980).
16. Cuccovia, I. M., Quina, F. H., and Chaimovich, H., *Tetrahedron* **38**, 917 (1982).
17. Chaimovich, H., Alcixo, R. M. V., Cuccovia, I. M., Zanette, D., and Quina, F. H., in "Solution Behaviour of Surfactants" (K. L. Mittal and E. J. Fendler, Eds.), Vol. 2, p. 949. Plenum Press, New York and London, 1982.
18. Herrmann, U., and Fendler, J. H., *Chem. Phys. Lett.* **64**, 270 (1970).
19. Jähnig, F., Harlos, K., Vogel, H., and Eibl, H., *Biochemistry* **18**, 1459 (1979).
20. Paradies, H. H., *Angew. Chem. Int. Ed. Engl.* **2**, 765 (1982).
21. Suurkuusk, J., Lentz, B. R., Barenholz, Y., Biltonen, R. L., and Thompson, T. E., *Biochemistry* **15**, 1393 (1976).
22. Kantor, H. L., Mabrey, S., Prestegard, J. H., and Sturtevant, J. M., *Biochim. Biophys. Acta* **466**, 402 (1977).
23. Larrabee, A. L., *Biochemistry* **18**, 3321 (1979).
24. Schullery, S. E., Schmidt, C. F., Felgner, P., Tillack, T. W., and Thompson, T. E., *Biochemistry* **19**, 3919 (1980).
25. Sheetz, M. P., and Chan, S. I., *Biochemistry* **11**, 4573 (1972).
26. Huang, C., and Mason, J. T., *Proc. Natl. Acad. Sci. USA* **75**, 308 (1978).
27. Israelachvili, J. N., Mitchell, D. J., and Ninham, B. W., *Biochim. Biophys. Acta* **470**, 185 (1977).
28. Cornell, B. A., Middlehurst, J., and Separovic, F., *Biochim. Biophys. Acta* **598**, 405 (1980).
29. Cornell, B. A., Fletcher, G. C., Middlehurst, J., and Separovic, F., *Biochim. Biophys. Acta* **690**, 15 (1982).
30. Jobst, G., *Ann. Phys.* **78**, 157 (1925).
31. Koch, A. L., *Biochim. Biophys. Acta* **51**, 429 (1961).
32. Blok, M. C., Van Der Neut-Kok, E. C. M., Van Deenen, L. L. M., and De Gier, J., *Biochim. Biophys. Acta* **406**, 187 (1975).
33. Blok, M. C., van Deenen, L. L. M., and De Gier, J., *Biochim. Biophys. Acta* **433**, 1 (1976).
34. Johnson, S. M., and Buttress, N., *Biochim. Biophys. Acta* **307**, 20 (1973).
35. Papahadjopoulos, D., and Watkins, J. C., *Biochim. Biophys. Acta* **135**, 639 (1967).
36. Hopfer, U., Lehninger, A. L., and Lennarz, W. J., *J. Membrane Biol.* **2**, 41 (1970).
37. Nichols, J. W., and Deamer, D. W., *Proc. Natl. Acad. Sci. USA* **77**, 2038 (1980).
38. Nozaki, Y., and Tanford, C., *Proc. Natl. Acad. Sci. USA* **78**, 4324 (1981).
39. Clement, N. R., and Gould, J. M., *Biochemistry* **20**, 1534 (1981).

40. Traudinh, S., Prigent, Y., Lacopère, J. J., and Gary-Bobo, C., *Biochim. Biophys. Res. Commun.* **99**, 429 (1981).
41. Elamrani, K., and Blume, A., *Biochim. Biophys. Acta* **727**, 22 (1983).
42. Keana, J. F. W., Acarregui, M. J., and Boyle, S. L. M., *J. Amer. Chem. Soc.* **104**, 827 (1982).
43. Hasted, J. B., "Aqueous Dielectrics," pp. 139-140. Chapman and Hall, London, 1973.
44. Gouy, M. G., *J. Phys.* **9**, 457 (1910).
45. Chapman, D. L., *Phil. Mag.* **25**, 475 (1913).
46. Stein, W. D., and Nir, S., *J. Membrane Biol.* **5**, 246 (1971).
47. Träuble, H., *J. Membrane Biol.* **4**, 193 (1971).
48. Parsegian, A., *Nature (London)* **221**, 844 (1969).
49. Eisenman, G., Ciani, S. M., and Szabo, G., *Fed. Proc.* **27**, 1289 (1968).
50. Ohki, S., and Aono, O., *J. Colloid Interface Sci.* **32**, 270 (1970).
51. Kornberg, R. D., and McConnell, M. H., *Biochemistry* **10**, 1111 (1971).
52. Hubbell, W. L., and McConnell, H. M., *J. Amer. Chem. Soc.* **93**, 314 (1971).
53. Papahadjopoulos, D., Nir, S., and Ohki, S., *Biochim. Biophys. Acta* **266**, 561 (1971).
54. Van Zoelen, E. J. J., Henriques de Jesus, C., De Jonje, E., Mulder, M., Blok, M. C., and De Gier, J., *Biochim. Biophys. Acta* **511**, 335 (1978).
55. Kleiner, D., *Biochim. Biophys. Acta* **639**, 41 (1981).
56. Alger, J. R., and Prestegard, J. H., *Biophys. J.* **28**, 1 (1979).

Referência 8

micelles with n gegenions. Then the following equations result from the assumption ($K'_{n,i} = K'_{n,1}/i$) for the Poisson distribution¹⁶

$$[MG_i]_n = K'^i_{n,1} [G]^i [M]_n / [Z]^i i!; \quad [MG_i]_n = [MZ_{n-i} G_i] \quad (18)$$

$$[G_t] = [G] + \sum_{n=1}^{\infty} K'_{n,1} [G] [M]_n \exp(K'_{n,1} [G] / [Z]) / [Z] \quad (19)$$

$$[M_t]_n = [M]_n \exp(K'_{n,1} [G] / [Z]); \quad [M]_n = [MG_0]_n \quad (20)$$

$$\bar{G}_n = K'_{n,1} [G] / [Z] \quad (21)$$

$$P(i) = \bar{G}^i \exp(-\bar{G}) / i! \quad (22)$$

where $[M_t]_n$ is the concentration of micelles with n gegenions. Equations 18-22 correspond to eq 25-29 in the ref 16, respectively. Hence, if one assumes the following equality, the difference in theoretical equations between the association and the exchange reaction models disappears

$$K_{n,1} = K'_{n,1} / [Z] \quad (23)$$

However, an equation corresponding to eq 2 becomes from eq 19 and 20

$$([G_t] - [G]) / [G] = \bar{K}_1 [M_t] = (\bar{K}'_1 / [Z]) [M_t] \quad (24)$$

where the following operations for averaging are made for the 1st step

$$\bar{K}'_1 = \sum_{n=1}^{\infty} K'_{n,1} [M_t]_n / \sum_{n=1}^{\infty} [M_t]_n; \quad [M_t] = \sum_{n=1}^{\infty} [M_t]_n \quad (25)$$

What must be stressed here is that the constant of the first stepwise exchange reaction is $\bar{K}'_1$, not $\bar{K}_1$. According to eq 24 the plots of $([G_t] - [G]) / [G]$ against $[M_t]$ should not be linear, because $[Z]$ changes with $[M_t]$. On the contrary, the fact is an excellent linearity of the plots. This means that the above exchange reaction cannot apply to the real bindings. In other words, two kinds of gegenions can independently associate with and dissociate from micelles; there is no 1:1 correspondence between an association of first gegenion and a dissociation of second gegenion, or vice versa. The n value in the summation of eq 25 ranges from 1 to infinity. However, the region in which the n value has a practical importance is a small range of micellar aggregation number, as mentioned above.

Registry No. Cu(DS)₂, 7016-47-9; Zn(DS)₂, 22397-58-6; Mn(DS)₂, 38344-88-6; SDS, 151-21-3; KDS, 4706-78-9.

Supplementary Material Available: Raw data used in plotting the curves in Figures 2, 3, 4, and 7 (4 pages). Ordering information is given on any current masthead page.

Salt Effects on the Stability of Dioctadecyldimethylammonium Chloride and Sodium Dihexadecyl Phosphate Vesicles

A. M. Carmona-Ribeiro,* L. S. Yoshida, and H. Chaimovich

Departamento de Bioquímica, Instituto de Química, Universidade de São Paulo, CP 20780, São Paulo, S.P., Brasil (Received: October 25, 1984; In Final Form: February 20, 1985)

Salt effects on large or small sodium dihexadecyl phosphate (DHP) and dioctadecyldimethylammonium chloride (DODAC) vesicles were determined by turbidimetry, quantified and analyzed within the framework of the Derjaguin-Landau-Verwey-Overbeek (DLVO) theory for electrostatically stabilized colloids. The dependences between initial flocculation rates (V_0) and salt concentration (C) and between the critical coagulation concentration (ccc) and the sixth power of the counterion valency were those expected from the DLVO theory. No correlation was found between $d \log V_0 / d \log C$ and vesicle size. When the amphiphile concentration (C_a) is increased, V_0 increases linearly as the square of C_a , the ccc values remain constant, and the maximal flocculation rate increases. Excimer-monomer (E/M) ratios were obtained from the fluorescence emission of DHP or DODAC vesicles labeled with 2-(10-(1-pyrenyl)decanoyl)phosphatidylcholine (PyPC). The E/M ratios of small vesicles decrease as a function of time after salt addition, while, for large DHP vesicles, E/M ratios remain constant. Fusion only in aggregated small vesicles is strongly suggested by the data.

Introduction

Interface interactions determine a wide variety of phenomena such as adsorption, surface tension, interfacial energy, aggregation, and fusion of vesicles or cells. Theories describing colloid stability explain and predict certain features of the aggregation of phospholipid vesicles.¹⁻⁴ The form of the energy barrier between two particles can be calculated by the Derjaguin-Landau-Verwey-Overbeek (DLVO) theory.⁵⁻¹⁰ When the electrolyte content is increased, the repulsion and stability decrease, the attraction being

assumed as constant. Israelachvili and co-workers have confirmed the classical theories involving van der Waals and electrostatic forces at large distances.^{11,12} In water, a large distance is around and above 50 Å.^{13,14} At smaller distances, surface forces are dominated by those which induce liquid structure, the hydration forces, which depend on the nature of the surface, its counterions, and the ionic concentration.¹⁰

Destabilizing salt effects on synthetic amphiphile vesicles have been reported,¹⁵⁻¹⁹ but quantitative analysis of these effects has

- (1) M. v. Smoluchowsky, *Z. Phys. Chem., Stoechiom. Verwandtschaftsl.*, **92**, 129 (1917).
- (2) S. Nir and J. Bentz, *J. Colloid Interface Sci.*, **65**, 399 (1978).
- (3) V. A. Parsegian, *Annu. Rev. Biophys. Bioeng.*, **2**, 221 (1973).
- (4) L. Rydhag, P. Stenius, and L. Oedberg, *J. Colloid Interface Sci.*, **86**, 274 (1975).
- (5) E. J. W. Verwey and J. T. Overbeek, "Theory of Stability of Liophobic Colloids", Elsevier, Amsterdam, 1948.
- (6) J. T. G. Overbeek in "Colloid Science", H. R. Kruyt, Ed., Elsevier, Amsterdam, 1952, Chapter 8.
- (7) B. V. Derjaguin, *Trans. Faraday Soc.*, **36**, 730 (1940).
- (8) B. V. Derjaguin and L. Landau, *Zh. Eksp. Teor. Fiz.*, **11**, 802 (1941).
- (9) B. V. Derjaguin and L. Landau, *Zh. Eksp. Teor. Fiz.*, **15**, 662 (1945).
- (10) B. W. Ninham, *Pure Appl. Chem.*, **53**, 2135 (1981).

- (11) J. N. Israelachvili and G. E. Adams, *Nature (London)*, **262**, 774 (1976).
- (12) J. N. Israelachvili and G. E. Adams, *J. Chem. Soc., Faraday Trans. 1*, **74**, 975 (1978).
- (13) R. M. Pashley, *J. Colloid Interface Sci.*, **80**, 153 (1981).
- (14) R. M. Pashley, *J. Colloid Interface Sci.*, **83**, 531 (1981).
- (15) C. D. Tran, P. L. Klahn, A. Romero, and J. H. Fendler, *J. Am. Chem. Soc.*, **100**, 1672 (1978).
- (16) K. Kano, A. Romero, B. Djermouni, H. Ache, and J. H. Fendler, *J. Am. Chem. Soc.*, **101**, 4030 (1979).
- (17) R. A. Mortara, F. H. Quina, and H. Chaimovich, *Biochem. Biophys. Res. Commun.*, **81**, 1080 (1978).
- (18) A. M. Carmona-Ribeiro and H. Chaimovich, *Biochim. Biophys. Acta*, **733**, 172 (1983).

only recently been attempted.^{4,20} In this work, salt effects on large and small sodium dihexadecyl phosphate (DHP) or dioctadecyldimethylammonium chloride (DODAC) vesicles are analyzed and discussed in terms of the DLVO theory. A more complete picture of the stability of DODAC and DHP vesicles emerges from concepts currently used for interacting surfaces.

Material and Methods

Dihexadecylphosphoric acid and Sephadex G-25 were purchased from Sigma (St. Louis, Mo). Sodium dihexadecyl phosphate (DHP) was obtained by titration of a methanolic solution of dihexadecylphosphoric acid with sodium methoxide. The precipitated DHP was filtered and dried under vacuum. Dioctadecyldimethylammonium chloride (DODAC) was obtained from Herga (Industrias Químicas do Brasil). The purification and analysis of DODAC have been described elsewhere.²¹ 2-(10-(1-Pyrenyl)decanoyl)phosphatidylcholine (PyPC) was synthesized and purified as described.²² All other reagents were analytical grade. Deionized water doubly distilled in glass was used throughout.

Small DHP or DODAC vesicles were obtained by sonication,¹⁵⁻¹⁷ and large DHP or DODAC vesicles, by vaporization of a chloroformic solution of the amphiphile in hot (70 °C) aqueous solutions.^{18,19} The osmolalities were measured in an osmometer (Model 2007, Precision Systems, Inc., Sudbury, MA). Rigorous control of osmolality is essential for the interpretation of the data, since osmotic gradients induce turbidity changes due to swelling or shrinkage of the vesicles.^{18,19} The osmolality was adjusted in all cases with D-glucose. Salt concentration was determined by titration.²³

The stability of the vesicles after salt addition was characterized by measuring absorbance at 400 nm in a Beckman M-25 spectrophotometer. The time lag between mixing and the start of the register was usually smaller than 6 s. The initial flocculation rate (V_0) was calculated from the absorbance vs. time register. The inverse of V_0 was taken as directly related to the stability of the vesicles.²⁴

The amphiphile concentration (C_a) was determined by measuring total phosphate²⁵ for DHP dispersions and by chloride microtitration²³ for DODAC dispersions.

Vesicles containing around 7 mol % of the fluorescent probe 2-(10-(1-pyrenyl)decanoyl)phosphatidylcholine (PyPC) were prepared as described above, with the exception of small sonicated DODAC vesicles labeled with PyPC. In this case, 1 mL of a chloroformic solution of PyPC (0.2 mg/mL) was evaporated under a nitrogen stream to form a PyPC film on the bottom of the sonication tube. The PyPC film was dried under vacuum for 2 h; DODAC and water were then added and the vesicles obtained by sonication.

For the fluorescence experiments, vesicles labeled with PyPC (0.1 mL) and unlabeled vesicles (0.4 mL), with identical amphiphile concentrations, were mixed. The vesicles interaction was promoted by addition of a salt solution (1.0 mL). Aliquots (50 μ L) of this mixture were diluted to 2 mL of an isoosmolar solution without salt. This dilution served both to stop the vesicles interaction and to eliminate possible scattering artifacts. Fluorescence intensities and spectra were obtained in 1-cm path length quartz fluorescence cells at 22 °C on a Hitachi-Perkin-Elmer MPF-4 spectrometer operated in the energy mode. As a control for PyPC spontaneous exchange between labeled and

Figure 1. Absorbance changes as a function of time for sonicated DODAC vesicles (0.21 mM) prepared in D-glucose (0.428 M) due to addition of NaCl to a final concentration of 0.030 (a), 0.064 (b), and 0.113 M (c).

Figure 2. Amphiphile concentration (C_a) effect on the initial flocculation rate (V_0) of sonicated (A and C) and injected (B and D) vesicles prepared in D-glucose (0.428 M). The final NaCl concentration was 0.107 M. Open symbols refer to DHP, and solid symbols refer to DODAC.

unlabeled vesicles, isoosmolar solutions without salt were used.

Results

Under isoosmolar conditions, the addition of salt to DODAC and DHP vesicles produced turbidity increases which were time and concentration dependent (Figure 1). From data such as that presented in Figure 1, initial flocculation rates (V_0) were obtained and used for quantitative analysis of the amphiphile and salt concentration effects on the stability of the vesicles.

V_0 increased with the concentration of amphiphile (C_a) for both small and large vesicles and was linearly related to the square of the amphiphile concentration (Figure 2). For sonicated vesicles V_0 values were approximately 1 order of magnitude higher than those of large vesicles, indicating a higher stability for the latter. DODAC vesicles were more sensitive to an increase in amphiphile concentration than the corresponding DHP preparations.

From data obtained in both colloid and lipid vesicle systems, it was expected that the stability of DODAC and DHP vesicles should depend on the electrolyte concentration,^{4,24} and in fact, V_0 varied sharply with salt concentration (C). Expressing V_0 and C as logarithmic functions, we obtained the critical coagulation concentration (ccc) (Figure 3). Below the ccc, V_0 increases (stability decreases) as a function of C and $d \log V_0 / d \log C$ can

(19) A. M. Carmona-Ribeiro, L. S. Yoshida, A. Sesso, and H. Chaimovich, *J. Colloid Interface Sci.*, **100**, 433 (1984).

(20) L. Rydhag, K. Rosenquist, P. Stenius, and L. Oedberg, in "Physical Chemistry of Surfactants in Solution", Plenum Press, New York, 1983.

(21) I. M. Cuccovia, R. M. V. Aleixo, R. A. Mortara, P. Berci Filho, J. B. S. Bonilha, F. H. Quina, and H. Chaimovich, *Tetrahedron Lett.*, 3065 (1979).

(22) S. Schenkman, P. S. Araujo, R. Dijkman, F. H. Quina, and H. Chaimovich, *Biochim. Biophys. Acta*, **649**, 633 (1981).

(23) O. Schales and S. S. Schales, *J. Biol. Chem.*, **140**, 879 (1941).

(24) H. Reerink and J. T. G. Overbeek, *Discuss. Faraday Soc.*, **18**, 74 (1954).

(25) G. Houser, S. Fleischer, and A. Yamamoto, *Lipids*, **5**, 494 (1970).

Figure 3. Salt concentration effect (C) on the initial flocculation rate (V_0) of DODAC vesicles prepared in 0.43 M D-glucose and DHP vesicles prepared in 0.55 M D-glucose. C_a values (mM) are as follows: 0.10 (A), 0.23 (B), and 0.47 (C), for sonicated DHP vesicles; 0.09 (D), 0.18 (E), and 0.33 (F), for injected DHP vesicles; 0.11 (G), 0.21 (H), and 0.48 (I), for sonicated DODAC vesicles; 0.11 (J), 0.24 (K), and 0.55 (L), for injected DODAC vesicles.

TABLE I: Critical Coagulation Concentrations (ccc), Maximal Initial Flocculation Rates ($V_{0,max}$), and Slopes $d \log V_0/d \log C$ for the NaCl Effect on the Stability of DHP and DODAC Vesicles

vesicle type	C_a , mM	ccc, M	$10^3 \times V_{0,max}$	$d \log V_0/d \log C$
sonicated DHP	0.10	0.186	2.82	11.2
	0.23	0.178	12.59	12.9
	0.47	0.170	25.12	13.3
		av 0.178		av 12.5
injected DHP	0.09	0.207	0.75	10.6
	0.18	0.229	3.76	9.1
	0.33	0.229	7.50	10.3
		av 0.222		av 10.0
sonicated DODAC	0.11	0.129	0.71	2.3
	0.21	0.112	1.50	1.9
	0.48	0.126		2.1
		av 0.122		av 2.1
injected DODAC	0.11	0.140	0.09	3.8
	0.24	0.129	0.12	3.1
	0.55	0.155	0.35	3.4
		av 0.141		av 3.4

be calculated. Above the ccc, V_0 remained constant with increasing salt concentration, yielding maximum values for V_0 ($V_{0,max}$). The values of ccc, $V_{0,max}$, and $d \log V_0/d \log C$ are presented in Table I. For a given amphiphile and vesicle size, the ccc was independent of C_a and mean ccc values were calculated from the ccc at several amphiphile concentrations. Two types of correlations are apparent from data in Table I. The mean ccc values are higher for large vesicles independent of the nature of the amphiphile, and DHP vesicles exhibit a higher mean ccc. Since ccc values can be used as criteria of salt stability, it is evident that DHP vesicles are more stable than DODAC vesicles toward salt-induced coagulation. At salt concentrations below the ccc, the values of

TABLE II: NaCl and $CaCl_2$ Effects on Critical Coagulation Concentrations (ccc), Maximal Initial Flocculation Rates ($V_{0,max}$), and $d \log V_0/d \log C$ of DHP and DODAC Vesicles

vesicle type	ccc, M		$10^3 V_{0,max}$		$d \log V_0/d \log C$	
	NaCl	$CaCl_2$	NaCl	$CaCl_2$	NaCl	$CaCl_2$
sonicated 0.30 mM DHP	>0.168	0.002		5.8	7.1	3.1
injected 0.28 mM DHP	>0.168	0.003		4.4	6.2	2.8
sonicated 0.21 mM DODAC	0.132	0.089	2.5	2.0	3.3	2.4
injected 0.24 mM DODAC	0.138	0.102	0.4	0.3	3.4	6.3

$d \log V_0/d \log C$ for sonicated and injected DHP vesicles were practically identical. For sonicated DODAC vesicles, $d \log V_0/d \log C$ values were slightly smaller than those for injected DODAC vesicles (Table I). Thus, no correlation was found between $d \log V_0/d \log C$ and vesicle size.

The effects of Na^+ and Ca^{2+} on the stability of DHP vesicles were markedly different (Figure 4A,B). In contrast, these ions affected the stability of DODAC vesicles very similarly (Figure 4C,D), suggesting that coagulation is controlled by Cl^- . Ccc values for DHP vesicles in the presence of Ca^{2+} are 2 orders of magnitude smaller than those for Na^+ and $d \log V_0/d \log C$ values for Na^+ were higher than those for Ca^{2+} (Table II).

Several conditions which lead to vesicle aggregation produce vesicle fusion both above and below the phase transition temperature of the lipid bilayer.^{2,26-28} The variation of the excim-

(26) J. Lansman and D. H. Haynes, *Biochim. Biophys. Acta*, **394**, 335 (1975).

(27) D. Papahadjopoulos, W. J. Vail, C. Newton, S. Nir, K. Jacobson, G. Poste, and R. Lazo, *Biochim. Biophys. Acta*, **465**, 579 (1977).

Figure 4. CaCl_2 (O) and NaCl (●) effects on the initial flocculation rates (V_0) of sonicated (A) or injected (B) 0.3 mM DHP vesicles and sonicated (C) or injected (D) 0.2 mM DODAC vesicles prepared in 0.428 M D-glucose.

Figure 5. Excimer-monomer ratios (E/M) of PyPC as a function of time for partially labeled DODAC or DHP sonicated (A, B, and C) and injected (D) vesicles. Final concentrations of amphiphile and NaCl in the mixture of labeled and unlabeled vesicles are, respectively, the following: 0.26 mM DHP and 0.115 M (A); 0.28 mM DHP and 0.079 M (B); 0.49 mM DODAC and 0.056 M (C); 0.18 mM DHP and 0.115 M (D). Open symbols refer to points obtained after salt addition, and solid symbols refer to points obtained in absence of salt (dilution control).

er-monomer ratio (E/M) of the fluorescent probe PyPC has been used for studies of protein-induced fusion of small unilamellar lecithin vesicles.²² DODAC or DHP vesicles containing PyPC were mixed with unlabeled vesicles. In absence of salt, E/M values of these vesicles mixtures remained constant for hours (Figure

TABLE III: Calculated Stern Potential Values (ψ) of DHP and DODAC Vesicles

vesicle type	vesicle radius, ^a Å	d log V_0 /d log C ^b	γ ^c	ψ , mV
sonicated DHP	400	12.5	0.381	-41.2
injected DHP	1350	10.0	0.186	-19.3
sonicated DODAC	150	2.1	0.255	26.8
injected DODAC	2500	3.4	0.080	8.5

^a From ref 15-19. ^b From Table I in Results section. ^c γ value is defined in eq 2.

5). The addition of salt to large DHP vesicles did not change the E/M values as a function of time (Figure 5D). In contrast, the E/M values of both DHP- and DODAC-sonicated vesicles decreased significantly upon salt addition (Figure 5A-C).

Discussion

In this section, current concepts for interacting surfaces are applied to create a more complete picture of the stability of DODAC and DHP vesicles.

Long-range van der Waals attractions²⁹⁻³² combined with long-range electrostatic repulsion are the forces considered for development of the DLVO theory for colloid stability.⁵⁻⁹ If there was no electrostatic repulsion between two colloidal particles, every collision would lead to coagulation according to Brownian collisions (rapid coagulation of von Smoluchowski).¹ When the repulsive energy is not zero, the fraction of effective collisions is much smaller (slow coagulation). As the salt concentration (C) or the counterion valency (ν) increases, the maximum of the potential energy of interaction ($V_{\max}^x$) for a given interparticle distance (x) decreases and the stability (W) decreases, W being directly related to $1/V_0$.²⁴ When $V_{\max}^x$ is small compared to kT , the system coagulates. The ccc corresponds to the salt concentration where $V_{\max}^x$ vanishes. Thus, from the DLVO theory, the stability (W) of the vesicles depends on the salt concentration (C), the counterion valency (ν), the Stern potential (ψ), the vesicle radius (a), and the van der Waals constant (A).

To avoid the laborious computations of the exact DLVO theory, two approximate equations were derived.²⁴ One has the form given in the equation

$$W = \text{constant} \exp(V_{\max}^x/2kT) \quad (1)$$

indicating that the stability factor W is determined by the maximum in the potential curve. The other starts from approximate equations (2) and (3), leading to a linear relationship between $\log W$ and $\log C$ for certain combinations of ψ and A when a compensation of errors occurs²⁴

$$V_R^{aa} = 4.62 \times 10^{-6} (a\gamma^2/\nu^2) \exp(-KH_0) \quad (2)$$

$$V_A^{aa} = -Aa/(12H_0) \quad (3)$$

where V_R^{aa} is the repulsive energy of interaction between the overlapping double layers on two spherical particles of equal size in water at 25 °C, a is the vesicle radius, ν is the counterion valency, γ is given by

$$(\exp(\nu q\psi/2kT) - 1)/(\exp(\nu q\psi/2kT) + 1)$$

with q as the electronic charge, $1/K$ is a measure of the thickness of the diffuse double layer, H_0 is the shortest distance between particle surfaces, V_A^{aa} is the attraction potential between two spheres for short distances ($H_0 \ll a$), and A is the van der Waals constant.

From the approximated linear relationship between $\log W$ and $\log C$, at 25 °C in water, the slope is given by²⁴

$$d \log W/d \log C = -2.15 \times 10^7 a\gamma^2\nu^{-2} \quad (4)$$

(29) H. Kallmann and M. Willstaetter, *Naturwissenschaften*, **20**, 952 (1932).

(30) J. H. de Boer, *Trans. Faraday Soc.*, **32**, 21 (1936).

(31) H. C. Hamaker, *Physica (Amsterdam)*, **4**, 1058 (1937).

(32) J. N. Israelachvili, *Q. Rev. Biophys.*, **6**, 341 (1974).

(28) M. Wong and T. E. Thompson, *Biochemistry*, **21**, 4133 (1982).

Ratios between $d \log V_0/d \log C$ for Ca^{2+} and Na^+ (Table II) are 0.19 for sonicated DHP vesicles and 0.45 for injected DHP vesicles, in reasonable agreement with the expected ratio (0.25) calculated from eq 4.

$d \log V_0/d \log C$ values in Table I immediately lead to γ and to the Stern potential (ψ) by using values for a from the literature.¹⁵⁻¹⁹ Calculated values (Table III) are low as expected from the employed approximation (see Figure 4 in ref 24) and in comparison with some experimental data reporting surface potentials in surfactant vesicles.³³ However, these calculated values can be used for comparison. DHP vesicles exhibit higher Stern potentials than the correspondent DODAC preparations. This difference could be the source of the higher stability of DHP vesicles (Figure 3, Table I). From the higher values for sonicated vesicles, in comparison with those for the injected (Table III), the former should be more stable, in conflict with experimental data (Figure 3 and Table I). If one considers the van der Waals interaction (eq 3), as a and/or A increases, V_A^{aa} also increases in absolute value. Thus, injected vesicles having both a and A greater than the sonicated vesicles^{2,32} would be under the strongest attractive interaction, also in disagreement with experimental data. In conclusion, the framework of the DLVO theory does not explain the compared stability of small and large DODAC or DHP vesicles. Structural differences between small and large vesicles inducing distinct head-group repulsions and hydrophobic interactions between adjacent monomers in each vesicle type could lead to distinct hydration forces and surface repulsions at short ranges. For mica surfaces and lecithin multibilayers, adhesion in a primary minimum (as expected from the DLVO theory) is prevented by short-range repulsions due to the overlap of hydration "shells" of ions held in the compressed double layer.^{12,14,34} Secondary minimum coagulation, however, would still be possible for the injected vesicles.^{2,26} In fact, for large amphiphile vesicles, we could visually observe the formation of "clouds" some hours after salt addition which reverted to an homogeneously turbid dispersion by hand shaking as occurs for colloidal particles coagulating in the secondary minimum.³⁵

From eq 4, for different vesicle sizes one should obtain significantly different $d \log V_0/d \log C$ values. However, no correlation was found between $d \log V_0/d \log C$ and vesicle size (Figure 3 and Table I).

The ccc for two particles with Stern potential is given by

$$\text{ccc} = (\text{constant})\gamma^4/(A^2\nu^6) \quad (5)$$

From eq 5, the ccc is independent of the amphiphile concentration, as was in fact shown in the Results Section (Figure 3 and Table I). The significant increase in the ccc obtained for injected in comparison with sonicated DODAC or DHP vesicles could be ascribed to higher Stern potentials and/or smaller van der Waals constants (A). If ψ is higher for sonicated than for injected vesicles (Table III) and A values are expected to be larger for injected than for sonicated vesicles,^{2,26} from eq 5, ccc values for sonicated vesicles should be higher than those for the corresponding injected vesicles. This reasoning further corroborates the disagreement between predictions from the DLVO theory and the compared stability of DODAC or DHP vesicles of different size.

Equation 5 deduced from the DLVO theory constitutes an expression for the empirical rule of Schulze-Hardy for high potentials.^{4,36} This rule states that $(\text{ccc})\nu^6$ is a constant, the counterion valency having a very large effect on coagulation rates. As a consequence, the ratio between ccc values for Ca^{2+} and Na^+ for DHP vesicles should be 0.016. These ratios are ca. 0.018 for injected and 0.012 for sonicated DHP vesicles, as calculated from data on Table II, in good agreement with the expected ratio for high potentials. This agreement confirms that the calculated values are underestimated, since the rule works at high potentials. For vesicles of dimyristoylphosphatidylcholine (DMPC) containing

10% cetyltrimethylammonium bromide (CTAB), ccc values for different counterions agreed more fairly with eq 5,⁴ a suggestion that only 10% CTAB in the vesicle bilayer may generate a high potential.

V_0 is second order in concentration (Figure 2), as flocculation is essentially a "bivesicular" process, at least for its early steps. This result was also found for aggregation rates of dipalmitoylphosphatidylcholine vesicles at temperatures below the gel-liquid crystalline phase transition²⁸ or for the initial rates of fusion of small phosphatidylserine vesicles in the presence of Ca^{2+} .³⁷ In contrast, for phosphatidic acid vesicles in the presence of CaCl_2 , the rate of scattering increase varied linearly with vesicle concentration.²⁶ Oster³⁸ demonstrated that coagulation processes may be regarded as polymerization processes. Thus, rapid coagulation of colloids can be viewed essentially as a linear condensation polymerization, since any aggregate of colloid units may combine with any other aggregate. For linear condensation polymerization, light scattering increases linearly in time with a slope proportional to the rate constant of the elementary reaction and to the square of the concentration of the colloid substance. Recently, rate constants were calculated from measured flocculation rates of DMPC/CTAB and PC/DCP vesicles following the Oster equation,^{4,20} presented as

$$d\delta/dt = k'2B_0C_0^2/\rho^2 \quad (6)$$

where δ is the turbidity measured as absorbance, C_0 is the concentration (g/cm^3) of scattering particles with density ρ , and B_0 is an optical constant given by

$$B_0 = (32/3)\pi^3\rho^2n_0^2(dn/dC)^2(1/\lambda^4) \quad (7)$$

In this equation B_0 depends on the refraction index increment of the vesicles (dn/dC), the refractive index of water (n_0), the incident wavelength (λ), and the density of the scattering particles (ρ).

From (6), $d\delta/dt$ varies linearly with the square of the concentration for particles much smaller than the incident wavelength. We can apply this equation for the sonicated DODAC vesicles and calculate k' . Taking C_0 as $6.6 \times 10^{-5} \text{ g}/\text{cm}^3$, dn/dC as $0.142 \text{ cm}^3/\text{g}$,³⁹ and $d\delta/dt$ as $7.1 \times 10^{-4} \text{ s}^{-1}$ for a final salt concentration equal to or bigger than the ccc (Table I), a value of $4 \times 10^{-14} \text{ cm}^3/\text{s}$ may be obtained for k' . This value is ca. 3 orders of magnitude too low when compared to simple Smoluchowsky coagulation rates and within the range of the values obtained for vesicles of phosphatidic acid and phosphatidylserine.²⁶ Calculated k' values for DMPC/CTAB- or PC/DHP-sonicated vesicles of ca. $2 \times 10^{-15} \text{ cm}^3/\text{s}$ have been reported also for conditions in which electrostatic repulsion is removed (at the ccc).²⁰ Thus, flocculation rates are slower than those expected if the process were simply diffusion controlled when electrostatic repulsion is absent. Probably extra repulsive hydration forces^{40,41} occurring at close approach of vesicles are involved in the observed slower processes.

The effect of divalent cations on acidic phospholipid vesicles involves more than charge neutralization and aggregation. Ca^{2+} and other divalent and higher valent cations bind strongly to phosphatidic acid membranes, whereas the monovalent cations do not.^{26,42} Ca^{2+} may function as a cross-link between negatively charged vesicles. The threshold for the Ca^{2+} concentration dependence of the vesicles aggregation coincides with the point of neutralization of the membrane surface charge.²⁶ At Ca^{2+} concentrations between 0.5 and 2.0 mM, phosphatidylserine-sonicated vesicles fuse.²⁷ The ccc values for Ca^{2+} in DHP vesicles (Table II) coincided with the threshold for Ca^{2+} -induced aggregation of

(37) J. Wilschut, N. Duzgunes, R. Fraley, and D. Papahadjopoulos, *Biochemistry*, **19**, 6011 (1980).

(38) G. Oster, *J. Colloid Interface Sci.*, **2**, 291 (1947).

(39) P. N. Yi and R. C. Macdonald, *Chem. Phys. Lipids*, **11**, 114 (1973).

(40) D. M. Le Neveu, R. P. Rand, and V. A. Parsegian, *Nature (London)*, **259**, 601 (1976).

(41) A. C. Cowley, N. L. Fuller, R. P. Rand, and C. A. Parsegian, *Biochemistry*, **17**, 3163 (1978).

(42) D. H. Haynes, *J. Membrane Biol.*, **17**, 341 (1974).

(43) J. B. Birks, "Photophysics of Aromatic Molecules", Elsevier, New York, 1968.

(44) T. Foerster, *Angew. Chem., Int. Ed. Engl.*, **8**, 333 (1969).

(33) S. Lukac, *J. Phys. Chem.*, **87**, 5045 (1983).

(34) J. N. Israelachvili, *Faraday Discuss. Chem. Soc.*, **65**, 20 (1978).

(35) S. Okamoto and S. Hachisu, *J. Colloid Interface Sci.*, **43**, 30 (1972).

(36) J. T. G. Overbeek, *Pure Appl. Chem.*, **52**, 1151 (1980).

phosphatidylserine or phosphatidic acid vesicles²⁶ and is in the Ca^{2+} concentration range used for inducing fusion.²⁷

Since excimer formation is a bimolecular process, its concentration and relative emission intensity (E/M) are directly proportional to the microscopic PyPC concentration or inversely proportional to the microscopic volume confining the PyPC molecules.^{45,46} For a mixture of PyPC-labeled and unlabeled vesicles, the occurrence of salt-induced fusion implies an increase of the available volume of amphiphile in which PyPC can dilute and a correspondent decrease of E/M. E/M decrease with time suggests that sonicated vesicles undergo fusion due to added salt (Figure 5A-C). However, one could also expect an analogous

E/M decrease if PyPC exchanged from labeled to unlabeled vesicles, as reported in cases of spontaneous phospholipid transfer.⁴⁷ As E/M for the mixture of labeled and unlabeled vesicles remains constant in absence of salt (see controls in Figure 5), we consider our data as strong evidence of salt-induced fusion in sonicated amphiphile vesicles and absence of salt-induced fusion in large injected amphiphile vesicles.

Acknowledgment. The financial support from FAPESP, FINEP, and CNPQ is gratefully acknowledged. We are indebted to Drs. M. J. Politi and L. Sepulveda for careful reading of the manuscript.

Registry No. DHP, 2197-63-9; DODAC, 107-64-2; Na, 7440-23-5; Ca, 7440-70-2.

(45) A. K. Soutar, H. J. Pownall, A. S. Hu, and L. C. Smith, *Biochemistry*, **13**, 2828 (1974).

(46) H. J. Pownall and L. C. Smith, *J. Am. Chem. Soc.*, **95**, 3136 (1973).

(47) M. A. Roseman and T. E. Thompson, *Biochemistry*, **19**, 439 (1980).

Micellar Microfluidities at High Pressures

W. D. Turley and H. W. Offen*

Department of Chemistry and The Marine Science Institute, University of California, Santa Barbara, California 93106 (Received: February 4, 1985)

The monomer/excimer intensity ratio and excimer lifetime of dipyranylpropane (DPyP) dissolved in micellar media are used to infer the microfluidity of the solubilization site as a function of pressure to 300 MPa. The viscosity scale at 15.0 °C is calibrated with DPyP in isobutyl alcohol (5–90 cP) and used as a standard for the probe studies of SDS (19 cP), LDS (19 cP), CTAC (39 cP), and C_{12}E_6 (57 cP). Their respective activation volumes of viscous flow, $\Delta V_\eta^\ddagger$, are 13, 11, 24, and 25 cm^3/mol . The marked difference in fluidity and its pressure coefficient of anionic surfactants as opposed to cationic and nonionic micelles is explained in terms of the relative strengths of headgroup hydration.

Introduction

The utility of surfactant micelles lies in their ability to solubilize various types of molecules.^{1,2} The use of fluorescent probes represents an important method for examining the probe microenvironment in micellar aggregates.^{1–4} The present work focuses on the pressure dependence of the effective micellar microviscosity η inferred from the intramolecular excimer kinetics of 1,3-di(1-pyranyl)propane (DPyP).⁵

Kinetic analysis of diarylpropanes^{5–7} is based on the observation that electronic excitation of one aromatic chromophore in fluid media leads to fluorescence from two temporally and energetically distinct species—the monomer M^* and excimer E^* —as shown in Scheme I. The excited-state rate of association k_a and dissociation k_d are related to the fluorescence intensity ratio I_M/I_E , the excimer lifetime τ_E , and the radiative rate constants k_M and k_E of the respective M^* and E^* by

$$I_M/I_E = (k_M/k_E)(k_d + \tau_E^{-1})k_a^{-1} \quad (1)$$

This expression for two dynamically coupled species can be simplified in the low-temperature or high-viscosity ($\eta \geq 5$ cP) limit,⁵

SCHEME I

where the dissociation rate becomes negligible ($\tau_E^{-1} \gg k_d$) and excimer formation is viscosity controlled. If diffusion of the aromatic chromophores to achieve the excimer conformational state is rate limiting, the applicable equation is $k_a = c\eta$, where c is a constant that depends only on temperature for a given solute/solvent system.⁶ These two approximations lead to a simple relation between observables and η of the solubilization sites:

$$\tau_E I_M/I_E = (k_M/k_E)c\eta \quad (2)$$

Application of eq 2 to pressure studies requires the reasonable assumption that the ratio of radiative transition rates are pressure independent.⁸ Then the pressure coefficient of the microviscosity is

$$d \ln \eta / dP = d \ln (\tau_E I_M/I_E) / dP \quad (3)$$

and the activation volume of viscous flow is

$$\Delta V_\eta^\ddagger = RT[d \ln (\tau_E I_M/I_E) / dP]_T \quad (4)$$

Equations 2, 3, and 4 can be used to evaluate $\eta(0)$ and $\eta(P)$ of micelle-probe aggregates provided that the bulk viscosity and its

(1) Fendler, J. H. "Membrane Mimetic Chemistry"; Wiley: New York, 1982.

(2) Mittal, K. L.; Lindman, B., Eds. "Surfactants in Solution"; Plenum Press: New York, 1984; Vol. I.

(3) Turro, N. J.; Gratzel, M.; Braun, A. M. *Angew. Chem., Int. Ed. Engl.* **1980**, **19**, 675.

(4) Singer, L. A. In "Solution Behavior of Surfactants", Vol. 1, Mittal, K. A., Fendler, E. J., Eds.; Plenum: New York, 1982.

(5) Zachariasse, K. A. *Chem. Phys. Lett.* **1978**, **57**, 429.

(6) Birks, J. B. "Photophysics of Aromatic Molecules"; Wiley-Interscience: London, 1970.

(7) Zachariasse, K. A.; Duveneck, G.; Busse, R. *J. Am. Chem. Soc.* **1984**, **106**, 1045.

(8) Johnson, P. C.; Offen, H. W. *J. Chem. Phys.* **1972**, **56**, 1638. Offen, H. W.; Phillips, D. T. *J. Chem. Phys.* **1968**, **49**, 3995.

Referência 9

SALT-INDUCED AGGREGATION AND FUSION OF DIOCTADECYLDIMETHYLAMMONIUM CHLORIDE AND SODIUM DIHEXADECYLPHOSPHATE VESICLES

A. M. CARMONA-RIBEIRO AND H. CHAIMOVICH

Departamento Bioquímica, Instituto de Química, Universidade de São Paulo, CEP 20780, Brazil

ABSTRACT Small dioctadecyldimethylammonium chloride (DODAC) vesicles prepared by sonication fuse upon addition of NaCl as detected by several methods (electron microscopy, trapped volume determinations, temperature-dependent phase transition curves, and osmometer behavior). In contrast, small sodium dihexadecyl phosphate (DHP) vesicles mainly aggregate upon NaCl addition as shown by electron microscopy and the lack of osmometer behavior. Scatter-derived absorbance changes of small and large DODAC or DHP vesicles as a function of time after salt addition were obtained for a range of NaCl or amphiphile concentration. These changes were interpreted in accordance with a phenomenological model based upon fundamental light-scattering laws and simple geometrical considerations. Short-range hydration repulsion between DODAC (or DHP) vesicles is possibly the main energy barrier for the fusion process.

INTRODUCTION

Our understanding of structure and function of biological membranes has been greatly developed by using structures of defined chemical composition having just some of the properties of biological membranes (1). Another aspect that has emerged from membrane mimicking is a novel chemistry of both practical and theoretical utility (2, 3).

The general problem of stability of interacting membranes is relevant in many systems. The process of mutual approach of membranes leading to adhesion or fusion may be conveniently divided into three main stages: (a) mutual approach; (b) destabilization of the liquid film between the membranes and/or the membranes themselves; and (c) expansion of the area of membrane contact during adhesion or expansion of the hole during fusion (4). Phospholipid vesicles have been extensively used as models for membrane fusion studies. Interaction forces involved in the mutual approach of membranes have been used to describe salt-induced interaction of phospholipid (5–10) or surfactant (11) vesicles. Didodecyldimethylammonium bromide (DDAB) large vesicles fuse in the presence of micromolar concentrations of certain organic and inorganic divalent anions (12). We have suggested that small sonicated vesicles of dioctadecyldimethylammonium chloride (DODAC) and sodium dihexadecylphosphate (DHP) vesicles fuse upon addition of NaCl (11). Here we unequivocally demonstrate NaCl-induced fusion of small sonicated DODAC vesicles and present a model that describes vesicle aggregation and fusion. We also introduce a simple

method for kinetic or equilibrium detection of the fusion of small vesicles.

MATERIAL AND METHODS

Dioctadecyldimethylammonium chloride (DODAC) and sodium dihexadecylphosphate (DHP) were purified and analyzed as described (11). Sephadex G-25 (fine) was obtained from Sigma Chemical Co. (St. Louis, MO) and (¹⁴C)-sucrose from Schwarz-Mann (Orangeburg, NY) specific activity 480 mCi/mmol). All other reagents were analytical grade. Deionized water, doubly distilled in glass, was used throughout. DODAC and NaCl concentrations were determined by chloride microtitration (13) and DHP concentration by measuring phosphate (14).

Small DODAC or DHP vesicles were obtained by sonication (15, 16) and large DODAC or DHP vesicles by vaporization of a DODAC or DHP chloroformic solution in aqueous solutions (17, 18). Salt-induced turbidity changes of vesicle dispersions were followed by measuring absorbance at 400 nm in a Beckman M-25 Spectrophotometer. The time-lag between mixing and register start was usually <6 s. The scatter-derived absorbance will be referred to as absorbance hereafter. The osmolality of the aqueous solutions used to prepare the large vesicles and that of the salt solution added to the vesicles were kept equal. Osmotic gradients induce turbidity changes as a consequence of swelling or shrinking of the vesicles (17, 18). Osmolality values were adjusted in all cases with D-glucose and measured in an osmometer (model 2007; Precision Systems, Inc., Natick, MA).

For electron microscopy, aliquots of sonicated DODAC or DHP vesicles (2.5 mM) in the presence of NaCl (0.042 M) were mixed with an equal volume of ammonium molybdate (2%) or uranyl acetate (1%) for 30 s. After draining and drying, the sample was examined in a Zeiss EM-9 52 Electron Microscope (Carl Zeiss, Inc., Thornwood, NY) operated at 60 KV. From electron micrographs, histograms were obtained and mean external diameters calculated.

Changes of absorbance as a function of temperature were obtained in a Cary-14 Spectrophotometer (Applied Physics Corp., Monrovia, CA) equipped with a thermostatically controlled cell compartment coupled to a Haake F-12 Circulating Bath (Haake Buchler Instruments, Inc., Saddle Brook, NJ). Heating or cooling rates were usually 2°K/min.

Correspondence should be addressed to A. M. Carmona-Ribeiro.

FIGURE 1 Scatter-derived absorbance change induced by NaCl on small (*A*, *B*, *D*) and large (*C*, *D*) DODAC vesicles. In *A*, DODAC concentration is 1.8 mM and NaCl concentration (mM) is 42 (*a*); 64 (*b*); 85 (*c*); 132 (*d*). In *B*, NaCl concentration is 0.117 M and DODAC concentration (mM) is 0.47 (*a*); 0.85 (*b*); 1.03 (*c*); 1.13 (*d*). In *C*, DODAC concentration is 0.26 mM and NaCl concentration (mM) is: 0 (*a*); 30 (*b*); 64 (*c*); 100 (*d*); 130 (*e*); 171 (*f*). In *D*, absorbance changes for small (*a*) and large (*b*) DODAC vesicles (0.35 mM) are compared in 110 mM NaCl. Large DODAC vesicles were prepared in 0.428 M D-glucose.

Temperatures were measured inside the cuvette with a copper-constantan thermocouple connected to a millivoltmeter. The response of this measuring system is linear between 0 and 100°C with a relative error of <0.01%.

Determination of internal volume (*V*) for sonicated DODAC vesicles upon salt addition was performed using (¹⁴C)-sucrose as a marker (19). A typical experiment was as follows. Sonicated DODAC vesicles (10 mM) were prepared in 0.02 M sucrose. 5.0 ml of this preparation was mixed with 5 ml of a 0.02 M sucrose and 0.100 M NaCl solution containing (¹⁴C)-sucrose. The time immediately after mixing was taken as zero time. Aliquots (0.02 ml) were then filtered through a Sephadex G-25 column (28 × 1 cm) previously saturated with sonicated DODAC vesicles and equilibrated with a 0.05 M NaCl and 0.02 M sucrose solution. Radioactivity and DODAC recoveries were higher than 93% in all cases. Radioactivity of the elution profiles was used for the calculation of *V* as described (17).

For determination of osmotic properties, vesicles in the presence of salt were added to aqueous solutions of NaCl (or NaCl plus erythritol) and the time-dependent absorbance changes were registered at 400 nm.

The initial shrinking rate (*v_s*%) was defined as: $v_s\% = (100/A_0) \cdot (\Delta A/\Delta t)$, where $\Delta A/\Delta t$ is the initial slope of the curve of absorbance vs. time and *A₀*, the initial absorbance. The reciprocal of the total absorbance change was normalized by using the parameter $A_0/\Delta A = A_0/(A_f - A_0)$ where *A₀* is the measured absorbance 24 h after the addition of vesicles to hypo- or hyperosmotic solutions. Mean values for these parameters were obtained from at least two independent experiments and errors are expressed as mean-square deviations.

RESULTS

To demonstrate the fusogenic effect of salt solutions on small sonicated DODAC vesicles, we characterized some physical and functional properties of the dispersions obtained upon NaCl addition. These properties were compared with those of the nontreated preparation and those of large DODAC vesicles obtained by chloroform vaporization (17).

FIGURE 2 Electron microscopy of a positively stained preparation of sonicated DODAC vesicles (2.5 mM) 48 h after NaCl addition (0.042 M) (*A*). The distribution of vesicles size is displayed in *B*.

FIGURE 3 Temperature-induced phase transition of sonicated DODAC vesicles (2.5 mM) 7 h after NaCl addition (0.042 M). Cooling or heating is indicated by arrows.

FIGURE 4 Spectra of sonicated DODAC vesicles (5 mM) prepared in water before (A) and 40 min (B), 3 h (C), 6 h (D), or 24 h (E) after NaCl addition (0.042 M). Final DODAC concentration in the sample containing salt is 2.5 mM.

Time-dependent absorbance changes obtained after salt addition for a range of salt (Fig. 1 A) or DODAC concentration (Fig. 1 B) were similar in shape to those obtained for large vesicles (Fig. 1 C). The change in absorbance as a function of time had three phases. There was an increase followed by a plateau then a decrease. For sonicated vesicles, this time-sequence was slower when compared with large DODAC vesicles (Fig. 1 D). DODAC dispersions remained homogeneous for at least 72 h after salt addition and began to become gelatinous after one month with no visible flocculation or precipitation.

Preparations of DODAC sonicated vesicles were examined by electron microscopy with both negative and positive staining 48 h after NaCl addition. The micrographs showed closed vesicular structures (Fig. 2 A) with a mean external diameter of 1.3 μm (range 0.2–4.0 μm , Fig. 2 B). This preparation will be hereafter referred to as fused DODAC vesicles.

The change of absorbance as a function of temperature for sonicated DODAC vesicles incubated for 7 h with a NaCl solution is shown in Fig. 3. The absorbance decreases monotonically with temperature. A sharp change in slope was observed around the (heating) transition temperature (38.6°C) and a pronounced hysteresis upon cooling (Fig. 3). A reproducible difference was detected between the absorbances observed before and after the heating-cooling cycle. The final absorbance after the temperature cycle remained unaltered (room temperature) for at least 24 h.

Absorbance (A) of sonicated DODAC vesicles prepared in water varied linearly with λ^{-4} (Fig. 4 A). This wavelength dependence remained unchanged for at least 24 h after preparation of the vesicles (13). After salt addition, A varied linearly with λ^{-2} (Fig. 4 B–E).

If a water-soluble impermeable marker such as (^{14}C)-sucrose (17) is added to sonicated DODAC vesicles, fusion can lead to an entrapment of the marker. This entrapment (V) can be used as a measure of fusion with time (19–21). For sonicated DODAC vesicles, V values increased as a function of time after salt addition (Fig. 5 A). This figure includes the V value calculated from electron micrographs (Fig. 2), taking 40 $\text{\AA}^2$ as the mean area per DODAC monomer (22). For comparison, we also followed the corresponding absorbance changes after salt addition (Fig. 5 B, full circles) and plotted the slope of the curves depicted in Fig. 4 B–E as a function of time (Fig. 5 B, open circles). Curves on Fig. 5 B are similar in shape. In comparison with the curve of V as a function of time (Fig. 5 A), the attainment of the plateau is faster for curves on Fig. 5 B.

Osmotic gradients induced time-dependent absorbance changes in fused DODAC vesicles. When intravesicular NaCl concentration was higher than the external concentration, absorbance decreased with time (Fig. 6 A–F). When the concentration of impermeable solutes was higher in the external solution absorbance increased with time (Fig. 6 G–P).

FIGURE 5 Internal volume (V) of sonicated DODAC vesicles (2.5 mM) prepared in 20 mM sucrose as a function of time after NaCl addition (0.042 M) (○) (A). (^{14}C)-sucrose was added to the sonicated vesicles prior to salt addition. At zero time, salt was added to the sonicated vesicles. Gel chromatography separated entrapped and free (^{14}C)-sucrose for calculation of V values. For comparison, the V value obtained from electron microscopy assuming $40 \text{ \AA}^2$ as the area per monomer is also shown (◊) (A). Time-dependent absorbance changes of the same preparation of sonicated DODAC vesicles after salt addition (0.042 M) (●) and slopes against time for "spectra" from Fig. 4 (○) are in B.

The initial shrinking rate (v_0 %) varied linearly with NaCl concentration gradient ($\Delta C \text{ NaCl}$) displaying two different slopes depending on the sign of the concentration gradient (Fig. 7 A). This difference could be associated with an increased permeability of the DODAC vesicle under hypotonic condition. The actual osmotic gradient would be smaller than that at time zero due to partial leakage of the contents of the vesicles as in the case of phospholipid liposomes (23). A linear relationship between v_0 % and ΔO (ER. + NaCl) was obtained for the absorbance changes induced by NaCl and erythritol (Fig. 7 C). $A_0/\Delta A$ varied linearly with the reciprocal of the NaCl concentration gradient (Fig. 7 B) or the NaCl plus erythritol osmolarity gradient (Fig. 7 D).

For sonicated DHP vesicles, time-dependent absorbance changes after salt addition for a range of salt (Fig. 8 A) or DHP concentration (Fig. 8 B) are similar to those for large DHP vesicles (Fig. 8 C). In these curves, the absorbance always increases as a function of time and no plateau is attained. Visual observations after 24 or 48 h showed particles in suspension that later precipitated.

DHP dispersions obtained after NaCl addition to sonicated vesicles, when examined by electron microscopy, showed small sonicated vesicles, aggregates of small sonicated vesicles and some large vesicles (Fig. 9). To confirm that the DHP vesicle population 24 h after salt addition was composed mainly of aggregates of small vesicles, we tested the dispersion behavior toward osmotic gradients. During the first 10 min following the osmotic shock no significant absorbance change could be detected. 24 h thereafter the absorbance was much higher than that expected if only shrinkage or swelling had occurred.

FIGURE 6 Time-dependent absorbance changes due to osmotic gradients imposed to DODAC vesicles at least 24 h after salt addition (0.057 M). Sonicated DODAC vesicles (10 mM) were prepared in water, NaCl was added to a final concentration of 0.057 M and, after 2 h, aliquots of this mixture (0.2 ml) were added to 0.8 ml water (A) or NaCl solution with the following concentration (in mM): 8 (A); 16 (B); 27 (C); 37 (D); 46 (E); 53 (F); 60 (G). For shrinking, 0.2 ml of the mixture containing the vesicles were added to 0.8 ml aqueous solutions containing 57 mM NaCl and erythritol in the following concentrations (in mM): 20 (H); 40 (I); 50 (J); 60 (K); 70 (L); 80 (M); 90 (N); 100 (O); 120 (P).

DISCUSSION

DODAC and DHP vesicles are suitable models for interfacial effects on chemical reactions (3), for permeability and transport studies (17, 18) and for membrane interactions induced by salts (11). In this section we show that the study of DODAC or DHP vesicles contributes to a better understanding of the relationship between monomer-structure and intervesicular interactions leading to membrane fusion.

It has been often pointed out that light scattering or turbidity measurements are unable to distinguish between vesicle aggregation and fusion (24–26). An analysis of the relative contributions of these processes using fundamental light-scattering theory in the context of a simple model for aggregation and fusion is lacking. Light scattering can follow the sample transformation without perturbation, an advantage which is absent for methods relying on the analysis of aliquots (electron microscopy, trapped volume, ultracentrifugation or gel chromatography) (26). Here we will try to overcome interpretative difficulties of turbidity data and propose a phenomenological model that is able to account for the scatter-derived absorbance changes observed for sonicated and large DODAC or DHP vesicles after NaCl addition (Fig. 1 and Fig. 8).

For spherical particles with a diameter D very small compared with the wavelength of light ($D \ll \lambda$), the

FIGURE 7 Osmometer behavior of a preparation containing sonicated DODAC vesicles in the presence of salt. Osmotic gradients were generated by NaCl or by NaCl plus erythritol and time-dependent absorbance changes as those in Fig. 6 were recorded permitting calculations of osmometer parameters. Initial swelling or shrinking rates (v_o %) were plotted against the NaCl concentration gradient (ΔC NaCl) (A) or against the NaCl plus erythritol osmolarity gradient (ΔO NaCl + ER) (C). The extension of swelling or shrinkage ($A_o/\Delta A$) was plotted against the reciprocal of the concentration gradient of NaCl ($1/\Delta C$ NaCl) (B) or against the reciprocal of the osmolarity gradient of NaCl plus erythritol ($1/\Delta O$ NaCl + ER) (D).

scatter-derived absorbance (A) is given by (27)

$$A = [32\pi^3 / (2.3 \cdot 3 \cdot n_o^2)] \cdot (dn/dc)^2 \cdot q^2 \nu \lambda^{-4} \quad (1)$$

where dn/dc is the specific refractive index increment, n_o the refractive index of the medium, q the anhydrous mass of the scattering particle, ν the number of particles per unit volume, and λ the wavelength of the incident radiation. Assuming that dn/dc does not change upon aggregation, the ratio between the absorbance before (A_o) and after (A_i) aggregation is simply

$$A_i/A_o = i, \quad (2)$$

where i is the number of vesicles in the aggregate. Upon aggregation, the absorbance will increase as long as the effective size of the aggregate remains lower than λ .

If the particles are not small with respect to λ , phase changes of the light will occur within the particle and the scattered light will interfere (28). For particles of dimensions comparable to λ ($D \approx \lambda$) and index of refraction similar to that of the medium, the Jobst law can be applied (29)

$$A = (5.984/n_o^2) (dn/dc)^2 q^2 \nu R^{-2} \lambda^{-2}, \quad (3)$$

where R is the particle radius.

FIGURE 8 Scatter-derived absorbance changes induced by NaCl on small (A, B) and large (C) DHP vesicles. In A, DHP concentration is 0.57 mM and salt concentration (in mM) is 53.7 (a), 85.0 (b), and 132.2 (c). In B, NaCl concentration is 110 mM and DHP concentration (mM) is 0.23 (a); 0.34 (b), and 0.57 (c). In C, large DHP vesicles (0.62 mM) prepared in 0.430 M D-glucose are in NaCl concentrations (mM) of: 70 (a); 127 (b); 153 (c); and 171 (d).

In this range of particle size the absorbance is linearly dependent on λ^{-2} (Eq. 3 and Fig. 4 B–D). Under these conditions, an increase in the slope of the function relating A with λ^{-2} implies that the anhydrous mass increase is larger than that of the particle size (R). The A_i/A_o value will depend exclusively upon the ratio between the number of aggregated particles (i) and the square of the ratio between the radius of the original particle (r) and that of the aggregate (R')

$$A_i/A_o = i (r/R')^2. \quad (4)$$

From Eq. 4, a decrease in absorbance ($A_i < A_o$) is due to aggregation of large vesicles ($i < 4$). This interpretation is unambiguous since fusion of large vesicles in an aggregate with radius r (giving a fused vesicle with radius R') will necessarily imply $R' < r$ and $A_i > A_o$. Both statements can be easily verified using Eq. 4 and geometrical calculations of r and R' for symmetrical aggregated or fused forms.

On the other hand, if $D \ll \lambda$, Eq. 1 must be applied and the independence of A on the particle size implies that fusion of vesicles composing an aggregate cannot be detected and an absorbance increase necessarily results from aggregation of small vesicles.

Using the condition of validity of the Jobst approxima-

FIGURE 9 Electron microscopy of a positively stained preparation of sonicated DHP vesicles (2.5 mM) 48 h after NaCl addition (0.042 M).

tion ($D \approx \lambda$), the particle diameter must be within 4,000 to 8,000 Å (visible range). Fused vesicles possessing diameters in the range of 4,000 to 8,000 Å may be thought as the fusion product of n vesicles with radius r (in Å), where n varies between $(2,000/r)$ and $(4,000/r)$. Therefore, the attainment of the absorbance decrease phase (Fig. 1) due to aggregation of large vesicles depends on the mutual approach, adhesion and fusion of 169–679 small vesicles ($r = 150$ Å) or 4–16 large vesicles ($r = 2,000$ Å). This analysis makes reasonable the retarded absorbance decrease phase for sonicated DODAC vesicles after salt addition as compared with the rapid attainment of the same phase for the large DODAC vesicles (Fig. 1 *D*).

The possibilities of the model may be further explored by supposing small vesicles (~ 150 Å mean diameter) which only aggregate. An absorbance increase due to small vesicle aggregation will be observed (Eq. 2). As the aggregate size increases and the dimension of the aggregate compares with λ , further aggregation of small vesicles yielding a large aggregate would also result in an absorbance increase (from Eq. 4 and simple geometrical calculations of R' by supposing for example, the formation of cubic aggregates with 27, 64, 125 small vesicles with radius r). Thus, the absorbance, for a small vesicle system which undergoes only aggregation, increases progressively with time. Aggregates can grow until the formation of flocs in suspension yields first an oscillation in absorbance (Fig. 8 *C*, curve *d*) and finally precipitation as was shown for DHP vesicles after salt addition (Fig. 8 *A–C*).

Eqs. 1 and 3 can also be analyzed with respect to osmotic effects in vesicular systems. Swelling or shrinkage of small vesicles ($D \ll \lambda$) due to osmotic gradients do not alter the

absorbance (17, 30), as predicted from Eq. 1. On the other hand, particle size variations due to osmotic effects on large vesicles ($D \approx \lambda$) produce an absorbance increase due to shrinkage and an absorbance decrease due to swelling (31, 32). Small aggregated vesicles should not exhibit absorbance variations due to osmotic gradients but large fused vesicles should. On this basis, we propose a method for detection of fusion and discrimination between aggregation of small vesicles and fusion based on the determination of osmometer behavior. Sonicated DODAC vesicles, 24 h after salt addition, behaved as an osmometer towards osmotic gradients of NaCl or NaCl and erythritol (Fig. 6 and 7), a fact which strongly suggests the presence of large fused vesicles in significant proportions. The lack of osmotic response in sonicated DHP vesicles 24 h after salt addition shows that large fused vesicles are not present in significant proportions. Both conclusions were confirmed by the electron micrographs (Figs. 2 and 9). DHP sonicated vesicles fuse to some extent (11) after salt addition as shown by the decay of the excimer-monomer ratio of a lipid fluorescent probe (19). However, both the shape of the time-dependent absorbance changes after salt addition (Fig. 8) and the absence of osmometer behavior suggest that fusion is not frequent.

Below the phase transition temperature (T_c), the occurrence of aggregation only or of aggregation followed by fusion will depend on the vesicle concentration range. For dipalmitoylphosphatidylcholine (DPPC) (4°C) in the 0.5–5.0 mM (Pi) concentration range, small vesicles aggregate and the size of the aggregates increases linearly with time (33). In the 20–120 mM Pi range, small DPPC vesicle aggregation is followed by fusion (33). Analogously, fusion of sonicated DHP vesicles will possibly be more frequent at DHP concentrations higher than those used in this study. Furthermore, for DPPC vesicles (4°C and 0.5–5.0 mM Pi) aggregation increases with time until the aggregates precipitate (28), in close analogy with the behavior of our sonicated DHP vesicles after salt addition.

Eqs. 1 to 4, associated with simple geometrical calculation, lead to interpretations summarized in Table I.

Particles formed upon NaCl addition to sonicated DODAC vesicles are shown, by a variety of methods (Figs. 3 to 7 and reference 11), to be large vesicles. For this to be the case, it is necessary for aggregation to be faster than fusion. That this is in fact the case can be seen by comparing the time-course of the scatter-derived absorbance increase and of the slope increase (Fig. 5 *B*), which reflect both fusion and aggregation, with the time-course of fusion alone, as assessed by the internal volume (V) increase (Fig. 5 *A*). For example, 2 h after salt addition, V is ~ 9 l/mol, i.e., only $\sim 40\%$ of the value attained after 24 h, whereas the absorbance value is 0.47, i.e., 87% of the value at 24 h. Therefore, at 2 h, the small sonicated vesicles were aggregated, in addition to being $\sim 40\%$ fused.

Sonicated DODAC vesicles exhibit a relatively broad phase transition temperature range when compared with

TABLE I
PHENOMENA ASSOCIATED WITH TIME-DEPENDENT
ABSORBANCE CHANGES FOR INTERACTING VESICLES
IN ACCORDANCE WITH THE SCATTERING LAWS

Absorbance behavior	Vesicle diameter (D)	
	$D \ll \lambda$	$D \approx \lambda$
Increase	1. Aggregation of small vesicles.	1. Fusion of large vesicles composing a large aggregate.
	2. Small aggregates further aggregating small vesicles.	2. Fusion of small vesicles composing a large aggregate.
	3. Disaggregation of large vesicles from large vesicle aggregates.	3. Shrinkage of large vesicles.
Decrease	1. Disaggregation of small vesicles.	1. Aggregation of large vesicles ($i < 4$).
	2. Vesicles yielding micelles.	2. Swelling of large vesicles.
Invariacy	1. Fusion of small vesicles composing a small aggregate.	
	2. Shrinkage or swelling of small vesicles.	

large DODAC vesicles (17). Seven hours after salt addition both the phase transition temperature range and the T_c values of salt-treated sonicated DODAC vesicles (Fig. 3) were very similar to those of large DODAC vesicles (17), a fact that suggests size transformation. The difference in the absorbance value at 25°C before and after the heating-cooling cycle at 7 h (Fig. 3) could a priori be explained as disaggregation of small aggregated vesicles or as aggregation of large fused vesicles. For disaggregation to be the case, it could be expected that during the hours following the heating-cooling cycle the absorbance should again increase. Since the absorbance value remained unchanged for, at least, 24 h after the cycle (see Results), the absorbance decrease upon heating and cooling can be explained by a temperature induced fusion of small aggregated vesicles resulting in large vesicles that could aggregate during the cooling process.

The Derjaguin-Landau-Verwey-Overbeek (DLVO) theory for colloid stability is based upon long-range interactions due to electrostatic and electrodynamic (van der Waals) forces (34). It has been shown, both for lipids and synthetic amphiphiles, that the DLVO could not account for the relative stabilities of small and large vesicles in the presence of salt (6, 11). The DLVO neglects important short-range interparticle interactions such as those due to water structure around the polar headgroup (35). As a consequence, in spite of the role played by electrostatic repulsion and van der Waals attraction for the establishment of short-range interactions, hydration repulsion can be the limiting barrier for the fusion process. Since the DODAC molecule has two methyl groups in the cationic polar head, one may reasonably suppose that the surround-

ing hydration water is less tightly bound than that around a phosphate head group. Therefore, the hydration repulsive force at close approach due to water structure is lower for DODAC than for DHP vesicles in agreement with the higher frequency of fusion observed for the former. As the DPPC molecule possesses both the quaternary ammonium and the phosphate groups, one could speculate that the surrounding hydration shell of phospholipid bilayers (36) is more structured in comparison with that of the same separate groups in our synthetic bilayers, an hypothesis that partially accounts for the experimentally established stability of phospholipid vesicles in the range of 0.01–0.2 M monovalent salts.

In addition to the hydration shell around each head group, other factors can be involved in the short-range repulsion between phospholipid bilayers, such as a polymer-type steric repulsion due to the thermal mobility of the head groups (37). Since the head groups of DODAC or DHP are less bulky than those of lecithin, the steric short-range repulsion should be smaller for DODAC or DHP sonicated vesicles, leading to a higher frequency of adhesion and fusion under comparable conditions.

Finally, we will try to compose an energetic picture which could qualitatively describe interactions between DODAC or DHP vesicles during their mutual approach. First, calculated Stern potential values for sonicated DODAC vesicles are smaller than those for sonicated DHP vesicles (11), a fact that allows us to assume an energy barrier of repulsion (E) for sonicated DODAC vesicles that is lower than the corresponding barrier for sonicated DHP vesicles. Second, since aggregation occurs preferentially for sonicated DHP vesicles and fusion for sonicated DODAC vesicles, the supposition of a deeper attractive energy minimum (M) for sonicated DHP vesicles is justifi-

FIGURE 10 Hypothetical picture for the energy of interaction as a function of the separation distance (d) between sonicated DODAC or DHP vesicles. The positive sign refers to repulsive, and the negative, to attractive energies. The presence of energy barriers (E and H) and energy minimum (M) are justified in the text.

ied. Third, from simple structural considerations, the repulsive energy barrier (H) at zero separation distance due to hydration of the polar heads should be smaller for DODAC than for DHP. On these grounds, we propose the qualitative energy diagram in Fig. 10, which differentiates a configuration leading to fusion from another leading to aggregation.

We gratefully acknowledge Mr. Hélio Brandão and Dr. Antonio Sesso for the electron microscopy and Dr. Mari Armelin for careful revision of the manuscript. We thank also one of the referees for pertinent references and comments.

This work was supported by the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) and Financiadora de Estudos e Projetos (FINEP).

Received for publication 20 December 1985 and in final form 9 April 1986.

REFERENCES

- Gomperts, B. D. 1977. Models for membrane structure. In *The Plasma Membrane: Models for Structure and Function*. H. B. Jovanovich, editor. Academic Press, Inc., London. 1-49.
- Fendler, J. H. 1980. Surfactant vesicles as membrane mimetic agents: characterization and utilization. *Acc. Chem. Res.* 13:7-13.
- Chaimovich, H., J. B. S. Bonilha, D. Zanette, and I. M. Cuccovia. 1984. Analysis of the effect of micelles and vesicles on the reactivity of nucleophiles derived from the dissociation of weak acids. In *Surfactants in Solution*. K. L. Mittal and B. Lindman, editors. Plenum Publishing Corp., New York and London. 1121-1138.
- Dimitrov, D. S., D. V. Zhlev, and R. K. Jain. 1985. Stability of membrane systems modeled as multilayered viscoelastic films. *J. Theor. Biol.* 113:353-377.
- Nir, S., and J. Bentz. 1978. On the force between phospholipid bilayers. *J. Colloid Interface Sci.* 65:399-414.
- Nir, S., J. Bentz, and N. Duezguenes. 1981. Two modes of reversible vesicle aggregation: particle size and the DLVO theory. *J. Colloid Interface Sci.* 84:266-269.
- Bentz, J., and S. Nir. 1981. Mass action kinetics and equilibria of reversible aggregation. *J. Chem. Soc. Faraday Trans. I.* 77:1249-1275.
- Ohki, S., N. Duezguenes, and K. Leonards. 1982. Phospholipid vesicle aggregation: effect of monovalent and divalent ions. *Biochemistry*. 21:2127-2133.
- Rydhag, L., K. Rosenquist, P. Stenius, and L. Oedberg. 1983. Colloidal stability of liposomes. In *Physical Chemistry of Surfactants in Solution*. K. L. Mittal and E. J. Fendler, editors. Plenum Publishing Corp., New York.
- Ohki, S., S. Roy, H. Ohshima, and K. Leonards. 1984. Monovalent cation-induced phospholipid vesicle aggregation: effect of ion binding. *Biochemistry*. 23:6126-6132.
- Carmona-Ribeiro, A. M., L. S. Yoshida, and H. Chaimovich. 1985. Salt effects on the stability of dioctadecyldimethylammonium chloride and sodium dihexadecylphosphate vesicles. *J. Phys. Chem.* 89:2928-2933.
- Rupert, L. A. M., D. Hoekstra, and J. B. F. N. Engberts. 1985. Fusogenic behaviour of didodecyldimethylammonium bromide bilayer vesicles. *J. Am. Chem. Soc.* 107:2628-2631.
- Schales, O., and S. S. Schales. 1941. A simple and accurate method for the determination of chloride in biological fluids. *J. Biol. Chem.* 140:879-884.
- Houser, G., S. Fleischer, and A. Yamamoto. 1970. Two dimensional thin layer chromatography separation of polar lipids and determination of phospholipids by phosphorus analysis of spots. *Lipids*. 5:494-496.
- Tran, C. D., P. L. Klahn, A. Romero, and J. H. Fendler. 1978. Characterization of surfactant vesicles as potential membrane models. Effect of electrolytes, substrates and fluorescent probes. *J. Am. Chem. Soc.* 100:1622-1624.
- Mortara, R. A., F. H. Quina, and H. Chaimovich. 1978. Formation of closed bilayers from a single phosphate diester. Preparation and some properties of vesicles of dihexadecyl phosphate. *Biochem. Biophys. Res. Commun.* 81:1080-1086.
- Carmona-Ribeiro, A. M., and H. Chaimovich. 1983. Preparation and characterization of large dioctadecyldimethylammonium chloride liposomes and comparison with small sonicated vesicles. *Biochim. Biophys. Acta.* 733:172-179.
- Carmona-Ribeiro, A. M., L. S. Yoshida, A. Sesso, and H. Chaimovich. 1984. Permeabilities and stability of large dihexadecylphosphate and dioctadecyldimethylammonium chloride vesicles. *J. Colloid Interface Sci.* 100:433-443.
- Schenkman, S., P. S. Araujo, R. Dijkman, F. H. Quina, and H. Chaimovich. 1981. Effects of temperature and lipid composition on the serum albumin-induced aggregation and fusion of small unilamellar vesicles. *Biochim. Biophys. Acta.* 649:633-641.
- Morgan, C. G., H. Williamson, S. Fuller, and C. Hudson. 1983. Melittin induces fusion of unilamellar phospholipid vesicles. *Biochim. Biophys. Acta.* 732:668-674.
- Wilschut, J., and D. Papahadjopoulos. 1979. Ca^{2+} -induced fusion of phospholipid vesicles monitored by mixing of aqueous contents. *Nature (Lond.)*. 281:690-692.
- Herrmann, U., and J. H. Fendler. 1979. Low angle laser light scattering and photon correlation spectroscopy in surfactant vesicles. *Chem. Phys. Lett.* 64:270-274.
- Yoshikawa, W., H. Akutsu, and Y. Kyogoku. 1983. Light-scattering properties of osmotically active liposomes. *Biochim. Biophys. Acta.* 735:397-406.
- Chong, C. S., and K. Colbow. 1976. Light scattering and turbidity measurements on lipid vesicles. *Biochim. Biophys. Acta.* 436:260-282.
- Koch, A. 1961. Some calculations on the turbidity of mitochondria and bacteria. *Biochim. Biophys. Acta.* 51:429-441.
- Schmidt, C. F., D. Lichtenberg, and T. E. Thompson. 1981. Vesicle-vesicle interactions in sonicated dispersions of dipalmitoylphosphatidylcholine. *Biochemistry*. 20:4792-4797.
- Lord Rayleigh. 1918. On the scattering of light by shells. *Proc. R. Soc. Lond. A. Math. Phys. Sci.* 94:296-300.
- Mie, G. 1908. Beitrage zur optik trueber Medien steziell colloidalen Metaloesungen. *Ann. Physik.* 25:377-445.
- Joebst, G. 1925. Strahlung dielektrischer Kugel in Grenzsalle das Kugelmaterial und umgebendes Medium fast gleich Brechungsfindiceshaben. *Ann. Physik.* 78:157-166.
- Johnson, S. M., and N. Buttress. 1973. The osmotic insensitivity of sonicated liposomes and the density of phospholipid-cholesterol mixtures. *Biochim. Biophys. Acta.* 307:20-26.
- Reeves, J. P., and R. M. Dowben. 1970. Water permeability of phospholipid vesicles. *J. Membr. Biol.* 3:123-141.
- Bangham, A. D., J. de Gier, and G. D. Greville. 1967. Osmotic properties and water permeability of phospholipid liquid crystals. *Chem. Phys. Lipids.* 1:225-246.
- Wong, M., and T. E. Thompson. 1982. Aggregation of dipalmitoylphosphatidylcholine vesicles. *Biochemistry*. 21:4133-4139.
- Verwey, E. J., and J. T. G. Overbeek. 1948. Theory of the Stability of Lyophobic Colloids. Elsevier, New York.
- Ninham, B. 1981. Surface forces: the last 30 Å. *Pure Appl. Chem.* 53:2135-2147.
- Le Neveu, D. M., R. P. Rand, V. A. Parsegian, and D. Gingell. 1977. Measurement and modification of forces between lecithin bilayers. *Biophys. J.* 18:209-230.
- Marra, J., and J. Israelachvili. 1985. Direct measurements of forces between phosphatidylcholine and phosphatidylethanolamine bilayers in aqueous electrolyte solutions. *Biochemistry*. 24:4608-4618.

Referência 10

Dihexadecyl Phosphate Monolayers: Intralayer and Interlayer Interactions

Per Claesson,^{*,†,§} Ana Maria Carmona-Ribeiro,^{†,§} and Kazue Kurihara^{†,‡}

Departamento de Bioquímica, Instituto de Química, CP 20780, Universidade de São Paulo, São Paulo, Brazil; and Institute for Surface Chemistry, Box 5607, S-11486 Stockholm, Sweden, and Department of Physical Chemistry, The Royal Institute of Technology, S-10044 Stockholm, Sweden (Received: May 3, 1988)

The interactions between deposited monolayers of dihexadecyl phosphate (DHP), immersed in aqueous NaCl solutions, have been studied. At pH 5–6 double-layer forces, well-described by Poisson–Boltzmann theory, were observed. The magnitude of the repulsive forces demonstrated that less than 2% of the molecules were charged. At small separations ($D < 2$ nm) attractive forces, which decreased with increasing NaCl concentration and pH, were present. At higher pH, a repulsive hydration force overcame the van der Waals force completely. Intralayer forces were investigated by recording pressure–area isotherms for DHP and for dioctadecyldimethylammonium bromide (DODABr). DHP formed a gaseous, a liquid condensed, and a solid condensed phase. When the pH or the NaCl concentration was increased, the liquid condensed phase became more important; thereby the area per molecule for a given surface pressure increased. We suggest that this is due to sodium ions replacing protons as counterions, which gives rise to an intralayer hydration force. The DODA monolayer behaved differently, and the area per molecule for a given surface pressure decreased when the NaCl concentration was increased. The same types of forces that operate between headgroups within a monolayer of DHP are also of importance for the interactions between such monolayers. The magnitude of some of these forces depends on the type of counterion.

Introduction

The behavior and the properties of many colloidal and biochemical systems, such as micelles, vesicles, microemulsions, and particles stabilized by surfactants, are governed by the forces acting between and within surfactant layers. For instance, from a biochemical point of view membrane–membrane, vesicle–vesicle, and vesicle–membrane interactions are the basis for all phenomena involving membrane coupling such as adhesion, juxtaposition, and fusion.

Interactions between phospholipid model membranes have been extensively studied.^{1–14} The technique of Parsegian and co-workers,^{1–3} and that of Israelachvili and co-workers,^{12–15} have successfully been used for demonstrating the existence of repulsive forces not accounted for by the conventional DLVO theory. These forces are commonly referred to as hydration forces. They are believed to be caused by changes in water–headgroup interactions occurring as a result of decreasing aggregate or surface separation.

Synthetic membranes composed of dioctadecyldimethylammonium chloride (DODACl) or dihexadecyl phosphate (DHP) have previously been characterized and compared with phospholipid systems.^{16,17} Vesicles composed of DODACl or of DHP behave similar to those of phospholipids in many respects, but in contrast to phospholipid vesicles they are unstable in the presence of NaCl.¹⁸ In 50 mM NaCl solutions DODACl vesicles mainly fuse, whereas DHP vesicles mainly aggregate.¹⁹

Strongly repulsive electrostatic double-layer forces dominate the interaction between mica surfaces coated with deposited bilayers of dioctadecyldimethylammonium ions²⁰ or with adsorbed bilayers of dihexadecyldimethylammonium acetate.²¹ This investigation aimed at characterizing the forces acting between and within DHP layers and comparing them with those present in the corresponding DODA system. A Langmuir trough force balance was employed for determining the intralayer forces as a function of the mean molecular area in monolayers of DHP at an air–water interface. The effects of changing the pH and the NaCl concentration in the subphase were explored. The forces acting between two DHP monolayers deposited onto hydrophobed mica (mica coated with a deposited monolayer of DODA) were directly measured with a surface force apparatus of the type developed

at the Australian National University by Israelachvili and co-workers.¹⁵ These interactions were also determined for a range of NaCl concentrations and at different pH values.

The results show very clearly that the forces operating in and between DHP layers are quite different from those acting in and between layers of DODA. These differences rationalize why DHP vesicles mainly aggregate at high salt concentrations whereas DODA vesicles tend to fuse.

Materials and Methods

Chemicals. The dihexadecyl phosphate used for surface force measurements was obtained in acid form from Sigma Chemical Co. and used as received. The DHP (in acid form) and the DODABr used in the Langmuir trough experiments were obtained from Sogo and used as received. The NaCl was of ultrapure grade and obtained from Merck. Green muscovite mica was received from Brown Mica Co., Sydney.

The surface force measurements were carried out in Canberra whereas the surface balance experiments were performed in Stockholm, explaining why two different water purification pro-

- (1) LeNeveu, D. M.; Rand, R. P.; Parsegian, V. A. *Nature (London)* **1976**, 259, 601.
- (2) LeNeveu, D. M.; Rand, R. P.; Parsegian, V. A.; Gingell, D. *Biophys. J.* **1977**, 18, 209.
- (3) Cowley, A. C.; Fuller, N. L.; Rand, R. P.; Parsegian, V. A. *Biochemistry* **1976**, 17, 3163.
- (4) Nir, S.; Bentz, J. J. *Colloid Interface Sci.* **1978**, 65, 399.
- (5) Lis, L. J.; Parsegian, V. A.; Rand, R. P. *Biochemistry* **1981**, 20, 1761.
- (6) Lis, L. J.; Lis, W. T.; Parsegian, V. A.; Rand, R. P. *Biochemistry* **1981**, 20, 1771.
- (7) Rand, R. P. *Annu. Rev. Biophys. Bioeng.* **1981**, 10, 277.
- (8) Nir, S.; Bentz, J.; Düzgünes, N. J. *Colloid Interface Sci.* **1981**, 84, 266.
- (9) Lis, L. J.; McAlister, M.; Fuller, N.; Rand, R. P.; Parsegian, V. A. *Biophys. J.* **1982**, 37, 657.
- (10) Afzal, S.; Tesler, W. J.; Blessing, S. K.; Collins, J. M.; Lis, L. J. *J. Colloid Interface Sci.* **1984**, 97, 303.
- (11) Ohki, S.; Roy, S.; Ohshima, H.; Leonards, K. *Biochemistry* **1984**, 23, 6126.
- (12) Marra, J.; Israelachvili, J. N. *Biochemistry* **1985**, 24, 4608.
- (13) Marra, J. J. *Colloid Interface Sci.* **1986**, 109, 11.
- (14) Horn, R. G. *Biochim. Biophys. Acta* **1984**, 778, 224.
- (15) Israelachvili, J. N.; Adams, G. E. *J. Chem. Soc., Faraday Trans. 1* **1978**, 74, 975.
- (16) Carmona-Ribeiro, A. M.; Chaimovich, H. *Biochim. Biophys. Acta* **1983**, 733, 172.
- (17) Carmona-Ribeiro, A. M.; Yoshida, L. S.; Sesso, L. S.; Chaimovich, H. J. *Colloid Interface Sci.* **1984**, 100, 433.
- (18) Carmona-Ribeiro, A. M.; Yoshida, L. S.; Chaimovich, H. *J. Chem. Phys.* **1985**, 82, 2328.
- (19) Carmona-Ribeiro, A. M.; Chaimovich, H. *Biophys. J.* **1986**, 50, 621.
- (20) Marra, J. J. *J. Phys. Chem.* **1986**, 90, 2145.
- (21) Pashley, R. M.; McGuiggan, P. M.; Ninham, B. W.; Brady, J.; Evans, D. F. *J. Phys. Chem.* **1986**, 90, 1637.

*To whom correspondence should be addressed.

†Universidade de São Paulo.

‡Institute for Surface Chemistry and the Royal Institute of Technology.

§Temporary address: Department of Applied Mathematics, Research School of Physical Sciences, Institute of Advanced Studies, The Australian National University, G.P.O. Box 4, ACT 2600, Australia.

‡ Present address: Research Development Corporation of Japan, 332 Kamikoga, Chihushino, Fukuoka 818, Japan.

cedures and two sources of DHP were used.

Water for the surface force experiments was doubly distilled and further purified with an ELGA UHQ unit, bypassing the reverse osmosis cartridge. The water was also presaturated with DHP in the acid form and deaerated for at least 4 h before being used in the surface force apparatus.

The water used for surface balance measurements was first purified by decalcination, prefiltration and reverse osmosis. The final purification was carried out by a modified MilliQ unit, which included two mixed bed ion exchangers, an activated charcoal cartridge, a 0.2- μm in-line filter cartridge, an Organex cartridge, and a final 0.2- μm filter. All filters were Zetapore products, whereas all other cartridges were from Millipore.

Surface Balance Measurements. Surface pressure-area isotherms were recorded by means of a computerized Langmuir trough system developed by KSV Chemicals, Helsinki. The trough is made from a single piece of Teflon whereas the barrier material is Delrin. This choice of materials ensures that no molecules leak under the slightly hydrophilic barrier.

The compression of the monolayer was carried out at a speed of 5 mm/min, which corresponds to a reduction in area per molecule of about 1.4 $\text{\AA}^2/\text{min}$. The surface pressure was continuously recorded with a precision of 0.01 mN/m, and the data were stored by the computer. Each isotherm was measured on a newly spread monolayer. For different monolayers spread from the same DHP stock solution the area per molecule at surface pressures larger than a few mN/m was reproducible to about 0.5 $\text{\AA}^2$. Between different stock solutions the variation was slightly larger.

Surface Preparations. Mica surfaces were first coated with a monolayer of DODA and then with a monolayer of DHP by means of a Langmuir-Blodgett technique. The all-Teflon trough used in Canberra was different from that used for the recording of surface pressure-area isotherms. It has been described by Marra.²² It was not equipped with any facility for keeping the surface pressure constant during deposition, but the barrier was moved with a predetermined speed in order to keep the surface pressure nearly constant.

First, a deposition of DODA was carried out from a pure water subphase at a surface pressure of 25 mN/m. The advancing water contact angle after deposition was 93–96°. Previous surface force measurements have shown that DODA-coated mica surfaces prepared under these conditions are uncharged.^{23,24} On each hydrophobic surface a monolayer of DHP was deposited at a surface pressure of 20 mN/m, again from a pure water subphase. The orientation of the second layer was such that the hydrophilic phosphate groups were directed toward the solution. The transfer ratio for this deposition was about 0.95, giving an area per deposited molecule of 43–45 $\text{\AA}^2$. The bilayer-coated surfaces were kept under water while transferred to and mounted in the surface force apparatus.

The absence of any facility for keeping the surface pressure constant during deposition is not very important for deposition of positively charged DODA ions onto negatively charged mica surfaces.²⁵ By contrast, the properties of the subsequent layer of DHP are likely to be sensitive to deposition conditions. The reasons are that the πA isotherm is very steep and that no electrostatic attraction exists between the DHP molecules and the substrate. This may explain why the magnitude of the double-layer force, measured between deposited layers of DHP, varied somewhat between different experiments (up to 30%). For this reason we have chosen to report force curves obtained from only one experiment. However, it should be emphasized that the variations of the force and surface charge density with pH and salt concentration were the same in all experiments.

Surface Force Measurements. Interlayer interactions were measured with a standard surface force apparatus.¹⁵ This ap-

Figure 1. Force between layers of DHP deposited onto hydrophobed mica surfaces measured as a function of surface separation in a range of NaCl solutions. The solid lines are calculated force curves using the double-layer parameters given in Table I and a nonretarded Hamaker constant of 1×10^{-20} (J). The symbols represent forces measured in the following solutions: $\blacktriangle$, 2×10^{-4} M NaCl; $\square$, 1×10^{-3} M NaCl; $\circ$, 1×10^{-2} M NaCl (all at pH 5.2–5.7). $\bullet$ represents forces measured in 1×10^{-2} M NaCl at pH 9.5. In all cases the interaction occurred under close to constant charge conditions. At pH 5.2–5.7 a van der Waals force gave rise to an attraction at small separations. However, at pH 9.5 no attraction was observed at any separation. Note the logarithmic force scale and that the force has been normalized by the local geometric mean radius. The dashed line represents that part of the force curve that cannot be explained by the DLVO theory.

paratus allows the force between two mica substrate surfaces in a crossed cylinder configuration to be determined as a function of surface separation. The surface separation is determined to within 0.2 nm by using fringes of equal chromatic order.¹⁵ The force (F_c) is determined from the deflection of a spring supporting one of the surfaces. The force between crossed cylinders normalized by the local geometric mean radius (R), which is 1–2 cm, is related to the free energy of interaction between flat surfaces (Ω_f) via the Derjaguin approximation:^{26,27}

$$F_c/R = 2\pi\Omega_f \quad (1)$$

Whenever the gradient of the force with respect to the surface separation (dF_c/dD) exceeds the spring constant, the mechanical system is in an unstable region. This instability causes the surface separation to change spontaneously until the next stable region has been reached.²⁸ For instance, under the action of a van der Waals force the surfaces may jump from a small separation to molecular contact.

Results

Interlayer Interactions. The forces measured between two deposited monolayers of DHP immersed in a range of aqueous NaCl solutions at pH 5.2–5.7 are illustrated in Figure 1. The zero separation distance ($D = 0$) was in all cases defined as the position of the surfaces reached under a high load ($F_c/R \geq 10$ mN/m). At large separations repulsive double-layer forces dominate the interaction. The measured Debye length was in each case in good agreement with the value expected from Poisson-Boltzmann (PB) theory. A numerical solution to the nonlinear PB equation was used for calculating double-layer forces.²⁹ It was found that the interaction occurred at close to constant charge conditions, whereas forces calculated by assuming constant potential were less repulsive at short separations than the measured

(22) Marra, J. J. *Colloid Interface Sci.* **1985**, *107*, 446.

(23) Claesson, P. M.; Christenson, H. K. *J. Phys. Chem.* **1988**, *92*, 1650.

(24) Christenson, H. K.; Claesson, P. M. *Science* **1988**, *239*, 390.

(25) Herder, P. C.; Claesson, P. M.; Blom, C. E. J. *Colloid Interface Sci.* **1985**, *119*, 155.

(26) Derjaguin, B. V. *Kolloid-Z.* **1934**, *69*, 155.

(27) Israelachvili, J. N. *Intermolecular Forces with Applications to Colloid and Biological Systems*; Academic: London, 1985.

(28) Horn, R. G.; Israelachvili, J. N. *J. Chem. Phys.* **1981**, *75*, 1400.

(29) Chan, D. Y. C.; Pashley, R. M.; White, L. R. *J. Colloid Interface Sci.* **1980**, *77*, 283.

TABLE I: Measured Adhesion Force and Some Parameters Used for Fitting Double-Layer Forces Calculated in the Nonlinear Poisson-Boltzmann Approximation to Measured Forces

[NaCl], M	pH	Debye length, nm	area/charge, nm ²	adhesion force, mN/m
2×10^{-4}	5.5	20	62	-5.4
1×10^{-3}	5.5	8.5	31	-3.7
1×10^{-2}	5.5	3.0	26	-3.3
1×10^{-2}	3			-7.2
1×10^{-2}	9.5	3.0	16	0

forces. The area per charge varied from about 62 nm² in 2×10^{-4} M NaCl solutions to 26 nm² in 10^{-2} M solutions (Table I). The area per molecule was about 0.44 nm², and hence less than 2% of the molecules were charged under these conditions. The low surface charge density is consistent with the low surface pressure observed in the gaseous phase with the Langmuir trough (Figure 3d).

The attractive van der Waals force was important at separations less than 2 nm, and at small enough separations the total force became attractive. In the continuum theory approximation^{30,31} to the van der Waals force, this system should behave much as mica coated with a 4-nm-thick hydrocarbon layer. The nonretarded Hamaker constant for mica interacting across water is 2.2×10^{-20} (J), whereas the value for hydrocarbon interacting across water is 0.5×10^{-20} (J).²⁷ In order to calculate the effective Hamaker constant for our system, these values were used in the approximate three-layer formula given by Ninham and Parsegian.³² For a surface separation of 2 nm the effective Hamaker constant was calculated to be 0.6×10^{-20} (J). The measured attraction was slightly stronger, and the best fit by DLVO theory was obtained when the effective Hamaker constant, for surface separations of 1–3 nm, was assumed to be slightly above 1×10^{-20} (J). The experiment by Marra²⁰ also indicated that the attraction between DODA bilayers deposited on mica surfaces is well-described by assuming a nonretarded Hamaker constant of 1×10^{-20} (J).

It is no surprise that DLVO theory fails to quantitatively account for the measured forces. The fundamental reason for this is that the van der Waals force and the double-layer force are interdependent.^{33–37} Hence, the basic assumption of additivity made in the DLVO theory is incorrect. For instance, the zero-frequency contribution to the van der Waals force is exactly balanced by polarization charges (image charges),³⁷ and ion-ion correlations in the diffuse double layer give rise to an attractive force contribution.^{33–37} This means that surface charge densities and Hamaker constants deduced from force measurements by a conventional DLVO analysis, as, e.g., in this paper, are apparent values. (The true values might not be too far off.) The main reason for using the DLVO concept in discussing forces is that it is well-known and the calculations are easy to carry out, thereby facilitating straightforward comparisons between systems and discussions between scientists. The limitations of the theory should, however, be kept in mind.

A further complication is that charge regulation³⁸ may take place at small separations. Hence, double-layer forces calculated in the PB approximation by assuming interaction at constant charge may overestimate the forces at small separations. In classical DLVO calculations these simplifications may partly be

Figure 2. Normalized force between monolayers of DHP deposited onto hydrophobed mica surfaces as a function of surface separation. The solution contains 1×10^{-2} M NaCl. The forces measured at a pH of about 5.5 are represented by $\circ$, whereas forces measured at pH 9.5 are represented by $\bullet$. No attraction is observed at pH 9.5, but a weak minimum exists at pH 5.5. The solid lines represent forces calculated from DLVO theory using the double-layer parameters given in Table I and a nonretarded van der Waals force. The upper and lower curves for each pH are calculated by assuming a Hamaker constant of 0.5×10^{-20} (J) and 1.0×10^{-20} (J), respectively. The J indicates a jump into an adhesive minimum, and the dashed line represents non-DLVO forces.

TABLE II: Surface Pressure (Π) at the Transition from the Liquid Condensed (LC) to the Solid Condensed Phase (SC)

[NaCl], M	pH	Π , mN/m	comment
	3.5		no LC phase
	5.5	8	
	9.0		no SC phase
	9.9		no SC phase
10^{-4}	5–6	8–9	
10^{-3}	5–6	10–11	
10^{-2}	5–6	20	
10^{-1}	5–6	28	
10^{-2}	3.1	2–3?	no LC phase?
10^{-2}	8.4		no SC phase
10^{-2}	9.1		no SC phase

compensated for by assuming a too small surface charge density and a too large Hamaker constant.³⁴

The magnitude of the adhesion force, measured at a separation of 0.1–0.3 nm from final contact, decreased with increasing salt concentration and increasing pH (Table I). At pH 5.2–5.7 the magnitude of the adhesion force, normalized by the radius of the surfaces, decreased from 5.4 mN/m in 2×10^{-4} M NaCl to 3.3 mN/m in 10^{-2} M solutions.

When the pH was increased to 9.5, the forces measured in 10^{-2} M NaCl solutions became repulsive at all surface separations (Figures 1 and 2). The double-layer force increased in magnitude, and at small separations a hydration force overcame the van der Waals force and prevented any adhesive contact. The influence of the hydration force could be noticed out to a separation of about 1 nm (Figure 2).

At pH 3 a weak double-layer force was detected. It was not possible to measure this force very accurately, but the force barrier was 0.3 ± 0.2 mN/m. The adhesion force was accurately measured to be -7.2 mN/m.

Intralayer Interactions. Some pressure-area isotherms for DHP on subphases containing different concentrations of NaCl, and with varying pH, are shown in Figure 3a–c. At a pH of 5–6 the liquid condensed phase becomes more pronounced as the NaCl concentration increases. For instance, on a 10^{-4} M NaCl subphase

(30) Dzyaloshinskii, I. E.; Lifshitz, E. M.; Pitaevskii, L. P. *Adv. Phys.* **1961**, *10*, 165.

(31) Ninham, B. W.; Parsegian, V. A.; Weiss, G. H. *J. Stat. Phys.* **1970**, *2*, 323.

(32) Ninham, B. W.; Parsegian, V. A. *J. Chem. Phys.* **1970**, *52*, 4578.

(33) Kjellander, R.; Marcelja, S. *Chem. Phys. Lett.* **1984**, *112*, 49.

(34) Kjellander, R.; Marcelja, S. *J. Chem. Phys.* **1985**, *82*, 2122.

(35) Guldbrand, L.; Jönsson, B.; Wennerström, H.; Linse, P. *J. Chem. Phys.* **1984**, *80*, 2221.

(36) Attard, P.; Kjellander, R.; Mitchell, D. J. *Chem. Phys. Lett.* **1987**, *139*, 219.

(37) Attard, P.; Kjellander, R.; Mitchell, D. J. *J. Chem. Phys.*, in press.

(38) Chan, D.; Healy, T. W.; White, L. R. *J. Chem. Soc., Faraday Trans. 1* **1976**, *72*, 2844.

Figure 3. Some pressure–area isotherms recorded for DHP monolayers. (a) illustrates the variation with NaCl concentration at pH 5–6. Line 1 represents the situation with no added salt, whereas lines 2, 3, 4, and 5 represent NaCl concentrations of 1×10^{-4} , 1×10^{-3} , 1×10^{-2} , and 0.1 M, respectively. (b) depicts the variation with pH with no added salt in the subphase (other than NaOH or HCl). Lines 6, 7, and 8 illustrate the situation at pH 3.5, 9.0, and 9.9, respectively. The variation with pH on 1×10^{-2} M NaCl subphases is shown in (c). Lines 9, 10, 11, and 12 illustrate the situation at pH 3.1, 5.5, 8.4, and 9.1, respectively. Three phases are observed. The monolayer is always in a gaseous state when the area per molecule is larger than $48 \text{ \AA}^2$. The surface pressure in this region is illustrated in (d). Lines 13, 14, and 15 represent the situation on subphases containing 1×10^{-4} M NaCl at pH 5–6, 1×10^{-2} M NaCl at pH 5–6, and 1×10^{-2} M NaCl at pH 9–10. A transition from the gaseous phase to the liquid condensed phase takes place except at low pH where the monolayer goes directly from the gaseous phase to the solid condensed phase. No solid condensed phase is present at high pH.

the liquid condensed phase exists when the area per molecule is between 43 and $41 \text{ \AA}^2$. This corresponds to a surface pressure change from 1 to 8 mN/m . By contrast, the DHP monolayer spread on a 0.1 M NaCl subphase is in the liquid condensed phase when the area per molecule is 47 to $42 \text{ \AA}^2$. The corresponding surface pressures are 2 and 28 mN/m , respectively (Figure 3a).

Generally, one observes that the liquid condensed phase appears at a larger area per molecule and that the transition to the solid condensed phase occurs at a higher surface pressure, as the NaCl concentration is increased at a constant pH (Table II). However, the area per molecule for the liquid condensed to solid condensed phase transition remains approximately constant at $41\text{--}42 \text{ \AA}^2$.

For a monovalent electrolyte the electrostatic contribution to the surface pressure (Π_{EL}) is given by³⁹

$$\Pi_{\text{EL}} = (2kT\sigma)[(S^2 + 1)^{1/2} - 1]/(eS) \quad (2)$$

where k is the Boltzmann constant, T is the absolute temperature, e is the charge of the proton, σ is the surface charge density, and S is a dimensionless parameter defined by

$$S = \sigma / (8kTn\epsilon_r\epsilon_0)^{1/2} \quad (3)$$

where n is the number density of ions in bulk solution, ϵ_r is the relative dielectric constant of water, and ϵ_0 is the dielectric permittivity of vacuum.

The calculated variation of Π_{EL} with the area per charge under different solution conditions is illustrated in Figure 4.

In the gaseous phase at an area per molecule of about $50 \text{ \AA}^2$, the surface pressure for DHP monolayers on a 10^{-4} M NaCl subphase is $0.12 \pm 0.05 \text{ mN/m}$. The corresponding value on a 10^{-2} M NaCl subphase is $0.17 \pm 0.05 \text{ mN/m}$ at pH 5–6 and 0.28 ± 0.05 at pH 9–10 (Figure 3d). If the surface pressure is caused by electrostatic interactions only, the corresponding area per charge

Figure 4. Electrostatic contribution to the surface pressure, Π_{EL} , as a function of the area per charge. Curves 1, 2, and 3 correspond to a monolayer spread on a 1×10^{-4} , 1×10^{-2} , and 1 M monovalent electrolyte subphase, respectively. The main figure depicts the situation at large areas per charge and the insert that at small areas.

would be $40\text{--}70 \text{ nm}^2$ on a 10^{-4} M NaCl subphase, $14\text{--}20 \text{ nm}^2$ on a 10^{-2} M NaCl subphase at pH 5–6, and $11\text{--}14 \text{ nm}^2$ at pH 9–10 (Figure 4). These values are in fair agreement with, but slightly lower than, those obtained by fitting DLVO theory to the forces measured between DHP layers using the surface force apparatus (Table I). We conclude that the area per charge is indeed large and that the contribution to the surface pressure from interactions between hydrocarbon chains is small at this area per molecule. At a slightly smaller mean molecular area, when the monolayer no longer is in the gaseous phase, compression of hydrocarbon chains gives an important contribution to the surface pressure.

Figure 5. Some pressure–area isotherms for DODA monolayers spread on water or aqueous NaCl subphases. On a pure water subphase a liquid expanded phase exists when the area per molecule is less than 120 Å^2 (line 1). The situation for 1×10^{-4} , 1×10^{-3} , and $1 \times 10^{-2} \text{ M}$ NaCl subphases is shown by lines 2, 3, and 4. When NaCl is present in the subphase, an intermediate and a liquid condensed phase are also observed. With increasing NaCl concentration these phases appear at lower surface pressures. Upon compression the transitions to the intermediate phase and to the liquid condensed phase occur at an area per molecule of $80\text{--}85$ and $55\text{--}60 \text{ Å}^2$, respectively.

The shape of the DHP pressure–area isotherm depends on the pH. This is illustrated for the case of no added salt in Figure 3b and for the case of 10^{-2} M NaCl solutions in Figure 3c. At high pH no solid condensed phase is observed. By contrast, hardly any liquid condensed phase is present at low pH. This demonstrates that the extension of the liquid condensed phase, like the strength of the hydration force and the adhesion force between DHP layers, depends critically upon whether the phosphate headgroup is neutralized by protons or surrounded by a hydrated counterion like sodium.

The situation for DODA is completely different. A monolayer of DODA spread on a water subphase containing no added salt is in the liquid expanded state when the area per molecule is less than 120 Å^2 (Figure 5). No condensed or intermediate phase was observed in this case. However, with NaCl in the subphase a transition from a liquid expanded to an intermediate phase takes place at an area per molecule of about 80 Å^2 , whereas a liquid condensed phase is present at an area per molecule less than about 55 Å^2 . A study by Marra²⁰ has shown that the type of counterion influences the DODA surface–pressure area isotherm and hence the intralayer forces. Marra²⁰ and Pashley et al.²¹ also showed that the interlayer forces did depend on the type of counterion in a similar way. For instance, with acetate counterions the repulsive double-layer force was considerably stronger than with bromide. This was assumed to be due to an extensive binding of bromide ions to the quaternary ammonium headgroup.²¹

Discussion

The Stability of DHP Layers at the Air–Water Interface and on Hydrophobic DODA-Coated Mica Surfaces. The recording of a surface pressure–area isotherm is strictly speaking a non-equilibrium measurement. In the case of DHP the dissolution into the subphase is very slow, and the monolayer was stable for hours. The pressure–area curves recorded on compression and on decompression were nearly identical, indicating that the barrier speed was sufficiently low to ensure equilibrium within the monolayer.

The reproducibility in each experiment was satisfying, and the deposited DHP layer was stable for more than a day. However, the DHP layer was affected by bringing the surfaces into contact several times at the same position. This gave rise to a slightly lower double-layer force at large separations and a higher repulsion at small separations. The explanation is that the DHP monolayer has been disrupted locally and possibly that disordered multilayer structures have formed around the contact area. The forces reported in Figures 1 and 2 were measured under conditions when

no such disturbances of the DHP layers had been observed.

It was not possible to induce fusion of the DHP layers under any condition studied. By contrast, Pashley et al.²¹ found that bromide ions decreased the cohesion within bilayers of dihexadecyldimethylammonium ions sufficiently for allowing the outer layer to be squeezed out. Further, fusion of adsorbed layers of phospholipids has been reported by Horn¹⁴ and by Marra and Israelachvili,¹² who also suggested a fusion mechanism. It was noted by Marra and Israelachvili that loosely packed layers of phospholipids do fuse, whereas tightly packed layers do not.¹²

The Surface Charge Density. The area per deposited DHP molecule is about 0.44 nm^2 whereas the area per surface charge is about 60 nm^2 in $2 \times 10^{-4} \text{ M}$ NaCl at pH 5.2–5.7. In this case protons neutralize the charge of almost all phosphate groups. Also at higher salt concentrations the fraction of charged molecules appears to be small (Table I).

An attempt was made to, in the Poisson–Boltzmann model, explain the variation of the surface charge density with the NaCl concentration and the pH using the adsorption model of Pashley.²¹ It turned out that (an apparent?) agreement between experiments and predictions of the model could be obtained assuming that protons and sodium ions adsorb to the surface. Without any sodium ion adsorption the model predicts a more rapid increase in surface charge density with increasing pH and ionic strength than observed experimentally.

Adhesion Forces and Hydration Forces. The adhesion force decreases gradually as the NaCl concentration is increased at a pH of about 5.5. This shows that it becomes less favorable to bring two surfaces from a far distance into molecular contact. The reasons for this is that the surface charge density on isolated surfaces increases and that more sodium ions replace protons as counterions when the surfaces are in molecular contact. Note that the electroneutrality condition, which states that the surface charges are balanced exactly by the excess of counterions outside the surfaces, must also be valid when the surfaces are in contact.

To understand these results, recall some earlier work by Pashley on mica–mica interactions in aqueous solutions.⁴⁰ It was demonstrated that when enough sodium ions are adsorbed on mica surfaces, a repulsive hydration force develops. In the case of mica, the hydration force appeared rather suddenly at a critical salt concentration. Below this critical hydration concentration (CHC), the adhesion between the mica surfaces was virtually not affected by the NaCl concentration, whereas the hydration force reached its full strength at a concentration not much above the CHC. This was explained by a pressure-induced ion exchange of adsorbed sodium ions for protons below the CHC, but not above.⁴⁰ This mechanism did later gain some support by an ESCA investigation, where the number of sodium ions adsorbed to mica surfaces after immersion in electrolyte solutions was determined.⁴¹

In the case of DHP surfaces the situation is different. In dilute salt solutions, where no long-range hydration forces are observed, the adhesion force decreases with increasing NaCl concentration. The decrease in adhesion is presumably due to dehydration of sodium counterions replacing protons at the DHP monolayer surface. Hence, in contrast to mica, no pressure-induced ion exchange takes place between DHP surfaces at low NaCl concentrations. This implies that sodium ions have a higher affinity to DHP than to mica and that one cannot define a CHC for this system. At high pH, when a large number of sodium ions have replaced protons at the DHP monolayer surface, the hydration force reaches a sufficient strength and range to overcome the van der Waals force completely (Figure 2).

The decrease in interlayer attraction correlates with the increase in intralayer repulsion observed with the Langmuir force balance as an expansion of the liquid condensed phase (Figure 3a). We suggest that both these phenomena are caused by the presence of hydrated sodium ions at the surface. Hence, the expansion of the liquid condensed phase can be viewed as a result of intralayer

(40) Pashley, R. M. *J. Colloid Interface Sci.* **1981**, *83*, 531.

(41) Claesson, P. M.; Herder, P. C.; Stenius, P.; Eriksson, J. C.; Pashley, R. M. *J. Colloid Interface Sci.* **1986**, *109*, 31.

hydration forces having the same molecular origin as interlayer hydration forces. These results are also consistent with the observation made by Ninham, Evans, and co-workers that the size and type of aggregate which surfactants form are influenced by the nature of the counterions.⁴²⁻⁴⁵ We are presently investigating intralayer hydration forces in more detail by using a range of counterions.

Fusion and Aggregation of DHP and DODA Vesicles. Vesicles of dioctadecyldimethylammonium chloride (DODACl) and of sodium dihexadecyl phosphate (NaDHP) are unstable in the presence of NaCl. In this respect they behave differently than phospholipid vesicles. DODACl and NaDHP vesicles also behave qualitatively different from each other. Thus DODACl vesicles tend to fuse in 50 mM NaCl whereas DHP vesicles aggregate mainly without fusing.¹⁹ The reason for this difference has to be found in the interaggregate and intraaggregate interactions. The pressure-area isotherm for DODA clearly shows that the head-group area (a) decreases with increasing NaCl concentration. This means that the preferred value of the surfactant parameter v/al , which determines the aggregate shape,^{46,47} increases. (v is the hydrocarbon chain volume, and l is the hydrocarbon chain length.) Hence, there is a tendency for DODA to form *larger* aggregates as the salt concentration increases. This favors fusion.

On the other hand, for DHP the area per molecule increases when NaCl is added to the solution. This will favor formation of smaller aggregates as the salt concentration increases, which counteracts any tendency for DHP vesicles to fuse.

Another difference between DODA and DHP is that the short-range interaction between layers of double-chained quaternary ammonium surfactants in most cases is strongly repulsive,^{20,21} whereas except at high pH the short-range interaction between DHP layers is attractive. Molecules in vesicles presumably have some freedom to diffuse in the bilayer and hence respond to the presence of a second aggregate nearby. Repulsion between aggregates, e.g., DODA vesicles, is an unfavorable situation, and molecules will tend to move away from the contact area,^{48,49} which

decreases the *local* packing density in the region of the vesicles facing each other. Attraction between two aggregates, like NaDHP vesicles, naturally causes the reverse to be true. Hence, stable intermembrane intermediates may form.⁵⁰ This is another factor that favors fusion of DODA vesicles but opposes fusion of vesicles composed of DHP.

Of course, other properties as well, like the fluidity of the membrane, are of importance for understanding the intricate molecular events leading to fusion.⁵⁰ However, from this investigation we have been able to pinpoint two factors that favor DODACl vesicle fusion but oppose fusion of DHP vesicles.

Conclusions

Less than 2% of all DHP molecules deposited onto hydrophobed mica surfaces are charged at pH 5–6. With increasing NaCl concentration the surface charge density increases and the adhesion force between the layers decreases. Not only protons but also to a lesser extent sodium ions bind to DHP. At high pH, when sodium is the dominating counterion, a repulsive force prevents the surfaces from coming into adhesive contact. This force, which is measurable out to a separation of 1 nm, has the same origin as the repulsive hydration forces observed between phospholipid bilayers and between mica surfaces carrying adsorbed hydrated cations.

The liquid condensed phase of a DHP monolayer spread on a water subphase expands when sodium ions replace protons as counterions. Hence, for a given surface pressure the area per molecule increases with increasing NaCl concentration. We suggest that this is caused by an intralayer hydration force. For DODA monolayers in the liquid expanded state the opposite behavior is observed.

The salt dependences of the intralayer and interlayer interactions for DODA and DHP are quite different from each other. These differences are consistent with the observation that DODA vesicles fuse, whereas DHP vesicles mainly aggregate upon addition of sodium chloride.

Acknowledgment. We thank Johan Berg for valuable discussions and demonstrations. We also thank Barry Ninham, Christina Herder, and Peter Herder for constructive criticism. Per Claesson acknowledges a travel grant from the Swedish Board for Technical Development. A. M. Carmona-Ribeiro acknowledges a visiting fellowship from the Australian National University and financial support from the Fundacao de Amparo a Pesquisa do Estado de Sao Paulo.

Registry No. DHP, 2197-63-9; DODABr, 3700-67-2; NaCl, 7647-14-5.

(50) Bentz, J.; Ellens, H. *Colloids Surf.* **1988**, *30*, 65.

(42) Ninham, B. W.; Evans, D. F.; Wei, G. J. *J. Phys. Chem.* **1983**, *87*, 5020.

(43) Ninham, B. W.; Hashimoto, S.; Thomas, J. K. *J. Colloid Interface Sci.* **1983**, *95*, 594.

(44) Talmon, D. F.; Evans, D. F.; Ninham, B. W. *Science* **1983**, *221*, 1047.

(45) Kachar, B.; Evans, D. F.; Ninham, B. W. *J. Colloid Interface Sci.* **1984**, *100*, 287.

(46) Israelachvili, J. N.; Mitchell, D. J.; Ninham, B. W. *J. Chem. Soc., Faraday Trans. 2* **1976**, *72*, 1525.

(47) Brady, J. E.; Evans, D. F.; Warr, G. G.; Grieser, F.; Ninham, B. W. *J. Phys. Chem.* **1986**, *90*, 1853.

(48) Israelachvili, J. N.; Sornette, D. *J. Phys. (Les Ulis, Fr.)* **1985**, *46*, 2125.

(49) Eriksson, J. C.; Ljunggren, S., submitted for publication in *Colloid Polym. Sci.*

Referência 11

Theoretical and Experimental Stabilities of Amphiphile Vesicles in the Presence of NaCl

A. M. Carmona-Ribeiro

Departamento de Bioquímica, Instituto de Química, Universidade de São Paulo, CP20780 São Paulo, SP, Brazil (Received: February 25, 1988; In Final Form: September 7, 1988)

Theoretical and experimental stability values (W) for dioctadecyldimethylammonium chloride (DODAC) or sodium dihexadecyl phosphate (DHP) vesicles in the presence of NaCl salt are compared. The Derjaguin-Landau-Verwey-Overbeek (DLVO) theory for the stability of charged spherical colloidal particles is used to calculate some theoretical stabilities for small or large synthetic amphiphile vesicles in the presence of NaCl. The agreement between theoretical and experimental W values is interpreted as merely fortuitous since both inherent problems with the model and other factors operating in real systems can influence the vesicle stability. By improving the model and the inputs, the theoretical stability is found to be much higher than the experimental one for small DHP vesicles.

Introduction

Interactions between phospholipid model membranes have been extensively studied.¹⁻¹³ At long ranges, the main interactions are electrostatic (for charged lipids) and electrodynamic (van der Waals). The pioneer techniques on direct measurements of forces between phospholipid bilayers in aqueous solutions have shown the existence of very strong repulsive forces at short ranges probably due to the structural ordering of water molecules induced by the phospholipidic headgroups.^{1,12} At low ionic strength and neutral pH, hydration forces were absent for interacting bilayers of dihexadecyldimethylammonium acetate (DHDA) or bromide (DHDAB) adsorbed on the anionic mica surfaces^{13,14} or for interacting monolayers of sodium dihexadecyl phosphate (DHP) adsorbed on DHDAB-coated mica surfaces.¹⁵

Vesicles composed of dioctadecyldimethylammonium chloride (DODAC) or sodium dihexadecyl phosphate (DHP)^{16,17} interact at low monovalent salt concentration.^{18,19} This contrasts with the high stability of phospholipid vesicles at comparatively higher monovalent salt.¹¹ Qualitatively, DODAC vesicles fuse upon addition of NaCl whereas DHP vesicles mainly aggregate,¹⁹ a behavior consistent with differences in packing of DHDA or DHP molecules in monolayer films and with differences in directly measured interaction forces.¹⁵ Since hydration forces are absent

when amphiphile surfaces interact at low¹³⁻¹⁵ ionic strength, this work explores the possibilities of the kinetical DLVO approach^{20,21} in explaining the experimental stability of DODAC or DHP vesicles for a range of NaCl concentration. The calculations demonstrate that additional destabilizing factors and/or improvements in the model are needed to explain the experimental stability of small DHP vesicles obtained by sonication.

The Theory for Colloids

The total potential energy (V) for the interaction between two colloidal particles is defined as

$$V = V_R + V_A \quad (1)$$

V_R is the term for the repulsive interaction between the electrical double layers associated with the particles explicitly given by Reerink and Overbeek²¹ (25 °C, in water)

$$V_R = 4.62 \times 10^{-6} (a\gamma^2/\nu^2) \exp(-\tau u) \quad (2)$$

where

$$\gamma = (\exp(z/2) - 1)/(\exp(z/2) + 1); \quad z = q_e \nu \psi_\delta / kT \quad (3)$$

ν is the valency of the gegenions

$$u = H_0/a \quad (4)$$

where H_0 is the shortest distance between particle surfaces (in Å) and a , the particle radius (in Å):

$$\tau = Ka \quad (5)$$

$$K = ((8\nu^2 q_e^2)/(\epsilon kT))^{1/2} \quad \text{or} \quad K = C_e^{1/2} \times 10^7 / (0.31) \text{ cm}^{-1}$$

with C_e as the salt concentration (in M).

A layer of gegenions adheres to the particle surface forming an ionic condenser, the Stern layer,²² the diffuse part of the double layer extending into the solution. This layer of gegenions adhering to the particle surfaces accounts for Stern potentials smaller than the corresponding surface potentials.

V_A is the van der Waals term for the nonretarded attraction potential between two equal spheres for short distances ($u \ll 1$, i.e., $H_0 \ll a$):²³

$$V_A = -A/12u \quad (6)$$

By introduction of eq 2-6 in eq 1, the total energy of interaction can be explicitly written as a function of a (in cm), ν , ψ_δ (in mV), C_e (in M), A (in ergs), and u at 25 °C:

$$V = 4.62 \times 10^{-6} a [(e^{-0.0195\nu\psi_\delta} - 1)/(e^{-0.0195\nu\psi_\delta} + 1)]^2 e^{(-C_e^{1/2} \times 10^7 u)/0.31} - A/12u \quad (7)$$

(20) Verwey, E. J. W.; Overbeek, J. T. G. *Theory of Stability of Lyophobic Colloids*; Elsevier: Amsterdam, 1948.

(21) Reerink, H.; Overbeek, J. T. G. *Discuss. Faraday Soc.* 1954, 18, 74.

(22) Debye, H.; Hückel, P. *Phys. Z.* 1923, 24, 185.

(23) Hamaker, H. C. *Physics (Amsterdam)* 1937, 4, 1058.

- (1) Le Neveu, D. M.; Rand, R. P.; Parsegian, V. A. *Nature* 1976, 259, 601.
- (2) Le Neveu, D. M.; Rand, R. P.; Parsegian, V. A.; Gingell, D. *Biophysical J.* 1977, 18, 209.
- (3) Cowley, A. C.; Fuller, N. L.; Rand, R. P.; Parsegian, V. A. *Biochemistry* 1978, 17, 3163.
- (4) Nir, S.; Bentz, J. J. *Colloid Interface Sci.* 1978, 65, 399.
- (5) Lis, L. J.; Parsegian, V. A.; Rand, R. P. *Biochemistry* 1981, 20, 1761.
- (6) Lis, L. J.; Lis, W. T.; Parsegian, V. A.; Rand, R. P. *Biochemistry* 1981, 20, 1771.
- (7) Rand, R. P. *Annu. Rev. Biophys. Bioeng.* 1981, 10, 277.
- (8) Nir, S.; Bentz, J.; Duzgunes, N. J. *Colloid Interface Sci.* 1981, 84, 266.
- (9) Lis, L. J.; McAlister, M.; Fuller, N.; Rand, R. P.; Parsegian, V. A. *Biophysical J.* 1982, 37, 657.
- (10) Afzal, S.; Tesler, W. J.; Blessing, S. K.; Collins, J. M.; Lis, L. J. *J. Colloid Interface Sci.* 1984, 97, 303.
- (11) Ohki, S.; Roy, S.; Ohshima, H.; Leonards, K. *Biochemistry* 1984, 23, 6126.
- (12) Marra, J.; Israelachvili, J. N. *Biochemistry* 1985, 24, 4608.
- (13) Pashley, R. M.; McGuigan, P. M.; Ninham, B. W.; Brady, J.; Evans, D. F. *J. Phys. Chem.* 1986, 90, 1637.
- (14) Marra, J. *J. Phys. Chem.* 1986, 90, 2145.
- (15) Claesson, P. M.; Carmona-Ribeiro, A. M.; Kurihara, K. *J. Phys. Chem.*, in press.
- (16) Carmona-Ribeiro, A. M.; Chaimovich, H. *Biochim. Biophys. Acta* 1983, 733, 172.
- (17) Carmona-Ribeiro, A. M.; Yoshida, L. S.; Sesso, A.; Chaimovich, H. *J. Colloid Interface Sci.* 1984, 100, 433.
- (18) Carmona-Ribeiro, A. M.; Yoshida, L. S.; Chaimovich, H. *J. Phys. Chem.* 1985, 89, 2328.
- (19) Carmona-Ribeiro, A. M.; Chaimovich, H. *Biophys. J.* 1986, 50, 621.

Figure 1. Schematic representation of an energetic profile for interaction between two charged spherical particles with the same radius according to DLVO theory. The energy of interaction $V(u)$ as a function of the reduced separation distance u has three inflection points: the primary minimum (not shown), the maximum ($V_{\max}$), and the secondary minimum (V_{sec}).

From this expression for the total energy of interaction, the stability factor W can be evaluated:^{20,21}

$$W = 2 \int_0^{\infty} \exp(V/kT) du / (u + 2)^2 \quad (8)$$

The value of the integral can be calculated by numerical or graphical integration.

To avoid this integration a rather good approximate equation was derived by Reerink and Overbeek²¹

$$W_A = 2\pi^{1/2}(\exp(V_m/kT)) / (p(U_m + 2)^2) \quad (9)$$

where

$$p = (-d^2V(u)/du^2)_{u=U_m} \times 1/2(kT)^{1/2}$$

and

$$d^2V(u)/du^2|_{u=U_m} = V_m(u)\gamma^2\tau^2 - A/(6U_m^3)$$

i.e., the approximate stability factor W_A is determined by the maximum V_m in the potential curve and U_m corresponds to the separation distance u where this maximum occurs.

Equation 6 has been recognized as poor unless $H_0 \ll a$.²³ For H_0 varying between 0 and 100 Å and particles with 150 Å mean radius one should use the complete Hamaker equation²³

$$V_A = -A/12\{4/(u^2 + 2u + 4) + 4/(u^2 + 2u) + 2 \ln [(u^2 + 2u)/(u^2 + 2u + 4)]\} \quad (10)$$

Computations

Estimates of the total potential energy (V) of interaction as a function of the separation distance (u) were computed for four radii values (150, 400, 1350, and 2500 Å). The corresponding Stern potential values were respectively 27, -38.8, -19.3, and 10 mV (from ref 18) and Hamaker constants varying between 0.8×10^{-14} and 1.4×10^{-13} ergs were used as parameters for fitting experimental data. An Apple II microcomputer was programmed

to tabulate the coordinates of (V, u) curves. The stability ratio W was obtained by numerical integration of eq 8. The approximate stability W_A was calculated from eq 9 and the point of maximum taken from the (V, u) curves. In all computations of V , W , and W_A only aqueous solutions of 1:1 electrolytes at 25 °C were considered. The dimensionless dielectric constant (ϵ) of water at 25 °C was taken at 78.54. The unit of K is cm^{-1} and if ψ_δ is expressed in millivolts and the radius a in centimeters, the unit of V is erg.

Experimental Stability Values (W_e)

The initial flocculation rates (v_0) for DODAC or DHP vesicles were taken from ref 18. v_0 values were obtained under isoosmolarity conditions, at the initial stages of the flocculation process and for low amphiphile concentration. This allows us to assume that just two spherical vesicles are interacting since osmotically induced strain or undulations in the bilayer are absent and, at its initial stages, flocculation is a bivesicular process. The linear dependence between v_0 and the square of the amphiphile concentration¹⁸ demonstrates the bivesicular character of the flocculation process at its initial stages.

Mean v_0 values and square deviations were calculated from, at least, four independent kinetics for each salt concentration. Mean v_0 values were taken as inversely proportional to the experimental stability values (W_e), the proportionality constant being calculated from rapid flocculations ($W = 1$).²¹

Application of the Theory for Colloids to Vesicles

The total energy of interaction (V) as a function of the distance of separation of particle surface as calculated by the DLVO theory^{4,20,21} is frequently represented by an energy profile as shown in Figure 1. At a very small separation the theory produces a deep potential well, the primary minimum, and at a larger separation, a shallower well, the secondary minimum. The energy barrier between the two minima has been used as a criterion of stability: the dispersion is stable if a potential barrier larger than the average kinetic energy of the particles exists.

The energetic profiles on Figure 2 as a function of the separation distance were obtained from theory²¹ by calculating V for given values of particle radius (a), monovalent salt concentration (C_e), Stern potential (ψ_δ), Hamaker constant (A), and separation distance (u). Particle radius values were experimentally measured by electron microscopy,^{16,17} Stern potential values were calculated from experimental data as described,^{18,21} and Hamaker constant values were adjusted to fit theoretical to experimental stability factors (Table I and Figure 3). Therefore, the only parameter which was varied to obtain fitting was the Hamaker constant. Most A values were considerably lower than those used for colloids and slightly less than typical hydrocarbon/water/hydrocarbon values (around 5×10^{-14} erg). In addition, for the large DODAC or DHP vesicles, the A value decreased as the electrolyte concentration increased in accordance with the reduction of the zero-frequency contribution to the van der Waals interaction of hydrocarbon across water by increasing NaCl concentration.²⁴

Figure 2. Energy of interaction $V(u)$ as a function of the separation distance (in Å) as calculated by the Reerink and Overbeek model for four particle radii: 150 Å (A), 400 Å (B), 1350 Å (C), and 2500 Å (D). Monovalent salt concentration (C_e) is increased from curves a to d and values for C_e , ψ_δ and A used for calculating the curves are listed in Table I.

Figure 3. Comparison between calculated (●) and experimental (○) stability factors (W) as a function of the monovalent salt concentration (C_e) for DHP and DODAC vesicles. DHP vesicles have 400 (A) and 1350 Å (B) mean radius and DODAC vesicles 150 (C) and 2500 Å (D). The bars are the mean square deviation obtained from at least four independent measurements taken from ref 18.

TABLE I: Values of Particle Radius (a), Stern Potential, (ψ_s), Monovalent Salt Concentration (C_e), and Hamaker Constant (A) Used for Calculations of the Energy Curves on Figure 2

C_e , M	curve	a , Å	ψ_s , mV	$10^{14}A$, erg	vesicle type
0.0025	a	150	27	3	small DODAC
0.0100	b				
0.0500	c				
0.1000	d				
0.080	a	400	-38.8	14	small DHP
0.100	b				
0.150	c				
0.200	d				
0.100	a	1350	-19.3	4	large DHP
0.137	b			3.6	
0.192	c			3.5	
0.241	d			3.4	
0.030	a	2500	10	1.2	large DODAC
0.050	b			1.1	
0.120	c			0.9	
0.200	d			0.8	

For particles with 150 Å mean radius practically no secondary minimum was obtained (Figure 1A) but for particles with 400, 1350, or 2500 Å mean radius, a significant secondary minimum could be calculated (Figure 2, B, C, and D). One must notice that, for particles with 2500 Å mean radius, with C_e values up to 0.08 M, and at room temperature, the secondary minimum has a depth equal to kT (ca. 4×10^{-14} erg) (Figure 2D). This allows us to assume that large DODAC vesicles would present, under the specified conditions above, a reversible aggregation at the secondary minimum. In fact, this prediction from theory was experimentally confirmed.¹⁸ Large DODAC vesicles to which salt was added to a final concentration up to 0.08 M, at room temperature, formed visible flocs. These flocs could be completely redispersed by hand-shaking.¹⁸

For 400- and 1350-Å particles the theory predicts that aggregation at the secondary minimum will be scarcely reversible up to 0.08 M NaCl and at room temperature, since it is up to 9×10^{-14} erg deep (Figure 2, B,C). Supporting this prediction, a irreversible aggregation of DHP vesicles was described.¹⁹ No fusion could be detected between DHP vesicles,¹⁹ another fact which corroborates the hypothesis of an aggregation at the secondary minimum.

The height of the energy barrier ($V_{\max}$) at salt concentrations able to induce vesicle fusion¹⁹ is equal to or higher than $5kT$ (Figure 2A) and too high to allow the occurrence of fusion. This probably reflects a failure of the theory at short ranges. However, aggregation phenomena for DHP or DODAC vesicles could be

Figure 4. Comparison between calculated (●) and experimental (○) stability factors (W) as a function of the monovalent salt concentration (C_e) for 150 Å mean radius DODAC vesicles. ψ_s is 27 mV and A is 1×10^{-14} erg. Adjustment was achieved by using the eq 10.

Figure 5. van der Waals interaction for two water (w) particles with radius R surrounded by a hydrocarbon (h) shell of thickness δ and separated by a distance Δ , given by eq 11. The equivalent Hamaker constant obtained from the relationships described in the text is that for two hydrocarbon shells interacting across water, i.e., 5×10^{-14} erg.

understood from the obtained depth for the secondary minimum and the height of the energy barrier in Figure 2B–D.

By use of eq 10 the energy curves and the stability values for the sonicated DODAC vesicles were recalculated (Figure 4). Although the adjustment between theoretical and experimental W values was improved for concentrations below 30 mM NaCl, up to this concentration theoretical values were slightly higher than the experimental ones. The comparison between calculations using eq 6 and 9 shows that the error introduced by using eq 6 instead of 9 is negligible since reasonable fittings were obtained in both cases.

Criticisms: Improving the Model and the Inputs

1. *The van der Waals Attraction.* Equations 6 and 10 for the van der Waals interaction for massive particles are not correct for vesicles which should be considered as water particles surrounded by a hydrocarbon shell. The Vold approach²⁵ for the van der Waals interaction between spherical particles with an adsorbed layer allows to take into account the interactions particle/particle (H_p), particle/shell (H_{ps}), and shell/shell (H_s) across an intervening medium attributing appropriate Hamaker constants for the different interactions. By use of the approximate relationships between the Hamaker constants $A_{ps}^2 = A_p A_s$, $A_{ms}^2 = A_m A_s$, and $(A_s^{1/2} - A_p^{1/2})^2 = A_{sps} = 5 \times 10^{-14}$ erg (for two hydrocarbon layers interacting across water) and by considering the interaction between two spherical water particles with a radius R surrounded by a hydrocarbon layer with a thickness δ and separated by a distance Δ (Figure 5) the van der Waals interaction energy is given by

$$V_A = (-5 \times 10^{-14}/12)(H_s + H_p - 2H_{ps}) \text{ erg} \quad (11)$$

with $H = H(x,y) = y/(x^2 + xy + x) + y/(x^2 + xy + x + y) + 2 \ln((x^2 + xy + x)/(x^2 + xy + x + y))$ and

$$H_{ps} = H(x,y) \quad \text{with } x = (\Delta + \delta)/2R \text{ and } y = (R + \delta)/R$$

$$H_s = H(x,y) \quad \text{with } x = \Delta/(2R + 2\delta) \text{ and } y = 1$$

$$H_p = H(x,y) \quad \text{with } x = (\Delta + 2\delta)/2R \text{ and } y = 1$$

The application of eq 11 requires some modifications in the inputs for computations of the theoretical stability W_A . The "hydrocarbon shell" thickness (δ) may be taken as 50 Å for

(24) Mahanty, I.; Ninham, B. W. *Dispersion Forces*; Academic Press: New York, 1976.

(25) Vold, M. J. *Colloid Sci.* **1961**, *16*, 1.

DODAC and as 46 Å for DHP vesicles. It must also be noticed that only one value of Hamaker constant (5×10^{-14} erg) is going to be employed for calculations. Additionally, there is no variation of the Hamaker constant with the vesicle radius or with the amphiphile nature and the Hamaker constant value is not going to be adapted to get coincidence between experimental and theoretical stabilities.

2. The Electrostatic Repulsion. **2.1. Interaction at Constant Potential.** If the time needed for the relaxation of the double layer (τ_{DL}) were large as compared to the time scale of the coagulation process (τ_c), one should have used calculations at constant charge instead of those we used at constant potential. To verify if calculations at constant surface charge would have been more suitable than those we performed at constant potential, τ_{DL} and τ_c were estimated for the quickest coagulation processes occurring for the vesicles, namely, those for small DODAC vesicles.¹⁸ Smoluchowski²⁶ calculated τ_{SMOL} following eq 12²⁷ where n_0 is the number

$$\tau_{SMOL} = 2 \times 10^{11} / n_0 \text{ cm}^{-3} \text{ s} \quad (12)$$

of particles per cm^3 . Assuming a mean radius of 150 Å,¹⁶ 60 Å² as the area per monomer,¹⁴ the ratio between the number of monomers in the inner and outer monolayer as 1:2,¹⁶ and a final DODAC monomers concentration of 0.11 mM at the critical coagulation concentration (so that the coagulation process will be slowed down by a factor equal to 1), the number of small DODAC vesicles per cm^3 (n_0) is 9.3×10^{12} and a mean Smoluchowski coagulation time (τ_{SMOL}) of 22 ms can be calculated. From experimental data we calculated a rate constant value for coagulation of $4 \times 10^{-14} \text{ cm}^3 \text{ s}^{-1}$ for the sonicated DODAC vesicles at the CCC,¹⁸ a value ca. 3 orders of magnitude lower than that evaluated for the Smoluchowski rate ($k' = 8kT/3\eta = 1.4 \times 10^{-11} \text{ cm}^3 \text{ s}^{-1}$). Correspondingly, one should expect a coagulation time value ca. 3 orders of magnitude higher than that of Smoluchowski and the true coagulation time (τ_c) would be 22 s.

The relaxation time of the double layer is given by²⁷

$$\tau_{DL} = 5 \times 10^{-1} / C_e \text{ s} \quad (13)$$

where C_e is in M. Taking C_e as 0.120 M, τ_{DL} is $41.7 \times 10^{-11} \text{ s}$, a value much smaller than the 22 s needed for coagulation. This suggests that calculations at constant potential as those we performed are actually more adequate for these amphiphile vesicles. This reasoning is valid for the four vesicle systems presented in this work, since flocculation rates for the small sonicated DODAC vesicles are the highest rates measured.

2.2. The Surface Potential. The Stern potential (ψ_s) is expected to decrease with increasing electrolyte concentration (C_e) and the approximation of Reerink and Overbeek²¹ simply ignores this (see Table I). The values of ψ_s used for computations in the first model were derived from experimental flocculation rates as a function of monovalent salt concentration and a priori recognized as low due to the employed approximation.^{18,21} However, they reflect the variations of ψ_s for different vesicle types and sizes. Different curvature radius and headgroup area and nature could lead to differences in screening and/or binding of counterions directly affecting the ψ_s value. The DODAC vesicles' interface is probably more loosely organized than that for DHP as suggested by the larger limiting area per monomer of DODAC¹⁴ in comparison with that for phosphatidic acids²⁸ or DHP,¹⁵ by the DODAC vesicles' ability to fuse under the same conditions where DHP vesicles just aggregate¹⁹ and by the higher gel to liquid crystalline phase transition temperature of DHP vesicles in comparison with DODAC vesicles.^{16,29}

Referring to the choice of some reliable absolute ψ_s values to be used for calculations, a complex issue is raised. Since the size of vesicles precludes the use of any direct electrode method for the determination of the surface potential, the dye pH indicators³⁰⁻³⁵ or spin labels³⁶ have been often applied in related systems.

Figure 6. Total energy of interaction $V(u)$ as a function of the separation distance, in Å, as calculated by eq 11 and 2 for interacting vesicles with 400 Å mean radius. The hydrocarbon thickness was considered to be 46 Å; the NaCl concentrations were 0.1 M (a), 0.15 M (b), and 0.2 M (c). Stern potential values were extrapolated from Lukac's data³⁸ shown in the insert with the arrows indicating the extrapolations from the experimental curve. The Hamaker constant was taken as 5×10^{-14} erg in the three cases. One should notice the magnitude of the resultant repulsion originating very high theoretical stability values.

Electrophoretic mobilities have also allowed some ζ potential determinations for liposomes.³⁷ By fitting directly measured double-layer forces in between charged surfaces with theory, the surface potential can be estimated.^{3,12-15} This theory is basically the DLVO based on the nonlinear Poisson-Boltzmann (PB) equation and the nonretarded van der Waals attraction. However, it has recently been shown via Monte Carlo simulations³⁸ and the anisotropic hypernetted chain (HNC) approximation³⁹ that the solutions of the PB equation are quantitatively, and sometimes even qualitatively, incorrect in describing the interaction between charged surfaces. At short ranges and in electrolyte solutions the PB equation overestimates the double-layer repulsion due to neglecting the ion-ion correlations.⁴⁰ This precludes our using data from the surface force apparatus¹³⁻¹⁵ as reliable estimates of the surface potential. Furthermore, even in case these estimates were reliable, one could not use them for vesicles since they are valid only for practically planar surfaces.

To complicate matters further, estimates of surface potentials in DODAC or DHP vesicles as a function of the monovalent salt concentration are scarce in the literature probably due to the low and strict range of salt concentrations where they are stable as well as to problems inherent to the methods available.⁴¹

After all, if one considers that the PB approach does an excellent

(26) Smoluchowski, M. V. Z. *Phys. Chem.* **1916**, *42*, 143.

(27) Overbeek, J. T. G. *J. Colloid Interface Sci.* **1977**, *58*, 408.

(28) Jaehnig, F.; Harlos, K.; Vogel, H.; Eibl, H. *Biochemistry* **1979**, *18*, 1459.

(29) Carmona-Ribeiro, A. M.; Sesso, A., manuscript in preparation.

(30) Hartley, G. S.; Roe, J. W. *Trans. Faraday Soc.* **1940**, *36*, 101.

(31) Mukerjee, P.; Banerjee, K. *J. Phys. Chem.* **1964**, *68*, 3567.

(32) Montal, M.; Gitler, C. *Bioenergetics* **1973**, *4*, 363.

(33) Fromherz, P. *Biochim. Biophys. Acta* **1973**, *323*, 326.

(34) Funasaki, N. *J. Colloid Interface Sci.* **1977**, *60*, 54.

(35) Lukac, S. *J. Phys. Chem.* **1983**, *87*, 5045.

(36) Cafiso, D. S.; Hubbell, W. L. *Annu. Rev. Biophys. Bioeng.* **1981**, *10*, 217.

(37) McLaughlin, S.; Bruder, A.; Chem, S.; Moser, C. *Biochim. Biophys. Acta* **1975**, *394*, 304.

(38) Gulbrand, L.; Joensson, H.; Wennerstrom, H.; Linse, P. *J. Chem. Phys.* **1984**, *80*, 2221.

(39) Kjellander, R.; Marcelja, S. *Chem. Phys. Lett.* **1984**, *112*, 49.

(40) Kjellander, R.; Marcelja, S. *Chem. Phys. Lett.* **1986**, *127*, 402.

(41) Drummond, C. J.; Grieser, F.; Healy, T. H. *Faraday Discuss. Chem. Soc.* **1986**, *81*, 95.

job in predicting the behavior of some single-chain amphiphiles⁴² and even of sonicated DHP vesicles,³⁵ that the range of short-ranged forces for DODAC or DHP is expected to be actually the shortest in a scale where all double-chain amphiphiles were included (they have quite small head groups, and in the vesicles, they are packed in the rigid gel phase), and that Lukac's Stern potential values for sonicated DHP vesicles are reliable, by using eq 2 and 11 one can recalculate the energy of interaction as a function of the separation distance at least for the sonicated DHP vesicles (Figure 6). The corresponding theoretical stability values (not shown) were found to be always much higher than the experimental ones. Thus, overwhelming destabilizing instead of stabilizing factors are to be called to explain this.

3. Factors Operating in Real Systems. In spite of being relatively successful at long ranges the DLVO is not able to describe quantitatively the experimental stability of small sonicated DHP vesicles, which are less stable than the DLVO predictions would suggest. Several factors operating in real systems should be considered: the shape of the interacting vesicles; the likelihood of several vesicles interacting at the same time; the hydration forces due to hydration shells of bound counterions at high ionic strength; intrinsic factors which destabilize the bilayer structure as the formation of hexagonal phase and the possibility of vesicle growth via monomer diffusion.

Bilayers in the fluid liquid-crystalline phase can deform, undulate, or vary their shape both when they approach contact or in individual noninteracting vesicles. However, DODAC or DHP vesicles are in the rigid gel phase and are probably more resistant to deformation. Evidences favoring this hypothesis are the non-responsiveness of small DODAC or DHP vesicles to osmotic gradients by shrinking or swelling,^{16,17} perfectly spherical aggregated DHP vesicles as seen by electron microscopy,¹⁷ and the obeying of the Rayleigh law by small DODAC vesicles and the Jobst law by the large DHP and DODAC vesicles,^{17,19} i.e., a spherical shape for these vesicles. Anyway, the vesicle shape can have a central role in determining the interaction forces between vesicles. For example, long-range and extremely strong van der Waals forces peculiar to cylindrical geometries have long been known to exist²⁴ and attractive energies are expected to be at least 1 order of magnitude larger for rod-shaped structures than for spheres and planes.⁴³ Electron micrographs of small sonicated DHP vesicles at neutral pH clearly show they are rod-shaped²⁹ in contrast with the spherical shape of the large DHP vesicles.¹⁷

This could account for the excessively high Hamaker constant in Table I for the small DHP vesicles as well as the extraattractive forces needed to explain experimental stabilities.

In the same way as previously shown for interacting mica surfaces,⁴⁴ when the pH is increased, hydration forces can develop between interacting DHP layers due to the hydration shells of bound counterions.¹⁵ However, at a moderately high ionic strength (0.1 M) and neutral pH, no hydration forces were detected at short ranges in accordance with the extraattraction needed to explain interactions between small DHP vesicles. Unfortunately, systematic surface force data for DODAC bilayers at moderate or high ionic strength are absent in the literature.

Intrinsic factors related to the stability of the bilayer structure as the formation of hexagonal phases⁴⁵ or vesicle growth via monomer diffusion⁴⁶⁻⁴⁸ might also be involved. Both processes are frequent for short-chain surfactants in the liquid-crystalline state having a high solubility in water. For the long-chain DODAC or DHP, which are probably below their phase transition temperature at the present conditions and rather insoluble in water, these processes are unlikely. Nevertheless, one cannot forget the fluidizing effect of salt on related systems²⁸ so that these possibilities cannot be dismissed.

Conclusions

By considering the differences between colloids and vesicles and employing an acceptable DLVO model for interacting vesicles without free parameters, theoretical stabilities are found to be much higher than the experimental ones for small DHP vesicles. This demonstrates that some destabilizing factors have to be called to account for the experimental reality and/or the DLVO model overestimates the electrostatic repulsion in the case of small DHP vesicles. The absence in the literature of reliable estimates of the surface potential as a function of the monovalent salt concentration for vesicles precludes further calculations for small or large DODAC and large DHP vesicles.

Acknowledgment. I thank J. A. Ribeiro for his help with computations, CNPq for a research fellowship, and FAPESP for financial support.

Registry No. DODAC, 107-64-2; DHP, 60285-46-3; NaCl, 7647-14-5.

(42) Ninham, B. W.; Evans, D. F. *Faraday Discuss. Chem. Soc.* **1986**, *81*, 1-18.

(43) Allen, M.; Evans, D. F.; Mitchell, D. J.; Ninham, B. W. *J. Phys. Chem.* **1987**, *91*, 2320.

(44) Pashley, R. M. *J. Colloid Interface Sci.* **1981**, *83*, 531.

(45) Rupert, L. A. M. Ph.D. Thesis, University of Groningen, The Netherlands, 1987.

(46) McLean, L. R.; Phillips, M. C. *Biochemistry* **1981**, *20*, 2893.

(47) Yeagle, P. L. *Biochim. Biophys. Acta* **1985**, *822*, 267.

(48) Johnson, N. W.; Kaier, E. W. *J. Colloid Interface Sci.* **1987**, *116*, 444.

Referência 12

Dihexadecylphosphate Bilayers: Interbilayer Interactions and Intrabilayer Structure

A. M. CARMONA-RIBEIRO

Departamento de Bioquímica, Instituto de Química, Universidade de São Paulo, CP 20780, São Paulo, Brazil

Received October 17, 1989; accepted February 16, 1990

The initial flocculation rate of sodium dihexadecylphosphate (v_0) (DHP) vesicles is determined as a measurement of intervesicles interaction for a range of NaCl concentrations (C) or pH values. Also, the stability of the membrane structure is assessed by turbidimetrically determining the gel-to-liquid-crystalline phase transition temperature (T_c) of the DHP vesicles. The stability of the vesicles with respect to interactions taken as $\log 1/v_0$ displays a bell-shaped profile as a function of pH for a given NaCl concentration. The temperature-induced phase transition of the bilayer occurs for a broad range of temperatures with midpoints between 59 and 73°C, depending on the pH value. The maximal T_c value is at pH 5.5–6.0, the apparent pK value of the phosphate polar head at the membrane/water interface. The pH range of maximal structural stability (T_c) coincides with the pH range of maximal colloidal stability ($1/v_0$) and with the maximal values of $d \log(1/v_0)/d \log C$, a parameter representative of the Stern Potential. In close analogy to phosphatidic acids and other lipids with a dissociative polar head, an increase in the total charge of the DHP headgroup from zero to half caused an increase in T_c . Factors governing the stability of the bilayer structure in the gel state seem to be the same as those governing the stability relative to interbilayer interactions. © 1990 Academic Press, Inc.

INTRODUCTION

A fundamental property of amphiphile molecules is their ability to organize themselves in water so that their hydrophilic polar or ionic head contacts water and their hydrophobic moieties interact with each other (1, 2). Depending on chemical structure, amphiphiles aggregate to form micelles, rods, bilayer vesicles, or inverted hexagonal phases (3, 4). The polymorphism accessible to many naturally occurring lipids such as phosphatidylethanolamine might be significant to some of the mechanisms of biological membrane fusion (5, 6) with bilayer destabilization proposed as initiator of the fusion event (7). On the other hand, biological membranes may contain up to 20% ionizable lipids, and surface charges represent a relevant regulation mechanism of the membrane structure and permeability triggered by pH or temperature changes (8–11). At neutral pH, sodium dihexadecylphos-

phate (DHP) forms vesicles (12, 13) suitable for permeation (13), aggregation (14), and intervesicle interaction (14, 16) studies. In spite of the absence of hydration forces and the presence of a strong adhesive force for DHP layers interacting at short ranges (17), no fusion was detected at pH 5.5–6.0 and 23°C either for DHP vesicles (14) or planar layers (17). A pH-dependent permeability of synthetic bilayers having a dissociative headgroup such as carboxylate or phosphate (18, 19) and a pH-dependent fusion of didodecylphosphate (DDP) vesicles induced by Ca^{2+} (20) have been reported. The dissociative polar head in the DHP bilayer allows one to vary membrane charge, surface potential, and membrane structure just by changing pH. This offers a unique opportunity to address the question of how membrane structure and intermembrane interactions are related in a simple membrane model system like the DHP vesicles. Furthermore, this line of investigation is relevant in

answering the more general question of how the chemical structure of the monomer determines the properties of its supramolecular assemblies. This paper presents data on flocculation and phase behavior of small bilayer vesicles prepared by ultrasonically dispersing DHP in water media. The overall strategy is to determine initial flocculation rates as a measurement of the extent of intervesicle interaction and to relate these data with the stability of the membrane in the gel state by turbidimetrically determining the phase behavior of the small DHP vesicles for a range of pH values. The results show very clearly that the highest colloidal stability of the vesicles occurs for the same pH range where the stability of the membrane structure in the gel state is maximal.

MATERIAL AND METHODS

Preparation of the DHP dispersion. Dihexadecylphosphoric acid (Sigma, St. Louis, MO) was used for the experiments in buffered solutions. Sodium dihexadecylphosphate (DHP) was obtained from the dihexadecylphosphoric acid as previously described (13) and used in the experiments performed with unbuffered solutions. DHP was ultrasonically dispersed in water solution (12, 13). DHP concentration was measured by inorganic phosphate determination (21). The bilayer nature of the DHP dispersion obtained by sonication has been confirmed both by X-ray scattering (22) and by electron microscopy (Carmona-Ribeiro, A. M., and Sesso, A., in preparation). The stability of DHP relative to hydrolysis, like that of some ether analogs of phosphatidic acids (23), is expected to be high, and problems with decomposition at high or low pH values, such as those observed for phosphatidic acids (PA) (10, 23), are presumably absent.

Buffers and solutions. NaCl aqueous solutions used to induce DHP vesicle flocculation (15) contained either 10 mM phosphate buffer or 10 mM borate buffer depending on the pH

value to be adjusted (smaller or higher than 8, respectively). Isoosmolarity between the inner and the outer vesicle compartment was kept by a complementary addition of D-glucose to NaCl solutions. The osmolarity of the NaCl/D-glucose solution added to the vesicles was equal to the osmolarity of the D-glucose solution in which the vesicles were prepared. The pH value of the saline solution was adjusted with a concentrated HCl or NaOH solution. The total Na^+ concentration was calculated, taking into account the NaCl concentration, the buffer concentration, and the NaOH eventually added for adjusting pH. NaCl concentration was determined by Cl^- titration (22).

Determination of the initial flocculation rate (v_0). One-tenth-milliliter DHP vesicles (2 mM) were added to 0.9 ml salt solution at a given pH. Turbidity at 400 nm was monitored as a function of time in a Varian DMS-70 spectrophotometer. From these curves, v_0 values (in s^{-1}) were calculated (15) and plotted as a function of Na^+ concentration for several pH values. For control, the pH was measured after each flocculation kinetics determination.

Determination of temperature effects. The vesicles dispersion for determination of temperature effects was prepared in water at a given pH value and for a final DHP concentration of 1 mM. The pH values of basic water media were maintained by preparing the vesicles under an N_2 atmosphere. It must be emphasized that the DHP vesicle dispersion for determination of temperature effects was always prepared in the absence of buffer due to its high instability in the presence of salt (15). The temperature of the vesicle dispersion inside a quartz cuvette was varied using a Polyttemp circulating bath (Model 80 T, constant temperature circulator, Polyscience Corp., Niles, IL, U.S.A.) connected to a thermostated cuvette holder in a Varian DMS-70 spectrophotometer. Temperature inside the cuvette was continuously monitored with a copper-constantan thermocouple connected to a galvanometer. The thermocouple was

previously calibrated, yielding a linear dependence between the galvanometer reading and the temperature (correlation coefficient of 0.9999). Turbidity at 260 nm was recorded as a function of temperature. An N_2 atmosphere inside the cuvette allowed the basic pH values to be kept constant. For control, the pH value was measured before and after each temperature cycle. The temperature variation rate was constant and equal to 1°C per minute.

RESULTS AND DISCUSSION

The Ordered-Fluid Phase Transition and the DHP Bilayer Structure

A characteristic property of biomembranes is the transition from the ordered to the fluid state of the lipid chains (25). This transition corresponds to abrupt changes in the fluidity and packing density of the hydrocarbon chains at a particular temperature, i.e., the phase transition temperature (T_c). The higher the T_c value, the higher the packing density of the hydrocarbon chains in the ordered state.

Measurement of turbidity has served as a simple and relatively accurate physical method of determining temperature-dependent optical properties of phospholipid (26, 27) or amphiphile (14, 28) membranes. The effect of temperature on the turbidity of DHP vesicles is pH-dependent (Fig. 1). DHP vesicles have a sensitive region below pH 6.5 and above pH 4.5. The release of the phosphate proton occurs in this region as determined by potentiometric titrations of DHP vesicles prepared in water (not shown), yielding an apparent pK value of 5.5–6.0 in fair agreement with previously reported data (19).

The high T_c values obtained for the small DHP vesicles (Fig. 1) indicate that its bilayer is one of the most tightly packed in a scale where all bilayers (including those of phospholipids) were placed. This high packing density is corroborated by the low area per monomer values measured under several experimental conditions for air/water DHP monolayers (17).

At low pH, the fully protonated DHP bilayer has a smaller T_c or packing density than the fully deprotonated one (Figs. 1G and 1A, respectively). This could be attributed to a tendency of DHP monomers to assume a truncated conical shape leading the hydrocarbon chains to bulge upon protonation of the headgroup. In fact, this would correspond to an increase in fluidity upon protonation and has been demonstrated for bilayers composed of didodecylphosphate (20).

At pH 4, multiple phase transitions can be observed (Fig. 1G). This might be due to the high polydispersity of the DHP dispersion formed at this pH as detected by light-scattering measurements (19) and electron microscopy (Carmona-Ribeiro, A. M., and Sesso, A., in preparation). Nevertheless, all the transitions detected occur around T_c values much lower than those observed at higher pH values, confirming the presence of a different DHP bilayer structure at low pH.

Turbidity curves induced by heating of DHP dispersions prepared at pH values outside the pK region present the abrupt decrease in turbidity characteristic of the gel-to-liquid-crystalline phase transition (Figs. 1A, 1B, 1C, 1F, 1G, 1H). A pronounced hysteresis, as currently obtained in similar systems (25, 29) is observed by comparison of the heating and cooling curves (Fig. 1). At and around the pK value (5.5–6.0), atypical turbidity vs temperature curves were obtained. An extensive, reversible and temperature-dependent turbidity increase abruptly occurs around 70–73°C and pH 5.5–6.0 (Figs. 1D and 1E). This turbidity increase possibly corresponds to vesicle aggregation at high temperatures. For similar systems, abrupt light-scattering increases like this have been associated with the formation of the isotropic phase (7, 30, 31). At high temperatures, linkages in between vesicles consist of an isotropic molecular arrangement so that the so-called intermembrane intermediate is formed (7). Therefore, aggregation is necessary although not sufficient for the formation of the isotropic phase, and the DHP vesicle

FIG. 1. Effect of temperature on the absorbance at 260 nm of DHP vesicles obtained in water at several pH values, under an N_2 atmosphere and in absence of buffer. Heating is represented by solid circles and cooling by open circles. pH values before and after the heating/cooling cycle are, respectively: 10.2 and 9.8 in A; 9.2 and 8.5 in B; 7.3 and 7.2 in C; 6.0 and 6.0 in D; 5.5 and 5.6 in E; 4.5 and 4.4 in F; 4.0 and 4.0 in G.

aggregation at the apparent pK and 70–73°C can be viewed as just an indication for the existence of DHP in the isotropic phase. However, a high capacity for hydrogen bonding, a high degree of hydrophobicity of the head-group, and a very tight packing were demonstrated to be typical properties of phospholipid analogs able to exist in the isotropic phase (30, 31). Since DHP molecules satisfy these requirements (17), the possibility of DHP existing in the isotropic phase is reinforced.

From curves in Fig. 4, the midpoints and ranges of the phase transition for the heating runs were plotted as a function of the bulk pH (Fig. 2). The T_c values are maximal around pH 5.5–6.0, decreasing slightly for higher pH values. At low pH, the smallest T_c values were

obtained. Inducing half a charge on DHP bilayers results in the formation of a bilayer structure with a T_c around 70°C, i.e., 10°C above the T_c of the fully protonated DHP bilayer (Fig. 2). This increase in the T_c value, also observed for PA (23) and for glycerophospholipids (32), can be explained by an overcompensation of the electrostatic destabilizing effect by intermolecular hydrogen bondings. The maximal stability of the DHP bilayer in the gel state occurs at the pH value where the maximal number of hydrogen bridges is expected to occur: at the pK . The pK for DHP is higher than the first pK for PA (ca. 3.5) (23) suggesting that the hydrophobicity of the DHP surface is higher than that of the PA surface. In fact, the occurrence of

FIG. 2. Mean phase transition temperature (T_c) of the DHP vesicles prepared in water as a function of pH. The vertical bars correspond to the T_c temperature range and the horizontal bars to the pH difference as measured before and after the heating/cooling cycle.

the dehydrated "high melting" solid phase (Figs. 1D and 1E), the absence of an hydration repulsion (17), the measurement of a strong adhesion (17), and the occurrence of vesicle aggregation at low salt (14) could reflect a low degree of water structuring on the DHP surface. Lipid hydration has been shown to outweigh the importance of the surface electrostatics by nearly one order of magnitude in determining the phase behavior (32). The lower the phase transition temperature, the higher the lipid hydration (32). Therefore, the DHP bilayer surface seems to be indeed poorly hydrated at the pH of water. A possible cause for this low degree of hydration is the lack in the DHP molecule of some hydrophilic groups present in the phospholipids such as the glycerol backbone and the glycerol/fatty acids ester bonds.

Up to pH 5.5–6.0, the T_c value decreases as a function of pH (Fig. 2). DHP monolayer data demonstrate a monolayer expansion due to an increase in the pH of the subphase in this same pH range (17). Both results are possibly due to binding of hydrated Na^+ ions. Thus, the DHP bilayer hydration would increase and, in the presence of 10 mM NaCl at pH 9.5, a repulsive intralayer and interlayer

hydration force would develop (17). Unfortunately, it is not feasible to measure turbidimetrically the phase behavior of small DHP vesicles as a function of Na^+ ion concentration at pH 9–10 due to a high extent of intervesicles interaction even at low salt concentrations. If this were feasible, at NaCl concentrations in the range 1–10 mM a much more pronounced T_c decrease as a function of pH would have to be expected due to the increase in the degree of hydrophilicity of the surface caused by the binding of hydrated Na^+ ions at the fully deprotonated bilayer. We are presently investigating salt effects on the phase behavior of another DHP bilayer model, the large DHP vesicles, since they are more stable in the presence of salt (13).

The Bilayer Stability Relative to Interbilayer Interactions and the Surface Charge

The DHP dispersion obtained by sonication is unstable in the presence of low-concentration NaCl solutions, i.e., intervesicle interaction begins to occur upon salt addition, leading the vesicles to aggregate at the pH of water (13–15). Here we take advantage of this property to measure the effect of the bilayer charge on the initial flocculation rates (v_0). The aggregation of DHP vesicles is measured as a turbidity increase as a function of time after salt addition. From this kinetics, a v_0 value can be determined.

The initial flocculation rate (v_0) of DHP vesicles obtained by sonication depends on Na^+ concentration and on the bulk pH value (Fig. 3). In general, $\log v_0$ varies linearly with $\log C$ as predicted by the DLVO theory of colloidal stability (33) (Fig. 3). v_0 increases with the salt concentration until the critical coagulation concentration (CCC) is attained. Up to the CCC, further increasing the salt concentration does not change the v_0 value. Salt is not very effective in inducing flocculation at low pH (Figs. 3A and 3B). This is further emphasized by the decrease in the CCC value with increasing pH (Fig. 4B). These results

FIG. 3. Initial flocculation rates (v_0) of DHP vesicles as a function of Na^+ concentration. pH values are: 2.8 in A; 3.8 in B; 5 in C; 6 in D; 7 in E; 7.8 in F; 8.6 in G; 9.6 in H. Vesicles are prepared in a 0.59 M D-glucose solution. Vesicle flocculation is obtained by adding 0.9 ml buffered NaCl solution to 0.1 ml of the vesicle dispersion. Flocculation is monitored as turbidity at 400 nm as a function of time. Final DHP concentration in the cuvette is 0.18 mM and final Na^+ ion concentration is indicated as C .

suggest that the DHP dispersion behaves like a colloid for the pH range where the bilayer bears some charge.

From the DLVO (33), the stability of the DHP dispersion should increase with increasing surface charge and potential. By calculating the slope of the $\log 1/v_0$ vs $\log C$ curves in Fig. 3, some estimates of the surface potential can be obtained (33). This allows one to verify

if the surface potential (and surface charge) is actually displaying the expected S-shaped titration curve with the apparent pK value of the phosphate at the surface as midpoint (the pK value should not be too far from 5.5–6.0). Figure 4A shows the dependence of the slope on the bulk pH. The slope first increases, reaches a maximal value at pH 5.0–8.0, and then decreases. It is clear that the surface charge and potential do not display the behavior expected for dissociative groups fixed in a solid colloidal particle. The bilayer rigidity varies as a function of pH (Fig. 2) and the surface potential increases when the membrane rigidity increases and vice versa, as depicted by the parallel runs of the T_c (Fig. 2) and the slopes (Fig. 4A) as a function of pH. Membrane structure determines the value the surface potential will assume. Nevertheless, the surface potential seems to determine the membrane stability relative to interactions, exactly as stated by the DLVO (33). In Fig. 5A, the vesicle stability relative to flocculation taken as $\log 1/v_0$ displays a bell-shaped profile as a function of pH, where the highest stability values occur for pH values varying between 5 and 8 at low salt. Thus, the colloidal stability is maximal (Fig. 5A) for the same pH range where the surface potential is maximal (Fig.

FIG. 4. Slopes ($d \log v_0 / d \log C$) and critical coagulation concentration (CCC) for DHP vesicles as a function of pH.

FIG. 5. Colloidal stability of DHP vesicles ($\log 1/v_0$) as a function of pH. The NaCl concentrations are: 0.1 M in A; 0.13 M in B; and 0.16 M in C.

4A). The colloidal stability of DHP vesicles at low salt runs in parallel to the electrostatic potential estimates as predicted by the DLVO. Of course, this qualitative agreement does not mean that the theory is able to account also for short-ranged interactions. Theoretical stability values for small DHP vesicles at the pH of water calculated using an adapted DLVO model are much higher than the experimental values taken from the flocculation data (16). This suggests the need for a short-ranged attractive force to explain the experimental data and is consistent with the attraction needed to form intermembrane intermediates and the isotropic phase at high temperatures (Figs. 1D and 1E).

By increasing the Na^+ ion concentration, the pH range of maximal stability decreases (Figs. 5B and 5C). The effect of increasing monovalent salt concentration on the dissociation of dissociative polar heads in phospholipid membrane is to facilitate the dissociation, thereby shifting the apparent pK' to lower values (9, 10). Consistently, the decrease in the pH range of maximal stability of DHP vesicles obtained by increasing salt can be due to the pK' shift of the phosphate polar head to lower pH values.

In summary, the DHP bilayer structure has its highest stability in the gel state for the same pH range where its colloidal stability and surface potential are maximal. This means that the bilayer structure determines the surface potential that determines to a certain extent

the colloidal stability. At the pH of water, a high packing density prevents DHP membranes from fusing whereas a low degree of hydrophilicity on the membrane due to intramembrane hydrogen bonding leads to adhesion and vesicle aggregation at very low salt concentrations.

ACKNOWLEDGMENTS

This work was supported by the Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP) and by the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq).

REFERENCES

1. Tanford, C., "The Hydrophobic Effect." Wiley Interscience, New York, 1982.
2. Attwood, D., and Florence, A. T., "Surfactant Systems—Their Chemistry, Pharmacy and Biology." Chapman and Hall, London/New York, 1983.
3. Israelachvili, J. N., Mitchell, D. J. and Ninham, B. W., *Biochim. Biophys. Acta* **470**, 185 (1977).
4. Cullis, P. R., Hope, M. J., and Tilcock, C. P. S., *Chem. Phys. Lipids* **40**, 127 (1986).
5. Cullis, P. R., and de Kruijff, B., *Biochim. Biophys. Acta* **559**, 399 (1979).
6. Siegel, D. P., *Biophys. J.* **45**, 399 (1984).
7. Ellens, H., Bentz, J., and Szoka, F. C., *Biochemistry* **25**, 4141 (1986).
8. Traeuble, H., and Eibl, H., *Proc. Natl. Acad. Sci. USA* **71**, 214 (1974).
9. Jaehnig, F., *Biophys. Chem.* **4**, 309 (1976).
10. Traeuble, H., Teubner, M., and Woolley, P., and Eibl, H., *Biophys. Chem.* **4**, 319 (1976).
11. Jaehnig, F., Harlos, K., Vogel, H., and Eibl, H., *Biochemistry* **18**, 1459 (1979).
12. Mortara, R. A., Quina, F. H., and Chaimovich, H., *Biochem. Biophys. Res. Commun.* **81**, 1080 (1978).
13. Carmona-Ribeiro, A. M., Yoshida, L. S., Sesso, A., and Chaimovich, H., *J. Colloid Interface Sci.* **100**, 433 (1984).
14. Carmona-Ribeiro, A. M., and Chaimovich, H., *Biophys. J.* **50**, 621 (1986).
15. Carmona-Ribeiro, A. M., Yoshida, L. S., and Chaimovich, H., *J. Phys. Chem.* **89**, 2928 (1985).
16. Carmona-Ribeiro, A. M., *J. Phys. Chem.* **93**, 2630 (1989).
17. Claesson, P., Carmona-Ribeiro, A. M., and Kurihara, K., *J. Phys. Chem.* **93**, 917 (1989).
18. Okahata, Y., and Seki, T., *J. Amer. Chem. Soc.* **106**, 8065 (1984).

19. Tricot, Y.-M., Furlong, D. N., Sasse, W. H. F., Daivis, P., Snook, I., and van Megen, W., *J. Colloid Interface Sci.* **97**, 380 (1984).
20. Rupert, L. A. M., van Breemen, J. F. L., Hoekstra, D., and Engberts, J. B. F. N., *J. Phys. Chem.* **92**, 4416 (1988).
21. Rouser, G., Fleischer, S., and Yamamoto, A., *Lipids* **5**, 494 (1970).
22. Paradies, H. H., *Biochem. Biophys. Res. Commun.* **101**, 1096 (1981).
23. Eibl, H., and Blume, A., *Biochim. Biophys. Acta* **553**, 476 (1978).
24. Schales, O., and Schales, S. S., *J. Biol. Chem.* **140**, 879 (1941).
25. Lee, A. G., *Biochim. Biophys. Acta* **472**, 237 (1977).
26. Yi, P. N., and MacDonald, R. C., *Chem. Phys. Lipids* **11**, 114 (1973).
27. Petersen, N. O., and Chan, S. I., *Biochim. Biophys. Acta* **509**, 111 (1978).
28. Carmona-Ribeiro, A. M., and Chaimovich, H., *Biochim. Biophys. Acta* **733**, 172 (1982).
29. McDonald, R. C., Simon, S. A., and Baer, E., *Biochemistry* **15**, 885 (1976).
30. Silviu, J. R., Brown, P., and O'Leary, T. J., *Biochemistry* **25**, 4249 (1986).
31. Brown, P. M., Steers, J., Hui, S. W., Yeagle, P. L., and Silviu, J. R., *Biochemistry* **25**, 4249 (1986).
32. Cevc, G., Seddon, J. M., and Marsh, D., *Faraday Discuss. Chem. Soc.* **81**, 179 (1986).
33. Reerink, H., and Overbeek, J. Th. G., *Disc. Faraday Soc.* **18**, 74 (1954).

Referência 13

pH Effects on Properties of Dihexadecyl Phosphate Vesicles

A. M. Carmona-Ribeiro* and S. Hix

Departamento de Bioquímica, Instituto de Química, Universidade de São Paulo, CP 20780 São Paulo, Brazil
(Received: June 4, 1990)

Vesicle size (S) and bilayer structure and function as well as colloidal stability were assessed in large dihexadecyl phosphate (DHP) vesicles on the basis of turbidity spectra, phase transition temperature (T_c) determinations, water permeation rates (v_s), and initial flocculation rates (v_0), respectively, for a range of NaCl concentration (C) or pH. S , T_c , v_s , and $\log 1/v_0$ display a bell-shaped profile as a function of pH for a given C . In water, the maximal S , T_c , v_s , and $\log 1/v_0$ values occur at pH 6.0–7.0, i.e., around the apparent pK value of the phosphate polar head at the membrane/water interface. The results emphasize the importance of the extent of surface hydration in determining the total surface area of the dispersion and thereby the vesicle size and phase behavior.

Introduction

A general question concerning amphiphiles in solution is how the chemical structure of the monomer and the solution composition determine the properties and the nature of the aggregates.^{1,2} In the case of dodecyltrimethylammonium bromide, for example, the size and nature of the aggregates formed in solution change with the added counterion.³

At the pH of water and low levels of monovalent salt, sodium dihexadecyl phosphate (DHP) forms small^{4,5} or large⁶ one-bilayer vesicles depending on the dispersion method. At an air–water interface and for a given surface pressure, a DHP monolayer expands upon increasing NaCl concentration (0–10 mM) or pH in the subphase.⁷ On the other hand, adsorbed DHP monolayers⁷ or DHP vesicles⁸ do not fuse but simply aggregate at pH 6.5 upon NaCl addition. For small DHP vesicles (SV), a high packing density of the DHP bilayer in the gel state and a high hydrophobicity of the bilayer surface could explain both the nonoccurrence of fusion and the vesicle adhesion observed upon NaCl addition at the pH of water.⁹

An extrarepulsive hydration force at pH 9–10 was measured by use of a surface force apparatus in a previous work of ours with adsorbed DHP layers on mica.⁷ At pH values probably close to the pK_a , an extraattractive force (not to be attributed to van der Waals attraction) was both measured by use of the surface force apparatus⁷ and predicted from DLVO calculation for vesicles.¹⁰

Here we take advantage of the pH and salt-dependent hydrophilicity of the DHP bilayer surface^{7,10} to address the question of how surface hydration and bilayer properties are related in a simple model system like the DHP vesicles. The restrictions imposed by the SV system, such as the osmotic nonresponsiveness^{6,11} and the low colloidal stability in the presence of a monovalent salt,¹² are circumvented by using large vesicles (LV). In LV, NaCl effects (0–20 mM) on the membrane properties can be determined without the occurrence of flocculation for a range of pH. Furthermore, LV turbidity spectra yield linear dependencies between turbidity and the inverse square of the wavelength of light, permitting an estimate of any vesicle size variation in a straightforward manner for a range of pH and C . The results clearly show the importance of the degree of surface hydration in determining the total surface area of the dispersion and thereby the vesicle size, the phase behavior, and the colloidal stability at low salt.

Material and Methods

Preparation of Large DHP Vesicles. Dihexadecylphosphoric acid was from Sigma (St. Louis, MO) and used as provided by the manufacturer for the experiments in buffered solutions. Sodium dihexadecyl phosphate (DHP) was obtained from the dihexadecylphosphoric acid as previously described⁶ and used in the experiments performed with unbuffered solutions. Large DHP vesicles were prepared by injecting a chloroformic DHP solution

into a water solution at 75 °C.⁶ DHP concentration was measured by inorganic phosphate determination.¹³ The bilayer nature of the DHP dispersion was confirmed by electron microscopy for a range of pH and C .¹⁴

Determination of Vesicle Size Variation. Large DHP vesicles were prepared at several pH and NaCl concentration values. The effect of medium on size was monitored by recording vesicle turbidity (A) as a function of the wavelength of the incident light (λ).⁶ Large DHP vesicles scatter light in accordance to the Joebst law¹⁵ at the pH of water.⁶ Taking advantage of the linear dependence between A and the inverse of λ^2 , a slope value can be calculated and normalized for different DHP concentrations. The normalized slope (NS) was taken as a vesicle size parameter indicative of size changes. Turbidity (A) is given by¹⁶

$$A = \text{constant} \cdot q^2 \delta / (V^{2/3} \lambda^2) \quad (1)$$

where q is the anhydrous mass of the particle, δ is the number of particles per unit volume, V is the particle volume, and λ is the wavelength of the incident radiation. Since the DHP concentration (P_1) is

$$P_1 = \delta n \quad (2)$$

where n is the number of DHP molecules per vesicle per Avogadro number, the NS value for A vs $1/\lambda^2$ is

$$\text{NS} = \text{SLOPE} / P_1 = \text{constant} \cdot q^2 (nV^{2/3}) \quad (3)$$

Considering that q is given by $n\text{MW}$ where MW is the molecular weight of DHP, NS is

$$\text{NS} = \text{constant} \cdot n\text{MW}^2 / V^{2/3} \quad (4)$$

From eq 4, it is clear that the variation in NS depends on the variation in n .

Determination of Temperature Effects. The vesicle dispersion

(1) Attwood, D.; Florence, A. T. *Surfactant Systems—Their Chemistry, Pharmacy and Biology*; Chapman and Hall: London, 1983.

(2) Tadros, T. F. *Surfactants*; Academic Press: London, 1984.

(3) Ninham, B. W.; Evans, D. F.; Wei, G. J. *J. Phys. Chem.* **1983**, *87*, 5020.

(4) Mortara, R. A.; Quina, F.; Chaimovich, H. *Biochem. Biophys. Res. Commun.* **1978**, *81*, 1080.

(5) Tricot, Y.; Furlong, D. N.; Sasse, W. H. F.; Daivis, P.; Snook, I.; Van Megen, W. *J. Colloid Interface Sci.* **1984**, *97*, 380.

(6) Carmona-Ribeiro, A. M.; Yoshida, L. S.; Sesso, A.; Chaimovich, H. *J. Colloid Interface Sci.* **1984**, *100*, 433.

(7) Claesson, P. M.; Carmona-Ribeiro, A. M.; Kurihara, V. *J. Phys. Chem.* **1989**, *93*, 917.

(8) Carmona-Ribeiro, A. M.; Chaimovich, H. *Biophys. J.* **1986**, *50*, 621.

(9) Carmona-Ribeiro, A. M. *J. Colloid Interface Sci.* **1990**, *139*, 343.

(10) Carmona-Ribeiro, A. M. *J. Phys. Chem.* **1989**, *93*, 2630.

(11) Carmona-Ribeiro, A. M.; Chaimovich, H. *Biochim. Biophys. Acta* **1983**, *733*, 172.

(12) Carmona-Ribeiro, A. M.; Yoshida, L. S.; Chaimovich, H. *J. Phys. Chem.* **1985**, *89*, 2928.

(13) Rouser, G.; Fleischer, S.; Yamamoto, A. *Lipids* **1970**, *5*.

(14) Hix, S.; Sesso, A.; Carmona-Ribeiro, A. M. Manuscript in preparation.

(15) Joebst, G. *Ann. Phys.* **1925**, *78*, 157.

(16) Koch, A. L. *Biochim. Biophys. Acta* **1961**, *51*, 429.

* To whom all correspondence should be addressed.

(LV) for the determination of temperature effects was prepared either in water or in buffer at a given pH value and for a final DHP concentration of 0.05–0.2 mM. The buffer solution used was HAc/NaAc, $\text{NaH}_2\text{PO}_4/\text{Na}_2\text{HPO}_4$, or $\text{NaOH}/\text{H}_3\text{BO}_3$ (0–20 mM total buffer concentration). The turbidity of LV prepared in buffer remained constant over 24 h for the entire pH range tested (3.0–10.0). The temperature of LV dispersion inside a quartz cuvette was varied by using a Polytemp circulating bath (Model 80T, constant-temperature circulator, Polyscience Corp., Niles, IL) connected to a thermostat-regulated cuvette holder in a Varian DMS-70 spectrophotometer. The temperature inside the cuvette was continuously monitored with an NiCr/Ni thermocouple connected to a previously calibrated digital thermometer (0.1 °C accuracy). Turbidity at 400 nm was recorded as a function of temperature. For samples prepared in the absence of buffer, basic pH values inside the cuvette were kept constant by using an N_2 atmosphere. For control, the pH value was measured before and after each temperature cycle. The temperature variation rate was constant and equal to 1 °C min^{-1} .

Determination of Vesicle Shrinkage. Equal volumes of buffered LV (ca. 0.2 mM) and of a 0.8 M *meso*-erythritol buffered solution were rapidly (5–10 s) mixed in a cuvette. Turbidity at 400 nm was recorded in a Varian DMS-70 spectrophotometer as a function of time at 25 °C. From the turbidity kinetics, initial shrinking rates (v_s) were calculated as¹⁷

$$v_s = dA/dt|_{t \rightarrow 0} \times 1/P_i \text{ s}^{-1} \text{ M}^{-1}$$

where dA/dt is the slope of the turbidity (A) vs time (t) curve at $t \sim 0$ normalized so as to be nondependent on vesicle concentration (P_i).

The total extent of shrinkage (E) was taken as¹⁶

$$E = (A_f - A_0)/(A_f A_0 P_i) \text{ M}^{-1}$$

where A_f corresponds to the turbidity value 24 h after mixing the vesicles and the *meso*-erythritol solution, A_0 is the initial absorbance value at $t \sim 0$, and P_i is the DHP concentration determined as inorganic phosphate.

Buffers used were NaAc/HAc, $\text{NaH}_2\text{PO}_4/\text{Na}_2\text{HPO}_4$, or tris(hydroxymethyl)aminomethane (3–20 mM). At a given pH and buffer concentration, at least four independent shrinkage kinetics were obtained. v_s and E were calculated and plotted as mean values with the standard deviation as an error bar.

Determination of Colloidal Stability. A 0.1-mL sample of DHP vesicles (2 mM) was added to 0.9 mL of NaCl solution at a given pH. Turbidity at 400 nm as a function of time was followed in a Varian DMS-70 spectrophotometer. From these curves, initial flocculation rates (v_0), in s^{-1} , were calculated.¹¹ To avoid osmotic effects, the isoosmolarity between the inner and outer vesicle compartments was maintained for each salt addition. The vesicles were prepared in a D-glucose solution, and isoosmolal buffered NaCl/D-glucose solutions were added to obtain flocculation. For control, the pH was measured after each flocculation kinetics determination. The total Na^+ concentration (C) was calculated taking into account the NaCl concentration, the buffer concentration, and the NaOH eventually added to adjust pH. $\log 1/v_0$ was plotted as a function of $\log C$ for a range of pH values.

Results

Large DHP vesicles obtained by chloroform vaporization (LV) in water are spherical with a mean external diameter of 270 nm and scatter light according to the Joebst law.⁶ In spite of the importance of medium composition in determining the nature and size of amphiphile aggregates,^{1–3} this dependence is still undetermined for LV. For a range of pH and C , we determined here whether LV obeys the Joebst law for light scattering.¹⁴ Turbidity spectra of LV at pH 3–10 and $C = 0$ –20 mM NaCl are linear functions of $1/\lambda^2$ (not shown). Therefore, the particles in the DHP dispersion are large with a mean external diameter in the range 220–700 nm¹⁵ for the range of pH and C tested. Any variation

Figure 1. Normalized slope (NS) from turbidity spectra obtained for LV as a function of pH. NaCl concentrations are 3 (A), 8 (B), and 16 mM (C).

in the normalized slope (NS) of this linear function indicates variation in particle size (see Material and Methods). For vesicles prepared at very low C (0–3 mM NaCl), NS displays a bell-shaped profile as a function of pH, being maximal around pH 5.5–6.5 (Figure 1A). The apparent pK of the phosphate polar head (pK_a) for LV in water was determined from potentiometric titrations and found to be 5.8–6.5 (not shown). Thus, the largest vesicle size at $C = 3$ mM occurs at half-dissociation of the phosphate polar head (Figure 1A).

The increase in particle size from pH 3.5 to 5–6 at low salt concentrations (Figure 1A) is confirmed by entrapment (ENT) (not shown) and phase behavior data. After proper annealing¹⁸ at pH 4.8 and 8 mM NaCl, the NS and ENT values increase, approximately attaining the values observed at pH 6.0–7.0 (not shown).

Up to pH 6.5, the vesicle size decreases to a minimal value at pH 9–10 (Figure 1A–C).

As illustrated in Figure 1A–C, the effect of NaCl (3–20 mM) is to decrease the maximal vesicle size.

Optical properties of phospholipid²³ or amphiphile^{6,8,9,11} vesicles such as turbidity are temperature-dependent, serving as a simple and nondisturbing physical method for determining the phase behavior of LV for a range of pH and C .

The effect of temperature on LV turbidity is pH-dependent (Figure 2). In water, at pH 4.8, there is a completely reversible turbidity change associated with the main transition and a small anomalous change several degrees below the main transition that has a real hysteresis (Figure 2A). At pH 6.5, a rather anomalous temperature effect on turbidity is observed (Figure 2B). Upon heating, an extensive and reversible turbidity increase abruptly occurs around 70–73 °C. The extent of this temperature-induced vesicle aggregation (Figure 2B) can be remarkably reduced by the addition of NaCl at low concentrations (3–20 mM) (Figure 3).

In water, at pH 8.0, the usual steep decrease in turbidity upon heating characterizes the main transition whereas the pronounced hysteresis (Figure 2C) typically compares with that observed for

(18) Lawaczek, R.; Kainosho, M.; Chan, S. I. *Biochim. Biophys. Acta* **1976**, *443*, 313.

(19) Yi, P. N.; MacDonald, R. C. *Chem. Phys. Lipids* **1973**, *11*, 114.

(20) Elamrani, K.; Blume, A. *Biochim. Biophys. Acta* **1983**, *727*, 22.

(21) Bittman, R.; Blau, L. *Biochemistry* **1972**, *11*, 4831.

(22) Reerink, H.; Overbeek, J. T. G. *Discuss. Faraday Soc.* **1954**, *18*, 74.

(23) Traueble, H.; Teubner, M.; Woolley, P.; Eibl, H. *Biophys. Chem.* **1976**, *4*, 319.

(24) Traueble, H.; Eibl, H. *Proc. Natl. Acad. Sci. U.S.A.* **1974**, *71*, 214.

(17) Bangham, A. D.; De Gier, J.; Greville, G. D. *Chem. Phys. Lipids* **1967**, *1*, 225.

Figure 2. Effect of temperature on LV turbidity at 400 nm at three different pH values: 4.8 (A), 6.5 (B), and 8.0 (D). Heating is represented by full circles and cooling by empty circles. DHP concentrations are 0.10 (A), 0.32 (B), and 0.34 mM (C). LV were prepared in water, and pH 8.0 was maintained by using an N_2 atmosphere.

charged bilayers.²⁵ Some LV aggregation is suggested by the lower turbidity value obtained after the heating/cooling cycle (Figure 2C). When NaCl (3–20 mM) was added to the LV preparation medium, this difference in turbidity completely vanished (not shown).

From curves such as those in Figure 2, the midpoints of the main phase transition (T_c) for the heating runs were plotted as a function of the bulk pH (Figure 4). The T_c values are maximal around pH 5.8–6.4 (Figure 4). Inducing half a charge on DHP bilayers results in the formation of a bilayer structure with a T_c around 73 °C, i.e., 11 °C above the T_c of the fully protonated DHP bilayer in LV (Figure 4).

Up to pH 6.5, the T_c value remarkably decreases as a function of pH (Figure 4). The LV size decreases in this same pH range (Figure 1A). At pH 8.0, by preparing LV samples at pH 8.0 and different C values (3–12 mM) and measuring the midpoint of the phase transition (T_c), it was observed that the T_c value decreases from 69 °C at 3 mM NaCl to 62 °C at 12 mM NaCl (not shown).

When the effects of pH on LV size (Figure 1A) and on T_c (Figure 4) are compared, the conclusion is that T_c increases with increasing NS.

The water permeability (P) in a one bilayer vesicle with a radius r is given by²⁰

$$P = kr/3 \quad (5)$$

Figure 3. Effect of temperature on LV turbidity at 400 nm at pH 6.5 and 3 different NaCl concentrations: 0 (A), 3.3 (B), and 14.1 mM (C). Heating is represented by full circles and cooling by empty circles. DHP concentration is around 0.1 mM.

Figure 4. Effect of pH on the phase transition temperature (T_c) of LV prepared in water. T_c values were taken from heating curves such as those in Figures 2 and 3.

where k is the water permeation rate constant. Upon imposing an osmotic gradient to the LV walls, the initial shrinking rate (v_s) can be directly related to P .^{17,21}

v_s values for LV display a bell-shaped profile as a function of pH, being maximal at pH 6.0–7.0 (Figure 5). The effect of decreasing the bilayer packing (T_c) is to increase k , thereby increasing P .²¹ If v_s were commanded just by k , an inverted-bell-shaped curve would describe the behavior of v_s as a function of pH. This is not the case in Figure 5. Thus, P seems to reflect variations in r . Remembering the vesicle size behavior as a function of pH (Figure 1), it is clear that v_s is mainly controlled by vesicle size (Figure 1).

The effect of increasing salt is to decrease the maximal v_s value (Figure 5A–C). This can be associated with the decrease in the

Figure 5. Effect of pH on the initial shrinking rate (v_s) of LV at three different NaCl concentrations: 3 (A), 9 (B), and 16 mM (C).

maximal vesicle size upon increasing salt (Figure 1A–C).

The total extent of shrinkage (E) is a parameter directly related to vesicle size.¹⁷ E displays a bell-shaped profile as a function of pH with maximal values around the pK_a (not shown), confirming the behavior of vesicle size as a function of pH obtained from the turbidity spectra (Figure 1).

At the pH of water and at a given C , LV are more stable than SV as inferred from flocculation data obtained for both dispersions.^{6,12} For LV, extensive and observable intervesicle interactions leading to flocculation begin after the addition of up to 60–80 mM NaCl at the pH of water.^{6,12} Here we extend the determination of the colloidal stability of LV for a range of pH.

The initial flocculation rate (v_0) for LV depends on pH and C (Figure 6). v_0 increases with C until the critical coagulation concentration (ccc) is attained (Figure 6). Up to the ccc, a further increase in C does not change v_0 . As predicted by the DLVO²¹ $\log v_0$ is a linear function of $\log C$ (Figure 6). In general, the LV system behaves as colloidal particles flocculating upon salt addition at rates lower than those previously reported for SV.⁹

Salt is not very effective in inducing flocculation at low pH (Figure 6A–C). This is further emphasized by the large and sometimes nonobservable ccc values at low pH (Figure 6A–C). At moderately high ionic strength (0.16–0.25 M NaCl), the highest values of colloidal stability occur at low pH (Figure 6A–C).

The colloidal stability of LV taken as $\log 1/v_0$ displays a bell-shaped profile as a function of pH (Figure 7A). When C is increased, the pH range of maximal stability decreases (Figure 7A).

From the DLVO,²² the colloidal stability should decrease with decreasing surface charge and potential. By calculating the slope of the $\log 1/v_0$ vs $\log C$ curves for LV (Figure 6), some estimates of the surface potential can be obtained. The slope decreases as a function of pH at $C = 0.16$ – 0.25 M (Figure 7B) following the profiles of the colloidal stability as a function of pH (Figure 7A). Therefore, at moderate NaCl concentrations, colloidal stability decreases when the surface potential decreases.

Discussion

In this section we address two major questions: (1) the significance of specific pH and salt effects on the properties of DHP vesicles; (2) how these effects compare with those previously reported for bilayers with ionizable headgroups.^{23–27} It is demonstrated that theories based on electrostatics do not predict the dependence of DHP vesicle properties on pH and monovalent salt at low concentration (0–20 mM).

The increase in size from pH 3.5 to 5–6 at low C (Figure 1A,B) might be related to the low stability of the DHP dispersion prepared by chloroform vaporization at low C and pH.

On the other hand, at low pH and C , the nature of the LV dispersion is not unequivocally determined. Although the Joebst law is obeyed by LV under these conditions, i.e., particles are spheres, the existence of multilamellar vesicles¹⁶ cannot be dismissed. Furthermore, light-scattering data for DHP ultrasonically dispersed in water (SV) at low pH show a two-peak distribution for particle size.⁵ This may actually be related to the existence of two different kinds of particles in solution at low pH. We are presently investigating the nature and morphology of the DHP dispersion by electron microscopy for a range of pH and C .¹⁴ For dipalmitoylphosphatidylcholine (DPPC) multilamellar liposomes, a similar phase behavior was reported and interpreted as arising from the interactions between close and concentric bilayer sheets.²⁸ In fact, the introduction of charged lipophilic substances in the DPPC multilamellar liposomes induced the pretransition to disappear by expanding the distance of the adjoining bilayers and reducing the interactions between them.²⁸ Thus, it is possible that the DHP dispersion at low pH and in water is composed of multilamellar liposomes. Due to the fully protonated state of the DHP bilayer at pH 4.8 and the absence of Na^+ (dispersion prepared in water), the distance between adjoining DHP bilayers would be indeed small.

The maximal number of intrabilayer hydrogen bridges occurs at half-dissociation. For some phospholipid analogues, a high capacity for hydrogen bonding has been associated with a high degree of hydrophobicity at the bilayer surface.^{29,30} Around the pK_a , the DHP vesicle surface should have the highest degree of hydrophobicity.^{7,9} Therefore, the smallest exposure of polar heads and the lowest total surface area of the DHP dispersion are to be expected at the pK_a . Consistently, the largest particle vesicle size and T_c were observed at the pK_a . Above the pK_a at low C , the vesicle size and T_c decrease (Figures 1 and 4) whereas at the air–water interface, a DHP monolayer expands upon a pH increase at low C (0.1–10 mM) in the subphase.⁷ After the pK_a region, both sets of results possibly indicate binding of hydrated Na^+ ions and an increase in DHP surface hydration occurring at pH 9–10. In fact, a short-ranged repulsive hydration force was directly measured between DHP surfaces at pH 9–10.⁷ Associated with this surface hydration, the smallest vesicle size as well as the lowest phase transition temperature was determined (Figures 1 and 4).

It is of interest to recall that the decrease in the packing density of the DHP bilayer (T_c) from pH 7.0–7.5 to pH 9–10 (Figure 4) cannot be associated with an increase in the charge density since after the pK_a region the charge density is supposed to be constant.

For lipids, the higher the degree of surface hydration, the lower the T_c .²⁷ When surface hydration is increased, a repulsive intralayer and interlayer hydration force⁷ may reduce vesicle size (Figure 1), increase the total exposed surface area, expand the DHP monolayer,⁷ and reduce the tightness in the packing of DHP monomers in the DHP bilayer (Figure 4).

For DPPC single lamellar liposomes a direct relationship between the midpoint of the phase transition and liposome size was described and interpreted as due to packing constraints associated with the curvature radius.²⁸ One of these constraints could well be the degree of surface hydration that is able to change vesicle size and T_c .

The maximal vesicle size (Figure 1) and the extent of LV aggregation at 70–73 °C (Figure 3) decrease with increasing NaCl concentration at pH values around the pK_a . In a DHP monolayer at pH 5.5, upon increasing NaCl concentration in the subphase the area per monomer increases.⁷ All those effects may be due to Na^+ adsorption. Na^+ is possibly able to disrupt some intermolecular hydrogen bridges, adsorbing and rendering the surface more hydrophilic. For phospholipid analogues, the steep increase

(26) Blume, A.; Eibl, H. *Biochim. Biophys. Acta* **1979**, *558*, 13.

(27) Cevc, G.; Seddon, J. M.; Marsh, D. *Faraday Discuss. Chem. Soc.* **1986**, *81*, 179.

(28) Takemoto, H.; Inoue, S.; Yasunaga, T.; Sukigara, M.; Toyoshima, Y. *J. Phys. Chem.* **1981**, *85*, 1032.

(29) Silvius, J. R.; Brown, P.; O'Leary, T. J. *Biochemistry* **1986**, *25*, 4249.

(30) Brown, P. M.; Steers, J.; Hui, S. W.; Yeagle, P. L.; Silvius, J. R. *Biochemistry* **1986**, *25*, 4259.

Figure 6. Effect of Na^+ concentration (C) on the initial flocculation rate (v_0) of LV at several pH values: 3 (A), 4 (B), 5 (C), 6 (D), 7 (E), 7.8 (F), 9 (G), and 10.2 (H). Final DHP concentration in the cuvette is 0.20 mM. Curves are plotted as $\log v_0$ against $\log C$.

Figure 7. Effect of pH on the colloidal stability of LV taken as $\log 1/v_0$ (A). Na^+ concentration is 0.16 (a), 0.20 (b), and 0.25 M (c). In (B), the effect of pH on the surface potential estimates taken as the slopes of the $\log v_0$ against $\log C$ curves in Figure 6.

in light scattering occurring at a high temperature has been associated with a low degree of surface hydration and a high capacity for hydrogen bonding.^{29,30} The occurrence of this "high-melting" solid phase was also observed for SV at and around the pK_a and attributed to temperature-induced vesicle aggregation with formation of the isotropic phase at linkages between vesicles at 70–73 °C.⁹ Here we observe an interesting effect of NaCl at low concentration, namely, the ability to reduce the high-temperature vesicle aggregation (Figure 3) and maximal vesicle size (Figure 1) possibly via hydrated Na^+ adsorption with increase in the area per monomer at the bilayer surface and increase in total surface area of the DHP dispersion.

Curiously, precisely at the pK_a , when the surface hydrophilicity of the membrane is lowest, the highest water permeability is observed (Figure 5). The gross structural intrabilayer organization determined by vesicle size seems to be more important in determining water permeability than surface hydrophilicity.

The low colloidal stability of DHP vesicles in the presence of salt at the pH of water remarkably contrasts with that of phospholipid systems.^{6,12} This high sensitivity to monovalent salt can be better understood from some properties exhibited by phosphatidic acids (PA), for example. The first pK_a of PA bilayers

is ca. 3.5 in water.²⁶ The pK_a values of DHP bilayers in form of SV or LV are respectively 5.5–6.0⁹ and 5.8–6.3 in water. Therefore, the pK_a value for LV or SV is higher than the first pK_a for PA bilayers. The fact that the phosphate dissociation occurs first for PA suggests that the PA bilayer surface is more hydrated than DHP bilayers which favors the ionized state for the PA phosphate. Hydrophilic polar groups such as the glycerol backbone and the ester linkages present in PA but absent in the DHP could be associated with the higher surface hydration of PA. The sensitivity of the DHP bilayer surface to salt could be related to a low degree of surface hydration.

At low C (0–20 mM) and pH, Na^+ ion adsorption increases the surface charge density in adsorbed DHP monolayers reducing adhesion,⁷ whereas at $\text{pH} \geq \text{pK}_a$, Na^+ increases the degree of surface hydration reducing vesicle aggregation (Figure 3 and Results section). In both cases, Na^+ ion at low concentration promotes, by different mechanisms, an increase in the colloidal stability of DHP systems.

At a moderate C range (0.16–0.25 M), the largest colloidal stability is observed at low pH ranges (Figure 7A). The main reason for this could be a facilitation in phosphate dissociation caused by salt, similar to that described for PA at higher C ranges.²³ Upon increasing C , the pK_a region would be shifted to lower pH values. The colloidal stability is high (Figure 7A) for the same pH range where the surface potential is high (Figure 7B). At least qualitatively, the surface potential can be considered as an important factor determining the colloidal stability of LV at moderate ionic strength, in accordance with DLVO predictions.²²

The phosphatidic acid (PA) system is apparently a matter of controversy in the literature.^{22–25} The controversy is related to effects of charge alteration in the phase transition temperature (T_c) and addresses the question of whether an increase in surface charge always results in a decrease of the T_c . Three major points clarify significantly the PA confusion: (1) the fatty acid ester of the secondary hydroxyl of glycerol is labile,²⁵ and titration curves (T_c vs pH) obtained by Traueble et al.²³ correspond to titration of some hydrolysis product; (2) stable ether analogues of PA²⁴ yield maximal T_c values at the first or at the second pK_a of the phosphate polar head;^{25,26} (3) just before the first pK_a , the T_c value increases upon a bilayer charge increase.^{25–27}

The results for LV in water corroborate those in refs 25–27, since the pK_a range in water from titration closely corresponds to the maximal T_c values (Figure 4). Very recently, we demonstrated for SV that the surface potential (and surface charge) does not display the expected sigmoidal titration curve with the apparent pK value of the phosphate at the surface as midpoint.⁹ Membrane structure determines the value that the surface potential will assume.⁹ Here we point out an important factor controlling

membrane structure at low salt (0–20 mM), namely, the surface hydration. Surface hydration is quite different at the pK_a or at pH 9–10.⁷ When it is high (at pH 9–10), the T_c is low and the vesicle is small. When it is low (at the pK_a), the T_c is high and the vesicle is large. Electrostatics seems to display a secondary role in determining the membrane structure at low salt because upon an increase in surface charge the T_c can increase (before the pK_a) or decrease (after the pK_a).

Conclusions

At low salt (0–20 mM NaCl), pH and salt effects on DHP vesicle properties such as size, phase behavior, and colloidal stability can be associated with a variable degree of surface hydration at the DHP bilayer.

At half-dissociation of the phosphate polar head in water ($pK_a = 5.8$ – 6.3), the degree of surface hydration is lowest due to the occurrence of the maximal number of hydrogen bridges between phosphate polar heads. As a consequence, the surface area exposed to the water solution is lowest, generating the largest vesicle size and the highest phase transition temperature (T_c) over the entire range of pH tested (3–10). At pK_a , monovalent salt (3–20 mM) can disrupt hydrogen bridges, increasing surface hydration, generating smaller vesicles, and extensively decreasing vesicle aggregation at 70–73 °C.

At pH $> pK_a + 1$, hydrated Na^+ ion adsorption (0.1–20 mM) plays an important role in providing surface hydration, reducing vesicle size, T_c , and the extent of intervesicle aggregation at room temperature.

At pH $< pK_a - 1$, the presence of multilamellar structures in water ($C = 0.1$ mM) is strongly suggested by phase behavior data.

Precisely when the DHP membrane possesses its highest degree of surface hydrophobicity (at the pK_a), water permeability is maximal. With increasing C , at the pK_a , maximal water permeability decreases due to a decrease in vesicle size.

Surface hydration is apparently not important in determining the colloidal stability of LV at moderate monovalent salt concentration (0.16–0.25 M), but ion adsorption significantly reduces vesicles aggregation, increasing colloidal stability at very low salt (0–20 mM).

Acknowledgment. Financial support from the Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP), the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), and the Third World Academy of Sciences (TWAS) is gratefully acknowledged. S. Hix is the recipient of a CNPq undergraduate fellowship. We thank one of the referees for deep comments and valuable criticism.

Registry No. Na(DHP), 60285-46-3; NaCl, 7647-14-5.

Protein Molecular Dynamics Constrained to Slow Modes. Theoretical Approach Based on a Hierarchy of Local Modes with a Set of Holonomic Constraints: The Method and Its Tests on Citrate Synthase

Jean Durup

Laboratoire de Physique Quantique (UA 505 of the CNRS), Université Paul Sabatier, 118, route de Narbonne, 31062 Toulouse, France (Received: April 23, 1990)

We present a new approach for theoretical simulation of protein dynamics constrained to slow modes in order to allow for an increase of the integration time step by an order of magnitude. It consists in building a hierarchy of internal vectors, according to the tree principle (generalization of Jacobi coordinates), and in using their polar coordinates, referred for each vector to the two vectors immediately lower in the tree, as local modes. Then the higher frequency modes are either fixed to their equilibrium values (holonomic constraints) or, as regards the cyclic coordinates which have no equilibrium value, affected with a friction coefficient. Conditions in the design of the tree are derived which allow the holonomic constraints to be applied as a correction after each step without needing an iteration. The method is tested on citrate synthase, a dimeric enzyme with 2×437 residues. The CHARMM-20 programs were used for topology, energy calculation, and basic dynamics. Comparisons between exact dynamics, 1-fs time step, and constrained dynamics, 8-fs time step, are presented in terms of the variations of some randomly chosen unconstrained coordinates along a 2.8-ps run, and in terms of statistics on average values and RMS fluctuations of all unconstrained coordinates in the same time lag. Finally, prospects for improvement and extension of this method are discussed.

I. Introduction

Although the functional properties of proteins are now fairly well understood from the structural viewpoint, the molecular specificities responsible for their remarkable dynamical behaviors are still largely a matter of speculation. The fact that catalytic proteins, i.e., enzymes, in particular, have been developed during primeval evolution through a gradual selection among 10^{100} – 10^{1000} possible nonfunctional polypeptides should warn us that very abnormal and specific structural or dynamical features might be of decisive importance in the control of biochemical processes.

These considerations explain why the theoretical approach of protein dynamics pioneered and developed by the group of Karplus and by his followers,^{1–6} resting upon a description as accurate as

possible of the macromolecule at the atomic level, is now accepted by most researchers in the field. Whenever comparisons were

(1) McCammon, J. A.; Gelin, B. R.; Karplus, M. *Nature* **1977**, *267*, 585. McCammon, J. A.; Wolynes, P. G.; Karplus, M. *Biochemistry* **1979**, *18*, 927. van Gunsteren, W. F.; Karplus, M. *Biochemistry* **1982**, *21*, 2259. Swaminathan, S.; Ichiye, T.; van Gunsteren, W.; Karplus, M. *Biochemistry* **1982**, *21*, 5230.

(2) Brooks, B. R.; Bruccoleri, R. E.; Olafson, B. D.; States, D. J.; Swaminathan, S.; Karplus, M. *J. Comput. Chem.* **1983**, *4*, 187.

(3) van Gunsteren, W. F.; Berendsen, H. J. C.; Hermans, J.; Hol, W. G. J.; Postma, J. P. M. *Proc. Natl. Acad. Sci. U.S.A.* **1983**, *80*, 4315.

(4) Levitt, M. *J. Mol. Biol.* **1983**, *168*, 595. Warshel, A. *Proc. Natl. Acad. Sci. U.S.A.* **1984**, *81*, 444. Levitt, M.; Sharon, R. *Proc. Natl. Acad. Sci. U.S.A.* **1988**, *85*, 7557.

Referência 14

formation has been observed by circular dichroism and X-ray studies.^{1,14} The α -CD molecule takes an unsymmetrically collapsed conformation to fit the cavity to two water molecules contained in the cavity. When the water molecules are replaced by other guest molecules in the cavity, the collapsed conformation changes into a circular unstrained conformation. β - and γ -CD's, however, are not strained^{1,15} and therefore, conformational volume changes of β - and γ -CD's upon inclusion-complex formation should be small. In the β -CD cavity, an average of 6.5 water molecules are involved.¹⁶ The vacancy of the β -CD cavity is almost fully occupied by DTBN;¹ hence, all water molecules in β -CD may be excluded from the CD cavity upon inclusion-complex formation. The 6.5 water molecules excluded give rise to the increase in volume by $\Delta V_{\text{push}} = 117 \text{ cm}^3 \text{ mol}^{-1}$. Accordingly, the sum of $\Delta V_{\text{desolv}} + \Delta V_{\text{includ}}$ becomes $-112 \text{ cm}^3 \text{ mol}^{-1}$ by eq 6. DTBN in CD rotates about the symmetry axis of CD as mentioned above, thus DTBN in CD is regarded as a cylinder in shape whose height and diameter were tentatively estimated to be 0.88 and 0.62 nm, respectively, with the aid of the Corey-Pauling-Koltun (CPK) model. Since the height of β - and γ -CD's is 0.80 nm,¹ DTBN should stick out of CD by 0.08 nm. The van der Waals volume of this part is then calculated to be $14.5 \text{ cm}^3 \text{ mol}^{-1}$. It is generally known that the partial molar volume of a species in solution is about twice as large as its van der Waals volume. Thus, the partial molar volume of the excluded part is estimated to be $29 \text{ cm}^3 \text{ mol}^{-1}$. If we subtract this from the partial molar volume of DTBN in water, $159 \text{ cm}^3 \text{ mol}^{-1}$, obtained experimentally in this work, ΔV_{includ} becomes $-130 \text{ cm}^3 \text{ mol}^{-1}$. Thus, the volume change due to the desolvation of DTBN upon inclusion ΔV_{desolv} is estimated to be $18 \text{ cm}^3 \text{ mol}^{-1}$ by using eq 6. In the cation/crown-ether complex,¹⁷

the reaction volumes ΔV in water have been estimated to be $8\text{--}25 \text{ cm}^3 \text{ mol}^{-1}$, and Harada et al.¹⁸ have pointed out that the main contribution to ΔV is the removal of water molecules from around guest ions. In our case, DTBN interacts with surrounding water molecules through dipole-dipole interaction as well as hydrogen bonding. Hence, the above value of ΔV_{desolv} seems to be reasonable, though the above estimations were carried out on the basis of some assumptions.

In case of γ -CD, 12 water molecules are originally situated in its cavity,¹⁹ and all or a part of the water molecules are excluded according to the size of the guest molecule upon complex formation. In the present case, $\Delta V = 20 \text{ cm}^3 \text{ mol}^{-1}$ and $\Delta V_{\text{includ}} + \Delta V_{\text{desolv}} = -112 \text{ cm}^3 \text{ mol}^{-1}$. Therefore, on the average 7.3 out of 12 water molecules are replaced by DTBN upon complex formation.

In conclusion, DTBN is shown to be included more tightly in β -CD than in γ -CD on the basis of the values of $1/\tau_{\perp}^{\text{rel}}$ and K for the inclusion complexes. The water molecules remaining in the γ -CD cavity upon complex formation interact with the $>\text{N}-\text{O}$ group of DTBN as well as the inner wall of CD through hydrogen bonding. This will cause a restriction of free rotation about the symmetry axis, resulting in the reduced $1/\tau_{\perp}^{\text{rel}}$ in the γ -CD cavity. The presence of polar water molecules in the CD cavity will favor the ionic form (b) in the following canonical resonance structures, which causes an increase in the nitrogen hyperfine coupling constant of DTBN in γ -CD.²⁰

Registry No. α -CD, 10016-20-3; β -CD, 7585-39-9; γ -CD, 17465-86-0; DTBN, 2406-25-9; water, 7732-18-5.

- (14) Rees, D. A. *J. Chem. Soc. B* 1980, 877.
 (15) Holand, H.; Hald, L. H.; Kvamner, O. R. *J. Solution Chem.* 1981, 10, 775.
 (16) Lingner, K.; Saenger, W. *Angew. Chem.* 1978, 90, 738.
 (17) Høiland, H.; Ringseth, J. A.; Vilkingstad, E. *J. Solution Chem.* 1978, 7, 515. Høiland, H.; Ringseth, J. A.; Brun, T. S. *J. Solution Chem.* 1979, 8, 779.

- (18) Harada, S.; Sahara, H.; Nakagawa, T. *Bull. Chem. Soc. Jpn.* 1983, 56, 3833.
 (19) MacLennan, J. M.; Stezowski, J. J. *Biochem. Biophys. Res. Commun.* 1980, 92, 926.
 (20) Kawamura, T.; Matsunami, S.; Yonezawa, T. *Bull. Chem. Soc. Jpn.* 1966, 40, 1111. Zger, S. A.; Freed, J. H. *J. Chem. Phys.* 1982, 77, 3344.

Bilayer Structure and Stability in Dihexadecyl Phosphate Dispersions

doc 85

A. M. Carmona-Ribeiro,*[†] C. E. Castuma,[‡] A. Sesso,[§] and S. Schreier[†]

Departamento de Bioquímica, Instituto de Química, Universidade de São Paulo, CP 20780 São Paulo, Brazil, Instituto de Investigaciones Bioquímicas, UNLP, CONICET, Facultad de Ciencias Médicas, 60 y 120, 1900, La Plata, Argentina, and Departamento de Patología, Faculdade de Medicina, Universidade de São Paulo, São Paulo, Brazil (Received: December 13, 1990)

The pH-dependent structure and stability of dihexadecyl phosphate (DHP) bilayer dispersions are determined by using fluorescent and spin probes. The hydrocarbon core region is probed by diphenylhexatriene (DPH) while the polar head/water interface is probed by either *trans*-parinaric acid (TPA) or an iminoxyl derivative of stearamide spin-label (SSL). At 5 mM NaCl, the fluorescence depolarization (FP) of DPH or TPA goes through a maximum as pH increases for vesicles prepared and maintained at a given pH value. On the other hand, if the pH is changed after vesicle preparation, as during titration, FP against pH is a typical sigmoidal curve. SSL in large vesicles (LV) exhibits a higher degree of spectral anisotropy than in small vesicles (SV) which is interpreted as a deeper penetration of SSL in the LV bilayer due to its higher fluidity. The SV bilayer fluidity increases as a function of time after sonication. From electron microscopy, the number of bilayer fragments in the SV dispersion decreases with increasing pH. For LV up to pH 6, however, the bilayer structure remained unchanged as a function of time. The degree of ionization of the headgroups is suggested to be important in determining the nonstationary bilayer structures occurring during titration. Nevertheless, for stationary DHP bilayers, headgroup hydration and hydrogen-bonding ability are important factors in determining the bilayer structure and vesicle size.

Introduction

Bilayers bearing ionizable polar heads have been a matter of some controversy in the literature.¹⁻⁷ The effect of increasing

charge on the bilayer structure has been proposed to be a decrease in the bilayer packing density in earlier work^{1,2} and an increase

* To whom correspondence should be addressed.
[†] Instituto de Química, Universidade de São Paulo.
[‡] Instituto de Investigaciones Bioquímicas.
[§] Faculdade de Medicina, Universidade de São Paulo.

(1) Traueble, H.; Teubner, M.; Woolley, P.; Eibl, H. *Biophys. Chem.* 1976, 4, 319.
 (2) Traueble, H.; Eibl, H. *Proc. Natl. Acad. Sci. U.S.A.* 1974, 71, 214.
 (3) Eibl, H.; Blume, A. *Biochim. Biophys. Acta* 1979, 553, 476.
 (4) Blume, A.; Eibl, H. *Biochim. Biophys. Acta* 1979, 558, 13.

followed by a decrease in more recent papers.³⁻⁷ Pure electrostatics would predict a sigmoidal shape for the bilayer packing as a function of pH with the bulk pH value at half-dissociation (pK_a) as the midpoint. The sigmoidal shape would be expected for pH values between $pK_a - 1$ and $pK_a + 1$ whereas the bilayer packing would have to remain constant for pH values outside this region. However, we recently demonstrated that other factors like the ability to form hydrogen bonds or the change in the degree of the surface hydration because of binding of hydrated counterions can completely alter the picture predicted from the surface electrostatics.⁶⁻⁸ For the dihexadecyl phosphate (DHP) system, several bilayer properties like the phase behavior, the vesicle size, and the interbilayer interactions vary remarkably depending on the ionization of the headgroups.^{6,7} At the pK_a , the phase transition temperature (T_c) and vesicle size were found to be a maximum.⁷ At full dissociation (high pH), the equilibrium bilayer structure in the large DHP vesicles has a larger total surface area not only because all the polar heads are charged but also due to binding of hydrated Na^+ at the negatively charged headgroups.^{7,8} Thus, if the surface hydration is higher, the total surface area is larger, the vesicle is smaller, and the T_c is lower.⁷ Furthermore, in contrast to predictions from pure electrostatics, at the pK_a of the phosphate polar head, the tightest packing was observed.⁷ Here we further explore the DHP system to clarify the effect of charge on the bilayer structure. Our overall approach is to monitor the pH effect on bilayer fluidity and packing density as seen from fluorescent and spin probes incorporated in the DHP bilayer. The DHP bilayer systems were obtained either by sonication (SV) or by chloroform vaporization (LV), and fluorescence depolarization (FP) or electron spin resonance (ESR) spectra of probes in the DHP bilayer were determined as a function of pH and time. The sigmoidal shape of FP obtained during titration is consistent with an electrostatic control of fluidity, but the stearamide ESR spectra for bilayer structures under titration before and after annealing clearly demonstrate that bilayer structures obtained during titration are nonstationary. The fluidity profile as a function of pH obtained for stationary DHP bilayers requires electrostatics, hydrogen bonding, and surface hydration to be understood. In addition, due to the polemic nature of amphiphile dispersions obtained by sonication,⁹⁻¹¹ we also monitored the nature and shape of the SV dispersion as a function of pH using electron microscopy. The bilayer structure in the SV dispersion is found to be time dependent over the entire range of pH tested (pH 3–10). The reason for this is the existence of a decreasing but significant amount of bilayer fragments in the SV dispersion at increasing pH values. On the other hand, the LV bilayer structure is shown to be stable up to pH 6 at low NaCl concentration. This further establishes the chloroform vaporization method^{12,13} as the most reliable procedure for the production of closed steady-state amphiphile bilayers.

Material and Methods

Preparation of the DHP Dispersions. Dihexadecylhydrogen phosphate was from Sigma (St. Louis, MO), and it was used as provided for the experiments in buffered solutions. Sodium dihexadecyl phosphate (DHP) was obtained from dihexadecylhydrogen phosphate as previously described¹³ and used in the experiments performed with unbuffered solutions. LV were

prepared by injecting a solution of DHP in chloroform into an aqueous solution at 75 °C.¹³ SV were prepared by sonication as previously described.¹⁰ The DHP concentration was measured by inorganic phosphate determination.¹⁴ The buffers used were NaAc/HAc, NaH_2PO_4/Na_2HPO_4 , or $H_2BO_3/NaOH$ at a final Na^+ concentration of 5 mM. For the titration experiments SV or LV were prepared in unbuffered solutions.

Incorporation of Hydrophobic Probes in the DHP Bilayer. The chemical structure of the markers used is represented below. 1,6-Diphenyl-1,3,5-hexatriene (DPH) (Molecular Probes, Inc., Eugene, OR) monitors the structure at the bilayer core.¹⁵ *trans*-Parinaric acid (TPA) (Molecular Probes) senses the structure at the bilayer surface.¹⁶ The iminoxyl derivative of stearamide (SSL), a kind gift from Dr. Ian C. P. Smith, National Research Council, Canada (synthesized by Dr. H. Dugas from the University of Montreal), detects also the polar headgroup region.^{17,18} The incorporation either of fluorescent or spin labels in the bilayer was done by adding the label to the solution of DHP in chloroform used either to prepare LV¹³ or to make a DHP plus label film on the bottom of a sonication tube to prepare SV.¹⁰ For the FP measurements, the molar ratio marker/DHP was about 1:200 or less for DPH and 1:150 for TPA. For the ESR experiments, the molar ratio was about 1:100 or less.

Fluorescence Depolarization Measurements. The FP measurements were carried out at 25 °C with an Aminco Bowman equipped with two glan prisms for polarization measurements. Samples were excited with vertical polarized light, while vertical ($I_{||}$) and horizontal ($I_{\perp}$) emission intensities were recorded. After appropriate blank and instrumental corrections, the fluorescence depolarization (FP) was calculated.¹⁹ The instrumental anisotropy (G) was determined by measuring vertical and horizontal emission values using horizontally polarized light.²⁰

$$FP = [(I_{||}/I_{\perp})G - 1]/[(I_{||}/I_{\perp})G + 1]$$

For DPH and TPA, the excitation wavelengths were 357 and 320 nm, respectively. The emission was recorded at 400 and 430 nm, respectively. For measurements of FP in unbuffered samples, a nitrogen atmosphere maintained a stable pH. In order to check the stability of the bilayer structure, FP values were measured before and after a heating/cooling cycle. Samples under test were heated to 75 °C, a temperature above the phase transition,^{6,7} and allowed to cool to room temperature (about 25 °C).

Measurements of ESR Spectra. ESR measurements were carried out at 25 °C with a Bruker ER 200D-SRC spectrophotometer. To minimize vesicle aggregation, the maximum amphiphile concentration used was 2 mM DHP. However, the

(5) Cevc, G.; Seddon, J. M.; Marsh, D. *Faraday Discuss. Chem. Soc.* **1986**, *81*, 179.

(6) Carmona-Ribeiro, A. M.; Hix, S. J. *Phys. Chem.* **1991**, *95*, 1812.

(7) Carmona-Ribeiro, A. M. *J. Colloid Interface Sci.* **1990**, *139*, 343.

(8) Claesson, P. M.; Carmona-Ribeiro, A. M.; Kurihara, K. *J. Phys. Chem.* **1989**, *93*, 917.

(9) Tran, C. D.; Klahn, P. L.; Romero, A.; Fendler, J. H. *J. Am. Chem. Soc.* **1978**, *100*, 1622.

(10) Mortara, R. A.; Quina, F. H.; Chaimovich, H. *Biochem. Biophys. Res. Commun.* **1978**, *81*, 1080.

(11) Pansu, R. B.; Arrio, B.; Roncin, J.; Faure, J. J. *Phys. Chem.* **1990**, *94*, 196.

(12) Carmona-Ribeiro, A. M.; Chaimovich, H. *Biochim. Biophys. Acta* **1983**, *733*, 172.

(13) Carmona-Ribeiro, A. M.; Yoshida, L. S.; Sesso, A.; Chaimovich, H. *J. Colloid Interface Sci.* **1984**, *100*, 433.

(14) Rouser, G.; Fleischer, S.; Yamamoto, A. *Lipids* **1970**, *5*, 494.

(15) Shinitzky, M.; Barenholz, Y. *J. Biol. Chem.* **1974**, *249*, 2652.

(16) Sklar, L.; Miljanich, G. P.; Dratz, E. A. *Biochemistry* **1979**, *18*, 1707.

(17) Schreier, S.; Polnaszek, C. F.; Smith, I. C. P. *Biochim. Biophys. Acta* **1978**, *515*, 375.

(18) Koltover, V. K.; Reichman, A. A.; Yasaitis, A. A.; Blumenfeld, L. A. *Biochim. Biophys. Acta* **1971**, *234*, 306.

(19) Castuma, C. E.; Brenner, R. R. *Biochem. J.* **1989**, *258*, 723.

(20) Azumi, T.; Maclynn, S. J. *Chem. Phys.* **1962**, *37*, 2413.

Figure 1. Fluorescence depolarization of DHP dispersions as a function of pH. The full circles indicate measurements in dispersions obtained by chloroform vaporization (LV) whereas the empty ones indicate measurements in dispersions obtained by sonication (SV). All circles represent measurements done 4–8 h after DHP dispersion. In (A), the fluorescent probe is TPA and in (B), DPH. The samples were prepared and maintained in buffer solutions at 5 mM Na⁺. The final probe and amphiphile concentrations are respectively 0.0027 and 0.1 mM. Each point represents the mean value plus/minus the mean square deviation for five independent experiments run in duplicate. Single FP values measured for SV 1.5 h after sonication are represented by triangles. Measurements after annealing the SV samples are represented by squares.

spectrometer sensitivity does not permit detection of less than 0.005 mM nitroxide spin label undergoing restricted motion. Therefore, experiments were restrained between the limits 1–2 mM DHP and 0.005–0.04 mM SSL. For titration of unbuffered samples, a syringe setup and a pH electrode adaptor were coupled to the extremities of the flat quartz cell. Each cell extremity was therefore isolated from air during spectra acquisition. Acid or base addition was followed by a rapid homogenization of the sample and pH measurement. Some contamination due to SSL hydrolysis to stearic acid and tempamine was detected and quantified by integration as corresponding to 4%. In most experiments however the signal due to tempamine was so insignificant that it was not necessary to remove it by either spectral subtraction or dialysis.

Transmission Electron Microscopy. Equal volumes of 1 mM DHP SV vesicles 2–4 h after sonication in water at a given pH value and 1% ammonium molybdate were mixed on a copper grid previously covered with a parlodium film. Twenty seconds thereafter, the sample was dried and observed with a Philips 301 electron microscope operated at 80 kV. Magnifications were carefully calibrated with catalase crystals which have periodic fringes separated by a reticule distance of 85.5 Å. This allowed discrimination between thicknesses corresponding to one or two DHP bilayers.

Results

Dependence of the DHP Bilayer Fluidity on the Dispersion Method, on pH, and on Time. The fluorescence depolarization (FP) of *trans*-parinaric acid (TPA)¹⁶ and diphenylhexatriene (DPH)^{15,19} has been used to measure the fluidity of phospholipid²¹ and amphiphile bilayers.²² FP as a function of pH has a max-

Figure 2. Titration of DHP dispersions obtained by chloroform vaporization (O) or by sonication (□) in water at 1 mM Na⁺ and pH 6.9. The fluorescence depolarization of diphenylhexatriene in the bilayers is plotted as a function of pH. Titration was performed in the absence of buffer and under nitrogen. The time lag between each pH change was about 10 min. The final concentration of fluorescent label and amphiphile is respectively 0.0027 and 0.1 mM. Each point represents the mean value plus/minus the mean square deviation for five independent experiments run in duplicate.

imum at about pH 7 for LV and at about pH 6 for SV both in 5 mM NaCl (Figure 1). This maximum for LV corresponds to the pH region around 7.8 that is the pK_a value of the phosphate polar head at 5 mM salt as determined from potentiometric titrations of LV (not shown). FP values were also time dependent over the entire range of pH for SV. However, for LV the time dependency occurred only below pH 6 in 5 mM salt. In fact, by accelerating the attainment of equilibrium using a heating/cooling cycle, it was observed that the FP values for the SV dispersion could be considerably reduced (Figure 1). In contrast, the LV fluidity was not affected by temperature cycles up to pH 6. The bell-shaped dependence of fluidity on pH is replaced by a sigmoidal one when, instead of preparing different samples, each one at a given pH, the pH is varied in a given sample after preparation. The titration procedure yields sigmoidal curves like those in Figure 2. By heating and cooling at different pH values during titration, the nonequilibrium character of the bilayer structure thus obtained was also demonstrated by use of ESR spectroscopy.

The Nonequilibrium Bilayer Structure from a Spin Label Probing the Bilayer/Water Interface. The structural organization of the LV and SV DHP bilayer was also monitored by ESR spectroscopy of spin-labeled samples.¹⁷ The high packing density of DHP bilayers in the gel state⁸ has not allowed entrapment of bulky labels like the oxazolidyl derivatives of stearic acid with the nitroxide located at different positions along the hydrocarbon chain. Fortunately, incorporation of stearamide spin label (SSL), a highly hydrophobic marker with a bulky polar head, was achieved so that the bilayer structure at the bilayer/water interface could be assessed. Nevertheless, at low pH values, the DHP bilayer has its highest packing density^{6–8} so that because of its bulky headgroup the SSL molecule tends to be expelled from the bilayer with a consequent increase in mobility of the nitroxide located at the SSL polar head. This increase in mobility is to be seen in the ESR signal as an increased spectral isotropy and reflects an increase in packing density of the polar heads. This increase in packing density of the polar heads is also to be observed as a decrease in fluidity when the pH is lowered during titration (Figure 2).

The ESR signal of SSL in LV exhibits a higher spectral anisotropy than labeled SV (Figure 3). This result was obtained for the entire range of pH and is consistent with measurements that demonstrate a higher fluidity for LV than for SV compared just after preparation (Figure 1).

The SSL spectra in LV in 5 mM NaCl are markedly affected by the pH at which the LV are buffered (Figure 4). At pH 4.8 or 9.8, the spin label undergoes a more restricted motion than at pH 7. This correlates well with the highest fluidity at pH 4.8 or 9.8 (Figure 1) and is understandable from a higher penetration

(21) Duezguenes, N.; Wilschut, K.; Hong, R.; Fraley, C.; Perry, D. S.; Friend, D. S.; James, T. L.; Papahadjopoulos, D. *Biochim. Biophys. Acta* 1983, 732, 289.

(22) Rupert, L. A. M.; van Breemen, J. F. L.; Hoekstra, D.; Engberts, J. B. F. *N. J. Phys. Chem.* 1988, 92, 4416.

Figure 3. ESR spectra of SSL in LV (A) or in SV (B) prepared in phosphate buffer at 5 mM Na⁺ and pH 7. Final DHP and SSL concentrations are 1 and 0.005 mM, respectively.

Figure 4. ESR spectra of SSL in LV prepared at three different pH values. LV were prepared and maintained in different buffer solutions at 5 mM Na⁺ for a final DHP and SSL concentration of 1 and 0.005 mM, respectively.

of the label in the bilayer leading to a restriction in motion of the nitroxide. The low-intensity lines at pH 4.8 (Figure 4) can be associated with some flocculation that occurs at low pH and 1–2 mM DHP. This observation is consistent with the low stability of DHP bilayers (LV or SV) at pH values lower than 6 (see previous subsection).

The transient structures that occur in titration of an unbuffered LV or SV sample are illustrated by some ESR spectra in Figures 5 and 6, respectively. Their transient character is demonstrated by using the heating/cooling procedure coupled to ESR spectroscopy of SSL labeled samples before and after the temperature cycle (Figure 7). No significant spectral change occurs after a temperature cycle (*T*) in the LV sample prepared and maintained at pH 7 (Figure 7), but different spectra are observed before and after *T* for samples whose pH was changed from 7 to 3 or to 9.7 (Figure 7). The imposition of *T* in Figure 7 renders the spectra quite similar to those obtained in samples prepared and maintained at a given pH (compare spectra after *T* in Figure 7 with the corresponding ones in Figure 4). In contrast to the LV behavior, after undergoing a *T* cycle, labeled SV yield different spectra even if they are prepared and maintained at a given pH value (not shown). This agrees with the unstable character of the SV dispersion detected as a time-dependent fluidity over the entire range of pH (Figure 1). To understand further why the SV dispersion is unstable, unbuffered SV samples prepared at three different

Figure 5. Titration of a SSL labeled LV sample prepared at pH 6.7 in water. The time lag between each pH change was about 20 min. pH values are indicated besides each ESR spectrum. The final DHP and SSL concentrations are 2 and 0.04 mM, respectively.

Figure 6. Titration of a SSL labeled SV sample prepared at pH 6.2 in water. The time lag between each pH change was about 20 min. pH values are indicated besides each spectrum. The final DHP and SSL concentrations are 2 and 0.04 mM, respectively.

Figure 7. ESR spectra of a SSL labeled LV sample prepared at pH 7 in water. The structural stability of the LV bilayer was checked for three different pH values by annealing after the pH change. The pH value is on the right. DHP and SSL concentrations are respectively 2 and 0.04 mM.

pH values were observed by electron microscopy. At pH 3–4, most of the structures observed (about 50%) have a thickness of 53 Å (Figure 8A). Round shapes have quite undefined limits, and some layering of bilayers to form multibilayers occurs (Figure

Figure 8. Electron micrographs of SV samples prepared at different pH values in water taken 2–4 h after DHP dispersion by sonication. Negative staining allows visualization of the particles on a dark field. SV are at pH 3–4 (A, top), 6.3 (B, middle), and 8 (C, bottom).

8A). At pH 6.3 (Figure 8B) or pH 8 (Figure 8C) particles in dispersion are smaller than those observed at pH 3–4 (Figure 8A). In addition, the frequency of round structures increases whereas that of 53-Å thickness structures decreases as a function of pH.

Discussion

In this section two questions are addressed: (1) whether the degree of ionization of the polar heads by itself is able to determine the stationary bilayer structure; (2) whether it is relevant to titrate bilayers thereby generating nonstationary structures at a given vesicle size.

The dependence of the DHP bilayer fluidity on pH (Figure 1) is reminiscent of the dependence of the phase transition tem-

perature (T_c) on pH both for LV and for SV.^{6,7} The T_c and the vesicle size have a maximum at the pH corresponding to half-dissociation.^{6,7} Also, the vesicle size closely followed the T_c behavior as a function of pH for the LV dispersion.⁶ Therefore, the changes in fluidity (Figure 1) are associated with changes in vesicle size and in T_c . When the vesicle size increases, the T_c increases and the bilayer fluidity decreases. From pure electrostatics, the profile of the fluidity as a function of pH would be a sigmoidal increase as a function of increasing bilayer charge. At half-dissociation of the polar heads, an intermediate fluidity value would occur. At pH < $pK_a - 1$, the fluidity would be constant as a function of pH having a minimum value, whereas at pH > $pK_a + 1$, the fluidity would have a constant maximum value. Contrary to these expectations, we have recently demonstrated that, at half-dissociation, the bilayer packing (T_c)⁶ as well as the bilayer fluidity (Figure 1) is not an intermediate value but, in fact, is at a maximum over the entire range of pH values investigated. Because the maximum number of intrabilayer hydrogen bridges occurs at the pK_a , the surface hydrophilicity is considerably reduced; the area per monomer decreases and thereby also the total surface area; the T_c , the vesicle size, and the fluidity decrease. Again, the simple electrostatics model would predict a constant fluidity at pH > $pK_a + 1$. However, an increasing fluidity (Figure 1) and decreasing T_c and vesicle size⁶ were observed at this pH region, showing that other factors are significant. One of these factors can well be an increasing hydrated sodium ion adsorption at the bilayer surface with increasing pH at the pH > $pK_a + 1$ region. Thereby the surface hydration⁸ and the bilayer fluidity would increase (Figure 1). Examining Figure 2, it could be argued that this is the sigmoidal profile expected from electrostatics. In fact, assuming that the vesicle size was maintained during the titration only the changes due to ionization of the headgroups are expected to be observed (Figure 2). Upon increasing pH, the bilayer charge increases, the area per monomer increases as observed in DHP monolayers,⁸ the bilayer fluidity decreases (Figure 2), and the restriction in motion of SSL increases (Figures 5 and 6) due to its increased penetration in the bilayer being titrated. At pH values outside the pK_a region, no significant fluidity change occurs as predicted by the simple electrostatic model. However, the transient character of the bilayer structures obtained during titration (Figure 7) suggests that the factors determining stationary bilayer structures cannot be merely electrostatic. A possible implication of these results is that, depending on pH and ionic strength, different vesicle sizes are obtained for stationary bilayers,⁶ each one with a different pK_a . We are presently investigating the effect of vesicle size on the pK_a value of LV.

It is important to emphasize that the structures successively obtained during titration are not stationary ones. A straightforward procedure is suggested to establish whether a given bilayer structure or amphiphile aggregate is under stationary conditions. It consists basically of annealing the dispersion and monitoring a given structural property P. If P is the same before and after annealing, then the dispersion can be considered as stationary. This offers also a check of the ability of dispersion methods yielding stationary dispersions. The sonication procedure, for example, is clearly not adequate in producing stable DHP bilayers. In contrast, the chloroform vaporization method produced stable DHP bilayers up to pH 6.

The instability of the SV dispersion can be explained by the presence of bilayer fragments in the SV dispersion (Figure 8). The high-energy input during sonication is not only able to induce amphiphile dispersion and vesicle formation but also is well-known as a usual procedure for membrane disruption and, more recently, as an energy source for the generation of free radicals or to the induction of chemical reactions. Evidence of the presence of bilayer fragments in our electron micrographs cannot be seen as definitive because the electronic stain might have induced structural changes. However, other results also favor the bilayer fragment hypothesis: (1) the higher fluidity of LV in comparison with SV just after sonication (Figure 1); (2) the rapid increase in SV turbidity as a function of time after sonication in water;¹³ (3) the osmotic nonresponsiveness of dioctadecyldimethyl-

ammonium chloride SV and DHP SV;^{12,13} (4) the huge theoretical DLVO stability of SV in comparison with the low experimental one obtained from flocculation data;^{23,24} (5) the higher T_c values obtained for SV⁷ than for LV⁶ 2 h after dispersion. Concerning points 1 and 5, gel LV and SV of phospholipids behave differently. For phospholipids, not only the fluidity of SV is higher than that of LV²¹ but also the T_c for LV is always higher than the T_c for SV.²⁵ In fact, the planar structure of a DHP bilayer fragment could be more tightly packed than the bilayer structure in a large vesicle. Also, the hydrophobic interaction between edges of bilayer fragments could well account for the extra attractive interaction that was demonstrated between SV at different salt concentrations.^{23,24}

A pH change in a DHP bilayer dispersion in the gel state does not seem to change the surface hydration and thereby the vesicle

size. For the gross structural changes to occur, the bilayer for which the surface charge has changed has to undergo a phase transition. Further investigations about the effect of the bilayer state on the shape of the titration curves are desirable.

Conclusions

Including a charge in a bilayer structure by titration initializes the gross structural changes that will take place to produce a stationary bilayer structure at a given pH and ionic strength. The attainment of the stationary structure involves changes in the surface hydration able to lead to changes in vesicle size. The DHP bilayer can exist as a stationary structure only up to pH 6. The stationary character of a bilayer dispersion can be checked by monitoring a structural property before and after an annealing procedure. The DHP dispersion obtained by sonication is not stationary at any pH value.

Acknowledgment. Dr. Thelma M. Hardman, Dr. Brian R. Midmore, and the referees are gratefully acknowledged for valuable criticisms. This work was financially supported by CNPq, FAPESP, FINEP, TWAS, CONICET, and BID.

(23) Carmona-Ribeiro, A. M.; Yoshida, L. S.; Chaimovich, H. *J. Phys. Chem.* **1985**, *89*, 2328.

(24) Carmona-Ribeiro, A. M. *J. Phys. Chem.* **1989**, *93*, 2630.

(25) Takemoto, H.; Inoue, S.; Yasunaga, T.; Sukigara, M.; Toyoshima, Y. *J. Phys. Chem.* **1981**, *85*, 1032.

Referência 15

Synthetic Bilayer Adsorption onto Polystyrene Microspheres

A. M. Carmona-Ribeiro*[†] and B. R. Midmore

Department of Chemistry, University of Reading, Reading RG6 2AD, U.K.

Received August 9, 1991. In Final Form: November 25, 1991

Small and large vesicles prepared from dihexadecyl phosphate (DHP) and dioctadecyldimethylammonium chloride (DODAC) or bromide (DODAB) interact with oppositely charged polystyrene microspheres forming bilayer covered latices. Adsorption isotherms are of the Langmuir type, and for the three amphiphiles studied the limiting areas at the polystyrene/water interface are consistent with bilayer deposition. At limiting adsorption, electrokinetic properties of the covered particles were very similar to those of the vesicles. An innovative procedure for detecting bilayer adsorption based on determination of the mean z -average diameter of particles in the latex/vesicle mixtures as a function of amphiphile concentration is described.

Introduction

The interaction in solution of surfactants with polymers is of fundamental importance to several processes in chemical, petrochemical, and pharmaceutical industries to achieve a desired solubility, rheology, or colloidal stability.¹ As interfacial agents, the behavior of single-chain and double-chain surfactants with short hydrocarbon chains has been both intensively and extensively studied.² However, the potential of amphiphiles with long alkyl double chains as interface agents able to modify surface properties remains hitherto unexplored.

Dioctadecyldimethylammonium chloride (DODAC) or bromide (DODAB) and dihexadecyl phosphate (DHP) form vesicles in aqueous solution. These vesicles are perfect spherical shells composed of a bilayer enclosing an aqueous compartment.³⁻⁶ They are highly charged, in the rigid gel state at room temperature, and have a much smoother interface than other colloids. Because of these properties, they behave as model colloids and have been used to test theory.⁷⁻¹² However, their polydispersity is still a disadvantage when they are compared with highly homodisperse systems as polystyrene latices.¹³ On the other hand, the possibly hairy, rough, or conducting surface of these latices often represents a problem.¹⁴⁻¹⁶ Thus, the

interaction between synthetic amphiphile vesicles and oppositely charged polystyrene latices is interesting from two different points of view: (1) the production of homodisperse and smooth bilayer covered polystyrene microspheres to represent a model colloid and (2) the use of these surfactants with long alkyl double chains as interfacial agents that modify surface properties.

Here we demonstrate deposition of synthetic amphiphile bilayers onto polystyrene microspheres from adsorption isotherms and microelectrophoresis measurements. Also, photon correlation spectroscopy (PCS) was successfully used to detect bilayer adsorption on small particles.

Material and Methods

Dioctadecyldimethylammonium chloride (DODAC) was obtained from Herga Industrias Químicas S.A. (R. J., Brazil) and purified as previously described.⁸ Dioctadecyldimethylammonium bromide (DODAB) and dihexadecylphosphoric acid were obtained from Fluka Chemie AG (Switzerland) and were used without further purification. Sodium dihexadecyl phosphate was obtained from DHP⁴ and used in the experiments performed with unbuffered solutions. DODAC and DODAB concentrations were obtained by solubilization of a dye-amphiphile complex in nonionic micelles.¹⁷ DHP concentration was determined by inorganic phosphorus analysis.¹⁸

Charged polystyrene microspheres described as ultraclean by the supplier were obtained from Interfacial Dynamics Corp. and used as supplied. These uniform latex microspheres were stabilized either by negative sulfate charges (SP) or by positive amidine functional groups (AP). The main characteristics of the microspheres as specified from the supplier are given in Table I. Mean diameters were obtained by the supplier using the electron microscope. All other reagents were analytical grade and were used without further purification. Water was Milli-Q quality.

Large DODAC or DHP vesicles (LV) were prepared by injecting a chloroformic solution of DODAC or DHP into an aqueous salt solution at a given pH.^{3,4} Small DODAC, DODAB, or DHP vesicles (SV) were prepared by sonication of the amphiphile in water or HEPES buffer solution at pH 7.0 and 3.2 mM NaCl.^{19,20}

[†] On leave from the Departamento de Bioquímica, Instituto de Química, Universidade de São Paulo, CP 20780, São Paulo, Brazil.

* To whom correspondence should be addressed.

(1) Robb, I. D. In *Surfactant Science Series*; Lucassen-Reynders, E. H., Ed.; Marcel Dekker: New York, 1981; Vol. 11, pp 107-142.

(2) Attwood, D.; Florence, A. T. *Surfactant Systems—Their Chemistry, Pharmacy and Biology*; Chapman and Hall: London, New York, 1983.

(3) Carmona-Ribeiro, A. M.; Chaimovich, H. *Biochim. Biophys. Acta* 1983, 733, 172-179.

(4) Carmona-Ribeiro, A. M.; Yoshida, L. S.; Sesso, A.; Chaimovich, H. *J. Colloid Interface Sci.* 1984, 100, 433-443.

(5) Carmona-Ribeiro, A. M.; Yoshida, L. S.; Chaimovich, H. *J. Phys. Chem.* 1985, 89, 2928-2933.

(6) Carmona-Ribeiro, A. M.; Chaimovich, H. *Biophys. J.* 1986, 50, 621-628.

(7) Claesson, P. M.; Carmona-Ribeiro, A. M.; Kurihara, K. *J. Phys. Chem.* 1989, 93, 917-922.

(8) Carmona-Ribeiro, A. M. *J. Phys. Chem.* 1989, 93, 2630-2634.

(9) Carmona-Ribeiro, A. M. *J. Colloid Interface Sci.* 1990, 139, 343-349.

(10) Carmona-Ribeiro, A. M.; Hix, S. *J. Phys. Chem.* 1991, 95, 1812-1817.

(11) Carmona-Ribeiro, A. M.; Castuma, C. E.; Sesso, A.; Schreier, S. *J. Phys. Chem.* 1991, 95, 5362-5366.

(12) Carmona-Ribeiro, A. M.; Midmore, B. R. *J. Phys. Chem.*, submitted for publication.

(13) Ottewill, R. H.; Shaw, J. N. *Trans. Faraday Soc.* 1966, 62, 154-163.

(14) Midmore, B. R. & Hunter, R. J. *J. Colloid Interface Sci.* 1988, 122, 521-529.

(15) Van der Put, A. G.; Bijsterbosch, B. H. *J. Colloid Interface Sci.* 1983, 92, 499-518.

(16) Zukoski, C. F.; Saville, D. A. *J. Colloid Interface Sci.* 1986, 114, 32-44.

(17) Stelmo, M.; Chaimovich, H.; Cuccovia, I. M. *J. Colloid Interface Sci.* 1987, 117, 200-204.

(18) Rouser, G.; Fleischer, S.; Yamamoto, A. *Lipids* 1970, 5, 494-496.

Table I. Properties of the Sulfate Polystyrene (SP) and Amidine Polystyrene (AP) Microspheres in Water^a

latex	mean diameter/nm	N_p/cm^{-3}	area per charge group/ $\text{\AA}^2$	SSA/($\text{cm}^2 \text{g}^{-1}$)
SP150	121	8.0×10^{11}	1835	470 017
SP285	264	1.5×10^{11}	1252	215 424
AP97	76	3.5×10^{12}	370	748 316
AP850	762	1.2×10^{10}	127	74 635

^a Final number densities (N_p) in the mixtures with the vesicles and specific surface areas (SSA) are shown.

LV and SV vesicles were centrifuged at 10^4 g for 1 h at 15°C to remove any multilamellar liposomes. The supernatant containing the unilamellar vesicles was used within 1 h of the preparation.

Polystyrene microspheres were always diluted using the same solution in which the vesicles were prepared. Interaction between vesicles and microspheres was induced by adding the vesicles to the polystyrene. The final number densities of microspheres appear in Table I. Mixtures were thermostated at $25 \pm 0.1^\circ \text{C}$ for 2 h for the system DODAB SV/SP150, 72 h for DHP SV/AP97, 2 h for DODAC LV/SP285, and 24–144 h for DHP LV/AP850. Thereafter mixtures were centrifuged at 20000g for 1 h at 15°C to separate particles from vesicles. The supernatant was used for determining amphiphile concentrations, and the precipitate was used to determine particle mobilities. The adsorption isotherms of DODAC and DHP onto the microspheres were obtained from DODAC and DHP analysis in the supernatant and in the vesicle preparation. The sensitivity of the methods employed for amphiphile analysis was quite satisfactory: $10 \mu\text{M}$ DODAC or DODAB¹⁷ and $0.01 \mu\text{M}$ DHP.¹⁸ Total surface area on the polystyrene was calculated from the mass fraction and the specific surface area (Table I). Adsorption was expressed as the number of amphiphile molecules adsorbed per square meter of polystyrene.

Electrophoretic mobilities (EM) of the particles before and after amphiphile adsorption were measured using a Rank Bros microelectrophoresis apparatus with a flat cell at 25°C . ζ potentials were calculated from measured sizes and mobilities using the theory of O'Brien and White.²¹

Particle sizes were measured using a Malvern 4700c PCS apparatus employing a Coherent Innova 90 laser. The size quoted throughout is the mean harmonic z -average diameter (D_z) of at least 15 independent measurements at 25°C .

Results

1. Adsorption of Bilayers from Large Vesicles onto Oppositely Charged Microspheres. Figure 1 shows a typical adsorption isotherm for DODAC adsorption from large DODAC vesicles onto sulfate polystyrene microspheres in 0.1 mM NaCl aqueous solution at 25°C . The mean curve drawn from the experimental points can be linearized using the Langmuir model:

$$c/(x/m) = c/(x/m)_{\text{max}} + 1/K(x/m)_{\text{ads}} \quad (1)$$

where x/m is the number of moles of amphiphile adsorbed per square meter of polystyrene, c is the free amphiphile concentration, and K is the adsorption constant. By plotting $c/(x/m)$ against c , a straight line is obtained, indicating that the adsorption can be fitted to the Langmuir model. The limiting adsorption ($(x/m)_{\text{max}}$) can be obtained from the slope and the adsorption constant (K) from the intercept of the straight line with the $c/(x/m)$ axis.

Figure 1. Isotherm for the adsorption of DODAC bilayers from large DODAC vesicles onto sulfate polystyrene microspheres in 0.1 mM NaCl aqueous solution at 25°C . D_z for DODAC LV is 256 nm and for the polystyrene microspheres is 285 nm . The inset shows the same adsorption isotherm in a logarithmic scale for the abscissa.

$$\text{slope} = 1/(x/m)_{\text{max}} \quad (2)$$

$$\text{intercept} = 1/K(x/m)_{\text{max}} = \text{slope}/K \quad (3)$$

From the linearization of the curve in Figure 1, $(x/m)_{\text{max}}$ is $5.8 \mu\text{mol}$ of DODAC/ m^2 of polystyrene and K is $29 \text{ dm}^3 \text{ mol}^{-1}$. At limiting adsorption, the area per DODAC molecule adsorbed is 0.286 nm^2 (Table II). This is half the usual area per monomer in DODAC monolayers at the air–water interface,⁷ suggesting bilayer deposition onto the polystyrene surface.

The effect of the ratio of the number densities of the vesicles (N_v) and polystyrene particles (N_p) on the mean z -average diameter in mixtures of DODAC LV and SP polystyrene particles of nearly the same size is shown in Figure 2. For $N_v < N_p$, aggregation is responsible for the high D_z values measured (Figure 2), but for $N_v > N_p$, a stable D_z value of 300 nm was measured. One week after mixing, flocculation was visible for samples in which $N_v/N_p < 1$ and absent when $N_v/N_p > 1$.

Figure 3 shows the adsorption isotherm for the adsorption of DHP from DHP vesicles onto AP microspheres in 1 mM NaCl solution at pH 9–10 and 25°C . The number of DHP bilayers adsorbed is found to be dependent on time. The number of DHP molecules adsorbed per square meter of polystyrene at the plateau values corresponds to 0.208 , 0.115 , 0.076 , and $0.057 \text{ nm}^2/\text{DHP molecule}$ on the polystyrene surface, i.e., to 1, 2, 3, and 4 DHP bilayers adsorbed, respectively. The adsorption parameters for the first adsorption step are in Table II.

2. Adsorption of Bilayers from Small Vesicles onto Oppositely Charged Polystyrene Microspheres. Figure 4A shows the adsorption isotherm of DODAB bilayers from small sonicated DODAB vesicles onto sulfate polystyrene microspheres in water at 25°C . From the linearization of the curve using the Langmuir model, $(x/m)_{\text{max}}$ is $6.6 \mu\text{mol}$ of DODAC/ m^2 of polystyrene and K is $25 \text{ dm}^3 \text{ mol}^{-1}$ (Table II). The corresponding area per molecule at limiting adsorption is 0.252 nm^2 (Table I), indicating that the DODAB bilayer adsorbs onto the polystyrene with a slightly higher packing density than the DODAC bilayer (Table II). The electrophoretic mobilities of the particles precipitated by centrifugation at 20000g as a function of DODAB concentration in the supernatant are shown in Figure 4B. At $3 \times 10^{-5} \text{ M}$ DODAB, about half the particles move to the positive electrode and half to the negative electrode. Above $5 \times$

(19) Tran, C. D.; Klahn, P. L.; Romero, A.; Fendler, J. H. *J. Am. Chem. Soc.* **1978**, *100*, 1622–1624.

(20) Mortara, R. A.; Quina, F. H.; Chaimovich, H. *Biochim. Biophys. Res. Commun.* **1978**, *81*, 1080–1086.

(21) O'Brien, R. W.; White, L. R. *J. Chem. Soc., Faraday Trans. 2* **1978**, *74*, 1607–1626.

Table II. Limiting Adsorption and Adsorption Constants for the Adsorption of the Ionic Double-Chain Amphiphile Bilayers from Large and Small DODAC, DODAB, and DHP Vesicles (V) in Aqueous Solutions onto Oppositely Charged Polystyrene Microspheres at 25 °C^a

V, D_z /nm	ζ_v /mV	latex	ζ_L /mV	$(x/m)_{\max}/$ (10^{-6} mol m $^{-2}$)	$K/(10^4)$ dm 3 mol $^{-1}$	$A_{\text{ads}}/\text{nm}^2$	ζ_{LV} /mV	A/nm^2	A/A_{ads}
DODAC LV in 0.1 mM NaCl, 256	87.4 ¹²	SP285	-28.3	0.58	29	0.286		0.56 ⁷	1.96
DHP LV in 1 mM NaCl at pH 9-10, 248	-66.8 ¹²	AP850	+49.5	0.89	75	0.187		0.44 ⁷	2.35
DHP LV in HEPES at pH 7, 340	-78.0 ¹²	AP850	69.4				-72.0		
DODAB SV in water, 85	78.4 ²⁴	SP150	-55.0	0.66	25	0.252	+41.3	0.57, ²⁵	2.26
								1 mM NaBr	
DHP SV in HEPES at pH 7 and 3.2 mM NaCl, 89	-77.9 ²⁴	AP97	+33.0	0.91	0.51	0.183	-79.8	0/415, ⁷ pH 6	2.27

^a ζ potentials for vesicles (ζ_v), latexes (ζ_L), or covered latexes (ζ_{LV}) were either from the literature or calculated from mobility measurements using the O'Brien and White theory. Areas per molecule adsorbed at maximum adsorption (A_{ads}) are compared with limiting areas per monomer for pressure/area isotherms from the literature (A). A/A_{ads} is the number of amphiphile monolayers adsorbed.

Figure 2. Mean z-average diameter (D_z) as a function of the ratio between number densities for vesicles and polystyrene microspheres (N_v/N_p). Large DODAC vesicles ($D_z = 256$ nm) were mixed with sulfate polystyrene latices ($D_z = 285$ nm) 4 h before D_z determination. N_v is 8.05×10^{10} cm $^{-3}$ as calculated from D_z and assuming $55 \text{ \AA}^2/\text{DODAC monomer}$. One week after mixing, flocculation was visible for samples in which $N_v/N_p < 1$ and absent when $N_v/N_p > 1$.

10^{-5} M DODAB, all the particles move toward the negative electrode (Table I).

Figure 5A shows the isotherm for the adsorption of DHP from small sonicated DHP vesicles onto AP microspheres in 10 mM HEPES buffer, pH 7.0 and 3.2 mM NaCl at 25 °C. Again, the mean curve in Figure 5A can be linearized using the Langmuir model and adsorption parameters obtained (Table II). The area per molecule adsorbed at $(x/m)_{\max}$ is also consistent with bilayer deposition (Table II). Figure 5B shows the variation of the mean mobilities (EM) of the particles in the precipitate as a function of DHP concentration in the supernatant. The ζ potential for the precipitated particles is the same as that for the DHP vesicles (Table II).

3. Photon Correlation Spectroscopy Detecting Bilayer Adsorption on Small Particles. The effect of the ratio N_v/N_p on the mean z-average diameter (D_z) for particles in mixtures of DODAB sonicated vesicles and small SP particles follows the same pattern described for the mixtures of large DODAC vesicles and large SP particles (Figure 6). There is aggregation when $N_v/N_p < 3$, and D_z attains a constant value when $12 > N_v/N_p > 3$ (Figure 6). When $N_v/N_p = 3$, the total surface area for the vesicles is equal to the total surface area of the polystyrene particles. In Figure 6, D_z values level off at $N_v/N_p = 3.5$ –

Figure 3. Isotherm for the adsorption of DHP bilayer from large DHP vesicles ($D_z = 248$ nm) onto polyamide polystyrene microspheres ($D_z = 850$ nm) in aqueous solution at 25 °C. The closed symbols were at pH 9–10 in 1 mM NaCl and the open diamonds at pH 7.0 in 10 mM HEPES buffer. Different symbols identify the time the mixtures were left interacting: 24 h (●); 48 h (▲); 72 h (■); 144 h (◆, ◇).

4.0 for two different dilutions of the particle/vesicle mixtures. The difference between the D_z value for the most diluted mixture and the D_z value for the polystyrene alone is 8.5 nm, a value compatible with a bilayer adsorbed onto the 150-nm polystyrene microspheres (Figure 6).

The effect of dilution on the D_z of particles in a mixture of small DODAB vesicles and small SP microspheres was further investigated at $N_v/N_p = 7$ (Table III). When the dilution is increased from 1:2 to 1:40, D_z first decreases and then remains constant and equal to 159 nm which compares with 150 nm for the uncovered particles (Table III). Thus, a 9-nm layer on the SP particle is indicated. When the N_v/N_p ratio is very high (7000), D_z for the mixture further decreases.

For the DHP (89 nm)/AP (97 nm) system, the effect of the free DHP concentration and N_v/N_p on D_z for the mixtures is shown in Figure 7. D_z levels off at $N_v/N_p \approx 1.2$ (Figure 7) in good agreement with the value of 1.2 calculated from where the surface areas are equal. The

Figure 4. Isotherm for the adsorption of DODAB bilayers from small sonicated DODAB vesicles ($D_z = 85$ nm) onto sulfate polystyrene microspheres ($D_z = 150$ nm) in water at 25 °C (A). After centrifugation of the mixtures, electrophoretic mobilities of the particles in the precipitate were measured as a function of DODAB in the supernatant (B). The inset shows the same adsorption isotherm using a logarithmic scale for the abscissa.

difference in D_z for the covered and uncovered polystyrene particles is again 9 nm and therefore consistent with DHP bilayer deposition.

Discussion

In this section the following questions are addressed: (1) the significance of the adsorption measurements; (2) whether the adsorption occurs via entire bilayer or via monomer adsorption; (3) the limitations and advantages of PCS to detect bilayer adsorption; (4) the potential of synthetic amphiphile vesicles as flocculants or stabilizers.

1. Significance of the Adsorption Measurements. At maximum adsorption, the areas per DHP molecule adsorbed onto the microspheres are lower than those for DODAC or DODAB (Table II) and consistent with the tighter packing of the DHP bilayer.^{3,10} The adsorption constant (K) is a measure of the affinity the amphiphile has for the surface. The largest affinity is that between the DHP LV and the AP 850 at pH 9–10 (Table II) due probably to the larger radius of curvature of the deposited bilayer.^{10–12} Due to packing restrictions, the DHP molecules associated to form vesicles tend to return to the lamellar state.^{10–12} This same tendency could also explain the deposition of several bilayers onto each large microsphere (Figure 3) as the curvature radius increases with each bilayer addition. The absence of DHP multibilayer adsorption onto the small microspheres (AP97) (Figure 5) may be a result of the same trend. The lowest adsorption constant was obtained for the DHP SV/AP97 system (Table II). Again the highly curved surface of the microspheres does not favor the adsorption of the DHP SV

Figure 5. Isotherm for the adsorption of DHP bilayer from small sonicated DHP vesicles ($D_z = 89$ nm) onto polyamidine polystyrene microspheres ($D_z = 97$ nm) in 10 mM HEPES (pH 7.0, 3.2 mM NaCl) at 25 °C (A). Vesicles and latices were left interacting for 72 h. After centrifugation of the mixtures, electrophoretic mobilities of precipitate particles were measured as a function of DODAB concentration in the supernatant (B).

Figure 6. Effect of the number density ratio (N_v/N_p) on the mean z -average diameter (D_z) of DODAB sonicated vesicles ($D_z = 85$ nm) and sulfate polystyrene microspheres ($D_z = 150$ nm) at two different dilutions: (●) $N_p = 8.32 \times 10^{11} \text{ cm}^{-3}$ and (○) $N_p = 2.08 \times 10^{11} \text{ cm}^{-3}$. The total surface area for the vesicles is approximately equal to the total polystyrene surface when $N_v/N_p = 3$. Leveling off occurs at 3.5–4.0 for both curves.

bilayers. In addition, the SV dispersion consists of planar bilayer fragments and small vesicles,¹¹ with the adsorption of the planar stable fragments made very difficult due to the packing restrictions preventing curvature. For DHP LV adsorption at pH 7, the multibilayer adsorption is favored when compared with that at pH 10 (Figure 3). This is possibly due to the formation of *interbilayer* hydrogen bridges at pH 7 that do not occur at pH 10 and

Table III. Effect of Dilution of the Dispersions before Mixing on the Mean D_z for Particles in a Mixture of Sonicated DODAB Vesicles plus Sulfate Polystyrene Microspheres (121 nm) at $N_v/N_p = 7$ and Effect of the Number Density Ratio (N_v/N_p) on the Mean D_z at 0.2 mM DODAC^a

[DODAB]/mM	$N_p/(10^{12} \text{ cm}^{-3})$	mean D_z/nm	polydispersity index	N_v/N_p
1	1.04	172.0 ± 1.0	0.065 ± 0.024	7
0.4	0.416	158.0 ± 2.3	0.066 ± 0.034	7
0.2	0.208	161.8 ± 2.0	0.061 ± 0.020	7
0.05	0.052	159.6 ± 2.0	0.044 ± 0.022	7
0.20	0.052	155.9 ± 2.5	0.065 ± 0.029	28
0.20	0.0002	139.6 ± 2.0	0.060 ± 0.020	7000
0	1.04	150.0 ± 0.8	0.043 ± 0.031	0

^a The last line shows the mean D_z value for single microspheres.

Figure 7. Mean z -average diameter (D_z) as a function of free DHP in the supernatant (A) and as a function of the ratio between number densities for vesicles and microspheres (N_v/N_p) (B). Sonicated DHP vesicles prepared in 10 mM HEPES buffer at pH 7.0 and 3.2 mM NaCl ($D_z = 89$ nm) were mixed with polyamide polystyrene latex ($D_z = 99.4$ nm) and left for 72 h before D_z measurements. $N_p = 7 \times 10^{11} \text{ cm}^{-3}$. The total surface area for vesicles is approximately equal to the surface area for microspheres when $N_v/N_p = 1.2$, the value at which D_z levels off.

to the lower surface potential of DHP vesicles at pH 7 leading to a lower interbilayer electrostatic repulsion.¹² The affinity (K) of DODAC or DODAB bilayers for the SP microspheres (Table II) is higher than the value of $18 \times 10^4 \text{ dm}^3 \text{ mol}^{-1}$ obtained for the adsorption of dimyristoylphosphatidylcholine from sonicated vesicles onto negatively charged barium titanate glass beads.²² This is consistent with the electrostatic attraction between the vesicle and the oppositely charged polystyrene not present in the neutral phospholipid/glass beads system.

2. Entire Bilayer Adsorption vs Monomer Adsorption. As previously demonstrated for the adsorption of phospholipid from vesicles onto glass balls,²² if the adsorption were via monomer adsorption, the time required would be about 10^8 h in contrast to the few seconds necessary for adsorption via vesicle diffusion and entire bilayer attachment. Since we detected bilayer adsorption at 2–144 h after mixing vesicles and microspheres, the mechanism via monomer adsorption is impossible. Furthermore, the electrophoretic mobilities measured for the precipitated particles were found to be quantized, i.e., close to either that of the covered or that of the uncovered particles.

3. PCS for Detection of Bilayer Adsorption. The determination of bilayer adsorption using PCS requires the following: a large difference in the scattering power of the bilayer vesicles and the supporting particles; an

intermediate dilution of vesicles and of particles before mixing to minimize aggregation during mixing; substrate particles with small sizes so that an increase in size corresponding to two bilayers thickness (about 10 nm) can be detected.

The synthetic amphiphile vesicles are poor light scatterers when compared with massive polystyrene particles of the same size. If N_v/N_p is small in a mixture of vesicles and polystyrene of the same size, then the intensity of light scattered by the vesicles is negligible compared with that scattered by the microspheres. If N_v/N_p is large, then in spite of being poor scatterers, the vesicles can contribute significantly to the total intensity of light scattered due to their comparatively higher number density. In Table III, at high N_v/N_p values, the decrease in size obtained can be understood, at least in part, from the significant contribution of the vesicles to the total light scattered and to the mean z -average diameter of the mixture. The Joebst law for the light scattering by particles with sizes approximately equal to the wavelength of the incident radiation²³ states that the turbidity (S) is given by:

$$S = (\text{constant})q^2NV^{-2/3}\lambda^{-2} \quad (4)$$

where q is the anhydrous mass of the particle, N is the number density, V is the particle volume, and λ is the wavelength of the incident radiation. Using eq 4, the relative contribution of vesicles and polystyrene particles to the total light scattered by the mixture can be estimated. Assuming 285-nm diameter for spherical particles and vesicles, 1.055 g cm^{-3} as the polystyrene density, $0.6 \text{ nm}^2/\text{amphiphile monomer}$, and 600 g as the molecular weight for the amphiphile, the ratio intensity of light scattered by polystyrene (S_p) and intensity of light scattered by vesicles (S_v) can be estimated as a function of N_v/N_p :²³

$$S_p/S_v = 228(N_p/N_v) \quad (5)$$

If $N_p/N_v = 1/100$, then $S_p/S_v = 2.3$ in good agreement with the value of 2.8 actually measured by us experimentally. If $N_p/N_v = 1/10$, then $S_p/S_v = 23$, indicating that the scattering by the vesicles is negligible.

From the comparison between sizes for polystyrene microspheres as given by the supplier (Table I) and measured by PCS, it was found that the polystyrene dispersions were slightly aggregated before mixing.

At very high N_v/N_p ratios, the amphiphile bilayers decrease the mean D_z values measured (Table III) possibly due to both disaggregation and the vesicles contribution to the mean D_z of the mixture. At intermediate N_v/N_p values (lower than 10) and diluted vesicles mixed with diluted microspheres, bilayer adsorption on small microspheres was detected (Figures 6 and 7; Table III). The dilution of the dispersions before mixing at a given N_v/N_p ratio is important to prevent some local concentrated regions during mixing where aggregation can occur.

The main limitation of PCS is its lack of sensitivity to detect bilayer adsorption onto large particles as the absolute error in D_z increases with particle size. Thus, variations in size as small as 10 nm are increasingly difficult to detect as the diameter of the bilayer support increases.

4. Synthetic Amphiphile Vesicles as Flocculants or Stabilizers. DODAC, DODAB, and DHP are effective flocculants or stabilizers for oppositely charged particles dispersed in solution (Figures 2, 6, and 7), and in contrast to other single-chain surfactants they are effective at very low concentrations. The border between their action as

(23) Joebst, G. *Ann. Phys. (N.Y.)* 1925, 78, 157–166.

(24) Cuccovia, I. M.; Feitosa, E.; Chaimovich, H.; Sepulveda, L.; Reed, W. *J. Phys. Chem.* 1990, 94, 3722–3725.

(25) Marra, J. *J. Phys. Chem.* 1986, 90, 2145–2148.

(22) Jackson, S.; Reboiras, M. D.; Lyle, I. G.; Jones, M. N. *Faraday Discuss. Chem. Soc.* 1986, 85, 291–301.

flocculants or stabilizers is defined by the equality between the total surface areas for each dispersion. When there is more vesicle than polystyrene surface, the vesicles act as stabilizers, and when there is more polystyrene than vesicle surface, they act as flocculants. Another advantage over the surfactants with short hydrocarbon chains is the irreversible character of the bilayer adsorption as shown by the high adsorption constants obtained (Table II).

Conclusions

Synthetic amphiphile bilayers from one-bilayer vesicles deposit with high adsorption constants onto oppositely charged polystyrene microspheres. At maximum adsorption, the areas per amphiphile molecule adsorbed, the electrokinetic properties, and the sizes measured for the covered particles are consistent with bilayer deposition. The deposition process is controlled not only by the interaction forces between the vesicle and the microsphere but also by the restrictions of packing acting inside the bilayer. Vesicles tending to return to the lamellar state, i.e., to a state with larger curvature radius, have a high affinity for larger microspheres depositing as multibilay-

ers and a low affinity for smaller microspheres depositing as single bilayers. Interbilayer hydrogen bridges were found to enhance the multibilayer deposition.

At number density ratios (N_v/N_p) smaller than 10 using small substrate particles (mean D_s smaller than 200 nm), photon correlation spectroscopy can be employed to detect bilayer adsorption if vesicles and particles are first diluted to prevent aggregation during the mixing process. Charged synthetic amphiphile vesicles composed of double-chain surfactants with long alkyl double chains may find several uses in chemical, petrochemical, and pharmaceutical industry due to their effective action as flocculants or stabilizers of oppositely charged dispersions at very low amphiphile concentration (10–200 μM).

Acknowledgment. We thank Dr. Thelma Hardman for making available the research facilities in her laboratory. Financial support from CNPq and BID is gratefully acknowledged.

Registry No. DODAC, 107-64-2; DODAB, 3700-67-2; DHP, 2197-63-9; polystyrene, 9003-53-6; sodium dihexadecyl phosphate, 60285-46-3.

Referência 16

THE SURFACE POTENTIAL IN CHARGED SYNTHETIC AMPHIPHILE VESICLES

A. M. CARMONA-RIBEIRO* & B. R. MIDMORE

Department of Chemistry, University of Reading, Reading, RG6 2AD, UK

* On leave from the Departamento de Bioquímica, Instituto de Química, Universidade de São Paulo, CP 20780, São Paulo, Brazil.

To whom correspondence should be addressed.

Running Title: Salt and pH effects on size and zeta-potentials of ionic amphiphile vesicles.

ABSTRACT

The surface potential in large dioctadecyldimethylammonium chloride (DODAC) and dihexadecylphosphate (DHP) bilayer vesicles at low ionic strength is determined. Salt and pH effects on vesicle size and zeta-potential are investigated using photon correlation spectroscopy and particle microelectrophoresis, respectively. A large effect of amphiphile concentration on electrophoretic mobilities (EM) is described. The EM values were extrapolated to zero amphiphile concentration and the zeta-potential was calculated using the O'Brien & White theory. The larger the harmonic z-average diameter (D_z), the smaller is the zeta-potential. pH and salt effects on size are in agreement with calculations from monolayer data using a self-assembly model. Zeta-potentials determined for a range of salt and pH qualitatively agree with Stern potentials estimated from flocculation data using the DLVO.

INTRODUCTION

Depending on its hydrophobic/hydrophilic balance, amphiphile molecules assembly in aqueous solution to form a variety of structures as micelles, rods, discs, vesicles, liposomes or hexagonal phases (1, 2). Amphiphiles with long alkyl double chains like the phospholipids (3) or some synthetic amphiphiles (4- 8) form bilayer systems when dispersed in water solution. Nature, size and properties of amphiphile assemblies in aqueous solution depend not only of the composition of the dispersion medium (9- 11) but also of the dispersion method itself (12, 13). Dioctadecyldimethylammonium chloride (DODAC) and dihexadecylphosphate (DHP) dispersions obtained by sonication with tip are composed of small nonspherical particles (14) that are unstable in water (8), fuse upon addition of monovalent salt (15), appear as bilayer fragments when observed by electron microscopy (13) and do not respond to osmotic gradients (7). The sonicated dispersions are possibly composed in part of bilayer fragments (12, 13). DODAC and DHP dispersions obtained by chloroform vaporization (7, 8) are composed of large spheric particles (8) that are more stable in water and in monovalent salt than the sonicated dispersions (8, 15), behave as osmometers (7, 8) and display permeabilities towards ionic and nonionic substances that are very similar to those presented by equivalent phospholipid systems in the gel state (8). Thus, the chloroform vaporization method produces large DODAC and DHP vesicles (7, 8) ie perfect spheric shells composed of a closed bilayer enclosing an aqueous compartment (7, 8, 15, 16). They are highly charged and in the rigid gel state at room temperature having a much smoother interface than other colloids. Because of these properties, they are suitable colloid models to check theories developed for smooth, spherical and uniformly charged colloidal particles. The colloidal stability of synthetic amphiphile

vesicles has already been described (10, 11, 15, 17) but the surface potential for these systems has not yet been determined for a range of electrolyte concentration and pH (17, 18). The determination of surface potential for amphiphile vesicles is further complicated by the dependence of vesicle size on electrolyte concentration and pH (11) which has not been quantified. Here we show how vesicle size and surface potential are affected by pH and electrolyte concentration in the preparation medium of the vesicles. The experimental strategy is to determine the harmonic z-average diameter by photon correlation spectroscopy and the electrophoretic mobility by particle microelectrophoresis as a function of NaCl concentration or pH. Mobility and size of large DODAC and DHP vesicles are influenced by surfactant concentration, NaCl concentration and pH. The electrophoretic mobility increases with increasing amphiphile concentration so that mobilities were extrapolated to zero amphiphile concentration in order to avoid this complication. From the mobilities at infinite amphiphile dilution using the O'Brien & White theory (19), it was possible to obtain zeta-potentials for large amphiphile vesicles over a range of electrolyte concentration and pH. It is demonstrated that size and zeta-potential display an inverse relationship in fair qualitative agreement with the DLVO equation relating sizes and Stern potentials (20). By estimating Stern potential from vesicle flocculation data (11) using the DLVO (20) the effect of pH on the Stern potential is theoretically determined and demonstrated to be in qualitative agreement with the effect of pH on the zeta-potential. Finally, the effect of electrolyte concentration on vesicle size is shown to be qualitatively consistent with salt effects on pressure/area isotherms for DODAC and DHP (21) and with the size variations expected theoretically from a self-assembly model (22).

MATERIAL AND METHODS

Diocetadecyldimethylammonium chloride (DODAC) was obtained from Herga Industrias Químicas do Brasil. Dihexadecylphosphoric acid (DHPH) was obtained from Sigma (St. Louis, MO.) and was used to prepare sodium dihexadecylphosphate (DHP) as described in (8). All other reagents were analytical grade and were used without further purification. Water was MILLI-Q quality.

Large DODAC and DHP vesicles (LV) were prepared by injecting a chloroform solution of DODAC or DHP into an aqueous solution, 75 °C, at a given NaCl concentration and pH value (7, 8). The final amphiphile concentration was kept deliberately low (0.5 mM for DODAC and 0.2 mM for DHP) to minimize aggregation or fusion (15, 16) and the sizes were found to be stable over 24 h. Filtration procedures after injection were avoided so that the self-assembly of the amphiphiles could be better compared with predictions from the model (22). Electrophoretic mobilities (EM) were determined at 25 °C using a Rank Brothers Mark II particle microelectrophoresis apparatus. Sizes were measured simultaneously to mobilities for each vesicle sample using a Malvern 4700c PCS apparatus employing a Coherent Innova 90 laser. The size quoted throughout is the mean harmonic z-average diameter (D_z) of at least 5 independent measurements at 25 °C.

Because of the large effect of amphiphile concentration on EM (see Results section), mobilities were determined as a function of amphiphile concentration for each NaCl concentration or pH value. The EM values used for calculating zeta-potentials (ζ) were those extrapolated to zero amphiphile concentration (EM_0). EM_0 values were transformed into reduced mobilities (E) (19). E values were used to calculate the zeta-potentials (19).

RESULTS

1. *The effect of amphiphile concentration on electrophoretic mobilities.*

There is an increase in the electrophoretic mobility of the large amphiphile vesicles as a function of amphiphile concentration. This effect was observed both for large DODAC (Fig. 1) and DHP vesicles. Therefore, to calculate zeta-potentials from mobilities, extrapolations to zero amphiphile concentration were necessary (Fig. 1 B, insert). From EM_0 and D_z , zeta-potentials were calculated using the theory of O'Brien and White (19) (see next subsection). The amphiphile concentration effect depends of the volume fraction of the vesicles. However, the different curves obtained in water and 10 mM NaCl indicate that the electrostatic factor is also important (Fig. 2).

2. *The effect of NaCl on zeta-potentials and sizes of the large vesicles.*

The complete Poisson-Boltzmann (PB) equation for the surface potential around a sphere relates the surface potential with the particle radius and the electrolyte concentration (23). Although it is impossible to provide an analytical solution for the nonlinear PB equation in the spherical case and at high potentials, there are a number of analytical approximations available (23). To calculate the surface charge from the zeta-potential obtained for the vesicles (Table 1), an analytical approximation suggested by Loeb, Wiersema and Overbeek (24) was used.

$$\sigma = - \epsilon (kT/ze) K \left\{ 2 [\sinh \Psi_\delta ze/(2kT)] + (4/Ka) [\tanh \Psi_\delta ze/(4kT)] \right\} \quad (1)$$

where σ is the charge density in $\mu\text{C}/\text{cm}^2$, Ψ_δ is the zeta-potential in volts, a is the vesicle radius equal to $D_z/2$ in cm and K is the inverse of the Debye length in cm^{-1} . The final expression for the charge density at

25 °C is given below.

$$\sigma = - 5.87 C^{1/2} (1/z) \{ 2 \sinh 19.46 \Psi_{\delta} + [4/(Ka)] \tanh 9.73 \Psi_{\delta} \} \quad (2)$$

where C is the electrolyte concentration in molar and σ is obtained in C/m^2 . From the zeta-potentials, charge densities were calculated for the large DODAC and DHP vesicles using eq 2 (Table 1). It must be mentioned that the neglect of ion-ion correlation in the Poisson-Boltzmann theory leads to an overestimation of the charge densities (25). When the surface potential varies, the charge density remains constant and viceversa (Table 1).

Both for large DODAC and DHP vesicles a maximum in the zeta-potential and a minimum in vesicle size occur at very low NaCl concentration (0.1 mM for DODAC and 0.4 mM for DHP) (Fig. 3). This maximum is particularly evident for the DHP vesicles at pH 10 (Fig. 3 C, D). Intralayer hydration and charge repulsion always increase the area per monomer (21). When salt is added, sodium ions replace protons and they possibly originate an intrabilayer hydration force. In Fig. 3 C and B, at $C < 0.4$ mM, this hydration repulsion increases with salt concentration more rapidly than the charge repulsion decreases. At $C > 0.4$ mM, the reverse occurs.

3. The pH effect on zeta-potentials and sizes of the large DHP vesicles.

Reerink and Overbeek (20) derived several decades ago a well-known relationship between particle radius (r), Stern Potential (Ψ_{δ}) and slope of the stability curve (dlog W/dlog C):

$$d \log W / d \log C = -2.15 \times 10^7 r \Psi_{\delta}^2 / z^2 \quad (3)$$

where W is the stability of the colloidal dispersion usually taken as the inverse of the initial flocculation rate divided by the inverse of the maximum flocculation rate ; C is the monovalent salt concentration, r is the particle radius (in cm), z is the valency of the counterion, and $Y = (e^{y/2}-1)/(e^{y/2}+1)$ where $y = z \Psi_{\delta} e/(kT)$ with Ψ_{δ} as the Stern potential.

At 3 mM NaCl, on increasing pH, the zeta-potential for large DHP vesicles increases (Fig. 4 A) and the size decreases (Fig. 4 B). From the slopes of the flocculation curves for DHP LV at pH values varying between 6 and 10 (11), Stern potentials for the large DHP vesicles were calculated as a function of pH using eq 3 (Fig. 4 A, insert). In spite of being too low in comparison with the zeta-potentials (Fig. 4 A), the Stern potential values reflect consistently the variation of zeta-potential as a function of pH.

4. The inverse relationship between surface potential and vesicle size.

The larger the vesicle size, the lower the surface potential (Figs. 3 and 4). Eq 3 states that an increase in r should lead to a decrease in Ψ_{δ} if the slope of the flocculation curves is constant as a function of the vesicle radius. Although systematic data on flocculation of synthetic amphiphile vesicles as a function of vesicle size are not available in the literature, for vesicles of extreme radii as the sonicated and the large ones, flocculation data analysis yielded practically the same slope (15, 17). Assuming a constant slope, the validity of eq 3 can be verified for synthetic amphiphile vesicles by plotting $1/r$ against Y^2 . Straight lines were thus obtained for both vesicle types (Fig. 5).

From eq 3 the slope of each straight line in Fig. 5, $dY^2/d(1/r)$, is inversely proportional to $d\log W/d\log C$. Therefore, the $d(1/r)/dY^2$ ratio for

DODAC and DHP vesicles (Fig. 5) can be compared with the $d\log W/d\log C$ ratio from flocculation data (15, 17). For large DODAC and DHP vesicles prepared in D-glucose solution, the ratio between the slopes of the flocculation curves is 3.3:13.3, ie 0.25 (15, 17). For large DODAC and DHP vesicles prepared in diluted salt solutions (DHP vesicles at pH 10), the $d(1/r)/dY^2$ ratio calculated from the straight lines is 31.3:131.7, ie 0.24 (Fig 5). Thus, the ratios are practically equal.

5. Testing a self-assembly model.

From a self-assembly model (22) with geometric parameters for the DODAC and DHP molecules (1) and limiting areas per monomer from monolayer data at a surface pressure of 25 mN/m (21), critical radii for vesicles (R^C) can be calculated and compared with $D_z/2$. In spite of being much lower than the sizes measured, R^C values as a function of salt concentration reflect the effect of salt on sizes of DODAC LV (Fig 6).

For DHP LV prepared in absence of buffer, on increasing salt concentration the bulk pH decreases (Tab 2) as expected from the salt-induced shift of the phosphate half-dissociation to lower pH values (10, 11).

From the hydrocarbon chain length, l , (1), the volume of the hydrocarbon chains, v , (1) and D_z values measured for the DHP LV, it is possible to calculate areas per monomer a_T (Tab 2). These areas can be compared with those estimated at 25 mN/m from pressure/area isotherms (21) (Tab 2). The effect of increasing the electrolyte concentration is an increase (0-1 mM NaCl) followed by a decrease in the area per monomer (1- 10 mM NaCl) (Tab 2). The same increase in charge density from 0- 0.2 mM NaCl that decreased vesicle size (Fig. 7) is responsible for the increase in area

per monomer observed in DHP monolayers from 0- 0.1 mM NaCl (Tab 2).

a and a_T are very similar for DHP vesicles (Tab 2).

a and a_T for DODAC vesicles decrease as the electrolyte concentration (C) increases (Tab 2).

DISCUSSION

This section will address mainly 3 questions: 1) whether reliable surface potentials can be obtained for the vesicles from microelectrophoresis and the O'Brien and White theory for colloids; 2) whether a satisfactory theory for the radius dependence of colloidal stability exists; 3) whether the self-assembly model accounts for the self-assembly of DODAC and DHP.

1. Reliability of ζ

When the combined volume of particles in solution is appreciable, as in the case of the large vesicles, the electrophoretic mobilities may be affected by overlap of ionic atmospheres and hydrodynamic interactions. If the only interaction responsible for the observed increase in mobilities with amphiphile concentration (Fig. 1) were the hydrodynamic one, at a given vesicle volume fraction the same mobility should be measured in water and in 10 mM NaCl. Since this was not the case (Fig. 2), the electrostatic factor seems also to be important in determining the effect. By using cylindrical and flat cells identical results were obtained (Fig. 1 A) proving that cell geometry is not a factor. It must be emphasized that the increase in mobilities as a function of increasing amphiphile concentration cannot be explained in terms of obstruction effects like solvent backflow or viscoelectric phenomena since these forces would result in lower mobilities at higher volume fractions and amphiphile

concentrations. At present, our theoretical understanding of these complications is still insufficient so that extrapolations to infinite amphiphile dilution were performed to determine mobilities of isolated vesicles. This procedure generated mobility values that could be fitted in the O'Brien & White theory to obtain zeta-potential values (Table 1, Figs. 3 and 4). The choice of this theory for performing the calculations was based on one of its fundamental results, namely, the independence of the electrophoretic mobility of the colloidal particle of its dielectric properties. Thus, no restrictions are apparent in using the theory for charged spheric rigid shells surrounding an internal aqueous compartment. All EM_0 values were found to lie below the maximum EM value predicted by the theory and the charge densities calculated from ζ (Table 1) were found to agree well with those determined earlier using the surface force apparatus and nonlinear Poisson-Boltzmann (21, 26). At 10 mM NaCl and pH 9.5, an area per charge of 16 nm^2 , ie a charge density of 0.010 C/m^2 was calculated from nonlinear Poisson-Boltzmann and surface force measurements between adsorbed DHP layers (5). This is consistent with a charge density of 0.011 C/m^2 calculated from eq 2 and mobility measurements for DHP LV at 5 mM NaCl and pH 10 (Tab 1). For the DODAC system the charge density at 2 mM KBr is 0.018 C/m^2 from the surface force measurements (26) and 0.014 C/m^2 at 5 mM NaCl from the mobility measurements (Tab 1). This small difference may be due to experimental error in both techniques and/or to the slightly different experimental conditions.

2. The radius dependence of the colloidal stability.

One of the most controversial results of the DLVO theory is the relationship between slope of the stability curve, Stern potential and particle radius (eq 3). From eq 3, it can be inferred that, if $d\log W/d\log C$ were constant, an inverse relationship between r and Ψ_δ should be

observed. Alternatively, if Ψ_δ is constant, $d\log W/d\log C$ should increase in absolute value with increasing size.

From flocculation data for DODAC vesicles, vesicle radii extreme as 20 and 150 nm resulted in $d\log W/d\log C$ values that were equal inside the limits of the experimental error (15). As a result, estimated Ψ_δ from the slope of the stability curves and eq 3 were higher for the small vesicles (15) in accordance with their higher conductivity and free flow electrophoretic mobility (18). Using eq 3, the slope taken from flocculation data (11) and the size measurements (Fig. 4A), Ψ_δ values were calculated for DHP LV at different pH values (Fig. 4 A, insert). Ψ_δ increase was consistent with the zeta-potential increase as a function of pH (Fig. 4). Also, the inverse relationship between vesicle size and zeta-potential (Fig. 3- 5) possibly resulting from a constant slope for different vesicle sizes (11, 15) is in qualitative agreement with eq 3.

The constant slope for different particle sizes resulting in a decrease in Stern potential with an increase in particle size has also been described for several colloids such as AgI, gold, selenium sols and also, for polystyrene homo disperse spheres with sizes varying over an order of magnitude (20, 27). Because no simple explanation for the inverse relationship between size and surface potential is available, doubt has been cast upon the validity of equation 3 (20, 27). For AgI, a system where Ψ_δ is constant, the expected increase in slope with particle radius was not observed (20). However, the nonsphericity of the AgI particles may be a complicating factor. For the perfectly spheric polystyrene particles, although direct mobility measurements were performed as a

function of particle size (27) later work has revealed many problems regarding the association of the mobility values with the surface potential. The polystyrene interface is hairy and there is possibly Stern layer conduction (28- 30). The decrease of the surface potential with increasing particle radius could be a real phenomenon with the constant slope as a mere consequence of this inverse relationship. For the vesicles, at least, we present strong experimental evidence for such a relationship (Figs 3- 5). The complete demonstration of the validity of eq 3, however, still depends on the systematic determination of the vesicle size effect on the slopes of the flocculation curves.

It is well-known that eq 3 is only an approximation of the exact DLVO theory (20). In spite of offering a good qualitative idea of the behavior of the surface potential as a function of pH (Fig 4 A), eq 3 yields values of Stern potential that are generally too low (15, 20). This might explain why the Stern potential calculated from flocculation data and eq 3 is so much lower than the zeta-potential (Fig 4 A and 4 A insert).

3. *The self-assembly of DODAC and DHP.*

Long alkyl double-chained amphiphiles as the phospholipids have an area per monomer in the lamellar phase that is very similar to their limiting areas per monomer in pressurized monolayers at the air-water interface (1, 3). At 25 mN/m DODAC and DHP molecules at an air/water interface occupy practically their limiting areas per monomer as determined from pressure/area isotherms for DODAC and DHP monolayers at the air/water interface (21). Thus the areas per monomer at 25 mN/m for a range of salt and pH in the subphase (21) were used for calculations (Tab 2). The change in area per monomer that is observed when the salt concentration is increased is

similar for vesicles and insoluble monolayers of DODAC and DHP. The decrease in a and a_T for DODAC vesicles with increasing electrolyte concentration (Tab 2) is consistent with the increase in the vesicle size (Fig. 6) and with vesicle fusion upon salt addition (16). The large difference in absolute value for measured and calculated vesicle radii (Fig. 6) could be seen as a failure of the self-assembly model (22). However, one should recall that only the minimum vesicle radius for a given vesicle forming amphiphile is predicted by the model so that there are no restrictions to the formation of larger vesicles. Regarding the prediction of the variation in vesicle size with salt concentration from the molecular geometry, the model is successful (Fig. 6, Tab 2).

The comparison between a and a_T (Tab. 2) allows some insight on the thermodynamic stability of the vesicle structure. a_T equal to a for the DHP system implies in its existence at the limit between a vesicle and a lamellar phase composed of planar bilayers. This means that any further decrease in the area per monomer caused either by decreasing pH or by increasing salt concentration (21) would probably lead to thermodynamically unstable vesicles returning to the lamellar phase. This is not the case for the DODAC vesicles since $D_z/2$ is larger than the critical radius (Fig. 6) allowing a geometrically nonconstrained existence for the DODAC vesicles within a more extensive range of variation in electrolyte concentration.

CONCLUSIONS

There is an inverse relationship between size and zeta-potential for the large DODAC and DHP vesicles. This correlation may be the explanation for the constant slope observed in stability curves for vesicles of extreme sizes. The DLVO theory predicts an inverse relationship between particle

size and surface potential as obtained for the synthetic amphiphile vesicles. The effect of NaCl concentration on vesicle size is consistent with salt effects on DODAC and DHP monolayers and with the predictions of the self-assembly model. The pH effect on the zeta-potential of the DHP vesicles is consistent with the estimations of Stern potential from flocculation data using the DLVO theory.

ACKNOWLEDGMENTS

Financial support from CNPq, FAPESP and BID is gratefully acknowledged. We thank Dr Thelma Hardman for making available the research facilities in her laboratory and one of the reviewers for valuable criticisms and suggestions.

REFERENCES

- 1 Tanford, C. *The Hydrophobic Effect*, Wiley Interscience: New York, 1982.
- 2 Israelachvili, J. N. *Intermolecular and Surface Forces*: Academic Press, London, 1985.
- 3 Bangham, A. D. (Editor) *Liposome Letters*: Academic Press, London, 1983.
- 4 Kunitake, T.; Okahata, Y.; Tamaki, K.; Kumamaru, F.; Takayanagi, M. *Chem. Lett.* 1977, 387.
- 5 Mortara, R. A.; Quina, F. H.; Chaimovich, H. *Biochem. Biophys. Res. Commun.* 1978, 81, 1080.
- 6 Kano, K.; Romero, A.; Djermouni, B.; Ache, H.; Fendler, J. H. *J. Am. Chem. Soc.* 1979, 101, 4030.
- 7 Carmona-Ribeiro, A.M.; Chaimovich, H. *Biochim. Biophys. Acta* 1983, 733, 172.
- 8 Carmona-Ribeiro, A.M.; Yoshida, L.S.; Sesso, A.; Chaimovich, H. *J. Colloid Interface Sci.* 1984, 100, 433.
- 9 Ninham, B. W.; Evans, D. F.; Wei, G. J. *J. Phys. Chem.* 1983, 87, 5020.
- 10 Carmona-Ribeiro, A.M. *J. Colloid Interface Sci.* 1990, 139, 343.
- 11 Carmona-Ribeiro, A.M.; Hix, S. *J. Phys. Chem.* 1991, 95, 1812.
- 12 Pansu, R.B.; Arrio, B.; Roncin, J.; Faure, J. *J. Phys. Chem.* 1990, 94, 796.
- 13 Carmona-Ribeiro, A.M.; Castuma, C.E.; Sesso, A.; Schreier, S. *J. Phys. Chem.* 1991, 95, 5361.
- 14 Herrmann, H.; Fendler, J.H. *Chem. Phys. Letters* 1979, 64, 270.
- 15 Carmona-Ribeiro, A.M.; Yoshida, L.S.; Chaimovich, H. *J. Phys. Chem.* 1985, 89, 2328.

- 16 Carmona-Ribeiro, A.M.; Chaimovich, H. *Biophys.J.* 1986, 50, 621.
- 17 Carmona-Ribeiro, A.M. *J.Phys.Chem.* 1989, 93, 2630.
- 18 Cuccovia, I.M.; Feltosa, E.; Chaimovich, H.; Sepulveda, L.; Reed, W. *J.Phys.Chem.* 1990, 94, 3722.
- 19 O'Brien, R.; White, L.R. *J.Chem.Soc.Faraday Trans.2* 1978, 74, 1607.
- 20 Reerink, H.; Overbeek, J.T.G. *Discuss.Faraday Soc.* 1954, 18, 95.
- 21 Claesson, P.M.; Carmona-Ribeiro, A.M.; Kurihara, K. *J.Phys.Chem.* 1989, 93, 917.
- 22 Israelachvili, J.N.; Mitchell, D.J.; Ninham, B.W. *J.Chem.Soc.Faraday Trans.2* 1976, 72, 1525.
- 23 White, L.R. *J.Chem.Soc.Faraday Trans. 2* 1977, 73, 577.
- 24 Loeb, A.L.; Wiersema, P.H.; Overbeek, J.Th.G. *The electrical double layer around a spherical particle* M.I.T. Press, Cambridge, Mass.
- 25 Attard, P.; Mitchell, J.; Ninham, B. W. *J.Chem.Phys.* 1980, 89, 4358.
- 26 Pashley, R.M.; MacGuiggan, P.M.; Ninham, B.W.; Brady, J.; Evans, D.F. *J.Phys.Chem.* 1986, 90, 1637.
- 27 Ottewill, R.H.; Shaw, J.N. *Trans.Faraday Soc.* 1966, 42, 154.
- 28 Midmore, B.R.; Hunter, R.J. *J.Colloid Interface Sci.* 1988, 122, 521.
- 29 Van der Put, A.G.; Bijsterbosch, B.H. *J.Colloid Interface Sci.* 1983, 92, 499.
- 30 Zukoski, C.F.; Saville, D.A. *J.Colloid Interface Sci.* 1986, 114, 32.

FIGURE CAPTIONS

FIG. 1 The effect of DODAC concentration on electrophoretic mobilities (EM) of large DODAC vesicles prepared in water (A) or at 10 mM NaCl (B). Circles indicate measurements done with the flat cell and squares, with the cylindrical cell. The insert shows the extrapolation to zero DODAC concentration yielding the mobility value, EM_0 , that is used to calculate the zeta-potential of large DODAC vesicles at 10 mM NaCl.

FIG. 2 The effect of the volume fraction (ϕ) on mobilities (EM) for the large DODAC vesicles prepared in water ($\square$) or at 10 mM NaCl ($\circ$). The volume fraction was calculated assuming a spherical shape, D_z equal to 280 nm in water and 420 nm at 10 mM NaCl and areas per monomer from the literature (.66 and .52 nm², in water and at 10 mM NaCl, respectively).

FIG. 3 The NaCl concentration effect on diameters (A, C) and zeta-potentials (B, D) of large DODAC (A, B) or DHP (C, D) vesicles. DHP vesicles were prepared and kept in buffer at pH 10.

FIG. 4 The effect of pH on zeta-potentials (A) and diameters (B) of large DHP vesicles at 3 mM NaCl. From the DLVO theory relating radius and Stern potential to the slope of the stability curve reported previously, Ψ_d was calculated as a function of pH (A, insert). In spite of being too low, the Ψ_d values reflect the variation of zeta-potential as a function of pH (A).

FIG. 5 The inverse relationship between sizes and zeta-potentials for charged synthetic amphiphile vesicles seen as a plot of $1/r$ against ζ^2 (eq 3). Variation of potentials and sizes was achieved by varying the NaCl concentration in the medium where DODAC (○) or DHP vesicles (▲) were prepared. For the DHP system, varying sizes and potentials by varying the pH in the preparation medium at a constant ionic strength also yielded a linear plot (■).

FIG. 6 The comparison between measured (o) and calculated (o) DODAC vesicle radii as a function of NaCl concentration (C). The calculated radii were obtained from $R^C = 1/(1 - v/(a l))$ where l is the hydrocarbon chain length equal to 2.3005 nm and v is the volume of the chains equal to $.969 \text{ nm}^3$ for the DODAC molecule. Areas per monomer were taken from reference 21.

Fig. 7 Mean z-average diameter (D_z) as a function of NaCl concentration for large DHP vesicles. DHP LV were prepared in unbuffered aqueous solutions at pH 6- 8 (Table 2).

TABLE 1

The effect of NaCl concentration (C) on mean zeta-average diameters (D_z) and zeta-potentials of large DODAC or DHP vesicles. DHP vesicles were prepared in buffer at pH 10. P is the polydispersity index. K is the inverse of the Debye length in cm^{-1} . EM_0 is the electrophoretic mobility extrapolated to zero amphiphile concentration and E is the reduced mobility (19). Charge densities, σ , were calculated from Poisson-Boltzmann (eq 2) and zeta-potentials, ζ .

Amphiphile	C	D_z	P	$KD_z/2$	$10^4 XEM_0$	$10^4 XE$	ζ	$10^3 X\sigma$
	(mM)	(nm)			($\text{cm}^2 \text{V}^{-1} \text{s}^{-1}$)		(mV)	(C/m^2)
DODAC	0	277	.321	0	4.4	3.3	84.8	-
	0.1	263	.305	4.3	3.75	2.81	87.4	3.5
	0.2	321	.345	7.5	3.9	2.9	77.1	3.8
	0.3	353	.335	10.4	3.87	2.9	73.2	4.1
	0.4	367	.391	12.5	4.0	3.0	72.0	4.7
	0.5	392	.366	14.0	4.22	3.17	65.5	4.6
	1.0	405	.337	21.3	4.1	3.08	64.3	6.2
	5.0	412	.327	47.8	4.38	3.28	64.3	13.5
	10.0	397	.415	68.0	4.85	3.64	65.5	19.5
DHP	0	245	.390	0	3.8	2.85	-70.7	-
	0.2	243	.380	5.6	5.6	3.0	-54.0	-3.4
	0.3	228	.427	6.5	3.7	2.78	-74.9	-4.5
	0.4	200	.382	6.6	4.2	3.15	-107.0	-8.4
	0.5	208	.418	7.6	4.2	3.15	-92.5	-8.2
	0.6	222	.433	8.9	4.0	3.00	-82.2	-7.4
	1.1	229	.418	12.5	3.8	2.80	-66.8	-7.0
	2.1	253	.470	19.0	3.2	2.40	-48.8	-6.2
	5.1	242	.466	28.4	3.7	2.78	-54.0	-10.8

TABLE 2

Experimental (a) and theoretical (a_T) areas per monomer in DHP LV prepared at several NaCl concentrations (C) in unbuffered solutions. a values were from monolayer data for DODAC or DHP at 25 mN/m (21). a_T was obtained from $a_T = (1/(1 - 2/D_z))v/l$ where l and v are .205 nm and .862 nm³ for DHP and .230 nm and .969 nm³ for DODAC, respectively.

Amphiphile	C /M	pH	$D_z/2$ /nm	a_T /nm ²	a /nm ²
DHP	0	7.8	185	.4255	.425
	0.0001	7.3	149	.4266	.427
	0.0010	6.9	149	.4266	.426
	0.0100	6.2	172	.4258	.421
DODAC	0		120	.4297	.660
	0.0001		128	.4292	.535
	0.0010		192	.4266	.520
	0.0100		198	.4264	.520

FIG. 5
CARMONA-RIB
& MIDMORE

Fig 6

The Journal of Physical Chemistry

PUBLISHED BY
THE AMERICAN CHEMICAL SOCIETY

Department of Chemistry and Biochemistry
University of California
Los Angeles, California 90024-1569
Phone: (310) 825-1352
FAX: (310) 206-6628; BITNET: JPHYSICHEM@UCLACH
INTERNET: JPHYSICHEM@UCLACH.UCLA.EDU

Mostafa A. El-Sayed, *Editor-in-Chief*

December 29, 1991

Dr. A. M. Carmona-Ribeiro
Department of Chemistry
University of Reading
Reading, RG6 2AD, UK

Dear Dr. Carmona-Ribeiro:

Reference is made to the manuscript (Number JP911219Q) entitled "The Surface Potential in Charged Synthetic Amphiphile Vesicles."

Your manuscript has been accepted for publication in the Journal of Physical Chemistry and should appear in the first issue of April, 1992. You should receive galleys approximately four weeks prior to publication. Further correspondence regarding this manuscript should be addressed to Dr. Marianne Brogan, Books & Journals Division, Editorial Department, American Chemical Society, P.O. Box 3330, Columbus, Ohio 43210 (Tel: 614-447-3696).

Enclosed is a letter from the ACS Publications Office concerning the JPC policy on page charges. Please read this carefully noting particularly that your institution will be billed for the page charges unless you apply for a waiver.

Sincerely yours,

M. A. El-Sayed
Editor-in-Chief

MAE/sds
Enc.

Referência 17

SYNTHETIC AMPHIPHILE VESICLES

A. M. Carmona-Ribeiro^{*}

Chemistry Department
Reading University
READING RG6 2AD UK

^{*} On leave from Departamento de Bioquímica, Instituto de Química, Universidade de São Paulo, CP 20780 , São Paulo, Brazil.

ABSTRACT

At the border between colloids and biomimetic systems, synthetic closed bilayers offer many possibilities of understanding the physical chemistry of biomembranes and the relationship between chemical structure of the monomer and the properties of its aggregates. During the last decade, dioctadecyldimethylammonium chloride and dihexadecylphosphate vesicles have had their properties scrutinized, but their full potential is so far unexplored. Developments have focused on physical and functional properties of vesicles, on the responsiveness of the vesicles to their environment and on intervesicular interactions leading to aggregation or fusion. The behavior of vesicles as ideal colloids with a perfectly regular structure at the interface or as colloid stabilizers or flocculants is among the most recent advances in basic research which should lead to new uses and physical insights. A critical overview regarding the importance and the perspective of synthetic amphiphile vesicles for biomimetic and surface chemistry is presented.

SYNOPSIS

- 1 Background
 - 1a Liposomes and model membranes.
 - 1b Molecular geometry and bilayers.
- 2 Physical and functional properties.
 - 2a Preparation methods.
 - 2b Captured volume, size and shape.
 - 2c Physical state and molecular packing.
 - 2d Inner compartment and functional properties.
- 3 (Meta)stability
 - 3a Stability of the bilayer structure and colloidal stability.
 - 3b Aggregation and fusion.
 - 3c Surface potential and theoretical stabilities.
- 4 Uses.
 - 4a Storage of light energy and catalysis.
 - 4b Bilayer adsorption.
 - 4c Flocculation and stabilization of dispersions.
 - 4d Production of model colloids.

1 BACKGROUND

1a Liposomes and model membranes.

At the beginning of the 20th century the plasma membrane seemed to be a thin structure impregnated with lecithin and cholesterol, which was likely to contain protein (1). Quantitative data on thickness and composition scarcely entered into this description. There were no ideas on the structural role of lipids or molecular organization. 1925 yielded the classical result of Gorter and Grendel: "Every chromocyte is surrounded by a layer of lipoids, of which the polar groups are directed to the inside and to the outside in much the same way as Bragg supposes the molecules to be oriented in a crystal of fatty acid, and as the molecules of a soap bubble are, according to Perrin" (1). Further, Grendel in 1929 presented a value of 3.1 nm for such a lipid layer based on calculations using the molecular lengths of lipid molecules and on his assumption that this membrane lipid layer must be bimolecular, because it comes in contact with water on both sides. A model of membrane structure, however, appeared only in 1935 in a paper by Davson and Danielli. The arrangement of the lipid molecules in the monolayers at the two membrane-water interfaces was derived from their work on lipid films at the air-water interface. Even today, with the maturation of the bilayer paradigm, a crucial question is whether or not the lipids in plasma membranes always take on the bimolecular leaflet form.

When lipids are mixed with water, organized lamellar structures called myelin figures are formed. "Water enters the masses of phospholipids which do not usually take a spherical shape but often assume the most irregular and striking forms, reminding one irresistibly of the structure of cells", as described by Matthews in 1920. The liposomes of Bangham are a particular kind of myelin figure that forms closed systems and is composed of coherent concentric spheroidal bilayers (1). They can

be obtained simply by vortexing the phospholipids in aqueous solution. A great deal of research has been devoted to the physico-chemical properties of liposomes since their discovery in 1965 (1). From these early days up to the present, the development and diversification of the liposome "membrane" model is fascinating. They have been used as model membranes for anaesthetic action, "test-beds" for membrane enzymes, carriers of encapsulated materials into cells, barriers to be invaded by viruses and bacteria, antigens to be attacked by anti-lipid antibodies, reconstituted membranes and presynaptic vesicles, binders for drugs, substrates for testing ionic channels, "food" for phospholipases and drug delivery systems (1).

Since 1976, other membrane mimetic systems composed of ionic synthetic amphiphiles with long alkyl double chains (Fig. 1) have been proposed (2- 6). New developments and uses for these systems are, however, dependent on further characterising the dispersions obtained in solution. In this respect, both the history and uses of the liposomes are very instructive. Methods employed for obtaining phospholipid dispersions can be successfully adapted to obtain vesicles of synthetic amphiphiles (2- 6). These highly-charged synthetic bilayers offer a challenge for colloid scientists. They are useful to test theoretical models for colloidal stability, electrical double layer and amphiphile self-assembly. Furthermore, they accomplish the usual tasks that single-chained surfactants do at much lower concentrations (7). Here about a decade of research on dioctadecyldimethylammonium chloride (DODAC) and dihexadecylphosphate (DHP) vesicles will be reviewed, focusing on the importance of good characterization of the dispersions in solution, on their analogies with and differences from phospholipid systems and on their uses.

1b Molecular geometry, bilayers and vesicles.

Amphiphilic molecules as surfactants and lipids associate in aqueous solution and at interfaces to form a variety of structures (8). The hydrocarbon chains are hydrophobic and the polar heads are hydrophilic so that the hydrophobic interaction induces the molecules to associate exposing the polar heads to the aqueous phase (Fig.2).

At thermodynamic equilibrium, the geometry of an amphiphile molecule in the aggregate is given by the optimal area a , ie the surface area occupied per headgroup when the total interfacial energy per molecule in the aggregate is a minimum; the volume of the hydrocarbon chain or chains, v ; and the maximum effective length that the chains can assume, l . From the self-assembly model and the molecular geometry, it is possible to determine the nature, shape and size of the aggregate into which the amphiphilic molecules can pack (8).

Amphiphiles that form bilayers are those with small headgroup areas, a , or those with bulky hydrocarbon regions. The value of v/al must lie close to 1 and the critical packing shape has to be cylindrical (Fig. 3). For a bilayer to curve, the amphiphiles in the outer monolayer must be able to pack into truncated cones and the value of v/al has to be smaller than 1 but larger than $1/2$. It must be noticed that the self-assembly model does not predict tilting or interdigitation of the hydrocarbon chains. Furthermore, the model was constructed to predict the nature and shape of amphiphile aggregates in thermodynamic equilibrium, a condition often circumvented by nature and its metastable states.

The geometry of DODAC and DHP molecules is in Tab. 1. A complete test of the self-assembly model using these synthetic amphiphile vesicles would

depend on determining the bilayer thickness and the areas per monomer in the bilayer vesicles for a range of experimental conditions. Assuming that the area per monomer a in DODAC and DHP vesicles is close to the limiting area per monomer at 25 mN/m of DODAC and DHP monolayers at an air-water interface (9), v/al was calculated for a range of electrolyte concentrations (10). As the pH decreases or the electrolyte concentration increases, the a value decreases and v/al becomes close to 1 indicating formation of planar bilayers (7, 10). The DHP molecule has an interesting geometry which depends on pH and ionic strength, and yields values of v/al close to 1. Using areas per DODAC molecule at 25 mN/m obtained for DODAC monolayers (9), the calculated values of v/al for the DODAC system increase from 0.64 in water to 0.81 in 0.01 M NaCl (10) (Tab. 1). Since the pH and salt concentration determine the area per monomer, they also determine the vesicle size (10, 11).

2 PHYSICAL AND FUNCTIONAL PROPERTIES

2a Preparation methods

The physical and functional properties of synthetic amphiphile vesicles vary with the method of preparation. Three principal methods have been employed to prepare DODAC and DHP dispersions in aqueous solutions: 1. sonication in a bath (2); 2. sonication with a tip (3, 4); 3. chloroform vaporization (5, 6). The first method yields multilamellar vesicles (2), the second yields small unilamellar vesicles (3, 4) and/or bilayer fragments (12, 13) and the third, large unilamellar vesicles (5, 6). With the exception of the large unilamellar vesicles obtained by chloroform vaporization, the dispersions obtained by the other two methods are not yet properly characterized. For a given medium composition, the three methods will yield different vesicle sizes. It is not possible for all of them to be in thermodynamic equilibrium (12). At least two of these preparations must correspond to metastable states.

Turbidity measurements as a function of time after vesicle preparation in water revealed that the dispersions obtained by sonication were not stable in contrast to the apparent stability over days of the dispersion prepared by chloroform vaporization (6).

2b Captured volume, size and shape

The captured volume is one important parameter of vesicle preparations which is a function of the vesicle size only. It is defined as the volume enclosed by a given amount of amphiphile and expressed in litres entrapped per mole of total amphiphile (l mol^{-1}). It is usually measured using a water-soluble marker which labels the inner aqueous compartment without being adsorbed or included in the vesicle bilayer. Entrapped and free marker can be separated by dialysis (14) or by gel chromatography (5, 6). The captured volume for sonicated DODAC dispersions in water is close to zero (0.13 l mol^{-1}) (5) and lower than that for equivalent phospholipid dispersions ($0.5\text{--}0.8 \text{ l mol}^{-1}$) (15, 16). The captured volume for large DODAC and DHP vesicles obtained by chloroform vaporization is 9.7 (5) and 13.6 l mol^{-1} (6), respectively and compares fairly well with that for large unilamellar phospholipid vesicles obtained by reversed phase vaporization, 11.7 l mol^{-1} (17).

Vesicle sizes are determined by electron microscopy or photon correlation spectroscopy (PCS). After sonication (10 minutes for DODAC and 20 minutes for DHP), the DODAC and DHP dispersions are composed of nonspherical particles of around 45 and 59 nm mean hydrodynamic radius, respectively (18). The DODAC and DHP dispersions obtained by chloroform vaporization in water are composed of spherical particles (6, 19) of around 120 and 185 nm mean hydrodynamic radius, respectively (10). The ratio between the hydrodynamic radius and the radius of a sphere with the molecular weight of a sonicated vesicle is close to 3 indicating a

pronounced deviation from a spherical shape (18). The turbidity of the dispersions obtained by chloroform vaporization is well described by the Joebst law for light-scattering by spherical particles (6, 19, 20). Thus, dispersions obtained by sonication produce small nonspherical particles whereas those obtained by chloroform vaporization yield large spherical ones. Vesicle size depends on electrolyte concentration and pH (10, 11). By increasing the NaCl concentration in the preparation medium from 0.1 to 0.5 mM, large DODAC vesicles almost doubled in size (10). By increasing pH from 6.2 to 8, large DHP vesicles decreased in size from 325 to 240 nm diameter (10).

2c Physical state and bilayer packing

At room temperature DODAC and DHP bilayers are in the rigid gel state (5, 11, 21). As the temperature increases, so does the rotational freedom of the hydrocarbon chains. The transition from the ordered gel phase to the fluid liquid-crystalline state corresponds to an abrupt and large decrease in the packing density at a particular temperature, the phase transition temperature (T_c). The higher the packing density of the hydrocarbon chains in the gel state, the higher T_c (22). T_c values measured for DODAC vesicles were around 38 °C (5) whereas those measured for DHP vesicles varied between 57 and 72 °C depending on pH (11, 21). For the DHP system, vesicle size and bilayer packing (T_c) vary as a function of pH and electrolyte concentration. Both were at maximum at pH values corresponding to half-dissociation of the phosphate polar heads (11). The largest DHP vesicles and the most tightly packed DHP bilayers were obtained at half-dissociation (11).

The DHP system may also exist in the isotropic or hexagonal phase (11, 21). At high temperature, very low NaCl concentration and pH at and around the pH of half-dissociation for the phosphate polar head, an abrupt and large increase in turbidity can be associated with vesicle

aggregation and formation of the isotropic phase at the contact points in between vesicles (11, 21).

The determination of pressure-area isotherms using DODAC and DHP monolayers at the air-water interface provides an alternative method of studying the molecular packing in DODAC and DHP bilayer vesicles (9). A high packing density in the vesicle bilayer usually corresponds to low limiting areas per monomer in the monolayer. At a surface pressure of 25 mN/m with water as subphase, the areas per monomer for DODAC and DHP are 0.66 and 0.41 nm², respectively (9). This large difference correlates well with the large difference in T_c found DODAC and DHP bilayers in water. A tight molecular packing in the gel state may also be associated with a low permeability towards several ionic and nonionic substances. Permeabilities of large DHP vesicles are systematically smaller than those of large DODAC vesicles (6).

2d Inner aqueous compartment and functional properties

Closed bilayer systems are one of the most explored models of membranes. They usually perform such basic membrane functions as compartmentalization and passive transport of solutes and water. The captured volume of the sonicated synthetic amphiphile dispersions is close to zero (see section 2b). Thus, incorporation of water soluble substrates and permeability studies have been restricted in the sonicated dispersions. In contrast, large DODAC or DHP vesicles obtained by chloroform vaporization entrap sizeable amounts of water soluble substrates and exhibit ideal osmotic behaviour (5, 6). In Fig. 4 the shrinkage of large DODAC vesicles due to an osmotic gradient of sucrose is seen as an increase in turbidity. The initial shrinking rate varies linearly with the applied osmotic gradient, as for an osmometer (5, 6). This ideal osmotic behaviour towards sucrose solutions suggests that

large DODAC and DHP vesicles are nonpermeant towards sucrose. Thus, radioactive sucrose could be used to determine the captured volume and cold sucrose could be used as a standard to determine relative permeabilities towards other solutes (5, 6). Most nonionic and ionic solutes were nonpermeant. Large DODAC and DHP vesicles at room temperature are impermeable even towards urea and permeable only to water, ethylene glycol and ammonium acetate (6). Synthetic amphiphile vesicles and phospholipid liposomes in the gel state behave similarly regarding passive transport (6).

3 (META)STABILITY

3a Stability of the bilayer structure and colloidal stability

A system is thermodynamically stable when its free energy is at a minimum. The invariance of a given structural property after an annealing procedure identifies a structure that is apparently time-independent, ie stationary (13). Stationary bilayer structures were obtained for DHP only for dispersions prepared by chloroform vaporization at pH values above 6. The DHP dispersions prepared by sonication were not stationary at any pH value (13).

Two other simple methods of determining the stability (stationary character) of synthetic amphiphile dispersions are turbidity determination as a function of time after vesicle preparation (5, 6) and determination of stability curves (23, 24), ie initial flocculation rates (v_0) as a function of monovalent salt concentration (Fig. 5). Sonicated DODAC or DHP vesicles prepared in water are not stable in contrast to the large vesicles prepared by chloroform vaporization. In the presence of NaCl, the same trend was observed (Tab. 2). Critical coagulation concentration (CCC) values were lower for the sonicated vesicles indicating a lowered stability in the presence of salt (24). However, Stern potential values estimated from the slope of the

stability curves were higher for the sonicated vesicles. Thus, the dispersion that is less stable has the highest electrostatic repulsion inbetween vesicles (24). There must be an extra attraction between the sonicated vesicles that does not occur for the large vesicles, which overwhelms the electrostatic repulsion and yields the lower stability measured. In fact, assuming only an electrostatic repulsion and a van der Waals attraction between vesicles, the calculated theoretical stability value for the sonicated DHP dispersion is much higher than the stability value obtained experimentally (25). An extra-attractive interaction force is needed to explain the low experimental stability of the sonicated DHP vesicles. This extra attraction can be due to an hydrophobic attraction between the edges of the bilayer fragments produced during sonication (13) or to a return to the lamellar state due to salt addition (10) or to both.

3b Aggregation and fusion

DODAC and DHP dispersions prepared by sonication with a tip undergo fusion upon NaCl addition (19, 24). For the large DHP vesicles, however, fusion was never observed (9, 19, 24). Thus, fusion in the sonicated dispersions could be associated with the formation of vesicles or coalescence of the bilayer fragments present in these dispersions (12, 13). Nevertheless, the contraction of DODAC monolayers at the air-water interface as the NaCl concentration increases in the subphase indicates a trend towards the formation of larger vesicles upon salt addition (9). In contrast, the DHP monolayer shows the opposite trend and expands as the NaCl concentration increases in the subphase (9). Furthermore, the high packing density of the DHP bilayer in the gel state does not favour fusion (11, 21). Adhesion, however, has often been described for the DHP system (6, 9, 11, 21). The adhesive force between DHP monolayers adsorbed on hydrophobic mica at 0.1- 0.3 nm separation distance

decreases as the NaCl concentration increases (pH 5.5) (9). A salt-induced shift of the half-dissociation of the phosphate polarheads to lower pH values might be increasing the charge density and the electrostatic repulsion (11, 21) and/or salt might disrupt adhesion originating from the formation of interbilayer hydrogen bonds at half-dissociation. Both phenomena would be observed as a decrease in the adhesive force as the salt concentration increases (9). At high temperature, very low NaCl concentration and pH 6.5 (close to the phosphate half-dissociation), a reversible temperature-induced vesicle aggregation has been described (11). The extent of aggregation markedly decreased as the NaCl concentration was increased. This was explained also by a salt-induced disruption of the interbilayer hydrogen bridges (11). Some direct measurements of adhesion between DHP layers are given in Table 3.

Whereas the DODAC system seems to be stabilized solely by electrostatic repulsion (26), fusing when this repulsion decays due to salt addition (19,26), the DHP system has a tightly packed bilayer (11, 21) which may adhere through hydrogen bonds at half-dissociation (11) or return to the lamellar state at higher salt concentration (7).

3c The surface potential and theoretical stabilities

Whether the classical DLVO theory for the stability of spherical charged colloidal particles is sufficient to describe the stability of the DODAC vesicles is still an open question (25). To further address this issue, theoretical stabilities must be calculated using the adapted DLVO model for vesicles and compared with the experimental ones (24, 25). However, Stern potentials are needed for calculating the electrostatic term of the interaction energy. We are presently measuring electrophoretic mobilities of large DODAC vesicles over an extensive range of NaCl

concentration to check the suitability of the DLVO forces to describe the interaction between synthetic amphiphile vesicles.

4 USES

4a Storage of solar energy and catalysis of chemical reactions

Charged synthetic amphiphile vesicles concentrate oppositely charged substrates at their bilayer/water interface. Consequently they may be used to separate charges generated by photoionization of certain molecules (28, 29) or they may catalyze certain chemical reactions by concentrating one of the reactants at the bilayer/water interface (28, 30). In this respect, synthetic amphiphile vesicles are as useful as ionic micelles and can be considered as a model system for the storage of light energy allowing compartmentalization of electron donors and acceptors relative to water (29). In addition, the selective incorporation of a charged substrate at the inner or outer vesicle surface and the permeation control of the other reactant can be used to modulate the reaction rate (30).

4b Bilayer adsorption

DODAC and DHP bilayers from DODAC and DHP dispersions prepared by sonication or by chloroform vaporization deposit onto oppositely charged polystyrene microspheres with high adsorption constants (7). At maximum adsorption, the areas per amphiphile molecule adsorbed, the electrokinetic properties and the sizes measured for the covered particles are consistent with bilayer deposition (7). The deposition process is controlled not only by the interaction forces between the vesicle and the microsphere but also by the restrictions of packing inside the bilayer. Vesicles tending to return to the lamellar state, ie to a state with larger curvature radius, have a high affinity for larger microspheres depositing as multibilayers and a low affinity for smaller

microspheres depositing as single bilayers. Interbilayer hydrogen bonds were found to enhance the multibilayer deposition.

4c Flocculation and stabilization of dispersions

DODAC and DHP vesicles are effective flocculants or stabilizers for oppositely charged dispersions in solution (Fig. 6). In contrast to single chained surfactants, they are effective at very low concentration (10-200 μM) (7). The division between their action as flocculants or stabilizers is defined by the equality between the total surface area for each dispersion. When there is more vesicle than particle surface area available then the vesicles act as stabilizers and when there is more particle than vesicle surface they act as flocculants. Another advantage over surfactants with short hydrocarbon chains is the irreversible character of the bilayer adsorption as shown by the high adsorption constants obtained (7).

4d Production of model colloids

Large DODAC or DHP vesicles are perfectly rigid spherical shells composed of a bilayer enclosing an aqueous compartment. They are highly charged and have a much smoother interface than other colloids. Because of these properties, they behave as model colloids and have been used to test theoretical models (24, 25). However their polydispersity is still a disadvantage compared with highly monodisperse systems as polystyrene microspheres (7). On the other hand, the possibly hairy, rough or conducting surface of these microspheres often represents a problem. Thus, the interaction between synthetic amphiphile vesicles and oppositely charged polystyrene microspheres leads to the production of an ideal model colloid: homodisperse and smooth bilayer covered polystyrene microspheres (Tab 4). From this system, significant new developments in colloid science can be expected.

Acknowledgments

I thank Dr Thelma M Herrington for valuable criticisms. Financial support from BID is gratefully acknowledged.

REFERENCES

- 1 Bangham, A D (1983) Editor *Liposome Letters*, Academic Press, London.
- 2 Kunitake, T ,Okahata, Y, Tamaki, K, Kumamaru, F & Takayanagi, M (1977) *Chem. Lett.* 387.
- 3 Mortara, R A, Quina, F H & Chaimovich, H (1978) *Biochem. Biophys. Res. Commun.* **81** 1080.
- 4 Kano, K ,Romero, A ,Djermouni, B , Ache, H & Fendler, J H (1979) *J. Am. Chem. Soc.* **101** 4030.
- 5 Carmona-Ribeiro, A M & Chaimovich, H (1983) *Biochim. Biophys. Acta* **733** 172.
- 6 Carmona-Ribeiro, A M , Yoshida, L S, Sesso, A & Chaimovich, H (1984) *J. Colloid Interface Sci.* **100** 433.
- 7 Carmona-Ribeiro, A M & Midmore, B R (1992) *Langmuir*, submitted.
- 8 Israelachvili, J N (1985) *Intermolecular and Surface Forces* Academic Press, London.
- 9 Claesson, P M ,Carmona-Ribeiro, A M & Kurihara, K (1989) *J. Phys. Chem.* **93** 917.
- 10 Carmona-Ribeiro, A M & Midmore, B R (1992) *J. Phys. Chem.*, submitted.
- 11 Carmona-Ribeiro, A M & Hix, S (1991) *J. Phys. Chem.* **95** 1812.
- 12 Pansu, R B ,Arrio, B , Roncin, J & Faure, J (1990) *J. Phys. Chem.* **94** 796.
- 13 Carmona-Ribeiro, A M , Castuma, C E ,Sesso, A & Schreier, S (1991) *J. Phys. Chem.* **95** 5362.
- 14 Augusto, O & Carmona-Ribeiro, A M (1989) *Biochemical Education* **17** (4) 209.

- 15 Deamer, D W & Bangham, A D (1976) *Biochim. Biophys. Acta* **443** 629.
- 16 Papahadjopoulos, D & Vail, W J (1978) *Ann. N.Y. Acad. Sci.* **308** 259.
- 17 Szoka, F & Papahadjopoulos, D (1978) *Proc. Natl. Acad. Sci. USA* **75** 4194.
- 18 Herrmann, H & Fendler, J H (1979) *Chem. Phys. Lett.* **64** 270.
- 19 Carmona-Ribeiro, A M & Chaimovich, H (1986) *Biophys. J.* **50** 621.
- 20 Koch, A L (1961) *Biochim. Biophys. Acta* **51** 429.
- 21 Carmona-Ribeiro, A M (1990) *J. Colloid Interface Sci.* **139** 343.
- 22 Lee, A G (1977) *Biochim. Biophys. Acta* **472** 237.
- 23 Reerink, H & Overbeek, J Th G (1954) *Disc. Faraday Soc.* **18** 74.
- 24 Carmona-Ribeiro, A M , Yoshida, L S & Chaimovich, H (1985) *J. Phys. Chem.* **89** 2928.
- 25 Carmona-Ribeiro, A M (1989) *J. Phys. Chem.* **93** 2630.
- 26 Pashley, R M , MacGuiggan, P M , Ninham, B W , Brady, J & Evans, D F (1986) *J. Phys. Chem.* **90** 1637.
- 27 Cuccovia, I M, Feitosa, E, Chaimovich, H, Sepulveda, L, Reed, W (1990) *J. Phys. Chem.* **94** 3722.
- 28 Fendler, J H (1980) *Acc. Chem. Res.* **13** 7.
- 29 Bratt, P, Kang, Y S, Kevan, L, Nakamura, H, & Matsuo, T (1991) *J. Phys. Chem.* **95** 6399.
- 30 Cuccovia, I M, Kawamuro, M K, Krutman, M A K & Chaimovich, H (1989) *J. Am. Chem. Soc.* **111** 365.

TABLE 1: Calculation of the geometric parameter v/al for DODAC and DHP assuming that the area per monomer a is equal to the limiting area per monomer at 25 mN/m in DODAC and DHP monolayers. C is the NaCl concentration. (Adapted from references 9 and 10).

Amphiphile	C/M	pH	v/nm^3	l/nm	a/nm^2	v/al
DODAC	0	6.5	.969	2.3	.66	.64
	.0001				.54	.79
	.0010				.52	.81
	.0100				.52	.81
DHP	0	7.8	.862	2.1	.425	.99
	.0001	7.3			.427	.985
	.0010	6.9			.426	.988
	.0100	6.2			.421	1

TABLE 2 : The stability of DODAC and DHP dispersions at pH 6.5 in the presence of NaCl. C_a is the amphiphile concentration; CCC is the critical coagulation concentration; v_{Omax} is the maximal initial flocculation rate and Ψ_δ is the estimated Stern potential. These estimated Stern potentials are too low in comparison with the real surface potentials measured by electrophoresis (10, 27) due to the DLVO approximation used for the calculations but they reflect well the differences between the dispersions. (Adapted from references 10, 24 and 27).

Dispersion	C_a /mM	CCC/M	$10^3 v_{Omax}$	Ψ_δ /mV	
				Calculated	Measured
Sonicated DHP	.23	.178	12.6	-41.2	-97.3
Injected DHP	.18	.229	3.8	-19.3	-70.7
Sonicated DODAC	.21	.112	1.5	26.8	85.0
Injected DODAC	.24	.129	.1	8.5	84.8

TABLE 3 : Direct measurements of adhesion forces between DHP layers adsorbed onto hydrophobic mica at room temperature as a function of NaCl concentration (C) and pH. The force (F) is normalized to the curvature radius on mica and expressed as mN/m . (Adapted from reference 9).

C /M	pH	F / mN m^{-1}
.0002	5.5	-5.4
.0010	5.5	-3.7
.0100	5.5	-3.3
.0100	3.0	-7.2
.0100	9.5	0

TABLE 4 : The homodisperse and smooth bilayer covered polystyrene microspheres as produced by mixing sonicated DODAC dispersions and sulphate polystyrene microspheres (150 nm of mean z-average diameter). The ratio between the vesicles number density and the microspheres number density is 7 ($N_v/N_p = 7$). At $N_v/N_p = 7$, the limiting diameter (D_z) for the mixture upon increasing dilution before mixing is 160 nm, ie the size of the microsphere plus about 10 nm what suggests surfactant deposition of one-bilayer thickness. (Adapted from reference 7).

C_a /mM	$N_p / 10^{12} \text{ cm}^{-3}$	mean D_z /nm	Polydispersity index
1	1.04	172.0	0.065
.4	.42	158.0	0.066
.2	.21	162.0	0.061
.05	.05	160.0	0.044
0	1.04	150.0	0.043

Figure Captions

Figure 1: The chemical structure of some double-chained ionic amphiphiles which form bilayers in water solution. Dioctadecyldimethylammonium chloride (DODAC) is a quaternary ammonium salt with a cationic polar head and chloride as counterion. Dihexadecylphosphate (DHP) is a derivative of phosphoric acid with an anionic phosphate polar head and protons or sodium ion as counterions. The acidic character of the DHP polar head originates a myriad of pH effects on DHP bilayer properties (11).

Figure 2: Association of amphiphilic molecules at air-air, air-water and water-water interfacial regions. The hydrocarbonic chains interact preferentially with each other and with air whereas the polar heads expose themselves to water.

Figure 3: Molecular geometry of amphiphiles and lipids that are able to form planar bilayers and bilayer vesicles. A truncated conic molecular shape is required to form a curved bilayer and a cylindric one to form a planar bilayer. (Adapted from reference 8).

Figure 4: Time-dependent absorbance changes after addition of 0.05 M KCl (—) or 0.1 M sucrose (- - -) to large DODAC vesicles (A) or to DODAC dispersions obtained by sonication (B). The turbidity increase after sucrose addition is due to vesicle shrinkage (A) and is not observed for DODAC or DHP dispersions prepared by sonication (B). The absence of osmotic response for the sonicated dispersions may suggest the inexistence of an inner aqueous compartment or the impossibility of detecting changes in size using the turbidimetric method. The increase in the turbidity of the sonicated dispersion due to KCl addition is due either to vesicle fusion or bilayer fragments coalescence (19, 24). (Adapted from reference 5).

Figure 5: The stability of synthetic amphiphile vesicles in water (A, B, C, D) or in NaCl solutions (E, F). The turbidity of DODAC (A) or DHP (B) dispersions in water varies as a function of time after dispersion whereas DODAC (C) and DHP (D) dispersions obtained by chloroform vaporization are stationary. By adding salt solutions to vesicle samples under conditions of isoosmolarity, the turbidity increases as a function of time indicating a low colloidal stability of the vesicles in the presence of salt (E). Quantification of stability is possible by measuring the initial flocculation rate (v_0) from the turbidity vs. time curve at a given NaCl concentration, C. As C increases, v_0 increases and the stability ($1/v_0$) decreases (from curve a to c, in E). Thus, stability curves like that in (F) were obtained for synthetic amphiphile dispersions in reference 24. The profile of $\log v_0$ against $\log C$ is as predicted by the DLVO theory for the stability of colloidal dispersions (F). Important parameters defining the colloidal stability are indicated in the stability curve in (F): the critical coagulation concentration (CCC); the slope of the flocculation curve ($\lg \theta$); the maximal initial flocculation rate ($v_{0\max}$). (Adapted from references 6 and 24).

Figure 6: Synthetic amphiphile vesicles acting as flocculants or as stabilizers of oppositely charged polystyrene microspheres. Large DODAC vesicles with a mean z-average diameter (D_z) of 256 nm were mixed with sulphate polystyrene latices (285 nm) 4 h before the D_z determination. One week after mixing, flocculation was visible for samples in which the ratio between number densities for vesicles and polystyrene microspheres (N_v/N_p) was smaller than 1 and absent when $N_v/N_p > 1$. (Adapted from reference 7).

DODAC

DHP

22 January 1992

Our ref: KJW/sjc:413

Thomas Graham House
Science Park
Milton Road
Cambridge CB4 4WF
United Kingdom

Tel: Cambridge (0223) 420066
Telex: 818293 ROYAL
Fax: Cambridge (0223) 423623
Dialcom/Telecom Gold 84: BUR 500

Dr A M Carmona-Ribeiro
Departamento de Bioquímica
Instituto de Química
Universidade de São Paulo
CP 20780 - São Paulo SP
Brazil

Dear Dr Carmona-Ribeiro

Chemical Society Reviews

Thank you for your letter of 16 January. I am pleased to inform you that your article 'Synthetic Amphiphile Vesicles' will be published in *Chemical Society Reviews*, 1992 (Vol.21) Issue 4 which will appear towards the end of November this year. I apologise that we are unable to publish it earlier but we have been overwhelmed by the number of reviews received for the new-style *Chemical Society Reviews*.

Proofs of your article will be despatched (airmail) to you on 10 September. Please let me know if you anticipate having difficulty in correcting and returning your proofs within a few days of their receipt.

An honorarium of £300 will be paid into your bank account in Reading on publication of the review.

Thank you for contributing to *Chemical Society Reviews*.

Yours sincerely

K J Wilkinson
Senior Editor, Reviews
Books & Reviews Department

c.c. Dr Thelma M Hardman

Referência 18

PHOSPHOLIPID ADSORPTION ONTO POLYSTYRENE MICROSPHERES

A. M Carmona-Ribeiro* and T. M. Herrington‡

* Departamento de Bioquímica, Instituto de Química,
Universidade de São Paulo, Caixa Postal 20780,
01498 São Paulo SP- Brazil

To whom all correspondence should be addressed

‡Department of Chemistry, University of Reading,
Reading, RG6 2AD United Kingdom

ABSTRACT

Small unilamellar phospholipid vesicles and polystyrene microspheres interact in aqueous solution to form homodisperse and stable phospholipid covered latexes. Firstly the bilayer attaches to the latex. Secondly, the hydrophobic attraction between the phospholipid bilayer and the hydrophobic polystyrene surface induces coverage with one phospholipid monolayer. Thereafter phospholipid bilayer(s) deposit onto the monolayer covered latex. These results may be of importance for reconstitution of protein function, studies on cell surface recognition and building-up of immunological kits and biosensors.

INTRODUCTION

The interaction between phospholipid bilayers and other biological or non-biological surfaces is poorly understood.

DPPC and PI from vesicles adsorb onto negatively charged glass beads as a monolayer with their head groups uppermost (1). Unilamellar PC vesicles disrupt and adhere to a mica surface to form a bilayer coating (2). Phospholipid monolayers with lipid haptens inserted can be supported by hydrophobic glass and are useful for specific adherence of macrophages and cell surface recognition studies but cannot serve as hosts for transmembrane proteins (3). Charged synthetic amphiphile bilayers composed of DODAC or DHP (4) are electrostatically attracted to oppositely charged polystyrene microspheres forming homodisperse bilayer covered lattices (5).

Here phospholipid adsorption onto polystyrene microspheres is investigated by two complementary techniques: 1) determination of phospholipid adsorption isotherms onto the latexes; 2) Dz measurement for the mixtures as a function of A_v/A_p . Maximal adsorption levels in the isotherms and Dz values for the vesicle/microsphere mixtures are consistent with hydrophobic adsorption of one monolayer of phospholipid onto the polystyrene followed by deposition of phospholipid bilayer(s).

MATERIAL AND METHODS

ASO from soybean (type IV-S) and PC, DPPC and DHP of the highest purity available were from Sigma Chem. Co..

Phospholipid and DHP concentration were determined by inorganic phosphorus (Pi) analysis (6). Charged polystyrene microspheres described as ultraclean by the supplier were obtained from Interfacial Dynamics Corporation and used as such. Microspheres with 3 different stabilizing charged groups were used: SU, CA and AM. Water was MILLI-Q quality.

ASO SV were prepared by sonication (7) and PC, DPPC or DPPC/DHP (10:1) SV, by ethanolic injection (8, 9) followed by 6 h dialysis and 1h centrifugation at 20000 g and 15 °C to eliminate any eventual multilamellae.

Interaction between SV and microspheres was induced by adding SV to the latex. Final Np for the mixtures is in Tab. 1. Mixtures were thermosttated at 25 °C for 72 h. Thereafter they were centrifuged at 20000 g for 1 h at 15 °C to separate particles from SV. The supernatant was used to determine phospholipid concentration. Adsorption isotherms of phospholipids onto the microspheres were obtained from Pi analysis in the supernatant and in the SV preparation. The sensitivity of the Pi analysis is 0.01 μ M Pi (6). Ap was calculated from the mass fraction and the specific surface area (5). Adsorption was expressed as the number of amphiphile molecules adsorbed per m² polystyrene.

Isotherms were linearized using the Langmuir model to determine adsorption constants k (1, 5). Limiting adsorption, $(x/m)_{\max}$, was determined from the plateau levels on the isotherms.

Particle size was measured using PCS. The equipment was a Malvern 4700c apparatus employing a Coherent Innova 90 laser. The size quoted throughout is D_z of at least 15 independent measurements at 25 °C.

RESULTS

Fig. 1 shows typical adsorption isotherms for phospholipid adsorption from SV onto SU, CA and AM microspheres in water at 25 °C. From the plateau regions on the curves, which are regions of maximal adsorption, the area per phospholipid molecule adsorbed can be calculated. For PC, this area is .69 nm² for the first plateau (Fig. 1A; Tab. 1). This is very close to the usual limiting area per monomer in PC monolayers at the air-water interface suggesting monolayer deposition onto the AM polystyrene surface. For the second plateau, the area per PC molecule adsorbed is .21 nm² (Fig. 1; Tab. 1). This is consistent with the area expected for deposition of one PC monolayer onto the bare latex and one PC bilayer onto the monolayer covered microspheres.

Limiting adsorption was obtained at areas per phospholipid molecule adsorbed which were consistent with one, three or five phospholipid monolayers adsorbed (Fig. 1 A- D; Tab. 1).

The effect of the ratio between total surface areas for SV and for polystyrene (A_v/A_p) on the particle size (D_z) in mixtures of phospholipid SV and AM polystyrene particles of nearly the same size is in Fig. 2. The final aggregation state of the mixture depends on A_v/A_p .

At $A_v/A_p = 0.5$, a minimum D_z value is obtained in all cases showing latex stabilization due to adsorption of the phospholipid monolayer. In contrast, at $A_v/A_p = 1$, D_z is a maximum showing that bilayer adsorption leads to aggregation (Fig. 2). For $A_v/A_p > 1$, D_z progressively decreases as a function of A_v/A_p indicating further stabilization of the mixtures as more phospholipid is added (Fig. 2). In particular, for PC adsorption, D_z levels off from $A_v/A_p = 1.5$, ie from the situation where one monolayer plus one bilayer deposit onto the latex (Fig. 2 b).

Dispersions which are very diluted before mixing are convenient to detect the early stages of the adsorption. For diluted DPPC/DHP SV and AM microspheres, there is bilayer adsorption (Fig.1 D, insert; last line on Tab. 1) accompanied by massive aggregation as detected by PCS (not shown).

The effect of dilution of the dispersions before mixing on D_z for a mixture of ASO SV and AM latex at $A_v/A_p = 9.8$ is in Tab. 2. As the diameter of the monolayer covered latex particle is 96.4 nm, a D_z value of 125.6 nm is consistent with two bilayers deposited onto the monolayer covered latex

if the separation distance between layers is taken as 2- 3 nm and the bilayer thickness is taken as 5 nm (Tab. 2). By mixing more diluted dispersions at the same A_v/A_p ratio, only one bilayer adsorption on the monolayer covered latex occurs as depicted from a D_z value of 108- 110 nm (Tab. 2). The mixtures have low polydispersity indexes and are very stable (Tab. 2).

DISCUSSION

The highest phospholipid affinity for the latex surface, k , occurs for the positively charged amidine polystyrene, AM, possibly due to the specific interaction between amidine on the latex and the phosphate of the phospholipid (Fig. 1; Tab. 1). Consistently, for the negatively charged carboxylate polystyrene, CA, which is more hydrophilic, lower phospholipid affinity was obtained. Moreover, the less hydrated DPPC bilayer (10) interacted with higher affinity for the latexes than the more hydrated PC bilayer (Fig. 1, Tab. 1). Thus, surface hydration shields the hydrophobic attraction between the phospholipid hydrocarbon chains and the latex surface decreasing k for monolayer coverage.

At high dilution before mixing, the zeroth step in the adsorption process is entire phospholipid bilayer attachment to the latex (Fig. 1 D, insert). A similar adsorption mechanism involving the entire bilayer instead of monomer adsorption was demonstrated for DPPC and PI adsorption onto glass balls (1) and for DODAC and DHP bilayer adsorption onto oppositely charged polystyrene microspheres (5).

After bilayer attachment, the hydrophobic attraction between the latex surface and the hydrocarbon chains of the phospholipids destroys the bilayer structure as the hydrocarbon tails turn themselves to the latex surface. At this stage, exposure of hydrophobic regions in the collapsed bilayer causes massive and stable aggregation if $A_v/A_p = 1$. When the total surface area for vesicles and polystyrene is the same and each microsphere is covered with a collapsed bilayer it is difficult to deliver the phospholipid excess to other latex particles which are also covered with a collapsed bilayer mainly if the lipids are in the rigid gel state. Consequently, D_z is maximum for all mixtures studied though this maximum is much lower for phospholipids in the liquid-crystalline state, eg PC and ASO (Fig. 2). In contrast, at $A_v/A_p = 0.5$ and monolayer coverage, D_z is minimum (Fig. 2). In this case, the problem of the collapsed bilayer attached to a latex particle is promptly solved by adhesion of other bare latex particles which become monolayer covered and stabilized.

We have been observing the stabilizing effect of the monolayer covering step by studying latex deflocculation at $A_v/A_p = 0.5$. Deflocculation is instantaneous for the positively charged amidine latex added of the neutral PC SV and takes about 10 minutes for SU latex added of the equally negatively charged ASO SV (to be reported elsewhere). The latter instance definitely shows the hydrophobic attraction overcoming the electrostatic repulsion. In this respect, the

mechanism for phospholipid adsorption onto polystyrene microspheres is completely different from that for adsorption of synthetic amphiphile bilayers onto oppositely charged latexes (5). For the extremely rigid and well-packed DODAC and DHP bilayers in the gel state (4), electrostatic attraction overwhelms the hydrophobic one and monolayer coverage is absent (5). At bilayer coverage ($A_v/A_p = 1$), D_z levels off (5). Thus, the zeroth and unique step in the synthetic bilayer adsorption mechanism onto oppositely charged microspheres is a stable bilayer attachment due to electrostatic attraction (5) whereas phospholipids adsorb onto latexes through an unstable bilayer attachment with the hydrophobic attraction directing monolayer deposition.

ACKNOWLEDGMENTS

We thank CNPq and BID for financial support.

REFERENCES

- 1) Jackson, S., Reboiras, M. D., Lyle, I. G. and, Jones, M. N. (1986) Faraday Discuss. Chem. Soc. **85**, 291- 301.
- 2) Horn, R. G. (1984) Biochim. Biophys. Acta **778**, 224- 228.
- 3) Lin, L.-c., Weis, R.M., and McConnell, H.M. (1982) Nature **296**, 164- 165.
- 4) Carmona-Ribeiro, A.M. (1992) Chem.Soc.Reviews **21**(4), in press.
- 5) Carmona-Ribeiro, A.M., and Midmore, B.R. (1992) Langmuir, in press.
- 6) Houser, G., Fleischer, S., and Yamamoto, A. (1970) Lipids **5**, 494-496.
- 7) Huang, C. H. (1969) Biochemistry **8**, 344- 352.
- 8) Batzri, S., and Korn, E. D. (1973) Biochim. Biophys. Acta **298**, 1015-1019.
- 9) Kremer, J. N. H., Esker, M. W., Pathmamanoharan, C., and Wiersema, P. H. (1977) Biochemistry **16**, 3932- 3935.
- 10) Rand, R. P. (1981) Annu. Rev. Biophys. Bioeng. **10**, 277- 314.

ABREVIATIONS: DPPC, dipalmitoylphosphatidylcholine; PI, phosphatidylinositol; PC, phosphatidylcholine; DODAC, dioctadecyldimethylammonium chloride; DHP, sodium dihexadecylphosphate; Dz, mean zeta-average diameter; PCS, photon correlation spectroscopy; ASO, asolecithin; Av and Ap, total surface areas for the population of vesicles and bare microspheres, respectively; k, adsorption constant; (x/m)_{max}, maximal adsorption; Nv, Np, number density of vesicles or microspheres; SV, small unilamellar vesicles; AM, amidine ; SU, sulfate ; CA, carboxylate polystyrene; V, vesicles; L, latex; P, polydispersity index; C_{sup}, phospholipid concentration in the supernatant; nads, number of molecules adsorbed per m² polystyrene.

Table 1

Limiting adsorption and adsorption constants for the adsorption of PC (A), ASO (B), DPPC (C), DPPC/DHP 10:1 (D) from small vesicles, V, in water onto AM (a), SU (b), CA (c) polystyrene microspheres, L, at 25 °C. Areas per molecule adsorbed at maximum adsorption (A_{ads}) are compared with limiting areas per monomer for pressure/area isotherms from the literature (A). A/A_{ads} is the number of phospholipid monolayers adsorbed. Sizes measured by PCS, D_z , are given for vesicles and latexes.

V, D_z /nm	L, D_z /nm	N_p / 10^{11} cm^{-3}	$(x/m)_{max}$ / $10^{17} \text{ molecules m}^{-2}$	K / $10^4 \text{ dm}^3 \text{ mol}^{-1}$	A_{ads} / nm^2	A / nm^2	A/A_{ads}
PC62	AM104	7	14.5 47.0	5 11	.69 .21	.70	1 3.3
	SU85	7.4	14.5	3	.69		1
	CA283	0.5	14.5	1	.69		1
ASO108	AM104	7	14.3 44.0 77.0	70 7 6	.70 .23 .13		1 3.1 5.4
DPPC74	AM104	7	20.0 63.6 106.6	56 - -	.50 .16 .094	.50	1 3.2 5.3
	SU85	7.4	20.0	1	.50		1
	CA283	0.5	20.0	1	.50		1
DPPC/ DHP60	AM104	7 1.2	100.0 40.0	51 1825	.10 .25	.50	5 2

Table 2

The effect of dilution of the dispersions before mixing on Dz for particles in a mixture of AS0112 and AM104 at Av/Ap = 9.8. On the last two lines, AM104 is covered with one monolayer of PC (Av/Ap = 0.6), Dz decreases from 104 to 96.4 nm and then, ASO SV are added (Av/Ap = 9) and the Dz is determined again being consistent with one bilayer deposition (108.2 nm = 96.4 nm monolayer covered latex + 10 nm bilayer coverage + 1.8 nm water between phospholipid layers). All Dz are constant as a function of time after mixing (0.2- 92 h).

Phospholipid /mM	Np /10 ¹¹ cm ⁻³	Dz /nm	P	Av/Ap
ASO/0.74	8.75	125.6 ± 0.7	.135 ± .020	9.8
0.37	4.38	110.3 ± 0.8	.091 ± .035	
0.148	1.75	109.8 ± 1.4	.094 ± .019	
0.074	0.88	110.9 ± 1.1	.102 ± .024	
0.037	0.44	107.6 ± 1.1	.106 ± .038	
PC/ 0.037	8.75	96.4 ± 1.4	.054 ± .036	0.6
ASO/0.37	4.38	108.2 ± 0.8	.078 ± .040	9.0

FIGURE LEGENDS

Fig. 1: Isotherms for the adsorption of PC (A), ASO (B), DPPC (C) and DPPC/DHP 10:1 (D) from small vesicles in water onto AM (a), SU (b) and CA (c) polystyrene microspheres in water at 25 °C. Dz for vesicles is 62, 108, 74 and 60 nm, respectively. For microspheres, Dz is 104 (AM), 85 (SU) and 283 (CA). The insert shows an isotherm obtained for very diluted dispersions before mixing. Number density for microspheres and isotherm analysis are in Tab. 1.

Fig. 2: Dz as a function of A_v/A_p for ASO (a), PC (b) and DPPC/DHP 10:1 (c) mixed with AM104 polystyrene microspheres at N_p equal to 7 (a), 8.75 (b) and 1.2×10^{11} particles per cm^3 (c). Dz for vesicles are those quoted on Fig. 1. Dz for AM microspheres is 104 nm. The arrows indicate the minimum Dz at $A_v/A_p = 0.5$.

CARMONA-
-RIBERO.
HEADINGTON

FIG. 2

Referência 19

**The low colloidal stability of charged synthetic amphiphile
vesicles and the DLVO**

A. M. Carmona-Ribeiro

**Departamento de Bioquímica, Instituto de Química,
Universidade de São Paulo, CP 20780, São Paulo, SP, Brazil**

ABSTRACT

Zeta-potentials for dioctadecyldimethylammonium chloride (DODAC) and sodium dihexadecylphosphate (DHP) vesicles as a function of NaCl concentration (C_e) are obtained over an extensive range of salt concentration (10^{-4} - 10^{-1} M) using particle microelectrophoresis. These estimates of the surface potential at a fixed vesicle size allow the calculation of theoretical stabilities for the vesicles (W) using a DLVO model adapted for interacting spherical vesicles without free parameters. W values thus obtained are much higher than the experimental stabilities from vesicle flocculation under identical experimental conditions (W_e). It is concluded that destabilizing factors not accounted for by the theory are needed to explain the experimental reality and/or the DLVO is overestimating the electrostatic repulsion.

INTRODUCTION

Charged synthetic amphiphile vesicles are becoming increasingly important as biomimetic systems (1- 5), convenient media for charge separation and storage of light energy (6- 8), model colloids for testing theory (9, 10) and interfacial agents able to cover and modify surfaces at very low concentration (11). Nevertheless, some of their fundamental properties are still controversial as, for example, the nature of the interaction forces between synthetic amphiphile bilayers.

Whereas some authors report the absence of hydration forces for interacting bilayers of dihexadecyldimethylammonium acetate (DHDAAC) or bromide (DHDAAB) adsorbed on anionic mica surfaces (12, 13), others criticize the surface force technique at short ranges and determine extrarepulsive hydration forces between nonadsorbed DHDAAC bilayers (14). On the other hand, at pH 5.5- 6, an extraattractive interaction force between adsorbed dihexadecylphosphate (DHP) layers on mica was determined (19) and independently confirmed for DHP vesicles using the kinetic DLVO approach (9). Here we further explore this approach to clarify the nature of the interaction forces acting between synthetic amphiphile bilayers over an extensive range of NaCl concentration.

Firstly the zeta-potential for large DODAC and DHP vesicles is determined as a function of NaCl concentration at a fixed vesicle size using particle microelectrophoresis. Thus, the electrostatic term of the DLVO interaction energy can be calculated and added to the van der Waals interaction energy for spherical water particles covered with an hydrocarbon shell interacting across water as the intervening medium. Secondly, theoretical DLVO stabilities (W) calculated from the interaction energy profile as a function of the separation distance are compared with the experimental stabilities (W_e) obtained previously (3, 9) from vesicle flocculation data. W values are much higher than W_e indicating the existence of

extraattractive interactions between interacting DODAC and DHP vesicles and/or an overestimation of the repulsive energy.

MATERIAL AND METHODS

Diocetadecyldimethylammonium chloride (DODAC) was obtained from Herga Industrias Quimicas do Brasil. Dihexadecylphosphoric acid (DHPH) was obtained from Sigma (St.Louis, MO.) and was used to prepare sodium dihexadecylphosphate (DHP) as described in (2). All other reagents were analytical grade and were used without further purification. Water was MILLI-Q quality.

Large DODAC and DHP vesicles (LV) were prepared by injecting a chloroform solution of DODAC or DHP into an aqueous D- glucose solution, 75 °C, at pH 5-5.5. DODAC and DHP LV were prepared at a D-glucose concentration (C_G) equal to 0.43 and 0.55 M, respectively. The reason for using these C_G values was the need to reproduce precisely the experimental conditions under which NaCl- induced vesicle flocculation was kinetically studied and initial flocculation rates were determined as a function of NaCl concentration (C_e) (3).

In order to prevent and control any change in size due to vesicle shrinkage, swelling or aggregation, vesicle size was measured before and after addition of composite isoosmolar solutions of NaCl and D-glucose. The size was found to be constant as a function of time or NaCl concentration up to 0.045 and 0.069 M NaCl, for DODAC and DHP LV, respectively. Sizes were measured using a Malvern 4700c PCS apparatus employing a Coherent Innova 90 laser and correcting the diffusion coefficients for the viscosity and refractive index of the vesicles medium. The size quoted is the mean harmonic z-average diameter (D_z) of at least 15 independent measurements at 25 °C.

Electrophoretic mobilities (EM) were determined at 25 °C using a Rank Brothers Mark II particle microelectrophoresis apparatus in the configuration for the flat cell.

DHP and DODAC concentration (C) were determined by inorganic phosphorus analysis (16) and by amphiphile-dye solubilization in micelles (17), respectively. Because of the large effect of amphiphile concentration (C) on EM (10), mobilities had to be determined as a function of amphiphile concentration at a given NaCl concentration. The extrapolation of the EM against C curve to zero amphiphile concentration yielded the EM_0 value. EM_0 was corrected to take into account the viscosity of the vesicles medium (EM_{oc}). EM_{oc} was then transformed into the reduced mobility, E (15). From E and D_z , the zeta-potential was calculated using the O'Brien and White theory (15). From the zeta-potential and D_z , the charge density was calculated using an approximated solution of the nonlinear Poisson-Boltzmann equation for the case of spherical double-layers at high potentials (10).

COMPUTATIONS

The total potential energy (V) of interaction as a function of the reduced separation distance (u) was calculated from vesicle radius ($D_z/2$), Stern potential (zeta-potential), NaCl concentration (C_e) and 5×10^{-14} ergs as the Hamaker constant (A) using the DLVO model adapted for vesicles without free parameters (9). The computer was programmed to tabulate and plot the coordinates of ($V/kT, u$) curves. The unit of k is cm^{-1} and if the zeta-potential is expressed in millivolts and the vesicle radius in cm, the unit of V is erg. The dimensionless dielectric constant of water at 25°C was taken as 78.54. The stability ratio (W) was obtained by numerical integration of eq 8 in reference 9. In all computations of V and W only aqueous solutions of 1:1 electrolyte at 25°C were considered. The experimental stability values (W_e) were calculated from vesicle flocculation data (3) using the DLVO algorithm previously adapted for interacting vesicles (9).

RESULTS

The electrophoretic mobility of large and charged synthetic amphiphile vesicles (EM) increases with the amphiphile concentration (C) (10). For the vesicles prepared in D-glucose solution and added of an isoosmolal NaCl/D-glucose solution, an analogous amphiphile concentration effect was obtained (Fig. 1 A, insert). Thus, the extrapolation to zero amphiphile concentration in the EM against C curve was necessary for each NaCl solution added to the vesicles.

The NaCl concentration effect on the zeta-potential of synthetic amphiphile vesicles is shown in Fig. 1 and Tab. 1. The zeta-potential decays exponentially with the concentration of the NaCl solution added to the vesicles for both vesicle types (Fig. 1). The zeta-potential varies linearly with the logarithm of the NaCl concentration (Fig. 1).

Under the present experimental conditions which keep the vesicle size fixed, both surface potential and surface charge vary as a function of the added electrolyte concentration (Tab. 1). Nevertheless, by increasing the electrolyte concentration from 10^{-4} to 0.045 M the surface potential decreases from 85 to 46 mV and from -78 to -51 mV, for DODAC and DHP vesicles respectively (Fig. 1 and Tab. 1). The range of surface potential variation is narrower for the DHP vesicles. The opposite occurs for the variation range of the surface charge which is narrower for DODAC than for DHP (Tab. 1).

The determination of the zeta-potential dependence on the electrolyte concentration over a range of C_e where vesicle flocculation is absent allows to extrapolate the surface potential to the region of higher NaCl concentration where flocculation occurs (Fig. 1). From the estimated surface potential, the potential energy of interaction as a function of the separation distance between vesicles was calculated using the DLVO model (Fig. 2). The resultant repulsive energy has maxima which are very high and vary between 80 and 250 kT depending on NaCl concentration (Fig. 2). As a consequence, theoretical stabilities (W) are very high or even too high to be calculated (Tab. 2). W values are much higher than the experimental stability values (W_e) (Tab. 2).

DISCUSSION

The nature of extraattractive and extrarepulsive interaction forces between surfaces is still a challenge in surface chemistry (18- 20). In spite of being often found acting between several kinds of surfaces, they are poorly understood and the theoretical models hitherto developed are often restricted explaining just a few isolated cases.

In particular, it is difficult to envisage how protusion or bilayer undulations could explain the "hydration" forces between bilayers in the rigid gel state as those

composed of dihexadecyldimethylammonium acetate (14) or dihexadecylphosphate at high pH (19). It seems more reasonable in these cases to assume that binding of hydrated ions originates the extrarepulsive force at short ranges and very low electrolyte concentration. On the other hand, very little is known about extraattractive interactions. The Poisson-Boltzmann approach breaks down in concentrated electrolyte solutions due to the neglect of ion-ion correlations (30). Furthermore, direct measurements of interaction forces between amphiphile bilayers in the gel state at moderate and high electrolyte concentration are absent in the literature.

For DODAC, an extrarepulsive force at low electrolyte concentration (14) not only disappears completely but also gives place to an opposite extraattractive interaction as the NaCl concentration is increased (Tab. 2). Possibly this is why bilayer membranes that occur in nature are not composed of synthetic amphiphiles as DODAC and DHP. Their instability at moderate and high electrolyte concentration precludes their existence as stable closed bilayers suitable for compartmentalization and other essential membrane functions.

In this work we provide the literature with reliable estimates of the surface potential in DODAC and DHP vesicles as a function of NaCl concentration over an extensive salt concentration range. This reliability can be checked by comparing our values for the surface charge density (Tab. 1) with those obtained previously by using the surface force apparatus. At pH 5.5 and 0.2 mM NaCl, 62 nm^2 per charge or 2.6 mC/m^2 were obtained from nonlinear Poisson-Boltzmann and direct surface force measurements (19). At pH 5.5 and 0.1 mM NaCl we estimated 2.7 mC/m^2 (Tab. 1). At pH 5.5 and 1 mM NaCl, the surface force technique yielded 31 nm^2 per charge (5.2 mC/m^2) and our mobility measurements for DHP vesicles under similar experimental conditions gave 6.6 mC/m^2 (Tab. 1).

At 2 mM KBr, for DHDA bilayers adsorbed on mica, a value of 18 mC/m^2 was reported (12). We calculated 10 mC/m^2 for DODAC vesicles at 2.7 mM NaCl.

While the DHP system at pH 5.5 presented extraattractive interaction forces at low (19) and moderate to high electrolyte concentration (Tab. 2), the DODAC system presented extrarepulsion at low electrolyte concentration ranges (14) and extraattraction at higher ranges (Tab. 2). The mechanism accounting for this change in behavior could perhaps involve exposure of hydrophobic regions in the bilayer. Indeed, the hydrophobic attraction measured between DHDA monolayers adsorbed on mica is about 100 times stronger than normal van der Waals attraction and causes water structure to decay rapidly in the vicinity of the hydrophobic regions (22). For the DODAC system, an unusually low value for the bilayer thickness at low ionic strength has been reported from measurements using the surface force apparatus (12, 14) and ellipsometry measurements of DODAC LB films (21). It is possible that interdigitation and/or tilting of the DODAC bilayer are hiding the methyl groups of the polar heads so that at high electrostatic repulsion (low ionic strength), the adsorption of hydrated counterions at the quaternary ammonium polar head can account for the existence of an hydration shell at the bilayer/water interface. When the electrolyte concentration increases and thereby the electrostatic repulsion decreases, the proximity of the polarheads could make interdigitation and tilting more difficult leading to a return to the usual bilayer structure and thickness in which exposure of hydrocarbonic chains would not be as rare as desirable for a functional bilayer structure.

A decrease in the area per monomer in a given bilayer structure is known to decrease the range and magnitude of extrarepulsive forces (18, 23). For example, the decrease in area per monomer that occurs when a phospholipid bilayer undergo a transition from the liquid-crystalline to the gel state significantly decreases

magnitude and range of extrarepulsive short-range forces. DODAC and DHP bilayers are in the gel state and have the smallest area per monomer in a scale where all double chained amphiphiles were included. As electrolyte concentration increases an additional decrease of the area per monomer could cause opening of hydrophobic areas. Consistently DODAC vesicles fuse due to salt addition (50 mM NaCl) (4) and an extraattractive interaction force is indicated from our present results (Tab. 2).

When the electrostatic potential of a spherical or planar charged surface is relatively high, a general, analytical solution of the Poisson-Boltzmann equation does not exist (24, 25). For a single planar surface in a symmetrical electrolyte, the nonlinear Poisson-Boltzmann equation can be integrated analytically in two steps (26) yielding the surface potential as a function of electrolyte concentration at a given charge density (25). The surface potential decreases exponentially as a function of electrolyte concentration, ie, the surface potential is a linear function of the logarithm of the electrolyte concentration (25). For the large vesicles, we obtained a straight line for the dependence of the zeta-potential as a function of the logarithm of the electrolyte concentration (Fig. 2). Because the vesicles are large, their curvature is small allowing the assumption of a planar bilayer to be not very far from the real situation. The surface potential in sonicated DHP vesicles plotted as a function of C_e also yields a straight line (27, 9) possibly because sonication is a disruptive process which generates planar bilayer fragments (28).

In summary, DODAC vesicles fuse and DHP vesicles aggregate upon NaCl addition to a final concentration of 50 mM (4, 9, 19) and there is in both cases an extraattractive interaction force which is not accounted for by the DLVO. The nature of this extraattraction may be completely different in each case. Fusion might be induced by hydrophobic attraction and aggregation, by the formation of

interbilayer hydrogen bridges (at pH 5.5- 6, the DHP bilayer is not very far from half-dissociation of the phosphate polar heads, see reference 29). Furthermore, the Poisson-Boltzmann approach breaks down in concentrated electrolyte solutions due to the neglect of ion-ion correlations (30, 31). Knowledge about the nature of the extraattractive interaction is at the present time not sufficient to decide between the possible interpretations.

ACKNOWLEDGMENTS

CNPq, FAPESP and BID are gratefully acknowledged.

REFERENCES

- 1 Carmona-Ribeiro, A. M., Chaimovich, H. **Biochim. Biophys. Acta** **733** 172 (1983).
- 2 Carmona-Ribeiro, A. M., Yoshida, L. S., Sesso, A., Chaimovich, H. **J. Colloid Interface Sci.** **100** 433 (1984).
- 3 Carmona-Ribeiro, A. M., Yoshida, L. S., Chaimovich, H. **J. Phys. Chem.** **89** 2328 (1985).
- 4 Carmona-Ribeiro, A. M., Chaimovich, H. **Biophys. J.** **50** 621 (1986).
- 5 Carmona-Ribeiro, A. M. **Chem. Soc. Reviews** **21**(4), in press (1992).
- 6 Fendler, J. H. **Acc. Chem. Res.** **13** 7 (1980).
- 7 Bratt, P., Kang, Y.S., Kevan, L., Nakamura, H., Matsuo, T. **J. Phys. Chem.** **95** 6399 (1991).
- 8 Kotchevar, A. T., Kreevoy, M. M. **J. Phys. Chem.** **95** 10345 (1991).
- 9 Carmona-Ribeiro, A. M. **J. Phys. Chem.** **93** 2630 (1989).
- 10 Carmona-Ribeiro, A. M., Midmore, B. R. **J. Phys. Chem.** **96**, in press (1992).
- 11 Carmona-Ribeiro, A. M., Midmore, B. R. **Langmuir**, in press (1992).
- 12 Pashley, R. M., McGuiggan, P. M., Ninham, B. W., Brady, J., Evans, D. F. **J. Phys. Chem.** **90** 1637 (1986).

- 13 Marra, J. J. **Phys. Chem.** **90** 2145 (1986).
- 14 Parsegian, V. A., Rand, R. P., Fuller, N. L. **J. Phys. Chem.** **95** 4777 (1991).
- 15 O'Brien, R., White, L.R. **J. Chem. Soc. Faraday Trans. 2** **74** 1674 (1978).
- 16 Rouser, G., Fleischer, S., Yamamoto, A. **Lipids** **5** 494 (1970).
- 17 Stelmo, M., Chaimovich, H., Cuccovia, I. M. **J. Colloid Interface Sci.** **117** 200 (1987).
- 18 Israelachvili, J. N. **Chem. Scr.** **25** 7 (1985).
- 19 Claesson, P. M., Carmona-Ribeiro, A. M., Kurihara, K. **J. Phys. Chem.** **93** 917 (1989).
- 20 Israelachvili, J. N. **J. Phys. Chem.** **96** 520 (1992).
- 21 Bonosi, F., Gabrielli, G. **Colloids & Surfaces** **52** 277 (1991).
- 22 Pashley, R. M., McGuiggan, P. M., Ninham, B. W. Evans, D. F. **Science** **229** 1088 (1985).
- 23 Rand, R. P. **Annu. Rev. Biophys. Bioeng.** **10** 277 (1981).
- 24 Hunter, R. J. "Foundations of Colloid Science", Clarendon Press, Oxford (1989).
- 25 Cevc, G., Marsh, D. "Phospholipid Bilayers- Physical Principles and Models", Wiley-Interscience, New York (1987).
- 26 Verwey, E. J. W., Overbeek, J. Th. G. "Theory of the Stability of Lyophobic Colloids", Elsevier, Amsterdam (1948).
- 27 Lukac, S. J. **Phys. Chem.** **87** 5045 (1983).
- 28 Carmona-Ribeiro, A. M., Castuma, C. E., Sesso, A., Schreier, S. J. **Phys. Chem.** **95** 5362 (1991).
- 29 Carmona-Ribeiro, A. M., Hix, S. J. **Phys. Chem.** **95** 1812 (1991).
- 30 Attard, P., Mitchell, J., Ninham, B. W. **J. Chem. Phys.** **89** 4358 (1980).
- 31 Claesson, P. M., Herder, P.C., Blom, C. E., Ninham, B.W. **J. Colloid Interface Sci.** **118** 68 (1987).

Table 1 The effect of NaCl concentration (C_e) on zeta-potentials (zeta) and charge densities (sigma) of large DODAC or DHP vesicles. Due to the experimental approach, there is no variation of vesicle size as a function of C_e . DODAC and DHP vesicles are prepared in 0.43 and 0.55 M D-glucose solutions, respectively. In order to vary NaCl concentration, isoosmolar composite solutions of NaCl and D-glucose were added to the vesicle dispersions. Large DHP vesicles are at pH 5- 5.5. D_z is 235 and 344 nm for large DODAC and DHP vesicles, respectively. The polydispersity index is 0.316 and 0.402 , respectively. EM_0 is the electrophoretic mobility extrapolated to zero amphiphile concentration. Mobilities are corrected considering the viscosity of the composite solution relative to the viscosity of water (η / η_0) ie, $EM_{oc} = (\eta / \eta_0)EM_0$. K is the inverse of the Debye length, in cm^{-1} . Sigma is calculated from zeta-potentials using nonlinear Poisson-Boltzmann (10).

Amphiphile	C_G	η / η_0	C_e	$KD_z/2$	EM_0	EM_{oc}	E	Zeta	Sigma
	(M)		(mM)		$(10^4 cm^2 V^{-1} s^{-1})$			(mV)	$(10^3 Cm^{-2})$
DODAC	0.430	1.23	0.09	4	3.0	3.7	2.8	85	3.2
	0.425	1.23	2.70	23	3.0	3.7	2.8	62	10.0
	0.413	1.22	9.00	42	3.3	4.0	3.0	58	16.0
	0.394	1.21	18.00	54	3.3	4.0	3.0	55	21.0
	0.379	1.20	27.00	73	3.1	3.7	2.8	49	22.0
	0.346	1.18	45.00	94	3.0	3.5	2.6	46	26.0
DHP	0.550	1.33	0.09	6	2.8	3.7	2.8	-78	2.7
	0.548	1.31	0.90	17	3.2	4.2	3.1	-69	6.6
	0.534	1.28	9.00	55	3.3	4.2	3.2	-59	16.0
	0.501	1.20	27.00	96	3.6	4.3	3.2	-57	26.0
	0.464	1.11	45.90	125	3.8	4.2	3.2	-54	32.0
	0.436	1.04	62.10	145	3.9	4.1	3.1	-51	35.0

Table 2 Theoretical (W) and experimental (W_e) stabilities of large DODAC and DHP vesicles as a function of salt concentration (C_e). W_e values (9) were calculated from initial flocculation rates for NaCl-induced vesicle flocculation (3). W values were calculated from the zeta-potentials (Fig.1), the vesicle size ($D_z/2$), the Hamaker constant for two hydrocarbon layers interacting across water (5×10^{-14} ergs), the bilayer thickness (δ) and the NaCl concentration (C_e) using the DLVO model adapted for vesicles (9). The maximum energy in the interaction energy profile as a function of the separation distance between vesicles is $V_{\max}$. TH means "too high".

Amphiphile	Delta (Å)	$D_z/2$ (cm)	C_e (M)	$\log C_e$	Zeta (mV)	$V_{\max}/kT$	W_e	W
DODAC	50	123×10^{-7}	0.040	-1.4	48	179	$10^{2.8}$	TH
			0.063	-1.2	45	145	$10^{2.2}$	TH
			0.100	-1.0	42	116	$10^{1.5}$	10^{47}
			0.159	-0.8	40	90	$10^{1.0}$	10^{36}
DHP	46	172×10^{-7}	0.100	-1.0	-50.4	242	$10^{3.1}$	TH
			0.126	-0.9	-49.3	221	$10^{2.0}$	TH
			0.159	-0.8	-48.5	202	$10^{0.7}$	TH
			0.200	-0.7	-47.6	183	10^0	TH

FIGURE CAPTIONS

Fig. 1: The zeta-potential of large DODAC (A) and DHP (B) vesicles as a function of the logarithm of the NaCl concentration (C_e). The large DHP vesicles are at pH 5 -5.5. The insert shows the electrophoretic mobility (EM) of large DODAC vesicles as a function of DODAC concentration (C). The large DODAC vesicles prepared in D-glucose 0.43 M were added to a composite isoosmolal D-glucose/NaCl solution. For the curve in the insert, EM was measured at a final D-glucose (C_G) and NaCl concentration (C_e) of 0.346 and 0.045 M, respectively. EM at $C = 0$, ie EM_0 is obtained by extrapolation EM_0 (insert). From EM_0 corrected to the viscosity of the solution, the zeta-potential is calculated using the theory of O'Brien & White (15). The arrows indicate the points taken from the experimental curve for calculating the total energy of interaction as a function of the separation distance between interacting vesicles using the DLVO model adapted for vesicles (9).

Fig. 2: The total energy of interaction (V) as a function of the separation distance (in Å) as calculated for interacting DODAC (A) and DHP (B) vesicles with 123 and 172 mean radius, respectively. The hydrocarbon thickness was taken as 50 (DODAC) and 46 Å (DHP), the Hamaker constant, as 5.4×10^{-14} ergs in all cases and the NaCl concentration, as 0.04 (a); 0.0631 (b); 0.1 (c, e); 0.126 (f); 0.1585 (d, g); 0.200 M (h). Stern potential values were assumed to be equal to the zeta-potentials in Fig. 1. One should notice that the magnitude of the resultant repulsive energy is about two orders of magnitude above kT . Thus very high theoretical stabilities are predicted.

Referência 28

Experimental Section

Introducing Model Membranes and Lipoperoxidation

OHARA AUGUSTO and ANA MARIA CARMONA-RIBEIRO

*Departamento de Bioquímica
Instituto de Química
Universidade de São Paulo
São Paulo CP 20780
Brazil*

Introduction

Aiming at a simplification and better understanding of the complex structure-function relationships occurring in biological membranes, several model membranes have been proposed.¹ Liposomes, for example, are multilayered spheroidal structures composed of concentric and closed phospholipid bilayers separated by internal aqueous spaces.^{2*} They are readily prepared and have been used for several purposes,³ among them, the study of lipoperoxidation, a process requiring lipids with polyunsaturated hydrocarbon chains.⁴ For example, asolectin phospholipids from soybean, commonly used in reconstituted systems,³ have 55% linoleic acid chains.⁵

Lipid peroxidation and free radicals may play a role in several physiological and pathological processes such as inflammation, aging, carcinogenesis and drug toxicity.⁶ The properties and reactions of free radicals, however, are rarely dealt with in most undergraduate textbooks. The simple and inexpensive experiment described here provides an easy introduction to both biological membranes and lipid peroxidation. In addition, it demonstrates one of the possible deleterious effects of lipid peroxidation, ie, an increase in membrane permeability leading to leakage of liposomal contents. The overall strategy of the experiment is to incorporate a chromophore, methylene blue, into asolectin liposomes. The entrapment can be visualized and measured by comparing the effects of dialysis on an aqueous solution of methylene blue (MB) and on liposomes prepared in this same MB solution (LMB). After dialysis, the dialysate LMB is submitted to chemically induced lipid peroxidation along with appropriate controls. Visual examination as well as quantitative measurements provide a clear connection between lipid peroxidation and increase in MB leakage.

Up-to-date concepts such as entrapment efficiency, bilayer, model membrane, membrane functionality, oxygen toxicity, lipoperoxidation and free radical generation can be discussed and the student is motivated to consider research in rapidly developing areas such as membrane mimetic chemistry, liposomes as drug carriers, oxygen toxicity and free radical reactions.

Background

Biological membranes often contain lipids with polyunsaturated hydrocarbon chains and are bathed in an oxygen- and metal-rich fluid. Such conditions favor lipid peroxidation since metal ions help to circumvent the spin restriction of the reaction between oxygen and organic molecules. Several lines of evidence suggest that lipid peroxidation is a free-radical process involving a chain

reaction in three distinct stages: initiation, propagation and termination (Eqns 1–5):

Initiation of lipid peroxidation can be promoted by enzymes or chemical agents.⁴ Among the chemicals are reducing agents such as vitamin C (ascorbic acid) in the presence of transition metal ions such as Fe(III). Although not completely elucidated, the mechanism by which ascorbate and complexed Fe(III)⁺ initiate lipid peroxidation appears to involve generation of the extremely reactive hydroxyl radical or an equivalent oxidant species such as the perferryl ion.⁴ Both species are able to abstract a hydrogen atom from the fatty acid chain initiating the free radical chain reaction (Eqns 1–5) that leads to the uptake of oxygen, rearrangement of the double bonds in the unsaturated lipids and eventually destruction of the membrane lipids, with the production of a variety of breakdown products including aldehydes, alcohols and hydrocarbons.** Malondialdehyde, although a minor product of lipid peroxidation, is commonly used as a marker of the process due to its reactivity with thiobarbituric acid (TBA) producing a chromophore with strong absorbance at 535 nm (molar absorptivity of $1.56 \times 10^5 \text{ M}^{-1} \text{ cm}^{-1}$).⁷

Experimental Procedure

Liposomes containing methylene blue were obtained by vortexing 10 mM asolectin in an aqueous solution of 0.2 mM methylene blue. Asolectin was from Sigma Chemical Company in the form of soybean L- α -phosphatidylcholine (PC), Type IV-S, and methylene blue (MB) from Merck-Darmstadt. All other reagents were analytical grade and twice-distilled water was used throughout.

Four samples (0.5 ml) of liposomes prepared in the methylene blue solution (LMB) and a control sample (0.5 ml) of the MB solution were dialysed against 2 l water for 30 min with vigorous stirring. Before dialysis, 0.2 ml samples of LMB or MB were reserved for the determination of MB entrapment in the liposomes. After dialysis, the absorbance at 660 nm of one of the four LMB dialysates, the control sample and the reserved aliquots was determined by diluting 0.2 ml of each in a mixture of 2 ml water and 1 ml 1% Triton X-100 (see Table 1). MB entrapment (ENT) can be taken as:

$$\text{ENT} = (1/C_L)(A_2/A_1 - A_{2c}/A_{1c})$$

where A_1 and A_2 are absorbances at 660 nm of diluted LMB before and after dialysis, respectively; A_{1c} and A_{2c} are absorbances at 660 nm of diluted 0.2 mM MB solution before and after dialysis, respectively; and C_L is the molar asolectin concentration.

The other three dialysis bags containing LMB dialysates were used to test lipoperoxidation. The ascorbic acid/Fe(III)/ethylenediaminetetraacetic acid (EDTA) system was used to promote lipid peroxidation in the asolectin liposomes containing methylene blue. A first control tube (CI) was prepared by adding one of

*L: here is linoleic acid (or linoleyl) radical that has lost hydrogen at C11, the carbon between the two double bonds. LOO: here is represented mainly by equal amounts of 9- and 13-hydroperoxy radicals of linoleic acid. The major termination reaction is Eqn 5 under these conditions. Porter, N A and Wujek, D G (1984) *J Amer Chem Soc* **102**, 5597

+Oxidation of linoleate by ascorbate plus Fe(III) ions has been reported by Yamamoto, K, Takahashi, M and Niki, E (1985) *Chem Lett* 1149

The breakdown of linoleic acid hydroperoxides in the presence of ferric ions are epoxy-hydroxy linoleic acids. Gardner, H W and Kleiman, R (1981) *Biochim Biophys Acta* **665, 113

*Liposomes are classified into MLV (multilamellar vesicles), SUV (small unilamellar vesicles) and LUV (large unilamellar vesicles). In this experiment MLV are used. The entrapment discussed later is in the order of MLV. Szoka, F, and Papahadjopoulos, D (1980) *Ann Rev Biochem* **9**, 467

the three bags containing the LMB dialysate to 9.5 ml water. A *second control tube* (CII) was prepared by adding the second bag to 7.5 ml water and 2.0 ml of 10 mM ascorbic acid. The *test tube* (T) was prepared by addition of the third bag to 2 ml 10 mM ascorbic acid, 2 ml 1 mM Fe(III)/EDTA and 5.5 ml water. The tubes were then incubated (30 min, 40°C) in a reciprocating water bath. After incubation, the absorbance at 660 nm was determined for samples of the external solution in CI, CII and T (see Table 2). These samples were returned to their respective tubes and the bags disrupted inside the tubes. The bag content and the external solution were homogenized and 1 ml aliquots were taken from each tube and added to 2 ml TBA. A tube containing 1 ml water and 2 ml TBA can be prepared as the blank. The four tubes were incubated for 15 min at 100°C. After cooling, absorbances were determined at 535 nm (see Table 3).

Results and Discussion

The results obtained in a typical experiment are shown in Tables 1–3. MB is retained by the liposomes thus permitting an entrapment (ENT) calculation (Table 1). The calculated value of

Table 1 Methylene blue (MB) entrapment (ENT) in asolectin liposomes

Sample	Absorbance at 660 nm after sample dilution		ENT (l/mol)
	Before dialysis	After dialysis	
LMB	0.756	0.191	21.8
MB 0.2 mM	0.740	0.023	

approx. 22 l/mol is much higher than the value of 1.5 l/mol expected for egg PC/phosphatidic acid liposomes.² Since the membrane has about 20% cardiolipin which is negatively charged⁵ and MB is positively charged, the MB entrapment probably corresponds to liposome incorporation into the internal aqueous compartments as well as the electrostatic adsorption of MB at the liposome bilayer. To measure the entrapment efficiency of the liposome population, a marker retained solely in the internal aqueous compartment is necessary.⁸ This can be achieved by using bromocresol green (see below). Although less precise, the experiment with methylene blue is more impressive as its high ENT value permits easy visualization of liposome retention or leakage.

MB leakage from the liposomes submitted to lipoperoxidation can be easily inferred by visual inspection of the three tubes (Table 2). A bleaching of MB is observed from tube CI to T.

Table 2 Methylene blue leakage under conditions of lipoperoxidation

Tubes	Absorbance at 660 nm
Control I (water plus LMB dialysate bag)	0.070
Control II (water plus LMB dialysate bag plus ascorbic acid)	0.030
Test (water plus LMB dialysate bag plus ascorbic acid plus Fe (III)/EDTA)	0.109

Part of it is due to reduction of MB by ascorbic acid and this can be concluded from examination of tube CII. This reduction-mediated bleaching can be discussed with the students and does not invalidate the experiment. Indeed, quantitative measurement of the methylene blue outside the dialysis bags in the three tubes (Table 2) clearly demonstrates that leakage is higher in the liposomes submitted to lipid peroxidation by ascorbic acid/Fe(III)/EDTA. This is further emphasized when the extent of lipid peroxidation in the three tubes is measured by the characteristic pink colour obtained with the thiobarbituric acid method (Table 3).

Table 3 Chemically induced lipoperoxidation in asolectin liposomes

Sample	Sample volume (ml)	TBA volume* (ml)	Absorbance at 535 nm
Water	1	2	0
Control I	1	2	0.015
Control II	1	2	0.058
Test	1	2	0.166

*TBA is an aqueous solution containing 1% Triton X-100, 0.25 M HCl and 0.375% thiobarbituric acid

With slight changes, this same experiment can be performed using bromocresol green (BG). BG is a pH indicator with an apparent pK of 4.7, so that the liposomes preparation, dialysis and incubation must be done in the presence of a buffer to avoid colour changes. The pH value of choice should be at least one unit above 4.7 to guarantee that BG will be negatively charged and located in the aqueous internal compartments of the liposomes. Furthermore, incubation cannot be performed at 40°C because of the occurrence of a temperature-induced leakage of BG.² At pH 7, entrapment was 2.5–5.0 l/mol. Other results were equivalent to those obtained with MB, but visual observation of BG retention or leakage was hampered.

Acknowledgements

We thank the Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP), the Financiadora de Estudos e Projetos (FINEP) and the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) for financial support.

References

- Gomperts, B D (1977) 'The Plasma Membrane: Models for Structure and Function', Academic Press, London
- Bangham, A D, Standish, M M and Watkins, J C (1965) *J Mol Biol* **13**, 238–252
- Bangham, A D (1983) Editor, *Liposome Letters*, Academic Press, London
- Chatterjee, S N and Agarwal, S (1988) *Free Radicals Biol Med* **4**, 51–72
- Kagawa, Y and Racker, E (1966) *J Biol Chem* **241**, 2467–2474
- Halliwell, B and Gutteridge, J M C (1985) *Free Radicals in Biology & Medicine* Clarendon Press, Oxford
- Buege, J A and Aust, S D (1978) *Methods in Enzymology* **52**, 302
- Carmona-Ribeiro, A M and Chaimovich, H (1983) *Biochim Biophys Acta* **733**, 172–179

§ Peroxidation and radical reactions of linoleyl phosphatidylcholine in bilayers are different from those in a solution. Porter, N A and Lehman, L S (1982) *J Amer Chem Soc* **104**, 4731