BIBLIOTECA
Faculdade de Ciências Farmacêuticas
Universidade de São Paulo

UNIVERSIDADE DE SÃO PAULO
FACULDADE DE CIÊNCIAS FARMACÊUTICAS
Departamento de Análises Clínicas e Toxicológicas

MODIFICAÇÕES OXIDATIVAS DAS LIPOPROTEÍNAS E SUAS
IMPLICAÇÕES NA ATEROGÊNESE

DULCINEIA SAES PARRA ABDALLA

Tese apresentada ao Departamento de
Análises Clínicas e Toxicológicas da
Faculdade de Ciências Farmacêuticas da
Universidade de São Paulo para o
concurso de Livre Docência

São Paulo

1995

13969

ÍNDICE

	Página
1. INTRODUÇÃO	1
2. PROCESSOS ENVOLVIDOS NAS MODIFICAÇÕES OXIDATIVAS DAS LIPOPROTEÍNAS	
2.1 Modificações das lipoproteínas	2
2.2 Possíveis fontes geradoras de espécies oxidantes das lipoproteínas in vivo	6
3. INTERAÇÕES DAS LIPOPROTEÍNAS OXIDADAS COM DIFERENTES TIPOS CELULARES ENVOLVIDOS NA ATEROGÊNESE	
3.1 Considerações gerais sobre a formação e evolução do ateroma	8
3.2 Biodistribuição da β -VLDL oxidada na aterogênese experimental e suas interações com monócitos e macrófagos	17
3.3 Interações das lipoproteínas oxidadas com o endotélio vascular	21
4. EFEITOS DAS MODIFICAÇÕES OXIDATIVAS EM PROCESSOS DO METABOLISMO INTRAVASCULAR DAS LIPOPROTEÍNAS	
4.1 Conceitos gerais sobre o metabolismo de lipoproteínas	25
4.2 Efeitos da modificação oxidativa das lipoproteínas na esterificação do colesterol plasmático	29
4.3 Efeitos da modificação oxidativa das lipoproteínas nos processos de transferência lipídica	35
5. HIPERLIPIDEMIAS E MODIFICAÇÕES OXIDATIVAS DAS LIPOPROTEÍNAS	
5.1 Considerações gerais sobre estresse oxidativo	37
5.2 Oxidação das lipoproteínas e estresse oxidativo nas hiperlipoproteinemias	38
6. COMENTÁRIOS FINAIS	44
7. REFERÊNCIAS BIBLIOGRÁFICAS	45
RESUMO	64
ABSTRACT	65
ANEXOS	
TRABALHOS QUE DERAM ORIGEM À TESE	

"Se a Ciência é a constelação de fatos, teorias e métodos coletados na literatura, então os cientistas são os indivíduos que têm se esforçado, com sucesso ou não, para contribuir com um ou outro elemento desta particular constelação"

Thomas S. Kuhn

*"Quando o **um** é posto em confronto com todos os outros, o **um** é visto como algo que os permeia a todos, ao mesmo tempo abraçando a todos em si mesmo."*

Mahayana

*Para ver um mundo num grão de areia
E um céu numa flor silvestre,
Segure o infinito na palma de sua mão,
E a eternidade numa hora.*

William Blake

Aos meus pais, por terem me dado a chance de participar desta experiência fascinante que é a vida.

Ao José Roberto, pela compreensão, carinho e apoio em todos os momentos desta longa jornada.

Ao Rafael, minha contribuição mais significativa para o mundo, por compreender meus períodos de ausência e respeitar minha opção de vida.

MEUS MAIS PROFUNDOS E SINCEROS AGRADECIMENTOS

AGRADECIMENTOS

Aos estudantes de Pós-Graduação Décio S. Barbosa, Diana M. Fries, Edson L. Silva, Fernanda B. Araujo, Jefferson Miara, Patricia Moriel e Yi H. Chang pela dedicação ao trabalho de bancada, sem o qual não teria sido possível a realização desta tese.

À amiga Ana Campa pelo apôio irrestrito e incondicional e constante incentivo. Aos amigos Regina e Jair pela amizade sincera.

Aos amigos Hugo P. Monteiro, Rui Curi e Raul C. Maranhão pela colaboração na nossa carreira científica.

Aos estudantes de Pós-Graduação Aricio, Julisa, Najila e Ricardo, por terem optado pela nossa orientação e aos estudantes Adriana, Alexandra, Ana Paula, Daniela, Edimar, Flavia, Maria Isabel e Patricia por sua importante colaboração.

Aos colegas do Departamento de Análises Clínicas e Toxicológicas e de outros Departamentos da Faculdade de Ciências Farmacêuticas da USP, pelo respeito e apôio.

A todos que, direta ou indiretamente, contribuíram para o nosso desenvolvimento científico e humano.

1. INTRODUÇÃO

Nos últimos vinte anos um número crescente de evidências experimentais têm indicado que a modificação oxidativa da lipoproteína de baixa densidade e de outras lipoproteínas é uma etapa importante e, possivelmente, obrigatória na patogênese da aterosclerose. Considerando essa hipótese, torna-se evidente que novos parâmetros devam ser analisados para avaliar-se o risco do desenvolvimento da aterosclerose; ou seja, além de considerar-se os níveis dos lípides plasmáticos é importante também que o grau de oxidação desses lípides seja avaliado. As diferenças individuais ou populacionais na evolução da aterosclerose podem estar relacionadas às defesas antioxidantes do organismo, que atuam impedindo as modificações oxidativas das lipoproteínas e, conseqüentemente, a aterogênese. Isso teria implicações para a prevenção da doença, assim como para o seu tratamento. Alguns estudos epidemiológicos já revelaram que existe uma correlação inversa entre os níveis plasmáticos, ou a ingestão, de alguns antioxidantes e a incidência de doenças coronarianas e enfarte do miocárdio. Estudos epidemiológicos prospectivos ainda deverão ser realizados para se investigar a relação entre o grau de oxidação das lipoproteínas *in vivo* e a evolução da aterosclerose, para que estratégias de intervenção sejam planejadas e aplicadas nas diferentes populações. Em paralelo a esses estudos, necessitamos entender quais seriam as implicações das modificações oxidativas das lipoproteínas em relação às diferentes vias metabólicas dessas partículas, tanto a nível intracelular como no compartimento plasmático.

Nos últimos cinco anos temos investigado alguns aspectos da modificação oxidativa das lipoproteínas relevantes para a aterogênese. Nesta tese abordaremos os trabalhos desenvolvidos nessa linha de investigação, os quais consideram os seguintes pontos:

1. Quais seriam os processos geradores de espécies oxidantes, radicalares ou não, que participariam na modificação oxidativa das lipoproteínas;
2. De que forma as lipoproteínas modificadas poderiam interagir com os diferentes tipos celulares envolvidos na aterogênese;
3. Quais seriam as implicações da modificação das lipoproteínas no metabolismo intravascular dessas partículas;
4. Qual a influência das hiperlipoproteinemias na modificação das lipoproteínas e nos sistemas de defesa antioxidante do organismo.

2. PROCESSOS ENVOLVIDOS NAS MODIFICAÇÕES OXIDATIVAS DAS LIPOPROTEÍNAS

2.1 MODIFICAÇÕES DAS LIPOPROTEÍNAS

Alterações estruturais das lipoproteínas, bem como alterações que resultem na diminuição da afinidade de ligação ou produção dos seus receptores celulares, podem ser importantes causas do processo aterosclerótico (Rapacz et al., 1986; Rudel et al., 1986). O conceito de que modificações pós-secretoras das lipoproteínas seriam importantes no processo da aterogênese foi formulado por Brown, Goldstein e col. (1979) (Brown & Goldstein, 1983) reunindo duas importantes evidências. A primeira é que pacientes com hipercolesterolemia familiar e animais deficientes em receptores funcionais da **lipoproteína de baixa densidade (LDL)**, denominados receptores B/E, apresentam lesões ateromatosas contendo **células espumosas (foam cells)** derivadas de macrófagos. A segunda evidência é que macrófagos expostos a altas concentrações de LDL nativa não se transformam em células espumosas, devido ao processo de retroalimentação negativa exercido pelo colesterol intracelular (Brown & Goldstein, 1983). Uma vez que a LDL nativa não é capaz de transformar macrófagos em células espumosas, os autores postularam a hipótese de que a LDL da circulação sanguínea deva sofrer alguma espécie de modificação pós-secretora que a tornaria mais aterogênica e passível de ser reconhecida por receptores diferentes do receptor B/E. Esse outro receptor, denominado de receptor *scavenger*, presente nos macrófagos reconhece e internaliza partículas de LDL acetiladas por um processo rápido, saturável e não controlado por esteróis; ou seja, o receptor *scavenger* não é hiporregulado pelo colesterol intracelular, como ocorre com o receptor B/E, o que permite o acúmulo intracelular de colesterol e a formação das células espumosas. Demonstrou-se que o fator determinante para o reconhecimento da LDL acetilada pelo receptor *scavenger* é o aumento das cargas negativas da apolipoproteína dessa partícula (apo B₁₀₀) (Brown, Goldstein e col. 1979). No entanto, para que a LDL acetilada seja reconhecida por esse receptor é necessário que 80% a 90% dos resíduos lisina da apo B₁₀₀ sejam acetilados, representando uma acetilação em cerca de 300 resíduos de lisina/molécula (Brown, Goldstein e col. 1979). Pelo fato desse alto grau de acetilação não ocorrer *in vivo* outras

modificações da LDL foram investigadas. Fogelman e col. (1980) demonstraram que a formação de bases de Schiff envolvendo os resíduos lisina da apo B₁₀₀ e malonaldeído (MDA, um dos produtos finais da lipoperoxidação) resulta na formação de uma partícula de LDL mais negativa e que é internalizada e degradada por macrófagos originando células espumosas. Em contraste com a acetilação, a derivatização de apenas 20% dos grupos ϵ -amino dos resíduos lisina da apo B₁₀₀ com MDA já permite o reconhecimento da MDA-LDL pelos receptores *scavenger* dos macrófagos (Haberland et al., 1984). A oxidação das lipoproteínas pode ocorrer em graus variáveis, originando diferentes partículas modificadas oxidativamente (Figura 1) e resulta em modificações de diversas características físico-químicas dessas partículas (Jurgens et al., 1987) (Quadro 1). Dentre essas ressalta-se o aumento das suas cargas negativas, devido à derivatização dos resíduos lisina da apo B₁₀₀ por aldeídos formados durante a peroxidação lipídica, o que facilita o reconhecimento das lipoproteínas pelo receptor *scavenger* e a formação de lisofosfatidilcolina, que atua como agente quimiotático para monócitos. A liberação de citocinas por macrófagos (Akeson et al., 1991), monócitos (Thomas et al., 1994; Terkeltaub, 1994) e linfócitos (Frostegård et al., 1992), assim como de fatores de crescimento pelas células musculares lisas (Stiko-Rham et al., 1992) e células mesangiais (Tan et al., 1994) também é estimulada pela LDL oxidada. Além disso, a LDL oxidada estimula a diferenciação de monócitos a macrófagos residentes (Frostegård et al., 1990), o que impediria que os monócitos retornassem do espaço subendotelial para os vasos sanguíneos. De um modo geral, a LDL oxidada apresenta propriedades pró-inflamatórias e citotóxicas (Steinberg e col., 1989; Ross, 1993; Regnström & Nilsson, 1994; O'Brien & Chait, 1994). As modificações estruturais da LDL oxidada alteram sua imunorreatividade e imunogenicidade (Bon et al., 1983; Virella et al., 1993), induzindo uma resposta autoimune com a formação de auto-anticorpos. Diversos trabalhos demonstram a presença de anticorpos, que reconhecem epítomos presentes na apo B₁₀₀ oxidada, no plasma de indivíduos com aterosclerose (Paruns & Mitchinson, 1987; Salonen et al., 1992; Virella et al., 1993). A LDL pode ser modificada oxidativamente por células endoteliais, células de músculo liso, macrófagos, plaquetas e outros tipos celulares (Parthasarathy, 1994; Malle & Sattler, 1994).

Nos últimos 10 anos, a intensa pesquisa na área de aterogênese tem demonstrando a importância das modificações oxidativas da LDL no processo aterosclerótico e a presença de

Figura 1: Partículas representativas da LDL oxidada. 1: LDL nativa, 2: LDL contendo hidroperóxidos lipídicos e apolipoproteína B intacta, 3: LDL minimamente oxidada (MM-LDL) contendo hidroperóxidos lipídicos, aminoácidos modificados e apolipoproteína B intacta, 4: LDL contendo hidroperóxidos lipídicos e hidroxi-derivados, aminoácidos modificados e apolipoproteína B degradada. CE = ésteres de colesterol, CE-OOH = hidroperóxidos de ésteres de colesterol, PL = fosfolípides, PL-CHO, CI-CHO = produtos de oxidação (hidroxi-derivados) de fosfolípides e ésteres de colesterol, LPC = lisofosfatidilcolina, Lys-X = lisinas modificadas (Parthasarathy, 1994).

Quadro 1: Alterações físico-químicas causadas pela oxidação da lipoproteína de baixa densidade (Parthasarathy, 1994)

Propriedade	Causa (s)	Consequência (s)
Aumento da carga negativa	Modificação dos resíduos lisina pela formação de bases de Schiff Oxidação direta de aminoácidos	Captação por macrófagos
Diminuição da captação pelo receptor da LDL	Modificação dos resíduos lisina	Desconhecidas
Aumento da captação pelo receptor <i>scavenger</i>	Modificações de lisinas e outros aminoácidos, causando aumento da carga negativa da partícula	Aumento do colesterol celular Alterações no metabolismo celular
Diminuição do conteúdo de antioxidantes	Oxidação	Aumento da oxidação da lipoproteína
Diminuição dos ácidos graxos polinsaturados (PUFAs)	Oxidação	Perda da biodisponibilidade dos PUFAs
Aumento dos produtos de oxidação do colesterol	Oxidação	Citotoxicidade Metabolismo alterado do colesterol
Aumento de lisofosfatidilcolina e produtos derivados da oxidação dos fosfolípides	Ativação da fosfolipase A ₂ Clivagem de ácidos graxos oxidados presentes nos fosfolípides	Quimiotaxia de monócitos e células T; inibição da síntese de NO; indução de moléculas de adesão
Aumentos de produtos derivados da oxidação dos ácidos graxos, incluindo aldeídos	Oxidação	Indução da síntese e secreção de interleucina-1; quimiotaxia e proliferação de células de músculo liso
Fragmentação da apo B	Oxidação de resíduos de aminoácidos	<i>Crosslinking</i> dos fragmentos; alteração da imunorreatividade
Perda de lisina, histidina e prolina	Formação de bases de Schiff; oxidação direta	Aumento da carga negativa; alteração da imunorreatividade
Aumento da fluorescência da apoproteína	Modificação dos resíduos lisina	Captação por macrófagos; alteração da imunorreatividade
Formação de material semelhante ao ceróide	Complexos lípide-proteína oxidados	Acúmulo de ceróide nas células
Alteração de propriedades imunológicas	Alterações da apolipoproteína, incluindo formação de adutos lisina	Formação de imuno complexos, levando ao aumento da captação por macrófagos

partículas de LDL oxidada *in vivo* (Palinski et al., 1989; Boyd et al., 1989; Yla-Herttuala et al., 1989; Hodis et al., 1994). No entanto, uma questão extremamente relevante ainda não esclarecida é quais seriam os processos que induziriam a oxidação das lipoproteínas *in vivo*.

2.2 POSSÍVEIS FONTES GERADORAS DE ESPÉCIES OXIDANTES DAS LIPOPROTEÍNAS *IN VIVO*

As células poderiam participar da oxidação das lipoproteínas pelos seguintes mecanismos: 1) geração de espécies reativas de oxigênio ou outros oxidantes, 2) participação indireta através do fornecimento de substratos que geram oxidantes extracelulares, 3) participação na propagação da peroxidação lipídica no meio e 4) depleção de antioxidantes da lipoproteína (Parthasarathy, 1994). Varias espécies radicalares e não radicalares podem estar envolvidas na oxidação das lipoproteínas, tais como: ânion radical superóxido (O_2^-) (Steinbrecher, 1988; Heinecke et al., 1986; Hiramatsu et al., 1987; Cathcart et al., 1989), peróxido de hidrogênio (Montgomery et al., 1986; Wieland et al., 1993), espécies ferril (Rice-Evans et al., 1993), hipoclorito (O'Connell et al., 1993; Hazell et al., 1994), radical tirosil (Savenkova et al., 1994), hidroperóxidos lipídicos (Parthasarathy et al., 1989; Rankin et al., 1991), tiol/radical tiil (17; Heinecke et al., 1993; Sparrow & Olszewski, 1993) e peroxinitrito (Darley Usmar et al., 1992; Graham et al., 1993; Freeman et al., 1994). O tabagismo, que é um importante fator de risco para a aterosclerose, também pode promover a oxidação da LDL (Scheffler et al., 1990) através dos radicais livres formados pela combustão do cigarro (radicais alquil, alcóxil e peróxil na fase gasosa e radicais semiquinona, hidroxila e superóxido na fase particulada) (Evans & Pryor, 1994).

A oxidação da LDL por culturas de células provenientes da musculatura lisa de artérias humanas, veias e capilares, ou células mononucleares isoladas do sangue de indivíduos normais, é diretamente proporcional à produção do ânion radical superóxido e inibida pela adição da enzima superóxido dismutase (SOD; E.C. 1.15.1.1.) (Heinecke et al., 1986; Hiramatsu et al., 1987; Heinecke et al., 1987; Hinsbergh et al., 1986). No entanto, essa oxidação pelo ânion radical superóxido é dependente da presença de metais de transição, tais como cobre e ferro (Heinecke et al., 1984). Os metais de transição, em especial o ferro, têm sido implicados na geração de radicais livres *in vitro* e *in vivo* (Aust et al., 1985; Halliwell & Gutteridge, 1985). A concentração desses

metais na forma livre é extremamente baixa no compartimento plasmático pelo fato de estarem complexados às suas proteínas de transporte, as quais impedem que os mesmos participem de ciclos redox em condições fisiológicas (Halliwell & Gutteridge, 1985). Intracelularmente, o ferro está armazenado na forma de Fe^{3+} na ferritina, que é uma proteína multimérica (24 subunidades) de alto peso molecular (450 kDa) com grande capacidade de armazenamento de ferro (4.500 mol de ferro/mol de ferritina) (Munro HN, 1978). Vários sistemas geradores de espécies radicalares podem liberar ferro da ferritina, através da redução de Fe^{3+} a Fe^{2+} , sugerindo que essa proteína possa ser uma fonte de ferro para promover peroxidação lipídica e lesão celular (Biemond et al, 1988; Winterbourn et al, 1991). O ânion radical superóxido também pode liberar ferro da ferritina em pH fisiológico (Biemond et al, 1986; Bolann & Ulvik, 1987), gerando espécies oxidantes com capacidade de iniciar a peroxidação lipídica (Monteiro & Winterbourn, 1988; Winterbourn et al., 1990). Os neutrófilos constituem uma fonte potencial para a produção de radical superóxido *in vivo*. Esses leucócitos contêm o complexo enzimático da NADPH oxidase, que pode ser ativado por estímulos solúveis e particulados, gerando grandes quantidades de radical superóxido no processo conhecido como *burst oxidativo* (Jesaitis et al, 1991).

Neutrófilos ativados pelo acetato de forbol miristato (PMA) e o peptídeo formilado formil-metionil-leucil-fenilalanina (FMLP), promovem uma extensa oxidação da LDL na presença de ferritina (**TRABALHO 1**). A oxidação da LDL é inibida pela apotransferrina, desferroxamina e superóxido dismutase, indicando a dependência do ferro liberado da ferritina pelo radical superóxido produzido pelos neutrófilos estimulados. A catalase e a ceruloplasmina não inibem a oxidação da LDL pelo sistema neutrófilos ativados/ferritina, sendo que o ácido úrico e o ácido ascórbico exercem uma proteção apenas parcial. Além disso, a ceruloplasmina estimulou a oxidação da LDL. Esse efeito pró-oxidante da ceruloplasmina em relação à LDL também foi observado recentemente em estudos *in vitro* (Ehrenwald et al., 1994). A LDL modificada por neutrófilos estimulados e ferritina é captada por macrófagos, derivados de monócitos isolados do sangue periférico de humanos, originando células espumosas (**TRABALHO 2**). Os neutrófilos são as primeiras células que aderem às superfícies de regiões da aorta mais suscetíveis ao desenvolvimento do ateroma (Zand et al, 1988). Uma observação importante é que os sítios vasculares que apresentam ateromas contêm maior quantidade de ferritina, comparados a regiões

normais adjacentes dos vasos (Juckett MB et al 1993). Além disso, o aumento do conteúdo de ferro das células endoteliais torna-as mais suscetíveis à citólise induzida por espécies oxidantes liberadas por neutrófilos ativados (Balla et al 1990). Segundo Balla e col. (1992) a ferritina poderia atuar como um "antioxidante" no endotélio, pelo fato da exposição prolongada das células endoteliais ao grupo heme *in vitro* aumentar o conteúdo de ferritina das células, tornando-as menos suscetíveis à citólise. No entanto, isso não seria compatível com dados epidemiológicos que indicam um aumento de cinco vezes para o risco de enfarte do miocárdio em indivíduos com níveis elevados de LDL e ferritina (Salonen et al, 1992), assim como, a associação entre a ingestão de ferro-heme na dieta e o risco aumentado de enfarte do miocárdio (Ascherio et al. 1994). Além disso, o acúmulo de ferritina na hemocromatose hereditária e nos indivíduos politransfundidos também está associado a lesões tissulares (Hershko & Weatherall, 1988). A citoproteção observada por Balla et al (1992) poderia ocorrer por mecanismos independentes da ferritina, uma vez que a regulação da IRE-BP (*iron responsive element-binding protein*) pelo ferro pode alterar a síntese de outras proteínas (Ward et al., 1994). Nesse quadro geral, a adesão dos neutrófilos a regiões do endotélio vascular ricas em ferro favoreceria a lesão das células endoteliais ricas em ferritina e a posterior liberação de parte do conteúdo de ferro da ferritina, permitindo a oxidação da LDL. As interações das lipoproteínas oxidadas com células presentes na luz dos vasos, assim como, com as células que compõem a parede arterial é de fundamental importância para o processo aterosclerótico.

3. INTERAÇÕES DAS LIPOPROTEÍNAS OXIDADAS COM OS DIFERENTES TIPOS CELULARES ENVOLVIDOS NA ATEROGÊNESE

3.1 CONSIDERAÇÕES GERAIS SOBRE A FORMAÇÃO E EVOLUÇÃO DO ATEROMA

A aterosclerose é uma doença multifatorial, sendo o principal fator na patogênese dos enfartes do miocárdio e cerebral, gangrena e perda funcional das extremidades, contribuindo para 50 % da mortalidade nos Estados Unidos, Europa e Japão. Esse processo, que em condições normais poderia ser considerado como uma resposta protetora a estímulos lesivos ao endotélio e a células musculares lisas da parede arterial, consiste na formação de lesões fibro-gordurosas e fibrosas, precedidas e acompanhadas por inflamação (Ross, 1993) (Figura 2). As lesões avançadas

Figura 2: Hipótese da resposta do endotélio à lesão na aterosclerose (Ross, 1993).

da aterosclerose (ateromas ou placas fibrosas) resultam de uma resposta inflamatória-fibroproliferativa excessiva, que pode resultar na oclusão das artérias (Ross, 1993). Estudos epidemiológicos têm identificado vários fatores de risco para a aterosclerose incluindo a hiperlipidemia (Srinivasan et al, 1976; Castelli, 1992), o tabagismo (McGill, 1988), a hipertensão (Kannel, 1975), o diabetes mellitus (Fuller et al., 1980) e a obesidade (Herbert et al., 1983). No entanto, esses fatores contribuem com menos de 50% da incidência da aterosclerose, indicando que um grande número de fatores de risco ainda não foram identificados, ou, que os mecanismos pelos quais esses fatores de risco induzem a doença ainda não são completamente compreendidos (Rengnström & Nilsson, 1994).

As lesões ateromatosas acumulam-se na íntima principalmente nas bifurcações e ramificações das artérias. A localização das placas é determinada principalmente pelo *shear stress* (força de cisalhamento) na parede arterial (Reinhart, 1994). Áreas submetidas a elevado *shear stress* (paredes internas ou divisoras do fluxo em uma bifurcação) apresentam um espessamento mínimo da íntima (Ku et al., 1985), enquanto que as paredes externas de uma bifurcação têm o menor *shear stress* (que oscila entre valores positivos e negativos) e apresentam maior espessamento da íntima, devido ao aumento da proliferação das células musculares lisas (Ku et al., 1985; Kraiss et al, 1991). O *shear stress* modula várias alterações funcionais dos vasos, podendo induzir a liberação da prostaciclina (Frangos et al., 1984) e óxido nítrico (Rubanyi et al., 1986), além de promover a formação do radical ascorbil (Laurindo et al., 1994). Alterações funcionais e morfológicas das células endoteliais precedem a formação das lesões, mas os eventos iniciais da aterogênese não levam necessariamente à denudação do endotélio e as lesões iniciais desenvolvem-se em locais com endotélio íntegro. As citocinas IL-1 (interleucina-1), TNF- α (fator de necrose tumoral- α), IFN- γ (interferon- γ) e IL-2 (interleucina-2), assim como os fatores estimuladores de colônias (CSFs) são moduladores da resposta inflamatória que ocorre quando o endotélio é exposto a um agente lesivo (Ross, 1993) (Figura 3). A adesão de monócitos às células endoteliais, mediada pela interação de moléculas de adesão presentes nos monócitos e nas células endoteliais, é um dos eventos iniciais na formação das lesões. Os linfócitos T também aderem e penetram no endotélio nas lesões iniciais. A expressão das moléculas de adesão é mediada por citocinas, tais como, a IL-1, IL-4 (interleucina-4), IFN- γ e TNF- α liberadas por monócitos/macrófagos e linfócitos T

Figura 3: Interações do endotélio com macrófagos, plaquetas e células musculares lisas na aterosclerose. PD-ECGF: fator de crescimento de células endoteliais derivados das plaquetas; EGF: fator de crescimento epidérmico; TGF- β : fator β de crescimento tumoral; VEGF: fator de crescimento endotelial vascular; bFGF: fator básico de crescimento de fibroblastos; IL-1: interleucina-1; TNF α : fator α de necrose tumoral; PGE: prostaglandina E; PDGF: fator de crescimento derivado das plaquetas; M-CSF: fator estimulador de colônias de macrófagos; GM-CSF: fator estimulador de colônias de granulócitos; MCP-1: proteína-1 quimiotática para monócitos; NO: óxido nítrico; PGI₂: prostaglandina I₂; ox LDL: LDL oxidada (Ross, 1993).

Figura 4: Fenótipos contrátil e proliferativo das células musculares lisas e suas interações com diversos tipos celulares presentes nas artérias na aterosclerose. IGF-1: fator de crescimento-1 semelhante à insulina; IFN γ : interferon γ ; HB-EGF: fator de crescimento epidérmico ligante de heparina; TXA $_2$: tromboxane A $_2$. As outras abreviações são as mesmas descritas na figura 3 (Ross, 1993).

(Regnström & Nilsson, 1994). Após a adesão à superfície endotelial, os monócitos migram para o espaço subendotelial, onde diferenciam-se em macrófagos. A transição para placas fibrosas ocorre à medida que as células musculares lisas migram da camada média para a íntima da artéria e mudam seu fenótipo de contrátil para proliferativo, produzindo grandes quantidades de colágeno, elastina, proteoglicanos e glicoproteínas (Regnström & Nilsson, 1994; O'Brien & Chait, 1994) (Figura 4). Mitógenos secretados localmente, tais como, o fator derivado de plaquetas (PDGF), fator básico de crescimento de fibroblastos (bFGF), fator de crescimento epidérmico ligante de heparina (HB-EGF), fator β de transformação de crescimento (TGF β), IL-1 e TNF- α atuam no recrutamento e na ativação da proliferação das células musculares lisas (Ross, 1993). Esses fatores de crescimento são produzidos em maior intensidade nas lesões ateroscleróticas e muitos deles também têm atividade quimiotática. Nos estágios mais avançados da aterogênese, à medida que as lesões tornam-se maiores, as placas fibrosas tornam-se vascularizadas e apresentam canais semelhantes a capilares e a vênulas. Essa vascularização pode ter um papel crucial nas fases terminais da aterosclerose, quando alterações na estrutura das placas, tais como fissuras ou ulcerações, levam à hemorragia a partir do lúmen arterial, ou desses pequenos vasos, ocasionando a formação de um trombo terminal. No processo de reparo dessas lesões, ocorre o acúmulo de diferentes constituintes da matriz celular que alteram a deformabilidade da lesão e a parede arterial. O TGF β induz a formação de tecido conectivo, induzindo a expressão dos genes da fibronectina, colágeno tipos I, III, IV e V, glicosaminoglicanos e proteoglicanos (Ross, 1993). A progressão das lesões ateroscleróticas é frequentemente complicada por alterações das placas fibrosas devido à hemorragia, calcificação, ruptura da placa, necrose e trombose mural. Espécies oxidantes e proteases liberadas pelos macrófagos poderiam contribuir para a instabilidade da placa fibrosa, alterando o tecido conectivo da superfície das mesmas (Regnström & Nilsson, 1994).

Quanto à morfologia, as lesões precoces podem ser divididas em três tipos distintos (Stary et al., 1994). As lesões tipos I e II representam pequenos depósitos lipídicos na camada íntima arterial, sendo que o tipo II inclui as estrias gordurosas (lesões precoces); a tipo III representa um estágio intermediário entre as lesões precoces e as lesões avançadas. As lesões do tipo I têm sido descritas em artérias coronárias de crianças e são constituídas por pequenos grupos isolados de macrófagos (**estrias gordurosas**), contendo um acúmulo citoplasmático de lipídeos (**células**

espumosas). As lesões do tipo II caracterizam-se pelas estrias gordurosas maiores e visíveis com corantes lipofílicos (sudanolífilia), estratificadas em camadas adjacentes e pela presença de macrófagos, mastócitos (Sary, 1990) e linfócitos T (Katsuda et al., 1992). Nessa fase aparecem algumas células espumosas derivadas de células musculares lisas e pequena quantidade de gotículas lipídicas (ricas em ésteres de colesterol) e vesículas extracelulares (ricas em colesterol livre e fosfolípidos) finamente dispersas (Katz & Small, 1980). Os lípidos presentes nas lesões tipo II consistem primariamente de ésteres de colesterol (77% do conteúdo lipídico total, sendo 35% oleato de colesterol e 26% linoleato de colesterol), colesterol e fosfolípidos (Insull et al., 1966). Nas artérias coronárias as lesões do tipo II subdividem-se em dois grupos: IIa e IIb. As lesões do subgrupo IIa são mais propensas a evoluírem para lesões avançadas, localizam-se em regiões específicas de espessamento da íntima e caracterizam-se pela presença de células musculares lisas, abundante matriz intercelular, acúmulo de lipoproteínas e de macrófagos, células espumosas localizadas mais profundamente na íntima e por gotículas e partículas gordurosas extracelulares. As lesões do grupo IIb encontram-se em regiões da íntima não espessadas, contêm poucas células musculares lisas e são resistentes à progressão, ou seja, só evoluem para lesões avançadas quando associadas a altos níveis de lipoproteínas aterogênicas (Sary et al., 1994). A evolução das lesões tipo II depende muito das forças mecânicas que atuam nas paredes dos vasos, em especial o *shear stress*, que aumenta o tempo de interação (tempo de residência) entre as lipoproteínas e a parede arterial e, conseqüentemente, a difusão transendotelial das mesmas (Glagov et al., 1988). Além disso, quando ocorrem altos níveis de lipoproteínas, como na hipercolesterolemia familiar, as lesões tipo II evoluem rapidamente para as formas avançadas, mesmo em localizações da artéria não suscetíveis à progressão. A lesão do tipo III, também conhecida como lesão de transição, intermediária ou pré-ateroma, constitui a ponte morfológica e funcional entre as lesões do tipo II e o ateroma (Sary et al., 1994). As placas fibrosas (ateromas) caracterizam-se por apresentarem uma parte central contendo lípidos extracelulares, *debris* celulares e células espumosas derivadas de macrófagos, a qual é revestida por células musculares lisas, linfócitos T e tecido fibroso extracelular. O depósito denso de gotículas de gordura extracelular contendo ésteres de colesterol (com predomínio de linoleato de colesterol) ocorre, provavelmente, devido ao acúmulo extracelular de lipoproteínas que interagem com os proteoglicanos e a elastina da matriz extracelular (Guyton &

Klemp, 1994). As células musculares lisas são predominantes nas placas fibrosas e sua proliferação é o evento determinante da extensão da placa.

Na infância (2 aos 15 anos), cerca de 99% das lesões tipo II ocorrem na artéria aorta (arco aórtico, aorta torácica descendente e aorta abdominal), sendo que na puberdade e idade adulta as lesões estendem-se por toda a extensão da aorta (Strong & McGill, 1969) e atingem as artérias coronárias (Eggen & Solberg, 1968). A partir da terceira década de vida, as lesões evoluem lentamente para lesões avançadas (ateromas). A íntima da artéria coronária descendente anterior esquerda é a região mais atingida pelas lesões do tipo II ricas em células espumosas e também o local onde as lesões avançadas tendem a ocorrer mais precocemente (Stary et al., 1994). As razões para essa cronologia de evolução ainda não estão estabelecidas.

Em 1952, Glavind et al. demonstraram pela primeira vez, um acúmulo de lípides peroxidados nas paredes de artérias ateroscleróticas e sugeriram a possibilidade de que esses compostos fossem importantes na aterogênese. Posteriormente, Harman (1957) propôs que a aterosclerose seria o resultado de três processos primários: 1) oxidação dos constituintes das lipoproteínas, 2) ancoramento desse material oxidado à parede arterial e 3) uma reação inflamatória induzida por esses produtos oxidados. Mais recentemente, Steinberg e col. (1989) propuseram uma hipótese sobre o desenvolvimento das estrias gordurosas, considerando os altos níveis de LDL que ocorrem na hipercolesterolemia e a presença de LDL oxidada na parede arterial. A participação da LDL oxidada na aterogênese poderia ocorrer pelos seguintes processos: a) recrutamento de monócitos circulantes através de fatores quimiotáticos presentes apenas na LDL oxidada e sua transformação em macrófagos residentes; b) inibição da motilidade dos macrófagos residentes da camada íntima pela LDL oxidada; c) aumento da captação da LDL oxidada pelos macrófagos residentes, levando à formação de células espumosas e d) citotoxicidade da LDL oxidada, levando à perda da integridade endotelial (Figura 5).

A presença da LDL oxidada em lesões ateroscleróticas de humanos e coelhos tem sido demonstrada, através de técnicas imunohistoquímicas (Haberland et al., 1988; Palinski et al., 1989; Boyd et al., 1989; Yla-Herttuala et al., 1989) em artérias retiradas cirurgicamente. LDL isolada de lesões fibro-gordurosas presentes nas artérias possui propriedades físicas e biológicas características da LDL oxidada *in vitro* (Palinski et al., 1989). Anticorpos monoclonais que reconhecem adutos

Figura 5: Mecanismos pelos quais a LDL oxidada poderia contribuir para a aterosclerose (Steinberg et al., 1989).

de resíduos lisina da apolipoproteína B com aldeídos derivados da peroxidação da fração lipídica da LDL, em especial malonaldeído e 4-hidroxinonenal, têm sido utilizados nesses estudos imunocitoquímicos (Palinski et al., 1990). Nas estrias gordurosas, os epítomos reconhecidos por esses anticorpos encontram-se predominantemente associados aos macrófagos, aparecendo em menor proporção nas células de músculo liso e muito pouco na matriz extracelular. Nas lesões de transição e na lesões fibrosas precoces a coloração positiva está associada fortemente aos macrófagos, porém a coloração de material particulado nas células de musculares lisas e uma coloração difusa da matriz da íntima também ocorre. Nas lesões mais avançadas, existe menor positividade para os macrófagos, extensa coloração extracelular e aumento da coloração das células de músculo liso da camada média arterial (Rosenfeld, 1990).

3.2 BIODISTRIBUIÇÃO DA β -VLDL OXIDADA NA ATEROGÊNESE EXPERIMENTAL E SUAS INTERAÇÕES COM MONÓCITOS E MACRÓFAGOS

Os coelhos são altamente suscetíveis à hipercolesterolemia induzida pela dieta rica em colesterol, desenvolvendo lesões arteroscleróticas predominantemente no arco aórtico e aorta torácica (Jokinen et al., 1985) e depósitos de colesterol em células do sistema monocítico fagocitário do fígado, baço, medula óssea e glândulas supra-renais (Pryor et al., 1961). A hipercolesterolemia induzida em coelhos pela dieta rica em colesterol resulta de um acúmulo de colesterol de origem exógena (Thompson & Zilversmit, 1983). Esses animais são incapazes de aumentar a excreção de esteróis e o fígado secreta lipoproteínas contendo grandes quantidades de ésteres de colesterol, as quais têm um tempo de residência aumentado no plasma devido ao mecanismo de retroalimentação negativa dos receptores da LDL (Shore et al., 1974; Kroon et al., 1985). A dieta rica em colesterol induz alterações marcantes das lipoproteínas plasmáticas em coelhos, havendo um aumento acentuado dos níveis de uma **lipoproteína de densidade muito baixa (VLDL)** com maior diâmetro, rica em ésteres de colesterol e apolipoproteína E. A mobilidade eletroforética desta VLDL muda da região pré-beta para a beta, o que determina sua denominação como β -VLDL. Essa partícula é a principal carreadora de colesterol nesses animais (Simionescu et al., 1986), estando envolvida diretamente no desenvolvimento da aterosclerose (Goldstein et al., 1980). Em coelhos, a fração lipoprotéica com densidade menor que 1,019 g/ml

(que contém a β -VLDL) é constituída por remanescentes de quilomícrons (Ross & ZilverSmith, 1977) e por partículas de origem hepática (Daugherty et al., 1986). A β -VLDL é captada pela interação da apo E com o receptor da VLDL, presente nas células de diversos tecidos (mais abundante no coração, músculos esqueléticos e tecido adiposo) (Takahashi et al, 1994), assim como com o receptor B/E e a proteína relacionada ao receptor da LDL (LRP) (Dergunov & Rosseneu, 1994). A associação entre β -VLDL e aterogênese também é relevante para humanos portadores de hiperlipoproteinemia tipo III, que é uma condição altamente aterogênica (Tatami et al, 1981; Mahley & Rall, 1989)

Pelo fato da β -VLDL ser considerada aterogênica *per se* são poucos os estudos que avaliam o efeito da oxidação no metabolismo dessa partícula (Parthasarathy et al., 1989). Recentemente, iniciamos um estudo de aterogênese experimental, com coelhos tratados com dieta rica em colesterol, com o objetivo de avaliarmos a oxidação da β -VLDL *in vivo* e as possíveis alterações da sua distribuição tissular e captação por células do sistema fagocítico monocitário, decorrentes da oxidação dessa partícula com cobre e peroxinitrito.

A modificação das lipoproteínas com peroxinitrito seria passível de ocorrer *in vivo* pelo fato da formação de peroxinitrito já ter sido demonstrada em células importantes para o processo da aterogênese, tais como, células endoteliais de aorta (Kooy & Royall, 1994), macrófagos e neutrófilos ativados com PMA (Ischiropoulos et al., 1992; Carreras et al., 1994). A maior formação de radical superóxido (Ohara et al., 1993) e óxido nítrico (Minor et al., 1990; Verbeuren et al., 1993) já foi demonstrada em aortas de coelhos tratados com dieta rica em colesterol. Pelo fato desses dois radicais livres reagirem entre si rapidamente formando peroxinitrito (Huie & Padmaja, 1993), é altamente provável que a sua geração também ocorra em maior intensidade nas artérias dos coelhos hipercolesterolêmicos. O peroxinitrito e sua forma protonada (ácido peroxinitroso, ONOOH) são potentes oxidantes que podem oxidar lipoproteínas (Graham et al., 1993; Freeman et al., 1994) e lípidos de membranas (Radi et al.; 1991), *in vitro*. A interação do peroxinitrito com proteínas resulta na formação de nitrotirosinas devido à adição de um grupo nitro (NO_2) na posição orto da tirosina (Beckman et al., 1992). Essa reação ocorre espontaneamente, podendo ser catalisada por metais de transição e superóxido dismutase (Ischiropoulos et al., 1992a; Beckman et al., 1992). Nitrotirosina e seus metabólitos (ácidos 3-nitro-4-hidroxifenilacético e 3-nitro-4-

hidroxifenilpropionico) já foram detectados na urina de humanos (Oshima et al., 1990) e a presença de nitrotirosina já foi demonstrada no endotélio, células espumosas (macrófagos) e células inflamatórias associadas a lesões ateroscleróticas de artérias coronárias humanas (Beckman et al., 1994).

A participação do peroxinitrito na patogênese da aterosclerose seria importante tanto para a modificação das lipoproteínas, quanto para as alterações do relaxamento dos vasos sanguíneos dependente do endotélio. A administração de superóxido dismutase encapsulada em lipossomos (White et al., 1994), ou derivatizada com polietilenoglicol (Mügge et al., 1991), a coelhos hipercolesterolêmicos melhora o relaxamento vascular dependente do endotélio, indicando que o comprometimento do vasorrelaxamento na hipercolesterolemia poderia ocorrer devido à inativação do óxido nítrico pela reação com o ânion radical superóxido, formando o ânion peroxinitrito (White et al., 1994). Essa hipótese seria possível uma vez que ocorre um aumento da produção de ânion radical superóxido (Ohara et al., 1993) em coelhos hipercolesterolêmicos. Além do efeito no relaxamento vascular, a perda do efeito inibitório do óxido nítrico sobre o crescimento celular poderia resultar, indiretamente, no aumento da proliferação das células musculares lisas, o que contribuiria para a evolução da lesão ateromatosa.

Em coelhos normocolesterolêmicos e hipercolesterolêmicos a β -VLDL modificada com peroxinitrito, *in vitro*, apresenta menor tempo de permanência no plasma ($T_{1/2}$) e maior taxa fracional de depuração, comparada às partículas nativa e oxidada com cobre (**TRABALHO 3**). Essa remoção plasmática seria compatível com uma captação pelo receptor *scavenger* ou pela proteína relacionada ao receptor da LDL, que não são regulados por retroalimentação negativa e apresentam maior afinidade pelas lipoproteínas oxidadas que o receptor B/E. No entanto, a rápida remoção da β -VLDL modificada *ex-vivo* por peroxinitrito e injetada nos coelhos também poderia indicar uma proteólise mais intensa dessa lipoproteína, o que foi sugerido pelo acúmulo de radioatividade na bexiga dos animais que receberam essa partícula marcada com ^{99m}Tc (**TRABALHO 3**). O perfil de remoção plasmática das β -VLDL nos coelhos hipercolesterolêmicos mostra um $T_{1/2}$ maior para todas as lipoproteínas, tanto nativa quanto oxidada, em comparação aos animais controles. Isto provavelmente reflete uma inibição competitiva entre as lipoproteínas oxidadas *ex vivo*, injetadas endovenosamente nos coelhos, e o

grande número de partículas lipoprotéicas presentes no plasma dos animais hipercolesterolêmicos. Neste sentido, antes do início da dieta aterogênica níveis indetectáveis ou muito baixos de hidroperóxidos de ésteres de colesterol e de triglicérides, respectivamente, são encontrados na VLDL isolada dos coelhos, enquanto que esses níveis aumentam nas β -VLDL isoladas 60 dias após o início da dieta rica em colesterol (**TRABALHO 4**). Esses resultados indicam uma formação crescente de hidroperóxidos lipídicos na principal lipoproteína aterogênica dos coelhos, em paralelo ao desenvolvimento da aterosclerose induzida pela dieta rica em colesterol.

Nos animais hipercolesterolêmicos, a β -VLDL oxidada com cobre apresenta a menor remoção plasmática, comparada com a β -VLDL nativa e modificada com peroxinitrito (**TRABALHO 3**). Este dado é totalmente inédito e contrasta com dados anteriormente publicados enfatizando que as lipoproteínas oxidadas seriam rapidamente eliminadas do compartimento intravascular (Mahley et al., 1979; Asai et al., 1991). Conseqüentemente, na vigência de hipercolesterolemia um maior número de partículas lipoprotéicas oxidadas estariam presentes na circulação, o que aumentaria a probabilidade destas partículas interagirem com células endoteliais e macrófagos. A distribuição tecidual da β -VLDL marcada com ^{99m}Tc demonstra que apenas nos coelhos normocolesterolêmicos as partículas oxidadas com cobre são mais captadas pelo fígado, embora todas as lipoproteínas apresentem menores níveis de captação hepática nos coelhos hipercolesterolêmicos em comparação aos controles (**TRABALHO 3**). No grupo hipercolesterolêmico ocorre maior captação de β -VLDL oxidada com cobre pelo coração e artéria aorta, havendo maior concentração desta lipoproteína oxidada nas placas de ateroma. No entanto, a captação da β -VLDL modificada com peroxinitrito pela aorta é similar à da β -VLDL nativa tanto nos animais hipercolesterolêmicos como nos controles.

A captação da β -VLDL nativa e oxidada por monócitos isolados do sangue periférico dos coelhos normolipidêmicos captam maior quantidade de β -VLDL modificada com peroxinitrito, em relação à lipoproteína nativa (**TRABALHO 5**). Em contraste, monócitos isolados de coelhos alimentados com dieta rica em colesterol internalizam mais β -VLDL nativa em comparação à β -VLDL modificada com peroxinitrito. Em monócitos, a internalização da β -VLDL nativa pode ocorrer tanto pelo receptor B/E como pelo receptor relacionado à LDL (LRP), através da interação com a apolipoproteína E (Dergunov & Rosseneu, 1994). Em relação à captação de lipoproteínas

oxidadas, os monócitos expressam baixas quantidades das isoformas I e II do receptor *scavenger* (Geng et al., 1994), mas também expressam a glicoproteína CD 36 que internaliza lipoproteínas oxidadas (Endeman et al., 1993). Pelo fato da β -VLDL modificada com peroxinitrito ser muito mais degradada pelos monócitos dos coelhos hipercolesterolêmicos que a sua forma nativa ou oxidada com cobre, essa menor internalização aparente pode resultar da intensa proteólise intracelular dessa partícula e a conseqüente liberação dos aminoácidos resultantes para o meio extracelular. A β -VLDL oxidada com sulfato de cobre é muito pouco captada pelos monócitos tanto dos coelhos normais como dos hipercolesterolêmicos, o que está de acordo com a baixa expressão de receptores *scavenger* nessas células (Geng et al., 1994). Pelo fato da atividade do receptor B/E estar suprimida nos animais hipercolesterolêmicos devido ao alto nível do colesterol plasmático, a internalização da β -VLDL nativa pelos monócitos desses animais deve ocorrer pelo LRP, o qual não é regulado pelo conteúdo intracelular de esteróis. Portanto, pode-se considerar que os monócitos poderiam contribuir tanto para remoção da β -VLDL nativa como da β -VLDL modificada com peroxinitrito, o que representaria um mecanismo de defesa do organismo contra o efeito aterogênico dessas partículas. No entanto, pelo fato dos monócitos do sangue periférico serem precursores dos macrófagos residentes do espaço subendotelial, essas células já enriquecidas em colesterol também favoreceriam a formação das células espumosas. Esse quadro geral sugere que a modificação da β -VLDL por espécies oxidantes, incluindo o peroxinitrito, é importante para a patogênese da aterosclerose induzida em coelhos pela dieta rica em colesterol.

3.3 INTERAÇÕES DAS LIPOPROTEÍNAS OXIDADAS COM O ENDOTÉLIO VASCULAR

O endotélio desempenha uma função central no controle da fisiologia vascular e muitos fatores liberados pelas células endoteliais apresentam efeitos parácrinos e autócrinos (Lüscher et al., 1993; Davies & Hagen, 1994). Dentre as várias funções desempenhadas pelo endotélio temos: 1) provisão de uma superfície não trombogênica (pela produção de prostaciclina, óxido nítrico e sulfato de heparan); 2) barreira de permeabilidade pela qual ocorrem trocas e transporte ativo de substâncias para a parede arterial; 3) manutenção do tônus vascular pela liberação de vasodilatadores (prostaciclina e óxido nítrico) e vasoconstritores (endotelina); 4) formação e

secreção de fatores de crescimento e citocinas; 5) manutenção de colágeno e proteoglicanos da membrana basal; 6) provisão de uma superfície não aderente para leucócitos e 7) **capacidade de modificar lipoproteínas** (Ross, 1993).

Disfunções do endotélio têm sido observadas na vigência da hipercolesterolemia em animais e humanos (Creager et al., 1990; Drexler et al., 1991; Chowienzyk, 1992). A interação entre a LDL modificada e as células do endotélio vascular pode ter um papel importante na fisiopatologia da aterosclerose. A LDL minimamente oxidada estimula a expressão de moléculas de adesão (Berliner et al., 1990), proteínas de choque térmico (Zhu et al., 1994), fatores de crescimento (Rajavashisth et al., 1990), substâncias quimiotáticas (Cushing, S.D. et al., 1990), vasoconstritoras (Boulanger et al., 1992) e pró-coagulantes (Witztum & Steinberg, 1993) nas células endoteliais.

As células endoteliais produzem uma espécie radicalar importante — o óxido nítrico ($\bullet\text{NO}$) (Rosen et al., 1984; Moncada et al., 1988), reconhecido atualmente como sendo o fator de relaxamento derivado do endotélio (EDRF) (Palmer et al., 1987). A meia-vida do óxido nítrico varia de 4 a 50 segundos (Furchgott & Vanhoutte, 1989), sendo praticamente o dobro na presença de superóxido dismutase (Rubanyi & Vanhoutte, 1986; Gryglewski et al. 1986). Esse efeito da superóxido dismutase ocorre pela remoção do ânion radical superóxido, evitando a reação deste com o óxido nítrico para formar peroxinitrito (ONOO^-) (Goren et al., 1987), ou pela oxidação do ânion nitroxila (NO^-) a óxido nítrico (Murphy & Sies, 1991).

O óxido nítrico é sintetizado por uma família de enzimas denominadas óxido nítrico sintetases (NO sintetases), que catalisam a oxidação de um nitrogênio guanidínico da L-arginina para formar óxido nítrico e citrulina (Moncada, 1991). As NO sintetases constitutivas (isoformas I e II), presentes nas células endoteliais e nas células do tecido nervoso, requerem cálcio e calmodulina como cofatores (Förstemann et al., 1991; Bredt & Snider, 1992). A NO sintetase indutível (isoforma II) não requer cálcio e calmodulina como cofatores, produz continuamente grandes quantidades de óxido nítrico e é expressa, em resposta a citocinas ou endotoxina, em macrófagos (Marletta et al., 1988; Xie et al., 1992), células endoteliais vasculares (Radomski et al., 1990), células musculares lisas vasculares (Besse & Mülsch, 1990), neutrófilos (Nathan, 1992) e miócitos cardíacos (Schulz et al., 1992). A superóxido dismutase aumenta a produção de óxido nítrico pela

Figura 6: Papel do óxido nítrico no relaxamento vascular (J. Nutr. Biochem. 3: 438, 1992).

NO sintetase *in vitro*, provavelmente acelerando a conversão de um intermediário da via L-arginina-NO a óxido nítrico (Hobbs et al., 1994). Substancias vasorrelaxantes como a acetilcolina, serotonina, ATP e bradicinina iniciam um influxo de cálcio mediado por receptor, iniciando a produção e liberação extracelular de óxido nítrico (Anderson et al., 1884; Knowles & Moncada, 1994). Após sua liberação, o óxido nítrico interage com o grupo heme da guanilato ciclase, levando à produção de GMPc e ao relaxamento das células musculares lisas vasculares (Moncada et al., 1991) (Figura 6). Além de ser um potente vasorrelaxante, o óxido nítrico inibe a agregação plaquetária (Büsse et al., 1987) e a proliferação de células do músculo liso (Nakaki et al., 1990), além de modular a proliferação de células endoteliais microvasculares (Ziche et al., 1994). A decomposição do óxido nítrico resulta na formação de nitrito e nitrato, que são frequentemente utilizados como indicadores da produção de óxido nítrico pelas células (Kaku et al., 1994; Meulemans, A. & Delsene, F., 1994). Na vigência de hipercolesterolemia ocorrem deficiências do relaxamento vascular dependente do endotélio (Verbeuren et al., 1986; Jayakody et al., 1987; Ragazzi, 1989; Casino et al., 1994). Essa diminuição da resposta do endotélio a diversas substancias vasorrelaxantes tem sido atribuída à diminuição da síntese ou liberação de óxido nítrico pelo endotélio vascular. Isso poderia ocorrer devido à deficiência de arginina (Cooke et al., 1991; Creager et al., 1992; Hamon et al., 1994), por interações da lisofosfatidilcolina, presente nas lipoproteínas oxidadas, com a NO sintetase ou com as membranas celulares (Kugiyama et al., 1990; Yokoyama et al., 1990; Yang et al., 1994) ou, ainda, pelas alterações do fluxo sanguíneo (Hutcheson et al., 1994). No entanto, na maioria desses trabalhos o óxido nítrico (ou seus derivados) não é medido diretamente, avaliando-se apenas a reatividade vascular de artérias isoladas, ou *in vivo*, em resposta a diferentes agonistas para indicar o relaxamento vascular dependente do endotélio (Kugiyama et al., 1990; Yokoyama et al., 1990; Kiyotaka et al., 1990; Cooke et al., 1991; Creager et al., 1992; Flavahan et al., 1992; Cooke et al., 1991; Hamon et al., 1994; Hutcheson et al., 1994; Casino et al., 1994; Gilligan et al., 1994). A medida do acúmulo de GMPc também é utilizada para indicar a liberação de óxido nítrico (Ezaki et al., 1994). Porém, a LDL oxidada inibe a guanilato ciclase das células musculares lisas (Schmidt et al., 1992; Ezaki et al., 1994) e, portanto, a diminuição de GMPc não seria um parâmetro adequado para indicar a liberação de óxido nítrico.

Minor et al. (1990), mediram diretamente a liberação de óxido nítrico em aortas de coelhos hipercolesterolêmicos por quimiluminescência e verificaram um aumento na liberação de óxido nítrico, apesar do relaxamento dessas artérias estar diminuído em comparação aos controles. O enriquecimento das células de músculo liso com colesterol, promovido pela interação da LDL oxidada com essas células, potencializa a produção de óxido nítrico induzida por endotoxina e TNF- α (Pomerantz et al., 1993). A LDL oxidada também aumenta a liberação de óxido nítrico em culturas de células endoteliais de aorta de coelhos (Fries et al., 1995). Os mecanismos envolvidos nessa modificação da resposta do endotélio pelas lipoproteínas ainda não está esclarecido. O influxo de cálcio intracelular nas células endoteliais, induzido pela LDL oxidada, poderia resultar na ativação da NO sintetase constitutiva das células endoteliais (Nathan & Xie, 1994,). Além disso, a LDL oxidada promove a inibição de proteína tirosina fosfatases (Fries et al. 1995), que poderia ter sido causada seja por hidroperóxidos lipídicos presentes na LDL (Heffetz & Zick, 1989), ou pelo aumento dos níveis de cálcio intracelular (Ostegaard & Trowbridge, 1991). Pelo fato da fosforilação de tirosinas estar associada ao aumento da atividade da NO sintetase (Marczin et al., 1993), a inibição das proteína tirosina fosfatases também poderia contribuir para a modulação da NO sintetase (Fries et al., 1995). Portanto, a LDL oxidada poderia interagir com as células endoteliais modulando algumas vias de transdução de sinal importantes para os processos relacionados à aterosclerose.

4. EFEITOS DAS MODIFICAÇÕES OXIDATIVAS EM PROCESSOS DO METABOLISMO INTRAVASCULAR DAS LIPOPROTEÍNAS

4.1 CONCEITOS GERAIS SOBRE O METABOLISMO DAS LIPOPROTEÍNAS

As lipoproteínas são os transportadores dos lípidos na circulação sanguínea e linfática (Reichl D, 1994). Constituem-se em um conjunto de partículas esféricas e heterogêneas em relação a diversas características físico-químicas e metabólicas (Jackson et al., 1977). São constituídas por uma parte central (*core*) contendo lípidos apolares (triglicérides e ésteres de colesterol), a qual é envolvida por uma camada superficial formada por lípidos anfífilos (colesterol não esterificado e fosfolípidos) e proteínas (apolipoproteínas). A classificação dessas partículas baseia-se nas suas densidades e migrações eletroforéticas, que dependem da composição lipídica e proteica das mesmas (Quadro 2).

Quadro 2: Classificação, propriedades e composição das lipoproteínas séricas humanas (Jackson et al., 1977)

Parâmetro	Quilomícrons	VLDL	IDL	LDL	HDL
Densidade (g/ml)	< 1,006	<1,006	1,006 - 1,019	1,019 - 1,063	1,063 - 1,210
Pêso molecular	$(0,4-30) \times 10^9$	$(5-10) \times 10^6$	$(3,9-4,8) \times 10^6$	$2,75 \times 10^6$	$(1,75-3,6) \times 10^5$
Diâmetro (ηm)	> 70,0	25,0 - 70,0	22,0 - 24,0	19,6 - 22,7	4,0 - 10,0
Mobilidade eletroforética (Acetato ou agarose)	Origin	Pré-beta	Beta larga (entre beta e pré-beta)	Beta	alfa
Composição (% por Pêso)					
Colesterol livre	2	5 - 8	8	13	6
Colesterol esterificado	5	11 - 14	22	39	13
Fosfolípidos	7	20 - 23	25	17	28
Triglicérides	84	44 - 60	30	11	3
Proteína	2	4 - 11	15	20	50
Apolipoproteínas (%)					
A-I	7,4	traços	-----	-----	67
A-II	4,2	traços	-----	-----	22
B-100	-----	36,9	50 - 70	98	-----
B-48	22,5	traços	traços	-----	-----
C-I, C-II, C-III	66	49,9	5 - 10	traços	5 - 11
E-II, E-III, E-IV	-----	13,0	10 - 20	traços	1 - 2
D	-----	-----	-----	-----	traços
Síntese	Intestino	Fígado, intestino	Intravascular	Intravascular	Intestino, fígado

VLDL= lipoproteína de densidade muito baixa; IDL= lipoproteína de densidade intermediária; LDL= lipoproteína de baixa densidade; HDL= lipoproteína de alta densidade.

Figura 7: Ciclos endógeno e exógeno do metabolismo das lipoproteínas (Stanbury et al, 1983).

O transporte dos lipídes é dividido em dois ciclos: exógeno e endógeno (Figura 7). No ciclo exógeno, os lipídes da dieta são absorvidos pelas células da mucosa intestinal e secretados na linfa como **quilomícrons**, lipoproteínas ricas em triglicérides que apresentam as apolipoproteínas B₄₈, apoA-I e A-IV. Após entrarem na circulação sanguínea via ducto torácico, os quilomícrons são transformados em **remanescentes de quilomícrons** pela ação da Lipase Lipoprotéica (LPL; E.C. 3.1.1.34). A hidrólise dos triglicérides pela LPL resulta na liberação de ácidos graxos, que são armazenados nas células dos tecidos periféricos ou transportados ao fígado ligados à albumina (Tan et al., 1977). A hidrólise posterior dos triglicérides e fosfolípidos dos remanescentes de quilomícrons, provavelmente pela lipase hepática, resulta na formação de partículas ricas em ésteres de colesterol que adquirem apo E da **lipoproteína de alta densidade (HDL)**. Esses remanescentes são então captados principalmente pelo fígado através de ligação específica com receptores da apo E (Florén & Nilsson, 1977; Dergunov & Rosseneu, 1994).

O ciclo endógeno inicia-se com a síntese da VLDL pelo fígado (Luskey & Goldstein, 1974). A VLDL também é rica em triglicérides e apresenta as apolipoproteínas B₁₀₀, C-II, C-III e E na sua superfície. Semelhante aos quilomícrons, a VLDL é hidrolisada pela lipase lipoprotéica para formar a **lipoproteína de densidade intermediária (IDL)**, que pode ser captada pelo fígado através do receptor B/E ou da proteína relacionada ao receptor da LDL (Dergunov & Rosseneu, 1994), ou transformar-se finalmente em LDL, que é uma partícula mais rica em ésteres de colesterol e apresenta apenas apo B₁₀₀ em sua superfície. A LDL tem como principal função o transporte de colesterol para os tecidos extra-hepáticos, onde é utilizado na síntese de novas membranas celulares ou na formação de hormônios esteróides pela córtex da supra-renal e gônadas. Cerca de 70% do pool corpóreo da LDL é depurado pelo fígado por endocitose via receptor B/E, sendo o restante captado pelas células dos tecidos periféricos.

A HDL nascente é sintetizada e secretada pelo fígado e intestino delgado, ou produzida extracelularmente (Johnson WJ et al, 1991), na forma de partículas discoidais contendo fosfolípidos. Seus principais componentes protéicos são apo A-I, A-II e E. Na circulação, estas partículas tornam-se enriquecidas com colesterol de outras lipoproteínas e das células dos tecidos extra-hepáticos. A HDL atua também como aceptora de fosfolípidos e colesterol livre liberados da VLDL e quilomícrons, durante a lipólise dessas lipoproteínas nos capilares dos tecidos periféricos

(Eisenberg & Levy, 1975; Redgrave & Small, 1979; Gotto, 1983) O colesterol livre na superfície da HDL é então esterificado pela ação da lecitina colesterol acil transferase (LCAT; E.C. 2.3.1.43), que utiliza apo A-I presente na HDL como cofator. Os ésteres de colesterol mais apolares migram para o interior da partícula da HDL, tornando-a progressivamente esférica (HDL₃) e com maior diâmetro (HDL₂). A HDL₂ rica em ésteres de colesterol doa parte desses lípides para as lipoproteínas contendo apo B (VLDL e LDL), recebendo triglicérides. A HDL₂ rica em triglicérides sofre a ação das lipases hepática e lipoprotéica liberando parte de seu conteúdo lipídico para o fígado e convertendo-se novamente em HDL₃. A principal função fisiológica da HDL seria sua participação nesse processo conhecido como **transporte reverso do colesterol** (Glomset, 1968), que consiste na captação do colesterol efluente das células dos tecidos periféricos e seu transporte para o fígado onde é metabolizado e excretado pelas vias biliares (Brown & Goldstein, 1983) (Figura 8).

Outra importante lipoproteína envolvida no desenvolvimento de doenças cardiovasculares é a Lp(a). A Lp(a) não é derivada do catabolismo de outra lipoproteína, sendo sintetizada no fígado (Hixson et al., 1989). Essa lipoproteína tem tamanho e densidade semelhantes à LDL, mas apresenta uma apoproteína adicional, a apo (a), com alta homologia com o plasminogênio (McLean JW et al, 1987) e que se liga à apo B₁₀₀ por uma ligação dissulfeto (Gaubatz JW et al, 1982). Alguns estudos demonstram que a Lp(a) pode ligar-se aos receptores da LDL, porém com menor eficiência (Utermann, 1989). Um possível mecanismo aterogênico da Lp(a) seria a competição com o plasminogênio por receptores específicos, prejudicando a formação de plasmina e criando um estado pró-trombótico (Scanu, 1991).

4.2 EFEITOS DA MODIFICAÇÃO OXIDATIVA DAS LIPOPROTEÍNAS NA ESTERIFICAÇÃO DO COLESTEROL PLASMÁTICO

A influência da modificação oxidativa das lipoproteínas sobre o metabolismo intravascular dessas partículas tem sido muito pouco estudada. Essas investigações são de extrema importância, pelo fato de que evidências recentes confirmam a presença de lipoproteínas oxidadas no plasma (Jeng et al., 1993; Hodis et al. 1994). Alterações de processos importantes, tais como, a

Figura 8: Metabolismo da HDL e transferência de colesterol (Transporte reverso do colesterol). CETP: proteína de transferência de ésteres de colesterol, também denominada PTL 1; CL: colesterol livre; CE: colesterol esterificado; TG: triglicérides; LH: lipase hepática; LPL: lipase lipoprotéica; LCAT: lecitina colesterol acil transferase. 1: a HDL nascente é produzida pela secreção hepática e intestinal e pela formação extracelular no fluido intersticial; 2: maturação da HDL nascente em HDL₃ e HDL₂, através da esterificação do colesterol livre pela LCAT; 3: troca de éster de colesterol da HDL com triglicérides da LDL e VLDL, resultando na formação de HDL₂ rica em TG; 4: estas últimas são substrato para as lipases hepática e/ou lipoprotéica (Johnson et al., 1991).

esterificação intravascular do colesterol e as transferências lipídicas podem ocorrer em consequência da oxidação das lipoproteínas com implicações diretas para o processo da aterogênese.

A LCAT é uma enzima central no metabolismo extracelular das lipoproteínas plasmáticas. Essa enzima é responsável pela formação do colesterol esterificado nas lipoproteínas e controla, indiretamente, os níveis de colesterol livre e esterificado em várias células e tecidos. No transporte reverso do colesterol a LCAT tem o papel crítico de reduzir as concentrações de colesterol livre na HDL, gerando um gradiente de concentração que favorece o fluxo de colesterol, a partir das células para o plasma. As consequências fisiológicas da diminuição ou ausência da atividade da LCAT estão claramente demonstradas nas deficiências hereditárias dessa enzima (Glomset et al., 1983). Nos pacientes deficientes em LCAT os níveis de colesterol livre plasmático estão elevados em paralelo à presença de altos níveis de partículas esféricas pequenas e discoidais de HDL nascente, estruturas lamelares com densidade semelhante à LDL e eritrócitos com membranas anormais contendo alto teor de colesterol. Além disso, pacientes deficientes em LCAT apresentam uma predisposição à aterosclerose precoce e ao acúmulo de colesterol livre e fosfolípidos em vários tecidos (Glomset et al., 1983). Portanto, o estudo dos fatores ou eventos que possam afetar a atividade da LCAT plasmática são de extrema relevância para o metabolismo lipídico.

A esterificação do colesterol, presente em emulsões lipídicas ricas em triglicerídeos, pela LCAT do soro humano é afetada pela oxidação dessas emulsões (**TRABALHOS 6 e 7**). Isso indica que há uma menor ação catalítica da LCAT na presença da emulsão oxidada, seja por alterações estruturais da partícula que contem o substrato, ou por uma diminuição da atividade da enzima induzida por produtos derivados da lipoperoxidação presentes na emulsão. Aldeídos de cadeia longa, como o nonenal, assim como hidroperóxidos dos ácidos graxos constituintes dos fosfolípidos, triglicérides e ésteres de colesterol, ou ainda, produtos de oxidação do colesterol, eventualmente presentes na emulsão, poderiam contribuir para a inibição da esterificação do colesterol. Alterações da conformação dos fosfolípidos da superfície da emulsão lipídica, induzidas pela oxidação da partícula (Cullis & Krijff, 1979; Misik et al., 1992), poderiam afetar a interação hidrofóbica tanto da LCAT como das apolipoproteínas ativadoras da enzima, resultando em menor formação de ésteres de colesterol. Modificações no diâmetro das partículas oxidadas pode afetar a

atividade da LCAT devido a alterações na conformação das apolipoproteínas aderidas à superfície da emulsão (Jonas, 1991). Alterações dos lipídeos presentes na região central (*core*) da partícula também poderiam afetar o potencial catalítico da enzima. Sparks & Pritchard (1989) observaram que o conteúdo de triglicérides de partículas reconstituídas de HDL correlaciona-se negativamente com a velocidade aparente máxima de reação (V_{max}) da LCAT, enquanto que a afinidade da enzima pelo substrato (K_m aparente), correlaciona-se positivamente com o conteúdo lipídico e protéico da superfície da partícula. De modo geral, os autores sugerem que a composição da superfície pode afetar a interação da LCAT com a partícula, enquanto que os lipídeos do *core* poderiam regular o potencial catalítico da enzima.

Além da alteração da ligação interfacial da enzima com a partícula do substrato, produtos derivados da peroxidação lipídica poderiam estar interagindo com resíduos de aminoácidos do sítio ativo da enzima envolvidos no seu mecanismo catalítico. Apesar das escassas informações sobre a estrutura terciária da LCAT, o sítio ativo da enzima deve conter uma cavidade hidrofóbica com tamanho suficiente para acomodar ao menos uma molécula de fosfolípide (Jonas, 1991). Sabe-se que a LCAT pode ser inibida por modificação química dos resíduos ativos de serina, cisteína e histidina (Aron et al., 1978; Francone & Fielding, 1991; Zhou & Jauhianen, 1991). Os resíduos serina e histidina estariam envolvidos na primeira etapa catalítica da LCAT (atividade semelhante à da fosfolipase A_2) e os dois resíduos cisteína (Cys₃₁ e Cys₁₈₄) participariam na segunda etapa, ou seja, na esterificação do colesterol (Dolphin & Jauhianen, 1986) (Figura 9). Dessa forma, poder-se-ia supor que o sítio ativo da LCAT eventualmente pudesse ser exposto aos hidroperóxidos e aldeídos gerados na peroxidação do substrato lipídico, o que favoreceria a modificação oxidativa dos resíduos cisteína ou a derivatização dos resíduos serina e histina (Schaich, 1980; Furukawa et al., 1985), resultando em menor atividade da enzima. Essa hipótese é reforçada por dados já descritos demonstrando a inibição da LCAT por peróxido de hidrogênio e fosfolípidos oxidados, *in vitro* (Mickel & Folds, 1970; Mickel et al., 1972). Experimentos *in vivo* realizados em indivíduos expostos à atmosfera de 100% de oxigênio (Mickel et al., 1971) mostraram inibição da LCAT, provavelmente em decorrência da geração exacerbada de espécies reativas de oxigênio nessa condição o que, potencialmente, poderia induzir a oxidação das lipoproteínas do compartimento plasmático ou afetar diretamente a enzima. Além disso, macacos submetidos à dieta rica em ácidos

graxos poliinsaturados e deficiente em vitamina E, o que favoreceria a lipoperoxidação, também apresentaram menor atividade de LCAT (Mickel et al., 1975). Além dos produtos de peroxidação oriundos dos ácidos graxos, alguns oxisteróis derivados da oxidação do colesterol também poderiam, potencialmente, inibir a LCAT. A esterificação do colesterol intracelular pela enzima acil-CoA:colesterol aciltransferase (ACAT) é inibida por oxisteróis derivados do colesterol, como o 22-hidroxicolesterol (Bates et al., 1983; Drevon et al., 1980). A inibição da esterificação intracelular do colesterol em decorrência da peroxidação lipídica tem sido descrita em células endoteliais (Jialal & Chait, 1989), células de hepatoma-G₂ (Stein et al., 1985) e em macrófagos (Zhang et al., 1990). Outro fator que poderia estar contribuindo para uma menor atividade da LCAT sobre a emulsão peroxidada é o fato dessa enzima apresentar especificidade em relação ao ácido graxo presente no substrato (Sgouta, 1972). A velocidade com a qual os ácidos graxos são transferidos pela LCAT, da posição sn-2 da molécula do fosfolípide para o colesterol apresenta a seguinte ordem: linoleato > oleato > araquidonato > ácidos graxos saturados (Sgouta, 1972). Embora não tenha sido realizado o perfil dos ácidos graxos da emulsão antes e após a oxidação, sabe-se que ocorre diminuição dos ácidos insaturados durante o processo de lipoperoxidação (Seligman et al., 1979). Assim, a menor atividade da LCAT sobre a emulsão oxidada poderia estar relacionada também ao menor teor de ácidos graxos insaturados presentes na mesma. A inibição da LCAT por lípidos oxidados pode ser relevante para a aterogênese, tanto pela menor formação de ésteres de colesterol nas partículas de HDL como pelo menor efluxo do colesterol livre, a partir das células. Portanto, a inibição da LCAT por substratos lipídicos oxidados poderia causar a diminuição do transporte reverso do colesterol e, conseqüentemente, o acúmulo de colesterol no interior das células.

Figura 9: Mecanismo de ação da lecitina colesterol acil transferase (LCAT) (Glomset, 1983).

4.3 EFEITOS DA MODIFICAÇÃO OXIDATIVA DAS LIPOPROTEÍNAS NOS PROCESSOS DE TRANSFERÊNCIA LIPÍDICA

Durante a lipólise dos quilomícrons e da VLDL ocorre a transferência de fosfolípidos dessas lipoproteínas para a HDL, mediada principalmente pela proteína de transferência de fosfolípidos (PTL II), acompanhada da transferência de colesterol livre dessas e outras lipoproteínas e também das células (Tall et al., 1987). Esses processos fornecem, portanto, os dois substratos necessários à síntese de ésteres de colesterol pela LCAT. A proteína de transferência de ésteres de colesterol (PTL I) redistribui esses ésteres de colesterol dos seus sítios de síntese (subclasses de HDL) para as lipoproteínas ricas em triglicérides e para a LDL (Tall et al., 1987). Pelo fato de existirem receptores hepáticos específicos para os remanescentes de quilomícrons e para a LDL, a redistribuição dos ésteres de colesterol pela PTL I representa uma importante etapa no transporte centrípeto de colesterol, a partir dos tecidos periféricos para o fígado (Tall et al., 1987).

Não há diferenças significantes para os processos de transferência de fosfolípidos, triglicérides e ésteres de colesterol, quando se utiliza uma emulsão oxidada rica em triglicérides como partícula doadora de lípidos (**TRABALHO 6**). Esse dado indica que a peroxidação dos lípidos constituintes da emulsão não afeta a interação hidrofóbica dessa partícula com as proteínas de transferência, ou a formação de um complexo ternário com as lipoproteínas aceptoras (LDL e HDL). Em contraste, a modificação oxidativa da HDL resulta em aumento da transferência de ésteres de colesterol da HDL oxidada para a LDL. Isso ocorre tanto com as lipoproteínas isoladas de indivíduos normolipidêmicos (**TRABALHO 7**). Marcel et al. (1989) demonstraram que a modificação oxidativa da HDL resulta em mudanças na imunorreatividade e *cross-linking* da apo A1. Uma das subfrações da HDL (com migração na região α_2) é particularmente sensível à oxidação (Kunitake et al., 1985), sendo a fração de HDL que contem a maior parte da PTL I do plasma (Marcel et al., 1990). O fato de não se observar um efeito inibitório no processo de transferência de ésteres de colesterol, decorrente da oxidação da HDL, indica que a HDL oxidada manteve sua capacidade de associação com a PTL I. Apesar das alterações observadas nas apolipoproteínas da HDL após sua oxidação, a interação interfacial entre essas e a PTL I parece ser preservada. A oxidação de resíduos metionina na face apolar da estrutura helicoidal das apo A1 e

A2 reduz a hidrofobicidade da proteína e sua afinidade pelos lipídeos (Anantharamaiah et al., 1988). A oxidação da HDL resulta também na formação de dímeros de apo A1 e apo A2 e polímeros de apo A1 (Rosenfeld et al., 1991; Zawadski et al., 1991), provavelmente pela ação de radicais hidroxila em resíduos de triptofano o que facilitaria o *cross-linking* dos monômeros dessas apolipoproteínas (Salmon et al., 1992). Além disso, no processo de peroxidação lipídica ocorre derivatização de grupos epsilon-amino dos resíduos de lisina presentes nas lipoproteínas, o que promove um aumento das cargas negativas dessas partículas (Haberland et al., 1984). Sabe-se que o aumento das cargas negativas nas lipoproteínas aumenta a associação da PTL I às mesmas (Pattnaik et al., 1979).

Alterações nos processos de transferência lipídica são relevantes para o processo da aterogênese. Duas interpretações podem ser dadas ao papel da LTP I na aterosclerose. Em primeiro lugar, a transferência de ésteres de colesterol da HDL para as lipoproteínas contendo apo B pode representar uma etapa relevante do transporte reverso do colesterol. Em favor à essa interpretação estão alguns dados mostrando que pacientes hiperlipidêmicos suscetíveis à aterosclerose apresentam valores muito baixos ou ausentes para a transferência de ésteres de colesterol no plasma (Fielding et al., 1984; Fielding, 1984). Por outro lado, a PTL I transfere ésteres de colesterol da HDL aos remanescentes de lipoproteínas que, potencialmente, promovem a deposição de ésteres de colesterol na placa ateromatosa (Patsch et al., 1983; Zilversmith, 1979). Coincidentemente espécies que apresentam altos níveis de PTL I (humanos, coelhos e macacos) são suscetíveis à aterosclerose enquanto espécies com baixos níveis (cães, porcos e ratos) são relativamente resistentes (Tall et al., 1987). Considerando-se essa segunda hipótese, poder-se-ia admitir que a inibição da PTL I resultaria em aumento marcante nos níveis de colesterol da HDL e menor formação de partículas remanescentes aterogênicas. Além disso, partículas maiores de HDL teriam menor probabilidade de promover a deposição de ésteres de colesterol nas células espumosas arteriais (Mahley & Innerarity, 1983). Sob esse enfoque, a inibição do processo de transferência de ésteres de colesterol resultaria em um perfil menos aterogênico das lipoproteínas plasmáticas enquanto que, ao contrário, condições que estimulassem esse processo poderiam ser desfavoráveis ao organismo (Tall et al., 1987).

Considerando que a oxidação de substratos lipídicos inibe a esterificação do colesterol pela LCAT e que o processo de transferência de ésteres de colesterol da HDL para a LDL pode ser estimulado por modificações oxidativas da HDL, o transporte reverso do colesterol estaria alterado nesses casos, o que contribuiria para o aumento do risco para o desenvolvimento da aterosclerose, caso esses efeitos também ocorram *in vivo*.

5. HIPERLIPIDEMIAS E MODIFICAÇÕES OXIDATIVAS DAS LIPOPROTEÍNAS

5.1 CONSIDERAÇÕES GERAIS SOBRE ESTRESSE OXIDATIVO

A relação entre estresse oxidativo e diversas doenças tem sido bastante estudada nos últimos anos (Gutteridge, 1993). O estresse oxidativo ocorre quando existe um predomínio de espécies oxidantes, radicalares e não radicalares, devido ao desequilíbrio entre a geração destas espécies e os sistemas de defesa antioxidante. A suscetibilidade de um órgão ou sistema ao estresse oxidativo é uma função do equilíbrio geral entre fatores que induzam o estresse e aqueles que apresentam capacidade antioxidante. A lesão decorrente do estresse oxidativo pode ser descrita como uma consequência da insuficiência do potencial antioxidante do organismo, ou, da exacerbação de processos que induzam a produção de espécies oxidantes.

O sistema de defesa antioxidante primário do organismo compreende substâncias antioxidantes lipo e hidrossolúveis (alfa-tocoferol, beta-caroteno, ubiquinol, glutathione, urato e ascorbato) e enzimas como a superóxido dismutase, glutathione peroxidase e catalase (Sies, 1986), que eliminam espécies reativas de oxigênio. Enzimas proteolíticas, que eliminam preferencialmente proteínas modificadas oxidativamente, sistemas de reparo do DNA e certas fosfolipases atuam como sistemas de defesa antioxidante secundários, eliminando ou reparando moléculas modificadas por espécies reativas (Sies, 1986; Davies, 1986). Além disso, proteínas quelantes de metais de transição (albumina e transferrina) presentes no plasma sanguíneo também fazem parte do sistema de defesa antioxidante, evitando que metais de transição participem na formação de espécies radicalares.

5.2 OXIDAÇÃO DAS LIPOPROTEÍNAS E ESTRESSE OXIDATIVO NAS HIPERLIPOPROTEINEMIAS

No compartimento plasmático a oxidação das lipoproteínas poderia ser iniciada, potencialmente, por espécies reativas de oxigênio ou nitrogênio geradas por monócitos, neutrófilos, plaquetas, células endoteliais e de músculo liso (White et al, 1994; Malle & Sattler, 1994; **TRABALHO 1**). Estudos recentes indicam que alterações estruturais da LDL, possivelmente decorrentes da sua oxidação, possam ocorrer na circulação sistêmica (Jeng et al., 1993; Hodis et al., 1994). As lipoproteínas dos portadores de hiperlipoproteinemias são mais suscetíveis à oxidação *ex-vivo*, quando comparadas às dos indivíduos normolipidêmicos (Lavy et al., 1991; **TRABALHOS 8 e 9**), indicando que essas lipoproteínas também poderiam ser mais oxidáveis *in vivo*, desde que a geração de espécies oxidantes suplantasse a capacidade de defesa, ou seja, na vigência de um estresse oxidativo.

As hiperlipoproteinemias constituem um conjunto de desordens de origem genética, ou adquiridas, que predispoem os indivíduos ao desenvolvimento da aterosclerose. As hiperlipoproteinemias são classificadas fenotipicamente através do aumento dos lípides e das lipoproteínas plasmáticas (Quadro 3). Essas doenças podem ser primárias, ou, secundárias a doenças metabólicas e hormonais, dieta e ao uso de álcool ou drogas. Nas hiperlipoproteinemias ocorre um aumento do número de partículas de lipoproteínas na circulação, seja pelo aumento da síntese hepática ou intestinal, ou pela diminuição do seu catabolismo. O aumento do tempo de permanência das lipoproteínas no compartimento plasmático, devido a deficiências catabólicas, poderia favorecer a oxidação dessas partículas pelos diversos mecanismos já discutidos neste trabalho. Alterações qualitativas das lipoproteínas também podem alterar a sua suscetibilidade à oxidação. Frações mais densas da LDL estão associadas a um maior risco para o desenvolvimento da aterosclerose (Austin et al., 1988; Slyper, 1994) e são mais oxidáveis *ex-vivo* que outras frações lipoprotéicas (De Graaf et al., 1991; **TRABALHO 8**). Alterações da composição lipídica e dos ácidos siálicos das subfrações densas da LDL podem, potencialmente, torná-las mais aterogênicas (Orekov et al., 1989; **TRABALHO 8**). Portanto, o monitoramento do risco para o desenvolvimento da aterosclerose deveria ser feito não apenas pela quantificação dos lípides e lipoproteínas plasmáticas, como também pela avaliação do grau de oxidação das mesmas (Witztum,

Quadro 3: Classificação das hiperlipoproteinemias (Fredrickson & Levy, 1972).

Lipoproteína aumentada	Aparência do plasma*	Colesterol total	Triglicérides	LDL-colesterol	HDL-colesterol	Apolipo-proteína	Elektroforese	Fenótipo	Associação clínica
Quilomícrons	camada cremosa superior, infranadante claro ou ligeiramente turvo	normal a moderadamente elevado	muito elevado	normal	normal a diminuído	↑ B-48 ↑ A-IV ↓↑ C-II	intensa banda na origem	Tipo I	abdomem agudo pancreatite
LDL	claro, possível aumento da coloração alaranjada	elevado	normal	elevado	normal a diminuído	↑ B-100	aumento da banda beta	Tipo IIa	maior risco de DAC
LDL; VLDL	claro a ligeiramente turvo	elevado	elevado	elevado	normal a diminuído	↑ B-100	aumento das bandas beta e pré-beta	Tipo II b; Hiperlipoproteinemia combinada	maior risco de DAC
IDL	turvo à opaco; fina camada cremosa superior	elevado	elevado	normal a diminuído	normal a diminuído	↑ E-II ↓ E-III ↓ E-IV	banda beta larga	Tipo III	maior risco de DAC
VLDL	turvo a opaco	normal a ligeiramente elevado	elevado	normal	normal a diminuído	↑↓ C-II ↑ B-100	aumento da banda pré-beta	Tipo IV	maior risco de DAC
VLDL; quilomícrons	camada superior cremosa, infranadante turvo a opaco	moderadamente elevado	muito elevado	normal	normal a diminuído	↑↓ C-II ↑ B-100 ↑ B-48	banda intensa na origem e aumento da banda pré-beta	Tipo V	pancreatite, maior risco de DAC
HDL	claro	normal a moderadamente elevado	normal	normal	elevado	↑ A-I ↑ A-II	aumento da banda alfa	Hiperalfalipoproteinemia	menor risco de DAC

1994). A detecção de lipoproteínas oxidadas no plasma de indivíduos com hiperlipidemia, ou outros fatores de risco para a aterosclerose, assim como a determinação da oxidabilidade dessas partículas, poderia ter um valor preditivo para a progressão da doença.

A oxidação das lipoproteínas plasmáticas em indivíduos hiperlipidêmicos poderia ser facilitada pela maior produção de espécies reativas de oxigênio por leucócitos estimulados (Ludwig et al., 1982; Prónai et al., 1991; Lavy et al., 1991; Krause et al., 1993). Leucócitos mononucleares de pacientes com hiperlipidemia combinada (TRABALHO 9), ou hipertrigliceridemia (Hiramatsu & Arimori, 1988), liberam maiores quantidades de espécies reativas de oxigênio, quando comparados aos indivíduos normolipidêmicos. Uma correlação positiva entre os níveis plasmáticos de triglicérides totais e a produção basal de espécies reativas de oxigênio pelos leucócitos mononucleares e neutrófilos ocorre em indivíduos normo e hiperlipidêmicos (TRABALHO 9). Além disso, os triglicérides presentes na LDL e na VLDL correlacionam-se positivamente com a liberação de espécies reativas de oxigênio pelos leucócitos mononucleares não estimulados e neutrófilos estimulados com éster de forbol (PMA). Esses dados sugerem que lipoproteínas enriquecidas em triglicérides possam desempenhar um papel crucial na ativação de leucócitos. Dentre os possíveis mecanismos envolvidos nesse efeito pode-se discutir o papel dos ácidos graxos presentes nessas lipoproteínas. Os ácidos graxos *cis*-poliinsaturados estimulam a produção do ânion radical superóxido por neutrófilos humanos na seguinte ordem : araquidonate > linolenato > linoleato > oleato (Badwey et al., 1981). Dentre os ácidos graxos saturados, aqueles com 14 a 18 átomos de carbonos, quando combinados com 1-oleoil-2-acetilglicerol, também estimulam a liberação de radical superóxido (Ozawa et al., 1989). A interação dos ácidos graxos com a membrana plasmática dos leucócitos ocorre por ligações hidrofóbicas das suas cadeias acil com sítios hidrofóbicos da membrana. Os ácidos graxos presentes nos triglicérides das lipoproteínas poderiam atuar ao nível da transdução de sinal da NADPH oxidase resultando na sua ativação e maior produção de superóxido. Uma vez que ácidos graxos peroxidados podem ativar a fosfolipase A₂ (Kuijk et al., 1987), a liberação de ácido araquidônico oxidado das membranas dos leucócitos poderia contribuir para a ativação da NADPH oxidase. Além disso, é possível que ácidos graxos peroxidados, liberados pela lipólise das lipoproteínas ricas em triglicérides mediada pela

lipoproteína lipase, também poderiam contribuir para a ativação do *burst* oxidativo dos leucócitos, *in vivo*.

Emulsões oxidadas ricas em triglicérides incubadas com monócitos do sangue periférico humano aumentam a produção de radical superóxido por essas células, provavelmente devido aos ácidos graxos peroxidados (Görög, 1991). Uma maior produção basal de espécies reativas de oxigênio é observada nos leucócitos mononucleares isolados dos portadores de hiperlipidemia combinada, o que sugere que essas células poderiam estar previamente ativadas no sangue periférico (**TRABALHO 9**). Esses indivíduos apresentam uma LDL enriquecida em triglicérides, a qual é altamente suscetível à oxidação *ex-vivo*, indicando que essa fração poderia contribuir para a estimulação do *burst* oxidativo dos leucócitos. Além disso, esses portadores de hiperlipidemia combinada apresentam altos níveis de ferritina plasmática, o que poderia contribuir para a oxidação das lipoproteínas, de acordo com o mecanismo proposto no **TRABALHO 1**. Em conjunto, esses dados indicam que altos níveis de ferritina, quando associados ao aumento da produção de espécies reativas de oxigênio pelos leucócitos, poderiam favorecer a oxidação de lipoproteínas *in vivo*, o que explicaria a associação entre altos níveis de ferritina plasmática e o risco de doença cardíaca coronariana demonstrada por dados epidemiológicos recentes (Salonen et al, 1992).

A hipertrigliceridemia atua sinergisticamente para aumentar o risco de desenvolvimento da doença arterial coronariana associada à hipercolesterolemia (Gotto, 1992) e, recentemente, as lipoproteínas ricas em triglicérides foram consideradas como um fator de risco independente para a progressão dessa doença (Hodis et al., 1994). O aumento da produção de espécies reativas de oxigênio pelos leucócitos dos portadores de hiperlipidemia combinada e sua relação positiva com os níveis de triglicérides (**TRABALHO 9**), pode contribuir para o esclarecimento dos mecanismos envolvidos na aterogenicidade das lipoproteínas ricas em triglicérides.

A HDL correlaciona-se negativamente com a incidência da aterosclerose e seu efeito protetor tem sido relacionado à sua participação no transporte reverso do colesterol (Eisenberg, 19984). No entanto alguns estudos *in vitro* demonstram um efeito antioxidante da HDL (Parthasarathy et al., 1990). Pelo fato de haver uma correlação negativa entre os níveis de colesterol da HDL e a liberação basal de espécies reativas de oxigênio pelos leucócitos mononucleares e neutrófilos (**TRABALHO 9**), a menor liberação de espécies reativas de oxigênio

pelos leucócitos promovida pela HDL também poderia estar relacionada ao efeito protetor dessa lipoproteína no desenvolvimento da aterosclerose.

Estudos epidemiológicos mostram uma relação inversa entre antioxidantes plasmáticos (tocoferol, caroteno e ascorbato) e a incidência de doenças cardiovasculares (Enstrom et al., 1992; Gey et al., 1993; Rimm et al., 1993). A demanda para esses antioxidantes pode estar consideravelmente aumentada quando a quantidade de lipídeos plasmáticos a ser protegida por eles aumenta, como acontece nas hiperlipoproteinemias. O defeito do catabolismo da LDL e de outras lipoproteínas em certas hiperlipoproteinemias resulta em um prolongamento da meia-vida dessas partículas no plasma, o que aumenta a probabilidade das mesmas interagirem com espécies oxidantes geradas por diferentes células. Os antioxidantes lipossolúveis são transportados pelas lipoproteínas e as protege contra a ação de espécies oxidantes, inibindo a oxidação das mesmas. Além disso, os antioxidantes lipofílicos também podem suprimir o *burst oxidativo* de neutrófilos (Baader et al., 1988). Portadores de hiperlipidemia combinada apresentam uma diminuição dos níveis plasmáticos dos carotenóides β -caroteno e licopeno, o que poderia contribuir para a maior suscetibilidade da LDL desses indivíduos à oxidação *ex-vivo* (TRABALHO 9). O β -caroteno é um supressor de oxigênio singlete e um sequestrador de radicais livres (Krinsky, 1992). Altas doses de β -caroteno aumentam os níveis de HDL (Ringer et al., 1991) e a suplementação com esse antioxidante reduz as complicações coronarianas e outros eventos vasculares (Gaziano et al., 1990). Considerando que a maior parte dos carotenóides é transportada pela LDL, os níveis diminuídos de β -caroteno e licopeno observados nos portadores de hiperlipidemia combinada poderiam contribuir para a diminuição da resistência da LDL desses indivíduos à oxidação. Além da diminuição dos antioxidantes lipossolúveis, a atividade da superóxido dismutase (EC-SOD) no plasma desses pacientes também diminuiu, implicando em uma menor defesa contra a ação lesiva do radical superóxido produzido pelos leucócitos e outras células do compartimento vascular, ou, espécies dele derivadas como o peroxinitrito (TRABALHO 9). Uma das isoformas da EC-SOD, a fração C, liga-se à superfície das células endoteliais dos vasos sanguíneos, através da ligação com o sulfato de heparan da matriz extracelular (Karlsson & Marklund, 1988), sendo importante para a proteção do endotélio. A ocorrência de uma correlação negativa entre a EC-SOD e os níveis plasmáticos de colesterol e triglicérides sugere que esses lipídeos poderiam afetar a ligação ou a liberação da EC-

SOD do endotélio vascular (**TRABALHO 4**). Em coelhos hipercolesterolêmicos, as atividades plasmáticas das enzimas antioxidantes superóxido dismutase e glutathione peroxidase são semelhantes àsquelas de coelhos normolipidêmicos (**TRABALHO 4**), no entanto, a fração C da superóxido dismutase extracelular (EC-SOD) está diminuída nos coelhos hipercolesterolêmicos. Essa diminuição da fração C da EC-SOD poderia facilitar a formação de peroxinitrito, a partir do óxido nítrico e do radical superóxido gerados pelo endotélio, o qual poderia interagir com as lipoproteínas e com componentes do sistema de defesa antioxidante do plasma (Augusto et al., 1994; Gatti et al., 1994; **TRABALHO 3**).

Esse quadro geral sugere que, em determinados tipos de hiperlipoproteinemias, possa ocorrer um aumento da geração de espécies oxidantes, resultante provavelmente da estimulação dos leucócitos pelos ácidos graxos oxidados presentes nos lípides das lipoproteínas, o que promoveria um maior consumo de antioxidantes ou alteração das enzimas antioxidantes, originando uma situação de estresse oxidativo.

6. COMENTÁRIOS FINAIS

As evidências reunidas em um número cada vez maior de trabalhos científicos, alguns dos quais indicados nesta tese, indicam a importância da modificação oxidativa das lipoproteínas na patogênese da aterosclerose. As lipoproteínas oxidadas poderiam atuar em diversos níveis na aterogênese, ou seja, desde a transdução de sinal celular, que resultaria na produção de mediadores importantes para a fisiologia do endotélio e para o processo aterosclerótico, até a formação das células espumosas. Indiretamente, a modificação oxidativa das lipoproteínas poderia afetar o desenvolvimento da aterosclerose, interferindo em importantes vias do metabolismo intravascular dessas partículas, tais como o transporte reverso do colesterol. Associada às hiperlipoproteinemias e a outros fatores de risco para a aterosclerose, a modificação oxidativa das lipoproteínas poderia atuar sinergisticamente para aumentar o risco, ou, acelerar o processo da aterosclerose. O direcionamento futuro das nossas investigações será para o desenvolvimento de metodologias que nos permitam esclarecer melhor esses pontos, principalmente no sentido de correlacionarmos as diferentes modificações oxidativas das lipoproteínas e o desenvolvimento da aterosclerose em humanos.

7. REFERÊNCIAS BIBLIOGRÁFICAS

- ANANTHARAMAIAH, G.M.; HUGHES, T.A.; IGBAL, M.; GAVISH, A.; NEAMER, J.N.; MEDLEY, M.F.; SEGREST, J.P. Effect of oxidation on the properties of apolipoproteins A-I and A-II. J. Lipid Res. **29**: 309-318, 1988.
- ANDERSON, T.J.; MEREDITH, I.T.; GANZ, P.; SELWYN, A.P.; YEUNG, A.C. Nitric oxide and nitrovasodilators: similarities, differences and potential interactions. J. Am. Coll. Cardiol. **24**: 555-566, 1994.
- ARON, L.; JONES, S.; FIELDING, C.J. Human plasma lecithin:cholesterol acyltransferase. J. Biol. Chem. **253**: 7220-7226, 1978.
- ASAI, K.; HAYASHI, T.; FUNAKI, C.; KUZUYA, M.; NAITO, M.; KUZUYA, F. Comparison of plasma clearance of low density lipoprotein with β -very low density lipoprotein or acetoacetylated low density lipoprotein in cholesterol-fed rabbits. Biochem. Inter. **23**: 327-334, 1991.
- ASCHERIO, A.; WILLET, W.C.; RIMM, E.B.; GIOVANNUCCI, E.L.; STAMPFER, M.J. Dietary intake and risk of coronary disease among men. Circulation **89**: 969-974, 1994.
- AUGUSTO, O.; GATTI, R.M.; RADİ, R. Spin-trapping studies of peroxynitrite decomposition and of 3-morpholinocydonimine N-ethylcarbamide autoxidation: direct evidence for metal-independent formation of free radicals intermediates. Arch. Biochem. Biophys. **310**: 118-125, 1994.
- AUST, S.D.; MOREHOUSE, I.A.; THOMAS, C.E. Role of metals in oxygen radicals reaction. J. Free Rad. Biol. Med. **1**: 3-25, 1985.
- AUSTIN, M.A.; BRESLOW, J.L.; HENNEKENS, C.H.; BURING, J.E.; WILLET, W.C.; KRAUSS, R.M. Low density lipoprotein subclass patterns and risk of myocardial infarction. JAMA **260**: 1917-1921, 1988.
- BAADER, W.J.; HATZELMANN, A.; ULLRICH, V. The suppression of granulocyte functions by lipophilic antioxidants. Biochem. Pharmacol. **37**: 1089-1095, 1988.
- BADWEY, J.A.; CURNUTTE, J.T.; KARNOVSKY, M.L. Cis-polyunsaturated fatty acids induce high levels of superoxide production by human neutrophils. J. Biol. Chem. **256**: 12640-12647, 1981.
- BALLA, G.; VERCELLOTTI, G.M.; EATON, J.W.; JACOB, H.S. Iron loading of endothelial cells augments oxidant damage. J. Lab. Clin. Med. **116**: 546-554, 1990.
- BALLA, G.; JACOB, H.S.; ROSENBERG, M.; NATH, K.; APPLE, F.; EATON, J.W.; VERCELLOTTI, G.M. Ferritin: a cytoprotective antioxidant strategem of endothelium. J. Biol. Chem. **267**: 18148-18153, 1992.

- BATES, S.R.; JETT, C.M.; MILLER, J.E. Prevention of the hyperlipidemic serum on LDL-induced cellular cholesterol ester accumulation by 22-hydroxycholesterol and its analogue. Biochim. Biophys. Acta **753**: 281-293, 1983.
- BECKMAN, J.S.; YE, Y.Z.; ANDERSON, P.G.; CHEN, J.; ACCAVITTI, M.A.; TARPEY, M.M.; WHITE, C.R. Extensive nitration of protein tyrosines in human atherosclerosis detected by immunohistochemistry. Biol. Chem. Hoppe-Seyler **375**: 81-88, 1994.
- BECKMAN, J.S.; ISCHIROPOULOS, H.; ZHU, L.; VAN DER EOERD, M.; SMITH, C.; CHEN J.; HARRISON, J.; MARTIN, J.C.; TSAI, M. Kinetics of superoxide dismutase and iron catalyzed nitration of phenolics by peroxynitrite. Arch. Biochem. Biophys. **298**: 438-444, 1992.
- BERLINER, J.A.; TERRITO, M.C.; SEVANIAN, A. et al. Minimally modified low density lipoprotein stimulates monocyte endothelial interactions. J. Clin. Invest. **85**: 1260-1268, 1990.
- BIEMOND, P.; SWAAK, A.J.G.; BEINDORF, C.M.; KOSTER, J.F. Superoxide-dependent and -independent mechanisms of iron mobilization from ferritin by xanthine oxidase. Biochem. J. **239**: 169-173, 1986.
- BIEMOND, P.; SWAAK, A.J.G.; VAN EIJK, H.G.; KOSTER, J.F. Superoxide dependent iron release from ferritin in inflammatory diseases. J. Free Rad. Biol. Med. **4**: 185-198, 1988.
- BOLANN, B.J.; ULVIK, R.J. Release of iron from ferritin by xanthine oxidase. Role of superoxide. Biochem. J. **243**: 55-59, 1987.
- BON, G.B.; CAZZOLATO, G.; AVOGARRO, P. Changes of apolipoprotein B molecular weight and immunoreactivity in malondialdehyde-modified low density lipoproteins. Artery, **12**: 74-80, 1983.
- BOULANGER, C.M.; TANNER, F.C.; BÉA, M.L.; HANHN, A.W.A.; WERNER, A.; LÜSCHER, T.F. Oxidized low density lipoproteins induce mRNA and release of endothelin from human and porcine endothelium. Circ. Res. **70**: 1191-1197, 1992.
- BOYD, H.C.; GOWN, A.M.; WOLFBAUER, G.; CHAIT, A. Direct evidence for a protein recognized by a monoclonal antibody against oxidatively modified LDL in atherosclerotic lesions from Watanabe heritable hyperlipidemic rabbit. Am. J. Pathol. **135**: 815-826, 1989.
- BREDT, D.S. & SNYDER, S.H. Isolation of nitric oxide synthetase, a calmodulin-requiring enzyme. Proc. Natl. Acad. Sci. USA **87**: 682-685, 1992.
- BROWN, M.S. & GOLDESTINE, J.L. - Lipoprotein receptors in the liver: control signs for plasma cholesterol traffic. J.Clin.Invest. **72**: 743-747, 1983.
- BUSSE, R. & MÜLSCH, A. Induction of nitric oxide synthase by cytokines in vascular smooth muscle cells. FEBS Lett. **175**: 87-90, 1990.
- BÜSSE, R.; LÜCKHOFF, A.; BASSENGE, E. Endothelial-derived relaxant factor inhibits platelet activation. Naunyn-Schmiedeberg's Arch. Pharmacol. **336**: 566-574, 1987.

- DAUGHERTY, A.; SCHONFELD, G.; SOBEL, B.E.; LANGE, L.G. Metabolism of very low density lipoproteins after cessation of cholesterol feeding in rabbits. J. Clin. Invest. **77**: 1108-1115, 1986.
- DEGRAAF J.; HAK-LEMMERS, H.L.M.; HECTORS, M.P.C.; DEMACKER, P.N.M.; HENDRIKS, J.C.M.; STALENHOEF, A.F.H. Enhanced susceptibility to in vitro oxidation of heavy low density subfraction in healthy subjects. Arterioscler. Thromb. **11**: 298-306, 1991.
- DERGUNOV, A.D. & ROSSENEU, M. The significance of apolipoprotein E structure to the metabolism of plasma triglyceride-rich lipoproteins. Biol. Chem. Hoppe-Seyler **375**: 485-495, 1994.
- DOLPHIN, P.J. & JAUHIAINEN, M. Human plasma lecithin cholesterol acyltransferase: An elucidation of the catalytic mechanism. J. Biol. Chem. **261**: 7032-7043, 1986.
- DREVON, C.A.; WEINSTEIN, D.B.; STEINBERG, D. Regulation of cholesterol ester esterification and biosynthesis in monolayer cultures of normal adult rat macrophages. J. Biol. Chem. **255**: 9128-9137, 1980.
- DREXLER, H.; ZEIHNER, A.M.; MEINZER, K.; JUST, H. Correction of endothelial dysfunction in coronary microcirculation of hypercholesterolemic patients by L-arginine. Lancet **338**: 1546-1550, 1991.
- EGGEN, D.A. & SOLBERG, L.A. Variation of atherosclerosis with age. Lab. Invest. **18**: 571-579, 1968.
- EHRENWALD, E.; CHISOLM, G.M.; FOX, P.L. Intact human ceruloplasmin oxidatively modifies low density lipoprotein. J. Clin. Invest. **93**: 1493-1501, 1994.
- EISENBERG, S. High-density lipoprotein metabolism. J. Lipid Res. **26**: 1017-1032, 1984.
- EISENBERG, S. & LEVY, R.I. - Lipoprotein metabolism. Adv. Lipid Res. **13**: 1-89, 1975.
- ENDEMANN, G.; STANTON, L.W.; MADDEN, K.S.; BRYANT, C.M.; WHITE, R.T.; PROTTER, A.A. CD 36 is a receptor for oxidized low density lipoprotein. J. Biol. Chem. **268**: 11811-11816, 1993.
- ENSTROM, J.E.; KANIM, L.E.; KLEIN, M.A. Vitamin C intake and mortality among a sample of the United States population. Epidemiology **3**: 194-202, 1992.
- EVANS, M.D. & PRYOR, W.A. Cigarette smoking, emphysema, and damage to α_1 -proteinase inhibitor. Am. J. Physiol. **266**: L593-L611, 1994.
- EZAKI, M.; IKEDA, M.; TOMITA, I.; TOMITA, T. Inhibition of EDRF release by native low-density lipoprotein from cultured porcine endothelial cells through intracellular mechanisms. J. Cardiovas. Pharmacol. **24**: 552-558, 1994.
- FIELDING, C.J.; REAVEN, G.M.; LIV, G.; FIELDING, P.E. Increased free cholesterol in plasma low and very low density lipoproteins in non-insulin dependent diabetes mellitus: its role in the inhibition of cholesterol ester transfer. Proc. Natl. Acad. Sci. USA **81**: 2512-2516, 1984.

- FIELDING, C.J. The origin and properties of free cholesterol potential gradients in plasma, and their relation to atherogenesis. J. Lipid Res. **25**: 1624-1628, 1984.
- FLORÉN, C.H. & NILSSON, A. - Binding, interiorization and degradation of cholesteryl ester-labelled chylomicron-remnant particles by rat hepatocyte monolayers. Biochem.J. **168**: 483-94, 1977.
- FOGELMAN, A.M.; SHECHTER, I.; SEAGER, J.; HOKOM, M.; CHILD, J.S.; EDWARDS, P.A. Malondialdehyde alteration of low density lipoproteins leads to cholesteryl ester accumulation in human monocyte-macrophages. Proc. Natl. Acad. Sci. USA **77**: 2214-2218, 1980.
- FÖRSTERMANN, U.; POLLOCK, J.S.; SCHMIDT, H.H.H.W.; HELLER, M.; MURAD, F. Calmodulin-dependent endothelium-derived relaxing factor/nitric oxide synthase activity is present in the particulate and cytosolic fractions of bovine aortic endothelial cells. Proc. Natl. Acad. Sci. USA **88**: 1788-1792, 1991.
- FRANCONE, O. & FIELDING, C.J. Structure-function relationships in human lecithin cholesterol acyltransferase. Site-directed mutagenesis at serine residues 181 and 216. Biochemistry **30**: 1074-1077, 1991.
- FRANGOS, J.A.; ESKIN, S.G.; MCINTIRE, L.V.; IVES, C.L. Flow effects on prostacyclin production by cultured human endothelial cells. Science **227**: 1477-1479, 1984.
- FREDRICKSON, D.S. & LEVY, R.I. Familial hyperlipoproteinemias. In: Stanbury, J.B.; Wyngaarden, J.B.; Fredrickson, D.S. (eds.) The metabolic basis of inherited diseases, 3a. edição, McGraw-Hill Book Co., New York, 1972, pp. 545-614.
- FRIES, D.M.; PENHA, R.G.; D'AMICO, E.A.; ABDALLA, D.S.P.; MONTEIRO, H.P. Oxidized low density lipoprotein stimulates nitric oxide release by rabbit aortic endothelial cells. Biochem. Biophys. Res. Commun. 1995 (no prelo).
- FRÖSTEGÅRD, J.; NILSSON, J.; HAEGERSTRAND, A.; HAMSTEM, A.; WIGZELL, H.; GIDLUND, M. Oxidized low density lipoprotein induces differentiation and adhesion of human monocytes and the monocytic cell line U 937. Proc. Natl. Acad. Sci. USA **87**: 904-908, 1990.
- FRÖSTEGÅRD, J.; WU, R.; GISCOMBE, R.; HOLM, G.; LEFVERT, A.K.; NILSSON J. Induction of T-cell activation by oxidized low density lipoprotein. Arterioscler. Thromb. **12**: 461-468, 1992.
- FUKUZAWA, K.; KISHIKAWA, K.; TOKUMURA, A.; TSUKATANY, H.; SHIBUYA, S. Fluorescent pigments by covalent binding of lipid peroxidation byproducts to protein and aminoacid. Lipids **20**: 854-861, 1985.
- FULLER, J.N.; SHIPLER, M.J.; ROSE, G.; JARRET, R.J.; KEEN, N. Coronary heart disease risk and impaired glucose tolerance, The Whitcath Study. Lancet **1**: 1373-1379, 1980.
- FURCHGOTT, R.F. & VANHOUTTE, P.M. Endothelial-derived relaxing and contracting factors. FASEB J. **3**: 2007-2018, 1989.

- GALLE, J.; MÜLSCH, A.; BUSSE, R.; BASENGE, E. Effects of native and oxidized low density lipoprotein on formation and inactivation of endothelium-derived relaxing factor. Arterioscler. Thromb. **11**: 198-203, 1991.
- GATTI, R.M.; RADI, R.; AUGUSTO, O. Peroxynitrite-mediated oxidation of albumin to the protein-thiyl free radical. FEBS Lett. **348**: 287-290, 1994.
- GAUBATZ, J.W.; HEIDMAN, C.; GOTTO, A.M.J.; MORRISETT J.D.; DAHLEN, G.H. Human plasma lipoprotein (a): structural properties. J. Biol. Chem. **258**: 4582-4589, 1982.
- GAZIANO, J.M.; MASON, J.A.E.; RIDKER, P.M.; BURING, J.E.; HENNEKENS, C.H. Beta carotene therapy for chronic stable angina. Circulation **82** (supl. 3) III201-III206, 1990.
- GENG, Y.J.; KODAMA, T.; HANSSON, G.K. Differential expression of scavenger receptor isoforms during monocyte-macrophage differentiation and foam cell formation. Arterioscler. & Thromb. **14**: 798-806.
- GEY, K.F.; STÄHELIN, H. B.; EICHHOLZER, M. Poor plasma status of carotene and vitamin C is associated with higher mortality from ischemic heart disease and stroke: Basel prospective study. Clin Invest. **71**: 3-6, 1993.
- GILLIGAN, D.M.; GUETTA, V.; PANZA, J.A.; GARCIA, C.E.; QUYYUMI, A.A.; CANNON, R.O. Selective loss of microvascular endothelial function in human hypercholesterolemia. Circulation **90**: 35-41, 1994.
- 13969 GLAGOV, S.; ZARINS, C.; GIDDENS, D.P.; KU, D.N. Hemodynamics and atherosclerosis: insights and prespectives gained from studies of human arteries. Arch. Pathol. Lab. Med. **112**: 1018-1031, 1988.
- GLAVIND, J.; HARTMANN, S.; CLEMMESSEN, J.; JESSEN, K.E.; DAM, H. Studies on the role of lipoperoxides in human pathology. II. The presence of peroxidized lipids in the atherosclerotic aorta. Acta Pathol. Microbiol. Immunol. Scand. **30**: 1-6, 1952.
- GLOMSET, J.A. The plasma lecithin:cholesterol acyltransferase reaction. J. Lipid Res. **9**: 155-167, 1968.
- GLOMSET, J.A.; NORUM, K.P.; GJONE, E.; Familial lecithin:cholesterol acyltransferase deficiency. In: Stanbury, J.B.; Wyngaarden, J.B.; Fredrickson, J. (eds.). The metabolic basis of inherited diseases, McGraw Hill, New York, 1983, pp. 643-654.
- GOLDSTEIN, J.L.; HO, Y.K.; BROWN, M.S.; INNERARITY, T.L.; MAHLEY, R.W. Cholesterol ester accumulation in macrophages resulting from receptor-mediated uptake and degradation of hypercholesterolemic canine β -very low density lipoproteins. J. Lipid Res. **25**: 1939-1848, 1980.
- GÖRÖG, P. Activation of human blood monocytes by oxidized polyunsaturated fatty acids: a possible mechanism for the generation of lipid peroxides in the circulation. Int. J. Exp. Pathol. **72**: 227-235, 1991.

- GORREN, A.C.F.; DE BOER, E.; WEVER, R. The reaction of nitric oxide with copper proteins and the photodissociation of copper-NO complexes. Biochim. Biophys. Acta **916**: 38-47, 1987.
- GOTTO, A.M. Jr. - High density lipoproteins : biochemical and metabolic factors. Am.J.Cardiol. **52**: 2-4, 1983.
- GOTTO, A.M. Hypertriglyceridemia: risks and perspectives. Am. J. Cardiol. **70**: 19H-26H, 1992.
- GRYGLEWSKI, R.J.; PALMER, R.M.J.; MONCADA, S. Superoxide anion is involved in the breakdown of endothelium-derived vascular relaxing factor. Nature **320**: 454-456, 1986.
- GUTTERIDGE, J.M.C. Free radicals in disease processes: a compilation of cause and consequence. Free Radical Res. Commun. **19**:141-158, 1993.
- GUYTON, J.R. & KLEMP, K.F. Development of the atherosclerotic core region. Chemical and ultrastructural analysis of microdissected atherosclerotic lesions from human aorta. Arterioscler. Thromb. **14**: 1305-1314, 1994.
- HABERLAND, M.E.; OLCH, C.L.; FOGELMAN, A.M. Role of lysines in mediating interaction of modified low density lipoproteins with the scavenger receptor of human monocyte macrophages. J. Biol. Chem. **259**: 11305-11311, 1984.
- HABERLAND, M.E.; FONG, D.; CHENG, L. Malondialdehyde-altered protein occurs in atheroma of Watanabe heritable hyperlipidemic rabbits. Science **241**: 215-218, 1988.
- HALLIWELL, B. & GUTTERIDGE, J.M.C. The importance of free radicals and catalytic metal ions in human diseases. Mol. Asp. Med. **8**: 89-193, 1985.
- HAMON, M.; VALLET, B.; BAUTERS, C.; WERNERT, N.; MCFADDEN, E.P.; LABLANCHE, J.M.; DUPUIS, B.; BERTRAND, M.E. Long-term oral administration of L-arginine reduces intimal thickening and enhances neoendothelium-dependent acetylcholine-induced relaxation after arterial injury. Circulation **90**: 1357-1362, 1994.
- HARMAN, D. Atherosclerosis: a hypothesis concerning the initiating steps in pathogenesis. J. Gerontol. **12**: 199-202, 1957.
- HAZELL, L.J.; VAN DEN BERG, J.J.M.; STOCKER, R. Oxidation of low-density lipoprotein by hypochlorite causes aggregation that is mediated by modification of lysine residues rather than lipid oxidation. Biochem. J. **302**: 297-404, 1994.
- HEFFETZ, D. & ZICK, Y. H₂O₂ potentiates phosphorylation of novel putative substrates for the insulin receptor kinase in intact Fao cells. J. Biol. Chem. **264**: 10126-10132, 1989.
- HEINECKE, J.W.; BAKER, L.; ROSEN, H.; CHAIT, A. Superoxide-mediated modification of low density lipoprotein by arterial smooth muscle cells. J. Clin. Invest. **77**: 757-761, 1986.
- HEINECKE, J.W.; ROSEN, H.; SUZUKI, L.A.; CHAIT, A. The role of sulfur-containing amino acids in superoxide production and modification of low density lipoprotein by arterial smooth muscle cells. J. Biol. Chem. **262**: 10098-10103, 1987.

- HEINECKE, J.H.; ROSEN, H.; CHAIT, A. Iron and copper promote modification of low density lipoprotein by human arterial smooth muscle cells in culture. J. Clin. Invest. **74**: 1890-1894, 1984.
- HEINECKE, J.W.; KAWAMURA, M.; SUZUKI, L.; CHAIT, A. Oxidation of low density lipoprotein by thiols: superoxide-dependent and -independent mechanisms. J. Lipid Res. **34**: 2051-2061, 1993.
- HERBERT, H.B.; FEINLEIB, M.; MCNAMARA, M.C.; CASTELLI, W.P. Obesity as an independent risk factor for cardiovascular disease: a 26-year follow-up of participants in the Framingham Heart Study. Circulation **67**: 968-977, 1983.
- HERSHKO, C.; WEATHERALL, D.J. Iron-chelating therapy. CRC Crit. Rev. Clin. Lab. Sci. **26**: 303-345, 1988.
- HINSBERGH, V.W.M.; SCHEFFER, M.; HAVEKES, L.; KEMPEN, H.J.M. Role of endothelial cells and their products in the modification of low-density lipoprotein. Biochim. Biophys. Acta. **878**: 49-64, 1986.
- HIRAMATSU, K.; ROSEN, H.; HEINECKE, J.W.; WOLFBauer, G.; CHAIT, A. Superoxide initiates oxidation of low density lipoprotein by human monocytes. Arteriosclerosis, **7**: 55-60, 1987.
- HIRAMATSU, K. & ARIMORI, A. Increased superoxide production by mononuclear cells of patients with hypertriglyceridemia and diabetes. Diabetes **37**: 832-837, 1988.
- HIXSON, J.E.; BRITTEN, M.L.; MANIS, G.S.; RAINWATER, D. Apolipoprotein (a) apo (a) glycoprotein isoforms result from size differences in apo (a) mRNA in baboons. J. Biol. Chem. **264**: 6013-6016, 1989.
- HOBBS, A.J.; FUKUTO, J.M.; IGNARRO, L.J. Formation of free nitric oxide from L-arginine by nitric oxide synthase: direct enhancement of generation by superoxide dismutase. Proc. Natl. Acad. Sci. USA **91**: 10992-10996, 1994.
- HODIS, H.N.; KRAMSCH, D.M.; AVOGARO, P.; BITTOLO-BON, G.; CAZZOLATO, G.; HWANG, J.; PETERSON, H.; SEVANIAN, A. Biochemical and cytotoxic characteristics of an in vivo circulating oxidized low density lipoprotein (LDL -). J. Lipid Res. **35**: 669-677, 1994.
- HODIS, H.N.; MACK, W.J.; AZEN, S.P.; ALAUPOVIC, P.; POGODA, J.M.; LABREE, L.; HEMPHILL, L.C.; KRAMSCH, J.C.; BLANKENHORN, D.H. Triglyceride- and cholesterol-rich lipoproteins have a differential effect on mild moderate and severe lesion progression as assessed by quantitative coronary angiography in a controlled trial of lovastatin. Circulation **90**: 42-49, 1994.
- HUIE, R.E. & PADMAJA, S. The reaction of nitric oxide with organic peroxy radicals. Free Rad. Res. Commun. **18**: 195-199, 1993.
- HUTCHESON I.R.; SMITH, J.A.; GRIFFITH, T.M. Abolition of flow-dependent EDRF release before that evoked by agonists in hypercholesterolemic rabbits. Br. J. Pharmacol. **113**: 190-194, 1994.
- ISCHIROPOULOS, H.; ZHU, L.; BECKMAN J.S. Peroxynitrite formation from macrophage-derived nitric oxide. Arch. Biochem. Biophys. **298**: 446-451, 1992.

- ISCHIROPOULOS, H.; ZHU, L.; CHEN, J.; TSAI, M.; MARTIN, J.M.; SMITH, C.D.; BECKMAN, J.S. Peroxynitrite-mediated tyrosine nitration catalysed by superoxide dismutase. Arch. Biochem. Biophys. 298: 431-437, 1992a.
- JACKSON, R.R.; MORRISETT, J.D.; GOTTO, A.M. Lipoproteins and lipid transport: structural and functional concepts. In: Rifkind, B.M. & Levy, R.I. (eds.) Hyperlipidemia: Diagnosis and Therapy, Grune & Stratton, New York, 1977, pp.1-16.
- JAYAKODY, L.; SENARATNE, M.P.; THOMPSON, A.B.; KAPPAGODA, C.T. Endothelium-dependent relaxation in experimental atherosclerosis in the rabbit. Circ. Res. 60: 251-264, 1987.
- JENG, J.R.; SHIEH, S.M.; CHIU, H.C. Decreased expression of a particular epitope on plasma apolipoprotein B of patients with coronary artery disease. Biochim. Biophys. Acta 1181: 287 - 294, 1993.
- JESAITIS, A. J.; QUINN, M.T.; MUKHERJEE, G.; WARD, P.A.; DRATZ, E.A. Death by oxygen: Radical views. New Biol. 3: 651-655, 1991.
- JIALAL, I. & CHAIT, A. Differences in the metabolism of oxidatively modified low density lipoprotein and acetylated low density lipoprotein by human endothelial cells: inhibition of cholesterol esterification by oxidatively modified low density lipoprotein. J. Lipid Res. 30: 1561-1568, 1989.
- JOHNSON, W.J.; MAHLBERG, F.H.; ROTHBLAT, G.H.; PHILLIPS, M.C. Cholesterol transport between cells and high-density lipoproteins. Biochim. Biophys. Acta 1085:273-298, 1991.
- JOKINEN, M.P.; HOLM, J.; SHALLI, O.; BONDJERO, G.; HANSSON, G.K. Regional accumulation of T cells, macrophages, and smooth muscle cells in human atherosclerotic plaque. Arteriosclerosis 6: 131-138, 1986.
- JONAS, A. Lecithin:cholesterol acyltransferase in the metabolism of high density lipoproteins. Biochim. Biophys. Acta 1084: 205-220, 1991
- JUCKETT, M.B.; BALLA, J.; BALLA, G.; BURKE, B.; JACOB, H.S.; VERCELLOTTI, G.M. Ferritin protects endothelial cells from oxidized LDL mediated cytotoxicity. Clin. Res. 41: 162 A, 1993.
- JURGENS, G.; HOFF, H.F.; CHISOLM, G.M.; HERMANN, E. Modification of human serum low density lipoprotein by oxidation - characterization and pathophysiological implications. Chem. Phys. Lipids, 45: 315-336, 1987.
- KAKU, S.; TANAKA, M.; MURAMATSU, M.; OTOMO, S. Determination of nitrite by high performance chromatography system with electrochemical detector: measurement of nitric oxide synthase activity in rat cerebellum cytosol. Biomed. Chromatogr. 8: 14-18, 1994.
- KANNEL, W.B. Role of blood pressure in cardiovascular disease. The Framingham Study. Angiology 26: 1-14, 1975.
- KARLSSON, K. & MARKLUND, S.L. Extracellular superoxide dismutase by human cell lines. Biochem. J. 255: 223-231, 1988.

- KATSUDA, S.; BOYD, H.C.; FLIGNER, C.; ROSS, R.; GROWN, A.M. Human atherosclerosis, part III: immunocytochemical analysis of the cell composition of lesions of young adults. Am. J. Pathol. 140: 907-914, 1992.
- KATZ, S.S. & SMALL, D.M. Isolation and partial characterization of the lipid phases of human atherosclerotic plaques. J. Biol. Chem. 255: 9753-9759, 1980.
- KNOWLES, R.G. & MONCADA, S. Nitric oxide synthases in mammals. Biochem. J. 298: 249-258, 1994.
- KOOY, N.W. & ROYALL, J.A. Agonist induced peroxynitrite production from endothelial cells. Arch. Biochem. Biophys. 310: 352-359, 1994.
- KRAISS, L.W.; KIKMAN, T.R.; KOHLER, T.R.; ZIERLER, B.; CLOWES, A.W. Shear stress regulates smooth muscle proliferation and neointimal thickening in porous polytetrafluoroethylene grafts. Arterioscler. Thromb. 11: 1844-1852, 1991.
- KRAUSE, S.; POHL, A.; POHL, C.; LIEBRENZ, A.; RÜHLING, K.; LÖSCHE, W. Increased generation of reactive oxygen species in mononuclear blood cells from hypercholesterolemic patients. Thrombosis Res. 71: 237-240, 1993.
- KREUZER, J. & HODENBERG, E. The role of apolipoproteins in lipid metabolism and atherogenesis: aspects in man and mice. J. Hypertension 12: 113-118, 1994.
- KRINSKY, N.I. Mechanism of action of biological antioxidants. Proc. Soc. Exp. Biol. Med. 200: 248-256, 1992.
- KROON, P.A.; THOMPSON, G.M.; CHAO, Y. β -VLDL in cholesterol-fed rabbits are of hepatic origin. Atherosclerosis 56: 323-326, 1985.
- KU, D.N.; GIDDENS, D.P.; ZARINS, C.K.; GLAGOV, S. Pulsatile flow and atherosclerosis in the human carotid bifurcation. Positive correlation between plaque location and low and oscillatory shear stress. Arteriosclerosis 5: 293-302, 1985.
- KUGIYAMA, K.; KERNS, S.A.; MORRISETT, J.D.; ROBERTS, R.; HENRY, P.D. Impairment of endothelium-dependent arterial by lysolecithin in modified low density lipoproteins. Nature 344: 160-162, 1990.
- KUIJK, F.J.G.M.; SEVANIAN, A.; HANDELMAN, G.J.; DRATZ, E.A. A new role for phospholipase A₂: protection of membranes from lipid peroxidation damage. TIBS 12: 31-37, 1987.
- KUNITAKE, S.T.; LA SALA, K.J.; KANE, J.P. Apolipoprotein A-1-containing lipoproteins with pre-beta electrophoretic mobility. J. Lipid Res. 26: 549-555, 1985.
- LAURINDO, F.R.M.; PEDRO, M.A.; BARBEIRO, H.V.; PILEGGI, F.; CARVALHO, M.H.C.; AUGUSTO, O.; LUZ, P.L. Vascular free radical release. Ex vivo and in vivo evidence for a flow-dependent endothelial mechanism. Cir. Res. 74: 700-709, 1994.

- LAVY, A.; BROOK, G.J.; DANKNER, G.; AMOTZ, A.B.; AVIRAM, M. Enhanced in vitro oxidation of plasma lipoproteins derived from hypercholesterolemic patients. Metabolism 40: 794-799, 1991.
- LUDWIG, P.W.; HUNNINGHAKE, D.B.; HOIDAL J.R. Increased leucocyte oxidative metabolism in hyperlipoproteinemia. Lancet 2: 348-350, 1982.
- LÜSCHER, T.F.; TANNER, F.C.; TSCHUDI, M. R.; NOLL, G. Endothelial dysfunction in coronary artery disease. Annu. Rev. Med. 44: 395-418, 1993.
- LUSKEY, K.L.; GOLDSTEIN, J.L. - Stimulation of the synthesis of very low density lipoprotein in rooster liver by estradiol. J.Biol.Chem. 249: 5939-5947, 1974.
- MAHLEY, R.W. & RALL, S.C., JR. Type III hyperlipoproteinemia (dysbetalipoproteinemia): the role of apolipoprotein E in normal and abnormal lipoprotein metabolism. In: Scriver, C.R.; Beaudet, A.L.; Sly, W.S.; Valle, D. (eds). The metabolic basis of inherited disease, New York, N.Y., McGraw-Hill Book Co., 1989, pp. 1195-1214.
- MAHLEY, R.W.; INNERARITY, T.L.; WEISGRABER, K.H.; OH, S.Y. Altered metabolism (in vivo and in vitro) of plasma lipoproteins after selective chemical modification of lysine residues of the apoproteins. J. Clin. Invest. 64: 743-750, 1979.
- MAHLEY, R.W. & INNERARITY, T.L. Lipoprotein receptors and cholesterol homeostasis. Biochem. Biophys. Acta 737: 197-222, 1983.
- MALLE, E. & SATTler, W. Platelets and the lipoproteins: native, modified and platelet modified lipoproteins. Platelets 5: 70-83, 1994.
- MARCEL, Y.L.; JEWER, L.; LEBLOND, L.; WEECH, P.K.; MILNE, R.W. Lipid peroxidation changes the expression of specific epitopes of apolipoprotein A1. J. Biol. Chem. 264: 19942-19950, 1989.
- MARCEL, Y.L.; MCPHERSON, L.; HOGHE, M.; CZARRNECKA, H.; ZAWADSKI, Z.; WEECH, P.K.; WHITLOCK, M.E.; TALL, A.R.; MILNE, R.W. Distribution and concentration of cholesteryl ester transfer protein in plasma of normolipidemic subjects. J. Clin. Invest. 85: 10-17, 1990.
- MARCZIN, N.; PAPAPETROPOULOS, A.; CATRAVAS, J.D. Tyrosine kinase inhibitors suppress endotoxin- and IL-1 β -induced NO synthesis in aortic smooth muscle cells. Am. J. Physiol. 265: H1014-H1018 (1993).
- MCGILL, H.C. The cardiovascular pathology of smoking. Am. Heart J. 115: 250-257, 1988.
- MCLEAN, J.W.; TOMLINSON, J.E.; KUANG, W.J.; EATON, D.L.; CHEN, E.Y.; FLESS, G.M.; SCANU, A.M.; LAWN, R.M. cDNA sequence of human apolipoprotein (a) is homologous to plasminogen. Nature 330: 132-137, 1987.
- MEULEMANS, A. & DELSENNE, F. Measurement of nitrite and nitrate levels in biological samples by capillary electrophoresis. J. Chromatogr. B 660: 401-404, 1994.

- MICKEL, S.H. & FOLDS JR., E.L. The effect of hydrogen peroxide on serum lecithin-cholesterol acyltransferase activity. Lipids **5**: 663-664, 1970.
- MICKEL, S.H.; FOULDS, JR, E.L.; CLARK, D.A.; LARKIN, D. Effects of pure oxygen atmosphere in vivo on plasma lecithin-cholesterol acyltransferase reaction hydrogen peroxide and peroxidized lecithin. Lipids **7**: 121-124, 1972.
- MICKEL, S.H.; HILL, P.L.; HAYES, K.C. Depression of lecithin-cholesterol acyltransferase esterification in vitamin E-deficient monkeys. Am. J. Clin. Nutr. **28**: 994-999, 1975.
- MINOR, R.L.; MYERS, P.R.; GUERRA, R.; BATES, J.N.; HARRISON, D.G. Diet-induced atherosclerosis increases the release of nitrogen oxides from rabbit aorta. J. Clin. Invest. **86**: 2109-2116, 1990.
- MISIK, V.; ONDRIAS, K.; BALGAVY, P. Effect of lipid peroxidation on molecular arrangement of phospholipids in liposomes prepared from egg phosphatidylcholine or total brain lipids. A ³¹P-NMR study. Gen. Physiol. Biophys. **11**: 317-325, 1992.
- MITCHELL, J.A.; WARNER, T.D.; HUANG, Z.J.; FÖSTERMANN, U.; MURAD, F. Native LDL inhibits the release of endothelial derived relaxing factor by reducing the activity of endothelial nitric oxide synthase. J. Vasc. Res. **29**: 169-176, 1992.
- MONCADA, S.; PALMER, R.M.J.; HIGGS, E.A. Nitric oxide: physiology, pathophysiology, and pharmacology. Pharmacol. Rev. **43**: 109-142, 1991.
- MONCADA, S.; PALMER, R.M.J.; HIGGS, E.A. The discovery of nitric oxide as the endogenous nitrovasodilator. Hypertension **12**: 365-372, 1988.
- MONTEIRO, H.P. & WINTERBOURN, C.C. The superoxide-dependent iron transfer from ferritin to transferrin and lactoferrin. Biochem. J. **256**: 923-930, 1988.
- MÜGGE, A.; ELWELL, J.H.; PETERSON, T. E.; HOFMEYER, T.G.; HEISTAD, D.D.; HARRISON, D.G. Chronic treatment with polyethylene-glycolated superoxide dismutase partially improves endothelium-dependent vascular relaxations in cholesterol-fed rabbits. Circ. Res. **69**: 1293-1300, 1991.
- MUNRO, H.N.; LINDER, M. Ferritin: structure, biosynthesis and role in iron metabolism. Physiol. Rev. **58**: 317-332, 1978.
- MUNRO, J.M.; VAN DER WALT, J.D.; MUNRO, C.S.; CHALMERS, J.A.; COX, E.L. An immunohistochemical analysis of human aortic fatty streaks. Hum. Pathol. **18**: 375-380, 1987.
- MURPHY, M.E. & SIES, H. Reversible conversion of nitroxyl anion to nitric oxide by superoxide dismutase. Proc. Natl. Acad. Sci. USA **88**: 10860-10864, 1991.
- NAKAKI, T.; NAKAYAMA, M.; KATO, R. Inhibition by nitric oxide and nitric oxide-producing vasodilators of DNA synthesis in vascular smooth muscle cells. Eur. J. Pharmacol. **189**: 347-353, 1990.

- NATHAN, C. Nitric oxide as a secretory product of mammalian cells. FASEB J. 6: 3051-3064, 1992.
- NATHAN, C. & XIE, Q. Regulation of biosynthesis of nitric oxide. J. Biol. Chem. 269: 13725-13728, 1994.
- NEWMAN, W.P.; FREEDMAN, D.S.; VOORS, A.W.; GARD, P.D.; SINIVASAN, S.R.; CRESANTA, J.L.; WILLIAMSON, G.D.; WEBBER, L.S.; BERENSON, G.S. Relation of serum lipoprotein levels and systolic blood pressure to early atherosclerosis: The Bogalusa Heart Study. N. Engl. J. Med. 314: 138-144, 1986.
- O'BRIEN, K.D. & CHAIT, A. The biology of the artery wall in atherogenesis. Med. Clin. North Am. 78: 41-67, 1994.
- O'CONNELL, A.M.; GIESEG, S.P.; STANLEY, K.K. Hypochlorite oxidation causes cross-linking of Lp(a). Biochim. Biophys. Acta 1225: 180-186, 1994.
- OHARA, Y.; PETERSON, T.E.; HARRISON, D.G. Hypercholesterolemia increases endothelial superoxide anion production. J. Clin. Invest. 91: 2546-2551, 1993.
- OHSHIMA, H.; FRIESEN, M.; BROUET, I.; BARTSCH, H. Nitrotyrosine as a new marker for endogenous nitrosation and nitration of proteins. Fund. Chem. Tox. 28: 647-652, 1990.
- OREKOV, A.N.; TERTOV, V.V.; MUKHIN, D.N.; MIKHAILENKO, I.A. Modification of low density lipoprotein by desialylation causes lipid accumulation in cultured cells. Biochem. Biophys. Res. Commun. 162: 206-211, 1989.
- OSTERGAARD, H.L. & TROWBRIDGE, I.S. Negative regulation of CD 45 protein tyrosine phosphatase activity by ionomycin in T-cells. Science 253: 1423-1425, 1991.
- OZAWA, J.A.; OHTSUKA, T. OKAMURA, N.; ISHIBASHI, S. Synergism between protein kinase C activator and fatty acids in stimulating superoxide anion production in guinea pig polymorphonuclear leukocytes. Arch. Biochem. Biophys. 273: 491-496, 1989.
- PALINSKI, W.; ROSENFELD, M.E.; YLÄ-HERTTUALA, S. et al. Low density lipoprotein undergoes oxidative modification in-vivo. Proc. Natl. Acad. Sci. USA 86: 1372-1376, 1989.
- PALINSKI, W.; YLÄ-HERTTUALA, S.; ROSENFELD, M.E. ET AL. Antisera and monoclonal antibodies specific for epitopes generated during oxidative modification of low density lipoprotein. Arteriosclerosis 10: 325-335, 1990.
- PARTHASARATHY, S.; QUINN, M.T.; SCHWENKE, D.C.; CAREW, T.E.; STEINBERG, D. Oxidative modification of beta-very low density lipoprotein. Potential role in monocyte recruitment and foam cell formation. Arteriosclerosis 9: 398-404, 1989.
- PARTHASARATHY, S.; WIELAND, E.; STEINBERG, D. A role for endothelial cell lipoxygenase in the oxidative modification of low density lipoprotein. Proc. Natl. Acad. Sci. USA 86: 1046-1050, 1989.
- PARTHASARATHY, S.; BARNETT, J.; FONG, L.G. High-density lipoprotein inhibits the oxidative modification of low density lipoprotein. Biochem. Biophys. Acta 1044: 275-282, 1990.

- PARTHASARATHY, S. Modified lipoproteins in the patogenesis of atherosclerosis. R.G. Landes Co., Austin, 1994, 131 pp.
- PARUMS DV, MITCHINSON MJ. Serum antibodies to oxidized LDL and ceroid in chronic periaortitis. Am. J. Pathol. 151: 57A-65A, 1987.
- PATSCH, J.R.; KARLIN, J.B.; SCOTT, L.M.; SMITH, L.C.; GOTTO, A.M. Inverse relationship between blood levels of high density lipoprotein subfraction 2 and magnitude of postprandial lipemia. Proc. Natl. Acad. Sci. USA 80: 1449-1453, 1983.
- POMERANTZ, K. B.; HAJJAR, D.P.; LEVI, R.; GROSS, S.S. Cholesterol enrichment of arterial smooth muscle cells upregulates cytokine-induced nitric oxide synthesis. Biochem. Biophys. Res. Commun. 191: 103-109, 1993.
- PRIOR, J.T.; KURTZ, D.M.; ZIEGLER, D.D. The hypercholesterolemic rabbit. An aid to understanding arteriosclerosis in man? Arch. Pathol. 71: 672-679, 1961.
- PRÓNAI, L.; HIRAMATSU, K.; SAIGUSA, Y.; NAKAZAWA, H. Low superoxide scavenging activity associated with enhanced superoxide generation by monocytes from male hypertriglyceridemia with and without diabetes. Atherosclerosis 90: 39-47, 1991.
- RADI, R.; BECKMAN J.S.; BUSH, K.M.; FREEMAN, B.A. Peroxynitrite-induced membrane lipid peroxidation: The cytotoxic potential of superoxide and nitric oxide. Arch. Biochem. Biophys. 288: 481-487, 1991.
- RADOMSKI, M.W.; PALMER, R.M.J.; MONCADA, S. Glucocorticoids inhibit the expression of an inducible, but not the constitutive, nitric oxide synthase in vascular endothelial cells. Proc. Natl. Acad. Sci. USA 87: 10043-10047, 1990.
- RAGAZZI, E.; CHINELLATO, A.; DE BIASI, M.; PANDOLFO, L.; PROSDOCIMI, M.; NORIDO, F.; CAPARROTTA, L.; FASSINA, G. Endothelium-dependent relaxation, cholesterol content and high energy metabolite balance in Watanabe hyperlipidemic rabbit aorta. Atherosclerosis 80: 125-134, 1989.
- RAJAVASHISTH, T.B.; ANDALIBI, A.; TERRITO, M.C.; BERLINER, J.A.; NAVAB, M.; FOGELMAN, A.M.; LUSIS, A.J. Induction of endothelial cell expression of granulocyte and macrophage colony-stimulating factors by modified low density lipoproteins. Nature 344: 254-259, 1990.
- RANKIN, S.M.; PARTHASARATHY, S.; STEINBERG, D. Evidence for a dominant role of lipoxygenase(s) in the oxidation of LDL by mouse peritoneal macrophages. J. Lipid Res. 32: 449-456, 1991.
- RAPACZ, J.; HASLER - RAPACZ, J.; TAYLOR, K.M.; CHECOVICH, W.J.; ATTLE, A.D. - Lipoprotein mutations in pigs are associated with elevated plasma cholesterol and artherosclerosis. Science 234: 1573-1577, 1986.
- REDGRAVE, T.G. & SMALL, D.M. - Quantitation of the transfer of surface phospholipid of chylomicrons to the high density lipoprotein fraction during catabolism of chylomicrons in the rat. J.Clin.Invest. 64 : 162-171, 1979.

- REGNSTRÖM, J. & NILSSON, J. Lipid oxidation and inflammation-induced intimal fibrosis in coronary heart disease. J. Lab. Clin. Med. 124: 162-168, 1994.
- REINHART, W.R. Shear-dependence of endothelial functions. Experientia 50: 87-93, 1994.
- RICE-EVANS, C., GREEN, E.; PAGANGA, G.; COOPER, C.; WRIGGLESWORTH, J. Oxidised low density lipoproteins induce iron release from activated myoglobin. FEBS Lett. 326: 177-182, 1993.
- RIMM, E.B.; STAMPFER, M.J.; ASCHERIO, A.; GIOVANUCCI, E.; COLDITZ, G.A.; WILLET, W.C. N. Vitamin E consumption and the risk of coronary heart disease. N. Eng. Med. 328: 1450-1456, 1993.
- RINGER, T.V.; DELOOF, M.J.; WINTERROWD, G.E.; FRANCON, S.R.; GAYLOR, S.K.; RYAN, J.A.; SANDERS, M.E.; HUGHES, G.S. Beta-carotene's effects on serum lipoproteins and immunologic indices in humans. Am. J. Clin. Nutr. 53: 688-697, 1991.
- ROSEN, G.M. & FREEMAN, B.A. Detection of superoxide generated by endothelial cells. Proc. Natl. Acad. Sci. USA 81: 7268-7273.
- ROSENFELD, M.E.; PALINSKI, W.; YLÄ-HERTTUALA, S.; BUTLER, S.; WITZTUM, J.L. Distribution of oxidation specific lipid-protein adducts and apolipoprotein B in atherosclerosis lesions of varying severity from WHHL rabbits. Arteriosclerosis 10: 336-349, 1990.
- ROSENFELD, M.; KHOO, J.C.; MILLER, E.; PARTHASARATHY, S.; PALINSKI, W.; WITZTUN, J.L. Macrophage-derived foam cells freshly isolated from rabbit atherosclerotic lesions degrade modified lipoproteins, promote oxidation of low density lipoproteins, and contain oxidation-specific lipid protein adducts. J. Clin. Invest. 87: 90-99, 1991.
- ROSS, A.C. & ZILVERSMITH, D.B. Chylomicron remnant cholesteryl esters as the major constituent of very low density lipoproteins in plasma of cholesterol-fed rabbits. J. Lipid Res. 18: 169-181, 1977.
- ROSS, R. The pathogenesis of atherosclerosis: a perspective for the 1990s. Nature 362: 801-809, 1993.
- RUBANY, G. & VANHOUTTE, P.M. Oxygen-derived free radicals, endothelium, and responsiveness of vascular smooth muscle. Am. J. Physiol. 250: H815-H821, 1986.
- RUBANYI, G.M.; ROMERO, J.C.; VANHOUTTE, P.M. Flow-induced release of endothelium-derived relaxing factor. Am. J. Physiol. 250: H1145-H1149, 1986.
- RUDEL, L.L.; PARKS, J.S.; JOHNSONS, F.L.; BABIAK, J. Low density lipoprotein in atherosclerosis. J. Lipid Res. 27: 465-474, 1986.
- SALMON, S.; SANTUS, R.; MAZINRE, J.C.; AUBAILLY, M.; HAIGLE, J. Modified apolipoprotein pattern after irradiation of human high-density lipoproteins by ultraviolet B. Biochim. Biophys. Acta 1128: 167-173, 1992.

- SALONEN, J.T.; NYYSSÖNEN, K.; KORPELA, H.; TUOMILETHO, J.; SEPPÄNEN, R.; SALONEN, R. High stored iron levels are associated with excess risk of myocardial infarction in eastern finnish men. Circulation 86: 803-811, 1992.
- SALONEN, J.T.; NYYSSÖNEN, K.; KORPELA, H.; TUOMILEHTO, J.; SEPPÄNEN, R.; SALONEN, R. High stored iron levels are associated with excess risk of myocardial infarction in eastern finnish men. Circulation 86: 803-811, 1992.
- SALONEN, J.T.; YLA-HERTTUALA, S.; YAMAMOTO, R.; BUTLER, S.; KORPELA, H.; SALONEN, R.; NYYSSÖNEN, K.; PALINSKI, W.; WITZUM, J. Autoantibody against oxidised LDL and progression of carotid atherosclerosis. Lancet 339: 883-891, 1992.
- SARAN, M.; MICHEL, C.; BORS, W. Reaction of NO with O_2^- . Implications for the action of endothelium derived relaxing factor (EDRF). Free Rad. Res. Commun. 10: 221-226, 1990.
- SAVENKOVA, M.I.; MUELLER, D.M.; HEINECKE, J.W. Tyrosyl radical generated by myeloperoxidase is a physiological catalyst for the initiation of lipid peroxidation in low density lipoprotein. J. Biol. Chem. 269: 20394-20400, 1994.
- SCANU, A.M. Lp(a) as a marker for coronary heart disease risk. Clin. Cardiol. 14: I-35-39, 1991.
- SCHAICH, K.M. Free radical initiation in proteins and monoamino acids by ionizing and ultraviolet radicals and lipid oxidation. CRC Crit. Rev. Food Sci. 13: 89-130, 1980.
- SCHEFFLER, E.; HUBER, L.; FRUHBIS, J.; SCHULZ, I.; ZIEGLER, R.; DRESEL, H.A. Alteration of plasma low density lipoprotein from smokers. Atherosclerosis 82: 261-265, 1990.
- SCHMIDT, K.; KLATT, P.; GRAIER, W.F.; KOSTNER, G.M.; KUKOVETZ, W.R. High-density lipoprotein antagonizes the inhibitory effects of oxidized low-density lipoprotein and lysolecithin on soluble guanylyl cyclase. Biochem. Biophys. Acta 182: 302-308, 1992.
- SCHULZ, R.; NAJA, E.; MONCADA, S. Induction and potential biological relevance of a Ca^{2+} -independent nitric oxide synthase in the myocardium. Br. J. Pharmacol. 105: 575-580, 1992.
- SELIGMAN, M.L.; MITAMURA, J.; SHERA, N.; DEMOPOULOS, H.B. Corticosteroid (methylprednisolone) modulation of photoperoxidation by ultraviolet light in liposomes. Photochem. Photobiol. 29: 549-558, 1979.
- SGOUTA, D.S. Fatty acid specificity of plasma phosphatidylcholine:cholesterol acyltransferase. Biochemistry 11: 293-296, 1972.
- SIES, H. Biochemistry of oxidative stress. Angw. Chem. I 25: 1058-1071, 1986.
- SLYPER, A.H. Low-density lipoprotein density and atherosclerosis. Unraveling the connection. JAMA 272: 305-308, 1994.
- SPARROW, C.P. & OLSZEWSKI, J. Cellular oxidation of low density lipoprotein is caused by thiol production in media containing transition metal ions. J. Lipid Res. 34: 1219-1228, 1993.

- SIMIONESCU, N.; VASILE, E.; LUPU, F.; POPESCU, G.; SIMIONESCU, M. Prelesional events in atherogenesis. Am. J. Pathol. **123**: 109-125, 1986.
- SHORE, V.G.; SHORE, B.; HART, R.G. Changes in apolipoproteins and properties of rabbit very low density lipoproteins on induction of cholesterolemia. Biochemistry **13**: 1579-1584, 1974.
- SRINIVASAN, S.R.; RADHAKRISHNA-MURTHY, B.; SMITH, C.C.; WOLF, R.H.; BERENSON, G.S. Serum lipid and lipoprotein responses of six nonhuman primate species to dietary changes in cholesterol levels. J. Nutr. **106**: 1757-1767, 1976.
- STANBURY, J.B.; WYNGAARDEN, J.B.; FREDRICKSON, J. (eds.). The metabolic basis of inherited diseases, McGraw Hill, New York, 1983, p.
- STARY, H.C. The sequence of cell and matrix changes in atherosclerotic lesions of coronary arteries in the first forty years of life. Eur. Heart J. **11** (supl. E): 3-19, 1990.
- STARY, H.C.; CHANDLER, A.B.; GLAGOV, S.; GUYTON, J.R.; INSULL, W.; ROSENFELD, M.E.; SCHAFFER, S.A.; SCHWARTZ, C.J.; WAGNER, W.D.; WISSLER, R.W. A definition of initial, fatty streak, and intermediate lesions of atherosclerosis. A report from the committee on vascular lesions of the council on atherosclerosis, American Heart Association. Circulation **89**: 2462-2478, 1994.
- STEIN, Y.; STEIN, O.; OLIVERCRONA, T.; HALPERIN, G. The origin and properties of free cholesterol potential gradients in plasma, and their relation to atherogenesis. Biochem. Biophys. Acta **834**: 336-345, 1985.
- STEINBERG, D.; PARTHASARATHY, S.; CAREW, T.E.; KHOO, J.C.; WITZTUM, J.L. Beyond cholesterol - Modifications of low-density lipoprotein that increase its atherogenicity. New Engl. J. Med. **320**: 915-924, 1989.
- STEINBRECHER, U.P. Role of superoxide in endothelial-cell modification of low density lipoproteins. Biochim. Biophys. Acta **959**: 20-30, 1988.
- STIKO-RHAM, A.; HULTGARDH-NILSSON, A.; REGNSTRÖM, J.; HAMSTEN, A.; NILSSON, J. Native and oxidized LDL enhances production of PDGF AA and the surface expression of PDGF receptors in cultured human smooth muscle cells. Arterioscler. Thromb. **12**: 1099-1108, 1992.
- STRONG, J.P. & MCGILL, H.C. The pediatric aspects of atherosclerosis. J. Atheroscler. Res. **9**: 251-256, 1969.
- TAN, M.H.; SATA, T.; HAVEL, R.J. The significance of lipoprotein lipase in the rat skeletal muscles. J.Lipid.Res. **18**: 363-370, 1977.
- TAKAHASHI, S.; KAWARABAYASI, Y.; NAKAI, T.; SAKAI, J.; YAMAMOTO, T. Rabbit very low density lipoprotein receptor: A low density lipoprotein receptor-like protein with distinct ligand specificity. Proc. Natl. Acad. Sci. USA **89**: 9252-9256, 1994.
- TALL, A.; SWENSON, T.; HESLER, C.; GRANOT, E. Mechanisms of facilitated lipid transfer mediated by plasma lipid peroxidation transfer proteins. In: Gotto, A.M. (ed.) Plasma lipoproteins, Elsevier Science Publ. B.V., Amsterdam, 1987, pp. 277-297.

- TAN, M.S.; LEE, Y.J.; SHIN, S.J.; TSAI, J.H. Regulation of heparin binding-epidermal growth factor like growth factor gene expression by LDL and oxidized-LDL in rat mesangial cells. Biochem. Biophys. Res. Commun. 202: 1001-1008, 1994.
- TATAMI, R.; MABUCHI, H.; UEDA, K.; UEDA, R.; HABA, T.; KAMETANI, T.; ITO, S.; KOIZUMI, J.; OHTA, M.; MIYAMOTO, S.; NAKAYAMA, H.; OIWAKE, H.; GENDA, A.; TAKEDA, R. Intermediate-density lipoprotein and cholesterol-rich very low density lipoprotein in angiographically determined coronary artery disease. Circulation 64: 1174-1184, 1981.
- TERKELTAUB, R.; BANKA, C.L.; SOLAN, J.; SANTORO, D.; BRAND, K.; CURTISS, L.K. Oxidized LDL monocytic cell expression of interleukin-8, a chemokine with T-lymphocyte chemotactic activity. Arterioscler. Thromb. 14: 47:53, 1994.
- THOMAS, C.E.; JACKSON, R.L.; OHLWEILER, D.F.; KU, G. Multiple lipid oxidation products in low density lipoproteins induce interleukin-1 beta release from human blood mononuclear cells. J. Lipid Res. 35: 417-427, 1994.
- THOMPSON, K.H. & ZIVERSMIT, D.B. Plasma VLDL in cholesterol-fed rabbits: chylomicron remnants of liver lipoproteins. J. Nutr. 113: 2002-2010, 1983.
- UTERMANN, G. The mysteries of lipoprotein (a). Science 246: 904-910, 1989.
- VERBEUREN, T.J.; JORDAENS, F.H.; ZONNEKEYN, L.L.; VAN HOVE, C.E.; COENE, M.C.; HERMAN, A.G. Effect of hypercholesterolemia on vascular reactivity in the rabbit. Circ. Res. 58: 552-554, 1986.
- VERBEUREN, T.J.; BONHOMME, E.; LAUBIE, M.; SIMONET, S. Evidence for induction of nonendothelial NO synthase in aortas of cholesterol-fed rabbits. J. Cardiovasc. Pharmacol. 21: 841-845, 1993.
- VIRELLA, G.; VIRELLA, I.; LEMAN, R.B.; PRYOR M.B.; VIRELLA M.F.L. Anti-oxidized low density lipoprotein antibodies in patients with coronary heart disease and normal healthy volunteers. Int. J. Clin. Lab. Res. 23: 95 - 101, 1993.
- WARD, R.J.; KÜHN, L.C.; KALDY, P.; FLORENCE, A.; PETERS, T.J.; CRICHTON, R.R. Control of cellular iron homeostasis by iron-responsive elements *in vivo*. Eur. J. Biochem. 220: 927-931, 1994.
- WHITE, C.R.; BROCK, T.A.; CHANG, L.Y.; CRAPO, J.; BRISCOE, P.; KU, D.; BRADLEY, W.A.; GIAN TURCO, S.H.; GORE, J.; FREEMAN, B.A.; TARPEY, M.M. Superoxide and peroxynitrite in atherosclerosis. Proc. Natl. Acad. Sci. USA 91: 1044-1048, 1994.
- WIELAND, E.; PARTHASARATHY, S.; STEINBERG, D. Peroxidase-dependent metal-independent oxidation of low density lipoprotein in vitro: a model for in vivo oxidation? Proc. Natl. Acad. Sci. USA 90: 5929-5933, 1993.
- WINTERBOURN C.C.; MONTEIRO, H.P.; GALILEE, C.F. Ferritin-dependent lipid peroxidation by stimulated neutrophils: inhibition by myeloperoxidase-derived hypochlorous acid but not by endogenous lactoferrin. Biochim. Biophys. Acta 1055: 179-185, 1990.

- WINTERBOURN C.C.; VILLE, G.F.; MONTEIRO, H.P. Ferritin, lipid peroxidation and redox-cycling xenobiotics. Free Rad. Res. Comms. **12-13**: 107-114, 1991.
- WITZTUM, J.L. & STEINBERG, D. Role of oxidized LDL in atherogenesis. Br. Heart J. **69** (supl.): 512-522, 1993.
- WITZTUM, J.L. The oxidation hypothesis of atherosclerosis. Lancet **344**: 793-795, 1994.
- WODITSCH, I.; HOHLFELD, T.; STROBACH, H.; SCHÖR, J. Oral cicaprost protects from hypercholesterolemia-induced impairment of coronary vasodilation. Agents Actions **37** (supl.): 297-304, 1992.
- XIE, Q.W.; CHO, H.J.; CALAYCA, J.; MUMFORD, R.A.; SWIDEREK, K.M.; LEE, T.D.; DING, A.; TROSO, T.; NATHAN, C. Cloning and characterization of inducible nitric oxide synthase from mouse macrophages. Science **256**: 225-228, 1992.
- YANG, X.; CAI, B.; SCIACCA, R.R.; CANNON, P.J. Inhibition of inducible nitric oxide synthase in macrophages by oxidized low density lipoproteins. Circ. Res. **74**: 318-328, 1994.
- YLA-HERTTUALA, C. e cols. Evidence for the presence of oxidatively modified low density lipoprotein in atherosclerosis lesions of rabbit and man. J. Clin. Invest. **84**: 1086-1095, 1989.
- YOKOYAMA, M.; HIRATA, K.; MIYAKE, R.; AKITA, H.; YUICHI, I.; FUKUZAKI, H. Lysophosphatidylcholine: essential role in the inhibition of endothelium-dependent vasorelaxation by oxidized lipoprotein. Biochem. Biophys. Res. Commun. **168**: 301-308, 1990.
- ZAND, T.; NUNNARI, J.J.; HOFFMAN A.H.; SAVILONIS, B.J.; MACWILLIAMS, B.; MAJNO, G.; JORIS, I. Endothelial adaptations in aortic stenosis. Correlation with flow parameters. Am. J. Pathol. **133**: 407-18, 1988.
- ZAWADSKI, Z.; MILNE, R.; MARCEL, Y.L. Cu^{2+} -mediated oxidation of dialysed plasma: effects on low and high density lipoproteins and cholesterol ester transfer protein. J. Lipid Res. **32**: 243-240, 1991.
- ZHANG, H.; BASRA, H.J.K.; STEINBRECHER, V.P. Effects of oxidatively modified low density lipoprotein on cholesterol esterification in culture macrophages. J. Lipid Res. **31**: 1361-1369, 1990.
- ZHOU, G. & JAUHIANEN, M. Human plasma lecithin:cholesterol acyltransferase. Preparation and use of immobilized p-aminophenyl-arsenoxide as a catalytic site directed covalent ligand in enzyme purification. J. Chromatography **568**: 69-83, 1991.
- ZHU, W.; ROMA, P.; PELLEGATTA, F.; CATAPANO, A.L. Oxidized-LDL induce the expression of heat shock protein 70 in human endothelial cells. Biochem. Biophys. Res. Commun. **200**: 389-394, 1994.
- ZICHE, M.; MORBIDELLI, L.; MASISNI, E.; GRANGER, H.; GEPPETTI, P.; LEDDA, F. Nitric oxide promotes DNA synthesis and cyclic GMP formation in endothelial cells from postcapillary venules. Biochem. Biophys. Res. Commun. **192**: 1198-1203, 1993.
- ZILVERSMITH, D.B. Atherogenesis: a postprandial phenomenon. Circulation **60**: 473-485, 1979.

RESUMO

As modificações oxidativas das lipoproteínas representam um evento crucial para a aterogênese. Nesta área de investigação alguns dos principais pontos a serem esclarecidos são os seguintes: 1) quais são os processos geradores de espécies oxidantes envolvidos na oxidação das lipoproteínas, *in vivo*; 2) como as lipoproteínas oxidadas interagem com as células que atuam no processo aterosclerótico; 3) quais os efeitos das modificações oxidativas das lipoproteínas nas vias metabólicas dessas partículas, que estariam relacionadas à aterogênese e 4) qual é a influência das hiperlipoproteinemias na modificação oxidativa das lipoproteínas e nos sistemas de defesa antioxidante do organismo. As contribuições para o esclarecimento de cada um dos itens citados foram :

1) A oxidação da lipoproteína de baixa densidade por neutrófilos estimulados, na presença de ferritina, pode ser um processo potencial para a oxidação das lipoproteínas, *in vivo*.

2) A interação de células endoteliais com a lipoproteína de baixa densidade oxidada induz a liberação de óxido nítrico. A maior geração deste radical livre pode facilitar a formação de peroxinitrito, que é um potente agente oxidante.

3) A β -lipoproteína de densidade muito baixa modificada por peroxinitrito é internalizada por monócitos/macrófagos, podendo ,potencialmente, contribuir para a formação do ateroma. No entanto, estudos de biodistribuição e dados biocinéticos indicam que as lipoproteínas modificadas com peroxinitrito são rapidamente eliminadas do plasma, enquanto que a oxidação catalizada por cobre resulta em partículas lipoprotéicas que se acumulam nas lesões ateroscleróticas. A oxidação das lipoproteínas afeta processos importantes do metabolismo intravascular dessas partículas, tais como, a esterificação do colesterol plasmático e os processos de transferência de ésteres de colesterol entre as lipoproteínas plasmáticas. Isto pode resultar em alterações do transporte reverso do colesterol com acúmulo de colesterol nos tecidos periféricos e enriquecimento de lipoproteínas aterogênicas com ésteres de colesterol, o que teria implicações diretas para a aterogênese.

4) Em indivíduos com hiperlipidemia combinada ocorre menor resistência da lipoproteína de baixa densidade e suas subfrações à oxidação, diminuição de componentes do sistema de defesa antioxidante e maior geração de espécies reativas de oxigênio por monócitos e neutrófilos do sangue periférico. Na hipercolesterolemia induzida em coelhos por dieta rica em colesterol, há um aumento acentuado de lipoproteínas oxidadas no plasma sanguíneo, em paralelo ao aumento do colesterol plasmático e diminuição da isoenzima da superóxido dismutase extracelular ligada ao endotélio vascular. Em conjunto, esses dados indicam a presença de um estresse oxidativo nas hiperlipoproteinemias.

ABSTRACT

Lipoprotein oxidative modification is a crucial event in development of atherosclerosis. Some of the important points to be elucidate in this issue are : 1) which are the processes that possibly participate in lipoprotein oxidation, *in vivo*; 2) how can oxidized lipoproteins interact with cells participating in atherosclerotic process; 3) which are the effects of lipoprotein oxidation on steps of lipoprotein's intravascular metabolism related to atherogenesis; 4) what is the influence of hyperlipoproteinemias on lipoprotein oxidative modification and the antioxidant defense system ? The following findings may contribute to elucidate these points:

1) Low density lipoprotein oxidation by stimulated neutrophils, in the presence of ferritin, may be a potential *in vivo* process for lipoprotein oxidative modification.

2) Oxidized low density lipoprotein induces the release of nitric oxide by endothelial cells. The greater generation of this free radical by endothelium may facilitate peroxynitrite formation, that is a strong oxidant agent.

3) Peroxynitrite-modified β -very low density lipoprotein is internalized by monocyte/macrophages and may potentially contribute to foam cell generation. However, biodistribution studies and biokinetic data indicate that peroxynitrite-modified lipoproteins are rapidly cleared from plasma while copper-oxidized lipoproteins accumulate in atherosclerotic lesions. Lipoprotein oxidation affects important processes of lipoprotein intravascular metabolism, such as cholesterol esterification and cholesteryl ester transfer among lipoproteins. These effects may alter the reverse cholesterol transport promoting cholesterol deposition in peripheral tissues and increasing the cholesteryl ester content of atherogenic lipoproteins.

4) In subjects with combined hyperlipidemia the low density lipoprotein, as well as its heavy and light subfractions are less resistant to oxidation. Moreover, some components of their antioxidant defense system are decreased and the monocytes and neutrophils isolated from their peripheral blood release high amounts of reactive oxygen species. In addition, a sharp increase of oxidized lipoproteins occurs in blood plasma of cholesterol-fed rabbits as their plasma cholesterol levels rise and the superoxide dismutase isoenzyme attached to endothelial cells decreases. Taken together these data suggest the occurrence of an oxidative stress in hyperlipoproteinemias.

ANEXOS

TRABALHOS QUE DERAM ORIGEM À TESE

Abdalla, D.S.P.; Campa, A. & Monteiro, H.P. Low density lipoprotein oxidation by stimulated neutrophils and ferritin. *Atherosclerosis* 97: 149-159, 1992. (TRABALHO 1)

Abdalla, D.S.P.; Costa-Rosa, L.F.B.; Monteiro, H.P.; Campa, A. & Curi, R. Human macrophage metabolism of low density lipoprotein oxidized by stimulated neutrophils and ferritin. *Atherosclerosis* 107: 157-163, 1994. (TRABALHO 2)

Silva, E.L.; Coelho, I.J.C.; Meneguetti, J.C.; & Abdalla, D.S.P. Biodistribution and fractional clearance rate of copper- and peroxynitrite-oxidized ^{99m}Tc - β -VLDL in cholesterol-fed rabbits. *J. Lipid Res.* (submetido) (TRABALHO 3)

Silva, E.L.; Moriel, P.; Chang, Y.H. & Abdalla, D.S.P. Plasma antioxidant enzymes and oxidized lipoproteins in hypercholesterolemic rabbits. *Biochem. Mol. Biol. Inter.* (submetido) (TRABALHO 4)

Silva, E.L. & Abdalla, D.S.P. Uptake of peroxynitrite modified β -VLDL by monocytes of normolipidemic and hypercholesterolemic rabbits. *Biochem. Biophys. Res. Commun.* (submetido) (TRABALHO 5)

Miara, J.; Maranhão, R.C. & Abdalla, D.S.P. Esterification of cholesterol and lipid transfer in human serum. Effect of lipid peroxidation. In *Oxygen Radicals*. Yagi, K.; Kondo, E., Niki, E. & Yoshikawa, T. (eds.). Elsevier Science Publishers B.V., Amsterdam, 1992, pp. 293-96. (TRABALHO 6)

Maranhão, R.C.; Miara, J. & Abdalla, D.S.P. Lipid transfer and cholesterol esterification in oxidized emulsions and lipoproteins. *Biochim. Biophys. Acta* (submetido)
(TRABALHO 7)

Barbosa, D.S.; Maranhão, R.C.; Araújo, F.B.; Chang, Y.H.; Hirata, M.H. & Abdalla, D.S.P. Sialic acid and oxidizability of low density lipoprotein subfractions of hyperlipidemic patients. *Clin. Biochem.* (em publicação) **(TRABALHO 8)**

Araujo, F.B.; Barbosa, D.S.; Chang, Y.H.; Maranhão, R.C. & Abdalla, D.S.P. Evaluation of oxidative stress in patients with hyperlipidemia. *Atherosclerosis* (em publicação)
(TRABALHO 9)

TRABALHO 1

ATHERO 04920

Low density lipoprotein oxidation by stimulated neutrophils and ferritin

Dulcinea S.P. Abdalla^a, Ana Campa^a and Hugo P. Monteiro^b

^a*Faculdade de Ciências Farmacêuticas, Universidade de São Paulo and* ^b*Fundação Hemocentro de São Paulo, SP (Brazil)*

(Received 22 November, 1991)

(Revised, received 4 August, 1992)

(Accepted 5 August, 1992)

Summary

Low density lipoprotein (LDL) oxidation mediated by phorbol myristate acetate (PMA)- and formylmethionylleucylphenylalanine (FMLP)-stimulated human neutrophils was enhanced by 70% in the presence of ferritin. Iron released from ferritin by the superoxide anion generated in the respiratory burst of stimulated neutrophils is shown to be involved in lipoprotein oxidation. Ascorbate (100 μ M), superoxide dismutase (10 μ g/ml) and uric acid (430 μ M) showed inhibitory effects of 10%, 70% and 50% on LDL oxidation, respectively. Ceruloplasmin (2.7 μ M) potentiated LDL oxidation by stimulated neutrophils and ferritin, both alone and in the presence of methionine. Methionine (1 mM) and catalase (30 μ g/ml) increased LDL oxidation by stimulated neutrophils and ferritin. These data suggest that LDL oxidation by stimulated neutrophils and ferritin may be relevant in inflammation when both neutrophils and ferritin are increased.

Key words: Ferritin; Neutrophils; LDL oxidation; Lipid peroxidation; Antioxidant

Introduction

Oxidation of low density lipoprotein (LDL) has been strongly implicated in the pathogenesis of atherosclerosis [1–3]. Arterial endothelial cells, monocytes and monocyte-derived macrophages can oxidatively modify LDL [4–7]. Modified LDL

is taken up rapidly by macrophages by means of the scavenger receptor, and this process has been proposed to explain the origin of the lipid-laden foam cells, the initial stage of atherosclerotic lesions [4,8,9]. Human neutrophils can also oxidize LDL, making it cytotoxic [10]. These cells are well known for their ability to produce superoxide (O_2^-) and other reactive oxygen species as an essential feature of their microbicidal and cytotoxic actions [11,12]. Although neutrophils could damage biological targets through lipid peroxida-

Correspondence to: Dr. Dulcinea S.P. Abdalla, Faculdade de Ciências Farmacêuticas, Universidade de São Paulo, CP 66355, CEP 05389, São Paulo, SP, Brazil. Fax: 55-11-815-3575.

tion [10,13–15] they do not contain appreciable amounts of endogenous iron that can catalyse lipid peroxidation [15,16]. Transition metals, particularly iron, and a reducing agent such as O_2^- are absolute requirements for the occurrence of lipid peroxidation in *in vitro* biological systems [17,18]. Therefore, lipid peroxidation mediated by neutrophils was observed only when exogenous iron was added to the system [15].

Such requirements imply that for neutrophils to promote lipid peroxidation under physiological conditions they must acquire an exogenous source of iron catalysts. Iron bound to lactoferrin or transferrin does not promote lipid peroxidation [19,20] and ferritin has been suggested as a possible iron source [21]. Biemond et al. [22] were the first to show that O_2^- generated by stimulated human neutrophils can release iron from ferritin. Furthermore, Winterbourn et al. [23] have demonstrated that human neutrophils stimulated with phorbol myristate acetate (PMA) or formylmethionylleucylphenylalanine (FMLP) released iron from ferritin and were able to use ferritin iron to promote lipid peroxidation. Products derived from fatty acid peroxidation have been associated with several disease states [24], and modification of LDL by cellular systems involves lipid peroxidation [3,5,6,8]. Thus, the present study was undertaken to examine the ability of stimulated neutrophils to oxidize LDL in the presence of ferritin as the source of iron. We showed that these cells extensively oxidize LDL in the presence of ferritin. Antioxidants such as catalase, methionine and ceruloplasmin were not protective against LDL oxidation by stimulated neutrophils and ferritin. In addition, ascorbate and uric acid, two important plasma antioxidants [25,26], afforded only partial protection.

Materials and Methods

Materials

Ferritin from horse spleen (46% iron saturated), superoxide dismutase (SOD) from bovine erythrocytes, ceruloplasmin from human plasma, aprotinin from bovine lungs, butylated hydroxytoluene (BHT), Ficoll, Dextran, PMA, FMLP, ascorbate, methionine, cholesteryl linoleate, ferrozine, EDTA, diethylenetetraminepentacetic acid

(DTPA), benzamidine and phenylmethylsulphonyl fluoride (PMSF) were from Sigma Chemical Co. (St. Louis, MO). Catalase was from Pharmacia (Uppsala, Sweden), Hypaque was from Sydney Ross Co. (S. Paulo, Brazil) and desferrioxamine from Ciba-Geigy (Basel, Switzerland). All other chemicals were of the purest analytical grade and were purchased from Merck (Darmstadt, Germany).

Ceruloplasmin was diluted in phosphate-buffered saline (PBS), pH 7.3 and passed through a Sepharose 6B column in the FPLC system (Pharmacia-LKB Biotechnology Products, Uppsala, Sweden) to separate the pure protein from other contaminants. Protein purity was determined from the A_{610}/A_{280} ratio, based on the value of $A_{610}/A_{280} = 0.044$ for 100% ceruloplasmin [27]. The estimated purity of the preparation was 99%. In some experiments, ferritin was purified on a Sephadex G-50 column eluted with PBS (pH 7.3), containing 1 mM EDTA. Solutions were prepared in highly purified water obtained via reverse osmosis in the Milli Rho System and subsequently deionized in the Milli Q System, yielding resistivity values of 18.1 M Ω /cm. Contaminant iron in the buffers was determined using ascorbate as a test for the presence of catalytic metals [28] and was found to be $<0.3 \mu\text{M}$.

Human neutrophil isolation

Neutrophils were isolated from freshly donated, heparinized peripheral blood by Ficoll-Hypaque centrifugation, dextran centrifugation of red cells and hypotonic lysis of remaining red cells [29]. Neutrophils were suspended in PBS (pH 7.3), containing CaCl_2 (1 mM), MgCl_2 (0.5 mM) and glucose (1 mg/ml).

LDL preparation

LDL was isolated from blood plasma from healthy donors in the presence of EDTA (1 mg/ml), benzamidine (2 mM) and PMSF (1 mM) to prevent both autooxidation and protein degradation by plasma endogenous proteases. The lipoprotein was isolated from plasma by ultracentrifugation. Plasma density was adjusted to 1.019 with KBr. After centrifugation for 24 h at 40 000 rev./min and 4°C in an ultracentrifuge using a RP 55T rotor (Hitachi Kobi Co. Ltd., Tokyo, Japan)

the upper layer was removed. The pooled infranant was adjusted to 1.063 density by adding KBr and was purified by ultracentrifugation at the same density and further dialyzed against 0.9% NaCl plus 1 mM DTPA overnight. Total protein content of the LDL preparation was determined according to the Lowry procedure [30], using albumin as standard and 0.05% deoxycholate. The final protein concentration of LDL was 5–7 mg/ml. The purity of the LDL fraction was assayed by agarose gel electrophoresis [31].

Iron release from ferritin

Ferrozine (200 μ M), various concentrations of ferritin and neutrophils ($1-4 \times 10^6$ cells/ml) were incubated in PBS at 37°C. The cells were stimulated with PMA (100 ng/ml) or FMLP (0.25 μ M) and absorbance at 562 nm was continuously monitored up to 30 min. Concentrations of Fe^{2+} -ferrozine were determined using $\epsilon_{562} = 26\,900 \text{ M}^{-1} \text{ cm}^{-1}$ [32]. All kinetic measurements were performed using a Beckman DU-70 spectrophotometer.

Superoxide production by stimulated neutrophils

Initial rates and total production of O_2^- for 30 min by stimulated neutrophils (1×10^6 cells/ml) were determined by measuring reduction of cytochrome *c* (100 μ M) at 550 nm using $\epsilon_{550} \text{ reduced-oxidized} = 21\,100 \text{ M}^{-1} \text{ cm}^{-1}$ in the presence or absence of 30 $\mu\text{g/ml}$ SOD at 37°C in PBS [33].

Lipid peroxidation

Neutrophils ($1 \times 10^6/\text{ml}$) stimulated with PMA (100 ng/ml) or with FMLP (0.25 μ M) were incubated with LDL (1 mg/ml) and ferritin (200 $\mu\text{g/ml}$) in a final volume of 1 ml, at 37°C, in a Dubnoff shaker for 17 h. In some experiments, different additives were added as described in the figure legends and tables. Lipid peroxidation in the incubation mixtures was monitored by two different procedures.

TBARS assay

LDL peroxidation was determined in 0.25-ml aliquots from the incubation mixtures by the thiobarbituric acid reactive substances (TBARS) procedure [34]. BHT (0.03%) was added to the

TBA reagent to prevent spurious lipid peroxidation during the assay. Malondialdehyde equivalents were estimated using $\epsilon_{532} = 1.56 \times 10^5 \text{ M}^{-1} \text{ cm}^{-1}$ [35].

Lipid hydroperoxides

Formation of cholesteryl ester hydroperoxides (CE-OOH) in LDL was followed by high performance liquid chromatography (HPLC) with UV detection. The HPLC system consisted of an HPLC pump (Waters, Milford, MA; model 510), a spectrophotometric detector (Waters, Milford, MA; model Lambda-Max 481), an injector (Rheodyne, Cotati, CA; model 7125), a data module (Waters, Milford, MA; model 745 B) and a C8 Inertsil column (GL Sciences Inc., Japan; 4.6 i.d. $\times$ 150 mm; 5 μm particle size). Cholesteryl linoleate hydroperoxide (Ch18:2OOH) to be used as HPLC standard was prepared by methylene blue-sensitized photooxidation of cholesteryl linoleate [36]. The hydroperoxide was purified on a C18 column (4.6 i.d. $\times$ 25 mm, 5 μm particle size, Waters) using methanol. Ch18:2OOH concentration was calculated by the weight of the purified fraction. The concentration of cholesteryl esters in the samples was determined by comparison of the peak areas monitored at 235 nm with that of Ch18:2OOH. Samples (1 ml of the incubation mixtures) were extracted with 10 ml of methanol/hexane (1:10). After evaporation of the hexane layer, the residue was dissolved in methanol and the presence of CE-OOH was investigated by HPLC with methanol/water (97:3, flow rate = 1 ml/min) as the mobile phase. The cholesterol content of methanol extracts was determined by an enzymatic method using commercial kits (Cholesterol CHODPAP, Merck S/A, RJ, Brazil) and CE-OOH levels are reported in relation to cholesterol content in each sample extract.

Results

Iron release from ferritin by stimulated neutrophils

PMA stimulated neutrophils released iron from ferritin in the form of an Fe^{2+} -ferrozine complex. Initial rates remained constant during the first 10 min, then gradually declined to zero by about 20 min (Fig. 1). FMLP-stimulated neutrophils also released iron from ferritin (Fig. 1), but to a lesser

Fig. 1. Time course of iron release from ferritin by PMA- and FMLP-stimulated neutrophils. The reaction mixtures containing ferrozine (200 μ M), ferritin (200 μ g/ml), neutrophils (1×10^6 /ml) and PMA (100 ng/ml) or FMLP (0.25 μ M) were incubated at 37°C in spectrophotometer cuvettes. Results are means $\pm$ S.D. of duplicates from 3 typical assays. Initial rates of O_2^- production by PMA- and FMLP-stimulated neutrophils were 5.0–7.2 and 1.8–2.4 nmol O_2^- /min per 10^6 cells/ml, respectively.

Fig. 2. Effect of cell number on LDL oxidation and on iron release from ferritin by PMA-stimulated neutrophils. The reaction mixtures containing 200 μ g/ml ferritin, neutrophils, 100 ng/ml PMA, 200 μ M ferrozine (iron release) or 1 mg/ml of LDL protein (LDL oxidation) were incubated at 37°C for 30 min (iron release) or 17 h (LDL oxidation). Results are means $\pm$ S.D. of duplicates performed with the same neutrophil preparation.

Fig. 3. Effect of ferritin concentration on LDL oxidation and on iron release from ferritin by PMA-stimulated neutrophils. Reaction mixtures containing ferritin, 1×10^6 neutrophils/ml, 100 ng/ml PMA, 200 μ M ferrozine (iron release) or 1 mg/ml of LDL protein (LDL oxidation) were incubated at 37°C for 30 min (iron release) or 17 h (LDL oxidation). Results are means $\pm$ S.D. of triplicates.

extent than detected with PMA. Initial rates of iron release remained constant only during the first 5 min, declining to zero thereafter. The kinetics of neutrophil-dependent iron release

TABLE 1

EFFECT OF ADDITIVES ON IRON RELEASE FROM FERRITIN BY PMA-STIMULATED NEUTROPHILS

The reaction mixtures containing ferritin (200 μ g/ml), ferrozine (200 μ M), PMA (100 ng/ml), neutrophils (1×10^6 /ml) and the indicated additives were incubated at 37°C for 30 min. Results are means $\pm$ S.D. of duplicates from 2 or 3 typical experiments.

Addition	Iron release (nmol Fe ²⁺ /ml per 1×10^6 cells)	Percent of control
None	2.22 $\pm$ 0.64	100
Methionine 1 mM	4.32 $\pm$ 0.88	162
Catalase 30 μ g/ml	3.65 $\pm$ 0.15	137
Ascorbate 100 μ M	1.30 $\pm$ 0.41	49
Uric acid 430 μ M	0.45 $\pm$ 0.05	17
SOD 10 μ g/ml	0	0
Ceruloplasmin 400 μ g/ml	0	0

observed with PMA and FMLP paralleled the oxidative burst and O_2^- production produced by both stimuli. Superoxide generation by the cells and ferritin-iron mobilization were maximal within the first hour of incubation. Within this period, PMA-stimulated neutrophils produced 65.9 ± 12.9 nmol $O_2^-/10^6$ cells and mobilized 4.9 ± 2.2 nmol $Fe^{2+}/10^6$ cells ($n = 15$) and FMLP-stimulated cells generated 7.1 ± 1.7 nmol $O_2^-/10^6$ cells with mobilization of 0.8 ± 0.2 nmol $Fe^{2+}/10^6$ cells ($n = 12$). The efficiency of iron release per generated O_2^- was 7.4% and 11.2% for PMA- and FMLP-stimulated neutrophils, respectively. In agreement with previous observations [23], there was a dependence of ferritin iron release on cell number (Fig. 2) and ferritin concentration (Fig. 3).

The effect of plasma antioxidants and catalase on iron release from ferritin by stimulated neutrophils was also determined (Table 1). Addition of catalase to PMA-stimulated neutrophils consistently increased the amount of Fe^{2+} -ferrozine in solution [23]. Methionine, an effective scavenger of hypochlorous acid (HOCl), also increased the yield of detectable Fe^{2+} -ferrozine, probably by protecting released ferrous iron

against reoxidation by HOCl. Ascorbate, uric acid and ceruloplasmin, at concentrations normally found in plasma, inhibited iron release from ferritin by PMA-stimulated neutrophils. Ascorbate can be oxidized by O_2^- [37] and consequently will compete with ferritin iron for O_2^- generated by stimulated phagocytes, leading to a decrease in the amount of iron released. Uric acid can form coordination complexes with iron [38] and compete with ferrozine for iron, resulting in lesser amounts of detectable Fe^{2+} -ferrozine in solution. Although ceruloplasmin scavenges O_2^- [39], its inhibitory effect on ferritin iron release has been reported to be greater than its scavenging capacity [39]. Thus, ceruloplasmin inhibition can be attributed to its ferroxidase activity and to its ability to reincorporate released ferritin-iron back into ferritin [39]. Superoxide dismutase completely inhibited iron release from ferritin by PMA-stimulated neutrophils (Table 1). In contrast, benzamidine (2 mM) and PMSF (1 mM), used as protease inhibitors during LDL isolation, inhibited iron release by 75% and 5%, respectively.

TABLE 2

LDL OXIDATION BY PMA- AND FMLP-STIMULATED NEUTROPHILS

Concentrations in the systems were: 1 mg/ml of LDL protein, 1×10^6 /ml neutrophils, 100 ng/ml PMA, 0.25 μ M FMLP and 200 μ g/ml ferritin. Reaction mixtures were incubated in a shaker bath, under air, at 37°C for 17 h. Results are means $\pm$ S.D. of duplicates from 3 experiments performed with different LDL and neutrophil preparations. LDL oxidation was measured as described in Materials and Methods. N.D., not determined.

System	TBARS nmol MDA equivalents/mg LDL protein	CE-OOH nmol/mg cholesterol
LDL	0.324 $\pm$ 0.054	159 $\pm$ 21
LDL/ferritin	0.673 $\pm$ 0.105	208 $\pm$ 27
LDL/neutrophils	0.356 $\pm$ 0.076	194 $\pm$ 22
LDL/neutrophils/PMA	0.778 $\pm$ 0.074	178 $\pm$ 25
LDL/neutrophils/ PMA/ferritin	1.640 $\pm$ 0.255	307 $\pm$ 51
LDL/neutrophils/ FMLP	0.311 $\pm$ 0.036	N.D.
LDL/neutrophils/ FMLP/ferritin	1.090 $\pm$ 0.403	N.D.

LDL oxidation by stimulated neutrophils

LDL peroxidation was slightly increased by PMA- and FMLP-stimulated neutrophils or by ferritin alone. However, there was a substantial increase in TBARS and CE-OOH formation when LDL was oxidized by stimulated neutrophils in the presence of ferritin (Table 2). LDL oxidation increased with increasing cell number (Fig. 2), ferritin concentration (Fig. 3) and LDL concentration (1.05 ± 0.60 , 2.10 ± 0.08 , 2.60 ± 0.10 and 3.00 ± 0.15 nmol MDA equivalents/mg LDL protein for 1, 2, 3 and 4 mg LDL/ml, respectively).

Addition of apotransferrin to PMA-stimulated neutrophils plus ferritin resulted in a concentration dependent inhibition of LDL oxidation (Table 3). Desferrioxamine (1 mM) added to the complete system (stimulated neutrophils, ferritin and LDL) immediately prior to incubation or 1 h after starting incubation inhibited LDL oxidation by 75% and 65%, respectively. This inhibition indicates a requirement for iron release from ferritin in the oxidation of LDL by this system. Addition of ferrous iron at equimolar concentrations to those released from ferritin by PMA-stimulated neutrophils induced less formation of TBARS from LDL (complete system, 2.22 ± 0.6 and 5 μ M

TABLE 3

EFFECT OF APOTRANSFERRIN ON LDL OXIDATION BY PMA STIMULATED NEUTROPHILS PLUS FERRITIN

The basic reaction mixtures containing 1 mg/ml LDL protein, 1×10^6 /ml neutrophils, 100 ng/ml PMA and 200 μ g/ml ferritin were incubated in a shaker bath under air at 37 °C for 17 h. Results are means $\pm$ S.D. of duplicates from 2 experiments performed with different LDL and neutrophil preparations. LDL oxidation was measured as described in Materials and Methods.

Addition	TBARS nmol MDA equivalents/ mg LDL protein	% inhibition
None	2.311 $\pm$ 0.668	
Apotransferrin		
0.5 mg/ml	1.970 $\pm$ 0.386	15
1.0 mg/ml	1.849 $\pm$ 0.351	20
2.0 mg/ml	1.775 $\pm$ 0.443	23
3.0 mg/ml	1.673 $\pm$ 0.065	28

Fig. 4. Effect of additives on LDL oxidation by PMA-stimulated neutrophils in the presence of ferritin. The basic reaction mixtures containing 1 mg/ml LDL protein, 1×10^6 neutrophils/ml, 200 μ g/ml ferritin and 100 ng/ml PMA were incubated in a shaker, under air, at 37°C for 17 h. Results represent means $\pm$ S.D. of duplicates from 2 typical experiments performed with different neutrophil and LDL preparations. Control system: LDL + neutrophils + PMA + ferritin. LDL oxidation was measured as described under Materials and Methods. Concentrations of additives were catalase (30 μ g/ml), desferrioxamine (100 μ M), SOD (10 μ g/ml), ascorbate (100 μ M) and uric acid (430 μ M).

Fe^{2+} , 0.70 ± 0.34 nmol MDA equivalents/mg LDL protein), indicating that iron by itself is not a good catalyst of LDL oxidation.

Catalase enhanced LDL oxidation by PMA-stimulated neutrophils plus ferritin (Fig. 4). Although neutrophil stimulation with PMA preferentially releases secondary granules (rich in lactoferrin), with a minor release of primary granules (rich in myeloperoxidase) [40], this was sufficient to convert some of the hydrogen peroxide produced during stimulation into HOCl. Hypochlorous acid has been reported to chlorinate lipids in phospholipid liposomes, so that peroxidation is inhibited [23]. Therefore, catalase scavenging hydrogen peroxide reduced HOCl formation and enhanced LDL lipid peroxidation. Methionine also enhanced LDL oxidation by scavenging HOCl (Table 4). The effects of SOD, ascorbate and uric acid were also reported (Fig. 4). These three antioxidants provided protection against LDL oxidation by PMA-stimulated neutrophils plus ferritin in accordance with their

inhibitory actions on iron release from ferritin. Ceruloplasmin, in contrast to its effect on iron release, enhanced LDL oxidation (Table 4). Ceruloplasmin and methionine presented additive effects on LDL oxidation by stimulated neutrophils plus ferritin (Table 4). Copper ions released from ceruloplasmin during the incubation period are probably responsible for the observed enhancement in LDL oxidation. For comparison LDL oxidation by copper(II) is shown in Table 4. Furthermore, lipoprotein-depleted plasma (LDP, 10 mg of protein/ml) reduced LDL oxidation by 50% (control, 1.640 nmol MDA equivalents/mg protein and LPD, 0.810 nmol MDA/mg protein).

Discussion

In this investigation we showed that LDL lipid peroxidation promoted by stimulated neutrophils was enhanced in the presence of ferritin. This enhancement was dependent on O_2^- -induced fer-

TABLE 4

EFFECT OF METHIONINE AND CERULOPLASMIN ON LDL OXIDATION BY PMA-STIMULATED NEUTROPHILS PLUS FERRITIN

The basic reaction mixtures (complete system), containing 1 mg/ml LDL protein, 200 μ g/ml ferritin, 200 μ M ferrozine, 100 ng/ml PMA and 1×10^6 cells/ml were incubated at 37°C for 17 h. Where indicated, 1 mM methionine and 2.7 μ M ceruloplasmin were added. Results are means $\pm$ S.D. of duplicates of the same neutrophil and LDL preparations in each series of experiments, but TBARS and CE-OOH were determined with different neutrophil and LDL preparations. LDL oxidation was measured as described in Materials and Methods. N.D., not determined.

System	TBARS nmol MDA equivalents/ mg LDL protein	CE-OOH nmol/mg choles- terol
Complete	2.210 $\pm$ 0.008	307 $\pm$ 51
Complete/methionine	2.674 $\pm$ 0.228	470 $\pm$ 71
Complete/ceruloplasmin	2.395 $\pm$ 0.325	544 $\pm$ 82
Complete/methionine/ ceruloplasmin	5.039 $\pm$ 0.153	N.D.
LDL/cells/methionine/ ceruloplasmin	3.027 $\pm$ 0.180	N.D.
LDL/5 μ M Cu ²⁺	5.076 $\pm$ 0.172	N.D.

ritin iron release, as can be concluded from inhibitions caused by SOD and by the iron chelator desferrioxamine. The initiator species in the process was not defined but iron has proved to be important. In its reduced form, iron is an efficient catalyst of hydroxyl radical formation. However, the observed stimulatory effects of catalase on LDL peroxidation by stimulated neutrophils and ferritin ruled out any role of hydroxyl radical as the initiator species in this process. Alternatively, ferrous and ferric iron may form a ternary complex with oxygen, which has been proposed to initiate lipid peroxidation [17]. The sharp inhibition of CE-OOH production by desferrioxamine suggests that iron did not act as a secondary initiator, catalyzing the decomposition of lipid hydroperoxides. In contrast to a previous report [41], neutrophils in the absence of a triggering stimulus did not promote significant LDL lipid peroxida-

tion. This observation emphasizes the role of oxygen radicals generated by stimulated neutrophils in the promotion of LDL oxidation.

Ferritin is a high molecular weight iron-storage protein as well as an acute phase reactant. In clinical disorders of iron balance, the levels of ferritin reflect the levels of body iron stores [42]. In inflammation, hepatic diseases and increased red cell turnover, serum levels of ferritin are elevated [43]. Although under normal conditions the predominant form of circulating ferritin is a glycosylated iron-poor isoform, other isoforms can be released from damaged tissues or from phagocytic cells, increasing the amount of isoforms with higher iron content in blood plasma [44]. Furthermore, in chronic inflammatory diseases and in infectious diseases, the levels of ferritin in inflamed tissues and in phagocytic cells are also elevated [45,46]. In our studies we found that ferritin by itself is not a good promoter of LDL oxidation. Iron ions in the absence of oxygen radicals are also inefficient ([47], this work). Additionally, release of ferritin iron requires a reducing species, represented in our system by O₂⁻. Thus, an interaction between oxygen radicals produced by stimulated neutrophils and iron released from ferritin is needed to promote LDL oxidation.

In a physiologic inflammatory response, neutrophils do not continuously degrade host tissues. Influx of neutrophils, along with their subsequent triggering, is carefully regulated and ceases when the initiating stimulus is eliminated [48]. However, if inflammatory stimuli are chronically directed against host tissues or are not properly down-regulated, neutrophils can be a continuous source of damaging oxygen radicals. In such a situation, ferritin released from damaged or ingested cells may be accessible to O₂⁻ produced by neutrophils and iron release would be favored. Consequently, LDL oxidation may occur at these inflammatory sites. The importance of metal ions, particularly iron, as catalysts of lipid peroxidation in vitro is well known [17,18]. Additionally, evidence supporting a role for iron in oxidative damage in biological systems has been accumulated in the literature: hepatic lipid peroxidation in vivo was observed in rats with chronic iron overload [49]. Houghlum et al. [50] showed the presence of malondialdehyde and 4-hydroxynon-

enal protein adducts in plasma and liver of rats with iron overload, and recently, Burkitt and Mason [51] presented evidence on the *in vivo* generation of hydroxyl radical in acute iron overload induced in rats. Although these data represent a strong support for a role of iron in *in vivo* lipid peroxidation, additional studies with different systems would be necessary to reinforce this point.

The mechanism by which LDL becomes oxidized *in vivo* is not well understood. In hyperlipidemic subjects, neutrophils present higher oxidative metabolism, and the rate of SOD-inhibitable O_2^- production correlates positively with LDL cholesterol concentrations [52]. Thus, the neutrophils of hyperlipidemic subjects, by producing higher amounts of oxidants, could release iron from ferritin present in the surrounding tissues or in the circulation, leading to elevated plasma levels of oxidized LDL.

The occurrence of lipoprotein oxidation in blood plasma has been questioned due to the presence of water-soluble antioxidants in this fluid. However, under our experimental conditions, physiological concentrations of ascorbate and uric acid, two important antioxidants present in blood plasma [25,26] did not fully prevent LDL oxidation. In relation to ascorbate, our results contrast with others in the literature showing a strong inhibition of LDL oxidation by ascorbate [53,54]. The observed poor efficiency of ascorbate in inhibiting LDL oxidation is possibly a result of iron catalysed oxidation of ascorbate [55]. Iron released from ferritin could react with ascorbate generating dehydroascorbate and other products of ascorbate oxidation. Such products would be less efficient than ascorbate itself, as antioxidants against LDL oxidation. Inhibition by SOD also occurred, but it is known that the activity of this enzyme in extracellular fluid is low [56]. The iron-binding protein apotransferrin, also present in blood plasma, was poorly inhibitory. Recently, Winterbourn and Molloy [57] have shown that transferrin incubated with the myeloperoxidase- H_2O_2 -Cl system underwent destruction of the iron binding site. Thus, myeloperoxidase released by stimulated neutrophils may impair the ability of transferrin to take up the iron released from ferritin and prevent lipid peroxidation [20].

Ceruloplasmin, also considered to be a plasma antioxidant, enhanced LDL oxidation by stimulated neutrophils and ferritin, possibly because of the release of copper ions from this protein. Again, oxidants generated from the myeloperoxidase system may be responsible for oxidative inactivation of the copper binding sites.

In conclusion, the system described here may be associated with oxidative modifications of lipoproteins in hyperlipidemic patients, especially in chronic inflammatory situations where there is an increase in production of oxidants by inflammatory cells with consumption of extracellular antioxidants and an increase in ferritin levels.

Acknowledgements

We gratefully acknowledge FBB (Fundação Banco do Brasil) and FAPESP (Fundação de Amparo à Pesquisa do Estado de São Paulo) for financial support. We also wish to thank Dr. Christine Winterbourn for helpful comments on the manuscript and Mr. A.C. Magnanelli for purification of ceruloplasmin by FPLC.

References

- 1 Steinberg, D., Parthasarathy, S., Carew, T.E. and Witztum, J., Beyond cholesterol. Modifications of low-density lipoprotein that increase its atherogenicity, *N. Engl. J. Med.*, 320 (1989) 915.
- 2 Steinbrecher, U.P., Zhang, H. and Lougheed, M., Role of oxidatively modified LDL in atherosclerosis, *Free Rad. Biol. Med.*, 9 (1990) 155.
- 3 Jurgens, G., Hoff, H.F., Chisolm, G.M. and Esterbauer, H., Modifications of human serum low density lipoprotein by oxidation — characterization and pathophysiological implications, *Chem. Phys. Lipids*, 45 (1987) 315.
- 4 Henriksen, T., Mahoney, E.M. and Steinberg, D., Enhanced macrophage degradation of low density lipoprotein previously incubated with cultured endothelial cells: recognition by receptor for acetylated low density lipoproteins, *Proc. Natl. Acad. Sci. USA*, 78 (1981) 6499.
- 5 Morel, D.W., Dicorleto, P.E. and Chisolm, G.M., Endothelial and smooth muscle cells alter low density lipoprotein *in vitro* by free radical oxidation, *Arteriosclerosis*, 4 (1984) 357.
- 6 Hiramatsu, K., Rosen, H., Heinecke, J.M., Wolfbauer, G. and Chait, A., Superoxide initiates oxidation of low density lipoprotein by human monocytes, *Arteriosclerosis*, 7 (1987) 55.
- 7 Parthasarathy, S., Printz, D.J., Boyd, D., Joy, L. and

- Steinberg, D., Macrophage oxidation of low density lipoprotein generates a modified form recognized by the scavenger receptor, *Arteriosclerosis*, 6 (1986) 505.
- 8 Fogelman, A.M., Shechter, I., Seager, J., Hokon, M., Child, J.S. and Edwards, P.A., Malondyaldehyde alteration of low density lipoprotein leads to cholesteryl-ester accumulation in human monocyte-macrophages, *Proc. Natl. Acad. Sci. USA*, 77 (1980) 2214.
- 9 Henricksen, T., Mahoney, E.M. and Steinberg, D., Enhanced macrophage degradation of biologically modified low density lipoprotein, *Arteriosclerosis*, 3 (1983) 149.
- 10 Cathcart, M.K., Morel, D.W. and Chisolm, G.M., Monocytes and neutrophils oxidise low density lipoprotein making it cytotoxic, *J. Leukocyte Biol.*, 38 (1985) 341.
- 11 Babior, B.M., Oxygen-dependent microbial killing by phagocytes, *N. Engl. J. Med.*, 298 (1978) 659.
- 12 Fantone, J.C. and Ward, P., Role of oxygen-derived free radicals and metabolites in leukocyte-dependent inflammatory reactions, *Am. J. Pathol.*, 107 (1984) 397.
- 13 Claster, S., Chiu, D.T., Quintanilha, A. and Lubin, B., Neutrophils mediate lipid peroxidation in human red cells, *Blood*, 64 (1984) 1079.
- 14 Sepe, S.M. and Clark, R.A., Oxidant membrane injury by the neutrophil myeloperoxidase system. II Injury by stimulated neutrophils and protection by lipid-soluble antioxidants, *J. Immunol.*, 134 (1985) 1896.
- 15 Thomas, M.J., Shirley, P.S., Hendrick, C.C. and DeChatelet, L.R., Role of free radical processes in stimulated human polymorphonuclear leukocytes, *Biochemistry*, 25 (1986) 8042.
- 16 Britigan, B.E., Rosen, G.M., Chai, Y. and Cohen, M.S., Do human neutrophils make hydroxyl radical? Determination of free radicals generated by human neutrophils activated with a soluble or particulate stimulus using electron paramagnetic resonance spectroscopy, *J. Biol. Chem.*, 261 (1986) 4426.
- 17 Minotti, G. and Aust, S.D., The requirement for iron(III) in the initiation of lipid peroxidation by iron(II) and hydrogen peroxide, *J. Biol. Chem.*, 262 (1987) 1098.
- 18 Aust, S.D., Morehouse, L.A. and Thomas, C.E., Role of metals in oxygen radicals reactions, *J. Free Rad. Biol. Med.*, 1 (1985) 3.
- 19 Gutteridge, J.M.C., Paterson, S.K., Segal, A.W. and Halliwell, B., Inhibition of lipid peroxidation by the iron-binding protein lactoferrin, *Biochem. J.*, 199 (1981) 259.
- 20 Monteiro, H.P. and Winterbourn, C.C., The superoxide-dependent iron transfer from ferritin to transferrin and lactoferrin, *Biochem. J.*, 256 (1988) 923.
- 21 Gutteridge, J.M.C., Halliwell, B., Treffry, A., Harrison, P.M. and Blake, D., Effect of ferritin-containing fractions with different iron loading on lipid peroxidation, *Biochem. J.*, 209 (1983) 557.
- 22 Biemond, P.E., Van Eijik, H.G., Swaak, A.J.G. and Koster, J.F., Iron mobilization from ferritin by superoxide derived from stimulated polymorphonuclear leukocytes: possible mechanism in inflammatory diseases, *J. Clin. Invest.*, 73 (1984) 1576.
- 23 Winterbourn, C.C., Monteiro, H.P. and Galilee, C.F., Ferritin dependent lipid peroxidation by stimulated neutrophils: inhibition by myeloperoxidase-derived hypochlorous acid but not by endogenous lactoferrin, *Biochim. Biophys. Acta*, 1055 (1990) 179.
- 24 Yagi, K., Lipid peroxides and human diseases, *Chem. Phys. Lipids*, 45 (1987) 337.
- 25 Wasil, M., Halliwell, B., Hutchison, D.C.S. and Baum, H., The antioxidant action of human extracellular fluids, *Biochem. J.*, 243 (1987) 219.
- 26 Frei, B., Stocker, R. and Ames, B., Antioxidants defenses and lipid peroxidation in human blood plasma, *Proc. Natl. Acad. Sci. USA*, 85 (1988) 9748.
- 27 Al-Timini, D.J. and Dormandy, T.L., The inhibition of lipid autoxidation by human ceruloplasmin, *Biochem. J.*, 168 (1977) 283.
- 28 Buettner, G.R., In the absence of catalytic metals ascorbate does not autoxidize at pH 7.0: ascorbate as a test for catalytic metals, *J. Biochem. Biophys. Methods*, 16 (1988) 27.
- 29 Boyum A., Isolation of mononuclear cells and granulocytes from human blood: isolation of mononuclear cells by one g centrifugation and of granulocytes by combining centrifugation and sedimentation at 1 g, *Scand. J. Clin. Lab. Invest.*, 21 (Suppl. 97) (1968) 77.
- 30 Lowry, O.H., Rosebrough, N.J., Farr, A.L. and Randall, R.J., Protein measurement with the Folin reagent, *J. Biol. Chem.*, 193 (1951) 265.
- 31 Noble, R.P., Electrophoretic separation of plasma lipoproteins in agarose gels, *J. Lipid Res.*, 9 (1968) 693.
- 32 Stookey, L.L., Ferrozine — a new spectrophotometric reagent for iron, *Anal. Chem.*, 42 (1970) 779.
- 33 Babior, B., Kipnes, R.S. and Curnutte, J.T., Biological defense mechanisms: the production of superoxide by leukocytes, a potential bactericidal agent, *J. Clin. Invest.*, 52 (1973) 741.
- 34 Winterbourn, C.C., Gutteridge, J.M.C. and Halliwell, B., Doxorubicin-dependent lipid peroxidation at low partial pressures of O₂, *J. Free Rad. Biol. Med.*, 1 (1985) 43.
- 35 Buege, J.A. and Aust, S.D., Microsomal lipid peroxidation, *Methods Enzymol.*, 52 (1978) 302.
- 36 Terao, J., Asano, I. and Matsushita, S, Preparation of hydroperoxy and hydroxy derivatives of rat liver phosphatidylcholine and phosphatidylethanolamine, *Lipids*, 20 (1985) 312.
- 37 Halliwell, B. and Gutteridge, J.M.C., *Free Radicals in Biology and Medicine*, Clarendon Press, Oxford, 1985, p. 20.
- 38 Sevanian, A., Davies, K.J.A. and Hochstein, P., Conservation of Vitamin C by uric acid in blood, *J. Free Rad. Biol. Med.*, 1 (1985) 117.
- 39 Samokyszyn, V.M., Miller, P.M., Reif, P.W. and Aust, S.D., Inhibition of superoxide and ferritin-dependent lipid peroxidation by ceruloplasmin, *J. Biol. Chem.*, 264 (1989) 21.
- 40 Wang-Iverson, P., Pryswansky, J.K., Spitsnagel, J.K. and Cooney, M.H., Bactericidal activity of phorbol myristate

- acetate treated human polymorphonuclear leukocytes, *Infect. Immun.*, 22 (1978) 945
- 41 Cathcart, M.K., Morel, D.W. and Chisolm, G.M., Human polymorphonuclear leukocyte oxidation and degradation of LPL, *Arteriosclerosis*, 6 (1986) 499a.
 - 42 Jacobs, A., Miller, F., Worwood, M., Beamish, M.R. and Wardrop, C.A., Ferritin in the serum of normal subjects and patients with iron deficiency and iron overload, *Br. Med. J.*, 4 (1972) 206.
 - 43 Lipschitz, D.A., Cook, J.D. and Finch, C.A., A clinical evaluation of serum ferritin as an index of iron stores, *N. Engl. J. Med.*, 290 (1974) 1213.
 - 44 Worwood, M., Serum ferritin, *Clin. Sci.*, 70 (1986) 215.
 - 45 Cartwright, G. E., The anemia of chronic disorders, *Semin. Hematol.*, 3 (1966) 351.
 - 46 Wohler, F. and Otte, W., Zur pathologie des speichereisens II. Mitteilung: Ferritin und Hemosideringehalt menschlicher organe bei entzündlichen erkrankungen, geschwulsten und krankheiten des blutbildenden systems, *Acta Hematol. (Basel)*, 26 (1961) 81.
 - 47 Heinecke, J.W., Baker, L., Rosen, H. and Chait, A., Superoxide-mediated modification of low density lipoprotein by arterial smooth muscle cells, *J. Clin. Invest.*, 77 (1986) 757.
 - 48 Clark, R.A., Extracellular effects of the myeloperoxidase-hydrogen peroxide-halide system, *Adv. Inflamm. Res.*, 5 (1983) 107.
 - 49 Bacon, B.R., Tavit, A.S., Brittenhan, J.N., Park, C.H. and Recknagel, R.O., Hepatic lipid peroxidation in vivo in rats with chronic iron overload. *J. Clin. Invest.*, 71 (1983) 429.
 - 50 Houghlum, K., Filip, M., Witztum, J.L. and Chojkier, M., Malondialdehyde and 4-hydroxynonenal protein adducts in plasma and liver of rats with iron overload, *J. Clin. Invest.*, 86 (1990) 1991.
 - 51 Burkitt, M.J. and Mason, R.P., Direct evidence for in vivo hydroxyl radical generation in experimental iron overload: an ESR spin-trapping investigation, *Proc. Natl. Acad. Sci. USA*, 88 (1991) 8440
 - 52 Ludwig, P.W., Hunninghake, D.B. and Hoidal, J.R., Increased leukocyte oxidative metabolism in hyperlipoproteinaemia, *Lancet* i (1982) 348.
 - 53 Jialal, I., Vega, G.L. and Grundy, S.M., Physiologic levels of ascorbate inhibit the oxidative modification of low density lipoprotein, *Atherosclerosis*, 82 (1990) 185.
 - 54 Harats, P., Naim, M.S., Dabach, Y., Hollander, G., Havivi, E., Stein, O. and Stein, Y., Effect of vitamin C and E supplementation on susceptibility of plasma lipoproteins to peroxidation induced by acute smoking, *Atherosclerosis*, 85 (1990) 47.
 - 55 Khan, M.T. and Martell, A.E., Metal ions and metal chelate catalysed oxidation of ascorbic acid by molecular oxygen. I: Cupric and ferric ion catalysed oxidation, *J. Am. Chem. Soc.*, 89 (1967) 4176.
 - 56 Halliwell, B. and Gutteridge, J.M.C., The antioxidants of human extracellular fluids, *Arch. Biochem. Biophys.*, 280 (1990) 1.
 - 57 Winterbourn, C.C. and Molloy, A.L., Susceptibilities of lactoferrin and transferrin to myeloperoxidase-dependent loss of iron binding capacity, *Biochem. J.*, 250 (1988) 613.

TRABALHO 2

Human macrophage metabolism of low density lipoprotein oxidized by stimulated neutrophils and ferritin

Dulcineia Saes Parra Abdalla^{*a}, Luiz Fernando Bicudo Pereira Costa-Rosa^b,
Hugo Pequeno Monteiro^c, Ana Campa^a, Rui Curi^b

^aFaculdade de Ciências Farmacêuticas, ^bInstituto de Ciências Biomédicas, Universidade de São Paulo,
São Paulo, S.P., Brazil

^cFundação Hemocentro de São Paulo, São Paulo, S.P., Brazil

(Received 5 November 1992; revision received 7 February 1994; accepted 8 February 1994)

Abstract

The metabolism of low density lipoprotein (LDL) oxidized with phorbol myristate acetate (PMA) stimulated neutrophils plus ferritin (LDLox) by human monocyte-derived macrophage (HMDM) was studied. Binding of ¹²⁵I-labeled LDLox to HMDM and further uptake and degradation were higher than for native ¹²⁵I-labeled LDL. LDLox seems to be taken up by HMDM through the scavenger receptor as indicated by competition studies with unlabeled native and autoxidized LDL. An increased concentration of cellular cholesteryl esters was observed in HMDM exposed to LDLox. Oxidative modification of LDL increased its electrophoretic migration on agarose gel and also the fragmentation of apolipoprotein B. Data suggest that LDLox is incorporated by human macrophages and can potentially induce foam-cell formation.

Key words: Oxidized LDL; Neutrophils; Ferritin; Human macrophages; Foam cells

1. Introduction

Monocyte-derived macrophages and macrophage-derived foam cells are found in the atherosclerotic lesions of animals and humans [1]. The transformation of macrophages into foam cells requires the intracellular accumulation of lip-

ids delivered mainly by plasma low density lipoprotein (LDL) [2]. Native LDL is taken up by B/E receptors in macrophages at a relatively low rate without cholesteryl ester accumulation [2]. However, macrophages can accumulate large amounts of cholesteryl esters by taking up chemically modified or oxidized LDL by the scavenger receptor [3]. Oxidatively modified LDL has been isolated in vivo from atheromatous lesions [1] and interstitial inflammatory fluid [4]. LDL can be oxidatively modified in vitro by arterial endothelial cells,

* Corresponding author, Faculdade de Ciências Farmacêuticas, Universidade de São Paulo, C.P. 66083, CEP 05389-970, São Paulo, S.P., Brazil. Fax: 55-11-813-2197.

monocytes and macrophages [5–7]. Neutrophils can oxidize LDL, making it cytotoxic [8]. These cells are able to produce superoxide radicals and other oxidant species when activated by diverse stimuli [9]. Lipid peroxidation mediated by neutrophils is exacerbated in the presence of an exogenous source of iron [10]. Since neutrophils do not contain appreciable amounts of endogenous iron, an available exogenous source of iron is necessary to catalyze this process. Iron bound to ferritin has been suggested as such a source [11]. Recently we showed [12] that LDL oxidation by stimulated neutrophils is greatly enhanced in the presence of ferritin. Superoxide radicals generated by the stimulated cells can release iron from ferritin and these ferrous ions can combine with oxygen to generate the final oxidant species. In the present study we investigated the metabolism of LDL oxidized with stimulated neutrophils plus ferritin (LDLox) by human monocyte-derived macrophages and compared the results with those obtained for homologous unmodified LDL and air-oxidized LDL.

2. Materials and methods

2.1. Materials

Sodium [125 I] (185 mBq, 5.0 mCi, 350–600 mCi/mmol, IMS 300) was purchased from Amersham International (Amersham, UK). Ficoll, dextran, sodium diatrizoate, phorbol 12-myristate 13-acetate (PMA), ferritin, bovine serum albumin (BSA), trypsin, aprotinin, HEPES, agarose, cholesterol oxidase, cholesterol ester hydrolase, dialysis tubing and butylated hydroxytoluene (BHT) were obtained from Sigma Chemical Co. (St. Louis, MO). Culture dishes and tubes were obtained from Corning Glass (Corning, NY). High performance liquid chromatography grade methanol, hexane and salts of analytical grade were obtained from Merck (Darmstadt, Germany). Eagle's medium and other cell culture supplies were purchased from Cultilab (Campinas, Brazil). Millex-GV, non-protein filters for lipoproteins and medium filtering were from Millipore S.A. (Rio de Janeiro, Brazil).

2.2. Methods

Macrophage culture. Human monocytes were isolated from freshly heparinized blood on a Ficoll-Hypaque discontinuous gradient [13]. Blood was donated by healthy subjects aged 27–32 years. The mononuclear cell preparations were washed three times with phosphate-buffered saline, pH 7.2, containing antibiotics. The cell suspension was 95% viable, as determined by Trypan blue exclusion. Mononuclear cells were resuspended to 10^4 cells/ml in Eagle's medium supplemented with 10% (v/v) fetal calf serum, 100 U/ml penicillin and 100 μ g/ml streptomycin. One milliliter of cell suspension was seeded onto 16 mm plastic wells and monocytes were allowed to adhere for 4 h. After incubation, nonadherent cells were eliminated with three phosphate buffered saline (PBS) washes and adhered monocytes were cultured with 1 ml of supplemented culture medium. The cells were fed every 3 days and were used for experiments within 9 days of plating. Two days before the assays, cells were incubated in 1 ml of culture medium containing 5 mg protein/ml lipoprotein-deficient human serum (LPDS) in order to induce LDL receptor expression. All the incubations were performed at 37°C in a 5% CO₂ humid atmosphere.

Lipoprotein and lipoprotein-deficient serum. Human LDL (d 1.019–1.063 g/ml) was prepared by sequential ultracentrifugation (4°C) of plasma from healthy, fasted male volunteers in tubes containing 1 mg/ml EDTA and 2 mM benzamidine [14]. After centrifugation ($1500 \times g$), plasma was separated and 1 mM PMSF and 40 μ M BHT were added. Labelling of LDL with [125 I]iodine was performed by the iodine monochloride method modified for lipoproteins [15]. The specific activity of 125 I-lipoprotein was between 20 and 50 counts/min per ng of protein. Ninety percent of the radioactivity was precipitated with 25% TCA (w/v) and less than 5% of the total was lipid extractable. LPDS (d > 1.215 g/ml) was prepared by ultracentrifugation of the infranatant obtained after sequential lipoprotein isolation.

Lipoprotein oxidation by neutrophils. Neutrophils were isolated from heparinized human blood by Ficoll-Hypaque centrifugation, dextran centrifugation of red cells and hypotonic lysis of the re-

maining red cells [13]. Neutrophils were suspended in PBS, pH 7.2, containing CaCl_2 (1 mM), MgCl_2 (0.5 mM) and glucose (1 mg/ml). Neutrophils ($1 \times 10^6/\text{ml}$) stimulated with PMA (100 ng/ml), LDL (1 mg/ml) and ferritin (200 $\mu\text{g}/\text{ml}$), in a final volume of 1 ml, were incubated in a shaker for 17 h at 37°C. Autoxidation of LDL was performed under the same conditions by incubating only the lipoprotein. Lipid hydroperoxide formation from lipoprotein oxidation was monitored by high performance liquid chromatography as previously described [16].

Lipoprotein binding, uptake and degradation. HMDM, preincubated for 48 h in culture medium with LPDS, were washed three times with PBS before adding medium containing ^{125}I -LDL and LDL oxidized by PMA-stimulated neutrophils plus ferritin (^{125}I -LDLox) or autoxidized ^{125}I -LDL at the concentrations indicated in the figures. For the competition experiments, a 20-fold excess of unlabeled LDL or LDLox was added to the medium. After 6 h of incubation at 37°C, the medium was removed and the extent of degradation of labeled lipoproteins was determined by measuring the amount of [^{125}I]-labeled trichloroacetic-soluble, non-iodide material (^{125}I -labeled moniodotyrosine) [17]. Parallel dishes without cells were included as a control for nonspecific fragmentation of labeled lipoprotein. The radioactivity in these dishes was subtracted from that of dishes with cells for final calculations. The cells were washed once with 2 ml of 50 mM Tris-HCl, pH 7.4, containing 150 mM NaCl and 2 mg/ml BSA (buffer A) and twice with the same buffer without albumin (buffer B). Cells were treated with 4 mg/ml dextran sulfate, 10 mM HEPES, 50 mM NaCl, pH 7.4 (buffer C) under slow agitation, at 37°C for 1 h. The dextran solution was transferred to a tube and the radioactivity corresponding to the cell surface-bound lipoprotein was counted. Adhered cells were extracted with 1 ml of 0.1 N NaOH and the radioactivity corresponding to the amount of lipoprotein internalized by the cells was counted. Total cell protein was measured in aliquots from the cell extraction.

The assay for binding of [^{125}I]-labeled lipoproteins to the cell surface was performed at 4°C [17].

HMDM preincubated in medium with LPDS for 48 h were washed once with cold buffer A and twice with cold buffer B. Prior to the experiment, the cells were incubated for 1 h at 4°C with culture medium and the [^{125}I]-labeled lipoproteins were then added at the concentrations indicated in the figures with or without unlabeled competing lipoproteins. After 2 h of incubation at 4°C, the cells were rapidly washed three times with cold buffer B and 1 ml of buffer C was added. Cells were incubated with 1 ml of buffer C for 1 h at 4°C. After incubation, the buffer was transferred to tubes and the radioactivity bound to the cell surface was counted. Cells were washed three times with cold buffer A and dissolved with 1 ml of 0.1 N NaOH for protein measurement. Nonspecific binding was defined as the amount of radiolabeled lipoprotein bound in the presence of a 10-fold excess of unlabeled lipoprotein. This count was subtracted from the total binding radioactivity obtained in the absence of unlabeled lipoproteins and specific cell-surface binding was calculated.

Other methods. The cholesteryl ester content of HMDM was measured after cell lysis with 0.1 N NaOH and lipid extraction with 2:1 chloroform-methanol (v/v). After solvent evaporation, the extracted lipids were dissolved in methanol and the mass of total and esterified cholesterol was determined by the fluorimetric method of Gamble et al. [18]. Protein was determined by a modification of the Lowry method with BSA as the standard [19]. Sodium dodecyl sulphate polyacrylamide gel electrophoresis was carried out as described by Laemmli [20]. For these experiments, 2 mM PMSF was added to the incubation systems to avoid proteolytic degradation of apo B by neutrophil proteases. Electrophoretic mobility was determined by agarose gel electrophoresis according to Noble [21].

3. Results

3.1. Lipoprotein binding, internalization and degradation

Binding studies indicate that native LDL and LDL oxidized by PMA-stimulated neutrophils plus ferritin (LDLox) do not compete for the same cell-surface receptor as shown in competition

Fig. 1. Competitive inhibition of the binding of ^{125}I -LDL to human monocyte-derived macrophages. Lipoprotein binding to macrophages was measured after 2 h incubation of ^{125}I -LDL alone or in the presence of unlabeled native or autooxidized LDL. Assays were run in duplicate and results (percent of binding) are mean of three experiments.

experiments with unlabeled lipoproteins (Fig. 1). It can be seen that unlabeled air-oxidized LDL inhibits about 40% of the binding of ^{125}I -LDL to the macrophage surface. The uptake and degradation of ^{125}I -LDL, ^{125}I -native LDL and autooxidized ^{125}I -LDL by HMDM are shown in Figs. 2 and 3, respectively. Internalization of LDLox was twice that of native LDL and 25% higher than au-

Fig. 2. Uptake of ^{125}I -LDL, cell-oxidized ^{125}I -LDL and autooxidized ^{125}I -LDL by human monocyte derived macrophages (HMDM). Internalization of labeled lipoproteins was determined as described in Materials and methods. Each point is the mean of duplicate determinations of three experiments.

Fig. 3. Lipoprotein degradation by human monocyte-derived macrophages. Cells were incubated with ^{125}I -labeled lipoprotein at the concentrations indicated. After 6 h at 37°C , medium was removed and the TCA-soluble radioactivity was measured as described in Materials and methods. Each point is the mean of duplicate determinations of three experiments.

toxidized LDL. Lipoprotein degradation was dependent on LDL, LDLox and autooxidized LDL concentrations in the medium. At the protein concentration range studied, ^{125}I -LDLox was degraded to a higher extent than ^{125}I -LDL and autooxidized ^{125}I -LDL. LDLox presented a higher level of lipid hydroperoxides than autooxidized LDL (LDLox = 307 ± 51 and autooxidized LDL = 159 ± 21 nmol of lipid hydroperoxides/mg cholesterol), indicating that the higher the oxidative modification, the greater is the uptake and degradation of lipoprotein. In addition, HMDM exposed to LDLox presented a 30% higher content of cholesteryl ester than those exposed to native LDL (LDLox = 51 ± 6 ng of cholesteryl ester/mg of cell protein and native LDL = 31 ± 5 ng of cholesteryl ester/mg of cell protein, for cells exposed to $10 \mu\text{g/ml}$ of either LDLox or native LDL).

3.2. Electrophoretic pattern

LDLox migrated as a single discrete band during agarose electrophoresis, indicating a homogeneous population of oxidized particles with increased electrophoretic mobility (migration distances for native, cell-oxidized and autooxidized

Fig. 4. Sodium dodecyl sulphate polyacrylamide gel electrophoresis of native and oxidized LDL. Samples (20 µg/lane) were run on 2.5% to 20% gradient gel and stained with Coomassie blue. Standard proteins used for reference (with their molecular weights) were thyroglobulin (330 kDa), ferritin (half unit, 220 kDa), phosphorylase (94 kDa), bovine serum albumin (67 kDa), ovalbumin (43 kDa), carbonic anhydrase (30 kDa), trypsin inhibitor (20 kDa), ferritin (subunit, 18.5 kDa) and alpha-lactalbumin (14 kDa). Reaction mixtures were incubated at 37°C for 17 h with shaking in a water bath, in the presence of 2 mM PMSF. Lane (a), high molecular weight standard; (b) LDL (1 mg/ml); (c) ferritin (200 µg/ml); (d) PMA-stimulated neutrophils (1×10^6 /ml); (e) LDL (1 mg/ml) + ferritin (200 µg/ml); (f) LDL + neutrophils + PMA (100 ng/ml); (g) LDL + PMA-stimulated neutrophils + ferritin; (h) low molecular weight standard.

LDL were 1.8 cm, 2.5 cm and 2.3 cm, respectively). This is in accordance with previous data showing that oxidatively modified LDL has an increased negative charge that causes its greater migration on agarose gels [22,23]. Fragmentation of apolipoprotein B during incubation of LDL with PMA-stimulated neutrophils plus ferritin also occurred (Fig. 4). Sodium dodecyl sulphate polyacrylamide gel electrophoresis of LDLox showed an extensive loss of Coomassie blue staining in the apo B₁₀₀ region (lane g, compared with lane b) and the appearance of lower molecular weight bands.

Incubation of LDL with PMA-stimulated neutrophils (lane f) or ferritin (lane e) alone promoted less fragmentation than the complete system (stimulated neutrophils plus ferritin). The appearance of these lower molecular weight polypeptides was

not due to the proteolytic action of enzymes released by neutrophils because incubations were done in the presence of the protease inhibitor PMSF.

4. Discussion

Oxidatively or chemically modified LDL can cause cholesteryl ester accumulation in macrophages following its uptake into the cells via alternative pathways (scavenger receptors) [2,3,7]. Unlike the classical LDL receptor which is down-regulated by intracellular cholesterol content, the scavenger receptor is not similarly regulated and uptake of oxidized LDL by this pathway can cause foam cell formation. Recently, other receptors for modified LDL have been described [22,23]. Oxidized LDL seems to be internalized by macro-

phages through two classes of receptors; one is a high-affinity receptor and the other is a low-affinity receptor that could also bind acetyl-LDL [22,23]. In the present study we have seen that air-oxidized LDL partially compete with LDLox for receptor binding. Addition of excess unlabeled native LDL did not affect the binding of ^{125}I -LDLox, suggesting that LDLox is taken up into HMDM by a receptor different from the B/E receptor. Both LDLox and autoxidized LDL showed increased electronegativity observed in agarose gel. This is a common feature of all forms of modified LDL taken up by the scavenger receptor [24,25] and this increase in net negative charge is attributed to derivatization of epsilon amino groups of apo B lysine residues [25]. Since autoxidized LDL is less electronegative than LDLox, it could bind to macrophages through the low-affinity receptor and not displace LDLox from the high-affinity receptors. Moreover, oxidation of lipoprotein may be important to expose normally hidden residues in apo B, essential for scavenger receptor recognition, and the extent of LDL oxidative modification would determine the number and type of these exposed residues and consequently the difference in the binding of LDLox and autoxidized LDL to macrophages.

LDL oxidatively modified by PMA-stimulated neutrophils plus ferritin (LDLox) caused accelerated uptake and degradation by HMDM with greater intracellular formation of cholesteryl esters. These results are similar to those obtained with LDL modified by other cellular systems [4,5,7]. Uptake and degradation of LDLox was higher than that of native and autoxidized LDL, possibly indicating that the intensity of oxidative modification is important in modulating the uptake of lipoprotein by alternative pathways. Fragmentation of apo B, reflecting nonenzymatic cleavage, is also a characteristic of oxidatively modified LDL [24,26] and may be caused by cell-derived active oxygen species (hydrogen peroxide, hydroxyl radical, ferryl ion or other oxidant species). The different patterns of fragmentation for autoxidized LDL and LDLox could also affect the metabolism of these lipoproteins by macrophages.

It is interesting to note that iron ions promote a slight apo B fragmentation in accordance with our previous data [12], showing a small oxidation of

LDL mediated by iron alone. This indicates that the presence of a binary system formed by stimulated neutrophils and ferritin is crucial in originating the oxidant species that initiate LDL oxidation. It has been reported that oxidation of LDL by leukocytes makes it cytotoxic [8]. Also, LDL isolated from interstitial inflammatory fluid showed an accelerated uptake by resident mouse peritoneal macrophages [4]. In a previous study [12] we showed that oxidation of LDL by neutrophils is potentiated by the presence of ferritin. Our hypothesis is that superoxide produced by stimulated neutrophils can reductively release iron from ferritin present in blood and derived from cells, which may catalyze the generation of more oxidant species initiating lipoprotein oxidation. Accordingly, recent epidemiological studies [27] have shown that high levels of stored iron, assessed by elevated serum ferritin concentration, are a risk factor for coronary heart disease.

In the present study, we showed that LDLox exhibits high uptake and degradation by HMDM. This process could be relevant in vivo, mainly in terms of focal inflammation where there is an accumulation of polymorphonuclear cells potentially able to produce large amounts of active oxygen species. These cells close to the vascular endothelium may damage endothelial cells, allowing plasma lipoproteins to enter the subendothelial space. In addition, ferritin from destroyed target cells (endothelial) [28] would be available as a source of iron that may participate in LDL oxidation. Thus, in this scenario we have many elements that are important for the atherogenic process: (a) endothelial damage, (b) lipoprotein oxidation, (c) enhanced uptake of LDL by macrophages and (d) accumulation of cellular cholesteryl esters permitting foam cell formation. Of course, in vivo systems are extremely complex and a diversity of factors should be regarded as participating in this process. The in vitro data shown here may be helpful for future work in atherogenesis that considers situations of high iron availability associated with an inflammatory focus.

5. Acknowledgments

This work was supported by a grant from FAPESP (Fundação de Amparo à Pesquisa do Estado de São Paulo). We wish to thank Mr.

Paulo Roberto Omoto for helpful technical assistance and Ms. Elettra Greene for revising the manuscript.

6. References

- Rosenfeld, M., Khoo, J.C., Miller, E., Parthasarathy, S., Palinski, W and Witztum, J.L., Macrophage derived foam cells freshly isolated from rabbit atherosclerotic lesions degrade modified lipoproteins, promote oxidation of low density lipoproteins, and contain oxidation-specific lipid-protein adducts, *J. Clin. Invest.*, 87 (1991) 90.
- Brow, M.S. and Goldstein, J.L., Lipoprotein metabolism in the macrophage: implications for cholesterol deposition in atherosclerosis, *Annu. Rev. Biochem.*, 52 (1983) 223.
- Goldstein, J.L., Ho, Y.K., Basu, S.K., and Brown, M.S., Binding sites on macrophages that mediate uptake and degradation of acetylated low density lipoprotein, producing massive cholesterol deposition, *Proc. Natl. Acad. Sci. USA*, 76 (1979) 333.
- Raymond, T.L., Reynolds, S.A. and Swanson, J.A., Lipoproteins of the extravascular space: enhanced macrophage degradation of low density lipoprotein from interstitial inflammatory fluid, *J. Lipid Res.*, 26 (1985) 1356.
- Henriksen, T., Mahoney, E.M. and Steinberg, D., Enhanced macrophage degradation of low density lipoprotein previously incubated with cultured endothelial cells: recognition by receptor for acetylated low density lipoproteins, *Proc. Natl. Acad. Sci. USA*, 78 (1981) 6499.
- Hiramatsu, K., Rosen, H., Heinecke, J.M., Wolfbauer, G. and Chait, A., Superoxide initiates oxidation of low density lipoprotein by human monocytes, *Arteriosclerosis*, 7 (1987) 55.
- Parthasarathy, S., Printz, D.J., Boyd, D., Joy, L. and Steinberg, D., Macrophage oxidation of low density lipoprotein generates a modified form recognized by the scavenger receptor, *Arteriosclerosis*, 6 (1986) 505.
- Cathcart, M.K., Morel, D.W. and Chisolm, G.M., III, Monocytes and neutrophils oxidize low density lipoprotein making it cytotoxic, *J. Leukocyte Biol.*, 38 (1985) 341.
- Britigan, B.E., Rosen, G.M., Chai, Y. and Cohen, M.S., Do human neutrophils make hydroxyl radical? Determination of free radicals generated by human neutrophils activated with soluble or particulate stimulus using electron paramagnetic resonance spectroscopy, *J. Biol. Chem.*, 261 (1986) 4426.
- Thomas, M.J., Shirley, P.S., Hendrich, C.C. and DeChatelet, L.R., Role of free radical processes in stimulated human polymorphonuclear leukocytes, *Biochemistry*, 25 (1986) 8042.
- Winterbourn, C.C., Monteiro, H.P. and Galilee, C.F., Ferritin-dependent lipid peroxidation by stimulated neutrophils: inhibition by myeloperoxidase-derived hypochlorous acid but not by endogenous lactoferrin, *Biochem. Biophys. Acta*, 1055 (1990) 179.
- Abdalla, D.S.P., Campa, A. and Monteiro, H.P., Oxidation of LDL by stimulated neutrophils and ferritin, *Atherosclerosis*, 97 (1992) 149.
- Boyum, A., Isolation of mononuclear and granulocytes from human blood: isolation of mononuclear cells by one g centrifugation and of granulocytes by combining centrifugation and sedimentation at 1 g, *Scand. J. Lab. Invest.*, 21 (1968) 77.
- Havel, R.J., Eder, H.A. and Bragdon, J.H., The distribution and chemical composition of ultracentrifugally separated lipoproteins in human serum, *J. Clin. Invest.*, 34 (1955) 1345.
- Shephard, J., Belford, D.K. and Morgan, H.G., Radioiodination of human low density lipoprotein: a comparison of four methods, *Clin. Chim. Acta*, 66 (1976) 97.
- Terao, J., Shibata, S.S. and Matsushita, S., Selective quantification of arachidonic acid hydroperoxides and their hydroxy derivatives in reverse-phase high performance liquid chromatography, *Anal. Biochem.*, 169 (1988) 415.
- Goldstein, J.L. and Brown, M.S., Receptor-mediated endocytosis of low density lipoprotein in cultured cells, *Methods Enzymol.*, 98 (1983) 241.
- Gamble, W., Vaughan, M., Kruth, H.S. and Avigan, J., Procedure for determination of free and total cholesterol in micro- or nanogram amounts suitable for studies with cultured cells, *J. Lipid Res.*, 19 (1978) 1068.
- Markwell, M.A.K., Haas, S.N., Bieber, L.L. and Tolbert, N.E., A modification of the Lowry procedure to simplify protein determination in membrane and lipoprotein samples, *Anal. Biochem.*, 87 (1978) 206.
- Laemmli, U.K., Cleavage of structural proteins during the assembly of the head bacteriophage T4, *Nature*, 227 (1970) 680.
- Nobel, R.P., Electrophoretic separation of plasma lipoproteins in agarose gels, *J. Lipid Res.*, 9 (1968) 693.
- Arai, H., Kita, T., Yokode, M., Narumiya, S. and Kawai, C., Multiple receptors for modified low density lipoproteins in mouse peritoneal macrophages: different uptake mechanisms for acetylated and oxidized low density lipoprotein, *Biochem. Biophys. Res. Commun.*, 159 (1989) 1375.
- Sparrow, C.P., Parthasarathy, S. and Steinberg, D., A macrophage receptor that recognizes oxidized low density lipoprotein but not acetylated low density lipoprotein, *J. Biol. Chem.*, 264 (1989) 2599.
- Montgomery, R.R., Nathan, C.F. and Cohn, Z.A., Effects of reagent and cell-generated hydrogen peroxide on the properties of low density lipoproteins, *Proc. Natl. Acad. Sci. USA*, 83 (1986) 6631.
- Haberland, M.E., Olch, C.L. and Fogelman, A.M., Role of lysines in mediating interaction of modified low density lipoproteins with the scavenger receptor of human macrophages, *J. Biol. Chem.*, 259 (1984) 11305.
- Schuh, J., Fairclough, G.F., Jr., and Haschemeyer, R.H., Oxygen-mediated heterogeneity of apo-low density lipoprotein, *Proc. Natl. Acad. Sci. USA*, 75 (1978) 3173.
- Salonen, J.K., Nyyssönen, K., Korpela, H., Tuomilehto, J., Seppänen, R. and Salonen, R., High stored iron levels are associated with excess risk of myocardial infarction in eastern Finnish men, *Circulation*, 86 (1992) 803.
- Gannon, D.E., Varani, J., Phan, S.H. et al., Source of iron in neutrophil-mediated killing of endothelial cells, *Lab. Invest.*, 57 (1987) 37.

TRABALHO 3

BIODISTRIBUTION AND FRACTIONAL CLEARANCE RATE OF COPPER- AND PEROXYNITRITE-OXIDIZED ^{99m}Tc - β -VLDL IN CHOLESTEROL-FED RABBITS

Edson L. Silva^a, Iris J.C. Coelho^b, José C. Meneghetti^b, Dulcineia S. P. Abdalla^{1,a}

^a*Department of Clinical and Toxicological Analysis, Faculty of Pharmaceutical Sciences (FCFUSP), University of São Paulo, São Paulo, S.P., Brazil.*

^b*Heart Institute (INCOR), Faculty of Medicine, University of São Paulo, São Paulo, Brazil.*

Running Title: Biodistribution and FCR of native and oxidized ^{99m}Tc - β -VLDL

Abbreviations: β -VLDL, β -very low density lipoprotein; ^{99m}Tc - β -VLDL, native β -very low density lipoprotein labeled with ^{99m}Tc ; ^{99m}Tc - β -VLDL-Per, peroxyntirite-oxidized β -very low density lipoprotein labeled with ^{99m}Tc ; ^{99m}Tc - β -VLDL-Ox, copper-oxidized β -very low density lipoprotein labeled with ^{99m}Tc ; FCR, fractional clearance rate.

¹Correspondence address:
Dr. Dulcineia S.P. Abdalla
Faculdade de Ciências Farmacêuticas
Universidade de São Paulo
Av. Prof. Lineu Prestes, 580 - Cidade Universitária
CP 66083, CEP 05389-970 - São Paulo, S.P., Brasil
FAX: (5511) 813-2197

Abstract

The oxidative modification of lipoproteins enhances their atherogenicity and is a critical event in development of atherosclerotic lesions. Oxidized β -Very low density lipoprotein (β -VLDL) may play an important role in the atherogenic process of cholesterol-fed rabbits and in patients with type-III hyperlipoproteinemia. The biodistribution and plasma clearance of native, copper-oxidized and peroxynitrite-modified ^{99m}Tc -labeled β -VLDL were studied in cholesterol-fed and control rabbits. The radiolabeled peroxynitrite-modified β -VLDL (^{99m}Tc - β -VLDL-Per) was cleared faster than either native (^{99m}Tc - β -VLDL) or copper-oxidized (^{99m}Tc - β -VLDL-Ox) lipoproteins in both cholesterol-fed and control rabbits. In contrast, the fractional clearance rate of ^{99m}Tc - β -VLDL-Ox was lower than that of ^{99m}Tc - β -VLDL and ^{99m}Tc - β -VLDL-Per in cholesterol-fed rabbits. In these animals the uptake of ^{99m}Tc - β -VLDL-Ox by liver, spleen and heart was greater than that of both ^{99m}Tc - β -VLDL and ^{99m}Tc - β -VLDL-Per. Biodistribution data indicate that aorta of cholesterol-fed rabbits accumulated more ^{99m}Tc - β -VLDL-Ox in comparison to that of control rabbits and, autoradiography shows the accumulation of ^{99m}Tc - β -VLDL-Ox in atherosclerotic lesions of cholesterol-fed rabbit aorta. The liver uptake of radiolabeled lipoproteins was lower in cholesterol-fed rabbits than in controls indicating either a down regulation of hepatic receptors or a competition between the endogenous pool of native and oxidized lipoproteins and the respective radiolabeled lipoproteins injected. In summary, the results indicate that under the experimental conditions used here the peroxynitrite-modified β -VLDL is rapidly cleared from plasma and does not accumulate in aorta of cholesterol-fed rabbits. In contrast, copper-oxidized ^{99m}Tc - β -VLDL is slowly cleared from blood and accumulates in atherosclerotic lesions.

Supplementary key words: β -very low density lipoprotein, copper-oxidized, peroxynitrite, biodistribution, fractional clearance rate, atherogenesis, atherosclerosis, hypercholesterolemia, cholesterol-feeding, rabbit.

INTRODUCTION

β -Very low density lipoprotein (β -VLDL) is an atherogenic lipoprotein that accumulates in plasma of cholesterol-fed rabbits and in patients with type-III hyperlipoproteinemia (1). There is increasing evidence to indicate that the oxidative modification of lipoproteins enhances their atherogenicity and is a critical event in development of atherosclerotic lesions (2). The oxidation of lipoproteins may be induced by oxidant species released by endothelial and smooth muscle cells (3), monocytes (4), neutrophils (5) and macrophages (6). Reactive oxygen species are mediators in the atherosclerotic process and other vascular diseases (7). Superoxide radical (O_2^-), hydrogen peroxide and transition metals may induce the modification of low density lipoprotein (LDL) and β -VLDL (4, 8).

The arteries of hypercholesterolemic rabbits produce a high amount of superoxide radical (9) and nitric oxide ($\cdot NO$) (10). These two free radicals can react with each other to form peroxynitrite ($ONOO^-$) (11) that is a strong oxidant and can oxidize LDL (12) and β -VLDL (13). Peroxynitrite is also a potential mediator of vascular tissue injury (14) and may be involved in the impairment of the endothelium-dependent relaxation observed in vessels from atherosclerotic subjects and animals (15).

Technetium-99m-lipoprotein labeling is a powerful tool for the noninvasive exploration of a variety of disorders of lipoprotein metabolism in patients and animal models (16). ^{99m}Tc -lipoprotein is taken up well by the actively evolving atherosclerotic lesions and acts as an intracellularly trapped ligand giving an accurate measure of lipoprotein uptake by tissues (17). ^{99m}Tc -lipoprotein imaging is useful in studies on the accumulation of lipoproteins by organs and atherosclerotic plaques (18).

In hypercholesterolemia the uptake of lipoproteins by the LDL receptor is reduced due to a down-regulation mechanism (19). However, native and oxidized β -VLDL can be taken up by other receptors which expression is not regulated by the sterol content of the cells (20) making them available to participate in the uptake of this lipoprotein even during hypercholesterolemia.

Considering that β -VLDL is an atherogenic lipoprotein and that its oxidation may even increase its atherogenicity it is important to investigate the effect of the oxidative modification of this lipoprotein, induced by different oxidant species, on its metabolism *in vivo*. The goal of the present study was to compare the biodistribution and the fractional clearance rate of native, copper-oxidized and peroxynitrite-modified ^{99m}Tc - β -VLDL in cholesterol-fed and control rabbits.

MATERIALS AND METHODS

Materials

Cholesterol, benzamidine, phenylmethylsulphonyl fluoride (PMSF), aprotinin, butylated hydroxytoluene (BHT), EDTA and the KODAK X-OMAT XRP-5 were purchased from Sigma Chem. Co. (St. Louis, MO.). Sephadex G25 was from Pharmacia (Uppsala, Sweden). The enzymatic reagents for cholesterol and triglyceride determinations were kindly donated by CELM (São Paulo, S.P., Brazil). All other chemicals were of the purest analytical grade available. Rompun[®] was kindly donated by Bayer Co. (São Paulo, S.P., Brasil).

Animals

Male New Zealand white rabbits weighing 2.5-3.0 kg were housed in individual cages at room temperature with a 12 h light/dark cycle. The control animals were fed a standard pelleted chow (Purina, Campinas, S.P., Brazil) and the experimental group was fed the standard chow containing 1% cholesterol, during 2 months. The water was available *ad libitum* to both groups. The plasma cholesterol and triglycerides of control rabbits and rabbits after two months cholesterol feeding were determined by enzymatic methods and were the following: control group ($\bar{x} \pm \text{S.D.}$), cholesterol = 33 ± 5 mg/dl and triglycerides = 50 ± 8 mg/dl; cholesterol-fed group ($\bar{x} \pm \text{S.D.}$), cholesterol = 1182 ± 172 mg/dl and triglycerides = 260 ± 72 mg/dl.

Lipoprotein isolation and oxidation

Blood was collected into tubes containing EDTA (1 mg/ml) and centrifuged at $1000 \times g$ for 10 min. at 4°C to isolate plasma. Immediately after isolation, 2 mM benzamidine, 1 mM PMSF, 1 mg/ml aprotinin and 0.1 mM BHT were added to plasma to inhibit proteolytic degradation of apolipoproteins and lipid peroxidation. β -VLDL (density < 1.019 g/ml) was isolated from plasma by sequential ultracentrifugation as described (21). After isolation the lipoprotein was dialyzed against 6 liters of 150 mM NaCl, 1 mM EDTA, 3 mM NaN_3 in Tris buffer pH 7.4. β -VLDL (1

mg of protein/ml) was exposed to 200 μM CuSO_4 or to 400 μM peroxyxynitrite, synthesized as described (14), by 18 h at 37 °C in a shaker bath. These concentrations of both oxidant agents were chosen based on the greatest production of thiobarbituric acid-reactive substances (TBARS) produced and the maximal electrophoretic migration observed in agarose gel.

Lipoprotein labeling with Technetium-99m

Native and oxidized β -VLDL were labeled with $^{99\text{m}}\text{Tc}$ according to Lees et al. (22). Briefly, β -VLDL (4-8 mg of protein) was mixed with 40-60 mCi of $^{99\text{m}}\text{Tc}$ -pertechnetate (TcO_4^-) and 10 mg of sodium dithionite, which was dissolved just before use in 0.5 M glycine bufer pH 9.8, and the mixture was incubated for 30 min. $^{99\text{m}}\text{Tc}$ - β -VLDL was separated from free $^{99\text{m}}\text{Tc}$ by Sephadex G 25 chromatography. The purified $^{99\text{m}}\text{Tc}$ - β -VLDL was dialyzed against 150 mM NaCl, 1 mM EDTA, 3 mM NaN_3 in Tris buffer pH 7.4, at 4 °C by 2 h and used for the experiments.

Biodistribution and plasma clearance studies

Overnight fasted rabbits received native or oxidized $^{99\text{m}}\text{Tc}$ - β -VLDL in a bolus (1-3 ml) corresponding to 0.5-1.5 mg of protein (5-10 mCi) by i.v. in the marginal ear vein. Blood samples were drawn in heparinized tubes 5, 15, 30, 60, 120, 160 and 240 min. post injection from the central ear artery. Plasma was isolated by centrifugation at 1000 x g for 10 min. and the radioactivity was measured with a gamma counter (Logic, Abbott Corp., USA). The fractional clearance rate (FCR), defined as the fraction of the intravascular pool cleared per hour, was calculated from biexponential curves obtained from the radioactivity found in plasma after injection fitted by the least square procedure. The percent of plasma radioactivity was calculated considering the value obtained at 5 min. as 100%. A compartmental model (23) was utilized for the analysis of plasma decaying curves of the $^{99\text{m}}\text{Tc}$ -lipoproteins as described elsewhere (24) and FCR of the $^{99\text{m}}\text{Tc}$ -lipoproteins from the intravascular compartment was estimated according to Matthews (25). After 6 h the rabbits were sacrificed by exsanguination, under ketamine/Rompun

anesthesia, and the organs were removed. After rinsing the tissues with 150 mM NaCl, the radioactivity was counted per g wet of tissue. The radioactivity, expressed as a percent of the injected dose, was calculated for liver, kidney, heart, ileo-psoas muscle, adrenal gland, lung, spleen, aorta and bone marrow.

Imaging studies

The biodistributions of ^{99m}Tc - β -VLDL, ^{99m}Tc - β -VLDL-Per and ^{99m}Tc - β -VLDL-Ox were imaged with a gamma camera (model ON 100, Ohio Nuclear) equipped with a high resolution collimator and stored in 128 x 128 matrices in a dedicated computer system. The planar images were acquired in anterior view of the total body 6 h after the injection of ^{99m}Tc -labeled lipoproteins.

Autoradiography

The aortas were removed and washed with saline. Further they were opened, covered with a plastic wrap, placed on X-ray film (KODAK X-OMAT XRP-5) and stored by 48 h in a cassette to perform autoradiography.

Statistical analysis

Data are expressed as mean $\pm$ standard deviation. All comparisons were made using ANOVA and the Bonferroni t-test. Statistical significance is defined as $p < 0.05$.

RESULTS

Biodistribution studies

Comparison among lipoproteins

Biodistribution data for injected native and oxidized ^{99m}Tc - β -VLDL was obtained in seven male overnight fasted New Zealand White control (Table 1) and cholesterol-fed rabbits (Table 2) 6 h after injection. Percent injected dose per g tissue of ^{99m}Tc -lipoproteins was high not only in organs with high LDL receptor activity, such as liver and adrenal gland, but also in kidney, spleen and bone marrow. In the control group the uptake of ^{99m}Tc - β -VLDL by heart was greater than that of oxidized lipoproteins, while ^{99m}Tc - β -VLDL-Per was more taken up by kidney as compared to native and copper-oxidized ^{99m}Tc -labeled β -VLDL (Table 1). In this group the uptake of ^{99m}Tc - β -VLDL-Ox by liver and lung was greater than that of ^{99m}Tc - β -VLDL-Per and, by adrenal was higher than both ^{99m}Tc - β -VLDL and ^{99m}Tc - β -VLDL-Per (Table 1). In cholesterol-fed rabbits ^{99m}Tc - β -VLDL-Ox was taken up in greater intensity than the other two particles by liver, heart, muscle, adrenal gland, lung, spleen and aorta (Table 2).

Comparison between groups

A lower uptake of ^{99m}Tc - β -VLDL by liver, adrenal gland and bone marrow was found in cholesterol-fed rabbits as compared to controls ($p < 0.05$, Tables 1 and 2). ^{99m}Tc - β -VLDL-Ox was more taken up by kidney, heart, muscle, lung and aorta of cholesterol-fed rabbits ($p < 0.05$). In the latter group the uptake of ^{99m}Tc - β -VLDL-Per by liver, kidney, muscle and spleen was lower than in controls ($p < 0.05$). Kidney had the highest uptake either for native or oxidized ^{99m}Tc -labeled β -VLDL in both cholesterol-fed and control groups (Tables 1 and 2).

Fractional clearance rate

In both control and hypercholesterolemic groups the plasma clearance of ^{99m}Tc - β -VLDL, ^{99m}Tc - β -VLDL-Ox and ^{99m}Tc - β -VLDL-Per reflects a biexponential decay mode (Fig. 1 and Fig. 2). Decay curves show that plasma clearance of ^{99m}Tc - β -VLDL and ^{99m}Tc - β -VLDL-Ox is slower than that of ^{99m}Tc - β -VLDL-Per in both groups. However, in the cholesterol-fed rabbits the plasma clearance of ^{99m}Tc - β -VLDL-Ox was slower than that of ^{99m}Tc - β -VLDL (Fig. 2). These data are confirmed by the fractional clearance rates (FCR) showed in Table 3. Furthermore, in control rabbits the half-life of ^{99m}Tc - β -VLDL-Per ($T_{1/2} = 1.30 \pm 0.24$ h) is the shortest one as compared to ^{99m}Tc - β -VLDL ($T_{1/2} = 2.13 \pm 0.20$ h) and ^{99m}Tc - β -VLDL-Ox ($T_{1/2} = 1.79 \pm 0.23$ h). The same occurred in cholesterol-fed rabbits: ^{99m}Tc - β -VLDL-Per ($T_{1/2} = 2.11 \pm 0.29$ h), ^{99m}Tc - β -VLDL ($T_{1/2} = 2.92 \pm 0.38$ h) and ^{99m}Tc - β -VLDL-Ox ($T_{1/2} = 4.98 \pm 1.59$ h).

Imaging studies

The gamma camera images of the biodistribution of ^{99m}Tc -labeled lipoproteins in one control rabbit and one cholesterol-fed rabbit, obtained 6 h after injection is shown in Fig. 3. The liver was the predominant uptake site for ^{99m}Tc - β -VLDL (Fig. 3A), ^{99m}Tc - β -VLDL-Per (Fig. 3C) and ^{99m}Tc - β -VLDL-Ox (Fig. 3E) in control rabbit. The uptake of all ^{99m}Tc -labeled lipoproteins by liver of control rabbit was greater than that of cholesterol-fed rabbit. In the latter, there was a great uptake of ^{99m}Tc - β -VLDL by kidneys (Fig. 3B). The oxidized lipoproteins also appeared in the bladder of both animals indicating a greater proteolysis of these particles. The accumulation of radioactivity in bladder occurred mainly for ^{99m}Tc - β -VLDL-Per (Fig. 3D) in cholesterol-fed rabbits.

Accumulation of ^{99m}Tc - β -VLDL-Ox in atherosclerotic lesions

Fig. 4 shows the accumulation of ^{99m}Tc - β -VLDL-Ox in the aorta of one control rabbit (Fig. 4A) and one cholesterol-fed rabbit (Fig. 4B). The aorta of the cholesterol-fed rabbit accumulates ^{99m}Tc - β -VLDL-Ox radioactivity mainly in the aortic arch where a great number of atherosclerotic lesions occurs (Fig. 4B). In control rabbits the accumulation of ^{99m}Tc - β -VLDL-Ox is barely seen (Fig. 4A). This result is according to the whole aorta radioactivity showed in Tables 1 and 2.

DISCUSSION

The biodistribution and fractional clearance rate were different among native, copper-oxidized and peroxyxynitrite-modified β -VLDL in cholesterol-fed and control rabbits. Consistent with previous reports in the presence of a down-regulation of LDL receptors in the cholesterol-fed rabbits the uptake of native β -VLDL by liver and adrenal gland is decreased (16, 26, 27). A relevant and new finding observed here was the intense accumulation of ^{99m}Tc - β -VLDL-Ox in atherosclerotic lesions of aortas from cholesterol-fed rabbits. The accumulation of oxidized LDL had been previously shown in atheromas of hypercholesterolemic rabbits and atherosclerotic subjects (28). However, until we know, the *in vivo* accumulation of oxidized β -VLDL-Ox in arteries of cholesterol-fed rabbits had not been described yet. The foam cells present in atheromas express scavenger receptors that recognize oxidized lipoproteins which has increased negative charges (29). In fact, the oxidation of LDL and β -VLDL results in particles with increased negative charge, due to derivatization of ϵ -amino group of lysine residues on apolipoprotein by aldehydes released from peroxidized lipids, allowing the recognition of the oxidized lipoproteins by the scavenger receptor (30, 31). Moreover, the FCR of ^{99m}Tc - β -VLDL-Ox indicates that this particle is removed from plasma slower than native β -VLDL in cholesterol-fed rabbits. This contrasts with data previously reported showing that the acetoacetylation of the LDL apoprotein leads to a marked acceleration of its plasma clearance (27), as well as with our data for ^{99m}Tc - β -VLDL-Per.

Although it is reported that peroxyxynitrite-modified LDL is recognized by the scavenger receptor (32) and nitrotyrosine has been detected in atherosclerotic lesions of human coronary arteries (33), we did not observe accumulation of ^{99m}Tc - β -VLDL-Per in the atherosclerotic lesions of cholesterol-fed rabbits. The distinct biodistribution between copper- and peroxyxynitrite-oxidized β -VLDL may be due to differences in the oxidation products formed in lipoprotein particles in both situations. Nitration of the *ortho* position on tyrosine and thiol oxidation are the major modifications resulting from peroxyxynitrite attack on proteins (33, 34). Furthermore,

hydroxyhydroperoxyl-, and nitrogen-containing adducts of ONOO^- besides alkyl hydroperoxide isomers (35). Thus, the oxidation of lipoproteins by peroxynitrite may produce differently modified lipoprotein particles as compared to those originated from copper-catalyzed oxidation. The lack of $^{99\text{m}}\text{Tc}$ - β -VLDL-Per accumulation in atheromas in our study may be also a consequence of its fast disappearance from plasma as indicated by the short $T_{1/2}$ of this particle in intravascular compartment.

A high uptake of $^{99\text{m}}\text{Tc}$ -labeled β -VLDL by kidneys was observed. The presence of apolipoprotein E on the surface of β -VLDL allows that this lipoprotein is taken up by other receptors, besides the LDL receptor, such as the LDL receptor-related protein (LRP), the VLDL receptor and the glycoprotein-330 (20). In kidney the epithelial cells of plasma membrane in proximal tubule and glomerulus have a high expression of the glycoprotein-330 (Gp330), a recently identified receptor that mediates the endocytosis of β -VLDL (20). GP-330 is also present in type-2 pneumocytes and in the epithelium lining the epididymis and yolk sac (20). Here it was observed that the accumulation of $^{99\text{m}}\text{Tc}$ - β -VLDL in kidneys was not decreased in cholesterol-fed rabbits indicating that the receptor mediating its internalization is not down regulated by the sterol content of cells. Although lipoproteins are not expected to be filtered by kidneys due to their size they may bind to GP-330 present on glomerulus plasma membrane. Another possibility for the presence of radioactivity in kidneys is the presence of free $^{99\text{m}}\text{Tc}$ released from labeled lipoproteins *in vivo*. However, it is reported that both in animals and humans only 5% to 12% of injected $^{99\text{m}}\text{Tc}$ activity was excreted in the urine within a 24-hour period following an injection of $^{99\text{m}}\text{Tc}$ -LDL, indicating that $^{99\text{m}}\text{Tc}$ -labeled lipoprotein acts as an intracellular ligand (16). In the case of oxidized lipoproteins it is possible that an increased proteolysis may occur and the resulting $^{99\text{m}}\text{Tc}$ -labeled low molecular weight peptides may be filtered by kidneys. This is indicated by scintigraphic images of the peroxynitrite-modified β -VLDL that shows radioactivity accumulation in the bladder of rabbits. Furthermore, the FCR of $^{99\text{m}}\text{Tc}$ - β -VLDL-Per shows its rapid elimination from plasma.

A sharp difference between cholesterol-fed and control rabbits was observed for the uptake of ^{99m}Tc -labeled- β -VLDL by the heart. While in controls the highest uptake occurred for native β -VLDL, in cholesterol-fed rabbits the copper-oxidized β -VLDL was more taken up by heart. These differences may be dependent on the different receptor expression in heart in situations of hypercholesterolemia. Heart, as well as skeletal muscle cells, express the VLDL receptor that recognizes β -VLDL (20). Although, the differentiation between the VLDL receptor and the scavenger receptor can not be made in the present study, we may suggest that in the heart of normolipidemic rabbits there is a predominant VLDL receptor activity, while in cholesterol-fed rabbits a greater expression of scavenger receptor occurs in this organ. The accumulation of ^{99m}Tc -labeled- β -VLDL by tissues with low LDL receptor activity, such as spleen and bone marrow, indicates the uptake into the reticulo-endothelial system possibly by the scavenger receptor, LRP or phagocytosis.

At the present stage of investigation it may be concluded that copper-catalyzed oxidation of β -VLDL induces changes in the biodistribution and plasma clearance of this lipoprotein leading to its accumulation in the heart and atherosclerotic lesions, that can be a crucial event in the atherogenic process. In contrast, β -VLDL modified *ex vivo* by peroxynitrite does not accumulate in atherosclerotic lesions, although this does not exclude the possibility that since this modification occurs *in vivo* in the subendothelial space it can be important in atherogenesis. Finally, it can be concluded that the biokinetic behavior of oxidized lipoproteins may be altered according to the nature of oxidative modifications induced by different oxidant species.

ACKNOWLEDGMENTS

The authors wish to acknowledge the financial support from Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP). Grant nº 92/4406-3 to D.S.P.A.

REFERENCES

1. Mahley, R.W. 1983. Development of accelerated atherosclerosis: concepts derived from cell biology and animal model studies. *Arch. Pathol. Lab. Med.* **107**: 393-399.
2. Witztum, J.L. 1994. The oxidation hypothesis of atherosclerosis. *Lancet* **344**: 793-795.
3. Morel, D.W., DiCorleto, P.E., and Chilson, G.M. 1984. Endothelial and smooth muscle cells alters low density lipoprotein in vitro by free radical oxidation. *Arteriosclerosis* **4**: 357-364.
4. Hiramatsu, K., Rosen, H. Heinecke, J.W., Wolfbauer G., and Chait, A. 1987. Superoxide initiates oxidation of low density lipoprotein by human monocytes. *Arteriosclerosis* **7**: 55-60.
5. Abdalla, D.S.P., Campa, A., and Monteiro, H.P. 1992. Low density lipoprotein oxidation by stimulated neutrophils and ferritin. *Atherosclerosis* **97** : 149-159.
6. Jessup, W., Darley-USmar, V.M., O'Leary, V.J., and Bedwell, S. 1991. 5-lipoxygenase is not essential in macrophage-mediated oxidation of low-density lipoprotein. *Biochem. J.* **278**: 163-169.
7. Halliwell, B. and Gutteridge, J.M.C. 1985. The importance of free radicals and catalytic metal ions in human diseases. *Mol. Aspec.Med.* **8**: 89-193.
8. Parthasarathy, S.1994. Mechanisms of oxidation of LDL. *In* Modified Lipoproteins in the Pathogenesis of Atherosclerosis. S. Parthasarathy, editor. R.G. Landes Company, Austin,TX. 91-119.
9. Ohara, Y., Peterson, T.E., and Harrison, D.G. 1993. Hypercholesterolemia increases endothelial superoxide anion production. *J. Clin. Invest.* **91**: 2546-2551.
10. Minor, R.L., Myers, P.R., Guerra, R., Bates, J.N., and Harrison, D.G. 1990. Diet-iduced atherosclerosis increases the release of nitrogen oxides from rabbit aorta. *J. Clin. Invest.* **86**: 2109-2116.
11. Padmaja, S. and Huie, R.E. 1993. The reaction of nitric oxide with organic peroxy radicals. *Biochem. Biophys. Res. Commun.* **195**: 539-544.

12. Darley-USmar, V.M., Hogg, N., O'Leary, V.J, Wilson, M.T., and Moncada, S. 1992. The simultaneous generation of superoxide and nitric oxide can initiate lipid peroxidation in human low density lipoprotein. *Free Rad. Res. Commun.* **17**: 9-20.
13. White, R.C., Brock, T.A., Chang, L.Y., Crapo, J., Briscoe, P., Ku, D., Bradley, W.A., Gianturco, S.H., Gore, J., and Freeman, B.A. 1994. Superoxide and peroxynitrite in atherosclerosis. *Proc. Natl. Acad. Sci. USA* **91**: 1044-1048.
14. Beckman, J.S., Beckman, T.W., Chen, J., Marshall, P.A., and Freeman, B.A. 1990. Apparent hydroxyl radical production by peroxynitrite: Implications for endothelial injury from nitric oxide and superoxide. *Proc. Natl. Acad. Sci. USA* **87**: 1620-1624.
15. Harrison, D.G. 1994. Endothelial dysfunction in atherosclerosis. *Basic Res. Cardiol.* **89** (suppl. 1): 87-102.
16. Vallabhajosula, S. and Goldsmith, S. 1990. ^{99m}Tc-Low density lipoprotein: intracellularly trapped radiotracer for noninvasive imaging of low density lipoprotein metabolism in vivo. *Sem.Nucl. Med.* **1**: 68-79.
17. Leitha, T., Hermann, M., Hüttinger, M., Angelberger, P., and Dudczack, R. 1993. Technetium-99m labelled LDL as a tracer for quantitative LDL scintigraphy. I. Tracer purification, in vitro and in vivo long-term stability, in vitro validation and biodistribution. *Eur. J. Nucl. Med.* **20**: 667-673.
18. Lees, A.M., Lees, R.S., Schoen, F.J., Isaacsohn, J.L., Fischman, A.J., McKusick, K.A., and Strauss, W.1988. Imaging human atherosclerosis with ^{99m}Tc-labeled low density lipoproteins. *Arteriosclerosis* **8**: 461-470.
19. Brown, M.S. and Goldstein, J.L. 1986. A receptor-mediated pathway for cholesterol homeostasis. *Science* **232**: 34-47.
20. Moestrup, S.K. 1994. The α_2 -macroglobulin receptor and epithelial glycoprotein-330: two giant receptors mediating endocytosis of multiple ligands. *Biochim. Biophys. Acta* **1197**: 197-213.

21. Havel, R.J., Eder, H.A., and Bradgon, H.J. 1984. The distribution and chemical composition of ultracentrifugally separated lipoproteins in human serum. *J. Clin. Invest.* **34**: 1345-1353.
22. Lees, R.S., Garabedian, H.D., and Lees, A.M. 1985. Technetium-99m low density lipoprotein preparation and biodistribution. *J. Nucl. Med.* **26**: 1056-1062.
23. Mesquita, C.H. 1994. Manual do Programa ANACOMP - Análise compartimental auxiliada por computador. *Publicações IPEN* - Instituto de Pesquisas Energéticas e Nucleares, São Paulo, S.P.
24. Maranhão, R.C., Cesar, T.B., Pedroso-Mariani, S.R., Hirata, M.H., and Mesquita, C.H. 1993. Metabolic behaviour in rats of a nonprotein microemulsion resembling low-density lipoprotein. *Lipids* **28**: 691-696.
25. Matthews, C.M.E. 1957. The theory of tracer experiments with ^{131}I -labeled plasma proteins. *Phys. Med. Biol.* **2**: 36-56.
26. Kovanen, P.T., Brown, M.S., Basu, K., Bilheimer, D.W., and Goldstein, J.L. 1981. Saturation and suppression of hepatic lipoprotein receptors: a mechanism for the hypercholesterolemia of cholesterol-fed rabbits. *Proc. Natl. Acad. Sci. USA.* **78**: 1396-1400.
27. Asai, K., Hayashi, T., Funaki, C., Kuzuya, M., Naito, M., and Kuzuya, F. 1991. Comparison of plasma clearance of low density lipoprotein with β -very low density lipoprotein or acetoacetylated low density lipoprotein in cholesterol-fed rabbits. *Biochem. Inter.* **23**: 327-334.
28. Ylä-Herttuala, S., Palinski, W., Rosenfeld, M.E., Parthasarathy, S., Carew, T.E., Butler, S., Witztum, J.L., and Steinberg, D. 1989. Evidence for the presence of oxidatively modified low density lipoprotein in atherosclerotic lesions of rabbit and man. *J. Clin. Invest.* **84**: 1086-1095.
29. Geng, Y., Kodama, T., and Hansson, G.K. 1994. Differential expression of scavenger receptor isoforms during monocyte-macrophage differentiation and foam cell formation. *Arterioscler. Thromb.* **14**: 798-806.

30. Haberland, M.E., Olch, C.L., and Fogelman, A.M. 1984. Role of lysines in mediating interaction of modified low density lipoproteins with the scavenger receptor of human monocyte macrophages. *J. Biol. Chem.* **259**: 11305-11311.
31. Parthasarathy, S., Quinn, M.T., Scwenke, D.C., Carew, T.E., and Steinberg, D. 1989. Oxidative modification of beta-very low density lipoprotein. *Arteriosclerosis* **9**: 398-404.
32. Graham, A., Hogg, N., Kalyanaraman, B., O'Leary, V., Darley-Usmar, V., and Moncada, S. 1993. Peroxynitrite modification of low density lipoprotein leads to recognition by the macrophage scavenger receptor. *FEBS Lett.* **330**: 181-185.
33. Beckman, J.S., Ye, Y.Z., Anderson, P.G., Chen, J., Accavitti, M.A., Tarpey, M.M., and White, C.R. 1994. Extensive nitration of protein tyrosines in human atherosclerosis detected by immunohistochemistry. *Biol. Chem. Hoppe-Seyler* **375**: 81-88.
34. Gatti, R.M., Radi, R., and Augusto, O. 1994. Peroxynitrite-mediated oxidation of albumin to the protein-thiyl free radical. *FEBS Lett.* **348**: 287-290.
35. Rubbo, H., Radi, R., Trujillo, M., Telleri, R., Kalyanaraman, B., Barnes, S., Kirk, M., and Freeman, B.A. Nitric oxide regulation of superoxide and peroxynitrite-dependent lipid peroxidation. *J. Biol. Chem.* **269**: 26066-26075.

Table 1: Biodistribution of ^{99m}Tc - β -VLDL in control rabbits.

Tissue	^{99m}Tc - β -VLDL % inj. dose	^{99m}Tc - β -VLDL-Ox % inj. dose	^{99m}Tc - β -VLDL-Per % inj. dose
Liver	0.272 ± 0.080	$0.522 \pm 0.283^*, **$	0.221 ± 0.098
Kidney	0.540 ± 0.050	0.433 ± 0.104	$0.899 \pm 0.274^*, **$
Heart	$0.031 \pm 0.010^{**}$	0.019 ± 0.007	0.018 ± 0.006
Muscle	0.002 ± 0.000	0.002 ± 0.001	0.003 ± 0.001
Adrenal gland	0.082 ± 0.020	$0.176 \pm 0.030^*, **$	0.060 ± 0.067
Lung	0.048 ± 0.008	$0.066 \pm 0.028^{**}$	0.033 ± 0.011
Spleen	0.115 ± 0.040	0.218 ± 0.135	0.214 ± 0.080
Aorta	0.025 ± 0.014	0.016 ± 0.010	0.024 ± 0.011
Bone marrow	0.093 ± 0.021	0.067 ± 0.022	0.087 ± 0.050

Values are mean $\pm$ standard deviation of results obtained from seven rabbits and are expressed in percentage of the injected radioactivity dose per gram of tissue. β -VLDL-Ox, copper-oxidized lipoprotein; β -VLDL-Per, peroxy-nitrite-modified lipoprotein. *Different as compared to native β -VLDL; ** compared to β -VLDL-Ox or β -VLDL-Per.

Table 2: Biodistribution of ^{99m}Tc - β -VLDL in cholesterol-fed rabbits.

Tissue	^{99m}Tc - β -VLDL % inj. dose	^{99m}Tc - β -VLDL-Ox % inj. dose	^{99m}Tc - β -VLDL-Per % inj. dose
Liver	0.108 ± 0.081	$0.262 \pm 0.148^*, **$	0.101 ± 0.064
Kidney	0.521 ± 0.208	0.919 ± 0.483	0.620 ± 0.187
Heart	0.027 ± 0.007	$0.055 \pm 0.033^*, **$	0.015 ± 0.003
Muscle	0.003 ± 0.002	$0.004 \pm 0.001^{**}$	0.002 ± 0.000
Adrenal gland	0.044 ± 0.015	$0.088 \pm 0.040^*, **$	0.029 ± 0.020
Lung	0.061 ± 0.045	$0.184 \pm 0.073^*, **$	0.052 ± 0.066
Spleen	0.090 ± 0.059	$0.372 \pm 0.150^*, **$	0.073 ± 0.050
Aorta	0.023 ± 0.008	$0.036 \pm 0.017^{**}$	0.017 ± 0.006
Bone marrow	0.065 ± 0.018	0.054 ± 0.032	0.080 ± 0.025

Values are mean $\pm$ standard deviation of results obtained from seven rabbits and are expressed in percentage of the injected radioactivity dose per gram of tissue. β -VLDL-Ox, copper-oxidized lipoprotein; β -VLDL-Per, peroxy-nitrite-modified lipoprotein. * Different as compared to native β -VLDL; ** compared to β -VLDL-Per.

Table 3: Removal of native and oxidized ^{99m}Tc -labeled β -VLDL from plasma of cholesterol-fed and control rabbits

^{99m}Tc -Lipoprotein	Fractional clearance rate (FCR, h^{-1})	
	Cholesterol-fed	Control
^{99m}Tc - β -VLDL	$0.24 \pm 0.03^{\#}$	0.33 ± 0.03
^{99m}Tc - β -VLDL-Ox	$0.15 \pm 0.05^{\#}$	0.39 ± 0.06
^{99m}Tc - β -VLDL-Per	$0.33 \pm 0.05^{\#,*,**}$	$0.55 \pm 0.13^{*,**}$

Values are mean $\pm$ standard deviation of results obtained from seven rabbits. β -VLDL-Ox, copper-oxidized lipoprotein; β -VLDL-Per, peroxynitrite-modified lipoprotein. $^{\#}$ Different as compared to control; * Compared to native β -VLDL; ** compared to β -VLDL-Ox.

FIGURE LEGENDS

FIGURE 1: Plasma decay curves of ^{99m}Tc - β -VLDL, ^{99m}Tc - β -VLDL-Ox, and ^{99m}Tc - β -VLDL-Per in control rabbits. Values are mean $\pm$ standard error mean of results obtained from seven rabbits. β -VLDL-Ox, copper-oxidized lipoprotein; β -VLDL-Per, peroxy-nitrite-modified lipoprotein.

FIGURE 2: Plasma decay curves of ^{99m}Tc - β -VLDL, ^{99m}Tc - β -VLDL-Ox, and ^{99m}Tc - β -VLDL-Per in cholesterol-fed rabbits. Values are mean $\pm$ standard error mean of results obtained from seven rabbits. β -VLDL-Ox, copper-oxidized lipoprotein; β -VLDL-Per, peroxy-nitrite-modified lipoprotein.

FIGURE 3: Biodistribution of ^{99m}Tc -labeled β -VLDL 6 hours after injection of tracer. **A**, ^{99m}Tc - β -VLDL in control rabbit; **B**, ^{99m}Tc - β -VLDL in cholesterol-fed rabbit; **C**, ^{99m}Tc - β -VLDL-Ox in control rabbit; **D**, ^{99m}Tc - β -VLDL-Ox in cholesterol-fed rabbit; **E**, ^{99m}Tc - β -VLDL-Per in control rabbit; **F**, ^{99m}Tc - β -VLDL-Per in cholesterol-fed rabbit. 1, liver; 2, kidney; 3, bladder.

FIGURE 4: Autoradiography for ^{99m}Tc - β -VLDL-Ox accumulation in the aorta. **A**, control rabbit; **B**, cholesterol-fed rabbit. The arrow indicates the accumulation of ^{99m}Tc - β -VLDL-Ox in the atherosclerotic lesions in the aorta arch from the cholesterol-fed rabbit.

FIGURE 3

FIGURE 4

FIGURE 3

TRABALHO 4

PLASMA ANTIOXIDANT ENZYMES AND OXIDIZED LIPOPROTEINS IN HYPERCHOLESTEROLEMIC RABBITS

Edson L. Silva, Patricia Moriel, Yi H. Chang and Dulcinea S.P. Abdalla*

Department of Clinical and Toxicological Analyses, Faculdade de Ciências Farmacêuticas, Universidade de São Paulo, São Paulo, S.P., C.P. 66083, CEP 05389-970, Brasil

SUMMARY: The activities of superoxide dismutase (SOD, EC 1.15.1.1), glutathione peroxidase (GPx, EC 1.11.1.9) and the levels of α -tocopherol and oxidized lipoproteins were investigated in the plasma of New Zealand rabbits either before or after cholesterol-diet induced hypercholesterolemia. Plasma SOD activity increased while GPx activity decreased after 60 days of cholesterol feeding. However, in the cholesterol-fed rabbits the release of superoxide dismutase fraction C from vasculature by heparin was lower as compared to control rabbits. The levels of triglyceride hydroperoxides increased in low density and high density lipoproteins after feeding rabbits with the cholesterol-rich diet during 60 days. Also, a trend for increasing cholesteryl ester hydroperoxides was observed in β -very low density and high density lipoproteins. Increase of α -tocopherol concentration (μ M) was observed in very low density and low density lipoprotein fractions, but after normalization of these results by the cholesterol content of lipoprotein particles only the α -tocopherol content of low density lipoprotein remained higher after 60 days of cholesterol feeding. Data suggest that low glutathione peroxidase and superoxide dismutase fraction C activities may facilitate the intravascular lipoprotein oxidation by oxidant species generated by the endothelium or blood cells.

INTRODUCTION

The oxidative modification of lipoproteins may possibly be an additional risk factor for the development of atherosclerosis (1). Therefore, it is important to verify if during the course of hyperlipidemias oxidized lipoproteins may accumulate in blood plasma. An increased residence time of lipoproteins in blood vessels occurs in hyperlipidemias and may facilitate the oxidation of these particles by oxidant species produced by endothelial cells, leukocytes and smooth muscle cells (2, 3, 4). A high susceptibility of lipoproteins of hyperlipidemic

* To whom correspondence should be addressed

subjects to *ex vivo* oxidation has been reported (5, 6). In this regard, the intravascular antioxidant defense system is important to protect lipoproteins against oxidation. The extracellular SOD (EC-SOD) is a tetrameric Cu- and Zn-containing glycoprotein that exists in both the interstitium of tissues and the vascular system (7). EC-SOD in extracellular fluids is in equilibrium between plasma phase and heparan sulfate proteoglycan in the glycocalyx of the endothelial cell surfaces and in the connective tissue matrix (7). The EC-SOD in plasma comprises three fractions: A which lacks affinity, B with weak affinity and C with strong affinity for heparin and that is bound to endothelium cell surface (7). The EC-SOD C is important to protect endothelium against superoxide radicals and other radicals derived from it (8). An excess generation of radical species by endothelial cells and leukocytes occurs in hyperlipidemic patients (9) and cholesterol-fed rabbits (10, 11). Since these free radicals can oxidize lipoproteins (2, 3, 4), it is important to monitor the plasma antioxidant status and the presence of oxidized lipoproteins in blood plasma mainly in hyperlipidemic states. In this work we investigated the activities of plasma antioxidant enzymes and the oxidation of lipoproteins in the course of hypercholesterolemia induced in rabbits by a cholesterol-rich diet.

MATERIAL AND METHODS

Chemicals: Cholesteryl linoleate, trilinolein, glutathione reductase, reduced glutathione, *tert*-butylhydroperoxide, NADPH, hypoxanthine, xanthine oxidase, α -naphthylendiamine, hydroxylamine, sulphanilic acid and α -tocopherol were purchased from Sigma Chem. Co. (St. Louis, M.O., U.S.A.). Chloroform, acetonitrile, methanol, hexane and heptane were from Merck Chem. Co. (Rio de Janeiro, R.J., Brazil). Enzymatic reagent kits for cholesterol and triglyceride determination were a gift of CELM Co. (São Paulo, S.P., Brazil). Heparin (Liquemine®) was from Roche (São Paulo, S.P., Brazil).

Animals: Male New Zealand white rabbits weighing 2.7-3.0 kg were kept on standard chow (controls) or chow containing 1% cholesterol during 2 months. Blood was collected from animals after 12 h fasting in heparinized tubes for enzyme determinations or in tubes containing 1 mg/ml EDTA for lipoprotein isolation.

Analytical Methods: The activity of SOD was determined by the detection of nitrite with sulphanilic acid and hypoxanthine/xanthine oxidase as the superoxide radical generator system, at 25 °C, according to (12); the incubation systems contained: 66 μ mol/ml hydroxylamine, 3.3 μ mol/ml hypoxanthine, 0.5 U/ml xanthine oxidase and sample (10 mg of protein) in borate buffer pH 8.2. GPx activity was determined following the oxidation of

NADPH at 340 nm, in a system containing 1 mM reduced glutathione, 0.5 U/ml glutathione reductase, 0.2 mM *tert*-butylhydroperoxide and sample (100 µg of protein) in phosphate buffer pH 7.4, at 37 °C, as described by (13). Both measurements were done in a spectrophotometer Hitachi U-3210 (Hitachi Co., Tokyo, Japan). The release of EC-SOD C from vascular endothelium was evaluated by the intravenous heparin injection (2000 UI/Kg) in control and hypercholesterolemic rabbits after 12 h fasting. Blood was collected before and 5, 10 and 15 min. after heparin injection. SOD activity was measured as described above. The concentration of α -tocopherol was determined by reverse phase liquid chromatography with electrochemical detection according to (14), at potential = + 600 mV, using a C-18 CG-Nucleosil column, (4.6 x 250) mm, 5 µm (CG Co., São Paulo, Brazil), a mobile phase constituted by methanol:acetonitrile:chloroform (47:47:6, v/v/v) containing 20 mM LiClO₄, at a flow rate of 1 ml/min. Lipoproteins were isolated from plasma by sequential ultracentrifugation (15). Lipid hydroperoxydes of lipoprotein fractions were detected by reverse phase liquid chromatography according to (16) at a potential = - 300 mV, using a C-8 Inertsil column (4.6 x 150 mm), 5 µm (GL Sciences, Inc., Japan); for the quantification of these compounds the absorbance of each peak at 235 nm was considered. Cholesteryl linoleate and trilinolein hydroperoxides were used as standards. A Class LC-10 chromatographic system (Shimadzu Corp., Tokyo, Japan) with LC10AD pumps, a CBM-10A module, a SPD-M10A. photodiode array detector and a L-ECD-6A electrochemical detector was used. The determinations of cholesterol and triglycerides in plasma and lipoproteins were done with enzymatic reagents (CELM Co., São Paulo, Brazil). Statistical comparisons were done by the Wilcoxon test and p-values below 0.05 were considered to be significant.

RESULTS AND DISCUSSION

The total cholesterol, triglyceride and protein concentrations of lipoprotein fractions are shown in Table 1. It can be seen a sharp increase of cholesterol, triglycerides and protein of very low density lipoprotein fraction (VLDL) caused by the cholesterol-rich diet. This is due to the accumulation of β -VLDL which is a VLDL particle enriched with cholesterol and with an electrophoretic migration in the β region in the cellulose acetate electrophoresis (17). The cholesterol and protein content of the low density lipoprotein (LDL) were also increased. A decrease of the high density lipoprotein (HDL) was observed after cholesterol feeding. These data are in accordance with other reports which describe the lipid alterations of cholesterol-fed rabbits (18). Table 2 shows the activities of SOD and GPx in the plasma of rabbits before, 30 and 60 days after feeding rabbits with the cholesterol diet. During cholesterol feeding an enhancement of EC-SOD and a decrease of GPx activities were observed. Similar changes of these antioxidant enzymes have been reported in the heart of

hypercholesterolemic rabbits (19) and quails (20) and also in aortas of cholesterol-fed rabbits (21). However, the release of EC-SOD C by the intravenous heparin injection was reduced in hypercholesterolemic rabbits as compared to controls (Figure 1A and 1B). This finding suggest that the binding of EC-SOD C to heparan sulfate proteoglycan may be affected in hypercholesterolemia. It is described that during the progression of atherosclerosis the heparan sulfate proteoglycan decreases (22) and, in this case, less EC-SOD C would be anchored to the cell surfaces and connective tissue matrix. Furthermore, the increased number of lipoprotein particles present in the blood vessels of hypercholesterolemic rabbits could affect the binding of EC-SOD C to the endothelial surface. In fact, the apolipoprotein E, present in β -VLDL, has two heparin binding domains (23) and could interact with heparan sulfate proteoglycan leading to a decreased EC-SOD C binding. Independently of the mechanism involved in this process, a decreased amount of EC-SOD C on the vasculature would result in less protection of endothelium against superoxide radicals, as well as the inactivation of nitric oxide by its reaction with superoxide radical with formation of peroxynitrite that it is a strong oxidant (24). Accordingly, the inactivation of nitric oxide by superoxide radical may be related to the impairment of vascular relaxation that occurs in hypercholesterolemic subjects (25) and in cholesterol-fed rabbits (8).

The increased production of superoxide radical and nitric oxide in the arteries of hypercholesterolemic rabbits has been described (10, 11) and may facilitate the formation of peroxynitrite and other strong oxidants derived from it (26). Since these oxidant species may oxidize lipoproteins (2, 27), these oxidized lipoproteins could be detected in cholesterol-fed rabbits. In fact, we observed an increase of triglyceride hydroperoxides in LDL and HDL, as well as, a trend for increased amounts in β -VLDL of cholesterol-fed rabbits as compared to their own basal values (Figure 2). A trend for enhancing cholesteryl ester hydroperoxides in HDL and β -VLDL of these rabbits was also observed (Figure 3). The total concentration of α -tocopherol (expressed in μ M) increased either in β -VLDL or LDL fractions after 60 days of cholesterol feeding (Table 3). Since α -tocopherol is incorporated into lipoproteins in the liver, these findings are a consequence of the great amount of β -VLDL produced by the liver of hypercholesterolemic rabbits and the further transformation of part of this lipoprotein fraction in LDL by intravascular lipolysis (17). However, when α -tocopherol values are

expressed in relation to the cholesterol content of each lipoprotein particle (Table 3), it decreases in β -VLDL suggesting a greater consumption of this antioxidant probably due to the peroxidation of these particles by oxidant species produced by the arteries of hypercholesterolemic rabbits (28). Although α -tocopherol concentration in LDL was high after cholesterol feeding, this antioxidant did not completely prevented the oxidation of this lipoprotein. In fact, it is proposed that peroxidation is propagated within lipoproteins by reaction of the α -tocopherol radical with polyunsaturated fatty acids in these particles (29). This mechanism may explain the oxidation of LDL in the cholesterol-fed rabbits even with the enhancement of α -tocopherol caused by the cholesterol rich-diet. Moreover, high α -tocopherol levels are associated with endothelial dysfunction and enhanced intimal proliferation in cholesterol-fed rabbits (30). In conclusion, these results suggest that during hypercholesterolemia a low EC-SOD C activity may contribute for a high generation of oxidant species by the endothelium that may induce lipoprotein oxidation.

ACKNOWLEDGMENTS

The authors wish to acknowledge the financial support from Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP). Grant nº 92/4406-3 to DSPA.

REFERENCES

1. Steinberg, D., Parthasarathy, S., Carew, T.E., Khoo, J.C., and Witztum, J.L. (1989) *New Engl. J. Med.* 320, 915-924.
2. Morel, D.W., DiCorleto, P.E., and Chilsom, G.M. (1984) *Arteriosclerosis* 4, 357-364.
3. Cathcart, M.K., Morel, D.W. and Chilsom, G.M. (1985) *J. Leukocyte Biol.* 38, 341-348.
4. Abdalla, D.S.P., Campa, A., and Monteiro, H.P. *Atherosclerosis* (1992) 97, 149-159.
5. Lavy, A., Brook, G.J., Danker, G., Amor, A.B., and Aviram, M. (1991) *Metabolism* 40, 794-799.
6. Barbosa, D.S.; Maranhão, R.C., Araujo, F.B., Chang, Y.H., Hirata, M.H. and Abdalla, D.S.P. (1995) *Clin. Biochem.* (*in press*).
7. Karlsson, K., and Marklund, S.L. (1987) *Biochem. J.* 242, 55-59.
8. Mügge, A., Elwell, J.H., Peterson, T.E., Hofmeyer T.G., Heistad, D.D., and Harrison, D.G. (1991) *Circ. Res.* 69, 1293-1300.
9. Araujo, F.B., Barbosa, D.S., Maranhão, R.C., and Abdalla, D.S.P. (1994) in *Frontiers of Reactive Oxygen Species in Biology and Medicine* (Asada, K., and Yoshikawa, T.) pp. 465-467, Elsevier Science B.V., Amsterdam, The Netherlands.
10. Minor, R.L.Jr., Myers, P.R., Guerra, R. Jr., Bates, J.N., and Harrison D.G. (1990) *J. Clin. Invest.* 86, 2109-2116

11. Ohara, Y., Peterson, T.E., and Harrison, D.G. (1993) *J. Clin. Invest.* 91, 2546-2551.
12. Oianagui, Y. (1984) *Anal. Biochem.* 142, 290-296.
13. Sinet, P.M., Heikkila, R.E., and Cohen, G. (1980) *J. Neurochem.* 34, 1421-1428.
14. Thurnham, D.I., Smith, E., and Flora, P.S. (1988) *Clin. Chem.* 34, 377-381.
15. Havel, R.J., Eder, H.A., and Bradgon, H.J. (1984) *J. Clin. Invest.* 34, 1345-1353.
16. Terao, J., Shibata, S.S., and Matsushita, S. (1988) *Anal. Biochem.* 169, 415-423.
17. Simionescu, N., Vasile, E., Lupu, F., Popescu, G., and Simionescu, M. (1986) *Am. J. Pathol.* 123, 109-125, 1986.
18. Thompson, K.H., and Zilversmit, D.B. (1983) *J. Nutr.* 113, 2002-2010.
19. Lapenna, D., Boccio, G., Porreca, E., Pennelli, A., Mezzetti, A., Gioia, S., Marzio, L., Ilio, C., and Cuccurullo, F. (1992) *Free Rad. Res. Comms.* 17, 87-96.
20. Godin, D. and Dahlman, D.M. (1993) *Res. Commun. Chem. Pathol. Pharmacol.* 79, 151-165.
21. Boccio, G., Lapenna, D., Porreca, E., Pennelli, A., Savini, F., Felicianin, P., Ricci, G., and Cuccurullo, F. (1990) *Atherosclerosis* 81, 127-135.
22. Stary, H.C., Chandler, A.B., Glagov, S., Guyton, J.R., Insull, W., Rosenfeld, M.E., Schaffer, S.A., Schwartz, C.J., Wagner, W.D., and Wissler, R.W. (1994) *Circulation* 89, 2462-2478.
23. Weisgraber, K.H., Rall, S.C., Jr., Mahley, R.W., Milne, R.W., Marcel, Y.L., and Sparrow, J.T. (1986) *J. Biol. Chem.* 261, 2068-2076.
24. Huie, R.E., and Padmaja, S. (1993) *Free Rad. Res. Comms.* 18, 195-199.
25. Creager, M.A., Cooke, J.P., Mendelsohn, M.E., Gallagher, S.J., Coleman, S.M., Loscalzo, J., and Dzau V.J. (1990) *J. Clin. Invest.* 86, 228-234.
26. Augusto, O., Gatti, R.M., and Radi, R. (1994) *Arch. Biochem. Biophys.* 310, 118-125.
27. White, C.R., Brock, T.A., Chang, L.Y., Crapo, J., Briscoe, P., Ku, D., Bradley, W.A., Gianturco, S.H., Gore, J., Freeman, B.A., and Tarpey, M.M. (1994) *Proc. Natl. Acad. Sci. USA* 91, 1044-1048.
28. Hogg, N., Darley-Usmar, V.M.; Wilson, M.T., and Moncada, S. (1993) *FEBS Lett.* 326, 199-203.
29. Bowry, V.W., Ingold, K.U., and Stocker, R. (1992) *Biochem. J.* 288, 341-344.
30. Keaney, J.F.Jr., Gaziano, J.M., Xu, A., Frei, B., Curran-Celentano, J., Shwaery, G.T., Loscalzo, J., and Vita, J.A. (1994) *J. Clin. Invest.* 93, 844-851.

Figure Legends

Figure 1: Release of EC-SOD C from vasculature to the plasma by intravenously injected heparin. Enzymatic activities were measured 60 days after feeding the rabbits with 1% cholesterol-diet. A, EC-SOD activity before and after heparin injection. B, percentage of EC-SOD fraction C released by heparin considering the basal EC-SOD activity as 100%. Results represent mean $\pm$ S.E.M. obtained from 17 rabbits. * Different from value before heparin injection.

Figure 2: Triglyceride hydroperoxides in lipoprotein fractions before and after 60 days of cholesterol feeding. Results represent mean $\pm$ S.E.M. obtained from 14 rabbits. * different from basal values. Light gray columns, basal values; dark gray columns, after 60 days of cholesterol feeding.

Figure 3: Cholesteryl ester hydroperoxides in lipoprotein fractions before and 60 days after cholesterol feeding. Results represent mean $\pm$ S.E.M. obtained from 14 rabbits. Light gray columns, basal values; dark gray columns, after 60 days of cholesterol feeding.

Abbreviations: LDL, low density lipoprotein; VLDL, very low density lipoprotein; HDL, high density lipoprotein; SOD, superoxide dismutase; EC-SOD C, superoxide dismutase fraction C; GPx, glutathione peroxidase.

Table 1 - Cholesterol, triglycerides and protein concentration of lipoprotein fractions.

Cholesterol-diet	Cholesterol (mg/dl)	Triglycerides (mg/dl)	Protein (mg/ml)
Basal			
VLDL	3.7 ± 0.7	41.7 ± 3.8	0.03 ± 0.02
LDL	8.3 ± 1.5	40.9 ± 3.8	0.10 ± 0.06
HDL	10.4 ± 1.0	33.9 ± 3.5	1.27 ± 0.17
30 days			
β-VLDL	567.3 ± 43.1*	133.7 ± 71.6	11.48 ± 1.97*
LDL	176.4 ± 42.2*	14.1 ± 3.4*	0.48 ± 0.10*
HDL	23.2 ± 3.2 *	10.6 ± 2.4*	0.87 ± 0.15*
60 days			
β-VLDL	911.8 ± 89.0*	235.2 ± 50.5*	16.91 ± 1.74*
LDL	97.0 ± 16.4	25.1 ± 7.0	0.50 ± 0.07*
HDL	16.8 ± 1.8	22.1 ± 1.5*	0.52 ± 0.06*

The results are mean ± S.E.M. from 15 rabbits; *different from basal values.

Table 2 - Plasma superoxide dismutase and glutathione peroxidase activities of hypercholesterolemic and control rabbits.

	Superoxide Dismutase (U/mg of protein)	Glutathione Peroxidase (U/mg of protein)
Basal	0.58 ± 0.01	16.4 ± 1.3
30 days	0.64 ± 0.02	12.4 ± 0.9
60 days	0.68 ± 0.02	10.7 ± 0.6

The results are mean $\pm$ S.E.M. obtained from 17 rabbits.

Table 3 - α -Tocopherol content of lipoprotein fractions before and 60 days after cholesterol feeding

Lipoprotein	Basal		60 days	
	μM	$\eta\text{mol/mg cholesterol}$	μM	$\eta\text{mol/mg cholesterol}$
VLDL	7 ± 2	40 ± 7	$267 \pm 43^*$	$22 \pm 4^*$
LDL	18 ± 5	23 ± 4	$134 \pm 29^*$	$58 \pm 11^*$
HDL	70 ± 22	81 ± 34	59 ± 12	143 ± 32

The results are mean $\pm$ S.E.M. obtained from 14 rabbits; *different from basal values.

Figure 1A

Figure 1B

Figure 2

Figure 3

TRABALHO 5

UPTAKE OF PEROXYNITRITE MODIFIED β -VLDL BY MONOCYTES OF NORMOLIPIDEMIC AND HYPERCHOLESTEROLEMIC RABBITS

Edson L. Silva & Dulcinea S.P. Abdalla*

Faculty of Pharmaceutical Sciences, University of São Paulo, São Paulo, S.P., Brazil

Summary: The objective of this study was to investigate the uptake of peroxynitrite-modified β -VLDL by monocytes isolated from blood of normolipidemic rabbits (control) and from rabbits fed with a cholesterol-rich diet. The uptake of this modified lipoprotein by monocytes was compared to that of copper-oxidized and non oxidized β -VLDL. 125 I-iodine labeled lipoproteins were incubated with monocytes and intracellular radioactivity was measured after 4 h incubation at 37 °C. Degradation of radiolabeled lipoproteins by monocytes was determined by the trichloroacetic-soluble radioactivity released to the incubation medium. Monocytes of controls internalized peroxynitrite-modified β -VLDL in a greater extent than native and copper-oxidized lipoproteins. In contrast, an enhanced uptake of native β -VLDL by monocytes of hypercholesterolemic rabbits was found. Copper-oxidized β -VLDL was less internalized than native and peroxynitrite-modified β -VLDL by monocytes of both hypercholesterolemic rabbits and controls. Degradation of peroxynitrite-modified β -VLDL by monocytes of hypercholesterolemic animals was greater than that of monocytes from controls. These results indicate that hypercholesterolemia may induce changes on the uptake and degradation of peroxynitrite-modified β -VLDL by monocytes.

* To whom correspondence should be addressed: Faculdade de Ciências Farmacêuticas, Universidade de São Paulo, CP 66083, CEP 05389-970, São Paulo, S.P., Brasil.

Fax: (5511)- 813-2197

β -Very low density lipoprotein (β -VLDL) is a cholesteryl ester enriched lipoprotein that accumulates in the plasma of cholesterol-fed rabbits and in patients with type-III hyperlipoproteinemia (1). This lipoprotein is atherogenic since it is able to induce the transformation of macrophages in foam cells. However, oxidative modifications of β -VLDL may increase its atherogenicity (2). In hypercholesterolemic rabbits an increased production of nitric oxide (3) and superoxide radicals (4) by the arteries has been reported. These two free radicals can react with each other to form peroxynitrite (5). This latter specie was already detected in endothelial cells (6), macrophages (7) and neutrophils (8). Peroxynitrite is a strong oxidant and can oxidize low density lipoprotein (LDL) (9) and β -VLDL (10), possibly making them more atherogenic (9, 10). LDL oxidized by peroxynitrite is recognized by the scavenger receptors and stimulates cholesterol esterification by macrophages (11), a critical event in foam cell formation. Since monocytes can enter the subendothelial space and are the precursors of macrophages in the intima of arteries, we investigated the uptake of peroxynitrite-modified (Per β -VLDL), as well as copper-oxidized (Ox β -VLDL) and native β -VLDL by monocytes isolated from blood of normolipidemic and hypercholesterolemic rabbits. Our data suggest that hypercholesterolemia may affect the uptake and intracellular degradation of Per β -VLDL by monocytes.

MATERIALS AND METHODS

Animals: Male New Zealand white rabbits weighing 2.7-3.0 kg were kept on standard chow (controls) or chow containing 1% cholesterol during 2 months. After this period, blood was collected from animals in heparinized tubes for monocyte isolation or tubes containing EDTA (1 mg/ml) for β -VLDL separation.

Lipoprotein isolation and oxidation: β -VLDL (density <1.019) was isolated from freshly drawn plasma of hypercholesterolemic rabbits by sequential ultracentrifugation (12). Labeling of β -VLDL with 125 -I-iodine was done according to (13). β -VLDL was incubated at a concentration of 1 mg/ml (50 mM phosphate buffer, pH 7.4) in 200 μ M copper sulfate or 400 μ M for 18 h, at 37 °C. Oxidation of β -VLDL was assessed by measurement of thiobarbituric acid-reactive substances (TBARS) (14) and electrophoretic mobility in 0.5 % agarose gel.

Isolation of monocytes: Mononuclear cells were isolated by Ficoll-Hypaque density gradient. Monocytes were recovered by adherence on plastic dishes for 30 min. and maintained in RPMI-1640 medium supplemented with 10 % fetal calf serum, 100 U/ml penicillin G, and 100 µg/ml streptomycin during 12 h, at 5% CO₂. Viability of cells was > 95% as determined by trypan blue exclusion and monocyte purity was > 90%.

Uptake and degradation assays: Monocytes were incubated with radiolabeled Per-, Ox- or native β -VLDL at indicated concentrations for 4 h, at 37 °C, to determine uptake and degradation of ¹²⁵I-lipoproteins. The medium was removed and the extent of lipoprotein degradation was determined by measuring the amount of ¹²⁵I-labeled trichloroacetic-soluble non-iodide material (15). The radioactivity of dishes without cells containing the same amounts of radiolabeled lipoproteins, corresponding to fragmentation, was subtracted from those with cells for final calculations. Cells were washed once with 50 mM Tris-HCL, 150 mM NaCl, 2 mg/ml BSA, pH 7.4 and, twice with the same buffer without albumin. Adhered cells were lysed with 0.1N NaOH and the radioactivity corresponding to the amount of radiolabeled lipoproteins internalized by the cells was determined. Total cell protein was measured according to (16).

RESULTS AND DISCUSSION

Cell internalization of native β -VLDL occurs by the B/E receptor, as well as by the LDL receptor-related protein (LRP) and the VLDL receptor through the binding of apolipoprotein E (17). Monocytes express both the B/E and the LRP receptors (18). In relation to the uptake of oxidized lipoproteins, monocytes express the I and II isoforms of scavenger receptor (19) and also the CD 36, a 88 kDa glycoprotein that was recently identified as an OxLDL receptor (20). In this study Per β -VLDL was more intensively internalized than native and Ox β -VLDL by monocytes of control rabbits (Figure 1). In contrast, monocytes isolated from hypercholesterolemic rabbits internalized two fold less Per β -VLDL and twice more native β -VLDL than cells from controls, at the maximal lipoprotein concentration studied (60 µg of protein/ml) (Figure 2). The low uptake of Ox β -VLDL by cells either of hypercholesterolemic rabbits or of controls was expected, since this particle is internalized by the scavenger receptors (2) and the expression of these receptors on the surface of peripheral blood monocytes is low (19). In the cholesterol-fed rabbit, B/E receptor activity is both suppressed and saturated by the high concentration of β -VLDL

(21). Therefore, this receptor should not contribute to the increased internalization of native β -VLDL by monocytes of hypercholesterolemic rabbits. It is possible that during hypercholesterolemia monocytes express other receptors that are able to bind and internalize β -VLDL and are not down regulated by cholesterol. The monocyte is the only blood cell that express LRP receptors (18). Furthermore, LRP expression is not regulated by the sterol content of the cells (22) and could be mediating the uptake of native β -VLDL by monocytes of hypercholesterolemic rabbits. A 190 kDa protein expressed in human monocytes is also able to bind β -VLDL and may be another likely candidate receptor (23).

A relevant finding was the substantially high degradation of Per β -VLDL by monocytes of hypercholesterolemic rabbits in comparison to that of cells from controls (Figure 3). This suggests that the lower amount of intracellular radioactivity found in monocytes of hypercholesterolemic rabbits resulted from an intense proteolysis of Per β -VLDL by these cells during incubation. Accordingly, an increased degradation rate of low density lipoprotein by monocytes isolated of patients with coronary artery disease was previously reported (24). When the total lipoprotein degradation is considered, i.e., the sum of extracellular fragmentation plus the intracellular proteolysis, it is possible to see that Per β -VLDL degradation is greater than that of native β -VLDL in hypercholesterolemic rabbits (Figure 4) as compared to controls (Figure 5). Therefore, if this occurs during hyperlipidemia in vivo, monocytes may contribute to remove Per β -VLDL formed intravascularly or in the subendothelial space. Since Per β -VLDL is possibly atherogenic (9), the removal of this particle by monocytes may be a mechanism to protect the body against atherosclerosis.

ACKNOWLEDGMENTS

The authors wish to acknowledge the financial support from Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP). Grant no.92/4406-3 to DSPA.

REFERENCES

1. Mahley, R.W. (1978) In Disturbances in lipid metabolism (A.M. Gotto and Ontko, J.A., Eds.), pp. 181-201. Am. Physiol. Soc., Bethesda, MD.
2. Parthasarathy, S., Quinn, M.T., Scwenke, D.C., Carew, T.E., and Steinberg, D. (1989) *Arteriosclerosis* 9, 398-404.
3. Minor, R.L., Myers, P.R., Guerra, R., Bates, J.N., and Harrison, D.G. J. (1990) *Clin. Invest.* 86, 2109-2116.
4. Ohara, Y., Petersom, T.E., and Harrison, D.G. (1993) *J. Clin. Invest.* 91, 2546-2551.
5. Huie, R.E. and Padmaja, S. (1993) *Free Rad. Res. Comms.* 18, 195199.
6. Marletta, M. (1989) *Trends Biochem. Sci.* 14, 488-492.
7. Hibbs, J.B., Taintor, R.R., Vavrin, Z., and Rachlin, E.M. (1988) *Biochem. Biophys. Res. Commun.* 157, 87-94.
8. Carreras, M.C., Pargament, G.A., Catz, S., Poderoso, J.J., and Boveris, A. (1994) *FEBS Lett.* 341, 65-68.
9. Darley-USmar, V.M., Hogg, N., O'Leary, V.J., Wilson, M.T., and Moncada, S. (1992) *Free Rad. Res. Comms.* 17, 9-20.
10. White, C.R., Brock, T.A., Chang, L.Y., Crapo, J., Briscoe, P., Ku, D., Bradley, W.A., Gianturco, S.H., Gore, J., Freeman, B.A., and Tarpey, M.M. (1994) *Proc. Natl. Acad. Sci. USA* 91, 1044-1048.
11. Graham, A., Hogg, N., Kalyanaraman, B., O'Leary, V., Darley-USmar, V., Moncada, S. (1993) *FEBS Lett.* 330, 181-185.
12. Havel, R.J., Eder, H.A., Bradgon, H.J. (1984) *J. Clin. Invest.* 34, 1345-1353.
13. Bilheimer, D.W., Eisenberg, S., Levy, R.I. (1972) *Biochim. Biophys. Acta* 260, 212-221.
14. Winterbourn, C.C., Gutteridge, J.M.C., Halliwell, B. (1985) *Free Rad. Biol. Med.* 1, 43-49.
15. Goldstein, J.L. and Brown, M.S. (1983) *Methods Enzymol.* 98, 241-260.
16. Lowry, O.H., Rosebrough, N.J., Farr, A.L., and Randall, R.J. (1951) *J. Biol. Chem.* 193, 265-269.
17. Dergunov, A.D. and Rosseneu, M. (1994) *Biol. Chem. Hoppe-Seyler* 375, 485-495.
18. Moestrup, S.K., Kaltoft, K., Petersen, C.M., Pedersen, S., Gliemann, J., and Christensen, E.I. (1990) *Exp. Cell Res.* 190, 195-203.
19. Geng, Y.J., Kodama, T., and Hansson, G.K. (1994) *Arterioscler. Thromb.* 14, 798-806.
20. Endemann, G., Stanton, L.W., Madden, K.S., Bryant, C.M., White, R.T., and Protter, A.A. (1993) *J. Biol. Chem.* 268, 11811-1816.
21. Kovanen, P.T., Brown, M.S., Basu, S.K., Bilheimer, D.W., and Goldstein, J.L. (1981) *Proc. Natl. Acad. Sci. USA* 78, 1396-1400.
22. Mahley, R.W., Hui, D.Y., Innerarity, T.L., and Weisgraber, K.H. (1981) *J. Clin. Invest.* 68, 1197-1206.
23. Gianturco, S.H., Lin, A.H.Y., Hwang, S.L.C., Young, J., Brown, S.A., Via, D.P., and Bradley, W.A. (1988) *J. Clin. Invest.* 82, 1633-1643.
24. Shi, F., Hurst, P.G., and McNamara, D.J. (1990) *Atherosclerosis* 85, 127-137.

Figure legends

- Figure 1.** Uptake of ^{125}I - β -VLDL (nat), copper-oxidized ^{125}I - β -VLDL (ox) and peroxynitrite-modified ^{125}I - β -VLDL (per) by monocytes isolated from control rabbits. Internalization of labeled lipoproteins was determined as described in Material and Methods. Each point is the mean of triplicate determinations of two experiments.
- Figure 2.** Uptake of ^{125}I - β -VLDL (nat), copper-oxidized ^{125}I - β -VLDL (ox) and peroxynitrite-modified ^{125}I - β -VLDL (per) by monocytes isolated from hypercholesterolemic rabbits. Internalization of labeled lipoproteins was determined as described in Material and Methods. Each point is the mean of triplicate determinations of two experiments.
- Figure 3.** Intracellular degradation of peroxynitrite-modified ^{125}I - β -VLDL by monocytes isolated from hypercholesterolemic rabbits and controls. Degradation of labeled lipoproteins was determined as described in Material and Methods. Each point is the mean of triplicate determinations of two experiments. HCH, hypercholesterolemic rabbits.
- Figure 4.** Total degradation (sum of intracellular degradation plus extracellular fragmentation) of peroxynitrite-modified (per), copper-oxidized (ox) and native (nat) ^{125}I - β -VLDL by monocytes isolated from hypercholesterolemic rabbits. Values obtained for cell degradation were added to those of control dishes without cells and containing the same amount of labeled lipoproteins. Each point is the mean of triplicate determinations of two experiments.
- Figure 5.** Total degradation (sum of intracellular degradation plus extracellular fragmentation) of peroxynitrite-modified (per), copper-oxidized (ox) and native (nat) ^{125}I - β -VLDL by monocytes isolated from control rabbits. Values obtained for cell degradation were added to those of control dishes without cells and containing the same amount of labeled lipoproteins. Each point is the mean of triplicate determinations of two experiments.

Abbreviations: β -VLDL, β -very low density lipoprotein; LDL, low density lipoprotein; VLDL, very low density lipoprotein; Per β -VLDL, peroxynitrite-modified β -VLDL; Ox β -VLDL, copper-oxidized β -VLDL; TBARS, thiobarbituric acid-reactive substances; LRP, LDL receptor-related protein.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

TRABALHO 6

Esterification of cholesterol and lipid transfer in human serum. Effect of lipid peroxidation.

J. Miara^a, R.C. Maranhão^b and D.S.P. Abdalla^a

^aFaculdade de Ciências Farmacêuticas da Universidade de São Paulo, C.P. 66355, São Paulo, SP, Brazil

^bInstituto do Coração, Fundação Zerbini, São Paulo, SP, Brazil

SUMMARY

The effect of lipid peroxidation on cholesterol esterification and transfer of cholesteryl esters, triacylglycerols, and phospholipids among lipoprotein classes was investigated *in vitro* utilizing an emulsion model of triacylglycerol-rich lipoprotein, labelled with radioactive lipids. Peroxidation of the emulsion resulted in diminished cholesterol esterification but lipid transfer was unaffected. In separate experiments, oxidation of low density lipoprotein decreased the transfer of cholesteryl esters.

INTRODUCTION

Peroxidation of lipoproteins can modify their lipid content (1). Also, changes in surface charge and fluidity of these oxidized particles have been demonstrated (1,2). On the other hand, HDL inhibits the cytotoxic effects of oxidized LDL and the presence of HDL also inhibits the oxidative modification of LDL (3). This effect is probably due to a rapid exchange of oxidized lipids from LDL to unoxidized HDL (3). Currently, we evaluated, *in vitro*, the effect of lipid peroxidation on lipid transfer between an emulsion model of triacylglycerol-rich lipoproteins and HDL or LDL from blood serum (4). The effect of lipid peroxidation on cholesterol esterification was also verified. In addition, the transfer of cholesteryl ester from peroxidized LDL to native HDL was also studied.

METHODS

The radiolabeled triacylglycerol-rich emulsions were prepared as previously described (4). These particles were oxidized with Fe²⁺/ascorbate (100:50 μ M) and lipid peroxidation was followed by the TBA test (5). Lipid transfer was measured after incubation of radiolabeled emulsions, oxidized or not, with blood serum and further separation of lipoproteic fractions by ultracentrifugation (6). Cholesteryl ester transfer between radiolabeled LDL, oxidized or not, and native HDL was evaluated by incubating both fractions and

further separating them by precipitation with heparin-manganese (7). Cholesterol esterification was followed by incubating blood plasma with radiolabeled emulsions and measuring the amount of cholesteryl esters formed after fractionation of the incubation mixtures by TLC. Data were analysed by ANOVA.

RESULTS AND DISCUSSION

Transfer of cholesteryl esters, triacylglycerols and phospholipids from emulsions to serum LDL and HDL was not affected by the previous peroxidation of the emulsion. (Table 1).

Table 1

Transfer of cholesteryl ester, triacylglycerol and phospholipid between triacylglycerol-rich emulsions and lipoproteins

Emulsion	15 minutes		60 minutes	
	LDL	HDL	LDL	HDL
<u>Control^b</u>				
CE	2.28 $\pm$ 0.87 ^a	3.10 $\pm$ 1.20	5.06 $\pm$ 5.72	5.23 $\pm$ 2.23
TG	1.26 $\pm$ 1.13	1.11 $\pm$ 0.69	1.99 $\pm$ 0.94	1.25 $\pm$ 0.67
PL	5.94 $\pm$ 2.84	16.04 $\pm$ 2.76	9.90 $\pm$ 2.71	17.07 $\pm$ 2.31
<u>Oxidized^c</u>				
CE	2.21 $\pm$ 1.20	2.60 $\pm$ 1.54	2.13 $\pm$ 1.94	6.01 $\pm$ 1.94
TG	1.21 $\pm$ 0.93	1.63 $\pm$ 2.75	1.52 $\pm$ 0.82	0.99 $\pm$ 0.47
PL	6.62 $\pm$ 2.84	14.54 $\pm$ 4.04	7.78 $\pm$ 2.05	17.82 $\pm$ 3.27

Emulsions containing ¹⁴C-cholesteryl oleate, ³H-phosphatidyl choline or ³H-glycerol trioleate were incubated with human blood serum. After incubation for 15 and 60 minutes, lipoproteins from incubation mixtures were isolated by ultracentrifugation. Lipids from lipoprotein fractions were extracted and separated by thin layer chromatography. Radioactivity of each lipid class was measured in lipoprotein fractions and the percentages of transfer were calculated. (a) Results are means and standard deviation from duplicates of 11 experiments and are expressed as percentage of transfer. (b) Control emulsion not oxidized. (c) Emulsion oxidized with Fe²⁺/ascorbate (100: 50 μ M). CE, cholesteryl esters; TG, triacylglycerols; PL, phospholipids.

In contrast, transfer of cholesteryl esters from oxidized LDL to native HDL decreased, as compared to control non oxidized LDL (7.9 $\pm$ 1.8% versus 11.5 $\pm$ 2.7%, respectively).

This effect may be attributed to oxidative modification of surface components on HDL that could affect the interaction of cholesteryl ester transfer protein with the lipoprotein particles. In addition, we found that cholesterol esterification of oxidized emulsion was also diminished (Table 2).

Table 2
Cholesterol esterification

System	Time (minutes)			
	30	90	120	180
Control (n=9)	nmoles of cholesteryl ester/minute			
mean	0.023	0.027	0.025	0.024
SD	0.05	0.006	0.007	0.004
Oxidized (n=9)				
mean	0.017 ^a	0.018 ^a	0.017 ^a	0.016 ^a
SD	0.002	0.004	0.002	0.002

Results are means of duplicates from 9 experiments with different blood samples. Plasma was incubated with non oxidized (control) or oxidized ³H-cholesterol labelled emulsions, in a shaker, at 37°C. Aliquots were withdrawn at indicated times and the cholesteryl ester formation was measured. (a) p 0,005.

This reaction is catalysed by lecithin: cholesterol acyltransferase (LCAT, EC 2.3.1.43) (8). Since there are two cysteinyl residues in the active site of this enzyme (9), it is conceivable that lipid peroxidation products could have interacted with these residues, thereby inhibiting the enzymatic activity. It is known that oxidized LDL inhibits the intracellular esterification of cholesterol presumably by translocating a lipid component from the oxidized LDL particles across the cell membrane to the cytoplasm (10). In our system the oxidized lipids could have been transferred from the emulsion to HDL wherein the inhibited the LCAT activity. In conclusion, our data lead to the assumption that triacylglycerol-rich lipoproteins from intestinal or hepatic origin (chylomicron and VLDL) when containing oxidized lipids may interfere with the activity of LCAT. This may be physiologically relevant, since peroxidized dietary lipids can be absorbed by the intestine and incorporated in chylomicrons (11).

ACNOWLEDGEMENT: We thank Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) for financial support.

REFERENCES

- 1 Jurgens G, Hoff HF, Chilson GM and Esterbauer H. *Chem Phys Lipids* 1987; 45: 315-336.
- 2 Aviram M, Danker G and Brook G. *Arteriosclerosis* 1990; 10: 559-562.
- 3 Cathcart MK, Chilson GM, Morel DW and McNally AK. *J Leuk Biol* 1986; 40: 273-279.
- 4 Redgrave TG and Maranhão RC. *Biochim Biophys Acta* 1985 ; 835: 104-112.
- 5 Winterbourn CC, Gutteridge JM and Halliwell B. *J Free Rad Biol Med* 1985; 1: 43-49.
- 6 Fielding CJ, Reaven GMM, Liv G and Fielding PE. 1984; *Proc Natl Acad Sci USA* 81: 2512-2516.
- 7 Burstein M and Scholnick H R. *Adv. Lipid. Res.* 1973; 2: 67-108.
- 8 Fielding CJ. 1986; *Methods Enzymol* 129: 783-990.
- 9 Jauhianen M and Dolphin PJP. 1986; *J Biochem* 261: 7032 - 7043.
- 10 Jialal I and Chait A. *J Lipid Res* 1989; 30: 1561-1568.
- 11 Naruszewicz M, Wosny E, Mirkiewicz E, Nowicka G and Szostak WB. *Atherosclerosis* 1987; 66: 45-53.

TRABALHO 7

LIPID TRANSFER AND CHOLESTEROL ESTERIFICATION IN OXIDIZED
EMULSIONS AND LIPOPROTEINS.

Raul Cavalcante Maranhão ^{a,b}, Jefferson Miara^a, Dulcineia Saes Parra Abdalla^{*,a}

*^aDepartment of Clinical and Toxicological Analyses, Faculty of Pharmaceutical
Sciences (FCFUSP), University of São Paulo, São Paulo, S.P., Brazil.*

*^bHeart Institute (INCOR, FMUSP), Faculty of Medicine, University of São Paulo, São
Paulo, S.P., Brazil.*

*Correspondence address:

Dr. Dulcineia Saes Parra Abdalla

Faculdade de Ciências Farmacêuticas

Universidade de São Paulo

Av. Prof. Lineu Prestes, 580 - Cidade Universitária

CP 66083, CEP 05389-970 - São Paulo, S.P., Brazil

FAX: (5511) 813-2197

Abstract

The effects of lipoprotein oxidation on cholesterol esterification and lipid transfer between lipoprotein classes were evaluated. An emulsion that mimics triglyceride-rich lipoproteins, LDL and HDL were labeled with radioactive lipids and aliquots were oxidized by iron/ascorbate or copper (II). Cholesterol esterification was measured by incubating the oxidized and the non-oxidized control emulsion with human plasma. Esterification of cholesterol of the oxidized emulsion was diminished compared with that of the control (oxidized: 0.016 ± 0.002 , control: 0.024 ± 0.004 nmol/ml/min; $p < 0.05$). A negative correlation was found between the amount of triglycerides in the oxidized emulsion and the plasma cholesterol esterification activity. Lipid transfer from the oxidized emulsion or LDL to other lipoproteins was not different compared to the non-oxidized respective controls. However, oxidation of HDL increased the transfer of its [^{14}C]-cholesteryl oleate to LDL. Therefore, oxidative modification may disturb these two important processes of the intravascular lipoprotein metabolism.

Keywords: Lipid transfer; Cholesterol esterification; Lipid peroxidation; Oxidation; Lipoproteins; Emulsions.

1. Introduction

Several in vitro studies have shown that the oxidative modification of lipoproteins may be an important mechanism leading to atherogenesis [1]. Although there is evidence that plasma lipoprotein oxidation may occur in vivo [2], the effects of oxidation on the biochemical events of the lipoprotein intravascular metabolism have been scarcely investigated. In this regard, cholesterol esterification and transfer of lipids among lipoproteins are very important processes occurring in the plasma that can substantially influence the composition and the metabolic fate of those particles [3, 4, 5].

Cholesterol esterification is catalyzed by lecithin:cholesterol acyl transferase (LCAT, EC 2.3.1.4.3) using apo A1 as a co-factor [6]. It occurs mainly in HDL particles, since this lipoprotein fraction contains most of the plasma apo A1 [6]. Cholesterol esterification is important for the compartmentalization of intravascular cholesterol because the esterified cholesterol molecules move from the surface to the lipoprotein core, where they become less diffusible to the cell membranes [5]. The transfer of phospholipids (PL), cholesteryl esters (CE) and triglycerides (TG) among lipoprotein classes is facilitated by the lipid transfer proteins (LTPs) [3]. There are at least two lipid transfer protein classes: cholesteryl ester transfer protein (CETP or LTP I) transfers CE, TG and PL, whereas the phospholipid transfer protein exclusively transfers PL [3].

In the present study, the hypothesis that oxidative modification of lipoproteins could interfere with plasma cholesterol esterification and lipid transfer was tested. To facilitate manipulations of particle composition and the labeling with radioactive lipids, an artificial emulsion system previously shown to mimic the metabolic behavior of triglyceride-rich emulsions [7-11] was used in addition to native plasma HDL and LDL. This emulsion system had been previously validated in in vivo studies on lipid transfer [12]. To assay for cholesterol esterification and lipid transfer, oxidized and non-oxidized

emulsions and plasma lipoproteins labeled with radioactive lipids were incubated with human plasma. Results showed that oxidative modification may influence both processes.

2. Material and Methods

2.1. Material

Cholesterol, cholesteryl oleate and triolein were obtained from Nu-Check Prep.. Phosphatidylcholine was obtained from egg yolk as previously described [13]. Phosphatidylcholine (L- α -dipalmitoyl [choline-methyl- ^3H]) and (1- ^{14}C)-cholesteryl oleate were obtained from Du Pont New Products (New England). Tri-(9,10 <n> - ^3H)-glycerol oleate, (7 <n> - ^3H)-cholesterol and (1- ^{14}C)-cholesterol were purchased from Amersham. All other reagents were of the highest grade commercially available.

2.2. Emulsion Preparation

Emulsions constituted by 69 weight % triolein, 2% cholesterol, 6% cholesteryl oleate and 23% phosphatidylcholine were prepared as described elsewhere [7]. Briefly, lipids and radioisotopes were mixed and evaporated under nitrogen. The residue was resuspended in 1.101 g/ml sodium chloride solution and sonicated at 70-80 watts (Branson sonicator, model 450), under nitrogen flux, during 30 minutes. Emulsions were purified by ultracentrifugation with discontinuous saline gradient and the layer floating at 1.006 g/ml was utilized. Emulsions were labeled with ^3H -phosphatidylcholine, ^3H -triglycerol oleate and ^{14}C -cholesteryl oleate, respectively for phospholipid, triglyceride and cholesteryl ester transfer experiments. For cholesterol esterification assays, emulsion was labeled with ^3H -cholesterol.

2.3. Oxidation

Emulsions were oxidized with 100 μM Fe^{2+} / 50 μM ascorbate in a shaker bath, at 37 $^{\circ}\text{C}$, during 1 h. Oxidation was interrupted by adding 1 mM BHT. Emulsions were repurified by ultracentrifugation as previously described [7] and dialyzed against a solution containing 150 mM NaCl, 10 mM Tris and 1 mM desferrioxamine, pH 7.4. Blood of normolipidemic volunteers was withdrawn with EDTA (1 mg/ml) after 12 h fasting and

lipoproteins were isolated from plasma by sequential ultracentrifugation as described elsewhere [14]. LDL and HDL (500 μ g of protein) were oxidized by incubation with 5 μ M (LDL) or 50 μ M (HDL) copper sulfate using a shaker bath, at 37 °C, during 6 h (LDL) or 24 h (HDL). After oxidation, lipoproteins were dialyzed against the same solution described above. Lipid peroxidation in all systems was monitored by measuring the thiobarbituric acid-reactive substances (TBARS) [15].

2.4. Cholesterol esterification

Aliquots (1.5 ml) of plasma freshly isolated from normolipidemic volunteers were incubated with oxidized and control emulsions (0.58 mg of triglycerides/ ml of plasma) labeled with 3 H- or 14 C- cholesterol, in a shaker bath, at 37 °C. After 30, 90, 120 and 180 min., 100 μ l aliquots were removed and immediately extracted according to Folch [16]. The residues were resuspended in 100 μ l of chloroform:methanol 2:1 (v/v) and lipids were separated by thin layer chromatography using silica gel 60 H plates and hexane:ethyl ether:acetic acid, 70:30:1 (v/v/v) as solvent. Free cholesterol (FC) and cholesteryl ester (CE) bands were removed, dissolved in appropriate liquid scintillation solution (5 g PPO, 0.5 g POPOP, 333 ml Triton X-100 and toluene q.s.p. 1 liter) and radioactivity was measured. The percentage of radioactivity present in CE band was calculated in relation to total radioactivity (CE + FC) and the amount of CE was determined considering the free cholesterol concentration initially present in the emulsion. Cholesterol esterification ratio was determined considering the amount of CE formed per unit of time and is expressed as nmol/ml/min.

2.5. Lipid transfer

Oxidized and control emulsions (1.05 mg of triglycerides) were incubated with 3 ml of plasma (as source of lipid transfer proteins, LDL and HDL) in a shaker bath, at 37 °C. After 15 and 60 min., 1.5 ml aliquots were removed, kept in ice, and lipoproteins were

isolated by ultracentrifugation using a discontinuous saline density gradient, at 4 °C (TH 641 rotor, Sorvall-Combi ultracentrifuge, model OTD, Du Pont, USA) [17]. One hundred μ l aliquot of each isolated fraction (VLDL + emulsion, LDL, HDL and infranatant) was extracted using Folch method [16] and radioactivity was measured in a liquid scintillation spectrometer (Beckman, model LS-100C). Percent of lipid transfer to LDL and HDL was calculated in relation to total radioactivity (sum of all fractions). In the experiments of CE transfer between LDL to HDL, native or oxidized lipoproteins were firstly isolated by sequential ultracentrifugation and labeled with ^{14}C -cholesteryl oleate according to Lasuncion et al. [18]. Systems containing LDL (10 mg of protein), HDL (100 μ g of protein) and lipoprotein-depleted plasma (as source of lipid transfer proteins, density > 1.21 g/ml) were incubated in a shaker bath, at 37 °C, during 3 h. Labeled LDL or HDL were used as CE donors, respectively, to HDL or LDL. After incubation, KBr was added to adjust density to 1.063 g/ml and LDL was isolated after ultracentrifugation at 40.000 rpm (rotor RP 55-T, Hitachi), at 4 °C, during 24 h. The upper (LDL) and lower (HDL) layers were extracted by Folch method [16] and radioactivity was measured in a liquid scintillation spectrometer. Percentage of CE transfer was calculated in relation to total radioactivity of each system.

2.6. SDS-PAGE

HDL electrophoresis was performed in SDS-polyacrylamide gel as described by Laemmli [19]. Aliquots (10 μ g) of HDL were loaded on SDS-PAGE (10-20%) and stained with Coomassie brilliant blue. High molecular weight standard (Pharmacia, Sweden) was: phosphorylase B (94 kDa), albumin (67 kDa), ovalbumin (43 kDa), carbonic anhydrase (30 kDa), trypsin inhibitor (20.1 kDa) and α -lactalbumin (14.4 kDa).

2.7. *Other assays*

Cholesterol and triglycerides were measured by enzymatic reagents in an autoanalyzer (ABA-VP, Abbot Lab.) and protein was determined according to Lowry *et al.* [20].

2.8. *Statistical Analysis*

Data of lipid transfer and cholesterol esterification of emulsion experiments were analyzed by variance analysis and differences were established by multiple comparisons [21]. CE transfer between native and oxidized lipoproteins were analyzed by Wilcoxon test and correlations were determined by Spearman test using the PRIMER software .

3. Results

Cholesterol Esterification

Cholesterol esterification by human plasma was inhibited when oxidized emulsions were used as substrate (Table 1, system A). When emulsions were made with increasing proportions of triglycerides and incubated with plasma to test for esterification, no correlation between emulsion TG mass and esterification. However, when the emulsions were oxidized, an inverse correlation appeared between these two variables ($r = -0.98$, $p = 0.0045$, by Spearman test, Fig. 1). Effect upon esterification values were observed only with TBARS below 0.35 nmol/mg of TG (Table 1, system B). Oxidation changed neither the emulsion lipid composition (Table 2), nor the apolipoprotein adsorption to the emulsion particles (tested by SDS-PAGE; not shown).

Lipid Transfer

Oxidation of the emulsion did not affect the transfer of PL, TG and CE from the emulsion to the other lipoprotein plasma fractions even at high TBARS levels (table 3). For a more complete understanding of the effect of oxidation, we also examined other CE shifts between plasma lipoprotein classes. Even when TBARS were very high (50 ± 2 nmol/mg of protein), CE transfer from oxidized LDL to the HDL fraction was similar to that of non-oxidized LDL (Fig. 2).

In contrast, the opposite flux, e.g., CE transfer from oxidized HDL to the LDL fraction, was increased as compared to the control experiment with non-oxidized HDL (Fig. 2). It is known that at 4 °C CETP is blocked. When non-oxidized LDL was incubated with plasma at 4 °C, the CE transfer from LDL to HDL was considerably diminished compared to the experiment performed at 37 °C (from 11.2 % at 37 °C to 2.8% at 4 °C:). SDS-PAGE showed that after HDL oxidation the apo AI band (28.3 kDa) was no longer

visible at the stained gels. Instead, new bands corresponding to higher molecular weight proteins appeared (Fig. 3).

Table 1
Cholesterol esterification on oxidized and control emulsions catalysed by human plasma.

Incubation Systems	Incubation Time (minutes)			
	30	90	120	180
System A				
Control (n= 9)	0.023 ± 0.005	0.027 ± 0.006	0.025 ± 0.007	0.024 ± 0.004
Oxidized (n= 9)	0.017 ± 0.002*	0.018 ± 0.004*	0.017 ± 0.002*	0.016 ± 0.002*
System B				
Control (n=5)	0.010 ± 0.004	0.015 ± 0.002	0.016 ± 0.003	0.017 ± 0.002
Oxidized (n=5)	0.010 ± 0.002	0.017 ± 0.004	0.014 ± 0.004	0.014 ± 0.004

Results are expressed as mean ± standard deviation of cholesterol esterification ratio (nmol/ml/min.). The set of experiments of system A was done with an oxidized emulsion containing 3 nmol of TBARS/mg of triglyceride and for system B, oxidized emulsion contained 0.3 nmol of TBARS/mg of triglyceride. In each experiment the control and oxidized emulsions were incubated with plasma of the same subject.

*Values different from control, $p < 0.05$.

Table 2
Lipid composition of control and oxidized emulsions.

	Control emulsion (mg/ml)	Oxidized emulsion (mg/ml)
Total lipids	10.50 $\pm$ 0.95	9.90 $\pm$ 1.00
Triglycerides	8.20 $\pm$ 0.78	8.00 $\pm$ 0.70
Total cholesterol	0.60 $\pm$ 0.05	0.50 $\pm$ 0.04
Phospholipids	1.70 $\pm$ 0.15	1.40 $\pm$ 0.12

Results are expressed as mean $\pm$ standard deviation of 3 experiments.

Table 3

Transfer of cholesteryl esters (CE), triglycerides (TG) and phospholipids (PL) from oxidized and control emulsions to LDL and HDL.

	Incubation Time			
	15 minutes		60 minutes	
	LDL	HDL	LDL	HDL
CE Transfer				
Control	5.0 ± 1.2	5.1 ± 1.9	6.4 ± 1.8	9.5 ± 3.9
Oxidized	4.6 ± 2.3	4.6 ± 2.4	5.8 ± 3.7	10.6 ± 3.1
TG Transfer				
Control	2.3 ± 1.9	1.7 ± 1.0	3.4 ± 0.9	2.7 ± 1.1
Oxidized	2.3 ± 1.9	1.4 ± 0.7	3.2 ± 1.6	1.6 ± 0.7
PL Transfer				
Control	11.8 ± 5.3	27.2 ± 5.0	15.7 ± 5.0	29.6 ± 4.0
Oxidized	13.2 ± 5.4	24.4 ± 5.9	15.6 ± 3.8	31.1 ± 1.2

Results are expressed as mean ± standard deviation of percentage of CE, TG and PL transfer from control and oxidized emulsions of eleven experiments with different plasma samples. In each experiment the control and oxidized emulsions were incubated with plasma of the same subject. TBARS of oxidized emulsions was 3 nmol/mg of triglycerides.

4. Discussion

Triglyceride-rich emulsions with defined lipid composition can mimic the steps of the intravascular metabolism of chylomicrons or VLDL [7-12]. The emulsion used in this study was also previously shown to be a valid tool for studying the lipid transfer process [12]. In contact with the plasma, the emulsion acquires from the native lipoproteins all the plasma apolipoproteins, except apo B [7]. A major part of the esterified cholesterol in human plasma is formed by the LCAT-HDL interaction. The free cholesterol from cells and phospholipids transferred from TG-rich lipoproteins to HDL supplies both substrates for cholesteryl ester synthesis by LCAT [3]. Since plasma levels of HDL-cholesterol are inversely correlated to the risk of CAD, it is presumed that an increased rate of cholesterol esterification might be beneficial, because it would help to remove cholesterol from peripheral cells to the liver [5]. In addition, this process could also promote the removal of cytotoxic oxidized lipids from cells.[22]. Inhibition of LCAT activity would decrease the reverse cholesterol transport, resulting in accumulation of the intracellular cholesterol of peripheral tissues, thus favoring atherogenesis.

Three hypothetical mechanisms can account for the decrease of the cholesterol esterification rate elicited by oxidation observed in this study. Firstly, the emulsion structure could be modified by oxidation to such extent that LCAT action would be prevented. Nevertheless, we observed that after oxidation the composition of the emulsion did not change. In addition, Apo A1 which is the co-factor for LCAT activity adhered to the oxidized emulsion as much as to the control emulsion. However, it is possible that non-detected changes in the emulsion structure could have occurred. Secondly, the lipid peroxidation-derived compounds generated in the emulsion particles could modify the LCAT molecule. Thirdly, oxidation could have altered the structure of apo A1 acquired by the emulsion. An important finding in this study was the strong inverse correlation between cholesterol esterification rate and the proportion of triglycerides in the emulsion. As this correlation was observed only with oxidized emulsions, it suggests that the

triglyceride moiety was an important source of lipid peroxidation products after exposure to the oxidant system.

Among the different lipid peroxidation products involved in the effects observed in our study, malondialdehyde and other low molecular weight products cannot be accounted for, since they were removed by the dialysis procedure previously to the incubation experiments. However, hydroperoxides originated by the lipid peroxidation of cholesteryl esters, triglycerides and phospholipids, as well as oxisterols derived from free cholesterol could have interfered with the cholesterol esterification process. During lipid peroxidation, the unsaturated fatty acid acyl chains originate polar hydroperoxides that can contribute to an increased effective diameter of the hydrophobic portion of phospholipid molecules [23]. The dynamic conformation of the phospholipid molecule with peroxidized fatty acids becomes more conical, with a larger effective diameter of its hydrocarbonated portion as compared with the polar head, thus favoring a non-lamellar arrangement [24].

Conformational changes affecting the emulsion surface could disturb the interaction of LCAT with the particle. Furthermore, modification of triglycerides into the emulsion core may also affect the catalytic potential of the enzyme [25]. Data previously reported [25] show that the triglyceride content of the lipoprotein is negatively correlated with the maximal apparent rate of LCAT, while the enzyme affinity for the substrate is positively correlated with the proteins and lipids on the lipoprotein surface. Another possible explanation for the diminished LCAT activity by the oxidation resides in the consumption of the emulsion unsaturated fatty acids by the oxidation process [26], since the rate of unsaturated fatty acid transfer to cholesterol by LCAT is greater than that of saturated fatty acids [27]. As previously reported, LCAT may be inhibited *in vitro* by hydrogen peroxide and oxidized phospholipids [28, 29]. Decreased LCAT activity was also detected in subjects exposed to high oxygen tension [30] and in monkeys fed vitamin E-deficient diet [31]. Accordingly, our data show that substrate oxidation inhibits cholesterol esterification by human plasma LCAT and may explain why certain *in vivo* conditions in

which lipoprotein oxidation is presumably facilitated are accompanied by decreased LCAT activity.

In contrast with the results of cholesterol esterification experiments, the lipid transfer processes were unchanged by emulsion oxidation. This implies that the hydrophobic interactions of the emulsion particles with the lipid transfer proteins and the acceptor lipoproteins (LDL and HDL) were not affected by emulsion peroxidation. However, the shift of cholesteryl esters from HDL to LDL was facilitated by HDL oxidation. This suggests that the interaction of cholesteryl ester with either CETP or LDL was somehow improved by oxidation. The HDL₂-subfraction contains the major part of plasma CETP [32] and is particularly susceptible to oxidation [33]. Accordingly, the apo A1 band of HDL was no longer visible after HDL oxidation while other apolipoprotein bands with higher molecular weight appeared, suggesting occurrence of protein aggregation. This aggregation may be due to apo A1 and A2 dimerization presumptively by the action of hydroxyl radicals on tyrosine residues of these proteins, which facilitates cross-linkings [34]. Another oxidative modification occurring in lipoproteins is the derivatization of lysine E-amino groups by aldehydes with a consequent increasing of their net negative charge [35]. It has been reported that the increase in lipoprotein negative charge enhances its association with CETP [36]. This effect may be important in the case of HDL and could be crucial for increasing cholesteryl ester transfer from HDL to LDL. In contrast, it is reported that CETP did not bind to native LDL but can form complexes with oxidized LDL [37]. However, in our studies no difference was observed between native and oxidized LDL in relation to cholesteryl ester transfer to HDL.

In conclusion, oxidation of LCAT substrates could decrease cholesterol esterification and potentially interfere with the removal of cholesterol and cytotoxic oxidized lipids from cells. Furthermore, the oxidative modification of HDL may increase the transfer of cholesteryl ester to LDL enhancing its atherogenic potential. Considering that oxidative

modification of lipoproteins may occur in the organism [1,2], the confirmation of these effects in *in vivo* experiments should be sought in future studies.

Acknowledgment: This work was supported by a grant from CNPq (Conselho Nacional de Pesquisa e Desenvolvimento).

5. References

- 1 Steinberg D, Parthasarathy S, Carew TE, Khoo JC, Witztum JL. Beyond Cholesterol: Modifications of low-density lipoprotein that increases its atherogenicity. *New Engl J Med* 1989; 320: 915-924.
- 2 Hodis HN, Kramsch DM, Avogaro P, Bittolo-Bon G, Cazzolato G, Hwang J, Peterson H, Sevanian A. Biochemical and citotoxic characteristics of an in vivo circulating oxidized low density lipoprotein (LDL -). *J Lipid Res* 1994; 35: 669-677.
- 3 Tall A, Swenson T, Hesler C, Granot E. Mechanisms of facilitated lipid transfer proteins. In Gotto AM, ed. *Plasma lipoproteins*. Amsterdam: Elsevier Science B.V., 1987; 277-297.
- 4 Eisenberg SJ. Preferential enrichment of larger sized very low density lipoprotein populations with transferred cholesteryl esters. *Lipid Res* 1985; 26: 487-484.
- 5 Davis RA, Helgerved P, Dveland S, Drevon LA. Evidence that reverse cholesterol transport occurs in vivo and requires lecithin-cholesterol acyltransferase. *Biochim. Biophys. Acta* 1982; 689: 410-414.
- 6 Fielding CJ. Mechanisms of action of lecithin:cholesterol acyltransferase. *Methods enzymol* 1986; 129:783-790.
- 7 Redgrave TG, Maranhão RC. Metabolism of protein-free lipid emulsion models of chylomicron in rats. *Biochim Biophys Acta* 1985; 835: 104-112.
- 8 Maranhão RC, Tercyak AM, Redgrave T. Effects of cholesterol content on the metabolism of protein free emulsion modelism of lipoproteins. *Biochim. Biophys. Acta* 1986; 875: 247-255.
- 9 Hirata MH, Oliveira HCF, Quintão ECR, Redgrave, TG. The effects of Triton WR-1339, protamine sulfate and heparin on the plasma removal of emulsion models of chylomicrons and remnants in rats. *Biochim Biophys Acta* 1987; 917: 334-346.

- 10 Oliveira HCF, Hirata MH, Redgrave TG, Maranhão RC. Competition between chylomicrons and their remnants for plasma removal: a study with artificial emulsion models of chylomicrons. *Biochim Biophys Acta* 1988; 958: 211-217.
- 11 Redgrave TG, Maranhão RC, Tercyak AM, Lincoln EC, Brunegaber H. Uptake of artificial remnant lipoprotein emulsions by the perfused rat liver. *Lipids* 1988; 23:101-105.
- 12 Maranhão RC, Roland IA, Hirata MY. Effects of Triton WR-1339 and heparin on transfer of surface lipids from triglyceride-rich emulsions to high density lipoproteins in rats. *Lipids* 1990; 25: 1-5.
- 13 Singleton WS, Gray MS, Brown ML, White JL. Chromatographically homogenous lecithin from egg phospholipids. *J Amer Oil Chem Soc* 1965; 42: 53-56.
- 14 Havel RJ, Efer HA, Bragdon JH. The distribution and chemical composition of ultracentrifugally separated lipoproteins. *J Clin Invest* 1955; 34: 1345-1354.
- 15 Winterbourn CC, Gutteridge JMC, Halliwell B. Doxorubicin-dependent lipid peroxidation at low pressures of O₂. *J Free Rad Biol Med* 1985; 1: 43-49.
- 16 Folch J, Lee M, Stanley GHA. A simple method for the isolation and purification of total lipids from animal tissues. *J Biol Chem* 1957; 226: 497-503.
- 17 Redgrave TG, Roberts DCK, West CE. Separation of plasma lipoprotein by density-gradient ultracentrifugation. *Anal Biochem* 1975; 65: 42-49.
- 18 Lasuncion MA, Iglesias A, Skottova N, Orozco E, Herrera E. High density lipoprotein subpopulation as substrates for the transfer cholesteryl esters to very low density lipoprotein. *Biochem. J.* 1990; 270: 441-449.
- 19 Laemmli UK. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 1970; 227: 680-685.
- 20 Lowry OH, Rosebrough NJ, Farr AL, Randall RJ. Protein measurement with Folin phenol reagent. *J Biol Chem* 1951; 193: 265-275.
- 21 Scott AJ, Knott MA. A cluster analysis method for grouping means in analysis of variance. *Biometrics* 1974; 30: 507-512.

- 22 Alomar Y, Negre-Salvaire A, Levade T, Valdiguie S, Salvayre R. Oxidized HDL are much less cytotoxic to lymphoblastoid cells than oxidized LDL. *Biochim Biophys Acta* 1992; 1128: 163-166.
- 23 Cullis PK, De Kruijff B. Lipid polymorphism and the functional roles of lipids in biological membranes. *Biochim Biophys Acta* 1979; 559: 339-420.
- 24 Mysik V, Ondrias K, Balgavy P. Effect of lipid peroxidation on molecular arrangement of phospholipids in liposomes prepared from egg yolk phosphatidylcholine or total brain lipids. A ^{31}P -NMR study. *Gen Physiol Biophys* 1992; 11: 317-325.
- 25 Sparks DL, Pritchard PH. The neutral lipid composition and size of recombinant high density lipoproteins regulates lecithin:cholesterol acyltransferase activity. *Biochem Cell Biol* 1989; 67: 358-364.
- 26 Seligman ML, Mitamura J, Shera N, Demopoulos HB. Corticosteroid (methylprednisolone) modulation of photoperoxidation by ultraviolet light in liposomes. *Photochem Photobiol* 1979; 29: 549-558.
- 27 Sgouta DS. Fatty acid specificity of plasma phosphatidylcholine: cholesterol acyltransferase. *Biochemistry* 1972; 11: 293-296.
- 28 Mickel SH, Folds Jr EL. The effect of hydrogen peroxide on serum lecithin-cholesterol acyltransferase activity. *Lipids* 1970; 5: 663-664.
- 29 Mickel SH, Foulds Jr EL, Larkin EC. Inhibition of plasma lecithin-cholesterol acyltransferase reaction by hydrogen peroxide and peroxidized lecithin. *Lipids* 1972; 7: 121-124.
- 30 Mickel SH, Foulds Jr EL, Clrak DA, Larkin EC. Effects of pure oxygen atmosphere in vivo on plasma lecithin-cholesterol acyltransferase reaction. *Lipids* 1971; 6: 740-744.
- 31 Mickel SH, Hill PL, Hayes KC. Depression of lecithin-cholesterol acyltransferase esterification in vitamin E-deficient monkeys. *Amer J Clin Nutr* 1975; 28: 994-999.

- 32 Marcel YL, Mcpherson L, Hogue M, Czarnicka H, Zawadski Z, Weech PK, Whitlock ME, Tall AR, Milne RW. Distribution and concentration of cholesteryl ester transfer protein in plasma of normolipidemic subjects. *J Clin Invest* 1990; 85: 10-17.
- 33 Kunitake ST, La Sala KJ, Kane JP. Apolipoprotein A1-containing lipoproteins with pre-beta electrophoretic mobility. *J Lipid Res* 1985; 26: 549-555.
- 34 Salmon S, Santus R, Mazière JC, Aubailly M, Haigle J. Modified apolipoprotein pattern after irradiation of human high-density lipoproteins by ultraviolet B. *Biochim Biophys Acta* 1992; 1128: 167-173.
- 35 Haberland ME, Olch CL, Folgeman AM. Role of lysines in mediating interaction of low density lipoproteins with scavenger receptor of human monocyte macrophages. *J Biol Chem* 1984; 259: 11305-11311.
- 36 Pattnaik NM, Zilversmitt DB. Interaction of cholesteryl ester exchange protein with human plasma lipoproteins and phospholipid vesicles. *J Biol Chem* 1979; 254: 2782-2786.
- 37 Zadawski Z, Milne R, Marcel YL. Cu²⁺-mediated oxidation of dialysed plasma: effects on low and high density lipoproteins and cholesteryl ester transfer protein. *J Lipid Res* 1991; 32: 243-250.

FIGURE LEGENDS:

Fig. 1: Correlation between triglyceride concentration of oxidized emulsions and cholesterol esterification activity of human plasma. Oxidized emulsions labelled with ^{14}C -cholesterol were incubated with HDL (240 μg of protein) and 200 μl of lipoprotein-depleted plasma ($d > 1.21 \text{ g/ml}$) for 90 minutes, at 37 $^{\circ}\text{C}$. Cholesterol esterification was determined as described in material and methods. Data are means of 3 experiments.

Fig. 2: Transfer of cholesteryl esters (CE) between native and oxidized lipoproteins. Native or oxidized donor lipoproteins (LDL and HDL, respectively) were labelled with ^{14}C -cholesteryl oleate and incubated with the native acceptor lipoprotein (HDL and LDL, respectively). CE transfer was determined as described in material and methods. Data represent mean and standard deviation of 5 experiments.

Fig. 3: SDS-PAGE of native and oxidized HDL. Aliquots of HDL (10 μg of protein/lane) were loaded on gel (10-20 %) and stained with Coomassie brilliant blue. Lane 1, native HDL; lane 2, oxidized HDL; lane 3, low molecular weight standard. Molecular mass is indicated in kDa. Apo, apolipoprotein.

Figure 1

Figure 2

Figure 3

TRABALHO 8

SIALIC ACID AND OXIDIZABILITY OF LOW DENSITY LIPOPROTEIN SUBFRACTIONS OF HYPERLIPIDEMIC PATIENTS.

Décio S. Barbosa^a, Raul C. Maranhão^{a,b}, Fernanda B. Araújo^a, Yi H. Chang^a, Mario H. Hirata^a, Dulcineia S.P. Abdalla^a.

Department of Clinical and Toxicological Analyses, Faculty of Pharmaceutical Sciences^a (FCFUSP) and Heart Institute^b (InCor, FMUSP), University of São Paulo, São Paulo, S.P., Brazil

Short Running Title: Sialic acids and oxidizability of LDL subfractions.

Category: Clinical Investigation

Correspondence address:

Dr. Dulcineia Saes Parra Abdalla

Faculdade de Ciências Farmacêuticas

Universidade de São Paulo

Av. Prof. Lineu Prestes, 580 - Cidade Universitária

CP 66083, CEP 05389-970 - São Paulo, SP, Brasil

FAX (5511) 813-2197

¹Décio S. Barbosa is presently at the Department of Clinical Analyses, Faculty of Pharmacy, University of Londrina^c, Paraná, Brazil.

SUMMARY

Objectives: Low density lipoprotein (LDL) does not constitute an homogenous fraction, and it is known that the heavy LDL subfraction is potentially more atherogenic than the light one. Because concentration of LDL subfractions tend to be different in hyperlipidemias, it was verified whether these subfractions can also differ in sialic acid and neutral sugar content, as well as their resistance to oxidation.

Design and Methods: Two subfractions of low density lipoprotein (light LDL, density 1.019-1.034 g/ml and heavy LDL, density 1.034-1.063 g/ml) were isolated from the plasma of 17 patients with hypercholesterolemia, 11 with combined hyperlipidemia, 7 with hypertriglyceridemia and 19 normolipidemic subjects. The content of sialic acids and neutral sugars of apo B was determined, respectively, by the periodate-thiobarbituric acid method and by reaction with phenol. The oxidation of LDL subfractions was determined by exposure to 5 μ M copper (II) followed by the measurement of lipid hydroperoxides production by high performance liquid chromatography with electrochemical detection.

Results: The study groups did not differ in the neutral sugar content of LDL subfractions. However, compared to normolipidemic subjects, the sialic acid concentration of both LDL subfractions was lower in patients with hypercholesterolemia and hypertriglyceridemia and higher in those with combined hyperlipidemia ($p < 0.05$). In the hypercholesterolemia and combined hyperlipidemia groups, the lipid hydroperoxide content (μ M) of heavy LDL was higher than in normolipidemic subjects ($p < 0.05$).

Conclusions: The heavy LDL subfraction was more susceptible to oxidation in the patients with combined hyperlipidemia compared to controls and the other hyperlipidemic groups. The effect of sialic acids on heavy LDL oxidizability seems to vary according to the type of hyperlipidemia.

Keywords: hyperlipidemia, hypercholesterolemia, hypertriglyceridemia, low density lipoprotein subfractions, lipoprotein oxidation, sialic acids, atherosclerosis.

INTRODUCTION

Low density lipoprotein (LDL) is a heterogeneous population of particles ranging from 1.019 to 1.063 g/ml which vary in physical chemical, kinetic and immunological properties. The LDL subfraction profile is influenced by genetic inheritance, sex and age [1]. The determination of the LDL subfraction pattern may be important because certain subspecies seem more atherogenic than others. These subfractions may also differ relative to intravascular metabolism: dense LDL particles from normolipidemic subjects may have lower binding affinity to cellular LDL receptors than the lighter ones, which results in diminished clearance from plasma [2, 3]. In addition, dense LDL of healthy subjects is more susceptible to oxidation *in vitro* [4] and may be more intensively taken-up by macrophages through the scavenger receptor, a process that has been associated with generation of foam cells [5].

The sialic acid content of LDL from patients with coronary heart disease is apparently lower than that of healthy subjects and a negative correlation was found between the capacity of LDL to stimulate intracellular lipid accumulation and its sialic acid content [6]. The metabolic changes into the intravascular compartment caused by accumulation of lipoproteins in the plasma conceivably result in compositional modifications of the LDL subfractions. To examine this issue in respect to sialic acid and neutral sugar content, as well as oxidizability, we isolated two subfractions of low density lipoprotein (light LDL, density 1.019-1.034 g/ml and heavy LDL, density 1.034-1.063 g/ml) from the plasma of dyslipidemic and normolipidemic subjects. Compositional changes of LDL subfractions in hyperlipidemia were found that may be related to their pathophysiological role.

MATERIAL AND METHODS

MATERIALS

Potassium bromide, methanol, chloroform, crystallized phenol, sodium chloride and cyclohexanone were purchased from Merck (Darmstadt, Germany). Cholesterol and triglyceride enzymatic reagent kits were obtained from Boehringer-Mannheim (Mannheim, Germany). Thiobarbituric acid, methyl linoleate, neuraminidase and N-acetylneuraminic acid were purchased from Sigma Chem. Co. (St. Louis, MO, USA). All other reagents were of the highest purity available.

Study groups

Hypercholesterolemic group (HCH; total plasma cholesterol > 6.20 mmol/L): 17 patients, 7 male and 10 female; aged (mean $\pm$ standard deviation) 55 ± 8 yr; total cholesterol (mean $\pm$ standard deviation): 7.67 ± 1.53 mmol/L; triglycerides (mean $\pm$ standard deviation): 1.77 ± 0.37 mmol/L; phospholipids (mean $\pm$ standard deviation): 2.89 ± 0.40 mmol/L.

Hypertriglyceridemic group (HTG; total triglycerides > 2.90 mmol/L): 7 patients, 5 male and 2 female; aged 51 ± 9 yr; cholesterol: 5.02 ± 1.09 mmol/L; triglycerides: 4.72 ± 3.15 mmol/L; phospholipids: 2.52 ± 0.60 mmol/L.

Combined Hyperlipidemic group (CHL; cholesterol > 6.20 and triglycerides > 2.90 mmol/L): 11 patients, 7 male and 4 female; phenotype IIb; aged 52 ± 9 yr; cholesterol 7.67 ± 0.67 mmol/L; triglycerides: 5.65 ± 2.78 mmol/L; phospholipids: 2.87 ± 0.68 mmol/L).

Control group (NL; cholesterol < 6.20 mmol/L, triglycerides < 2.90 mmol/L): 19 healthy volunteers, 7 male and 12 female; aged 42 ± 7 yr; cholesterol: 4.86 ± 1.12 ; triglycerides: 1.29 ± 0.60 mg/dl; phospholipids: 2.01 ± 0.42 mg/dl.

Patients were selected from the outpatient clinic of the Heart Institute of the University of São Paulo Medical School Hospital. Control group was selected from laboratory staff members. Patients under hypolipidemic drugs had the treatment discontinued for at least one month before

sampling. Secondary causes of hyperlipoproteinemia, including liver, kidney or thyroid diseases, as well as diabetes mellitus, gross obesity, or use of excess alcohol were excluded. The protocol and all procedures followed were approved by the Ethical and Scientific Committee of the Heart Institute.

METHODS

Isolation of LDL subfractions

Freshly drawn venous blood from overnight-fasted hyperlipidemic and healthy subjects was collected into heparinized tubes, from which plasma was separated by low speed centrifugation (1000 g, 15 minutes at 4 °C). Plasma sample densities were adjusted to 1.210 g/ml using solid KBr and lipoprotein fractions were isolated by saline discontinuous density gradients made at room temperature according to (4) with some modifications. Briefly, gradients were formed by addition of 3.0 ml of plasma and 0.5, 1.0, 1.0, 1.0, 2.0, 2.0 and 1 ml, respectively, of NaCl solutions, pH 7.4, with densities (g/ml) 1.182, 1.060, 1.056, 1.045, 1.034, 1.024 and 1.019 in tubes of a Hitachi RPS 40T swinging bucket rotor (Hitachi Co. Japan). The samples were centrifuged in a Hitachi mod. 70P-72 for 24 h at 15 °C and 284,000 g. To ascertain the position of LDL subfractions along the density gradient tube, 20 µl 15 g/L aqueous solution of Coomassie Brilliant Blue R was added to one of the tubes. With the use of reference tube for each subject, fractions were removed by aspiration from top to bottom of tubes as follow: 1.0 ml (VLDL, $d < 1.019$ g/ml), 3.0 ml (light LDL, $1.019 < d < 1.034$ g/ml), 4.0 ml (dense LDL, $1.034 < d < 1.060$ g/ml) and 3.5 ml of infranatant. Each subfraction was dialyzed at 4°C for 24 h against 4 L of 0.9 % (w/v) NaCl solution, pH 7.4. Subfraction densities were determined by refractometry from background salt tubes included during centrifugation. The purity of lipoprotein fractions was verified by 10-20% SDS-polyacrylamide gel electrophoresis. After gels staining with Coomassie Blue only the apo B band was visible for LDL subfraction samples. The contamination of heavy LDL subfractions with LP(a) was also investigated. Apo (a) levels, determined by RIE (Pharmacia, Uppsala, Sweden),

were not statistically different among groups (mean $\pm$ standard deviation: NL, 0.36 ± 0.27 mU/ml; HCH, 0.29 ± 0.16 mU/ml; HTG, 0.15 ± 0.05 mU/ml; CHL, 0.21 ± 0.17 mU/ml).

Chemical Analysis

Lipid and protein analysis

Cholesterol and triglyceride of plasma and LDL subfractions were determined by enzymatic kits (Boehringer-Mannheim, Mannheim, Germany). The protein content of fractions was measured by the method of Lowry et al. [7]. Phospholipids of plasma and lipoprotein fractions were determined according to Bartlett [8].

Sialic acids and neutral sugars

Apolipoproteins were extracted from 1.0 ml of each lipoprotein fraction by adding 9.0 ml of methanol:chloroform 6:5 (v/v). After 2h at 4 °C, apolipoproteins were sedimented by centrifugation at 1,100 g for 5 min. The supernatant was discarded in another assay tube and the sediment was washed once with the same mixture. After centrifugation, the supernatant was joined to the previous one and the pellet containing apolipoproteins was dried under nitrogen. Protein recovery after delipidation was 85%-90%. Neutral sugars of LDL subfractions were determined according to Dubois et al. [9]. The amount of neutral sugars associated to apolipoprotein B was calculated by measuring the total amount of neutral sugars in each LDL subfraction and subtracting the values measured in supernatants obtained after apolipoprotein extraction. Sialic acids were released from apolipoproteins by hydrolysis with 0.1 N sulfuric acid at 80 °C, for 1 h and measured according to Warren [10].

Lipoprotein oxidation

Protein concentration of lipoprotein fractions was adjusted to 0.25 mg/ml with PBS, pH 7.4 and 1 ml was incubated with 5 μ M cupric sulfate using a shaker water bath during 12h, at 37 °C. Then, 0.5 ml aliquots were extracted with methanol (2 ml) and hexane (10 ml). Hexane layer was

removed and evaporated under nitrogen. The lipid residue was dissolved with 0.45 ml of methanol and lipid hydroperoxides (LOOH) were determined by reversed phase HPLC with electrochemical detection [11]. Separation was performed using a 5 μ m C8 Inertsil (150 x 4.6 mm, GL Sciences, Japan) analytical column and isocratic elution, at a flow rate of 1.0 ml/min. The solvent system used was methanol/water (95:5, v/v) with 20 mM lithium perchlorate and 2 mM potassium chloride. The peak profile was monitored with a 1049 HP detector (Hewlett Packard, Germany) set at +900 mV. The lipoprotein LOOH was calculated by comparison to cholesteryl linoleate and trilinolein hydroperoxides standards.

Desialylation of lipoprotein fractions

To verify the effect of desialylation on the lipoprotein susceptibility to oxidation, two different samples of each LDL subfraction were randomly chosen among the samples of each group of patients. These samples were divided in two aliquots; one of them was treated with neuraminidase (0.03 units/ mg of apolipoprotein) to remove sialic acids as previously described by Sattler et al [12]. Both neuraminidase-treated and non-treated lipoprotein fractions (0.3 mg of protein/ ml) were oxidized with 5 μ M cupric sulfate in 100 mM Tris-HCl, pH 7.2, for 6 or 12 h, at 37 °C. The effect of N-acetyl-neuraminic acid (NANA) on copper-induced peroxidation of emulsions constituted by 69% triolein, 2% cholesterol, 6% cholesteryl oleate and 23% egg yolk phosphatidylcholine was evaluated after incubation with 5 μ M copper sulfate for 3h, at 37 °C. Lipid peroxidation was evaluated by measuring the thiobarbituric acid reactive substances (TBARS) as described by Winterbourn et al. [13] and malondialdehyde (MDA) equivalents were calculated by MDA absorption coefficient ($\epsilon_{532}=1.56 \times 10^4 \text{ M}^{-1} \text{ cm}^{-1}$).

Statistical Analysis

The significance of differences between groups was analyzed by Kruskal-Wallis test followed by Dunn test for multiple comparisons. Variation of data of lipoprotein fractions among groups were first analyzed by adjusting to a linear multivariate model. The statistical significance was

considered at the $p < 0.05$ level. Correlation coefficients were determined according to Spearman. Analyses were done using the SAS statistical program package [14].

RESULTS

Comparison of heavy and light LDL subfractions

The content of triglycerides, cholesterol, phospholipids and protein of light and heavy LDL of the study groups is shown in Table 1. The composition of these fractions expressed as mean weight % is also depicted in Figure 1. Since the LDL particle has only one protein molecule (apo B 100), it can be stated that there was a greater number of heavy LDL particles than of the light LDL, since there was more protein in the heavy than in the light subfraction (Fig. 1). The light LDL had a greater content of triglycerides and phospholipids per mg of protein than the heavy one (Table 1). Both subfractions had similar contents of sialic acids per mg of protein and neutral sugars (Table 2).

Comparison of the normolipidemic with the dyslipidemic groups

Table 1 shows that the protein content of heavy LDL subfraction was larger in hypercholesterolemic and combined hyperlipidemia patients than in controls. In addition, both patient groups had also more light LDL protein than controls. In hypertriglyceridemics the protein of both subfractions was similar to that of controls. The cholesterol content in heavy and light LDL is larger in hypercholesterolemics than in controls, but it did not differ in hypertriglyceridemics and combined hyperlipidemia patients. The triglyceride concentration in both LDL subfractions was increased in hypertriglyceridemics and combined hyperlipidemia patients compared to controls. The phospholipid content of both subfractions was similar between the hyperlipidemic groups and controls. In Table 2 it is shown that the sialic acid concentration (by mg of protein) of both subfractions was increased in patients with combined hyperlipidemia and

decreased in hypercholesterolemics and hypertriglyceridemics as compared to controls. The content of neutral sugars was similar between the hyperlipidemic groups and controls (Table 2).

Oxidation of LDL subfractions

A possible way to evaluate the oxidizability of lipoproteins is to induce their oxidation with copper. This parameter was evaluated here considering that it may reflect the susceptibility of lipoproteins to oxidation *in vivo*. Since most of lipid hydroperoxides in oxidized LDL are cholesteryl ester hydroperoxides [15], the concentration of hydroperoxides was expressed by LDL cholesterol content. Lipid hydroperoxides were not detected in lipoprotein samples before copper induced peroxidation. In Table 3, it is shown the lipid hydroperoxide values of LDL subfractions expressed by the cholesterol content of each fraction ($\mu\text{mol}/\text{mg}$ of cholesterol) and also as the absolute amount of LOOH (μM). The light LDL subfraction found in hypercholesterolemics and in patients with combined hyperlipidemia was more resistant to oxidation than that from controls. In contrast, the heavy LDL from the combined hyperlipidemia patients was more sensitive to oxidation than that from controls. In the former group, the heavy LDL subfraction was more susceptible to oxidation than the light one.

A possible correlation between the lipid components of LDL subfractions and their susceptibility to oxidation was investigated through correlation data analyses. There was a negative correlation between lipid hydroperoxides and cholesterol of light LDL from combined hyperlipidemia patients ($r = -0.71$, $p = 0.048$), as well as between lipid hydroperoxides and phospholipids of light LDL from hypertriglyceridemics ($r = -0.54$, $p = 0.048$). No correlation between sialic acids and lipid hydroperoxides was observed in light and heavy LDL. Regarding the two desialylated LDL subfractions samples (Tab. 4), it was observed loss of oxidation resistance in those from HCH, HTG and NL groups but not in those from the CHL group.

Antioxidant effect of NANA.

NANA did not show antioxidant effect on copper-induced oxidation of triglyceride-rich emulsions, even at concentrations much higher than those found in LDL (TBARS values for different NANA concentrations, expressed as % of control without NANA, were: 50 μ M, 109%; 100 μ M, 105% and 150 μ M, 107%). In control experiments, where the same concentrations of NANA were used in the absence of emulsion, it was verified that NANA did not react with thiobarbituric acid at our experimental conditions (data not shown).

DISCUSSION

In our study, heavy LDL was increased in both hypercholesterolemic and combined hyperlipidemia patients and it was the major LDL subfraction in all groups but hypercholesterolemics. Dejager et al [16] also showed predominance of heavy LDL subfractions (density > 1.039 g/ml) in hyperlipidemics compared to normal subjects. Our finding of higher triglyceride content in the light LDL reflects the TG enrichment of LDL particles that ultimately results in diminishing their density [16]. The pathophysiologic process which determines the increase in the heavy LDL subfraction in hyperlipidemic patients is still unclear. It may be related to either enhanced production or decreased catabolism of these particles. Epidemiologically, it seems well established that the population with predominance of heavy LDL is more exposed to risk of developing atherosclerosis, independently of age, sex and relative weight [3, 17, 18, 19]. Thus, our finding of increased heavy LDL can at least in part contribute to the increased atherosclerosis risk linked to hypercholesterolemia and combined hyperlipidemia.

Increased susceptibility to oxidative modification may be involved in the atherogenicity of heavy LDL subfractions [4, 20, 21]. In this study, this oxidizability increase was observed only in combined hyperlipidemia patients. The results of lipid peroxidation of different samples of LDL subfractions showed that they had a different oxidation time course and lipid peroxidation production (TBARS). This may be due to their different contents of PUFA or lipophilic antioxidants. The lipoprotein carbohydrate content, particularly the sialic acids may be also an

important factor in lipoprotein atherogenicity. In this regard, LDL with less sialic acid induces a greater intracellular cholesterol accumulation [6]. In addition, when the lipoprotein is desialylated its uptake by macrophages becomes greater [6]. Our finding that both LDL subfractions had lower sialic acid/protein ratios in hypercholesterolemics and hypertriglyceridemics could then contribute for the increased atherogenicity related to those conditions. A contrasting situation was, however, found in combined hyperlipidemia patients, wherein despite of the high sialic content of heavy LDL this subfraction was the most susceptible to oxidation among all the groups. Interestingly, desialylation of heavy LDL of two combined hyperlipidemia patients resulted in increased resistance to oxidation. In contrast, in the samples of two patients of hypercholesterolemic, hypertriglyceridemic and control groups desialylation lead to increased oxidizability of the subfraction. Despite the small number of samples analyzed, this suggests that sialic acid may play a protective role against heavy LDL oxidation in hypercholesterolemia and hypertriglyceridemia, but not in combined hyperlipidemia. When we investigated the antioxidant effect of a typical sialic acid — N-acetyl-neuraminic acid (NANA) — on copper-induced peroxidation of a triglyceride-rich emulsion, no antioxidant effect of NANA was found. Therefore, it seems that free sialic acids do not act as free radical scavenger or chain breaking antioxidants in a lipidic system. This is suggestive that their binding to apolipoproteins may be necessary for protection against lipoprotein oxidation. Generally, sialic acids delay the protein uptake by the specific liver and macrophage receptors [22]. Apparently, the more sialylated the protein the longer its plasma half life [22]. Thus, sialic acid-enriched lipoproteins may remain longer in the plasma, thus favoring exposure to oxidant species generated by endothelial cells and leukocytes [23].

In conclusion, our findings show a predominance of heavy LDL particles in all groups of hyperlipidemic patients, but the subfraction showed high oxidizability only in the combined hyperlipidemia group. This effect of sialic acids on heavy LDL susceptibility to oxidation seems to vary according to the type of hyperlipidemia.

ACKNOWLEDGMENTS: Supported by funds provided by the Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP) and Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq).

REFERENCES

1. Austin MA, Brunzell JD, Fitch WL, Krauss RM. Inheritance of low density lipoprotein subclass patterns in familial combined hyperlipidemia. *Arteriosclerosis* 1990; 10: 520-530.
2. Nigon F, Lesnik P, Rouis M, Chapman MJ. Discrete subspecies of human low density lipoproteins are heterogeneous in their interaction with the cellular LDL receptor. *J Lipid Res* 1991; 32: 1741-1753.
3. Austin MA, Breslow JL, Hennekens CH, Buring JE, Willet WC, Krauss RM. Low density lipoprotein subclass patterns and risk of myocardial infarction. *JAMA* 1988; 260: 1917-1921.
4. De Graaf J, Hak-Lemmers HLM, Hectors MPC, Demacker PNM, Hendriks JCM, Stalenhoef AFH. Enhanced susceptibility to in vitro oxidation of heavy low density lipoprotein subfraction in healthy subjects. *Arterioscler Thromb* 1991; 11: 298-306.
5. Steinberg D, Parthasarathy S, Carew TE, Khoo JC, Witztum JL. Modifications of low-density lipoprotein that increases its atherogenicity. *N Engl J Med* 1989; 320: 915-924.
6. Orekhov AN, Tertov VV, Mukhin DN, Mikhailenko IA. Modification of low density lipoprotein by desialylation causes lipid accumulation in cultured cells. *Biochem Biophys Res Commun* 1989; 162: 206-211.
7. Lowry OH, Rosebrough NJ, Farr AL, Randall RJ. Protein measurement with Folin phenol reagent. *J Biol Chem* 1951; 193: 265-275.
8. Bartlett GR. Phosphorus assay in column chromatography. *J Biol Chem* 1959; 466-468.
9. Dubois M, Gilles KA, Hamilton JK, Rebers PA, Smith F. Colorimetric method for determination of sugars and related substances. *Anal Chem* 1956; 28: 350-356.
10. Warren L. The thiobarbituric acid assay of sialic acids. *J Biol Chem* 1959; 234: 1971-1975.
11. Terao J, Shibata SS, Matsushita S. Selective quantification of arachidonic acid hydroperoxides and their hydroxy derivatives in reverse-phase high performance liquid chromatography. *Anal Biochem* 1988; 169: 415-423.

12. Sattler W, Kostner GM, Waeg G, Esterbauer H. Oxidation of lipoprotein Lp (a). A comparison with low-density lipoproteins. *Biochim. Biophys. Acta* 1991; 1081: 65-74.
13. Winterbourn CC, Gutteridge JM, Halliwell B. Doxorubicin-dependent lipid peroxidation at low partial pressures of O₂. *J Free Rad Biol Med* 1985; 1:43-49.
14. SAS Institute Inc, SAS/STAT User's guide, version 6, fourth edition, vol. 1 and 2, Cary, NC, 1989.
15. Stocker R, Bowry VW, Frei B. Ubiquinol-10 protects human low density lipoprotein more efficiently against lipid peroxidation than does α -tocopherol. *Proc Natl Acad Sci USA* 1991; 88: 16461650.
16. Dejager S, Bruckert E, Chapman MJ. Heavy low density lipoprotein subspecies with diminished oxidative resistance predominate in combined hyperlipidemia. *J Lipid Res* 1993; 34: 295-308.
17. Marzetta CA, Foster DM, Brunzell J. Relationships between LDL density and kinetic heterogeneity in subjects with normolipidemia and familial combined hyperlipidemia using density gradient ultracentrifugation. *J Lipid Res* 1989; 30: 1307-1317.
18. Slack J, Mills JL. Anomalous low density lipoproteins in familial hyperbetalipoproteinemia. *Clin Chim Acta* 1970; 29: 15-25
19. Sniderman A, Shapiro S, Marpole D, Skinner B, Teng B, Kwiterovich PO. Association of coronary atherosclerosis with hyperapobetalipoproteinemia (increased protein but not normal cholesterol levels in human plasma low density [β] lipoproteins). *Proc Natl Acad Sci USA* 1980; 77: 604-608.
20. De Graaf J, Hendriks JCM, Demacker PNM, Stalenhoef AFH. Identification of multiple heavy LDL subfractions with enhanced susceptibility to in vitro oxidation among hypertriglyceridemic subjects. Normalization after clofibrate treatment. *Arterioscler Thromb* 1993; 13: 712-719.
21. Tribble DL, Holl LG, Wood PD, Krauss RM. Variations in oxidative susceptibility among six low density lipoprotein subfractions of differing density and particle size. *Atherosclerosis* 1992; 93: 189-199.

22. Paulson JC. Glycoproteins: what are the sugar chains for? TIBS 1989; 14: 272-276.
23. Abdalla DSP, Campa A, Monteiro HP. Low density lipoprotein oxidation by stimulated neutrophils and ferritin. Atherosclerosis 1992; 97: 149-156.

TABLE 1. Mean concentration of protein and lipids in LDL subfractions of hyperlipidemic patients and controls.

	Protein (g/L)	Total cholesterol (mmol/L)	Triglycerides (mmol/L)	Phospholipids (mmol/L)
<hr/>				
LDL				
(1.019 < d > 1.034)				
Control	0.38 ± 0.10	1.11 ± 0.09	0.32 ± 0.03	0.34 ± 0.03
HCH	0.60 ± 0.23 *	2.12 ± 0.21 *	0.52 ± 0.07	0.56 ± 0.05
HTG	0.32 ± 0.05	0.94 ± 0.05	0.82 ± 0.05 *	0.33 ± 0.01
CHL	0.46 ± 0.13 *	1.41 ± 0.14	1.23 ± 0.19 *	0.51 ± 0.06
LDL				
(1.034 < d > 1.060)				
Control	0.50 ± 0.14 **	1.12 ± 0.09 **	0.16 ± 0.02 **	0.33 ± 0.03 **
HCH	0.72 ± 0.17 *	1.99 ± 0.10 *, **	0.28 ± 0.03 **	0.50 ± 0.03 **
HTG	0.51 ± 0.08 **	1.06 ± 0.01	0.51 ± 0.01 *, **	0.31 ± 0.01 **
CHL	0.74 ± 0.21 *, **	1.68 ± 0.01 **	0.59 ± 0.09 *, **	0.54 ± 0.08 **

Values are means ± SD of duplicate determinations of each component; control subjects, n = 19; HCH, hypercholesterolemic patients, n = 17; HTG, hypertriglyceridemic patients, n = 7; CHL, patients with combined hyperlipidemia, n = 11. * Values significantly different as compared to controls (p < 0.05).

**Values significantly different from light LDL.

TABLE 2: Mean concentration of sialic acids and neutral sugars in LDL subfractions of hyperlipidemic patients and controls.

	Sialic acids (nmol/mg of protein)	Sialic acids (μ M)	Neutral sugars (μ g/mg of protein)
<hr/>			
LDL			
(1.019 < d > 1.034)			
Control	10.73 $\pm$ 6.80	3.85 $\pm$ 0.57	150.92 $\pm$ 84.49
HCH	9.80 $\pm$ 7.80 *	6.32 $\pm$ 1.45	166.15 $\pm$ 78.09
HTG	9.46 $\pm$ 6.74 *	2.92 $\pm$ 0.77	153.91 $\pm$ 91.13
CHL	13.23 $\pm$ 6.97 *	5.58 $\pm$ 0.78	150.50 $\pm$ 104.16
LDL			
(1.034 < d > 1.060)			
Control	15.63 $\pm$ 6.58	8.18 $\pm$ 0.94**	130.55 $\pm$ 84.49
HCH	14.30 $\pm$ 10.87 *	10.50 $\pm$ 2.21	171.52 $\pm$ 66.6
HTG	8.84 $\pm$ 5.25 *	6.76 $\pm$ 1.55**	129.76 $\pm$ 66.68
CHL	19.79 $\pm$ 10.76 *	13.04 $\pm$ 1.63*,**	119.29 $\pm$ 71.97

Values are means $\pm$ SD of duplicate determinations of each component; control subjects, n = 19; HCH, hypercholesterolemic patients, n = 17; HTG, hypertriglyceridemic patients, n = 7; CHL, patients with combined hyperlipidemia, n = 11; *Values significantly different as compared to controls. **Values significantly different when compared to light LDL (1.019 < d > 1.034) of the same group.

TABLE 3: Mean concentration of lipid hydroperoxides (LOOH) of LDL subfractions of hyperlipidemic patients and controls.

	LOOH ($\mu\text{mol/mg}$ of cholesterol)		LOOH (μM)	
	LDL 1.019<d>1.034	LDL 1.034<d>1.060	LDL 1.019<d>1.034	LDL 1.034<d>1.060
Control	0.94 ± 0.41	0.87 ± 0.43	380 ± 56	344 ± 56
HCH	$0.65 \pm 0.48^*$	0.87 ± 0.27	498 ± 119	$877 \pm 253^*$
HTG	1.51 ± 1.18	1.03 ± 0.65	760 ± 504	498 ± 180
CHL	$0.76 \pm 0.40^*$	$1.28 \pm 0.74^*$	358 ± 64	$688 \pm 189^{*,**}$

Values are means $\pm$ SE of duplicate determinations of each component; light LDL (1.019 < d > 1.034); heavy LDL (1.034 < d > 1.060); control subjects, n = 19; HCH, hypercholesterolemic patients, n = 17; HTG, hypertriglyceridemic patients, n = 7; CHL, patients with combined hyperlipidemia, n = 11. *Values significantly different as compared to controls. **Values significantly different as compared to light LDL of the same group.

TABLE 4: Effect of desialylation on susceptibility of LDL subfractions to oxidation by 5 μ M copper (II).

		6 hr incubation				24 hr incubation			
		NATIVE		DES	Ratio ^c	NATIVE		DES	Ratio ^c
					D/N				D/N
		Tbars ^a	Tbars ^b	Tbars ^b		Tbars ^a	Tbars ^b	Tbars ^b	
Control	LDL _L	1.23	10.31	7.96	0.8	0.69	5.83	7.95	1.4
	LDL _H	0.71	6.63	8.72	1.3	0.45	4.20	2.38	0.6
HCH	LDL _L	0.75	3.63	8.40	2.3	0.60	2.90	1.96	0.7
	LDL _H	0.61	3.65	9.27	2.5	0.34	2.04	2.62	1.3
HTG	LDL _L	0.53	4.65	80.15	17.3	0.23	2.05	25.24	12.3
	LDL _H	0.82	6.13	25.10	4.1	0.50	3.74	12.06	3.2
CHL	LDL _L	0.53	4.51	21.31	4.7	0.49	4.20	7.90	1.9
	LDL _H	1.45	11.58	2.96	0.3	1.04	8.28	1.36	0.2

^a μ mol of MDA equivalents/nmol of sialic acid; ^b μ mol of MDA equivalents/mg of protein; ^cratio between Tbars^b values of desialylated (D) and native (N) samples. DES, desialylated lipoproteins. LDL_L, light LDL ($1.019 < d < 1.034$); LDL_H, heavy LDL ($1.034 < d < 1.060$). Results are triplicate of two samples per group. Measurements were done as described in material and methods.

FIGURE LEGEND

FIGURE 1 - Composition of the heavy and light LDL subfractions in the groups of control (NL, n = 19), hypercholesterolemic (HCH, n = 17), hypertriglyceridemic (HTG, n = 7) and combined hyperlipidemic (CHL, n = 11). PT = protein; CL = cholesterol; TG = ; PL = phospholipids. Values were calculated from duplicate determinations and represent mean weight % of each component.

TRABALHO 9

EVALUATION OF OXIDATIVE STRESS IN PATIENTS WITH HYPERLIPIDEMIA.

Fernanda B. Araujo^a, Décio S. Barbosa^a, Chang Y. Hsin^a, Raul C. Maranhão^{a,b} and Dulcineia S.P. Abdalla^a

^a*Department of Clinical and Toxicological Analyses, Faculty of Pharmaceutical Sciences (FCFUSP) and ^bHeart Institute (Instituto do Coração HC-FMUSP), University of São Paulo, São Paulo (Brazil).*

Keywords: hyperlipidemia; oxidative stress; leukocyte respiratory burst; free radicals; antioxidants.

Correspondence address:

Dr. Dulcineia S.P. Abdalla
Faculdade de Ciências Farmacêuticas
Universidade de São Paulo
Av. Prof. Lineu Prestes, 580, Cidade Universitária
C.P. 66083, CEP 05389-970, SÃO PAULO, S.P., BRAZIL.
FAX (5511) 813-2197

SUMMARY

An antioxidant defense system consisting of enzymes and non-enzymatic compounds prevents oxidative damage of lipoproteins in the plasma. When the activity of this system decreases or the reactive oxygen species (ROS) production increases, an oxidative stress may occur. Since fatty acids and triglyceride-rich emulsions can stimulate leukocytes to produce ROS, it is conceivable that raised plasma triglyceride-rich lipoproteins such as very low density lipoprotein (VLDL) may overload the antioxidant system. To test this hypothesis, we selected 14 patients with combined hyperlipidemia (HLP), in whom low density lipoprotein (LDL) and VLDL levels are elevated, as well as 18 hypercholesterolemic patients (HCH) with increased LDL levels and 19 controls (NL) to examine the trend for an imbalance between the production of oxidative species and the antioxidant defense system as challenged by increased plasma lipids. With this goal, plasma lipoprotein lipid fractions were determined and correlated with the release of ROS by leukocytes monitored by luminol-enhanced chemiluminescence. Plasma β -carotene, α -tocopherol, lycopene and the lipoprotein lipid hydroperoxides were determined by high pressure liquid chromatography with electrochemical detection. HLP had lower plasma superoxide dismutase (SOD) activity (0.04 and 0.11 U/mg protein; $p < 0.05$) as well as lower concentrations of lycopene (0.1 and 0.2 nmol/mg cholesterol; $p < 0.05$) and β -carotene (0.8 and 2.7 nmol/mg cholesterol; $p < 0.05$) in the plasma, as compared to NL. Moreover, HLP showed the highest ROS production by resting mononuclear leukocytes (MN) among the three study groups. When the results of the subjects of the three groups were taken together, the plasma triglyceride concentration was positively correlated to ROS release by resting polymorphonuclear leukocytes (PMN, $r = 0.38$, $p = 0.04$) and MN ($r = 0.56$, $p < 0.005$). Moreover, ROS release by resting MN was positively correlated with VLDL ($r = 0.47$, $p = 0.02$) and LDL ($r = 0.57$, $p = 0.01$) triglycerides. There was also a positive correlation between ROS release by stimulated PMN and VLDL ($r = 0.44$, $p = 0.03$) as well as LDL ($r = 0.53$, $p = 0.01$) triglycerides. High density lipoprotein (HDL) cholesterol showed a negative correlation with ROS release by resting MN ($r = -0.48$, $p = 0.02$) and resting PMN ($r = -$

0.49, $p=0.01$). VLDL susceptibility to copper (II) oxidation was not different among the three groups. Regarding LDL, there was an increased oxidizability in HLP group. Plasma ferritin which may act as a source of catalytic iron for lipid peroxidation, was found to be greater in HLP and HCH than in controls ($p<0.05$). These results suggest that oxidative stress is more likely to occur in HLP than in NL and HCH, since in HLP the release of ROS by leukocytes was greater, while some components of their antioxidant defense system were also decreased. Our finding that the leucocyte ROS production is positively correlated with either VLDL or LDL triglycerides sheds light on a new aspect of the leukocyte activation and oxidative stress in hyperlipidemia.

INTRODUCTION

Plasma lipoproteins are protected against oxidative modification by the antioxidant defense system of the organism. This system is constituted by the enzymes SOD, glutathione peroxidase (GPx) and catalase, as well as hydrophilic antioxidants such as ascorbate, reduced glutathione and urate. Lipoprotein particles carry lipophilic antioxidants, such as tocopherols and carotenoids. All these react with ROS or block free radical chain reactions (1). ROS, which comprises superoxide radical, hydrogen peroxide, hydroxyl radical and other oxygen-centered radicals, are generated in aerobic cells during reduction of molecular oxygen by enzymatic reactions, electron transport chains, and autoxidation of diverse substances (1). These species are maintained at very low steady-state concentration by the antioxidant system, but when their production enhances they may overcome the scavenger capacity of the antioxidant system, resulting in an oxidative stress and damage to biological targets. In this regard, *in vitro* studies demonstrate that lipoproteins can be oxidized by ROS released by the leukocyte respiratory burst in the presence of transition metals, such as iron and copper (2, 3, 4). Moreover, ferritin iron released by superoxide radicals produced by stimulated leukocytes can enhance lipoprotein oxidation by leukocytes (4).

It is then conceivable that when lipoproteins accumulate in the plasma, as in primary hyperlipidemias, the likelihood of them suffering oxidative modification is greater. In this case, both the oxidizable lipid mass and the residence time of the lipoprotein particles in the plasma increase, and unless the antioxidant system is capable of matching this challenge, lipoproteins may become prone to oxidative modification. The chain of events which follows lipoprotein oxidation and how it can ultimately lead to atherogenesis has been demonstrated in *in vitro* experiments by several investigators (5, 6).

In this study, we hypothesized whether there could exist an imbalance in the antioxidant defense system resulting from the accumulation of LDL or VLDL in the plasma. With this goal, the production of ROS by MN and PMN leukocytes, the levels of lipid soluble antioxidants and

ferritin in the plasma, as well as the activities of SOD and GPx were evaluated in subjects with primary hypercholesterolemia, in whom the levels of LDL are elevated, or with combined hyperlipidemia, in whom VLDL levels are also increased. Our results showed that the oxidative stress is more likely to occur in HLP and that elevated triglyceride levels may trigger leukocyte ROS production.

MATERIALS AND METHODS

Materials

Ficoll was purchased from Pharmacia LKB Biotechnology (Uppsala, Sweden). Hypoxanthine, xantine oxidase, catalase, CuZn-SOD, hydroxylamine, α -naphtyldiamine, reduced glutathione, glutathione reductase, t-butyl hydroperoxide, NADPH, sulphanilic acid, luminol, sodium diatrizoate, phorbol miristate acetate, sodium dodecylsulphate, Tris-HCl, α -tocopherol, β -carotene type III and lycopene were products of Sigma Chem. Co. (St. Louis, MO, USA). Chloroform, acetonitrile, methanol, hexane, heptane, enzymatic reagent kits for determination of cholesterol, triglycerides, as well as all the other salts used in buffer preparation were purchased from Merck (Rio de Janeiro, R.J., Brazil). Cobas Mira EIA ferritin kit was purchased from Roche Diagnostic Systems (Hoffmann-La Roche AG, Basel, Switzerland).

Subjects

The study protocol was approved by the Scientific and Ethical Committee of the Heart Institute. Subjects were selected from the outpatient clinic of the Heart Institute (Instituto do Coração, HC-FMUSP). They were classified in two groups:

HCH group: 10 women and 8 men; age: 55 ± 8 years; total plasma cholesterol >240 mg/dl and total triglycerides <250 mg/dl.

HLP group: 6 women and 8 men; age: 52 ± 9 years; total plasma cholesterol >240 mg/dl, total triglycerides >250 mg/dl.

The control group (**NL group**) had 13 women and 6 men; age: 42 ± 7 years, all healthy volunteers from the laboratory staff.

Exclusion criteria: Secondary causes of hyperlipoproteinemia, including liver diseases, renal impairment, hypothyroidism, diabetes mellitus, present smoking habit, gross obesity, history of excess alcohol intake, and infectious or inflammatory diseases. Subjects were not using antilipidemic drugs or antioxidant supplementation for at least one month prior to blood sampling.

Blood samples were withdrawn after 12 h fasting into tubes containing heparin (20 units/ml of blood).

Analytical Methods

MN and PMN leukocytes were isolated from heparinized blood by gradient centrifugation in Ficoll-Hypaque and sedimentation in dextran, as previously described (7). ROS release by resting (basal chemiluminescence) and phorbol miristate acetate (PMA)-stimulated MN and PMN cells was measured by luminol-sensitized chemiluminescence, using a 1900 TR Packard liquid scintillation analyzer (Packard Instr. Co., IL, USA) set up at the chemiluminescence mode. Light emission was followed for up to 60 min. and the integrated chemiluminescence was calculated for this period of time. The activity of plasma SOD was determined by the detection of nitrite with sulphanilic acid and hypoxanthine/xanthine oxidase as superoxide radical generator system, at 25 °C, according to Oyanagui (8); incubation systems contained: 66 $\mu\text{mol/ml}$ hydroxylamine, 3.3 $\mu\text{mol/ml}$ hypoxanthine, 0.5 U/ml xanthine oxidase and sample (10 mg of protein) in borate buffer pH 8.2. The activity of GPx was determined following oxidation of NADPH at 340 nm, in a system containing 1 mM reduced glutathione, 0.5 U/ml glutathione reductase, 0.2 mM t-butyl hydroperoxide and sample (100 μg of protein) in phosphate buffer pH 7.4, at 37 °C, as described by Sinet et al. (9). Both measurements were done in a spectrophotometer Hitachi U-3210 (Hitachi Co., Tokyo, Japan). The levels of α -tocopherol, β -carotene and lycopene in blood plasma were measured by HPLC with electrochemical detection according to Thurnan et al. (10), at potential = +600 mV, using a C-18 CG-Nucleosil column, (4.6 x 250) mm, 5 μm (CG Co., São Paulo, Brazil), a mobile phase constituted by methanol:acetonitrile:chloroform (47:47:6, v/v/v) containing 20 mM LiClO_4 , at a flow rate of 1 ml/min. Lipoproteins were isolated from plasma by ultracentrifugation in saline density gradient as described elsewhere (11). Oxidation of lipoproteins was induced by incubation of aliquots (250 μg of protein) with 5 μM copper sulfate in a shaker, at 37 °C, for 12 h. The extension of lipid peroxidation was estimated by determining the lipid hydroperoxide content of lipoprotein fractions by HPLC according to Terao et al. (12),

at a potential = +900 mV, using a C-8 Inertsil column (4.6 x 150) mm, 5 μ m (GL Sciences Inc., Japan), and methanol:water (98:2, v/v) as mobile phase (flow rate = 1 ml/min.). Cholesteryl linoleate hydroperoxide was used as standard. The chromatographic system used was a 510 pump, a 745 B data integrator and Rheodyne injector from Waters (Millipore Corp., MA, USA) and a electrochemical detector HP 1049A (Hewlett Packard, Walbronn, Germany). Ferritin was determined by using a COBAS Core equipment (Roche Diag. Sys., Hoffmann-La Roche AG, Basel, Switzerland). Determinations of cholesterol and triglycerides in blood plasma and lipoprotein fractions were done with commercial enzymatic reagents using an autoanalyser Technicon, model R4100. Data were analyzed by the test of Kruskal-Wallis followed by the Dunn test for multiple comparisons or Bonferroni t-test using the Statistical Analyses System (SAS) (13). Correlation analyses were done by Spearman test in each group (14) and Pearson test when variables were correlated considering all studied individuals as a single group (13).

RESULTS

Total plasma and lipoprotein fraction lipids.

Table 1 shows the total and lipoprotein lipids of the studied groups. Total plasma cholesterol, as well as VLDL-cholesterol and LDL-cholesterol were higher in both hyperlipidemic groups than in controls. Increased triglycerides were found in plasma, VLDL and LDL of HLP patients as compared to controls. As expected, in the HLP group HDL-cholesterol was lower and HDL-triglycerides were higher than those of NL and HCH ($p < 0.05$, Bonferroni t-test). Also, an inverse correlation between total triglycerides and HDL-cholesterol was found ($r = -0.67$, $p = 0.00003$, Pearson test) in all groups. Plasma phospholipids were higher in HCH and HLP as compared to controls ($p < 0.05$, Bonferroni t-test).

Release of ROS by mononuclear and polymorphonuclear leukocytes

The means $\pm$ S.E.M. of integrated luminol-enhanced chemiluminescence obtained for resting and stimulated leukocytes are shown in Fig. 1 and Fig. 2. SOD and catalase promoted 90% and 70% inhibition of PMN and MN cells luminol-enhanced chemiluminescence, respectively, indicating that both superoxide radical and hydrogen peroxide were present in the incubation systems. Release of ROS by resting MN from HLP was significantly greater than that from HCH patients and controls ($p < 0.05$, Dunn test) (Fig. 1). In stimulated MN of HLP there was an enhanced ROS release as compared to HCH group ($p < 0.05$, Dunn test). Resting and stimulated PMN of HLP released more ROS than those of controls and HCH ($p < 0.05$, Dunn test) (Fig. 2).

The correlational analyses including the values from all studied subjects (Pearson test), showed a significant positive correlation between plasma total triglyceride levels and ROS release by resting MN ($r = 0.56$, $p < 0.005$; Fig. 3), resting PMN ($r = 0.66$, $p < 0.05$; Fig. 4) and stimulated PMN ($r = 0.49$, $p < 0.05$). The statistical significance of these correlations still holds even when the values of the 2 subjects with extreme high plasma triglyceride are not considered (resting MN: $r = 0.74$, $p = 0.00001$; resting PMN: $r = 0.85$, $p = 0.0002$). Moreover, ROS release by resting MN was

positively correlated with VLDL- ($r = 0.47$, $p = 0.02$) and LDL- ($r = 0.57$, $p = 0.01$) triglycerides. There was also a positive correlation between ROS release by stimulated PMN and VLDL- ($r = 0.44$, $p = 0.03$) as well as LDL- ($r = 0.53$, $p = 0.01$) triglycerides. In contrast, plasma total cholesterol levels were not correlated with ROS production by leukocytes. However, HDL-cholesterol showed a negative correlation with ROS release by resting MN ($r = -0.48$, $p = 0.02$) (Fig. 5) and resting PMN ($r = -0.49$, $p = 0.01$) (Fig. 6). No correlation was found between ROS production by stimulated MN and either triglycerides or cholesterol of LDL and VLDL, whereas a positive correlation with HDL-triglycerides ($r = 0.41$, $p = 0.05$) was observed. Chemiluminescence of resting MN was inversely correlated to their GPx activities in NL and HCH groups (NL: $r = -0.94$, $p = 0.005$; HCH, $r = -0.71$, $p = 0.05$, Spearman test).

Antioxidant enzymes

The means $\pm$ S.E.M. of SOD and GPx activities in plasma and leukocytes are shown in table 2. Compared to NL, SOD plasma activity was low in HLP ($p < 0.05$, Bonferroni t-test). The activity of this enzyme was inversely correlated with total cholesterol and total triglyceride levels when the values of all subjects are taken together ($r = -0.36$, $p < 0.05$ for cholesterol; $r = -0.42$, $p < 0.005$ for triglycerides, Pearson test). No differences were observed for intracellular SOD and GPx activities among groups. A positive correlation between SOD and GPx activities of resting MN (and PMN leukocytes was found only in NL group (PMN: $r = 0.74$, $p = 0.04$; MN: $r = 0.71$, $p = 0.04$, Spearman test).

Lipid soluble antioxidants and lipoprotein oxidation

Table 3 shows the means $\pm$ S.E.M. of α -tocopherol, β -carotene and lycopene plasma levels. They were expressed by plasma total cholesterol content since these antioxidants are carried in plasma lipoproteins; micromolar concentrations in lipoprotein particles would not take into account the increased number of these particles occurring in hyperlipidemia or redistribution of antioxidants within the lipoproteins. HLP β -carotene and lycopene plasma levels were significantly lower

compared to NL ($p < 0.05$, Bonferroni t-test). Plasma levels of α -tocopherol were similar in HCH, HLP and NL.

The lipid hydroperoxide content of lipoproteins measured after incubation with copper is also shown in table 3. No lipid hydroperoxides were detected in lipoprotein fractions before copper-catalysed oxidation in our experimental conditions. VLDL susceptibility to oxidation was similar among study groups. Copper-catalysed oxidation resulted in higher LDL lipid hydroperoxides in HLP group than in HCH and controls ($p < 0.05$, Bonferroni t-test).

Ferritin plasma levels

Since our previous in vitro study (4) showed that the presence of ferritin enhances lipoprotein oxidation by stimulated leukocytes, we decided to measure the plasma ferritin to verify if increased levels occur in dislipidemic subjects. The plasma ferritin levels of HLP were higher than those of HCH and NL (mean $\pm$ S.E.M.: HLP, 266 ± 67 ng/ml; HCH, 140 ± 25 ng/ml; NL, 64 ± 13 ng/ml; $p < 0.05$; Bonferroni t-test).

DISCUSSION

Our data indicates that an imbalance in the antioxidant defense system seems to result from the accumulation of LDL or VLDL in the course of hyperlipidemias. This imbalance may result from the greater ROS production by leukocytes induced by hyperlipidemia and the consumption of plasma antioxidants, such as β -carotene, lycopene and EC-SOD leading to an oxidative stress. The positive correlation between total triglycerides and leukocyte ROS production found in our study suggests that triglyceride-rich lipoproteins may have a crucial role in leukocyte activation. Previous studies have shown that ROS production was increased in exclusively hypertriglyceridemic patients. (15, 16). Our data reinforce and expand that observation by establishing a correlation between plasma total triglycerides, as well as LDL-TG and LDL-TG, and ROS in patients with hypertriglyceridemia plus hypercholesterolemia. Moreover, studies suggesting that cholesterol is an important stimulus of leukocyte ROS release (17, 18) also included patients that showed triglyceride levels higher than their respective NL groups.

The mechanism determining the enhanced respiratory burst leading to ROS release by NADPH oxidase in the membrane of leukocytes from hyperlipidemic subjects is not known, but some possibilities may be considered. It is known that cis-polyunsaturated fatty acids stimulate the production of superoxide radical by human neutrophils in the following order: arachidonate > linolenate > linoleate > oleate (19). Among saturated fatty acids, those with carbon numbers 14 to 18 are the greatest stimulators of superoxide production in combination with 1-oleoyl-2-acetyl glycerol (20). It is proposed that the interaction of fatty acids and plasma membrane of leukocytes occurs through hydrophobic binding of the fatty acyl chain to the hydrophobic sites of plasma membrane (21). Signal transduction events involved in the activation of NADPH oxidase multi-protein complex include the phosphorylation of one of its protein components by protein kinase C and translocation from the cytosol to the membrane (22). Arachidonate appears to be an immediate activator of the NADPH oxidase independent on phosphorylation (23).

It is reported that the *in vitro* co-incubation of triglyceride-rich emulsions with leukocytes enhances ROS release by MN (24) and PMN (25). Moreover, oxidized triglyceride-rich emulsions

promote a ten-fold increase of superoxide production by human blood MN as compared to non-oxidized emulsions, this effect being attributed to the peroxidized fatty acids (26). In HLP subjects triglyceride-rich LDL was more susceptible to oxidation and could have the same effect on leukocytes. Free radicals generated in the leukocyte respiratory burst could induce lipoprotein oxidation, specially in those HLP patients with higher plasma ferritin levels. The rationale for this assumption is that LDL oxidation induced by stimulated neutrophils is enhanced in the presence of ferritin (4). This is a consequence of ferritin-iron release by superoxide radicals produced by stimulated leukocytes (4, 26). Iron released from ferritin is then available for the catalysis of lipoprotein peroxidation. The high plasma ferritin levels found in HLP could potentially contribute to enhance lipoprotein oxidation in vivo, when associated to the increased ROS production by leukocytes. This is also a possible mechanistic explanation for a previously epidemiological report showing that high plasma ferritin, reflecting high stored iron levels, is a risk factor for coronary heart disease (27). Since it is known that peroxidized fatty acids can activate phospholipase A₂ (28), a release of arachidonic acid from leukocyte membrane by this enzyme could contribute to the activation of NADPH oxidase. In addition, it is possible that peroxidized free fatty acids generated by lipolysis of triglyceride-rich lipoproteins by lipoprotein lipase, could contribute to activation of leukocyte ROS production. Although the molecular mechanisms involved are not yet elucidated, leukocytes of HLP seems to be previously primed by stimuli present in their blood, since we found increased basal ROS production by their isolated MN leukocytes.

Hypertriglyceridemia seems to synergistically enhance the risk for developing coronary artery disease (CAD) associated with hypercholesterolemia (29, 30). Recently, triglyceride-rich lipoproteins have been shown to be independent risk factors for progression of CAD once LDL-cholesterol is aggressively lowered (31). Our data showing increased ROS production by leukocytes in HLP can thus be contributive for the understanding of the underlying mechanisms for these relationships, since ROS are important initiators of the atherogenic lipoprotein oxidation process (32). In addition ROS can also directly damage the arterial wall (33).

The demand for the antioxidant defenses is conceivably increased when the amount of plasma lipids to be protected from oxidation is increased. Defective catabolism of LDL and other lipoprotein particles in certain hypercholesterolemic and hypertriglyceridemic states (34, 35) results in a prolonged half-life of these particles in blood, increasing the possibility for lipoproteins to be exposed to ROS generated by leukocytes. Decreased plasma SOD activity found in HLP may result in less lipoprotein protection against superoxide radicals generated by leukocytes and other cells in the blood. The major SOD isoenzyme in plasma is extracellular SOD (EC-SOD), which differs from the intracellular isoenzyme in many aspects (36). EC-SOD is a secretory tetrameric Cu- and Zn-containing heterogeneous glycoprotein constituted of three fractions (A, B and C) with different heparin affinity (36). EC-SOD fractions A and B primarily are found unassociated in plasma, while fraction C is bound to endothelial cell surfaces in the vasculature (37) since it has a strong affinity for heparan sulfate proteoglycan. It is therefore conceivable that the activities of intracellular SOD and EC-SOD are regulated by different factors. Moreover, the normal activity of leukocyte SOD found in HLP may be due to an increased enzyme production in response to the high leukocyte ROS generation. We found an inverse correlation between plasma SOD activity and cholesterol as well as triglyceride levels. This suggests that lipids may affect binding of EC-SOD to the endothelial cell surface or, alternatively, the release of this enzymic fraction from vasculature. In relation to GPx, we observed only a trend for decreasing plasma activity. Extracellular GPx also differs from the intracellular isoenzyme in many structural and kinetic properties (38). Since GPx removes hydrogen peroxide produced by the SOD-catalysed reaction, an imbalance between both enzymes can be harmful to cells. Indeed, in the control group the activities of both enzymes were positively correlated. In contrast, in the hyperlipidemic groups this correlation did not stand, suggesting that the functional relationship between both enzymes was disrupted.

In addition to the antioxidant enzymes, the lipophilic antioxidants carried by lipoproteins are important for scavenging ROS and inhibiting free radical-induced damage. Lipophilic antioxidants can suppress PMN respiratory burst (39), in addition to inhibiting lipoprotein

oxidation (40, 41, 42). Although α -tocopherol was not significantly different between groups, HLP patients showed low plasma levels of β -carotene and lycopene, potentially rendering plasma lipoproteins more susceptible to oxidation. In fact, increased LDL oxidation was observed in HLP indicating that the triglyceride-rich LDL is the most oxidizable lipoprotein fraction in this group. β -carotene is a quencher of singlet oxygen and a radical-trapping antioxidant (43). It has been reported that high-doses of β -carotene enhance HDL levels (44), and an inverse association of myocardial infarction with adipose-tissue β -carotene, but not with α -tocopherol was also described (45). In addition, supplementation with β -carotene has been shown to reduce not only major coronary events but all major vascular events (46), although it should be noted that administration of β -carotene has variable effects on protection of LDL from ex-vivo oxidation (49). On the other hand, the high intake of α -tocopherol has also been associated to a lower risk of coronary heart disease (47), α -tocopherol consistently protects LDL from copper-catalysed oxidation (48, 49, 50). However, there is also strong evidence that carotenoids and other lipid soluble antioxidants do play an important protective role against lipoprotein oxidation (51, 52, 53, 54). Considering that most plasma carotenoids are transported in LDL, the lower plasma carotenoid concentration observed in HLP may contribute to the decreased LDL resistance to oxidation found in this study.

HDL is negatively correlated with the incidence of atherosclerosis. The protective effect of HDL could be linked to its participation in the reverse transport of cholesterol (55) from the arterial wall. However, in some in vitro studies an antioxidant effect of HDL has been postulated (56). Our observation that HDL-cholesterol had a negative correlation with ROS release by resting MM and PMN leukocytes is unprecedented and adds suggestive evidence for the latter hypothesis. Since ROS can damage the arterial wall, the inverse correlation we found suggests a novel protective role for HDL.

In conclusion, the results suggest that oxidative stress is more likely to occur in HLP than in NL and HCH, since in HLP the release of ROS by leukocytes was greater while some components of the plasma antioxidant defense system were decreased. Our finding that leukocyte

ROS production is positively correlated with either VLDL or LDL triglycerides may be important for the understanding of the mechanisms behind leukocyte activation and oxidative stress in hyperlipidemia. Further direct testing of these findings is warranted.

ACKNOWLEDGEMENTS

This study was supported by grants of FAPESP (Fundação de Amparo à Pesquisa do Estado de São Paulo) and CNPq (Conselho Nacional de Desenvolvimento Científico e Tecnológico).

REFERENCES

- 1 . Halliwell B, Gutteridge JMC. The importance of free radicals and catalytic metal ions in human diseases. *Molec. Aspects Med.*, 8 (1985) 89.
2. Hiramatsu K, Rosen H, Heinecke JW, Wolfbauer G, Chait A. Superoxide initiates oxidation of low density lipoprotein by human monocytes. *Arteriosclerosis*, 7 (1987) 55.
3. Cathcart MK, Morel DV, Chilsolm GM. Monocytes and neutrophils oxidize low density lipoprotein making it cytotoxic. *J Leukocyte Biol.*, 38 (1985) 341.
4. Abdalla DSP, Campa A, Monteiro HP. Low density lipoprotein oxidation by stimulated neutrophils and ferritin. *Atherosclerosis*, 97 (1992) 149.
5. Lunec J. Oxygen radicals: their measurement and role in major diseases. *J. Int. Fed. Clin. Chem.*, 4 (1992) 58.
- 6 . Parthasarathy S, Rankin SM. Role of oxidized low density lipoprotein in atherogenesis. *Prog. Lipid Res.*, 31 (1992) 127.
7. Boyum A. Isolation of mononuclear cells and granulocytes from human blood: isolation of mononuclear cells by one g centrifugation and of granulocytes by combining centrifugation and sedimentation at 1 g. *Scand. J. Lab. Invest.*, 21 (1968) (Suppl.97) 77.
8. Oyanagui Y. Reevaluation of assay methods and establishment of kit for superoxide dismutase activity. *Anal. Biochem.*, 142 (1984) 290.
9. Sinet PM, Michelson, AM, Bazin A, Lejecine J, Jerome H. Increase in glutathione peroxidase activity in erythrocytes from trisomy 21 subjects. *Biochem. Biophys. Res. Commun.*, 67 (1975) 910.
10. Thurnan DI, Smith E, Flora PS. Concurrent liquid-chromatographic assay of retinol, α -tocopherol, β -carotene, α -carotene, lycopene and β -cryptoxanthin. *Clin. Chem.*, 34 (1988) 337.
11. Redgrave TG, Roberts DCK, West, CE. Separation of plasma lipoprotein by density-gradient ultracentrifugation. *Anal. Biochem.*, 65 (1975) 42.

12. Terao J, Shibata SS, Matsushita S. Selective quantification of arachidonic acid hydroperoxides and their hydroxy derivatives in reverse-phase high performance liquid chromatography. *Anal. Biochem.*, 169 (1988) 415.
13. SAS Institute Inc., SAS/STAT user's guide, version 6, 4th edn, Cary, NC, SAS Institute Inc., 1989.
14. Rosner B. *Fundamentals of Biostatistics*, 2nd edn, Boston: Duxbury Press, 1986.
15. Hiramatsu K, Arimori S. Increased superoxide production by mononuclear cells of patients with hypertriglyceridemia and diabetes. *Diabetes*, 37 (1988) 832.
16. Prónai L, Hiramatsu K, Saigusa Y, Nakazawa H. Low superoxide scavenging activity associated with enhanced superoxide generation by monocytes from male hypertriglyceridemia with and without diabetes. *Atherosclerosis*, 90 (1991) 49.
17. Ludwig PW, Hunninghake DB, Hoidal JR. Increased leukocyte oxidative metabolism in hyperlipoproteinemia. *Lancet*, 2 (1982) 348.
18. Krause S, Pohl C, Liebrecht A, Rühling K, Lösche W. Increased generation of reactive oxygen species in mononuclear blood cells from hypercholesterolemic patients. *Thrombosis Res.*, 71 (1993) 237.
19. Badwey JA, Curnutte JT, Karnovsky ML. cis-polyunsaturated fatty acids induce high levels of superoxide production by human neutrophils. *J. Biol. Chem.*, 256 (1981) 12640.
20. Ozawa M, Ohtsuka T, Okamura N, Ishibashi S. Synergism between protein kinase C activator and fatty acids in stimulating superoxide anion production in guinea pig polymorphonuclear leukocytes. *Arch. Biochem. Biophys.*, 273 (1989) 491.
21. Kakinuma K. Effects of fatty acids on the oxidative metabolism of leukocytes. *Biochim. Biophys. Acta*, 348 (1974) 76.
22. Morel F, Doussière J and Vignais PV., The superoxide-generating oxidase of phagocytic cells. Physiological, molecular and pathological aspects. *J. Biochem.*, 201 (1991) 523.
25. Henderson LM, Moule SK, Chappell JB. The immediate activator of the NADPH oxidase is arachidonate not phosphorylation. *Eur. J. Biochem.*, 211 (1993) 157.

24. Görög P. Activation of human blood monocytes by oxidized polyunsaturated fatty acids: a possible mechanism for the generation of lipid peroxides in the circulation. *Int. J. Exp. Pathol.*, 72 (1991) 227.
25. Bellinati-Pires R, Waitzberg DL, Salgado MM, Carneiro-Sampaio MMS. Functional alterations of human neutrophils by medium chain triglyceride emulsions: evaluation of phagocytosis, bacterial killing, and oxidative activity. *J. Leukocyte Biol.*, 53 (1993) 404.
26. Biemond, P.E.; Van Eijik, H.G.; Swaak, A.J.G.; and Koster, J.F., Iron mobilization from ferritin by superoxide derived from stimulated polymorphonuclear leukocytes: possible mechanism in inflammatory diseases, *J. Clin. Invest.*, 73 (1984) 1576.
27. Salonen JT, Nyyssönen K, Korpela H, Tuomilehto J, Seppänen R, Salonen R. High stored iron levels are associated with excess risk of myocardial infarction in eastern Finnish men. *Circulation*, 86 (1992) 803.
28. Kuijk, FJGM, Sevanian A, Handelman GJ and Dratz EA., A new role for phospholipase A₂: protection of membranes from lipid peroxidation damage. *TIBS*, 12 (1987) 31.
29. Assmann G, Schulte H. Role of triglycerides in coronary artery disease: lessons from the prospective cardiovascular Munster study. *Am. J. Cardiol.*, 70 (1992) 10H.
30. Gotto AM. Hypertriglyceridemia: risks and perspectives. *Am. J. Cardiol.*, 70 (1992) 19H.
31. Hodis HN, Mack WJ, Azen SP, Alaupovic P, Pogoda JM, LaBree L, Hemphill LC, Kramsch, Blankenhorn DH. *Circulation*, 90 (1994) 42.
32. Cathcart, M.K.; Morel, D.W. and Chisolm, G.M., Monocytes and neutrophils oxidize low density lipoprotein making it cytotoxic. *J. Leukocyte Biol.*, 38 (1985) 341.
33. Korpela H. Increased myocardial and hepatic iron concentration in pigs with microangiopathy (Mulberry heart disease) as a risk factor of oxidative damage. *Ann. Nutr. Metab.*, 34 (1990) 193.
34. Goldstein JL, Brown MS. The low density lipoprotein pathway and its relation to atherosclerosis. *Annu. Rev. Biochem.*, 46 (1977) 897.
35. Hiramatsu K, Bierman EL, Chait A. Metabolism of low density lipoprotein from patients with hypertriglyceridemia and diabetes. *Diabetes*, 34 (1985) 8.

36. Marklund SL, Holme E, Hellner L. Superoxide dismutase in extracellular fluids. *Clin. Chim. Acta*, 126 (1982) 41.
37. Karlsson K, Marklund SL. Extracellular superoxide dismutase by human cell lines. *Biochem. J.*, 255 (1988) 223.
38. Avissar N, Whitin JC, Allen PZ, Palmer IS, Cohen HJ. Antihuman plasma glutathione peroxidase antibodies: immunologic Investigations to determine plasma glutathione peroxidase protein and selenium content in plasma. *Blood*, 73 (1989) 318.
39. Baader WJ, Hatzelmann A, Ullrich V. The suppression of granulocyte functions by lipophilic antioxidants. *Biochem. Pharmacol.*, 37 (1988) 1089.
40. Suarna C, Hood RL, Dean RT, Stocker R. Comparative antioxidant activity of tocotrienols and other natural lipid-soluble antioxidants in a homogenous system, and in rat and human lipoproteins. *Biochim. Biophys. Acta*, 1166 (1993) 163.
41. Reaven PD and Witztum JL. Comparison of supplementation of RRR- α -tocopherol and racemic α -tocopherol in humans. *Arterioscler. Thromb.*, 13 (1993) 601.
42. Jialal I, Norkus EP, Cristol L, Grundy SM. Beta-carotene inhibits the oxidative modification of low-density lipoprotein. *Biochim. Biophys. Acta*, 1086 (1991) 134.
43. Krinsky NI. Mechanism of action of biological antioxidants. *Proc. Soc. Exp. Biol. Med.*, 200 (1992) 248.
44. Ringer TV, DeLoof MJ, Winterrowd GE, et al. Beta-carotene's effects on serum lipoproteins and immunologic indices in humans. *Am. J. Clin. Nutr.*, 53 (1991) 688.
45. Kardinaal AFM, Kok FJ, Ringstad J, Gomez-Aracena J, Mazaev VP, Kohlmeier L, Martin BC, Aro, A., Kark JD, Delgado-Rodriguez M, Riemersa RA, van't Veer, P, Huttunen JK, Martin-Moreno JM. Antioxidants in adipose tissue and risk of myocardial infarction: the EURAMIC study. *Lancet*, 342 (1993) 1379.
46. Gaziano JM, Mason JAE, Ridker PM, Buring JE, Hennekens CH. Beta carotene therapy for chronic stable angina. *Circulation*, 82 (suppl. 3) (1990) III-201.

47. Rimm EB, Stampfer MJ, Ascherio A, Giovannucci E, Colditz GA, Willett WC. Vitamin E consumption and the risk of coronary heart disease in men. *N. Engl. J. Med.*, 328 (1993) 1450.
48. Jialal I, Grundy S, The effect of dietary supplementation with alpha-tocopherol on the oxidative modification of low density lipoprotein. *J. Lipid Res.*, 33 (1992) 899.
49. Princen HMG, Van Poppel G, Vogelezang C, Bluytenhak R, Kok FJ. Supplementation of vitamin E but not beta-carotene in vivo protects LDL from lipid peroxidation. *Arterioscler. Thromb.*, 12 (1992) 554.
50. Jialal I, Grundy SM. Effect of combined supplementation with α -tocopherol, ascorbate, and beta carotene on low-density lipoprotein oxidation. *Circulation*, 88 (1993) 2780
51. Dieber-Rotheneder M, Puhl H, Waeg G, Striegl G, Esterbauer H. Effect of oral supplementation with D- α -tocopherol on the vitamin E content of human low density lipoproteins and resistance to oxidation. *J. Lipid Res.*, 32 (1991) 1325.
52. Jessup W, Rankin SM, De Whalley CV, Hoult JRS, Scott J, Leake DS. α -Tocopherol consumption during low density lipoprotein. *Biochem. J.*, 265 (1990) 399.
53. Jialal I, Grundy SM. Effect of dietary supplementation with α -tocopherol on the oxidative modification of low density lipoprotein. *J. Lipid Res.*, 33 (1992) 899.
54. Babi AV, Gebicki JM, Sullivan DR. Vitamin E content and low density lipoprotein oxidizability induced by free radicals. *Atherosclerosis*, 81 (1990) 175.
55. Eisenberg S. High density lipoprotein metabolism. *J. Lipid Res.*, 26 (1984) 1017.
56. Parthasarathy S, Barnett J, Fong LG. High-density lipoprotein inhibits the oxidative modification of low density lipoprotein. *Biochim. Biophys. Acta*, 1044 (1990) 275.

TABLE 1

DATA OF HYPERLIPIDEMIC PATIENTS AND CONTROLS

mg/dL	NL (<i>n</i> = 19)	HCH (<i>n</i> = 18)	HLP (<i>n</i> = 14)
Total cholesterol	187 ± 8	295 ± 14*	295 ± 16*
VLDL-cholesterol	28 ± 1	68 ± 3*	101 ± 5*
LDL-cholesterol	87 ± 3	159 ± 5*	166 ± 6*
HDL-cholesterol	72 ± 2	69 ± 1	28 ± 1*
Total triglycerides	112 ± 12	154 ± 7	492 ± 65*
VLDL-triglycerides	84 ± 6	94 ± 6	287 ± 16*
LDL-triglycerides	20 ± 1	53 ± 3*	195 ± 12*
HDL-triglycerides	8 ± 1	7 ± 1	15 ± 2*
Total phospholipids	157 ± 8	226 ± 8*	224 ± 14*

Results are mean ± S.E.M. *Values between groups significantly different from controls (NL), (*p* < 0.05).
HCH : patients with hypercholesterolemia ; HLP: patients with combined hyperlipidemia.

TABLE 2

LEUKOCYTE AND PLASMA ENZYMATIC ACTIVITIES OF HYPERLIPIDEMIC PATIENTS AND CONTROLS

Leukocytes	NL (n = 19)	HCH (n = 18)	HLP (n = 14)
Mononuclear			
SOD	5.71±1.41	5.41±1.19	1.67±0.63
GPx	4.26±0.91	4.01±0.85	7.70±1.56
Polymorphonuclear			
SOD	3.08±0.76	1.47±0.33	2.14±0.33
GPx	3.72±1.07	0.53±0.12	3.08±1.02
Plasma			
SOD	0.12±0.01	0.10±0.02	0.06±0.01*
GPx	8.85±1.48	5.96±0.91	5.63±1.35

Results are means±SEM of enzymatic activities (U/mg of protein). NL, controls; HCH, patients with hypercholesterolemia; HLP, patients with combined hyperlipidemia. * p<0.05 compared to controls.

TABLE 3
PLASMA LIPID SOLUBLE ANTIOXIDANTS AND LIPOPROTEIN LIPID HYDROPEROXIDES

	NL (n = 19)	HCH (n = 18)	HLP (n = 14)
α -Tocopherol	4.66 $\pm$ 1.03	3.81 $\pm$ 0.66	3.14 $\pm$ 0.29
β -Carotene	3.96 $\pm$ 1.16	1.83 $\pm$ 0.33	1.04 $\pm$ 0.31*
Lycopene	0.69 $\pm$ 0.22	0.22 $\pm$ 0.05	0.16 $\pm$ 0.03*
Lipid hydroperoxides			
VLDL	0.47 $\pm$ 0.02	0.45 $\pm$ 0.01	0.47 $\pm$ 0.03
LDL	0.67 $\pm$ 0.02	0.66 $\pm$ 0.02	0.86 $\pm$ 0.03*, **

Results are means $\pm$ SEM and are expressed as μ mol/mg of cholesterol. NL, controls; HCH, patients with hypercholesterolemia; HLP, patients with combined hyperlipidemia. * $p < 0.05$ compared to controls; ** $p < 0.05$ compared to HCH. Lipid hydroperoxides were determined after lipoprotein oxidation with copper as described in Materials and Methods.

FIGURE LEGENDS

Fig. 1: Release of reactive oxygen species (ROS) by resting and PMA-stimulated mononuclear cells. Cells were isolated from heparinized blood and chemiluminescence of incubation systems containing 1×10^4 cells/ml, 5 μ M luminol and 50 ng/ml PMA (only for stimulated cells), in PBS pH 7.4 was followed for 1 h in a liquid scintillation counter. Results are expressed as mean $\pm$ S.E.M. of integrated chemiluminescence. NL, control group; HCH, hypercholesterolemic group; HLP, combined hyperlipidemia group. * $p < 0.05$ as compared to NL and HCH; $\star$ $p < 0.05$ as compared to HCH.

Fig. 2 : Release of reactive oxygen species (ROS) by resting and PMA-stimulated polymorphonuclear cells. Cells were isolated from heparinized blood and the chemiluminescence of incubation systems containing 1×10^4 cells/ml, 5 μ M luminol and 50 ng/ml PMA (only for stimulated cells), in PBS pH 7.4 was followed for 1 h in a liquid scintillation counter. Results are expressed as mean $\pm$ S.E.M. of integrated chemiluminescence. * $p < 0.05$ as compared to NL and HCH.

Fig. 3 : Correlation between the luminol-enhanced chemiluminescence of resting mononuclear leukocytes and the concentration of total plasma triglycerides .

Fig. 4: Correlation between the luminol-enhanced chemiluminescence of resting polymorphonuclear leukocytes and the concentration of total plasma triglycerides.

Fig. 5: Correlation between HDL-cholesterol and the luminol-enhanced chemiluminescence of resting mononuclear leukocytes.

Fig. 6: Correlation between HDL-cholesterol and the luminol-enhanced chemiluminescence of resting polymorphonuclear leukocytes.

FIGURE 1

FIGURE 2

FIGURE 3

Fig. 3

FIGURE 4

FIGURE 5

FIGURE 6