

8. REFERÊNCIAS BIBLIOGRÁFICAS

- AB'SABER, A. (2006) **Brasil: Paisagens de Exceção – o litoral e o pantanal mato-grossense patrimônios básicos**. Cotia: Ateliê Editorial. 182p.
- AHAGON, N., Tanaka, Y. e Ujié, H. (1993) *Florisphaera profunda*, a possible nannoplankton indicator of late Quaternary changes in sea-water turbidity at the northwestern margin of the Pacific. *Marine Micropaleontology*, **22**, 255-273.
- AIZAWA, C., Oba, T., Okada, H. (2004) Late Quaternary paleoceanography deduced from coccolith assemblages in a piston core recovered off the central Japan coast. *Marine Micropaleontology*, **52**, 277-297.
- ALLABY, A., e Allaby, M. (1999) **Dictionary of Earth Sciences**. 2 ed. Oxford: Oxford University Press. 619p.
- ANDRULEIT, H., Rogalla, U., Stäger, S. (2004) From living communities to fossil assemblages: origin and fate of coccolithophores in the northern Arabian Sea. *Micropaleontology*, **50** (1), 5-21.
- ANTUNES, R. L. (1997) **Introdução ao estudo dos nanofósseis calcários**. Rio de Janeiro: UFRJ. 115p.
- ARZ, H.W., Gerhardt, S., Pätzold, J. e Röhl, U. (2001) Millennial-scale changes of surface- and deep-water flow in the western tropical Atlantic linked to Northern Hemisphere high-latitude climate during the Holocene. *Geology*, **29**, 239-242.
- ARZ, H.W., Pätzold, J. e Wefer, G. (1999a) The deglacial history of the western tropical Atlantic as inferred from high resolution stable isotope records off northeastern Brazil. *Earth and Science Letters*, **167**, 105-117.

- ARZ, H.W., Pätzold, J. e Wefer, G. (1999b) Climatic changes during the last deglaciation recorded in sediment cores from the northeastern Brazilian Continental Margin. *Geo-Marine Letters*, **19**, 209-218.
- ARZ, H.W., Pätzold, J. e Wefer, G. (1998) Correlated Millennial-Scale Changes in surface hydrography e terrigenous sediment yield inferred from last-glacial marine deposits off northeastern Brazil. *Quaternary Research*, **50**, 157-166.
- BARETTA-BEKKER, J.G., Duursma, E.K. & Kuipers, B.R. (Eds.) (1998). **Encyclopedia of Marine Sciences**. 2. ed. Berlin: Springer-Verlag. 357p.
- BAUMANN, K.-H., Andruleit, H., Böeckel, B., Geisen, M., Kinkel, H. (2005) The significance of extant coccolithophores as indicators of ocean water masses, surface water temperature, and paleoproductivity: a review. *Paläontologische Zeitschrift*, **79** (1), 93-112.
- BAUMANN, K.-H., Böeckel, B., Donner, B., Gerhardt, S., Heinrich, R., Vink, A., Volbers, A., Willems, H. e Zonneveld, K.A.F. (2004) Contribution of Calcareous Plankton Groups to the Carbonate Budget of South Atlantic Surface Sediments. In: WEFER, G. MULITZA, S. e RATMEYER, V. (Ed.). **The South Atlantic in the Late Quaternary**. Berlin: Springer-Verlag. p. 81-99.
- BAUMANN, K.-H., Cepek, M. e Kinkel, H. (1999) Coccolithophores as indicator of ocean water masses, surface-water temperature, e paleoproductivity from the South Atlantic. In: FISCHER, G., WEFER, G. (Ed.). **Use of proxies in Paleoceanography: Examples from the South Atlantic**. Berlin: Springer-Verlag. p. 117-144.
- BEAUFORT, L., Heussner, S. (2001) Seasonal dynamics of calcareous nannoplankton on a West European continental margin: the Bay of Biscay. *Marine Micropaleontology*, **43**, 27-55.
- BEAUFORT, L., Lancelot, Y., Camberlin, P., Cayre, O., Vincent, E., Bassinot, F. e Labeyrie, L. (1997). Insolation cycles as a major control of equatorial Indian Ocean primary production. *Science*, **278**, 1451-1454.

- BEHLING, H., Arz, H.W., Pätzold, J. e Wefer, G. (2002) Late Quaternary vegetational and climate dynamics in southeastern Brazil, inferences from marine cores GeoB 3229-2 and GeoB 3202-1. *Palaeogeography, palaeoclimatology, Palaeoecology*, **179**, 227-243.
- BEHLING, H., Arz, H.W., Pätzold, J. e Wefer, G. (2000) Late Quaternary vegetational and climate dynamics in northeastern Brazil, inferences from marine core GeoB 3104-1. *Quaternary Science Reviews*, **19**, 981-994.
- BERGER, W.H. e Wefer, G. (1996) Central Themes of South Atlantic Circulation. In: WEFER, G., BERGER, W. H., SIEDLER, G. e WEBB, D. J. (Ed.). **The South Atlantic: present and past circulation**. Berlin: Springer-Verlag. p. 1-11.
- BOECKEL, B., Baumann, K-H., Henrich, R., Kinkel, H. (2006) Coccolith distribution patterns in South Atlantic and Southern Ocean surface sediments in relation to environmental gradients. *Deep-Sea Research I*, **53**, 1073-1099.
- BOECKEL, B. e Baumann, K.-H. (2004) Distribution of coccoliths in surface-sediments of the south-eastern South Atlantic Ocean: ecology, preservation and carbonate contribution. *Marine Micropaleontology*, **51**, 301-320.
- BOLLMANN, J. (1997) Morphology and biogeography of *Gephyrocapsa* coccolith in Holocene sediments. *Marine Micropaleontology*, **29**, 319-350.
- BOWN, P.R. e Young, J.R. (1998a) Introduction. In: BOWN, P.R. (Ed.). **Calcareous Nannofossil Biostratigraphy**. Kluwer Academic Publishers. p. 1-15.
- BOWN, P.R. e Young, J.R. (1998b) Techniques. In: BOWN, P.R. (Ed.). **Calcareous Nannofossil Biostratigraphy**. Kluwer Academic Publishers. p. 16-28.
- BOWN, P.R., Young, J.R. (1997) Mesozoic calcareous nannoplankton classification. *Journal of Nannoplankton Research*, **19**, 21-36.
- BRADLEY, R.S. (1999) Marine sediments and corals. In: BRADLEY, R.S. **Paleoclimatology – reconstructing climates of the Quaternary**. Burlington: Elsevier Academic Press. p.191-283.

- BRAND, L.E. (1994) Physiological ecology of marine coccolithophores. In: WINTER, A. & SIESSER, W. (Ed.). **Coccolithophores**. Cambridge: Cambridge University Press, p. 39-49.
- BROCCOLI, J. (2000) Tropical Cooling at the Last Glacial Maximum: an atmosphere-mixed layer model simulation. *Journal of Climate*, **13**, 951-976.
- BROECKER, W.S. (1998) The end of the present interglacial: how and when?. *Quaternary Science Reviews*, **17**, 689-694.
- BROECKER, W.S. (1986) Oxygen Isotope Constraints on Surface Ocean Temperatures. *Quaternary Research*, **26**(1), 121-134.
- BROECKER, W.S. (1982) Ocean chemistry during glacial time. *Geochimica et Cosmochimica Acta*, **46**, 1689-1705.
- BROECKER, W.S. e Takahashi, T. (1978) The relationship between lysocline depth and in situ carbonate ion concentration. *Deep Sea Research Part I Oceanographic Research Papers*, **25**, 65-95.
- BROERSE, A.T.C., Ziveri, P., van Hinte, J.E., Honjo, S. (2000a) Coccolithophore export production, species composition, and coccolith-CaCO₃ fluxes in the NE Atlantic (34°N 21°W and 48°N 21°W). *Deep-Sea Research II*, **47**, 1877-1905.
- BROERSE, A.T.C., Brummer, G.-J.A., van Hinte, J.E. (2000b) Coccolithophore export production in response to monsoonal upwelling off Somalia (northwestern Indian Ocean). *Deep-Sea Research II*, **47**, 2179-2205.
- BROWN, C.E. (1998) **Applied Multivariate Statistics in Geohydrology and Related Sciences**. Berlin: Springer-Verlag. 248p.
- BROWN, J. e Wright, J. (2001) **Ocean Chemistry and Deep-Sea Sediments**. Oxford: Butterworth-Heinemann. 134 p.
- BRUMMER, G.J.A. e Van Eijden, A.J.M. (1992) "Blue-ocean" paleoproductivity estimates from pelagic carbonate mass accumulation rates. *Marine Micropaleontology*, **19**, 99-117.

- BUCCHERI, G., Capretto, G., Di Donato, V., Esposito, P., Ferruzza, G., Pescatore, T., Ermolli, E. R., Senatore, M. R., Sprovieri, M., Bertoldo, M., Carella, D. e Madonia, G. (2002) A high resolution record of the last deglaciation in the southern Tyrrhenian Sea: environmental and climatic evolution. *Marine Geology*, **186**, 447-470.
- BUCCIANTI, A. e Esposito, P. (2004) Insights into Late Quaternary calcareous nannoplankton assemblages under the theory of statistical analysis for compositional data. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **202**, 209-227.
- BURNETT, J.A., Young, J.R. e Bown, P.R. (2000) Calcareous nannoplankton and global climate change. In: CULVER, S.J., RAWSON, P.F. (Eds.), **Biotic Response to Global Change: The Last 145 Million Years**. Cambridge: Cambridge University Press, pp.35-50.
- BUTZIN, M., Prange, M. e Lohmann, G. (2005) Radiocarbon simulations for the glacial ocean: the effects of wind stress, Southern Ocean sea ice and Heinrich events. *Earth and Planetary Science Letters*, **235**, 45-61.
- CANTOLLA, A.U. (2003) **Historia Del Clima de la Tierra**. Servicio Central de Publicaciones Del Gobierno Vasco. 306p.
- CAO, L., Fairbanks, R.G., Butzin, M. e Naik, N. (2007) The marine radiocarbon reservoir age. *Radiocarbon*, in prep.
- CHANG, P. (1998) Unusual tropical Atlantic Ocean temperatures cause climate swings in Brazil and West Africa. *Quarterdeck*, **6**, (The Atlantic Ocean).
- CHANG, P., Ji, L. e Li, H. (1997) A decadal climatic variation in the Atlantic Ocean from thermodynamic air-sea interactions. *Nature*, **385**, 516-518.
- CHARLSON, R.J., Lovelock, J.E., Andreae, M.O., Warren, S.G. (1987) Oceanic phytoplankton, atmospheric sulphur, cloud albedo and climate. *Science*, **326**, 655-661.
- CLIMAP. (1981) Seasonal Reconstructions of the Earth's Surface at the Last Glacial Maximum. *Geological Society of America Map and Chart Series*, **MC-36**, 1-18.
- COLLING, A. (2001) **Ocean Circulation**. Oxford: Butterworth-Heinemann. 286 p.

- COSTA, K. B. (2000). **Variações Paleoceanográficas na Porção Oeste do Atlântico Sul entre o Último Máximo Glacial e o Holoceno: Isótopos Estáveis do Oxigênio e Carbono e Razão Cd/Ca em Foraminíferos Bentônicos**. In: Instituto de Geociências, Universidade Federal do Rio Grande do Sul, Porto Alegre. Tese de Doutorado, pp. 250.
- CURRY, W.B. (1996) Late Quaternary circulation in the western equatorial Atlantic. In: WEFER, G., BERGER, W. H., SIEDLER, G. e WEBB, D. J. (Ed.). **The South Atlantic: present and past circulation**. Berlin: Springer-Verlag. p. 577-598.
- CURRY, W.B., Duplessy, J.C., Labeyrie, L.D. e Shackleton, N.J. (1988) Changes in the distribution of $\delta^{13}\text{C}$ of deep water ΣCO_2 between the last glaciation and the Holocene. *Paleoceanography*, **3**(3), 317-341.
- DITTERT, N., Baumann, K.-H., Bickert, T., Heinrich, R., Huber, R., Kinkel, H. e Meggers, H. (1999) Carbonate Dissolution in the Deep-Sea: Methods, Quantification and Paleoceanographic Application In: FISCHER, G., WEFER, G. (Ed.). **Use of proxies in Paleoceanography: Examples from the South Atlantic**. Berlin: Springer-Verlag. p. 255-284.
- DUNCOMBE RAE, C.M. (1991) Agulhas retroflection rings in the South Atlantic ocean: an overview. *South African Journal of Marine Science*, **11**, 327-344.
- DUPLESSY, J.C., Labeyrie, L. e Waelbroeck, C. (2002) Constraints on the ocean oxygen isotopic enrichment between the Last Glacial Maximum and the Holocene: Paleoceanographic implications. *Quaternary Science Reviews*, **21**, 315-330.
- DUPLESSY, J.C., Shackleton, N.J., Fairbanks, R.G., Labeyrie, L., Oppo, D. e Kallel, N. (1988) Deepwater source variations during the last climatic cycle and their impact on global deepwater circulation. *Paleoceanography*, **3**(3), 343-360.
- EMILIANI, C. (1955) Pleistocene Temperatures. *Journal of Geology*, **63**, 538-578.
- ERBA, E. e Tiraboschi, D. Nannofossils as paleothermometers of the mid-Cretaceous Greenhouse. In: BIENNIAL INTERNATIONAL NANNOPLANKTON ASSOCIATION

- CONFERENCE, 11, 2006, Lincoln. **Abstracts**. Lincoln: University of Nebraska, 2006. p.41-42.
- EVANS, D.L. e Signorini, S.S. (1985) Vertical structure of the Brazil Current. *Nature*, **315**, 48-50.
- FAIRBANKS, R.G. (1989) Glacio-eustatic sea level record 0-17,000 years before present: influence of glacial melting rates on Younger Dryas event and deep ocean circulation. *Nature*, **342**(6250), 637-642.
- FAIRBANKS, R.G., Mortlock, R.A., Chiu, T.-C., Cão, L., Kaplan, A., Guilderson, T.P., Fairbanks, T.W., Bloom, A.L., Grootes, P.M. e Nadeau, M.-J. (2005) Radiocarbon calibration curve spanning 0 to 50,000 years BP based on paired $^{230}\text{Th}/^{234}\text{U}/^{238}\text{U}$ and ^{14}C dates on pristine corals. *Quaternary Science Reviews*, **24** (16-17), 1781-1796.
- FARMER, E.C., DeMenocal, P.B. e Marchitto, T.M. (2005) Holocene and deglacial ocean temperature variability in the Benguela upwelling region: Implications for low-latitude atmospheric circulation. *Paleoceanography*, **20**, PA2018.
- FLORES, J.-A., Gersonde, R., Sierro, F. J. (1999) Pleistocene fluctuations in the Agulhas Current Retroflection based on the calcareous plankton record. *Marine Micropaleontology*, **37**, 1-22.
- FRENZ, M. e HENRICH, R. (2007) Carbonate dissolution revealed by silt grain-size distribution: comparison of Holocene and Last Glacial Maximum sediments from the pelagic South Atlantic. *Sedimentology*, **54** (2), 391–404.
- GATES, W.L. (1976) Modeling the Ice-Age Climate. *Science*, **191**, 1138-1144.
- GEOSECS, Geochemical Ocean Sections Study. Disponível em: <<http://ingrid.ldeo.columbia.edu/SOURCES/.GEOSECS>> Acesso em: 19 novembro 2006.
- GERSONDE, R., Abelmann, A., Cortese, G., Becquey, S., Bianchi, C., Brathauer, U., Niebler, H.-S., Zielinski, U. e Pätzold, J. (2004) The Late Pleistocene South Atlantic and Southern Ocean Surface – A Summary of Time-Series Studies. In: WEFER, G. MULITZA, S. e

- RATMEYER, V. (Ed.). **The South Atlantic in the Late Quaternary**. Berlin: Springer-Verlag. p. 499-529.
- GILDOR, H., Follows, M.J. (2002) Two-way interactions between ocean biota and climate mediated by biogeochemical cycles. *Israel Journal of Chemistry*, **42**, 15-27.
- GIRAUDEAU, J. (1992) Coccolith paleotemperature and paleosalinity estimates in the Caribbean Sea for the middle-late Pleistocene (DSDP Leg 68 – Hole 502B). *Memorie Di Scienze Geologiche*, **43**, 375-387.
- GONÇALVES, H.M.S., Dehnhardt, B.A. (1999) Interpretação do testemunho T-101, (32°05'S; 47°35'W) com base em nanofósseis calcários – Cone do Rio Grande, Rio Grande do Sul, Brasil. *Revista Brasileira de Ciências*, **29** (4), 527-532.
- GORDON, A.L. (1996) Comment on the South Atlantic's Role in the Global Circulation. In: WEFER, G., BERGER, W. H., SIEDLER, G. e WEBB, D. J. (Ed.). **The South Atlantic: present and past circulation**. Berlin: Springer-Verlag. p. 121-124.
- GRDC, Global Runoff Data Center, Alemanha. Disponível em: <<http://grdc.bafg.de>> e <<http://www.grdc.sr.unh.edu>> Acesso em: 20 outubro 2006.
- GREEN, J.C., Jordan, R.W. (1994) Systematic history and taxonomy. In: GREEN, J.C., LEADBEATER, B.S.C. (Ed.). **The Haptophyte Algae**. The Systematics Association Special Volume No. 51, Oxford: Clarendon Press, p. 1-21.
- HAGINO, K., Okada, H., Matsuoka, H. (2005) Coccolithophore assemblages and morphotypes of *Emiliana huxleyi* in the boundary zone between the cold Oyashio and warm Kuroshio currents off the coast of Japan. *Marine Micropaleontology*, **55**, 19-47.
- HAGINO, K., Okada, H. (2004) Floral response of coccolithophores to progressive oligotrophication in the South Equatorial Current, Pacific Ocean. In: *Global Environmental Change in the Ocean and on Land* (Ed. M. Shiyomi *et al.*), pp. 121-132.
- HAGINO, K., Okada, H., Matsuoka, H. (2000) Spatial dynamics of coccolithophore assemblages in the Equatorial Western-Central Pacific Ocean. *Marine Micropaleontology*, **39**, 53-72.

- HALE, W. e Pflaumann, U. (1999) Sea-Surface temperature Estimations using a Modern Analog Technique with Foraminiferal Assemblages from Western Atlantic Quaternary Sediments. In: FISCHER, G., WEFER, G. (Ed.). **Use of proxies in Paleoceanography: Examples from the South Atlantic**. Berlin: Springer-Verlag. p. 69-90.
- HARRIS, S. E. e Mix, A.C. (1999) Pleistocene precipitation balance in the Amazon Basin recorded in deep sea sediments. *Quaternary Research*, **51**, 14-26.
- HASTENRATH, S. (1990) Prediction of Northeast Brazil rainfall Anomalies. *Journal of Climate*, **3**, 893-904.
- HAY, W.W. (1977) Calcareous Nannofossils. In: RAMSAY, A. T. S. (Ed.). **Oceanic Micropaleontology**. London: Academic Press. p. 1055-1200.
- HENRIKSSON, A. S. (2000) Coccolithophore response to oceanographic changes in the equatorial Atlantic during the last 200,000 years. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **156**, 161-173.
- HERRLE, J.O. (2003) Reconstructing nutricline dynamics of mid-Cretaceous oceans: evidence from calcareous nannofossils from the Niveau Paquier black shale (SE France). *Marine Micropaleontology*, **47**, 307-321.
- HERRLE, J.O., Pross, J., Friedrich, O., Köbler, P. e Hemleben, C. (2003) Forcing mechanisms for mid-Cretaceous black shale formation: evidence from the Upper Aptian and Lower Albian of the Vocontian Basin (SE France). *Palaeogeography, Palaeoclimatology, Palaeoecology*, **190**, 399-426.
- HOLMES, M.E., Eichner, C., Struck, U. e Wefer, G. (1999) Reconstruction of Surface Ocean Nitrate Utilization Using Stable Nitrogen Isotopes in Sinking Particles and Sediments. In: FISCHER, G., WEFER, G. (Ed.). **Use of proxies in Paleoceanography: Examples from the South Atlantic**. Berlin: Springer-Verlag. p. 447-468.
- HOLMES, M.E., Lavik, G., Fischer, G. e Wefer, G. (2004) Nitrogen Isotopes in Sinking Particles and Surface Sediments in the Central and Southern Atlantic. In: WEFER, G. MULITZA, S.

e RATMEYER, V. (Ed.). **The South Atlantic in the Late Quaternary**. Berlin: Springer-Verlag. p. 143-165.

IBGE, Instituto Brasileiro de Geografia e Estatística - Boletins. Disponível em: <<http://www.ibge.gov.br>> Acesso em: 30 agosto 2006.

IMBRIE, J., Boyle, E.A., Clemens, S.C., Duffy, A., Howard, W.R., Kukla, G., Kutzbach, J., Martinson, D.G., McIntyre, A., Mix, A.C., Molfino, B., Morley, J.J., Peterson, L.C., Pisias, N.G., Prell, W.L., Raymo, M.E., Shackleton, N.J. e Toggweiler, J.R. (1992) On the structure and origin of major glaciation cycles.1.Linear responses to Milankovitch Forcing. *Paleoceanography*, **7**(6).

IMBRIE, J., Hays, J.D., Matinson, D.G., McIntyre, A., Mix, A.C., Morley, J.J., Pisias, N.G., Prell, W.L. e Shackleton, N.J. (1984) The orbital theory of Pleistocene climate: support from a revised chronology of the marine d18O record. In: *Milankovitch and Climate, Part 1* (Ed. by A. L. B. et al.), pp. 269-305. D. Reidel Publishing Co.

IMBRIE, J., Mix, A.C. e Matinson, D.G. (1993) Milankovitch theory viewed from Devil's Hole. *Nature*, **363**, 531-533.

JENNERJAHN, T.C., Ittekkot, V., Arz, H.W., Behling, H., Pätzold, J. e Wefer, G. (2004) Asynchronous terrestrial and marine signals of climate change during Heinrich events. *Science*, **306**, 2236-2239.

KARWATH, B., Janofske, D., Tietjen, F., Willems, H. (2000) Temperature effects on growth and cell size in the marine calcareous dinoflagellate *Thoracosphaera heimii*. *Marine Micropaleontology*, **39**, 43-51.

KELLY, D.C., Norris, R.D. e Zachos, J.C. (2003) Deciphering the paleoceanographic significance of Early Oligocene *Braarudosphaera* chalks in the South Atlantic. *Marine Micropaleontology*, **49**, 49-63.

- KIM, J.-H. e Schneider, R. R. (2003) Low-latitude control of interhemispheric sea-surface temperature contrast in the tropical Atlantic over the past 21 kyears: the possible role of SE trade winds. *Climate Dynamics*, **21**, 337-347.
- KINKEL, H., Baumann, K.-H., Cepek, M. (2000) Coccolithophores in the equatorial Atlantic Ocean: response to seasonal and Late Quaternary surface water variability. *Marine Micropaleontology*, **39**, 87-112.
- KUCERA, M. (2003) Numerical approach to microfossil proxy data. Lecture notes for Summer School *Paleoceanography: theory and field evidence*. IAMC Geomare 2003, pp.66-90.
- KUCERA, M. e Malmgren, B.A. (1998) Logratio transformation of compositional data – a resolution of the constant sum constraint. *Marine Micropaleontology*, **34**, 117-120.
- LABEYRIE, L. D., Duplessy, J. C. e Blanc, P. L. (1987) Variations in mode of formation and temperature of oceanic seep waters over the past 125.000 years. *Nature*, **327**, 477-482.
- LALLI, C.M. e Parsons, T.R. (1993) **Biological Oceanography: an introduction**. Oxford: Pergamon Press. 301p.
- LEDRU, M. P., Bertaux, J., Sifeddine, A. e Suguio, K. (1998) Absence of Last Glacial Maximum records in lowland tropical forests. *Quaternary Research*, **49**, 233-237.
- LITTLE, M.G., Schneider, R., Kroon, D., Price, B., Summerhayes, C. e Segl, M. (1997) Trade wind forcing of upwelling, seasonality, e Heinrich events as a response to sub-Milankovitch climate variability. *Paleoceanography*, **12**, 568-576.
- LOVELOCK, J.E. (1987) **Gaia, um novo olhar sobre a vida na Terra**. Lisboa: Edições 70. 169p.
- LOVELOCK, J.E. (1991) **As eras de Gaia – a biografia da nossa Terra viva**. Rio de Janeiro: Editora Campus. 236p.
- LOVELOCK, J.E. (2006) **The revenge of Gaia**. London: Penguin Books Allen Lane. 177p.
- MARGULIS, L., Sagan, D. (2002) **O que é vida?** Rio de Janeiro: Jorge Zahar Editor. 289p.
- MARTIN, L., Bertaux, J., Corrège, T., Ledru, M.P., Mourguiart, P., Sifeddine, A., Soubiès, F., Wirmann, D., Suguio, K. e Turcq, B. (1997) Astronomical Forcing of Contrasting rainfall

changes in tropical south America between 12.400 e 8.800 cal yr B. P. *Quaternary Research*, **47**, 117-122.

MASLIN, M.A., Shackleton, N.J. e Pflaumann, U. (1995) Surface water temperature, salinity and density changes in the northeast Atlantic during the last 45.000 years: Heinrich events, deep water formation e climatic rebounds. *Paleoceanography*, **10**, 527-544.

MATSUOKA, H. e Okada, H. (1989) Quantitative analysis of Quaternary nanoplankton in the subtropical north western pacific ocean. *Marine Micropaleontology*, **14**, 97-118.

MAURRASSE, F.J-M.R., Lamolda, M.A., Aguado, R., Peryt, D., Sen, G. (2005) Spatial and temporal variations of the Haitian K/T boundary record: implications concerning the event or events. *Journal of Iberian Geology*, **31** (1), 113-133.

MIRANDA, L.B. e Castro Filho, B.M. (1982) Geostrophic flow conditions of the Brazil Current at 19°S. *Ciencia Interamericana*, **22**, 44-48.

MIX, A.C., Bard, E. and Schneider, R. (2001). Environmental processes of the ice age: land, oceans, glaciers (EPILOG). *Quaternary Science Reviews*, **20**, 627-657.

MOLFINO, B. e McIntyre, A. (1990) Precessional Forcing of Nutricline Dynamics in the Equatorial Atlantic. *Science*, **249**, 766-769.

MOSTAJO, E. L. (1986) Estudio de dos testigos submarinos del oceano Pacífico Sudecuatorial. *Revista Española de Micropaleontología*, 18 (3): 433-442.

MÜLLER, P.J. e Fischer, G. (2004) C37-Alkenone as Paleotemperature Tool: Fundamentals Based on Sediment traps and Surface Sediments from the South Atlantic Ocean. In: WEFER, G. MULITZA, S. e RATMEYER, V. (Ed.). **The South Atlantic in the Late Quaternary**. Berlin: Springer-Verlag. p. 167-193.

MÜLLER, P.J., Cepek, M., Ruhland, G. e Schneider, R.R. (1997) Alkenone and coccolithophorid species changes in late Quaternary sediments from the Walvis Ridge: Implications for the alkenone paleotemperature method. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **135**, 71-96.

- NYBAKKEN, J. W. (2001) Introduction to the marine environment. In: NYBAKKEN, J. W. **Marine Biology an ecological approach**. 5. ed. São Francisco: Benjamin Cummings. p. 1-37.
- ODUM, E.P. (2004) **Fundamentos de Ecologia**. 7. ed. Lisboa: Fundação Calouste Gulbenkian. 927p.
- OKADA, H. (1992) Biogeographic control of modern nannofossil assemblages in surface sediments of Ise Bay, Mikawa Bay e Kumano-Nada, off coast of central Japan. *Memorie Di Scienze Geologiche*, **43**, 431-449.
- OKADA, H. (1983). Modern nannofossil assemblages in sediments of coastal and marginal seas along the western Pacific ocean. *Utrecht Micropaleontology Bulletin*, **30**, 171-187.
- OKADA, H. e McIntyre, A. (1979) Seasonal distribution of modern coccolithophorids in the western North Atlantic. *Marine Biology*, **54**, 319-328.
- OKADA, H., McIntyre, A. (1977) Modern coccolithophores of the Pacific and North Atlantic Oceans. *Micropaleontology*, **23** (1), 1-55.
- OKADA, H. e Honjo, S. (1973) The distribution of oceanic coccolithophorids in the Pacific. *Deep-Sea Research*, **20**, 355-374.
- OLIVEIRA, P.E., Behling, H., Ledru, M.-P., Barberi, M., Bush, M., Salgado-Labouriau, M.L., Garcia, M.J., Medeanic, S., Barth, O.M., Barros, M.A., Scheel-Ybert, R. (2005) Paleovegetação e Paleoclimas do Quaternário do Brasil. In: SOUZA, C.R.G., SUGUIO, K., OLIVEIRA, A.M.S. e OLIVEIRA P.E. (Ed.). **Quaternário do Brasil**. Ribeirão Preto: Editora Holos. p. 52-74.
- OLIVEIRA, P. E., Barreto, A. M. F. e Suguio, K. (1999) Late Pleistocene/Holocene climatic and vegetational history of the Brazilian caatinga: the fossil dunes of the middle São Francisco River. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **152**, 319-337.
- PARKER, W.C. e Arnold, A.J. (1999) Quantitative methods of data analysis in foraminiferal ecology. In: GUPTA, B.K.S. (Ed.), **Modern Foraminifera**. Kluwer Academic Publishers. p. 71-89.

- PETERSON, L. C., Haug, G. H., Hughen, K. A., Röhl, U. (2000) Rapid changes in the hydrologic cycle of the Tropical Atlantic during the last glacial. *Science*, **290**, 1947-1951.
- PETERSON, R. G. e Stramma, L. (1991) Upper-level circulation in the South Atlantic. *Proceedings in Oceanography*, **26**, 1-73.
- PONTE, F.C. e Asmus, H.E. (2004) As bacias marginais brasileiras: estágio atual de conhecimento. *Boletim de Geociências da Petrobrás*, **12** (2), 385-420.
- PRENTICE, I.C. (1986) Multivariate methods for data analysis. In: BERGLUND, B.E. (Ed.), **Handbook of Holocene Palaeoecology and Palaeohydrology**. Chichester: Ed. Willey. p. 775-797.
- RAHMSTORF, S. (2002) Ocean circulation and climate during the past 120,000 years. *Nature*, **419**, 207-214.
- RENAUD, S. Ziveri, P., Broerse, A.T.C. (2002) Geographical and seasonal differences in morphology and dynamics of the coccolithophore *Calcidiscus leptoporus*. *Marine Micropaleontology*, **46**, 363-385.
- ROTH, P. (1994) Distribution of coccoliths in oceanic sediments. In: WINTER, A. & SIESSER, W. (Eds), **Coccolithophores**. Cambridge: Cambridge University Press, p. 199-218.
- RÜHLEMANN, C., Müller, P.J. e Schneider, R.R. (1999) Organic Carbon and Carbonate as Paleoproductivity Proxies: Examples from High and Low Productivity Areas of the Tropical Atlantic. In: FISCHER, G., WEFER, G. (Ed.). **Use of proxies in Paleoceanography: Examples from the South Atlantic**. Berlin: Springer-Verlag. p. 315-344.
- RÜHLEMANN, C., Frank, M., Hale, W., Mangini, A., Mulitza, S., Müller, P.J. e Wefer, G. (1996) Late Quaternary productivity changes in the western equatorial Atlantic: evidence from 230TH-normalized carbonate and organic carbon accumulation rates. *Marine Geology*, **135**, 127-152.

- SACHS, J.P., Schneider, R.R., Eglinton, T.I., Freeman, K.H., Ganssen, G., McManus, J.F. e Oppo, D.W. (2000) Alkenones as paleoceanographic proxies. *Geochemistry, Geophysics, Geosystems*, **1**, 2000GC000059.
- SANT'ANNA NETO, J.L. e Nery, J.T. (2005) Variabilidade e Mudanças Climáticas no Brasil e seus Impactos Regionais. In: SOUZA, C.R.G., SUGUIO, K., OLIVEIRA, A.M.S. e OLIVEIRA P.E. (Ed.). **Quaternário do Brasil**. Ribeirão Preto: Editora Holos. p. 28-51.
- SANTOS, R.C.R. e Babinski, N.A. (1996) Relatório de viagem dos trabalhos de pesquisas petrolíferas desenvolvidas no período de 26/01 a 05/03/96 pelo projeto conjunto da PETROBRAS/CENPES e GERG da Texas A e M University (USA) em águas jurisdicionais da plataforma continental brasileira. PETROBRAS/DIVEX/SEGEQ, Rio de Janeiro.
- SCHMIDT, D.N. (2002) **Size Variability in Planktic Foraminifers**. In: Swiss Federal Institute of Technology, Zürich. Tese de Doutorado, pp. 121.
- SCHRAGG, D. P., Hampt, G. e Murray, D. W. (1996) Pore fluid constraints on the temperature and isotopic composition of the glacial ocean. *Science*, **272**, 1930-1932.
- SEIDOV, D. e Maslin, M. (2001) Atlantic Ocean heat piracy and the bipolar climate see-saw during Heinrich and Dansgaard-Oeschger events. *Journal of Quaternary Sciences*, **16**, 321-328.
- SHACKLETON, N.J. (1987) Oxygen isotopes, ice volume and sea level. *Quaternary Science Reviews*, **6**, 183-190.
- SHACKLETON, N.J. (1977) $\delta^{13}\text{C}$ in *Uvigerina*: Tropical rainforest history and the Equatorial Pacific carbonate dissolution cycles. In: ANDERSEN, N. R. e MALAHOFF, A. (Ed.). **The Fate of Fossil Fuel CO₂ in the Oceans**. New York: Plenum. p. 401-427.
- SIESSER, W.G. (1994) Historical background of coccolithophore studies. In: WINTER, A. & SIESSER, W. (Eds.). **Coccolithophores**. Cambridge: Cambridge University Press, p. 1-12.

- SIESSER, W.G., Winter, A. (1994) Composition and morphology of coccolithophore skeletons. In: WINTER, A. & SIESSER, W. (Eds.). **Coccolithophores**. Cambridge: Cambridge University Press, p. 51-62.
- SILVEIRA, I.C.A., Miranda, L.B., Brown, W.S. (1994) On the origins of the North Brazil Current. *Journal of Geophysical Research*, **99** (c11), 22,501-22,512.
- STEINMETZ, J.C. (1994) Stable isotopes in modern coccolithophores. In: WINTER, A. & SIESSER, W. (Ed.). **Coccolithophores**. Cambridge: Cambridge University Press, p.219-229.
- STRAMMA, L. (1989) The Brazil Current transport south of 23°S. *Deep-Sea Research*, **36**, 639-646.
- STRAMMA, L., Ikeda, Y., Peterson, R.G. (1990) Geostrophic transport in the Brazil Current region north of 20°S. *Deep-Sea Research, Part A*, **37**, 1875-1886.
- SUGUIO, K. (1999) **Geologia do Quaternário e Mudanças Ambientais** (Passado + Presente = Futuro?). São Paulo: Editora Paulo's. 366p.
- TAKANO, Y., Hagino, K., Tanaka, Y., Horiguchi, T., Okada, H. (2006) Phylogenetic affinities of an enigmatic nanoplankton, *Braarudosphaera bigelowii* based on the SSU rDNA sequences. *Marine Micropaleontology*, **60**, 145-156.
- THOMSON, J., Crudeli, D., de Lange, G.J., Slomp, C.P., Erba, E., Corselli, C., Calvert, S.E. (2004) *Florisphaera profunda* and the origin and diagenesis of carbonate phases in eastern Mediterranean sapropel units. *Paleoceanography*, **19**, PA3003.
- TOKUTAKE, L.R. (2005) **Bioestratigrafia de Nanofósseis Calcários e Estratigrafia de Isótopos (C e O) do Talude médio, Quaternário, porção N da Bacia de Campos, ES**. Dissertação de Mestrado. Universidade Federal do Rio Grande do Sul – UFRGS. Instituto de Geociências. Curso de Pós Graduação em Geociências, Porto Alegre, RS, Brasil. 96p.
- TOLEDO, F.A.L. (2000) **Variações Paleoceanográficas nos últimos 30.000 anos no oeste do Atlântico Sul: isótopos de oxigênio, assembléias de foraminíferos planctônicos e**

nanofósseis calcários. Tese de doutorado. Universidade Federal do Rio Grande do Sul – UFRGS. Instituto de Geociências. Curso de Pós Graduação em Geociências, Porto Alegre, RS, Brasil. 245p.

TOLEDO, F.A.L. (1995) Nanofósseis Calcários Quaternários da Bacia de Campos - Sistemática e Interpretação Paleoambiental. In: Instituto de Geociências, Universidade Federal do Rio Grande do Sul, Porto Alegre. Dissertação de Mestrado, pp. 159.

TOLEDO, F.A.L., Costa, K.B. e Pivel, M.A.G. (2007a) Salinity changes in the western tropical South Atlantic during the last 30 kyr. *Global and Planetary Change*, DOI:10.1016/j.gloplacha.2007.01.001

TOLEDO, F.A.L., Cachão, M., Costa, K.B. e Pivel, M.A.G. (2007b) Planktonic foraminifera, calcareous nannoplankton and ascidian variations during the last 25 kyr in the Southwestern Atlantic: a paleoproductivity signature? *Marine Micropaleontology*, doi: 10.1016/j.marmicro.2007.03.001.

TOLEDO, F.A.L., Ayup-Zouain, R.N. e Costa, K.C. (1999) Análise fatorial (modo-Q) em estudos quantitativos de nanofósseis calcários do Quaternário superior em um testemunho da Bacia de Campos, RJ-Brasil. *Pesquisas*, **26**(2), 62-72.

TREMOLADA, F., Erba, E., van de Schootbrugge, B., Mattioli, E. (2006) Calcareous nannofossil changes during the late Callovian-early Oxfordian cooling phase. *Marine Micropaleontology*, **59**, 197-209.

TRIANANTAPHYLLOU, M.V., Ziveri, P., Tselepides, A. (2004) Coccolithophore export production and response to seasonal surface water variability in the oligotrophic Cretan Sea (NE Mediterranean). *Micropaleontology*, **50** (1), 127-144.

VAROL, O. (2006) Didemnid ascidian spicules from the Arabian Peninsula. *Journal of Nannoplankton Research*, **28** (1), 35-55.

VITAL, H., Esteves, L.S., Araújo, T.C.M., Patchineelam, S.M. (2005) Oceanografia Geológica e Geofísica da Plataforma Continental Brasileira. In: SOUZA, C.R.G., SUGUIO, K.,

OLIVEIRA, A.M.S. e OLIVEIRA P.E. (Ed.). **Quaternário do Brasil**. Ribeirão Preto: Editora Holos. p. 153-175.

WEFER, G., Berger, W.H., Bijma, J. e Fischer, G. (1999) Clues to Ocean History: a Brief Overview of Proxies. In: FISCHER, G., WEFER, G. (Ed.). **Use of proxies in Paleoceanography: Examples from the South Atlantic**. Berlin: Springer-Verlag. p. 1-68.

WELDEAB, S., Schneider, R.R. e Kölling, M. (2006) Deglacial sea surface temperature and salinity increase in the western tropical Atlantic in synchrony with high latitude climate instabilities. *Earth and Planetary Science Letters*, **241**, 699-706.

WELLS, P., Okada, H. (1996) Holocene and Pleistocene glacial paleoceanography off southeastern Australia, based on foraminifers and nannofossil in Vema cored hole V18-222. *Australian Journal of Earth Sciences*, **43**, 509-523.

WINTER, A., Jordan, R.W., Roth, P.H. (1994). Biogeography of the living coccolithophores in ocean waters. In: WINTER, A. & SIESSER, W. (Eds.). **Coccolithophores**. Cambridge: Cambridge University Press, p.161-178.

WOA98, World Ocean Atlas, 1998. Disponível em: <http://ingrid.ldeo.columbia.edu/SOURCES/.NOAA/.NODC/.WOA98> Acesso em: 5 novembro 2006.

WOLFF, T., Grieger, B., Hale, W., Dürkoop, A., Mulitza, S., Pätzold, J. e Wefer, G. (1999) On the Reconstruction of Paleosalinities. In: FISCHER, G., WEFER, G. (Ed.). **Use of proxies in Paleoceanography: Examples from the South Atlantic**. Berlin: Springer-Verlag. p. 207-228.

WOLFF, T., Mulitza, S., Arz, H., Pätzold, J. e Wefer, G. (1998) Oxygen isotopes versus CLIMAP (18 ka) temperatures: A comparison from the tropical Atlantic. *Geology*, **26**, 675-678.

YOUNG, J.R. (1994) Functions of coccoliths. In: WINTER, A. & SIESSER, W. (Ed.). **Coccolithophores**. Cambridge: Cambridge University Press, p. 63-82.

- YOUNG, J. R., Geisen, M., Cros, L., Kleijne, A., Sprengel, C., Probert, I. e Ostergaard, J. (2003) A guide to extant coccolithophore taxonomy. *Journal of Nannoplankton Research*, Special Issue (1), 125p.
- YOUNG, J.R., Bown, P.R. (1997a) Higher classification of calcareous nannoplankton. *Journal of Nannoplankton Research*, **19**, 15-20.
- YOUNG, J.R., Bown, P.R. (1997b) Cenozoic calcareous nannoplankton classification. *Journal of Nannoplankton Research*, **19**, 36-47.
- ZABEL, M.T., Wagner, T. e DeMenocal, P. (2004) Terrigenous Signals in Sediments of the Low Latitude Atlantic – Indications to Environmental Variations during the Late Quaternary: Part II: Lithogenic Matter. In: WEFER, G. MULITZA, S. e RATMEYER, V. (Ed.). **The South Atlantic in the Late Quaternary**. Berlin: Springer-Verlag. p. 323-345.
- ZABEL, M., Bickert, T., Dittert, L. e Haese, R. R. (1999) Significance of the sedimentary Al:Ti ratio as an indicator for variations in the circulation patterns of the equatorial North Atlantic. *Paleoceanography*, **14**, 789-799.
- ZONNEVELD, K.A.F., Höll, C., Janofske, D., Karwath, B., Kerntopf, B., Rühlemann, C., Willems, H. (1999) Calcareous Dinoflagellate Cysts as Paleo-Environmental Tools. In: FISCHER, G., WEFER, G. (Ed.). **Use of proxies in Paleoceanography: Examples from the South Atlantic**. Berlin: Springer-Verlag. P. 145-164.