

MARINA TONETTI BOTANA

The role of *Symbiodinium* membrane lipids in response to heat shock: implications for coral bleaching

Thesis presented to the Instituto Oceanográfico of the Universidade de São Paulo, in partial fulfillment of the requirements for obtaining the degree of Master in Sciences, Oceanography Program, concentration area of Biological Oceanography.

Supervisor: Prof. Dr. Paulo Y. G. Sumida

São Paulo

2019

Universidade de São Paulo
Instituto Oceanográfico

**The role of *Symbiodinium* membrane lipids in
response to heat shock: implications for coral
bleaching**

Marina Tonetti Botana

Thesis presented to the Instituto Oceanográfico of the
Universidade de São Paulo, in partial fulfillment of the
requirements for obtaining the degree of Master in Sciences,
Oceanography Program, concentration area of Biological
Oceanography.

Evaluated in:

_____	_____
Prof(a). Dr.(a)	Grade
_____	_____
Prof(a). Dr.(a)	Grade
_____	_____
Prof(a). Dr.(a)	Grade

“Look up at the stars and not down at your feet. Try to make sense of what you see and wonder about what makes the universe exist. Be curious. Science is not only a discipline of reason, but also, one of romance and passion”

Steve Hawking

“Commit yourself to the process. Not the goal. The final score take care of itself”

James Clear

ACKNOWLEDGEMENTS

I would like to thank my advisor Paulo Sumida for giving me the opportunity to pursue my masters in his lab. He allowed me to work on something I am very passionate for and even supported my parallel career as a triathlete. I feel privileged for having him in my life as a tutor and friend. My gratitude also goes to Dr. Marcos Yoshinaga who first presented to me this fascinating world of lipids and without whom this work would never have been possible. I thank all his tutoring and friendship, both pivotal for guiding this moment of my career. Additionally, immense thanks to Prof. Sayuri Miyamoto who opened the doors of her lab to my project and from whom I also could learn a lot about lipids metabolism and oxidative stress. My deep gratitude extends to Dra. Flavia Saldanha-Correa who helped me with the experiments and showed me other cool applications of studying lipids in other phytoplankton phylotypes. Thanks to Dr. Marius Müller for the skype talks adding ideas on the ecological implications of my results.

Regarding my learning process I also thanks Dr. Matthias Kellerman and Prof. Ray Valentine for kindly teaching me about the universal principle of bioenergetics and how the existence of life is utterly dependent on lipid membranes. My great appreciation also to Dr. Florence Schubotz for sending valuable lipid standards and enabling better results.

For all their gentle help in the lab I thank Alex Inague, Adriano Brito and Debora Kutner. Also, Thomas Banha and Arthur Güth for the many talks about my experiment design and application of lipids in coral reefs ecology. Thanks to everyone from Sumida's and Sayuri's laboratories for our daily routines full of scientific learnings and lots of funny stories in the last two years.

Special thanks to my training buddies and triathlon coaches who also were part of my routine from Sunday to Sunday in the last years and showed me that everything is possible. Thanks for teaching me that *interest is different from commitment* and that we must commit every day to the process of doing things that we really believe and are passionate for, because that is what make us successful in the long term. That is what make us grow as better athletes, scientists and human beings.

Last, but not least, I would mainly thank my parents for being supportive along the way and believing in my career as a scientist. Since I was little, I was a very curious for ocean things and they gave me an environment in which my curiosity could grow into passion and then I could turn it into my something useful for nature and society. I would never be able to put into words how grateful I feel for being your daughter.

This study was financed in part by the Coordenação de Aperfeiçoamento de Nível Superior (CAPES) – Finance Code 001.

RESUMO

Recifes de coral do mundo inteiro vêm sendo devastados pelo fenômeno de branqueamento, o qual as evidências indicam que seja causado pelo stress oxidativo promovido pelo aquecimento global e eventos catastróficos de El Niño. A grande variabilidade genética da Família Symbiodiniacea também é sugerida como determinante da susceptibilidade do coral hospedeiro porque cada espécie possui limites fisiológicos específicos, tanto no modo de vida livre como em simbiose. Neste estudo apresentamos pela primeira vez o sucesso da utilização das técnicas de lipidômica (i.e, caracterização dos lipídeos globais em um determinado organismo) oferecendo suporte para as investigações moleculares de investigação dos mecanismos relacionados ao stress térmico em espécies de endosimbiontes de coral. *Symbiodinium minutum* foi sensível às temperaturas elevadas, enquanto *S. microadriaticum* e *S. goreau* apresentaram distintos níveis de termo tolerância. Os fenótipos lipídicos das espécies após o stress, incluindo o transportador de elétrons do fotossistema II – plastoquinona – sugerem que cada um apresentou uma estratégia diferente para sobreviver. Além disso, os lipídeos específicos do cloroplasto com ácidos graxos poliinsaturados (PUFA) formado, principalmente, por espécies com ômega 3 (n-3) foram essenciais para manter a bioenergética celular à longo prazo (10 dias após stress) em todos os *Symbiodinium* spp. A capacidade de manter altas concentrações de n-3 na membrana dos cloroplastos determinou a sobrevivência dos *S. microadriaticum* e *S. goreau*. Os dados apresentados nesta dissertação revelam, pela primeira vez, o aumento de ácidos graxos oxidados na membrana do cloroplasto e também na forma livre (FFA) em resposta aos dados de stress oxidativo causados pelo calor. O estudo das membranas lipídicas é fundamental para melhor compreensão da bioenergética dos simbiontes e para determinar a vulnerabilidade da relação de simbiose com o coral aos estressores climáticos em um futuro com temperaturas mais elevadas.

Palavras-chave: *Symbiodinium*, recifes de coral, lipidômica, stress térmico, stress oxidativo.

ABSTRACT

Coral reefs around the world have been largely devastated by the phenomenon of “coral bleaching”, which causes have been reported to be strongly related to oxidative stress promoted by climate change drivers, including mainly global warming and catastrophic El Niño events. Genetic variability in coral endosymbionts from the Family Symbiodiniacea was also suggested as determinant of host susceptibility to stress because they present distinct physiological boundaries when in free living or in symbiosis. Here we present for the first time the successful use of lipidomics (*i.e.*, the global characterization of lipids in a given organism) supporting molecular investigation in the oxidative mechanisms related to thermal stress in coral endosymbionts phylotypes. *Symbiodinium minutum* was thermal sensitive, whereas *S. microadriaticum* and *S. goreau* presented different levels of thermal tolerance. Their lipid phenotypes after stress, including the photosystem electron transporter - plastoquinone - suggested they had different survival strategies. In addition, chloroplast specific lipids with polyunsaturated fatty acids (PUFAs) mainly formed by omega 3 (n-3) seemed to be essential to sustain *Symbiodinium* cells bioenergetics in the long term (10 days after stress). *S. microadriaticum* and *S. goreau* capability of keeping high n-3 concentrations in the chloroplast membranes determined their survival. The present thesis reports, for the first-time, upregulation of oxidized lipids derived from precursor chloroplast membranes and free fatty acids (FFA) in response to oxidative stress damage caused by heat. The study of lipid membranes is of paramount importance to better understand the bioenergetics of symbionts and to determine the host/endosymbiont vulnerability to climate change stressors in a warmer future.

Key words: *Symbiodinium*, coral reefs, lipidomics, thermal stress, oxidative stress

LIST OF ACRONYMS AND ABBREVIATIONS

ARA - arachidonic acid

AC - *Symbiodinium* phylotype A1 control sample

AT – *Symbiodinium* phylotype A1 temperature stressed sample

ATP – adenosine triphosphate

BC - *Symbiodinium* phylotype B1 control sample

BT - *Symbiodinium* phylotype B1 temperature stressed sample

CC - *Symbiodinium* phylotype C1 control sample

CE – cholesterol ester

Cer - ceramide

Chl-a -chlorophyl-a

CL - cardiolipin

CT - *Symbiodinium* phylotype C1 temperature stressed sample

DAG – diacylglycerol

DGCC - 1,2-diacylglycerol-3-(O-carboxyhydroxymethylcholine)

DGDG – digalactosyldiacylglycerol

DGTS - diacylglyceroltrimethylhomoserine

DHA - docosahexaenoic acid

EPA - eicosapentaenoic acid

ER - endoplasmic reticulum

ESI-TOFMS – electron spray ionization time of flight mass spectrometer

FFA - free fatty acids

Gluc Acid – glucuronic acid

HUFA – High unsaturated fatty acids

LC - liquid chromatography

MGDG - monogalactosyldiacylglycerol

MUFA - monounsaturated fatty acid

MS – mass spectrometer

PC – phosphatidylcholine

PE - phosphatidylethanolamine

PG – phosphatidylglycerol

PI - phosphatidylinositol

PL – polar lipids

PUFA - polyunsaturated fatty acid

RPLC – reverse phase liquid chromatography

ROS - reactive oxygen species

SFA - saturated fatty acid

SQDG – sulfoquinovosyldiacylglycerol

SM - sphingomyelin

TAG - triacylglycerol

UHPLC - ultra-high-performance liquid chromatography

LIST OF FIGURES

List is presented in the order they appeared in the text.

Figure 1.1: Bleached corals of the genus *Mussismilia hispida*, endemic from the tropical waters of Brazil, observed in Ubatuba (São Paulo State north shore) in the summer of 2019.....2

Figure 1.2: Exemplification of how membrane lipids are affected by fatty acid composition and temperature. Figure shows how polar lipids with unsaturated fatty acids attribute high motion and low viscosity to the membrane. Plus, elevated temperatures also promote higher membrane's motion and fluidity. Combination of elevated temperatures and high polyunsaturated fatty acids can compromise membrane's electron transport chain viability.....4

Figure 1.3: Electron transport chain resulting in ATP synthesis in the thylakoid membranes of chloroplasts. Protein complexes photosystem II (PSII), cytochrome b_6-f (cyt b_6-f) and photosystem I (PSI) are highlighted. Plastoquinone pool responsible for electron transport from PSII to cyt b_6-f is circled in red.....5

Figure 2.1: Diversity of lipid species identified in *Symbiodinium* phylotypes A1, B1 and C1 sorted into their respective lipid classes. Abbreviations: AMINO = aminolipids, GLYCO = glycolipids, SPHINGO = sphingolipids, PHOSPHO = phospholipids. Detailed description of lipid subclasses abbreviations is shown in Table S2.1.....17

Figure 2.2: Lipid molecular species profiles of *Symbiodinium* phylotypes A1, B1 and C1 control samples. Cultures grew at 22 °C; 80 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$; 12L:12D cycle. Bars show the mean abundance of each lipid class per cells and their respective standard errors. Storage lipids encompass cholesterol esters and triacylglycerol; membrane lipids group sphingolipids, phospholipids, glycolipids and aminolipids; Diacylglycerol (DAG) and free fatty acids (FFA) are presented alone. * represents statistically significantly different ($p < 0.05$) mean values inside their respective lipid groups. Fig.S1.1 exhibits separately details about storage lipids and membrane lipids. AC, BC and CC stand for control samples of A1, B1 and C1, respectively.....18

Figure 2.3: Plastoquinone concentration in *Symbiodinium* phylotypes A1, B1 and C1; and its respective molecular structure. * indicates that only BC is significantly different ($p < 0.05$) from other mean values in AC and CC. AC, BC and CC stand for control samples of A1, B1 and C1, respectively.....22

Figure 2.4: Heatmap of the most significantly different lipid molecular species between *Symbiodinium* phylotypes A1, B1 and C1. In total, 40 lipid molecular species are shown in rows and

samples in column. Statistical significance was evaluated by one-way ANOVA followed by Turkey's post-hoc test ($p < 0.05$) using Metaboanalyst. Distance was measured in Pearson, and Ward's clustering algorithm. Each colored cell on heatmap corresponds to normalized concentrations. Log transformation was used for data normalization value. Red color indicates upregulation, whereas blue means downregulation. AC, BC and CC stand for control samples of A1, B1 and C1, respectively.....24

Figure 2.5: Distribution of main omega 3 (n-3) fatty acids found in *Symbiodinium* phylotypes A1, B1 and C1 on their respective lipid polar heads. a) octadecatetraenoic acid (18:4) distribution; b) octadecapentaenoic acid (18:5) distribution; c) docosahexaenoic acid distribution; d) relative percentage of main omega 3 (n-3) fatty acids to total lipids in the 3 studied phylotypes. Means showing * are significantly different ($p < 0.05$) from other mean values inside their respective lipid groups. Means showing # are exclusively significantly different from each other (same $p < 0.05$). Vertical lines represent the standard error. AC, BC and CC stand for control samples of A1, B1 and C1, respectively.....26

Figure S2.1: Membrane and storage lipid subclasses in *Symbiodinium* phylotypes A1, B1 and C1. a) main membrane lipids subclasses. Sphingolipids are not represented here because their concentrations were lower than 10^{-4} ; b) storage lipid subclasses. Means showing an * are significantly different ($p < 0.05$) from other mean values inside their respective lipid subclasses. Vertical lines represent the standard error. AC, BC and CC stand for control samples of A1, B1 and C1, respectively.....32

Figure S2.2: Pigments in *Symbiodinium* phylotypes A1, B1 and C1. a) most abundant pigments; b) minor representative pigments. Means were not significantly different. Vertical lines represent the standard error. AC, BC and CC stand for control samples of A1, B1 and C1, respectively.....33

Figure S2.3: Most abundant lipid subclasses in *Symbiodinium* phylotypes A1, B1 and C1. a) monogalactosyldiacylglycerol (MGDG); b) digalactosyldiacylglycerol (DGDG); c) 1,2-diacylglycerol-3-(O-carboxyhydroxymethylcholine) – DGCC; concentrations of oxo-DHA were significantly lower in phylotype C1, plus concentrations of DHA-OH were significantly higher in B1; d) phosphatidylcholine (PC); e) triacylglycerol (TAG); f) free fatty acids (FFA). Means showing an * are significantly different ($p < 0.05$) from other mean values inside their respective lipid subclasses. Means showing # are exclusively significantly different from each other (same $p < 0.05$). Vertical lines represent the standard error. AC, BC and CC stand for control samples of A1, B1 and C1, respectively.....34

Figure S2.4: Minor abundant phospholipids in *Symbiodinium* phylotypes A1, B1 and C1. a) phosphatidylglycerol (PG); b) phosphatidyletanolamina; c) phosphatidylinositol. Means were not significantly different. Vertical lines represent the standard error. AC, BC and CC stand for control samples of A1, B1 and C1, respectively.....35

Figure S2.5: Minor compounds significantly different in *Symbiodinium* phylotypes A1, B1 and C1. a) diacylglyceroltrimethylhomoserine (DGTS); b) Sphingolipids; c) Cholesterol ester (CE). Means showing an * are significantly different ($p < 0.05$) from other mean values inside their respective lipid subclasses.

Means showing # are exclusively significantly different from each other (same $p < 0.05$). Vertical lines represent the standard error. AC, BC and CC stand for control samples of A1, B1 and C1, respectively.....36

Figure S2.6: Other minor compounds not significantly different in *Symbiodinium* phylotypes A1, B1 and C1. a) diacylglycerol (DAG); b) Cholesterol; c) sulfoquinovosyldiacylglycerol (SQDG); d) Glucuronic acid (Gluc Acid). Vertical lines represent the standard error. AC, BC and CC stand for control samples of A1, B1 and C1, respectively.....37

Figure S2.7: Chlorophyll-a molecule fragments break spectrum. Molecule draw and exact mass calculations were performed with ChemDraw.....38

Figure S2.8: Monogalactosyldiacylglycerol molecule ionized with ammonium fragments break spectrum. High peak of 333 indicates double 18:4 fatty acids and 161 represents sugar head. Molecule draw and exact mass calculations were performed with ChemDraw.....38

Figure S2.9: 1,2-diacylglycerol-3-(O-carboxyhydroxymethylcholine) molecule fragments break spectrum. 562 represents loss of DHA fatty acids and 544 represents loss of DHA plus water. Molecule draw and exact mass calculations were performed with ChemDraw.....39

Figure S2.10: Diacylglyceroltrimethylhomoserine molecule fragments break spectrum. 236 represents polar head and 500 represents loss of 18:1 fatty acids. Molecule draw and exact mass calculations were performed with ChemDraw.....39

Figure S2.11: Phosphatidylcholine molecule ionized with acetate fragments break spectrum. 168 represents its polar head and each indicated arrow on figure indicate fatty acid losses. Molecule draw and exact mass calculations were performed with ChemDraw.....40

Figure S2.11: Phytoceramide molecule fragments ionized with acetate break spectrum. Molecule draw and exact mass calculations were performed with ChemDraw.....40

Figure S2.12: Triacylglycerol molecule ionized with ammonium fragments break spectrum. Fatty acids were determined based on exact mass loss. Higher peak of 577 indicates double loss of 16:0 and 551 peak indicates loss of 18:1. Molecule draw and exact mass calculations were performed with ChemDraw.....41

Figure S2.13: Cholesterol ester molecule ionized with ammonium fragments break spectrum. Molecule draw and exact mass calculations were performed with ChemDraw.....41

Figure 3.1: Growth rate curves of control (AC, BC, CC) and temperature stressed *Symbiodinium* phylotypes cultures (AT, BT, CT) with its respective μ values and standard deviation. Regression curves were obtained with n=3 (triplicates). a) Growth rate curves of phylotype A1; b) Growth rate curves of phylotype B1; c) Growth rate curves of phylotype C1. Images below graphs show final culture appearance. Note that lighter coloration of cultures denotes lower cell densities.....50

Figure 3.2: Lipid remodeling data indicating distinct phylotype response time that are correspondent to growth rate curves behavior. a) number of significant modified lipid species overtime after heat shock event comparing samples from stressed cultures against samples from control (C x T). X axis presents time in hours after heat shock event. Data points were based on the cultures sampling made 4 h, 24 h (1 day) and 240 hours (10 days) after stress; b) same significant modified compounds sorted into the main lipid groups and the same sampling times. Bars above zero show species that were “upregulated” compared to control; whereas bars below zero show lipids that were “downregulated” compared to control. The * represents no upregulation of glycolipids in phylotype B1 after 10 days. Differences were evaluated through Volcano-Plot analysis (FC = 1.5 and p < 0.05; FDR adjusted). Detailed data and molecular lipid compounds are presented on Table S3.1.....52

Figure 3.3: Total variation of omega 3 polyunsaturated fatty acids over time. Relative percentage of all omega 3 (n-3) fatty acids (DHA, EPA, DPA, 18:3, 18:4 & 18:5) was considered for each *Symbiodinium* phylotype considering 4- and 24-hours variations as short term and 10 days as long-term effect. Variations were calculated based on values from stressed minus control lipid data in each sampling time. Bars show standard error values obtained from triplicates.....53

Figure 3.4: Correlation of polyunsaturated fatty acids (PUFA) in the most abundant membrane lipids and plastoquinone. Respective R² and p values are represented together in each graph. Glycolipids were calculated considering monogalactosyldiacylglycerol (MGDG), digalactosyldiacylglycerol (DGDG), sulfoquinovosyldiacylglycerol (SQDG) and glucuronic acid (GlucA). 1,2-Diacylglycerol-3-(O-carboxyhydroxymethylcholine) was the only aminolipid associated with PUFAs, as well as phosphatidylcholine (PC) for phospholipids. All lipid data is represented in ng per cell.....55

Figure 3.5: Pigments ratios that changed significantly after heat shock. Ratios are represented in logarithmic scale to evidence its up or downregulation comparing temperature stressed cultures (T) and control cultures (C) in three distinct sampling times: a) 4 hours; b) 24 hours; c) 10 days after heat shock event. * represent samples statistically significant altered based on Volcano Plot analysis (FC = 1.5 and p < 0.05) and details are on Table S3.1.....56

Figure 3.6: Ratios of total concentrations of oxidized fatty acids and their precursors from polar lipids (PLs) and free fatty acids (FFAs). 15 molecules represented by free fatty acids (6) and polar lipids (8) were identified and monitored during our experiment. Ratios considered total concentration of each identified fatty acid in either free or membrane associated forms. a) short term (4h and 24h after heat

shock); b) long term (10 days after heat shock). Bars represent average concentrations with respective standard errors. Temperature stressed samples values are represented by (T), while control samples are represented by (C).....58

Figure 3.7: Heatmap of oxylipins ratios that changed significantly after heat shock. In total, 12 oxylipins ratios are shown in rows and samples in columns. Statistical significance was evaluated by one-way ANOVA followed by Turkey's post-hoc test ($p < 0.05$) using Metaboanalyst. Distance was measured in Pearson, and Ward's clustering algorithm. Each colored cell on heatmap corresponds to normalized concentrations. Log transformation was used for data normalization values. Red color indicates upregulation, whereas blue means downregulation of oxylipins ratios. AC, BC and CC samples represent controls. AT, BT and CT represent temperature stressed samples. 4h, 24h and 10 days represent sampling times after heat shock for both stressed (T) and control (C) samples. Middle black rectangle groups phylotype B1 samples after heat shock and green rectangle highlights single higher concentrations of FFA oxylipins.....60

Figure F1: Summary of *Symbiodinium* phylotypes cell's proposed mechanisms caused by upregulation of ROS. Temperature alters membrane viscosity and enable scape of high energy electrons from electron transport chain. This might be responsible for upregulation of ROS that will concomitantly lead to the peroxidation of most vulnerable omega-3 polyunsaturated fatty acids and activate quenching mechanisms. We propose that phylotype B1 did not survive because it could not avoid lipid peroxidation of essential ω -3 in the membranes, neither synthesize *de novo* epoxidized lipids. On the other hand, it was not true for phylotypes A1 and C1. Both kept efficient energy output, although we suggest that they followed distinct strategies already discussed in chapter 3. Protein complexes photosystem II (PSII), cytochrome b_6-f (cyt b_6-f) and photosystem I (PSI) were highlighted. Plastoquinone pool responsible for electron transport from PSII to cyt b_6-f was also highlighted in red. Scape of high energy electron leading to ROS was highlighted in dark red.....70

LIST OF TABLES

Table S2.1: Internal standards used for quantification of sphingolipids, phospholipids, storage lipids and plastoquinone described in the material and methods section.....42

Table S2.2: Additional standards used for quantification of glycolipids, aminolipids and pigments described in the material and methods section.....42

Table S2.3: All identified lipid compounds including their exact mass and retention time in both positive and negative modes. File is in excel spreadsheet available in Research Gate profile website.....INDEX

Table S3.1: Description of lipids and pigments compounds either up or downregulated after heat shock for *Symbiodinium* phylotypes A1, B1 and C1 after 4h, 24h and 240h. File is in excel spreadsheet available in Research Gate profile website.....INDEX

SUMMARY

CHAPTER 1: GENERAL INTRODUCTION

1.1 Coral reefs and their symbionts in the context of global warming.....	1
1.2 Coral bleaching and the oxidative stress theory.....	2
1.3 The effect of high temperature in membrane's viscosity leading to free radical and ROS formation in the chloroplast.....	3
1.4 Lipid membrane profiles and symbionts fate after thermal stress.....	6
1.4.1 Lipidomics as tool for better explanation of thermal sensibility.....	7
2. Aimed goals and scope of this thesis.....	8

CHAPTER 2: Lipidomics as a throughput method for profiling lipids and pigments in *Symbiodinium* phylotypes (A1, B1 & C1): physiological and ecological applications

1. INTRODUCTION.....	10
2. MATERIALS AND METHODS.....	12
2.1 Experiment design.....	12
2.2 Internal standards.....	12
2.3 Other standards.....	12
2.4 Lipid extraction.....	13
2.5 Lipidomics analysis.....	13
2.6 Data processing.....	14
2.7 Statistical analysis.....	15
3. RESULTS.....	16
3.1 Description of total lipids and pigments from <i>Symbiodinium</i> phylotypes A1, B1 and C1.....	18
3.1.1 Pigments.....	19
3.1.2 Membrane lipids.....	19

3.1.2.1 Glycolipids.....	19
3.1.2.2 Aminolipids.....	20
3.1.2.3 Phospholipids.....	20
3.1.2.4 Sphingolipids.....	20
3.1.2.5 Cholesterol.....	21
3.1.3 Storage lipids.....	21
3.1.4 Other compounds.....	21
3.2 Main distinctions in <i>Symbiodinium</i> phylotypes are revealed in membrane lipids composition.....	22
4. DISCUSSION.....	27
Supporting information.....	32

CHAPTER 3: Concentration of omega 3 fatty acids in membrane lipids determine *Symbiodinium* phylotypes (A1, B1 & C1) fate after temperature stress

1. INTRODUCTION.....	43
2. MATERIALS AND METHODS.....	47
2.1 Experiment design.....	47
2.1 Population growth rates.....	47
2.3 Standards, lipid analysis and data processing.....	48
2.4 Statistical analysis.....	48
3. RESULTS.....	49
3.1. Heat shock experiment revealed two resistant <i>Symbiodinium</i> spp. and drastic changes in lipids and pigments profiles.....	49
3.2 Changes in lipid composition after heat shock.....	50
3.3 Polyunsaturated fatty acids in membrane lipids were pivotal for <i>Symbiodinium</i> spp. long term survival.....	53
3.4 Changes in pigments and oxylipins as evidences of oxidative stress caused by heat shock.....	55

4. DISCUSSION.....	61
4.1. Chloroplast bioenergetics and oxidative stress: a benefit – risk story.....	61
4.2 Ecological implications.....	66
4.2.1 Lipid ecology of zooxanthellae in coral reefs symbiosis and global warming perspectives.....	66
Supporting Information.....	68
FINAL REMARKS.....	69
Importance of phytoplankton lipids in the global carbon budget and bottom-up effects in the marine food chains.....	72
Additional considerations.....	74
FUTURE PERSPECTIVES.....	76
REFERENCES.....	77

CHAPTER 1: GENERAL INTRODUCTION

1.1 Coral reefs and their symbionts in the context of global warming

Coral reefs are one of the most productive ecosystems on Earth, representing one of the most diverse marine environments (Grigg *et al.*, 1984). They hold high biodiversity and ecosystem services important for sustaining higher trophic level organisms and providing goods for large numbers of people (Martinez *et al.*, 2007; Alves de Lima *et al.*, 2010; Knowlton *et al.*, 2010). Corals are defined as ecosystem engineering organisms responsible for controlling the availability of resources to other organisms through physical changes in biotic and abiotic materials (Jones *et al.*, 1994). Therefore, studying them is pivotal to understand marine community structure and production of resources to humankind.

Tropical calcium carbonate reefs are built by stony corals (Scleractinia) associated with dinoflagellates of the family *Symbiodinidaceae* perceived as a mutualistic symbiosis. Photosynthetically produced endosymbiont metabolites are exchanged with coral hosts guaranteeing their survival and growth even in nutrient-poor waters (Muscatine & Porter, 1977; Leggat *et al.*, 2003). In addition, the family *Symbiodinidaceae* is divided into nine clades (A-I) (Pochon & Gates, 2010) and multiple phlotypes within each clade (Thornhill *et al.*, 2014), which shows distinct degrees of host-specificity and different tolerances to environmental conditions (Toller *et al.*, 2001; Baker, 2003; Chen *et al.*, 2003; Coffroth & Santos, 2005; LaJeunesse, 2005; Berkelmans & van Oppen, 2006; Goulet, 2006). Recent research has focused on the strength of this mutualistic relationship in response to predicted climate alteration (Davy & Cook, 2001; Cervino *et al.*, 2004; Barneah *et al.*, 2007; Hoegh-Guldberg *et al.*, 2007), which is the main culprit causing extensive coral reef degradation (Hoegh-Guldberg & Smith, 1989; Graham *et al.* 2008; Wilkinson, 2008).

1.2 Coral bleaching and the oxidative stress theory

Coral bleaching is defined by visible whitening of the coral as a result of decreasing densities of *Symbiodinium* spp. cells and/or declines in their photosynthetic pigments (Fig.1.1). Since corals are extremely dependent on endosymbiont metabolites, their lack leads to coral degradation, eventually followed by coral death (Glynn, 1996; Brown, 1997). A recognized biochemical explanation for the phenomenon was first proposed by Lesser (1997) and coined by Downs *et al.* (2002) as “the oxidative stress theory of coral bleaching”. They proposed that excessive light and temperature cause physiological stress in the symbiont in combination with increased production of reactive oxygen species (ROS). ROS (i.e., hydrogen peroxide, superoxide, hydroxyl and singlet oxygen - Valko *et al.* 2007) can trigger the oxidation of essential biomolecules such as proteins and lipids in both coral and *Symbiodinium* spp. cells, thereby leading to disruption of symbiotic association.



Figure 1.1: Bleached corals of the genus *Mussismilia hispida*, endemic from the tropical waters of Brazil, observed in Ubatuba (São Paulo State north shore) in the summer of 2019.

The concept of ROS playing a key role in bleaching events has been an important subject of research in the last decades. The increased ROS production in symbiont cells may be caused by combination of temperature, light and even other stressors (e.g., alterations in

carbonate chemistry) (Tchernov *et al.*, 2004; Smith *et al.*, 2005; Ragni *et al.*, 2010; Roberty *et al.*, 2015; Goyen, 2017). Today, despite the evidences, the specific causes triggering excessive ROS production leading to bleaching and harmful impacts on host physiology and impairment of symbiosis are still a matter of debate and further investigation.

1.3 The effect of high temperature in the viscosity of membranes leading to free radical and ROS formation in the chloroplast

The membranes of both cells and organelles are universally formed by lipids. Lipid composition directly affects membrane motion and fluidity, which are determined by the size and saturation levels of fatty acids chains of polar lipids. Polyunsaturated fatty acids (PUFA) tend to increase membrane's fluidity, whereas saturated fatty acids (SFA) have the opposite effect. Besides, membrane characteristics can also be altered by abiotic factors, such as temperature. Higher temperatures tend to increase spaces between fatty acids chains increasing membrane fluidity (Fig. 1.2). Therefore, organisms must adjust their membrane composition with variations in abiotic factors in order to maintain cellular functions, a process known as homeoviscous stability control (Sinensky *et al.*, 1974; Cossins *et al.*, 1978; Kellerman *et al.*, 2016).




Figure 1.2: Exemplification of how membrane lipids are affected by fatty acid composition and temperature. Figure shows how polar lipids with unsaturated fatty acids attribute high motion and low viscosity to the membrane. Plus, elevated temperatures also promote higher membrane's motion and fluidity. Combination of elevated temperatures and high polyunsaturated fatty acids can compromise membrane's electron transport chain viability. Source: adapted from bio.libretexts.org.

Controlling membrane composition is pivotal for survival of all life forms, from bacteria to plants to mammals. The lipid composition of energy transducing membranes (i.e., cytoplasmic membranes of bacteria, thylakoid membranes of chloroplasts and mitochondrial inner membranes) is extremely specialized and must perform two essential functions: 1) control permeability of ions such as protons and sodium; and 2) tighten the electron transport at the membrane level. In chloroplasts (Fig. 1.3), the high-energy electrons coming from the water split at photosystem II (PS II) and must be safely transported within membranes by a plastoquinone to energize protein complexes or proton pumps (cytochrome b6f and PSI), a process known as electron transport chain. These pumps generate a proton gradient (high in the lumen and low in the cytoplasm) needed to promote adenosine triphosphate (ATP) synthesis (Fig.1.3). In retrospect, this ion gradient can only be established by controlling permeability of energy transducing membranes. All living forms must always adapt and keep membrane motion and fluidity stable in order to make ATP or energy, thus, lipids and bioenergetics are inseparable.




Figure 1.3: Electron transport chain resulting in ATP synthesis in the thylakoid membranes of chloroplasts. Protein complexes photosystem II (PSII), cytochrome b_6-f (cyt b_6-f) and photosystem I (PSI) are highlighted. Plastoquinone pool responsible for electron transport from PSII to cyt b_6-f is circled in red. Source: Adapted from Wada & Murata, 2009.

Thylakoid membranes of chloroplasts are very unique in that they are composed of glycolipids and specially at protein complexes such as the PS II. They are also decorated with several pigments including carotenoids and chlorophylls (Wada & Murata *et al.*, 2009). The arrangement or conformation of lipids and pigments has been honed by Darwinian evolution to prevent the leakage of both protons and electrons at the membrane level. Any leakage of protons through the membrane may prevent the generation of proton gradients needed for ATP synthesis (e.g., Kellerman *et al.*, 2016; Yoshinaga *et al.*, 2016). On the other hand, leakage of high-energy electrons may lead to free radical generation, which combined with oxygen leads to the formation of ROS (Polle, 1996). Both free radicals and ROS, if not contained by the cell's antioxidant machinery (including carotenoids and antioxidant enzymes such as catalases), can cause oxidation of biomolecules: proteins, pigments and lipids (Augusto & Miyamoto, 2011; Yin *et al.* 2011). Lipid peroxidation is the substitution of a bis-allylic hydrogen in the lipid structure by a free radical, yielding the formation of a lipid radical ($L\cdot$). This lipid radical can easily react with oxygen, generating a lipid peroxy radical ($LOO\cdot$) and undergo complex cyclic reactions that might propagate through the membrane through a process best known as chain reaction (Niki, 2009; Yin *et al.*, 2011). In this context, therefore, fatty acids containing high unsaturation levels, such as polyunsaturated fatty acids (PUFAs),

are the main targets of radical and ROS attack in their double bounds and propagate lipid oxidation even further unless stopped by an antioxidant agent.

The combination of high temperatures and high concentration of PUFAs in the chloroplasts of *Symbiodinium* is likely a strong trigger for coral bleaching, and the associated “oxidative theory of coral bleaching” (Lesser, 1997; Downs *et al.*, 2002). Drastic changes in membrane permeability and fluidity are expected to occur as a consequence of high temperatures. Altering the thylakoid membrane conformation most likely promotes leakage of protons and electrons, thereby leading to decreased energy production and increased oxidative stress, respectively. It is, however, unknown whether the disruption of symbiosis occurs by a decreased supply of metabolites to the coral host, death of symbionts or simply symbionts themselves representing a potential threat to the host due to high ROS production.

1.4 Lipid membrane profiles of symbionts and their fate after thermal stress

Strong evidence suggests that *Symbiodinium* spp. can modulate the lipid composition of cells and organelles membranes in order to keep homeoviscous stability and adapt to environmental alterations (D'amico *et al.*, 2006). Higher abundance of SFAs relative to PUFAs has been reported to enhance physical stability of thylakoid membranes of *Symbiodinium* sp. in response to thermal stress (Tchernov *et al.*, 2004; Bachok *et al.*, 2006; Tolosa *et al.*, 2011). The rationale is that *Symbiodinium* spp. thylakoid membranes are enriched in PUFA, which are highly susceptible to oxidative damage by free radicals and ROS (Wada, 1994; Lesser, 2006; Catalá, 2009). However, *Symbiodinium* spp. may protect the photosynthetic membranes against ROS and thus acquire thermal tolerance altering the ratio saturated/unsaturated fatty acids. Tchernov *et al.* (2004) have even suggested that thermal tolerance is not associated with a single monophyletic phylotype, but rather with the level of saturation of their membrane lipids.

The above-mentioned studies marked the initial investigations of the role of lipids in coral bleaching. They were essential to establish that bulk fatty acid composition is crucial for survival and supports the “oxidative theory of coral bleaching” (Lesser, 1997; Downs *et*

al, 2002). In this dissertation, we generated data based on the global lipidome of some *Symbiodinium* phylotypes (e.g., glycolipids, phospholipids, aminolipids and storage lipids), including their pigments, in response to thermal stress. Thus, we not only report data on fatty acids, but also the lipid molecular species containing these fatty acids. That is, we are able to pinpoint whether thermal stress affects thylakoid membranes by characterizing their specific glycolipids rather than bulk fatty acids derived from other pool of lipids such as the triglycerides or phospholipids. Such detailed and comprehensive lipid analysis could only be achieved by recent analytical developments in mass spectrometry and the advance of lipidomics (Jones *et al.*, 2012; Yao *et al.*, 2015; Nygren *et al.*, 2017).

1.4.1 Lipidomics as tool to better characterize microalgae lipids

Lipidomics is a fairly recent technique that evolved from metabolomics in its own research field (Tomita & Nishioka, 2006; German *et al.*, 2007). Previous lipid analytical techniques allowed qualitative information about polar lipids such as acquired by thin-layer chromatography or quantitative analysis of bulk fatty acids by gas chromatography. Contrasting with past lipid analytical techniques, lipid characterization by liquid chromatography coupled to mass spectrometry (LC-MS) enabled precise characterization and quantification of every lipid molecular species present in a given sample (German *et al.*, 2007; Oresic *et al.*, 2008), including molecules that are specific markers of chloroplast, such as glycolipids and plastoquinones. For example, a great diversity of glyco and amino membrane lipids in *Symbiodinium* spp. and other dinoflagellates has been described by LC-MS analysis, including few alterations when growing in distinct temperatures (Leblond *et al.*, 2000, 2006, 2010, 2015; Gray *et al.*, 2009; Dahmen *et al.*, 2013; Flaim *et al.*, 2014; Anesi *et al.*, 2015, 2016). These studies were mostly focused on a specific class of polar lipid such as glycolipids or aminolipids, and not aimed at characterizing the global lipidome together with specialized lipids, such as plastoquinone and pigments using LC-MS. Besides the culture-based investigations, important data have been generated in environmental studies reported by Van Mooy *et al.* (2006, 2009 and 2010), Moutin *et al.* (2007), Schubotz *et al.* (2009), Xie *et al.* (2014) and Becker *et al.* (2018). These include not only data from phospho, glyco and

aminolipids, but also storage lipids and quinone molecules (e.g., ubiquinone in Becker *et al.*, 2018). These studies describe the lipidome of the water column of the oceans, where a diverse variety of phytoplankton occurs, in response to diel temperature oscillations and distinct nutrient conditions. The present study is, to the best of our knowledge, the first attempt to characterize the global lipidome of a microalga, describing not only membrane lipids, but also storage lipids, pigments and plastoquinone from *Symbiodinium* phylotypes. This information will be linked to cell physiology data to further knowledge on their thermal sensitivity.

2. GOALS AND SCOPE OF THIS THESIS

The overall goal of this thesis is to investigate how the lipidomes of *S. microadriaticum* (phylotype A1), *S. minutum* (phylotype B1) and *S. goreau* (phylotype C1) most prevalently associated with scleractinian corals are related to their thermal sensitivity. Description of *Symbiodinium* spp. lipids and pigments is very scarcely found in the literature, let alone their lipid alterations with stress events. For best comprehension all studied *Symbiodinium* spp. will be referred as their phylotypes A1, B1 and C1. Therefore, this thesis is divided into two further chapters:

Chapter 2:

Description and quantification of global lipidome and pigment profiles of *Symbiodinium* phylotypes A1, B1 and C1 growing under optimum conditions of temperature, light and nutrients. Here, the goals are to examine whether differences in lipidome and pigments profiles between *Symbiodinium* phylotypes can predict their thermal tolerance to heat stress events based on previous studies (e.g., Tchernov *et al.*, 2004).

Chapter 3:

Monitor *Symbiodinium* phylotypes (A1, B1 and C1) growth rates and lipidome/pigment alterations after a heat shock event (4h under 34°C) describing short-term (4 and 24 h) and long-term phenotypical alterations (10 days).

Main assumption: “High temperatures known to change thylakoid membrane stability and enabling scape of high energy electrons generate higher concentration of ROS leading to lipid peroxidation accompanied by reduction of ATP production. Both facts affect *Symbiodinium* spp. growth rates and reflect in drastic changes in their lipidomes and pigments profiles”.

Specific hypotheses:

- 1) *Symbiodinium* spp. growth rates are negatively affected by heat shock;
- 2) Changes in lipids and pigments are different between *Symbiodinium* phylotypes after heat shock;
- 3) Lipidome and pigments profiles of *Symbiodinium* phylotypes after stress are good indicators of oxidative stress caused by heat shock experiment;
- 4) Lipid peroxidation preferably occurs in polyunsaturated fatty acids (PUFA) from chloroplast membranes.

FINAL REMARKS

In Chapter 2 we verified the use of lipidomics as a precise tool for description, quantification and monitoring of total lipids and pigments of *Symbiodinium* phylotypes A1, B1 and C1 growing under optimum conditions of temperature, light and nutrients. We also showed that statistically significant differences among phylotypes were mainly determined by their membrane lipids. Although A1 and B1 were more similar based in the heatmap analysis, A1 was more alike to C1 considering their higher concentrations of omega-3 polyunsaturated fatty acids and plastoquinone.

In Chapter 3 we analyzed variations in *Symbiodinium* growth rates and both short and long terms lipidomes after a heat shock event summarized in Figure F1. Phylotype B1 was not heat shock resistant and its high decrease in biomass and cell death occurred after a downregulation of essential membrane omega-3 fatty acids and all identified pigments from chloroplasts. Thus, lipidome and pigments changes of B1 could not guarantee its cell's energetic requirements. However, phylotypes A1 and C1 both resisted mainly because they were capable of maintaining higher concentrations of essential omega-3 fatty acids in the thylakoid membranes and supply cells energetic requirements in the long term.

Combined information from both chapters demonstrated lipidomics as a functional tool to comprehend cell physiological alterations caused by thermal stress by a unified concept. We noticed that functional thylakoid membrane lipid structure cannot vary much in order to feasibly supply the cell bioenergetic demands. Therefore, the fate of organisms is likely to be determined by the cell antioxidant machinery, which were not analyzed in the present study, but that might protect membranes against ROS and keep efficient energy output even under stress conditions.




Figure F1: Summary of *Symbiodinium* phylotypes cell's proposed mechanisms caused by upregulation of ROS. Temperature alters membrane viscosity and enable scape of high energy electrons from electron transport chain. This might be responsible for upregulation of ROS that will concomitantly lead to the peroxidation of most vulnerable omega-3 polyunsaturated fatty acids and activate quenching mechanisms. We propose that phylotype B1 did not survive because it could not avoid lipid peroxidation of essential $\omega-3$ in the membranes, neither synthesize *de novo* epoxidized lipids. On the other hand, it was not true for phylotypes A1 and C1. Both kept efficient energy output, although we suggest that they followed distinct strategies already discussed in chapter 3. Protein complexes photosystem II (PSII), cytochrome b_6-f (cyt b_6-f) and photosystem I (PSI) were highlighted. Plastoquinone pool responsible for electron transport from PSII to cyt b_6-f was also highlighted in red. Scape of high energy electron leading to ROS was highlighted in dark red. Source: Adapted from Wada & Murata, 2009.

Finally, we attempted to elucidate specific hypothesis delineated in this dissertation/thesis:

1) "*Symbiodinium spp.* growth rates are negatively affected by heat shock"

Growth rates were all reduced after heat shock. Besides, response was different between phylotypes. *S. minutum* (B1) did not survive stress, whereas *S. microadriaticum* (A1) and *S. goreau* (C1) were resistant to heat shock but decreased their growth rates.

2) *“Changes in lipids and pigments are different between Symbiodinium phylotypes after heat shock”*

Symbiodinium spp. lipidomes and pigments profiles were differently altered by heat shock. However, downregulation of n-3 fatty acids 4 hours after stress was common in all phylotypes. After 24 hours and in the long term considered as 10 days, each phylotype had a specific survival strategy and fate summarized in figure 4. This demonstrates the essential role of omega-3 fatty acids for cellular energy as suggested by Valentine and Valentine (2004).

3) *“Lipidome and pigments profiles of Symbiodinium spp. after stress are good indicators of oxidative stress caused by heat shock experiment”*

We considered variations in the pigments ratios as indicators of chloroplast “health status” and chlorophyll decrease strongly indicated damage in the chloroplasts caused by heat shock. This data together with variations in epoxy/de-epoxy xanthophylls suggest that damage was likely caused by oxidative stress but could not prove it. However, oxidized fatty acids in the free form and connected with polar lipids analytically proved damage provoked by oxidative stress.

4) *“Lipid peroxidation preferably occurs in polyunsaturated fatty acids (PUFAs) from chloroplast membranes”*

Lipid peroxidation was evidenced by the presence of oxidized fatty acids mostly derived from omega 3 polyunsaturated fatty acids in free (FFA) and membrane associated (PL) forms of MGDG, DGDG, DGCC and PC. Glycolipids (MGDG and DGDG) are well known structural thylakoid membrane lipids, plus, evidences discussed in chapter 3 strongly indicated DGCC presence also in the chloroplast. Therefore, chloroplast membranes were main targets of lipid peroxidation. Plus, significant changes in DGCC and DGDG compounds suggest that they might be specifically located closer to chloroplast ROS

formation sources when compared to MGDG because they did not change significantly after heat shock.

Importance of phytoplankton lipids in the global carbon budget and bottom-up effects in the marine food chains

Symbiodinium sp. takes part of the Dinophyceae class and many lipid compounds characterized and monitored in our work have been previously reported in the phenotypes of relatives from the same class and other microalgal classes that inhabit oligotrophic ocean gyres (e.g., Cryptophyceae, Haptophyceae) (Van Mooy *et al.*, 2009, 2010; Shemi *et al.*, 2016). We suggest our temperature-related alterations in the lipidome of *Symbiodinium* spp. are similar to those that may occur in phytoplankton communities considering a universal biochemical principle of lipid composition in bioactive membranes.

Becker *et al.* (2018) showed that in the North Pacific subtropical gyre (NPSG) nearly half of the relative abundance of organisms was composed of Dinophyceae and Haptophyceae (ca. 25.3% each). This region comprises 40% of Earth's total surface area, representing the world's largest biome (Emmerson *et al.*, 1997; Sarmiento *et al.*, 2004). Local and global alterations in phytoplankton communities were noticed with elevated temperatures and other climate change stressors in the past years leading to alterations in global primary production and carbon sinking (Sarmiento *et al.*, 2004; Behrenfeld *et al.*, 2006; Schmittner *et al.*, 2008; Nagelkerken & Connell, 2010), but the physiological mechanisms responsible for such alterations were poorly discussed. We suggest that changes are likely to happen because of phytoplankton photosynthetic structure vulnerability to oxidative stress considering that they are the major DHA producers in the biosphere (Valentine, 2009). High concentrations of DHA were reported by Becker *et al.* (2018) in the NPSG community associated with DGCC betaine polar head. Their most abundant DGCC had C38:6 – 800.6035 m/z - acyl chains of palmitic acid (16:0) and docosahexaenoic acid (DHA), which was also the most abundant feature in our samples. Photosynthesis and phytoplankton growth in phenotypes with high DHA concentration can be impaired through mechanisms already discussed in our work.

Furthermore, the high abundance of DHA in *Symbiodinium* phylotypes and in other flagellate microalgae (Leblond *et al.*, 2000, 2006, 2015; Gray *et al.*, 2009; Awai *et al.*, 2012; Armada *et al.*, 2013; Dahmen *et al.*, 2013; Anesi & Guella, 2015; Anesi *et al.*, 2016) sustains the idea proposed by Valentine (2009): phytoplankton represents the global stock of DHA production in the marine ecosystem. Life cycle of zooplankton species depend on the nutritional quality of phytoplankton, which is defined by lipid content (Søreide *et al.*, 2010). Ingestion of DHA and other omega 3 enables development of their sensorial mechanisms essential for survival, growth and reproduction (Müller-Navarra *et al.*, 2000; Falkowski & Oliver, 2007). Higher in the food chains, other organisms also evolved with the same dependence in n-3 and their populations may either decrease or present individuals with neurological and sensorial deficiencies if n-3 consumption is low (Davis *et al.*, 1992; Budge *et al.*, 2001; Jonasdottir *et al.*, 2002). Consequently, we highlight the importance of DHA ingestion in the marine food chains and how they are also likely to be impacted by global warming. Behrenfeld *et al.* (2006) showed that from 1999 to 2004 the average global primary productivity dropped by about 200 tons a year. Local changes had a decrease as high as 50% (see also Bopp *et al.*, 2013). If ocean temperatures keep increasing progressively as predicted by climate models (Hansen *et al.*, 2010; Rogelj *et al.*, 2012; Cabré *et al.*, 2015), omega 3 producing phytoplankton might follow an opposite way and promote a bottom-up effect in all marine food chains that could potentially lead to ecological collapse of the whole ecosystem

Omega-3 presence in microalgae membranes mediate cell death cascades with temperature changes in the environment (Valentine, 2009). It was highly evidenced in our experiments with *Symbiodinium* spp. and generated valuable data for coral reef symbiosis and bleaching research. Besides, the concomitant omega-3 presence in phytoplankton populations with large distribution patterns enhances our insights of large detrimental effects in marine food chains following ocean warming. Therefore, we highlight the importance of the present study in mechanistically explaining a universal biochemical principle for all living creatures, from algae cells to more complex organisms. A principle useful to understanding cell physiology and also how slight modifications can impact the whole environment.

REFERENCES

- Alves de Lima, L., Migliolo, L., Barreiro e Castro, C., de Oliveira Pires, D., López-Abarrategui, C., Ferreira Goncalves, E., ... & Campos Dias, S. (2013). Identification of a novel antimicrobial peptide from Brazilian coast coral *Phyllogorgia dilatata*. *Protein and peptide letters*, 20(10), 1153-1158.
- Andras, J. P., Kirk, N. L., & Drew Harvell, C. (2011). Range-wide population genetic structure of *Symbiodinium* associated with the Caribbean Sea fan coral, *Gorgonia ventalina*. *Molecular Ecology*, 20(12), 2525-2542.
- Anesi, A., & Guella, G. (2015). A fast liquid chromatography-mass spectrometry methodology for membrane lipid profiling through hydrophilic interaction liquid chromatography. *Journal of chromatography A*, 1384, 44-52.
- Anesi, A., Obertegger, U., Hansen, G., Sukenik, A., Flaim, G., & Guella, G. (2016). Comparative analysis of membrane lipids in psychrophilic and mesophilic freshwater dinoflagellates. *Frontiers in plant science*, 7, 524.
- Anthony KRN, Hoogenboom MO, Maynard JA, Grottoli AG, Middlebrook R (2009) Energetics approach to predicting mortality risk from environmental stress: a case study of coral bleaching. *Funct Ecol* 23:539-550.
- Armada, I., Hachero-Cruzado, I., Mazuelos, N., Ríos, J. L., Manchado, M., & Cañavate, J. P. (2013). Differences in betaine lipids and fatty acids between *Pseudoisochrysis paradoxa* VLP and *Diacronema vlkianum* VLP isolates (Haptophyta). *Phytochemistry*, 95, 224-233.
- Augusto, O., & Miyamoto, S. (2011). Oxygen radicals and related species. *Principles of free radical biomedicine*, 1, 19-42.
- Awai, K., Matsuoka, R., & Shioi, Y. (2012). Lipid and fatty acid compositions of *Symbiodinium* phylotypes. In *Proceedings of the 12th International Coral Reef Symposium*.
- Bachok, Z., Mfilinge, P., & Tsuchiya, M. (2006). Characterization of fatty acid composition in healthy and bleached corals from Okinawa, Japan. *Coral Reefs*, 25(4), 545-554.
- Barneah O. Brickner, M. Hooge (2007) Three party symbiosis: acoelomorph worms, corals and unicellular algal symbiont in Eilat (Red Sea). *Marine Biology* 151: 1215

- Baker A. (2003) Flexibility and specificity in coral-algal symbiosis: diversity, ecology, and biogeography of Symbiodinium. *Annual Review Ecology Evolution Systems*, 34: 661-689
- Bastien, O., Botella, C., Chevalier, F., Block, M. A., Jouhet, J., Breton, C., ... & Maréchal, E. (2016). New insights on thylakoid biogenesis in plant cells. In *International review of cell and molecular biology* (Vol. 323, pp. 1-30). Academic Press.
- Bates, P. D., Fatihi, A., Snapp, A. R., Carlsson, A. S., & Lu, C. (2012). Acyl editing and headgroup exchange are the major mechanisms that direct polyunsaturated fatty acid flux into triacylglycerols. *Plant physiology*, pp-112.
- Babiychuk E, Müller F, Eubel H, Braun HP, Frentzen M AND Kushnir S (2003) Arabidopsis phosphatidylglycerophosphate synthase 1 is essential for chloroplast differentiation but is dispensable for mitochondrial function. *Plant J* 33: 899–909
- Becker, K. W., Collins, J. R., Durham, B. P., Groussman, R. D., White, A. E., Fredricks, H. F., ... & Van Mooy, B. A. (2018). Daily changes in phytoplankton lipidomes reveal mechanisms of energy storage in the open ocean. *Nature communications*, 9(1), 5179.
- Behrenfeld, M. J., O'Malley, R. T., Siegel, D. A., McClain, C. R., Sarmiento, J. L., Feldman, G. C., ... & Boss, E. S. (2006). Climate-driven trends in contemporary ocean productivity. *Nature*, 444(7120), 752.
- Berkelmanns, R. & Van Oppen, M. J. H. (2006) Flexible partners in coral symbiosis: a 'nugget of hope' for coral reefs in an era of climate change. *Proc. R. Soc. B* 273, 2305–2312
- Bligh, E. G., & Dyer, W. J. (1959). Canadian journal of biochemistry and physiology. A rapid method of lipid extraction and purification, 37, 911-917.
- Block, M. A., Dorne, A. J., Joyard, J., & Douce, R. (1983). Preparation and characterization of membrane fractions enriched in outer and inner envelope membranes from spinach chloroplasts. II. Biochemical characterization. *Journal of Biological Chemistry*, 258(21), 13281-13286.
- Bopp, L., Resplandy, L., Orr, J. C., Doney, S. C., Dunne, J. P., Gehlen, M., ... & Tjiputra, J. (2013). Multiple stressors of ocean ecosystems in the 21st century: projections with CMIP5 models. *Biogeosciences*, 10, 6225-6245.
- Boudière, L., Botté, C. Y., Saidani, N., Lajoie, M., Marion, J., Brehelin, L., ... & Bastien, O. (2012). Galvestine-1, a novel chemical probe for the study of the glycerolipid homeostasis system in plant cells. *Molecular BioSystems*, 8(8), 2023-2035.
- Boudière, L., Michaud, M., Petroustos, D., Rébeillé, F., Falconet, D., Bastien, O., ... & Block, M. A. (2014). Glycerolipids in photosynthesis: composition, synthesis and trafficking. *Biochimica et Biophysica Acta (BBA)-Bioenergetics*, 1837(4), 470-480.g

Brown BE (1997) Coral bleaching: causes and consequences. *Coral Reefs* 16: S129–S138

Budge, S. M., Parrish, C. C., & McKenzie, C. H. (2001). Fatty acid composition of phytoplankton, settling particulate matter and sediments at a sheltered bivalve aquaculture site. *Marine Chemistry*, 76(4), 285-303.

Budin, I., de Rond, T., Chen, Y., Chan, L. J. G., Petzold, C. J., & Keasling, J. D. (2018). Viscous control of cellular respiration by membrane lipid composition. *Science*, 362(6419), 1186-1189.

Cabr e, A., Marinov, I., & Leung, S. (2015). Consistent global responses of marine ecosystems to future climate change across the IPCC AR5 earth system models. *Climate Dynamics*, 45(5-6), 1253-1280.

Ca avate, J. P., Armada, I., R os, J. L., & Hachero-Cruzado, I. (2016). Exploring occurrence and molecular diversity of betaine lipids across taxonomy of marine microalgae. *Phytochemistry*, 124, 68-78.

Cardol, P., Forti, G., & Finazzi G. (2011). Regulation of electron transport in microalgae. *Biochimica et Biophysica Acta (BBA)-Bioenergetics*, 1807(8), 912-918.

Catal a A (2009) Lipid peroxidation of membrane phospholipids generates hydroxy-alkenals and oxidized phospholipids active in physiological and/or pathological conditions. *Chemical Physical Lipids* 157:1–11

Cervino, J.M., Hayes, R., Goreau, T.J., Smith, G.W. (2004) Zooxanthellae regulation in yellow blotch/band and other coral diseases contrasted with temperature related bleaching: in situ destruction vs expulsion. *Symbiosis* 37: 63–85

Chen C.A., Lam K.K., Nakano Y., Tsai W.S. (2003) Stable association of stresstolerant zooxanthellae, Symbiodinium clade D, with the low-temperature tolerant coral *Oulastrea crispata*, (Scleractinia: Faviidae) in subtropical non-reefal coral communities. *Zoology Studies* 42: 540-550

Coffroth M.A., Lewis C., Santos S., Weaver J. (2006) Environmental populations of symbiotic dinoflagellates in the genus *Symbiodinium* can initiate symbioses with reef cnidarians. *Current Biology* 16: 985–987

Cooper, T. F., Lai, M., Ulstrup, K. E., Saunders, S. M., Flematti, G. R., Radford, B., & van Oppen, M. J. (2011). Symbiodinium genotypic and environmental controls on lipids in reef building corals. *PLoS One*, 6(5), e20434.

Cossins, A. R., & Prosser, C. L. (1978). Evolutionary adaptation of membranes to temperature. *Proceedings of the National Academy of Sciences*, 75(4), 2040-2043.

D'Amico, S., Collins, T., Marx, J. C., Feller, G., & Gerday, C. (2006). Psychrophilic microorganisms: challenges for life. *EMBO reports*, 7(4), 385-389.

D Lambreva, M., Russo, D., Polticelli, F., Scognamiglio, V., Antonacci, A., Zobnina, V., ... & Rea, G. (2014). Structure/function/dynamics of photosystem II plastoquinone binding sites. *Current Protein and Peptide Science*, 15(4), 285-295.

Da Costa, E., Silva, J., Mendonça, S., Abreu, M., & Domingues, M. (2016). Lipidomic approaches towards deciphering glycolipids from microalgae as a reservoir of bioactive lipids. *Marine drugs*, 14(5), 101.

Dahmen, J. L., Khadka, M., Dodson, V. J., & Leblond, J. D. (2013). Mono-and digalactosyldiacylglycerol composition of dinoflagellates. VI. Biochemical and genomic comparison of galactolipid biosynthesis between *Chromera velia* (Chromerida), a photosynthetic alveolate with red algal plastid ancestry, and the dinoflagellate, *Lingulodinium polyedrum*. *European journal of phycology*, 48(3), 268-277.

Davis, M. W., & Olla, B. L. (1992). Comparison of growth, behavior and lipid concentrations of walleye pollock *Theragra chalcogramma* larvae fed lipid-enriched, lipid-deficient and field-collected prey. *Marine ecology progress series. Oldendorf*, 90(1), 23-30.

Daum G & Vance JE (1997) Import of lipids into mitochondria. *Prog Lipid Res* 36: 103–130

Davy S.V.K., Cook C.V. (2001) The influence of 'host release factor' on carbon release by zooxanthellae isolated from fed and starved *Aiptasia pallida*. *Comparative Biochemistry and Physiology Part A* 129: 487-494.

Demmig-Adams, B., & Adams III, W. W. (1996). *The role of xan-thophyll cycle carotenoids in the protection of photosyn*(Doctoral dissertation, thesis. *Trends in Plant Science* 1, 21-6).

Díaz-Almeyda, E., Thomé, P. E., El Hafidi, M., & Iglesias-Prieto, R. (2011). Differential stability of photosynthetic membranes and fatty acid composition at elevated temperature in *Symbiodinium*. *Coral Reefs*, 30(1), 217-225.

Díaz-Almeyda, E. M., Prada, C., Ohdera, A. H., Moran, H., Civitello, D. J., Iglesias-Prieto, R., & Medina, M. (2017). Intraspecific and interspecific variation in thermotolerance and photoacclimation in *Symbiodinium* dinoflagellates. *Proceedings of the Royal Society B: Biological Sciences*, 284(1868), 20171767.

Dietz, K. J., Turkan, I., & Krieger-Liszkay, A. (2016). Redox-and reactive oxygen species-dependent signaling into and out of the photosynthesizing chloroplast. *Plant Physiology*, 171(3), 1541-1550.

Downs CA, Fauth JE, Halas JC, Dustan P, Bemiss J, et al. (2002) Oxidative stress and seasonal coral bleaching. *Free Radical Biological Medicine* 33: 533–543.

Eichenberger, W., & Gribi, C. (1997). Lipids of *Pavlova lutheri*: cellular site and metabolic role of DGCC. *Phytochemistry*, 45(8), 1561-1567.

Emerson, S., Quay, P., Karl, D., Winn, C., Tupas, L., & Landry, M. (1997). Experimental determination of the organic carbon flux from open-ocean surface waters. *Nature*, 389(6654), 951.

Evans, T. W., Wörmer, L., Lever, M. A., Lipp, J. S., Lagostina, L., Lin, Y. S., ... & Hinrichs, K. U. (2017). Size and composition of subseafloor microbial community in the Benguela upwelling area examined from intact membrane lipid and DNA analysis. *Organic Geochemistry*, 111, 86-100.

Falkowski, P. G., & Oliver, M. J. (2007). Mix and match: how climate selects phytoplankton. *Nature reviews microbiology*, 5(10), 813.

Flaim, G., Obertegger, U., Anesi, A., & Guella, G. (2014). Temperature-induced changes in lipid biomarkers and mycosporine-like amino acids in the psychrophilic dinoflagellate *P. eridinium aciculiferum*. *Freshwater biology*, 59(5), 985-997.

Fitt, W. K., Brown, B. E., Warner, M. E., & Dunne, R. P. (2001). Coral bleaching: interpretation of thermal tolerance limits and thermal thresholds in tropical corals. *Coral reefs*, 20(1), 51-65.

Gates RD, Baghdasarian G, Muscatine L (1992) Temperature stress causes host cell detachment in symbiotic cnidarians: implications for coral bleaching. *Biological Bull* 182: 324–332

German, J. B., Gillies, L. A., Smilowitz, J. T., Zivkovic, A. M., & Watkins, S. M. (2007). Lipidomics and lipid profiling in metabolomics. *Current opinion in lipidology*, 18(1), 66-71.

Glynn, P. W. 1991. Coral reef bleaching in the 1980s and possible connections with global warming. *Trends Ecol. Evol.* 6: 175-179

Glynn, P. W. 1996. Coral reef bleaching: facts, hypotheses and implications. *Glob Change Biol* 2:495-509. Glynn, P. W. 1996. Coral reef bleaching: facts, hypotheses and implications. *Glob Change Biol* 2:495-509.

Goss R, Lohr M, Latowski D, Grzyb J, Vleler A, Wilhelm C & Strzalka K (2005) Role of hexagonal structure-forming lipids in diadinoxanthin and violaxanthin solubilization and de-epoxidation. *Biochemistry* 44: 4028–4036

Goulet, T. L., Simmons, C., & Goulet, D. (2008). Worldwide biogeography of *Symbiodinium* in tropical octocorals. *Marine Ecology Progress Series*, 355, 45-58.

Gordon, B. R., Martin, D. E., Bamberg, K. R., & Motti, C. A. (2018). Chemical imaging of a Symbiodinium sp. cell using synchrotron infrared microspectroscopy: a feasibility study. *Journal of microscopy*, 270(1), 83-91.

Goyen, S., Pernice, M., Szabó, M., Warner, M. E., Ralph, P. J., & Suggett, D. J. (2017). A molecular physiology basis for functional diversity of hydrogen peroxide production amongst Symbiodinium spp. (Dinophyceae). *Marine Biology*, 164(3), 46.

Graham, N.A.J., McClanahan, T.R., Macneil, M.A., Wilson, S.K., Polunin, N.V.C., Jennings, S. et al. (2008) Climate warming, marine protected areas and the ocean-scale integrity of coral reef ecosystems. *PLoS ONE* 3: e3039.

Gray, C. G., Lasiter, A. D., Li, C., & Leblond, J. D. (2009). Mono- and digalactosyldiacylglycerol composition of dinoflagellates. I. Peridinin-containing taxa. *European Journal of Phycology*, 44(2), 191-197.

Grottoli, A. G., Rodrigues, L. J., & Juarez, C. (2004). Lipids and stable carbon isotopes in two species of Hawaiian corals, *Porites compressa* and *Montipora verrucosa*, following a bleaching event. *Marine Biology*, 145(3), 621-631.

Grottoli AG, Rodrigues LJ, Palardy JE (2006) Heterotrophic plasticity and resilience in bleached corals. *Nature* 440: 1186-1189.

Grigg, R. W., Polovina, J. J., & Atkinson, M. J. (1984) Model of a coral reef ecosystem. *Coral Reefs*, 3(1), 23-27.

Guillard, R. (1975) *Culture of phytoplankton for feeding marine invertebrates*. Culture of Marine Invertebrate Animals, Springer, New York, p. 29–60.

Gunderson, A. R., Armstrong, E. J., & Stillman, J. H. (2016). Multiple stressors in a changing world: the need for an improved perspective on physiological responses to the dynamic marine environment. *Annual Review of Marine Science*, 8, 357-378.

Gurr, M.I., Harwood, J.L., and Frayn, K.N. (2002) *Lipid Biochemistry. An Introduction*, 5th ed. Blackwell, Oxford, 320 pp.

Guschina, I. A., & Harwood, J. L. (2006) Lipids and lipid metabolism in eukaryotic algae. *Progress in lipid research*, 45(2), 160-186.

Guschina, I. A., & Harwood, J. L. (2009) Algal lipids and effect of the environment on their biochemistry. In *Lipids in aquatic ecosystems* (pp. 1-24). Springer, New York, NY.

Hansen, J., Ruedy, R., Sato, M., & Lo, K. (2010) Global surface temperature change. *Reviews of Geophysics*, 48(4).

Harland, A. D., Fixter, L. M., Davies, P. S., & Anderson, R. A. (1991) Distribution of lipids between the zooxanthellae and animal compartment in the symbiotic sea

anemone *Anemonia viridis*: Wax esters, triglycerides and fatty acids. *Marine Biology*, 110(1), 13-19.

Harland, A. D., Navarro, J. C., Davies, P. S., & Fixter, L. M. (1993). Lipids of some Caribbean and Red Sea corals: total lipid, wax esters, triglycerides and fatty acids. *Marine Biology*, 117(1), 113-117.

Havaux, M., & Tardy, F. (1996). Temperature-dependent adjustment of the thermal stability of photosystem II in vivo: possible involvement of xanthophyll-cycle pigments. *Planta*, 198(3), 324-333.

Havaux, M. (2014). Carotenoid oxidation products as stress signals in plants. *The Plant Journal*, 79(4), 597-606.

Hazel, J. R. (1995). Thermal adaptation in biological membranes: is homeoviscous adaptation the explanation? *Annual review of physiology*, 57(1), 19-42.

Hendry, George AF, Jennifer D. Houghton, and Stanley B. Brown. "The degradation of chlorophyll—a biological enigma." *New Phytologist* 107.2 (1987): 255-302.

Hoegh-Guldberg O, Smith GJ (1989) The effect of sudden changes in temperature, light and salinity on the population density and export of zooxanthellae from the reef corals *Stylophora pistillata* Esper and *Seriatopora hystrix* Dana. *J Exp Marine Biology Ecology* 129: 279–303.

Hoegh-Guldberg, O. AND Bruno, J.F. (2010) The impact of climate change on the world's marine ecosystems. *Science* 328: 1523–1528

Hu Q, Sommerfeld M, Jarvis E, Ghirardi M, Posewitz M, Seibert M & Darzins A (2008) Microalgal triacylglycerols as feedstocks for biofuel production: perspectives and advances. *Plant J* 54: 621–639

Holzwarth AR, Miloslavina Y, Nilkens M, Jahns P (2009) Identification of two quenching sites active in the regulation of photosynthetic light-harvesting studied by time-resolved fluorescence. *Chem Phys Lett* 483:262–267

Hughes, T. P., Kerry, J. T., Álvarez-Noriega, M., Álvarez-Romero, J. G., Anderson, K. D., Baird, A. H., ... & Bridge, T. C. (2017). Global warming and recurrent mass bleaching of corals. *Nature*, 543(7645), 373-377.

Imbs, A. B., Demidkova, D. A., Latypov, Y. Y., & Pham, L. Q. (2007). Application of fatty acids for chemotaxonomy of reef-building corals. *Lipids*, 42(11), 1035-1046.

Imbs, A. B., & Yakovleva, I. M. (2012). Dynamics of lipid and fatty acid composition of shallow-water corals under thermal stress: an experimental approach. *Coral Reefs*, 31(1), 41-53.

- Jahns, P., & Holzwarth, A. R. (2012). The role of the xanthophyll cycle and of lutein in photoprotection of photosystem II. *Biochimica et Biophysica Acta (BBA)-Bioenergetics*, 1817(1), 182-193
- Jonasdottir, S., Gudfinnsson, H., Gislason, A., & Astthorsson, O. (2002). Diet composition and quality for *Calanus finmarchicus* egg production and hatching success off south-west Iceland. *Marine Biology*, 140(6), 1195-1206.
- Jones, C. G., Lawton, J. H. & Shachak, M. 1994. Organisms as ecosystem engineers. *Oikos* 69:373-86.
- Jones M. R. (2007) Lipids in photosynthetic reaction centres: structural roles and functional holes. *Progress in Lipid Research* 46: 56–87
- Jones, J., Manning, S., Montoya, M., Keller, K., & Poenie, M. (2012). Extraction of algal lipids and their analysis by HPLC and mass spectrometry. *Journal of the American Oil Chemists' Society*, 89(8), 1371-1381.
- Kato, M., Sakai, M., Adachi, K., Ikemoto, H., & Sano, H. (1996). Distribution of betaine lipids in marine algae. *Phytochemistry*, 42(5), 1341-1345.
- Kellermann, M. Y., Yoshinaga, M. Y., Valentine, R. C., Wörmer, L., & Valentine, D. L. (2016). Important roles for membrane lipids in haloarchaeal bioenergetics. *Biochimica et Biophysica Acta (BBA)-Biomembranes*, 1858(11), 2940-2956.
- Khozin-Goldberg, I., & Cohen, Z. (2006). The effect of phosphate starvation on the lipid and fatty acid composition of the fresh water eustigmatophyte *Monodus subterraneus*. *Phytochemistry*, 67(7), 696-701.
- Klueter, A., Crandall, J. B., Frederick, I., Archer, F. I., Teece, M. A., & Coffroth, M.A. (2015). Taxonomic and environmental variation of metabolite profiles in marine dinoflagellates of the genus *Symbiodinium*. *Metabolites* 5, 74–99. doi: 10.3390/metabo5010074
- Kneeland, J., Hughen, K., Cervino, J., Hauff, B., & Eglinton, T. (2013). Lipid biomarkers in *Symbiodinium* dinoflagellates: new indicators of thermal stress. *Coral Reefs*, 32(4), 923-934.
- Knowlton, N., Brainard, R. E., Fisher, R., Moews, M., Plaisance, L., & Caley, M. J. (2010). Coral reef biodiversity. *Life in the world's oceans: diversity distribution and abundance*, 65-74.
- Krinsky, N. I. (1989). Antioxidant functions of carotenoids. *Free Radical Biology and Medicine*, 7(6), 617-635

Krueger, T., Becker, S., Pontasch, S., Dove, S., Hoegh-Guldberg, O., Leggat, W., ... & Davy, S. K. (2014). Antioxidant plasticity and thermal sensitivity in four types of *Symbiodinium* sp. *Journal of phycology*, 50(6), 1035-1047.

Krueger, T., Fisher, P. L., Becker, S., Pontasch, S., Dove, S., Hoegh-Guldberg, O., ... & Davy, S. K. (2015a). Transcriptomic characterization of the enzymatic antioxidants FeSOD, MnSOD, APX and KatG in the dinoflagellate genus *Symbiodinium*. *BMC evolutionary biology*, 15(1), 48.

Krueger, T., Hawkins, T. D., Becker, S., Pontasch, S., Dove, S., Hoegh-Guldberg, O., ... & Davy, S. K. (2015b). Differential coral bleaching—Contrasting the activity and response of enzymatic antioxidants in symbiotic partners under thermal stress. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, 190, 15-25.

Künzler, K., & Eichenberger, W. (1997). Betaine lipids and zwitterionic phospholipids in plants and fungi. *Phytochemistry*, 46(5), 883-892.

LaJeunesse, T. (2002). Diversity and community structure of symbiotic dinoflagellates from Caribbean coral reefs. *Marine Biology*, 141(2), 387-400.

LaJeunesse, T.C. (2005) 'Species' radiations of symbiotic dinoflagellates in the Atlantic and Indo-Pacific since the Miocene-Pliocene transition. *Molecular Biology Evolution* 22: 570-581.

LaJeunesse, T. C., Parkinson, J. E., Gabrielson, P. W., Jeong, H. J., Reimer, J. D., Voolstra, C. R., & Santos, S. R. (2018). Systematic revision of Symbiodiniaceae highlights the antiquity and diversity of coral endosymbionts. *Current Biology*, 28(16), 2570-2580.

Leblond, J. D., & Chapman, P. J. (2000). Lipid class distribution of highly unsaturated long chain fatty acids in marine dinoflagellates. *Journal of Phycology*, 36(6), 1103-1108.

Leblond, J. D., Anderson, B., Kofink, D., Logares, R., Rengefors, K., & Kremp, A. (2006). Fatty acid and sterol composition of two evolutionarily closely related dinoflagellate morphospecies from cold Scandinavian brackish and freshwaters. *European Journal of Phycology*, 41(3), 303-311.

Leblond, J. D., Dahmen, J. L., & Evens, T. J. (2010). Mono- and digalactosyldiacylglycerol composition of dinoflagellates. IV. Temperature-induced modulation of fatty acid regiochemistry as observed by electrospray ionization/mass spectrometry. *European Journal of Phycology*, 45(1), 13-18.

Leblond, J. D., Khadka, M., Duong, L., & Dahmen, J. L. (2015). Squishy lipids: Temperature effects on the betaine and galactolipid profiles of a C 18/C 18 peridinin-containing dinoflagellate, *Symbiodinium microadriaticum* (Dinophyceae), isolated from the mangrove jellyfish, *Cassiopea xamachana*. *Phycological research*, 63(3), 219-230.

Leggat W, Buck BH, Grice AM, Yellowlees D (2003) The impact of bleaching on the metabolic contribution of dinoflagellate symbionts to their giant clam host. *Plant Cell Environ*, 26:1951-1961.

Leggat, W., Hoegh-Guldberg, O., Dove, S., & Yellowlees, D. (2007). Analysis of an EST library from the dinoflagellate (*Symbiodinium* sp.) symbiont of reef-building corals 1. *Journal of Phycology*, 43(5), 1010-1021.

Lesser M P (1996) Elevated temperatures and ultraviolet radiation cause oxidative stress and inhibit photosynthesis in symbiotic dinoflagellates. *Limnology Oceanography* 41: 271–283.

Lesser M. P. (1997) Oxidative stress causes coral bleaching during exposure to elevated temperatures. *Coral reefs*, 16(3), 187-192.

Lesser M. P. (2006) Oxidative stress in marine environments: Biochemistry and physiological ecology. *Annual Review Physiology* 68:253–278.

Lesser, M. P. (2011). Coral bleaching: causes and mechanisms. In *Coral reefs: an ecosystem in transition* (pp. 405-419). Springer, Dordrecht.

Lesser, M. P. (2019). Phylogenetic signature of light and thermal stress for the endosymbiotic dinoflagellates of corals (Family Symbiodiniaceae). *Limnology and Oceanography*.

Manning, M. M., & Gates, R. D. (2008). Diversity in populations of free-living *Symbiodinium* from a Caribbean and Pacific reef. *Limnology and Oceanography*, 53(5), 1853-1861.

Mansour, M. P., Volkman, J. K., Jackson, A. E., & Blackburn, S. I. (1999). The fatty acid and sterol composition of five marine dinoflagellates. *Journal of Phycology*, 35(4), 710-720.

Mansour, J. S., Pollock, F. J., Díaz-Almeyda, E., Iglesias-Prieto, R., & Medina, M. (2018). Intra-and interspecific variation and phenotypic plasticity in thylakoid membrane properties across two *Symbiodinium* clades. *Coral Reefs*, 37(3), 841-850.

Martínez, M., Intralawan, A., Vázquez, G., Pérez-Maqueo, O., Sutton, P. & Landgrave, R. 2007. The coasts of our world: Ecological, economic and social importance. *Ecol Econ* 63:254-72

Matthews, J. L., Oakley, C. A., Lutz, A., Hillyer, K. E., Roessner, U., Grossman, A. R., ... & Davy, S. K. (2018). Partner switching and metabolic flux in a model cnidarian–dinoflagellate symbiosis. *Proceedings of the Royal Society B*, 285(1892), 20182336.

Mcclanahan, T. R., Ateweberhan, M., Graham, N. A. J., Wilson, S. K., Sebastian, C. R., Guillaume, M. M., & Bruggemann, J. H. (2007). Western Indian Ocean coral communities: bleaching responses and susceptibility to extinction. *Marine Ecology Progress Series*, 337, 1-13.

McGinty, E. S., Pieczonka, J., & Mydlarz, L. D. (2012). Variations in reactive oxygen release and antioxidant activity in multiple Symbiodinium types in response to elevated temperature. *Microbial Ecology*, 64(4), 1000-1007.

Mizusawa, N., & Wada, H. (2012). The role of lipids in photosystem II. *Biochimica et Biophysica Acta (BBA)-Bioenergetics*, 1817(1), 194-208

Moberg, F., & Folke, C. (1999). Ecological goods and services of coral reef ecosystems. *Ecological economics*, 29(2), 215-233

Mongrand, S., Bessoule, J. J., Cabantous, F., & Cassagne, C. (1998). The C16: 3\C18: 3 fatty acid balance in photosynthetic tissues from 468 plant species. *Phytochemistry*, 49(4), 1049-1064.

Moutin, T., Karl, D. M., Duhamel, S., Rimmelin, P., Raimbault, P., Van Mooy, B. A., & Claustre, H. (2007). Phosphate availability and the ultimate control of new nitrogen input by nitrogen fixation in the tropical Pacific Ocean. *Biogeosciences Discussions*, 4(4), 2407-2440.

Müller-Navarra, D. C., Brett, M. T., Liston, A. M., & Goldman, C. R. (2000). A highly unsaturated fatty acid predicts carbon transfer between primary producers and consumers. *Nature*, 403(6765), 74.

Muscantine L., Porter J.W. (1977) Reef corals: mutualistic symbioses adapted to nutrient-poor environments. *BioSciences* 27: 454–460.

Murphy D. J. (1986) The molecular organisation of the photosynthetic membranes of higher plants. *Biochim Biophys Acta* 864: 33–94

Nagelkerken, I., & Connell, S. D. (2015). Global alteration of ocean ecosystem functioning due to increasing human CO₂ emissions. *Proceedings of the National Academy of Sciences*, 112(43), 13272-13277.

Nygren, H., Seppänen-Laakso, T., & Rischer, H. (2017). Liquid Chromatography-Mass Spectrometry (LC-MS)-Based Analysis of Molecular Lipids in Algae Samples.

Nanjo, Y., Mizusawa, N., Wada, H., Slabas, A. R., Hayashi, H., & Nishiyama, Y. (2010). Synthesis of fatty acids de novo is required for photosynthetic acclimation of *Synechocystis* sp. PCC 6803 to high temperature. *Biochimica et Biophysica Acta (BBA)-Bioenergetics*, 1797(8), 1483-1490.

Niedzwiedzki, D. M., Jiang, J., LO, C. S., & Blankenship, R. E. (2014). Spectroscopic properties of the Chlorophyll a–Chlorophyll c2–Peridinin-Protein-Complex (acpPC) from the coral symbiotic dinoflagellate *Symbiodinium*. *Photosynthesis research*, 120(1-2), 125-139.

Niki E (2009) Lipid peroxidation: Physiological levels and dual biological effects. *Free Radic Biol Med* 47:469–484. doi: 10.1016/j.freeradbiomed.2009.05.032

Nisar, N., Li, L., Lu, S., Khin, N. C., & Pogson, B. J. (2015). Carotenoid metabolism in plants. *Molecular plant*, 8(1), 68-82.

Ohlrogge JB & Browse J (1995) Lipid biosynthesis. *Plant Cell* 7: 957–97

Olivotto I, Planas M, Simões N, et al (2011) Advances in breeding and rearing marine ornamentals. *Journal of the World Aquaculture Society* 42:135–166.

Orešič, M., Hänninen, V. A., & Vidal-Puig, A. (2008). Lipidomics: a new window to biomedical frontiers. *Trends in biotechnology*, 26(12), 647-652.

Parkinson, J. E., Baumgarten, S., Michell, C. T., Baums, I. B., LaJeunesse, T. C., & Voolstra, C. R. (2016). Gene expression variation resolves species and individual phylotypes among coral-associated dinoflagellates within the genus *Symbiodinium*. *Genome biology and evolution*, 8(3), 665-680.

Picciani, N., e Seiblitiz, I. G. D. L., de Paiva, P. C., e Castro, C. B., & Zilberberg, C. (2016). Geographic patterns of *Symbiodinium* diversity associated with the coral *Mussismilia hispida* (Cnidaria, Scleractinia) correlate with major reef regions in the Southwestern Atlantic Ocean. *Marine biology*, 163(11), 236.

Parkinson, J. E., Coffroth, M. A., & LaJeunesse, T. C. (2015). New species of Clade B *Symbiodinium* (Dinophyceae) from the greater Caribbean belong to different functional guilds: *S. aenigmaticum* sp. nov., *S. antillogorgium* sp. nov., *S. endomadracis* sp. nov., and *S. pseudominutum* sp. nov. *Journal of phycology*, 51(5), 850-858.

Pochon X., Gates R.D. (2010) A new *Symbiodinium* clade (Dinophyceae) from soritid foraminifera in Hawaii. *Molecular Phylogenetics and Evolution* 56: 492–497.

Polle, A. (1996). Mehler reaction: friend or foe in photosynthesis? *Botanica Acta*, 109(2), 84-89.

Ragni, M et al. PSII photoinhibition and photorepair in *Symbiodinium* (Pyrrhophyta) differs between thermally tolerant and sensitive phylotypes. *Marine Ecology Progress Series*, v. 406, p. 57-70, 2010.

Ramel, F., Birtic, S., Ginies, C., Soubigou-Taconnat, L., Triantaphylidès, C., & Havaux, M. (2012). Carotenoid oxidation products are stress signals that mediate gene responses to singlet oxygen in plants. *Proceedings of the National Academy of Sciences*, *109*(14), 5535-5540.

Raven, J. A., & Allen, J. F. (2003). Genomics and chloroplast evolution: what did cyanobacteria do for plants? *Genome biology*, *4*(3), 209.

Rees, S. A., Opdyke, B. N., Wilson, P. A., & Fifield, L. K. (2005). Coral reef sedimentation on Rodrigues and the Western Indian Ocean and its impact on the carbon cycle. *Philosophical Transactions of the Royal Society of London A: Mathematical, Physical and Engineering Sciences*, *363*(1826), 101-120.

Roberty, S., Fransolet, D., Cardol, P., Plumier, J. C., & Franck, F. (2015). Imbalance between oxygen photoreduction and antioxidant capacities in Symbiodinium cells exposed to combined heat and high light stress. *Coral Reefs*, *34*(4), 1063-1073.

Rodrigues, L. J., Grottoll, A. G., & Pease, T. K. (2008). Lipid class composition of bleached and recovering *Porites compressa* Dana, 1846 and *Montipora capitata* Dana, 1846 corals from Hawaii. *Journal of Experimental Marine Biology and Ecology*, *358*(2), 136-143.

Rogelj, J., Meinshausen, M., & Knutti, R. (2012). Global warming under old and new scenarios using IPCC climate sensitivity range estimates. *Nature climate change*, *2*(4), 248.

Rowan, R., Knowlton, N., Baker, A., & Jara, J. (1997). Landscape ecology of algal symbionts creates variation in episodes of coral bleaching. *Nature*, *388*(6639), 265.

Salvat, B. (1992). Coral reefs: a challenging ecosystem for human societies. *Global Environmental Change*, *2*(1), 12-18.

Saragosti, E., Tchernov, D., Katsir, A., & Shaked, Y. (2010). Extracellular production and degradation of superoxide in the coral *Stylophora pistillata* and cultured Symbiodinium. *PLoS One*, *5*(9), e12508.

Sarmiento, J. L., Slater, R., Barber, R., Bopp, L., Doney, S. C., Hirst, A. C., ... & Soldatov, V. (2004). Response of ocean ecosystems to climate warming. *Global Biogeochemical Cycles*, *18*(3).

Sato, N., Hagio, M., Wada, H., & Tsuzuki, M. (2000). Environmental effects on acidic lipids of thylakoid membranes.

- Sato, N., Suda, K., & Tsuzuki, M. (2004). Responsibility of phosphatidylglycerol for biogenesis of the PSI complex. *Biochimica et Biophysica Acta (BBA)-Bioenergetics*, 1658(3), 235-243.
- Schubotz, F., Wakeham, S. G., Lipp, J. S., Fredricks, H. F., & Hinrichs, K. U. (2009). Detection of microbial biomass by intact polar membrane lipid analysis in the water column and surface sediments of the Black Sea. *Environmental Microbiology*, 11(10), 2720-2734.
- Schmittner, A., Oschlies, A., Matthews, H. D., & Galbraith, E. D. (2008). Future changes in climate, ocean circulation, ecosystems, and biogeochemical cycling simulated for a business-as-usual CO₂ emission scenario until year 4000 AD. *Global biogeochemical cycles*, 22(1).
- Seemann, J., Sawall, Y., Auel, H., & Richter, C. (2013). The use of lipids and fatty acids to measure the trophic plasticity of the coral Stylophora subseriata. *Lipids*, 48(3), 275-286.
- Shemi, A., Schatz, D., Fredricks, H. F., Van Mooy, B. A., Porat, Z., & Vardi, A. (2016). Phosphorus starvation induces membrane remodeling and recycling in *Emiliana huxleyi*. *New Phytologist*, 211(3), 886-898.
- Sheppard, C. R. (1998). Biodiversity patterns in Indian Ocean corals, and effects of taxonomic error in data. *Biodiversity and Conservation*, 7(7), 847-868.
- Shi, X. M., & Chen, F. (2002). High-Yield Production of Lutein by the Green Microalga *Chlorella protothecoides*in Heterotrophic Fed-Batch Culture. *Biotechnology progress*, 18(4), 723-727.
- Shimajima, M., Ohta, H., & Nakamura, Y. (2009). Biosynthesis and function of chloroplast lipids. In *Lipids in Photosynthesis* (pp. 35-55). Springer, Dordrecht.
- Silverstein, R. N., Cunning, R., & Baker, A. C. (2017). Tenacious D: Symbiodinium in clade D remain in reef corals at both high and low temperature extremes despite impairment. *Journal of Experimental Biology*, 220(7), 1192-1196.
- Sinensky, M. (1974). Homeoviscous adaptation—a homeostatic process that regulates the viscosity of membrane lipids in *Escherichia coli*. *Proceedings of the National Academy of Sciences*, 71(2), 522-525.
- Smith D. J., Suggett DJ, Baker NR (2005) Is photoinhibition of zooxanthellae photosynthesis the primary cause of thermal bleaching in corals? *Global Change Biology* 11: 1–11.

- Søreide, J. E., Leu, E. V. A., Berge, J., Graeve, M., & Falk-Petersen, S. T. I. G. (2010). Timing of blooms, algal food quality and *Calanus glacialis* reproduction and growth in a changing Arctic. *Global change biology*, *16*(11), 3154-3163.
- Spencer, T., Teleki, K. A., Bradshaw, C., & Spalding, M. D. (2000). Coral bleaching in the southern Seychelles during the 1997–1998 Indian Ocean warm event. *Marine Pollution Bulletin*, *40*(7), 569-586.
- Suzuki, N., & Mittler, R. (2006). Reactive oxygen species and temperature stresses: a delicate balance between signaling and destruction. *Physiologia plantarum*, *126*(1), 45-51.
- Takahashi, S., Whitney, S., Itoh, S., Maruyama, T., & Badger, M. (2008). Heat stress causes inhibition of the de novo synthesis of antenna proteins and photobleaching in cultured Symbiodinium. *Proceedings of the National Academy of Sciences*, *105*(11), 4203-4208.
- Tchernov, D., Gorbunov, M. Y., DE Vargas, C., Yadav, S. N., Milligan, A. J., Häggblom, M., & Falkowski, P. G. (2004). Membrane lipids of symbiotic algae are diagnostic of sensitivity to thermal bleaching in corals. *Proceedings of the National Academy of Sciences of the United States of America*, *101*(37), 13531-13535
- Teece, M. A., Estes, B., Gelsleichter, E., & Lirman, D. (2011). Heterotrophic and autotrophic assimilation of fatty acids by two scleractinian corals, *Montastraea faveolata* and *Porites astreoides*. *Limnology and Oceanography*, *56*(4), 1285-1296.
- Terao, J., Minami, Y., & Bando, N. (2010). Singlet molecular oxygen-quenching activity of carotenoids: relevance to protection of the skin from photoaging. *Journal of clinical biochemistry and nutrition*, *48*(1), 57-62.
- Thornhill, D. J., Lewis, A. M., Wham, D. C., & LaJeunesse, T. C. (2014). Host-specialist lineages dominate the adaptive radiation of reef coral endosymbionts. *Evolution*, *68*(2), 352-367.
- Toller W.W., Rowan R., Knowlton N. (2001) Zooxanthellae of the *Montastrea annularis* species complex: patterns of distribution of four taxa of Symbiodinium on different reefs and across depths. *Biological Bulletin* 201: 348-359.
- Tomita, M., & Nishioka, T. (Eds.). (2006). *Metabolomics: the frontier of systems biology*. Springer Science & Business Media.
- Tolosa I, Treignier C, Grover R, Ferrier-Page S C (2011) Impact of feeding and short-term temperature stress on the content and isotopic signature of fatty acids, sterols, and alcohols in the scleractinian coral *Turbinaria reniformis*. *Coral Reefs* 30:763–774

- Tong, H., Cai, L., Zhou, G., Yuan, T., Zhang, W., Tian, R., ... & Qian, P. Y. (2017). Temperature shapes coral-algal symbiosis in the South China Sea. *Scientific reports*, 7, 40118.
- Towle, E. K., Palacio-Castro, A. M., Baker, A. C., & Langdon, C. (2017). Source location and food availability determine the growth response of *Orbicella faveolata* to climate change stressors. *Regional Studies in Marine Science*, 10, 107-115.
- Treignier, C., Grover, R., Ferrier-Pages, C., & Tolosa, I. (2008). Effect of light and feeding on the fatty acid and sterol composition of zooxanthellae and host tissue isolated from the scleractinian coral *Turbinaria reniformis*. *Limnology and Oceanography*, 53(6), 2702-2710.
- Valentine, R. C., & Valentine, D. L. (2004). Omega-3 fatty acids in cellular membranes: a unified concept. *Progress in lipid research*, 43(5), 383-402.
- Valentine, R. C., & Valentine, D. L. (2009). *Omega-3 fatty acids and the DHA principle*. CRC press.
- Van Mooy, B. A., Rocap, G., Fredricks, H. F., Evans, C. T., & Devol, A. H. (2006). Sulfolipids dramatically decrease phosphorus demand by picocyanobacteria in oligotrophic marine environments. *Proceedings of the National Academy of Sciences*, 103(23), 8607-8612.
- Van Mooy, B. A., Fredricks, H. F., Pedler, B. E., Dyhrman, S. T., Karl, D. M., Koblížek, M., ... & Rappé, M. S. (2009). Phytoplankton in the ocean use non-phosphorus lipids in response to phosphorus scarcity. *Nature*, 458(7234), 69.
- Van Mooy, B. A., & Fredricks, H. F. (2010). Bacterial and eukaryotic intact polar lipids in the eastern subtropical South Pacific: water-column distribution, planktonic sources, and fatty acid composition. *Geochimica et Cosmochimica Acta*, 74(22), 6499-6516.
- Van Oppen, M. J., Palstra, F. P., Piquet, A. M. T., & Miller, D. J. (2001). Patterns of coral–dinoflagellate associations in *Acropora*: significance of local availability and physiology of Symbiodinium phylotypes and host–symbiont selectivity. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 268(1478), 1759-1767.
- Vigh, L., Escribá, P. V., Sonnleitner, A., Sonnleitner, M., Piotta, S., Maresca, B., ... & Harwood, J. L. (2005). The significance of lipid composition for membrane activity: new concepts and ways of assessing function. *Progress in lipid research*, 44(5), 303-344.
- Vigh, L., Nakamoto, H., Landry, J., GOMEZ-MUNOZ, A. N. T. O. N. I. O., Harwood, J. L., & Horvath, I. (2007). Membrane regulation of the stress response from prokaryotic models to mammalian cells. *Annals of the New York Academy of Sciences*, 1113(1), 40-51.

- Vogel, G., & Eichenberger, W. (1992). Betaine lipids in lower plants. Biosynthesis of DGTS and DGTA in *Ochromonas danica* (Chrysophyceae) and the possible role of DGTS in lipid metabolism. *Plant and cell physiology*, 33(4), 427-436.
- Wada H (1994) Contribution of membrane lipids to the ability of the photosynthetic machinery to tolerate temperature stress. *Proc Natl Acad Sci USA* 91:4273–4277
- Wada, H., & Murata, N. (Eds.). (2009). *Lipids in Photosynthesis*. Springer Netherlands.
- Wang, L., Shen, W., Kazachkov, M., Chen, G., Chen, Q., Carlsson, A. S., ... & Zou, J. (2012). Metabolic interactions between the Lands cycle and the Kennedy pathway of glycerolipid synthesis in *Arabidopsis* developing seeds. *The Plant Cell*, tpc-112.
- Warner, M. E., Fitt, W. K., & Schmidt, G. W. (1999). Damage to photosystem II in symbiotic dinoflagellates: a determinant of coral bleaching. *Proceedings of the National Academy of Sciences*, 96(14), 8007-8012.
- Weis, V. M. (2008). Cellular mechanisms of Cnidarian bleaching: stress causes the collapse of symbiosis. *Journal of Experimental Biology*, 211(19), 3059-3066.
- Wenk, M. R. (2005). The emerging field of lipidomics. *Nature reviews Drug discovery*, 4(7), 594-610.
- Widomska, J., Welc, R., & Gruszecki, W. I. (2019). The effect of carotenoids on the concentration of singlet oxygen in lipid membranes. *Biochimica et Biophysica Acta (BBA)-Biomembranes*.
- Wilkinson C (2008) Status of coral reefs of the world: 2008. Global Coral Reef Monitoring Network and Reef and Rainforest Research Centre, Townsville
- Wohlers, J., Engel, A., Zöllner, E., Breithaupt, P., Jürgens, K., Hoppe, H. G., ... & Riebesell, U. (2009). Changes in biogenic carbon flow in response to sea surface warming. *Proceedings of the national academy of sciences*, 106(17), 7067-7072
- Wu W. G. & Chi L. M. (1991) Conformational change of cholesterol side chain in lipid bilayers. *J Chem Soc* 113: 4683–4685
- Yamamoto H. Y. (2006) Functional roles of the major chloroplast lipids in the violaxanthin cycle. *Planta* 224: 719–724
- Yamashiro, H., Oku, H., & Onaga, K. (2005). Effect of bleaching on lipid content and composition of Okinawan corals. *Fisheries Science*, 71(2), 448-453.
- Yang, S. Y., Keshavmurthy, S., Obura, D., Sheppard, C. R., Visram, S., & Chen, C. A. (2012). Diversity and distribution of Symbiodinium associated with seven common coral species in the Chagos Archipelago, Central Indian Ocean. *PLoS One*, 7(5), e35836.

Yao, L., Gerde, J. A., Lee, S. L., Wang, T., & Harrata, K. A. (2015). Microalgae lipid characterization. *Journal of agricultural and food chemistry*, 63(6), 1773-1787.

Yin, H.; Xu, L.; Porter, N. A. (2011) Free Radical Lipid Proxidation: Mechanisms and Analysis. *Chem. Rev.*, 111 (10), 5944–5972.

Yoshinaga, M. Y., Kellermann, M. Y., Valentine, D. L., & Valentine, R. C. (2016). Phospholipids and glycolipids mediate proton containment and circulation along the surface of energy-transducing membranes. *Progress in lipid research*, 64, 1-15.

Zhukova, N. V., & Titlyanov, E. A. (2003). Fatty acid variations in symbiotic dinoflagellates from Okinawan corals. *Phytochemistry*, 62(2), 191-195.