

REFERÊNCIAS BIBLIOGRÁFICAS

REFERÊNCIAS BIBLIOGRÁFICAS

- 1 Smith R. The scientific basis of health services. *BMJ* 1995;311:961-2.
- 2 Office of Technology Assessment of the Congress of the United States. Assessing the efficacy and safety of medical technologies. Washington. 1978 . (online). Available: http://www.wws.princeton.edu/~ota/ns20/year_f.html 1978.
- 3 Office of Technology Assessment of the Congress of the United States. The Impact of randomized clinical trials on health policy and medical practice. Washington. Government Printing Office, 1983. http://www.wws.princeton.edu/~ota/ns20/year_f.html
- 4 George DL. Life expectancy, health and the ABPI. *Lancet* 1986;346
- 5 Freeman HE. *Handbook of medical sociology*. Englewood Cliffs: NJ Prentice Hall; 1988.
- 6 Lenfant C. Shattuck lecture--clinical research to clinical practice--lost in translation? *N Engl J Med* 2003;349:868-74.

- 7 Guyer B, Freeman MA, Strobino DM, Sondik EJ. Annual summary of vital statistics: trends in the health of Americans during the 20th century. *Pediatrics* 2000;106:1307-17.
- 8 Lee SK, Mcmillan DD, Ohisson A, Pendray M, Synnes A, Whyte R, Chien LY, Sale J. Variations in practice and outcomes in the Canadian NICU network: 1996-1997. *Pediatrics* 2000;106:1070-9.
- 9 Soll RF, Andruscavage L. The principles and practice of evidence-based neonatology. *Pediatrics*. 1999;103:215-24.
- 10 Sueta CA, Massing MW, Chowdhury M, Biggs DP, Simpson RJ Jr. Undertreatment of hyperlipidemia in patients with coronary artery disease and heart failure. *J Card Fail*. 2003;9:36-41.
- 11 Stafford RS, Radley DC. The underutilization of cardiac medications of proven benefit, 1990 to 2002. *J Am Coll Cardiol* 2003;41:56-61.
- 12 Hewitt M, Simone JV. Ensuring quality cancer care. Editors National Cancer Policy Board, Institute of Medicine and National Research Council, 1999. p. 79 <http://www.nap.edu/books/0309064805/html>
- 13 Gupta M. A critical appraisal of evidence-based medicine: some ethical considerations. *J Eval Clin Pract*. 2003;9:111-21.
- 14 Sackett DL, Rosenberg WM, Gray JA, Haynes RB, Richardson WS. Evidence based medicine: what it is and what it isn't. *BMJ* 1996;13:312:71-2.
- 15 University of Illinois of Chicago UIC. Is all evidence created equal? <http://www.uic.edu/depts/lib/lhsp/resources/levels.html>.
- 16 Wong JH, Findlay JM, Suarez-Almazor ME. Regional performance of carotid endarterectomy. Appropriateness, outcomes, and risk factors for complications. *Stroke*. 1997;28:891-8.
- 17 Straus SE, McAlister FA. *CMAJ*. 2000;163:837-41. Evidence-based medicine: a commentary on common criticisms.
- 18 Evidence-based medicine Working Group. Appraise using JAMA Guide. <http://medicine.ucsf.edu/resources/guidelines/users.html>

- 19 Smith R. What clinical information do doctors need? *BMJ* 1996;313:1062-8.
- 20 Harbour R, Miller J. A new system for grading recommendations in evidence based guidelines. *BMJ*. 2001;323:334-6.
- 21 Berlin JA. The role of meta-analysis in the regulatory process for foods, drugs, and devices. *JAMA* 1999;281:830.
- 22 Bergman DA. Evidence-based guidelines and critical pathways for quality improvement. . *Pediatrics* 1999;103:225-32.
- 23 Ely JW, Osheroff JA, Ebell MH, Bergus GR, Levy BT, Chambliss ML, Evans ER. Analysis of questions asked by family doctors regarding patient care. *BMJ*. 1999; 319:358-61.
- 24 Guyatt GH, Meade MO, Jaeschke RZ, Cook DJ, Haynes RB. Practitioners of evidence based care. Not all clinicians need to appraise evidence from scratch but all need some skills. *BMJ* 2000;320:954-5.
- 25 Lomas J, Anderson GM, Dominick-Pierre K, Vayda E, ENKIN MW, HANNAH WJ. Do practice guidelines guide practice? The effect of a consensus statement on the practice of physicians. *N Engl J Med* 1989;321:1306-11.
- 26 Mccoll A, Smith H, White P, Field J. General practitioner's perceptions of the route to evidence based medicine: a questionnaire survey. *BMJ* 1998;316:361-5.
- 27 Al-Ansary LA, Khoja TA. The place of evidence-based medicine among primary health care physicians in Riyadh region, Saudi Arabia. *Fam Pract* 2002;9:537-42.
- 28 Hossne WS. In: SEgre M, Cohen C. *Bioética*. 2ª ed. São Paulo: Edusp; p.75.
- 29 Culver CM. In:Serge M, Cohen C. *Bioética*. 2ª ed. São Paulo: Edusp; p.63.

- 30 Sackett DL, Richardson WS, Rosenberg W, Haynes RB. *Evidence-based medicine. How to practice and Teach EMB*. New York: Churchill Livingstone; 1996.
- 31 Kerridge I, Lowe M, Henry D. Ethics and evidence based medicine. *BMJ* 1998; 316:1151.
- 32 Christakis DA, Davis R, Rivara FP. Pediatric evidence-based medicine: past, present, and future. *J Pediatr* 2000;136:383-9.
- 33 Wesley ELY, Bernard GR. Transfusions in critically ill patients. *N Engl J Med* 1999;340:467-8.
- 34 [No authors listed] Ventilation with lower tidal volumes as compared with traditional tidal volumes for acute lung injury and the acute respiratory distress syndrome. The Acute Respiratory Distress Syndrome Network. *N Engl J Med*. 2000 ;342:1301-8.
- 35 Kilbride HW, Powers R, Wirtschafter DD, Sheehan MB, Charsha DS, LaCorte M, Finer N, Goldmann DA.
Evaluation and development of potentially better practices to prevent neonatal nosocomial bacteremia. *Pediatrics*. 2003 ;111:e504-18.
- 36 Vincent JL. Which therapeutic interventions in critical care medicine have been shown to reduce mortality in prospective, randomized, clinical trials? A survey of candidates for the Belgian Board examination in Intensive Care Medicine. *Crit Care Med* 2000;28:1616-20.
- 37 Ellis J, Mulligan I, Rowe J, Sackett DL. Inpatient general medicine is evidence based. A-Team, Nuffield Department of Clinical Medicine. *Lancet* 1995;346:407-10.
- 38 Gill P, Dowell AC, Neal RD, Smith N, Heywood P, Wilson AE. Evidence based general practice: a retrospective study of interventions in one training practice. *BMJ* 1996;312:819-21.

- 39 Kenny SE, Shankar KR, Rintala R, Lamont GL, Lloyd DA. Evidence-based surgery: interventions in a regional paediatric surgical unit. *Arch Dis Child* 1997;76:50-3.
- 40 Rudolf MC, Lyth N, Bundle A, Rowland G, Kelly A, Bosson S, Garner M, Guest P, Khan M, Thazin R, Bennett T, Damman D, Cove V, Kaur V. A search for the evidence supporting community paediatric practice. *Arch Dis Child* 1999;80:257-61.
- 41 Tsuruoka K, Tsuruoka Y, Yoshimuri M, Sekigushi S, Mise J, Asai Y, Nago N, Igarashi M. Evidence based general practice. Drug treatment in general practice in Japan is evidence based. *BMJ* 1996;313:114.
- 42 Godlee F. Getting evidence into practice. Needs the right resources and the right organisation. *BMJ* 1998;317:6.
- 43 Evid-based Ped: the future is now. *J Pediatr* 2000;13:282-4
- 44 Hewitt M, Simone JV. Ensuring quality cancer care. Editors National Cancer Policy Board, Institute of Medicine and National Research Council, 1999. p. 96 <http://www.nap.edu/books/0309064805/html>
- 45 American Academy of Pediatrics. Guidelines for developing admission and discharge policies for the pediatric intensive care unit.. Committee on Hospital Care and Section of Critical Care. Society of Critical Care Medicine. Pediatric Section Admission Criteria Task Force. *Pediatrics*. 1999 103:840-2.
- 46 Guyatt GH, Sackett DL, Sinclair JC, Hayward R, Cook DJ, Cook RJ. Users' guides to the medical literature Users' guides to the medical literature. IX. A method for grading health care recommendations. Evidence-Based Medicine Working Group. *JAMA* 1995;13:274:1800-4.
- 47 Boldt J. The good, the bad, and the ugly: should we completely banish human albumin from our intensive care units? *Anesth Analg* 2000; 91:887-95.

- 48 Marik PE. *Handbook of evidence-based critical care*. New York: Springer; 2001. p.113
- 49 Alderson P, Schierhout G, Roberts I, Bunn F. Colloids versus crystalloids for fluid resuscitation in critically ill patients (Cochrane Review). In: The Cochrane Library. Oxford: Update Software. Issue 4, 2002.
- 50 Choi PT, Yip G, Quinonez LG, Cook DJ. Crystalloids vs. colloids in fluid resuscitation: a systematic review. *Crit Care Med* 1999;27:200-10.
- 51 Schierhout G, Roberts I. Fluid resuscitation with colloid or crystalloid solutions in critically ill patients: a systematic review of randomised trials. *BMJ* 1998;316:961-4.
- 52 [No authors listed]. Human albumin administration in critically ill patients: systematic review of randomised controlled trials. Cochrane Injuries Group Albumin Reviewers. *BMJ* 1998;317:235-40.
- 53 Task Force of the American College of Critical Care Medicine, Society of Critical Care Medicine. Practice parameters for hemodynamic support of sepsis in adult patients in sepsis.. *Crit Care Med* 1999;27:639-60.
- 54 Gentilini P, Casini-Raggi V, Di Fiore G, Romanelli RG, Buzzelili G, Pinzani M, La Villa G, Laffi G. Albumin improves the response to diuretics in patients with cirrhosis and ascites: results of a randomized, controlled trial. *J Hepatol* 1999;30:639-45.
- 55 Magder S, Lagonidis D. Effectiveness of albumin versus normal saline as a test of volume responsiveness in post-cardiac surgery patients. *J Crit Care* 1999;14:164-71.

- 56 Sort P, Navasa M, Arroyo V, Aldeguer X, Planas R, Ruiz-Del-Arbol L, Castells L, Vargas V, Soriano G, Guevara M, Gines P, Rodes J. Effect of intravenous albumin on renal impairment and mortality in patients with cirrhosis and spontaneous bacterial peritonitis. *N Engl J Med* 1999;5:341:403-9.
- 57 Fliser D, Zurbruggen I, Mutschler E, Bischoff I, Nussberger J, Franek E, Ritz E. Coadministration of albumin and furosemide in patients with the nephritic syndrome. *Kidney Int.* 1999;55:629-34.
- 58 Boldt J, Muller M, Mentges D, Papsdorf M, Hempelmann G. Volume therapy in the critically ill: is there a difference? *Intensive Care Med* 14:28-36,1998.
- 59 Rubin H, Carlson S, De Meo M, Ganger D, Craig RM. Randomized, double-blind study of intravenous human albumin in hypoalbuminemic patients receiving total parenteral nutrition. *Crit Care Med* 1997;25:249-52.
- 60 Gines A, Fernandez-Esparrach G, Monescilo A, Vila C, Domenech E, Abecasis R, Angeli P, Ruiz-Del-Arbol L, Planas R, Sola R, Gines P, Terg R, Inglada L, Vaque P, Salerno F, Vargas V, Clemente G, Quer JC, Jimenez W, Arroyo V, Rodes J. Randomized trial comparing albumin, dextran 70, and polygeline in cirrhotic patients with ascites treated by paracentesis. *Gastroenterology* 1996;111:1002-10.
- 61 Akcicek F, Yalniz T, Basci A, Ok E, Mees EJ. Diuretic effect of frusemide in patients with nephrotic syndrome: is it potentiated by intravenous albumin? *BMJ.* 1995;310:162-3.
- 62 Golub R, Sorrento JJ JR, Cantu R JR, Nieman DM, Moideen A, Stein HD. Efficacy of albumin supplementation in the surgical intensive care unit: a prospective, randomized study. *Crit Care Med* 1994;22:613-9.

- 63 Stockwell MA, Soni N, Riley B. Colloid solutions in the critically ill. A randomised comparison of albumin and polygeline. 1. Outcome and duration of stay in the intensive care unit. *Anaesthesia* 1992;47:3-6.
- 64 Stockwell MA, Scott A, Day A, Riley B, Soni N. Colloid solutions in the critically ill. A randomised comparison of albumin and polygeline 2. Serum albumin concentration and incidences of pulmonary oedema and acute renal failure. *Anaesthesia* 1992;47:7-9.
- 65 Wojtysiak SL, Brown RO, Roberson D, Powers DA, Kudsk KA. Effect of hypoalbuminemia and parenteral nutrition on free water excretion and electrolyte-free water resorption. *Crit Care Med* 1992;20:164-9.
- 66 Greissman A, Silver P, Nimkoff L, Sagy M. Albumin bolus administration versus continuous infusion in critically ill hypoalbuminemic pediatric patients. *Intensive Care Med* 1996;22:495-9.
- 67 Haws RM, Baum M. Efficacy of albumin and diuretic therapy in children with nephrotic syndrome. *Pediatrics*. 1993 91:1142-6.
- 68 Marik PE. *Handbook of evidence-based critical care*. New York: Springer; 2001. p.110.
- 69 Schaer GL. Norepinephrine plus low-dose dopamine: a beneficial combination. *Crit Care Med*. 1998;26:205.
- 70 Olsen NV, Lund J, Jensen PF, Espersen K, Kanstrup IL, Plum I, Leyssac PP. Dopamine, dobutamine, and dopexamine. A comparison of renal effects in unanesthetized human volunteers. *Anesthesiology* 1999;79:685-94.
- 71 Katz DV Valor terapêutico da dopamina na proteção da função renal na sepse grave, choque séptico e pós-operatório de cirurgia cardíaca em pediatria [dissertação]. São Paulo: Faculdade de Medicina da Universidade de São Paulo; 2000.

- 72 Bellomo R, Chapman M, Finfer S, Hickling K, Myburgh J. Low-dose dopamine in patients with early renal dysfunction: a placebo-controlled randomised trial. Australian and New Zealand Intensive Care Society (ANZICS) *Clinical Trials Group* 2000;356:2139-43.
- 73 Lassnigg A, Donner E, Grubhofer G, Presteri E, Drumi W, Hiesmayr M. Lack of renoprotective effects of dopamine and furosemide during cardiac surgery. *J Am Soc Nephrol* 2000;11:97-104.
- 74 Ichai C, Soubielle J, Carles M, Giunti C, Grimaud D. Comparison of the renal effects of low to high doses of dopamine and dobutamine in critically ill patients: a single-blind randomized study. *Crit Care Med* 2000;28:921-8.
- 75 Abizaid AS, Clark CE, Mintz GS, Dosa S, Popma JJ, Pichard AD, Satler LF, Harvey M, Kent KM, Leon MB. Effects of dopamine and aminophylline on contrast-induced acute renal failure after coronary angioplasty in patients with preexisting renal insufficiency. *Am J Cardiol* 1999;83:260-3, A5.
- 76 Marik PE, Iglesias J. Low-dose dopamine does not prevent acute renal failure in patients with septic shock and oliguria. NORASEPT II Study Investigators. *Am J Med* 1999;107:387-90.
- 77 Lema G, Urzua J, Jalil R, Canessa R, Moran S, Sacco C, Medel J, Irrarrazaval M, Zalaquett R, Fajardo C, Meneses G. Renal protection in patients undergoing cardiopulmonary bypass with preoperative abnormal renal function. *Anesth Analg* 1998;86:3-8.
- 78 Abu-Saad HH, Bours GJ, Stevens B, Hamers JP. Assessment of pain in the neonate. *Semin Perinatol* 1998;22:402-16.
- 79 Cote CJ, Karl HW, Notterman DA, Weinberg JA, McCloskey C. Adverse sedation events in pediatrics: analysis of medications used for sedation. *Pediatrics*. 2000 Oct;106(4):633-44.
- 80 Petrack EM, Christopher NC, Kriwinsky J. Pain management in the emergency department: patterns of analgesic utilization. *Pediatrics* 1997;99:711-4.

- 81 Porter FL, Wolf CM, Gold J, Lotsoff D, Miller JP. Pain and pain management in newborn infants: a survey of physicians and nurses. *Pediatrics* 1997;100:626-32.
- 82 Stein-Parbury J, Mckinley S. Patients experiences of being in an intensive care unit: a select literature review. *Am J Crit Care* 2000; 9:20-7.
- 83 Partridge JC, Wall SN. Analgesia for dying infants whose life support is withdrawn or withheld. *Pediatrics* 1997;99:76-9.
- 84 Marik PE. *Handbook of evidence-based critical care*. New York: Springer; 2001. p.399.
- 85 Lowrie L, Weiss AH, Lacombe C. The pediatric sedation unit: a mechanism for pediatric sedation. *Pediatrics*. 1998 ;102:E30.
- 86 Ng E, Taddio A, Ohlsson A. Intravenous midazolam infusion for sedation of infants in the neonatal intensive care unit. *Cochrane Database Syst Rev* CD002052, 2000.
- 87 Ostermann ME, Keenan SP, Seiferling RA, Sibbald WJ. Sedation in the intensive care unit: a systematic review. *Sedation in the intensive care unit: a systematic review*. 2000;283:1451-59.
- 88 Recommandations pour la pratique clinique. Sédation, analgésie et curarisation en réanimation. *An Fr Anesth Réanim* 2000;19:98-105.
- 89 Practice guidelines for sedation and analgesia by non-anesthesiologists. A report by the American Society of Anesthesiologists task force on sedation and analgesia by non-anesthesiologists. *Anesthesiology* 1996;84:459-71.
- 90 Shapiro Ba, Warren J, Egol Ab, Greenbaum Dm, Jacobi J, Nasraway Sa, Schein Rm, Spevetz A, Stone Jr. Practice parameters for intravenous analgesia and sedation for adult patients in the intensive care unit: an executive summary. *Crit Care Med* 1995; 23:1596-600.

- 91 No authors listed. American Academy of Pediatrics Committee on Drugs. Guidelines for monitoring and management of pediatric patients during and after sedation for diagnostic and therapeutic procedures. *Pediatrics* 1992; 89:1110-5.
- 92 Wathen JE, Roback MG, Mackenzie T, Bothner JP.
Does midazolam alter the clinical effects of intravenous ketamine sedation in children? A double-blind, randomized, controlled, emergency department trial. *Ann Emerg Med.* 2000;36:579-88.
- 93 Marcus RJ, Victoria BA, Rushman SC, Thompson JP.
Comparison of ketamine and morphine for analgesia after tonsillectomy in children. *Br J Anaesth.* 2000;84:739-42.
- 94 Anand KJ, Hickey PR. Halothane-morphine compared with high-dose sufentanil for anesthesia and postoperative analgesia in neonatal cardiac surgery. *N Engl J Med.* 1992 ;326:1-9.
- 95 Kennedy RM, Porter FL, Miller JP, Jaffe DM. Comparison of fentanyl/midazolam with ketamine/midazolam for pediatric orthopedic emergencies. *Pediatrics* 1998;102:956-63.
- 96 Marx CM, Stein J, Tyler MK, Nieder ML, Shurin SB, Blumer JL.
Ketamine-midazolam versus meperidine-midazolam for painful procedures in pediatric oncology patients. *J Clin Oncol.* 1997;15:94-102.
- 97 Parkinson L, Hughes J, Gill A, Billingham I, Ratcliffe J, Choonara I.
A randomized controlled trial of sedation in the critically ill. *Paediatr Anaesth* 1997;7:405-10.
- 98 Anand KJ, Hickey PR. Halothane-morphine compared with high-dose sulfentanil for anesthesia and postoperative analgesia in neonatal cardiac surgery. *N Engl J Med* 1992;326:1-9.
- 99 Hertzog JH, Dalton HJ, Anderson BD, Shad AT, Gootenberg JE, Hauser GJ. Prospective evaluation of propofol anesthesia in the pediatric intensive care unit for elective oncology procedures in ambulatory and hospitalized children. *Pediatrics* 2000;106:742-7.

- 100 Parker RI, Mahan RA, Giugliano D, Parker MM. Efficacy and safety of intravenous midazolam and ketamine as sedation for therapeutic and diagnostic procedures in children. *Pediatrics* 1997; 99:427-31.
- 101 Miller. Faltam Autores completos *Anesthesia*. 5th ed. Churchill Livingstone. 2000. p.1087.
- 102 Krauss B, Green SM. Sedation and analgesia for procedures in children. *N Engl J Med* 2000;342:938-45.
- 103 Ambrose C, Sale S, Howells R, Bevan C, Jenkins I, Weir P, Murphy P, Wolf A. Intravenous clonidine infusion in critically ill children: dose-dependent sedative effects and cardiovascular stability. *Br J Anaesth* 2000;84:794-6.
- 104 Lieh-Lai MW, Kauffman RE, Uy HG, Danjin M, Simpson PM. A randomized comparison of ketorolac tromethamine and morphine for postoperative analgesia in critically ill children. *Crit Care Med* 1999; 27:2786-91.
- 105 Levy GJ, Strauss RG, Hume H, Schloz L, Albanese MA, Blazina J, Werner A, Sotelo-Avila C, Barrasso C, Blanchette V, et al. National survey of neonatal transfusion practices: I. Red blood cell therapy. *Pediatrics* 1993;91:523-9.
- 106 Westphal RG. Handbook of transfusion medicine. 3th ed. Washington:American Red Cross Blood services. 1996.
- 107 Lee GR, Lukens J, Greer JP, Foerster J, Paraskevas F, Rodges GM. *Wintrobe's clinical hematology*. 10th ed. Baltimore: Williams and Wilkins;1998. p.827-40.
- 108 Ringer SA, Richardson DK, Sacher RA, Keszler M, Churchill WH. Variations in transfusion practice in neonatal intensive care. *Pediatrics* 1998;101:194-200.

- 109 Hebert PC, Wells G, Martin C, Tweeddale M, Marshall J, Blauchman M, Pagliarello G, Schweitzer I, Calder L. A Canadian survey of transfusion practices in critically ill patients. Transfusion Requirements in Critical Care Investigators and the Canadian Critical Care Trials Group. *Crit Care Med* 1998;26:482-7.
- 110 Wardrop CAJ, Holland BM, Jones G. Consensus in red cell transfusion. *BMJ* 1995;311:962-3.
- 111 Algora Weber M, Fernandez Montoya A, Gomez Viillagran JL, Martin Vega C, Prats Arrojo I, Puig Alcaraz N, Pujol Bosch M, Vesga Carasa Ma. Guide for transfusion of red blood cells, platelets and labile plasma products. Joint directive of the Spanish society of blood transfusion. *Med Clin (Barc)* 1999;113:471-4.
- 112 Simon TL, Alverson DC, Aubuchon J, Cooper ES, Dechristopher PJ, Glenn GC, Gouid, SA, Harrison CR, Milam JD, Moise KJ JR, Rodwig FR JR, Sherman LA, Shulman IA, Stehling L. Practice parameter for the use of red blood cell transfusions: developed by the Red Blood Cell Administration Practice Guideline Development Task Force of the College of American Pathologists. *Arch Pathol Lab Med* 1998; 122:130-8.
- 113 American Academy Of Pediatrics Section on Anesthesiology Evaluation and Preparation of Pediatric Patients Undergoing Anesthesia (RE9633). *Pediatrics* 1996;98:502-8.
- 114 Slaughter TF, Greenberg CS. Practice guidelines and treatment of patients with von Willebrand's disease. *Anesthesiology* 1996;85:441-2.
- 115 Fetus and Newborn Committee, Canadian Paediatric Society. Guidelines for transfusion of erythrocytes to neonates and premature infants. *CMAJ* 1997;147:1781-92.
- 116 Welch HG, Meehan KR, Goodnough LT. Prudent strategies for elective red blood cell transfusion. *Ann Intern Med* 1992;1:116:393-402.

- 117 Hebert PC, Wells G, Blajchman MA, Marshall J et al. Canadian Critical Care Trials Group. A multicenter, randomized, controlled clinical trial of transfusion requirements in critical care. Transfusion requirements in critical care investigators. *N Engl J Med* 1999;340:409-17.
- 118 Brooks SE, Marcus DM, Gillis D, Pirie E, Johnson C, Bhatia J. The effect of blood transfusion protocol on retinopathy of prematurity: A prospective, randomized study. *Pediatrics* 1999;104:514-8.
- 119 Adams RJ, Mckie VC, Hsu L, Files B, Vichinsky E, Pegelow C, Abboud M, Gallagher D, Kutiar A, Nichols FT, Bonds DR, Brambila D. Prevention of a first stroke by transfusions in children with sickle cell anemia and abnormal results on transcranial Doppler ultrasonography. *N Engl J Med* 1998;339:5-11.
- 120 Vichinsky EP, Haberkem CM, Neumayr L, Earles AN, Black D, Koshy M, Pegelow C, Abboud M, Ohene-Frempong K, Iyer RV. A comparison of conservative and aggressive transfusion regimens in the perioperative management of sickle cell disease. The Preoperative Transfusion in Sickle Cell Disease Study Group. *N Engl J Med* 1995;333:206-13.
- 121 Meyer J, SIVE A, Jacobs P. Empiric red cell transfusion in asymptomatic preterm infants. *Acta Paediatr* 1993;82:30-4.
- 122 Lam NP. National survey of stress ulcer prophylaxis. *Crit Care Med* 1999; 27:98.
- 123 Kupfer Y, Cappeli MS, Tessier S. High risk bleeding, part ii acute gastrointestinal bleeding in the intensive care unit The Intensivist's perspective. *Gastroenterol Clin North Am* 2000;29:275-307.
- 124 Neff M, Metry JM, Frick P, Anliker M, Knoblauch M. The microvasculature of the small-intestinal mucosa of the rat: quantification of hemodynamic effects of topically applied cimetidine, ranitidine, somatostatin, and vasopressin. *Scand J Gastroenterol* 1985;(Supl.112): 6-11.

- 125 Kaszaki J, Boros M, Szabo A, Nagy S. Role of histamine in the intestinal flow response following mesenteric ischemia. *Shock* 1994;2:413-20.
- 126 Abraham E, Matthay MA, Dinarello CA, Vincent JL, Cohen J, Opal SM, Glauser M, Parsons P, Fisher CJ Jr, Repine JE. Consensus conference definitions for sepsis, septic shock, acute lung injury, and acute respiratory distress syndrome: time for a reevaluation. *Crit Care Med*. 2000;28:232-5.
- 127 Cantu TG, Korek JS. Central nervous system reactions to histamine-2 receptor blockers. *Ann Intern Med* 1991;114:1027-34.
- 128 Messori A, Trippoli S, Vaiani M, Gorini M, Corrado A. Bleeding and pneumonia in intensive care patients given ranitidine and sucralfate for prevention of stress ulcer: meta-analysis of randomized controlled trials *BMJ* 2000;321:1103.
- 129 Hanisch EW, Encke A, Naujoks F, Windolf J. A randomized, double-blind trial for stress ulcer prophylaxis shows no evidence of increased pneumonia. *Am J Surg* 176:453-7,1998.
- 130 Pickworth KK, Falcone RE, Hoogeboom JE, Santanello SA. Occurrence of nosocomial pneumonia in mechanically ventilated trauma patients: a comparison of sucralfate and ranitidine. *Crit Care Med* 1993;21:1856-62.
- 132 Prod'hom G, Leuenberger P, Koerfer J, Blum A, Chiolerio R, Schaller MD, Perret C, Spinnier O, Biondel J, Siegrist H, et al. Nosocomial pneumonia in mechanically ventilated patients receiving antacid, ranitidine, or sucralfate as prophylaxis for stress ulcer. A randomized controlled trial. *Ann Intern Med* 1994;120:653-62.
- 133 Reusser P, Gyr K, Scheidegger D, Buchmann B, Buser M, Zimmerli W. Prospective endoscopic study of stress erosions and ulcers in critically ill neurosurgical patients: current incidence and effect of acid-reducing prophylaxis. *Crit Care Med* 1990;18:270-4.

- 134 Ruiz-Santana S, Ortiz E, Gonzalez B, Bolanos J, Ruiz-Santana AJ, Manzano JL. Stress-induced gastroduodenal lesions and total parenteral nutrition in critically ill patients: frequency, complications, and the value of prophylactic treatment. A prospective, randomized study. *Crit Care Med* 1991;19:887-91.
- 135 Zinner MJ, Zuidema GD, Smith PL. The prevention of upper gastrointestinal tract bleeding in patients in an intensive care unit. *Surg Gynecol Obstet* 1981;153:214-20.
- 136 Tryba M. Research on stress ulcer prophylaxis: wrong questions, wrong answers? *Crit Care Med* 1999;27:16-7.
- 137 Cook DJ, Reeve BK, Guyatt GH. Stress ulcer prophylaxis in critically ill patients. Resolving discordant meta-analyses. *JAMA* 1996;275:308-14.
- 138 Cook DJ, Fuller HD, Guyatt GH, Marshall JC, Leasa D, Hall R, Winton TL, Rutledge F, Todd TJ, Roy P, et al. Risk factors for gastrointestinal bleeding in critically ill patients. Canadian Critical Care Trials Group. *N Engl J Med* 1994;330:377-81.
- 139 Cook D, Heyland D, Griffith L, Cook R, Marshall J, Pagliarello J. Risk factors for clinically important upper gastrointestinal bleeding in patients requiring mechanical ventilation. Canadian Critical Care Trials Group. *Crit Care Med* 1999;27:2812-7.
- 140 Tryba M. Prophylaxis of stress ulcer bleeding. A meta-analysis. *J Clin Gastroenterol* 1991;13(Suppl 2):S44-55.
- 141 Therapeutic guidelines on stress ulcer prophylaxis. *Am J Health Syst Pharm* 1999;56:347-79.
- 142 Cook D, Heyland D, Griffith L, Cook R, Marshall J, Pagliarello J. Risk factors for clinically important upper gastrointestinal bleeding in patients requiring mechanical ventilation. Canadian Critical Care Trials Group. *Crit Care Med* 1999;27:2812-7.

- 143 Cook D, Guyatt G, Marshall J, Leasa D, Fuller H, Hall R, Peters S, Rutledge F, Griffith L, Mcleilan A, Wood G, Kirby A. A comparison of sucralfate and ranitidine for the prevention of upper gastrointestinal bleeding in patients requiring mechanical ventilation. Canadian Critical Care Trials Group. *N Engl J Med* 1998; 19:338:791-7.
- 144 Aanpreung P, Vanprapar N, Susiva C, Parkpreaw C, Boonyachart C. A randomized clinical trial comparing the efficacy of ranitidine and famotidine on intragastric acidity in critically ill pediatric patients. *J Med Assoc Thai* 1998;81:185-9.
- 145 Calvet X, Baigorri F, Duarte M, Saura P, Royo C, Joseph D, Mas A, Artigas A. Effect of ranitidine on gastric intramucosal pH in critically ill patients. *Intensive Care Med* 1998;24:12-7.
- 146 Ephgrave KS, Kleiman-Wexler R, Pfaller M, Booth BM, Reed D, Werkmeister L, Young. Effects of sucralfate vs antacids on gastric pathogens: results of a double-blind clinical trial. *S Arch Surg* 1998;133:251-7.
- 147 Ortiz JE, Sottile FD, Sigel P, Nasraway SA. Gastric colonization as a consequence of stress ulcer prophylaxis: a prospective, randomized trial. *Pharmacotherapy* 1998;18:486-91.
- 148 Levy MJ, Seelig CB, Robinson NJ, Ranney JE. Comparison of omeprazole and ranitidine for stress ulcer prophylaxis. *Dig Dis Sci* 1997;42:1255-9.
- 149 Kuusela AL, Ruuska T, Karikoski R, Laippata P, Ikonen RS, Janas M, Maki M. A randomized, controlled study of prophylactic ranitidine in preventing stress-induced gastric mucosal lesions in neonatal intensive care unit patients. *Crit Care Med* 1997;25:346-51.

- 150 Thomason MH, Payseur ES, Hakenewerth AM, Norton HJ, Mehta B, Reeves TR, Moore-Swartz MW, Robbins PI. Nosocomial pneumonia in ventilated trauma patients during stress ulcer prophylaxis with sucralfate, antacid, and ranitidine. *J Trauma* 1996;41:503-8.
- 151 Laggner AN, Tryba M, Georgopoulos A, Lenz K, Grimm G, Graninger W, Schneeweiss B, Drumi W. Oropharyngeal decontamination with gentamicin for long-term ventilated patients on stress ulcer prophylaxis with sucralfate? *Wien Klin Wochenschr* 1994;106:15-9.
- 152 Maier RV, Mitchell D, Gentiello L. Optimal therapy for stress gastritis. *Ann Surg* 1994;220:353-60.
- 153 Behrens R, Hofbeck M, Singer H, Scharf J, Rupprecht T. Frequency of stress lesions of the upper gastrointestinal tract in paediatric patients after cardiac surgery: effects of prophylaxis. *Br Heart J* 1994;72:186-9.
- 154 Cloud ML, Offen W. Continuous infusions of nizatidine are safe and effective in the treatment of intensive care unit patients at risk for stress gastritis. The Nizatidine Intensive Care Unit Study Group. *Scand J Gastroenterol* 1994;(Suppl 206):29-34.
- 155 Eddleston JM, Pearson RC, Holland J, Tooth JA, Vohra A, Doran BH. Prospective endoscopic study of stress erosions and ulcers in critically ill adult patients treated with either sucralfate or placebo. *Crit Care Med* 1994;22:1949-54.
- 156 Grau JM, Casademont J, Fernandez-Sola J, Cardellach F, Urbano-Marquez A. Prophylaxis of gastrointestinal tract bleeding in patients admitted to a general hospital ward. Comparative study of sucralfate and cimetidine. *Scand J Gastroenterol* 1993;28:244-8.

- 157 Pickworth KK, Falcone RE, Hoogeboom JE, Santanello SA. Occurrence of nosocomial pneumonia in mechanically ventilated trauma patients: a comparison of sucralfate and ranitidine. *Crit Care Med* 1993;21:1856-62.
- 158 Ryan P, Dawson J, Teres D, Celoria G, Navab F. Nosocomial pneumonia during stress ulcer prophylaxis with cimetidine and sucralfate. *Arch Surg* 1993;128:1353-7.
- 159 Vargas V, Castro J, Del Solar F, Sanhueza H, Aguila R, Luppi M, Fernandez D, Rojas M. Comparative study of famotidine vs ranitidine in critically ill patients in mechanical ventilation. *Rev Med Chil* 1993;121:746-51.
- 160 Martin LF, Booth FV, Karistadt RG, Silverstein JH, Jacobs DM, Hampsey J, Bowman SC, D'ambrosio CA, Rockhold FW. Continuous intravenous cimetidine decreases stress-related upper gastrointestinal hemorrhage without promoting pneumonia. *Crit Care Med* 1993;21:19-30.
- 161 Metz CA, Livingston DH, Smith JS, Larson GM, Wilson TH. Impact of multiple risk factors and ranitidine prophylaxis on the development of stress-related upper gastrointestinal bleeding: a prospective, multicenter, double-blind, randomized trial. The Ranitidine Head Injury Study Group. *Crit Care Med* 1993;21:1844-9.
- 162 Lopez-Herce J, Dorao P, Elola P, Delgado MA, Ruza F, Madero R. Frequency and prophylaxis of upper gastrointestinal hemorrhage in critically ill children: a prospective study comparing the efficacy of almagate, ranitidine, and sucralfate. The gastrointestinal hemorrhage Study Group. *Crit Care Med* 1992;20:1082-9.
- 163 Mahul P, Auboyer C, Jospe R, Ros A, Guerin C, El Khouri Z, Galliez M, Dumont A, Gaudin O. Prevention of nosocomial pneumonia in intubated patients: respective role of mechanical subglottic secretions drainage and stress ulcer prophylaxis. *Intensive Care Med* 1992;18:20-5.

- 164 Martin LF, Booth FV, Reines HD, Deysach LG, Kochman RL, Erhardt LJ, Geis GS. Stress ulcers and organ failure in intubated patients in surgical intensive care units. *Ann Surg* 1992; 215:332-7.
- 165 Apte NM, Karnad DR, Medhekar TP, Tilve GH, Morye S, Bhave GG. Gastric colonization and pneumonia in intubated critically ill patients receiving stress ulcer prophylaxis: a randomized, controlled trial. *Crit Care Med* 1992;20:590-3.
- 166 Eddleston JM, Vohra A, Scott P, Tooth JA, Pearson RC, McCloy RF, Morton AK, Doran BH. A comparison of the frequency of stress ulceration and secondary pneumonia in sucralfate- or ranitidine-treated intensive care unit patients. *Crit Care Med* 1991; 19:1491-6.
- 167 Simms HH, Demaria E, McDonald L, Peterson D, Robinson A, Burchard KW. Role of gastric colonization in the development of pneumonia in critically ill trauma patients: results of a prospective randomized trial. *J Trauma* 1991;31:531-6.
- 168 Estruch R, Pedrol E, Castells A, Masanes F, Marrades RM, Urbano-Marquez A. Prophylaxis of gastrointestinal tract bleeding with magaldrate in patients admitted to a general hospital ward. *Scand J Gastroenterol* 1991;26:819-26.
- 169 Lamothe PH, Rao E, Serra AJ, Castellano J, Woronick CL, McNicholas KW, Lemole GM. Comparative efficacy of cimetidine, famotidine, ranitidine, and mylanta in postoperative stress ulcers. Gastric pH control and ulcer prevention in patients undergoing coronary artery bypass graft surgery. *Gastroenterology* 1991; 100:1515-20.
- 170 Kitler ME, Hays A, Enterline JP, Allo M, Zuidema GD.
Preventing postoperative acute bleeding of the upper part of the gastrointestinal tract. *Surg Gynecol Obstet.* 1990 Nov;171(5):366-72.

- 171 Flaherty J, Nathan C, Kabins SA, Weinstein RA. Pilot trial of selective decontamination for prevention of bacterial infection in an intensive care unit. *J Infect Dis* 1990;162:1393-7.
- 172 Reusser P, Gyr K, Scheidegger D, Buchmann B, Buser M, Zimmerli W. Prospective endoscopic study of stress erosions and ulcers in critically ill neurosurgical patients: current incidence and effect of acid-reducing prophylaxis. *Crit Care Med* 1990;18:270-4.
- 173 Zinner MJ, Rypins EB, Martin LR, Jonasson O, Hoover EL, Swab EA, Fakouhi TD. Misoprostol versus antacid titration for preventing stress ulcers in postoperative surgical ICU patients. *Ann Surg*. 1989 Nov;210(5):590-5.
- 174 Garvey BM, McCambley JA, Tuxen DV. Effects of gastric alkalization on bacterial colonization in critically ill patients. *Crit Care Med*. 1989 Mar;17(3):211-6.
- 175 Chaibou M, Tucci M, Dugas MA, Farrell CA, Proulx F, Lacroix J. Clinically significant upper gastrointestinal bleeding acquired in a pediatric intensive care unit: a prospective study. *Pediatrics* 1998;102:933-8.
- 176 Harrison AM, Lugo RA, Vernon DD. Gastric pH control in critically ill children receiving intravenous ranitidine. *Crit Care Med* 1998;26:1433-6.
- 177 Ellison RT, Perez-Perez G, Welsh CH, Blaser MJ, Riester KA, Cross AS, Donta ST, Peduzzi P. Risk factors for upper gastrointestinal bleeding in intensive care unit patients: role of helicobacter pylori. Federal Hyperimmune Immunoglobulin Therapy Study Group. *Crit Care Med* 1996;24:1974-81.
- 178 Cook DJ, Reeve BK, Scholes LC. Histamine-2-receptor antagonists and antacids in the critically ill population: stress ulceration versus nosocomial pneumonia. *Infect Control Hosp Epidemiol* 1994; 15:437-42.

- 179 Cochran EB, Phelps SJ, Tolley EA, Stidham GL. Prevalence of, and risk factors for, upper gastrointestinal tract bleeding in critically ill pediatric patients. *Crit Care Med* 1992;20:1519-23.
- 180 Kuusela AL, Maki M, Ruuska T, Laippala P. Stress-induced gastric findings in critically ill newborn infants: frequency and risk factors. *Intensive Care Med* 2000;26:1501-6.
- 181 Zeitsman D, Rowland M, Shanavas Z, Kerstein MD. Is the incidence of hemorrhagic stress ulceration in surgical critically ill patients affected by modern antacid prophylaxis? *Am Surg* 1996;62:1010-3.
- 182 Ben-Menachem T, McCarthy BD, Fogel R, Schiffman RM, Patell RV, Zarowitz BJ, Nerenz DR, Bresalier RS. Prophylaxis for stress-related gastrointestinal hemorrhage: a cost effectiveness analysis. *Crit Care Med* 1996;24:338-45.
- 183 Kuusela AL. Long-term gastric pH monitoring for determining optimal dose of ranitidine for critically ill preterm and term neonates. *Arch Dis Child Fetal Neonatal*. 1998;78:151-3.
- 184 Baghale AA, Moitahedzadeh M, Levine RL, Fromm RE JR, Guntupali KK, Opekun AR JR. Comparison of the effect of intermittent administration and continuous infusion of famotidine on gastric pH in critically ill patients: results of a prospective, randomized, crossover study. *Crit Care Med* 1995;23:687-91.
- 185 Heiselman DE, Hulisz DT, Fricker R, Bredle DL, Black LD. Randomized comparison of gastric pH control with intermittent and continuous intravenous infusion of famotidine in ICU patients. *Am J Gastroenterol* 1995;90:277-9.
- 186 Zandstra DF, Stoutenbeek CP. The virtual absence of stress-ulceration related bleeding in ICU patients receiving prolonged mechanical ventilation without any prophylaxis. A prospective cohort study. *Intensive Care Med* 1994;20:335-40.

- 187 Osteyee JL, Banner W Jr. Effects of two dosing regimens of intravenous ranitidine on gastric pH in critically ill children. *Am J Crit Care* 1994;3:267-72.
- 188 Gedeit RG, Weigle CG, Havens PL, Werlin SL. Control and variability of gastric pH in critically ill children. *Crit Care Med* 1993; 21;1850-5.
- 189 Geus WP, Vinks AA, Smith SJ, Westra P, Lamers CB. Comparison of two intravenous ranitidine regimens in a homogeneous population of intensive care unit patients. *Aliment Pharmacol Ther* 1993;7:451-7.
- 190 Moore JG, Clemmer TP, Taylor S, Bishop AL, Maggio S. Twenty-four-hour intragastric pH patterns in ICU patients on ranitidine. *Dig Dis Sci* 1992;37:1802-9.
- 191 Heyland D, Bradley C, Mandeli LA. Effect of acidified enteral feedings on gastric colonization in the critically ill patient. *Crit Care Med* 1992;20:1388-94.
- 192 Tsuruoka K, Tsuruoka Y, Yoshimura M, Imai K, Sekiguchi S, Mise J, Asai Y, Nago N, Igarashi M. Evidence based general practice. Drug treatment in general practice in Japan is evidence based. *BMJ*. 1996;313:114;
- 193 Chikwe J. Evidence based general practice. Findings of study should prompt debate. *BMJ*. 1996;313:114;
- 194 Geddes JR, Game D, Jenkins NE, Peterson LA, Pottinger GR, Sackett DL.
What proportion of primary psychiatric interventions are based on evidence from randomised controlled trials? *Qual Health Care*. 1996;5:215-7.

- 195 Eisenberg MJ, Califf RM, Cohen EA, Adelman AG, Mark DB, Topol EJ. Use of evidence-based medical therapy in patients undergoing percutaneous coronary revascularization in the United States, Europe, and Canada. Coronary Angioplasty Versus Excisional Atherectomy Trial (CAVEAT-I) and Canadian Coronary Atherectomy Trial (CCAT) investigators. *Am J Cardiol.* 1997;79:867-72.
- 196 Jemec GB, Thorsteinsdottir H, Wulf HC. Evidence-based dermatologic out-patient treatment. *Int J Dermatol* 1998;37:850-4.
- 197 Rudolf MC, Lyth N, Bundle A, Rowland G, Kelly A, Bosson S, Garner M, Guest P, Khan M, Thazin R, Bennett T, Damman D, Cove V, Kaur V. A search for the evidence supporting community paediatric practice. *Arch Dis Child* 1999 ;80:257-61.
- 198 Shakib S, Phillips PA. The Australian Centre for Evidence-based Clinical Practice generic audit tool: Auditmaker for health professionals. *J Eval Clin Pract.* 2003;9:259-63.
- 199 Baker R, Fraser RC. Development of review criteria: linking guidelines and assessment of quality. *BMJ.* 1995;311:370-3.
- 200 Baker R, Fraser RC. Goals and methods of audit should be reappraised. Evidence of effectiveness of audit in improving care is growing. *BMJ.* 1996;313:1405.
- 201 Merritt TA, Palmer D, Bergman DA, Shiono PH. Clinical practice guidelines in pediatric and newborn medicine: implications for their use in practice. *Pediatrics.* 1997;99:100-14.
- 202 Alderson P, Bunn F, Lefebvre C, Li WP, Li L, Roberts I, Schierhout G. Human albumin solution for resuscitation and volume expansion in critically ill patients. *Cochrane Database Syst Rev* 2002;(1):CD001208.
- 203 Petros A, Schindler M, Pierce C, Jacobe S, Mok Q. Human albumin administration in critically ill patients. Evidence needs to be shown in paediatrics. *BMJ* 1998;317:882.

- 204 Nel MR. Human albumin administration in critically ill patients. Critical analysis of original studies has to take place. *BMJ* 1998;317:882
- 205 Nadel S, Marriage S, De Munter C, Britto J, Habibi P, Levin M. Human albumin administration in critically ill patients. Review did not provide recommendations for alternative treatment. *BMJ* 1998;317:882-3.
- 206 Allison SP, Lobo DN. Debate: Albumin administration should not be avoided. *Crit Care* 2000;4:147-50.
- 207 Bunn F, Alderson P, Hawkins V. Colloid solutions for fluid resuscitation. *Cochrane Database Syst Rev* 2003;(1):CD001319.
- 208 Alderson P, Schierhout G, Roberts I, Bunn F. Colloids versus crystalloids for fluid resuscitation in critically ill patients. *Cochrane Database Syst Rev* 2000;(2):CD000567.
- 209 Wilkes MM, Navickis RJ. Patient survival after human albumin administration. A meta-analysis of randomized, controlled trials. *Ann Intern Med* 2001;135:149-64.
- 210 Cox P, Sheridan R, editors. Current concepts in pediatric critical care.. Proceedings of the 31st Critical Care Congress; 2002 Jan 25-26; San Diego, United States.
- 211 Sun X, Iles M, Weissman C. Physiologic variables and fluid resuscitation in the postoperative intensive care unit patient. *Crit Care Med* 1993;21:555-61.
- 212 Blunt MC, Nicholson JP, Park GR. Serum albumin and colloid osmotic pressure in survivors and nonsurvivors of prolonged critical illness. *Anaesthesia* 1998;53:755-61.
- 213 Carcillo JA, Fields AI; American College of Critical Care Medicine Task Force Committee Members. Clinical practice parameters for hemodynamic support of pediatric and neonatal patients in septic shock. *Crit Care Med*. 2002;30:1365-78

- 214 Niermeyer S, Kattwinkel J, Van Reempts P, Nadkarni V, Phillips B, Zideman D, Azzopardi D, Berg R, Boyle D, Boyle R, Burchfield D, Carlo W, Chameides L, Denson S, Fallat M, Gerardi M, Gunn A, Hazinski MF, Keenan W, Knaebel S, Milner A, Perlman J, Saugstad OD, Schleien C, Solimano A, Speer M, Toce S, Wiswell T, Zaritsky A. International Guidelines for Neonatal Resuscitation: An excerpt from the Guidelines 2000 for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care: International Consensus on Science. Contributors and Reviewers for the Neonatal Resuscitation Guidelines. *Pediatrics*. 2000;106:E29.
- 215 Chalasani N, Gorski JC, Horlander JC Sr, Craven R, Hoen H, Maya J, Brater DC. Effects of albumin/furosemide mixtures on responses to furosemide in hypoalbuminemic patients. *J Am Soc Nephrol* 2001;12:1010-6.
- 216 Kirchner KA, Voelker JR, Brater DC. Intratubular albumin blunts the response to furosemide-A mechanism for diuretic resistance in the nephrotic syndrome. *J Pharmacol Exp Ther*. 1990;252:1097-101.
- 217 Chalasani N, Gorski JC, Horlander JC Sr, Craven R, Hoen H, Maya J, Brater DC. Effects of albumin/furosemide mixtures on responses to furosemide in hypoalbuminemic patients. *J Am Soc Nephrol* 2001;12:1010-6.
- 218 Flancbaum L, Dick M, Choban PS, Dasta JP. Effects of low-dose dopamine on urine output in oliguric, critically ill, renal transplant patients. *Clin Transplant* 1998;12:256-9.
- 219 Hoogenberg K, Smit AJ, Girbes AR. Effects of low-dose dopamine on renal and systemic hemodynamics during incremental norepinephrine infusion in healthy volunteers. *Crit Care Med* 1998;26:260-5.
- 220 Kellum JA, M Decker J. Use of dopamine in acute renal failure: a meta-analysis. *Crit Care Med* 2001;29:1526-31.

- 221 Galley HF. Renal-dose dopamine: will the message now get through? *Lancet* 2000;356:2112-3.
- 222 Romano G, Ferraccioli G. Usefulness of dopamine treatment in non-oliguric renal failure. *Lancet*. 2001;357:960.
- 223 Committee on Fetus and Newborn and Fetus and Newborn Committee. Prevention and Management of Pain and Stress in the Neonate. *Pediatrics* 2000; 105:454 - 61.
- 224 Anand KJ, Arnold JH. Opioid tolerance and dependence in infants and children. *Crit Care Med* 1994;22:334-42.
- 225 Castro MCFZ, Guinsburg R, Almeida MFB, Peres CA; Yanaguibashi G; Kopelman BI. Perfil da indicação de analgésicos opióides em recém-nascidos em ventilação pulmonar mecânica. *J Pediatr (Rio J)* 2003; 79(1): 41
- 226 Partridge JC, Wall SN. Analgesia for dying infants whose life support is withdrawn or withheld. *Pediatrics* 1997;99:76-9.
- 227 John M, Freeman, Eileen P, Vining G. Oranges and Apples: Sedation and Analgesia. *Pediatrics* 2000;106:1519.
- 228 Pokela ML, Olkkola KT, Seppala T, Koivisto M. Age-related morphine kinetics in infants. *Dev Pharmacol Ther* 1993;20:26-34.
- 229 Surgenor SD, Hampers MJ, Corwin HL. Optimizing red blood cell transfusion practice. In Vincent JL editor. Yearbook of intensive care and emergency medicine. Berlin: Springer; 2001. p.117
- 230 Maier RF, Sonntag J, Walka MM, Liu G, Metze BC, Obladen M. Changing practices of red blood cell transfusions in infants with birth weights less than 1000 g. *J Pediatr* 2000;136:220-4.
- 231 Hebert PC, Blajchman MA, Cook DJ, Yetisir E, Wells G, Marshall J, Schweitzer I. Transfusion Requirements in Critical Care Investigators for the Canadian Critical Care Trials Group. Do blood transfusions improve outcomes related to mechanical ventilation? *Chest* 2001;119:1850-7.

- 232 Spence RK, Carson JA, Poses R, McCoy S, Pello M, Alexander J, Popovich J, Norcross E, Camishion RC. Elective surgery without transfusion: influence of preoperative hemoglobin level and blood loss on mortality. *Am J Surg* 1990;159:320-4.
- 233 Dietrich KA, Conrad SA, Hebert CA, Levy GL, Romero MD. Cardiovascular and metabolic response to red blood cell transfusion in critically ill volume-resuscitated nonsurgical patients. *Crit Care Med* 1990;18:940-4.
- 234 Mathru M, Kleinman B, Blakeman B, Sullivan H, Kumar P, Dries DJ. Myocardial metabolism and adaptation during extreme hemodilution in humans after coronary revascularization. *Crit Care Med* 1992;20:1420-5.
- 235 Lorente JA, Landin L, De Pablo R, Renes E, Rodriguez-Diaz R, Liste D.
Effects of blood transfusion on oxygen transport variables in severe sepsis. *Crit Care Med*. 1993;21:1312-8.
- 236 Nelson AH, Fleisher LA, Rosenbaum SH. Relationship between postoperative anemia and cardiac morbidity in high-risk vascular patients in the intensive care unit. *Crit Care Med* 1993;21:860-6.
- 237 Mann R, Heimbach DM, Engrav LH, Foy H. Changes in transfusion practices in burn patients. *J Trauma* 1994;37:220-2.
- 238 Carson JL, Duff A, Poses RM, Berlin JA, Spence RK, Trout R, Noveck H, Strom BL. Effect of anaemia and cardiovascular disease on surgical mortality and morbidity. *Lancet* 1996;348:1055-60.
- 239 Schonhofer B, Wenzel M, Geibel M, Kohler D. Blood transfusion and lung function in chronically anemic patients with severe chronic obstructive pulmonary disease. *Crit Care Med* 1998;26:1824-8.
- 240 Adams RJ. Lessons from the Stroke Prevention Trial in Sickle Cell Anemia (STOP) study. *J Child Neurol* 2000;15:344-9.

- 241 Salem-Schatz SR, Avorn J, Soumerai SB. Influence of clinical knowledge, organizational context, and practice style on transfusion decision making. Implications for practice change strategies. *JAMA* 1990;264:476-83.
- 242 Vincent JL, Baron JF, Reinhart K, Gattinoni L, Thijs L, Webb A, Meier-Hellmann A, Nollet G, Peres-Bota D. ABC (Anemia and Blood Transfusion in Critical Care) Investigators. Anemia and blood transfusion in critically ill patients. *JAMA* 2002;288:1499-507.
- 243 Soubasi V, Kremenopoulos G, Diamandi E, Tsantali C, Tsakiris D. In which neonates does early recombinant human erythropoietin treatment prevent anemia of prematurity? Results of a randomized, controlled study. *Pediatr Res* 1993;34:675-9.
- 244 Williamson P, Griffiths G, Norfolk D, Levene M. Blood transfusions and human recombinant erythropoietin in premature newborn infants. *Arch Dis Child Fetal Neonatal Ed* 1996;75:F65-8.
- 245 Corwin HL, Gettinger A, Rodriguez RM, Pearl RG, Gubler KD, Enny C, Colton T, Corwin MJ. Efficacy of recombinant human erythropoietin in the critically ill patient: a randomized, double-blind, placebo-controlled trial. *Crit Care Med* 1999;27:2346-50.
- 246 Tryba M, Cook D. Current guidelines on stress ulcer prophylaxis. *Drugs* 1997 ;54:581-96.
- 247 Fennerty MB. Pathophysiology of the upper gastrointestinal tract in the critically ill patient: rationale for the therapeutic benefits of acid suppression. *Crit Care Med* 2002;30:S351-5.
- 248 Cook DJ, Griffith LE, Walter SD, Guyatt GH, Meade MO, Heyland DK, Kirby A, Tryba M, Canadian Critical Care Trials Group. The attributable mortality and length of intensive care unit stay of clinically important gastrointestinal bleeding in critically ill patients. *Crit Care* 2001;5:368-75.

- 249 Freedman MD. Stress ulcer prevention and histamine-2-receptor antagonists: leaps of faith or back to the drawing board? *Crit Care Med* 1999;27:13-4.
- 250 Abraham E. Acid suppression in a critical care environment: state of the art and beyond. *Crit Care Med* 2002;30:S349-50.
- 251 MacLaren R, Jarvis CL, Fish DN. Use of enteral nutrition for stress ulcer prophylaxis. *Ann Pharmacother* 2001;35:1614-23.
- 252 Marik PE. *Handbook of evidence-based critical care*. New York: Springer; 2001. p.421
- 253 Singer M, Little R. ABC of intensive care. Cutting edge. *BMJ*. 1999;319:501-4.
- 254 Portaria no. 3432 de 1998, Pub. D.O. da União Séc I, no.154 (Aug. 08, 1998) p.109-10.
- 255 Souza DC. Avaliação da estrutura das unidades de terapia intensiva pediátrica e neonatal da cidade de São Paulo [dissertation]. São Paulo (SP): Faculdade de Medicina da Universidade de São Paulo; 2003.
- 256 Williamson LM, Heptonstall J, Soldan K. A SHOT in the arm for safer blood transfusion. *BMJ* 1996;313:1221-2.
- 257 Jamtvedt G, Young JM, Kristoffersen DT, Thomson O'Brien MA, Oxman AD.
Audit and feedback: effects on professional practice and health care outcomes.
Cochrane Database Syst Rev. 2003;(3):CD000259.
- 258 Burch K, Rhine W, Baker R, Litman F, Kaempf JW, Schwarz E, Sun S, Payne NR, Sharek PJ. Implementing potentially better practices to reduce lung injury in neonates. *Pediatrics*. 2003;111:e432-6.
- 259 Sueta CA, Massing MW, Chowdhury M, Biggs DP, Simpson RJ Jr. Undertreatment of hyperlipidemia in patients with coronary artery disease and heart failure. *J Card Fail* 2003;9:36-41.

- 260 Osborne CA, Hooper R, Swift CG, Jackson SH. Explicit, evidence-based criteria to assess the quality of prescribing to elderly nursing home residents. *Age Ageing* 2003;32:102-8.
- 261 Pazart L, Janvier G, Durocher A, Carpentier F, Marty J. [A study of practice patterns of albumin administration in surgical patients by anesthetists]. *Ann Fr Anesth Reanim* 1996;15:424-7.
- 262 Lam NP, Le PD, Crawford SY, Patel S. National survey of stress ulcer prophylaxis. *Crit Care Med*. 1999;27:98-103.
- 263 Moyer VA, Gist AK, Elliott EJ. Is the practice of paediatric inpatient medicine evidence-based? *J Paediatr Child Health*. 2002;38(4):347-51.
- 264 Djulbegovic B, Loughran TP Jr, Hornung CA, Kloecker G, Efthimiadis EN, Hadley TJ, Englert J, Hoskins M, Goldsmith GH. The quality of medical evidence in hematology-oncology. *Am J Med* 1999;106:198-205.
- 265 Lai TY, Wong VW, Leung GM. Is ophthalmology evidence based? A clinical audit of the emergency unit of a regional eye hospital. *Br J Ophthalmol* 2003 ;87(4):385-90.
- 266 Saxena S, Weiner JM, Rabinowitz A, Fridey J, Shulman IA, Carmel R. Transfusion practice in medical patients. *Arch Intern Med* 1993;153:2575-80.
- 267 Corwin HL, Parsonnet KC, Gettinger A. RBC transfusion in the ICU. Is there a reason? *Chest* 1995;108:767-71.
- 268 Koyama H, Matsui K, Goto M, Sekimoto M, Maeda K, Morimoto T, Hira K, Fukui T. In-patient interventions supported by results of randomized controlled trials in Japan. *Int J Qual Health Care* 2002;14:119-25.

- 269 Tsuruoka K, Tsuruoka Y, Yoshimura M, Imai K, Sekiguchi S, Mise J, Asai Y, Nago N, Igarashi M. Evidence based general practice. Drug treatment in general practice in Japan is evidence based. *BMJ*. 1996;313:114
- 270 Howes N, Chagla L, Thorpe M, McCulloch P. Surgical practice is evidence based. *Br J Surg* 1997;84:1220-3.
- 271 Slim K, Lescure G, Voitellier M, Ferrandis P, Le Roux S, Dumas PJ, Lere JM, Patouillard P, Caburet A, Baudet B, Rolet JP, Prat M, Pezet D, Chipponi J. [Is laparoscopic surgical practice "factual" (evidence based)? Results of a prospective regional survey]. *Presse Med* 1998;27:1829-33.