

**UNIVERSIDADE DE SÃO PAULO
ESCOLA DE ENFERMAGEM**

CHRISTIANE BORGES DO NASCIMENTO CHOFAKIAN

**CONTRACEPTIVE DISCONTINUATION AND ITS
RELATION TO EMERGENCY CONTRACEPTION USE
AMONG UNDERGRADUATE WOMEN**

**SÃO PAULO
2017**

CHRISTIANE BORGES DO NASCIMENTO CHOFAKIAN

**CONTRACEPTIVE DISCONTINUATION AND ITS
RELATION TO EMERGENCY CONTRACEPTION USE
AMONG UNDERGRADUATE WOMEN**

Tese apresentada ao Programa de Pós-Graduação em Enfermagem da Escola de Enfermagem da Universidade de São Paulo para obtenção do título de Doutor em Ciências.

Área de concentração: Cuidados em Saúde

Orientadora:

Prof^a. Dr^a. Ana Luiza Vilela Borges

Supervisora do doutorado sanduíche:

Prof^a. Dr^a. Caroline Moreau

**São Paulo
2017**

AUTORIZO A REPRODUÇÃO E DIVULGAÇÃO TOTAL OU PARCIAL DESTE TRABALHO, POR QUALQUER MEIO CONVENCIONAL OU ELETRÔNICO, PARA FINS DE ESTUDO E PESQUISA, DESDE QUE CITADA A FONTE.

Assinatura: _____ Data: ____ / ____ / _____

Catálogo na Publicação (CIP)
Biblioteca “Wanda de Aguiar Horta”
Escola de Enfermagem da Universidade de São Paulo

Chofakian, Christiane Borges do Nascimento
Contraceptive discontinuation and its relation to emergency contraception use among undergraduate women / Christiane Borges do Nascimento Chofakian. – São Paulo, 2017.

277 p.

Tese (Doutorado) - Escola de Enfermagem da Universidade de São Paulo.

Orientadora: Prof^a Dr^a Ana Luiza Vilela Borges

Área de concentração: Cuidados em saúde

1. Contraception 2. Contraceptive discontinuation
3. Undergraduate students 4. Emergency contraception
5. Sexual and Reproductive Health

I. Título.

Nome: Christiane Borges do Nascimento Chofakian

Título: Contraceptive discontinuation and its relation to emergency contraception use among undergraduate women

Tese apresentada ao Programa de Pós-Graduação em Enfermagem da Escola de Enfermagem da Universidade de São Paulo para obtenção do título de Doutor em Ciências.

Aprovado em: ___/___/___

BANCA EXAMINADORA

Prof.^a Dr.^a Ana Luiza Vilela Borges Instituição: _____

Julgamento: _____ Assinatura: _____

Prof. Dr. _____ Instituição: _____

Julgamento: _____ Assinatura: _____

Prof. Dr. _____ Instituição: _____

Julgamento: _____ Assinatura: _____

Prof. Dr. _____ Instituição: _____

Julgamento: _____ Assinatura: _____

Prof. Dr. _____ Instituição: _____

Julgamento: _____ Assinatura: _____

DEDICATION

To my dear husband Carlos José Chofakian, who has been a constant source of steadfast support. For your unconditional love, company, patience and encouragement, which were essential to the progress and finalization of this thesis. Always present in my life, specially in the moments when I most needed it. Your prayers and dedicated love enabled me to remain committed throughout my academic journey. I am truly thankful for having you in my life.

To my beloved family, who have always loved me unconditionally and whose good examples of determination and perseverance have taught me to work hard for the things that I aspire to achieve.

ACKNOWLEDGMENTS

A dissertation is not the outcome of the struggles of one person. Many people contributed to its development. At the moment, I take the opportunity to acknowledge those who have made some impact on my doctoral journey and achievement.

Firstly, I would like to thank to the One who makes all things possible, and for enabling me to complete this thesis, Jesus Christ – to him I give all glory, honor and praise. I now understand the depth of this journey.

I must thank my husband, Carlos José Chofakian, who supported me through the entire doctoral process, since my master degree. He has always been my truly love and my best friend, always offering me your “shoulder”, holding me in the most difficult times, being my safe harbor, and never let me give up even when I thought I should. He always provided me encouragement, was patient, and always got me to laugh even when it was a most difficult time. My true love, you are undoubtedly the best gift that God gave me. Thank you so much for sharing in my struggles and successes each day – you got me through this. I will always love you in all my lifes.

I am sincerely thankful to my advisor, my inspiration, teacher and mentor, Profa. Ana Luiza Vilela Borges. My understanding of sexual and reproductive health has broadened because of you. You always emphasized the value of hard work and dedication to academic excellence. Regardless of the challenges I faced during graduate school you were always available to offer good advices, inspiration and constructive criticism over the past six years, all of which permitted me to persist focused as a student. It is impossible to repay you for all you have done for me. For me your wisdom transcends the academic teachings, for me you represent the balance between personal and professional life, and you have all my respect for it. More than my supervisor, I consider you as a great friend that I intend to take for the rest of my life, outside the academic walls.

I would also like to sincerely thank to Profa. Caroline Moreau. I believe that you never imagined that there was someone in Brazil who always read your papers. Meeting

you and having the opportunity to work with you in Baltimore was something I never imagined in my life. It was, without a doubt, one of the greatest professional experiences that I had in my life. You showed me a world that I never imagined existed. You represent to me a strong woman, clever, with a lot of energy and determination in what you do. I will never forget your support, for exposing me to cutting edge techniques in statistics, and I will always be grateful for it. I hope one day transmit all the knowledge learned from you, and give someone the same opportunity you gave me one day. You have a friend who really admires you in Brazil.

I extend my appreciation to Profa. Elizabeth Fujimori for the inspiration and unlimited encouragement. You have been following my academic journey and my growth for some years, and I am privileged to share this entire process with you. Thank you for sharing your teachings, for your friendship, for listening me in many times, for all invaluable support, help, and advice throughout the entire process of researching.

I would like to truly thank to my father Sergio and my mother Cecilia Maria even though she is no longer among us, for teaching me that anything is impossible and giving me the confidence to believe it. You have been a tower of strength and words cannot fully express my gratitude. There are no words to express my love for you. To Tatiana and Simone, thank you for always encouraging me and being the best sisters I could ever ask for. Even though we were miles apart, you always accompanying me on this journey and always supported me. I truly love you both.

I would like to thank to my husband's family, specially to my in-laws Ana Maria e Karekin, and to my sister-in-law's Renata and Ana Vitória, who always believed in me, for supporting me in all moments, for your assistance and encouragement. Thank you very much for your generosity, commitment, understanding, care and support that you gave me through this path.

I am also grateful to all my friends for their interest and insights on my dissertation. Special thanks go to the NEPESC Group, my friends at School of Nursing, and the friends that I made at Johns Hopkins University, for all support in particular and different moments in my life. I would particularly like to thank Osmary, Karina, Ana

Paula, Áurea, Luciana, Claudia, Clariana, Natália, Patrícia, all of you inspired me, your support helped propel me to the finish line.

Finally, I particularly wish to thank the 1,679 young women who participated in my study and so openly and shared their lives with me. This study would not have been possible without their willingness to participate.

EPIGRAPH

*“We are shaped by our thoughts; we become what we think.
When the mind is pure, joy follows like a shadow that never leave”.*

(Buddha)

Chofakian CBN. Contraceptive discontinuation and its relation to emergency contraception use among undergraduate women [thesis]. São Paulo: Escola de Enfermagem, Universidade de São Paulo; 2017.

ABSTRACT

Introduction: Contraceptive use dynamics is relevant to undergraduate students as they present high educational and professional aspirations, which affects reproductive intention. Also, they are mostly single, so they alternate the contraception according to their relationships. In that case, emergency contraception is an option, mainly in situations of discontinuation. However, little is known about discontinuation and its relation to the use of emergency contraception in Brazil.

Objective: To analyze the frequency and correlates of contraceptive discontinuation within 12-months; to assess emergency contraception use after discontinuation, and evaluate discontinuation after emergency contraception use. **Methods:** We conducted a 12-month retrospective cohort study on a sample of undergraduate women at University of São Paulo, Brazil. Students were selected by simple random sampling without replacement (n=1,679). Data were collected online using a self-administered questionnaire. In Stata 14.2, we used multinomial and multivariate logistic regression, and generalized estimating equation to analyze the data. **Results:** First, we observed that women with casual relationships, lower socioeconomic status, enrolled in Human and Health Sciences programs, with less years of sexual experience, with multiple sexual partners in lifetime, and who use less effective method were more likely to discontinuation one or several times. Second, pill and condom users who had casual relationships, with lower socioeconomic status, and who had previous pregnancy were more likely to discontinue and to abandon or switch to a less effective method. Third, a significant proportion of women did not use emergency contraception after discontinuing or abandoning contraception. Fourth, emergency contraception was mostly used after inconsistent use of contraception. Fifth, women who used contraception prior to emergency contraception use, had no religion, were in stable relationships, and had only one sexual partner were more likely to use contraception after emergency contraception. Sixth, women with stable relationships, from lower socioeconomic status, enrolled in Human Sciences programs, and who had one sexual partner were more likely to switch to a less effective method after emergency contraception use. Lastly, few women presented gaps in contraception within 30-days after emergency contraception use. **Conclusions:** Discontinuation does differ by type of method. Emergency contraception is underutilized after discontinuation. Partnership has an important influence on contraceptive use dynamics. Also, educational background, socioeconomic status, and number of lifetime sexual partners are important characteristics that should be considered when implementing family planning programs focused on young women.

Keywords: Contraception; Contraceptive discontinuation; Undergraduate students; Emergency contraception; Sexual and Reproductive Health.

Chofakian CBN. Contraceptive discontinuation and its relation to emergency contraception use among undergraduate women [thesis]. São Paulo: Escola de Enfermagem, Universidade de São Paulo; 2017.

RESUMO

Introdução: A dinâmica do uso dos métodos contraceptivos é importante entre os universitários, pois estes apresentam altas aspirações educacionais e profissionais, o que afeta a intenção reprodutiva. Ainda, por serem na maioria solteiros, os jovens alternam os métodos de acordo com o tipo de relacionamento. Neste contexto, a anticoncepção de emergência é uma opção, sobretudo nos casos de descontinuidades. Porém, pouco se sabe sobre as descontinuidades e sua relação com o uso da anticoncepção de emergência no Brasil. **Objetivo:** Analisar a frequência e os determinantes da descontinuidade contraceptiva em um período de 12 meses; avaliar o uso da anticoncepção de emergência após as descontinuidades, e avaliar as descontinuidades após o uso da anticoncepção de emergência. **Métodos:** Estudo de coorte retrospectivo, realizado com amostra probabilística de mulheres universitárias da Universidade de São Paulo. As alunas foram selecionadas por amostragem aleatória simples sem reposição (n=1.679). Os dados foram coletados através de um questionário de autopreenchimento respondido online. No Stata 14.2, os dados foram analisados por meio de regressão logística multinomial e multivariada, e equações de estimação generalizadas. **Resultados:** Primeiro, observou-se que as jovens com relacionamentos casuais, com menor nível socioeconômico, matriculadas nos cursos de Humanas e Ciências da Saúde, com menos anos de experiência sexual, com múltiplos parceiros sexuais e que usavam métodos menos eficazes, apresentaram maior probabilidade de descontinuar uma ou várias vezes. Segundo, as usuárias de pílula e preservativo que tinham relacionamentos casuais, com menor nível socioeconômico e que tiveram gravidez anterior, foram mais propensas a descontinuar, abandonar ou mudar para um método menos eficaz. Terceiro, uma proporção significativa de mulheres não usou anticoncepção de emergência após descontinuarem ou abandonarem o método. Quarto, a anticoncepção de emergência foi mais utilizada após inconsistências no uso do método. Quinto, as jovens que usaram um método antes do uso da anticoncepção de emergência, sem religião, com um relacionamento estável, e que tiveram um parceiro sexual na vida, foram mais propensas a usar contracepção após a anticoncepção de emergência. Sexto, as jovens com relacionamento estável, de nível socioeconômico mais baixo, matriculadas nos cursos de Humanas e que tiveram um parceiro sexual na vida tiveram maior probabilidade de mudar para um método menos eficaz após o uso da anticoncepção de emergência. Por fim, poucas jovens apresentaram descontinuidades dentro de 30 dias após o uso da anticoncepção de emergência. **Conclusões:** A descontinuidade contraceptiva difere por tipo de método. A anticoncepção de emergência é subutilizada após as descontinuidades. A parceria influencia na dinâmica do uso de contraceptivos. Ainda, aspectos educacionais, nível socioeconômico e número de parceiros sexuais são características importantes a serem consideradas na implementação de programas de planejamento familiar focados em mulheres jovens.

Descritores: Anticoncepção; Descontinuidade contraceptiva; Estudantes universitários; Anticoncepção de emergência; Saúde Sexual e Reprodutiva.

LIST OF ILLUSTRATIONS

FIGURES

- Figure 1 – Conceptual framework illustrating contraceptive discontinuation and their relation to the use of emergency contraception.....58
- Figure 2 – Map showing the distribution of the campuses at University of São Paulo, Brazil in São Paulo State62
- Figure 3 – Flowchart of organizational structure of the women interviewed, according to sexual behavior and contraception use. São Paulo, 2015.....64
- Figure 4 – Number of women discontinuing their method in the last 12-months, by measure of discontinuation and type of method used at the start of the contraceptive calendar (baseline). São Paulo, 2015**Error! Bookmark not defined.**
- Figure 5 – Number of monthly of contraception that were discontinued during the calendar year, by measure of discontinuation and type of method used at the start of the contraceptive calendar (baseline). São Paulo, 2015**Error! Bookmark not c**
- Figure 6 – Probability of contraceptive discontinuation for any reason by method type (Kaplan-Meier estimates). São Paulo, Brazil, 2015**Error! Bookmark not defined.**
- Figure 7 – Probability of contraceptive discontinuation for method-related reasons by method type (Kaplan-Meier estimates). São Paulo, Brazil, 2015**Error! Bookmark not defi**
- Figure 8 – Probability of abandonment or switching to less effective method reasons by method type (Kaplan-Meier estimates). São Paulo, Brazil, 2015**Error! Bookmark not c**
- Figure 9 – Using of emergency contraception in context or not of contraceptive discontinuation for method-related reasons. São Paulo, 2015**Error! Bookmark not define**
- Figure 10 – Emergency contraception use in context of use and non-use of contraception. São Paulo, 201.....**Error! Bookmark not defined.**

FRAMES

- Frame 1 – Summary of three chapters51
- Frame 2 – Description of the survey questionnaire65
- Frame 3 – Coding of the dependent variables69
- Frame 4 – Coding of the dependent variables assessing EC use after contraceptive discontinuation71
- Frame 5 – Coding of the dependent variables.....72

LIST OF TABLES

- Table 1 – Sociodemographic and sexual and reproductive health characteristics of undergraduate women in the sample. São Paulo, Brazil – 2015 **Error! Bookmark not defined.**
- Table 2 – Sociodemographic and sexual and reproductive health characteristics of undergraduate women in the sample according to the type of contraceptive method used at baseline. São Paulo, Brazil – 2015 **Error! Bookmark not defined.**
- Table 3 – Proportion of undergraduate women according to frequency of discontinuation for any reason and for method-related reasons within 12 months, by sociodemographic and sexual and reproductive health characteristics. São Paulo, Brazil - 2015 **Error! Bookmark not defined.**
- Table 4 – Correlates of contraceptive discontinuation for any reason and for method-related reasons, according to frequency of discontinuation within 12 months (comparison between one and more than one episode of discontinuation). São Paulo, Brazil – 2015 **Error! Bookmark not defined.**
- Table 5 – Multinomial logistic regression of cumulative instances of contraceptive discontinuation for any reason, by sociodemographic and socioeconomic characteristics, educational background, sexual/reproductive and contraceptive behavior. São Paulo, Brazil – 2015 . **Error! Bookmark not defined.**
- Table 6 – Proportion of all months of contraceptive discontinuation according to the type of discontinuation, by type of contraceptive method. São Paulo, Brazil - 2015..... **Error! Bookmark not defined.**
- Table 7 – Correlates of contraceptive discontinuation according to the type of discontinuation, by type of contraceptive method. São Paulo, Brazil – 2015 **Error! Bookmark not defined.**
- Table 8 – Proportion of undergraduate women discontinuing oral pill according to the type of discontinuation. São Paulo – 2015..... **Error! Bookmark not defined.**
- Table 9 – Proportion of undergraduate women discontinuing male condom according to the type of discontinuation. São Paulo – 2015 **Error! Bookmark not defined.**
- Table 10 – Cox proportional hazards model of contraceptive discontinuation according to the type of discontinuation, by type of contraceptive method. São Paulo, Brazil – 2015 **Error! Bookmark not defined.**
- Table 11 – Emergency contraception usage characteristics of undergraduate women in the sample. São Paulo, Brazil – 2015 ... **Error! Bookmark not defined.**

- Table 12 – Sociodemographic and sexual and reproductive health characteristics of undergraduate women according to using of emergency contraception following discontinuation for method-related reasons and method abandonment. São Paulo, Brazil – 2015**Error! Bookmark not defined.**
- Table 13 – Sociodemographic and sexual and reproductive health characteristics of undergraduate women by context of EC use (inconsistent use of contraception or in the context of non-regular contraception). São Paulo, Brazil – 2015.....**Error! Bookmark not defined.**
- Table 14 – Sociodemographic, sexual and reproductive health characteristics of São Paulo university students according to the use of contraception within 30 days after emergency contraception use. São Paulo, Brazil – 2015.....**Error! Bookmark not defined.**
- Table 15 – Correlates of using contraception within 30 days after emergency contraception use. São Paulo, Brazil – 2015.....**Error! Bookmark not defined.**
- Table 16 – Proportion of university students according to contraceptive switching patterns after EC use, by sociodemographic and sexual and reproductive health characteristics. São Paulo, Brazil - 2015**Error! Bookmark not defined.**
- Table 17 – Multinomial logistic regression of contraceptive switching patterns after emergency contraception use, by sociodemographic and socioeconomic characteristics, educational background, sexual/reproductive and contraceptive behavior. São Paulo, Brazil – 2015 .**Error! Bookmark not defined.**
- Table 18 – Sociodemographic, sexual and reproductive health characteristics of São Paulo university students according to the consistent use of contraception after using emergency contraception. São Paulo, Brazil – 2015.....**Error! Bookmark not defined.**
- Table 19 – Reasons for discontinuation the contraceptive method by method type. São Paulo, Brazil – 2015195

LIST OF ABBREVIATIONS

ABEP	Associação Brasileira de Empresas de Pesquisa
AIC	Akaike Information Criterion
CDC	Centers for Disease Control and Prevention
DHS	Demographic and Health Survey
EC	Emergency Contraception
GEE	Generalized Estimating Equations
IBGE	Brazilian Institute of Geography and Statistics
IUD	Intrauterine Device
LARC	Long-Acting Reversible Contraception
LH	Luteinizing Hormone
LNG	Levonorgestrel
NSFG	National Survey of Family Growth
SES	Socioeconomic Status
SRH	Social and Reproductive Health
STATA	Statistical Software
STD	Sexually Transmitted Diseases
UK	United Kingdom
US	United States
USP	University of São Paulo
WHO	World Health Organization

TABLE OF CONTENTS

1. CHAPTER ONE	29
INTRODUCTION	31
1 BACKGROUND SECTION.....	33
1.1 CONTRACEPTIVE CONTEXT IN BRAZIL	33
1.2 SEXUAL AND REPRODUCTIVE HEALTH IN YOUTH POPULATIONS IN BRAZIL	35
1.2.1 Unintended Pregnancies and Contraceptive Behaviors among youth in Brazil.....	35
1.2.2 Special needs of university students.....	36
1.3 CONTRACEPTION DISCONTINUATION AND INCONSISTENT USE.....	38
1.3.1 Discontinuation rates	38
1.3.2 Switching versus abandonment in need of contraception.....	40
1.3.3 Reasons for discontinuation.....	42
1.3.4 Contribution of discontinuation and inconsistent use to unintended pregnancies	43
1.4 EMERGENCY CONTRACEPTION.....	44
1.4.1 Emergency contraception to prevent pregnancy	44
1.4.2 Emergency contraception use in relation to method discontinuation.....	47
1.5 RATIONALE FOR STUDY.....	49
1.6 STRUCTURE OF STUDY.....	50
2. CHAPTER TWO	53
DISSERTATION AIMS, HYPOTHESIS AND CONCEPTUAL FRAMEWORK	55
2.1 DISSERTATION AIMS.....	55
2.2 HYPOTHESIS.....	55
2.3 THEORETICAL FRAMEWORK.....	56
3. CHAPTER THREE	59
METHOD SECTION	61
3.1 OBJECTIVE OF THE SURVEY	61
3.2 SURVEY SITE	61
3.3 SAMPLING (FLOWCHART) AND DATA COLLECTION PROCEDURE	62
3.4 QUESTIONNAIRE	65
3.5 MEASURES.....	67
3.5.1 Unit of analysis: contraceptive discontinuation	67
3.5.2 Outcome Variables	68
3.5.3 Independent Variables.....	72
3.6 STATISTICAL ANALYSIS	74
3.6.1 Descriptive results.....	74
3.6.2 Contraceptive discontinuation analysis.....	75
3.6.3 Emergency contraception analysis based on contraceptive calendar (12-months of observation).....	78
3.6.4 Emergency contraception analysis based on EC calendar (2-months of observation, 30 days before and 30 days after EC use).....	79

4. CHAPTER FOUR.....81

CONTRACEPTIVE DISCONTINUATION (CONTRACEPTIVE CALENDAR)..... 83

4.1 BACKGROUND OF THE STUDY	83
4.1.1 Introduction	83
4.1.2 Aims of study	85
4.2 METHODOLOGY.....	86
4.2.1 Study population and survey site	86
4.2.2 Sampling	86
4.2.3 Data collection and study instruments	87
4.2.4 Unit of analysis.....	87
4.2.5 Measures	88
4.2.6 Statistical analysis.....	90
4.3 RESULTS	ERROR! BOOKMARK NOT DEFINED.
4.3.1 Descriptive statistics	Error! Bookmark not defined.
4.3.2 Characteristics of women who discontinue contraception	Error! Bookmark not defined.
4.3.3 Monthly discontinuation of contraception.....	Error! Bookmark not defined.
4.3.4 Cumulative probabilities of discontinuation of new episodes of contraception: Kaplan-Meier.....	Error! Bookmark not defined.
4.4 DISCUSSION.....	ERROR! BOOKMARK NOT DEFINED.
4.4.1 Contraceptive discontinuation according to frequency of discontinuation.....	Error! Bookmark not defined.
4.4.2 Monthly discontinuation of contraception.....	Error! Bookmark not defined.
4.4.3 Cumulative probabilities of discontinuation of new episodes of contraception.....	Error! Bookmark not defined.
4.5 CONCLUSION	92

5. CHAPTER FIVE94

EMERGENCY CONTRACEPTION AND CONTRACEPTIVE DISCONTINUATION (CONTRACEPTIVE CALENDAR)..... 96

5.1 BACKGROUND OF THE STUDY	96
5.1.1 Introduction	96
5.1.2 Aims of study	98
5.2 METHODOLOGY.....	98
5.2.1 Study population and survey site	99
5.2.2 Sampling	99
5.2.3 Data collection and study instruments	100
5.2.4 Unit of analysis.....	101
5.2.5 Measures	101
5.2.6 Statistical analysis.....	104
5.3 RESULTS OF USING EMERGENCY CONTRACEPTION BASED ON CONTRACEPTIVE CALENDAR.....	ERROR! BOOKMARK NOT DEFINED.
5.3.1 Characteristics of women according to the last time that emergency contraception was used in lifetime and in the 12-months prior to the survey	Error! Bookmark not defined.
5.3.2 Emergency contraception in the context of contraceptive discontinuation for method-related reasons	Error! Bookmark not defined.
5.3.3 Characteristics of women who used emergency contraception following contraceptive discontinuation for method-related reasons and method abandonment	Error! Bookmark not defined.
5.3.4 Characteristics of women who used emergency contraception following inconsistent use of contraception or in the context of non-regular contraception.....	Error! Bookmark not defined.
5.4 DISCUSSION.....	ERROR! BOOKMARK NOT DEFINED.
5.5 CONCLUSION	105

6. CHAPTER SIX	107
CONTRACEPTIVE PATTERNS IN THE MONTH BEFORE AND AFTER EMERGENCY CONTRACEPTION USE (EC CALENDAR).....	109
6.1 BACKGROUND OF THE STUDY.....	109
6.1.1 Introduction	109
6.1.2 Aims of study	110
6.2 METHODOLOGY.....	111
6.2.1 Study population and survey site	111
6.2.2 Sampling.....	111
6.2.3 Data collection and study instruments	112
6.2.4 Measures	112
6.2.5 Statistical analysis.....	113
6.3 RESULTS OF USING EMERGENCY CONTRACEPTION BASED ON EMERGENCY CONTRACEPTION DAILY CALENDAR.....	ERROR! BOOKMARK NOT DEFINED.
6.3.1 Characteristics of women who used contraception after using emergency contraception	Error! Bookmark not defined.
6.3.2 Correlates of using contraception within 30 days after emergency contraception use: results of multivariate analysis	Error! Bookmark not defined.
6.3.3 Characteristics of women according to contraceptive switching patterns after emergency contraception use.....	Error! Bookmark not defined.
6.3.4 Correlates of contraceptive switching patterns after emergency contraception use: results of multinomial analysis	Error! Bookmark not defined.
6.3.5 Characteristics of women who had gaps after using emergency contraception.....	Error! Bookmark not defined.
6.4 DISCUSSION.....	ERROR! BOOKMARK NOT DEFINED.
6.4.1 Use of contraception after EC use	Error! Bookmark not defined.
6.4.2 Contraceptive switching patterns after emergency contraception use.....	Error! Bookmark not defined.
6.4.3 Gaps in contraception after emergency contraception use.....	Error! Bookmark not defined.
6.5 CONCLUSION	114
7. CHAPTER SEVEN.....	117
7.1 SUMMARY OF FINDINGS	119
7.2 STRENGTHS AND LIMITATIONS.....	122
7.3 RECOMMENDATIONS FOR FUTURE RESEARCHES.....	126
7.4 PUBLIC HEALTH IMPLICATIONS	127
REFERENCES.....	129
APPENDICES.....	155
APPENDIX A – CONSENT FORM	157
APPENDIX B – QUESTIONNAIRE (ENGLISH VERSION)	159
APPENDIX C – QUESTIONÁRIO (VERSÃO EM PORTUGUÊS).....	171
APPENDIX D – INDEPENDENCE MODEL CRITERION (QIC).....	185
APPENDIX E – REASONS FOR CONTRACEPTIVE DISCONTINUATION	195
ANNEX	197
ANNEX – APPROVAL OF THE RESEARCH ETHICS COMMITTEE	199

PREFACE

I am a midwife, I graduated at University of São Paulo, School of Arts, Sciences and Humanities in 2009. I started appreciating the reality of Sexual and Reproductive Health (SRH) of young women when I was an undergraduate student, in which I had the opportunity to participate on a project about contraceptive methods, women's health, and prenatal care with young women from a community in the city of São Paulo, Brazil. In addition, I worked with young women during the university internship. In these works, I identified areas for further research to better understand the problems of young people. So, I quickly fitted well and identified with female young people.

After I finalized my undergraduate course, I immediately started my master degree at University of São Paulo, School of Nursing in 2010, and Profa. Ana Luiza Vilela Borges welcomed me with "open arms". I realized a dream to work with contraception, especially with emergency contraception among adolescent people. In my master's degree, I again had the opportunity to be close to adolescents through lectures given at schools on contraceptive and sexual behaviors. My admiration for the subject and for this group only increased even more. In my PhD, I was already sure that I would like to work with young people and contraception, and obviously with emergency contraception. At this time, Profa. Ana Luiza talked to me about a study that she and other authors conducted with young students at University of São Paulo, in which they had observed inconsistency in the contraceptive use. She presented me the contraceptive calendar and told me about her idea of doing something similar in relation to emergency contraception. The idea was sounded awesome for me and I was enthusiastic on contraceptive discontinuation issues surrounding sexual and reproductive health.

In 2015, I received a scholarship from a Brazilian government program, and I went to Baltimore to work with Profa. Caroline Moreau at Johns Hopkins University in 2016. She supported me in all this process and shared her knowledge of sexual and contraceptive behavior, and taught me how to analyze the contraceptive calendar, which is used in Demographic and Health Surveys.

All these challenges motivated me throughout the study process. This study opened my eyes to the reality of fertility, sexuality and contraceptive needs of young people and how we can help them. Although I did not have direct contact with the young women who participated in my study, I had the opportunity to receive a lot of messages from them through E-mail, Facebook and WhatsApp. This approach helped me contextualize the problem of the study population. It was a privilege and grateful for me to be able to help these young women in some way. Certainly, I get much more than I give.

1.CHAPTER ONE

CHAPTER ONE

INTRODUCTION

This document presents a study on contraceptive discontinuation among undergraduate women at University of São Paulo, Brazil. The present work builds on two studies conducted among Brazilian university students at University of São Paulo in 2002 and 2009 (Pirota, Schor 2004; Borges et al., 2010). The study by Pirota, Schor (2004) noted high proportions of contraceptive use, concomitantly with high proportions of pregnancies ended by abortion, which reveals that the contraceptive use was inadequate and inconsistent in this group. On the other hand, the study by Borges et al. (2010) pointed out that the emergency contraception (EC) use was a marker of the contraceptive discontinuation. Therefore, both studies noted a worrying aspect related to sexual and reproductive health among university women: the contraceptive use is frequent, but occurs in a discontinuous way. Although relevant, none of the previous studies measured contraceptive discontinuation and EC use in the context of contraceptive discontinuation. Thus, this study advances in relation to the two other studies mentioned, since it aims to understand the contraceptive practices among university students, which represents a specific group with high educational and professional perspectives, and which frequently use effective methods to postpone reproductive life (Longo et al., 2002; Sant'Anna et al., 2008; Veloso et al., 2014). The investigation of contraceptive practices among university students is justified by the fact that the level of knowledge and contraceptive access are supposedly higher among women with higher educational level. However, apparently, these are not the only aspects for the success of preventing an unintended pregnancy, since, even with knowledge and easy access to contraception, contraceptive discontinuation among university students can lead to unwanted outcomes, such as unintended pregnancy.

To summarize, one of the issues which is consistently raised in studies of contraceptive use in Brazil is the inconsistency and gaps in contraception use, and the lack of EC use after contraceptive discontinuation, all contributing to high proportions of pregnancies that are unintended. The specific focus of the study is to

evaluate frequency and correlates of contraceptive discontinuation and how it relates to subsequent use of emergency contraception. More specifically, we will consider the following research questions:

- How often do undergraduate and young women discontinue their method of contraception in a 12-months period?
- What are the determinants of contraceptive discontinuation in a 12-months period and more specifically, what are the differences between woman who did not discontinue their method, woman who discontinue once, and those who discontinued their method several times in the period of 12-months?
- What is the monthly probability of method discontinuation for method-related reasons and of method abandonment over a 12-month period?
- What is the probability and correlates of using EC after contraceptive discontinuation?
- What are the patterns of contraceptive use after using EC?
- What determines the contraceptive discontinuation within a month following the use of EC?

The thesis is presented in a seven-chapter format where a variety of data and methods are employed to fully explore the topic. This introductory chapter presents information on the background context in relation to contraceptive discontinuation and EC in Brazil. We then present the specific aims of the study and outline the structure of the thesis.

1 BACKGROUND SECTION

1.1 CONTRACEPTIVE CONTEXT IN BRAZIL

Brazil has recently undergone a rapid fertility transition with total fertility rates dropping below replacement, from 2.5 in 1996 to 1.8 in 2006 (Brasil, 2008a). At the same time Brazil, has seen a dramatic increase in modern contraceptive prevalence rates, estimated at 77% in the last national survey conducted in 2006 (Brasil, 2008a). While contraceptive prevalence rate is one of the proximate determinants of fertility and is used to monitor the effectiveness of Family Planning Programs in reducing unwanted fertility, prevalence alone is no longer sufficient in informing programs aimed at reducing the proportion of pregnancies reported as unintended, which remains high despite high contraceptive coverage (Boogarts, 1978; Lo et al., 2014). By implication, then, it is important to assess the dynamics of contraceptive use, beyond prevalence.

In Brazil, unmet need is estimated at 6% but 46% of pregnancies are reported as unintended (Brasil, 2008a; Berquó, Lima, 2008). This situation is comparable to other countries with high contraceptive coverage, such as the United States (U.S.), France or the United Kingdom (U.K.), where 33% to 45% of pregnancies are classified as unintended (Bajos et al., 2014; Moreau et al., 2014; Wellings et al., 2015; Finer, Zolna, 2016). Most of these pregnancies are due to inconsistent use or gaps in the use of contraception (Goulard et al., 2006; Frost, Darroch, 2008), which raises the issue of method choice/access and dynamics of use. In addition, it is important to underline that unintended pregnancy has been associated with increased risks of maternal morbidity, and health behaviors during pregnancy that are associated with adverse maternal and infant health outcomes, such as delayed prenatal care, premature birth, lower rates of breastfeeding, risk of miscarriage, having low birth weight infants, exposure to suicide and depression, and negative physical and mental health effects for children (Mayer et al., 1997; Santelli et al., 2003; Sereshti et al., 2005; Gipson, Koenig, Hindin, 2008; Cheng et al., 2009; CDC, 2011a; CDC, 2011b).

In Brazil, for limiting needs, sterilization is used by one third of women, but for spacing needs, the contraceptive landscape is dominated by pills or condoms, which are associated with high discontinuation rates and moderate failure rates (Brazil,

2008a; Kost et al., 2008). In the last Demographic and Health Survey (DHS) survey conducted in 2006, a quarter of Brazilian women were relying on the contraceptive pill (24.7%) and 12% used male condoms for the prevention of pregnancy (Brasil, 2008a). While very effective if used consistently, these methods are subject to failure because of inconsistent or errors of use (failure rates among pill and condom users are 9% and 18%, respectively) (Trussell, 2011). In Brazil, the lack of trained professionals and limited contraceptive options (Heilborn et al., 2009; Vieira, Souza, 2011) contribute to skewed contraceptive distribution favoring user dependent methods (Moura, Silva, Galvão, 2007). Long-acting reversible contraception (LARC) such as implant or intrauterine device (IUD) are more effective (failure rates are 0.05% and 0.8%, respectively) (Trussell, 2011), but they are seldom used in Brazil (0.1% and 1.9%, respectively) (Brasil, 2008a). While similar contraceptive patterns are observed in other contexts such as U.K., Ireland and Uganda (Lader, 2009; Lally et al., 2015; Nsubuga et al., 2016), recent efforts in a number of countries, including the U.S. and U.K, are seeing upward trends in the use of LARC mainly among young people, which are believed to have contributed towards lowering fertility rates among youth (National Institute for Health and Clinical Excellence, 2014; Secura et al., 2014; Ricketts, Klingler, Schwalberg, 2014; Kavanaugh, Jerman, Finer, 2015; Daniels et al., 2015; Lotke, 2015).

Suboptimal efficiency of contraceptive practices in Brazil (favoring moderate instead of highly effective methods) is a missed opportunity to ensure successful reproductive planning. A better understanding of contraceptive use dynamics, including discontinuation and use of back up options to reduce the risk of pregnancy after unprotected sex is an important step in informing more effective strategies to reduce unintended pregnancies. In the context of legal restrictions to abortion, the prevention of unintended pregnancy is a primary prevention strategy to reduce unsafe abortion, which is estimated at an annual average of 994,465 abortions, and responsible for 12% of maternal mortality in Brazil (Faúndes et al., 2007; Martins-Melo et al., 2014), and thus improve women's health.

1.2 SEXUAL AND REPRODUCTIVE HEALTH IN YOUTH POPULATIONS IN BRAZIL

1.2.1 Unintended Pregnancies and Contraceptive Behaviors among youth in Brazil

The higher rates of unintended pregnancies across most developed countries occur among young women aged 20–24 (Diamond-Smith, Moreau, Bishai, 2014; Wellings et al., 2015; Coetzee, Ngunyulu, 2015; Finer, 2016; Moreau, Bohet, 2016). Similarly, in Brazil the highest burden of unintended pregnancies falls on youth (ages 15 to 24) who are at the peak of their fertile years. Youth population (20 – 24 years old) represent 17.9% of the population in Brazil but account for 25% of unintended pregnancies (Berquó, Lima, 2008; IBGE, 2010). In case of unintended pregnancies, young women are also more likely to have an induced abortion, with peak of lifetime experience observed between 20 and 24 years (24% in this age group) (Diniz, Medeiros, 2010).

While increase in contraceptive prevalence over time (increase of 11% between 1986 and 1996, and 4% between 1996 and 2006) has potentially reduced unintended pregnancies (decrease from 22% in 1996 to 18% in 2006) among all women including young women (Brasil, 1997; Brasil, 2008a); social changes towards delaying the first birth are extending the number of years young women are exposed to the risk of unintended pregnancies. Although the average age at first birth has remained over the years (21 years) (Brasil, 2008a) and age at first intercourse has remained rather stable (15 years old) until 2006, recent trends are noticeable (Brasil, 2008a). Specifically, the proportion of women, mainly those with higher educational level, who have their first birth at 30 years or more has grown substantially in the last decade (14% increase), representing now 32% of Brazilian women (Brasil, 2008a; Brasil, 2014; IBGE, 2015), which means there is a growing number of years between first sex and first birth among women who are in the most fertile years of the reproductive life span (Brasil, 2014).

In the face of exposure to unintended pregnancy risk, a growing proportion of Brazilian youth (77.1%) use modern contraceptive, mainly hormonal contraception (pill) (44%), condom (16.8%), and injectable (9.2%) (Brazil, 2008a). These data reveal a breakthrough in the use of contraception in a 10-year period, as previous

1996 DHS estimates indicated that only 66% of Brazilian youth were using modern contraceptive, especially hormonal contraception (39.5%), condom (5%), and injectable (3.6%). The 2006 also reveals the sharp decline in the prevalence of female sterilization among young women (from 11.4% in 1996 to 4% in 2006) (Brasil, 2008a). While overall coverage has increased, the method mix indicates that women in Brazil favor methods that have moderate to high typical use failure rates which in turn contribute to unintended pregnancies and induced abortion (Frost, Singh, Finer, 2007a; Bradley, Croft, Rutstein, 2011; Sedgh et al., 2012).

A study published by Brazil's Ministry of Health based on a literature review from 1987 to 2008, noted that more than half of Brazilian young women who had induced abortions reported use of contraceptive methods at the time of conception (in particular the contraceptive pill), reflecting inconsistent or errors of use, in addition to frequent discontinuation (Brasil, 2008b). This is in line with other studies in high contraceptive prevalence countries indicating that most unintended pregnancies are attributable to incorrect or inconsistent contraceptive use (Jones, Darroch, Henshaw, 2002; Bajos et al., 2006; Schünmann, Glasier, 2006; Rash et al., 2007; Frost, Darroch, Remez, 2008; Moreau et al, 2010; Moreau et al, 2012; Wellings et al., 2015). A study conducted among French women aged 13-50 undergoing an elective abortion, observed that 66% of these women were using contraception at the time of conception (Moreau et al., 2010). Jones, Darroch & Henshaw (2002) indicated that 53.3% of young women aged 20-24 in the U.S. who had induced abortion were using a method in the month of conception, and they were the most likely of any age-group to have been using the pill (15%) when they became pregnant.

These data illustrate the need not only to consider overall contraceptive prevalence, but more specifically, the dynamics of contraceptive use including discontinuation which is reported to be frequent in youth populations (47.3% of young women aged 20–24 in the U.S. discontinue contraception) (Pazol et al., 2015).

1.2.2 Special needs of university students

Undergraduate students represent a specific group among young people, with high educational and professional aspirations. Delaying childbearing generates a special demand for effective contraception until they reach professional stability (Longo et al., 2002; Sant'Anna et al., 2008; Veloso et al., 2014). While pregnancy

rates are lower in this population compared to their less educated peers (Finer, Zolna, 2016), a higher proportion of these pregnancies are unintended (60%), mostly related to contraceptive discontinuation (Bryant 2009; Trieu et al., 2011). A study conducted among undergraduate students in the U.S. estimated that continuous use of contraception could prevent about 12 million unwanted pregnancies every year (Ersek et al., 2011). In Brazil, studies conducted among undergraduate students in two states (Minas Gerais and Paraná) found that 29.3% and 65.9% of students, respectively, had used contraceptive methods inconsistently despite high levels of knowledge about contraception (Dessunti, Kings, 2012; Silva, Camargo, Iwamoto, 2014). Young women with higher education try to prevent unintended pregnancies more than others (Mosher, Jones, Abma, 2012), but while they use contraception whether the contraception fail, they are more likely to have an induced abortion (Finer, Zolna, 2011). In this scenario, young people with higher education are 3.8 times more likely to have an induced abortion than less educated youth, which contributes to potential maternal morbidity in the context of restrictive laws on abortion (Menezes, Aquino, 2009). High rates of contraceptive discontinuation resulting in unintended pregnancies amongst students at higher education institutions are also reported in other countries, including Asia, Africa, and the U.S. (Trieu et al., 2011; Mehra et al., 2012; Somba et al., 2014; Coetzee, Ngunyulu, 2015; Wang et al., 2015).

It is important to note that there are few people in Brazil attending tertiary educational. It is estimated that only 12.6% of Brazilians have a university degree (IBGE, 2015). Therefore, Brazil presents a relevant social and educational disparity, which needs to be considered in the context of sexual and reproductive health, as high educational level is related to the process of women postponing marital unions, using contraceptive methods and tending to have fewer unintended pregnancies (Pirota, Schor, 2004). Unintended pregnancies represent a barrier to achieving academic success, besides impacting a life project encompassing the pursuit of economic autonomy, professional achievement and expectation to maintain or improve socioeconomic status in relation to the family of origin. Thereby, an unintended pregnancy has considerable social and psychological repercussions in the lives of university students (Pirota, Schor, 2004).

In this context, investigating highly educated women allows us to understand the contraceptive practices of a “privileged group”, which is motivated to prevent a

pregnancy and less likely to face access barriers relative to lower socioeconomic population.

1.3 CONTRACEPTION DISCONTINUATION AND INCONSISTENT USE

1.3.1 Discontinuation rates

Contraceptive discontinuation is defined as switching a method of contraception or stopping a method altogether. Indicators typically discern discontinuation for any reason or for method-related reasons, the later excluding stopping a method to get pregnant or because women are no longer exposed to the risk of pregnancy (sterile or no partner) (Trussell, Vaughan, 1999).

In the U.S., overall, 68% of all methods are discontinued for any reason by the end of 12 months, and discontinuation rates for any reason range from 81.9% (for male condom users) to 47.4% (for oral pill users) (Vaughan et al., 2008). Moreau et al. (2009a) found lower contraceptive discontinuation rates among French women although similar patterns by method were also observed, with one-year discontinuation rates ranging, from 30.7% (for oral pill users) to 15% (for IUD users). In Brazil, 41.1% of all methods are discontinued for any reason within one year, and the discontinuation rate for any reason is higher for injectable (61.8%), withdrawal (60.1%), male condom (59.8%), periodic abstinence (46.3%), oral contraceptive (44.3%), and IUD (5%) (Ali, Cleland, Shah, 2012; Ferreira, 2014).

Discontinuation for method-related reasons pertains to situations when women stop the method despite having still a need for contraception. Reasons usually include dissatisfaction or problems with the method because of side-effects, health concerns, inconvenience of use, difficulties in access, high cost, or partner's opposition (Bradley, Schwandt, Khan, 2009). Discontinuation for method-related reasons is a more relevant indicator of discontinuation than discontinuation for any reasons in terms of pregnancy prevention, as it does not include discontinuation due to a desire to get pregnant or because of partner breakup (Ali, Cleland, Shah, 2012). Moreover, discontinuation for method-related reasons deserves specific attention, since it suggests a mismatch between women's desire and their practice that can potentially be resolved with better counselling (Hooper, 2010).

Considering all methods, 47.1% of all methods are discontinued for method-related reasons within 12-months in the U.S., compared with 27.8% in Brazil (Vaughan et al., 2008; Ali, Cleland, Shah, 2012). Of particular concern are the high discontinuation rates of user dependent methods for method-related reasons. In the U.S., this estimate ranges from 57% (for male condom users) to 33% (for oral pill users), while 8% of IUDs and 13.1% of implants are discontinued for method-related reasons (Vaughan et al., 2008; Mansour, 2008; Castle, Askew, 2015). Similarly, Moreau et al. (2009) observed that contraceptive discontinuation for method-related reasons among user dependent methods in France, ranged from 38.9% (for withdrawal user) to 21.5% (for oral pill users), while discontinuation of IUDS for method-related reason was only 10.7%. In Brazil, the method-related discontinuation within 12-months range from a high of 42.6% for male condom to 37.3% for injectable and 33.1% for oral pill – the most widely used methods among Brazilian youth (Ali, Cleland, Shah, 2012; Alano et al, 2012; Silva et al., 2010; Silva Camargo, Iwamoto, 2014).

Many factors are related to discontinuation, including external factors (such as quality of family planning services, family planning service environment, guidelines, among others), but also individual and partner factors (such as demographic characteristics, partner dissatisfaction, fertility motivations, among others) (Curtis, Blanc, 1997; Ali, Cleland, 1999; Curtis, Evens, Sambisa, 2011; Barden-O'Fallon, 2011a; Moreau et al., 2013). Prior studies indicate that women who are younger, who are black, and have higher parity are more likely to discontinue their methods (Ali, Cleland, 1995; Curtis, Blanc, 1997; Grady, Billy, Klepinger, 2002). Moreover, fertility desires, representing women's motivations to prevent a pregnancy is also related to contraceptive discontinuation (Blanc, 2001; Moreau et al., 2013).

Partners are also involved in method selection and continuation, although their level of influences is not totally comprehended (Barden-O'Fallon, 2011a). Earlier research on contraceptive discontinuation highlighted that lack of planning and sporadic sexual intercourse, as well as relationship dynamics and short duration of the relationship were factors contributing to use versus nonuse of contraception and contraceptive discontinuation (Manlove et al., 2003; Frost, Singh, Finer, 2007b; Sessler et al., 2009). Length of the relationship in particular is found to be associated with a tendency to relax contraceptive use when the partner is known and the relationship becomes stable (Aquino et al., 2003; Pirotta, Schor, 2004; Manlove,

2007). Studies about the link between relationship status and contraceptive use is frequently analyzed in terms of motivation (Frost, Singh, Finer, 2007b; Sassler et al., 2009). According to Manlove, Ryan & Franzetta, 2003, unmarried women use contraception less consistently in relationships that women report to be more serious. The authors suggest that motivation to avoid pregnancy is lower when the relationship develops more commitment. However, Reed et al. (2014) point out that the differences in consistency by relationship type or length probably occur because is more challenging to be consistent for a long than a short time. This also occurs in the beginning of a relationship, where there is need for negotiating contraception, mostly condoms because of STI considerations (Aquino et al., 2003; Manlove, 2007; Osei et al., 2014).

1.3.2 Switching versus abandonment in need of contraception

Not all women who discontinue contraception have an unmet need for contraception, since many women switch to another contraceptive, reducing their risk of unintended pregnancy. Nevertheless, it is important to note that switching in itself contributes to heighten risk of an unintended pregnancy, since failures are higher at the start of use of a method when women are still familiarizing themselves with the regimen (Peterson et al., 1998; Leite, Gupta, 2007; Ranjit et al., 2001; Trussell, 2004; Moreau et al., 2007).

After discontinuation, most women switch rather than abandon their method, and in most cases switching occurs within three months; so, switching methods is the primary reason for discontinuation (Vaughan et al., 2008; Moreau et al., 2009a; Ali, Cleland, Shah, 2012). Grady, Billy & Klepinger (2002) observed that 40% of married women in the U.S. switch methods (stop a method to adopt a new contraceptive) within a two-year period. In a national telephone survey of a sample of nearly 2,000 women 18-44 years in the U.S., Frost, Singh & Finer (2007b) found that nearly 25% of women switched their contraception in 12-months. In contrast, a national cohort study of women aged 18–49 in England, noted that 11.3% of women switched their method in 12-months, and only 1.8% switched for method-related reasons (Wellings et al., 2015). In addition, switching to another option for method-related reasons varies by contraceptive method. According to a national study conducted among women aged 18–44 in France, the frequency of switching for method-related reasons

within 12 months is higher among withdrawal users (38.9%), followed by condom (38.3%), fertility awareness (34.5%), oral pills users (21.5%), and IUD (10.7%).

In Brazil, 21.6% of women switched their methods for method-related reasons within 12 months, and rates of switching for method-related reasons were higher for injectable (38.8%), followed by male condoms (37.6%), traditional methods (29.6%), and oral pills (16.1%) (Leite, Gupta, 2007). In addition, a cross-national study that selected nationally representative samples of women aged 18–44 in eight countries (including Brazil), observed that Brazil was the second country with the highest rate of switching from one hormonal contraceptive to another (25%) (switching rates ranged from 12% in France to 28% in Russia) (Hooper, 2010). After discontinuing contraception for method-related reasons, Brazilian women were more likely to switch to any modern reversible method (21.3%) than traditional method (20.4%) (Ali, Cleland, Shah, 2012).

The probability of abandonment of contraception for method-related reasons is highest in the U.S. (25%), compared with Brazil (10.34%), France (6.0%) and England (0.4%) (Leite, Gupta, 2007; Vaughan et al., 2008; Moreau et al., 2009a; Wellings et al., 2015). Similarly, to switching, contraceptive abandonment for method-related reasons varies by method. In France and Brazil, highly effective methods users (pill, injectable, IUD or other hormonal methods) were more likely to abandon for method-related reasons contraceptive use than traditional methods users (Leite, Gupta, 2007; Moreau et al., 2009a). Besides side-effects, high rates of method abandonment in the Brazilian context may reflect lack of contraceptive options for women who stop their method (Castle, Askew, 2015), and may also reflect method stock-out (Bradley, Schwandt, Khan, 2009).

Abandonment and switching are more common among young women (Blanc et al., 2009, Barden-O'Fallon, 2011b). According to Grady et al. (2002), unmarried women in the U.S., younger and more highly educated had high rates of switching to the condom, and were more likely to abandon the oral pill, putting them at greater risk of unintended pregnancy. Similarly, a study conducted with women in England, found that stoppers and switchers were younger, better educated and more likely to be single (Wellings et al., 2015). In Brazil, 20.7% of young women aged 20–24 switched their method and almost 10% abandoned the method within the first year of use (Leite, Gupta, 2007). High rates of switching and abandonment among youth are probably related to the specific challenges they face in managing regular

contraception across short-term and less stable relationships with periods of infrequent intercourse (Pirota et al., 2004; Beadnell et al., 2005; Borges et al., 2010).

1.3.3 Reasons for discontinuation

The reasons for discontinuation varies by method. Reasons for discontinuation of condoms include the partner's opposition, vaginal irritation, latex allergies, desire to switch to an easier or more reliable method, and the perception that condoms are uncomfortable and interfere with sexual pleasure (Brunner Huber et al., 2006; Ersek et al., 2011; Davidoff-Gore, Luke, Wawire, 2011; Zembe et al., 2012; Mehra et al., 2014). Studies have suggested that use of condom is greater than the use of other methods at first intercourse, in the beginning of a relationship, and among those whose sexual intercourse is casual rather than stable – situations that are more common among youth (Santelli et al., 1997; Manlove, Ryan, Franzetta, 2003; Bastos et al., 2008; Raine et al., 2010; Beltzer et al., 2011). However, as relationships become more stable, many couples switch to hormonal methods, and the main reason to switch is the “supposed mutual trust”, which indicates that they become less concerned about the prevention of sexually transmitted diseases (STDs) (Bastos et al., 2008; D'Antona et al., 2009; Shih et al., 2011; Goldstein, Upadhyay, Raine, 2013; Mehra et al., 2014).

The main reasons for discontinuation of hormonal contraceptives include “(non)compliance” with regular use (difficulty remembering, absence from home and hence lack of pills, loss of health insurance, depletion of supplies, lack of time, and difficulty keeping track of the ring), side-effects (headache, breast pain, yeast infections, nausea, weight gain, low libido, changes in menstrual flow) and health concerns (fear of hormone/cancer) (Jones, Darroch, Henshaw, 2002; Caruso et al., 2005; Leite, Gupta, 2007; Harvey et al., 2009; Castle, Askew, 2015; Inoue, Barratt, Richters, 2015).

The most common reason for discontinuing traditional methods is the desire to switch to a more effective method (Leite, Gupta, 2007; Vaughan et al., 2008; D'Antona et al., 2009; Bradley, Schwandt, Khan, 2009). Another common reason is pregnancy due to higher failure rates of these methods as compared to modern methods (Kost et al., 2008).

1.3.4 Contribution of discontinuation and inconsistent use to unintended pregnancies

Unintended pregnancies are generally attributed to many complex and interconnected aspects, such as partnership and autonomy of women/gender relations; quality of health care; sexual and reproductive experience; factors related to contraception, which include: contraceptive failure, contraceptive nonuse, and incorrect or inconsistent use of contraceptive methods, among others (Langer et al., 2002; Prietsch et al., 2011; Finer, Zolna, 2011; Theme-Filha et al., 2016). In countries with moderate to high contraceptive prevalence, the majority of unintended pregnancies are the result of contraceptive discontinuation or failures due to inconsistent use of contraception (Jain, 1999; Trussell, 1999; Blanc, Curtis, Croft, 2002; Casterline, El-Zanaty, El-Zeini, 2003; Kost et al., 2008). A study in the US estimated that nearly 50% of the 3.1 million unintended pregnancies occurred to women who were not using contraception at the time of conception but had used contraception in the past (Finer, Henshaw, 2006).

In the same way, Blanc et al. (2002) noted that in 14 of 15 developed countries examined, more than half of unintended pregnancies were due to gap in use of contraception or failure due to errors of use, including in Brazil, Colombia, Jordan, and Peru. According to the authors of that study, the total fertility rate would drop by 20 to 48% in these 15 countries, by preventing births resulting from contraceptive discontinuation. Discontinuation for reasons other than a desire to get pregnant was also strongly associated with unintended pregnancies in Guatemala (17%) (Barden-O'Fallon, Speizer, White, 2008).

Additional research has associated discontinuation with induced abortion (Jain, 1999; Casterline, El-Zanaty, El-Zeini, 2003). Creanga et al. (2007) estimated that 60% of pregnancies resulting from contraceptive discontinuation among Romanian women resulted in induced abortion. In Armenia and Kazakhstan, these rates reached 80% (Bradley, Schwandt, Khan, 2009). In France, Bajos et al. (2006) observed that 50% of women who had an abortion had switched their contraception in the six months before the event. In this context, it is important to note that this association may be different in countries with legal restrictions on abortion, such as Brazil.

Given the strong association between contraceptive discontinuation and unintended pregnancy, a better understanding of factors that favor sustainable use of contraception and reduce the risk of pregnancy after method discontinuation is critical to reduce levels of unintended pregnancy (Curtis, Evens, Sambisa, 2011).

1.4 EMERGENCY CONTRACEPTION

1.4.1 Emergency contraception to prevent pregnancy

Emergency contraception (EC) prevents women from becoming pregnant after unprotected sexual intercourse and may be used as a backup option, in particular in cases of contraceptive discontinuation with potential gaps in use (Raine et al., 2005; Borges et al., 2010).

EC levonorgestrel (LNG) pills are the only EC products available in Brazil until 2016, introduced in the public health programs in 2001 (Brasil, 2010; World Health Organization, 2012). LNG EC prevents pregnancy by delaying or inhibiting ovulation if is given before the Luteinizing Hormone (LH) peak, that is 24 to 48h before ovulation (Brache et al., 2010). LNG EC is estimated to prevent between 52% and 94% of pregnancies (Arowojolu et al., 2002; Hamoda et al., 2004; Creinin et al., 2006; Glasier et al., 2010), if given up to 120 hours after unprotected intercourse. While there are other EC methods that are more effective such, as the intrauterine device, mifepristone EC and ulipristal acetate, these methods were not available at the time of this research (Glasier et al., 2010; Cleland et al., 2012; Trussell, Raymond, Cleland, 2014).

Levels of use of EC are mostly a function of access conditions, and prescription of EC is one of the most important determinant to its use. In some countries, EC is obtainable only after a prescription, which can be a challenging and time-wasting, especially on holidays and weekends, because the primary health care is closed. In addition, some women think that request EC from health professionals is embarrassing, and others do not have access to primary health care services (Polis et al., 2010; Westley, Bass, Puig, 2016). In some countries, such as Brazil, women can buy EC at the pharmacy without a prescription (Borges et al., 2010; Tavares, Foster, 2016). In this context, a systematic review of the literature based on clinical trials comparing advanced provision to prescription only access shows substantial

increases in usage in non-prescription status conditions (Polis et al., 2010). These results are confirmed at the population level where levels of use increase when prescription requirements are abandoned.

National estimates of EC use are difficult to compare as studies refer to different time frames (lifetime or annual incidence), different study populations (all women of reproductive age, young women, abortion patients) and different time periods extending over 15 years. Following the switch to non-prescription status, 17% of women 15-44 years old in France had ever used EC in 2004 (Moreau, Trussell, Bajos, 2006), 11% of women 15–49 years old has done so in the U.S. in the period 2006–2010 (NHS Contraceptive Services, 2011/12), while 16% of young women (average age 19.5 years) in Italy had ever used EC despite the prescription requirement (Tafari et al., 2012), or annual incidence (7% of women aged 16–49 in the U.K. had used EC in the past year, 2008 – 2009) (Daniels, Jones, Abma, 2013).

In Brazil, according to 2006 DHS Survey, 12% of all women ever used EC, with the highest use rate observed among youths aged 20-24 (18.5%) (Brazil, 2008a). EC use is significantly more frequent among Brazilian undergraduate students, ranging from 42.0% to 50.4% lifetime use (Borges et al, 2010; Silva et al., 2010; Alano et al., 2012). Although a substantial proportion of women have already used EC, EC is not used enough to have a public health impact in many countries, that is, EC is not utilized as often as needed to curve unintended pregnancy risk (Hu et al., 2005; Glasier et al., 2010; Palermo, Bleck, Westley, 2014; Santos et al., 2014; Nappi et al., 2014). Studies conducted in Scotland, France and Denmark over the last 15 years, observed that nine out of 10 women undergoing an abortion do not use EC to try to reduce the risk of becoming pregnant, a figure that has remained constant despite greater use of EC in the general population, (Sørensen et al., 2000; Moreau et al., 2005; Cameron et al., 2012). In Scotland, the availability of free EC from pharmacies did increase the proportion of EC use among women seeking an abortion (Cameron et al., 2012).

A number of studies have identified a number of barriers to EC use such as lack of knowledge or misperceptions of pregnancy risk (Moreau et al., 2005; Alano et al., 2012; Westley et al., 2013; Santos et al., 2013). A study based on literature review of global access to EC noted that the majority of women in high-income countries were aware of EC, but that this was not the case in middle- and low-income countries (Westley et al., 2013). However, a study carried out with high school students in

Brazil, observed that the level of knowledge about EC did not interfere in the method use (Nascimento, 2012). Santos & colleagues (2014) observed that no pregnancy risk awareness was associated with non-use of EC among women, on average 25 years old, in Brazil. However, awareness alone is not synonymous of using EC among women who need it (ESHRE Capri Workshop Group, 2015). Misinformation about the timeframe for use, and misperceptions about mode of action were associated with nonuse of EC among women undergoing an abortion in France (Moreau et al., 2005).

Research conducted in the U.K., France and Denmark consistently shows that underestimation of the chance of becoming pregnant is one of the reasons for nonuse of EC (Sørensen et al., 2000; Moreau et al., 2005; Lakha Glasier, 2006). According to Moreau et al. (2005) almost 61% of women undergoing an abortion in France indicated that they did not use EC because they did not think could get pregnant. Similarly, another study conducted among women aged 16–46 in five European countries (France, Germany, Italy, Spain, and the U.K.) noted that the majority of women did not use EC when they needed it because they did not perceive themselves to be at risk of pregnancy (Nappi et al., 2014). In Brazil, study by Santos et al. (2014) observed that women who were not aware of pregnancy risk were three times more likely to non-use of EC than women who had pregnancy risk awareness.

In the country, other barriers to EC access and use prevail, including, stock outs in the public health sector, moral and cultural aspects, cost, inadequate disclosure of the places where it can be acquired, and concerns about EC deterring from regular contraception (Hardy et al., 2001; Costa et al, 2008; Alano et al., 2012; Faúndes et al, 2015; Tavares, Foster, 2016). In addition, studies have shown that the availability of EC in the Brazilian scenario is not fully consolidated in primary health care, mainly due to moral and religious factors, and the lack of knowledge both from health professionals and users (Costa, Guilhem, Silver, 2006; Souza, Brandão, 2009; Figueiredo, Paula, 2012). A study carried out in São Paulo state, whose municipalities received remittances of EC from the Ministry of Health, shows that almost half of the primary health care services do not dispense EC (Figueiredo, Paula, 2012). The authors also point out a lack of technical preparation of health professional to deal with EC, i.e., misconceptions such as a belief that an "abusive" use of EC might occur in case the method is highly available (Souza, Brandão, 2009; Figueiredo, Paula, 2012).

It is important to emphasize that not only in Brazil but also in other countries such as France, the U.S. and the U.K., EC users are mostly young and unmarried (Martin, 2004; Moreau et al., 2006; Lakha F, Glasier, 2006; Escobar et al., 2008; Cameron et al., 2011; Daniels, Jones, Abma, 2013); and the main reasons cited for EC use are: contraceptive abandonment, switching or failure of the contraceptive, i.e., contraceptive discontinuities (Moreau et al., 2006; Cameron et al., 2011).

Certainly, EC is a backup option to prevent unintended pregnancies, in particular in cases of contraceptive discontinuation with potential gaps in use (Raine et al., 2005; Borges et al., 2010).

1.4.2 Emergency contraception use in relation to method discontinuation

EC is mostly used to compensate for discontinuation and errors of the use of regular methods (mainly because of a ruptured condom and missed oral contraceptives), (Falk et al., 2001; Sorhaindo et al., 2002; Moreau et al, 2006). In this scenario, EC is considered a marker of inconsistent or errors of use of contraception rather than a regular method for women who are not using any other options (Jones, Darroch, Henshaw, 2002; Moreau et al., 2009a; Nappi et al., 2014).

Resumption of use of contraception use immediately after EC use is critical in limiting the risk of unintended pregnancy since EC failure rates are higher if there are several acts of unprotected intercourse in the same month (Glasier et al., 2011; Trussell, Raymond, Cleland, 2014). The risk is two- to three-fold higher among women who have other episodes of sexual intercourse in the same cycle that EC was used (Cheng, Che, Gülmezoglu, 2012). Moreover, EC can be the precursor of the regular contraceptive use; a concept referred to as “bridging” (Cameron, Glasier, 2010). In this context, Michie et al. (2014) noted that almost half (44%) of women in Edinburgh who not using a hormonal method of contraception at the time of EC wished to use an effective method. In this context, while EC is not an effective option to replace regular contraception for women who have regular intercourse (WHO, 2004), it may still act as a bridge to regular contraception among the 32.2% sexually active Brazilian women at risk of pregnancy who do not use a regular method (Brasil, 2010; Mosher, Jones, 2010; Trussell, Raymond, Cleland, 2014).

A number of studies indicate greater use of regular contraception after EC exposure; however, studies examining the type of contraceptive use after EC exposure (highly effective method or non–highly effective method) report inconsistent results. A prospective study of EC users in France show that while 8% of women switched from very effective to lower effective or nonuse of contraception after using EC, 20% of women moved in the other direction, that is they switched from less effective or no method to very effective method use after using EC (Moreau et al., 2009b). Also, in another study in France, Moreau et al. (2006) observed that 84% of women resumed use of a regular contraceptive after EC use, and almost 50% of women used highly effective method following EC use. Other study conducted in Jamaica, highlighted that 55% of women adopted an ongoing method of contraception following their first use of EC (Sorhaindo et al., 2002). In Minnesota, study found that 68% of women who were using non–highly effective method, switched to hormonal contraception after EC use (Sidebottom et al., 2008).

Other studies however, find the reverse is true. A case control study conducted among high risk women (16-24 years old) in California, indicate that women provided advanced supplies of EC were more likely than those in the control group (not given EC) to use less effective and less consistent contraception (Raine et al., 2000). Cameron et al. (2011) observed that despite the high use of contraception after the EC use among Scottish women (88%), the use of effective ongoing contraception after EC (even from a specialist service) was not high. A case control study conducted among women aged 15–20 in Pennsylvania found higher reported condom use (77%) in the advance EC group at follow-up (MacLean, 2004). Similarly, a case control study among Brazilian women aged 18-49, found a significant increase in regular contraceptive use among women who used EC; however almost 64% of women chose non–highly effective methods for ongoing contraception (Costa et al., 2008).

It is important to highlight that after EC use; some women do not resume use of regular contraception. In the French cohort of EC users, Moreau et al. (2009b) observed that 11% of these women did not use contraception six months after EC use. Similarly, 12% of women in Scotland did not use a regular contraception after EC use (Cameron et al., 2011). One factor that may probably contribute to the lack of bridging to prescription contraception is that women access EC in the pharmacy are not given proper counseling about more effective contraception (ESHRE Capri

Workshop Group, 2015). According to Glasier et al (2010), less than half (43%) of pharmacists supplying EC gave counseling about the importance of using contraception after EC. Although EC is prescribed only by doctors and nurses in Brazil, it is sold without prescription in pharmacies. Study based on literature review which aimed to assess the policies for EC provision in pharmacies in the Brazilian context, observed that EC is also sold without any counseling by the pharmacists, since there is no specific policy in this regard (Paiva, Brandão, 2012). Even though counseling on EC by pharmacists is relevant, it is important to acknowledge that they also have doubts and misinformation (Brandão et al., 2016).

One randomized trial conducted in Jamaica gave a discount coupon for obtaining oral contraception in pharmacy for all women requesting EC (Chin-Quee et al., 2010). The intervention had no positive impact on use of effective contraception after using EC, as only 1% of coupons were redeemed, and only 13% of women in the intervention group and 11% of the control group used the oral pill for 2 months or more after requesting EC from a pharmacy. More than two-thirds of the women in the intervention group indicated that they did not want to use the oral pill. In a pilot study conducted in Edinburgh, women selected in pharmacies were randomized according to three groups: standard care (pharmacists provided EC and verbal advice about the importance of using contraception after EC), one-month supply of progestogen-only oral contraceptive pills (POP), or an invitation to present the empty EC packet to a family planning clinic for contraceptive counseling. The proportion of women using effective contraceptive methods at follow-up was significantly greater in both the POP group and family planning clinic group, compared with standard care (Michie et al., 2014). The authors suggest that pharmacy-based interventions can increase the uptake of effective contraception after using EC.

1.5 RATIONALE FOR STUDY

The issue of contraceptive discontinuation is an important contributing factor in unintended pregnancy and induced abortion, and hence in the mortality and morbidity associated to reproduction. University students represent a singular group with both high need for contraception, high prevalence of contraceptive use, but high rates of contraceptive discontinuation resulting in frequent unintended pregnancy. In

case of unprotected sexual intercourse, many of them use EC; however, there is little information about the circumstances of EC use after method discontinuation and the role of EC in bridging to more effective contraception in this population

No study has been done in Brazil focused on exploring the contraceptive discontinuation and their relation to the use of EC. The latest data on contraceptive discontinuities dates back to the 1996 DHS survey (Brazil, 2008a) with little insights on how overall practices and user dynamics have evolved ever since. These estimates are now over 20 years old and may not be an accurate reflection of young Brazilian women's current experience in using reversible contraceptive methods.

The present work based on two studies carried out with university students at University of São Paulo (Pirotta, Schor 2004; Borges et al., 2010) showing the use of contraception in this group appeared to be discontinuous, but this was not measured. We extend on these findings, by exploring contraceptive discontinuation, recourse to EC and its effect on bridging to more effective methods among college students.

1.6 STRUCTURE OF STUDY

This thesis is presented in seven chapters which are briefly detailed in this section. The first chapter reviews the literature and establishes the theoretical framework, presents the study rationale and outlines the topics that are addressed in the thesis.

The second chapter presents the methods used across studies, including the study population, survey site, and data collection procedures. We also present a detail description of the measures that are used in each chapter and the statistical analysis that was followed.

Each of the next three chapters (chapter 4, 5 and 6) presents a specific aim of the thesis. The chapters are organized by summarizing the important literature relevant to each specific topic, the method used for the specific analysis, description and discussion of specific topic of interest (discontinuation, EC use after discontinuation, contraceptive patterns of use surrounding EC). A summary of the three chapters and their associated data and methods is provided in Frame 1 below.

Ultimately, the thesis is concluded in chapter 7 by a summary of the results of the study as a whole, along with policy implications and suggestions for further studies.

Frame 1 – Summary of three chapters

CHAPTER	4	5	6
Data	Retrospective cohort study (longitudinal survey data).	Retrospective cohort study (longitudinal survey data).	Retrospective cohort study (longitudinal survey data).
Method	Generalized estimating equations and multinomial logistic regression.	Descriptive analysis	Multivariate logistic regression, and multinomial logistic regression.
Study site	University of São Paulo, Brazil.	University of São Paulo, Brazil.	University of São Paulo, Brazil.
Population	Women aged 18-24 who were enrolled in undergraduate course of all campuses at University of São Paulo, Brazil.	Women aged 18-24 who were enrolled in undergraduate course of all campuses at University of São Paulo, Brazil.	Women aged 18-24 who were enrolled in undergraduate course of all campuses at University of São Paulo, Brazil.
Sample	<p>Women level sample: 1,679 women who used a contraceptive method within 12-months prior to the study.</p> <p>Monthly discontinuation of contraception sample: 18,469 observation periods (months).</p> <p>Contraceptive episode level sample: 1,230 new episodes of contraception use.</p>	<p>Women level sample: 916 women who used EC in lifetime, and 449 women who used EC within 12-months prior to the study.</p> <p>Monthly level sample: 842 months of discontinuation for method-related reasons, 428 months of method abandonment, 485 instances of EC in the last 12-months.</p>	<p>Women level sample: 916 women who used EC in lifetime, and 691 women who used contraception after EC use.</p>

2.CHAPTER TWO

CHAPTER TWO

DISSERTATION AIMS, HYPOTHESIS AND CONCEPTUAL FRAMEWORK

2.1 DISSERTATION AIMS

Our specific aims are as follows:

- 1) Assess frequency and correlates of method discontinuation among undergraduate women in São Paulo within a 12-months period, according to the number of times women discontinued.
- 2) Assess frequency and correlates of method-specific contraceptive discontinuation among undergraduate women in São Paulo within a 12-months period.
- 3) Evaluate frequency of EC use after contraceptive discontinuation among undergraduate women in São Paulo. Assess differences in contraceptive patterns following discontinuation for women who use EC after discontinuation and those who do not use.
- 4) Describe contraceptive patterns (frequency and correlates) within a month after using EC among undergraduate women in São Paulo.

2.2 HYPOTHESIS

- 1) We hypothesize that the frequency of contraceptive discontinuation within a 12-months period varies by women's sociodemographic and partnership characteristics; and that women who discontinue once and several times differ according to these criteria.

2) We hypothesize that the frequency of contraceptive discontinuation within a 12-months period varies by type of method, and it is frequent among user dependent methods.

3) We hypothesize that women take rarely advantage of EC after discontinuing contraception, and that EC use varies across sociodemographic and partnership characteristics.

4) We hypothesize that women resume use of effective contraception within a month after EC use, and they do not have changes in contraception after EC use. In addition, we hypothesize that women have gaps in contraception within a month after EC use, which vary across sociodemographic and partnership characteristics.


2.3 THEORETICAL FRAMEWORK

Figure 1 is a diagrammatic representation of the conceptual framework of contraceptive discontinuation dynamics and EC use on which this study is based. The figure demonstrates the process of contraceptive use as it occurs through time. First, at step 1 an individual can make the choice to use a specific contraceptive method, and therefore, an episode of use starts. This choice is influenced by a number of factors not only external to the individual such as supply environment, but also individual characteristics such as sociodemographic characteristics, type of relationship, the motivation to avoid pregnancy, among others.

Step 2 represents the length of the episode which, in the case of this study, can be anywhere from 1 to 12 months long. The line between step 1 and step 2 denotes the decision of the user to either continue their method, to switch to a different method or to abandon their method. Again, numerous factors may influence this decision and the central aim of this study is to understand which determinates are relating to contraceptive discontinuation among young women in Brazil. Also, step 2 exemplifies the importance of using EC at the time of switching or abandoning the method, which can be for any reasons or for method-related reasons. For those individuals who discontinue methods they are immediately returned to step 1 and the cycle begins again.

Step 3 represents the possible consequences of contraceptive discontinuation and no EC use, such as unintended pregnancy, consequently induced abortion, and finally the maternal mortality. Each individual may complete this cycle of periods of use and non-use many times and may take similar or distinct pathway.

Figure 1 – Conceptual framework illustrating contraceptive discontinuation and their relation to the use of emergency contraception



3.CHAPTER THREE

CHAPTER THREE

METHOD SECTION

3.1 OBJECTIVE OF THE SURVEY

This 12-month retrospective cohort study was conducted among a random sample of undergraduate women from University of São Paulo, Brazil, between June and July, 2015. The main objective of the study was to analyze contraceptive discontinuation and EC use. Women between the ages of 18 and 24 years old and enrolled in regular undergraduate course in all campuses of University of São Paulo, Brazil were eligible to participate in the study. They were selected by simple random sampling without replacement, and they were chosen electronically based on a list of all the email addresses of undergraduate students enrolled in the Electronic Computer Center of the University of São Paulo, organized in alphabetical order.

3.2 SURVEY SITE

The University of São Paulo is considered the largest institution of higher education and research in Brazil, under the responsibility of the São Paulo state government. The University of São Paulo offers undergraduate courses in all areas of knowledge. All courses are free and last from four to six years. There are eight campuses at University of São Paulo: one in São Paulo, capital, and seven in the state, in the cities of Bauru, Piracicaba, Pirassununga, Ribeirão Preto, São Carlos, Santos and Lorena (Figure 2). Altogether, there are 42 teaching units, offering 300 undergraduate courses in which 59,081 undergraduates were registered in 2015. According to data from the Statistical Yearbook of 2014, the university had 27,396 female students.

Figure 2 – Map showing the distribution of the campuses at University of São Paulo, Brazil in São Paulo State



3.3 SAMPLING (FLOWCHART) AND DATA COLLECTION PROCEDURE

The sample included 1,026 undergraduate women, however after the pre-test (n=50), only 10% of women answered the questionnaire; a response rate that was far lower than found in other surveys that used online questionnaires in Brazil (Bastos et


al., 2008; Hooper, 2010). Therefore, considering the low response rate in the pre-test, the sample size was tripled, and thus 3,078 women were randomly selected and invited to participate in the study.

After obtaining ethical approval to conduct the study from the Research Ethics Committee at University of São Paulo School of Nursing (CAAE Number: 39142514.4.0000.5392) (ANNEX I) and an authorization from the pro-rector at University of São Paulo, an electronic message was sent to all students who were selected to participate in the study. The research project (objectives, methodology, questions and time to answer the questions) was described in the email, and the student who had an interest in participating in the study could click on the study to complete a consent form (APPENDIX A).

In total, 2,182 undergraduate students (71%) accessed the online questionnaire. However, 453 students who answered the questionnaire were not included because they did not meet inclusion criteria. In particular, students under the age of 18 years old (n=2), students who had never had sexual intercourse (n=358), and students who did not use any contraceptive method in the 12 month prior to the survey (n=93) were excluded. The first group were excluded because it was necessary to obtain a permission from parents or guardians to participate in study, the second group because they were not at risk of pregnancy in the last 12 months, while the latter group was excluded because they could not discontinue a method. In order to analyze only women who were at risk of becoming pregnant, other situations such as female sterilization and "want to become pregnant" were considered, however no woman reported these conditions. In addition, 50 women refused to participate in the study for different reasons, such as "did not want to answer questions related to sexuality" (n=10), some women felt "inadequate to participate because they were lesbians" (n=12), some women answered that the theme of the research was against their religion (n=5), others did not report the reason (n=23). Therefore, the final study population comprised 1,679 undergraduate students between 18 and 24 years of age (Figure 3).

Students received reminder emails weekly to participate in the study (on average 10 times). Moreover, 14 students did not finish answering the questionnaire and had to answer the questions again at another time because uncompleted questionnaires were canceled.

Figure 3 – Flowchart of organizational structure of the women interviewed, according to sexual behavior and contraception use. São Paulo, 2015



3.4 QUESTIONNAIRE

The data were collected online through *Google Form*, using a self-administered questionnaire that was pre-tested (APPENDICE B AND C). The questionnaire contained sociodemographic questions; questions about sexual and reproductive antecedents, questions about the use of contraceptive methods in the 12 months prior to the survey; and issues about the use of contraceptives surrounding the use of emergency contraception in the past 12 months. On average, it took 5-7 minutes to complete the questionnaire.

Questionnaire was structured with closed questions, and divided into modules:

Frame 2 – Description of the survey questionnaire

MODULES	OBJECTIVES
Sociodemographic context	Identify the sociodemographic profile of the individual, such age, race/ethnicity, religion, marital status.
Socioeconomic context	Identify the socioeconomic level of the individual. The socioeconomic status was calculated according to Brazil's Criterion of Economic Classification based on four questions about household possessions (refrigerator, television, computer, etc..) educational level of the head of household, and public services near the residence (paved street and piped water) (ABEP, 2015).
Scholar context	questions about school environment, such as the field of study, part time or full time enrollment, campus and scholarship.
Sexual and reproductive behavior	questions explored past sexual and reproductive behavior, such as age at first sexual intercourse, pregnancy history, age at first pregnancy, previous abortions, among others.
Contraceptive behavior	Questions were based on current and last use of contraception.

(continua)

(continuação)

MODULES	OBJECTIVES
Contraceptive calendar	<p>First, a question identified women who had used contraceptives in the 12 months prior to the study, reasons for non-use contraception were asked of women who had not used contraception during that time period.</p> <p>Second, women who reported using contraception in the past 12-months were asked to complete a contraceptive calendar on a month-by-month basis (Bradley, Schwandt, Khan, 2009). For each month, they indicated the type of contraception used, reasons for discontinuation their method, as well as the type of relationship, and occurrence of pregnancy. In addition, they were asked about the use of EC for each month.</p> <p>Monthly use of contraception was assessed based on the question: <i>“What type of contraceptive were you using in each month during the 12 previous months?”</i> This question was followed by a number of contraceptive methods from which women could select just one method for each month or they could indicate that they had not used a method. Additionally, women were asked about the use of EC for each month based on the questions: <i>“Did you use emergency contraception in the past 12-months?”</i> <i>“In positive case, what months did you use it?”</i></p>
Emergency contraception	<p>Questions explored circumstances of last EC use, including the type of relationship when EC was used, who bought the medicine, prescription, site purchase location, time between sexual intercourse and use of EC, among others.</p>
Emergency contraception calendar	<p>The contraceptive circumstances surrounding last EC use were also explored 30 days before and 30 days after the last use. Women were asked what type of contraception they mostly used within 30 days before and 30 days after the last time they used EC.</p> <p>The specific questions included: <i>“Did you use contraception during the 30 days before/after of using emergency contraception?”</i>, <i>“What type of contraception did you use in most of these 30 days (before and after EC use)?”</i>, <i>“Did you use the method continuously?”</i> If women answered no to this later question, an EC calendar was made available, asking about use or non-use of contraceptive method for each day (30-days). The main question of the EC calendar was: <i>“In these 30 days before/after using the emergency contraception, which days did you use the contraceptive method?”</i></p> <p>Questions only allowed “no/yes” answers so it was only possible to evaluate nonuse or abandonment of contraception before/after EC use, not switching among methods.</p>

(conclusão)

3.5 MEASURES

To explore discontinuation, each woman had a full one-year period of observation; however, we could not assess method duration at the time of discontinuation, because the date that women started using contraception was not collected in this study. For this reason, this study focuses on frequency of discontinuation within a 12-months period.

With respect to the EC calendar, each woman had two-months of observation (30-days before EC use and 30-days after EC use). Similarly, to the contraceptive calendar, the start date of using contraception before EC was not available, therefore we explore the prevalence of discontinuation one month before and one month after using EC.

3.5.1 Unit of analysis: contraceptive discontinuation

We considered three levels of analysis of contraceptive discontinuation, women level analysis, monthly discontinuation of contraception analysis, and contraceptive episode level analysis.

- Women level sample: we considered the full sample of 1,679 women for which we assessed the frequency of method discontinuation within 12 months.
- Monthly discontinuation of contraception sample: we partitioned the contraceptive calendar into monthly observations. Each woman contributed up to 12-months observations. A total of 18,469 months of observations were identified and are used as the analytical sample to investigate the factors related to monthly discontinuation of specific contraceptives.
- Contraceptive episode level sample: we partitioned the contraceptive calendar into segments of contraceptive use, defined as continuous months of use of the same method. Each woman contributes one or several episodes of contraception, based on the number of times she switches within the calendar year. A total of 3,937 episodes of use were identified. Among these, 1,230 were started during the calendar (new episodes), and are used as the analytical sample to investigate probabilities of discontinuation of new episodes of contraception over time.

3.5.2 Outcome Variables

3.5.2.1 Contraceptive discontinuation

The dependent variables, i.e., variable whose value depends upon one or more other variables (response variable) for this research are constructed from the following central question which was asked of all women in the selected sample:

“What type of contraceptive were you using in each month during the 12 previous months?”

The response was based on the method women used in each particular month. We inferred method discontinuation by identifying when methods differed in two consecutive months (“month n” versus “month n+1”). We defined a monthly dichotomous discontinuation variable coded yes/no based on information that a method was not used in the following month.

In this study, the term “discontinuation” included women who switched to another method and those who abandoned their methods. Women were considered switchers if they initiated a method within a month of discontinuing their previous method. On the other hand, we considered discontinuation to be abandonment if the women did not use any other method during the follow-up period or resumed use of any method after experiencing at least one month of nonuse.

We defined three types of contraceptive discontinuation, i.e., discontinuation for any reason, discontinuation for method-related reasons (excludes discontinuation among women who don't need contraception, because they are seeking to get pregnant, are sterile or no longer gave a partner), and abandonment or switching to less effective method while still at risk to become pregnant (excludes discontinuation among women who don't need contraception and women who switched to more effective methods). We considered the following switched to represent a decline in contraceptive effectiveness: male condom to withdrawal/ fertility-awareness/ spermicide; oral pill to male condom/ withdrawal/ fertility-awareness/ spermicide/ diaphragms; IUD to male condom/ injectable/ oral pill/ spermicides/ vaginal ring/ patch/ diaphragms/ withdrawal/ fertility-awareness; injectable to male condom/ oral pill/ spermicides/ diaphragms/ vaginal ring/ patch/ withdrawal/ fertility-awareness; and withdrawal/ fertility-awareness to spermicide (CDC, 2013).

Altogether, three different dependent dichotomous variables were created at the monthly level: “discontinuation for any reason”, “discontinuation for method-related reasons” and “abandonment or switching to less effective method”.

Month discontinuation information was also summarized at the women level by a dichotomous variable “any discontinuation in the 12 months of observation” coded yes if a woman had at least one month of discontinuation. Additionally, a three category variable was created at the women level distinguishing women who had never continued a method during the 12 month, women who had discontinued one time and women who had discontinued several times during the 12 months of observation.

All the dependent variables are detailed below in Frame 3.

Frame 3 – Coding of the dependent variables

DEPENDENT VARIABLES	CODE
Discontinuation for any reason	<p>0: non-discontinuation (woman always used the same method of contraception during the 12-months).</p> <p>1: all situations of discontinuation.</p>
Discontinuation for method-related reasons	<p>0: non-discontinuation for method related reasons (woman always used the same method of contraception or only discontinued their method because they were no longer at risk of unintended pregnancy (seeking pregnancy, sterile or had no partner).</p> <p>1: women who discontinued for method-related reasons (discontinued their method while still at risk of unintended pregnancy, including switching to more effective method, or switching to less effective method or abandonment).</p>
Abandonment or switching to less effective method	<p>0: non-abandon (woman always used the same method of contraception or only discontinued their method because they were no longer at risk of unintended pregnancy (seeking pregnancy, sterile or had no partner) or switched to more effective methods.</p> <p>1: women who abandoned their methods or switched to less effective method while still at risk.</p>

(continua)

(continuação)

DEPENDENT VARIABLES	CODE
Frequency of discontinuation for any reason	<p>0: woman did not discontinue their method over the 12-months period.</p> <p>1: woman discontinued their method for any reason one time over the 12-months period.</p> <p>2: woman discontinued their method for any reason more than once over the 12-months period.</p>
Frequency of discontinuation for method-related reasons	<p>0: woman did not discontinue their method.</p> <p>1: woman discontinued their method for method-related reasons one time over the 12-months period.</p> <p>2: woman discontinued their method for method-related reasons more than once over the 12-months period.</p>

(conclusão)

3.5.2.2 Emergency contraception based on contraceptive calendar

The coding of the variables related to EC are presented in Frame 3.

Emergency contraception following discontinuation for method-related reasons, and method abandonment

To identify instances of EC use following discontinuation for method-related reasons and method abandonment, we used data collected in the 12-months contraceptive calendar. In the calendar, women could choose EC as one of the contraceptive used in a particular month. Based on this information, we identified EC use in the month following contraceptive discontinuation for method-related reasons and method abandonment. In addition, we identified EC use following inconsistent use or in the context of non-regular contraception.

The dependent variables related to EC were defined as dichotomous measures (no/yes), which are detailed below in Frame 4.

Frame 4 – Coding of the dependent variables assessing EC use after contraceptive discontinuation

DEPENDENT VARIABLES	CODE
Emergency contraception use following discontinuation for method-related reasons	<p>0: woman did not use EC in the month following discontinuation for method-related reasons.</p> <p>1: woman used EC in the month following discontinuation for method-related reasons</p>
Emergency contraception use following method abandonment	<p>0: woman did not use EC in the month following method abandonment.</p> <p>1: woman used EC in the month following method abandonment.</p>
Emergency contraception use following inconsistent use	<p>0: woman used EC in a month of inconsistent use of contraception (switching and continuous use).</p> <p>1: woman did not use EC in a month of inconsistent use of contraception; woman used EC following non-regular contraception (abandonment and non-regular contraception).</p>

3.5.2.3 Contraceptive patterns of use in the month after emergency contraception use based on emergency contraception calendar data

To identify contraceptive utilization after using EC, we used a second contraceptive calendar, referred to the EC calendar, exploring contraceptive use within 30 days before and 30 days after the last instance of EC use. Only women who responded positively to ever having used EC: *“Have you ever used emergency contraception?”*, were selected to answer the EC calendar. The dependent variable is based on responses to the following questions:

“What method did you most use within 30 days after the last time you used emergency contraception?”

“Did you use the method continuously?”

Our outcome variable measures contraceptive use over the full period of 30 days after EC use, the last time EC was use.

The dependent variables related to EC were defined as a dichotomous measure (no/yes), which are detailed below in Frame 5.

Frame 5 – Coding of the dependent variables

DEPENDENT VARIABLES	CODE
Contraceptive use after using emergency contraception in the last time	<p>0: woman did not use contraception within 30 days after EC use in the last time.</p> <p>1: woman used contraception within 30 days after EC use in the last time.</p>
Contraceptive switching patterns after emergency contraception use	<p>0: woman did not change the contraceptive method after EC use in the last time; or women did not use a method before and after EC use.</p> <p>1: women switching to more effective method after EC use in the last time; or women were not using a method within 30 days before EC and started to use a method after EC use.</p> <p>2: women switching to less effective method after EC use in the last time; or women were using a method within 30 days before EC and stopped to use it after EC use.</p>
Gaps after using emergency contraception in the last time	<p>0: woman did not have gaps within 30 days after using EC in the last time.</p> <p>1: woman had gaps within 30 days after using EC in the last time.</p>

3.5.3 Independent Variables

Variables were selected on the basis of our conceptual framework related to contraceptive discontinuation and emergency contraception and previous literature reporting on the determinants of contraceptive discontinuation and EC use.

In all analysis, we considered the following sociodemographic and sexual health information

- Sociodemographic variables included:
 - age (18–19 or 20–24);
 - race/ethnicity (white, black, other);
 - religious affiliation (no religion, roman catholic, kardecista spiritualism doctrine, evangelical, other religion);
 - relationship status (none; steady relationship, such as with boyfriend or living with a partner; and casual relationship).
 - wage labor (no/yes);
 - socioeconomic status (A/B, C – D/E) (ABEP, 2015). We stress that we classified the socioeconomic status as high (A/B), and middle/low (C – D/E).
 - Educational information: field of study (human sciences, health sciences, exact sciences); type of enrollment (full-time or part-time); campus site (São Paulo, countryside).

- Sexual and reproductive health variables included:
 - time since first sexual intercourse (≤ 1 year, 2-3 years, ≥ 5 years);
 - number of sexual partners (1 partner, 2-3 partners, ≥ 4 partners);
 - pregnancy prior to 12-months calendar
 - current use of contraceptive method (no/yes) & type of current contraceptive method (highly effective method or non-highly effective method). Highly effective method includes hormonal contraception and an intrauterine device, while non-highly effective method includes condoms, spermicides, fertility-awareness and withdrawal);
 - use of contraceptive method in the last sexual intercourse (no/yes) & type of contraceptive method used in the last sexual intercourse (highly effective method or non-highly effective method);
 - Ever use of EC.

Time varying information collected at each month (12-month) calendar was also considered in the analysis of monthly discontinuation of contraception and EC use following method discontinuation:

- Monthly use of contraception described in 4 categories (oral pill, male condom, other methods, no contraception);
- Monthly relationship status defined in 3 categories: steady, casual relationships, none; and change in relationship status yes/no from previous month;

- Reasons for discontinuation if a method was discontinued (side effects, wanted a more effective method, criticism of the partner, too expensive or difficult to obtain, another reason).

In addition to the information described above, specific information about EC was considered in analysis of EC use following discontinuation. It included:

- Knows anyone who has used EC (no/yes);
- Past use of EC, including number of times (once, twice, three or more times) and timing of last use (up to two years ago, three or more years ago, do not remember);
- Ever had unprotected intercourse and not used EC (no/yes);
- Contraceptive method used at the time of EC use (oral pill, condom, withdrawal/fertility-awareness, injectable, other methods, no contraception).

Descriptive information about EC was also used to illustrate the circumstances of EC use. For each instance of EC use, we considered the following information:

- Reasons for using EC (by sheer insecurity, forgot to use some contraceptive method, the contraceptive method that was using failed, I did not want to use any contraceptive method, I was a victim of sexual violence);
- Person who took the initiative to use EC (woman, partner, both);
- prescription for EC (no/yes);
- EC source (pharmacy; boyfriend who brought; private doctor/health insurance; and national health system, which included primary health care and the Unified Health System, SUS).
- Timing of EC use after unprotected intercourse (<24 hours, 1 – 5 days; ≥6 days, do not remember).

3.6 STATISTICAL ANALYSIS

3.6.1 Descriptive results

Descriptive statistics of the study sample were conducted according to the sociodemographic characteristics, educational background, and sexual/reproductive/contraceptive indicators. In addition, we conducted a descriptive

analysis of the sample according to type of contraceptive method used at baseline (oral pill, male condom, and other methods, i.e., methods less used among Brazilian women such as: injectable, withdrawal/fertility-awareness, patch, intrauterine device, diaphragms, implants, vaginal ring, female condoms, and spermicide). “Baseline”, was defined as the earliest month in the 12-months calendar, i.e., the month farthest back in time. For women who reported using dual protection (oral pill and male condom), we prioritized the most effective method based on failure rates (Trussell, 2011; CDC, 2013), that is, they were included in the group “oral pill users”.

3.6.2 Contraceptive discontinuation analysis

Analyses of contraceptive discontinuation were performed in several steps: women level analysis, monthly discontinuation of contraception analysis, and contraceptive episode level analysis.

Women level analysis

Using women as the unit of observation, this analysis focused on frequency of discontinuation in the 12 months of observation.

1) To compare women who never discontinued their contraception over the last 12 months, to those who discontinued once and those who discontinued more than once, a bivariate analysis was conducted to investigate women’s sociodemographic and sexual reproductive health characteristics according to the frequency of method discontinuation within the last 12-months. We consider two measures of discontinuation as previously described: discontinuation for any method, and discontinuation for method-related reasons. Abandonment or switching to less effective method was not contemplated in this analysis due to the small number of women who had switched or abandoned their method more than once (n=20). Pearson’s Chi-square test was used to assess the relationships between women’s characteristics and method discontinuation.

2) The analysis of the determinants of contraceptive discontinuation according to the number of times a method was discontinued, was carried out using multinomial logistic regression, as the dependent variable was defined in three categories: no discontinuation (reference), 1 discontinuation, and > 1

discontinuation (high rank of discontinuation). Multivariate analysis included the following independent variables: age, and race/ethnicity as well as factors that were the most likely associated with method discontinuation in bivariate analysis (p value <0.10) The criteria for including variables in the multinomial analysis were not only those that showed significance in the bivariate analysis, but also variables based on conceptual framework, that is, factors identified by previous studies as related to contraceptive use and discontinuation (as age, race/ethnicity, and religion). *Previous pregnancy* was not included in the multinomial analysis due to the small number of women reporting a previous pregnancy. The final model was chosen by using the Akaike Information Criterion (AIC), which measures the relative quality of statistical models, i.e., how well a model fits a dataset. Adjusted odds ratios and 95% confidence intervals were estimated.

Monthly discontinuation of contraception analysis

Using calendar months as the unit of observation, this analysis focused on monthly probability of contraceptive discontinuation.

1) Since the risk of pregnancy is higher among women who discontinue contraception but are still at risk of an unintended pregnancy, we evaluated reasons for method discontinuation for all monthly instances of discontinuation, by distinguishing: discontinuation for any reason, discontinuation for method-related reasons, and abandonment or switch to a less effective method. These analyzes were stratified by method type (oral pill and male condom – main contraception utilized among Brazilian women). There were too few episodes of other barrier methods to perform a stratified analysis for this third category.

In the absence of a starting date relative to the method reported the first month of the calendar, it was not possible to estimate duration of use, and therefore we focused our analysis on monthly probabilities of discontinuation irrespective of the time women had been on the method. We exclude the last month of calendar data since information about discontinuation the following month is unavailable (“right-censored”).

We evaluated the factors related to each of the “measures” of discontinuation for pills and for condoms using Chi-square tests (with a significance threshold adopted was 5% ($p \leq 0.05$)).

Because women contributed 12 monthly observations, i.e., analysis of monthly discontinuation need to take into account correlations between months contributed by the same women. Accordingly, we used Generalized Estimating Equations (GEE). GEE is utilized in longitudinal data, and provides adjusted estimates for multiple observations contributed by the same individual (Liang, Zeger, 1986). Specifically, GEE is based on generalized linear models, and includes a *working correlation matrix* between observations belonging to the same individual. The specification of the correlation structure was conducted considering three possible structures: exchangeable, unstructured, and auto-regressive. An exchangeable correlation structure means the correlation between observations belonging to the same individual remains the same over time. An unstructured correlation structure means that a different correlation value is assumed for each observation within the group, and the auto-regressive correlation structure is used when the data are correlated over time (Ballinger, 2004; Agranonik, 2009). The correlation structure which was more suited to the data and adopted in the final GEE model was *exchangeable* with logit link transformation function. The selection of the most appropriate correlation structure, was based on the Quasi-likelihood under the Independence Model Criterion (QIC) (APPENDIX D). Adjusted odds ratios and 95% confidence intervals were estimated. We also included factors showing stronger associations in the bivariate analysis.

Contraceptive episode level analysis

Using a new contraceptive episode (segment of use of a particular method without stopping) as the unit of observation, this analysis focused on cumulative probability of discontinuation of episodes started in the last 12 months

- 1) Kaplan-Meier life-table were conducted to estimate the probabilities of contraceptive discontinuation (discontinuation for any reason, discontinuation for method-related reasons, and for abandonment or switching to less effective

method) over time for new users. The analyzes were stratified by method type (oral pill or male condom). Only episodes that started during the calendar (new episodes), are included since the duration of use of methods started before the calendar is unknown (left censoring). Episodes that are still ongoing at the time of the survey were included and treated as right censored. The Log-Rank test was used to assess the differences in discontinuation rates by women's characteristics (Log-Rank test).

2) Cox proportional hazards models were used to estimate factors associate with method discontinuation in multivariate analysis. Separate models stratified by method type (oral pill or male condom), were run for: discontinuation for any reason, discontinuation for method-related reasons, and for abandonment or switching to less effective method. Shoenfeld Residuals test were used to assess the assumption that Hazard Ratios were proportionate over time for all covariates. Adjusted hazard ratios and 95% confidence intervals were estimated.

3.6.3 Emergency contraception analysis based on contraceptive calendar (12-months of observation)

We first explored the circumstances of EC use in last time it was used, among all women EC users and among who last used EC in the last 12-months.

Monthly discontinuation of contraception analysis

We explore factors associated with using EC following contraceptive discontinuation for method-related reasons, and conducted the same analysis focused on EC use after method abandonment during the 12-months prior to the survey. The comparison was between women used EC after discontinuing their method for method-related reasons versus those who did not use EC after discontinuing their method. Similarly, we distinguished women who used EC after method abandonment and those who did not use EC after method abandonment.

Next, we investigated the correlates of EC use in the context of inconsistent use of contraception (it includes instances of EC use following method switching and instances of EC use following inconsistent use of contraception that was not

discontinued), *versus* EC use in the context of non-regular contraception (includes instances of EC use following method abandonment and instances of EC use following month where no contraception was used).

We do not explore EC use after discontinuation for any reason, since some woman stop contraception because they want to become pregnant or do not have partner, in which case EC is not needed.

Pearson's Chi-square tests were conducted to assess the relationships between EC use and each independent variable.

3.6.4 Emergency contraception analysis based on EC calendar (2-months of observation, 30 days before and 30 days after EC use)

Women level analysis

1) First, a bivariate analysis was conducted to investigate women's sociodemographic and sexual reproductive health characteristics according to the use of contraception within 30 days after EC use, using the EC calendar data. Highlights that we consider use of contraception within 30 days before EC as a key predictor of post EC contraceptive. The analysis of the determinants of using contraception within 30 days after EC use, was carried out using multivariate logistic regression. The criteria for including variables in the multivariate analysis were not only those that showed significance in the bivariate analysis, but also variables based on conceptual framework, that is, factors identified by previous studies as related to contraceptive use after EC use (as age, race/ethnicity, religion, socioeconomic status, and time since first sexual intercourse). Hosmer and Lemeshow's Test was used to assess how well the model fits the data. Adjusted odds ratios and 95% confidence intervals were estimated.

2) Second to analyze how EC affects contraceptive trajectories, we assessed changes in contraceptive use before and after EC use, using the EC calendar. We distinguished women who had no change in contraception after EC use (reference), women who switched to more effective method after EC use, and women who switched to less effective method after EC use. We consider use of

contraception within 30 days before EC as a key predictor of post EC contraceptive. After conducting bivariate analysis, multinomial regression models were performed (dependent variable defined as a 3-categorical variable). Adjusted odds ratios and 95% confidence intervals were estimated. The final model was chosen based on prior knowledge and based on Akaike Information Criterion (AIC).

3) Finally, we investigated gaps in contraceptive use within 30-days post EC use, among all women who used contraception after EC, using EC calendar.

We stress that in all analysis, it was considered the last time that EC was used in the EC calendar. Pearson's Chi-square tests were conducted to identify the determinants of contraceptive use after EC and changes in contraceptive use after EC. All analyses were conducted using Stata 14.2 (Stata Corporation, College Station, TX, USA).

4.CHAPTER FOUR

CHAPTER FOUR

CONTRACEPTIVE DISCONTINUATION (CONTRACEPTIVE CALENDAR)

4.1 BACKGROUND OF THE STUDY

4.1.1 Introduction

Brazil has recently undergone a rapid fertility transition with total fertility rates dropping below replacement, from 2.5 in 1996 to 1.8 in 2006 (Brasil, 2008a). At the same time, Brazil has seen a dramatic increase in modern contraceptive prevalence rates, estimated at 77% in the last national survey conducted in 2006 (Brasil, 2008a). However, despite advances in the use of contraception and low unmet need for modern contraception (6%), almost 46% of pregnancies are reported as unintended in Brazil (Brasil, 2008a; Berquó, Lima, 2008), a situation that is comparable to other countries with high contraceptive coverage (Bajos et al., 2014, Moreau et al., 2014; Wellings et al., 2015; Finer, Zolna, 2016).

The highest burden of unintended pregnancies across most developed countries, including Brazil, falls on youth (ages 15 to 24) who are at the peak of their fertile years (Berquó, Lima, 2008; IBGE, 2010; Diamond-Smith, Moreau, Bishai, 2014; Wellings et al., 2015; Coetzee, Ngunyulu, 2015; Finer, 2016; Moreau, Bohet, 2016). Youth population (20 – 24 years old) represent 17.9% of the population in Brazil, but account for 25% of unintended pregnancies (Berquó, Lima, 2008; IBGE, 2010). In case of unintended pregnancies, young women are also more likely to have an induced abortion, with peak of lifetime experience observed between 20 and 24 years (24% in this age group in Brazil) (Diniz, Medeiros, 2010). Social changes towards delaying the first birth are extending the number of years young women are exposed to the risk of unintended pregnancies. Specifically, the proportion of women at first birth aged 30 years has grown in the last decade in Brazil, which means that the gap between age at first sex (which has remained stable) and age at first is growing in some segments of the population (Brasil, 2008a; Brasil, 2014; IBGE,

2015). As a result, there is a growing population of youth exposed to a potential risk of unintended pregnancy.

Most unintended pregnancies are due to inconsistent use or gaps in use of contraception (Goulard et al., 2006; Frost, Darroch, 2008), which raises the issue of method choice and dynamics of use. The contraceptive landscape in Brazil is dominated by oral pills and male condoms that are associated with high discontinuation rates and moderate failure rates (Brazil, 2008a; Kost et al., 2008; Trussell, 2011). In this context, contraceptive discontinuation is defined as switching a method of contraception or stopping a method altogether. Indicators typically discern discontinuation for any reason or for method-related reasons. The latter is potentially more concerning because women are still exposed to the risk of pregnancy when they stop their method (sterile or no partner) (Trussell, Vaughan, 1999). Altogether, 47.1% of all methods are discontinued for method-related reasons within 12-months in the U.S., compared with 27.8% in Brazil (Vaughan et al., 2008; Ali, Cleland, Shah, 2012). Discontinuation rates methods for method-related reasons are particularly high in the case of user dependent methods (Ali, Cleland, Shah, 2012; Alano et al., 2012; Silva et al., 2010; Silva Camargo, Iwamoto, 2014). Many factors are related to discontinuation, including external factors (such as quality of family planning services, family planning service environment, guidelines, among others), and internal factors (such as demographic characteristics, partner, and fertility motivations) (Curtis, Blanc, 1997; Ali, Cleland, 1999; Curtis, Evens, Sambisa, 2011; Barden-O'Fallon, 2011a; Moreau et al., 2013).

Contraceptive use dynamics is of particular relevance to undergraduate students represent with high educational and professional aspirations, who are especially motivated to delay childbearing until they reach professional stability (Longo et al., 2002; Sant'Anna et al., 2008; Veloso et al., 2014). While overall pregnancy rates are lower in this population compared to their less educated peers (Finer, Zolna, 2016), a higher proportion of these pregnancies are unintended (60%), and are mostly related to contraceptive discontinuation (Bryant 2009; Trieu et al., 2011). A study conducted amongst undergraduate students in the U.S. estimated that continuous use of contraception could prevent about 12 million unwanted pregnancies every year among this population (Ersek et al., 2011). In Brazil, studies conducted among undergraduate students in Minas Gerais and Paraná states found that 29.3% and 65.9% of students respectively, had used contraceptive methods inconsistently

despite high levels of knowledge about contraception (Dessunti, Kings, 2012; Silva, Camargo, Iwamoto, 2014). As a result, young people with higher education are 3.8 times more likely to have an induced abortion than less educated youth, which contributes to potential maternal morbidity in the context of restrictive laws on abortion such as in Brazil (Pirodda, Schor, 2004; Menezes, Aquino, 2009; Dessunti, Kings, 2012; Silva, Camargo, Iwamoto, 2014).

In summary, a high proportion of young women are at risk of unintended pregnancies and, among those, university students represent a singular group with both high prevalence of contraception and high rates of contraceptive discontinuation. Investigating contraceptive use patterns among highly educated women allows us to understand such dynamics in a “privileged” population, which is less likely to face access barriers relative to lower socioeconomic populations. The frequency and extent of contraceptive gaps among young women in Brazil nowadays is not well documented. The latest data about contraceptive discontinuation in Brazil dates back to the DHS survey conducted in 1996 (Brazil, 2008a) with little insights on how overall practices and user dynamics have evolved since, and no insights on these issues among young women. The present study focuses on highly educated young women who are most motivated to prevent pregnancy.

4.1.2 Aims of study

The aim of this study is to assess the frequency and correlates of method-specific contraceptive discontinuation within a period of 12 months. We also aim to identify sociodemographic and reproductive health factors related to the frequency of discontinuation (0, 1 instance, >1 instance of method discontinuation) within the last 12 months. The research questions will address are:

- How often do women discontinue their method of contraception in a 12-months period and what are the differences between woman who did not discontinue their method in the period of 12 months, woman who discontinued their method once, and those who had several episodes of discontinuation in the period of 12 months?
- What is the monthly probability of method discontinuation and the correlates of such events?

- What is the cumulative probability of discontinuation of new episodes of contraception and the correlates of such events?

4.2 METHODOLOGY

This section outlines the sample, data collection procedures, measures and the statistical method used for the analysis.

4.2.1 Study population and survey site

A 12-month retrospective cohort study was conducted on a sample of undergraduate women from University of São Paulo, Brazil (the largest institution of higher education and research in Brazil), between June and July, 2015. Women between the ages of 18 and 24 years old enrolled in regular undergraduate course in any campus at University of São Paulo, Brazil, were eligible to participate in the study. They were selected by simple random sampling without replacement, and they were chosen electronically based on a list of all the email addresses of undergraduate students enrolled in the Electronic Computer Center of the University of São Paulo, organized alphabetically. The study obtained ethical approval from the Research Ethics Committee at University of São Paulo School of Nursing (CAAE Number: 39142514.4.0000.5392) (ANNEX I), an electronic message was sent to all students who were selected to participate in the study. The research project was described in the email, and the student who had an interest in participating in the study could click on the study to complete a consent form (APPENDIX A).

4.2.2 Sampling

A total of 3,078 women were randomly selected and invited to participate in the study. These women received an electronic message describing the research project and inviting the student to participate after completing a consent form. Students received reminder emails weekly to participate in the study (on average 10 times).

A total of 2,182 undergraduate students (71%) answered the online questionnaire. However, the final study population comprised 1,679 undergraduate

students between 18 and 24 years of age, since 50 women refused to participate in the study, and 453 students were not eligible for the study (< 18 years old (n=2), had never had sexual intercourse (n=358), and had not used any contraceptive method in the last 12 months (n=93)).

4.2.3 Data collection and study instruments

The data were collected online through *Google Form*, using a self-administered questionnaire that was previously pre-tested and took, on average, 5–7 minutes to complete (APPENDICES B AND C). The survey questionnaire collected information on sociodemographic characteristics, sexual and reproductive antecedents, and use of contraceptive methods in the 12 months prior to the survey. Women who reported that they used contraception in the past 12-months, completed a contraceptive calendar on a month-by-month basis (Bradley, Schwandt, Khan, 2009). In the contraceptive calendar the main methods were listed, and young women could select just one answer, that is, they were asked about the main contraception that they used per month. In addition, for each month in the calendar, the student provided information about the reasons for discontinuation, occurrence of pregnancy and their relational status with a partner.

4.2.4 Unit of analysis

We consider three levels of analysis for this chapter, women level analysis, monthly discontinuation of contraception analysis, and the contraceptive episode level analysis.

- Women level sample: as described above, we consider a sample of 1,679 women for which we assess the prevalence and frequency of method discontinuation within 12 months.
- Monthly discontinuation of contraception sample: we partition the contraceptive calendar into 12-months of contraceptive use. Each woman contributes up 12 months of observation, based on number of months she used contraception. A total of 18,469 months of use were identified and are

used as the analytical sample to investigate the factors related to method-specific monthly probability of discontinuation.

- Contraceptive episode level sample: we partition the contraceptive calendar into segments of contraceptive use, defined as continuous months of use of the same method. Each woman contributes one or several episodes of contraception, based on the number of times she switches within the calendar year. A total of 3,937 episodes of use were identified. Among these, 1,230 were started during the calendar (new episodes), and are used as the analytical sample to investigate cumulative probabilities of discontinuation of new episodes of contraception.

4.2.5 Measures

Contraceptive discontinuation

We define three measures of discontinuation: “discontinuation for any reason”, “discontinuation for method related reasons”, and “abandonment or switched to less effective method”. These indicators are measured for each episode of use of contraception:

- “Discontinuation for any reason” refers to any instance when women stopped using their method regardless of the reason and regardless of what they did next (switched to another method or abandoned contraception).
- “Discontinuation for method-related reason” related to any instance when women stopped using their method while still in need of contraception (with a partner, not pregnant, not trying to conceive, not sterile), regardless of what they did next (switched to another method or abandoned contraception).
- “Abandonment or switched to less effective method” refers to any instance when women stopped using their method while still in need of contraception (with a partner, not pregnant, not trying to conceive, not sterile) and switched to a less effective method or abandoned contraception. We considered that women had switched to a less effective method if they initiated a method that was less effective than their previous method and if they were using the new method within a month of discontinuing the previous one. On the other hand,

we considered that women had abandoned their method if they did not resume contraception within a month of discontinuation.

These three measures represent different levels of risk relative to unintended pregnancy, from lower to higher. Discontinuation for any reason includes women who stop their method because they are no longer at risk of pregnancy (no partner, sterile) or because they want to become pregnant. Discontinuation for method-related reasons only considers as a discontinuation an instance when women stop their method but are still exposed to the risk of unintended pregnancy (women who are no longer at risk are not discontinuing for method-related reasons, and therefore coded “0” for this measure). Finally, “Abandonment or switched to less effective method” only counts as discontinuation instances when women stop their method while still at risk and switch to a less effective method or to no method. Women who are no longer at risk or who switch to a more effective method are not switching to a less effective method or abandoning their method, and therefore coded “0” for this variable. An example of “abandonment or switched to less effective method” would be when women who switch from condom to withdrawal/ fertility-awareness/ spermicide; or who switch from oral pill to condom/ withdrawal/ fertility-awareness/ spermicide/ diaphragms. Method effectiveness is determined based on typical use failure rates that are published in the literature (Trussell, 2011).

We also define a summary measure for each woman indicating if she has discontinued any method in the 12-month period of observation. This summary measure is defined in 3 categories (any discontinuation, discontinued once, and discontinued>1), and it is an indicator measure for each individual woman.

Women’s sociodemographic and reproductive health history measures

We consider women’s sociodemographic characteristics as predictors of contraceptive discontinuation based on previous research findings. Specifically, we include age (18–19 or 20–24); race/ethnicity (white, black, other); religious affiliation (none, roman catholic, kardecista spiritualism doctrine, evangelical, other); socioeconomic status (high (A/B) – middle/low (C/D/E)) (ABEP, 2015).

Because this study focuses on university students, we will consider their educational characteristics. This information includes field of study (human sciences, health sciences, exact sciences); period of study (full-time or part-time); and campus (São Paulo, countryside).

Sexual and reproductive health indicators are shown to be strongly associated with contraceptive discontinuation. In our study, we consider the type of relationship (steady, such as with boyfriend or living with a partner, casual relationship); time since first sexual intercourse (≤ 1 year, 2-3 years, ≥ 5 years); number of lifetime sexual partners (1 partner, 2-3 partners, ≥ 4 partners); pregnancy history (no/yes); current contraceptive method (highly effective method or non-highly effective method).

4.2.6 Statistical analysis

Women level analysis

We first described the sociodemographic and sexual and reproductive indicators of the study sample and describe the distribution and correlates of women's contraceptive behaviors at baseline (first month of contraceptive calendar, recorded 12 months prior to the survey).

We then investigated women's sociodemographic and sexual reproductive health characteristics according to frequency of method discontinuation within the last 12 months. We distinguished women who had discontinued their method once, more than once or who never discontinued (reference). Method discontinuation was defined according to the two measures: discontinuation for any method, discontinuation for method related reasons. We did not investigate the factors related to abandonment or switching to less effective method as only 20 women had switched or abandoned their method more than once. After conducting bivariate analysis, multinomial regression models were performed (dependent variable defined as a 3-categorical variable). The multinomial analysis was adjusted for age, and race/ethnicity. Adjusted odds ratios and 95% confidence intervals were estimated. The final model was chosen based on prior knowledge and based on Akaike Information Criterion (AIC).

Monthly discontinuation of contraception analysis

We investigated the correlates of monthly discontinuation for any reason, method related reasons and due to abandonment or switching to less effective contraception using bivariate and logistic regression techniques. We also conducted this analysis stratified by type of contraceptive method (oral pill or male condom). There were too few months of use of other barrier methods to perform a stratified analysis for this third category. Because women could contribute several months of contraceptive use, such months were not independent of each other. Therefore, after fitting the best logistic model, we conducted the analysis using Generalized Estimating Equation (GEE) modelling that accounts for intra-correlation among the months belonging to the same individual. We used an exchangeable correlation structure with logit link transformation function, i.e., the correlation among the observations of the individuals was based on the correlation matrix, and the value of QIC (APPENDIX D). Adjusted odds ratios and 95% confidence intervals were estimated.

Contraceptive episode level analysis

Kaplan-Meier life-table were conducted to estimate the probabilities of contraceptive discontinuation over time for new users. Separate analysis stratified by method type (oral pill or condom), were run for: discontinuation for any reason, discontinuation for method-related reasons, and for abandonment or switching to less effective method. Only episodes that started during the calendar (new episodes) were considered in this analysis, i.e., episodes started before the 1st month prior to the survey were excluded as the exact date of start was not known (left censoring); whereas the episodes which were ongoing at the time of survey (right censoring) were included and treated as censored. The Log-Rank test was used to assess the differences in discontinuation rates by women's characteristics (Log-Rank test). Lastly, Cox proportional hazards models were performed to evaluate the independent effect of each factor in multivariate analysis. A hazard ratio (HR) below 1.0 means that increases in the covariate reduces the hazard ("risk" or "likelihood") of discontinuation, while a HR above 1.0 increase the probability of discontinuation. Schoenfeld Residuals test were

used to assess the assumption that HRs were proportionate over time for all covariates. Adjusted hazard ratios and 95% confidence intervals were estimated. All analyses were conducted using Stata 14.2 (Stata Corporation, College Station, TX, USA).

4.3 CONCLUSION

We conclude that few young women discontinued their method for method-related reasons; however, we stress that a relative proportion of women abandoned their method or switched to a less effective method. Additionally, we conclude that the frequency of discontinuation within 12-months is low, most of women who discontinued their method for any reason or for method-related reasons, discontinued just once during the 12-months.

We find that contraceptive discontinuation does differ by type of contraceptive method. Oral pill was the most contraceptive method used 12-months prior to the survey, following by male condom. However, young women who were using male condom were more likely to discontinued their method (any reason, method-related reasons and method abandonment); although we also observed (in analysis of new episodes of contraception) that oral pill users were more likely to abandonment or switching to a less effective method when compared with male condom users.

The variables most strongly associated with contraceptive discontinuation for any reason, discontinuation for method-related reasons, and abandonment or switching to a less effective method were type of relationship, number of partners, and sexual experience, which appeared in most analyzes. In addition, socioeconomic status was a significant variable, except for the analysis of new episodes of contraception.

Drawing insights from our outcomes, we believe that a larger comprehending of young's concerns and experience using contraception could help primary health care providers reshape counseling strategies to improve adequate contraceptive continuation. Further researches are needed to explore the issues that influence contraceptive dissatisfaction which leading to contraceptive discontinuation among young people. These issues include women's social and demographic characteristics, which have shown to be associated with contraceptive behaviors in Brazil (Pirota, Schor 2004; Borges et al., 2010).

While method abandonment is an important outcome to study because it leads to immediate risk of unintended pregnancy, a key limitation when analyzing this outcome is that women's choice of new method sometimes is not taken into account. In contrast, method switching is essential within women's reproductive life course because the rate of switching and type of switching negatively or positively affects reproductive health outcomes (Steele, Curtis, 2003). We noted that, on one hand, method switching may compromise women's protection and, as such, make them more susceptible to unintended pregnancy (Grady et al. 2002; Vaughan et al. 2008). Conversely, it may improve women's reproductive health, in general, if they switch from less to more effective contraception.

Contraceptive discontinuation is a particularly significant issue for adolescents and young women, since they tend to have more limited access, and face more barriers than older women in family planning programs, as well as more unpredictable and irregular sexual activity, and probably have lower knowledge about using contraceptive methods properly. Therefore, high quality family planning services are necessary, especially in a country where abortion is restricted, like Brazil. In this context, initiatives to improve young women's knowledge and access to all contraception are required. Adequate, accurate and detailed information on the importance of consistent contraceptive use, and issues around side-effects should be emphasized in order to minimize contraceptive discontinuation, whether during medical consultations or students discussion meetings around the units. Furthermore, researchers could work together with university students to try to solve the major problems about contraceptive and sexual behaviors in this population.

5.CHAPTER FIVE

CHAPTER FIVE

EMERGENCY CONTRACEPTION AND CONTRACEPTIVE DISCONTINUATION (CONTRACEPTIVE CALENDAR)

5.1 BACKGROUND OF THE STUDY

5.1.1 Introduction

Nearly 46% of pregnancies are reported as unintended in Brazil (Brasil, 2008a; Berquó, Lima, 2008). In this context, emergency contraception (EC) can prevent pregnancy after unprotected sexual intercourse (Raine et al., 2005). However, the options of EC in Brazil are limited. EC levonorgestrel pills are the only EC products available in Brazil, introduced in public health programs in 2001 (Brasil, 2010; WHO, 2012). While there are other EC methods that are more effective such, as the intrauterine device, mifepristone EC and ulipristal acetate, these methods were not available at the time of this study (Glasier et al., 2011; Cleland et al., 2012; Trussell, Raymond, Cleland, 2014). In addition, studies have shown that the institutionalization of EC in the Brazilian scenario is not fully consolidated in primary health care system, mainly due to moral and religious factors, and the lack of knowledge from health professionals and users (Costa, Guilhem, Silver, 2006; Souza, Brandão, 2009; Figueiredo, Paula, 2012).

National estimates of EC use are difficult to compare as studies refer to different time frames (lifetime or annual incidence), different study populations (all women of reproductive age, young women, abortion patients) and different time periods extending over 15 years. Following the switch to non-prescription status, 17% of women 15-44 years old in France had ever used EC in 2004 (Moreau, Bajos, Trussell, 2006), 11% of women 15–49 years old has done so in the US in the period 2006–2010 (NHS Contraceptive Services, 2011/12), while 16% of young women (average age 19.5 years) in Italy had ever used EC despite the prescription requirement (Tafari et al., 2012).

In Brazil, according to 2006 DHS, 12% of all women ever used EC, with the highest use rate observed among youths aged 20-24 (18.5%) (Brazil, 2008a). EC use is significantly more frequent among Brazilian undergraduate students, ranging from 42.0% to 50.4% lifetime use (Borges et al, 2010; Silva et al., 2010; Alano et al., 2012). These age patterns are also observed in other developing countries such as France, the U.S. and the U.K. (Martin, 2004; Gainer et al., 2005; Moreau et al., 2006; Lakha F, Glasier, 2006; Escobar et al., 2008; Cameron et al., 2011; Daniels, Jones, Abma, 2013).

Although a substantial proportion of women have already used EC, EC is not used enough to have a public health impact in many countries, that is, EC is not utilized as often as needed to curve unintended pregnancy risk (Hu et al., 2005; Glasier et al., 2010; Palermo, Bleck, Westley, 2014; Nappi et al., 2014). Studies conducted in Scotland, France and Denmark over the last 15 years, observed that nine out of 10 women undergoing an abortion do not use EC to try to reduce the risk of pregnancy, a figure that has remained constant despite greater use of EC in the general population, (Sørensen et al., 2000; Moreau et al., 2005; Cameron et al., 2012).

Researchers have shown that EC is mostly used to compensate for discontinuation and errors of use of regular methods (mainly because of a ruptured condom and missed oral contraceptives) (Falk et al., 2001; Sorhaindo et al., 2002; Moreau et al, 2006). In Brazil, discontinuation for method-related reasons range from 42.6% for male condom to 33.1% for oral pill – the most widely used methods among Brazilian youth (Ali, Cleland, Shah, 2012; Alano et al, 2012; Silva et al., 2010; Silva Camargo, Iwamoto, 2014). In case of contraceptive discontinuation with potential gaps in use, EC becomes especially important and may be used as a backup option (Borges et al., 2010).

A better understanding of EC use in relation to contraceptive use dynamics, including discontinuation is an important step in informing more effective strategies to reduce unintended pregnancies. In the context of legal restrictions to abortion, the prevention of unintended pregnancy is a primary prevention strategy to reduce unsafe abortion, which is estimated at an annual average of 994,465 abortions, and responsible for 12% of maternal mortality in Brazil (Faúndes et al., 2007; Martins-Melo et al., 2014), and thus improve women's health.

Notwithstanding, no study has been done in Brazil focused on exploring EC in relation to the contraceptive discontinuation. The latest data on contraceptive discontinuation dates back to the 1996 DHS survey (Brazil, 2008a) with little insights on how overall practices and user dynamics have evolved ever since.

5.1.2 Aims of study

This study focuses on the factors that are related to EC use. First we exam the context of EC use the last time it was used among women who ever used EC and then focus our analysis on recent use of EC reported in the 12-month calendar. We specifically focus our attention to EC use following contraceptive discontinuation, bridging this analysis with our previous exploration of contraceptive discontinuation among university students in São Paulo. Acknowledging that most women use EC in relation to inconsistent use of a regular method rather than in the context of nonuse of contraception, we also investigate the factors related to EC use following inconsistent use versus EC use related to nonuse of contraception. Our specific research questions are the following:

- What are the contraceptive circumstances and access conditions related to the last time of using EC among women who reported ever using the method?
- What are the contraceptive circumstances and access conditions related to the last time of using EC among women who reported using the method in the past 12-months?
- What are the factors related to EC use in the context of contraceptive discontinuation (for method related reasons of method abandonment)?
- What are the differences between women who use EC following inconsistent use of contraception, versus those who use EC in context of non-regular contraception?

5.2 METHODOLOGY

This section outlines the study population, methods of data collection, the measures and the statistical method used for the analysis.

5.2.1 Study population and survey site

We conducted a 12-months retrospective cohort study with a probabilistic sample of undergraduate female students from University of São Paulo, Brazil. Students were selected electronically by simple random sampling without replacement based on a list of all the email addresses of undergraduate students organized alphabetically. Eligibility criteria to participate in this study included being between 18 and 24 years old, and being enrolled in regular undergraduate course in any of the campuses of University of São Paulo, Brazil. All students who were selected received an electronic message, and those who agreed to participate in the study completed an electronic consent form (APPENDIX A). Further selection criteria included ever having sexual intercourse and having used contraception any time during the last 12 months since the study focuses on contraceptive discontinuation. The ethical approval to conduct this study was obtained from the Research Ethics Committee at University of São Paulo School of Nursing (CAAE Number: 39142514.4.0000.5392) (ANNEX I).

5.2.2 Sampling

Among the 3,078 undergraduate students who were randomly selected and invited to participate in the study, 2,182 (71%) answered the online questionnaire. However, the final study population included 1,679 undergraduate students, since 50 women refused to participate in the study, and 453 students were not eligible for the study (< 18 years old (n=2), had never had sexual intercourse (n=358), and had not used any contraceptive method in the last 12 months (n=93)).

In this analysis, we further identified 4 study samples:

- 1) The first sample comprises women who reported ever using EC (n=916), for research question 1.
- 2) The second sample comprises women who indicated using EC in the last 12-months (n=449) to address question 2.

3) The third sample comprises women who discontinued use of contraception, for research question 3. Building on our previous analysis of method discontinuation, we defined contraceptive discontinuation for method-related reasons as a time when women stopped using their method (switched to another method or abandoned contraception) while still in need of contraception (with a partner, not pregnant, not trying to conceive, not sterile). In addition, we defined method abandonment as women who stopped using their method while still in need of contraception, with no resumption of use within a month of discontinuation. Altogether, 464 women discontinued their contraception for method related reasons and a subsample of 151 of these women abandoned contraception.

4) Finally, we consider all instances of EC use following inconsistent use of contraception (n=432), and all instances of EC use in the context of non-regular contraception (n=53) to address question 4.

5.2.3 Data collection and study instruments

A self-administered questionnaire was constructed and made available through Google Form, which took on average between 5 and 7 minutes to complete (APPENDICES B AND C). The questionnaire collected information on women's sociodemographic characteristics, their sexual and reproductive antecedents, and the contraceptive method mostly used, in the 12-months prior to the survey. Women who reported using contraception in the past 12-months, answered a monthly contraceptive calendar, modeled off of the study by Bradley and colleagues conducted in 2009 (Bradley, Schwandt, Khan, 2009). In addition, we collected information on monthly use of EC in the last 12-months, that is, women were asked if they used EC in each month of the contraceptive calendar.

Independently of using EC in the last 12-months, women were also questioned if they had already used EC in lifetime. In positive case, they were specially asked to describe the circumstances of the last time that they used EC, including the type of relationship when EC was used, who bought the medicine, prescription, site purchase location, time between sexual intercourse and use of EC, among others.

5.2.4 Unit of analysis

Our unit of analysis was different based on our research objective:

- For question 1, we conducted a women level analysis exploring the circumstances in which they used EC for the last time in lifetime.
- For question 2, we conducted a women level analysis exploring the circumstances in which they used EC for the last time in the last 12-months.
- For question 3, we conduct a monthly use level analysis, where we partitioned the contraceptive calendar into 12-months of contraceptive use and specifically selected months of observations following discontinuation for method-related reasons or months of observations following method abandonment for analysis. We identified 842 months of observation that followed discontinuation for method-related reasons, and 428 months of observation that followed method abandonment. These observations are used as the analytical samples to investigate the factors related to EC use following contraceptive discontinuation for method-related reasons or method abandonment.
- For question 4, we also conduct a monthly use level analysis, by selecting all calendar months in which women reported using EC in order to compare instances of EC use resulting from inconsistent use of contraception *versus* instances of EC use related to non-use of contraception or method abandonment.

5.2.5 Measures

EC use

We define four measures of EC use: “EC use following discontinuation for method-related reasons”, “EC use following method abandonment”, “EC use following inconsistent use of contraception”, and “EC use in the context of non-regular contraception”. These indicators are measured for each episode of use of EC:

- “EC use following discontinuation for method-related reasons” refers to instances of using EC when women stopped using their method while still in need of contraception (with a partner, not pregnant, not trying to conceive, not sterile), regardless of what they did next (switched to another method or abandoned contraception).
- “EC use following abandonment” related to instances of using EC when women stopped using their method while still in need of contraception (with a partner, not pregnant, not trying to conceive, not sterile), and they did not resume contraception within a month of discontinuation.
- “EC use following inconsistent use of contraception” refers to instances of using EC when women switching their method while still in need of contraception (with a partner, not pregnant, not trying to conceive, not sterile); and instances of using EC when women use their method continuously. We consider this latter situation as “inconsistent use of contraception,” because we assume that for women to use EC when they use their method continuously, it is because they probably had inconsistencies in the use of their contraception.
- “EC use in context of non-regular contraception” refers to instances of using EC when women stopped using their method while still in need of contraception (with a partner, not pregnant, not trying to conceive, not sterile) and did not resume contraception within a month of discontinuation; and instances of using EC when women were not using any contraception.

Emergency contraception history

For our first and second research questions, to explore the circumstances of the last time that EC was used in lifetime and in the past 12-months, we considered the information based on previous research findings. Specially, we include knows anyone who have used EC (no/yes); past use of EC; ever had unprotected intercourse and not used EC (no/yes); used contraceptive method when using EC (no/yes); contraceptive method used at the time of EC use (oral pill, condom, withdrawal/fertility-awareness, injectable, other methods, no contraception); reasons for using EC (by sheer insecurity, forgot to use some contraceptive

method, the contraceptive method that was using failed, I did not want to use any contraceptive method, I was a victim of sexual violence); prescription for EC (no/yes); EC source (pharmacy, boyfriend who brought, private doctor/health insurance, and national health system, which included primary health care and the Unified Health System, SUS).

In addition, detailed informations about contraception, partner and time of using EC were contemplated, such as ever had unprotected intercourse and not used EC (no/yes); used contraceptive method when using EC (no/yes); contraceptive method used at the time of EC use (oral pill, condom, withdrawal/fertility-awareness, injectable, other methods, no contraception); type of relationship when used EC (steady, such as with boyfriend or living with a partner, casual relationship); and person who took the initiative to use EC (woman, partner, both); number of times EC was used (once, twice, three or more times); timing of last use (up to two years ago, three or more years ago, do not remember); time of EC use after unprotected sexual intercourse (<24 hours, 1 – 5 days, ≥6 days, do not remember).

For our third and fourth questions, to investigate the EC use following discontinuation for method-related reasons and method abandonment; and use of EC following inconsistent use of contraception *versus* in context of non-regular contraception, respectively; we considered the women's sociodemographic characteristics and sexual and reproductive health indicators that showed to be associated with EC use according to previous research:

Women's sociodemographic and reproductive health history measures

We consider women's sociodemographic characteristics as predictors of EC use based on previous research findings. Specifically, we include age (18–19 or 20–24); race/ethnicity (white, black, other); religious affiliation (none, roman catholic, kardecista spiritualism doctrine, evangelical, other); socioeconomic status (high (A/B) – middle/low (C/D/E)) (ABEP, 2015).

Because this study focuses on university students, we will consider their educational characteristics. This information includes field of study (human

sciences, health sciences, exact sciences); period of study (full-time or part-time); and campus (São Paulo, countryside).

Sexual and reproductive health indicators are found to be strongly associated with EC use. In our study, we considered the following correlates of EC use: type of relationship (steady, such as a boyfriend or living with a partner, casual relationship); time since first sexual intercourse (≤ 1 year, 2-3 years, ≥ 5 years); number of sexual partners (1 partner, 2-3 partners, ≥ 4 partners); pregnancy history (no/yes); current contraceptive method (highly effective method or non-highly effective method).

5.2.6 Statistical analysis

We only conducted descriptive analysis due to the small number of women reporting EC use in the last 12-months.

We first explored the circumstances of the last time that EC was used in lifetime, and also in the past 12-months among undergraduate students. Posteriorly, we investigated the correlates of using EC following contraceptive discontinuation for method-related reasons in the 12-months prior to the survey (which it includes instances of EC use following method switching and instances of EC use following method abandonment), and after the correlates of EC use following method abandonment in the 12-months prior to the survey (which it includes only instances of method abandonment). While women could theoretically report several episodes of EC use after method discontinuation, there were only few months of discontinuation reported by the same women, which reduces the issue of inter-correlation between observations.

In the last phase of the analysis we compared factors related to EC use following inconsistent use of contraception (including instances of EC use following method switching and instances of EC use following inconsistent use of the same contraception), to factors related to EC use in the context of non-regular contraception (which includes instances of EC use following method abandonment and instances of EC use following month where no contraception was used).

We performed Pearson's Chi-square tests to assess the relationships between EC use and each independent variable. All analyses were conducted using Stata 14.2 (Stata Corporation, College Station, TX, USA).

5.3 CONCLUSION

We conclude that the majority of undergraduate students in our study probably do not use EC in situations where the method is required, i.e., following discontinuation for method-related reasons and method abandonment, potentially reflecting a lack of pregnancy risk recognition.

Future studies, including qualitative work are indispensable to understand why undergraduate college students are not using EC following discontinuation for method-related reasons and method abandonment, specially in a country with restrictive abortion laws such as Brazil. Moreover, recognizing factors that affect with students' ability or choices to prevent unintended pregnancy are relevant in promoting a healthy college environment.

We stress that it is important for young women to be well knowledgeable about contraception. A study conducted with high school students in Brazil, noted that the majority of adolescents did not obtain information about EC in schools or with health professionals (Chofakian et al., 2014). According to the authors, not only health services but also school environment contributes little to the knowledge of sexuality and reproductive life. Reproductive health education as well as family planning counseling are the tools for responsible sexual behavior among youth (Keletzian et al., 2012). Also, appropriate and effective sexual health measures to support undergraduate students are strongly recommended and could stimulate responsible sexual behavior as a Public Health priority.

6.CHAPTER SIX

CHAPTER SIX

CONTRACEPTIVE PATTERNS IN THE MONTH BEFORE AND AFTER EMERGENCY CONTRACEPTION USE (EC CALENDAR)

6.1 BACKGROUND OF THE STUDY

6.1.1 Introduction

When used after unprotected sexual intercourse within time limits, emergency contraception (EC) can avoid an unintended pregnancy (Trussell, Raymond, Cleland, 2014). It is a backup option to prevent pregnancy, and a potential precursor to regular contraception; a concept referred to as “bridging” (Cameron, Glasier, 2010). Michie et al. (2014) noted that almost half (44%) of women in Edinburgh who were not using a hormonal method of contraception at the time of EC use indicated that they wished to use an effective method. In this context, while EC is not an effective option to replace more effective contraception for women who have regular intercourse (WHO, 2004), it may still facilitate transitions towards more effective contraception among the 32.2% sexually active Brazilian women at risk of pregnancy who do not use a regular method (Brasil, 2010; Mosher, Jones, 2010; Trussell, Raymond, Cleland, 2014).

A number of studies indicate greater use of regular contraception after EC exposure; however, studies examining the type of contraceptive used after EC (highly effective method or non–highly effective method) report inconsistent results. A prospective study of EC users in France shows that while 8% of women switched from very effective to lower effective or to no contraception after using EC, 20% of women moved in the other direction, that is they switched from less effective or no method to very effective method use after using EC (Moreau et al., 2009b). In another study in France, Moreau et al. (2006) observed that 84% of women resumed use of a regular contraceptive after EC use, and almost 50% of women used highly effective methods following EC use. A study conducted in Minnesota, highlighted that 68% of women who were using non–highly effective method, switched to hormonal

contraception after EC use (Sidebottom et al., 2008). Other studies however, find the reverse is true. A case control study conducted among high risk women (16-24 years old) in California, indicates that women provided advanced supplies of EC were more likely than those in the control group (not given EC) to use less effective and less consistent contraception (Raine et al., 2000). Cameron et al. (2001) observed that despite the high use of contraception after EC use among Scottish women (88%), the use of effective ongoing contraception after EC (even from a specialist service) was low. Even when EC bridges to more effective methods, a significant proportion of women remain at high risk of unintended pregnancy after using EC: 11% of women in the French study did not use contraception in the six months after EC use (Moreau et al.; 2009b).

Contraceptive discontinuation is an important contributor to unintended pregnancy and induced abortion in countries with high contraceptive prevalence, as it is associated with gaps in use and inconsistent use of new methods. Discontinuation is a particular issue among young college women in Brazil who are highly motivated to prevent a pregnancy, have high contraceptive coverage, but struggle in their daily management of contraception as exemplified in the high rates of method discontinuation in this population (Bryant 2009; Borges et al., 2010; Trieu et al., 2011; Dessunti, Kings, 2012; Silva, Camargo, Iwamoto, 2014; Somba et al., 2014). In case of unprotected sexual intercourse, a significant proportion (although not a majority) use EC; however, there is little information about their opportunity to bridge to more effective methods. In addition, it is noteworthy that some findings of the studies mentioned were controversial, denoting the need to understand this issue in Brazil.

6.1.2 Aims of study

The aim of this study is to describe contraceptive patterns (frequency and correlates) 1-month after using EC, among female university students in São Paulo. The research questions we will address are:

- Do women use contraception after using EC? What are the correlates of using contraception after using EC?
- Do women switch contraceptives after using EC and what are the switching patterns observed? What are the correlates of method switching after EC use?

- Are there contraceptive gaps after using EC? What are the correlates of continuous use of contraception after using EC?

6.2 METHODOLOGY

This section summarizes the sample, data collection procedures, measures and the statistical method used for the analysis.

6.2.1 Study population and survey site

This study is part of a larger project exploring contraceptive use dynamics and EC use over 1 year of observation among university students in São Paulo, Brazil. The study was designed as a 12-month retrospective cohort study, conducted among a probability sample of undergraduate women from University of São Paulo, Brazil (the largest institution of higher education and research in Brazil), between June and July, 2015. Women between the ages of 18 and 24 years old enrolled in regular undergraduate course in any campus at University of São Paulo, Brazil, were eligible to participate in the study. They were selected by simple random sampling without replacement, and they were chosen electronically based on a list of all the email addresses of undergraduate students enrolled in the Electronic Computer Center of the University of São Paulo, organized alphabetically. The research project was described in the email, and the student who had an interest in participating in the study could click on the study to complete a consent form (APPENDIX A). The study obtained Ethical approval from the Research Ethics Committee at University of São Paulo School of Nursing (CAAE Number: 39142514.4.0000.5392) (ANNEX I).

6.2.2 Sampling

A total of 3,078 women were randomly selected and invited to participate in the project. Students received reminder emails weekly to participate in the study (on average 10 times). A total of 2,182 undergraduate students (71%) answered the online questionnaire. However, the final study population comprised 1,679 undergraduate students between 18 and 24 years of age, since 50 women refused to

participate in the study, and 453 students were not eligible for the study (< 18 years old (n=2), had never had sexual intercourse (n=358), and had not used any contraceptive method in the last 12 months (n=93)). In the present study, we focus on the 916 women who ever used EC.

6.2.3 Data collection and study instruments

The data were collected online through *Google Form*, using a self-administered questionnaire that was previously pre-tested and took, on average, 5–7 minutes to complete (APPENDICES B AND C). Women who reported they had ever use EC completed an EC calendar describing contraceptive usage at the time they last used EC. Specifically, they indicated what contraception they used within 30 days prior to EC use and within 30 days after EC use. Women were questioned about any use of contraception before and after EC, and in case they responded positively, were asked about the method they used within 30-day time periods before and after EC use.

6.2.4 Measures

Our primary outcome measure (question 1) relates to any use of contraception within 30 days after EC use.

We consider use of contraception within 30 days before EC as a key predictor of post EC contraceptive use. We also used this information to build our secondary outcome measure (question 2), exploring change in contraceptive utilization between before and after EC use. Three category indicators of change in contraceptive behaviors post EC was considered:

- “No change in contraception” refers to women who had not changed their contraception between 30 days before and 30 days after using EC (this included women who did not use contraception before and after EC).
- “Switching to more effective method” refers to women who changed to more effective method after EC use. Women who started using a method after EC use were included in this category. The contraceptive effectiveness was based on a study by CDC (2013).

- “Switching to less effective method” refers to women who changed to less effective method after EC use. Women who stopped using a method after EC were included in this category.

Our third outcome variable (question 3) relates to gaps in contraceptive use within 30 days after EC use, which was defined based on contraceptive discontinuation or not within 30 days after EC use.

Women’s sociodemographic and reproductive health history measures

We consider women’s sociodemographic characteristics as predictors of contraceptive use after EC use, based on prior research findings. Specifically, we include age (18–19 or 20–24); race/ethnicity (white, black, other); religious affiliation (none, roman catholic, kardecista spiritualism doctrine, evangelical, other); socioeconomic status (high (A/B) – middle/low (C/D/E)) (ABEP, 2015).

Because this study focuses on university students, we will consider their educational characteristics. This information includes field of study (human sciences, health sciences, exact sciences); period of study (full-time or part-time); and type of campus (São Paulo, countryside).

Sexual and reproductive health indicators are shown to be strongly associated with contraceptive use after EC use. In our study, we consider the type of relationship (steady, casual relationship or living with a partner); time since first sexual intercourse (≤ 1 year, 2-3 years, ≥ 5 years); number of lifetime sexual partners (1 partner, 2-3 partners, ≥ 4 partners); pregnancy history (no/yes); use of contraception within 30 days prior to EC use (no/yes); and type of contraceptive method used within 30 days prior to EC use (oral pill, male condom, and other methods).

6.2.5 Statistical analysis

1) First, a bivariate analysis was conducted to investigate women’s sociodemographic and sexual reproductive health characteristics according to the use of contraception within 30 days after EC, using the EC calendar data. We considered use of contraception within 30 days before EC as a key predictor of

post EC contraceptive in this analysis. We then conducted multivariate logistic regression to identify the independent effects of these factors on contraceptive use post EC use. The criteria for including variables in the multivariate analysis were not only those that showed significance in the bivariate analysis, but also variables based on our conceptual framework, that is, factors identified by previous studies as related to contraceptive use after EC use (as age, race/ethnicity, religion, socioeconomic status). The multivariate analysis also included field of study, period of study, campus, time since first sexual intercourse, and previous pregnancy. Hosmer and Lemeshow's Test was used to assess how well the model fitted the data. Adjusted odds ratios and 95% confidence intervals were estimated.

2) Second to analyze how EC affects contraceptive trajectories, we assessed changes in contraceptive use before and after EC use, using the EC calendar. We distinguished women who had no change in contraception after EC use (reference), women who switched to more effective method after EC use, and women who switched to less effective method after EC use. After conducting bivariate analysis, multinomial regression models were performed (dependent variable defined as a 3-categorical variable). Adjusted odds ratios and 95% confidence intervals were estimated. The final model was chosen based on prior knowledge and based on Akaike Information Criterion (AIC).

3) Finally, we investigated gaps in contraceptive use within 30-days post EC use, among all women who used contraception after EC, using EC calendar.

We stress that in all analysis, we explore the circumstances of last EC use, which was the episode investigated in the EC calendar. Pearson's Chi-square tests were conducted to identify the determinants of contraceptive use after EC and changes in contraceptive use after EC. All analyses were conducted using Stata 14.2 (Stata Corporation, College Station, TX, USA).

6.3 CONCLUSION

First of all, we conclude that most women in our sample use contraceptive method within 30-days after EC use; also, the use of contraception after EC use

among women who were not using a method within 30 days prior to EC use was a relevant issue. Second, we conclude that EC is mainly used in situations of inconsistency use of contraception, since the majority of women in our study had no change in contraception after EC use. A positive finding was the switching to more effective methods among women who had changes in contraception after EC. Finally, we conclude that continuous use of contraceptives is not a rare event, since gaps in contraception following EC use was infrequent among our sample.

Our findings of EC – particularly related to EC as potentially “bridge” to regular contraception and the few gaps observed after EC use – may positively reflect in the issues of unintended pregnancy, and induced abortions. However, it is important that the medical community and the family planning services address and comprehend how youth use contraceptive method, in order to avoid inconsistency in its use, and consequently, an unnecessary use of EC.

Unintended pregnancies resulting from inconsistent use of contraception could be decreased by couples’ increased reliance on male condoms and by increased adherence to daily oral pill. Therefore, counseling to educate youth alike on the boundaries of EC and to stimulate the correct use of ongoing contraception, along with the availability of the method, should be strongly considered in primary health care in Brazil.

A recent pilot study in Scotland found that a simple intervention in which pharmacists provided women presenting for EC with a cycle of oral pills significantly increased the probability of the women using effective contraception following EC use (Michie et al., 2014). Also, a case control study conducted among high risk women (15-19 years old) in Sweden, indicate that women provided advanced supplies of EC were more likely than those in the control group (not given EC) to reported a shorter time interval from unprotected sexual intercourse to EC intake, without any evidence of decreased use of contraceptives (Ekstrand et al., 2013). Therefore, expanding EC access and implementing advanced supply of EC in the Unified Health System (SUS) could benefit several Brazilian women, specially those who are at risk to unintended pregnancy.

Furthermore, although our results show that the use and consistency of contraception after EC use among undergraduate students were consistent with other studies and was relevant, since there are no studies that approached this scenario in Brazil, the time analyzed (1-month before EC use and 1-month after EC

use) was not enough for an in-depth analysis. Therefore, we would recommend that future researches in Brazil address contraceptive use after accessing EC considering a longer length.

7.CHAPTER SEVEN

CHAPTER SEVEN

7.1 SUMMARY OF FINDINGS

This section summarizes the key findings of the current study. In addition, limitations of the study were discussed, as were implications of the findings and future research possibilities.

Trends over the last half century show worldwide changes in sexual and reproductive health behaviors with increases in modern contraceptive use in Brazil (Brasil, 2008a). However, the sexual and reproductive health of all women in Brazil continues to be a serious public health concern, since the unintended pregnancies are higher especially due to contraceptive discontinuation, and there are limitations in implementing sexual and reproductive rights in the country (Brasil, 2008a; Berquó, Lima, 2008).

During the course of this study, a number of research gaps became apparent. This is the first Brazil study to describe young women's patterns of contraceptive use and use of EC in context of contraceptive discontinuation, and these quantitative findings should be further explored with other groups of women throughout the Brazil. Our study was developing to expand other two prior researches (Pirota, Schor 2004; Borges et al., 2010) on contraceptive behavior undertaken with young people in Brazil.

Contraceptive calendar is applicable for this study because it contains contraceptive histories for a representative sample of women between the ages of 18 and 24 ($n= 1,679$). In addition, we created a "EC calendar" based on the contraceptive calendar, to comprehend the dynamics of contraceptive discontinuation before and after using EC.

In chapter 3, examine whether contraceptive discontinuation for any reason, for method-related reasons and abandonment or switching to a less effective method is a common event among undergraduate students, also we verify the frequency of discontinuation, and if contraceptive discontinuation differs by type of method contraceptive. We find that that few young women discontinued their method for any reason (35%), method-related reasons (28%); however, a relative proportion of women (18%) abandoned their method or switched to a less effective method.

Additionally, we conclude that the frequency of discontinuation within 12-months is low, most of women who discontinued their method for any reason or for method-related reasons, discontinued just once during the 12-months (24% and 19%, respectively). We find that contraceptive discontinuation does differ by type of contraceptive method. Oral pill was the most contraceptive method used 12-months prior to the survey, following by male condom. However, according to the analysis of monthly discontinuation, we find that young women who were using male condom (10%) were more likely to discontinued their method for method-related reasons than oral pill users (3%), and also to method abandonment (5% of condom users against 3% of pill users); though we also observed (in analysis of new episodes of contraception) that oral pill users (3%) were a bit more likely to abandonment or switching to a less effective method when compared with male condom users (2%).

The differences among the results can be explained by the exclusion of women who used the same contraception continuously, and the exclusion of the firsts discontinuation episode, when we analyzed the probability of discontinuation of new episodes of contraception. On one hand, the analysis of the new episodes of contraception presents an overview of the relative risks of discontinuation. Conversely, they possible overestimating the discontinuation rates due to these exclusions. Similarly, the analysis of monthly discontinuation has pros and cons. The analysis of monthly discontinuation does not present this problematic, but on the other hand this analysis was not left-censored, that is, women who had initiated their method before the calendar period started (considered in most cases consistent users) were maintained in the analysis. Contraceptive discontinuation between women who have used a contraceptive some years with new users is totally different, since the likelihood of new users discontinuing the method is much greater. The performance of both analysis was important to compare the findings, which further deepened the study.

In chapter 4, we focus on using EC following discontinuation for method-related reasons and method abandonment. Additionally, we investigated the use of EC following inconsistent use of contraception or EC use in the context of non-regular contraception. One notable aspect of the evaluation findings is that the majority of women used EC in context of continuous use of the method, in cases of errors in the method, or in the occurrence of possible contraceptive gaps when women switching for another method. It means that the use of EC in context of inconsistent use

(90.6%) was higher than in context of non-regular contraception (9.4%), which is a positive finding. However, when we focus only on women who had discontinued their method for method-related reasons and women who abandoned their method, we observed that a relevant proportion of women did not use EC in situations of pregnancy risk. Only 24% of women who are at risk to become pregnant used EC following discontinuation for method-related reasons, and 11% of women used EC following method abandonment. A relevant proportion of young women in our study probably do not recognize they be at risk of pregnancy, which lead to non-use of EC in situations where the method is required, i.e., following discontinuation for method-related reasons and method abandonment.

In chapter 5 investigates the use of contraception before and after EC use during 1-month, through “EC calendar”. We also examined if the contraceptive utilized was the same before and after EC use, that is, whether the woman really used EC in context of inconsistent use. In addition, we evaluated contraceptive gaps after EC use. Our findings show that the contraceptive discontinuation 1-month before and 1-month after using EC is not high (33% and 25%, respectively). We also revealed that EC is mainly used in situations of inconsistency use of contraception, since the majority of women in our study used the same method before and after EC used (72%), that is, few women use EC in context of unprotected sexual intercourse. Gaps in contraception following EC use was also low than predictable (7.5%).

The main point was that the type of relationship, socioeconomic status, sexual experience, sexual partners, and type of contraceptive were strong variables that were associate with contraceptive discontinuation and were common in most analyzes. As for type of relationship, although type of relationship relationship was strongly associate with contraceptive discontinuation, and with EC use in the context of discontinuation, this study cannot assess the mediating effects of relationship dynamics/quality on contraceptive discontinuation. Recent studies, especially those using young people samples, have observed the importance of these relationship dynamics as having both a direct and mediating effect on contraceptive use (Manning et al., 2009; Gibbs et al., 2014; Manlove et al., 2014). Perhaps diary data as collected by the Barber & colleagues (2011) would be best to capture quality of the relationship at the time of discontinuation.

This study address a variety types of contraceptive discontinuation (discontinuation for any reason, discontinuation for method-related reasons, and

method abandonment), that has been not recently investigated in Brazil, and provides an update to research by considering discontinuation by type of contraceptive method as well as EC in context of contraceptive discontinuation. Understanding contraceptive discontinuation is significant because it has negatively and positively reproductive consequences for women throughout their life course, especially among young women who the contraceptive trajectories are dynamic. In addition, from the life course perspective, comprehending contraceptive and reproductive behaviors among young people are important, because family (and union) formation has experienced a major transformation over the last decade with delays in marriage (Manning, Brown, 2014; IBGE, 2015). It means that a growing population of youth exposed to a potential risk of unintended pregnancy.

For the endpoint of this viability study, we are well aware of the limitations specially of using retrospective measures of contraceptive behaviors in this study, but we believe that our results can contribute to understand the contraceptive dynamic, and the use of EC among Brazilian young women in context of contraceptive discontinuation.

7.2 STRENGTHS AND LIMITATIONS

The most strengths of our study are the ability to comprehend the contraceptive discontinuation among undergraduate students, since the frequency and extent of contraceptive gaps among young women in Brazil nowadays is not well-known. In addition, this is the first study from Brazil to evaluate the EC following a contraceptive discontinuation, and is also original in evaluating contraceptive discontinuation before and after using EC, among young women who are at risk to become pregnant. There are few studies that assess discontinuation among Brazilian women, but none of them have already included EC. Therefore, our study can help to understand the scenario of contraceptive discontinuation among young women in Brazil, which it has significant implications for the prevention of unintended pregnancies and subsequences induced abortions.

We stress that this is the first study that addressed field of study in the context of contraceptive discontinuation. Most of the studies that contemplate university

students do not address the educational background, which has been shown to be associated with contraceptive discontinuation in our analyzes.

Our data provide a rich and detailed picture of contraceptive behavior among a cohort of female undergraduate students in Brazil. However, this study has some limitations which need to be considered while interpreting our results.

- The study was carried out only in one university, although contemplated all campuses of this university, it may not be representative of Brazil university female students, and therefore results may not be generalizable to other universities in Brazil.
- Because of the retrospective nature of the contraceptive method history calendar, may there were recall bias. However, starting with the National Survey of Family Growth (NSFG) and now in succeeding studies, the female interview uses a life history calendar as a remembrance aid for the pregnancy and contraceptive history (Bradley, Schwandt, Khan, 2009). Notwithstanding these methodological strategies, we recognize that the recollection of contraceptive methods may be problematic.
- Another limitation refers to the grouping of “switching to a less effective method” with “method abandonment”, a resource used due to the low number of women who abandoned their methods and therefore, which would make some analyzes impossible. Although women who abandon their method or switching to a less effective method represent women at high risk of unintended pregnancies, we understand that contraceptive behavior differs between the two categories.
- There is no corresponding contraceptive method history calendar for men. Therefore, equivalent analyses for men cannot be investigated. Future consideration to men is also necessary, since the involvement of the partner impact in the contraceptive behavior (Manning et al., 2009; Manlove et al., 2011; Wildsmith, Manlove, Steward-Streng, 2015).
- Another limitation of this study relates to chosen methodology in which left-censored episodes were not included in analysis of cumulative probabilities of new episodes of contraception. This causes the loss of a significant amount of data and potentially leads to over estimates of discontinuation rates because

long time users who use consistently throughout the entire observation period are excluded due to left-censoring.

- The questionnaire contained a predetermined list of contraceptive methods for undergraduate students to choose from. There are a variety of issues regarding to this that limit our understanding of their contraceptive use. Firstly, in the “Contraceptive Calendar” the main methods were listed, and young women could select just one answer, that is, they were asked about the main contraception that they used per month; so, it was possible that women used several methods in the same month, also it was possible that particular methods were under-reported or forgotten.
- The oral pill and male condoms are effective contraceptive methods when they are used properly. However, undergraduate students were not asked about this important aspect of contraceptive use. In addition, young women were not asked why they used specific contraceptive methods at each month.
- Some data collected refer to the time of research, which made difficult the time specific indicators of sociodemographic and socioeconomic. Our contraceptive calendar did not include a time-varying measure of sociodemographic and socioeconomic and so we were unable to capture changes in these segments which may be consequential for discontinuation. Religion is another variable used in the analyses. However, this variable is based on affiliation, this by no means indicates that women attended the religion or were religious during the time of observation. Consequently, results related to religious affiliation must be carefully interpreted.
- We did not evaluate contraceptive failures, since it was not our objective. However, we recognize that failure of contraceptive method is an important outcome to study because it can lead to immediate risk of unintended pregnancy.
- Only few reasons for discontinuation were collected. In actuality, there are numerous reasons for discontinuing a contraceptive method, so we recognize our analyses of reasons for discontinuation are probable overgeneralized.
- Our analysis is limited to two contraceptive methods (oral pill and male condom), since the other methods were poorly reported by our study

population. In Brazil, modern contraceptive methods are used on a limited scale, irrespective of women's age (Brasil, 2008a).

- Because of the environment of the study which included sensitivity matter and self-reporting, information bias could be introduced which may have affected reliability of the results. Response bias can exist whether university students who have grown exhausted of completing surveys, opt to satisfice rather than cognitively choose the most appropriate responses (Barge, Gehlbach, 2011). Satisficing can include young women choosing the intermediate response rather than making an authentic choice, or answering all sections with the same response, leading to false responses. Satisficing can lead to invalid inferences about the data. In addition, the probability that interviewed provided socially satisfactory responses is a limitation to the validity of the inferences made using the data collected. However, the fact that the study questionnaire was online, and consequently quick and easy to answer, may have minimized these aspects. Also, the use of self-reports by university students has been found to be a precise way to collect information (Pike, 2011).
- Only few reasons for contraceptive discontinuation were collected. In actuality, there are numerous reasons for discontinuing a contraceptive method, so we recognize our analyses of reasons for discontinuation are probable overgeneralized. Seventh, our analysis is limited to two contraceptive methods (oral pill and male condom), since the other methods were poorly reported by our study population. In Brazil, modern contraceptive methods are used on a limited scale (Brasil, 2008a).
- Our study did not collect data on reasons for use or non-use of EC in the "EC calendar".
- Although we questioned young women about the use of contraception 1-month before and after EC use, we did not know with accuracy if they had any unprotected sexual intercourse during this time, and therefore, we did not quantify the rate of unprotected sexual intercourse. Finally, the "EC calendar" was just limited to 1-month interval.

7.3 RECOMMENDATIONS FOR FUTURE RESEARCHES

The topics of contraceptive discontinuation and emergency contraception are extensive open for research. Because these topics have been and remain a controversial theme it might not be easy to conduct research, but it is important in helping to understand the contraceptive dynamics, especially among university students. The up-to-date analysis of women's contraceptive method behavior provides a solid platform for future research. Areas for possible research include, but are not limited to:

- A qualitative study addressing focus groups could be undertaken with women in reproductive ages or EC users to get the whole picture of their needs, opinions, and beliefs. This could guide future interventions to reduction unintended pregnancies.
- A longitudinal study of undergraduate students beginning as freshman through senior year to investigate sexual activity as well as EC use could be conducted.
- Studies could be conducted to investigate diverse education modes to determine how much education is required to provoke contraceptive behavior change. Are brochures and posters effective? Can the media be used to spread adequate information to women? How much acquaintance is needed to dismiss misconceptions about EC? Do on-line classes improve knowledge and behavior modification? Are presentations better than television programs? Are direct interventions better option than group interventions?
- It would be useful to examine the emerging questions around sexual pleasure, sources of contraceptive knowledge, and personality. This would provide a more complete understanding of the role of contraceptive discontinuation and method switching in women's lives in Brazil, which could help to develop suitable sexual and reproductive health interventions.
- Future studies should approach and discuss with men about the importance of consistent contraceptive use and the risks of unintended pregnancy. Prospective studies on contraceptive use patterns are also recommended.

- Targeting people who are important to the students: probably if parents, pharmacy workers and health professionals understand EC better, they may be more accepting its use. If they are more accepting of its use, then undergraduate students may be more likely to use it.
- Other possibility to consider are group discussions in student residence at university, or making sexual health classes obligatory and increasing the amount of time spent in those classes on teaching about contraception, and EC.

7.4 PUBLIC HEALTH IMPLICATIONS

The main public health implication of contraceptive discontinuation is the unintended pregnancy and the amount of health, economic and social consequences that it causes. We believe that the public health impact of contraception will not be concretized until all women who desire to prevent pregnancies are using their method of choice continuously and effectively.

The public health implications of this study comprise an important contribution to the dearth of literature about contraceptive discontinuation among young women in Brazil. At the public health programming level, outcomes from this study can lead to intervention efforts to improve unintended pregnancy prevention among young women.

Lastly, access to contraceptive methods needs to be improved in primary health care system in Brazil, and clinical counselling and public health education efforts could focus on informing young women about the importance of effective and consistent use of contraception. We believe that a larger comprehending of young's concerns and experience using contraception could help primary health care providers reshape counseling strategies to improve adequate contraceptive continuation. In addition, appropriate and effective sexual health measures to support undergraduate students are strongly recommended and could stimulate responsible sexual behavior as a Public Health priority.

REFERENCES

REFERENCES

Aiken AM, Gold MA, Parker AM. Changes in young women's awareness, attitudes, and perceived barriers to using emergency contraception. *J Pediatr Adolesc Gynecol*. 2005;18(1):25-32.

Alano GM, Costa LN, Mirand LR, Galato D. Conhecimento, consumo e acesso à contracepção de emergência entre mulheres universitárias no sul do Estado de Santa Catarina. *Cienc Saúde Coletiva*. 2012;17(9):2397-404.

Ali M, Cleland J. Contraceptive discontinuation in six developing countries: a cause-specific analysis. *Int Fam Plan Perspect*. 1995;64(3):92-97.

Ali M, Cleland J. Determinants of contraceptive discontinuation in six developing countries. *J Biosoc Sci*. 1999;31(3):343-60.

Ali MM, Cleland J, Shah IH. Causes and consequences of contraceptive discontinuation: evidence from 60 Demographic and Health Surveys. Geneva: World Health Organization; 2012. Available at: http://apps.who.int/iris/bitstream/10665/75429/1/9789241504058_eng.pdf.

American College Health Association. American college health association-national college health assessment II: reference group executive summary spring 2015. Hanover, MD: American College Health Association; 2015. Available at: [http://www.acha-ncha.org/docs/NCHA-II WEB SPRING 2015 REFERENCE GROUP EXECUTIVE SUMMARY.pdf](http://www.acha-ncha.org/docs/NCHA-II_WEB_SPRING_2015_REFERENCE_GROUP_EXECUTIVE_SUMMARY.pdf).

Anasel MG, Mlinga UJ. Determinants of contraceptive use among married women in Tanzania: Policy implication. *African Population Studies*. 2014;28(2):978-88.

Aquino EML, Heilborn ML, Knauth D, Bozon M, Almeida MC, Araújo J, et al. Adolescência e reprodução no Brasil: a heterogeneidade dos perfis sociais. *Cad Saúde Pública*. 2003;19 Suppl 2:S377-88.

Arowojolu AO, Okewole IA, Adekunle AO. Comparative evaluation of the effectiveness and safety of two regimens of levonorgestrel for emergency contraception in Nigerians. *Contraception*. 2002;66:269-73.

Associação Brasileira de Empresas de Pesquisa (ABEP). Critério de classificação econômica Brasil. Dados com base no levantamento socioeconômico. CCEB, 2015. Available at: <http://www.abep.org/criterio-brasil>.

Bajos N, Lamarche Vadel A, Gilbert F, Ferrand M, Moreau C. Cocon group. Contraception at the time of abortion: high-risk period or high-risk women? *Hum Reprod*. 2006;21:2862-7.

Bajos N, Le Guen M, Bohet A, Panjo H, Moreau C; FECOND group. Effectiveness of family planning policies: the abortion paradox. *PLoS One*. 2014;9(3):e91539. DOI: 10.1371/journal.pone.0091539. eCollection 2014.

Ballinger GA. Using Generalized Estimating Equations for longitudinal data analysis. *Org Research Methods*. 2004;7(2):127-50.

Barden-O'Fallon JL, Speizer IS, White JS. Association between contraceptive discontinuation and pregnancy intentions in Guatemala. *Rev Panam Salud Publica*. 2008;23(6):410-7.

Barden-O'Fallon J, Speizer IS, Cálix J, Rodriguez F. An analysis of contraceptive discontinuation among female, reversible method users in urban Honduras. *Stud Fam Plann*. 2011a;42(1):11-20.

Barden-O'Fallon J, Speizer I. What differentiates method stoppers from switchers? Contraceptive discontinuation and switching among Honduran women. *Int Perspect Sex Reprod Health*. 2011b;37(1):16-23. DOI: 10.1363/3701611.

Barge S, Gehlbach H. Using the theory of satisficing to evaluate the quality of survey data. *Research in Higher Education*. 2011;53(2):182-200. DOI:10.1007/s11162-011-9251-2.

Bastianelli C, Rosato E, Farris M, Benagiano G. Emergency contraception: a survey of 1773 women. *Eur J Contracept Reprod Health Care*. 2016;21(6):455-461.

Bastos MR, Borges ALV, Hoga LAK, Fernandes MP, Contin MV. Práticas contraceptivas entre jovens universitárias: o uso da anticoncepção de emergência. *Texto Contexto Enferm*. 2008;17(3):447-56.

Bataglião EML, Mamede FB. Conhecimento e utilização da contracepção de emergência por acadêmicos de enfermagem. *Esc. Anna Nery*. 2011;15(2):284-90.

Beadnell B, Morrison DM, Wilsdon A, Wells EA, Murowchick E, Hoppe M, Gillmore MR, Nahom D. Condom use, frequency of sex, and number of partners: multidimensional characterization of adolescent sexual risk-taking. *J Sex Res*. 2005; 42(3):192-202.

Beltzer N, Moreau C, Bajos N; groupe CSF. Prevention of unintended pregnancies: toward a dual protection at sexual debut? *Rev Epidemiol Sante Publique*. 2011; 59(1):15-21. DOI: 10.1016/j.respe.2010.10.006.

Belzer M, Yoshida E, Tejirian T, Tucker D, Chung K. Advanced supply of emergency contraception for adolescent mothers increased utilization without reducing condom or primary contraception use. *J Adolesc Health*. 2003;32(2):122-3.

Berquó E, Lima LP. Intenções reprodutivas e planejamento da fecundidade. In: Elza Berquó E, Garcia S, Lago T, organizadores. *Relatório Final da Pesquisa Nacional de Demografia e Saúde da Criança e da Mulher – PNDS 2006*. Brasília: Ministério da Saúde; 2008. p. 203-04.

Blanc AK. The effect of power in sexual relationships on sexual and reproductive health: an examination of the evidence. *Stud Fam Plann*. 2001;32(3):189-213.

Blanc AK, Curtis SL, Croft TN. Monitoring contraceptive continuation: links to fertility outcomes and quality of care. *Stud Fam Plann*. 2002;33(2):127-40.

Blanc AK, Tsui AO, Croft TN, Trevitt JL. Patterns and trends in adolescent's contraceptive use and discontinuation in developing countries and comparisons with adult women. *Int Perspect Sex Reprod Health*. 2009;35(2):63-71.

Bongaarts, J. A framework for analyzing the proximate determinants of fertility. *Popul Dev Rev*. 1978;4:105-132.

Borges ALV, Fujimori E, Hoga LAK, Contim M. Práticas contraceptivas entre jovens universitários: uso da anticoncepção de emergência. *Cad Saúde Pública*. 2010; 26(4):816-26. DOI:10.1590/S0102-311X2010000400023.

Brache V, Cochon L, Jesam C, Maldonado R, Salvatierra AM, Levy DP, Gainer E, Croxatto HB. Immediate preovulatory administration of 30 mg ulipristal acetate significantly delays follicular rupture. *Hum Reprod.* 2010;25:2256-63.

Bradley SEK, Schwandt HM, Khan S. Levels, Trends, and Reasons for Contraceptive Discontinuation. DHS: Demographic and Health Research Division: Report. United States; 2009. Available at: <http://www.measuredhs.com/pubs/AS20/AS20.pdf>.

Bradley SEK, Croft TN, Rutstein SO: DHS Analytical Studies No. 22. The Impact of Contraceptive Failure on Unintended Births and Induced Abortions: Estimates and Strategies for Reduction. Maryland: ICF Inc; 2011.

Brandão ER, Cabral CS, Silva MV, Bastos LL, Paiva SP, Oliveira NVBV, Szabo I. O olhar do farmacêutico sobre a contracepção de emergência. In: Figueiredo R, Borges ALV, Paula SHB, organizadores. 1 ed. São Paulo: Instituto de Saúde; 2016. p.186.

Brasil. Ministério da Saúde. Secretaria da Ciência, Tecnologia e Insumos Estratégicos. Departamento de Ciência e Tecnologia. Pesquisa Nacional de Demografia e Saúde 1996: relatório. Rio de Janeiro; 1997. Available at: <https://dhsprogram.com/pubs/pdf/FR77/FR77.pdf>.

Brasil. Ministério da Saúde. Secretaria da Ciência, Tecnologia e Insumos Estratégicos. Departamento de Ciência e Tecnologia. PNDS: 2006: Pesquisa Nacional de Demografia e Saúde da Criança e da Mulher: relatório. Brasília: Ministério da Saúde; 2008a. Available at: http://bvms.saude.gov.br/bvs/publicacoes/relatório_pnds_2006.pdf.

Brasil. Ministério da Saúde. Aborto e Saúde Pública: 20 anos de pesquisa no Brasil. Brasília: Ministério da Saúde; 2008b. Available at: http://bvms.saude.gov.br/bvs/publicacoes/livro_aborto.pdf.

Brasil. Ministério da Saúde. Saúde sexual e saúde reprodutiva. Brasília: Ministério da Saúde; 2010. Available at: http://189.28.128.100/dab/docs/publicacoes/cadernos_ab/abcd26.pdf.

Brasil. Ministério da Saúde. Secretaria de Vigilância em Saúde. Departamento de Análise de Situação em Saúde. Saúde Brasil 2013: Uma análise da situação de saúde e das doenças transmissíveis relacionadas à pobreza. Brasília: Ministério da Saúde; 2014. Available at: http://bvsmms.saude.gov.br/bvs/publicacoes/saude_brasil_2013_analise_situacao_saude.pdf.

Brunner Huber LR, Hogue CJ, Stein AD, et al. Contraceptive use and discontinuation: findings from the contraceptive history, initiation, and choice study. *Am J Obstet Gynecol*. 2006;194:1290-5.

Bryant, KD. Contraceptive use and attitudes among female college students. *ABNF J*. 2009;20(1):12-6.

Burdette AM, Haynes SH, Hill TD, Bartkowski JP. Religious variations in perceived infertility and inconsistent contraceptive use among unmarried young adults in the United States. *J Adolesc Health*. 2014;54(6):704-9. DOI: 10.1016/j.jadohealth.2013.11.002. Epub 2014 Jan 1.

Cameron ST, Glasier A, Johnstone A, Rae L. Ongoing contraception after use of emergency contraception from a specialist contraceptive service. *Contraception*. 2011;84(4):368-71. DOI: 10.1016/j.contraception.2011.02.012.

Cameron ST, Gordon R, Glasier A. The effect on use of making emergency contraception available free of charge. *Contraception*. 2012;86(4):366-9. DOI: 10.1016/j.contraception.2012.02.018.

Caruso S, Agnello C, Intelisano G, et al. Prospective study on sexual behavior of women using 30 microg ethinylestradiol and 3 mg drospirenone oral contraceptive. *Contraception*. 2005;72:19-23.

Casterline J, El-Zanaty F and El-Zeini L. Unmet need and unintended fertility: longitudinal evidence from Upper Egypt. *International Family Planning Perspectives*. 2003;29(4):158-66.

Castle S, Askew I. Contraceptive discontinuation: reasons, challenges, and solutions. Population Council, 2015. [report]. Available at: http://ec2-54-210-230-186.compute-1.amazonaws.com/wp-content/uploads/2015/12/FP2020_ContraceptiveDiscontinuati on_SinglePage_Final_12.08.15.pdf.

Cavazos-Rehg PA, Krauss MJ, Spitznagel EL, Schootman M, Peipert JF, Cottler LB, Bierut LJ. Type of contraception method used at last intercourse and associations with health risk behaviors among US adolescents. *Contraception*. 2010;82(6):549-55. DOI: 10.1016/j.contraception.2010.05.007.

Centers for Disease Control and Prevention (CDC). Unintended pregnancy prevention, 2011a. Available at: <http://www.cdc.gov/reproductivehealth/UnintendedPregnancy/index.htm>.

Centers for Disease Control and Prevention (CDC). Pregnancy risk assessment monitoring system (PRAMS): PRAMS and unintended pregnancy, 2011b. Available at: <http://www.cdc.gov/PRAMS/UP.htm>.

Centers for Disease Control and Prevention (CDC). U.S. selected practice recommendations for contraceptive use, 2013: adapted from the World Health Organization selected practice recommendations for contraceptive use, 2nd edition. *MMWR Recomm Rep*. 2013;62(RR-05):1-60. Available at: <http://www.cdc.gov/mmwr/pdf/rr/rr62e0614.pdf>.

Cheng D, Schwarz E, Douglas E, Horon I. Unintended pregnancy and associated maternal preconception, prenatal and postpartum behaviors. *Contraception* 2009; 79(3):194-98.

Chin-Quee DS, Wedderburn M, Otterness C, Janowitz B, Chen-Mok M. Bridging emergency contraceptive pill users to regular contraception: results from a randomized trial in Jamaica. *Contraception* 2010;81:133-9.

Chofakian CBN, Borges ALV, Fujimori E, Hoga LAK. Conhecimento sobre anticoncepção de emergência entre adolescentes do Ensino Médio de escolas públicas e privadas. *Cad. Saúde Pública*. 2014;30(7):1525-36.

Cleland K, Zhu H, Goldstuck N, Cheng L, Trussell J. The efficacy of intrauterine devices for emergency contraception: a systematic review of 35 years of experience. *Hum Reprod*. 2012;27(7):1994-2000. DOI: 10.1093/humrep/des140.

Coetzee MH, Ngunyulu RN. Assessing the use of contraceptives by female undergraduate students in a selected higher educational institution in Gauteng. *Curationis*. 2015;38(2):1535. DOI:10.4102/curationis.v38i2.1535.

Costa AM, Guilhem D, Silver LD. Planejamento familiar: autonomia das mulheres sob questão. *Rev Bras Saúde Mater Infant*. 2006;6(1):75-84. DOI: 10.1590/S1519-38292006000100009.

Costa NFP, Ferraz EA, Souza CT, Silva CFR, Almeida MG. Access to emergency contraception: old barriers and new questions. *Rev Bras Ginecol Obstet*. 2008;30(2):55-60.

Creanga AA, Acharya R, Ahmed S, Tsui AO. Contraceptive discontinuation and failure and subsequent abortion in Romania: 1994-99. *Stud Fam Plann*. 2007;38(1):23-34.

Creinin MD, Schlaff W, Archer DF, Wan L, Frezieres R, Thomas M, Rosenberg M, Higgins J. Progesterone receptor modulator for emergency contraception: a randomized controlled trial. *Obstet Gynecol*. 2006;108:1089-97.

Curtis SL, Blanc AK. Determinants of contraceptive failure, switching, and discontinuation: an analysis of DHS contraceptive histories. *DHS Analytical Reports*. Calverton, MD, USA: Macro International; 1997.

Curtis S, Evens E, Sambisa W. Contraceptive discontinuation and unintended pregnancy: an imperfect relationship. *IntPerspect Sex Reprod Health*. 2011;37(2):58-66.

Cwiak C, Howard B, Hsieh J, Ricciotti N, Sucato GS. Sexual and Contraceptive Behaviors among Adolescents Requesting Emergency Contraception. *J Pediatr Adolesc Gynecol*. 2016;29(6):643-647. DOI: 10.1016/j.jpag.2016.06.003.

D'Antona AO, Chelekis JA, D'Antona MF, Siqueira AD. Contraceptive discontinuation and non-use in Santarém, Brazilian Amazon. *Cad Saúde Pública*. 2009;25(9):2021-32.

Daniels K, Jones J, Abma JC. Use of emergency contraception among women aged 15-44: United States, 2006-2010. NCHS data brief no 112. Hyattsville, MD: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Health Statistics; 2013.

Daniels K, Daugherty J, Jones J, Mosher W. Current contraceptive use and variation by selected characteristics among women aged 15-44: United States, 2011-2013. *Natl Health Stat Report*. 2015; 86:1-15. Available at: <http://www.cdc.gov/nchs/data/nhsr/nhsr086.pdf>.

Davidoff-Gore A, Luke N, Wawire S. Dimensions of poverty and inconsistent condom use among youth in urban Kenya. *AIDS Care*. 2011;23(10):1282-90. DOI:10.1080/09540121.2011.555744.

Davies SL, DiClemente RJ, Wingood GM, Person SD, Dix ES, Harrington K, Crosby RA, Oh K. Predictors of inconsistent contraceptive use among adolescent girls: findings from a prospective study. *J Adolesc Health*. 2006;39(1):43-9.

Dessunti EM, Reis AOA. Vulnerabilidade às dst/aids entre estudantes da saúde: estudo comparativo entre primeira e última série. *Cienc Cuid Saude*. 2012;11:274-83. DOI: 10.4025/cienccuidsaude.v11i5.17738.

Diamond-Smith NG, Moreau C, Bishai DM. Reducing unintended pregnancies: a microsimulation of contraceptive switching, discontinuation, and failure patterns in France. *Stud Fam Plann*. 2014;45(4):429-41. DOI: 10.1111/j.1728-4465.2014.00006.x.

Diniz D, Medeiros, M. Aborto no Brasil: uma pesquisa domiciliar com técnica de urna. *Ciênc Saúde Coletiva*. 2010;15(1):959-66. DOI: 10.1590/S1413-81232010000700002.

Donadiki EM, Jimenez-Garcia R, Velonakis EG, Hernandez-Barrera V, Sourtzi P, Lopez de Andres A, Jimenez-Trujillo I, Pino CG, Carrasco-Garrido P. Factors related to contraceptive methods among female higher education students in Greece. *J Pediatr Adolesc Gynecol*. 2013; 26(6):334-9. DOI: 10.1016/j.jpag.2013.06.011.

Ekstrand M, Tydén T, Darj E, Larsson M. Twelve-month follow-up of advance provision of emergency contraception among teenage girls in Sweden-a randomized controlled trial. *Ups J Med Sci*. 2013;118(4):271-5.

Ersek JL, Brunner Huber LR, Thompson ME, Warren-Findlow J. Satisfaction and discontinuation of contraception by contraceptive method among university women. *Matern Child Health J*. 2011;15(4):497-506. DOI: 10.1007/s10995-010-0610-y.

Escobar B, Román C, Muños C, López F. Perfil biosocial de las usuarias que demandan la píldora de anticoncepción de emergencia en un centro de salud de Santiago, 2006-2007. *Rev Chil Obstet Ginecol.* 2008;73(4):223-7.

ESHRE CapriWorkshop Group. Emergency contraception. Widely available and effective but disappointing as a public health intervention: a review. *Hum Reprod.* 2015; 30(4):751-60. DOI: 10.1093/humrep/dev019.

Falk G, Falk L, Hanson U, Milsom I. Young women requesting emergency contraception are, despite contraceptive counseling, a high risk group for new unintended pregnancies. *Contraception.* 2001;64(1):23-7.

Faúndes A, Simoneti RM, Duarte GA, Andalaft-Neto J. Factors associated to knowledge and opinion of gynecologists and obstetricians about the Brazilian legislation on abortion. *Rev Bras Epidemiol.* 2007; 10(1): 6-18. DOI: 10.1590/S1415-790X2007000100002.

Ferreira JM, Nunes FR, Modesto W, Gonçalves MP, Bahamondes L. Reasons for Brazilian women to switch from different contraceptives to long-acting reversible contraceptives. *Contraception.* 2014;89(1):17-21. DOI: 10.1016/j.contraception.2013.09.012.

Figueiredo R, Puppo LR, Segri NJ. Comportamento sexual e preventivo de adolescentes de São Paulo: um estudo com estudantes do ensino médio. *BIS, Bol Inst Saúde.* 2008;(46):31-3.

Figueiredo R, Paula BHS. Distribuição da contracepção de emergência nos municípios do estado de São Paulo: avaliação de dispensa em serviços e fluxos estabelecidos. *Reprod Clim.* 2012; 27(2):64-72.

Finer LB, Henshaw SK. Disparities in rates of unintended pregnancy in the United States, 1994 and 2001. *Perspect Sex Reprod Health.* 2006; 38:90-6.

Finer LB, Zolna MR. Unintended pregnancy in the United States: incidence and disparities, 2006. *Contraception.* 2011;84(5):478-85. DOI: 10.1016/j.contraception.2011.07.013.

Finer LB, Zolna MR. Declines in unintended pregnancy in the United States, 2008-2011. *N Engl J Med.* 2016;374(9):843-52. DOI: 10.1056/NEJMsa1506575.

Free C., Lee RM, Ogden J. Young women's accounts of factors influencing their use and non-use of emergency contraception: in-depth interview study. *BMJ*. 2002;325:1393-6.

Free C, Ogden, J. Emergency contraception use and non-use in young women: the application of a contextual and dynamic model. *Br J Health Psychol*. 2005;10(2):237-53.

Freedman LR, Stulberg DB. The research consortium on religious healthcare institutions: studying the impact of religious restrictions on women's reproductive health. *Contraception*. 2016;94(1):6-10. DOI: 10.1016/j.contraception.2016.03.015.

Frost JJ, Singh S, Finer LB. Factors associated with contraceptive use and nonuse, United States, 2004. *Perspect Sex Reprod Health*. 2007a;39(2):90-9.

Frost JJ, Singh S, Finer LB. U.S. women's one-year contraceptive use patterns, 2004. *Perspect Sex Reprod Health*. 2007b;39(1):48-55.

Frost JJ, Darroch JE, Remez L. Improving contraceptive use in the United States. Brief. New York: Guttmacher Institute; 2008. N.1. Available at: <http://www.guttmacher.org/pubs/2008/05/09/ImprovingContraceptiveUse.pdf>.

Frost JJ, Darroch JE. Factors associated with contraceptive choice and inconsistent method use, United States, 2004. *Perspect Sex Repro Health*. 2008b; 40:94-104.

Gainer E, Prudhomme M, Perriot Y, Leroux MC, Bouyer J. A prospective study of the provision of emergency contraception in French family planning centers. *Gynecol Obstet Fertil*. 2005;33(6):403-8.

Gibbs L, Manning WD, Longmore M, Giordano PC. Qualities of romantic relationships and consistent condom use among dating young adults. In: Bourgois L, Cauchois S, editors. *Contraceptives: Predictors of use, role of cultural attitudes & practices and levels of effectiveness*. Hauppauge, New York: Nova Science Publishers, Inc.; 2014. p.157-182.

Gipson JD, Koenig MA, Hindin MJ. The effects of unintended pregnancy on infant, child and parental health. A review of the literature. *Studies Fam Plann* 2008; 39(1): 18-38.

Glasier AF, Cameron ST, Fine PM, et al. Ulipristal acetate versus levonorgestrel for emergency contraception: a randomised non-inferiority trial and meta-analysis. *Lancet*. 2010; 375:555-62.

Gold MA, Wolford JE, Smith KA, Parker AM. The effects of advance provision of emergency contraception on adolescent women's sexual and contraceptive behaviors. *J Pediatr Adolesc Gynecol*. 2004;17(2):87-96.

Goldstein RL, Upadhyay UD, Raine TR. With pills, patches, rings, and shots: who still uses condoms? A longitudinal cohort study. *J Adolesc Health*. 2013;52(1):77-82. DOI: 10.1016/j.jadohealth.2012.08.001.

González E, Molina T, Montero A, Martínez V, Molina R. Factors associated with maintenance of contraception among adolescents. *Rev Méd Chile*. 2009.137(9):1187-192. <http://dx.doi.org/10.4067/S0034-98872009000900007>.

Goulard H, Moreau C, Gilbert F, Job-Spira N, Bajos N; Cocon Group. Contraceptive failures and determinants of emergency contraception use. *Contraception*. 2006;74(3):208-13.

Grady WR, Billy JO, Klepinger DH. Contraceptive method switching in the United States. *Perspect Sex Reprod Health*. 2002;34(3):135-45.

Gray E, McDonald P. Using a reproductive life course approach to understand contraceptive method use in Australia. *J Biosoc Sci*. 2010;42(1):43-57. DOI: 10.1017/S0021932009990381.

Hall KS, Moreau C, Trussell J. Lower use of sexual and reproductive health services among women with frequent religious participation, regardless of sexual experience. *J Womens Health (Larchmt)*. 2012;21(7):739-47. DOI: 10.1089/jwh.2011.3356.

Hamoda H, Ashok PW, Stalder C, et al. A randomized trial of mifepristone (10 mg) and levonorgestrel for emergency contraception. *Obstet Gynecol*. 2004;104:1307-13.

Hardy EE, Duarte GA, Osis MJD, Arce XE, Possan M. Anticoncepção de emergência no Brasil: facilitadores e barreiras. *Cad Saúde Pública*. 2001;17(4):1031-5.

Harvey C, Seib C, Lucke J. Continuation rates and reasons for removal among Implanon® users accessing two family planning clinics in Queensland, Australia. *Contraception*. 2009;80:527-32.

Heilborn ML, Portella AP, Brandão ER, Cabral CS, Grupo CONPuSUS. Contraception and family planning services as viewed by users of three clinics in the Unified National Health System, Rio de Janeiro State, Brazil. *Cad Saúde Pública*. 2009;25(2):S269-78.

Hooper DJ. Attitudes, awareness, compliance and preferences among hormonal contraception users: a global, cross-sectional, self-administered, online survey. *Clin Drug Investig*. 2010;30(11):749-63. DOI: 10.2165/11538900-000000000-00000.

Hu X, Cheng L, Hua X, Glasier A. Advanced provision of emergency contraception to postnatal women in China makes no difference in abortion rates: a randomized controlled trial. *Contraception*. 2005;72:111–6.

Inoue K, Barratt A, Richters J. Does research into contraceptive method discontinuation address women's own reasons? A critical review. *J Fam Plann Reprod Health Care*. 2015;41(4):292-9. DOI: 10.1136/jfprhc-2014-100976.

Instituto Brasileiro de Geografia e Estatística (IBGE). Censo demográfico 2010. Características gerais da população: resultados gerais da amostra. Rio de Janeiro; 2010. Available at: http://biblioteca.ibge.gov.br/visualizacao/periodicos/94/cd_2010_religiao_deficiencia.pdf.

Instituto Brasileiro de Geografia e Estatística (IBGE). Pesquisa Nacional de Saúde. Ciclos de vida. Brasil e grandes regiões. Rio de Janeiro; 2015. Available at: http://biblioteca.ibge.gov.br/visualizacao/periodicos/135/rc_2013_v40.pdf.

Jain A. Should eliminating unmet need for contraception continue to be a program priority? *International Family Planning Perspectives*, 1999,25(Suppl.):S39-S43;S49.

Jones RK, Darroch JE, Henshaw SK. Contraceptive use among US women having abortions in 2000-2001. *Perspect Sex Reproduct Health*. 2002;34:294-303.

Jones J, Mosher W, Daniels K. Current contraceptive use in the United States, 2006-2010, and changes in patterns of use since 1995. *Natl Health Stat Report*. 2012;(60):1-25.

Kavanaugh ML, Jerman J, Finer LB. Changes in Use of Long-Acting Reversible Contraceptive Methods Among U.S. Women, 2009-2012. *Obstet Gynecol.* 2015; 126(5):917-27. DOI: 10.1097/AOG.0000000000001094.

Keletzian E, Koumousidis A, Dimopoulos S, et al: Contraceptive consciousness and sexual behavior in three different female age groups in Greece: a retrospective study of the evolution during the last three decades. *Clin Exp Obstet Gynecol.* 2012; 39:160.

Kost K, Singh S, Vaughan B, Trussell J, Bankole A. Estimates of contraceptive failure from the 2002 National Survey of Family Growth. *Contraception.* 2008;77(1):10-21.

Lader D. Contraception and sexual health, 2008/09 [Opinions survey report no. 41]. Newport: Office for National Statistics; 2009. Available at: <http://www.ons.gov.uk/ons/rel/lifestyles/contraception-and-sexual-health/2008-09/index.html>.

Lakha F, Glasier A. Unintended pregnancy and use of emergency contraception among a large cohort of women attending for antenatal care or abortion in Scotland. *Lancet.* 2006;368:1782-7.

Lally K, Nathan-V Y, Dunne S, McGrath D, Cullen W, Meagher D, Coffey JC, Dunne C. Awareness of sexually transmitted infection and protection methods among university students in Ireland. *Ir J Med Sci.* 2015;184(1):135-42.

Lammers C., Ireland M., Resnick, M., Blum R. Influences on adolescent's decision to postpone onset of sexual intercourse: A survival analysis of virginity among youths aged 13 to 18 years. *J Adolesc Health.* 2000;26(1):42-8.

Langer A. El embarazo no deseado: impacto sobre la salud y la sociedad en América Latina y el Caribe. *Pan Am J Public Health.* 2002;11(3):192-205. DOI: 10.1590/S1020-49892002000300013.

Leite IC, Gupta N. Assessing regional differences in contraceptive discontinuation, failure and switching in Brazil. *Reprod Health.* 2007; 4:6.

Liang KY, Zeger SL. Longitudinal data analysis using generalized linear models. *Biometrika.* 1986;73:13-22.

Lindh I, Hognert H, Milsom I. The changing pattern of contraceptive use and pregnancies in four generations of young women. *Acta Obstet Gynecol Scand*. 2016; 95(11):1264-72. DOI: 10.1111/aogs.13003.

Lo SST, Fan SY, Ho PC, Glasier AF. Effect of advanced provision of emergency contraception on women's contraceptive behaviour: a randomized controlled trial. *Hum Reprod*. 2004;19(10):2404-10.

Longo LAFB. Juventude e contracepção: um estudo dos fatores que influenciam o comportamento contraceptivo das jovens brasileiras de 15 a 24 anos. *Rev Bras Estud Popul*. 2002;19(2):229-47.

Lotke PS. Increasing Use of Long-Acting Reversible Contraception to Decrease Unplanned Pregnancy. *Obstet Gynecol Clin North Am*. 2015;42(4):557-67. DOI: 10.1016/j.ogc.2015.07.008.

MacLean R. Teenagers given advance emergency contraception still use pill and condoms. *Perspect Sex Reprod Health*. 2004;36(3):134-5. DOI: 10.1111/j.1931-2393.2004.tb00205.x.

Manlove J, Ryan S, Franzetta K. Patterns of contraceptive use within teenagers' first sexual relationships. *Perspect Sex Reprod Health*. 2003;35(6):246-55.

Manlove J, Ryan S, Franzetta K. Contraceptive use and consistency in U.S. Teenagers' most recent sexual relationships. *Perspect Sex Reprod Health*. 2004; 36(6): 265-75. DOI: 10.1363/psrh.36.265.04.

Manlove J, Ryan S, Franzetta K. Contraceptive use patterns across teens' sexual relationships: the role of relationships, partners, and sexual histories. *Demography*. 2007;44(3):603-21.

Manlove J, Welti K, Barry M, Peterson K, Schelar E, Wildsmith E. Relationship characteristics and contraceptive use among young adults. *Perspect Sex Reprod Health*. 2011; 43:119-28. DOI:10.1363/4311911.

Manlove J, Welti K, Wildsmith E, Barry M. Relationship types and contraceptive use within young adult dating relationships. *Perspect Sex Reprod Health*. 2014; 46:41-50. DOI:10.1363/46e0514

Manning WD, Flanigan CM, Giordano PC, Longmore MA. Relationship dynamics and consistency of condom use among adolescents. *Perspect Sex Reprod Health*. 2009; 41(3):181-90. DOI:10.1363/4118109.

Manning WD, Brown SL, Payne KK. Two decades of stability and change in age at first union formation. *J. Marriage Fam*. 2014; 76:247-60.

Mansour D, Korver T, Marintcheva-Petrova M, Fraser IS. The effects of Implanon on menstrual bleeding patterns. *Eur J Contracept Reprod Health Care*. 2008;13(Suppl 1):13-28. DOI:10.1080/13625180801959931.

Martin A. La Anticoncepción de Emergencia en América Latina y el Caribe. *Rev Panam Salud Publica*. 2004;16(6):424-31.

Martins-Melo FR, Lima M da S, Alencar CH, et al. Temporal trends and spatial distribution of unsafe abortion in Brazil, 1996-2012. *Rev Saude Publica*. 2014; 48(3):508-520. DOI:10.1590/S0034-8910.2014048004878.

Mayer JP. Unintended childbearing, maternal beliefs, and delay of prenatal care. *Birth*. 1997;24(4):247-52.

Mehra D, Agardh A, Petterson KO, Östergren PO. Non-use of contraception: determinants among Ugandan university students. *Glob Health Action*. 2012; 5:18599. DOI: 10.3402/gha.v5i0.18599.

Mehra D, Östergren PO, Ekman B, Agardh A. Inconsistent condom use among Ugandan university students from a gender perspective: a cross-sectional study. *Glob Health Action*. 2014;10;7:229-42. DOI:10.3402/gha.v7.22942.

Meier AM. Adolescents' transition to first intercourse, religiosity, and attitudes about sex. *Soc Forces*. 2003;81(3):1031-52. DOI 10.1353/sof.2003.0039.

Menezes GMS, Aquino EML. Mortalidade materna na Bahia, 1998: relatório de pesquisa. Salvador: Instituto de Saúde Coletiva/Secretaria de Saúde do Estado da Bahia; 2001.

Meng CX, Gemzell-Danielsson K, Stephansson O, Kang JZ, Chen QF, Cheng LN. Emergency contraceptive use among 5677 women seeking abortion in Shanghai, China. *Hum Reprod*. 2009;24(7):1612-8.

Michie L, Cameron ST, Glasier A, Larke N, Muir A, Lorimer A. Pharmacy-based interventions for initiating effective contraception following the use of emergency contraception: a pilot study. *Contraception*. 2014;90(4):447-53. DOI: 10.1016/j.contraception.2014.05.004.

Moreau C, Bouyer J, Goulard H, Bajos N. The remaining barriers to the use of emergency contraception: perception of pregnancy risk by women undergoing induced abortions. *Contraception*. 2005;71(3):202-7.

Moreau C, Trussell J, Bajos N. The determinants and circumstances of use of emergency contraceptive pills in France in the context of direct pharmacy access. *Contraception*. 2006;74(6):476-82. Epub 2006 Sep 7.

Moreau C, Cleland K, Trussell J. Contraceptive discontinuation attributed to method dissatisfaction in the United States. *Contraception*. 2007;76(4):267-72.

Moreau C, Bouyer J, Bajos N, Rodríguez G, Trussell J. Frequency of discontinuation of contraceptive use: results from a French population-based cohort. *Hum Reprod*. 2009a;24(6):1387-92. DOI: 10.1093/humrep/dep027.

Moreau C, Trussell J, Michelot F, Bajos N. The effect of access to emergency contraceptive pills on women's use of highly effective contraceptives: results from a French national cohort study. *Am J Public Health*. 2009b;99(3):441-2. DOI: 10.2105/AJPH.2007.118935.

Moreau C, Trussell J, Desfreres J, Bajos N. Patterns of contraceptive use before and after an abortion: results from a nationally representative survey of women undergoing an abortion in France. *Contraception*. 2010;82:337-44. DOI: 10.1016/j.contraception.2010.03.011.

Moreau C, Trussell J, Bajos N. Contraceptive paths of adolescent women undergoing an abortion in France. *J Adolesc Health*. 2012;50(4):389-94. DOI: 10.1016/j.jadohealth.2011.07.013.

Moreau C, Hall K, Trussell J, Barber J. Effect of prospectively measured pregnancy intentions on the consistency of contraceptive use among young women in Michigan. *Hum Reprod*. 2013a;28(3):642-50. DOI: 10.1093/humrep/des421.

Moreau C, Bohet A, Trussell J, Bajos N; FECOND group. Estimates of unintended pregnancy rates over the last decade in France as a function of contraceptive behaviors. *Contraception*. 2014;89(4):314-21. DOI: 10.1016/j.contraception.2013.11.004.

Moreau C, Bohet A. Frequency and correlates of unintended pregnancy risk perceptions. *Contraception*. 2016;94(2):152-9. DOI: 10.1016/j.contraception.2016.02.029. Epub 2016 Mar 18.

Moreira-Almeida A, Pinsky I, Zaleski M, Laranjeira R. Envolvimento religioso e fatores sociodemográficos: resultados de um levantamento nacional no Brasil. *Rev Psiquiatr Clín*. 2010;37(1):12-15. DOI: 10.1590/S0101-60832010000100003.

Mosher WD, Jones J. Use of Contraception in the United States: 1982-2008. National Center for Health Statistics. *Vital Health Stat*. 2010; 23(29). Available at: http://www.cdc.gov/NCHS/data/series/sr_23/sr23_029.pdf.

Mosher WD, Jones J, Abma JC. Intended and unintended births in the United States: 1982–2010. National health statistics reports; no. 55. Hyattsville, MD: National Center for Health Statistics. 2012. Available at: <https://www.cdc.gov/nchs/data/nhsr/nhsr055.pdf>.

Moura ERF, Silva RM, Galvão RT. Dinâmica do atendimento em planejamento familiar no Programa de Saúde da Família no Brasil. *Cad Saúde Pública*. 2007; 23(4):961-70.

Nappi RE, Lobo Abascal P, Mansour D, Rabe T, Shojai R; Emergency Contraception Study Group. Use of and attitudes towards emergency contraception: a survey of women in five European countries. *Eur J Contracept Reprod Health Care*. 2014; 19(2):93-101. doi: 10.3109/13625187.2013.865164.

Nascimento CB. Conhecimento e uso da anticoncepção de emergência entre adolescentes estudantes do ensino médio. [Dissertação de Mestrado]. São Paulo: Escola de Enfermagem da Universidade de São Paulo; 2012.

National Institute for Health and Care Excellence. CG30: Long-acting reversible contraception [webpage on the Internet]. London, UK: NICE; 2014. Available from: <http://publications.nice.org.uk/long-actingreversible-contraception-cg30>. Accessed May 16, 2016.

Nelson AL. Recent use of condoms and emergency contraception by women who selected condoms as their contraceptive method. *Am J Obstet Gynecol.* 2006; 194(6):1710-5.

NHS Contraceptive Services: England, 2011/12. Community Contraceptive Clinics Health and Social Care Information Centre (HSCIC), Lifestyles statistics. October 2012.

Nguyen BT, Zaller N. Male access to over-the-counter emergency contraception: a survey of acceptability and barriers in Providence, Rhode Island. *Womens Health Issues.* 2009; 19(6):365-72.

Nsubuga H, Sekandi JN, Sempeera H, Makumbi FE. Contraceptive use, knowledge, attitude, perceptions and sexual behavior among female University students in Uganda: a cross-sectional survey. *BMC Womens Health.* 2016 Jan 27;16:6. doi: 10.1186/s12905-016-0286-6.

Osei IF, Mayhew SH, Biekro L, Collumbien M; ECAF Team. Fertility decisions and contraceptive use at different stages of relationships: windows of risk among men and women in accra. *Int Perspect Sex Reprod Health.* 2014; 40(3):135-43. doi: 10.1363/4013514.

Paiva SP, Brandão ER. Contracepção de emergência no contexto das farmácias: revisão crítica de literatura. *Physis.* 2012; 22(1): 17-34. doi: 10.1590/S0103-73312012000100002.

Palermo T, Bleck J, Westley E. Knowledge and Use of Emergency Contraception: A Multicountry Analysis. *Int Perspect Sex Reprod Health.* 2014; 40(2):79-86.

Patton EW, Hall KS, Dalton VK. How does religious affiliation affect women's attitudes toward reproductive health policy? Implications for the affordable care act. *Contraception.* 2015; 91(6):513-9. doi: 10.1016/j.contraception.2015.02.012.

Pazol K, Whiteman MK, Folger SG, Kourtis AP, Marchbanks PA, Jamieson DJ. Sporadic contraceptive use and nonuse: age-specific prevalence and associated factors. *Am J Obstet Gynecol.* 2015; 212(3):324.e1-8. doi: 10.1016/j.ajog.2014.10.004.

Peterson LS, Oakley D, Potter LS, Darroch JE. Women's efforts to prevent pregnancy: consistency of oral contraceptive use. *Fam Plann Perspect*. 1998; 30(1): 19–23.

Pike, GR. Using college students' self-reported learning outcomes in scholarly research. *New Directions for Institutional Research*. 2011; (150): 41-58. doi:10.1002/ir.388.

Pirotta KCM, Schor N. Intenções reprodutivas e práticas de regulação da fecundidade entre universitários. *Rev Saúde Pública*. 2004; 38:495-502.

Polis CB, Schaffer K, Blanchard K, et al. Advance provision of emergency contraception for pregnancy prevention (full review) *Cochrane Db Syst Rev*. 2010; 3. Available at: http://www.clacaidigital.info:8080/xmlui/bitstream/handle/123456789/194/Advance_provision_of_EC_for_pregnancy_prevention.pdf?sequence=1&isAllowed=y

Prietsch SOM, González-Chica DA, Cesar JA., Mendoza-Sassi RA. Gravidez não planejada no extremo Sul do Brasil: prevalência e fatores associados. *Cad. Saúde Pública*. 2011; 27(10): 1906-1916. doi: 10.1590/S0102-311X2011001000004.

Pulerwitz J, Amaro H, De Jong W, Gortmaker SL, Rudd R. Relationship power, condom use and HIV risk among women in the USA. *AIDS Care*. 2002; 14:789–800.

Raine T, Harper C, Leon K, Darney P. Emergency contraception: advance provision in a young, high-risk clinic population. *Obstet Gynecol*. 2000; 96(1):1-7.

Raine TR, Harper CC, Rocca CH, Fischer R, Padian N, Klausner JD, Darney PD. Direct access to emergency contraception through pharmacies and effect on unintended pregnancy and STIs: a randomized controlled trial. *JAMA*. 2005; 293(1):54-62.

Raine TR, Gard JC, Boyer CB, Haider S, Brown BA, Ramirez Hernandez FA, Harper CC. Contraceptive decision-making in sexual relationships: young men's experiences, attitudes and values. *Cult Health Sex*. 2010; 12(4):373-86. doi: 10.1080/13691050903524769.

Ranjit N, Bankole A, Darroch J, Singh S. Contraceptive failure in the first two years of use: differences across socioeconomic subgroups. *Fam Plann Perspect*. 2001; 33(1): 19–27.

Rash V, Knudsen LB, Gammeltoft T, et al. Contraceptive attitudes and contraceptive failure among women requesting induced abortion in Denmark. *Hum Reprod*. 2007; 22:1320–6.

Reed J, England P, Littlejohn K, Bass BC, Caudillo ML. Consistent and inconsistent contraception among young women: insights from qualitative interviews. *Family Relations*. 2014; 63(2): 244–58. doi:10.1111/fare.12058.

Richters J, Fitzadam S, Yeung A, Caruana T, Rissel C, Simpson JM, de Visser RO. Contraceptive practices among women: the second Australian study of health and relationships. *Contraception*. 2016; 94(5):548-555. DOI: 10.1016/j.contraception.2016.06.016.

Ricketts S, Klingler G, Schwalberg R. Game change in Colorado: widespread use of long- acting reversible contraceptives and rapid decline in births among young, low-income women. *Perspect Sex Reprod Health*. 2014;46:125-32.

Rocca CH, Schwarz EB, Stewart FH, Darney PD, Raine TR, Harper CC. Beyond access: acceptability, use and nonuse of emergency contraception among young women. *Am J Obstet Gynecol*. 2007;196(29):1-6.

Rocca CH, Harper CC, Raine-Bennett TR. Young women's perceptions of the benefits of childbearing: associations with contraceptive use and pregnancy. *Perspect Sex Reprod Health*. 2013;45(1):23-32. DOI:10.1363/4502313.

Ross J, Hardee K. Use of male methods of contraception worldwide. *J Biosoc Sci*. 2016;28:1-16.

Sant'Anna MJC, Carvalho KAM, Passarelli MLB, Coates V. Comportamento sexual entre jovens universitários. *Adolesc Saude*. 2008;5(2):52-6.

Santelli JS, Warren CW, Lowry R, Sogolow E, Collins J, Kann L, Kaufmann RB, Celentano DD. The use of condoms with other contraceptive methods among young men and women. *Fam Plann Perspect*. 1997;29(6):261-7. DOI: 10.1363/2926197.

Santelli J, Rochat R, Hatfield-Timajchy K, Gilbert BC, Curtis K, Cabral R, Hirsch JS, Schieve L; Unintended Pregnancy Working Group. The measurement and meaning of unintended pregnancy. *Perspect Sex Reprod Health*. 2003;35(2):94-101.

Santos OA, Borges, Borges ALV, Chofakian CBN, Pirotta KCM. Determinants of emergency contraception non-use among women in unplanned or ambivalent pregnancies. *Rev. esc. enferm. USP* [online]. 2014;48(spe):16-22. DOI: 10.1590/S0080-623420140000600003.

Sassler S, Miller A, Favinger SM. Planned parenthood: Fertility intentions and experiences among cohabiting couples. *J Fam Issues*. 2009;30:206-232.

Schünmann C, Glasier A. Measuring pregnancy intention and its relationship with contraceptive use among women undergoing therapeutic abortion. *Contraception*. 2006;73(5):520-4.

Secura GM, Madden T, McNicholas C, Mullersman J, Buckel CM, Zhao Q, Peipert JF. Provision of no-cost, long-acting contraception and teenage pregnancy. *N Engl J Med*. 2014;371(14):1316-23.

Sedgh G, Bankole A, Singh S, Eilers M. Legal abortion levels and trends by woman's age at termination. *Int Perspect Sex Reprod Health*. 2012;38(3):143-53.

Sereshti M, Delaram M, Rafieian M. Prevalence and causes of unwanted pregnancy from the perspective of pregnant women (Persian). *JRHS*. 2005;13(24):8-15.

Shih SL, Kebodeaux CA, Secura GM, Allsworth JE, Madden T, Peipert JF. Baseline correlates of inconsistent and incorrect condom use among sexually active women in the contraceptive CHOICE Project. *Sex Transm Dis*. 2011;38(11):1012-9. DOI: 10.1097/OLQ.0b013e318225f8c3.

Sidebottom A, Harrison PA, Donna A, Finnegan K. The varied circumstances prompting requests for emergency contraception at school-based clinics. *J Sch Health*. 2008;78(5):258-63.

Silva FC, Vitale MSS, Maranhão HS, Canuto MHA, Pires MS, Fisberg M. Diferenças regionais de conhecimento, opinião e uso de contraceptivo de emergência entre universitários brasileiros de cursos da área de saúde. *Cad Saúde Pública*. 2010;26(9):1821-31.

Silva LP, Camargo FC, Iwamoto HH. Comportamento sexual dos acadêmicos ingressantes em cursos da área da saúde de uma universidade pública. REAS. 2014;3(1):39-52.

Somba MJ, Mbonile M, Obure J, Mahande MJ. Sexual behaviour, contraceptive knowledge and use among female undergraduates' students of Muhimbili and Dar es Salaam Universities, Tanzania: a cross-sectional study. BMC Womens Health. 2014; 14:94. DOI: 10.1186/1472-6874-14-94.

Sorhaindo A, Beckera D, Fletcher H, Garcia SG. Emergency contraception among university students in Kingston, Jamaica: a survey of knowledge, attitudes, and practices. Contraception. 2002;66:261-8.

Sørensen MB, Pedersen BL, Nyrnberg LE. Differences between users and non-users of emergency contraception after a recognized unprotected intercourse. Contraception. 2000;62(1):1-3.

Souza RA, Brandão RE. Marcos normativos da anticoncepção de emergência e as dificuldades de sua institucionalização nos serviços de saúde. Physis. 2009; 19(4):1067-1086. DOI: 10.1590/S0103-73312009000400009.

Steele F, Curtis S. Appropriate methods for analyzing the effect of method choice on contraceptive discontinuation. Demography. 2003;40(1):1-22.

Tafari S, Martinelli D, Germinario C, Prato R. A survey of opinions on emergency contraception in young women in Southern Italy. Eur J Contracept Reprod Health Care 2012;17:164-9.

Tavares MP, Foster AM. Emergency contraception in a public health emergency: exploring pharmacy availability in Brazil. Contraception. 2016; 94(2):109-14. DOI: 10.1016/j.contraception.2016.04.006.

Theme-Filha MM, Baldisserotto ML, Fraga AC, Ayers S, da Gama SG, Leal MD. Factors associated with unintended pregnancy in Brazil: cross-sectional results from the Birth in Brazil National Survey, 2011/2012. Reprod Health. 2016;13(Suppl 3):118.

Trieu SL, Shenoy DP, Bratton S, Marshak HH. Provision of emergency contraception at student health centers in California community colleges. Womens Health Issues. 2011;21(6):431-7. DOI: 10.1016/j.whi.2011.04.011.

Trussell J, Rodriguez G, Ellertson C Updated estimates of the effectiveness of the Yuzpe regimen of emergency contraception. *Contraception*. 1999;59(3):147-51.

Trussell J. Contraceptive failure in the United States. *Contraception*. 2004;70(2):89-96.

Trussell J. Contraceptive efficacy. In: Hatcher RA, Trussell J, Nelson AL, Cates W, Stewart FH, Kowal D, editors. *Contraceptive Technology: Nineteenth revised edition*. Ardent Media; New York NY: 2007.

Trussell J. Contraceptive failure in the United States. *Contraception*. 2011;83(5):397-404. doi: 10.1016/j.contraception.2011.01.021.

Trussell J, Raymond E, Cleland K. Emergency contraception: A Last Chance to Prevent Unintended Pregnancy. *Contemporary Readings in Law and Social Justice*. 2014;6(2):7-38.

Vaughan B, Trussell J, Kost K, Singh S, Jones R. Discontinuation and resumption of contraceptive use: results from the 2002 National Survey of Family Growth. *Contraception*. 2008;78(4):271-83. DOI: 10.1016/j.contraception.2008.05.007.

Veloso, DLC, Peres VC, Lopes JSOC, Salge AKM, Guimarães JV. Anticoncepção de emergência: conhecimento e atitude de acadêmicos de enfermagem. *Rev Gaucha Enferm*. 2014;35(2):33-9.

Vieira EM, Souza L. A satisfação com o serviço de esterilização cirúrgica entre os usuários do Sistema Único de Saúde em um município paulista. *Rev Bras Epidemiol*. 2011;14(4):556-64.

Zembe YZ, Townsend L, Thorson A, Ekström AM. Predictors of inconsistent condom use among a hard to reach population of young women with multiple sexual partners in peri-urban South Africa. *PLoS One*. 2012;7(12):e51998. DOI: 10.1371/journal.pone.0051998.

Walsh TL, Frezieres RG, Peacock K, Nelson AL, Clark VA, Bernstein L. Evaluation of the efficacy of a nonlatex condom: results from a randomized, controlled clinical trial. *Perspect Sex Reprod Health*. 2003;35(2):79-86.

Wang H, Long L, Cai H, Wu Y, Xu J, Shu C, Wang P, Li B, Wei Q, Shang X, Wang X, Zhang M, Xiong C, Yin P. Contraception and Unintended Pregnancy among Unmarried Female University Students: A Cross-sectional Study from China. *PLoS One*. 2015;10(6):e0130212. DOI: 10.1371/journal.pone.0130212.

Wellings K, Brima N, Sadler K, Copas AJ, McDaid L, Mercer CH, McManus S, Stephenson J, Glasier A. Stopping and switching contraceptive methods: findings from Contessa, a prospective longitudinal study of women of reproductive age in England. *Contraception*. 2015;91(1):57-66. DOI: 10.1016/j.contraception.2014.09.008.

Westley E, Schwarz EB. Emergency contraception: global challenges, new opportunities. *Contraception*. 2012;85(5):429-31. DOI: 10.1016/j.contraception.2012.01.012. Epub 2012 Mar 2.

Westley E, Kapp N, Palermo T, Bleck J. A review of global access to emergency contraception. *Int J Gynaecol Obstet*. 2013;123(1):4-6. DOI: 10.1016/j.ijgo.2013.04.019. Epub 2013 Jul 13.

Westley E, Bass J, Puig C. A global assessment of emergency ontraception acessibility. In: Figueiredo R, Borges ALV, Paula SHB, organizadores. 1 ed. São Paulo: Instituto de Saúde; 2016. p.24-38.

Wildsmith E, Manlove J, Steward-Streng N. Relationship characteristics and contraceptive use among dating and cohabiting young adult couples. *Perspect Sex Reprod Health*. 2015;47(1):27-36. DOI: 10.1363/47e2515.

Williamson LM, Buston K, Sweeting H. Young women's perceptions of pregnancy risk and use of emergency contraception: findings from a qualitative study. *Contraception*. 2009;79(4):310-5. DOI: 10.1016/j.contraception.2008.10.014.

World Health Organization (WHO). Global Strategy for STI Prevention and Control Meeting. Geneva: WHO; 2004.

World Health Organization (WHO). Emergency contraception. Report. Geneva: WHO; 2012. Available at: <http://www.who.int/mediacentre/factsheets/fs244/en>.

APPENDICES

APPENDIX A – CONSENT FORM



UNIVERSITY OF SÃO PAULO
SCHOOL OF NURSING
DEPARTMENT OF PUBLIC HEALTH NURSING

(Obrigatório para Pesquisas Científicas em Seres Humanos- Resolução no 196 de 10/10/96- CNS)

Estamos realizando uma pesquisa com as graduandas da Universidade de São Paulo (USP) com o objetivo de avaliar as descontinuidades no uso de métodos contraceptivos, bem como a dinâmica contraceptiva antes e após o uso da anticoncepção de emergência. Na verdade, gostaríamos de conhecer sua história contraceptiva no último ano, ou seja, quais os métodos contraceptivos que você usou neste período e, caso tenha usado a anticoncepção de emergência, quais métodos foram utilizados antes e após o uso deste método. Mesmo que faça sexo apenas com mulheres e não tenha preocupação em evitar uma gravidez, é importante que responda todo o questionário. Queremos conhecer aspectos da saúde sexual e reprodutiva de todas as universitárias da USP. Esta pesquisa é coordenada pela Profa. Dra. Ana Luiza Vilela Borges da Escola de Enfermagem da Universidade de São Paulo.

O questionário é anônimo, ou seja, **o nome da entrevistada não aparece** em nenhum momento. Todas as informações são sigilosas e as pessoas não serão identificadas. O questionário deverá ser respondido *online* e, durará em média 30 minutos. Caso tenha qualquer dúvida sobre a pesquisa, esteja à vontade para entrar em contato com a Coordenação do Projeto. Você pode se recusar a participar da pesquisa inteira ou de qualquer parte dela e isto será bem compreendido pela entrevistadora. Este formulário *online* serve para comprovar que você está de acordo em participar desta pesquisa. Você deverá clicar em “aceite”, caso queira participar da pesquisa. Em seguida, será redirecionada para um *link* contendo o questionário *online*.

Em caso de dúvidas sobre a pesquisa, entre em contato com:

<p>Coordenadora do Projeto: Ana Luiza Vilela Borges Av. Dr. Enéas de Carvalho Aguiar, 419 CEP 05401-001 São Paulo/SP Tel: (11) 3061-7652 E-mail: alvilela@usp.br</p>	<p>Comitê de Ética em Pesquisa: Escola de Enfermagem da USP Av. Dr. Enéas de Carvalho Aguiar, 419 CEP 05401-001 São Paulo/SP Tel: (11) 3061 748/8858 E-mail: edipesq@usp.br</p>
--	--

Caso, **ACEITE** participar da pesquisa clique no botão ao lado

Caso, **NÃO ACEITE** participar da pesquisa clique no botão ao lado

APPENDIX B – QUESTIONNAIRE (ENGLISH VERSION)

SCHOOL DATA
What is the area of knowledge of your undergraduate degree? 1 () Health Sciences 2 () Exact Sciences 3 () Human Sciences
Your undergraduate degree belongs to which campus? 1 () São Paulo 2 () Countryside
What is the study period of your undergraduate degree? 1 () Full-time 2 () Part-time (morning) 3 () Part-time (afternoon) 4 () Part-time (evening)
Do you have a scholarship? 1 () No 2 () Yes
SOCIODEMOGRAPHIC INFORMATION
How old are you? _____ years
How do you rate your race? 1 () White (caucasian) 2 () Black (african american) 3 () Yellow (asian) 4 () Indian
What's your religion? 1 () No religion 2 () Roman Catholic 3 () Spiritism 4 () Evangelical 5 () Other
Do you work (wage labor)? 0 () No 1 () Yes 
Do you have a formal contract? 0 () No 1 () Yes

Which of the items below do you have in your home, and how many do you own?

Items	Quantity				
	0	1	2	3	4 ou +
Television	<input type="radio"/>				
Bathroom	<input type="radio"/>				
Automobile	<input type="radio"/>				
Housekeeper	<input type="radio"/>				
Washing machine	<input type="radio"/>				
DVD	<input type="radio"/>				
Refrigerator	<input type="radio"/>				
Freezer	<input type="radio"/>				

Who is the householder in your home?

- 1 () Father
 2 () Mother
 3 () Uncle/aunt
 4 () Grandfather/grandmother
 5 () Brother/sister
 6 () You
 7 () Outher: _____

Which school degree did the head of the family study or complete with approval?

Does your house have piped water?

- 0 () No
 1 () Yes

Does your house have piped water?

- 0 () No
 1 () Yes

SEXUAL BEHAVIOR

What's your marital status?

- 1 () Married or living with a partner
 2 () Single
 3 () Divorced
 5 () Widow
- 

What type of relationship do you have?

- 0 () None
 1 () Steady, as with boyfriend
 2 () Casual relationship (have a fling)

<p>Have you ever had sexual intercourse?</p> <p>0 () No (Questionnaire ends here)</p> <p>1 () Yes </p>
<p>How old were you when you first had sexual intercourse? _____ years</p>
<p>How many sexual partners have you had in your life? _____ sexual partners</p>
<p>Have you ever become pregnant?</p> <p>0 () No</p> <p>1 () Yes </p>
<p>How old were you at your first pregnancy? _____ years</p>
<p>How many times have you become pregnant? _____ times</p>
<p>How many children do you have? _____ children</p>
<p>Have you ever had an abortion?</p> <p>0 () No</p> <p>1 () Yes </p>
<p>How many abortions have you had? _____ abortions</p>
<p>Did you use a contraceptive method in the last year?</p> <p>0 () No </p> <p>1 () Yes</p>

Which was the reason for not using contraception in the past year?

- 0 () I didn't have sexual intercourse
 1 () Infrequent sexual intercourse
 2 () I became pregnant
 3 () I'd like to become pregnant
 4 () I did not want to use any contraceptive method
 5 () Partner opposition
 6 () Religious reasons
 7 () I did not know anything about contraception
 8 () I did not know where to get contraception
 9 () Health problems
 10 () Fear of side effects
 11 () Access difficulties
 12 () Cost of contraception
 13 () Inconvenient to use
 14 () Interfere with normal body functions
 15 () Husband / partner infertile
 16 () You or your partner has any condition that prevents pregnancy (hysterectomy, azoospermia)
 17 () I have sexual intercourse only with women
 18 () Other: _____

(The questionnaire ends for women who answered this question)

Currently, do you use some contraceptive method?

- 1 () No
 2 () Yes

Which contraceptive method are you using?

- 1 () Oral pill
 2 () Male condom
 3 () Injectable
 4 () Withdrawal/Fertility-awareness
 5 () IUD
 6 () Other

Did you use a contraceptive method in your last sexual intercourse?

- 1 () No
 2 () Yes

Which contraceptive method did you use in your last sexual intercourse?

- 1 () Oral pill
 2 () Male condom
 3 () Injectable
 4 () Withdrawal/Fertility-awareness
 5 () IUD
 6 () Other

Did you use emergency contraception in the last 12-months?

1 () No

2 () Yes

Which month did you use emergency contraception?

12

11 months ago

10 months ago

9 months ago

8 months ago

7 months ago

6 months ago

5 months ago

4 months ago

3 months ago

2 months ago

1 month ago

Emergency contraception

Did you become pregnant in the last 12 months?

1 () No

2 () Yes

Which month did you become pregnant? _____

What type of relationship (more frequently) did you have in each month, in the last 12-months?	Steady, as with boyfriend	Casual relationship	Living with a partner	None
1 month ago	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
2 months ago	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
3 months ago	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
4 months ago	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
5 months ago	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
6 months ago	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
7 months ago	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
8 months ago	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
9 months ago	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
10 months ago	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
11 months ago	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
12 months ago	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

EMERGENCY CONTRACEPTION
<p>Do you know anyone who has used the emergency contraception?</p> <p>0 () No 1 () Yes</p>
<p>Have you ever used emergency contraception?</p> <p>0 () No (The questionnaire ends here) 1 () Yes</p>
<p>Have you used the emergency contraception in the last 12 months?</p> <p>0 () No 1 () Yes</p>
<p>IF YES – What was the reason that led you to use the emergency contraception in the last time?</p> <p>1 () The contraceptive method used failed 2 () I forgot to use a contraceptive method 3 () I did not want to use any contraceptive method 4 () I was a victim of sexual violence 5 () By sheer insecurity 6 () Other</p>
<p>IF YES – In last time that you used the emergency contraception, the relationship was:</p> <p>1 () Steady, as with boyfriend 2 () Casual relationship (have fling)</p>
<p>IF YES – Who took the initiative in the emergency contraception use?</p> <p>1 () You 2 () Partner 3 () Both</p>
<p>IF YES – The emergency contraception was acquired with doctor's prescription?</p> <p>0 () No 1 () Yes</p>
<p>IF YES – Where did you get the emergency contraception?</p> <p>1 () Drugstore 2 () Health Center 3 () In particular doctor or medical insurance 4 () With friends or acquaintances 5 () Boyfriend who brought the contraception of emergency 6 () Relatives 7 () Other</p>
<p>How long after an unprotected sexual intercourse did you take the emergency contraception (in the last time)?</p> <p>1 () Least one day</p>

<p>2 () One day</p> <p>3 () Two days</p> <p>4 () Three days</p> <p>5 () Four days</p> <p>6 () Five day</p> <p>7 () Six days or more days</p> <p>8 () I do not know, I do not remember</p>
<p>How many times have you used the emergency contraception in your lifetime?</p> <p>1 () Once</p> <p>2 () Twice</p> <p>3 () 3 times</p> <p>4 () 4 times</p> <p>5 () 5 times or more times</p>
<p>Have you ever had unprotected sexual intercourse and did not use emergency contraception?</p> <p>1 () No</p> <p>2 () Yes</p>
<p>When was the last time you used the emergency contraception? ___/___/_____</p>

<p>Were you using some other contraceptive method when you used the emergency contraception in the last time?</p> <p>0 () No</p> <p>1 () Yes</p>
<p>Which contraceptive method were you using?</p> <p>1 () Oral pill</p> <p>2 () Male condom</p> <p>3 () Injectable</p> <p>4 () Withdrawal/Fertility-awareness</p> <p>5 () IUD</p> <p>6 () Other</p>
<p>→ 30 DAYS BEFORE THE USE OF EMERGENCY CONTRACEPTION</p>
<p>Did you use some contraceptive method 30 days BEFORE emergency contraception use?</p> <p>0 () No</p> <p>1 () Yes</p>
<p>Which contraceptive method did you use on most days?</p> <p>1 () Oral pill</p> <p>2 () Male condom</p> <p>3 () Male condom and oral pill</p>

4 () Injectable
 5 () Withdrawal/Fertility-awareness
 6 () IUD
 7 () Other

In these 30 days BEFORE emergency contraception use, did you use this method continuously (every day without exception)?
 0 () No 1 () Yes

In these 30 days BEFORE emergency contraception use, which days you used a contraceptive method?

30 DAYS BEFORE	Day	I did not use contraception	I used contraception
	30		
	29		
	28		
	27		
	26		
	25		
	24		
	23		
	22		
	21		
	20		
	19		
	18		
	17		
	16		
	15		
	14		
	13		
	12		
	11		
	10		
	9		
	8		
	7		
	6		
	5		
	4		
	3		
	2		
	1		
EMERGENCY CONTRACEPTIVE USE			

→ 30 DAYS AFTER THE USE OF EMERGENCY CONTRACEPTION

Did you use some contraceptive method during 30 days AFTER emergency contraception use?

0 () No

1 () Yes

Which contraceptive method did you use on most days?

1 () Oral pill

2 () Male condom

3 () Male condom and oral pill

4 () Injectable

5 () Withdrawal/Fertility-awareness

6 () IUD

7 () Other

Did you adopt a contraceptive method immediately after the use of emergency contraception?

0 () No

1 () Yes

How many days after using emergency contraception did you start contraceptive methods use?

1 () < 10 days

2 () 10 – 15 days

3 () ≥ 15 days

How long did you use this method continuously?

1 () < 1 month

2 () 1 month

3 () 1 -| 6 months

4 () 6 -| 12 months

5 () 1 -| 2 years

6 () ≥ 2 years

APPENDIX C – QUESTIONÁRIO (VERSÃO EM PORTUGUÊS)

DADOS ESCOLARES
<p>Qual é a área de conhecimento do seu curso de graduação?</p> <p>1 () Biológicas/Saúde</p> <p>2 () Exatas</p> <p>3 () Humanas</p>
<p>Seu curso de graduação pertence a qual <i>campus</i>?</p> <p>1 () São Paulo</p> <p>2 () São Carlos</p> <p>3 () Ribeirão Preto</p> <p>4 () Pirassununga</p> <p>5 () Bauru</p> <p>6 () Piracicaba</p> <p>7 () Santos</p> <p>8 () Lorena</p>
<p>Qual é o período de estudo de seu curso de graduação?</p> <p>1 () Integral</p> <p>2 () Matutino</p> <p>3 () Vespertino</p> <p>4 () Noturno</p>
INFORMAÇÕES SOCIODEMOGRÁFICAS
<p>Quantos anos você tem? _____ anos</p>
<p>Como você classifica sua cor ou raça?</p> <p>1 () Branca</p> <p>2 () Preta</p> <p>3 () Parda</p> <p>4 () Amarela (de origem asiática)</p> <p>5 () Indígena</p>
<p>Qual é a sua religião?</p> <p>1 () Católica</p> <p>2 () Evangélica</p> <p>3 () Espírita</p> <p>4 () Pentecostal</p> <p>5 () Judaica</p>

6 () Umbanda, candomblé, batuque
 7 () Budista
 8 () Nenhuma
 8 () Outra: _____

Você trabalha (trabalho assalariado)?
 0 () Não
 1 () Sim 

Tem carteira assinada?
 0 () Não
 1 () Sim

Dos itens abaixo, quais você possui em sua casa e quantos você possui de cada?

Variáveis	Quantidade				
	0	1	2	3	4 ou +
Televisão em cores	<input type="radio"/>				
Banheiro	<input type="radio"/>				
Automóvel	<input type="radio"/>				
Empregada mensalista	<input type="radio"/>				
Máquina de lavar	<input type="radio"/>				
Videocassete e/ou DVD	<input type="radio"/>				
Geladeira	<input type="radio"/>				
Freezer (aparelho independente ou parte da geladeira duplex)	<input type="radio"/>				

Quem é o chefe da sua família?
 1 () Pai
 2 () Mãe
 3 () Tio(a)
 4 () Avô(á)
 5 () Irmão(a)
 6 () Você
 7 () Outro: _____

Até que série na escola o chefe da família estudou ou completou com aprovação?
 1 () Analfabeto
 2 () Ensino fundamental 1 incompleto
 3 () Ensino fundamental 1 completo
 4 () Ensino fundamental 2 incompleto
 5 () Ensino fundamental 2 completo
 6 () Ensino médio incompleto

7 () Ensino médio completo 8 () Ensino superior incompleto 9 () Ensino superior completo 10 () Outro: _____
A sua casa tem água encanada? 0 () Não 1 () Sim
A rua onde você mora é pavimentada? 0 () Não 1 () Sim
AGORA, PERGUNTAREMOS SOBRE SUA VIDA AFETIVA E SEXUAL
No momento, você está: 1 () Casada/vivendo com o parceiro(a) (Pula a próxima pergunta) 2 () Solteira 3 () Separada/divorciada 5 () Viúva 6 () Outra: _____
Atualmente, qual o tipo de relacionamento que você tem? 0 () Nenhum 1 () fixo(a)/regular, como com namorado(a) 2 () eventual, como com ficante
Você já teve alguma relação sexual? 0 () Não (Se você respondeu não, a pesquisa se encerra. Agradecemos a sua participação) 1 () Sim
SE SIM: Com quantos anos teve a sua 1ª relação sexual? _____ anos
Quantos parceiros (as) sexuais já teve na vida? _____ parceiros
Você já engravidou anteriormente? 0 () Não (Pule as próximas perguntas sobre gravidez) 1 () Sim
Que idade tinha quando engravidou pela primeira vez? _____ anos
Quantas vezes você ficou grávida? _____ vezes
Quantos filhos você teve? _____ filho(s)

Você já teve um aborto?

0 () Não

1 () Sim **Quantos abortos você teve? _____ abortos****No último ano, você usou algum método contraceptivo?**0 () Não 1 () Sim **(Pula para o Quadro 2)****Qual o motivo para não ter utilizado método contraceptivo no último ano?**

0 () Não tinha relações sexuais

1 () Relações sexuais pouco frequentes

2 () Estava grávida

3 () Queria engravidar

4 () Não queria usar nenhum método contraceptivo

5 () Companheiro se opôs

6 () Motivos religiosos

7 () Não conhece nenhum método

8 () Não sabe onde obter

9 () Problemas de saúde/efeitos colaterais

10 () Medo de efeitos colaterais

11 () Dificuldades de acesso _____

12 () Custo

13 () Inconveniente de usar

14 () Interfere nas funções normais do corpo

15 () Marido/companheiro infértil

16 () Apresenta (você ou parceiro) alguma condição que impossibilita a gravidez (histerectomia, azoospermia, entre outras)

17 () Tem relações sexuais apenas com mulheres

18 () Outros: _____ (especifique)

19 () Não consegue se organizar para isso

(Após esta pergunta, a pesquisa se encerra. Agradecemos a sua participação).**Se sim:****Neste momento, perguntas a fim de avaliar a ocorrência de descontinuidades contraceptivas no último ano foram apresentadas.**

Você usou pílula do dia seguinte nos últimos 12 meses?

1 () Não

2 () Sim

Qual(ais) mês(es) você utilizou a pílula do dia seguinte?

1

mês

atrás

2

meses

atrás

3

meses

atrás

4

meses

atrás

5

meses

atrás

6

meses

atrás

7

meses

atrás

8

meses

atrás

9

meses

atrás

10

meses

atrás

11

meses

atrás

12

meses

atrás

Pílula do dia seguinte

Nos últimos 12 meses, você trocou de método?
 1 () Não 2 () Sim

Em qual mês você trocou de método e por que?
 Razões:
 1: eEfeitos colaterais
 2: Queria um método mais eficaz
 3: Desaprovação do parceiro
 4: Troquei de parceiro
 5: Muito caro ou dificuldade na obtenção
 6: Outro motivo

Em qual(ais) mês(es) você trocou de método e por que?	1	2	3	4	5	6
1 mês atrás	<input type="radio"/>					
2 meses atrás	<input type="radio"/>					
3 meses atrás	<input type="radio"/>					
4 meses atrás	<input type="radio"/>					
5 meses atrás	<input type="radio"/>					
6 meses atrás	<input type="radio"/>					
7 meses atrás	<input type="radio"/>					
8 meses atrás	<input type="radio"/>					
9 meses atrás	<input type="radio"/>					
10 meses atrás	<input type="radio"/>					
11 meses atrás	<input type="radio"/>					
12 meses atrás	<input type="radio"/>					

Você engravidou nos últimos 12 meses?

- 1 () Não
- 2 () Sim

Em qual mês você engravidou? _____

Nos últimos 12 meses, que tipo de relacionamento (MAIS FREQUENTE) você teve em cada mês?	Estável, como com namorado(a)	Eventual, como com ficante	Unida ou morava com o parceiro(a)	Nenhum
1 mês atrás	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
2 meses atrás	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
3 meses atrás	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
4 meses atrás	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
5 meses atrás	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
6 meses atrás	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
7 meses atrás	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
8 meses atrás	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
9 meses atrás	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
10 meses atrás	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
11 meses atrás	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
12 meses atrás	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

AGORA IREMOS FAZER ALGUMAS PERGUNTAS SOBRE O USO DA PÍLULA DO DIA SEGUINTE!!!

Você conhece alguém que já usou a pílula do dia seguinte?

0 () Não

1 () Sim

Você já usou alguma vez a anticoncepção de emergência (pílula do dia seguinte)?

0 () Não **(Se você respondeu não, a pesquisa se encerra. Agradecemos a sua participação)**

1 () Sim

SE SIM – Qual foi a razão que levou você ao uso da pílula do dia seguinte da última vez?

1 () O método que usava falhou

2 () Houve esquecimento de usar algum método

3 () Não quis usar nenhum método

4 () Foi vítima de violência sexual

5 () Por pura insegurança

6 () Outros: _____

SE SIM – Nessa última vez que usou a pílula do dia seguinte, o relacionamento era:

1 () fixo(a)/regular, como com namorado(a)

2 () eventual, como com ficante

SE SIM – Quem tomou a iniciativa do uso da pílula do dia seguinte?

1 () Você

2 () Parceiro

3 () Ambos

SE SIM – A pílula do dia seguinte foi adquirida com receita médica?

0 () Não

1 () Sim

SE SIM – Onde você adquiriu a pílula do dia seguinte?

1 () Farmácia

2 () No posto de saúde

3 () No médico particular ou do convênio

4 () Com amigos ou conhecidos

5 () Namorado que trouxe

6 () Com parente

7 () Outros? _____

Na última vez, a pílula do dia seguinte foi tomada quanto tempo após a relação sexual desprotegida?

1 ____ horas ou ____ dias

2 () Não sei, não me lembro

Quantas vezes você usou a pílula do dia seguinte no último ano?

1 _____ vezes

2 () Não sei, não me lembro

Você estava usando algum método contraceptivo quando você usou a pílula do dia seguinte na última vez?

0 () Não

1 () Sim

Qual método contraceptivo você estava usando?

1 () Pílula oral

2 () Camisinha masculina

3 () Pílula oral e camisinha masculina

4 () Injetável

5 () Coito interrompido/Tabelinha

6 () DIU

7 () Other

AGORA IREMOS PERGUNTAR SOBRE O USO DE MÉTODOS CONTRACEPTIVOS UM MÊS ANTES E UM MÊS APÓS O USO DA PÍLULA DO DIA SEGUINTE!!!

→ 30 DAYS BEFORE THE USE OF EMERGENCY CONTRACEPTION



Você usou algum método anticoncepcional 30 dias ANTES do us da pílula do dia seguinte?

- 0 () Não
- 1 () Sim

Calendário							
Meses	Seg	Ter	Qua	Qui	Sex	Sab	Dom
novembro	28	29	30	31	01	02	03
	04	05	06	07	08	09	10
	11	12	13	14	15	16	17
	18	19	20	21	22	23	24
dezembro	25	26	27	28	29	30	01
	02	03	04	05	06	07	08
	09	10	11	12	13	14	15
	16	17		19	20	21	22
	23	24		26	27	28	29

30 dias ANTES

Uso da pílula do dia seguinte

Qual método contraceptivo você mais usou na maioria dos dias?

- 1 () Pílula oral
- 2 () Camisinha masculina
- 3 () Pílula oral e camisinha masculina
- 4 () Injetável
- 5 () Coito interrompido/Tabelinha
- 6 () DIU
- 7 () Other


Nesses 30 dias ANTES de você usar a pílula do dia seguinte, você usou esse método continuamente (todos os dias sem exceção)?

0 () Não 1 () Sim

Nesses 30 dias ANTES de você usar a pílula do dia seguinte, quais dias você usou o método contraceptivo?

30 DIAS ANTES	Day	Não usou contracepção	Usou contracepção
	30		
	29		
	28		
	27		
	26		
	25		
	24		
	23		
	22		
	21		
	20		
	19		
	18		
	17		
	16		
	15		
	14		
	13		
	12		
	11		
	10		
	9		
	8		
	7		
	6		
	5		
	4		
	3		
	2		
	1		
USO DA PÍLULA DO DIA SEGUINTE			

→ 30 DIAS APÓS O USO DA PÍLULA DO DIA SEGUINTE



Você usou algum método contraceptivo 30 dias APÓS você usar a pílula do dia seguinte?

- 0 () Não
- 1 () Sim

A calendar titled "Calendário" showing the months of November and December. The days of the week are listed as Seg, Ter, Qua, Qui, Sex, Sab, Dom. The date 21st of November is highlighted in yellow, and a pink arrow points upwards from the text "Uso da pílula do dia seguinte" below the calendar to this date. A thought bubble next to a woman's head contains the text "30 dias DEPOIS".

Meses	Seg	Ter	Qua	Qui	Sex	Sab	Dom
novembro	28	29	30	31	01	02	03
	04	05	06	07	08	09	10
	11	12	13	14	15	16	17
	18	19	20	21	22	23	24
dezembro	25	26	27	28	29	30	01
	02	03	04	05	06	07	08
	09	10	11	12	13	14	15
	16	17	18	19	20	21	22
	23	24	25	26	27	28	29

Qual método contraceptivo você mais usou na maioria dos dias?

- 1 () Pílula oral
- 2 () Camisinha masculina
- 3 () Pílula oral e camisinha masculina
- 4 () Injetável
- 5 () Coito interrompido/Tabelinha
- 6 () DIU
- 7 () Other

Você iniciou imediatamente o uso deste método, APÓS usar a pílula do dia seguinte?

0 () Não

1 () Sim

Quantos dias DEPOIS de usar a pílula do dia seguinte, você iniciou o uso do método?

1 () < 10 dias

2 () 10 – 15 dias

3 () ≥ 15 dias

Você usou este método continuamente por quanto tempo?

1 () < 1 mês

2 () 1 mês

3 () 1 -| 6 meses

4 () 6 -| 12 meses

5 () 1 -| 2 anos

6 () ≥ 2 anos

APPENDIX D – INDEPENDENCE MODEL CRITERION (QIC)

a) Discontinuation for any reason

- Oral pill

Correlation matrix	Working correlation structure											
	Exchangeable											
	1.00											
0.22	1.00											
0.22	0.22	1.00										
0.22	0.22	0.22	1.00									
0.22	0.22	0.22	0.22	1.00								
0.22	0.22	0.22	0.22	0.22	1.00							
0.22	0.22	0.22	0.22	0.22	0.22	1.00						
0.22	0.22	0.22	0.22	0.22	0.22	0.22	1.00					
0.22	0.22	0.22	0.22	0.22	0.22	0.22	0.22	1.00				
0.22	0.22	0.22	0.22	0.22	0.22	0.22	0.22	0.22	1.00			
0.22	0.22	0.22	0.22	0.22	0.22	0.22	0.22	0.22	0.22	1.00		
0.22	0.22	0.22	0.22	0.22	0.22	0.22	0.22	0.22	0.22	0.22	1.00	
0.22	0.22	0.22	0.22	0.22	0.22	0.22	0.22	0.22	0.22	0.22	0.22	1.00
QIC	7645.70											

Correlation matrix	Working correlation structure											
	Unstructured											
	1.00											
0.02	1.00											
0.01	0.01	1.00										
0.01	0.06	0.02	1.00									
0.06	0.03	0.03	0.01	1.00								
0.00	0.05	0.00	0.05	0.00	1.00							
0.00	0.02	0.01	0.00	0.00	0.00	1.00						
0.02	-0.01	-0.01	0.04	0.02	-0.01	-0.01	1.00					
0.00	0.11	0.00	0.00	0.01	0.01	0.04	0.00	1.00				
0.02	0.05	-0.01	0.05	0.02	0.07	0.00	-0.01	-0.01	1.00			
0.06	0.04	-0.02	0.05	0.01	0.05	0.05	0.11	0.08	-0.02	1.00		
QIC	7655.56											

Correlation matrix	Working correlation structure										
	Autoregressive										
	1.00										
0.00	1.00										
0.00	0.00	1.00									
0.00	0.00	0.00	1.00								
0.00	0.00	0.00	0.00	1.00							
0.00	0.00	0.00	0.00	0.00	1.00						
0.00	0.00	0.00	0.00	0.00	0.00	1.00					
0.00	0.00	0.00	0.00	0.00	0.00	0.00	1.00				
0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	1.00			
0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	1.00		
0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	1.00	
0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	1.00
QIC	8053.35										

- Male Condom

Correlation matrix	Working correlation structure										
	Exchangeable										
	1.00										
0.11	1.00										
0.11	0.11	1.00									
0.11	0.11	0.11	1.00								
0.11	0.11	0.11	0.11	1.00							
0.11	0.11	0.11	0.11	0.11	1.00						
0.11	0.11	0.11	0.11	0.11	0.11	1.00					
0.11	0.11	0.11	0.11	0.11	0.11	0.11	1.00				
0.11	0.11	0.11	0.11	0.11	0.11	0.11	0.11	1.00			
0.11	0.11	0.11	0.11	0.11	0.11	0.11	0.11	0.11	1.00		
0.11	0.11	0.11	0.11	0.11	0.11	0.11	0.11	0.11	0.11	1.00	
QIC	9099.91										

Correlation matrix	Working correlation structure																		
	Unstructured																		
	1.00	0.11	1.00																
	0.10	0.05	1.00																
		0.12	0.11	0.11	1.00														
			0.23	0.13	0.13	0.08	1.00												
				0.06	0.16	0.10	0.07	0.07	1.00										
					0.15	0.22	0.16	0.11	0.16	0.09	1.00								
						0.07	0.08	0.07	0.21	0.06	0.11	0.07	1.00						
							0.15	0.11	0.15	0.02	0.17	0.03	0.08	0.03	1.00				
								0.00	0.13	0.12	0.03	0.13	0.13	0.10	0.07	0.03	1.00		
									0.10	0.11	0.16	0.01	0.14	0.22	0.09	0.18	0.13	0.07	1.00
QIC	9094.83																		

Correlation matrix	Working correlation structure													
	Autoregressive													
	1.00	-0.01	1.00											
	0.00	-0.01	1.00											
		0.00	0.00	-0.01	1.00									
			0.00	0.00	0.00	-0.01	1.00							
				0.00	0.00	0.00	0.00	-0.01	1.00					
					0.00	0.00	0.00	0.00	-0.01	1.00				
						0.00	0.00	0.00	0.00	-0.01	1.00			
							0.00	0.00	0.00	0.00	-0.01	1.00		
								0.00	0.00	0.00	0.00	-0.01	1.00	
									0.00	0.00	0.00	0.00	-0.01	1.00
QIC	9690.39													

b) Discontinuation for method-reason reasons

- Oral pill

Correlation matrix	Working correlation structure										
	Exchangeable										
	1.00										
	0.02	1.00									
	0.02	0.02	1.00								
	0.02	0.02	0.02	1.00							
	0.02	0.02	0.02	0.02	1.00						
	0.02	0.02	0.02	0.02	0.02	1.00					
	0.02	0.02	0.02	0.02	0.02	0.02	1.00				
	0.02	0.02	0.02	0.02	0.02	0.02	0.02	1.00			
	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	1.00		
0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	1.00		
0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	1.00	
0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	1.00
QIC	6944.11										

Correlation matrix	Working correlation structure										
	Unstructured										
	1.00										
	0.02	1.00									
	0.01	0.01	1.00								
	0.01	0.06	0.01	1.00							
	0.04	0.03	0.04	0.01	1.00						
	0.01	0.05	0.00	0.05	0.01	1.00					
	0.00	0.00	0.00	0.00	0.00	0.01	1.00				
	0.03	-0.01	-0.02	0.05	0.02	-0.01	-0.01	1.00			
	-0.01	0.12	0.00	0.00	0.02	-0.01	0.04	0.00	1.00		
0.00	0.05	0.00	0.05	0.00	0.07	0.00	0.00	0.00	1.00		
0.06	0.02	-0.02	0.07	0.02	0.07	0.06	0.13	0.08	-0.01	1.00	
QIC	6957.77										

Correlation matrix	Working correlation structure										
	Autoregressive										
	1.00										
	0.00	1.00									
	0.00	0.00	1.00								
	0.00	0.00	0.00	1.00							
	0.00	0.00	0.00	0.00	1.00						
	0.00	0.00	0.00	0.00	0.00	1.00					
	0.00	0.00	0.00	0.00	0.00	0.00	1.00				
	0.00	0.00	0.00	0.00	0.00	0.00	0.00	1.00			
0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	1.00			
0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	1.00		
0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	1.00	
QIC	7327.24										

- Male Condom

Correlation matrix	Working correlation structure										
	Exchangeable										
	1.00										
	0.09	1.00									
	0.09	0.09	1.00								
	0.09	0.09	0.09	1.00							
	0.09	0.09	0.09	0.09	1.00						
	0.09	0.09	0.09	0.09	0.09	1.00					
	0.09	0.09	0.09	0.09	0.09	0.09	1.00				
	0.09	0.09	0.09	0.09	0.09	0.09	0.09	1.00			
0.09	0.09	0.09	0.09	0.09	0.09	0.09	0.09	1.00			
0.09	0.09	0.09	0.09	0.09	0.09	0.09	0.09	0.09	1.00		
0.09	0.09	0.09	0.09	0.09	0.09	0.09	0.09	0.09	0.09	1.00	
QIC	8185.12										

Correlation matrix	Working correlation structure										
	Unstructured										
	1.00										
0.08	1.00										
0.09	0.04	1.00									
0.09	0.08	0.09	1.00								
0.21	0.10	0.11	0.06	1.00							
0.03	0.16	0.08	0.06	0.05	1.00						
0.14	0.19	0.14	0.10	0.14	0.07	1.00					
0.04	0.05	0.05	0.19	0.04	0.08	0.05	1.00				
0.13	0.07	0.12	0.01	0.17	0.02	0.08	0.02	1.00			
0.00	0.15	0.04	0.05	0.14	0.13	0.07	0.03	0.02	1.00		
0.09	0.12	0.16	0.02	0.10	0.18	0.08	0.11	0.12	0.05	1.00	
QIC	8191.82										

Correlation matrix	Working correlation structure										
	Autoregressive										
	1.00										
-0.01	1.00										
0.00	-0.01	1.00									
0.00	0.00	-0.01	1.00								
0.00	0.00	0.00	-0.01	1.00							
0.00	0.00	0.00	0.00	-0.01	1.00						
0.00	0.00	0.00	0.00	0.00	-0.01	1.00					
0.00	0.00	0.00	0.00	0.00	0.00	-0.01	1.00				
0.00	0.00	0.00	0.00	0.00	0.00	0.00	-0.01	1.00			
0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	-0.01	1.00		
0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	-0.01	1.00	
0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	-0.01	1.00
QIC	9113.63										

c) Abandonment or switching to less effective method

- Oral pill

Correlation matrix	Working correlation structure										
	Exchangeable										
	1.00										
0.00	1.00										
0.00	0.00	1.00									
0.00	0.00	0.00	1.00								
0.00	0.00	0.00	0.00	1.00							
0.00	0.00	0.00	0.00	0.00	1.00						
0.00	0.00	0.00	0.00	0.00	0.00	1.00					
0.00	0.00	0.00	0.00	0.00	0.00	0.00	1.00				
0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	1.00			
0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	1.00		
0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	1.00	
QIC	5504.50										

Correlation matrix	Working correlation structure										
	Unstructured										
	1.00										
-0.01	1.00										
-0.02	-0.03	1.00									
0.02	0.04	-0.02	1.00								
0.05	0.05	0.01	-0.01	1.00							
0.00	0.02	-0.03	0.03	-0.02	1.00						
-0.02	-0.02	-0.03	0.00	-0.02	-0.03	1.00					
0.01	0.02	0.00	0.09	0.01	0.03	-0.04	1.00				
-0.03	0.02	-0.04	-0.03	0.00	-0.03	0.02	-0.03	1.00			
0.00	0.02	-0.04	0.03	0.00	0.09	0.04	0.02	-0.03	1.00		
0.02	0.01	0.00	0.01	0.02	0.02	0.01	0.01	0.01	0.01	1.00	
QIC	5505.24										

Correlation matrix	Working correlation structure																				
	Autoregressive																				
	1.00	-0.02	1.00	0.00	-0.02	1.00	0.00	0.00	-0.02	1.00	0.00	0.00	0.00	0.00	-0.02	1.00	0.00	0.00	0.00	0.00	-0.02
QIC	5529.75																				

- Male Condom

Correlation matrix	Working correlation structure																										
	Exchangeable																										
	1.00	-0.01	1.00	-0.01	-0.01	1.00	-0.01	-0.01	-0.01	1.00	-0.01	-0.01	-0.01	-0.01	1.00	-0.01	-0.01	-0.01	-0.01	-0.01	1.00	-0.01	-0.01	-0.01	-0.01	-0.01	-0.01
QIC	5684.09																										


APPENDIX E – REASONS FOR CONTRACEPTIVE DISCONTINUATION

Table 19 – Reasons for discontinuation the contraceptive method by method type. São Paulo, Brazil – 2015

Variables	Oral pill users		Male condom users		Other methods users		p*
	N	%	N	%	N	%	
Reasons for Discontinuation for any reasons (n=567)							
0.011							
Side effects	196	68.8	02	0.9	35	62.5	
Wanted a more effective method	18	6.3	146	64.6	07	12.5	
Criticism of the partner	04	1.4	39	17.3	02	3.6	
Too expensive or difficult to obtain	04	1.4	01	0.4	02	3.6	
Another reason	63	22.1	38	16.8	10	17.9	
Discontinuation for method-related reasons (n=464)							
0.123							
Side effects	151	65.9	02	1.1	28	60.9	
Wanted a more effective method	17	7.4	117	61.9	07	15.2	
Criticism of the partner	03	1.3	34	18.0	01	2.2	
Too expensive or difficult to obtain	02	0.9	01	0.5	01	2.2	
Another reason	56	24.5	35	18.5	09	19.6	
Abandoned or switched to less effective method (n=305)							
0.829							
Side effects	146	67.6	01	1.7	21	67.7	
Wanted a more effective method	14	6.5	03	5.2	02	6.5	
Criticism of the partner	03	1.4	43	74.1	01	3.2	
Too expensive or difficult to obtain	02	0.9	0	0	01	3.2	
Another reason	51	23.6	11	19.0	06	19.4	

ANNEX

ANNEX – APPROVAL OF THE RESEARCH ETHICS COMMITTEE



PARECER CONSUBSTANCIADO DO CEP

DADOS DO PROJETO DE PESQUISA

Título da Pesquisa: DESCONTINUIDADES CONTRACEPTIVAS E SUA RELAÇÃO COM O USO DA ANTICONCEPÇÃO DE EMERGÊNCIA ENTRE UNIVERSITÁRIAS

Pesquisador: Christiane Borges do Nascimento Chofakian

Área Temática:

Versão: 1

CAAE: 39142514.4.0000.5392

Instituição Proponente: Escola de Enfermagem da Universidade de São Paulo - EEUSP

Patrocinador Principal: Financiamento Próprio

DADOS DO PARECER

Número do Parecer: 912.504

Data da Relatoria: 08/12/2014

Apresentação do Projeto:

As práticas contraceptivas de jovens universitárias apresentam dinâmica própria, na qual o alto uso de métodos contraceptivos coexiste com as descontinuidades e as irregularidades de seu uso; situações que constituem uma das indicações de uso da anticoncepção de emergência. No Brasil, há escassez de estudos que abordem as descontinuidades contraceptivas e sua relação com o uso da anticoncepção de emergência. Com a finalidade de compreender melhor a dinâmica contraceptiva, para além da prevalência e dos tipos de métodos utilizados, a proposta deste estudo é estimar a probabilidade de ocorrência de descontinuidades contraceptivas e analisar as práticas contraceptivas antes e após o uso da anticoncepção de emergência entre jovens universitárias. Para isto, será conduzido um estudo quantitativo, do tipo longitudinal, com jovens universitárias de todos os campi da Universidade de São Paulo. Por se tratar de questões de cunho íntimo, será usado um questionário de autopreenchimento respondido online. O questionário é composto por questões sociodemográficas; questões sobre o uso de métodos contraceptivos – para analisar as descontinuidades contraceptivas – nos 12 meses anteriores à pesquisa; e questões que versam sobre o uso de métodos contraceptivos um mês antes e um mês após o uso da anticoncepção de emergência, no caso de relato de uso anterior. A análise da probabilidade de ocorrência de descontinuidades contraceptivas por tipo de método e da prática contraceptiva antes

Endereço: Av. Dr Enéas de Carvalho Aguiar, 419
Bairro: Cerqueira Cesar **CEP:** 05.403-000
UF: SP **Município:** SAO PAULO
Telefone: (11)3061-7548 **Fax:** (11)3061-7548 **E-mail:** edipesq@usp.br